

LIFTINGS FOR ULTRA-MODULATION SPACES, AND ONE-PARAMETER GROUPS OF GEVREY TYPE PSEUDO-DIFFERENTIAL OPERATORS

AHMED ABDELJAWAD, SANDRO CORIASCO, AND JOACHIM TOFT

ABSTRACT. We deduce one-parameter group properties for pseudo-differential operators $\text{Op}(a)$, where a belongs to the class $\Gamma_*^{(\omega_0)}$ of certain Gevrey symbols. We use this to show that there are pseudo-differential operators $\text{Op}(a)$ and $\text{Op}(b)$ which are inverses to each others, where $a \in \Gamma_*^{(\omega_0)}$ and $b \in \Gamma_*^{(1/\omega_0)}$.

We apply these results to deduce lifting property for modulation spaces and construct explicit isomorphisms between them. For each weight functions ω, ω_0 moderated by GRS submultiplicative weights, we prove that the Toeplitz operator (or localization operator) $\text{Tp}(\omega_0)$ is an isomorphism from $M_{(\omega)}^{p,q}$ onto $M_{(\omega/\omega_0)}^{p,q}$ for every $p, q \in (0, \infty]$.

0. INTRODUCTION

The topological vector spaces V_1 and V_2 is said to possess lifting property if there exists a "convenient" homeomorphisms (a lifting) between them. For example, for any weight ω on $\mathbf{R}^d$, $p \in (0, \infty]$ and $s \in \mathbf{R}$ the convenient mappings $f \mapsto \omega \cdot f$ and $f \mapsto (1 - \Delta)^{s/2} f$ are homeomorphic from the (weighted) Lebesgue space $L_{(\omega)}^p$ and the Sobolev space H_s^p , respectively, into $L^p = H_0^p$, with inverses $f \mapsto \omega^{-1} \cdot f$ and $f \mapsto (1 - \Delta)^{-s/2} f$, respectively. (Cf. [30] and Section 1 for notations.) Hence, these spaces possess lifting properties.

It is often uncomplicated to deduce lifting properties between (quasi-)Banach spaces of functions and distributions, if the definition of their norms only differs by a multiplicative weight on the involved distributions, or on their Fourier transforms, which is the case in the previous homeomorphisms. Here recall that multiplications on the Fourier transform side are linked to questions on differentiation of the involved elements. A more complicated situation appear when there are some kind of interactions between multiplication and differentiation in the definition of the involved vector spaces.

An example where such interactions occur concerns the extended family of Sobolev spaces, introduced by Bony and Chemin in [3] (see also [34]). More precisely, let ω, ω_0 be suitable weight functions and g a suitable Riemannian metric, which are defined on the phase space $W \simeq T^*\mathbf{R}^d \simeq \mathbf{R}^{2d}$. Then Bony and Chemin introduced in [3] the generalised Sobolev space $H(\omega, g)$ which fits the Hörmander-Weyl calculus well in the sense that $H(1, g) = L^2$, and if a belongs to the Hörmander class $S(\omega_0, g)$, then Weyl operator $\text{Op}^w(a)$ with symbol a is continuous from $H(\omega_0 \omega, g)$ to $H(\omega, g)$. Moreover, they deduced group algebras, from which it follows that to each such weight ω_0 , there exist symbols a and b such that

$$\text{Op}^w(a) \circ \text{Op}^w(b) = \text{Op}^w(b) \circ \text{Op}^w(a) = I, \quad a \in S(\omega_0, g), \quad b \in S(1/\omega_0, g). \quad (0.1)$$

2010 *Mathematics Subject Classification.* Primary: 35S05, 47D06, 46B03, 42B35, 47B35, 46F05; Secondary: 46G15, 47L80.

Key words and phrases. Gelfand-Shilov spaces, Banach Function spaces, ultra-distributions, confinement of symbols, comparable symbols.

Here I is the identity operator on $\mathcal{S}'$. In particular, by the continuity properties of $\text{Op}^w(a)$ it follows that $H(\omega_0\omega, g)$ and $H(\omega, g)$ possess lifting properties with the homeomorphism $\text{Op}^w(a)$, and with $\text{Op}^w(b)$ as its inverse.

The existence of a and b in (0.1) is a consequence of solution properties of the evolution equation

$$(\partial_t a)(t, \cdot) = (b + \log \vartheta) \# a(t, \cdot), \quad a(0, \cdot) = a_0 \in S(\omega, g), \quad \vartheta \in S(\vartheta, g), \quad (0.2)$$

which involve the Weyl product $\#$ and a fixed element $b \in S(1, g)$. It is proved that (0.2) has a unique solution $a(t, \cdot)$ which belongs to $S(\omega\vartheta^t, g)$ (cf. [3, Theorem 6.4] or [34, Theorem 2.6.15]). The existence of a and b in (0.1) will follow by choosing $\omega = a_0 = 1$, $t = 1$ and $\vartheta = \omega_0$.

If g is the constant euclidean metric on the phase space $\mathbf{R}^{2d}$, then $S(\omega_0, g)$ equals $S^{(\omega_0)}(\mathbf{R}^{2d})$, the set of all smooth symbols a which satisfies $|\partial^\alpha a| \lesssim \omega_0$. We notice that also for such simple choices of g , (0.1) above leads to lifting properties that are not trivial. In fact, let ω and ω_0 be polynomially moderate weight on the phase space, and let $\mathcal{B}$ be a suitable translation invariant BF-space. Then it is observed in [26] that the continuity results for pseudo-differential operators on modulation spaces in [48, 50] imply that $\text{Op}^w(a)$ in (0.1) is continuous and bijective from $M(\omega_0\omega, \mathcal{B})$ to $M(\omega, \mathcal{B})$ with continuous inverse $\text{Op}^w(b)$. In particular, by choosing $\mathcal{B}$ to be the mixed norm space $L^{p,q}(\mathbf{R}^{2d})$ of Lebesgue type, then $M(\omega, \mathcal{B})$ is equal to the classical modulation space $M_{(\omega)}^{p,q}$, introduced by Feichtinger in [12]. Consequently, $\text{Op}^w(a)$ above lifts $M_{(\omega_0\omega)}^{p,q}$ into $M_{(\omega)}^{p,q}$.

An important class of operators in quantum mechanics and time-frequency analysis concerns Toeplitz, or localisation operators. The main issue in [26, 27] is to show that the Toeplitz operator $\text{Tp}(\omega_0)$ lifts $M_{(\omega_0\omega)}^{p,q}$ into $M_{(\omega)}^{p,q}$ for suitable ω_0 . The assumptions on ω_0 in [26] is that it should be polynomially moderate and satisfies $\omega_0 \in S^{(\omega_0)}$. In [27], it is only assumed that ω_0 is moderated by a GRS weight, but instead it is here required that ω_0 is *radial in each phase shift*, i.e. ω_0 should satisfy

$$\omega_0(x_1, \dots, x_d, \xi_1, \dots, \xi_d) = \vartheta(r_1, \dots, r_d), \quad r_j = |(x_j, \xi_j)|,$$

for some weight ϑ .

The approaches in [26, 27] are also different. In [27], the lifting properties for $\text{Tp}(\omega_0)$ is reached by using the links between modulation spaces and Bargmann-Fock spaces in combination of suitable estimates for a sort of generalised gamma-functions. The approach in [26] relies on corresponding lifting properties for pseudo-differential operators, as follows:

- (1) $\text{Tp}(\omega_0) = \text{Op}^w(c)$ for some $c \in S^{(\omega_0)}$;
- (2) by the definitions, it follows by straightforward computations that if $\vartheta = \omega_0^{\frac{1}{2}}$, then $\text{Tp}(\omega_0)$ is a homeomorphism from $M_{(\vartheta)}^{p,q}$ to $M_{(\vartheta)}^{p,q}$;
- (3) combining (0.1) with Wiener's lemma for $(S^{(1)}, \#)$ ensures that the inverse of $\text{Tp}(\omega_0)$ in (2) is a pseudo-differential operator $\text{Op}^w(b)$ with the symbol b in $S^{(1/\omega_0)}$;
- (4) by (1), (3) and duality,

$$T_1 \equiv \text{Op}^w(b) \circ \text{Tp}(\omega_0) \quad \text{and} \quad T_2 \equiv \text{Tp}(\omega_0) \circ \text{Op}^w(b)$$

are both the identity operator on $\mathcal{S}'(\mathbf{R}^d)$, since T_1 is the identity operator on $M_{(\vartheta)}^{p,q}$, T_2 is the identity operator on $M_{(1/\vartheta)}^{p,q}$, and $\mathcal{S} \subseteq M_{(\vartheta)}^{p,q} \cap M_{(1/\vartheta)}^{p,q}$.

- (5) by (4), $T_1 = T_2 = \text{Op}^w(1)$ is the identity operator on each $M_{(\omega)}^{p,q}$. Since

$$\text{Tp}(\omega_0) = \text{Op}^w(c) : M_{(\omega_0\omega)}^{p,q} \rightarrow M_{(\omega)}^{p,q} \quad \text{and} \quad \text{Op}^w(b) : M_{(\omega)}^{p,q} \rightarrow M_{(\omega_0\omega)}^{p,q}$$

are continuous (cf. [48, 50]) and inverses to each other, it follows that they are homeomorphisms.

In the first part of the paper we deduce an analog of (0.1) for the Gevrey type symbol classes $\Gamma_s^{(\omega_0)}$ and $\Gamma_{0,s}^{(\omega_0)}$ of orders $s \geq 1$, the set of all $a \in C^\infty$ such that

$$|\partial^\alpha a(X)| \lesssim h^{|\alpha|} \alpha!^s \omega(X)$$

for some $h > 0$, respectively for every $h > 0$, considered in [4]. That is, in Section 3 we show that there exist symbols a and b such that

$$\text{Op}^w(a) \circ \text{Op}^w(b) = \text{Op}^w(b) \circ \text{Op}^w(a) = I, \quad a \in \Gamma_s^{(\omega_0)}, \quad b \in \Gamma_s^{(1/\omega_0)}, \quad (0.3)$$

and similarly when $\Gamma_s^{(\omega_0)}$ and $\Gamma_s^{(1/\omega_0)}$ are replaced by $\Gamma_{0,s}^{(\omega_0)}$ and $\Gamma_{0,s}^{(1/\omega_0)}$, respectively.

For general ω_0 it is clear that $\Gamma_{0,s}^{(\omega_0)} \subseteq \Gamma_s^{(\omega_0)} \subseteq S^{(\omega_0)}$. On the other hand, for the weights ω_1, ω_2 and ω_3 in $\Gamma_{0,s}^{(\omega_1)}, \Gamma_s^{(\omega_2)}$ and $S^{(\omega_3)}$ we always assume that they belong to $\mathcal{P}_{E,s}(\mathbf{R}^{2d})$, $\mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$ and $\mathcal{P}(\mathbf{R}^{2d})$, respectively. That is, they should satisfy

$$\omega_1(X+Y) \lesssim \omega_1(X) e^{r_1|Y|^{\frac{1}{s}}}, \quad \omega_2(X+Y) \lesssim \omega_2(X) e^{r_2|Y|^{\frac{1}{s}}},$$

$$\text{and } \omega_3(X+Y) \lesssim \omega_3(X)(1+|Y|)^N,$$

for some $r_1 > 0$ and $N > 0$, and every $r_2 > 0$. Since it is clear that $\mathcal{P} \subseteq \mathcal{P}_{E,s}^0 \subseteq \mathcal{P}_{E,s}$, it follows by straightforward computations that there are admissible $a_1 \in \Gamma_{0,s}^{(\omega_1)}$ which are not contained in any admissible $\Gamma_s^{(\omega_0)}$ and $S^{(\omega_0)}$, and admissible $a_2 \in \Gamma_s^{(\omega_2)}$ which are not contained in any $S^{(\omega_0)}$.

As in [3], (0.3) is obtained by proving that the evolution equation

$$(\partial_t a)(t, \cdot) = (b + \log \vartheta) \# a(t, \cdot), \quad a(0, \cdot) = a_0 \in \Gamma_s^{(\omega)}, \quad \vartheta \in \Gamma_s^{(\vartheta)}, \quad (0.4)$$

analogous to (0.2), has a unique solution $a(t, \cdot)$ which belongs to $\Gamma_s^{(\omega \vartheta^t)}$ (and similarly when the $\Gamma_s^{(\omega)}$ -spaces are replaced by corresponding $\Gamma_{0,s}^{(\omega)}$ -spaces), given in Section 3.

In Sections 4 and 5 we use the framework in [26] in combination with (0.3) to extend the lifting properties in [26] in such ways that the involved weights are allowed to belong to $\mathcal{P}_{E,s}^0$ or in $\mathcal{P}_{E,s}$ instead of the smaller set $\mathcal{P}$ which is the assumption in [26].

Our main result, which is similar to [26, Theorem 0.1], can be stated as follows.

Theorem 0.1. *Let $s \geq 1$, $\omega, \omega_0 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$, $p, q \in (0, \infty]$ and let $\phi \in \mathcal{S}_s(\mathbf{R}^d)$. Then the Toeplitz operator $\text{Tp}_\phi(\omega_0)$ is an isomorphism from $M_{(\omega)}^{p,q}(\mathbf{R}^{2d})$ onto $M_{(\omega/\omega_0)}^{p,q}(\mathbf{R}^{2d})$.*

We note that, in contrast to [26, 27], our lifting properties also hold for modulation spaces which may fail to be Banach spaces, since p and q in Theorem 0.1 are allowed to be smaller than 1.

We will establish several related result. Firstly, the window function may be chosen in certain modulation spaces that are much larger than the Gelfand-Shilov space $\mathcal{S}_s$. Secondly, the theorem holds for a more general family of modulation spaces that includes the classical modulation spaces. Finally, we also establish isomorphisms given by pseudo-differential operators rather than Toeplitz operators.

In contrast to [27], we do not impose in Theorem 0.1 and in its related results in Section 5 that ω_0 should be radial in each phase shift (cf. e. g. [27, Theorem 4.3]). Summing up, our lifting results in Section 5 extend the lifting results in [26, 27].

The paper is organised as follows. In Section 1 we introduce some notation, and discuss modulation spaces and Gelfand-Shilov spaces of functions and distributions,

and pseudo-differential calculus. In Section 2 we introduce and discuss basic properties for confinements of symbols in $\Gamma_s^{(\omega_0)}$ and in $\Gamma_{0,s}^{(\omega_0)}$. These considerations are related to the discussions in [3, 34], but here adapted to symbols that possess Gevrey regularity, e. g. when the symbols belong to $\Gamma_s^{(\omega_0)}$ or $\Gamma_{0,s}^{(\omega_0)}$.

In contrast to the classical Hörmander symbol classes $S_{1,0}^r$ and the SG-classes $SG_{1,1}^{m,\mu}$, techniques on asymptotic expansions are absent for symbols in the classes $\Gamma_s^{(\omega_0)}$ or in $\Gamma_{0,s}^{(\omega_0)}$, and might be absent for symbols in the general Hörmander class $S(m, g)$. The approach with confinements is, roughly speaking, a sort of stand-in of these absent asymptotic expansion techniques.

In Section 3 we show that the (0.4) has a unique solution with the requested properties, which leads to (0.3). In Sections 4 and 5 we use the results from Section 3 to deduce lifting properties for modulation spaces under pseudo-differential operators and Toeplitz operators with symbols in $\Gamma_s^{(\omega_0)}$ or in $\Gamma_{0,s}^{(\omega_0)}$.

1. PRELIMINARIES

In this section we recall some basic facts on modulation spaces, Gelfand-Shilov spaces of functions and distributions and pseudo-differential calculus (cf. [11–16, 18, 21, 25, 30, 31, 34, 38, 42, 45, 47–51]).

1.1. Weight functions. A *weight* on $\mathbf{R}^d$ is a positive function $\omega \in L_{loc}^\infty(\mathbf{R}^d)$ such that $1/\omega \in L_{loc}^\infty(\mathbf{R}^d)$. If ω and v are weights on $\mathbf{R}^d$, then ω is called *moderate* or *v-moderate*, if

$$\omega(x+y) \leq C\omega(x)v(y), \quad x, y \in \mathbf{R}^d, \quad (1.1)$$

for some constant C . The set of all moderate weights on $\mathbf{R}^d$ is denoted by $\mathcal{P}_E(\mathbf{R}^d)$. The weight v on $\mathbf{R}^d$ is called *submultiplicative*, if it is even and (1.1) holds for $\omega = v$ and $C = 1$. From now on, v always denote a submultiplicative weight if nothing else is stated. In particular, if (1.1) holds and v is submultiplicative, then it follows by straightforward computations that

$$C^{-1} \frac{\omega(x)}{v(y)} \leq \omega(x+y) \leq C\omega(x)v(y), \quad (1.2)$$

$$v(x+y) \leq v(x)v(y) \quad \text{and} \quad v(x) = v(-x) \geq 1, \quad x, y \in \mathbf{R}^d.$$

If ω is a moderate weight on $\mathbf{R}^d$, then there is a submultiplicative weight v on $\mathbf{R}^d$ such that (1.1) and (1.2) hold (cf. [47, 48, 52]). Moreover if v is submultiplicative on $\mathbf{R}^d$, then

$$1 \lesssim v(x) \lesssim e^{r|x|} \quad (1.3)$$

for some constant $r > 0$ (cf. [24]). Here and in what follows we write $A(\theta) \lesssim B(\theta)$, $\theta \in \Omega$, if there is a constant $c > 0$ such that $A(\theta) \leq cB(\theta)$ for all $\theta \in \Omega$. In particular, if ω is moderate, then

$$\omega(x+y) \lesssim \omega(x)e^{r|y|} \quad \text{and} \quad e^{-r|x|} \leq \omega(x) \lesssim e^{r|x|}, \quad x, y \in \mathbf{R}^d \quad (1.4)$$

for some $r > 0$.

Next we introduce suitable subclasses of $\mathcal{P}_E$.

Definition 1.1. Let $s > 0$. The set $\mathcal{P}_{E,s}^0(\mathbf{R}^d)$ ($\mathcal{P}_{E,s}^0(\mathbf{R}^d)$) consists of all $\omega \in \mathcal{P}_E(\mathbf{R}^d)$ such that

$$\omega(x+y) \lesssim \omega(x)e^{r|y|^{\frac{1}{s}}}, \quad x, y \in \mathbf{R}^d, \quad (1.5)$$

holds for some (every) $r > 0$.

By (1.4) it follows that $\mathcal{P}_{E,s_1}^0 = \mathcal{P}_{E,s_2}^0 = \mathcal{P}_E$ when $s_1 < 1$ and $s_2 \leq 1$. For convenience we set $\mathcal{P}_E^0(\mathbf{R}^d) = \mathcal{P}_{E,1}^0(\mathbf{R}^d)$.

1.2. **Gelfand-Shilov spaces.** We let $\mathcal{F}$ be the Fourier transform given by

$$(\mathcal{F}f)(\xi) = \widehat{f}(\xi) \equiv (2\pi)^{-d/2} \int_{\mathbf{R}^d} f(x) e^{-i\langle x, \xi \rangle} dx$$

when $f \in L^1(\mathbf{R}^d)$. Here $\langle \cdot, \cdot \rangle$ denotes the usual scalar product on $\mathbf{R}^d$.

Definition 1.2. The Gelfand-Shilov space $\mathcal{S}_s^\sigma(\mathbf{R}^d)$ of Roumieu type ($\Sigma_s^\sigma(\mathbf{R}^d)$ of Beurling type), $\sigma > 0$, $s > 0$, consists of all $f \in \mathcal{S}(\mathbf{R}^d)$ such that

$$|f(x)| \lesssim e^{-r|x|^{\frac{1}{s}}} \quad \text{and} \quad |(\mathcal{F}f)(\xi)| \lesssim e^{-r|\xi|^{\frac{1}{\sigma}}}, \quad x, \xi \in \mathbf{R}^d \quad (1.6)$$

for some $r > 0$ (for all $r > 0$). We set $\mathcal{S}_s = \mathcal{S}_s^s$ and $\Sigma_s = \Sigma_s^s$.

The classes $\mathcal{S}_s^\sigma(\mathbf{R}^d)$ and related generalizations were widely studied, and used in the applications to partial differential equations, see for example [2, 6, 9, 28, 35, 38]. We recall the following characterisations of $\mathcal{S}_s^\sigma(\mathbf{R}^d)$.

Proposition 1.3. *Let $s, \sigma > 0$, $p \in [1, \infty]$ and let $f \in \mathcal{S}(\mathbf{R}^d)$. Then the following conditions are equivalent:*

- (1) $f \in \mathcal{S}_s^\sigma(\mathbf{R}^d)$ ($f \in \Sigma_s^\sigma(\mathbf{R}^d)$);
- (2) for some (every) $h > 0$ it holds

$$\|x^\alpha f\|_{L^p} \lesssim h^{|\alpha|} \alpha!^s \quad \text{and} \quad \|\xi^\beta \widehat{f}\|_{L^p} \lesssim h^{|\beta|} \beta!^\sigma, \quad \alpha, \beta \in \mathbf{N}^d;$$

- (3) for some (every) $h > 0$ it holds

$$\|x^\alpha f\|_{L^p} \lesssim h^{|\alpha|} \alpha!^s \quad \text{and} \quad \|\partial^\beta f\|_{L^p} \lesssim h^{|\beta|} \beta!^\sigma, \quad \alpha, \beta \in \mathbf{N}^d;$$

- (4) for some (every) $h > 0$ it holds

$$\|x^\alpha \partial^\beta f(x)\|_{L^p} \lesssim h^{|\alpha+\beta|} \alpha!^s \beta!^\sigma, \quad \alpha, \beta \in \mathbf{N}^d;$$

- (5) for some (every) $h, r > 0$ it holds

$$\|e^{r|\cdot|^{\frac{1}{s}}} \partial^\alpha f\|_{L^p} \lesssim h^{|\alpha|} (\alpha!)^\sigma \alpha \in \mathbf{N}^d.$$

Remark 1.4. Let $h, s, s_0, \sigma, \sigma_0 \in \mathbf{R}_+$ be such that $s + \sigma > 1$ and $s_0 + \sigma_0 \geq 1$, and let $\mathcal{S}_{s,h}^\sigma(\mathbf{R}^d)$ be the set of all $f \in C^\infty(\mathbf{R}^d)$ such that

$$\|f\|_{\mathcal{S}_{s,h}^\sigma} \equiv \sup \frac{|x^\beta \partial^\alpha f(x)|}{h^{|\alpha+\beta|} \alpha!^\sigma \beta!^s}$$

is finite. Here the supremum is taken over all $\alpha, \beta \in \mathbf{N}^d$ and $x \in \mathbf{R}^d$.

Obviously $\mathcal{S}_{s,h}^\sigma(\mathbf{R}^d)$ is a Banach space which increases as h , s and σ increase, and is contained in $\mathcal{S}(\mathbf{R}^d)$, the set of Schwartz functions on $\mathbf{R}^d$. Furthermore

$$\mathcal{S}_{s,h}^\sigma(\mathbf{R}^d) \quad \text{and} \quad \bigcup_{h>0} \mathcal{S}_{s_0,h}^{\sigma_0}(\mathbf{R}^d)$$

are dense in $\mathcal{S}(\mathbf{R}^d)$. Hence, the dual $(\mathcal{S}_{s,h}^\sigma)'(\mathbf{R}^d)$ of $\mathcal{S}_{s,h}^\sigma(\mathbf{R}^d)$ is a Banach space which contains $\mathcal{S}'(\mathbf{R}^d)$.

The spaces $\mathcal{S}_s^\sigma(\mathbf{R}^d)$ and $\Sigma_s^\sigma(\mathbf{R}^d)$ are the inductive and projective limits, respectively, of $\mathcal{S}_{s,h}^\sigma(\mathbf{R}^d)$ with respect to h . This implies that

$$\mathcal{S}_s^\sigma(\mathbf{R}^d) = \bigcup_{h>0} \mathcal{S}_{s,h}^\sigma(\mathbf{R}^d) \quad \text{and} \quad \Sigma_s^\sigma(\mathbf{R}^d) = \bigcap_{h>0} \mathcal{S}_{s,h}^\sigma(\mathbf{R}^d), \quad (1.7)$$

and that the topology for $\mathcal{S}_s^\sigma(\mathbf{R}^d)$ is the strongest possible one such that each inclusion map from $\mathcal{S}_{s,h}^\sigma(\mathbf{R}^d)$ to $\mathcal{S}_s^\sigma(\mathbf{R}^d)$ is continuous. Moreover, any of the conditions (2)–(5) in Proposition 1.3 induce the same topology for $\mathcal{S}_s^\sigma(\mathbf{R}^d)$ and $\Sigma_s^\sigma(\mathbf{R}^d)$.

The Gelfand-Shilov distribution spaces $(\mathcal{S}_s^\sigma)'(\mathbf{R}^d)$ and $(\Sigma_s^\sigma)'(\mathbf{R}^d)$ are the projective and inductive limit respectively of $(\mathcal{S}_{s,h}^\sigma)'(\mathbf{R}^d)$. Hence

$$(\mathcal{S}_s^\sigma)'(\mathbf{R}^d) = \bigcap_{h>0} (\mathcal{S}_{s,h}^\sigma)'(\mathbf{R}^d) \quad \text{and} \quad (\Sigma_s^\sigma)'(\mathbf{R}^d) = \bigcup_{h>0} (\Sigma_{s,h}^\sigma)'(\mathbf{R}^d). \quad (1.7)'$$

We have that $(\mathcal{S}_s^\sigma)'$ and $(\Sigma_s^\sigma)'$ are the topological duals of $\mathcal{S}_s^\sigma$ and Σ_s^σ , respectively (see [37]).

Remark 1.5. Let $s, \sigma > 0$. Then $\Sigma_s^\sigma(\mathbf{R}^d)$ is a Fréchet space with seminorms $\|\cdot\|_{\Sigma_{s,h}^\sigma}$, $h > 0$. Moreover, $\mathcal{S}_s^\sigma(\mathbf{R}^d) \neq \{0\}$ if and only if $s + \sigma \geq 1$, and $\Sigma_s^\sigma(\mathbf{R}^d) \neq \{0\}$ if and only if $s + \sigma \geq 1$ and $(s, \sigma) \neq (\frac{1}{2}, \frac{1}{2})$. Moreover, if $\varepsilon > 0$ and $s + \sigma \geq 1$, then

$$\Sigma_s^\sigma(\mathbf{R}^d) \subseteq \mathcal{S}_s^\sigma(\mathbf{R}^d) \subseteq \Sigma_{s+\varepsilon}^{\sigma+\varepsilon}(\mathbf{R}^d) \subseteq \mathcal{S}'(\mathbf{R}^d) \subseteq \mathcal{S}'(\mathbf{R}^d) \subseteq (\Sigma_{s+\varepsilon}^{\sigma+\varepsilon})'(\mathbf{R}^d) \subseteq (\mathcal{S}_s^\sigma)'(\mathbf{R}^d).$$

If in addition $(s, \sigma) \neq (\frac{1}{2}, \frac{1}{2})$, then

$$(\mathcal{S}_s^\sigma)'(\mathbf{R}^d) \subseteq (\Sigma_s^\sigma)'(\mathbf{R}^d).$$

The Gelfand-Shilov spaces are invariant or possess convenient mapping properties under several basic transformations. For example they are invariant under translations, dilations, and under (partial) Fourier transformations.

The Fourier transform $\mathcal{F}$ on $\mathcal{S}'(\mathbf{R}^d)$ extends uniquely to homeomorphisms on $\mathcal{S}'(\mathbf{R}^d)$, $\mathcal{S}'_s(\mathbf{R}^d)$ and $\Sigma'_s(\mathbf{R}^d)$, and restricts to homeomorphisms on $\mathcal{S}_s(\mathbf{R}^d)$ and $\Sigma_s(\mathbf{R}^d)$, and to a unitary operator on $L^2(\mathbf{R}^d)$.

We also recall some mapping properties of Gelfand-Shilov spaces under short-time Fourier transforms. Let $\phi \in \mathcal{S}'(\mathbf{R}^d)$ be fixed. For every $f \in \mathcal{S}'(\mathbf{R}^d)$, the *short-time Fourier transform* $V_\phi f$ is the distribution on $\mathbf{R}^{2d}$ defined by the formula

$$(V_\phi f)(x, \xi) = \mathcal{F}(f \overline{\phi(\cdot - x)})(\xi) = (f, \phi(\cdot - x) e^{i\langle \cdot, \xi \rangle}). \quad (1.8)$$

We recall that if $T(f, \phi) \equiv V_\phi f$ when $f, \phi \in \mathcal{S}_{1/2}(\mathbf{R}^d)$, then T is uniquely extendable to sequentially continuous mappings

$$T : \mathcal{S}'_s(\mathbf{R}^d) \times \mathcal{S}_s(\mathbf{R}^d) \rightarrow \mathcal{S}'_s(\mathbf{R}^{2d}) \bigcap C^\infty(\mathbf{R}^{2d}),$$

$$T : \mathcal{S}'_s(\mathbf{R}^d) \times \mathcal{S}'_s(\mathbf{R}^d) \rightarrow \mathcal{S}'_s(\mathbf{R}^{2d}),$$

and similarly when $\mathcal{S}_s$ and $\mathcal{S}'_s$ are replaced by Σ_s and Σ'_s , respectively, or by $\mathcal{S}$ and $\mathcal{S}'$, respectively (cf. [7, 52]). We also note that $V_\phi f$ takes the form

$$V_\phi f(x, \xi) = (2\pi)^{-d/2} \int_{\mathbf{R}^d} f(y) \overline{\phi(y - x)} e^{-i\langle y, \xi \rangle} dy \quad (1.8)'$$

when $f \in L^p_{(\omega)}(\mathbf{R}^d)$ for some $\omega \in \mathcal{P}_E(\mathbf{R}^d)$, $\phi \in \Sigma_1(\mathbf{R}^d)$ and $p \geq 1$. Here $L^p_{(\omega)}(\mathbf{R}^d)$, when $p \in (0, \infty]$ and $\omega \in \mathcal{P}_E(\mathbf{R}^d)$, is the set of all $f \in L^p_{loc}(\mathbf{R}^d)$ such that $f \cdot \omega \in L^p(\mathbf{R}^d)$.

1.3. Suitable function classes with Gelfand-Shilov regularity. The next result shows that for any $\omega \in \mathcal{P}_E(\mathbf{R}^d)$ one can find an equivalent weight ω_0 which satisfies similar regularity estimates as functions in Gelfand-Shilov spaces.

Proposition 1.6. *Let $\omega \in \mathcal{P}_E(\mathbf{R}^d)$ and $s > 0$. Then there is an $\omega_0 \in \mathcal{P}_E(\mathbf{R}^d) \cap C^\infty(\mathbf{R}^d)$ such that the following is true:*

- (1) $\omega_0 \asymp \omega$;
- (2) $|\partial^\alpha \omega_0(x)| \lesssim \omega_0(x) h^{|\alpha|} \alpha!^s \asymp \omega(x) h^{|\alpha|} \alpha!^s$ for every $h > 0$.

Proof. We may assume that $s < 1$. It suffices to prove that (2) should hold for some $h > 0$. Let $\phi_0 \in \Sigma_{1-s}^s(\mathbf{R}^d) \setminus 0$, and let $\phi = |\phi_0|^2$. Then $\phi \in \Sigma_{1-s}^s(\mathbf{R}^d)$, giving that

$$|\partial^\alpha \phi(x)| \lesssim h^{|\alpha|} e^{-c|x|^{\frac{1}{1-s}}} \alpha!^s,$$

for every $h > 0$ and $c > 0$. Now let $\omega_0 = \omega * \phi$.

We have

$$\begin{aligned} |\partial^\alpha \omega_0(x)| &= \left| \int \omega(y) (\partial^\alpha \phi)(x-y) dy \right| \\ &\lesssim h^{|\alpha|} \alpha!^s \int \omega(y) e^{-c|x-y|^{\frac{1}{1-s}}} dy \\ &\lesssim h^{|\alpha|} \alpha!^s \int \omega(x+(y-x)) e^{-c|x-y|^{\frac{1}{1-s}}} dy \\ &\lesssim h^{|\alpha|} \alpha!^s \omega(x) \int e^{-\frac{c}{2}|x-y|^{\frac{1}{1-s}}} dy \asymp h^{|\alpha|} \alpha!^s \omega(x), \end{aligned}$$

where the last inequality follows from (1.4) and the fact that ϕ is bounded by a super exponential function. This gives the first part of (2).

The equivalences in (1) follows in the same way as in [52]. More precisely, by (1.4) we have

$$\begin{aligned} \omega_0(x) &= \int \omega(y) \phi(x-y) dy = \int \omega(x+(y-x)) \phi(x-y) dy \\ &\lesssim \omega(x) \int e^{c|x-y|} \phi(x-y) dy \asymp \omega(x). \end{aligned}$$

In the same way, (1.4) gives

$$\begin{aligned} \omega_0(x) &= \int \omega(y) \phi(x-y) dy = \int \omega(x+(y-x)) \phi(x-y) dy \\ &\gtrsim \omega(x) \int e^{-c|x-y|} \phi(x-y) dy \asymp \omega(x), \end{aligned}$$

and (1) as well as the second part of (2) follow. $\square$

A weight ω_0 which satisfies Proposition 1.6 (2) is called *elliptic* or *s-elliptic*.

Definition 1.7. Let $s \geq 0$ and $\omega \in \mathcal{P}_E(\mathbf{R}^d)$. The class $\Gamma_s^{(\omega)}(\mathbf{R}^d)$ ($\Gamma_{0,s}^{(\omega)}(\mathbf{R}^d)$) consists of all $f \in C^\infty(\mathbf{R}^d)$ such that

$$|D^\alpha f(x)| \lesssim \omega(x) h^{|\alpha|} \alpha!^s, \quad x \in \mathbf{R}^d,$$

for some $h > 0$ (for every $h > 0$).

Evidently, by Proposition 1.6 it follows that the family of symbol classes in Definition 1.7 is not reduced when the assumption $\omega \in \mathcal{P}_E(\mathbf{R}^{2d})$ is replaced by the more restrictive assumption $\omega \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ or by $\omega \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$.

By similar arguments as in the proof of Proposition 1.6 we get the following analog of Proposition 2.3.16 in [33]. The details are left for the reader.

Proposition 1.8. Let $s > 1/2$, $\omega \in \mathcal{P}_E(\mathbf{R}^{2d})$, and $\phi \in \Sigma_s(\mathbf{R}^{2d})$. Then $\omega * \phi$ belongs to $\Gamma_{0,s}^{(\omega)}$.

The following definition is motivated by Lemma 2.6.13 in [33].

Definition 1.9. Let $s \geq 1$, $\omega \in \mathcal{P}_E(\mathbf{R}^d)$ and $\vartheta_0 = 1 + |\log \omega|$. Then c is called *comparable* to ω with respect to $s \geq 1$ if

- (1) $\|c - \log \omega\|_{L^\infty} < \infty$
- (2) $c \in \Gamma_s^{(\vartheta_0)}(\mathbf{R}^d)$ and $\partial^\alpha c \in \Gamma_s^{(1)}(\mathbf{R}^d)$, when $|\alpha| = 1$.

Proposition 1.10. *Let $\omega, v \in \mathcal{P}_E(\mathbf{R}^d)$ be such that v is submultiplicative and ω is v -moderate. Also let*

$$v_1(x) \equiv 1 + |\log v(x)| \quad \text{and} \quad \omega_1(x) \equiv 1 + |\log \omega(x)|.$$

Then v_1 is submultiplicative and ω_1 is v_1 -moderate, satisfying (1.1) with $1 + \log C \geq 1$ in place of C .

Proof. By (1.2) we get

$$\begin{aligned} v_1(x+y) &= 1 + \log v(x+y) \leq 1 + \log v(x) + \log v(y) \\ &\leq (1 + \log v(x))(1 + \log v(y)) = v_1(x)v_1(y), \end{aligned}$$

and

$$\begin{aligned} \omega_1(x+y) &= 1 + |\log \omega(x+y)| \\ &\leq 1 + \log C + |\log \omega(x)| + \log v(y) \\ &\leq (1 + \log C)(1 + |\log \omega(x)|)(1 + \log v(y)) \\ &\leq (1 + \log C)\omega_1(x)v_1(y), \end{aligned}$$

as claimed. $\square$

Lemma 1.11. *Let $s \geq 1$, $\omega \in \mathcal{P}_E(\mathbf{R}^d)$ and $\vartheta_0 = 1 + |\log \omega|$. Then the following is true:*

- (1) *there exists an elliptic weight $\omega_0 \in \mathcal{P}_E(\mathbf{R}^d) \cap \Gamma_s^{(\omega)}(\mathbf{R}^d)$ such that $\omega \asymp \omega_0$ and $1 + |\log \omega_0| \in \mathcal{P}_E(\mathbf{R}^d) \cap \Gamma_s^{(\vartheta_0)}(\mathbf{R}^d)$;*
- (2) *there exists an element c which is comparable with ω_0 .*

Proof. If $\omega_1(x)$ is equal to $\omega(x)$ when $\omega(x) \geq e$ or $\omega(x) \leq e^{-1}$, and 3 otherwise, then ω_1 is equivalent to ω . The result now follows from Proposition 1.6 and its proof, with ω_1 in place of ω . $\square$

1.4. Modulation spaces. Let $\phi \in \Sigma_1(\mathbf{R}^d) \setminus 0$, $p, q \in (0, \infty]$ and $\omega \in \mathcal{P}_E(\mathbf{R}^{2d})$ be fixed. Then the *modulation space* $M_{(\omega)}^{p,q}(\mathbf{R}^d)$ consists of all $f \in \Sigma'_1(\mathbf{R}^d)$ such that

$$\|f\|_{M_{(\omega)}^{p,q}} \equiv \left(\int \left(\int |V_\phi f(x, \xi) \omega(x, \xi)|^p dx \right)^{q/p} d\xi \right)^{1/q} < \infty \quad (1.9)$$

(with the obvious modifications when $p = \infty$ and/or $q = \infty$). We set $M_{(\omega)}^p = M_{(\omega)}^{p,p}$, and if $\omega = 1$, then we set $M^{p,q} = M_{(\omega)}^{p,q}$ and $M^p = M_{(\omega)}^p$.

The following proposition is a consequence of well-known facts in [12, 20, 21, 54]. Here and in what follows, we let p' denotes the conjugate exponent of p , i.e.

$$p' = \begin{cases} \infty & \text{when } p \in (0, 1] \\ \frac{p}{p-1} & \text{when } p \in (1, \infty) \\ 1 & \text{when } p = \infty. \end{cases}$$

Proposition 1.12. *Let $p, q, p_j, q_j, r \in (0, \infty]$ be such that $r \leq \min(1, p, q)$, $j = 1, 2$, let $\omega, \omega_1, \omega_2, v \in \mathcal{P}_E(\mathbf{R}^{2d})$ be such that ω is v -moderate, $\phi \in M_{(v)}^r(\mathbf{R}^d) \setminus 0$, and let $f \in \Sigma'_1(\mathbf{R}^d)$. Then the following is true:*

- (1) *$f \in M_{(\omega)}^{p,q}(\mathbf{R}^d)$ if and only if (1.9) holds, i.e. $M_{(\omega)}^{p,q}(\mathbf{R}^d)$ is independent of the choice of ϕ . Moreover, $M_{(\omega)}^{p,q}$ is a Banach space under the norm in (1.9), and different choices of ϕ give rise to equivalent norms;*

(2) if $p_1 \leq p_2$, $q_1 \leq q_2$ and $\omega_2 \leq C\omega_1$ for some constant C , then

$$\Sigma_1(\mathbf{R}^d) \subseteq M_{(\omega_1)}^{p_1, q_1}(\mathbf{R}^d) \subseteq M_{(\omega_2)}^{p_2, q_2}(\mathbf{R}^d) \subseteq \Sigma'_1(\mathbf{R}^d).$$

Proposition 1.12(1) allows us to be rather vague about to the choice of $\phi \in M_{(v)}^r \setminus 0$ in (1.9). For example, if $C > 0$ is a constant and Ω is a subset of Σ'_1 , then $\|a\|_{M_{(\omega)}^{p, q}} \leq C$ for every $a \in \Omega$, means that the inequality holds for some choice of $\phi \in M_{(v)}^r \setminus 0$ and every $a \in \Omega$. Evidently, for any other choice of $\phi \in M_{(v)}^r \setminus 0$, a similar inequality is true although C may have to be replaced by a larger constant, if necessary.

Let $s, t \in \mathbf{R}$. Then the weights

$$(x, \xi) \mapsto \langle \xi \rangle^s \langle x \rangle^t \quad \text{and} \quad (x, \xi) \mapsto \langle (x, \xi) \rangle^s, \quad x, \xi \in \mathbf{R}^d, \quad (1.10)$$

are common in the applications. It follows that they belong to $\mathcal{P}(\mathbf{R}^{2d})$ for every $s, t \in \mathbf{R}$. If $\omega \in \mathcal{P}(\mathbf{R}^{2d})$, then ω is moderated by any of the weights in (1.10) provided s and t are chosen large enough.

Remark 1.13. For modulation spaces of the form $M_{(\omega)}^{p, q}$ with fixed $p, q \in [1, \infty]$ the norm equivalence in Proposition 1.12(1) can be extended to a larger class of windows. In fact, assume that $\omega, v \in \mathcal{P}_E(\mathbf{R}^{2d})$ with ω being v -moderate and

$$1 \leq r \leq \min(p, p', q, q').$$

Let $\phi \in M_{(v)}^r(\mathbf{R}^d) \setminus \{0\}$. Then a Gelfand-Shilov distribution $f \in \Sigma'_1(\mathbf{R}^d)$ belongs to $M_{(\omega)}^{p, q}(\mathbf{R}^d)$, if and only if $V_\phi f \in L_{(\omega)}^{p, q}(\mathbf{R}^{2d})$. Furthermore, different choices of $\phi \in M_{(v)}^r(\mathbf{R}^d) \setminus \{0\}$ in $\|V_\phi f\|_{L_{(\omega)}^{p, q}}$ give rise to equivalent norms. (Cf. Theorem 3.1 in [51].)

1.5. A broader family of modulation spaces. As announced in the introduction we consider in Section 2 mapping properties for pseudo-differential operators when acting on a broader class of modulation spaces, which are defined by imposing certain types of translation invariant solid BF-space norms on the short-time Fourier transforms. (Cf. [11–16, 18].)

Definition 1.14. Let $\mathcal{B} \subseteq L_{loc}^r(\mathbf{R}^d)$ be a quasi-Banach of order $r \in (0, 1]$, and let $v \in \mathcal{P}_E(\mathbf{R}^d)$. Then $\mathcal{B}$ is called a *translation invariant Quasi-Banach Function space on $\mathbf{R}^d$* , or *invariant QBF space on $\mathbf{R}^d$* , if there is a constant C such that the following conditions are fulfilled:

(1) if $x \in \mathbf{R}^d$ and $f \in \mathcal{B}$, then $f(\cdot - x) \in \mathcal{B}$, and

$$\|f(\cdot - x)\|_{\mathcal{B}} \leq C v(x) \|f\|_{\mathcal{B}}; \quad (1.11)$$

(2) if $f, g \in L_{loc}^r(\mathbf{R}^d)$ satisfy $g \in \mathcal{B}$ and $|f| \leq |g|$, then $f \in \mathcal{B}$ and

$$\|f\|_{\mathcal{B}} \leq C \|g\|_{\mathcal{B}}.$$

If v belongs to $\mathcal{P}_{E, s}(\mathbf{R}^d)$ ($\mathcal{P}_{E, s}^0(\mathbf{R}^d)$), then $\mathcal{B}$ in Definition 1.14 is called an *invariant BF-space* of Roumieu type (Beurling type) of order s .

We notice that the quasi-norm $\|\cdot\|_{\mathcal{B}}$ in Definition 1.14 should satisfy

$$\|f + g\|_{\mathcal{B}} \leq 2^{\frac{1}{r}-1} (\|f\|_{\mathcal{B}} + \|g\|_{\mathcal{B}}) \quad f, g \in \mathcal{B}. \quad (1.12)$$

By Aoki and Rolewić in [1, 39] it follows that there is an equivalent quasi-norm to the previous one which additionally satisfies

$$\|f + g\|_{\mathcal{B}}^r \leq \|f\|_{\mathcal{B}}^r + \|g\|_{\mathcal{B}}^r \quad f, g \in \mathcal{B}. \quad (1.13)$$

From now on we suppose that the quasi-norm of $\mathcal{B}$ has been chosen such that both (1.12) and (1.13) hold true.

It follows from (2) in Definition 1.14 that if $f \in \mathcal{B}$ and $h \in L^\infty$, then $f \cdot h \in \mathcal{B}$, and

$$\|f \cdot h\|_{\mathcal{B}} \leq C \|f\|_{\mathcal{B}} \|h\|_{L^\infty}. \quad (1.14)$$

If $r = 1$, then $\mathcal{B}$ in Definition 1.14 is a Banach space, and the condition (2) means that a translation invariant QBF-space is a solid BF-space in the sense of (A.3) in [15]. The space $\mathcal{B}$ in Definition 1.14 is called an *invariant BF-space* (with respect to v) if $r = 1$, and Minkowski's inequality holds true, i. e.

$$\|f * \varphi\|_{\mathcal{B}} \leq C \|f\|_{\mathcal{B}} \|\varphi\|_{L^1_{(v)}}, \quad f \in \mathcal{B}, \varphi \in \Sigma_1(\mathbf{R}^d) \quad (1.15)$$

for some constant C which is independent of $f \in \mathcal{B}$ and $\varphi \in \Sigma_1(\mathbf{R}^d)$.

Example 1.15. Assume that $p, q \in [1, \infty]$, and let $L_1^{p,q}(\mathbf{R}^{2d})$ be the set of all $f \in L^1_{loc}(\mathbf{R}^{2d})$ such that

$$\|f\|_{L_1^{p,q}} \equiv \left(\int \left(\int |f(x, \xi)|^p dx \right)^{q/p} d\xi \right)^{1/q}$$

if finite. Also let $L_2^{p,q}(\mathbf{R}^{2d})$ be the set of all $f \in L^1_{loc}(\mathbf{R}^{2d})$ such that

$$\|f\|_{L_2^{p,q}} \equiv \left(\int \left(\int |f(x, \xi)|^q d\xi \right)^{p/q} dx \right)^{1/p}$$

is finite. Then it follows that $L_1^{p,q}$ and $L_2^{p,q}$ are translation invariant BF-spaces with respect to $v = 1$.

For translation invariant BF-spaces we make the following observation.

Proposition 1.16. Assume that $v \in \mathcal{P}_E(\mathbf{R}^d)$, and that $\mathcal{B}$ is an invariant BF-space with respect to v such that (1.15) holds true. Then the convolution mapping $(\varphi, f) \mapsto \varphi * f$ from $C_0^\infty(\mathbf{R}^d) \times \mathcal{B}$ to $\mathcal{B}$ extends uniquely to a continuous mapping from $L^1_{(v)}(\mathbf{R}^d) \times \mathcal{B}$ to $\mathcal{B}$, and (1.15) holds true for any $f \in \mathcal{B}$ and $\varphi \in L^1_{(v)}(\mathbf{R}^d)$.

The result is a straightforward consequence of the fact that C_0^∞ is dense in $L^1_{(v)}$.

The quasi-Banach space $\mathcal{B}$ above is usually a mixed quasi-normed Lebesgue space, given as follows. Let E be a non-degenerate parallelepiped in $\mathbf{R}^d$ which is spanned by the ordered basis $\kappa(E) = \{e_1, \dots, e_d\}$. That is,

$$E = \{x_1 e_1 + \dots + x_d e_d; (x_1, \dots, x_d) \in \mathbf{R}^d, 0 \leq x_k \leq 1, k = 1, \dots, d\}.$$

The corresponding lattice, dual parallelepiped and dual lattice are given by

$$\Lambda_E = \{j_1 e_1 + \dots + j_d e_d; (j_1, \dots, j_d) \in \mathbf{Z}^d\},$$

$$E' = \{\xi_1 e'_1 + \dots + \xi_d e'_d; (\xi_1, \dots, \xi_d) \in \mathbf{R}^d, 0 \leq \xi_k \leq 1, k = 1, \dots, d\},$$

and

$$\Lambda'_E = \Lambda_{E'} = \{\iota_1 e'_1 + \dots + \iota_d e'_d; (\iota_1, \dots, \iota_d) \in \mathbf{Z}^d\},$$

respectively, where the ordered basis $\kappa(E') = \{e'_1, \dots, e'_d\}$ of E' satisfies

$$\langle e_j, e'_k \rangle = 2\pi \delta_{jk} \quad \text{for every } j, k = 1, \dots, d.$$

Note here that the Fourier analysis with respect to general biorthogonal bases has recently been developed in [41].

The basis $e'_1, \dots, e'_d$ is called the *dual basis* of $e_1, \dots, e_d$. We observe that there is a matrix T_E such that $e_1, \dots, e_d$ and $e'_1, \dots, e'_d$ are the images of the standard basis under T_E and $T_{E'} = 2\pi(T_E^{-1})^t$, respectively.

In the following we let

$$\max \mathbf{q} = \max(q_1, \dots, q_d) \quad \text{and} \quad \min \mathbf{q} = \min(q_1, \dots, q_d)$$

when $\mathbf{q} = (q_1, \dots, q_d) \in (0, \infty]^d$, and χ_Ω be the characteristic function of Ω .

Definition 1.17. Let E be a non-degenerate parallelepiped in $\mathbf{R}^d$ spanned by the ordered set $\kappa(E) \equiv \{e_1, \dots, e_d\}$ in $\mathbf{R}^d$, $\mathbf{p} = (p_1, \dots, p_d) \in (0, \infty]^d$ and $r = \min(1, \mathbf{p})$. If $f \in L_{loc}^r(\mathbf{R}^d)$, then

$$\|f\|_{L_{\kappa(E)}^{\mathbf{p}}} \equiv \|g_{d-1}\|_{L^{p_d}(\mathbf{R})}$$

where $g_k(z_k)$, $z_k \in \mathbf{R}^{d-k}$, $k = 0, \dots, d-1$, are inductively defined as

$$g_0(x_1, \dots, x_d) \equiv |f(x_1 e_1 + \dots + x_d e_d)|,$$

and

$$g_k(z_k) \equiv \|g_{k-1}(\cdot, z_k)\|_{L^{p_k}(\mathbf{R})}, \quad k = 1, \dots, d-1.$$

The space $L_{\kappa(E)}^{\mathbf{p}}(\mathbf{R}^d)$ consists of all $f \in L_{loc}^r(\mathbf{R}^d)$ such that $\|f\|_{L_{\kappa(E)}^{\mathbf{p}}}$ is finite, and is called *E-split Lebesgue space (with respect to $\mathbf{p}$ and $\kappa(E)$)*.

Definition 1.18. Let $E_0 \subseteq \mathbf{R}^d$ be a non-degenerate parallelepiped with dual parallelepiped E'_0 , and $E \subseteq \mathbf{R}^{2d}$ be a parallelepiped spanned by the ordered set $\kappa(E) \equiv \{e_1, \dots, e_{2d}\}$. Then E is called a *phase-shift split parallelepiped* (with respect to E_0) if E is non-degenerate and d of the vectors in $\{e_1, \dots, e_{2d}\}$ span E_0 and the other d vectors is the corresponding dual basis which span E'_0 .

Next we consider the extended class of modulation spaces which we are interested in.

Definition 1.19. Assume that $\mathcal{B}$ is a translation invariant QBF-space on $\mathbf{R}^{2d}$, $\omega \in \mathcal{P}_E(\mathbf{R}^{2d})$, and that $\phi \in \Sigma_1(\mathbf{R}^d) \setminus 0$. Then the set $M_{(\omega)} = M(\omega, \mathcal{B})$ consists of all $f \in \Sigma'_1(\mathbf{R}^d)$ such that

$$\|f\|_{M_{(\omega)}} = \|f\|_{M(\omega, \mathcal{B})} \equiv \|V_{\phi} f \omega\|_{\mathcal{B}}$$

is finite.

Obviously, we have $M_{(\omega)}^{p,q}(\mathbf{R}^d) = M(\omega, \mathcal{B})$ when $\mathcal{B} = L_1^{p,q}(\mathbf{R}^{2d})$ (cf. Example 1.15). It follows that many properties which are valid for the classical modulation spaces also hold for the spaces of the form $M(\omega, \mathcal{B})$. For example we have the following proposition, which shows that the definition of $M(\omega, \mathcal{B})$ is independent of the choice of ϕ when $\mathcal{B}$ is a Banach space. We omit the proof since it follows by similar arguments as in the proof of Proposition 11.3.2 in [21]. (See also [36] for topological aspects of $M(\omega, \mathcal{B})$.)

Proposition 1.20. Let $\mathcal{B}$ be an invariant BF-space with respect to $v_0 \in \mathcal{P}_E(\mathbf{R}^{2d})$ for $j = 1, 2$. Also let $\omega, v \in \mathcal{P}_E(\mathbf{R}^{2d})$ be such that ω is v -moderate, $M(\omega, \mathcal{B})$ is the same as in Definition 1.19, and let $\phi \in M_{(v_0 v)}^1(\mathbf{R}^d) \setminus 0$ and $f \in \Sigma'_1(\mathbf{R}^d)$. Then $M(\omega, \mathcal{B})$ is a Banach space, and $f \in M(\omega, \mathcal{B})$ if and only if $V_{\phi} f \omega \in \mathcal{B}$, and different choices of ϕ gives rise to equivalent norms in $M(\omega, \mathcal{B})$.

We refer to [11–16, 18, 20, 21, 40, 54] for more facts about modulation spaces.

1.6. Pseudo-differential operators. We use the notation $\mathbf{M}(d, \Omega)$ for the set of $d \times d$ -matrices with entries in the set Ω . Let $s \geq 1/2$, $a \in \mathcal{S}_s(\mathbf{R}^{2d})$, and $A \in \mathbf{M}(d, \mathbf{R})$ be fixed. Then, the pseudo-differential operator $\text{Op}_A(a)$ is the linear and continuous operator on $\mathcal{S}_s(\mathbf{R}^d)$ given by

$$(\text{Op}_A(a)u)(x) = (2\pi)^{-d} \iint a(x - A(x - y), \xi) f(y) e^{i\langle x - y, \xi \rangle} dy d\xi \quad (1.16)$$

when $f \in \mathcal{S}_s(\mathbf{R}^d)$. For general $a \in \mathcal{S}'_s(\mathbf{R}^{2d})$, the pseudo-differential operator $\text{Op}_A(a)$ is defined as the continuous operator from $\mathcal{S}_s(\mathbf{R}^d)$ to $\mathcal{S}'_s(\mathbf{R}^d)$ with distribution kernel given by

$$K_{a,A}(x, y) = (2\pi)^{-d/2} (\mathcal{F}_2^{-1} a)(x - A(x - y), x - y). \quad (1.17)$$

Here $\mathcal{F}_2 F$ is the partial Fourier transform of $F(x, y) \in \mathcal{S}'_s(\mathbf{R}^{2d})$ with respect to the y variable. This definition makes sense, since the mappings

$$\mathcal{F}_2 \quad \text{and} \quad F(x, y) \mapsto F(x - A(x - y), y - x) \quad (1.18)$$

are homeomorphisms on $\mathcal{S}'_s(\mathbf{R}^{2d})$. In particular, the map $a \mapsto K_{a,A}$ is a homeomorphism on $\mathcal{S}'_s(\mathbf{R}^{2d})$.

Remark 1.21. For any $K \in \mathcal{S}'_s(\mathbf{R}^{d_1+d_2})$, let T_K be the linear and continuous mapping from $\mathcal{S}_s(\mathbf{R}^{d_1})$ to $\mathcal{S}'_s(\mathbf{R}^{d_2})$, defined by the formula

$$(T_K f, g)_{L^2(\mathbf{R}^{d_2})} = (K, g \otimes \bar{f})_{L^2(\mathbf{R}^{d_1+d_2})}. \quad (1.19)$$

It is well-known that if $t \in \mathbf{R}$, then it follows from e.g. [5, 32] that the Schwartz kernel theorem also holds in the context of Gelfans-Shilov spaces. That is, the mappings $K \mapsto T_K$ and $a \mapsto \text{Op}_t(a)$ are bijective from $\mathcal{S}'_s(\mathbf{R}^{2d})$ to the set of linear and continuous mappings from $\mathcal{S}_s(\mathbf{R}^d)$ to $\mathcal{S}'_s(\mathbf{R}^d)$. Similar facts hold true if $\mathcal{S}_s$ and $\mathcal{S}'_s$ are replaced by Σ_s and Σ'_s , respectively (or by $\mathcal{S}$ and $\mathcal{S}'$, respectively).

As a consequence of Remark 1.21 it follows that for each $a_1 \in \mathcal{S}'_s(\mathbf{R}^{2d})$ and $A_1, A_2 \in \mathbf{M}(d, \mathbf{R})$, there is a unique $a_2 \in \mathcal{S}'_s(\mathbf{R}^{2d})$ such that $\text{Op}_{A_1}(a_1) = \text{Op}_{A_2}(a_2)$. The relation between a_1 and a_2 is given by

$$\text{Op}_{A_1}(a_1) = \text{Op}_{A_2}(a_2) \quad \Leftrightarrow \quad a_2(x, \xi) = e^{i\langle (A_1 - A_2)D_\xi, D_x \rangle} a_1(x, \xi). \quad (1.20)$$

(Cf. [30].) Note here that the right-hand side makes sense, since it is equivalent to $\widehat{a}_2(\xi, x) = e^{i(A_1 - A_2)\langle x, \xi \rangle} \widehat{a}_1(\xi, x)$, and that the map $a \mapsto e^{i\langle Ax, \xi \rangle} a$ is continuous on $\mathcal{S}'_s$ when $A \in \mathbf{M}(d, \mathbf{R})$.

Let $A \in \mathbf{M}(d, \mathbf{R})$ and $a \in \mathcal{S}'_s(\mathbf{R}^{2d})$ be fixed. Then a is called a rank-one element with respect to A , if the corresponding pseudo-differential operator is of rank-one, i.e.

$$\text{Op}_A(a)f = (f, f_2)f_1, \quad f \in \mathcal{S}_s(\mathbf{R}^d), \quad (1.21)$$

for some $f_1, f_2 \in \mathcal{S}'_s(\mathbf{R}^d)$. By straightforward computations it follows that (1.21) is fulfilled, if and only if $a = (2\pi)^{d/2} W_{f_1, f_2}^A$, where W_{f_1, f_2}^A is the A -Wigner distribution defined by the formula

$$W_{f_1, f_2}^A(x, \xi) \equiv \mathcal{F}(f_1(x + A \cdot) \overline{f_2(x - (I - A) \cdot)})(\xi), \quad (1.22)$$

which takes the form

$$W_{f_1, f_2}^A(x, \xi) = (2\pi)^{-d/2} \int f_1(x + Ay) \overline{f_2(x - (I - A)y)} e^{-i\langle y, \xi \rangle} dy,$$

when $f_1, f_2 \in \mathcal{S}_s(\mathbf{R}^d)$. Here $I \in \mathbf{M}(d, \mathbf{R})$ is the identity matrix. By combining these facts with (1.20) it follows that

$$W_{f_1, f_2}^{A_2} = e^{i\langle (A_1 - A_2)D_\xi, D_x \rangle} W_{f_1, f_2}^{A_1}, \quad (1.23)$$

for each $f_1, f_2 \in \mathcal{S}'_s(\mathbf{R}^d)$ and $A_1, A_2 \in \mathbf{M}(d, \mathbf{R})$. Since the Weyl case is particularly important, we set $W_{f_1, f_2}^A = W_{f_1, f_2}$ when $A = \frac{1}{2}I$, i.e. W_{f_1, f_2} is the usual (cross-)Wigner distribution of f_1 and f_2 .

For future references we note the link

$$(\text{Op}_A(a)f, g)_{L^2(\mathbf{R}^d)} = (2\pi)^{-d/2} (a, W_{g, f}^A)_{L^2(\mathbf{R}^{2d})}, \quad a \in \mathcal{S}'_s(\mathbf{R}^{2d}) \quad \text{and} \quad f, g \in \mathcal{S}_s(\mathbf{R}^d) \quad (1.24)$$

between pseudo-differential operators and Wigner distributions, which follows by straightforward computations (see also e.g. [55]).

Next we discuss the Weyl product, the twisted convolution and related objects. Let $s \geq 1/2$ and let $a, b \in \mathcal{S}'_s(\mathbf{R}^{2d})$. Then the Weyl product $a \# b$ between a and b is the function or distribution which fulfills $\text{Op}^w(a \# b) = \text{Op}^w(a) \circ \text{Op}^w(b)$, provided

the right-hand side makes sense as a continuous operator from $\mathcal{S}_s(\mathbf{R}^d)$ to $\mathcal{S}'_s(\mathbf{R}^d)$. More generally, if $A \in \mathbf{M}(d, \mathbf{R})$, then the product $\#_A$ is defined by the formula

$$\text{Op}_A(a \#_A b) = \text{Op}_A(a) \circ \text{Op}_A(b), \quad (1.25)$$

provided the right-hand side makes sense as a continuous operator from $\mathcal{S}_s(\mathbf{R}^d)$ to $\mathcal{S}'_s(\mathbf{R}^d)$, in which case a and b are called *suitable* or *admissible*. We also use the notation $\#$ instead of $\#_A$ when $A = \frac{1}{2}I$ (i. e. in the Weyl case).

The Weyl product can also, in a convenient way, be expressed in terms of the symplectic Fourier transform and the twisted convolution. More precisely, let $s \geq 1/2$. Then the *symplectic Fourier transform* for $a \in \mathcal{S}_s(\mathbf{R}^{2d})$ is defined by the formula

$$(\mathcal{F}_\sigma a)(X) = \pi^{-d} \int a(Y) e^{2i\sigma(X, Y)} dY,$$

where σ is the symplectic form given by

$$\sigma(X, Y) = \langle y, \xi \rangle - \langle x, \eta \rangle, \quad X = (x, \xi) \in \mathbf{R}^{2d}, \quad Y = (y, \eta) \in \mathbf{R}^{2d}.$$

We note that $\mathcal{F}_\sigma = T \circ (\mathcal{F} \otimes (\mathcal{F}^{-1}))$, when $(Ta)(x, \xi) = a(\xi, x)$. In particular, $\mathcal{F}_\sigma$ is continuous on $\mathcal{S}_s(\mathbf{R}^{2d})$, and extends uniquely to a homeomorphism on $\mathcal{S}'_s(\mathbf{R}^{2d})$, and to a unitary map on $L^2(\mathbf{R}^{2d})$, since similar facts hold for $\mathcal{F}$. Furthermore, $\mathcal{F}_\sigma^2$ is the identity operator.

Let $s \geq 1/2$ and $a, b \in \mathcal{S}_s(\mathbf{R}^{2d})$. Then the *twisted convolution* of a and b is defined by the formula

$$(a *_\sigma b)(X) = \left(\frac{2}{\pi}\right)^{\frac{d}{2}} \int a(X - Y) b(Y) e^{2i\sigma(X, Y)} dY. \quad (1.26)$$

The definition of $*_\sigma$ extends in different ways. For example, it extends to a continuous multiplication on $L^p(\mathbf{R}^{2d})$ when $p \in [1, 2]$, and to a continuous map from $\mathcal{S}'_s(\mathbf{R}^{2d}) \times \mathcal{S}_s(\mathbf{R}^{2d})$ to $\mathcal{S}'_s(\mathbf{R}^{2d})$. If $a, b \in \mathcal{S}'_s(\mathbf{R}^{2d})$, then $a \# b$ makes sense if and only if $a *_\sigma b$ makes sense, and then

$$a \# b = (2\pi)^{-\frac{d}{2}} a *_\sigma (\mathcal{F}_\sigma b). \quad (1.27)$$

We also remark that for the twisted convolution we have

$$\mathcal{F}_\sigma(a *_\sigma b) = (\mathcal{F}_\sigma a) *_\sigma b = \check{a} *_\sigma (\mathcal{F}_\sigma b), \quad (1.28)$$

where $\check{a}(X) = a(-X)$ (cf. [46, 51, 53]). A combination of (1.27) and (1.28) gives

$$\mathcal{F}_\sigma(a \# b) = (2\pi)^{-\frac{d}{2}} (\mathcal{F}_\sigma a) *_\sigma (\mathcal{F}_\sigma b). \quad (1.29)$$

We now define the subspace of symbols in $\mathcal{S}'(\mathbf{R}^{2d})$ which give rise to $L^2(\mathbf{R}^d)$ -bounded pseudo-differential operators, which will be useful in the sequel.

Definition 1.22. The set $s_\infty^w(\mathbf{R}^{2d})$ consists of all $a \in \mathcal{S}'(\mathbf{R}^{2d})$ such that $\text{Op}^w(a)$ is linear and continuous on $L^2(\mathbf{R}^d)$, and we set

$$\|a\|_{s_\infty^w} \equiv \|\text{Op}^w(a)\|_{L^2(\mathbf{R}^d) \rightarrow L^2(\mathbf{R}^d)}.$$

Remark 1.23. By Remark 1.21 it follows that the map $a \mapsto \text{Op}^w(a)$ is an isometric bijection from $s_\infty^w(\mathbf{R}^{2d})$ to the set of linear continuous operators on $L^2(\mathbf{R}^d)$.

Remark 1.24. We remark that the relations in this subsection hold true after $\mathcal{S}_s$, $\mathcal{S}'_s$ and $s \geq \frac{1}{2}$ are replaced by Σ_s , Σ'_s and $s > \frac{1}{2}$ respectively, in each place.

Next we recall some algebraic properties and characterisations of $\Gamma_s^{(\omega)}(\mathbf{R}^{2d})$ and $\Gamma_{0,s}^{(\omega)}(\mathbf{R}^{2d})$, and begin with the following. We refer to [4] for its proof.

Proposition 1.25. *Let $s \geq 1$, $\omega_j \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$, $A_j \in \mathbf{M}(d, \mathbf{R})$ for $j = 1, 2$, and let $\omega_{0,r}(X, Y) = \omega_0(X) e^{-r|Y|^{\frac{1}{s}}}$ when $r > 0$. Then the following is true:*

- (1) If $a_1, a_2 \in \Sigma'_s(\mathbf{R}^{2d})$ satisfy $\text{Op}_{A_1}(a_1) = \text{Op}_{A_2}(a_2)$, then $a_1 \in \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$ if and only if $a_2 \in \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$.
- (2) $\Gamma_s^{(\omega_1)} \# \Gamma_s^{(\omega_2)} \subseteq \Gamma_s^{(\omega_1 \omega_2)}$.
- (3) $\Gamma_s^{(\omega_0)} = \bigcup_{r>0} M_{(1/\omega_0, r)}^{\infty, 1} = \bigcup_{r \geq 0} \mathcal{M}_{(1/\omega_0, r)}^{\infty, 1}$.

Proposition 1.26. *Let $s \geq 1$, $\omega_j \in \mathcal{P}_{E, s}(\mathbf{R}^{2d})$, $A_j \in \mathbf{M}(d, \mathbf{R})$ for $j = 1, 2$, and let $\omega_{0, r}(X, Y) = \omega_0(X) e^{-r|Y|^{\frac{1}{s}}}$ when $r > 0$. Then the following is true:*

- (1) If $a_1, a_2 \in \Sigma'_s(\mathbf{R}^{2d})$ satisfy $\text{Op}_{A_1}(a_1) = \text{Op}_{A_2}(a_2)$, then $a_1 \in \Gamma_{0, s}^{(\omega_0)}(\mathbf{R}^{2d})$ if and only if $a_2 \in \Gamma_{0, s}^{(\omega_0)}(\mathbf{R}^{2d})$.
- (2) $\Gamma_{0, s}^{(\omega_1)} \# \Gamma_{0, s}^{(\omega_2)} \subseteq \Gamma_{0, s}^{(\omega_1 \omega_2)}$.
- (3) $\Gamma_{0, s}^{(\omega_0)} = \bigcap_{r>0} M_{(1/\omega_0, r)}^{\infty, 1} = \bigcap_{r \geq 0} \mathcal{M}_{(1/\omega_0, r)}^{\infty, 1}$.

In time-frequency analysis one also considers mapping properties for pseudo-differential operators between modulation spaces or with symbols in modulation spaces. Especially we need the following two results, where the first one is a generalisation of [44, Theorem 2.1] by Tachizawa, and the second one is a weighted version of [21, Theorem 14.5.2]. We refer to [56] for the proof of the first two propositions and to [56] for the proof of the third one.

Proposition 1.27. *Assume that $A \in \mathbf{M}(d, \mathbf{R})$, $s \geq 1$, $\omega, \omega_0 \in \mathcal{P}_{E, s}^0(\mathbf{R}^{2d})$, $a \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$, and that $\mathcal{B}$ is an invariant BF-space on $\mathbf{R}^{2d}$ of Beurling type. Then $\text{Op}_A(a)$ is continuous from $M(\omega_0 \omega, \mathcal{B})$ to $M(\omega_0, \mathcal{B})$, and also continuous on $\mathcal{S}_s(\mathbf{R}^d)$ and on $\mathcal{S}'_s(\mathbf{R}^d)$.*

Proposition 1.28. *Assume that $A \in \mathbf{M}(d, \mathbf{R})$, $s \geq 1$, $\omega, \omega_0 \in \mathcal{P}_{E, s}(\mathbf{R}^{2d})$, $a \in \Gamma_{0, s}^{(\omega)}(\mathbf{R}^{2d})$, and that $\mathcal{B}$ is an invariant BF-space on $\mathbf{R}^{2d}$ of Roumieu type. Then $\text{Op}_A(a)$ is continuous from $M(\omega_0 \omega, \mathcal{B})$ to $M(\omega_0, \mathcal{B})$, and also continuous on $\Sigma_s(\mathbf{R}^d)$ and on $\Sigma'_s(\mathbf{R}^d)$.*

Proposition 1.29. *Assume that $p, q \in (0, \infty]$, $r \leq \min(p, q, 1)$, $\omega \in \mathcal{P}_E(\mathbf{R}^{2d} \oplus \mathbf{R}^{2d})$ and $\omega_1, \omega_2 \in \mathcal{P}_E(\mathbf{R}^{2d})$ satisfy*

$$\frac{\omega_2(X - Y)}{\omega_1(X + Y)} \leq C\omega(X, Y), \quad X, Y \in \mathbf{R}^{2d}, \quad (1.30)$$

for some constant C . If $a \in \mathcal{M}_{(\omega)}^{\infty, r}(\mathbf{R}^{2d})$, then $\text{Op}^w(a)$ extends uniquely to a continuous map from $M_{(\omega_1)}^{p, q}(\mathbf{R}^d)$ to $M_{(\omega_2)}^{p, q}(\mathbf{R}^d)$.

Finally we need the following result concerning mapping properties of modulation spaces under the Weyl product. The result is a special case of Theorem 0.3 in [10].

Proposition 1.30. *Assume that $\omega_j \in \mathcal{P}_E(\mathbf{R}^{2d} \oplus \mathbf{R}^{2d})$ for $j = 0, 1, 2$ satisfy*

$$\omega_0(X, Y) \leq C\omega_1(X - Y + Z, Z)\omega_2(X + Z, Y - Z), \quad (1.31)$$

for some constant $C > 0$ independent of $X, Y, Z \in \mathbf{R}^{2d}$. Then the map $(a, b) \mapsto a \# b$ from $\Sigma_1(\mathbf{R}^{2d}) \times \Sigma_1(\mathbf{R}^{2d})$ to $\Sigma_1(\mathbf{R}^{2d})$ extends uniquely to a mapping from $\mathcal{M}_{(\omega_1)}^{\infty, 1}(\mathbf{R}^{2d}) \times \mathcal{M}_{(\omega_2)}^{\infty, 1}(\mathbf{R}^{2d})$ to $\mathcal{M}_{(\omega_0)}^{\infty, 1}(\mathbf{R}^{2d})$.

In the proof of our main theorem we will need the following consequence of Proposition 1.30.

Proposition 1.31. *Let $s \geq 1$, $\omega_0, v_0, v_1 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d} \oplus \mathbf{R}^{2d})$ be such that ω_0 is v_0 -moderate. Set $\vartheta = \omega_0^{1/2}$, and*

$$\begin{aligned}\omega_1(X, Y) &= \frac{v_0(2Y)^{1/2}v_1(2Y)}{\vartheta(X+Y)\vartheta(X-Y)}, \\ \omega_2(X, Y) &= \vartheta(X-Y)\vartheta(X+Y)v_1(2Y), \\ v_2(X, Y) &= v_1(2Y).\end{aligned}\tag{1.32}$$

Then

$$\Gamma_s^{(1/\vartheta)} \# \mathcal{M}_{(\omega_1)}^{\infty,1} \# \Gamma_s^{(1/\vartheta)} \subseteq \mathcal{M}_{(v_2)}^{\infty,1}, \tag{1.33}$$

$$\Gamma_s^{(1/\vartheta)} \# \mathcal{M}_{(v_2)}^{\infty,1} \# \Gamma_s^{(1/\vartheta)} \subseteq \mathcal{M}_{(\omega_2)}^{\infty,1}. \tag{1.34}$$

The same holds with $\mathcal{P}_{E,s}$ and $\Gamma_{0,s}^{(1/\vartheta)}$ in place of $\mathcal{P}_{E,s}^0$ and $\Gamma_s^{(1/\vartheta)}$ respectively, at each occurrence.

Proof. Since $\Gamma_s^{(1/\vartheta)} = \bigcup_{r \geq 0} \mathcal{M}_{(\vartheta_r)}^{\infty,1}$ with $\vartheta_r(X, Y) = \vartheta(X)e^{r|Y|^{1/s}}$ (Proposition 1.25(3)), it suffices to argue with the symbol class $\mathcal{M}_{(\vartheta_r)}^{\infty,1}$ for some sufficiently large r instead of $\Gamma_s^{(1/\vartheta)}$.

Introducing the intermediate weight

$$\omega_3(X, Y) = \frac{v_1(2Y)\vartheta(X+Y)}{\omega_0(X-Y)}.$$

we will show that for suitable r

$$\omega_3(X, Y) \leq C\omega_1(X-Y+Z, Z)\vartheta_r(X+Z, Y-Z) \tag{1.35}$$

$$v_1(2Y) \leq C\vartheta_r(X-Y+Z, Z)\omega_3(X+Z, Y-Z). \tag{1.36}$$

Proposition 1.30 applied to (1.35) shows that $\mathcal{M}_{(\omega_1)}^{\infty,1} \# \Gamma_s^{(1/\vartheta)} \subseteq \mathcal{M}_{(\omega_3)}^{\infty,1}$, and (1.36) implies that $\Gamma_s^{(1/\vartheta)} \# \mathcal{M}_{(\omega_3)}^{\infty,1} \subseteq \mathcal{M}_{(v_2)}^{\infty,1}$, and (1.33) follows.

Since ϑ is $v_0^{1/2}$ -moderate and $v_0 \in \mathcal{P}_{E,s}^0$, we have

$$\vartheta(X-Y)^{-1} \leq v_0(2Z)^{1/2}\vartheta(X-Y+2Z)^{-1} \quad \text{and} \quad \vartheta(X+Y) \leq \vartheta(X+Z)e^{r|Y-Z|^{1/s}}$$

for suitable $r > 0$. Using these inequalities repeatedly in the following, a straightforward computation yields

$$\begin{aligned}\omega_3(X, Y) &= \frac{v_1(2Y)\vartheta(X+Y)}{\vartheta(X-Y)^2} \\ &\leq C_1 \frac{v_0(2Z)^{1/2}v_1(2Z)\vartheta(X+Z)e^{r|Y-Z|^{1/s}}}{\vartheta(X-Y+2Z)\vartheta(X-Y)} \\ &= C_1\omega_1(X-Y+Z, Z)\vartheta_r(X+Z, Y-Z),\end{aligned}$$

for some $C_1 > 0$ and $r > 0$.

Likewise we obtain

$$\begin{aligned}
v_1(2Y) &= \frac{\vartheta(X-Y)v_1(2Y)\vartheta(X-Y)}{\vartheta(X-Y)^2} \\
&\leq C_1 \frac{\vartheta(X-Y)v_0(2Y)^{1/2}v_1(2Y)\vartheta(X+Y)}{\vartheta(X-Y)^2} \\
&\leq C_2 \frac{\vartheta(X-Y+Z)e^{r|Z|^{\frac{1}{s}}}v_0(2(Y-Z))^{1/2}v_1(2(Y-Z))\vartheta(X+Y)}{\vartheta(X-Y+2Z)^2} \\
&= C_2\vartheta_r(X-Y+Z, Z)\omega_3(X+Z, Y-Z).
\end{aligned}$$

The twisted convolution relation (1.34) is proved similarly. Let

$$\omega_4(X, Y) = \vartheta(X-Y)v_1(2Y)$$

be the intermediate weight. Then the inequality

$$\begin{aligned}
\omega_4(X, Y) &= \vartheta(X-Y)v_1(2Y) \leq C\vartheta(X-Y+Z)e^{r|Z|^{\frac{1}{s}}}v_1(2(Y-Z)) \\
&= C\vartheta_r(X-Y+Z, Z)v_2(X+Z, Y-Z)
\end{aligned}$$

implies that $\Gamma_s^{(1/\vartheta)} \# \mathcal{M}_{(v_2)}^{\infty, 1} \subseteq \mathcal{M}_{(\omega_4)}^{\infty, 1}$.

Similarly we obtain

$$\begin{aligned}
\omega_2(X, Y) &\leq C\vartheta(X-Y)v_1(2Z)\vartheta(X+Z)e^{r|Z-Y|^{\frac{1}{s}}} \\
&= C\omega_4(X-Y+Z, Z)\vartheta(X+Z)e^{r|Z-Y|^{\frac{1}{s}}} \\
&= C\omega_4(X-Y+Z, Z)\vartheta_r(X+Z, Y-Z),
\end{aligned}$$

and thus $\mathcal{M}_{(\omega_4)}^{\infty, 1} \# \Gamma_s^{(1/\vartheta)} \subseteq \mathcal{M}_{(\omega_2)}^{\infty, 1}$.

The case $\mathcal{P}_{E,s}$ and $\Gamma_{0,s}^{(1/\vartheta)}$ in place of $\mathcal{P}_{E,s}^0$ and $\Gamma_s^{(1/\vartheta)}$ respectively, at each occurrence, is treated in similar ways and is left for the reader. $\square$

1.7. The Wiener Algebra Property. As a further crucial tool in our study of the isomorphism property of Toeplitz operators we need to combine these continuity results with convenient invertibility properties. The so-called Wiener algebra property of certain symbol classes asserts that the inversion of a pseudo-differential operator preserves the symbol class and is often referred to as the spectral invariance of a symbol class.

Proposition 1.32. *Let $A \in \mathbf{M}(d, \mathbf{R})$. Then the following are true:*

- (1) *If $s > 1$, $a \in \Gamma_{0,s}^{(1)}(\mathbf{R}^{2d})$ and $\text{Op}_A(a)$ is invertible on $L^2(\mathbf{R}^d)$, then $\text{Op}_A(a)^{-1} = \text{Op}_A(b)$ for some $b \in \Gamma_{0,s}^{(1)}(\mathbf{R}^{2d})$.*
- (2) *If $s \geq 1$, $a \in \Gamma_s^{(1)}(\mathbf{R}^{2d})$ and $\text{Op}_A(a)$ is invertible on $L^2(\mathbf{R}^d)$, then $\text{Op}_A(a)^{-1} = \text{Op}_A(b)$ for some $b \in \Gamma_s^{(1)}(\mathbf{R}^{2d})$.*
- (3) *If $s \geq 1$, $v_0 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$ is submultiplicative, $v(X, Y) \equiv v_0(Y)$, $X, Y \in \mathbf{R}^{2d}$, $a \in M_{(v)}^{\infty, 1}(\mathbf{R}^{2d})$ and $\text{Op}_A(a)$ is invertible on $L^2(\mathbf{R}^d)$, then $\text{Op}_A(a)^{-1} = \text{Op}_A(b)$, for some $b \in M_{(v)}^{\infty, 1}(\mathbf{R}^{2d})$.*
- (4) *If $s > 1$, $v_0 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ is submultiplicative, $v(X, Y) \equiv v_0(Y)$, $X, Y \in \mathbf{R}^{2d}$, $a \in M_{(v)}^{\infty, 1}(\mathbf{R}^{2d})$ and $\text{Op}_A(a)$ is invertible on $L^2(\mathbf{R}^d)$, then $\text{Op}_A(a)^{-1} = \text{Op}_A(b)$, for some $b \in M_{(v)}^{\infty, 1}(\mathbf{R}^{2d})$.*

Proof. The results follows essentially from [22, Corollary 5.5] or [23]. More precisely, Suppose $s > 1$, $a \in \Gamma_s^{(1)}(\mathbf{R}^{2d})$, $\text{Op}_A(a)$ is invertible on $L^2(\mathbf{R}^d)$, and let $v_r(X, Y) = e^{r|Y|^{frac{1}{s}}}$ when $r \geq 0$. Then $a \in M_{(v_r)}^{\infty,1}(\mathbf{R}^{2d})$ for some $r > 0$. By [22, Corollary 5.5], $\text{Op}(M_{(v_r)}^{\infty,1}(\mathbf{R}^{2d}))$ is a Wiener algebra, giving that $\text{Op}(a)^{-1} = \text{Op}(b)$ for some $b \in M_{(v_r)}^{\infty,1}(\mathbf{R}^{2d}) \subseteq \Gamma_s^{(1)}(\mathbf{R}^{2d})$. This gives (2) in the case $s > 1$.

If instead $s = 1$, then by [17, Theorem 4.4] there is an $r_0 > 0$ such that $\text{Op}(a)^{-1} = \text{Op}(b)$ for some $b \in M_{(v_{r_0})}^{\infty,1}(\mathbf{R}^{2d}) \subseteq \Gamma_1^{(1)}(\mathbf{R}^{2d})$, and (2) follows for general $s \geq 1$.

By similar arguments, (1), (3) and (4) follow. The details are left for the reader. $\square$

1.8. Toeplitz Operators. Fix a symbol $a \in \mathcal{S}(\mathbf{R}^{2d})$ and a window $\varphi \in \mathcal{S}(\mathbf{R}^d)$. Then the Toeplitz operator $\text{Tp}_\varphi(a)$ is defined by the formula

$$(\text{Tp}_\varphi(a)f_1, f_2)_{L^2(\mathbf{R}^d)} = (a V_\varphi f_1, V_\varphi f_2)_{L^2(\mathbf{R}^{2d})}, \quad (1.37)$$

when $f_1, f_2 \in \mathcal{S}(\mathbf{R}^d)$. Obviously, $\text{Tp}_\varphi(a)$ is well-defined and extends uniquely to a continuous operator from $\mathcal{S}'(\mathbf{R}^d)$ to $\mathcal{S}(\mathbf{R}^d)$.

The definition of Toeplitz operators can be extended to more general classes of windows and symbols by using appropriate estimates for the short-time Fourier transforms in (1.37).

We state two possible ways of extending (1.37). The first result follows from [8, Corollary 4.2] and its proof, and the second result is a special case of [57, Theorem 3.1]. We use the notation $\mathcal{L}(V_1, V_2)$ for the set of linear and continuous mappings from the topological vector space V_1 into the topological vector space V_2 . We also set

$$\omega_{0,t}(X, Y) = v_1(2Y)^{t-1} \omega_0(X) \quad \text{for } X, Y \in \mathbf{R}^{2d}. \quad (1.38)$$

Proposition 1.33. *Let $0 \leq t \leq 1$, $p, q \in [1, \infty]$, and $\omega, \omega_0, v_1, v_0 \in \mathcal{P}(\mathbf{R}^{2d})$ be such that v_0 and v_1 are submultiplicative, ω_0 is v_0 -moderate and ω is v_1 -moderate. Set*

$$v = v_1^t v_0 \quad \text{and} \quad \vartheta = \omega_0^{1/2},$$

and let $\omega_{0,t}$ be as in (1.38). Then the following are true:

- (1) *The definition of $(a, \varphi) \mapsto \text{Tp}_\varphi(a)$ from $\mathcal{S}(\mathbf{R}^{2d}) \times \mathcal{S}(\mathbf{R}^d)$ to $\mathcal{L}(\mathcal{S}(\mathbf{R}^d), \mathcal{S}'(\mathbf{R}^d))$ extends uniquely to a continuous map from $\mathcal{M}_{(1/\omega_{0,t})}^\infty(\mathbf{R}^{2d}) \times M_{(v)}^1(\mathbf{R}^d)$ to $\mathcal{L}(\mathcal{S}(\mathbf{R}^d), \mathcal{S}'(\mathbf{R}^d))$.*
- (2) *If $\varphi \in M_{(v)}^1(\mathbf{R}^d)$ and $a \in \mathcal{M}_{(1/\omega_{0,t})}^\infty(\mathbf{R}^{2d})$, then $\text{Tp}_\varphi(a)$ extends uniquely to a continuous map from $M_{(\vartheta\omega)}^{p,q}(\mathbf{R}^d)$ to $M_{(\omega/\vartheta)}^{p,q}(\mathbf{R}^d)$.*

Proposition 1.34. *Let $\omega, \omega_1, \omega_2, v \in \mathcal{P}(\mathbf{R}^{2d})$ be such that ω_1 is v -moderate, ω_2 is v -moderate and $\omega = \omega_1/\omega_2$. Then the following are true:*

- (1) *The mapping $(a, \varphi) \mapsto \text{Tp}_\varphi(a)$ extends uniquely to a continuous map from $L_{(\omega)}^\infty(\mathbf{R}^{2d}) \times M_{(v)}^2(\mathbf{R}^d)$ to $\mathcal{L}(\mathcal{S}(\mathbf{R}^d), \mathcal{S}'(\mathbf{R}^d))$.*
- (2) *If $a \in L_{(1/\omega)}^\infty(\mathbf{R}^{2d})$ and $\varphi \in M_{(v)}^2(\mathbf{R}^d)$, then $\text{Tp}_\varphi(a)$ extends uniquely to a continuous operator from $M_{(\omega_1)}^2(\mathbf{R}^d)$ to $M_{(\omega_2)}^2(\mathbf{R}^d)$.*

1.9. Weyl formulation of Toeplitz operators. We finish this section by recalling some important relations between Weyl operators, Wigner distributions, and Toeplitz operators. For instance, the Weyl symbol of a Toeplitz operator is the convolution between the Toeplitz symbol and a Wigner distribution. More precisely, if $a \in \mathcal{S}(\mathbf{R}^{2d})$ and $\varphi \in \mathcal{S}(\mathbf{R}^d)$, then

$$\text{Tp}_\varphi(a) = (2\pi)^{-d/2} \text{Op}^w(a * W_{\varphi, \varphi}). \quad (1.39)$$

Our analysis of Toeplitz operators is based on the pseudo-differential operator representation given by (1.39). Furthermore, any extension of the definition of Toeplitz operators to cases which are not covered by Propositions 1.33 and 1.34 is based on this representation. Here we remark that this leads to situations where certain mapping properties for the pseudo-differential operator representation make sense, whereas similar interpretations are difficult or impossible to make in the framework of (1.37) (see Remark 5.7 in Section 2). We refer to [50] or Section 2 for extensions of Toeplitz operators in context of pseudo-differential operators.

2. CONFINEMENT OF THE SYMBOL CLASSES $\Gamma_s^{(\omega)}(\mathbf{R}^d)$ AND $\Gamma_{0,s}^{(\omega)}(\mathbf{R}^d)$

In this section we introduce and discuss basic properties for confinements for symbols in $\Gamma_s^{(\omega_0)}$ and in $\Gamma_{0,s}^{(\omega_0)}$. These considerations follow lines similar to the discussions in [3, 34], but are here adapted to symbols that possess Gevrey regularity. In particular, this requires the deduction of various types of delicate estimates, concerning the compositions of symbols that are confined in different ways.

We recall that if σ is the (standard) symplectic form on $\mathbf{R}^{2d}$, namely,

$$\sigma(X, Y) = \langle y, \xi \rangle - \langle x, \eta \rangle, \quad X = (x, \xi) \in \mathbf{R}^{2d}, \quad Y = (y, \eta) \in \mathbf{R}^{2d},$$

then the *symplectic Fourier transform* $\mathcal{F}_\sigma$ is defined by

$$(\mathcal{F}_\sigma a)(X) = \hat{a}(X) \equiv \pi^{-d} \int a(Y) e^{2i\sigma(X, Y)} dY,$$

and the twisted convolution $*_\sigma$ is given by

$$(a *_\sigma b)(X) \equiv \left(\frac{2}{\pi}\right)^{\frac{d}{2}} \int a(X - Y) b(Y) e^{2i\sigma(X, Y)} dY,$$

for suitable $a, b \in \mathcal{S}'_{1/2}(\mathbf{R}^{2d})$. The twisted convolution is linked to the Weyl product by the formula

$$a \# b = (2\pi)^{-\frac{d}{2}} a *_\sigma (\mathcal{F}_\sigma b),$$

hence,

$$(a \# b)(X) = \pi^{-d} \int a(X - Y) \hat{b}(Y) e^{2i\sigma(X, Y)} dY.$$

We also note that $\mathcal{F}_\sigma^2$ is the identity map.

In what follows we let $a_Y = a(\cdot - Y)$ when $a \in \mathcal{S}'_{1/2}(\mathbf{R}^{2d})$ and $Y \in \mathbf{R}^{2d}$, and in analogous ways, $b_Y, \phi_Y, \varphi_Y, \psi_Y$ etc. are defined when $b, \phi, \varphi, \psi \in \mathcal{S}'_{1/2}(\mathbf{R}^{2d})$. For admissible a and b we have

$$(a \# b)_Y = a_Y \# b_Y. \quad (2.1)$$

We also recall that if $\varphi \in \mathcal{S}_s(\mathbf{R}^{2d})$, then there are functions $\phi, \psi \in \mathcal{S}_s(\mathbf{R}^{2d})$ such that $\varphi = \phi \# \psi$ (cf. [5, 58]). The same is true if $\mathcal{S}_s$ is replaced by Σ_s or by $\mathcal{S}$. In particular, by choosing φ such that $\int \varphi(X) dX = 1$, (2.1) gives the following.

Proposition 2.1. *Let $s \geq \frac{1}{2}$. Then there are $\phi, \psi \in \mathcal{S}_s(\mathbf{R}^{2d})$ such that*

$$\int_{\mathbf{R}^{2d}} \psi_Y \# \phi_Y dY = 1. \quad (2.2)$$

The same holds true with Σ_s or $\mathcal{S}$ in place of $\mathcal{S}_s$, provided $s > \frac{1}{2}$.

For independent translations in Weyl products we have the following.

Proposition 2.2. *Let $s \geq \frac{1}{2}$ and let $\phi, \psi \in \mathcal{S}_s(\mathbf{R}^{2d})$. Then*

$$(\phi_Y \# \psi_Z)(X) = \Psi(X - Y, X - Z)$$

for some $\Psi \in \mathcal{S}_s(\mathbf{R}^{2d} \times \mathbf{R}^{2d})$. The same holds true with Σ_s or $\mathcal{S}$ in place of $\mathcal{S}_s$.

Proof. We only prove the result when $\phi, \psi \in \mathcal{S}_s(\mathbf{R}^{2d})$. The other cases follow by similar arguments and are left for the reader.

We have

$$\begin{aligned} (\phi_Y \# \psi_Z)(X) &= \pi^{-d} \int \phi(X - Y - Y_1) \widehat{\psi}(Y_1) e^{2i\sigma(Y_1, Z)} e^{2i\sigma(X, Y_1)} dY_1 \\ &= \pi^{-d} \int \phi((X - Y) - Y_1) \widehat{\psi}(Y_1) e^{2i\sigma(X - Z, Y_1)} dY_1 = \Psi(X - Y, X - Z), \end{aligned}$$

where

$$\Psi(X, Z) = \pi^{-d} \int \phi(X - Y_1) \widehat{\psi}(Y_1) e^{2i\sigma(Z, Y_1)} dY_1.$$

We note that

$$\Psi = (\mathcal{F}_{\sigma, 2} \circ T)(\phi \otimes \widehat{\psi}),$$

where $(T\Phi)(X, Z) = \Phi(X - Z, Z)$ when $\Phi \in \mathcal{S}_s(\mathbf{R}^{2d} \times \mathbf{R}^{2d})$, and $\mathcal{F}_{\sigma, 2}\Phi$ is the partial symplectic Fourier transform of $\Phi(X, Z)$ with respect to the Z variable. Since $(\phi, \psi) \mapsto \phi \otimes \widehat{\psi}$ is continuous from $\mathcal{S}_s(\mathbf{R}^{2d}) \times \mathcal{S}_s(\mathbf{R}^{2d})$ to $\mathcal{S}_s(\mathbf{R}^{2d} \times \mathbf{R}^{2d})$, and T and $\mathcal{F}_{\sigma, 2}\Phi$ are continuous on $\mathcal{S}_s(\mathbf{R}^{2d} \times \mathbf{R}^{2d})$, it follows that $\Psi \in \mathcal{S}_s(\mathbf{R}^{2d} \times \mathbf{R}^{2d})$. $\square$

Since Ψ in Proposition 2.2 belongs to similar types of spaces as ϕ and ψ , it follows that estimates of the form

$$|D^\alpha \Psi(X, Y)| \lesssim h^{|\alpha|} \alpha!^s e^{-(|X|^{\frac{1}{s}} + |Y|^{\frac{1}{s}})/h}$$

hold true. In particular, the following is an immediate consequence of Proposition 2.2 and some standard manipulations in Gelfand-Shilov theory.

Corollary 2.3. *Let $s \geq \frac{1}{2}$. If $\phi, \psi \in \mathcal{S}_s(\mathbf{R}^{2d})$ ($\phi, \psi \in \Sigma_s(\mathbf{R}^{2d})$), then*

$$|D_X^\alpha D_Y^\beta D_Z^\gamma (\phi_Y \# \psi_Z)(X)| \lesssim h^{|\alpha+\beta+\gamma|} (\alpha! \beta! \gamma!)^s e^{-(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}})/h} \quad (2.3)$$

for some $h > 0$ (for every $h > 0$).

Proof. Obviously, as stated in the previous Proposition 2.2, we can write $\phi_Y \# \psi_Z$ as $\Psi(X - Y, X - Z)$ for some $\Psi \in \mathcal{S}_s(\mathbf{R}^{2d} \times \mathbf{R}^{2d})$. Thus

$$\begin{aligned} |D_X^\alpha D_Y^\beta D_Z^\gamma \Psi(X - Y, X - Z)| &= \left| D_X^\alpha \left(D_1^\beta D_2^\gamma \Psi \right) (X - Y, X - Z) \right| \\ &\leq \sum_{\delta \leq \alpha} \binom{\alpha}{\delta} \left| \left(D_1^{\beta+\delta} D_2^{\gamma+\alpha-\delta} \Psi \right) (X - Y, X - Z) \right| \\ &\leq h^{|\alpha+\beta+\gamma|} \sum_{\delta \leq \alpha} \binom{\alpha}{\delta} ((\beta+\delta)! (\gamma+\alpha-\delta)!)^s e^{-r(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}})}. \end{aligned}$$

Moreover, we have

$$\sum_{\delta \leq \alpha} \binom{\alpha}{\delta} ((\beta+\delta)! (\gamma+\alpha-\delta)!)^s \leq 2^{|\alpha|} 4^{s|\alpha+\beta+\gamma|} (\alpha! \beta! \gamma!)^s.$$

Indeed, using the fact that $\sum_{\delta \leq \alpha} \binom{\alpha}{\delta} = 2^{|\alpha|}$ and that $n! \leq 2^n (n-k)! k!$, which implies $(n+k)! \leq 2^{n+k} n! k!$, then

$$(\alpha + \beta + \gamma)! = \prod_{j=1}^d (\alpha_j + \beta_j + \gamma_j)! \leq \prod_{j=1}^d 4^{\alpha_j + \beta_j + \gamma_j} \alpha_j! \beta_j! \gamma_j! = 4^{|\alpha+\beta+\gamma|} \alpha! \beta! \gamma!.$$

Thus, inequality (2.3) holds with $2 \cdot 4^s h$ in place of h . $\square$

The next fundamental result is a consequence of Theorem 4.12 in [4].

Proposition 2.4. *Let $s \geq \frac{1}{2}$ and $\vartheta \in \mathcal{P}_E(\mathbf{R}^{2d})$. Then, the map $(\phi, a) \mapsto \phi \# a$ is continuous from $\Sigma_s(\mathbf{R}^{2d}) \times \Gamma_s^{(\vartheta)}(\mathbf{R}^{2d})$ to $\mathcal{S}_s(\mathbf{R}^{2d})$.*

The next lemma concerns uniform estimates of the Weyl product between elements in sets Ω_{j,Y_j} , $Y_j \in \mathbf{R}^{2d}$, $j = 1, 2$, such that related sets $\Omega_j^\cup$, given by

$$\Omega_j^\cup \equiv \bigcup_{Y \in \mathbf{R}^{2d}} \{a(\cdot + Y); a \in \Omega_{j,Y}\}, \quad (2.4)$$

are bounded in $\mathcal{S}_s(\mathbf{R}^{2d})$ or in $\Sigma_s(\mathbf{R}^{2d})$, $j = 1, 2$.

Lemma 2.5. *Let $s \geq \frac{1}{2}$ and let $\Omega_{j,Y} \subseteq \mathcal{S}(\mathbf{R}^{2d})$, $Y \in \mathbf{R}^{2d}$, $\Omega_j^\cup$ be as in (2.4), $a_j \in \Omega_j^\cup$ and choose $Y_j \in \mathbf{R}^{2d}$ such that $a_j \in \Omega_{j,Y_j}$, $j = 1, 2$. Then the following is true:*

- (1) *if $\Omega_j^\cup$ are bounded in $\mathcal{S}_s(\mathbf{R}^{2d})$, then there are constants $C > 0$ and $h > 0$ which are independent of $Y_j \in \mathbf{R}^{2d}$ and a_j , $j = 1, 2$, and such that*

$$|(a_1^{(\alpha_1)} \# a_2^{(\alpha_2)})(X)| \leq Ch^{|\alpha_1 + \alpha_2|} (\alpha_1! \alpha_2!)^s e^{-(|X - Y_1|^{\frac{1}{s}} + |X - Y_2|^{\frac{1}{s}} + |Y_1 - Y_2|^{\frac{1}{s}})/h} \quad (2.5)$$

and

$$|D^\alpha(a_1 \# a_2)(X)| \leq Ch^{|\alpha|} \alpha!^s e^{-(|X - Y_1|^{\frac{1}{s}} + |X - Y_2|^{\frac{1}{s}} + |Y_1 - Y_2|^{\frac{1}{s}})/h} \quad (2.6)$$

- (2) *if $\Omega_j^\cup$ are bounded in $\Sigma_s(\mathbf{R}^{2d})$, then for every $h > 0$, there is a constant $C > 0$ which is independent of $Y_j \in \mathbf{R}^{2d}$ and a_j , $j = 1, 2$, and such that (2.5) and (2.6) hold.*

Proof. We only prove (2). The assertion (1) follows by similar arguments and is left for the reader.

We have $D_X(a_1 \# a_2) = (D_X a_1) \# a_2 + a_1 \# (D_X a_2)$, which implies that the Leibnitz rule is valid for the Weyl product. Hence, (2.6) follows for every $h > 0$ if we prove that (2.5) holds for every $h > 0$.

Let $Y = Y_1$, $Z = Y_2$, $a = a_1$, $b = a_2$, $a_Y = a(\cdot + Y)$ and $b_Z = b(\cdot + Z)$. Then

$$\begin{aligned} a_1 \# a_2(X) &= \pi^{-d} \int a_Y((X - Y) - Y_1) \mathcal{F}_\sigma(b_Z(\cdot - Z))(Y_1) e^{2i\sigma(X, Y_1)} dY_1 \\ &= \pi^{-d} \int a_Y((X - Y) - Y_1) \mathcal{F}_\sigma(b_Z)(Y_1) e^{2i\sigma(X - Z, Y_1)} dY_1 \\ &= \mathcal{F}_\sigma(a_Y(X_1 - \cdot))(\mathcal{F}_\sigma b_Z)(X_2), \end{aligned}$$

where $X_1 = X - Y$ and $X_2 = X - Z$. That is $a_1 \# a_2(X) = G(X_1, X_2)$, where

$$G = G_{Y,Z} = (\mathcal{F}_{\sigma,2} \circ T)(a_Y \otimes b_Z).$$

Here $\mathcal{F}_{\sigma,2} F$ is the partial symplectic Fourier transform of $F(\cdot, X_2)$ with respect to the variable $X_2 \in \mathbf{R}^{2d}$, and

$$(TF)(X, Y) = F(X - Y, Y).$$

Since both $\mathcal{F}_{\sigma,2}$ and T are continuous on $\Sigma_s(\mathbf{R}^{2d} \times \mathbf{R}^{2d})$, it follows from the boundedness of the sets $\Omega_j^\cup$ that for every $h > 0$, there is a constant C_h which is independent of $a_j \in \Omega_j^\cup$ such that

$$|D_{X_1}^\alpha D_{X_2}^\beta G(X_1, X_2)| \leq C_h h^{|\alpha + \beta|} (\alpha \beta)^s e^{-(|X_1|^{\frac{1}{s}} + |X_2|^{\frac{1}{s}})/h}.$$

By straightforward computations it follows that

$$|D_{X_1}^\alpha D_{X_2}^\beta G(X_1, X_2)| = |(a^{(\alpha)} \# b^{(\beta)})|,$$

since

$$|Y_1 - Y_2|^{\frac{1}{s}} \leq c(|X - Y_1|^{\frac{1}{s}} + |X - Y_2|^{\frac{1}{s}})$$

for some constant c which only depends on s . The estimate (2.5) follows from these relations. $\square$

Lemma 2.6. *Let $s \geq \frac{1}{2}$, $\phi, \psi \in \Sigma_s(\mathbf{R}^{2d})$, $\omega, \vartheta \in \mathcal{P}_E(\mathbf{R}^{2d})$, $\phi_Y = \phi(\cdot - Y)$, and $\psi_Z = \psi(\cdot - Z)$. Then the following is true:*

- (1) *if $a \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$ ($a \in \Gamma_{0,s}^{(\omega)}(\mathbf{R}^{2d})$), then*

$$|D_X^\alpha D_Y^\beta (\phi_Y a)(X)| \lesssim h_1^{|\alpha|} h_2^{|\beta|} (\alpha! \beta!)^s e^{-|X-Y|^{\frac{1}{s}}/h_1} \min(\omega(X), \omega(Y)) \quad (2.7)$$

and

$$|D_X^\alpha D_Y^\beta (\phi_Y \# a)(X)| \lesssim h_1^{|\alpha|} h_2^{|\beta|} (\alpha! \beta!)^s e^{-|X-Y|^{\frac{1}{s}}/h_1} \min(\omega(X), \omega(Y)), \quad (2.8)$$

for some $h_1 > 0$ (for every $h_1 > 0$) and every $h_2 > 0$;

- (2) *if $a_1 \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$ and $a_2 \in \Gamma_s^{(\vartheta)}(\mathbf{R}^{2d})$ ($a_1 \in \Gamma_{0,s}^{(\omega)}(\mathbf{R}^{2d})$ and $a_2 \in \Gamma_{0,s}^{(\vartheta)}(\mathbf{R}^{2d})$), then*

$$\begin{aligned} & |D_X^\alpha D_Y^\beta D_Z^\gamma ((\phi_Y a_1) \# (\psi_Z a_2))(X)| \\ & \lesssim h_1^{|\alpha+\beta|} h_2^{|\gamma|} (\alpha! \beta! \gamma!)^s e^{-(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}} + |Y-Z|^{\frac{1}{s}})/h_1} \min_{X_1, X_2 \in \{X, Y, Z\}} (\omega(X_1) \vartheta(X_2)), \end{aligned}$$

for some $h_1 > 0$ (for every $h_1 > 0$) and every $h_2 > 0$.

Proof. We only consider the case when $a \in \Gamma_{0,s}^{(\omega)}(\mathbf{R}^{2d})$ and $b \in \Gamma_{0,s}^{(\vartheta)}(\mathbf{R}^{2d})$. The other cases follow by similar arguments and are left for the reader.

Let

$$\Psi(X, Y) = \phi(X - Y) a(X).$$

By Leibniz rule we get

$$\begin{aligned} |D_X^\alpha D_Y^\beta \Psi(X, Y)| & \leq \sum_{\gamma \leq \alpha} \binom{\alpha}{\gamma} |\phi^{(\alpha+\beta-\gamma)}(X - Y) a^{(\gamma)}(X)| \\ & \lesssim 2^{|\alpha|} \sup_{\gamma \leq \alpha} \left(h^{|\alpha+\beta|} ((\alpha + \beta - \gamma)! \gamma!)^s e^{-|X-Y|^{\frac{1}{s}}/h} \omega(X) \right) \\ & \leq (2^{1+s} h)^{|\alpha+\beta|} (\alpha! \beta!)^s e^{-|X-Y|^{\frac{1}{s}}/h} \omega(X) \lesssim (2^{1+s} h)^{|\alpha+\beta|} (\alpha! \beta!)^s e^{-|X-Y|^{\frac{1}{s}}/(2h)} \omega(Y), \end{aligned}$$

for every $h > 0$ which is chosen small enough. Here we have used the fact that for some $c > 0$

$$\omega(X) \lesssim \omega(Y) e^{c|X-Y|} \lesssim \omega(Y) e^{|X-Y|^{\frac{1}{s}}/(2h)},$$

since ω is a moderate function. This gives (2.7).

Next we prove (2). Let

$$\begin{aligned} \Omega_{1,Y} &= \left\{ X \mapsto \frac{D_Y^\beta (\phi_Y a)}{h^{|\beta|} \beta!^s \omega(Y)}; \beta \in \mathbf{N}^d \right\} \\ \Omega_{2,Z} &= \left\{ X \mapsto \frac{D_Z^\gamma (\psi_Z b)}{h^{|\gamma|} \gamma!^s \vartheta(Z)}; \gamma \in \mathbf{N}^d \right\}, \end{aligned}$$

and let $\Omega_1^\cup$ and $\Omega_2^\cup$ be as in (2.4). By (2.7) it follows that $\Omega_1^\cup$ and $\Omega_2^\cup$ are bounded in $\Sigma_s(\mathbf{R}^{2d})$. Hence, Lemma 2.5 shows that

$$\left| D_X^\alpha \left(\left(\frac{D_Y^\beta(\phi_Y a)}{h^{|\beta|} \beta! s \omega(Y)} \right) \# \left(\frac{D_Z^\gamma(\psi_Z b)}{h^{|\gamma|} \gamma! s \vartheta(Z)} \right) \right) (X) \right| \lesssim h^{|\alpha|} \alpha! s e^{-(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}} + |Y-Z|^{\frac{1}{s}})/h}$$

for every $h > 0$, or equivalently,

$$|D_X^\alpha D_Y^\beta D_Z^\gamma((\phi_Y a) \# (\psi_Z b))(X)| \lesssim h^{|\alpha| + |\beta| + |\gamma|} (\alpha! \beta! \gamma!)^s e^{-(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}} + |Y-Z|^{\frac{1}{s}})/h} \omega(Y) \vartheta(Z).$$

The assertion now follows from the latter estimate and the fact that ω and ϑ are moderate weights, giving that

$$\omega(Y) \lesssim \omega(X) e^{|X-Y|^{\frac{1}{s}}/(2h)} \lesssim \omega(Z) e^{(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}})/(2h)},$$

and similarly for ϑ .

Finally, the estimate in (2) also holds for $(\phi_Y b) \# (\psi_Z a)$ in place of $(\phi_Y a) \# (\psi_Z b)$, and (2.8) follows from this estimate by letting $\gamma = 0$, $b = 1$, $\vartheta = 1$, and then integrate with respect to Z . The proof is complete. $\square$

Lemmas 2.5 and 2.6 imply the following characterisation of $\Gamma_s^{(\omega)}(\mathbf{R}^{2d})$.

Proposition 2.7. *Let $s > 1/2$, $\omega \in \mathcal{P}_E(\mathbf{R}^{2d})$, $a \in \Sigma'_1(\mathbf{R}^{2d})$, $\phi \in \Sigma_s(\mathbf{R}^{2d})$ have non-vanishing integrals, and let $\phi_Y = \phi(\cdot - Y)$. Then the following conditions are equivalent:*

- (1) $a \in \Gamma_s^{(\omega)}$;
- (2) $\phi_Y a$ is smooth and satisfies (2.7) for some $h_1 > 0$ and every $h_2 > 0$;
- (3) $\phi_Y \# a$ is smooth and satisfies (2.8) for some $h_1 > 0$ and every $h_2 > 0$;
- (4)

$$|D_X^\alpha(\phi_Y a)(X)| \lesssim h_1^{|\alpha|} \alpha! s e^{-|X-Y|^{\frac{1}{s}}/h_1} \min(\omega(X), \omega(Y)), \quad (2.9)$$

for some $h_1 > 0$;

(5)

$$|D_X^\alpha(\phi_Y \# a)(X)| \lesssim h_1^{|\alpha|} \alpha! s e^{-|X-Y|^{\frac{1}{s}}/h_1} \min(\omega(X), \omega(Y)), \quad (2.10)$$

for some $h_1 > 0$.

Lemma 2.8. *Let $s \geq \frac{1}{2}$. Assume that $\{a_1(\cdot + Y, Y)\}_{Y \in \mathbf{R}^{2d}}$ and $\{a_2(\cdot + Z, Z)\}_{Z \in \mathbf{R}^{2d}}$ are bounded families in $\mathcal{S}_s(\mathbf{R}^{2d})$. Then, for any $X, Y, Z \in \mathbf{R}^{2d}$, $\alpha \in \mathbf{N}^{2d}$,*

$$|D_X^\alpha(a_1(\cdot + Y, Y) \# a_2(\cdot + Z, Z))(X)| \lesssim h^{|\alpha|} \alpha! s e^{-r(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}})}, \quad (2.11)$$

for some $h > 0$.

Proof. The result is a straightforward consequence of Corollary 2.3 and its proof, since the involved constants on the right hand side of (2.3) depend continuously on ϕ and ψ in $\mathcal{S}_s(\mathbf{R}^{2d})$. $\square$

We notice that, by straightforward computation, for some other $h, r > 0$, (2.11) gives, for any $X, Y, Z \in \mathbf{R}^{2d}$, $\alpha \in \mathbf{N}^{2d}$,

$$|D_X^\alpha(a_1(\cdot + Y, Y) \# a_2(\cdot + Z, Z))(X)| \lesssim h^{|\alpha|} \alpha! s e^{-r(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}} + |Y-Z|^{\frac{1}{s}})}. \quad (2.12)$$

2.1. A family related to $\Gamma_s^{(1)}$ and $\Gamma_{0,s}^{(1)}$. Let $I_R = [-R, R]$ and $E^0 = E_{h,s}^0 = L^\infty(I_R \times \mathbf{R}^{2d}; s_\infty^w(\mathbf{R}^{2d}))$, with the symbol subspace $s_\infty^w(\mathbf{R}^{2d})$ from Definition 1.22. We shall consider suitable decreasing family $\{E_{h,s}^n\}_{n=0}^\infty$ of Banach spaces. To this aim, let

$$G_n = \{(Y, T_1, \dots, T_n) \in \mathbf{R}^{2d(n+1)} : Y, T_j \in \mathbf{R}^{2d} \text{ with } |T_j| \leq 1, j = 1, \dots, n\}, n \in \mathbf{Z}_+.$$

We define $E_{h,s}^n$, $n \geq 1$, as the set of all $a \in E^0$ such that

$$\|a\|^{(n)} = \sup_{1 \leq k \leq n} \sup_{t \in I_R} \sup_{(Y, T_1, \dots, T_k) \in G_k} \frac{\|\langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle a(t, Y, \cdot)\|_{s_\infty^w}}{h^k(k!)^s} < \infty,$$

with the norm

$$\|a\|_{E_{h,s}^n} \equiv \|a\|_{E^0} + \|a\|^{(n)}.$$

We also let $E_{h,s}^\infty$ be the set of all

$$a \in \bigcap_{n \geq 0} E_{h,s}^n$$

such that

$$\|a\|_{E_{h,s}^\infty} \equiv \sup_{n \geq 0} \|a\|_{E_{h,s}^n}$$

is finite.

Lemma 2.9. *Let $n \geq 0$. Then $E_{h,s}^n$ is a Banach space.*

Proof. Let $\{a_j\}_{j \geq 0}$ be a Cauchy sequence in $E_{h,s}^n$, $n \geq 1$. By definition, this sequence clearly has a limit $a \in E^0$, and for some $X \mapsto b_k(t, Y, T_1, \dots, T_k, X) \in s_\infty^w(\mathbf{R}^{2d})$ we have

$$\lim_{j \rightarrow \infty} \sup \frac{\|\langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle a_j(t, Y, \cdot) - b_k(t, Y, T_1, \dots, T_k, \cdot)\|_{s_\infty^w}}{h^k(k!)^s} = 0,$$

where the supremum is taken over all

$$k \in \{1, \dots, n\}, \quad t \in I_R \quad \text{and} \quad (Y, T_1, \dots, T_k) \in G_k.$$

We need to prove that $a \in E_{h,s}^n$, and $a_j \rightarrow a$ in $E_{h,s}^n$.

The conditions here above are equivalent to

$$\lim_{j \rightarrow \infty} \sup_{t \in I_R} \sup_{Y \in \mathbf{R}^{2d}} \|a_j(t, Y, \cdot) - a(t, Y, \cdot)\|_{s_\infty^w} = 0 \quad (2.13)$$

and

$$\lim_{j \rightarrow \infty} \sup \frac{\|(-1)^k \langle T_1, D \rangle \cdots \langle T_k, D \rangle a_j(t, Y, \cdot) - b_k(t, Y, T_1, \dots, T_k, \cdot)\|_{s_\infty^w}}{h^k(k!)^s} = 0, \quad (2.14)$$

where the latter supremum should be taken over all

$$k \in \{1, \dots, n\}, \quad t \in I_R \quad \text{and} \quad (Y, T_1, \dots, T_k) \in G_k.$$

Since $s_\infty^w(\mathbf{R}^{2d})$ is continuously embedded in $\mathcal{S}'(\mathbf{R}^{2d})$, it follows from (2.13) and (2.14) that

$$X \mapsto (-1)^k \langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle a_j(t, Y, X)$$

has the limit

$$X \mapsto (-1)^k \langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle a(t, Y, X)$$

in $\mathcal{S}'(\mathbf{R}^{2d})$, and the limit

$$X \mapsto b_k(t, Y, T_1, \dots, T_k, X)$$

in $s_\infty^w(\mathbf{R}^{2d})$, and thereby in $\mathcal{S}'(\mathbf{R}^{2d})$, as j tends to ∞ . Hence

$$b_k(t, Y, T_1, \dots, T_k, X) = (-1)^k \langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle a(t, Y, X)$$

and it follows that $E_{h,s}^n$ is a Banach space for every $h > 0$, $s > 0$ and integer $n \geq 0$. $\square$

The spaces $E_{h,s}^\infty$ can be related to $\Gamma_s^{(1)}$ and $\Gamma_{0,s}^{(1)}$, as the following lemma shows. The details are left for the reader.

Lemma 2.10. *Let $a \in L^\infty(I_R \times \mathbf{R}^{2d}; s_\infty^w(\mathbf{R}^{2d}))$. Then $\{a(t, Y, \cdot)\}_{t \in I_R, Y \in \mathbf{R}^{2d}}$ is a uniformly bounded family in $\Gamma_s^{(1)}(\mathbf{R}^{2d})$ ($\Gamma_{0,s}^{(1)}(\mathbf{R}^{2d})$), if and only if*

$$\|a\|_{E_{h,s}^\infty} < \infty$$

for some $h > 0$ (for every $h > 0$).

Later on we also need the following result of differential equations with functions depending on a real variable with values in $E_{h,s}^\infty$. The proof is omitted since the result can be considered as a part of the standard theory of ordinary differential equations of first order in Banach spaces.

Lemma 2.11. *Suppose $s \geq 0$ and $T > 0$, and let $\mathcal{K}$ be an operator from $E_{h,s}^\infty$ to $E_{h,s}^\infty$ for every $h > 0$ such that*

$$\|\mathcal{K}a\|_{E_{h,s}^\infty} \leq C\|a\|_{E_{h,s}^\infty}, \quad a \in E_{h,s}^\infty, \quad (2.15)$$

for some constant C which only depend on $h > 0$. Then

$$\frac{dc(t)}{dt} = \mathcal{K}(c(t)), \quad c(0) \in E_{h,s}^\infty,$$

has a unique solution $t \mapsto c(t)$ from $[-T, T]$ to $E_{h,s}^\infty$ which satisfies

$$\|c(t)\|_{E_{h,s}^\infty} \leq \|c(0)\|_{E_{h,s}^\infty} e^{CT},$$

where C is the same as in (2.15).

3. ONE-PARAMETER GROUP OF ELLIPTIC SYMBOLS IN THE CLASSES $\Gamma_s^{(\omega)}(\mathbf{R}^d)$

In this section we show that for suitable s and ω_0 , there are elements $a \in \Gamma_s^{(\omega_0)}$ and $b \in \Gamma_s^{(1/\omega_0)}$ such that $a \# b = b \# a = 1$. This is essentially a consequence of Theorem 3.8, where it is proved that the evolution equation (0.4) has a unique solution $a(t, \cdot)$ which belongs to $\Gamma_s^{(\omega \vartheta^t)}$, thereby deducing needed semigroup properties for scales of pseudo-differential operators. Similar facts hold for corresponding Beurling type spaces (cf. Theorem 3.9).

First we have the following result on certain logarithms of weight functions.

Theorem 3.1. *Let $\omega \in \mathcal{P}_E(\mathbf{R}^{2d}) \cap \Gamma_{s_0}^{(\omega)}(\mathbf{R}^{2d})$, $s_0 \in (0, 1]$, $v \in \mathcal{P}_E(\mathbf{R}^{2d})$, be such that ω is v -moderate, $\vartheta(X) = 1 + \log v(X)$ and let*

$$c(X, Y) = \log \frac{\omega(X + Y)}{\omega(Y)}.$$

Then,

- $\{c(\cdot, Y)\}_{Y \in \mathbf{R}^{2d}}$ is a uniformly bounded family in $\Gamma_s^{(\vartheta)}(\mathbf{R}^{2d})$, $s \geq 1$;
- for $\alpha \neq 0$, $\{(\partial_X^\alpha c)(\cdot, Y)\}_{Y \in \mathbf{R}^{2d}}$ is a uniformly bounded family in $\Gamma_s^{(1)}(\mathbf{R}^{2d})$, $s \geq 1$.

For the proof of Theorem 3.1 we will need the following multidimensional version of the well-known Faà di Bruno formula for the derivatives of composed functions. It can be found, e.g., setting $q = p = 1$, $n = 2d$, in equations (2.3) and (2.4) in [29].

Lemma 3.2. *Let $f: \mathbf{R} \rightarrow \mathbf{R}$, $g: \mathbf{R}^{2d} \rightarrow \mathbf{R}$. Then, for any $\alpha \in \mathbf{N}^{2d}$, $\alpha \neq 0$,*

$$\frac{\partial^\alpha f(g(x))}{\alpha!} = \sum_{1 \leq k \leq |\alpha|} \frac{f^{(k)}(g(x))}{k!} \sum_{\substack{\beta_1 + \dots + \beta_k = \alpha \\ \beta_j \neq 0, j=1, \dots, k}} \prod_{1 \leq j \leq k} \frac{(\partial^{\beta_j} g)(x)}{\beta_j!}. \quad (3.1)$$

We will also need the next *factorial estimate*, for expressions involving decompositions of $\alpha \in \mathbf{N}^{2d}$, $\alpha \neq 0$, into the sum of k nontrivial multiindices β_j , $j = 1, \dots, k$, as in (3.1), and corresponding products of (powers of) factorials.

Lemma 3.3. *Let $s_0 \in (0, 1]$, $\alpha \in \mathbf{N}^{2d}$, $\alpha \neq 0$. Then, for a suitable $C_0 > 0$, depending only in d ,*

$$\sum_{1 \leq k \leq |\alpha|} \frac{1}{k} \sum_{\substack{\beta_1 + \dots + \beta_k = \alpha \\ \beta_j \neq 0, j=1, \dots, k}} \prod_{1 \leq j \leq k} (\beta_j!)^{s_0-1} \lesssim C_0^{|\alpha|}. \quad (3.2)$$

The proof of Lemma 3.3 can be found in the Appendix.

Proof of Theorem 3.1. We have to show that $c(\cdot, Y)$ satisfies $\Gamma_s^{(\vartheta)}$ estimates, uniformly with respect to $Y \in \mathbf{R}^{2d}$. If $c(X, Y) \geq 0$, then it follows by submultiplicativity of ω , that

$$\begin{aligned} c(X, Y) &= \log \omega(Y + X) - \log \omega(Y) \lesssim \log \omega(Y) + \log v(X) - \log \omega(Y) \\ &\leq \vartheta(X), \end{aligned}$$

for any $Y \in \mathbf{R}^{2d}$. Again by moderateness, when $c(X, Y) \leq 0$, recall that $\omega(X+Y) \geq \frac{\omega(Y)}{v(X)}$, so that

$$c(X, Y) \gtrsim \log \frac{\omega(Y)}{v(X)} - \log \omega(Y) \geq -\log v(X) \geq -\vartheta(X),$$

and we can conclude $|c(X, Y)| \lesssim \vartheta(X)$, $X \in \mathbf{R}^{2d}$. Now, for $\alpha \in \mathbf{N}^{2d}$, $\alpha \neq 0$, (3.1) gives

$$\partial_X^\alpha c(X, Y) = \alpha! \sum_{1 \leq k \leq |\alpha|} \frac{(-1)^{k+1}}{k [\omega(X+Y)]^k} \sum_{\substack{\beta_1 + \dots + \beta_k = \alpha \\ \beta_j \neq 0, j=1, \dots, k}} \prod_{1 \leq j \leq k} \frac{(\partial^{\beta_j} \omega)(X+Y)}{\beta_j!}$$

We can then estimate, in view of (3.2),

$$\begin{aligned} |\partial_X^\alpha c(X, Y)| &\lesssim \alpha! \sum_{1 \leq k \leq |\alpha|} \frac{1}{k [\omega(X+Y)]^k} \sum_{\substack{\beta_1 + \dots + \beta_k = \alpha \\ \beta_j \neq 0, j=1, \dots, k}} \prod_{1 \leq j \leq k} \frac{\omega(X+Y) h^{|\beta_j|} (\beta_j!)^{s_0}}{\beta_j!} \\ &= h^{|\alpha|} \alpha! \sum_{1 \leq k \leq |\alpha|} \frac{1}{k} \sum_{\substack{\beta_1 + \dots + \beta_k = \alpha \\ \beta_j \neq 0, j=1, \dots, k}} \prod_{1 \leq j \leq k} (\beta_j!)^{s_0-1} \lesssim (C_0 h)^{|\alpha|} (\alpha!)^s, \end{aligned}$$

which concludes the proof. $\square$

Proposition 3.4. *Assume $s > \frac{1}{2}$ and $\omega(X) \lesssim e^{r|X|^{\frac{1}{s}}}$ for some $r > 0$. Let $\{a(\cdot, Y)\}_{Y \in \mathbf{R}^{2d}}$ be a uniformly bounded family in $\Sigma_s(\mathbf{R}^{2d})$ and $\{c(\cdot, Z)\}_{Z \in \mathbf{R}^{2d}}$ be a bounded family in $\Gamma_s^{(\omega)}(\mathbf{R}^{2d})$. Then,*

$$\{a(\cdot, Y) \# c(\cdot, Z)\}_{Y, Z \in \mathbf{R}^{2d}} \text{ and } \{c(\cdot, Z) \# a(\cdot, Y)\}_{Y, Z \in \mathbf{R}^{2d}}$$

are bounded families in $\mathcal{S}_s(\mathbf{R}^{2d})$.

Proof. An immediate consequence of (2.8) in Lemma 2.6 is the following

$$|D_X^\alpha(\phi \# a)(X)| \leq Ch^{|\alpha|} \alpha!^s e^{-r|X|^\frac{1}{s}}, \quad (3.3)$$

which implies that, for $\phi \in \Sigma_s$ and $a \in \Gamma_s^{(\omega)}$, (3.3) holds if and only if $\phi \# a$ belongs to $\mathcal{S}_s$. Then by the proof of (3.3) it follows that the constant C, h and r can be chosen depending continuously on $\phi \in \Sigma_s(\mathbf{R}^{2d})$ and $a \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$. Hence if Ω_1 is a bounded family in $\Sigma_s(\mathbf{R}^{2d})$ and Ω_2 is a bounded family in $\Gamma_s^{(\omega)}(\mathbf{R}^{2d})$, then $\{\phi \# a\}_{\phi \in \Omega_1, a \in \Omega_2}$ is a bounded family in $\mathcal{S}_s(\mathbf{R}^{2d})$. $\square$

The following result can be found, e.g., in [46].

Lemma 3.5. *Let $a \in \mathcal{S}'(\mathbf{R}^{2d})$. Then,*

$$\|a\|_{s_\infty^w} \leq C \sum_{|\alpha| \leq d+1} \|\partial^\alpha a\|_{L^\infty} \quad (3.4)$$

and

$$\|a\|_{L^\infty} \leq C \sum_{|\alpha| \leq 2d+1} \|\partial^\alpha a\|_{s_\infty^w}, \quad (3.5)$$

for some constant $C > 0$ depending only on the dimension d .

Proposition 3.6. *Let $a \in \mathcal{S}'(\mathbf{R}^{2d})$, $s \geq \frac{1}{2}$ and set $b_{\alpha\beta}(X) = \partial^\alpha(X^\beta a(X))$ when $\alpha, \beta \in \mathbf{N}^{2d}$. Then the following conditions are equivalent:*

- (1) $a \in \mathcal{S}_s(\mathbf{R}^{2d})$;
- (2) for some $h > 0$ it holds

$$\|b_{\alpha\beta}\|_{L^\infty} \lesssim h^{|\alpha+\beta|} (\alpha! \beta!)^s, \quad \alpha, \beta \in \mathbf{N}^{2d};$$

- (3) for some $h > 0$ it holds

$$\|b_{\alpha\beta}\|_{s_\infty^w} \lesssim h^{|\alpha+\beta|} (\alpha! \beta!)^s, \quad \alpha, \beta \in \mathbf{N}^{2d}.$$

Proof. The equivalence between (1) and (2) is well-known. The proof of the equivalence of (2) and (3) follows by a straightforward application of Lemma 3.5. In fact, assume that (2) holds true. Then (3.4) gives

$$\begin{aligned} \|b_{\alpha\beta}\|_{s_\infty^w} &\leq C \sum_{|\gamma| \leq d+1} \|\partial^\gamma b_{\alpha\beta}\|_{L^\infty(\mathbf{R}^{2d})} \lesssim \sum_{|\gamma| \leq d+1} h^{|\alpha+\beta+\gamma|} ((\alpha+\gamma)! \beta!)^s \\ &= h^{|\alpha+\beta|} (\alpha! \beta!)^s \sum_{|\gamma| \leq d+1} h^{|\gamma|} (\gamma!)^s \left(\frac{(\alpha+\gamma)!}{\alpha! \gamma!} \right)^s \\ &\lesssim (2^s h)^{|\alpha+\beta|} (\alpha! \beta!)^s, \end{aligned}$$

with a constant depending only on d and h , since

$$\sum_{|\gamma| \leq d+1} h^{|\gamma|} (\gamma!)^s \left(\frac{(\alpha+\gamma)!}{\alpha! \gamma!} \right)^s \leq C_1 \cdot 2^{s(|\alpha|+d+1)} \leq C_2 2^{s|\alpha+\beta|},$$

where the constants C_1 and C_2 only depend on d and h . Then, (3) holds true, as claimed. The proof of the converse follows by similar argument, employing (3.5). $\square$

We have now the following.

Theorem 3.7. *Let $a \in \mathcal{S}'(\mathbf{R}^{2d})$ and $s > 0$. Then the following conditions are equivalent:*

- (1) $a \in \Gamma_s^{(1)}(\mathbf{R}^{2d})$;

(2) *there exists $h > 0$ such that*

$$\|\partial^\alpha a\|_{L^\infty(\mathbf{R}^{2d})} \lesssim h^{|\alpha|}(\alpha!)^s,$$

for all $\alpha \in \mathbf{Z}_+^{2d}$;

(3) *there exists $h > 0$ such that*

$$\|\partial^\alpha a\|_{s_\infty^w} \lesssim h^{|\alpha|}(\alpha!)^s,$$

for all $\alpha \in \mathbf{Z}_+^{2d}$;

(4) *there exists $h > 0$ such that*

$$\|\langle T_1, D_X \rangle \cdots \langle T_m, D_X \rangle a\|_{s_\infty^w} \lesssim h^m(m!)^s,$$

for any $T_1, \dots, T_m \in \mathbf{R}^{2d}$ such that $|T_j| \leq 1$, $j = 1, \dots, m$, $m \geq 1$.

Proof. The equivalence between (1) and (2) is well known. The equivalence of (2) and (3) is proved by an argument completely similar to the one employed in the proof of Proposition 3.6, using Lemma 3.5. It remains to prove only the equivalence with (4). Assume that (3) holds true, and let

$$T_j = \sum_{l=1}^d t_{jl} \mathbf{e}_l + \sum_{l=1}^d \tau_{jl} \varepsilon_l,$$

for the standard basis $(\mathbf{e}_l)_{l=1, \dots, d}$ of $\mathbf{R}^d$ and the dual basis $(\varepsilon_l)_{l=1, \dots, d}$. Recall that

$$\langle T_k, D_X \rangle a = \sum_{l=1}^d t_{kl} \frac{\partial a}{\partial x_l} + \sum_{l=1}^d \tau_{kl} \frac{\partial a}{\partial \xi_l},$$

so that the symbol $\langle T_1, D_X \rangle \cdots \langle T_m, D_X \rangle a$ is in the span of symbols of the form

$$\left(\prod_{j=1}^m t_j^{\beta_j} \tau_j^{\gamma_j} \right) \partial_x^\beta \partial_\xi^\gamma a, \quad \sum_{j=1}^m \beta_j = \beta, \sum_{j=1}^m \gamma_j = \gamma, |\beta| + |\gamma| = m,$$

where the summation contains at most $(2d)^m$ terms. Since $|T_j| \leq 1$, $j = 1, \dots, m$, by (3.4) we find

$$\begin{aligned} \|\langle T_1, D_X \rangle \cdots \langle T_m, D_X \rangle a\|_{s_\infty^w} &\leq (2d)^m \sup_{|\alpha|=m} \|\partial^\alpha a\|_{s_\infty^w} \\ &\lesssim \sup_{|\alpha|=m} \sum_{|\gamma| \leq d+1} h^{|\alpha+\gamma|} ((\alpha+\gamma)!)^s \\ &= \sup_{|\alpha|=m} h^{|\alpha|} (\alpha!)^s \sum_{|\gamma| \leq d+1} h^{|\gamma|} (\gamma!)^s \left(\frac{(\alpha+\gamma)!}{\alpha! \gamma!} \right)^s \\ &\lesssim (2^{s+1} h)^m (m!)^s, \end{aligned}$$

which proves that (4) holds true. As in the proof of Proposition 3.6, the converse implication is obtained by a completely similar argument, employing (3.5). $\square$

We are now ready to state and prove the first main result of this section, the following Theorem 3.8. It deals with the existence of one-parameter groups of pseudo-differential operators, obtained as solutions to suitable evolution equations.

Theorem 3.8. *Let $s \geq 1$, $\omega, \vartheta \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$ be such that $\omega \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$ and $\vartheta \in \Gamma_s^{(\vartheta)}(\mathbf{R}^{2d})$, and let $a_0 \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$, $b \in \Gamma_s^{(1)}(\mathbf{R}^{2d})$. Then, there exists a unique smooth map $(t, X) \mapsto a(t, X) \in \mathbf{C}$ such that $a(t, \cdot) \in \Gamma_s^{(\omega, \vartheta^t)}(\mathbf{R}^{2d})$ for all $t \in \mathbf{R}$, and*

$$\begin{cases} (\partial_t a)(t, \cdot) = (b + \log \vartheta) \# a(t, \cdot) \\ a(0, \cdot) = a_0. \end{cases} \quad (3.6)$$

If $\omega \equiv a_0 \equiv 1$, then $a(t, X)$ also satisfies

$$\begin{cases} (\partial_t a)(t, \cdot) = a(t, \cdot) \# (b + \log \vartheta) \\ a(0, \cdot) = a_0, \end{cases} \quad (3.7)$$

and

$$a(t_1, \cdot) \# a(t_2, \cdot) = a(t_1 + t_2, \cdot), \quad a(t, \cdot) \in \Gamma_s^{(\vartheta^t)}(\mathbf{R}^{2d}), \quad t, t_1, t_2 \in \mathbf{R}. \quad (3.8)$$

Proof. First suppose that a solution $a(t, X)$ of (3.6) exists. Then,

$$a(t, X) = a_0(X) + \int_0^t c(u, X) du$$

with

$$c(t, \cdot) = (b + \log \vartheta) \# a(t, \cdot) \in \Gamma_s^{(\omega(\log \vartheta) \vartheta^t)}(\mathbf{R}^{2d}),$$

in view of Theorem 3.1 and the properties of the Weyl product in the $\Gamma_s^{(\omega)}(\mathbf{R}^{2d})$ classes, see [4]. This implies that the map $t \mapsto a(t, \cdot)$ is C^1 from $[-R, R]$ into the symbol space

$$\Gamma_s^{(\omega(\log \vartheta)(\vartheta + \vartheta^{-1})^R)}(\mathbf{R}^{2d}).$$

Choose $s_0 < s$, and $\phi, \psi \in \mathcal{S}_{s_0}(\mathbf{R}^{2d})$ such that (2.2) holds true. Let

$$c_1(t, Y, \cdot) = \omega(Y)^{-1} \vartheta(Y)^{-t} \phi_Y \# a(t, \cdot). \quad (3.9)$$

By Lemma 2.6 (1) we have that, for any $Y \in \mathbf{R}^{2d}$, $t \mapsto c_1(t, Y, \cdot)$ is a C^1 map from $[-R, R]$ into $\mathcal{S}_s(\mathbf{R}^{2d})$. Moreover,

$$\partial_t c_1(t, Y, \cdot) = \omega(Y)^{-1} \vartheta(Y)^{-t} \phi_Y \# (b + \log \vartheta) \# a(t, \cdot) - c_1(t, Y, \cdot) \log \vartheta(Y).$$

Let

$$f(Y, X) = b(X) + \log \frac{\vartheta(X)}{\vartheta(Y)}.$$

Then,

$$\begin{aligned} (\partial_t c_1)(t, Y, \cdot) &= \omega(Y)^{-1} \vartheta(Y)^t \int \phi_Y \# f(Y, \cdot) \# \psi_Z \# \phi_Z \# a(t, \cdot) dZ \\ &= \int K_{Y,Z}(t, \cdot) \# c_1(t, Z, \cdot) dZ \end{aligned}$$

with

$$K_{Y,Z}(t, \cdot) = \frac{\omega(Z) \vartheta(Z)^t}{\omega(Y) \vartheta(Y)^t} \phi_Y \# f(Y, \cdot) \# \psi_Z, \quad (3.10)$$

and

$$c_1(0, Y, \cdot) = \omega(Y)^{-1} \phi_Y \# a_0.$$

We also need to consider the similar equation where $f(Y, \cdot)$ is replaced by $f(Z, \cdot)$, that is

$$\partial_t c_2(t, Y, \cdot) = \int \tilde{K}_{Y,Z}(t, \cdot) \# c_2(t, Z, \cdot) dZ, \quad (3.11)$$

where

$$\tilde{K}_{Y,Z}(t, \cdot) = \frac{\omega(Z) \vartheta(Z)^t}{\omega(Y) \vartheta(Y)^t} \phi_Y \# f(Z, \cdot) \# \psi_Z,$$

and

$$c_2(0, Y, \cdot) = c_1(0, Y, \cdot) = \omega(Y) \phi_Y \# a_0. \quad (3.12)$$

We consider the operators $\mathcal{K}$ and $\tilde{\mathcal{K}}$ on E^0 , defined by

$$(\mathcal{K}a)(t, Y, X) = \int (K_{Y,Z}(t, \cdot) \# a(t, Z, \cdot))(X) dZ,$$

and

$$(\tilde{\mathcal{K}}a)(t, Y, X) = \int (\tilde{K}_{Y,Z}(t, \cdot) \# a(t, Z, \cdot))(X) dZ,$$

and show that

$$\begin{aligned} \|\mathcal{K}a\|_{E_{h,s}^n} &\leq C(n+1)\|a\|_{E_{h,s}^n}, & \|\mathcal{K}a\|_{E_{h,s}^\infty} &\leq C\|a\|_{E_{h,s}^\infty}, \\ \|\tilde{\mathcal{K}}a\|_{E_{h,s}^n} &\leq C(n+1)\|a\|_{E_{h,s}^n} \quad \text{and} \quad \|\tilde{\mathcal{K}}a\|_{E_{h,s}^\infty} &\leq C\|a\|_{E_{h,s}^\infty} \end{aligned} \quad (3.13)$$

for some constant C , which is independent of h , n and s .

In order to prove this, it is convenient to let $\mathcal{P}_k$ be the family of all subsets of $\{1, \dots, k\}$, $k \geq 1$. For each $P \in \mathcal{P}_k$, $a \in s_\infty^w(\mathbf{R}^{2d})$, we set

$$H(a, P) = \begin{cases} a & \text{when } P = \emptyset, \\ \langle T_{j_1}, D_X \rangle \cdots \langle T_{j_l}, D_X \rangle a & \text{when } P = \{j_1 < \cdots < j_l\}, \quad l \leq k. \end{cases}$$

We now estimate

$$\frac{\|(\langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle \mathcal{K}a)(t, Y, \cdot)\|_{s_\infty^w(\mathbf{R}^{2d})}}{h^k(k!)^s}.$$

when $a \in E_{h,s}^n$. Since

$$\begin{aligned} &(\langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle \mathcal{K}a)(t, Y, X) \\ &= \langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle \int (K_{Y,Z}(t, \cdot) \# a(t, Z, \cdot))(X) dZ \\ &= \sum_{P \in \mathcal{P}_k} \int (H(K_{Y,Z}(t, \cdot), P) \# H(a(t, Z, \cdot), P^c))(X) dZ, \end{aligned}$$

we find

$$\begin{aligned} &\frac{\|(\langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle \mathcal{K}a)(t, Y, \cdot)\|_{s_\infty^w}}{h^k(k!)^s} \\ &\leq \sum_{l=0}^k \sum_{|P|=l} \binom{k}{l}^{-s} \int \frac{\|H(K_{Y,Z}(t, \cdot), P)\|_{s_\infty^w}}{h^l l!^s} \cdot \frac{\|H(a(t, Z, \cdot), P^c)\|_{s_\infty^w}}{h^{k-l}((k-l)!)^s} dZ \\ &\leq \sum_{l=0}^k \sum_{|P|=l} \|a\|_{E_{h,s}^{k-l}} \binom{k}{l}^{-s} \int \frac{\|H(K_{Y,Z}(t, \cdot), P)\|_{s_\infty^w}}{h^l l!^s} dZ \\ &\lesssim \|a\|_{E_{h,s}^n} \sum_{l=0}^k \sum_{|P|=l} \binom{k}{l}^{-1} \int \frac{\|H(K_{Y,Z}(t, \cdot), P)\|_{s_\infty^w}}{h^l l!^s} dZ, \quad (3.14) \end{aligned}$$

where the last inequality follows from the fact that $s \geq 1$ and $\|a\|_{E_{h,s}^n}$ increases with n .

We have now to estimate $\|H(K_{Y,Z}(t, \cdot), P)\|_{s_\infty^w}$, and study the different quantities on the right-hand side of (3.10). Since ω and ϑ belong to $\mathcal{P}_{E,s}^0$, it follows that for every $r > 0$,

$$\begin{aligned} \frac{\omega(Z) \vartheta(Z)^t}{\omega(Y) \vartheta(Y)^t} &= \frac{\omega(Z)}{\omega(Y)} \left(\frac{\vartheta(Z)}{\vartheta(Y)} \right)^t \lesssim e^{r|Y-Z|^{\frac{1}{s}}} \left(e^{r|Y-Z|^{\frac{1}{s}}} \right)^t \\ &\lesssim e^{r(1+t)|Y-Z|^{\frac{1}{s}}}, \quad Y, Z \in \mathbf{R}^{2d}. \quad (3.15) \end{aligned}$$

For the Weyl product in (3.10) we have

$$\begin{aligned}\phi_Y \# f(Y, \cdot) &= \phi(\cdot - Y) \# \left(b(\cdot) + \log \frac{\vartheta(\cdot)}{\vartheta(Y)} \right) \\ &= \left(\phi \# b(\cdot + Y) \right)_Y + \phi(\cdot - Y) \# \left(\log \frac{\vartheta(\cdot)}{\vartheta(Y)} \right) \\ &= \left(\phi \# b(\cdot + Y) \right)_Y + \left(\phi \# \log \frac{\vartheta(\cdot + Y)}{\vartheta(Y)} \right)_Y.\end{aligned}$$

By Theorem 3.1 and Proposition 3.4,

$$\left\{ \phi \# b(\cdot + Y) \right\}_{Y \in \mathbf{R}^{2d}} \quad \text{and} \quad \left\{ \phi \# \log \frac{\vartheta(\cdot + Y)}{\vartheta(Y)} \right\}_{Y \in \mathbf{R}^{2d}} \quad (3.16)$$

are uniformly bounded families in $\mathcal{S}_s(\mathbf{R}^{2d})$. Note that

$$a_2(Z, X) = \psi_Z(X) \quad \Rightarrow \quad \{a_2(Z, \cdot + Z)\}_{Z \in \mathbf{R}^{2d}} = \{\psi\}_{Z \in \mathbf{R}^{2d}},$$

which is evidently a uniformly bounded family in $\mathcal{S}_s(\mathbf{R}^{2d})$. Combining this last observation with the computations on $\phi_Y \# f(Y, \cdot)$ above, using Lemma 2.8 and (2.12), we finally obtain, for some $h, r_0 > 0$,

$$\begin{aligned}|D_X^\alpha (\phi_Y \# f(Y, \cdot) \# \psi_Z)(X)| &\lesssim h^{|\alpha|} \alpha!^s e^{-r_0(|X-Y|^{\frac{1}{s}} + |X-Z|^{\frac{1}{s}} + |Y-Z|^{\frac{1}{s}})}, \\ X, Y, Z &\in \mathbf{R}^{2d}, \alpha \in \mathbf{N}^{2d}.\end{aligned} \quad (3.17)$$

By Theorem 3.7, (3.15) and (3.17) we get for all $P \in \mathcal{P}_k$ $Y, Z \in \mathbf{R}^{2d}$ and some $r_0, h > 0$ that

$$\|H(K_{Y,Z}(t, \cdot), P)\|_{s_\infty^w} \leq C h^l l!^s e^{-r_0|Y-Z|^{\frac{1}{s}}}, \quad l = |P|,$$

where C is independent of k . Hence, (3.14) gives

$$\|\mathcal{K}a(t, Y, \cdot)\|_{s_\infty^w} \lesssim \|a\|_{E_{h,s}^n},$$

and

$$\begin{aligned}&\frac{\|\langle T_1, D_X \rangle \cdots \langle T_k, D_X \rangle \mathcal{K}a(t, Y, \cdot)\|_{s_\infty^w}}{h^k (k!)^s} \\ &\leq C_1 \|a\|_{E_{h,s}^n} \sum_{l=0}^k \sum_{|P|=l} \binom{k}{l}^{-1} \int e^{-r_0|Y-Z|^{\frac{1}{s}}} dZ \\ &= C_2 \|a\|_{E_{h,s}^n} \sum_{l=0}^k \sum_{|P|=l} \binom{k}{l}^{-1} = C_2 (k+1) \|a\|_{E_{h,s}^n}, \quad 1 \leq k \leq n,\end{aligned}$$

as claimed, where C_1 and C_2 are independent of k, n and h .

By a completely similar argument, an analogous result can be obtained for $\tilde{\mathcal{K}}$. In fact, by similar arguments that lead to (3.16) it follows that

$$\{b(\cdot + Z) \# \psi\}_{Z \in \mathbf{R}^{2d}} \quad \text{and} \quad \left\{ \log \frac{\vartheta(\cdot + Z)}{\vartheta(Z)} \# \psi \right\}_{Z \in \mathbf{R}^{2d}}$$

are bounded in $\mathcal{S}_s(\mathbf{R}^{2d})$, given that (3.17) holds with $f(Z, \cdot)$ in place of $f(Y, \cdot)$. This gives the first and third inequalities in (3.13). From these estimates we get

$$\|\mathcal{K}a\|_{E_{h,s}^{n+1}} \leq C \|a\|_{E_{h,s}^n} \quad \text{and} \quad \|\tilde{\mathcal{K}}a\|_{E_{h,s}^{n+1}} \leq C \|a\|_{E_{h,s}^n},$$

which give the other inequalities in (3.17).

We have proven that for any $T > 0$, then

$$\|\mathcal{K}_t\|_{E_{h,s}^n \rightarrow E_{h,s}^n} \leq C(n+1), \quad |t| \leq T,$$

where C is independent of n . As a consequence, since $\omega(Y)^{-1}\phi_Y\#a_0$ belongs to $E_{h,s}^\infty$, the equation

$$\frac{dc_1}{dt} = \mathcal{K}c_1, \quad c_1(0) = \omega(Y)^{-1}\phi_Y\#a_0 \quad (3.18)$$

has a unique solution on $\mathbf{R}$ belonging to $E_{h,s}^\infty$, in view of Lemma 2.11. By Proposition 2.7 it follows that $c_1(t, Y, \cdot) \in \Gamma_s^{(1)}(\mathbf{R}^{2d})$, uniformly in Y and for bounded t .

In order to prove the uniqueness of the solution a of (3.6), first we assume the existence and by what we have proven above i.e. that $c_1(t, Y, \cdot)$ in (3.9) satisfies (3.18) which implies the uniqueness of the solution of (3.6), since

$$a(t, \cdot) = \int_{\mathbf{R}^{2d}} \psi_Y\#\phi_Y\#a(t, \cdot) dY = \int_{\mathbf{R}^{2d}} \omega(Y)\vartheta(Y)^t \psi_Y\#c_1(t, Y, \cdot) dY. \quad (3.19)$$

To prove the existence of a solution of (3.6), we consider the solution $c_2(t, Y, \cdot)$ of (3.11) with the initial data (3.12), and we let

$$a(t, \cdot) = \int_{\mathbf{R}^{2d}} \omega(Y)\vartheta(Y)^t \psi_Y\#c_2(t, Y, \cdot) dY. \quad (3.20)$$

By Propositions 2.7 and 1.8, the family $\{\psi_Y\#c_2(t, Y, \cdot)\}_{Y \in \mathbf{R}^{2d}}$ belongs to $\mathcal{S}_s$ and $a(t, \cdot)$ belongs to $\Gamma_s^{(w\vartheta^t)}$. Moreover,

$$\begin{aligned} \frac{da(t, \cdot)}{dt} &= \int_{\mathbf{R}^{2d}} \omega(Y)\vartheta(Y)^t \log \vartheta(Y) \psi_Y\#c_2(t, Y, \cdot) dY \\ &\quad + \int_{\mathbf{R}^{2d}} \int_{\mathbf{R}^{2d}} \omega(Y)\vartheta(Y)^t \psi_Y\#\tilde{K}_{Y,Z}(t, \cdot)\#c_2(t, Z, \cdot) dY dZ \\ &= \int_{\mathbf{R}^{2d}} \omega(Z)\vartheta(Z)^t \log \vartheta(Z) \psi_Z\#c_2(t, Z, \cdot) dZ \\ &\quad + \int_{\mathbf{R}^{2d}} \int_{\mathbf{R}^{2d}} \omega(Z)\vartheta(Z)^t \psi_Y\#\phi_Y\#f(Z, \cdot)\#\psi_Z\#c_2(t, Z, \cdot) dY dZ \\ &= \int_{\mathbf{R}^{2d}} \omega(Z)\vartheta(Z)^t (b + \log \vartheta) \psi_Z\#c_2(t, Z, \cdot) dZ \\ &= (b + \log \vartheta)\#a(t, \cdot), \end{aligned}$$

with the initial data

$$a(0, \cdot) = \int_{\mathbf{R}^{2d}} \omega(Y)\psi_Y\#(\omega(Y)^{-1}\phi_Y\#a_0) dY = a_0,$$

which provide a solution of (3.6).

In order to prove the last part we consider the unique solution $a(t, \cdot)$ of (3.6) with the initial data $a(0, \cdot) \equiv 1$. If $\omega \equiv 1$, then for $u \in \mathbf{R}$ the mappings

$$t \mapsto a(t+u, \cdot) \quad \text{and} \quad t \mapsto a(t, \cdot)\#a(u, \cdot)$$

are both solutions of (3.6) with value $a(u, \cdot)$ at $t = 0$, and

$$a(t+u, \cdot) = a(t, \cdot)\#a(u, \cdot), \quad (3.21)$$

by the uniqueness property for the solution of (3.6).

Using (3.21) we have for all $t \in \mathbf{R}$, $a(t, \cdot)\#a(-t, \cdot) = 1$. Taking the derivative we get

$$0 = \frac{d}{dt}(a(t, \cdot)\#a(-t, \cdot)) = (b + \log \vartheta)\#a(t, \cdot)\#a(-t, \cdot) - a(t, \cdot)\#(b + \log \vartheta)\#a(-t, \cdot).$$

That is $(b + \log \vartheta) = a(t, \cdot)\#(b + \log \vartheta)\#a(-t, \cdot)$, implying the commutation for the sharp product of $a(t, \cdot)$ with $(b + \log \vartheta)$, and the result follows. $\square$

By similar argument as for the previous result we get the following.

Theorem 3.9. *Let $s \geq 1$, $\omega, \vartheta \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ be such that $\omega \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$ and $\vartheta \in \Gamma_s^{(\vartheta)}(\mathbf{R}^{2d})$, and let $a_0 \in \Gamma_{0,s}^{(\omega)}(\mathbf{R}^{2d})$, $b \in \Gamma_{0,s}^{(1)}(\mathbf{R}^{2d})$. Then, there exists a unique smooth map $(t, X) \mapsto a(t, X) \in \mathbf{C}$ such that $a(t, \cdot) \in \Gamma_s^{(\omega \vartheta^t)}(\mathbf{R}^{2d})$ for all $t \in \mathbf{R}$, and $a(t, \cdot)$ satisfies (3.6).*

Moreover, if $\omega \equiv a_0 \equiv 1$, then $a(t, X)$ also satisfies (3.7) and

$$a(t_1, \cdot) \# a(t_2, \cdot) = a(t_1 + t_2, \cdot), \quad a(t, \cdot) \in \Gamma_{0,s}^{(\vartheta^t)}(\mathbf{R}^{2d}), \quad t, t_1, t_2 \in \mathbf{R}.$$

4. LIFTING OF PSEUDO-DIFFERENTIAL OPERATORS AND TOEPLITZ OPERATORS ON MODULATION SPACES

In this section we apply the group properties in Theorems 3.8 and 3.9 to deduce lifting properties of pseudo-differential operators on modulation spaces. Thereafter we combine these results by the Wiener property of certain pseudo-differential operators with symbols in suitable modulation spaces to get lifting properties for Toeplitz operators with weights as their symbols.

We begin to apply Theorems 3.8 and 3.9 to get the following.

Theorem 4.1. *Let $s \geq 1$, $\mathbf{p} \in (0, \infty]^{2d}$, $A \in \mathbf{M}(d, \mathbf{R})$, $\omega \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$, and let $\mathcal{B}$ be an invariant BF-space on $\mathbf{R}^{2d}$, or $\mathcal{B} = L^{\mathbf{p},E}(\mathbf{R}^{2d})$ for some phase-shift split parallelepiped E in $\mathbf{R}^{2d}$. Then the following are true:*

- (1) *There exist $a \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$ and $b \in \Gamma_s^{(1/\omega)}(\mathbf{R}^{2d})$ such that*

$$\text{Op}_A(a) \circ \text{Op}_A(b) = \text{Op}_A(b) \circ \text{Op}_A(a) = \text{Id}_{S'_s(\mathbf{R}^d)}. \quad (4.1)$$

Furthermore, $\text{Op}_A(a)$ is an isomorphism from $M(\omega_0, \mathcal{B})$ onto $M(\omega_0/\omega, \mathcal{B})$, for every $\omega_0 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$.

- (2) *Let $a_0 \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$ be such that $\text{Op}_A(a_0)$ is an isomorphism from $M_{(\omega_1)}^2(\mathbf{R}^d)$ to $M_{(\omega_1/\omega)}^2(\mathbf{R}^d)$ for some $\omega_1 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$. Then $\text{Op}_A(a_0)$ is an isomorphism from $M(\omega_2, \mathcal{B})$ to $M(\omega_2/\omega, \mathcal{B})$, for every $\omega_2 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$. Furthermore, the inverse of $\text{Op}_A(a_0)$ is equal to $\text{Op}_A(b_0)$ for some $b_0 \in \Gamma_s^{(1/\omega)}(\mathbf{R}^{2d})$.*

Theorem 4.2. *Let $s > 1$, $\mathbf{p} \in (0, \infty]^{2d}$, $A \in \mathbf{M}(d, \mathbf{R})$, $\omega \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$, and let $\mathcal{B}$ be an invariant BF-space on $\mathbf{R}^{2d}$, or $\mathcal{B} = L^{\mathbf{p},E}(\mathbf{R}^{2d})$ for some phase-shift split parallelepiped E in $\mathbf{R}^{2d}$. Then the following are true:*

- (1) *There exist $a \in \Gamma_{0,s}^{(\omega)}(\mathbf{R}^{2d})$ and $b \in \Gamma_{0,s}^{(1/\omega)}(\mathbf{R}^{2d})$ such that*

$$\text{Op}_A(a) \circ \text{Op}_A(b) = \text{Op}_A(b) \circ \text{Op}_A(a) = \text{Id}_{\Sigma'_s(\mathbf{R}^d)}. \quad (4.2)$$

Furthermore, $\text{Op}_A(a)$ is an isomorphism from $M(\omega_0, \mathcal{B})$ onto $M(\omega_0/\omega, \mathcal{B})$, for every $\omega_0 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$.

- (2) *Let $a_0 \in \Gamma_{0,s}^{(\omega)}(\mathbf{R}^{2d})$ be such that $\text{Op}_A(a_0)$ is an isomorphism from $M_{(\omega_1)}^2(\mathbf{R}^d)$ to $M_{(\omega_1/\omega)}^2(\mathbf{R}^d)$ for some $\omega_1 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$. Then $\text{Op}_A(a_0)$ is an isomorphism from $M(\omega_2, \mathcal{B})$ to $M(\omega_2/\omega, \mathcal{B})$, for every $\omega_2 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$. Furthermore, the inverse of $\text{Op}_A(a_0)$ is equal to $\text{Op}_A(b_0)$ for some $b_0 \in \Gamma_{0,s}^{(1/\omega)}(\mathbf{R}^{2d})$.*

We only prove Theorem 4.2. Theorem 4.1 follows by similar arguments and is left for the reader.

Proof of Theorem 4.2. The existence of $a \in \Gamma_{0,s}^{(\omega)}(\mathbf{R}^{2d})$ and $b \in \Gamma_{0,s}^{(1/\omega)}(\mathbf{R}^{2d})$ such that (4.2) holds is guaranteed by Theorem 3.9. By [56, Theorems 2.3 and 2.6] it

follows that

$$\text{Op}_A(a) : M(\omega_0, \mathcal{B}) \rightarrow M(\omega_0/\omega, \mathcal{B}) \quad (4.3)$$

and

$$\text{Op}_A(b) : M(\omega_0/\omega, \mathcal{B}) \rightarrow M(\omega_0, \mathcal{B}) \quad (4.4)$$

are continuous. By (4.2) and the fact that $M(\omega_0, \mathcal{B})$ and $M(\omega_0/\omega, \mathcal{B})$ are contained in $\Sigma'_s(\mathbf{R}^{2d})$, it follows that (4.3) and (4.4) are homeomorphic, and (1) follows.

(2) It suffices to prove the result in the Weyl case, $A = \frac{1}{2}I$, in view of Proposition 1.25. By (1), we may find

$$a_1 \in \Gamma_{0,s}^{(\omega_1)}, \quad b_1 \in \Gamma_{0,s}^{(1/\omega_1)}, \quad a_2 \in \Gamma_{0,s}^{(\omega_1/\omega)}, \quad b_2 \in \Gamma_{0,s}^{(\omega/\omega_1)}$$

satisfying the following properties:

- $\text{Op}^w(a_j)$ and $\text{Op}^w(b_j)$ are inverses to each others on $\Sigma'_s(\mathbf{R}^d)$ for $j = 1, 2$;
- For arbitrary $\omega_2 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$, the mappings

$$\begin{aligned} \text{Op}^w(a_1) &: M_{(\omega_2)}^2 \rightarrow M_{(\omega_2/\omega_1)}^2, \\ \text{Op}^w(b_1) &: M_{(\omega_2)}^2 \rightarrow M_{(\omega_2\omega_1)}^2, \\ \text{Op}^w(a_2) &: M_{(\omega_2)}^2 \rightarrow M_{(\omega_2\omega/\omega_1)}^2, \\ \text{Op}^w(b_2) &: M_{(\omega_2)}^2 \rightarrow M_{(\omega_2\omega_1/\omega)}^2 \end{aligned} \quad (4.5)$$

are isomorphisms.

In particular, $\text{Op}^w(a_1)$ is an isomorphism from $M_{(\omega_1)}^2$ to L^2 , and $\text{Op}^w(b_1)$ is an isomorphism from L^2 to $M_{(\omega_1)}^2$.

Now set $c = a_2 \# a \# b_1$. Then by [4, Theorem 4.14], the symbol c satisfies

$$c = a_2 \# a \# b_1 \in \Gamma_s^{(\omega_1/\omega)} \# \Gamma_s^{(\omega)} \# \Gamma_s^{(1/\omega_1)} \subseteq \Gamma_s^{(1)}.$$

Furthermore, $\text{Op}^w(c)$ is a composition of three isomorphisms and consequently $\text{Op}^w(c)$ is boundedly invertible on L^2 .

By Proposition 1.32 (2), $\text{Op}^w(c)^{-1} = \text{Op}^w(c_1)$ for some $c_1 \in \Gamma_{0,s}^{(1)}$. Hence, by (1) it follows that $\text{Op}^w(c)$ and $\text{Op}^w(c_1)$ are isomorphisms on $M(\omega_2, \mathcal{B})$, for each $\omega_2 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$. Since $\text{Op}^w(c)$ and $\text{Op}^w(c_1)$ are bounded on every $M(\omega, \mathcal{B})$, the factorization of the identity $\text{Op}^w(c) \text{Op}^w(c_1) = \text{Id}$ is well-defined on every $M(\omega, \mathcal{B})$. Consequently, $\text{Op}^w(c)$ is an isomorphism on $M(\omega, \mathcal{B})$.

Using the inverses of a_2 and b_1 , we now find that

$$\text{Op}^w(a) = \text{Op}^w(b_2) \circ \text{Op}^w(c) \circ \text{Op}^w(a_1)$$

is a composition of isomorphisms from the domain space $M(\omega_2, \mathcal{B})$ onto the image space $M(\omega_2/\omega, \mathcal{B})$ (factoring through some intermediate spaces) for every $\omega_2 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ and every invariant BF-space $\mathcal{B}$. This proves the isomorphism assertions for $\text{Op}^w(a)$.

Finally, the inverse of $\text{Op}^w(a)$ is given by

$$\text{Op}^w(b_1) \circ \text{Op}^w(c_1) \circ \text{Op}^w(a_2).$$

which is a Weyl operator with symbol in $\Gamma_{0,s}^{(1/\omega)}$, and the result follows. $\square$

5. MAPPING PROPERTIES FOR TOEPLITZ OPERATORS

In this section we study the isomorphism properties of Toeplitz operators between modulation spaces. We first state results for Toeplitz operators that are well-defined in the sense of (1.37) and Propositions 1.33 and 1.34. Then we state and prove more general results for Toeplitz operators that are defined only in the framework of pseudo-differential calculus.

We start with the following result about Toeplitz operators with smooth symbols.

Theorem 5.1. *Let $s \geq 1$, $\omega, \omega_0, v \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$ be such that $\omega_0 \in \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$ and that ω_0 is v -moderate, and let $\mathcal{B}$ be an invariant BF-space on $\mathbf{R}^{2d}$ or $\mathcal{B} = L^{p,E}(\mathbf{R}^{2d})$ for some phase-shift split parallelepiped E in $\mathbf{R}^{2d}$. If $\phi \in M_{(v)}^1(\mathbf{R}^d)$, then $\text{Tp}_\phi(\omega_0)$ is an isomorphism from $M(\omega, \mathcal{B})$ to $M(\omega/\omega_0, \mathcal{B})$.*

In the next result we relax our restrictions on the weights but impose more restrictions on $\mathcal{B}$.

Theorem 5.2. *Let $s > 1$, $0 \leq t \leq 1$, $p, q \in [1, \infty]$, and $\omega, \omega_0, v_0, v_1 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ be such that ω_0 is v_0 -moderate and ω is v_1 -moderate. Set $v = v_1^\dagger v_0$, $\vartheta = \omega_0^{1/2}$ and let $\omega_{0,t}$ be the same as in (1.38). If $\phi \in M_{(v)}^1(\mathbf{R}^d)$ and $\omega_0 \in \mathcal{M}_{(1/\omega_0,t)}^\infty(\mathbf{R}^{2d})$, then $\text{Tp}_\phi(\omega_0)$ is an isomorphism from $M_{(\vartheta\omega)}^{p,q}(\mathbf{R}^d)$ to $M_{(\omega/\vartheta)}^{p,q}(\mathbf{R}^d)$.*

Before the proofs of Theorems 5.1 and 5.2 we state the following consequence of Theorem 5.2 which was the original goal of our investigations.

Corollary 5.3. *Let $s \geq 1$, $\omega, \omega_0, v_1, v_0 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ and that ω_0 is v_0 -moderate and ω is v_1 -moderate. Set $v = v_1 v_0$ and $\vartheta = \omega_0^{1/2}$. If $\phi \in M_{(v)}^1(\mathbf{R}^d)$, then $\text{Tp}_\phi(\omega_0)$ is an isomorphism from $M_{(\vartheta\omega)}^{p,q}(\mathbf{R}^d)$ to $M_{(\omega/\vartheta)}^{p,q}(\mathbf{R}^d)$ simultaneously for all $p, q \in [1, \infty]$.*

Proof. Let $\omega_1 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d}) \cap \Gamma_{0,s}^{(\omega_1)}(\mathbf{R}^{2d})$ be such that $C^{-1} \leq \omega_1/\omega_0 \leq C$, for some constant C . Hence, $\omega/\omega_0 \in L^\infty \subseteq M^\infty$. By Theorem 2.2 in [50], it follows that $\omega = \omega_1 \cdot (\omega/\omega_1)$ belongs to $M_{(\omega_2)}^\infty(\mathbf{R}^{2d})$, when $\omega_2(x, \xi, \eta, y) = 1/\omega_0(x, \xi)$. The result now follows by setting $t = 1$ and $q_0 = 1$ in Theorem 5.2. $\square$

In the proofs of Theorems 5.1 and 5.2 we consider Toeplitz operators as defined by an extension of the form (1.37). Later on we present extensions of these theorems (cf. Theorems 5.1' and 5.2' below) for those readers who accept to use pseudo-differential calculus to extend the definition of Toeplitz operators. Except for the interpretation of $\text{Tp}_\phi(\omega)$ the proofs of Theorems 5.1 and 5.2 are identical to those of Theorems 5.1' and 5.2'.

We need some preparations and start with the following lemma.

Lemma 5.4. *Let $s \geq 1$, $\omega, v \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ be such that $\vartheta = \omega^{1/2}$ is v -moderate. Assume that $\phi \in M_{(v)}^2$. Then $\text{Tp}_\phi(\omega)$ is an isomorphism from $M_{(\vartheta)}^2(\mathbf{R}^d)$ onto $M_{(1/\vartheta)}^2(\mathbf{R}^d)$.*

Proof. Recall from Remark 1.13 that for $\phi \in M_{(v)}^2(\mathbf{R}^d) \setminus \{0\}$ the expression $\|V_\phi f \cdot \vartheta\|_{L^2}$ defines an equivalent norm on $M_{(\vartheta)}^2$. Thus the occurring STFTs with respect to ϕ are well defined.

Since $\text{Tp}_\phi(\omega)$ is bounded from $M_{(\vartheta)}^2$ to $M_{(1/\vartheta)}^2$ by Proposition 1.34, the estimate

$$\|\text{Tp}_\phi(\omega)f\|_{M_{(1/\vartheta)}^2} \lesssim \|f\|_{M_{(\vartheta)}^2} \quad (5.1)$$

holds for all $f \in M_{(\vartheta)}^2$.

Next we observe that

$$(\text{Tp}_\phi(\omega)f, g)_{L^2(\mathbf{R}^d)} = (\omega V_\phi f, V_\phi g)_{L^2(\mathbf{R}^{2d})} = (f, g)_{M_{(\vartheta)}^{2,\phi}}, \quad (5.2)$$

for $f, g \in M_{(\vartheta)}^2(\mathbf{R}^d)$ and $\phi \in M_{(v)}^2(\mathbf{R}^d)$. The duality of modulation spaces (Proposition 1.12(3)) now yields the following identity:

$$\begin{aligned} \|f\|_{M_{(\vartheta)}^2} &\asymp \sup_{\|g\|_{M_{(\vartheta)}^2}=1} |(f, g)_{M_{(\vartheta)}^2}| \\ &\asymp \sup_{\|g\|_{M_{(\vartheta)}^2}=1} |(\text{Tp}_\phi(\omega)f, g)_{L^2}| \asymp \|\text{Tp}_\phi(\omega)f\|_{M_{(1/\vartheta)}^2}. \end{aligned} \quad (5.3)$$

A combination of (5.1) and (5.3) shows that $\|f\|_{M_{(\vartheta)}^2}$ and $\|\text{Tp}_\phi(\omega)f\|_{M_{(1/\vartheta)}^2}$ are equivalent norms on $M_{(\vartheta)}^2$.

In particular, $\text{Tp}_\phi(\omega)$ is one-to-one from $M_{(\vartheta)}^2$ to $M_{(1/\vartheta)}^2$ with closed range. Since $\text{Tp}_\phi(\omega)$ is self-adjoint with respect to L^2 , it follows by duality that $\text{Tp}_\phi(\omega)$ has dense range in $M_{(1/\vartheta)}^2$. Consequently, $\text{Tp}_\phi(\omega)$ is onto $M_{(1/\vartheta)}^2$. By Banach's theorem $\text{Tp}_\phi(\omega)$ is an isomorphism from $M_{(\vartheta)}^2$ to $M_{(1/\vartheta)}^2$. $\square$

We need a further generalization of Proposition 1.33 to more general classes of symbols and windows. Set

$$\omega_1(X, Y) = \frac{v_0(2Y)^{1/2}v_1(2Y)}{\omega_0(X+Y)^{1/2}\omega_0(X-Y)^{1/2}}. \quad (5.4)$$

Proposition 1.33'. *Let $s \geq 1$, $0 \leq t \leq 1$, $p, q, q_0 \in [1, \infty]$, and $\omega, \omega_0, v_0, v_1 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ be such that v_0 and v_1 are submultiplicative, ω_0 is v_0 -moderate and ω is v_1 -moderate. Set*

$$r_0 = 2q_0/(2q_0 - 1), \quad v = v_1^t v_0 \quad \text{and} \quad \vartheta = \omega_0^{1/2},$$

and let $\omega_{0,t}$ and ω_1 be as in (1.38) and (5.4). Then the following are true:

- (1) *The definition of $(a, \phi) \mapsto \text{Tp}_\phi(a)$ from $\Sigma_s(\mathbf{R}^{2d}) \times \Sigma_s(\mathbf{R}^d)$ to $\mathcal{L}(\Sigma_s(\mathbf{R}^d), \Sigma'_s(\mathbf{R}^d))$ extends uniquely to a continuous map from $\mathcal{M}_{(1/\omega_{0,t})}^{\infty, q_0}(\mathbf{R}^{2d}) \times M_{(v)}^{r_0}(\mathbf{R}^d)$ to $\mathcal{L}(\Sigma_s(\mathbf{R}^d), \Sigma'_s(\mathbf{R}^d))$.*
- (2) *If $\phi \in M_{(v)}^{r_0}(\mathbf{R}^d)$ and $a \in \mathcal{M}_{(1/\omega_{0,t})}^{\infty, q_0}(\mathbf{R}^{2d})$, then $\text{Tp}_\phi(a) = \text{Op}^w(a_0)$ for some $a_0 \in \mathcal{M}_{(\omega_1)}^{\infty, 1}(\mathbf{R}^{2d})$, and $\text{Tp}_\phi(a)$ extends uniquely to a continuous map from $M_{(\vartheta\omega)}^{p,q}(\mathbf{R}^d)$ to $M_{(\omega/\vartheta)}^{p,q}(\mathbf{R}^d)$.*

For the proof we need the following result, which follows from [49, Proposition 2.1] and its proof.

Lemma 5.5. *Assume that $s \geq 1$, $q_0, r_0 \in [1, \infty]$ satisfy $r_0 = 2q_0/(2q_0 - 1)$. Also assume that $v \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ is submultiplicative, and that $\kappa, \kappa_0 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d} \oplus \mathbf{R}^{2d})$ satisfy*

$$\kappa_0(X_1 + X_2, Y) \leq C\kappa(X_1, Y)v(Y + X_2)v(Y - X_2) \quad X_1, X_2, Y \in \mathbf{R}^{2d}, \quad (5.5)$$

for some constant $C > 0$. Then the map $(a, \phi) \mapsto \text{Tp}_\phi(a)$ from $\Sigma_s(\mathbf{R}^{2d}) \times \Sigma_s(\mathbf{R}^d)$ to $\mathcal{L}(\Sigma_s(\mathbf{R}^d), \Sigma'_s(\mathbf{R}^d))$ extends uniquely to a continuous mapping from $\mathcal{M}_{(\omega)}^{\infty, q_0}(\mathbf{R}^{2d}) \times M_{(v)}^{r_0}(\mathbf{R}^d)$ to $\mathcal{L}(\Sigma_s(\mathbf{R}^d), \Sigma'_s(\mathbf{R}^d))$. Furthermore, if $\phi \in M_{(v)}^{r_0}(\mathbf{R}^d)$ and $a \in \mathcal{M}_{(\kappa)}^{\infty, q_0}(\mathbf{R}^{2d})$, then $\text{Tp}_\phi(a) = \text{Op}^w(b)$ for some $b \in \mathcal{M}_{(\kappa_0)}^{\infty, 1}$.

Proof of Proposition 1.33'. We show that the conditions on the involved parameters and weight functions satisfy the conditions of Lemma 5.5.

First we observe that

$$v_j(2Y) \leq Cv_j(Y + X_2)v_j(Y - X_2), \quad j = 0, 1$$

for some constant C which is independent of $X_2, Y \in \mathbf{R}^{2d}$, because v_0 and v_1 are submultiplicative. Referring back to (5.4) this gives

$$\begin{aligned} \omega_1(X_1 + X_2, Y) &= \frac{v_0(2Y)^{1/2}v_1(2Y)}{\omega_0(X_1 + X_2 + Y)^{1/2}\omega_0(X_1 + X_2 - Y)^{1/2}} \\ &\leq C_1 \frac{v_0(2Y)^{1/2}v_1(2Y)v_0(X_2 + Y)^{1/2}v_0(X_2 - Y)^{1/2}}{\omega_0(X_1)} \\ &= C_1 v_1(2Y)^{1-t} \frac{v_0(2Y)^{1/2}v_1(2Y)^t v_0(X_2 + Y)^{1/2}v_0(X_2 - Y)^{1/2}}{\omega_0(X_1)} \\ &\leq C_2 v_1(2Y)^{1-t} \frac{v_1(X_2 + Y)^t v_1(X_2 - Y)^t v_0(X_2 + Y)v_0(X_2 - Y)}{\omega_0(X_1)}. \end{aligned}$$

Hence

$$\omega_1(X_1 + X_2, Y) \leq C \frac{v_1(2Y)^{1-t}v(X_2 + Y)v(X_2 - Y)}{\omega_0(X_1)}. \quad (5.6)$$

By letting $\kappa_0 = \omega_1$ and $\kappa = 1/\omega_{0,t}$, it follows that (5.6) agrees with (5.5). The result now follows from Lemma 5.5. $\square$

In the remaining part of the paper we interpret $\text{Tp}_\phi(a)$ as the extension of a Toeplitz operator provided by Proposition 1.33'. (See also Remark 5.7 below for more comments.)

Proposition 1.33' can be applied to Toeplitz operators with smooth weights as symbols.

Proposition 5.6. *Assume that $s \geq 1$, $\omega_0 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$ be such that $\omega_0 \in \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$, that $v \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$ is submultiplicative, and that $\omega_0^{1/2}$ is v -moderate. If $\phi \in M_{(v)}^2(\mathbf{R}^d)$, then $\text{Tp}_\phi(\omega_0) = \text{Op}^w(b)$ for some $b \in \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$.*

Proof. By Propositions 1.25 and 1.26 with $t = 0$ we have $\omega_0 \in \mathcal{M}_{(1/\omega_0, r_0)}^{\infty, 1}(\mathbf{R}^{2d})$ for some $r_0 \geq 0$, where $\omega_{0,r_0}(X, Y) = \omega_0(X)e^{-r_0|Y|^{1/s}}$. Furthermore, by letting $v_1(Y) = e^{r_0|Y|^{1/s}}$, and ω_1 in (5.4) we have

$$\omega_1(X, Y) \gtrsim \frac{e^{r_0|2Y|^{1/s}}v(2Y)^{1/2}}{\omega_0(X+Y)^{1/2}\omega_0(X-Y)^{1/2}} \gtrsim \frac{e^{r_0|Y|^{1/s}}}{\omega_0(X)}.$$

Proposition 1.33' implies that existence of some $b \in \mathcal{M}_{(1/\omega_0, r_0)}^{\infty, 1}(\mathbf{R}^{2d}) \subseteq \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$. $\square$

The following generalization of Theorem 0.1 is an immediate consequence of Theorem 4.1, Lemma 5.4 and Proposition 5.6, since it follows by straightforward computations that $\mathcal{S}_s \subseteq M_{(v)}^2$ when v satisfies the hypothesis in Proposition 5.6.

Theorem 0.1'. *Let $s \geq 1$, $\omega, \omega_0 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$, $\mathbf{p} \in (0, \infty]^{2d}$, $\mathcal{B}$ be an invariant BF-space on $\mathbf{R}^{2d}$ or $\mathcal{B} = L^{\mathbf{p}, E}(\mathbf{R}^{2d})$ for some phase-shift split parallelepiped E in $\mathbf{R}^{2d}$, and let $\phi \in \mathcal{S}_s(\mathbf{R}^d)$. Then the Toeplitz operator $\text{Tp}_\phi(\omega_0)$ is an isomorphism from $M(\omega, \mathcal{B})$ onto $M(\omega/\omega_0, \mathcal{B})$.*

Remark 5.7. As remarked and stated before, there are different ways to extend the definition of a Toeplitz operator $\text{Tp}_\phi(a)$ (from $\phi \in \Sigma_1(\mathbf{R}^d)$ and $a \in \Sigma_1(\mathbf{R}^{2d})$) to more general classes of symbols and windows. For example, Propositions 1.33 and 1.34 are based on the ‘‘classical’’ definition (1.37) of such operators and a straightforward extension of the L^2 -form on Σ_1 . Proposition 5.6 interprets $\text{Tp}_\phi(\omega)$

as a pseudo-differential operator. Let us emphasize that in this context the bilinear form (1.37) may not be well defined, even when $\phi \in M_{(v)}^2(\mathbf{R}^d)$ and $\omega \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$ satisfies $\omega \in \Gamma_s^{(\omega)}(\mathbf{R}^{2d})$. (See also [26, Remark 3.8].)

In Theorems 5.1' and 5.2' below, we extend the definition of Toeplitz operators within the framework of pseudo-differential calculus and we interpret Toeplitz operators as pseudo-differential operators. With this understanding, the stated mapping properties are well-defined.

The reader who is not interested in full generality or does not accept Toeplitz operators that are not defined directly by an extension of (1.37) may only consider the case when the windows belong to $M_{(v)}^1$ and stay with Theorems 5.1 and 5.2. For the more general window classes in Theorems 5.1' and 5.2' below, however, one should then interpret the involved operators as “pseudo-differential operators that extend Toeplitz operators”.

The following generalization of Theorem 5.1 is an immediate consequence of Theorem 4.1, Lemma 5.4 and Proposition 5.6.

Theorem 5.1'. *Let $s \geq 1$, $\omega, v, v_0 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$ be such that $\omega_0 \in \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$ and that ω_0 is v -moderate, and let $\mathcal{B}$ be an invariant BF-space on $\mathbf{R}^{2d}$ or $\mathcal{B} = L^{p,E}(\mathbf{R}^{2d})$ for some phase-shift split parallelepiped E in $\mathbf{R}^{2d}$. If $\phi \in M_{(v)}^2(\mathbf{R}^d)$, then $\text{Tp}_\phi(\omega_0)$ is an isomorphism from $M(\omega, \mathcal{B})$ to $M(\omega/\omega_0, \mathcal{B})$.*

Theorem 5.1' holds only for smooth weight functions. In order to relax the conditions on the weight function ω_0 , we use the Wiener algebra property of $\mathcal{M}_{(v)}^{\infty,1}$ instead of $\Gamma_s^{(1)}$. On the other hand, we have to restrict our results to modulation spaces of the form $M_{(\omega)}^{p,q}$ instead of $M(\omega, \mathcal{B})$.

Theorem 5.2'. *Let $s > 1$, $0 \leq t \leq 1$, $p, q, q_0 \in [1, \infty]$ and $\omega, \omega_0, v_0, v_1 \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$ be such that ω_0 is v_0 -moderate and ω is v_1 -moderate. Set $r_0 = 2q_0/(2q_0 - 1)$, $v = v_1^t v_0$, $\vartheta = \omega_0^{1/2}$ and let $\omega_{0,t}$ be the same as in (1.38). If $\phi \in M_{(v)}^{r_0}(\mathbf{R}^d)$ and $\omega_0 \in \mathcal{M}_{(1/\omega_{0,t})}^{\infty, q_0}$, then $\text{Tp}_\phi(\omega_0)$ is an isomorphism from $M_{(\omega)}^{p,q}(\mathbf{R}^d)$ to $M_{(\omega/\vartheta)}^{p,q}(\mathbf{R}^d)$.*

Proof. First we note that the Toeplitz operator $\text{Tp}_\phi(\omega_0)$ is an isomorphism from $M_{(\vartheta)}^2$ to $M_{(1/\vartheta)}^2$ in view of Lemma 5.4. With ω_1 defined in (5.4), Proposition 1.33' implies that there exist $b \in \mathcal{M}_{(\omega_1)}^{\infty,1}$ and $c \in \mathcal{S}'_s(\mathbf{R}^{2d})$ such that

$$\text{Tp}_\phi(\omega_0) = \text{Op}^w(b) \quad \text{and} \quad \text{Tp}_\phi(\omega_0)^{-1} = \text{Op}^w(c).$$

Let ω_2 be the “dual” weight defined as

$$\omega_2(X, Y) = \vartheta(X - Y)\vartheta(X + Y)v_1(2Y). \quad (5.7)$$

We shall prove that $c \in \mathcal{M}_{(\omega_2)}^{\infty,1}(\mathbf{R}^{2d})$. Let us assume for now that we have already proved the existence of such a symbol c . Then we may proceed as follows.

After checking (1.30), we can apply Proposition 1.29 and find that each of the mappings

$$\text{Op}^w(b) : M_{(\omega\vartheta)}^{p,q} \rightarrow M_{(\omega/\vartheta)}^{p,q} \quad \text{and} \quad \text{Op}^w(c) : M_{(\omega/\vartheta)}^{p,q} \rightarrow M_{(\omega\vartheta)}^{p,q} \quad (5.8)$$

is well-defined and continuous.

In order to apply Proposition 1.30, we next check condition (1.31) for the weights ω_1 , ω_2 , and

$$\omega_3(X, Y) = \frac{\vartheta(X + Y)}{\vartheta(X - Y)}.$$

In fact we have

$$\begin{aligned}
& \omega_1(X - Y + Z, Z) \omega_2(X + Z, Y - Z) \\
&= \left(\frac{v_0(2Z)^{1/2} v_1(2Z)}{\vartheta(X - Y + 2Z) \vartheta(X - Y)} \right) \cdot (\vartheta(X - Y + 2Z) \vartheta(X + Y) v_1(2(Y - Z))) \\
&= \frac{v_0(2Z)^{1/2} v_1(2Z) v_1(2(Y - Z)) \vartheta(X + Y)}{\vartheta(X - Y)} \\
&\gtrsim \frac{\vartheta(X + Y)}{\vartheta(X - Y)} = \omega_3(X, Y).
\end{aligned}$$

Therefore Proposition 1.30 shows that the Weyl symbol of $\text{Op}^w(b) \circ \text{Op}^w(c)$ belongs to $\mathcal{M}_{(\omega_3)}^{\infty,1}(\mathbf{R}^{2d})$, or equivalently, $b \# c \in \mathcal{M}_{(\omega_3)}^{\infty,1}$. Since $\text{Op}^w(b)$ is an isomorphism from $M_{(\vartheta)}^2$ to $M_{(1/\vartheta)}^2$ with inverse $\text{Op}^w(c)$, it follows that $b \# c = 1$ and that the constant symbol 1 belongs to $\mathcal{M}_{(\omega_3)}^{\infty,1}$. By similar arguments it follows that $c \# b = 1$. Therefore the identity operator $\text{Id} = \text{Op}^w(b) \circ \text{Op}^w(c)$ on $M_{(\omega\vartheta)}^{p,q}$ factors through $M_{(\omega/\vartheta)}^{p,q}$, and thus $\text{Op}^w(b) = \text{Tp}_\phi(\omega_0)$ is an isomorphism from $M_{(\omega\vartheta)}^{p,q}$ onto $M_{(\omega/\vartheta)}^{p,q}$ with inverse $\text{Op}^w(c)$. This proves the assertion.

It remains to prove that $c \in \mathcal{M}_{(\omega_2)}^{\infty,1}(\mathbf{R}^{2d})$. Using once again the basic result in Section 3 (cf. Theorem 3.9), we choose $a \in \Gamma_{0,s}^{(1/\vartheta)}(\mathbf{R}^{2d})$ and $a_0 \in \Gamma_{0,s}^{(\vartheta)}(\mathbf{R}^{2d})$ such that the map

$$\text{Op}^w(a) : L^2(\mathbf{R}^d) \rightarrow M_{(\vartheta)}^2(\mathbf{R}^d)$$

is an isomorphism with inverse $\text{Op}^w(a_0)$. By Propositions 1.25 and 1.26, $\text{Op}^w(a)$ is also bijective from $M_{(1/\vartheta)}^2(\mathbf{R}^d)$ to $L^2(\mathbf{R}^d)$. Furthermore, by Theorem 4.2 it follows that $a \in \mathcal{M}_{(\vartheta_r)}^{\infty,1}$ when $r \geq 0$, where

$$\vartheta_r(X, Y) = \vartheta(X) e^{r|Y|^{1/s}}.$$

Let $b_0 = a \# b \# a$. From Proposition 1.31 we know that

$$b_0 \in \mathcal{M}_{(v_2)}^{\infty,1}(\mathbf{R}^{2d}), \quad \text{where} \quad v_2(X, Y) = v_1(2Y) \quad (5.9)$$

is submultiplicative and depends on Y only. Since $\text{Op}^w(b)$ is bijective from $M_{(\vartheta)}^2$ to $M_{(1/\vartheta)}^2$ by Lemma 5.4 (2), $\text{Op}^w(b_0)$ is bijective and continuous on L^2 .

Since v_2 is submultiplicative and in $\mathcal{P}_{E,s}(\mathbf{R}^{2d})$, $\mathcal{M}_{(v_2)}^{\infty,1}$ is a Wiener algebra by Proposition 1.32. Therefore, the Weyl symbol c_0 of the inverse to the bijective operator $\text{Op}^w(b_0)$ on L^2 belongs to $\mathcal{M}_{(v_2)}^{\infty,1}(\mathbf{R}^{2d})$.

Since

$$\text{Op}^w(c_0) = \text{Op}^w(b_0)^{-1} = \text{Op}^w(a)^{-1} \text{Op}^w(b)^{-1} \text{Op}^w(a)^{-1},$$

we find

$$\text{Op}^w(a_0) = \text{Op}^w(b)^{-1} = \text{Op}^w(a) \text{Op}^w(c_0) \text{Op}^w(a),$$

or equivalently,

$$a_0 = a \# c_0 \# a, \quad \text{where} \quad a \in \Gamma_{0,s}^{(1/\vartheta)} \quad \text{and} \quad c_0 \in \mathcal{M}_{(v_2)}^{\infty,1}. \quad (5.10)$$

The definitions of the weights are chosen such that Proposition 1.31 implies that $a_0 \in \mathcal{M}_{(\omega_2)}^{\infty,1}$, and the result follows. $\square$

6. EXAMPLES ON BIJECTIVE PSEUDO-DIFFERENTIAL OPERATORS ON MODULATION SPACES

In this section we construct explicit isomorphisms between modulation spaces with different weights. Applying the results of the previous sections, these may be in the form of pseudo-differential operators or of Toeplitz operators.

Proposition 6.1. *Let $s \geq 1$, $\omega_0, \omega \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d})$, and let $\mathcal{B}$ be an invariant BF-space on $\mathbf{R}^{2d}$ or $\mathcal{B} = L^{p,E}(\mathbf{R}^{2d})$ for some phase-shift split parallelepiped E in $\mathbf{R}^{2d}$. For $\lambda = (\lambda_1, \lambda_2) \in \mathbf{R}_+^2$ let Φ_λ be the Gaussian*

$$\Phi_\lambda(x, \xi) = Ce^{-(\lambda_1|x|^2 + \lambda_2|\xi|^2)}.$$

- (1) $\omega_0 * \Phi_\lambda$ belongs to $\mathcal{P}_{E,s}^0(\mathbf{R}^{2d}) \cap \Gamma_{0,1}^{(\omega_0)}$ for all $\lambda \in \mathbf{R}_+^2$ and

$$\omega_0 * \Phi_\lambda \asymp \omega_0.$$

- (2) If $\lambda_1 \cdot \lambda_2 < 1$, then there exists $\nu \in \mathbf{R}_+^2$ and a Gauss function ϕ on $\mathbf{R}^d$ such that $\text{Op}^w(\omega_0 * \Phi_\lambda) = \text{Tp}_\phi(\omega_0 * \Phi_\nu)$ is bijective from $M(\omega, \mathcal{B})$ to $M(\omega/\omega_0, \mathcal{B})$ for all $\omega \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$.
- (3) If $\lambda_1 \cdot \lambda_2 \leq 1$ and in addition $\omega_0 \in \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$, then $\text{Op}^w(\omega_0 * \Phi_\lambda) = \text{Tp}_\phi(\omega_0)$ is bijective from $M(\omega, \mathcal{B})$ to $M(\omega/\omega_0, \mathcal{B})$ for all $\omega \in \mathcal{P}_{E,s}(\mathbf{R}^{2d})$.

Proof. The assertion (1) follows easily from the definitions.

(2) Choose $\mu_j > \lambda_j$ such that $\mu_1 \cdot \mu_2 = 1$. Then the Gaussian Φ_μ is a multiple of a Wigner distribution, precisely $\Phi_\mu = cW(\phi, \phi)$ with $\phi(x) = e^{-\mu_1|x|^2/2}$. By the semigroup property of Gaussian functions (cf. e.g., [19, 21]) there exists another Gaussian, namely Φ_ν , such that $\Phi_\lambda = \Phi_\mu * \Phi_\nu$. Using (1.39), this factorization implies that the Weyl operator with symbol $\omega_0 * \Phi_\lambda$ is in fact a Toeplitz operator, namely

$$\begin{aligned} \text{Op}^w(\omega_0 * \Phi_\lambda) &= \text{Op}^w(\omega_0 * \Phi_\nu * \Phi_\mu) \\ &= \text{Op}^w(\omega_0 * \Phi_\nu * cW(\phi, \phi)) \\ &= c(2\pi)^{d/2} \text{Tp}_\phi(\omega_0 * \Phi_\nu). \end{aligned}$$

By (1) $\omega_0 * \Phi_\nu \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d}) \cap \Gamma_{0,1}^{(\omega_0)}(\mathbf{R}^{2d})$ is equivalent to ω_0 . Hence Theorem 5.1' shows that $\text{Op}^w(\omega_0 * \Phi_\lambda)$ is bijective from $M(\omega, \mathcal{B})$ to $M(\omega/\omega_0, \mathcal{B})$. This proves (2).

(3) follows from (2) in the case $\lambda_1 \cdot \lambda_2 < 1$. If $\lambda_1 \cdot \lambda_2 = 1$, then as above $\Phi_\lambda = cW(\phi, \phi)$ for $\phi(x) = e^{-\lambda_1|x|^2/2}$ and thus

$$\text{Op}^w(\omega_0 * \Phi_\lambda) = \text{Tp}_\phi^w(\omega_0)$$

is bijective from $M(\omega, \mathcal{B})$ to $M(\omega/\omega_0, \mathcal{B})$, since $\omega_0 \in \mathcal{P}_{E,s}^0(\mathbf{R}^{2d}) \cap \Gamma_s^{(\omega_0)}(\mathbf{R}^{2d})$. $\square$

APPENDIX A. PROOF OF LEMMA 3.3

Lemma A.1. *Let $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbf{N}^d$. Then the number of elements in the set*

$$\Omega_{k,\alpha} \equiv \{ (\beta_1, \dots, \beta_k) \in \mathbf{N}^{kd}; \beta_1 + \dots + \beta_k = \alpha \} \quad (\text{A.1})$$

is equal to

$$\prod_{j=1}^d \binom{\alpha_j + k}{k}.$$

For the proof we recall the formula

$$\sum_{j=0}^k \binom{n+j}{j} = \binom{n+k+1}{k}, \quad (\text{A.2})$$

which follows by a standard induction argument.

Proof. Let N be the number of elements in the set (A.1), which is the searched number, and let N_j be the number of elements of the set

$$\{(\beta_1^0, \dots, \beta_k^0) \in \mathbf{N}^k; \beta_1^0 + \dots + \beta_k^0 = \alpha_j\}, \quad j = 1, \dots, d$$

By straightforward computations it follows that $N = N_1 \cdots N_d$, and it suffices to prove the result in the case $d = 1$, and then $\alpha = \alpha_1$.

In order to prove the result for $d = 1$, let $\gamma \in \mathbf{N}$,

$$S_1(\gamma) = \sum_{\beta=0}^{\gamma} 1 = \gamma + 1,$$

and define inductively

$$S_{j+1}(\gamma) = \sum_{\beta=0}^{\gamma} S_j(\beta), \quad j = 1, 2, \dots$$

By straightforward computations it follows that $N = N_1 = S_k(\alpha)$. We claim

$$S_j(\gamma) = \binom{\gamma+j}{j}, \quad j = 1, 2, \dots \quad (\text{A.3})$$

In fact, (A.3) is obviously true for $j = 1$. Assume that (A.3) holds for $j = n$, and consider $S_{n+1}(\gamma)$. Then (A.2) gives

$$\begin{aligned} S_{n+1}(\gamma) &= \sum_{\beta=0}^{\gamma} S_n(\beta) = \sum_{\beta=0}^{\gamma} \binom{\beta+n}{n} \\ &= \sum_{\beta=0}^{\gamma} \binom{\beta+n}{\beta} = \binom{\gamma+n+1}{\gamma} = \binom{\gamma+n+1}{n+1}, \end{aligned}$$

which gives (A.3) when $j = n + 1$. This proves (A.3), and the result follows. $\square$

Lemma A.2. *Let $\alpha \geq 1$ be an integer, $s_0 \in (0, 1]$, and let $\Omega_{k,\alpha}$ be the same as in (A.1). Then there is a constant C which is independent of α such that*

$$\sum_{k=1}^{\alpha} \frac{1}{k} \sum_{\beta \in \Omega_{k,\alpha}} (\beta!)^{s_0-1} \leq 16^{\alpha}.$$

Proof. By Lemma A.1 and the fact that $s_0 - 1 < 0$ we get

$$\sum_{k=1}^{\alpha} \frac{1}{k} \left(\sum_{\beta \in \Omega_{k,\alpha}} \beta!^{s_0-1} \right) \leq \sum_{k=1}^{\alpha} \left(\sum_{\beta \in \Omega_{k,\alpha}} 1 \right) = \sum_{k=1}^{\alpha} \binom{\alpha+k}{k} \leq (1+\alpha)2^{2\alpha} \leq 16^{\alpha}. \quad \square$$

REFERENCES

- [1] T. Aoki *Locally bounded topological spaces*, Proc. Japan Acad. **18** (1942), 588–594.
- [2] H.A. Biagioni, T. Gramchev *Fractional derivative estimates in Gevrey classes, global regularity and decay for solutions to semilinear equations in $\mathbf{R}^d$* . J. Differential Equations **194** (2003), 140–165.
- [3] J. M. Bony, J. Y. Chemin *Espaces Fonctionnels Associés au Calcul de Weyl-Hörmander*, Bull. Soc. math. France **122** (1994), 77–118.

- [4] M. Cappiello, J. Toft *Pseudo-differential operators in a Gelfand-Shilov setting*, Math. Nachr. **290** (2017), 738–755.
- [5] Y. Chen, M. Signahl, J. Toft *Factorizations and singular value estimates of operators with Gelfand-Shilov and Pilipović kernels*, J. Fourier Anal. Appl. (appeared online 2017).
- [6] J. Chung, S.Y. Chung, D. Kim *Characterization of the Gelfand-Shilov spaces via Fourier transforms*, Proc. Amer. Math. Soc. **124** (1996), 2101–2108.
- [7] E. Cordero, S. Pilipović, L. Rodino, N. Teofanov *Quasianalytic Gelfand-Shilov spaces with applications to localization operators*, Rocky Mt. J. Math. **40** (2010), 1123–1147.
- [8] E. Cordero, K. Gröchenig *Time-Frequency Analysis of Localization Operators*, J. Funct. Anal. (1) **205** (2003), 107–131.
- [9] E. Cordero, S. Pilipović, L. Rodino, N. Teofanov *Localization operators and exponential weights for modulation spaces*, Mediterr. J. Math. **2** (2005), 381–394.
- [10] E. Cordero, J. Toft, P. Wahlberg *Sharp results for the Weyl product on modulation spaces*, J. Funct. Anal. **267** (2014), 3016–3057.
- [11] H. G. Feichtinger *Banach convolution algebras of Wiener’s type*, in: Proc. Functions, Series, Operators in Budapest, Colloquia Math. Soc. J. Bolyai, North Holland Publ. Co., Amsterdam Oxford NewYork, 1980.
- [12] H. G. Feichtinger *Modulation spaces on locally compact abelian groups*. In Proceedings of “International Conference on Wavelets and Applications” 2002, pages 99–140, Chennai, India, 2003. Updated version of a technical report, University of Vienna, 1983.
- [13] H. G. Feichtinger *Atomic characterizations of modulation spaces through Gabor-type representations*, in: Proc. Conf. on Constructive Function Theory, Rocky Mountain J. Math. **19** (1989), 113–126.
- [14] H. G. Feichtinger *Wiener amalgams over Euclidean spaces and some of their applications*, in: Function spaces (Edwardsville, IL, 1990), Lect. Notes in pure and appl. math., **136**, Marcel Dekker, New York, 1992, pp. 123–137.
- [15] H. G. Feichtinger *Modulation spaces: Looking back and ahead*, Sampl. Theory Signal Image Process. **5** (2006), 109–140.
- [16] H. G. Feichtinger and K. H. Gröchenig *Banach spaces related to integrable group representations and their atomic decompositions, I*, J. Funct. Anal. **86** (1989), 307–340.
- [17] C. Fernandez, A. Galbis, J. Toft *Spectral invariance for matrix algebras*, J. Fourier Anal. Appl. **20** (2014), 362–383.
- [18] H. G. Feichtinger and K. H. Gröchenig *Banach spaces related to integrable group representations and their atomic decompositions, II*, Monatsh. Math. **108** (1989), 129–148.
- [19] G. B. Folland *Harmonic analysis in phase space*, Princeton Univ. Press, Princeton, 1989.
- [20] Y. V. Galperin, S. Samarah *Time-frequency analysis on modulation spaces $M_m^{p,q}$, $0 < p, q \leq \infty$* , Appl. Comput. Harmon. Anal. **16** (2004), 1–18.
- [21] K. H. Gröchenig *Foundations of Time-Frequency Analysis*, Birkhäuser, Boston, 2001.
- [22] K. H. Gröchenig *Composition and spectral invariance of pseudodifferential operators on modulation spaces*, J. Anal. Math. **98** (2006), 65–82.
- [23] K. H. Gröchenig *Time-frequency analysis of Sjöstrand’s class*, Rev. Mat. Iberoam. **22** (2006), 703–724.
- [24] K. Gröchenig *Weight functions in time-frequency analysis* in: L. Rodino, M. W. Wong (Eds) Pseudodifferential Operators: Partial Differential Equations and Time-Frequency Analysis, Fields Institute Comm., **52** 2007, pp. 343–366.
- [25] K. Gröchenig and T. Strohmer. *Pseudodifferential operators on locally compact abelian groups and Sjöstrand’s symbol class*, J. Reine Angew. Math., 613:121–146, 2007.
- [26] K. Gröchenig, J. Toft *Isomorphism properties of Toeplitz operators and pseudo-differential operators between modulation spaces*, J. Anal. Math. **114** (2011), 255–283.
- [27] K. Gröchenig, J. Toft *The range of localization operators and lifting theorems for modulation and Bargmann-Fock spaces*, Trans. Amer. Math. Soc. **365** (2013), 4475–4496.
- [28] K.H. Gröchenig, G. Zimmermann *Spaces of test functions via the STFT*, J. Funct. Spaces. Appl. **2** (2004), 25–53.
- [29] H. Gzyl *Multidimensional Extension of Faà di Bruno’s formula*, J. Math. Anal. Appl. **116** (1986), 450–455.
- [30] L. Hörmander *The Weyl calculus of pseudo-differential operators*, Comm. Pure Appl. Math. **32** (1979), 359–443.
- [31] L. Hörmander *The Analysis of Linear Partial Differential Operators*, vol I, III, Springer-Verlag, Berlin Heidelberg NewYork Tokyo, 1983, 1985.
- [32] Z. Lozanov-Crvenković, D. Perišić, M. Tasković *Gelfand-Shilov spaces structural and kernel theorems*, (preprint), arXiv:0706.2268v2.
- [33] N. Lerner *The Wick calculus of pseudo-differential operators and some of its applications*, CUBO, **5** (2003), 213–236.

- [34] N. Lerner *Metrics on the Phase Space and Non-Selfadjoint Pseudo-Differential Operators*, Birkhäuser Springer, (2010).
- [35] B.S. Mitjagin *Nuclearity and other properties of spaces of type $\mathcal{S}$* , Amer. Math. Soc. Transl., Ser. **93** (1970), 49–59.
- [36] C. Pfeuffer, J. Toft *Compactness properties for modulation spaces* (under preparation).
- [37] S. Pilipović *Tempered ultradistributions*, Boll. Un. Mat. Ital., VII Ser., **2** (1988), 235–251.
- [38] S. Pilipović, N. Teofanov *Pseudodifferential operators on ultramodulation spaces*, J. Funct. Anal. **208** (2004), 194–228.
- [39] S. Rolewicz *On a certain class of linear metric spaces*, Bull. Acad. Polon. Sci. Sér. Schi. Math. Astronom. Phys. **5** (1957), 471–473.
- [40] M. Ruzhansky, M. Sugimoto, N. Tomita, J. Toft *Changes of variables in modulation and Wiener amalgam spaces*, Math. Nachr. **284** (2011), 2078–2092.
- [41] M. Ruzhansky, N. Tokmagambetov *Nonharmonic analysis of boundary value problems*, Int. Math. Res. Notices **12** (2016), 3548–3615.
- [42] J. Sjöstrand *An algebra of pseudodifferential operators*, Math. Res. L. **1** (1994), 185–192.
- [43] J. Sjöstrand. *Pseudodifferential operators and weighted normed symbol spaces*, Serdica Math. J. **34** (2008), no. 1, 1–38.
- [44] K. Tachizawa *The boundedness of pseudo-differential operators on modulation spaces*, Math. Nachr. **168** (1994), 263–277.
- [45] N. Teofanov *Modulation spaces, Gelfand-Shilov spaces and pseudodifferential operators*, Sampl. Theory Signal Image Process, **5** (2006), 225–242.
- [46] J. Toft *Continuity Properties for non-Commutative Convolution Algebras with Applications in Pseudo-Differential Calculus*, Bull. Sci. Math. (2) **126** (2002), 115–142.
- [47] J. Toft *Continuity properties for modulation spaces with applications to pseudo-differential calculus, II*, Ann. Global Anal. Geom., **26** (2004), 73–106.
- [48] J. Toft *Continuity and Schatten properties for pseudo-differential operators on modulation spaces* in: J. Toft, M. W. Wong, H. Zhu (Eds) *Modern Trends in Pseudo-Differential Operators*, Operator Theory: Advances and Applications **172**, Birkhäuser Verlag, Basel, 2007, pp. 173–206.
- [49] J. Toft *Continuity and Schatten properties for Toeplitz operators on modulation spaces* in: J. Toft, M. W. Wong, H. Zhu (Eds) *Modern Trends in Pseudo-Differential Operators*, Operator Theory: Advances and Applications **172**, Birkhäuser Verlag, Basel, 2007, pp. 313–328.
- [50] J. Toft *Pseudo-differential operators with smooth symbols on modulation spaces*, Cubo. **11** (2009), 87–107.
- [51] J. Toft *Multiplication properties in pseudo-differential calculus with small regularity on the symbols*, J. Pseudo-Differ. Oper. Appl. **1** (2010), 101–138.
- [52] J. Toft *The Bargmann transform on modulation and Gelfand-Shilov spaces, with applications to Toeplitz and pseudo-differential operators*, J. Pseudo-Differ. Oper. Appl. **3** (2012), 145–227.
- [53] J. Toft *Multiplication properties in Gelfand-Shilov pseudo-differential calculus* in: S. Molahajlo, S. Pilipović, J. Toft, M. W. Wong (eds) *Pseudo-Differential Operators, Generalized Functions and Asymptotics*, Operator Theory: Advances and Applications **231**, Birkhäuser, Basel Heidelberg NewYork Dordrecht London, 2013, pp. 117–172.
- [54] J. Toft *Gabor analysis for a broad class of quasi-Banach modulation spaces*, in: S. Pilipović, J. Toft (eds) *Pseudo-Differential Operators and Generalized Functions*, Operator Theory: Advances and Applications Vol 245, Birkhäuser, Basel Heidelberg NewYork Dordrecht London, 2015, pp. 249–278.
- [55] J. Toft *Matrix parameterized pseudo-differential calculi on modulation spaces* in: M. Oberguggenberger, J. Toft, J. Vindas, P. Wahlberg (eds), *Generalized functions and Fourier analysis*, Operator Theory: Advances and Applications Vol 260, Birkhäuser/Springer, Basel, 2017, pp. 215–235.
- [56] J. Toft *Continuity of Gevrey-Hörmander pseudo-differential operators on modulation spaces*, (preprint) arXiv:1710.11366 (2017).
- [57] J. Toft, P. Boggiatto, *Schatten classes for Toeplitz operators with Hilbert space windows on modulation spaces*, Adv. Math. **217** (2008), no. 1, 305–333.
- [58] J. Toft, A. Khrennikov, B. Nilsson, S. Nordebo *Decompositions of Gelfand-Shilov kernels into kernels of similar class*, J. Math. Anal. Appl. **396** (2012), 315–322.

DIPARTIMENTO DI MATEMATICA “G. PEANO”, UNIVERSITÀ DEGLI STUDI DI TORINO
E-mail address: `ahmed.abdeljawad@unito.it`

DIPARTIMENTO DI MATEMATICA “G. PEANO”, UNIVERSITÀ DEGLI STUDI DI TORINO
E-mail address: `sandro.coriasco@unito.it`

DEPARTMENT OF MATHEMATICS, LINNÆUS UNIVERSITY, SWEDEN
E-mail address: `joachim.toft@lnu.se`