

MIXED-EFFECTS MODELS USING THE NORMAL AND THE LAPLACE DISTRIBUTIONS: A 2×2 CONVOLUTION SCHEME FOR APPLIED RESEARCH

BY MARCO GERACI[†]

University of South Carolina[†]

Abstract In statistical applications, the normal and the Laplace distributions are often contrasted: the former as a standard tool of analysis, the latter as its robust counterpart. I discuss the convolutions of these two popular distributions and their applications in research. I consider four models within a simple 2×2 scheme which is of practical interest in the analysis of clustered (e.g., longitudinal) data. In my view, these models, some of which are less known than others by the majority of applied researchers, constitute a ‘family’ of sensible alternatives when modelling issues arise. In three examples, I revisit data published recently in the epidemiological and clinical literature as well as a classic biological dataset.

1. Introduction. The normal (or Gaussian) distribution historically has played a prominent role not only as limiting distribution of a number of sample statistics, but also for modelling data obtained in empirical studies. Its probability density is given by

$$(1) \quad f_N(t) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left\{ -\frac{1}{2} \left(\frac{t - \mu}{\sigma} \right)^2 \right\},$$

for $-\infty < t < \infty$. The Laplace (or double exponential) distribution, like the normal, has a long history in Statistics. However, despite being of potentially great value in applied research, it has never received the same attention. Its density is given by

$$(2) \quad f_L(t) = \frac{1}{\sqrt{2}\sigma} \exp \left\{ -\sqrt{2} \left| \frac{t - \mu}{\sigma} \right| \right\}.$$

Throughout this paper, these distributions will be denoted by $\mathcal{N}(\mu, \sigma)$ and $\mathcal{L}(\mu, \sigma)$, respectively.

[†]Corresponding author: Marco Geraci, Department of Epidemiology and Biostatistics, Arnold School of Public Health, University of South Carolina, 915 Greene Street, Columbia SC 29209, USA. E-mail: geraci@mailbox.sc.edu

MSC 2010 subject classifications: Primary 62F99; secondary 62J05

Keywords and phrases: Crohn’s disease, linear quantile mixed models, meta-analysis, multilevel designs, random effects




FIGURE 1. (a) Left: The normal (solid line) and double exponential (dashed line) densities. The location parameter is set to 0 and the variance is set to 1. (b) Right: The normal-normal (solid line), normal-Laplace (dashed line), and Laplace-Laplace (dot-dash line) densities. The location parameter is set to 0 and the variance is set to 1.

In (1) and (2), μ and σ , where $-\infty < \mu < \infty$ and $\sigma > 0$, represent a location and a scale parameters, respectively. These two densities are shown in the left-hand side plots of Figure 1. The normal and Laplace distributions are both symmetric about μ and have variance equal to σ^2 . As compared to the normal one, the Laplace density has a more pronounced peak (a characteristic technically defined *leptokurtosis*) and fatter tails. Interestingly, the Laplace distribution can be represented as a scale mixture of normal distributions. Let $T \sim \mathcal{L}(\mu, \sigma)$, then (Kotz, Kozubowski and Podgórski, 2001)

$$T \stackrel{d}{=} \mu + \sigma\sqrt{E}Z,$$

where E and Z are independent standard exponential and normal variables, respectively. That is, the Laplace distribution emerges from heterogeneous normal sub-populations.

Both laws were proposed by Pierre-Simon Laplace: the double exponential in 1774 and the normal in 1778 (for an historical account, see Wilson, 1923). At Laplace's time, the problem to be solved was that of estimating μ

according to the linear model

$$T = \mu + \sigma \varepsilon,$$

where ε denotes the error term. This problem was encountered, for example, in astronomy and geodesy, where μ represented the ‘true’ value of a physical quantity to be estimated from experimental observations. It is well known that, under the Gaussian error law (1), the maximum likelihood estimate of μ is the sample mean but, under the double exponential error law (2), it is the sample median. The former is the minimiser of the least squares (LS) estimator, while the latter is the minimiser of the least absolute deviations (LAD) estimator.

The robustness of the LAD estimator in presence of large errors was known to Laplace himself. However, given the superior analytical tractability of the LS estimator (and therefore of the normal distribution), the mean regression model

$$T = x^\top \beta + \sigma \varepsilon, \quad \varepsilon \sim \mathcal{N}(0, 1),$$

quickly became the ‘standard’ tool to study the association between the location parameter of T (the response variable) and other variables of interest, x (the covariates).

In the past few years, theoretical developments related to least absolute error regression (Bassett and Koenker, 1978; Koenker and Bassett, 1978) have led to a renewed interest in the Laplace distribution and its asymmetric extension (Yu and Zhang, 2005) as pseudo-likelihood for quantile regression models of which median regression is a special case (see, among others, Yu and Moyeed, 2001; Yu, Lu and Stander, 2003; Geraci and Bottai, 2007). In parallel, computational advances based on interior point algorithms have made LAD estimation a serious competitor of LS methods (Portnoy and Koenker, 1997; Koenker and Ng, 2005). Another reason for the ‘comeback’ of the double exponential is related to its robustness properties which makes this distribution and distributions alike desirable in many applied research areas (Kozubowski and Nadarajah, 2010).

In statistical applications, the interest is often in processes where the source of randomness can be attributed to more than one ‘error’ (a hierarchy of errors is also established). For instance, this is the case of longitudinal studies where part of the variation is attributed to an individual source of heterogeneity (often called ‘random effect’), say ε_1 , independently from the noise, ε_2 , i.e.

$$T = \mu + \sigma_1 \varepsilon_1 + \sigma_2 \varepsilon_2,$$

where the distributions of ε_1 and ε_2 are often assumed to be symmetric about zero. It will be shown later that this model can be extended to include covariates associated with the parameter μ and the random effect ε_1 . For now, it suffices to notice that the linear combination of random errors leads to the study of convolutions. So let us define a convolution (Mood, Graybill and Boes, 1974).

DEFINITION 1. *If U and V are two independent, absolutely continuous random variables with density functions f_U and f_V , respectively, and $T = U + V$, then*

$$f_T(t) = f_{U+V}(t) = \int_{-\infty}^{\infty} f_V(t-u)f_U(u) du = \int_{-\infty}^{\infty} f_U(t-v)f_V(v) dv$$

is the convolution of f_U and f_V .

In Section 2, I consider convolutions based on the normal and the Laplace distributions within a simple and practical 2×2 scheme. In Section 3, I discuss inference when data are clustered, along with the implementation of estimation procedures using existing R (R Core Team, 2014) software (further technical details are provided in Appendix, along with a simulation study). In Section 4, I show some applications and, in Section 5, conclude with final remarks.

2. Convolutions. Let Y be a real-valued random variable with absolutely continuous distribution function $F(y) = \Pr\{Y \leq y\}$ and density $f(y) \equiv F'(y)$. The variable Y is observable and represents the focus of the analysis in specific applications (e.g., as the response variable in regression models). I consider four cases in which Y results from one of the four convolutions reported in Table 1. The letters ν and λ are used to denote normal and Laplace variates with densities (1) and (2), respectively. The subscripts 1 and 2 indicate, respectively, which of the two random variables plays the role of a random effect and which one is considered to be the noise. Here, the former may in general be associated with a vector of covariates and may represent an inferential quantity of interest; the latter is treated as a nuisance. Moreover, I assume independence between the components of the convolution throughout the paper.

A few remarks about notation are needed. The shorthand $\text{diag}(t)$ or $\text{diag}(t_1, \dots, t_n)$, where $t = (t_1, \dots, t_n)^\top$ is a $n \times 1$ vector, is used to denote the $n \times n$ diagonal matrix whose diagonal elements are the corresponding elements of t . The standard normal density and cumulative distribution functions will be denoted by ϕ and Φ , respectively.

TABLE 1
 2×2 convolution scheme for independent Gaussian (ν) and Laplacian (λ) random variables.

	Normal	Laplace
Normal	$\nu_1 + \nu_2$ (NN)	$\nu_1 + \lambda_2$ (NL)
Laplace	$\lambda_1 + \nu_2$ (LN)	$\lambda_1 + \lambda_2$ (LL)

2.1. *Normal-normal (NN) convolution.* The first convolution

$$(3) \quad Y = \nu_1 + \nu_2,$$

where $\nu_1 \sim \mathcal{N}(0, \sigma_1)$ and $\nu_2 \sim \mathcal{N}(0, \sigma_2)$, represents, in some respects, the simplest case among the four combinations defined in Table 1. Standard theory of normal distributions leads to

$$(4) \quad f_{NN}(y) = \frac{1}{\psi} \phi\left(\frac{y}{\psi}\right),$$

where $\psi^2 \equiv \text{var}(Y) = \sigma_1^2 + \sigma_2^2$.

Model (3) can be generalised to the regression model

$$(5) \quad Y = x^\top \beta + z^\top \nu_1 + \nu_2,$$

where x and z are, respectively, $p \times 1$ and $q \times 1$ vectors of covariates, and β is a $p \times 1$ dimensional vector of regression coefficients. If $q > 1$, then I assume $\nu_1 \sim \mathcal{N}_q(0, \Sigma_1)$, that is, a multivariate normal distribution with $q \times q$ variance-covariance matrix Σ_1 . It follows that

$$(6) \quad g_{NN}(y) = \frac{1}{\psi} \phi\left(\frac{y - x^\top \beta}{\psi}\right),$$

where $\psi^2 \equiv \text{var}(Y) = z^\top \Sigma_1 z + \sigma_2^2$.

Model (5) is known as a linear mixed effects (LME) model or, simply, as a mixed model (Pinheiro and Bates, 2000; Demidenko, 2013). There is a vast number of applications of LME models, especially for the analysis of clustered data in the social, life and physical sciences.

2.2. *Normal-Laplace (NL) convolution.* The second convolution consists of a normal and a Laplace components, that is

$$(7) \quad Y = \nu_1 + \lambda_2,$$

where $\nu_1 \sim \mathcal{N}(0, \sigma_1)$ and $\lambda_2 \sim \mathcal{L}(0, \sigma_2)$. The resulting density is given by (Reed, 2006)

$$(8) \quad f_{NL}(y) = \frac{1}{\sqrt{2}\sigma_2} \phi(y/\sigma_1) \{R(\kappa - y/\sigma_1) + R(\kappa + y/\sigma_1)\},$$

where $\kappa = \sqrt{2}\sigma_1/\sigma_2$ and R is the Mills ratio

$$R(t) = \frac{1 - \Phi(t)}{\phi(t)}.$$

The above distribution arises from a Brownian motion whose starting value is normally distributed and whose stopping hazard rate is constant. An extension of (8) to skewed forms can be obtained by letting λ_2 follow an asymmetric Laplace distribution (Reed, 2006). Applications of the NL convolution can be found in finance (Reed, 2007; Meintanis and Tsionas, 2010). See also the double Pareto-lognormal distribution, associated with $\exp(Y)$, which has applications in modeling size distributions (Reed and Jorgensen, 2004).

As in the previous section, I consider a generalisation of model (7) to the regression case

$$(9) \quad Y = x^\top \beta + z^\top \nu_1 + \lambda_2.$$

If $q > 1$, then I assume $\nu_1 \sim \mathcal{N}_q(0, \Sigma_1)$. It follows that $z^\top \nu_1$ is normal with mean zero and variance $z^\top \Sigma_1 z$. This leads to

$$(10) \quad g_{NL}(y) = \frac{1}{\sqrt{2}\sigma_2} \phi\left(\frac{y - x^\top \beta}{\sigma_1}\right) \left\{ R\left(\kappa - \frac{y - x^\top \beta}{\sigma_1}\right) + R\left(\kappa + \frac{y - x^\top \beta}{\sigma_1}\right) \right\},$$

where $\sigma_1 \equiv \sqrt{z^\top \Sigma_1 z}$ and, as defined above, $\kappa = \sqrt{2}\sigma_1/\sigma_2$. It is easy to verify that $\text{var}(Y) = \sigma_1^2 + \sigma_2^2$.

Model (9) is a median regression model with normal random effects, a special case of the linear quantile mixed models (LQMMs) discussed by Geraci and Bottai (2007, 2014). LQMMs have been used in a wide range of research areas, including marine biology (Muir et al., 2015; Duffy et al., 2015; Barneche et al., 2016), environmental science (Fornaroli et al., 2015), cardiovascular disease (Degerud et al., 2014; Blankenberg et al., 2016), physical activity (Ng et al., 2014; Beets et al., 2016), and ophthalmology (Patel et al., 2015; Patel, Geraci and Cortina-Borja, 2016).

2.3. *Laplace-normal (LN) convolution.* The Laplace-normal convolution is given by

$$(11) \quad Y = \lambda_1 + \nu_2,$$

where $\lambda_1 \sim \mathcal{L}(0, \sigma_1)$ and $\nu_2 \sim \mathcal{N}(0, \sigma_2)$. The LN appears in robust meta-analysis (Demidenko, 2013, p.266).

The LN convolution in (11), clearly, is the same as the NL convolution in (7) (so I omit writing its density). However, note that now the Laplace component is associated with the random effect, not with the error term; therefore, the two scale parameters σ_1 and σ_2 will appear swapped. The distinction becomes clear when considering the regression model

$$(12) \quad Y = x^\top \beta + z^\top \lambda_1 + \nu_2.$$

By analogy with the NL convolution, I assume that, for $q > 1$, λ_1 has a q -dimensional multivariate Laplace distribution (Kotz, Kozubowski and Podgórski, 2001, p.235).

DEFINITION 2. *An n -dimensional random variable T is said to follow a zero-centred multivariate Laplace distribution with parameter Σ , $T \sim \mathcal{L}_q(0, \Sigma)$, if its density is given by*

$$f_L(t) = 2(2\pi)^{-n/2} |\Sigma|^{-1/2} \left(t^\top \Sigma^{-1} t / 2 \right)^{\omega/2} K_\omega \left(\sqrt{2t^\top \Sigma^{-1} t} \right),$$

where Σ is an $n \times n$ nonnegative definite symmetric matrix, $\omega = (2 - n)/2$ and K_ω is the modified Bessel function of the third kind.

REMARK 2.1. *If $T \sim \mathcal{L}_q(0, \Sigma)$, then $\text{cov}(t) = \Sigma$ (Kotz, Kozubowski and Podgórski, 2001, p.249). For a diagonal matrix $\Sigma = \text{diag}(\varsigma_1, \dots, \varsigma_q)$, the coordinates of the multivariate Laplace are uncorrelated, but not independent. Therefore, the joint distribution of n independent univariate Laplace variates does not have the properties of the multivariate Laplace with diagonal variance-covariance matrix.*

For $n = 1$, the multivariate density defined above reduces to the univariate density (2) with $\sigma = \Sigma^{1/2}$. Moreover, a linear combination of the coordinates of the multivariate Laplace is still a Laplace (Kotz, Kozubowski and Podgórski, 2001, p.255). Indeed, if we assume $\lambda_1 \sim \mathcal{L}_q(0, \Sigma_1)$, then $z^\top \lambda_1 \sim \mathcal{L}(0, \sigma_1)$, where $\sigma_1 = \sqrt{z^\top \Sigma_1 z}$. Thus, the density of Y in Equation (12) is given by

$$(13) \quad g_{LN}(y) = \frac{1}{\sqrt{2}\sigma_1} \phi \left(\frac{y - x^\top \beta}{\sigma_2} \right) \left\{ R \left(\kappa - \frac{y - x^\top \beta}{\sigma_2} \right) + R \left(\kappa + \frac{y - x^\top \beta}{\sigma_2} \right) \right\},$$

where $\kappa = \sqrt{2}\sigma_2/\sigma_1$. Again, it is easy to verify that $\text{var}(Y) = \sigma_1^2 + \sigma_2^2$.

2.4. *Laplace-Laplace (LL) convolution.* The fourth and last convolution consists of two Laplace variates, i.e.

$$(14) \quad Y = \lambda_1 + \lambda_2,$$

where $\lambda_1 \sim \mathcal{L}(0, \sigma_1)$ and $\lambda_2 \sim \mathcal{L}(0, \sigma_2)$. It can be shown (Kotz, Kozubowski and Podgórski, 2001, p.35) that the density of Y is

$$(15) \quad f_{LL}(y) = \begin{cases} \frac{1}{4}s(1 + s|y|) \exp(-s|y|), & \text{if } s_1 = s_2 = s, \\ \frac{\kappa}{2\kappa^2 - 2} \{s_1 \exp(-s_2|y|) - s_2 \exp(-s_1|y|)\}, & \text{if } s_1/s_2 = \kappa \neq 1, \end{cases}$$

with $s_1 = \sigma_1/\sqrt{2}$ and $s_2 = \sigma_2/\sqrt{2}$.

For the regression model

$$(16) \quad Y = x^\top \beta + z^\top \lambda_1 + \lambda_2,$$

with $\lambda_1 \sim \mathcal{L}_q(0, \Sigma_1)$, I obtain

$$(17) \quad g_{LL}(y) = \begin{cases} \frac{1}{4}s(1 + s|y - x^\top \beta|) \exp(-s|y - x^\top \beta|), & \text{if } s_1 = s_2 = s, \\ \frac{\kappa}{2\kappa^2 - 2} \{s_1 \exp(-s_2|y - x^\top \beta|) - s_2 \exp(-s_1|y - x^\top \beta|)\}, & \text{if } s_1/s_2 = \kappa \neq 1. \end{cases}$$

with $s_1 = \sigma_1/\sqrt{2}$ and $\sigma_1 = \sqrt{z^\top \Sigma_1 z}$. The variance is given by $\text{var}(Y) = \sigma_1^2 + \sigma_2^2$.

Model (16) is a median regression model with ‘robust’ random effects, another special case of LQMMS (Geraci and Bottai, 2014).

2.5. *Some properties.* All the convolutions are symmetric, unimodal, twice differentiable and have continuous first and second derivatives (the NN and NL are also smooth). Also, they are log-concave since both the normal (1) and Laplace (2) densities are log-concave (Prékopa, 1973). The right-hand side plots of Figure 1 shows that, as compared to the NN density, the NL (LN) and LL densities are leptokurtic and have more weight in the tails, with the NL density sitting between the NN and LL distributions. Thus, the presence of the Laplace term in the convolution confers different degrees of robustness to the model depending on whether one or both random terms are assumed to be Laplacian. Also, notice that the marginal regression models are location–scale–shift models, since both the location and the scale of Y are functions of the covariates.

3. Inference. In this section, I briefly discuss inferential issues, with detailed mathematical derivations provided in Appendix.

Let $Y_i = (Y_{i1}, Y_{i2}, \dots, Y_{in_i})^\top$ be a multivariate $n_i \times 1$ random response vector, and x_{ij} and z_{ij} , be vectors of covariates for the j th observation, $j = 1, \dots, n_i$, in cluster i , $i = 1, \dots, M$. Each component of Y_i can be modelled using any of the convolutions discussed in Section 2 by assuming

$$Y_{ij} = x_{ij}^\top \beta + z_{ij}^\top \varepsilon_{1i} + \varepsilon_{2ij},$$

where the random effect ε_{1i} and the error term ε_{2ij} are either Gaussian or Laplacian according to the scheme in Table 1. The marginal models implied by these four convolutions have been defined in expressions (6), (10), (13), and (17). At the cluster level, I use the notation

$$(18) \quad Y_i = X_i \beta + Z_i \varepsilon_{1i} + \varepsilon_{2i},$$

where X_i and Z_i are, respectively, $n_i \times p$ and $n_i \times q$ design matrices. I assume that the vector of random effects ε_{1i} has variance-covariance matrix Σ_1 for all $i = 1, \dots, M$ and that the Y_i 's are independent of one another. The structure of Σ_1 is, for the moment, left unspecified. Also, I assume that $\text{cov}(\varepsilon_{2i})$ is a multiple of the identity matrix, although this assumption can be easily relaxed (see Section 3.4).

There are several approaches to mixed effects model estimation (see, for example, [Pinheiro and Bates, 2000](#); [Demidenko, 2013](#)), each approach having its own advantages and disadvantages. One approach is to work with the marginal likelihood of Y_i . Although independence between clusters can still be assumed, in general the Y_{ij} 's will be correlated within the same cluster. Therefore, parameter estimation based on the marginal likelihood requires knowing the joint distribution of $Y_{i1}, Y_{i2}, \dots, Y_{in_i}$. Under the NN convolution, Y_i is known to be multivariate normal. It is beyond the scope of this paper to derive the multivariate distribution of Y_i for the NL, LN and LL convolutions.

Likelihood-based estimation of location and scale parameters using the NN model has been largely studied. Therefore, I will focus on the NL, LN, and LL models. Since an important aspect in applied research is the availability of software to perform data analysis, here I consider two methods which can be applied using existing software. The first method is based on numerical integration and applies to specific NL and LL models, while the other method is based on a Monte Carlo Expectation-Maximisation (MCEM) algorithm and applies to NL, LN and LL models.

3.1. *Numerical integration.* Let the i th contribution to the marginal log-likelihood be

$$\ell(\beta, \Sigma_1, \sigma_2; y_i) = \log \int_{\mathbb{R}^q} g(y_i - X_i\beta - Z_i u_i) h(u_i) du_i,$$

where g denotes the density of the error term conditional on the random effect u_i and h denotes the density of the random effect. One can work with the numerically integrated likelihood

(19)

$$\tilde{\ell}(\beta, \Sigma_1, \sigma_2; y_i) = \log \sum_{k_1}^K \cdots \sum_{k_q}^K g\left(y_i - X_i\beta - Z_i (\Sigma_1)^{1/2} v_{k_1, \dots, k_q}\right) h(v_{k_1, \dots, k_q}),$$

with nodes $v_{k_1, \dots, k_q} = (v_{k_1}, \dots, v_{k_q})^\top$ and weights $h(v_{k_1, \dots, k_q})$, $k_l = 1, \dots, K$, $l = 1, \dots, q$, as an approximation to the marginal log-likelihood.

The maximisation of the approximate log-likelihood (19) can be time-consuming depending on the dimension of the quadrature q , the required accuracy of the approximation controlled by the number of nodes K , and, of course, the distribution $h(u)$. If Σ_1 is a diagonal matrix, then $h(v_{k_1, \dots, k_q}) = \prod_{l=1}^q h(v_{k_l})$. This greatly simplifies calculations since the q -dimensional integral can be carried out with q successive applications of one-dimensional quadrature rules. In the multivariate normal case, a non-diagonal covariance matrix can be rescaled to a diagonal one and the joint density factorises into q normal variates. However, this is not the case for the multivariate Laplace, at least not for the one defined in Section 2.3. Geraci and Bottai (2014) considered a steepest-descent approach combined with Gauss-Hermite and Gauss-Laguerre quadrature for, respectively, the NL and LL likelihoods. Standard errors were obtained by bootstrapping the clusters (block bootstrap).

Since Geraci's (2014) algorithms, which are implemented in the R package `lqmm`, can be applied to selected models only (namely, NL models with correlated or uncorrelated random effects and LL models with uncorrelated random effects), in the next section I develop an alternative, more general approach based on the EM algorithm.

3.2. *EM estimation.* Rather than working with the Laplace distribution directly, I consider its representation as a scale mixture of normal distributions. As noted before, if $T \sim \mathcal{L}(0, \sigma)$, then $T \stackrel{d}{=} \sigma \sqrt{W}V$, where W and V are, respectively, independent standard exponential and normal variates. This equivalence has been used in EM estimation of regression quantiles

(see, for example, [Lum and Gelfand, 2012](#); [Tian, Tian and Zhu, 2013](#)). Similarly, in the multivariate case, if $T \sim \mathcal{L}_q(0, \Sigma_1)$, then $T \stackrel{d}{=} \sqrt{W}V$, where W is, again, standard exponential and $V \sim \mathcal{N}_q(0, \Sigma_1)$. As shown in Appendix, the normal components in the scale mixture representation of the NL, LN, and LL models can be easily convolved (conditionally on W) and the resulting log-likelihood for the i th cluster becomes

$$\ell(\beta, \Sigma_1, \sigma_2; y_i, w_i) = \log g(y_i|w_i) + \log h(w_i),$$

where g is multivariate normal and h is standard exponential.

The proposed EM algorithm starts from the likelihood of the complete data (y_i, w_i) , where w_i represents the unobservable data. In the E-step, the expected value of the complete log-likelihood is approximated using a Monte Carlo expectation. As shown in expression (25) in Appendix, the M-step reduces to the maximum likelihood estimation of a linear mixed model with prior weights which can be carried out using fitting routines from existing software (e.g., `nlme` or `lme4` in R).

3.3. Modelling and estimation of Σ_1 . There are different possible structures for Σ_1 . The simplest is a multiple of the identity matrix, with constant diagonal elements and zero off-diagonal elements. Other structures include, for example, diagonal (variance components), compound symmetric (constant diagonal and constant off-diagonal elements), and the more general symmetric positive-definite matrix. These are all available in the `nlme` ([Pinheiro et al., 2014](#)), `lme4` ([Douglas et al., 2015](#)) and `lqmm` ([Geraci, 2014](#)) packages, as well as in SAS procedures for mixed effects models.

The variance-covariance matrix of the random effects, whether normal or Laplace, must be nonnegative definite. However, it is possible that, during MLE, the estimate $\hat{\Sigma}_1$ may be singular or veer off into the space of negative definite matrices. This problem does not occur in EM estimation if the starting matrix is nonnegative definite. However, the monotonicity property is lost when a Monte Carlo error is introduced at the E-step ([McLachlan and Krishnan, 2008](#)). There are at least three approaches one can consider ([Demidenko, 2013](#), p.88): (i) allow $\hat{\Sigma}_1$ to be negative definite during estimation and, if negative definite at convergence, replace it with a nonnegative definite matrix after the algorithm has converged; (ii) constrained optimisation; (iii) matrix reparameterisation ([Pinheiro and Bates, 2000](#)). As discussed in Appendix, I follow the latter approach.

3.4. Residual heteroscedasticity and correlation. The development of the EM algorithm discussed above is based on the assumption that the within-group errors are independent with common scale parameter σ_2 . As briefly

outlined in Section A.6 in Appendix, it is easy to extend the NL, LN, and LL models to the case of heteroscedastic and correlated errors. Commonly available mixed effects software provide capabilities for estimating residual variance and correlation parameters. For the sake of simplicity, I do not consider this extension any further in this paper.

4. Examples.

4.1. *Meta-analysis.* Here, I discuss an application in meta-analysis. The data consist of mean standard deviation scores of height at diagnosis in osteosarcoma patients which had been reported in five different studies (Figure 2) and were successively meta-analysed by Arora et al. (2011). Let Y denote the study-specific effect. For these data, I considered the NN model (DerSimonian and Laird, 1986)

$$Y_i = \mu + \nu_{1i} + \nu_{2i}, \quad i = 1, \dots, 5,$$

where $\nu_{1i} \sim \mathcal{N}(0, \tau)$ and $\nu_{2i} \sim \mathcal{N}(0, \sigma_i)$, and the LN model (Demidenko, 2013)

$$Y_i = \mu + \lambda_{1i} + \nu_{2i}, \quad i = 1, \dots, 5,$$

where $\lambda_{1i} \sim \mathcal{L}(0, \tau)$ and $\nu_{2i} \sim \mathcal{N}(0, \sigma_i)$.

In meta-analysis, the goal is to estimate an ‘overall’ or ‘pooled’ effect (μ) and the between-study variance or heterogeneity among study-specific effects (τ^2). The sampling variances σ_i^2 are assumed to be known.

Estimation for the osteosarcoma data was carried out using R software developed for standard (Viechtbauer, 2010) and robust (Demidenko, 2013) meta-analysis. The estimates (standard errors) of μ and τ^2 were, respectively, 0.260 (0.087) and 0.029 (0.027) for the NN model, and 0.246 (0.073) and 0.021 (0.033) for the LN model. The larger estimated overall effect and heterogeneity for the NN model are a consequence of the outlying effect size of study 5 (Figure 2) which skews the location μ and inflates the scale of the normal distribution. In contrast, the Laplace distribution is more robust to outliers and heavy tails. Indeed, the estimate of μ from the LN model was more precise as demonstrated by the lower standard error (as a consequence, the related test statistic has smaller p -value). A similar example is described by Demidenko (2013).

4.2. *Repeated measurements in clinical trials.* Ten Crohn’s disease patients with endoscopic recurrence were followed over time (Sorrentino et al., 2010). Colonoscopy was performed and surrogate markers of disease activity




FIGURE 2. Forest plot for five studies on the relationship between height at diagnosis and osteosarcoma in young people. Each study is represented by a block, with area proportional to its weight, centered at the effect point estimate. Horizontal grey lines depict 95% confidence intervals.

were collected on four occasions. One of the goals of this trial was to assess the association between fecal calprotectin (FC – mg/kg) and endoscopic score (ES – Rutgeerts). The data were analysed using a log-linear median regression model under the assumption of independence between measurements (Sorrentino et al., 2010). Here, I take the within-patient correlation into account and analyse the data using three of the four regression models discussed in Section 2: the NN model

$$\log Y_{ij} = \beta_0 + \beta_1 x_{ij} + \nu_{1i} + \nu_{2ij}, \quad j = 1, \dots, 4, \quad i = 1, \dots, 10,$$

the NL model

$$\log Y_{ij} = \beta_0 + \beta_1 x_{ij} + \nu_{1i} + \lambda_{2ij}, \quad j = 1, \dots, 4, \quad i = 1, \dots, 10,$$

and the LL model

$$\log Y_{ij} = \beta_0 + \beta_1 x_{ij} + \lambda_{1i} + \lambda_{2ij}, \quad j = 1, \dots, 4, \quad i = 1, \dots, 10,$$

where Y_{ij} and x_{ij} denote, respectively, FC and ES measurements on patient i at occasion j , $\nu_{1i} \sim \mathcal{N}(0, \tau)$, $\nu_{2ij} \sim \mathcal{N}(0, \sigma)$, $\lambda_{1i} \sim \mathcal{L}(0, \tau)$, and $\lambda_{2ij} \sim \mathcal{L}(0, \sigma)$. Therefore, the variance of the random effects is τ^2 , while the variance of the error term is σ^2 .

TABLE 2

Association between fecal calprotectin and endoscopic score in Crohn's disease patients. Estimates and standard errors (SE) of the fixed effects (β), variance of the random effects (τ^2), and intra-class correlation (ρ) from three models. The log-likelihood (ℓ) is reported in brackets.

	β_0	β_1	τ^2	ρ
<i>Normal-Normal ($\ell = -21.8$)</i>				
Estimate	3.293	0.910	0.031	0.191
SE	0.113	0.056	0.133	
<i>Normal-Laplace ($\ell = -22.2$)</i>				
Estimate	3.354	0.871	0.994	0.877
SE	0.135	0.051	0.046	
<i>Laplace-Laplace ($\ell = -14.2$)</i>				
Estimate	3.269	0.905	0.293	0.757
SE	0.114	0.035	0.053	

In this case, the parameters of interest are the slope β_1 and the intra-class correlation $\rho = \tau^2/(\tau^2 + \sigma^2)$, which measures how much of the total variance is due to between-individual variability. Estimation was carried out using the nlme (Pinheiro et al., 2014) and lqmm (Geraci, 2014) packages. The results are shown in Table 2. The estimates of the regression coefficients β




FIGURE 3. *QQ-plot of the first-level (left plot) and second-level (right plot) residuals from the normal-normal model for the Crohn's disease data.*

tallied across models. However, the estimates of τ^2 and ρ differed substantially, with values from the NN model much lower than those from the NL and LL models. First-level residuals (i.e., predictions of the random effects plus the error term) and second-level residuals (i.e., predictions of the error term only) from the NN model are shown in Figure 3. It is apparent that σ^2 may be inflated by an unusual second-level residual, to the detriment of τ^2 . As a consequence, the intra-class correlation appeared to be heavily underestimated by the NN model. The NL model improved upon the estimation of the scale parameters as it is more robust to outliers in the error term. However, the LL model gave the largest value of the log-likelihood, suggesting that the goodness of the fit is further improved by using a robust distribution for the random effects as well. Note also that the standard error of the slope was smallest for the LL model.

4.3. *Growth curves.* In a weight gain experiment, 30 rats were randomly assigned to three treatment groups: treatment 1, a control (no additive); treatments 2 and 3, which consisted of two different additives (thiouracil and thyroxin respectively) to the rats drinking water (Box, 1950). Weight (grams) of the rats was measured at baseline (week 0) and at weeks 1, 2, 3, and 4. Data on three of the 10 rats from the thyroxin group were subsequently removed due to an accident at the beginning of the study. Figure 4 shows estimated intercepts and slopes obtained from rat-specific




FIGURE 4. Ordinary least squares estimates of intercepts and slopes for individual growth curves in the rats weight gain data. The scatterplots on the top show the pairwise estimates with LOESS smoothing superimposed (dashed grey lines mark mean values). The plots on the bottom depict the estimated densities of intercepts (solid line) and slopes (dashed line) centred and scaled using their respective means and standard deviations.

LS regressions of the type

$$Y_{ij,k} = \beta_{0i,k} + \beta_{1i,k}x_j + \sigma_{i,k}\varepsilon_{ij,k}, \quad \varepsilon_{ij,k} \sim \mathcal{N}(0, 1),$$

where the response $Y_{ij,k}$ is weight measurement taken on rat $i = 1, \dots, M_k$ on occasion $j = 1, \dots, 5$ conditional on treatment group $k = 1, 2, 3$, and $x_j = j - 1$. (Note that $M_1 = M_2 = 10$ and $M_3 = 7$.) It is evident that the weight of rats treated with thiouracil grew slower than the controls', though at baseline the former tended to be heavier than the latter. In contrast, rats in the control and thyroxin groups had, on average, similar intercepts and slopes.

The Pearson's correlation coefficients of the estimated intercept-slope pairs $(\hat{\beta}_{0i,k}, \hat{\beta}_{1i,k})$ gave -0.26 ($k = 1$), -0.37 ($k = 2$), and -0.16 ($k = 3$), suggesting a negative association between baseline weight and growth rate in all treatment groups. However, the direction of the association in treatment group 3 is unclear. Interestingly, the Kendall rank correlation coefficient in the thyroxin group indicated a weak positive association (0.05), while the

Pearson's coefficient became strongly positive (0.97) after removing the two pairs with the largest slopes. Moreover, the distributions of intercepts and slopes showed the presence of skewness and bimodality. Therefore, some degree of robustness against departures from normality might be needed.

To model the heterogeneity within each treatment group, subject-specific random intercepts and slopes were included in the following four models: the NN model

$$Y_{ij,k} = \beta_{0,k} + \beta_{1,k}x_j + \nu_{1i,k}^{(1)} + \nu_{1i,k}^{(2)}x_j + \nu_{2ij,k},$$

$$j = 1, \dots, 5, \quad i = 1, \dots, M_k, \quad k = 1, 2, 3,$$

the NL model

$$Y_{ij,k} = \beta_{0,k} + \beta_{1,k}x_j + \nu_{1i,k}^{(1)} + \nu_{1i,k}^{(2)}x_j + \lambda_{2ij,k},$$

$$j = 1, \dots, 5, \quad i = 1, \dots, M_k, \quad k = 1, 2, 3,$$

the LN model

$$Y_{ij,k} = \beta_{0,k} + \beta_{1,k}x_j + \lambda_{1i,k}^{(1)} + \lambda_{1i,k}^{(2)}x_j + \nu_{2ij,k},$$

$$j = 1, \dots, 5, \quad i = 1, \dots, M_k, \quad k = 1, 2, 3,$$

and the LL model

$$Y_{ij,k} = \beta_{0,k} + \beta_{1,k}x_j + \lambda_{1i,k}^{(1)} + \lambda_{1i,k}^{(2)}x_j + \lambda_{2ij,k},$$

$$j = 1, \dots, 5, \quad i = 1, \dots, M_k, \quad k = 1, 2, 3,$$

where I assumed $(\nu_{1i,k}^{(1)}, \nu_{1i,k}^{(2)}) \sim \mathcal{N}_2(0, \Sigma_{1,k})$, $\nu_{2ij,k} \sim \mathcal{N}(0, \sigma_2)$, $(\lambda_{1i,k}^{(1)}, \lambda_{1i,k}^{(2)}) \sim \mathcal{L}_2(0, \Sigma_{1,k})$, and $\lambda_{2ij,k} \sim \mathcal{L}(0, \sigma_2)$, and the $\Sigma_{1,k}$'s, $k = 1, 2, 3$, are 2×2 symmetric matrices,

$$\Sigma_{1,k} = \begin{bmatrix} \varsigma_{11,k} & \varsigma_{12,k} \\ \varsigma_{12,k} & \varsigma_{22,k} \end{bmatrix}.$$

Further, I assumed that the random effects are uncorrelated between treatment groups.

The NL, LN, and LL models were estimated using the EM algorithm as detailed in Appendix with a Monte Carlo size equal to 100, fixed at each EM iteration, and a convergence tolerance of $5 \cdot 10^{-4}$. The four models gave similar estimates of the fixed effects (Table 3), although the trajectory in the thiouracil group resulting from the LN model tended to be less steep than the corresponding trajectory resulting from the other three models. However, this difference might be of little practical importance. In contrast, more substantial seemed to be the differences between the estimates of the correlation

TABLE 3

Rats weight gain data. Estimates and standard errors (SE) of the fixed effects (β) from four models. The log-likelihood (ℓ) is reported in brackets.

	$\beta_{0,1}$	$\beta_{0,2}$	$\beta_{0,3}$	$\beta_{1,1}$	$\beta_{1,2}$	$\beta_{1,3}$
<i>Normal-Normal</i> ($\ell = -444.4$)						
Estimate	52.880	57.700	52.086	26.480	17.050	27.143
SE	2.349	2.058	1.578	1.177	0.879	1.928
<i>Normal-Laplace</i> ($\ell = -448.4$)						
Estimate	52.934	57.568	52.928	26.383	17.208	26.791
SE	2.427	2.204	1.519	1.208	0.928	2.146
<i>Laplace-Normal</i> ($\ell = -551.6$)						
Estimate	53.069	58.392	51.104	25.620	16.794	26.665
SE	1.992	1.972	1.817	0.885	0.814	1.910
<i>Laplace-Laplace</i> ($\ell = -454.0$)						
Estimate	52.680	58.433	53.415	26.067	17.305	27.621
SE	1.960	1.762	1.041	0.924	0.748	1.353

TABLE 4

Rats weight gain data. Estimated correlation matrix of the random intercepts and slopes for each treatment group from three models. The log-likelihood (ℓ) is reported in brackets.

<i>Normal-Normal</i> ($\ell = -444.4$)						
	<i>Treatment 1</i>		<i>Treatment 2</i>		<i>Treatment 3</i>	
	Int.	Slope	Int.	Slope	Int.	Slope
Int.	1.000		1.000		1.000	
Slope	-0.145	1.000	-0.203	1.000	0.050	1.000
<i>Normal-Laplace</i> ($\ell = -448.4$)						
	<i>Treatment 1</i>		<i>Treatment 2</i>		<i>Treatment 3</i>	
	Int.	Slope	Int.	Slope	Int.	Slope
Int.	1.000		1.000		1.000	
Slope	-0.076	1.000	-0.133	1.000	0.634	1.000
<i>Laplace-Normal</i> ($\ell = -551.6$)						
	<i>Treatment 1</i>		<i>Treatment 2</i>		<i>Treatment 3</i>	
	Int.	Slope	Int.	Slope	Int.	Slope
Int.	1.000		1.000		1.000	
Slope	-0.117	1.000	-0.065	1.000	0.194	1.000
<i>Laplace-Laplace</i> ($\ell = -454.0$)						
	<i>Treatment 1</i>		<i>Treatment 2</i>		<i>Treatment 3</i>	
	Int.	Slope	Int.	Slope	Int.	Slope
Int.	1.000		1.000		1.000	
Slope	0.030	1.000	-0.294	1.000	0.876	1.000

matrices $D_{1,k}^{-1}\Sigma_{1,k}D_{1,k}^{-1}$, where $D = \text{diag}(\sqrt{\varsigma_{11,k}}, \sqrt{\varsigma_{22,k}})$, $k = 1, 2, 3$ (Table 4). It is interesting to note that there is disagreement on the magnitude and even direction of some of the estimates. Notably, $\hat{\varsigma}_{12,3}/(\hat{\varsigma}_{11,3} \cdot \hat{\varsigma}_{22,3})$ was smallest for the NN model but it was substantially larger for the NL and LL models. The best fit in terms of the log-likelihood was for the NN model, followed

closely by the NL model. The LL model and, especially, the LN model gave smaller log-likelihoods.

5. Final remarks. In the words of [Wilson \(1923, p.842\)](#) “No phenomenon is better known perhaps, as a plain matter of fact, than that the frequencies which I actually meet in everyday work in economics, in biometrics, or in vital statistics, very frequently fail to conform at all closely to the so-called normal distribution”. [Kotz and colleagues \(2001\)](#) echo Wilson’s observations on the inadequacy of the normal distribution in many practical applications and give a systematic exposition of the Laplace distribution, an unjustifiably neglected error law which can be “a natural and sometimes superior alternative to the normal law” ([Kotz, Kozubowski and Podgórski, 2001, p.13](#)).

My proposed 2×2 convolution scheme brings together the normal and Laplace distributions showing that these models represent a *family* of sensible alternatives as they introduce a varying degree of robustness in the modelling process. Estimation can be approached in different ways. The EM algorithm discussed in this paper takes advantage of the scale mixture representation of the Laplace distribution which provides the opportunity for computational simplification. In a simulation study with a moderate sample size (see [Section A.7](#) in Appendix), this algorithm provided satisfactory results in terms of mean squared error for the NL and LL models. The estimation of the LN model needed a relatively larger number of Monte Carlo samples to achieve reasonable bias, though the results were never fully satisfactory in terms of efficiency. Finally, model selection has been left out of consideration, but further research on this topic is needed, especially at smaller sample sizes. An interesting starting point is offered by [Kundu \(2005\)](#).

To reiterate the main point of this study, these convolutions have a large number of potential applications and, as demonstrated using several examples, may provide valuable insight into different aspects of the analysis.

Acknowledgements. This research has been supported by the National Institutes of Health – National Institute of Child Health and Human Development (Grant Number: 1R03HD084807-01A1).

APPENDIX

A.1. EM estimation. Here, I discuss maximum likelihood inference for β , Σ_1 , and σ_2 in normal-Laplace (NL), Laplace-normal (LN), and Laplace-Laplace (LL) models. In particular, I develop an estimation approach based

on the scale mixture representation of the Laplace distribution. If $T \sim \mathcal{L}(0, \sigma)$ then $T \stackrel{d}{=} \sigma \sqrt{W}V$, where W and V are, respectively, independent standard exponential and normal variates. Similarly, if $T \sim \mathcal{L}_q(0, \Sigma_1)$, then $T \stackrel{d}{=} \sqrt{W}V$, where W is, again, standard exponential and $V \sim \mathcal{N}_q(0, \Sigma_1)$ (Kotz, Kozubowski and Podgórski, 2001).

Let $Y_i = (Y_{i1}, Y_{i2}, \dots, Y_{in_i})^\top$ be a multivariate $n_i \times 1$ random vector, and x_{ij} and z_{ij} be, respectively, $p \times 1$ and $q \times 1$ vectors of covariates for the j th observation, $j = 1, \dots, n_i$, in cluster i , $i = 1, \dots, M$. Also, let X_i and Z_i be, respectively, $n_i \times p$ and $n_i \times q$ design matrices for cluster i . I assume that the random effects have variance-covariance matrix Σ_1 for all $i = 1, \dots, n$ and that the Y_i 's are independent of one another. The structure of Σ_1 is purposely left unspecified. The relative precision matrix $\sigma_2^2 \Sigma_1^{-1}$ is parameterised in terms of an unrestricted m -dimensional vector, $1 \leq m \leq q(q+1)/2$, of non-redundant parameters ξ (Pinheiro and Bates, 2000). The parameter to be estimated is then $\theta = (\beta^\top, \xi^\top, \sigma_2)^\top$ of dimension $(p+m+1) \times 1$. The $n \times n$ identity matrix will be denoted by I_n .

A.2. Normal-Laplace convolution. Let $w_i = (w_{i1}, \dots, w_{in_i})^\top$ be a $n_i \times 1$ vector of independent standard exponential variates and let $D_i = \text{diag}(w_i)$. The NL model can be written as

$$(20) \quad Y_i = X_i \beta + Z_i \nu_{1i} + D_i^{1/2} v_i,$$

where $\nu_{1i} \sim \mathcal{N}_q(0, \Sigma_1)$ and $v_i \sim \mathcal{N}_{n_i}(0, \sigma_2^2 I_{n_i})$. The model can be simplified by convolving ν_{1i} and v_i conditional on w_i , i.e., by integrating out the random effects

$$g(y_i, w_i) = \int_{\mathbb{R}^q} g(y_i, \nu_{1i} | w_i) h(w_i) d\nu_{1i} = g(y_i | w_i) h(w_i),$$

where $y_i | w_i \sim \mathcal{N}_{n_i}(X_i \beta, \Omega_i)$, $\Omega_i = Z_i \Sigma_1 Z_i^\top + \sigma_2^2 D_i$, and $h(w_i) = \prod_{j=1}^{n_i} \exp(-w_{ij})$.

A.3. Laplace-Normal convolution. The LN model can be written as

$$(21) \quad Y_i = X_i \beta + \sqrt{w_i} Z_i v_i + \nu_{2i},$$

where w_i is a standard exponential variate, $v_i \sim \mathcal{N}_q(0, \Sigma_1)$, and $\nu_{2i} \sim \mathcal{N}_{n_i}(0, \sigma_2^2 I_{n_i})$. The normal component of the random effects can be integrated out as follows

$$g(y_i, w_i) = \int_{\mathbb{R}^q} g(y_i, v_i | w_i) h(w_i) dv_i = g(y_i | w_i) h(w_i),$$

where $y_i | w_i \sim \mathcal{N}_{n_i}(X_i \beta, \Omega_i)$, $\Omega_i = w_i Z_i \Sigma_1 Z_i^\top + \sigma_2^2 I_{n_i}$, and $h(w_i) = \exp(-w_i)$.

A.4. Laplace-Laplace convolution. Let $w_i = (w_{i,1}, w_{i,2}^\top)^\top$ be a $(1 + n_i) \times 1$ vector of independent standard exponential variates, where $w_{i,2} = (w_{i1,2}, \dots, w_{in_i,2})^\top$, and let $D_i = \text{diag}(w_{i,2})$. The LL model can be written as

$$(22) \quad Y_i = X_i\beta + \sqrt{w_{i,1}}Z_i v_{1i} + D_i^{1/2}v_{2i},$$

where $v_{1i} \sim \mathcal{N}_q(0, \Sigma_1)$ and $v_{2i} \sim \mathcal{N}_{n_i}(0, \sigma_2^2 I_{n_i})$. As before, the joint density can be simplified to

$$g(y_i, w_i) = \int_{\mathbb{R}^q} g(y_i, v_{1i}|w_i) h(w_i) dv_{1i} = g(y_i|w_i) h(w_i),$$

where $y_i|w_i \sim \mathcal{N}_{n_i}(X_i\beta, \Omega_i)$, $\Omega_i = w_{i,1}Z_i\Sigma_1Z_i^\top + \sigma_2^2D_i$, and $h(w_i) = \exp(-w_{i,1}) \cdot$

$$\prod_{j=1}^{n_i} \exp(-w_{ij,2}).$$

A.5. The algorithm. The joint density $g(y, w)$ could be further integrated to obtain $g(y) = \int g(y|w) \cdot h(w) dw$. Except for the normal-normal (NN) model, the form of the marginal likelihood of Y_i does not seem to have an immediate known form. A numerical integration could have some appeal since this integral would reduce to a Gauss-Laguerre quadrature ($h(w)$ is standard exponential). However, since quadrature methods are notoriously inefficient if the dimension of the integral is large, I consider an alternative approach based on Monte Carlo EM (MCEM) estimation. In this case, the unobservable variable w is sampled from the conditional density $g(w|y)$. While the Monte Carlo sample size does not depend as much on dimensionality as quadrature methods do, convergence can be slower for MCEM than for quadrature-based methods (McLachlan and Krishnan, 2008).

The i th contribution to the complete data log-likelihood for the models (20)-(22) is given by

$$(23) \quad \ell(\theta; y_i, w_i) = \log g(y_i|w_i) + \log h(w_i).$$

Note that $h(w_i)$ does not depend on θ . The EM approach alternates between an

- (i) expectation step (E-step) $Q_i(\theta|\theta^{(t)}) = E_{w|y_i, \theta^{(t)}} \{ \ell(\theta; y_i, w_i) \}$, $i = 1, \dots, M$; and a
- (ii) maximisation step (M-step) $\theta^{(t+1)} = \arg \max_{\theta} \sum_i Q_i(\theta|\theta^{(t)})$,

where $\theta^{(t)}$ is the estimate of the parameter after t cycles. The expectation in step (i) is taken with respect to $h(w_i|y_i, \theta^{(t)}) \propto g(y_i|w_i, \theta^{(t)}) h(w_i)$, that

is, the distribution of the unobservable data w_i conditional on the observed data y_i and the current estimate of θ . Given that the latter density does not have an immediate known form, I consider a Monte Carlo approach and use the following numerical approximation

$$(24) \quad \tilde{Q}_i(\theta|\theta^{(t)}) = \frac{1}{K} \sum_{k=1}^K \left\{ \ell \left(\theta; y_i, w_{ik}^{(t)} \right) \right\},$$

where $w_{ik}^{(t)}$ is a vector of appropriate dimensions sampled from $h(w_i|y_i, \theta^{(t)})$ at iteration t . The number of samples K can be fixed at the same value for all iterations or may vary with t . The approximate complete data log-likelihood for all clusters (Q-function), averaged over $w|y$, is given by

$$(25) \quad \tilde{Q}(\theta|\theta^{(t)}) \equiv \sum_{i=1}^M \tilde{Q}_i(\theta|\theta^{(t)}) = \frac{1}{K} \sum_{k=1}^K \sum_{i=1}^M -\frac{n_i}{2} \log(2\pi) - \frac{1}{2} \log |\Omega_{ik}| \\ - \frac{1}{2} e_i^\top \Omega_{ik}^{-1} e_i + \log h \left(w_{ik}^{(t)} \right),$$

where $e_i = y_i - X_i\beta$, $\Omega_{ik} = \sigma_2^2 \Psi_{ik}$,

$$\Psi_{ik} = \begin{cases} Z_i \dot{\Sigma}_1 Z_i^\top + D_{ik} & \text{with } D_{ik} = \text{diag} \left(w_{ik}^{(t)} \right) \text{ for the NL model,} \\ w_{ik}^{(t)} Z_i \dot{\Sigma}_1 Z_i^\top + I_{n_i} & \text{for the LN model,} \\ w_{ik,1}^{(t)} Z_i \dot{\Sigma}_1 Z_i^\top + D_{ik} & \text{with } D_{ik} = \text{diag} \left(w_{ik,2}^{(t)} \right) \text{ for the LL model,} \end{cases}$$

and $\dot{\Sigma}_1 = \sigma_2^{-2} \Sigma_1$ is the scaled variance-covariance matrix of the random effects. Note that all the information given by $\theta^{(t)}$ is contained in Ω_{ik} which depends on $w_{ik}^{(t)}$ (the superscript (t) has been dropped from Ω_{ik} , Ψ_{ik} , and D_{ik} to ease notation). Furthermore, the parameter ξ is defined to be the vector of non-zero elements of the upper triangle of the matrix logarithm of U , where U is the $q \times q$ matrix obtained from the Cholesky decomposition $\dot{\Sigma}_1^{-1} = U^\top U$ (Pinheiro and Bates, 2000).

The Q-function (25) can be easily maximised with respect to β , ξ , and σ_2 using standard (restricted) maximum likelihood formulas for linear mixed models (Pinheiro and Bates, 2000; Demidenko, 2013). Indeed, the derivative

of (25) with respect to θ has the familiar form
(26)

$$\tilde{Q}_*(\theta|\theta^{(t)}) = \begin{pmatrix} \frac{1}{K} \sum_{k=1}^K \sum_{i=1}^M \sigma_2^{-2} X_i \Psi_{ik}^{-1} e_i \\ -\frac{1}{2K} \sum_{i=1}^M \sum_{k=1}^K Z_i^\top \Psi_{ik}^{-1} Z_i - \sigma_2^{-2} Z_i^\top \Psi_{ik}^{-1} e_i e_i^\top \Psi_{ik}^{-1} Z_i \\ -\frac{1}{2} N \sigma_2^{-2} + \frac{1}{2K} \sigma_2^{-4} \sum_{k=1}^K \sum_{i=1}^M e_i^\top \Psi_{ik}^{-1} e_i \end{pmatrix},$$

where $N = \sum_i^M n_i$. Since the system of equations $\tilde{Q}_*(\theta|\theta^{(t)}) = 0$ does not have a simultaneous closed-form solution, we must resort to an iterative algorithm (e.g., Newton–Raphson). Note, however, that for fixed Ψ_{ik} at iteration t , the Q-function is maximised by

$$\hat{\beta} = \left(\sum_{i=1}^M X_i^\top \Psi_{ik}^{-1} X_i \right)^{-1} \left(\sum_{i=1}^M X_i^\top \Psi_{ik}^{-1} y_i \right).$$

Thus, for the NL, LN, and LL models, the EM estimate $\hat{\beta}$ can be seen as the solution of the generalised least squares (GLS) with weights that depend on the sampled values $w_{ik}^{(t)}$. The variance-covariance of $\hat{\beta}$,

$$\text{cov}(\hat{\beta}) = \sigma_2^2 \left(\sum_{i=1}^M X_i^\top \Psi_{ik}^{-1} X_i \right)^{-1},$$

is a by-product of fitting routines from commonly available software.

Similarly, for fixed Ψ_{ik} at iteration t , the GLS estimate of σ_2^2 is

$$\hat{\sigma}_2^2 = \left(\sum_{i=1}^M y_i^\top \Psi_{ik}^{-1} y_i \right) - \left(\sum_{i=1}^M X_i^\top \Psi_{ik}^{-1} y_i \right)^\top \left(\sum_{i=1}^M X_i^\top \Psi_{ik}^{-1} X_i \right)^{-1} \left(\sum_{i=1}^M X_i^\top \Psi_{ik}^{-1} y_i \right).$$

The E-step is updated with $\theta^{(t+1)}$ and the algorithm stops when $\Delta_{h:t,t+1} \left\{ \tilde{Q}(\theta|\theta^{(h)}) \right\} < \delta$ or $\Delta_{h:t,t+1} \left\{ \theta_l^{(h)} \right\} < \delta$, $l = 1, \dots, p + m + 1$, where $\Delta_{h:t,t+1} \left\{ u^{(h)} \right\}$ is the (absolute or relative) change in u between iterations t and $t + 1$, and δ is an appropriately small constant. The starting values $\theta^{(0)}$ can be obtained from an LME model.

Finally, standard errors for $\hat{\theta}$ can be computed using the methods described in [McLachlan and Krishnan \(2008\)](#). See also the application of Rubin’s rules for multiple imputation to Monte Carlo EM samples ([Goetghebuer and Ryan, 2000](#); [Geraci and Farcomeni, 2016](#)).

A.6. Residual heteroscedasticity and correlation. In the previous sections, I assumed that the within-group errors are independent with common scale parameter σ_2 . Using the scale mixture representation, it is immediate to extend the NL, LN, and LL models to the case of heteroscedastic and correlated errors. In particular, let's assume $\lambda_{2i} \sim \mathcal{L}_{n_i}(0, \Sigma_{2i})$ for the NL and LL convolutions, and $\nu_{2i} \sim \mathcal{N}_{n_i}(0, \Sigma_{2i})$ for the LN convolution, with general Σ_{2i} , $i = 1, \dots, M$. Then the variance-covariance matrix in (25) can be written as

$$\Omega_{ik} = \begin{cases} Z_i \Sigma_1 Z_i^\top + w_{ik}^{(t)} \Sigma_{2i} & \text{for the NL model,} \\ w_{ik}^{(t)} Z_i \Sigma_1 Z_i^\top + \Sigma_{2i} & \text{for the LN model,} \\ w_{ik,1}^{(t)} Z_i \Sigma_1 Z_i^\top + w_{ik,2}^{(t)} \Sigma_{2i} & \text{for the LL model.} \end{cases}$$

A.7. Monte Carlo. In this section, I report on the results of a small simulation study. The purpose was to investigate the bias, variance, and mean squared error (MSE) of $\hat{\beta}$ and $\hat{\xi}$ for the NN, NL, LN, and LL models when data were generated according to the following four scenarios:

1. $Y_{ij} = x_{ij}^\top \beta + z_{ij}^\top \nu_{1i} + \nu_{2ij}$,
2. $Y_{ij} = x_{ij}^\top \beta + z_{ij}^\top \nu_{1i} + \lambda_{2ij}$,
3. $Y_{ij} = x_{ij}^\top \beta + z_{ij}^\top \lambda_{1i} + \nu_{2ij}$,
4. $Y_{ij} = x_{ij}^\top \beta + z_{ij}^\top \lambda_{1i} + \lambda_{2ij}$,

where $\beta = (\beta_0, \beta_1)^\top = (1, 2)^\top$, $x_{ij} = (1, x_{1ij})^\top$, $z_{ij} = x_{ij}$, with $x_{1ij} = \gamma_i + \zeta_{ij}$, $\gamma_i \sim \mathcal{N}(0, 1)$, and $\zeta_{ij} \sim \mathcal{N}(0, 1)$. The random effects were sampled from multivariate normal (ν_1) or Laplace (λ_1) distributions with variance-covariance

$$\Sigma_1 = \begin{bmatrix} 3 & 1 \\ 1 & 2 \end{bmatrix},$$

while the errors were drawn from normal (ν_2) or Laplace (λ_2) distributions with scale $\sigma_2 = 2$, independently. The unrestricted parameter for $\sigma_2^2 \Sigma_1^{-1}$ is given by $\xi = (\xi_1, \xi_2, \xi_3)^\top = (-0.183, 0.215, -0.398)^\top$.

A balanced design with $n = 5$ repeated measurements per cluster and $M = 100$ clusters was used. For each scenario, 100 datasets were replicated. NN models were fitted using MLE routines from the `nlme` package (Pinheiro et al., 2014). The NL, LN, and LL models were fitted using the EM algorithm discussed above. In particular, Monte Carlo samples for the E-step were drawn using an adaptive rejection Metropolis sampler (Gilks, Best and Tan, 1995) as implemented in the package HI (Petrus and Tardella, 2013). The number of samples was set to increase at each EM iteration as a multiple of

20, capped at 500, thus $K = \min\{20 \cdot t, 500\}$, $t = 1, 2, \dots$. The Q-function (25) was maximised using the MLE equations for NN models (Demidenko, 2013). The convergence criterion was defined as $\Delta_{h:t,t+1} \left\{ \tilde{Q}(\theta|\theta^{(h)}) \right\} < 0.001$ and the maximum number of EM iterations was set to 100.

The results of the simulation study are reported in Tables 5-7. The NL and LL showed some advantages in terms of bias as compared to the NN model in all considered scenarios, including when the data were generated from a NN model. However, in the latter case the lower bias was more than compensated by a larger variability which made the MSE for NL and LL models up to about 44% larger than that for the NN model. In contrast, the NN model was less competitive than the NL and LL models when data were generated according to these two scenarios, with losses up to about 40% in terms of MSE.

The LN model's performance was somewhat poor, even when the data were generated from a LN model. In a separate analysis using the same data (results not shown), the NL models were re-estimated with the number K of Monte Carlo samples fixed at 500 at all iterations. The relative bias decreased to values below or near 1 for both $\hat{\beta}$ and $\hat{\xi}$, whereas the relative MSE was still above 1.

The average estimation times (standard deviation) for the NL, LN, and LL models were, respectively, 6.6 (10.0), 17.7 (25.3), and 9.8 (11.0) minutes on a 64-bit operating system machine with 16 Gb of RAM and quad-core processor at 3.60 GHz. The average number of iterations to convergence for all these three models was 14 (standard deviation 12).

TABLE 5

The estimated bias of $\hat{\beta}$ and $\hat{\xi}$ for the normal-normal (NN) model is reported in brackets. The bias for the normal-Laplace (NL), Laplace-normal (LN), and Laplace-Laplace (LL) models is relative to the NN model.

	$\hat{\beta}_0$	$\hat{\beta}_1$	$\hat{\xi}_1$	$\hat{\xi}_2$	$\hat{\xi}_3$
<i>Scenario 1: NN data</i>					
NN	(-0.012)	(0.006)	(-0.036)	(0.007)	(-0.035)
NL	0.324	0.254	1.648	-0.418	1.977
LN	0.088	1.424	-1.823	8.010	-2.997
LL	0.840	-0.369	-0.402	0.784	-0.001
<i>Scenario 2: NL data</i>					
NN	(0.002)	(-0.016)	(-0.021)	(0.014)	(-0.062)
NL	-1.301	0.503	0.970	0.848	0.881
<i>Scenario 3: LN data</i>					
NN	(-0.005)	(0.020)	(-0.072)	(0.044)	(-0.042)
LN	1.731	1.232	-0.243	2.363	-1.822
<i>Scenario 4: LL data</i>					
NN	(-0.006)	(0.012)	(-0.029)	(0.023)	(-0.079)
LL	0.716	0.483	-0.504	0.829	0.345

TABLE 6

The estimated variance of $\hat{\beta}$ and $\hat{\xi}$ for the normal-normal (NN) model is reported in brackets. The variance for the normal-Laplace (NL), Laplace-normal (LN), and Laplace-Laplace (LL) models is relative to the NN model.

	$\hat{\beta}_0$	$\hat{\beta}_1$	$\hat{\xi}_1$	$\hat{\xi}_2$	$\hat{\xi}_3$
<i>Scenario 1: NN data</i>					
NN	(0.040)	(0.028)	(0.012)	(0.005)	(0.014)
NL	1.123	1.081	1.021	0.994	1.259
LN	1.362	1.842	2.207	2.709	1.665
LL	1.287	1.441	1.174	1.246	1.087
<i>Scenario 2: NL data</i>					
NN	(0.048)	(0.029)	(0.018)	(0.007)	(0.016)
NL	0.894	1.042	0.925	0.904	0.843
<i>Scenario 3: LN data</i>					
NN	(0.034)	(0.037)	(0.039)	(0.015)	(0.040)
LN	1.126	1.146	2.020	2.211	1.691
<i>Scenario 4: LL data</i>					
NN	(0.043)	(0.024)	(0.026)	(0.011)	(0.026)
LL	0.744	0.862	0.813	0.598	0.997

TABLE 7

The estimated mean squared error (MSE) of $\hat{\beta}$ and $\hat{\xi}$ for the normal-normal (NN) model is reported in brackets. The MSE for the normal-Laplace (NL), Laplace-normal (LN), and Laplace-Laplace (LL) models is relative to the NN model.

	$\hat{\beta}_0$	$\hat{\beta}_1$	$\hat{\xi}_1$	$\hat{\xi}_2$	$\hat{\xi}_3$
<i>Scenario 1: NN data</i>					
NN	(0.040)	(0.028)	(0.013)	(0.005)	(0.016)
NL	1.119	1.080	1.190	0.987	1.471
LN	1.357	1.842	2.319	3.242	2.248
LL	1.285	1.440	1.073	1.240	1.001
<i>Scenario 2: NL data</i>					
NN	(0.048)	(0.029)	(0.018)	(0.007)	(0.020)
NL	0.894	1.035	0.925	0.898	0.830
<i>Scenario 3: LN data</i>					
NN	(0.034)	(0.038)	(0.044)	(0.017)	(0.042)
LN	1.127	1.150	1.787	2.598	1.759
<i>Scenario 4: LL data</i>					
NN	(0.043)	(0.025)	(0.027)	(0.012)	(0.033)
LL	0.744	0.859	0.796	0.602	0.830

REFERENCES

- ARORA, R. S., KONTOPANTELOS, E., ALSTON, R. D., EDEN, T. O. B., GERACI, M. and BIRCH, J. M. (2011). Relationship between height at diagnosis and bone tumours in young people: A meta-analysis. *Cancer Causes and Control* **22** 681-688.
- BARNECHE, D. R., KULBICKI, M., FLOETER, S. R., FRIEDLANDER, A. M. and ALLEN, A. P. (2016). Energetic and ecological constraints on population density of reef fishes. *Proceedings of the Royal Society B: Biological Sciences* **283** 1-8.
- BASSETT, G. and KOENKER, R. (1978). Asymptotic theory of least absolute error regression. *Journal of the American Statistical Association* **73** 618-622.
- BEETS, M. W., WEAVER, R. G., TURNER-MCGRIEVY, G., MOORE, J. B., WEBSTER, C., BRAZENDALE, K. et al. (2016). Are we there yet? Compliance with physical activity standards in YMCA afterschool programs. *Childhood Obesity* **12** 237-246.
- BLANKENBERG, S., SALOMAA, V., MAKAROVA, N., OJEDA, F., WILD, P., LACKNER, K. J. et al. (2016). Troponin I and cardiovascular risk prediction in the general population: the BiomarCaRE consortium. *European Heart Journal* **37** 2428-2437.
- BOX, G. E. P. (1950). Problems in the analysis of growth and wear curves. *Biometrics* **6** 362-389.
- DEGERUD, E., LØLAND, K. H., NYGÅRD, O., MIDTTUN, Ø., UELAND, P. M., SEIFERT, R. et al. (2014). Vitamin D status was not associated with 'one-year' progression of coronary artery disease, assessed by coronary angiography in statin-treated patients. *European Journal of Preventive Cardiology* **22** 594-602.
- DEMIDENKO, E. (2013). *Mixed models: Theory and applications with R*, Second ed. John Wiley & Sons, Hoboken, NJ.
- DERSIMONIAN, R. and LAIRD, N. (1986). Meta-analysis in clinical trials. *Control Clinical Trials* **7** 177-88.
- DOUGLAS, B., MÄCHLER, M., BOLKER, B. and WALKER, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67** 1-48.
- DUFFY, L. M., OLSON, R. J., LENNERT-CODY, C. E., GALVÁN-MAGAÑA, F., BOCANEGRA-CASTILLO, N. and KUHNERT, P. M. (2015). Foraging ecology of silky sharks, *Carcharhinus falciformis*, captured by the tuna purse-seine fishery in the eastern Pacific Ocean. *Marine Biology* **162** 571-593.
- FORNAROLI, R., CABRINI, R., SARTORI, L., MARAZZI, F., VRACEVIC, D., MEZZANOTTE, V. et al. (2015). Predicting the constraint effect of environmental characteristics on macroinvertebrate density and diversity using quantile regression mixed model. *Hydrobiologia* **742** 153-167.
- GERACI, M. (2014). Linear quantile mixed models: The lqmm package for Laplace quantile regression. *Journal of Statistical Software* **57** 1-29.
- GERACI, M. and BOTTAI, M. (2007). Quantile regression for longitudinal data using the asymmetric Laplace distribution. *Biostatistics* **8** 140-154.
- GERACI, M. and BOTTAI, M. (2014). Linear quantile mixed models. *Statistics and Computing* **24** 461-479.
- GERACI, M. and FARCOMENI, A. (2016). Probabilistic principal component analysis to identify profiles of physical activity behaviours in the presence of nonignorable missing data. *Journal of the Royal Statistical Society C* **65** 51-75.
- GILKS, W. R., BEST, N. G. and TAN, K. K. C. (1995). Adaptive rejection metropolis sampling within Gibbs sampling. *Journal of the Royal Statistical Society Series C* **44** 455-472.
- GOETGHEBEUR, E. and RYAN, L. (2000). Semiparametric regression analysis of interval-censored data. *Biometrics* **56** 1139-1144.

- KOENKER, R. and BASSETT, G. (1978). Regression quantiles. *Econometrica* **46** 33-50.
- KOENKER, R. and NG, P. (2005). A Frisch–Newton algorithm for sparse quantile regression. *Acta Mathematicae Applicatae Sinica (English Series)* **21** 225-236.
- KOTZ, S., KOZUBOWSKI, T. J. and PODGÓRSKI, K. (2001). *The Laplace distribution and generalizations*. Birkhäuser, Boston, MA.
- KOZUBOWSKI, T. J. and NADARAJAH, S. (2010). Multitude of Laplace distributions. *Statistical Papers* **51** 127-148.
- KUNDU, D. (2005). Discriminating between normal and Laplace distributions. In *Advances in Ranking and Selection, Multiple Comparisons, and Reliability: Methodology and Applications* (N. Balakrishnan, H. N. Nagaraja and N. Kannan, eds.) 65-79. Birkhäuser Boston, Boston, MA.
- LUM, K. and GELFAND, A. (2012). Spatial quantile multiple regression using the asymmetric Laplace process. *Bayesian Analysis* **7** 235-258.
- McLACHLAN, G. and KRISHNAN, T. (2008). *The EM algorithm and extensions*. John Wiley & Sons, Hoboken, NJ.
- MEINTANIS, S. G. and TSIONAS, E. (2010). Testing for the generalized normal-Laplace distribution with applications. *Computational Statistics & Data Analysis* **54** 3174-3180.
- MOOD, A., GRAYBILL, F. and BOES, D. (1974). *Introduction to the theory of statistics*, Third ed. McGraw-Hill, New York, NY.
- MUIR, P. R., WALLACE, C. C., DONE, T. and AGUIRRE, J. D. (2015). Limited scope for latitudinal extension of reef corals. *Science* **348** 1135-1138.
- NG, S. W., HOWARD, A. G., WANG, H. J., SU, C. and ZHANG, B. (2014). The physical activity transition among adults in China: 1991-2011. *Obesity Reviews* **15** 27-36.
- PATEL, D. E., GERACI, M. and CORTINA-BORJA, M. (2016). Modelling normative kinetic perimetry isopters using mixed-effects quantile regression. *Journal of Vision* **16** 1-6.
- PATEL, D. E., CUMBERLAND, P. M., WALTERS, B. C., RUSSELL-EGGITT, I., CORTINA-BORJA, M. and RAHI, J. S. (2015). Study of optimal perimetric testing in children (OPTIC): Normative visual field values in children. *Ophthalmology* **122** 1711-1717.
- PETRIS, G. and TARDELLA, L. (2013). HI: Simulation from distributions supported by nested hyperplanes R package version 0.4. <http://CRAN.R-project.org/package=HI>.
- PINHEIRO, J. C. and BATES, D. M. (2000). *Mixed-effects models in S and S-PLUS*. Springer Verlag, New York.
- PINHEIRO, J., BATES, D., DEBROY, S., SARKAR, D. and R CORE TEAM (2014). nlme: Linear and nonlinear mixed effects models R package version 3.1-117. <http://CRAN.R-project.org/package=nlme>.
- PORTNOY, S. and KOENKER, R. (1997). The Gaussian hare and the Laplacian tortoise: Computability of squared-error versus absolute-error estimators. *Statistical Science* **12** 279-300.
- PRÉKOPA, A. (1973). Logarithmic concave measures and functions. *Acta Scientiarum Mathematicarum* **34** 334-343.
- REED, W. J. (2006). The normal-Laplace distribution and its relatives. In *Advances in Distribution Theory, Order Statistics, and Inference* (N. Balakrishnan, E. Castillo and J. M. Sarabia, eds.) 61-74. Birkhäuser, Boston, MA.
- REED, W. J. (2007). Brownian-Laplace motion and its use in financial modelling. *Communications in Statistics – Theory and Methods* **36** 473-484.
- REED, W. J. and JORGENSEN, M. (2004). The double Pareto-lognormal distribution—A new parametric model for size distributions. *Communications in Statistics – Theory and Methods* **33** 1733-1753.
- SORRENTINO, D., PAVIOTTI, A., TERROSU, G., AVELLINI, C., GERACI, M. and ZARIFI, D. (2010). Low-dose maintenance therapy with infliximab prevents postsurgical recurrence

- of Crohn's disease. *Clinical Gastroenterology and Hepatology* **8** 591-599.
- R CORE TEAM (2014). R: A Language and environment for statistical computing R Foundation for Statistical Computing, Vienna, Austria <http://www.R-project.org>.
- TIAN, Y., TIAN, M. and ZHU, Q. (2013). Linear quantile regression based on EM algorithm. *Communications in Statistics – Theory and Methods* **43** 3464-3484.
- VIECHTBAUER, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software* **36** 1-48.
- WILSON, E. B. (1923). First and second laws of error. *Journal of the American Statistical Association* **18** 841-851.
- YU, K. M., LU, Z. D. and STANDER, J. (2003). Quantile regression: Applications and current research areas. *Journal of the Royal Statistical Society D* **52** 331-350.
- YU, K. M. and MOYEED, R. A. (2001). Bayesian quantile regression. *Statistics & Probability Letters* **54** 437-447.
- YU, K. M. and ZHANG, J. (2005). A three-parameter asymmetric Laplace distribution and its extension. *Communications in Statistics – Theory and Methods* **34** 1867-1879.