

Multivariate Extremes Over a Random Number of Observations

Enkelejd Hashorva^{1,*}, Simone A. Padoan^{2,**} and Stefano Rizzelli^{3,†}

¹*Department of Actuarial Science, University of Lausanne, UNIL-Dorigny 1015 Lausanne, Switzerland.
e-mail: *enkelejd.hashorva@unil.ch*

²*Department of Decision Sciences, Bocconi University, via Roentgen 1, 20136 Milan, Italy.
e-mail: **simone.padoan@unibocconi.it*

³*Institute of Mathematics, École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland.
e-mail: †stefano.rizzelli@epfl.ch*

Abstract: The classical multivariate extreme-value theory concerns the modeling of extremes in a multivariate random sample, suggesting the use of max-stable distributions. In this work, the classical theory is extended to the case where aggregated data, such as maxima of a random number of observations, are considered. We derive a limit theorem concerning the attractors for the distributions of the aggregated data, which boil down to a new family of max-stable distributions. We also connect the extremal dependence structure of classical max-stable distributions and that of our new family of max-stable distributions. By means of an inversion method, we derive a semiparametric composite-estimator for the extremal dependence of the unobservable data, starting from a preliminary estimator of the extremal dependence of the aggregated data. Furthermore, we develop the large-sample theory of the composite-estimator and illustrate its finite-sample performance via a simulation study.

MSC 2010 subject classifications: Primary 62G32, 62G05, 62G20; secondary 60F05, 60G70.

Keywords and phrases: Extremal dependence, Extreme-value copula, Inverse problem, Multivariate max-stable distribution, Nonparametric estimation, Pickands dependence function.

1. Introduction and background

The multivariate extreme-value theory aims to quantify the probability of extreme events concerning multiple dependent observations. A commonly employed approach for modelling extremes in high dimensions is the componentwise maxima, where, for each of the involved variables, the partial maximum values are taken into account, e.g., yearly maxima, (e.g., [Falk et al., 2011](#), Ch. 4). Basic foundations of the componentwise maxima approach are here briefly introduced.

First, however, we specify the notation that we use throughout the paper. Given $\mathcal{X} \subset \mathbb{R}^n$, $n \in \mathbb{N}$, let $\ell^\infty(\mathcal{X})$ denote the spaces of bounded real-valued functions on \mathcal{X} . For $f : \mathcal{X} \mapsto \mathbb{R}$, let $\|f\|_\infty = \sup_{x \in \mathcal{X}} |f(x)|$. The arrows “ $\xrightarrow{\text{as}}$ ”, “ $\xrightarrow{\text{P}}$ ”, “ \rightsquigarrow ” denote convergence (outer) almost surely, convergence in (outer) probability and convergence in distribution of random vectors (see [van der Vaart, 2000](#), Ch. 2) or weak convergence of random functions in $\ell^\infty(\mathcal{X})$ (see [van der Vaart, 2000](#), Ch. 18–19), the distinction between the two will be clear from the context. For a non-decreasing function f , let f^\leftarrow denote the left-continuous inverse of f . The abbreviation $a \sim b$ stands for a is asymptotically equivalent to b . Finally, the multiplication, division and maximum operation between vectors is meant componentwise.

Let $\mathbf{X} = (X_1, \dots, X_d)$ be a d -dimensional random vector with distribution $F_{\mathbf{X}}$ and margins F_{X_j} , $j = 1, \dots, d$, and X_1, X_2, \dots be independent and identically distributed (iid) copies of \mathbf{X} . Assume that $F_{\mathbf{X}}$ is in the maximum-domain of attraction (simply domain of attraction) of a multivariate extreme-value distribution G , in symbols $F_{\mathbf{X}} \in \mathcal{D}(G)$. This means that there are sequences of constants $\mathbf{a}_n > \mathbf{0} = (0, \dots, 0)$ and $\mathbf{b}_n \in \mathbb{R}^d$ such that $(\max(X_1, \dots, X_n) - \mathbf{b}_n)/\mathbf{a}_n \rightsquigarrow \boldsymbol{\eta}$ as $n \rightarrow \infty$, where the distribution of $\boldsymbol{\eta}$ is a multivariate extreme-value distribution (e.g., [Falk et al., 2011](#), pp. 147-153) of the form

$$G(\mathbf{x}) = C_G(G_1(x_1), \dots, G_d(x_d)), \quad \mathbf{x} \in \mathbb{R}^d.$$

Precisely, G_j ’s are members of the generalized extreme-value distribution (GEV) (e.g., [Falk et al., 2011](#), p. 21) and C_G is an extreme-value copula, i.e.,

$$C_G(\mathbf{u}) = \exp(-L((-\ln u_1), \dots, (-\ln u_d))), \quad \mathbf{u} \in (0, 1]^d,$$

where $L : [0, \infty)^d \mapsto [0, \infty)$ is the so-called stable-tail dependence function (e.g., [Falk et al., 2011](#), pp. 177–179). G is a max-stable distribution, i.e., for $k = 1, 2, \dots$, there are norming sequences $\mathbf{a}'_k > \mathbf{0}$ and $\mathbf{b}'_k \in \mathbb{R}^d$ such that $G^k(\mathbf{a}'_k \mathbf{x} + \mathbf{b}'_k) = G(\mathbf{x})$, for all $\mathbf{x} \in \mathbb{R}^d$. The extreme-value copula expresses the dependence among extremes. Examples of parametric extreme-value copula models are: the Logistic or Gumbel, the Hüsler-Reiss and the Extremal- t , just to name a few. An extensive list of additional models is available in [Joe \(2015, Ch. 4\)](#). Since L is a homogeneous function of order 1, it can be conveniently represented as

$$L(\mathbf{z}) = (z_1 + \dots + z_d)A(\mathbf{t}), \quad \mathbf{z} \in [0, \infty)^d, \quad (1.1)$$

where $t_j = z_j/(z_1 + \dots + z_d)$ for $j = 1, \dots, d$. The function A , named Pickands (dependence) function, denotes the restriction of L on the d -dimensional unit simplex $\mathcal{S}_d := \{(v_1, \dots, v_d) \in [0, 1]^d : v_1 + \dots + v_d = 1\}$. It summarizes the extremal dependence among the components of $\boldsymbol{\eta}$, specifically it holds that $1/d \leq \max(t_1, \dots, t_d) \leq A(\mathbf{t}) \leq 1$, where the lower and upper bounds represent the cases of complete dependence and independence. A synthesis of the extremal dependence is provided by the extremal coefficient, that is $\theta(G) = dA(1/d, \dots, 1/d) \in [1, d]$. It can be interpreted as the (fractional) number of independent variables in a d -dimensional random vector with joint distribution G and common margins. An alternative summary index that measures the dependence among observations falling in the upper tail region is the coefficient of upper tail dependence (e.g., [Joe, 2015, Ch. 2.13](#)). In the bivariate case it is equal to

$$\lambda(F_X) = \lim_{u \uparrow 1} \mathbb{P}(X_1 > F_{X_1}^{-1}(u) | X_2 > F_{X_2}^{-1}(u)) = \lim_{u \uparrow 1} \mathbb{P}(X_2 > F_{X_2}^{-1}(u) | X_1 > F_{X_1}^{-1}(u)) \in [0, 1].$$

It is said that F_X exhibits independence or dependence in the upper tail whenever $\lambda(F_X) = 0$ or $\lambda(F_X) > 0$, respectively, with the case of complete dependence covered when $\lambda(F_X) = 1$. The coefficient $\lambda(F_X)$ is linked to the extremal coefficient by the relationship $\theta(G) = 2 - \lambda(F_X)$.

Nowadays, applications involving complex phenomena frequently deal with the analysis of aggregated data. This is especially true in big-data problems, where it may be convenient (or unavoidable) to work with aggregated data in order to reduce the computational cost. Examples of aggregated data are the total amounts and maximum amounts obtained on a random number of observations. Such aggregated data are realizations of the random vectors

$$\mathbf{S}_N = \left(\sum_{i=1}^N X_{i,1}, \dots, \sum_{i=1}^N X_{i,d} \right), \quad \mathbf{M}_N = \left(\max_{1 \leq i \leq N} X_{i,1}, \dots, \max_{1 \leq i \leq N} X_{i,d} \right), \quad (1.2)$$

where N is a discrete random variable defined on $\mathbb{N}_+ = \mathbb{N} \setminus \{0\}$. Assume hereafter that N with distribution F_N is independent of X_i 's. For dimension $d = 1$ the tail behaviour of S_N has been extensively studied in the literature (e.g., [Embrechts et al., 1997](#); [Robert and Segers, 2008](#)) and only few results are known on the extremes of M_N ([Barakat et al., 1990](#); [Silvestrov and Teugels, 1998](#)).

The first main purpose of this contribution is to extend the classical probabilistic theory on the extreme-values to the case when the latter are computed using replicates of \mathbf{M}_N (the random vector that represents aggregated data). In Section 5 we illustrate an application, in the context of big data on Internet traffic, which would benefit from such new theoretical developments. In the literature there are no results which can point to how different the extremal behaviour of F_{M_N} is with respect to F_X , some preliminary findings are available in [Hashorva et al. \(2017\)](#). In this work, with Theorem 2.1, we provide the attractor for the joint distribution of (\mathbf{M}_N, N) , where from the extremal behaviour of \mathbf{M}_N (our main focus) can be deduced, such as the tail behaviours of F_{M_N} .

Some more interesting results are obtained in the case where F_N is very heavy-tailed, which appears to be the more relevant one in a context of big-data, as more data are produced and aggregating them is beneficial. Specifically, when $\mathbb{P}(N > y) = y^{-\alpha} \mathcal{L}(y)$, with $y > 0$ and $\alpha \in (0, 1)$, where \mathcal{L} is a slowly varying function at infinity, and $F_X \in \mathcal{D}(G)$, we show that \mathbf{M}_N and N are asymptotically dependent. Furthermore, we find that in this case $F_{M_N} \in \mathcal{D}(G_\alpha)$, where G_α is a new max-stable distribution with an extreme-value copula C_{G_α} , given in (2.4), differing from C_G , the extreme-value copula of G . The coefficient $\alpha \in (0, 1)$ influences the extremal dependence structure of G_α . A practical implication of our finding is that the extreme properties of F_{M_N} can be recovered by knowing the extremal properties of F_X and the tail behaviour of N . Here are two examples. The joint upper-tail probability of \mathbf{M}_N can be approximated as

$$\mathbb{P} \left(F_{M_N^{(1)}}(M_N^{(1)}) > 1 - \frac{1}{y_1} \text{ or } \dots \text{ or } F_{M_N^{(d)}}(M_N^{(d)}) > 1 - \frac{1}{y_d} \right) \sim L^\alpha(\mathbf{y}^{-1/\alpha}),$$

for a large enough $y > 0$, where $M_N^{(j)} = \max_{1 \leq i \leq N} X_{i,j}$, $j = 1, \dots, d$ and L is the stable-tail dependence function of C_G (see Proposition 2.2). This means that by combining L and α , we can approximate the probability that at least one component among $M_N^{(1)}, \dots, M_N^{(d)}$ exceeds a high percentile of its own distribution. When $d = 2$, we also have

$$\lambda(F_{M_N}) = 2 - [2 - \lambda(F_X)]^\alpha. \quad (1.3)$$

The second purpose of this contribution concerns an inverse statistical problem in the context of extremes of aggregated data, which also motivates the study of the asymptotic joint distribution of (M_N, N) . Precisely, in applications where the variables M_N and N are observable in place of X , the interest may however be in inferring the extremal dependence of the distribution G . Theorem 2.1 provides the mathematical ground to address this issue, as it gives a joint (limiting) statistical model for the sample extremes of M_N and N . In particular, the sample maxima of the latter random vector and variable can be used to obtain estimators of $\lambda(F_{M_N})$ and α , respectively. Then, by exploiting the relation among extremal properties of F_{M_N} and F_X and the tail behaviour of N , and solving (1.3) for $\lambda(F_X)$, an estimator of the latter can be obtained. More generally, we focus on the Pickands function as it allows to derive the λ and θ coefficients and other related quantities. We define a new semi-parametric procedure for inferring A , which combines together preliminary estimators for α and A_α (the Pickands function relative to C_{G_α}). Specifically, we consider a likelihood- and moments-based estimator for α and we use three nonparametric estimators for A_α (existing in the literature).

The rest of the paper is organized as follows. In Section 2, we present our main theorem providing the attractor for the joint distribution of (M_N, N) . Different representations for the distribution G_α are derived in Section 3. In Section 4, by an inversion method, we define an estimator for inferring A . We establish its asymptotic properties (Theorem 4.1) and show its finite-sample performance by a simulation study. Finally, we discuss directions for future research in Section 5, including a real data example which appears as a promising field of application for our theoretical framework. The proofs are reported in the Appendix, whereas some technical details and additional simulation results are included in the supplementary material.

2. Main Results

First, recall that the members of the GEV distribution are: the α -Fréchet (heavy-tailed distribution), Gumbel (light-tailed distribution) and Weibull (short-tailed distribution), in symbols, $\Phi_\alpha(x) = \exp(-x^{-\alpha})$, with $x > 0$ and $\alpha > 0$, $\Lambda(x) = \exp(-e^{-x})$ with $x \in \mathbb{R}$ and $\Psi_\alpha(x) = \exp(-(-x)^{-\alpha})$ with $x < 0$. In the sequel, for a positive random variable, say V , we denote its Laplace transform by $L_V(s) = E(e^{-sV})$, $s > 0$. We also recall that a random variable, S , is positive (asymmetric) α -stable with index parameter $0 < \alpha < 1$ if its Laplace transform is $L_S(s) = e^{-s^{-\alpha}}$.

Let N be a random block size and M_N be a vector of componentwise maxima obtained with a randomly sized block of iid random vectors X_1, X_2, \dots with common distribution F_X , defined in (1.2). Note that

$$F_{M_N}(x) = \sum_{n=1}^{\infty} F_X^n(x) \mathbb{P}(N = n) = \sum_{n=1}^{\infty} \exp[-n\{-\ln F_X(x)\}] \mathbb{P}(N = n) = L_N(-\ln F_X(x)).$$

Assuming that $F_X \in \mathcal{D}(G)$ and $F_N \in \mathcal{D}(H)$, where either $H \equiv \Phi_\alpha$ or $H \equiv \Lambda$ since N is positive integer-valued (Robert and Segers, 2008), we establish new limit results concerning the attractor for the joint distribution of the random vector (M_N, N) and the tail behaviour of the random vector M_N .

2.1. Domains of Attraction

The first limit result provides the attractor for the joint distribution of the random vector (M_N, N) .

Theorem 2.1. Assume that $F_X \in \mathcal{D}(G)$ and $F_N \in \mathcal{D}(H)$ with $H \equiv \Phi_\alpha$ or $H \equiv \Lambda$. Then, there exist norming constants $c_n > 0, \kappa_n > 0, d_n \in \mathbb{R}^d, \varrho_n \in \mathbb{R}$ such that

$$\lim_{n \rightarrow \infty} \mathbb{P}^n \left(\frac{M_N - d_n}{c_n} \leq x, \frac{N - \varrho_n}{\kappa_n} \leq y \right) = Q(x, y),$$

where Q is defined as follows:

1. if $F_N \in \mathcal{D}(\Phi_\alpha)$, then

$$-\ln Q(\mathbf{x}, y) = \begin{cases} y^{-\alpha} e^{-y\sigma(\mathbf{x}, \alpha)} + \sigma^\alpha(\mathbf{x}, \alpha) \gamma(1 - \alpha, y\sigma(\mathbf{x}, \alpha)), & \alpha \in (0, 1) \\ -\ln G(\mathbf{x}) + y^{-\alpha}, & \alpha \geq 1 \end{cases} \quad (2.1)$$

for all $\mathbf{x} \in \mathbb{R}^d$ and $y > 0$, where $\sigma(\mathbf{x}, \alpha) = (-\ln G(\mathbf{x}))/\Gamma^{1/\alpha}(1 - \alpha)$ and Γ, γ denote the Euler Gamma and Lower Incomplete Gamma functions, respectively.

2. if $F_N \in \mathcal{D}(\Lambda)$, then

$$-\ln Q(\mathbf{x}, y) = -\ln G(\mathbf{x}) + e^{-y}, \quad \mathbf{x} \in \mathbb{R}^d, y \in \mathbb{R}. \quad (2.2)$$

The margins of Q are

$$\begin{cases} G_\alpha(\mathbf{x}) := \exp(-(-\ln G(\mathbf{x}))^\alpha), & \alpha \in (0, 1) \\ G(\mathbf{x}), & \alpha \geq 1 \end{cases} \quad (2.3)$$

and $\Phi_\alpha(y)$, when $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha > 0$. While they are equal to $G(\mathbf{x})$, $\mathbf{x} \in \mathbb{R}^d$, and $\Lambda(y)$, $y \in \mathbb{R}$, when $F_N \in \mathcal{D}(\Lambda)$. Specifically, the distribution G_α , $\alpha \in (0, 1)$, is max-stable with margins $G_{\alpha,1}, \dots, G_{\alpha,d}$, that are members of the GEV class, and extreme-value copula

$$C_{G_\alpha}(\mathbf{u}) = \exp(-L^\alpha((-\ln u_1)^{1/\alpha}, \dots, (-\ln u_d)^{1/\alpha})), \quad \mathbf{u} \in (0, 1]^d, \quad \alpha \in (0, 1), \quad (2.4)$$

where L is the stable-tail dependence function of the max-stable distribution G .

A probabilistic interpretation of the problem addressed in Theorem 2.1 is as follows. Let N_1, N_2, \dots be iid copies of N and set $N_n^+ = N_1 + \dots + N_n$, $N_n^\vee = \max(N_1, \dots, N_n)$, then we have

$$\mathbb{P}\left(\frac{\mathbf{M}_{N_n^+} - \mathbf{d}_n}{\mathbf{c}_n} \leq \mathbf{x}, \frac{N_n^\vee - \varrho_n}{\kappa_n} \leq y\right) = \mathbb{P}^n\left(\frac{\mathbf{M}_N - \mathbf{d}_n}{\mathbf{c}_n} \leq \mathbf{x}, \frac{N - \varrho_n}{\kappa_n} \leq y\right).$$

Loosely speaking, we are concerned with the asymptotic distribution of the random vector $(\mathbf{M}_{N_n^+}, N_n^\vee)$ appropriately normalized (a.n.). When $F_N \in \mathcal{D}(\Lambda)$ (light-tailed) or $F_N \in \mathcal{D}(\Phi_\alpha)$ (heavy-tailed), with $\alpha > 1$, then $\mu = E(N) < \infty$. Since $n^{-1}N_n^+$ converges to μ , then the asymptotic distributions of $\mathbf{M}_{N_n^+}$ and $\mathbf{M}_{[n\mu]}$ a.n. are approximately the same. When $F_N \in \mathcal{D}(\Phi_\alpha)$ (heavy-tailed), with $\alpha \in (0, 1)$, then $\tilde{\kappa}_n^{-1}N_n^+$ converges in distribution to a positive stable random variable S , where $\tilde{\kappa}_n := \kappa_n \Gamma^{1/\alpha}(1 - \alpha)$. Consequently, the asymptotic distributions of $\mathbf{M}_{N_n^+}$ and $\mathbf{M}_{[\tilde{\kappa}_n S]}$ a.n. coincide and are equal to G_α in the first line of (2.3), see Corollary 2.2, which is a location-scale mixture of the limiting max-stable distribution G , obtained with a deterministic block size. A similar result can be established in the case of $\alpha = 1$, which is not explicitly discussed here for the sake of brevity.

Corollary 2.2. Let $F_X \in \mathcal{D}(G)$, $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \in (0, 1)$ and $\mathbf{c}_n, \mathbf{d}_n$ and κ_n as in Theorem 2.1. Then $\tilde{\kappa}_n^{-1}N_n^+ \rightsquigarrow S$, as $n \rightarrow \infty$, where $\tilde{\kappa}_n = \kappa_n \Gamma^{1/\alpha}(1 - \alpha)$ and

$$\lim_{n \rightarrow \infty} \mathbb{P}(\mathbf{M}_{[\tilde{\kappa}_n S]} \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n) = G_\alpha(\mathbf{x}).$$

As for asymptotic dependence, we point out the following. When $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \geq 2$, or $F_N \in \mathcal{D}(\Lambda)$, N_n^+ and N_n^\vee a.n. converge to nondegenerate independent random variables (e.g. Anderson and Turkman, 1995, pp. 1-2). Intuitively, this explains the asymptotic independence between $\mathbf{M}_{N_n^+}$ and N_n^\vee (and, in turn, the extremal independence between \mathbf{M}_N and N). In the special case of $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \in (0, 2)$, N_n^+ and N_n^\vee a.n. are asymptotically dependent (e.g. Anderson and Turkman, 1995, pp. 1-2). Thus, when $\alpha \in [1, 2)$ we explain the asymptotic independence between $\mathbf{M}_{N_n^+}$ and N_n^\vee a.n. in a different way. From the derivations in Lemmas A.2-A.3 and Section A.1.3 one sees that the dependence structure of the limiting distribution $Q(\mathbf{x}, y)$ is determined by the limit of the conditional exceedance probability $1 - \mathbb{P}(\mathbf{M}_N \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n | N > \kappa_n y + \varrho_n)$, as $n \rightarrow \infty$. Note that, as n grows, the norming sequences $(\mathbf{c}_n, \mathbf{d}_n)$ and (κ_n, ϱ_n) affect this threshold exceedance probability in two opposing ways: the first ones increase the threshold $\mathbf{c}_n \mathbf{x} + \mathbf{d}_n$; the second ones force N , and thus \mathbf{M}_N , to be stochastically larger and larger. Clearly, a non-degenerate limit – i.e. in $(0, 1)$ – is obtained only if the two effects offset each other.

On one hand

$$F_{\mathbf{M}_{N_n^+}}(\mathbf{c}_n \mathbf{x} + \mathbf{d}_n) = \mathbb{L}_N^n(-\ln F_X(\mathbf{c}_n \mathbf{x} + \mathbf{d}_n))$$

and so \mathbf{c}_n and \mathbf{d}_n are affected by the behaviour of $1 - \mathbf{L}_N(1/y)$ as $y \rightarrow \infty$. On the other hand $F_{N_n^\vee}(\kappa_n y + \varrho_n) = F_N^n(\kappa_n y + \varrho_n)$ and so κ_n and ϱ_n depend on the tail properties of F_N . When $\alpha \in (0, 1)$, $\{1 - F_N(y)\}/\{1 - \mathbf{L}_N(1/y)\}$ converges to a nondegenerate limit as $y \rightarrow \infty$, while it converges to zero when $1 \leq \alpha < 2$. Accordingly, in the first case, the limit of the conditional probability of exceedances is positive, whereas it is zero in the second case, since $\mathbf{c}_n, \mathbf{d}_n$ are “too heavy” relative to κ_n, ϱ_n . As a result, the marginal distributions of $\mathbf{M}_{N_n^+}$ and N_n^\vee a.n. converge to non-degenerate limits but their dependence is wiped out as $n \rightarrow \infty$.

The dependence structure of the $(d+1)$ -dimensional distribution Q defined in (2.1) and (2.2) can be synthesized by means of its extremal coefficient.

Corollary 2.3. *When the expression of Q is given in (2.1), then the extremal coefficient is*

$$\theta(Q) \equiv \theta(G, \alpha) := \begin{cases} 1 - \mathcal{E}_{\Gamma^{1/\alpha}(1-\alpha)}(\theta(G)) + (\theta(G))^\alpha \mathcal{G}_{1-\alpha, \Gamma^{1/\alpha}(1-\alpha)}(\theta(G)), & \alpha \in (0, 1) \\ \theta(G) + 1, & \alpha \geq 1. \end{cases},$$

where $\mathcal{E}_b(\cdot)$ and $\mathcal{G}_{a,b}(\cdot)$ denote the Exponential and Gamma cumulative distribution functions, with shape and scale parameters a and b . In particular, for $\alpha \in (0, 1)$ we have

$$1 \leq \theta(G, \alpha) \leq 1 + \theta(G), \quad \lim_{\alpha \rightarrow 1^-} \theta(G, \alpha) = 1 + \theta(G). \quad (2.5)$$

When the expression of Q is given in (2.2), $\theta(Q) = \theta(G) + 1$.

From the first result in the left-hand side of (2.5) we deduce that the extremal coefficient of Q is larger than or equal to 1 (as expected) and bounded from above by $1 + \theta(G)$, representing the case where \mathbf{M}_N and N have no tail dependence. Moreover, the second result in the right-hand side of (2.5) highlights a continuous transition of the extremal dependence level from asymptotic dependence between \mathbf{M}_N and N (i.e. $\alpha \in (0, 1)$) to asymptotic independence (i.e. $\alpha \geq 1$), when F_N belongs to the α -Fréchet domain.

The extremal coefficient for the distribution G_α with $\alpha \in (0, 1)$, $\theta(G_\alpha)$, is given in (3.5). In the bivariate case, by (3.5) and the relationship between the extremal coefficient and the coefficient of upper tail dependence, i.e., $\theta(G) = 2 - \lambda(F_X)$, we obtain

$$\theta(G_\alpha) = [2 - \lambda(F_X)]^\alpha.$$

Since it is also true that $\theta(G_\alpha) = 2 - \lambda(F_{\mathbf{M}_N})$, we obtain the result (1.3).

2.2. Tail Behaviours

The second limit result establishes the tail behaviours of the random vector \mathbf{M}_N , which concerns the probability that at least one component of the random vector \mathbf{M}_N exceeds an increasingly large value.

In the sequel, for a given max-stable distribution G we denote by \tilde{G} a distribution with the same copula as G and common unit-Fréchet margins.

Proposition 2.4. *Assume that $F_X \in \mathcal{D}(G)$ and $F_N \in \mathcal{D}(H)$ with $H \equiv \Phi_\alpha$ or $H \equiv \Lambda$. For $\mathbf{y} \in (0, \infty)^d$ and $n \in \mathbb{N}_+$,*

1. *if $F_N \in \mathcal{D}(\Phi_\alpha)$ with $0 < \alpha \leq 1$, then*

$$\begin{aligned} 1 - \mathbb{P}(\mathbf{M}_N \leq U_X(n\mathbf{y})) &\sim \begin{cases} \Gamma(1-\alpha) \mathbb{P}\left(N > \frac{n}{-\ln \tilde{G}(\mathbf{y})}\right), & \alpha \in (0, 1) \\ \{1 - \mathbf{L}_N(1/n)\} \{-\ln \tilde{G}(\mathbf{y})\}, & \alpha = 1 \end{cases}, \quad n \rightarrow \infty \\ 1 - \mathbb{P}(\mathbf{M}_N \leq U_{\mathbf{M}_N}(n\mathbf{y})) &\sim \{-\ln \tilde{G}(\mathbf{y}^{1/\alpha})\}^\alpha n^{-1}, \quad n \rightarrow \infty \end{aligned}$$

2. *if $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha > 1$ or $F_N \in \mathcal{D}(\Lambda)$, then*

$$\begin{aligned} 1 - \mathbb{P}(\mathbf{M}_N \leq U_X(n\mathbf{y})) &\sim n^{-1} E(N) \{-\ln \tilde{G}(\mathbf{y})\}, \quad n \rightarrow \infty \\ 1 - \mathbb{P}(\mathbf{M}_N \leq U_{\mathbf{M}_N}(n\mathbf{y})) &\sim n^{-1} \{-\ln \tilde{G}(\mathbf{y})\}, \quad n \rightarrow \infty \end{aligned}$$

where $U_X(n\mathbf{y}), U_{\mathbf{M}_N}(n\mathbf{y}) \rightarrow \infty$ as $n \rightarrow \infty$, with U_X and $U_{\mathbf{M}_N}$ defined in Appendix A.4.

Set $p_j = (ny_j)^{-1}$, $j = 1, \dots, d$, and recall that L denotes the stable-tail dependence function of G . As $n \rightarrow \infty$, by Proposition 2.4, the probability that at least one component $\mathbf{M}_N^{(j)}$ of \mathbf{M}_N exceeds the $1 - p_j$ quantile of its own distribution is approximately $L((ny)^{-1})$, when $E(N) < \infty$, while it is approximately $L^\alpha((ny)^{-1/\alpha})$, when $E(N) = \infty$.

3. Representations of the model G_α

In this section we show that there are different constructions that yield a max-stable distribution with the same copula C_{G_α} in (2.4) of the distribution G_α . Furthermore, we derive the Pickands function corresponding to C_{G_α} .

Let S be a positive α -stable random variable with index parameter $0 < \alpha < 1$. Let \mathbf{Z} be a random vector with max-stable distribution \tilde{G} . Assume S and \mathbf{Z} to be independent. Define $\mathbf{R} = (SZ_1, \dots, SZ_d)$, then for every $\mathbf{y} > \mathbf{0}$,

$$\mathbb{P}(\mathbf{R} \leq \mathbf{y}) = \mathbb{E}(\tilde{G}^S(\mathbf{y})) = \mathbb{L}_S(-\ln \tilde{G}(\mathbf{y})) = \exp(-(-\ln \tilde{G}(\mathbf{y}))^\alpha) =: \tilde{G}_\alpha(\mathbf{y}). \quad (3.1)$$

The distribution \tilde{G}_α is a special case of G_α , that is max-stable with extreme-value copula C_{G_α} and common α -Fréchet margins. By (3.1), it follows easily that for any $\alpha_1, \alpha_2 \in (0, 1)$ we have $(\tilde{G}_{\alpha_1})_{\alpha_2} = (\tilde{G}_{\alpha_2})_{\alpha_1} = \tilde{G}_{\alpha_1\alpha_2}$. This means that the max-stable \tilde{G}_α can be obtained by (iterated) random scaling in various ways. In the particular case that the components of \mathbf{Z} are independent, the copula of \tilde{G}_α is

$$C_{\tilde{G}_\alpha}(\mathbf{u}) = \exp(-((- \ln u_1)^{1/\alpha} + \dots + (- \ln u_d)^{1/\alpha})^\alpha), \quad \mathbf{u} \in (0, 1]^d, \quad \alpha \in (0, 1), \quad (3.2)$$

which is the well-know Symmetric Logistic copula (e.g., Joe, 2015, p. 172). Therefore, the elements of \mathbf{Z} are dependent for any $\alpha \in (0, 1)$, and they become nearly independent as $\alpha \rightarrow 1$ and almost completely dependent as $\alpha \rightarrow 0$. Random scaling constructions similar to this one have been already discussed by Fougères et al. (2009) and Fougères et al. (2013).

The de Haan representation of max-stable processes (de Haan, 1984) provides a Poisson point process construction of a random vector with any max-stable distribution \tilde{G} . A question that arises here is: What is the spectral representation of a random vector \mathbf{R} defined by the random scaling construction? The next result establishes that the findings presented in Robert (2013) indeed provide the spectral representation of \mathbf{R} .

Proposition 3.1. *Let $\mathbf{Z}_1, \mathbf{Z}_2, \dots$ be iid copies of \mathbf{Z} , with distribution \tilde{G} , independent of P_1, P_2, \dots that are points of a Poisson process on $(0, \infty)$ with intensity measure $\alpha r^{-(\alpha+1)} dr$, $\alpha \in (0, 1)$. Define*

$$\mathbf{R} = \frac{1}{\Gamma(1-\alpha)} \left(\max_{i \geq 1} P_i Z_{i1}, \dots, \max_{i \geq 1} P_i Z_{id} \right).$$

Then, the distribution of \mathbf{R} is \tilde{G}_α .

Here we provide an alternative, more general proof than that given in Robert (2013). Specifically, ours does not rely on an unnecessary smoothness assumption. Next, we provide a characterization of G_α , as the attractor distribution for a general random scaling and centering construction.

Proposition 3.2. *Let $\mathbf{X}_1, \dots, \mathbf{X}_n$ be iid copies of the random vector \mathbf{X} , with distribution F_X . Assume $F_X \in \mathcal{D}(G)$. Let S be a positive α -stable random variable, $\alpha \in (0, 1)$. Assume S is independent of \mathbf{X} . Define*

$$\mathbf{w}_n := \frac{\mathbf{a}_{\lfloor nS \rfloor}}{\mathbf{a}_n}, \quad \mathbf{v}_n := \mathbf{b}_n - \mathbf{b}_{\lfloor nS \rfloor} \frac{\mathbf{a}_n}{\mathbf{a}_{\lfloor nS \rfloor}},$$

where \mathbf{a}_n and \mathbf{b}_n are the usual norming constants of F_X . Then,

$$\mathbf{a}_n^{-1} (\max(\mathbf{w}_n(\mathbf{X}_1 - \mathbf{v}_n), \dots, \mathbf{w}_n(\mathbf{X}_n - \mathbf{v}_n)) - \mathbf{b}_n) \rightsquigarrow G_\alpha, \quad n \rightarrow \infty.$$

A simple implication of Proposition 3.2 is the following. Transforming \mathbf{X} into \mathbf{Y} , a random vector with common unit-Fréchet marginal distributions, and setting $\mathbf{R} = S\mathbf{Y}$, implies that $F_{\mathbf{R}} \in \mathcal{D}(\tilde{G}_\alpha)$. To see this, note that $F_{\mathbf{Y}} \in \mathcal{D}(\tilde{G})$ with norming sequences $\mathbf{a}_n = \mathbf{1}_n$, $\mathbf{b}_n = \mathbf{0}$, and that, as $n \rightarrow \infty$, $S\mathbf{Y}$ and $\mathbf{w}_n\mathbf{Y} = (\lfloor nS \rfloor/n)\mathbf{Y}$ have approximately the same distribution. As a result, the attractors of $F_{\mathbf{R}}$ and F_{M_N} , when $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \in (0, 1)$, share the same extreme-value copula, C_{G_α} . We finally derive the explicit form of the Pickands function corresponding to the latter.

Proposition 3.3. *The Pickands function corresponding to the extreme-value copula C_{G_α} in (2.4) is*

$$A_\alpha(\mathbf{t}) = \|\mathbf{t}\|_{1/\alpha} A^\alpha \left((\mathbf{t}/\|\mathbf{t}\|_{1/\alpha})^{1/\alpha} \right), \quad \mathbf{t} \in \mathcal{S}_d, \quad \alpha \in (0, 1), \quad (3.3)$$

where A is the Pickands dependence function corresponding to C_G and

$$\|\mathbf{t}\|_{1/\alpha} = \left(\sum_{i=1}^d t_i^{1/\alpha} \right)^\alpha, \quad \mathbf{t} \in \mathcal{S}_d, \alpha \in (0, 1) \quad (3.4)$$

is the Pickands function corresponding to the Logistic copula.

As a direct consequence of Proposition 3.3, the following facts ensue. The smaller the parameter α , the more A_α represents a stronger dependence level than A . Since we have that $\|(1/d, \dots, 1/d)\|_\alpha = d^{\alpha-1}$, then, by the definition of the extremal coefficient in Section 1, we obtain

$$\theta(G_\alpha) = (\theta(G))^\alpha. \quad (3.5)$$

By solving for A in equation (3.3), we obtain the inverse relation between A_α and A , i.e.,

$$A^*(\mathbf{t}) := A\left((\mathbf{t}/\|\mathbf{t}\|_{1/\alpha})^{1/\alpha}\right) = (A_\alpha(\mathbf{t})/\|\mathbf{t}\|_{1/\alpha})^{1/\alpha}, \quad \mathbf{t} \in \mathcal{S}_d \quad (3.6)$$

and $A(\mathbf{t}) = A^*(\mathbf{t}^\alpha/\|\mathbf{t}^\alpha\|_1)$, providing the expression for the Pickands function in (1.1).

4. Inferring the Pickands function

In this section we introduce a new semiparametric procedure to estimate the Pickands function A in (1.1). Several nonparametric estimators are already available for A when a data sample from the limiting distribution for unaggregated data, G , is observable, see e.g., Klüppelberg and May (2006), Zhang et al. (2008), Berghaus et al. (2013), Cormier et al. (2014), Marcon et al. (2017) among others. Unlike the above references, we assume that only replicates of (M_N, N) are observable, from which sample extremes (maxima) are extracted. Then, we construct an estimator for A , exploiting an inversion method via (3.6). This is a substantial novelty in the extreme-value literature. It is common practice to assume that sample maxima are exactly coming from the limiting model \mathcal{Q} in (2.1), with $\alpha \in (0, 1)$, and provide an asymptotic validation of the proposed inferential procedure in such a setting. Via an extensive simulation study in Section 4.2, we show that, in practice, our method provides a good performance with data that are only approximately coming from \mathcal{Q} . Extending the asymptotic statistical theory to the latter case goes beyond the scope of the present already quite technical work. Observe that, since $\mathbf{t} \mapsto (\mathbf{t}/\|\mathbf{t}\|_\alpha)^{1/\alpha}$ is a bijective map, estimating A^* is equivalent to estimating A , thus, for simplicity, we hereafter focus on the former function.

4.1. A semiparametric composite-estimator

Let $(\boldsymbol{\eta}_1, \xi_1), (\boldsymbol{\eta}_2, \xi_2), \dots$, be iid random vectors with joint distribution in (2.1) with $\alpha \in (0, 1)$. Assume that a sample of n observations from such a sequence is available. An estimate of A^* is obtained by combining the results of a two-step procedure: we estimate α and A_α , we plug the estimates in (3.6). Precisely, ξ_1, \dots, ξ_n follows a α -Fréchet distribution. For estimating α we consider two well-known estimators: the Generalized Probability Weighted Moment (GPWM) (Guillou et al., 2014) and the Maximum Likelihood (ML). In the first case the estimator is

$$\widehat{\alpha}_n^{\text{GPWM}} := \left(k - 2 \frac{\widehat{\mu}_{1,k}}{\widehat{\mu}_{1,k-1}} \right)^{-1}, \quad (4.1)$$

for $k \in \mathbb{N}_+$, where

$$\widehat{\mu}_{a,b} = \int_0^1 H_n^\leftarrow(v) v^a (-\ln v)^b dv, \quad a, b \in \mathbb{N}$$

and

$$H_n(y) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(\xi_i \leq y), \quad y > 0. \quad (4.2)$$

In the second case the estimator is

$$\widehat{\alpha}_n^{\text{ML}} := \operatorname{argmax}_{\tilde{\alpha} \in (0, \infty)} \sum_{i=1}^n \ln \dot{\Phi}_{\tilde{\alpha}}(\xi_i), \quad (4.3)$$

where $\dot{\Phi}_{\tilde{\alpha}}(x) = \partial/\partial x \Phi_{\tilde{\alpha}}(x)$, $x > 0$.

The sequence η_1, \dots, η_n follows the distribution G_α . For estimating A_α we consider three well-know estimators: Pickands (P) (Pickands, 1981), Capéraà-Fougère-Genest (CFG) (Capéraà et al., 1997) and Madogram (MD) (Marcon et al., 2017). In the first case the estimator is

$$\begin{aligned} \widehat{A}_{\alpha,n}^{\text{P}} &:= \left(\frac{1}{n} \sum_{i=1}^n \widehat{\vartheta}_i(\mathbf{t}) \right)^{-1}, \\ \widehat{\vartheta}_i(\mathbf{t}) &= \min_{1 \leq j \leq d} \left\{ -\frac{1}{t_j} \ln \left(\frac{n}{n+1} G_{n,j}(\eta_{i,j}) \right) \right\} \end{aligned} \quad (4.4)$$

where for every $x \in \mathbb{R}$ and $j \in \{1, \dots, d\}$

$$G_{n,j}(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}(\eta_{i,j} \leq x). \quad (4.5)$$

In the second case the estimator is

$$\widehat{A}_{\alpha,n}^{\text{CFG}} := \exp \left(-\frac{1}{n} \sum_{i=1}^n \ln \widehat{\vartheta}_i(\mathbf{t}) - \varsigma \right), \quad (4.6)$$

where ς is the Euler's constant. Finally, in the third case the estimator is

$$\widehat{A}_{\alpha,n}^{\text{MD}} := \frac{\widehat{\nu}_n(\mathbf{t}) + c(\mathbf{t})}{1 - \widehat{\nu}_n(\mathbf{t}) - c(\mathbf{t})}, \quad (4.7)$$

$$\widehat{\nu}_n(\mathbf{t}) = \frac{1}{n} \sum_{i=1}^n \left(\max_{j=1, \dots, d} G_{n,j}^{1/t_j}(\eta_{i,j}) - \frac{1}{d} \sum_{j=1}^d G_{n,j}^{1/t_j}(\eta_{i,j}) \right), \quad (4.8)$$

where $u^{1/0} = 0$ for $0 < u < 1$ by convention and $c(\mathbf{t}) = \sum_{j=1}^d t_j / (1 + t_j)$.

For brevity we denote the estimators of α and A_α by $\widehat{\alpha}_n^\bullet$ and $\widehat{A}_{\alpha,n}^\circ$, respectively, where the symbols “ \bullet ” and “ \circ ” are representative of the labels “GPWM”, “ML” and “P”, “CFG”, “MD”, respectively. Then, plugging the estimators into equation (3.6) we obtain the following composite-estimator for A^\star ,

$$\widehat{A}_{\alpha,n}^{\star\bullet}(\mathbf{t}) := \left(\widehat{A}_{\alpha,n}^\circ(\mathbf{t}) / \|\mathbf{t}\|_{1/\widehat{\alpha}_n^\bullet} \right)^{1/\widehat{\alpha}_n^\bullet}, \quad \mathbf{t} \in \mathcal{S}_d. \quad (4.9)$$

Next, we establish the asymptotic theory of the composite-estimator in (4.9) defined by all the combinations of the GPWM and ML estimators for α with the P, CFG and MD estimators for A_α . Our results rely on the following assumptions.

Condition 4.1. For $j \in \{1, \dots, d\}$, let $\mathcal{U}_j = \{\mathbf{u} \in [0, 1]^d : 0 < u_j < 1\}$. Assume that:

- (i) for $j \in \{1, \dots, d\}$, the first-order partial derivative $\dot{C}_{G_\alpha;j}(\mathbf{u}) := \partial/\partial u_j C_{G_\alpha}(\mathbf{u})$ exists and is continuous in \mathcal{U}_j ;
- (ii) for $i, j \in \{1, \dots, d\}$, the second-order partial derivative $\ddot{C}_{G_\alpha;i,j}(\mathbf{u}) := \partial/\partial u_i \dot{C}_{G_\alpha;j}(\mathbf{u})$ exists and is continuous in $\mathcal{U}_i \cap \mathcal{U}_j$ and

$$\sup_{\mathbf{u} \in \mathcal{U}_i \cap \mathcal{U}_j} \max(u_i, u_j) |\ddot{C}_{G_\alpha;i,j}(\mathbf{u})| < \infty.$$

Theorem 4.1. For the estimators $\widehat{A}_{\alpha,n}^{\star\text{MD},\bullet}$, assume that Condition 4.1(i) holds true; assume that Condition 4.1(ii) is also satisfied for the estimators $\widehat{A}_{\alpha,n}^{\star\text{P},\bullet}$, $\widehat{A}_{\alpha,n}^{\star\text{CFG},\bullet}$. Finally, assume that the choice of $k \in \mathbb{N}_+$ in the GPWM-based estimator $\widehat{A}_{\alpha,n}^{\star\text{GPWM},\bullet}$ satisfies $\alpha > 1/(k-1)$. Then, as $n \rightarrow \infty$

$$\sqrt{n} \left\{ \widehat{A}_{\alpha,n}^{\star\bullet}(\mathbf{t}) - A^\star(\mathbf{t}) \right\}_{\mathbf{t} \in \mathcal{S}_d} \rightsquigarrow \{(\phi_{\bullet,\bullet}(\mathbb{C}_Q))(\mathbf{t})\}_{\mathbf{t} \in \mathcal{S}_d} \quad (4.10)$$

in $\ell^\infty(S_d)$, for an operator $\phi_{\circ,\bullet}$ into $\ell^\infty(S_d)$ and a zero-mean Gaussian process \mathbb{C}_Q , whose covariance function is

$$\text{Cov}(\mathbb{C}_Q(\mathbf{u}), \mathbb{C}_Q(\mathbf{v})) = C_Q(\min(\mathbf{u}, \mathbf{v})) - C_Q(\mathbf{u})C_Q(\mathbf{v}), \quad \mathbf{u}, \mathbf{v} \in [0, 1]^{d+1}, \quad (4.11)$$

where C_Q is the copula

$$C_Q(\mathbf{u}, \mathbf{v}) = Q(G_{\alpha,1}^{\leftarrow}(u_1), \dots, G_{\alpha,d}^{\leftarrow}(u_d), \Phi_\alpha^{\leftarrow}(\mathbf{v}))$$

and the minimum is taken componentwise. Moreover,

$$\|\widehat{A_n^{\circ,\bullet}} - A^*\|_\infty \xrightarrow{p} 0, \quad \|\widehat{A_n^{\text{MD,GPWM}}} - A^*\|_\infty \xrightarrow{as} 0, \quad n \rightarrow \infty.$$

Remark 4.1. For brevity, the explicit definitions of $\phi_{\circ,\bullet}$'s are postponed to Definitions A.4(iv)-(v), A.4(vii)–A.4(viii), and functional limit results are provided all at once. Although each estimators' combination has its own peculiarities, these can be framed within a fairly general theory, which might be of interest per se. Due to the high degree of technicality, we present such theory in the appendix, herein focusing on ready-to-use estimators' examples.

Remark 4.2. In Gudendorf and Segers (2012) and Marcon et al. (2017) modified versions of the estimators P, CFG and MD for A_α are proposed to guarantee that $\widehat{A_{\alpha,n}^\circ}(\mathbf{e}_j) = 1$ for all $n = 1, 2, \dots$ and $j = 1, \dots, d$ where $\mathbf{e}_j = (0, \dots, 0, 1, 0, \dots, 0)$. The results in Theorem 4.1 are also valid when such adjusted estimators are considered in place of (4.4), (4.6) and (4.7), respectively, due to asymptotic arguments developed in the aforementioned works.

Remark 4.3. By the identity in (2.4), Proposition 1 in Gudendorf and Segers (2012) guarantees, if the stable-tail dependence function L satisfies Assumption 2 therein, that C_{G_α} satisfies Condition 4.1.

4.2. Simulation study

We show the finite sample performance of the composite-estimator $\widehat{A_n^{\circ,\bullet}}$ in (4.9) through a simulation study. Hereafter we consider for the P, CFG and MD estimators, the adjusted versions mentioned in Remark 4.2.

Since it is not straightforward to simulate from the limit distribution Q , we study the performance of the composite-estimator $\widehat{A_n^{\circ,\bullet}}$ when it is used with data that are only approximately coming from the limiting distribution Q . Nevertheless, this is a more realistic scenario.

Specifically, we set $N = \lceil N' \rceil$, where we assume that N' follows a standard Pareto distribution with shape parameter $\alpha \in (0, 1)$. We simulate N observations of a two-dimensional random vector \mathbf{X} with a standard bivariate Student- t distribution with a fixed value of the correlation ρ and the degrees of freedom ν . We recall that a Student- t distribution is in the domain of attraction of a multivariate extreme-value distribution with an extreme-value copula that is the so-called Extremal- t (e.g., Joe, 2015, p. 189). In the bivariate case, the extremal coefficient of the Extremal- t copula is $\theta = 2T_{\nu+1}[\{(\nu+1)(1-\rho)/(1+\rho)\}^{1/2}]$, where $T_{\nu+1}$ is a univariate standard Student- t distribution with $\nu+1$ degrees of freedom. Next, with the simulated data we compute the observed value of the componentwise maxima \mathbf{M}_N in (1.2). We repeat these simulation steps $n' = 500$ times generating n' independent observations from the pair (N, \mathbf{M}_N) with which we compute an observation from the random variable $\xi = \max(N_1, \dots, N_{n'})$ and vector $\boldsymbol{\eta} = \max(\mathbf{M}_{N_1}, \dots, \mathbf{M}_{N_{n'}})$, where the later maximum is meant componentwise. We repeat these simulation steps n times, generating a data sample approximately drawn from the distribution Q , whose expression is given in the first line of (2.1) and where the expression of G can be deduced from Joe (2015, p. 189).

Then, we estimate α using the observations generated from the sequence ξ_1, \dots, ξ_n with the GPWM estimator $\widehat{\alpha_n^{\text{GPWM}}}$ in equation (4.1), with $k = 5$, and the ML estimator $\widehat{\alpha_n^{\text{ML}}}$ in (4.3). Afterwards, we estimate the Pickands dependence function A_α using the observations generated from the sequence $\boldsymbol{\eta}_1, \dots, \boldsymbol{\eta}_n$ with the P estimator $\widehat{A_{\alpha,n}^{\text{P}}}$ in (4.4), CFG estimator $\widehat{A_{\alpha,n}^{\text{CFG}}}$ in (4.6) and MD estimator $\widehat{A_{\alpha,n}^{\text{MD}}}$ in (4.7). Finally, we estimate A^* using the composite-estimator $\widehat{A_n^{\circ,\bullet}}$ in equation (4.9).

We repeat the simulation and estimation steps for different values of the model parameters α , ρ and ν and different sample sizes. Precisely, we consider $\alpha = 0.5, 0.633, 0.767, 0.9$ and, for the Student- t distribution, we consider the degrees of freedom $\nu = 1$ and 15 equally spaced values of the correlation ρ in $[-0.99, 0.99]$. With these parameters' values, the extremal coefficient θ (related to the Extremal- t copula) takes values in $[1, 2]$, where the lower and upper bounds represent the cases of complete dependence and independence. We also consider the




FIG 1. MISE, ISB and IV for 1000 samples of size 50 from an approximated distribution Q , obtained on the basis of the standard Pareto distribution for N and the bivariate Student- t distribution for X , for different values of the parameters α and ρ , v . The parameter α is estimated with the GPWM estimator in (4.1). The parameter θ is the extremal coefficient related to the corresponding extreme-value copula extremal- t .

sample sizes $n = 50, 100$. We repeat this experiment (the simulation and estimation considering different values of the parameters and the sample sizes) 1000 times and we compute a Monte Carlo approximation of the Mean Integrated Squared Error (MISE), i.e.,

$$\begin{aligned} \text{MISE}(\widehat{A}_n^{\circ\bullet}, A^*) &= E \left(\int_{S_d} [\widehat{A}_n^{\circ\bullet}(t) - A^*(t)]^2 dt \right) \\ &= \int_{S_d} [E(\widehat{A}_n^{\circ\bullet}(t)) - A^*(t)]^2 dt + \int_{S_d} E(\widehat{A}_n^{\circ\bullet}(t) - E(\widehat{A}_n^{\circ\bullet}(t)))^2 dt, \end{aligned}$$

where the first and second terms in the second line are known as Integrated Squared Bias (ISB) and Integrated Variance (IV) (Gentle, 2009, Ch. 6.3).

Figure 1 displays the results obtained with the GPWM-based estimators for the sample size $n = 50$. The MISE, ISB and IV ($\times 1000$) of the GPWM-based estimators are reported from the first to the third row. The solid black, dashed green and dotted red lines report the results obtained estimating A_α with P, CFG and MD estimators, respectively. The results for the different values of α are reported along the columns.

For each fixed value of α we see that IV is close to zero at the strongest dependence level ($\theta = 1$), then it increases with the decrease of the dependence level (θ increases approaching two). On the contrary, ISB takes the largest value at $\theta = 1$ and then it decreases with the decrease of the dependence level, for the cases $\alpha = 0.5, 0.633$. Overall, for the case $\alpha = 0.5$, MISE takes the largest value at $\theta = 1$ and then it decreases with the decrease of the dependence level. For the case $\alpha = 0.633$, MISE does not change much over the whole range of dependence levels, since ISB and IV compensate each other. While, for the cases $\alpha = 0.767, 0.9$, IV grows much more than ISB decreases, implying that MISE increases with the decrease of the dependence level. The smallest values of ISB and IV are obtained with the CFG-based and P-based estimator, respectively. Overall, on the basis of the MISE,




Fig 2. Ratio between MISE, ISB and IV computed estimating the function A^* by the estimator $\widehat{A}_n^{GPWM, \bullet}$ and $\widehat{A}_n^{ML, \bullet}$ in formula (4.9). The same setting as Figure 1 is considered.

the best performance is obtained with the CFG-based estimator, although there is little difference with the P-based estimator. As shown in the supplementary material, there is not much difference in the performance of the P-, CFG- and MD-based estimators already for the sample size $n = 100$.

Although in this experiment we consider synthetic data that only approximately come from the distribution Q , the results summarised by ISB, IV and MISE highlight the robustness of our method to model misspecification (for only approximately max-stable data). In particular, our composite-estimator displays a moderate distortion, despite that the uniform consistency guarantee of Theorem 4.1 does not directly extend to the present setting. Similar conclusions are obtained with sample size $n = 100$ (available in the supplementary material).

A comparison between the estimation results obtained with the GPWM- and ML-based estimators is reported in Figure 2. Precisely, from the first to the third row, the ratio between the MISE, ISB and IV computed estimating the function A^* by the GPWM- and ML-based estimators are displayed. For the case $\alpha = 0.5$, on the basis of the ISB, the GPWM- and ML-based estimators perform very similarly, when $1 \leq \theta \leq 1.5$, i.e. for strong up to moderate dependence levels. Instead, when $1.5 \leq \theta \leq 2$, that is for moderate up to weak dependence levels, the ML-based estimators outperform the GPWM-based estimators. However, for very weak dependence levels (θ close to 2) the GPWM-CFG estimator outperforms the ML-CFG estimator. On the basis of the IV, the ML-based estimators outperform the GPWM-based estimators for all cases. Concerning the configurations with $\alpha > 0.5$, the ML-based estimators considerably outperform GPWM-based estimators in terms of MISE. These conclusions are valid for all three P-, CFG- and MD-based estimators. Specifically, IV is smaller for the ML-based estimators and their better performances are obtained for weaker extremal dependence structures (when θ approaches 2). The ML-based estimators are much less biased than the GPWM-based estimators and the difference is much more pronounced for increasing values of α and weaker extremal dependence structures (when θ approaches 2), although for the P- and CFG-based estimators such a difference diminishes when θ is close to 2.

The study was performed using the R (R Core Team, 2014) packages Copula (Kojadinovic and Yan, 2010) and

evd (Stephenson, 2002).

5. Discussion

Here we shortly discuss directions for future research, from both an applied and a theoretical viewpoint. First, we illustrate a potential real data problem with Internet traffic. In this domain, our theoretical framework can be applied to improve and extend the existing methods for the study of Internet traffic data and to perform an extreme-value statistical analysis. Next, we provide some concluding remarks, including probabilistic and methodological extensions of this work.

Massive Internet traffic data. The analysis of Internet traffic data is crucial for improving the performance of large networks. Inferring internet congestion at different levels continues to receive increasing attention as new challenges are posed, for example, the booming demand for high-bandwidth contents (e.g., video streaming). Large scale collection and analysis of scientific data on Internet traffic is conducted by renowned research institutions, such as the Center for Applied Internet Data Analysis (CAIDA, <http://www.caida.org/home/>). CAIDA collects anonymized traffic traces from several monitors connected to commercial Internet backbones and large Internet service providers. A one-minute trace lists a huge amount of IP (Internet Protocol) packets, resulting in several gigabytes of compressed files. A first reduction of the data size is obtained resorting to flow records, i.e. measurements pertaining to coherent strings of packets, e.g., those stemming from the same traffic source, such as a single user (see van de Meent, 2006, Ch. 3). Though, statistical analysis at this level is still computationally too burdensome. Nonetheless, knowledge of the joint extremal behaviour of flow size (amount of data transmitted as part of a flow), say X_1 , and flow duration (time difference between the first and last packet of a flow), say X_2 , would improve the existing techniques for large network optimization. For instance, it would be of interest to forecast the through output rate X_1/X_2 , corresponding to extremely large flow sizes (see Markovich, 2007, Ch. 1.3.2, 1.3.5).

To infer the extremal dependence between X_1 and X_2 , it is first necessary to reduce the dimension of flow data by appropriately aggregating them. This is obtained by computing maxima of X_1 and X_2 over the random number of flows, N , occurring in a suitable time interval. The survival function can be reasonably expected to display a power-law behaviour, i.e. $\mathbb{P}(N > n) = \mathcal{L}(n)n^{-\alpha}$, with α smaller than one. This is consistent both with earlier theoretical findings, e.g. Bonald et al. (2001), and with the complexity of modern applications and backbone infrastructures. Different to the classical probabilistic description of Internet aggregate traffic (e.g., Taqqu et al., 1997; Willinger et al., 1997), in which the number of active sources (flows) is deterministically sent to infinity, we account for randomness in the number of flows. Still, stochastic modelling via Pareto tails with $\alpha < 1$ (hence $E(N) = +\infty$) appears a coherent refinement of the standard approach.

In this setting, Theorem 2.1 provides the mathematical basis for modeling the extremal dependence of aggregated-flow size and duration by means of the componentwise maxima approach. Furthermore, by the statistical inversion method in Section 4.1 we can infer the extremal dependence between the extreme single-flow size and duration. Due to the extremely complex nature of flow data, we defer the actual data analysis to a future specialized work.

Concluding remarks. The probabilistic and statistical modeling of aggregated data is an interesting and important topic. The total and maximum amounts derived on a random number of observations described by the random vectors S_N and M_N in (1.2) are two simple examples of aggregated data. There are not many results available on the extremal behaviour of S_N and M_N , apart from those on S_N in the univariate case. This contribution makes a first step by establishing the multivariate extreme-value theory for M_N . Investigating the joint upper-tail behaviour of M_N , under the hidden regular variation framework (e.g., Resnick, 2002; Maulik and Resnick, 2004; Mitra and Resnick, 2011), would represent a first extension of this work. Of particular interest would be investigating whether $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha \geq 1$ implies a stronger residual dependence than the case $F_N \in \mathcal{D}(\Lambda)$, as no difference emerges in terms of classical extremal dependence (see Theorem 2.1). In this paper we consider for simplicity the same random number N of independent copies for each components of $X = (X_1, \dots, X_d)$. Of greater generality would be to consider $N = (N_1, \dots, N_d)$, i.e. a different random number of independent copies for each component of X . In this context, it can be assumed that N belongs to the domain of attraction of a multivariate extreme value distribution. This provides a second extension of our work. A third extension could be the derivation of nonparametric estimators and their asymptotic results using threshold exceedances (for at least one component).

Finally, the extension of our results (probabilistic and inferential) to the case of \mathbf{S}_N represents a relevant open problem.

Appendix A: Proofs

A.1. Proof of Theorem 2.1

We start with some notation. Let $U_{X_j}(t) := F_{X_j}^{\leftarrow}(1 - 1/t)$, $t > 1$, $D_j(x) = G_j^{\leftarrow}(e^{-1/x})$, $x > 0$, $j = 1, \dots, d$, and

$$F_*(\cdot) = F_X(U_{X_1}(\cdot), \dots, U_{X_d}(\cdot)), \quad \tilde{G}(\mathbf{y}) = G(D_1(y_1), \dots, D_d(y_d)), \quad \mathbf{y} \in (0, \infty)^d.$$

In extreme-value theory, it is common practice to derive the attractor G of a distribution F_X by analyzing separately the behaviour of its margins and its dependence structure. The latter is typically investigated by focusing on F_* , which obtains from F_X by transforming the margins into unit-Pareto. In this way, we have that $F_* \in \mathcal{D}(\tilde{G})$, where \tilde{G} has common unit-Fréchet and the same extreme-value copula of G . Although different types of common marginal distributions can be considered (e.g., Falk et al., 2011, Ch. 4), with Pareto margins we can exploit the theory of regularly varying tails. Precisely, since $F_X \in \mathcal{D}(G)$, we have that (Resnick, 2007, Propositions 5.10, 5.15 and 5.17)

$$\lim_{n \rightarrow \infty} \frac{1 - F_*(n\mathbf{y})}{1 - F_*(n\mathbf{1})} = \frac{-\ln \tilde{G}(\mathbf{y})}{\theta(G)}, \quad \mathbf{y} \in (0, \infty)^d, \quad (\text{A.1})$$

where $-\ln \tilde{G}(\mathbf{y})$ is homogeneous of order -1 and $\theta(G) = \theta(\tilde{G}) = -\ln \tilde{G}(\mathbf{1})$. We also define $T_*(y) := F_*(y\mathbf{1})$, $y > 0$, a univariate non-decreasing function. Finally, we set $U_*(t) := T_*^{\leftarrow}(1 - 1/t)$ and $U_N(t) := F_N^{\leftarrow}(1 - 1/t)$, $t > 1$.

The proof is organized in three parts: the derivation of the norming constants, two preliminary results and the conclusion. For the sake of brevity, some of the technical derivations are deferred to the supplementary material.

A.1.1. Norming constants

Let the norming sequences κ_n, ϱ_n for F_N be defined in the standard way (e.g., Resnick, 2007, pp. 48-54). We recall that $F_X^n(\mathbf{a}_n\mathbf{x} + \mathbf{b}_n) \rightarrow G(\mathbf{x})$ as $n \rightarrow \infty$ if and only if $n(1 - F_X(\mathbf{a}_n\mathbf{x} + \mathbf{b}_n)) \rightarrow -\ln G(\mathbf{x})$ as $n \rightarrow \infty$, see e.g. Falk et al. (2011, Ch.4) for details. Analogously, here we focus on

$$n(1 - F_{M_N}(\mathbf{c}_n\mathbf{x} + \mathbf{d}_n)) = n(1 - L_N(-\ln p_n(\mathbf{x}))), \quad (\text{A.2})$$

where $p_n(\mathbf{x}) = F_X(\mathbf{c}_n\mathbf{x} + \mathbf{d}_n)$. Accordingly, the derivation of the norming constants \mathbf{c}_n and \mathbf{d}_n requires an analysis of the behaviour of $1 - L_N(s)$, as $s \downarrow 0$. Observe that $L_N(1/x)$, $x > 0$, is monotone nondecreasing. Set $U_{L_N}(x) := (L_N(1/\cdot))^{\leftarrow}(1 - 1/x)$, $x > 0$. Define $z_n := U_{L_N}(n/b)$, where $b := (\theta(G))^\alpha$ when $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \in (0, 1)$, and $b := (\theta(G))$ otherwise. Then, setting $m_n := U_*(z_n)$, as $n \rightarrow \infty$ we have $m_n \rightarrow \infty$ and $z_n \sim 1/\bar{T}_*(m_n)$, where $\bar{T}_*(y) = 1 - T_*(y)$, $y > 0$. Consequently, we also have

$$1 - L_N(\bar{T}_*(m_n)) \sim \frac{b}{n}, \quad n \rightarrow \infty. \quad (\text{A.3})$$

Next, we explicitly provide some asymptotic approximations which help to understand the results derived in the following subsections. When $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \in (0, \infty)$, then $\bar{F}_N(y) = 1 - F_N(y)$, $y > 0$, satisfies

$$\bar{F}_N(y) \sim \mathcal{L}(y)y^{-\alpha}, \quad y \rightarrow \infty, \quad (\text{A.4})$$

where \mathcal{L} is slowly varying. In particular, for any $\alpha \in (0, 1]$ we have that

$$1 - L_N(s) \sim \mathcal{L}^*(1/s)s^\alpha, \quad s \downarrow 0, \quad (\text{A.5})$$

for some slowly varying function \mathcal{L}^* , which is equal to $\Gamma(1 - \alpha)\mathcal{L}$ if $\alpha \in (0, 1)$, or satisfies $\lim_{x \rightarrow \infty} \mathcal{L}(x)/\mathcal{L}^*(x) = 0$, if $\alpha = 1$. See Section 3.1 of the supplementary material for details. When $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha > 1$, or $F_N \in \mathcal{D}(\Lambda)$, then $E(N) \in (0, \infty)$ and

$$1 - L_N(s) \sim E(N)s, \quad s \downarrow 0. \quad (\text{A.6})$$

Therefore, as $n \rightarrow \infty$ we have $z_n \sim E(N)n/b$ and

$$\bar{T}_*(m_n) \sim 1/z_n \sim \theta(G)/(nE(N)), \quad n \rightarrow \infty. \quad (\text{A.7})$$

We finally derive \mathbf{c}_n and \mathbf{d}_n . When $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha > 0$,

(i) if $F_{X_j} \in \mathcal{D}(\Phi_{\beta_j})$, then we set $d_{n,j} = 0$ and

$$c_{n,j} = \begin{cases} U_{X_j}(m_n), & \alpha \in (0, 1] \\ U_{X_j}(n)\{E(N)\}^{1/\beta_j}, & \alpha > 1, \end{cases}$$

(ii) if $F_{X_j} \in \mathcal{D}(\Lambda)$, then we set

$$\begin{aligned} c_{n,j} &= \begin{cases} \omega_j(d_{n,j}), & \alpha \in (0, 1] \\ \omega_j\{\Upsilon_j^{\leftarrow}(1 - 1/\delta_j n)\}, & \alpha > 1 \end{cases} \\ d_{n,j} &= \begin{cases} \Upsilon_j^{\leftarrow}(1 - 1/\delta_j m_n), & \alpha \in (0, 1] \\ c_{n,j} \ln E(N) + \Upsilon_j^{\leftarrow}(1 - 1/\delta_j n), & \alpha > 1, \end{cases} \end{aligned}$$

where Υ_j is the Von Mises function associated to \bar{F}_{X_j} , with $\bar{F}_{X_j}(x) = 1 - F_{X_j}(x)$ for $x \in \mathbb{R}$, ω_j its auxiliary function (e.g., [Resnick, 2007](#), pp. 40-43) and $\delta_j = \lim_{x \rightarrow \infty} \bar{F}_{X_j}(x)/\{1 - \Upsilon_j(x)\}$;

(iii) $F_{X_j} \in \mathcal{D}(\Psi_{\beta_j})$, then we set $d_{n,j} = x_{0,j}$, where $x_{0,j} = \sup\{x : F_{X_j}(x) < 1\}$, and

$$c_{n,j} = \begin{cases} \{U_{\tilde{X}_j}(m_n)\}^{-1}, & \alpha \in (0, 1] \\ \{U_{\tilde{X}_j}(n)E(N)^{1/\beta_j}\}^{-1}, & \alpha > 1, \end{cases}$$

where $\tilde{X}_j := 1/(x_{0,j} - X_j)$ and $U_{\tilde{X}_j} = F_{\tilde{X}_j}^{\leftarrow}(1 - 1/t)$, $t > 0$.

When $F_N \in \mathcal{D}(\Lambda)$, \mathbf{c}_n and \mathbf{d}_n are set equal to the sequences derived for the case $F_N \in \mathcal{D}(\Phi_\alpha)$, with $\alpha > 1$. With these norming constants, we obtain the following approximations as $n \rightarrow \infty$

$$U_{X_j}^{\leftarrow}(c_{n,j}x_j + d_{n,j}) \sim m_n D_j^{\leftarrow}(x_j) = \begin{cases} m_n x_j^{\beta_j}, & F_{X_j} \in \mathcal{D}(\Phi_{\beta_j}) \\ m_n e^{x_j}, & F_{X_j} \in \mathcal{D}(\Lambda) \\ m_n (-x_j)^{-\beta_j}, & F_{X_j} \in \mathcal{D}(\Psi_{\beta_j}) \end{cases} \quad (\text{A.8})$$

and

$$\begin{aligned} 1 - p_n(\mathbf{x}) &\sim 1 - F_*(m_n D_1^{\leftarrow}(x_1), \dots, m_n D_d^{\leftarrow}(x_d)) \\ &\sim \bar{T}_*(m_n)(-\ln G(\mathbf{x}))/\theta(G) \\ &\rightarrow 0. \end{aligned} \quad (\text{A.9})$$

A.1.2. Preliminary results

Given the identity in (A.2), our first preliminary result provides the attractor of F_{M_N} .

Lemma A.1. *If $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \in (0, 1]$, then we have*

$$\lim_{n \rightarrow \infty} n(1 - L_N(-\ln p_n(\mathbf{x}))) = (-\ln G(\mathbf{x}))^\alpha, \quad \mathbf{x} \in \mathbb{R}^d. \quad (\text{A.10})$$

If $F_N \in \mathcal{D}(\Lambda)$ or $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha > 1$, then (A.10) holds with $\alpha = 1$.

Proof. If $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \in (0, 1]$, using sequentially (A.9), (A.5) together with the properties of slowly varying functions, and (A.3), as $n \rightarrow \infty$ we obtain

$$\begin{aligned} n(1 - L_N(-\ln p_n(\mathbf{x}))) &\sim n(1 - L_N(1 - p_n(\mathbf{x}))) \\ &\sim n(1 - L_N(\bar{T}_*(m_n)(-\ln G(\mathbf{x}))/\theta(G))) \\ &\sim n(1 - L_N(\bar{T}_*(m_n)))(-\ln G(\mathbf{x}))^\alpha/b \\ &\sim (-\ln G(\mathbf{x}))^\alpha. \end{aligned} \quad (\text{A.11})$$

If $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha > 1$, or $F_N \in \mathcal{D}(\Lambda)$, we have $E(N) < \infty$, thus from (A.9) and (A.7) it follows that, in this case, as $n \rightarrow \infty$

$$1 - p_n(\mathbf{x}) \sim -\ln G(\mathbf{x})/\{nE(N)\}. \quad (\text{A.12})$$

Consequently, in view of (A.11) and the approximation in (A.6), as $n \rightarrow \infty$ we have

$$\begin{aligned} n(1 - L_N(-\ln p_n(\mathbf{x}))) &\sim nE(N)(1 - p_n(\mathbf{x})) \\ &\sim -\ln G(\mathbf{x}), \end{aligned} \quad (\text{A.13})$$

which completes the proof. \square

Next, we state an auxiliary result (see Section 3.2 of the supplementary material for the proof), that we use to establish our second preliminary result, characterizing the tail dependence between M_N and N .

Lemma A.2. *If $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha \in (0, 1)$, then we have (set $u_n(y) := \kappa_n y + \varrho_n$, $y > 0$)*

$$\lim_{n \rightarrow \infty} u_n(y) \bar{T}_*(m_n) = \lim_{n \rightarrow \infty} \frac{U_N(n/(-\ln H(y)))}{U_{L_N}(b/n)} = y \theta(G) \Gamma^{-1/\alpha} (1 - \alpha), \quad (\text{A.14})$$

$$\lim_{n \rightarrow \infty} \frac{\bar{F}_N(-c/\ln p_n(\mathbf{x}))}{\bar{F}_N(u_n(y))} = \lim_{n \rightarrow \infty} \frac{\bar{F}_N(cg(\mathbf{x})U_{L_N}(n/b))}{\bar{F}_N(u_n(y))} = y^\alpha c^{-\alpha} \sigma^\alpha(\mathbf{x}, \alpha), \quad \forall c > 0 \quad (\text{A.15})$$

with $\sigma(\mathbf{x}, \alpha)$ as in (2.1) and $g(\mathbf{x}) = \theta(G)/(-\ln G(\mathbf{x}))$. If $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha \geq 1$ or $F_N \in \mathcal{D}(\Lambda)$, the limits in (A.14)-(A.15) (with $y > 0$ or $y \in \mathbb{R}$, respectively) are equal to zero.

Lemma A.3. *If $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha \in (0, 1)$, then we have*

$$\lim_{n \rightarrow \infty} \mathbb{P}(M_N \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n | N > u_n(y)) = \pi(\mathbf{x}, y) - y^\alpha [(-\ln G(\mathbf{x}))^\alpha - \sigma^\alpha(\mathbf{x}, \alpha) \gamma(1 - \alpha, y\sigma(\mathbf{x}, \alpha))] < 1,$$

where $\pi(\mathbf{x}, y) := e^{-y\sigma(\mathbf{x}, \alpha)}$. If $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha \geq 1$ or $F_N \in \mathcal{D}(\Lambda)$, then the above limit is equal to 1.

Proof. Few algebraic steps yield

$$\mathbb{P}(M_N \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n | N > u_n(y)) = p_n^{u_n(y)}(\mathbf{x}) - \frac{1}{\bar{F}_N(u_n(y))} \int_0^{p_n^{u_n(y)}(\mathbf{x})} \bar{F}_N\left(\frac{\ln v}{\ln p_n(\mathbf{x})}\right) dv, \quad (\text{A.16})$$

see Section 3.3 of the supplementary material for details. Using (A.9) and Lemma A.2, as $n \rightarrow \infty$ we obtain

$$\begin{aligned} p_n^{u_n(y)}(\mathbf{x}) &\sim \exp\{-u_n(y)(1 - p_n(\mathbf{x}))\} \\ &\sim \exp\{-u_n(y) \bar{T}_*(m_n)(-\ln G(\mathbf{x}))/\theta(G)\} \\ &\sim \begin{cases} \pi(\mathbf{x}, y), & F_N \in \mathcal{D}(\Phi_\alpha), \alpha \in (0, 1) \\ 1, & F_N \in \mathcal{D}(\Phi_\alpha), \alpha \geq 1 \text{ or } F_N \in \mathcal{D}(\Lambda). \end{cases} \end{aligned}$$

Hence, if $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha \in (0, 1)$, by uniform convergence (Resnick, 2007, Proposition 0.5) and Lemma A.2, as $n \rightarrow \infty$ we also obtain

$$\begin{aligned} -\frac{1}{\bar{F}_N(u_n(y))} \int_0^{p_n^{u_n(y)}(\mathbf{x})} \bar{F}_N\left(\frac{\ln v}{\ln p_n(\mathbf{x})}\right) dv &\sim -\frac{\bar{F}_N(-1/\ln p_n(\mathbf{x}))}{\bar{F}_N(u_n(y))} \int_0^{\pi(\mathbf{x}, y)} (-\ln v)^{-\alpha} dv \\ &\sim -y^\alpha \sigma^\alpha(\mathbf{x}, \alpha) [\Gamma(1 - \alpha) - \gamma(1 - \alpha, y\sigma(\mathbf{x}, \alpha))] \\ &= -y^\alpha [(-\ln G(\mathbf{x}))^\alpha - \sigma^\alpha(\mathbf{x}, \alpha) \gamma(1 - \alpha, y\sigma(\mathbf{x}, \alpha))]. \end{aligned}$$

While, if $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha \geq 1$ or $F_N \in \mathcal{D}(\Lambda)$, for any arbitrarily small $\epsilon > 0$ and large enough n , we have $p_n^{u_n(y)}(\mathbf{x}) \in [1 - \epsilon, 1]$ and $-\ln v / (-\ln p_n(\mathbf{x})) > u_n(y)$, for all $v \in (0, 1 - \epsilon)$; therefore, an application of Lemma A.2 yields that, as $n \rightarrow \infty$,

$$\bar{F}_N \left(\frac{\ln v}{\ln p_n(\mathbf{x})} \middle| N > u_n(y) \right) = \frac{\bar{F}_N \left(\frac{-\ln v}{-\ln p_n(\mathbf{x})} \right)}{\bar{F}_N(u_n(y))} \rightarrow 0, \quad \forall v \in (0, 1 - \epsilon),$$

where $\bar{F}_N(t|N > u_n(y)) := \mathbb{P}(N > t|N > u_n(y))$, $t > u_n(y)$. Hence, by the dominated convergence theorem, as $n \rightarrow \infty$

$$\begin{aligned} \frac{1}{\bar{F}_N(u_n(y))} \int_0^{p_n^{u_n(y)}(\mathbf{x})} \bar{F}_N \left(\frac{\ln v}{\ln p_n(\mathbf{x})} \right) dv &= \int_0^{p_n^{u_n(y)}(\mathbf{x})} \bar{F}_N \left(\frac{\ln v}{\ln p_n(\mathbf{x})} \middle| N > u_n(y) \right) dt \\ &\leq \int_0^{1-\epsilon} \bar{F}_N \left(\frac{\ln v}{\ln p_n(\mathbf{x})} \middle| N > u_n(y) \right) dt + \epsilon \\ &\rightarrow \epsilon, \end{aligned}$$

Since ϵ is arbitrarily small, the term on the left-hand side must converge to zero. The proof is now complete. \square

A.1.3. Conclusion

As $n \rightarrow \infty$, we have

$$\begin{aligned} -n \ln \mathbb{P}(\mathbf{M}_N \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n, N \leq \kappa_n y + \varrho_n) &\sim n[1 - \mathbb{P}(\mathbf{M}_N \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n, N \leq \kappa_n y + \varrho_n)] \\ &= n[1 - F_{\mathbf{M}_N}(\mathbf{c}_n \mathbf{x} + \mathbf{d}_n)] + n\bar{F}_N(u_n(y))\mathbb{P}(\mathbf{M}_N \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n|N > u_n(y)). \end{aligned}$$

In view of (A.2), the limit of the first term on the right hand-side of the second line above is established in Lemma A.1. The second term is asymptotically equivalent to $-\log H(y)\mathbb{P}(\mathbf{M}_N \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n|N > u_n(y))$, where the limit of the conditional probability is established in Lemma A.3. Combining these results we obtain the limiting expressions in (2.1) and (2.2) and the proof is now complete.

A.2. Proof of Corollary 2.2

Let $\mathbf{c}_n, \mathbf{d}_n, \kappa_n, \varrho_n$ be the norming sequences defined in Section A.1 for the case $F_N \in \mathcal{D}(\Phi_\alpha)$, $\alpha \in (0, 1)$. In particular $\varrho_n = 0$, $n\bar{F}_N(\kappa_n) \sim 1$ and $\tilde{\kappa}_n \sim U_N(n\Gamma(1 - \alpha))$ as $n \rightarrow \infty$. The first result, i.e. $\tilde{\kappa}_n^{-1}N_n^+ \rightsquigarrow S$ as $n \rightarrow \infty$, now follows from Theorem 5.4.2 in Uchaikin and Zolotarev (2011).

We recall that $p_n(\mathbf{x}) = F_X(\mathbf{c}_n \mathbf{x} + \mathbf{d}_n)$. In the proof of Lemma A.3 it has been established that

$$p_n^{\kappa_n}(\mathbf{x}) \sim \exp \left(-\frac{-\ln G(\mathbf{x})}{\Gamma^{1/\alpha}(1 - \alpha)} \right), \quad n \rightarrow \infty.$$

Consequently, the second result now follows by noting that by the dominated convergence theorem

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{P}(\mathbf{M}_{[\tilde{\kappa}_n S]} \leq \mathbf{c}_n \mathbf{x} + \mathbf{d}_n) &= \lim_{n \rightarrow \infty} \int_0^\infty p_n^{[\tilde{\kappa}_n S]}(\mathbf{x}) dF_S(s) \\ &= \lim_{n \rightarrow \infty} \int_0^\infty (p_n^{\kappa_n}(\mathbf{x}))^{[\tilde{\kappa}_n S]/\kappa_n} dF_S(s) \\ &= \int_0^\infty \left(e^{-\frac{(-\ln G(\mathbf{x}))}{\Gamma^{1/\alpha}(1 - \alpha)}} \right)^{s\Gamma^{1/\alpha}(1 - \alpha)} dF_S(s) \\ &= \mathbb{L}_S(-\ln G(\mathbf{x})) = G_\alpha(\mathbf{x}). \end{aligned}$$

A.3. Proof of Corollary 2.3

Let Q_j , $j = 1, \dots, d+1$, be the one-dimensional marginal distributions of the max-stable distribution Q . We focus on the case $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha \in (0, 1)$ – the other cases are trivial. From the first line of (2.3) we have $Q_j(x_j) = \exp\{-[-\ln G_j(x_j)]^\alpha\}$, $j = 1, \dots, d$, from which it follows that

$$Q_j^\leftarrow(u_j) = G_j^\leftarrow(\exp\{-[-\ln u_j]^{1/\alpha}\}), \quad u_j \in (0, 1).$$

In particular, $Q_j^\leftarrow(e^{-1}) = G_j^\leftarrow(e^{-1})$, $j = 1, \dots, d$. By assumption, $Q_{d+1}(y) = \Phi_\alpha(y)$ and therefore $Q_{d+1}^\leftarrow(u_{d+1}) = \Phi_\alpha^\leftarrow(u_{d+1})$, with $\Phi_\alpha^\leftarrow(u) = (-\ln u)^{1/\alpha}$. Hence $Q_{d+1}^\leftarrow(e^{-1}) = 1$.

The extremal-coefficient is defined by

$$\theta(Q) = -\log Q(Q_1^\leftarrow(e^{-1}), \dots, Q_d^\leftarrow(e^{-1}), Q_{d+1}^\leftarrow(e^{-1})). \quad (\text{A.17})$$

Plugging in the expressions of $Q_j^\leftarrow(e^{-1})$, $j = 1, \dots, d+1$, into the right-hand side of (A.17), the expression of $\theta(Q)$ is then obtained.

As for the first result in (2.5), it immediately follows from the inequalities $\theta(G) \geq 1$ and $1 \leq 1 - \mathcal{E}_{\Gamma^{1/\alpha}(1-\alpha)}(\theta(G)) + \mathcal{G}_{1-\alpha, \Gamma^{1/\alpha}(1-\alpha)}(\theta(G))$, for $\alpha \in (0, 1)$. To obtain the second one, it is sufficient to note that, as $\alpha \rightarrow 1^-$, $1 - \mathcal{E}_{\Gamma^{1/\alpha}(1-\alpha)}(\theta(G)) \rightarrow 1$ and

$$\begin{aligned} (\theta(G))^\alpha \mathcal{G}_{1-\alpha, \Gamma^{1/\alpha}(1-\alpha)}(\theta(G)) &= \frac{(\theta(G))^\alpha}{\Gamma(1-\alpha)} \gamma\left(1-\alpha, \frac{\theta(G)}{\Gamma^{1/\alpha}(1-\alpha)}\right) \\ &= \frac{(\theta(G))^\alpha}{\Gamma(1-\alpha)} \left[\frac{\theta(G)}{\Gamma^{1/\alpha}(1-\alpha)} \right]^{1-\alpha} \sum_{k=0}^{\infty} \frac{\left[-\frac{\theta(G)}{\Gamma^{1/\alpha}(1-\alpha)} \right]^k}{k!(1-\alpha+k)} \\ &= \frac{\theta(G)}{\Gamma^{\frac{1}{\alpha}-1}(1-\alpha)} \left\{ \frac{1}{\Gamma(1-\alpha)} \frac{1}{1-\alpha} + \frac{1}{\Gamma(1-\alpha)} \sum_{k=1}^{\infty} \frac{\left[-\frac{\theta(G)}{\Gamma^{1/\alpha}(1-\alpha)} \right]^k}{k!(1-\alpha+k)} \right\} \\ &= \frac{\theta(G)}{1+o(1)} \left\{ \frac{1}{1+o(1)} + o(1) \right\} \end{aligned}$$

In the above display, we use the following convergence results: as $\alpha \rightarrow 1^-$, $\Gamma(2-\alpha) \rightarrow \Gamma(1) = 1$, $\Gamma(1-\alpha) \rightarrow \infty$,

$$\begin{aligned} \Gamma^{1/\alpha}(1-\alpha) &= \exp\left\{ \frac{1}{\alpha} \ln \Gamma(1-\alpha) \right\} \rightarrow \infty, \\ \Gamma^{\frac{1}{\alpha}-1}(1-\alpha) &= \exp\left\{ \left(\frac{1}{\alpha} - 1 \right) \ln \Gamma(2-\alpha) - \frac{1}{\alpha} (1-\alpha) \ln(1-\alpha) \right\} \rightarrow 1. \end{aligned}$$

A.4. Proof of Proposition 2.4

Let $U_X(\mathbf{t}) = (U_{X_1}(t_1), \dots, U_{X_d}(t_d))$, with $\mathbf{t} = (t_1, \dots, t_d) \geq 1$ and U_{X_j} , $j = 1, \dots, d$, as in Section A.1, and set $F_*(\mathbf{t}) = F_X(U_X(\mathbf{t}))$. Consider the case where $F_N \in \mathcal{D}(\Phi_\alpha)$ with $0 < \alpha \leq 1$. By exploiting (A.4)-(A.5) and arguments similar to those in the proof of Lemma A.1, we obtain that for $y > 0$ and as $n \rightarrow \infty$

$$\begin{aligned} 1 - \mathbb{P}(\mathbf{M}_N \leq U_X(n\mathbf{y})) &= 1 - \mathbb{L}_N(-\ln F_*(n\mathbf{y})) \\ &\sim \{1 - F_*(n\mathbf{y})\}^\alpha \mathcal{L}^*(\{1 - F_*(n\mathbf{y})\}^{-1}) \\ &\sim \{-\ln \tilde{G}(\mathbf{y})/n\}^\alpha \mathcal{L}^*(n\{-\ln \tilde{G}(\mathbf{y})\}^{-1}) \\ &\sim \begin{cases} \Gamma(1-\alpha) \mathbb{P}(N > -n/\ln \tilde{G}(\mathbf{y})), & \alpha \in (0, 1) \\ (-\ln \tilde{G}(\mathbf{y}))(1 - \mathbb{L}_N(1/n)), & \alpha = 1 \end{cases}. \end{aligned}$$

Furthermore, $U_{\mathbf{M}_N}(n\mathbf{y}) \sim U_X(U_{\mathbf{L}_N}(n\mathbf{y})^{1/\alpha})$ as $n \rightarrow \infty$, with $U_{\mathbf{L}_N}$ as in Section A.1.1 and, for $\mathbf{t} = (t_1, \dots, t_d) \geq 1$,

$$U_{\mathbf{M}_N}(\mathbf{t}) = (U_{M_N^{(1)}}(t_1), \dots, U_{M_N^{(d)}}(t_d)), \quad U_{M_N^{(j)}}(t_j) = F_{M_N^{(j)}}^\leftarrow(1 - 1/t_j), \quad j = 1, \dots, d.$$

As a consequence $1 - \mathbb{P}(M_N \leq U_{M_N}(ny)) \sim n^{-1} \{-\ln \tilde{G}(y^{1/\alpha})\}^\alpha$ as $n \rightarrow \infty$.

Consider the case where $F_N \in \mathcal{D}(\Phi_\alpha)$ with $\alpha > 1$ or $F_N \in \mathcal{D}(\Lambda)$. Again, with steps similar to those in Lemma A.1 we have that for $y > \mathbf{0}$ and as $n \rightarrow \infty$

$$\begin{aligned} 1 - \mathbb{P}(M_N \leq U(ny)) &\sim \mathbb{E}(N) \{1 - F_*(ny)\} \\ &\sim \mathbb{E}(N) (1 - F_*(n\mathbf{1})) \frac{-\ln \tilde{G}(y)}{\theta(G)} \\ &\sim n^{-1} \mathbb{E}(N) \{-\ln \tilde{G}(y)\}. \end{aligned}$$

In addition, $U_{M_N}(ny) \sim U_X(\mathbb{E}(N)ny)$ as $n \rightarrow \infty$. Therefore, $1 - \mathbb{P}(M_N \leq U_{M_N}(ny)) \sim n^{-1} \{-\ln \tilde{G}(y)\}$ as $n \rightarrow \infty$.

A.5. Proof of Proposition 3.1

Observe that $\mathbb{E}(Z_1^\alpha) = \Gamma(1 - \alpha)$ for every $0 < \alpha < 1$. Setting $k_\alpha = 1/\Gamma(1 - \alpha) > 0$ we have

$$\begin{aligned} \frac{1}{k_\alpha} &= \mathbb{E}(Z_1^\alpha) = \int_0^\infty \mathbb{P}\{Z_1 > t^{1/\alpha}\} dt \\ &= \int_0^\infty (1 - e^{-t^{1/\alpha}}) dt = \int_0^\infty (1 - e^{-v^{1/\alpha}}) v^{-2} dv. \end{aligned} \quad (\text{A.18})$$

For any positive (y_1, \dots, y_d) we have

$$\begin{aligned} -\ln \mathbb{P}(R_1 \leq y_1, \dots, R_d \leq y_d) &= k_\alpha \mathbb{E} \left(\max_{1 \leq j \leq d} Z_j^\alpha / y_j^\alpha \right) \\ &= k_\alpha \int_0^\infty \mathbb{P}\{\exists j \in \{1, \dots, d\} : Z_j^\alpha > v y_j^\alpha\} dv \\ &= \int_0^\infty \mathbb{P}\{\exists j \in \{1, \dots, d\} : v k_\alpha Z_j^\alpha > y_j^\alpha\} v^{-2} dv \\ &= \int_0^\infty \mathbb{P}\{\exists j \in \{1, \dots, d\} : Z_j > y_j (k_\alpha v)^{-1/\alpha}\} v^{-2} dv \\ &= \int_0^\infty \left[1 - \mathbb{P}\{\forall j \in \{1, \dots, d\} : Z_j \leq y_j (k_\alpha v)^{-1/\alpha}\} \right] v^{-2} dv. \end{aligned}$$

Now, we have

$$\mathbb{P}\{\forall j \in \{1, \dots, d\} : Z_j \leq y_j (k_\alpha v)^{-1/\alpha}\} = \exp\left(-L(1/y_1, \dots, 1/y_d) k_\alpha^{1/\alpha} v^{1/\alpha}\right),$$

where $L(z)$, with $z = 1/y$, is the stable-tail dependence function. Consequently, by (A.18) we obtain the final result

$$\begin{aligned} -\ln \mathbb{P}(R_1 \leq y_1, \dots, R_d \leq y_d) &= \int_0^\infty \left[1 - e^{-L(1/y_1, \dots, 1/y_d) k_\alpha^{1/\alpha} v^{1/\alpha}} \right] v^{-2} dv \\ &= \left(L(1/y_1, \dots, 1/y_d) \right)^\alpha k_\alpha \int_0^\infty \left[1 - e^{-v^{1/\alpha}} \right] v^{-2} dv \\ &= L^\alpha(1/y_1, \dots, 1/y_d) \end{aligned}$$

establishing the proof.

A.6. Proof of Proposition 3.2

Let $M_n := \max\{X_1, \dots, X_n\}$. By the max-stability of G , there exist maps $\mathcal{A} : (0, \infty) \mapsto (0, \infty)^d$ and $\mathcal{B} : (0, \infty) \mapsto \mathbb{R}^d$, such that $G^s(\mathbf{x}) = G(\mathcal{A}(s)\mathbf{x} + \mathcal{B}(s))$, $s > 0$. By equation (5.18) in Resnick (2007), for every $s > 0$ we have

$$\frac{a_{[ns]}}{a_n} \left(\frac{M_n - b_n}{a_n} - \frac{b_n - b_{[ns]}}{a_{[ns]}} \right) \doteq \frac{1}{\mathcal{A}(s)} \left(\frac{M_n - b_n}{a_n} - \mathcal{B}(s) \right) \rightsquigarrow G(\mathcal{A}(s) \cdot + \mathcal{B}(s)) = G^s, \quad n \rightarrow \infty.$$

where \doteq denotes asymptotic equivalence in distribution. Consequently, the final result follows from the equality $G_\alpha(\mathbf{x}) = E(G^S(\mathbf{x}))$, $\mathbf{x} \in \mathbb{R}^d$, and the dominated convergence theorem

$$\begin{aligned} G_\alpha(\mathbf{x}) &= \int_0^\infty G^S(\mathbf{x}) dF_S(s) = \lim_{n \rightarrow \infty} \int_0^\infty \mathbb{P}\left(\frac{a_{[ns]}}{a_n} \left(\frac{M_n - b_n}{a_n} - \frac{b_n - b_{[ns]}}{a_{[ns]}}\right) \leq \mathbf{x}\right) dF_S(s) \\ &= \lim_{n \rightarrow \infty} \mathbb{P}\left(a_n^{-1} [\max\{\mathbf{w}_n(X_1 - v_n), \dots, \mathbf{w}_n(X_n - v_n)\} - b_n] \leq \mathbf{x}\right). \end{aligned}$$

A.7. Proof of Proposition 3.3

We recall that for all $\mathbf{x} \in \mathbb{R}^d$ we have $G(\mathbf{x}) = C_G(G_1(x_1), \dots, G_d(x_d))$, where the extreme-value copula is of the form $C_G(\mathbf{u}) = \exp(-L((-\ln u_1), \dots, (-\ln u_d)))$, for $\mathbf{u} \in (0, 1]^d$, L is the stable tail dependence function of G and the corresponding Pickands dependence function satisfies (1.1). We also recall that, for $\alpha \in (0, 1)$, $G_\alpha(\mathbf{x}) = \exp(-(-\ln G(\mathbf{x}))^\alpha)$ is a max-stable distribution, with copula C_{G_α} given in (2.4).

The copula C_{G_α} must be of extreme-value type, i.e. of the form $C_{G_\alpha}(\mathbf{u}) = \exp\{-L_\alpha(-\ln u_1, \dots, -\ln u_d)\}$, $\mathbf{u} \in [0, 1]^d$, for a stable tail-dependence function L_α . We then deduce that

$$L_\alpha((-\ln u_1), \dots, (-\ln u_d)) = L^\alpha((-\ln u_1)^{1/\alpha}, \dots, (-\ln u_d)^{1/\alpha}), \quad (\text{A.19})$$

for all $\mathbf{u} \in [0, 1]^d$. By the homogeneity of the stable-tail dependence function, we have

$$L_\alpha(z_1, \dots, z_d) = (z_1 + \dots + z_d) A_\alpha(\mathbf{t}), \quad \mathbf{z} \in [0, \infty)^d,$$

where $t_j = z_j/(z_1 + \dots + z_d)$ for $j = 1, \dots, d$ and A_α is the Pickands function of G_α , $\alpha \in (0, 1)$. Combining (A.19) with (1.1), we obtain

$$L_\alpha(z_1, \dots, z_d) = \left(\sum_{j=1}^d z_j^{1/\alpha}\right)^\alpha \left\{A\left(\frac{\mathbf{z}^{1/\alpha}}{\sum_{j=1}^d z_j^{1/\alpha}}\right)\right\}^\alpha,$$

where A is the Pickands dependence function of G . Therefore, choosing $\mathbf{z} \in \mathcal{S}_d$ we finally obtain

$$A_\alpha(\mathbf{t}) = \left(\sum_{j=1}^d t_j^{1/\alpha}\right)^\alpha A^\alpha\left(\frac{\mathbf{t}^{1/\alpha}}{\sum_{j=1}^d t_j^{1/\alpha}}\right),$$

which is the result in (3.3).

A.8. Proof of Theorem 4.1

A.8.1. Notation and general setting

Empirical processes. Recall that $(\eta_1, \xi_1), \dots, (\eta_n, \xi_n)$, $n = 1, 2, \dots$, are iid random vectors with distribution Q in (2.1) with fixed $\alpha \in (0, 1)$. For $i = 1, 2, \dots$, and $j \in \{1, \dots, d\}$, let

$$V_i := \Phi_\alpha(\xi_i), \quad U_{i,j} := G_{\alpha,j}(\eta_{i,j}), \quad \widehat{U}_{i,j} := G_{n,j}(\eta_{i,j}), \quad (\text{A.20})$$

where Φ_α is the α -Fréchet distribution, $G_{\alpha,j}$ is the j -th margin of the distributions in the first line of (2.3) and $G_{n,j}$ is as in (4.5). Set $\mathbf{U}_i = (U_{i,1}, \dots, U_{i,d})$ and $\widehat{\mathbf{U}}_i = (\widehat{U}_{i,1}, \dots, \widehat{U}_{i,d})$. In the sequel, when the index i is omitted, we refer to a single observation. For every $\mathbf{u} \in [0, 1]^d$ and $v \in [0, 1]$, define the random copula functions

$$C_{Q,n}(\mathbf{u}, v) := \frac{1}{n} \sum_{i=1}^n \mathbb{1}(\mathbf{U}_i \leq \mathbf{u}, V_i \leq v), \quad C_{G_\alpha,n}(\mathbf{u}) = C_{Q,n}(\mathbf{u}, 1), \quad B_n(v) = C_{Q,n}(\mathbf{1}, v), \quad (\text{A.21})$$

where $\mathbf{1} = (1, \dots, 1)$, and the copula processes

$$\mathbb{C}_{Q,n}(\mathbf{u}, v) = \sqrt{n}(C_{Q,n}(\mathbf{u}, v) - C_Q(\mathbf{u}, v)), \quad \mathbb{C}_{G_\alpha,n}(\mathbf{u}) = \mathbb{C}_{Q,n}(\mathbf{u}, 1), \quad \mathbb{B}_n = \mathbb{C}_{Q,n}(\mathbf{1}, v). \quad (\text{A.22})$$

Let $\mathbb{C}_{G_a}(\mathbf{u}) := \mathbb{C}_Q(\mathbf{u}, 1)$, $\mathbf{u} \in [0, 1]^d$. The covariance function of \mathbb{C}_{G_a} is as in (4.11), with C_Q replaced by C_{G_a} . Furthermore, for every $\mathbf{u} \in [0, 1]^d$, define the empirical copula function and process

$$\widehat{C}_{G_a,n}(\mathbf{u}) = \frac{1}{n} \sum_{j=1}^n \mathbb{1}(\widehat{U}_i \leq \mathbf{u}), \quad \widehat{\mathbb{C}}_{G_a,n} = \sqrt{n}(\widehat{C}_{G_a,n} - C_{G_a}). \quad (\text{A.23})$$

In the sequel we view the above empirical processes as random signed measures, when appropriate (e.g., [van der Vaart and Wellner, 1996](#), Examples 1.7.4 and 1.10.6). We then use the notation $Mf := \int f dM$, for any signed measure M on the measurable space $(\mathbb{X}, \mathcal{X})$ and measurable function f . Furthermore, for asymptotically measurable sequences X_n, X'_n in $\ell^\infty(\mathbb{X})$, with $n \rightarrow \infty$, $X_n(\cdot) = X'_n(\cdot) + o_p(1)$ stands for $\sup_{x \in \mathbb{X}} |X_n(x) - X'_n(x)| = o_p(1)$, and we recall that $X_n(\cdot) \rightsquigarrow X(\cdot)$ is shorthand for $\{X_n(x)\}_{x \in \mathbb{X}} \rightsquigarrow \{X(x)\}_{x \in \mathbb{X}}$ in $\ell^\infty(\mathbb{X})$, as already stated in Section 1.

Weighting and selection maps. For all $f \in \ell([0, 1]^{d+1})$, let $g_\epsilon : \ell([0, 1]^{d+1}) \mapsto \ell([0, 1]^{d+1})$ be the weighting-map given by

$$(g_\epsilon(f))(z) = \begin{cases} w_\epsilon^{-1}(z)f(z), & z \in (0, 1]^{d+1} \setminus \{\mathbf{1}\} \\ 0, & \text{otherwise,} \end{cases} \quad (\text{A.24})$$

where, for any fixed $\epsilon \in [0, 1/2)$, w_ϵ denotes the weighting-function

$$w_\epsilon : [0, 1]^{d+1} \mapsto [0, 1] : z \mapsto \min_{1 \leq j \leq d+1} z_j^\epsilon \left(1 - \min_{1 \leq j \leq d+1} z_j \right)^\epsilon.$$

To keep the notation light, when $f(z)$ is computed at $z = (\mathbf{u}, 1)$, we occasionally still write $g_\epsilon(f)$. The difference in meaning will be clear from the context. For every $f \in \ell([0, 1]^{d+1})$, let $\pi_{1,\dots,d} : \ell([0, 1]^{d+1}) \mapsto \ell([0, 1]^d)$ and $\pi_{d+1} : \ell([0, 1]^{d+1}) \mapsto \ell([0, 1])$ be the selection maps defined by

$$(\pi_{1,\dots,d}(f))(\mathbf{u}) := f(u_1, \dots, u_d, 1), \quad (\pi_{d+1}(f)) := f(1, \dots, 1, v). \quad (\text{A.25})$$

Then, for every $\alpha \in (0, 1)$, $\epsilon \in [0, 1/2)$ and $\mathbf{u} \in (0, 1]^d \setminus \{\mathbf{1}\}$, let $\omega_{\epsilon,\mathbf{u}} : [0, 1]^d \mapsto \mathbb{R}$ be the weighted-function defined by

$$\omega_{\epsilon,\mathbf{u}}(\mathbf{v}) := \frac{\mathbb{1}(\mathbf{v} \leq \mathbf{u}) - C_{G_a}(\mathbf{u})}{(\pi_{1,\dots,d}(w_\epsilon))(\mathbf{u})}. \quad (\text{A.26})$$

For every $\mathbf{u}, \mathbf{v} \in [0, 1]^d$, set

$$\omega'_{\epsilon,\mathbf{u}}(\mathbf{v}) = \begin{cases} \omega_{\epsilon,\mathbf{u}}(\mathbf{v}), & \mathbf{u} \in (0, 1]^d \setminus \{\mathbf{1}\} \\ 0, & \text{otherwise.} \end{cases} \quad (\text{A.27})$$

Proof's overview. By [Genest and Segers \(2009\)](#) and [Gudendorf and Segers \(2012\)](#) we have that as $n \rightarrow \infty$

$$g_\epsilon(\mathbb{C}_{Q,n}) \rightsquigarrow g_\epsilon(\mathbb{C}_Q), \quad g_\epsilon(\mathbb{C}_{G_a,n}) \rightsquigarrow g_\epsilon(\mathbb{C}_{G_a}), \quad (\text{A.28})$$

in $\ell([0, 1]^{d+1})$ and $\ell([0, 1]^d)$, respectively. In particular, $g_\epsilon(\mathbb{C}_Q)$ is a zero-mean Gaussian process on $[0, 1]^{d+1}$ with covariance function

$$\text{Cov}\{(g_\epsilon(\mathbb{C}_Q))(\mathbf{u}), (g_\epsilon(\mathbb{C}_Q))(\mathbf{v})\} = \frac{C_Q(\mathbf{u} \wedge \mathbf{v}) - C_Q(\mathbf{u})C_Q(\mathbf{v})}{w_\epsilon(\mathbf{u})w_\epsilon(\mathbf{v})}, \quad \mathbf{u}, \mathbf{v} \in [0, 1]^{d+1}$$

and $g_\epsilon(\mathbb{C}_{G_a})$ is a zero-mean Gaussian process on the lower dimensional hypercube $[0, 1]^d$ with covariance function defined analogously. The convergence results in (A.28) and, more generally, the asymptotic properties of the copula processes $\mathbb{C}_{Q,n}$ and $\mathbb{C}_{G_a,n}$, are crucial for the derivation of the results presented in Section A.8. As argued in Appendix A.8.3, the P, CFG and MD estimators of A_α belong to a class of estimators $\widehat{A}_{\alpha,n}$ satisfying the following conditions.

Condition A.1. The estimator $\widehat{A}_{\alpha,n}$ of A_α allows the subsequent representation:

- (i) For a continuous linear map $\phi : \ell^\infty([0, 1]^d) \mapsto \ell^\infty(\mathcal{S}_d)$ and some $\epsilon \in [0, 1/2)$,

$$\sqrt{n}\{\ln \widehat{A}_{\alpha,n}(\cdot) - \ln A_\alpha(\cdot)\} = (\phi \circ g_\epsilon(\mathbb{C}_{G_{\alpha,n}}))(\cdot) + o_p(1),$$

where $\mathbb{C}_{G_{\alpha,n}}$ is as in (A.22);

- (ii) For some $m \in \mathbb{N}_+$ and $-\infty \leq a < b \leq \infty$ the map ϕ is of the form

$$(\phi(f))(y) = \sum_{i=0}^m \int_a^b f(\beta_{i,1}(z; t_1), \dots, \beta_{i,d}(z; t_d)) K_i(z; \mathbf{t}) dz. \quad (\text{A.29})$$

where, for $i = 1, \dots, m$, $j \in \{1, \dots, d\}$ and $\mathbf{t} \in \mathcal{S}_d$, $z \mapsto \beta_{i,j}(z, t_j)$ is a bijective and continuous function, while K_1, \dots, K_m are functions satisfying

$$\sup_{\mathbf{t} \in \mathcal{S}_d} \max_{0 \leq i \leq m} |K_i(z; \mathbf{t})| \leq K(z), \quad z \in (a, b), \quad -\infty \leq a < b \leq \infty,$$

for an integrable function K .

An estimator $\widehat{A}_{\alpha,n}$ allowing the above representations together with some suitable estimators $\widehat{\alpha}_n$ of α (that meet some appropriate conditions) enables to deduce a general theory on the weak convergence for composite-estimators \widehat{A}_n^* of A^* , based on the copula process $\mathbb{C}_{Q,n}$. Such a theory is established in Appendix A.8.2 and then applied to the specific cases of the GPWM and ML estimators of α , and the P, CFG and MD estimators for A_α , in Appendix A.8.3. Some auxiliary results are deferred to Appendix A.8.4. The arguments therein make extensive use of composition maps related to ϕ . Hence, to improve the readability of the remaining part of Section A.8 we conclude this subsection by providing a comprehensive list of such composition maps.

Definition A.4. Let ϕ be as in Condition A.1 and $\alpha \in (0, 1)$ denote the true parameter value for the distribution Q in (2.1). Then:

- (i) For a measurable functional $\tau : \ell^\infty([0, 1]) \mapsto \mathbb{R}$, the map $\phi_\tau : \ell^\infty([0, 1]^{d+1}) \mapsto \ell^\infty(\mathcal{S}_d)$ is defined via

$$(\phi_\tau(f))(\mathbf{t}) = \alpha^{-1}(\phi \circ \pi_{1,\dots,d}(f))(\mathbf{t}) + K_\alpha(\mathbf{t})[\tau \circ \pi_{d+1}(w_\epsilon f)], \quad (\text{A.30})$$

for all $f \in \ell^\infty([0, 1]^{d+1})$ and $\mathbf{t} \in \mathcal{S}_d$, with

$$K_\alpha(\mathbf{t}) = \alpha^{-2} \left\{ \|\mathbf{t}\|_{1/\alpha}^{-1/\alpha} \sum_{1 \leq j \leq d: t_j > 0} t_j^{1/\alpha} \ln t_j - \ln A_\alpha(\mathbf{t}) \right\}, \quad (\text{A.31})$$

- π_{d+1} as in (A.25) and w_ϵ as in (A.24). In particular, $\tau \circ \pi_{d+1}(w_\epsilon f)$ is a (real-valued) measurable functional;
- (ii) $\phi' : \ell^\infty([0, 1]^d) \mapsto \ell^\infty(\mathcal{S}_d)$ is defined, for all $f \in \ell^\infty([0, 1]^d)$ and $\mathbf{t} \in \mathcal{S}_d$, via

$$(\phi'(f))(\mathbf{t}) = \alpha^{-1}(\phi(f))(\mathbf{t}) + K_{m+1}(\mathbf{t})f(1, \dots, 1),$$

with K_α as in (A.31) and

$$K_{m+1}(\mathbf{t}) = K_\alpha(\mathbf{t}) - \alpha^{-1} \sum_{i=0}^m \int_a^b K_i(z; \mathbf{t}) dz. \quad (\text{A.32})$$

For a measurable function $\varphi : [0, 1] \mapsto \mathbb{R}$, $\omega'_{\epsilon,u}$ is as in (A.27) and $\delta(\mathbf{u}) = 1$, $\forall \mathbf{u} \in [0, 1]^d$,

$$g'_{\epsilon,\varphi} : M \mapsto \left\{ \int_{[0,1]^{d+1}} [\omega'_{\epsilon,u}(\mathbf{a}) + \delta(\mathbf{u})\varphi(b)] dM(\mathbf{a}, b) \right\}_{\mathbf{u} \in [0,1]^d} \quad (\text{A.33})$$

maps a signed measure M on $[0, 1]^{d+1}$ to an element of $\ell^\infty([0, 1]^d)$. With a little abuse of notation, we also apply this operator to the C_Q -Brownian bridge \mathbb{C}_Q in Theorem 4.1, yielding $g'_{\epsilon,\phi}(\mathbb{C}_Q)$, the zero-mean Gaussian process on $[0, 1]^d$ with covariance function in (A.39);

(iii) $\phi_{\text{MD}} : \ell^\infty([0, 1]^d) \mapsto \ell^\infty(\mathcal{S}_d)$ is a special case of the map ϕ , defined, for every $\mathbf{t} \in \mathcal{S}_d$, by

$$(\phi_{\text{MD}}(f))(\mathbf{t}) = \frac{(1 + A_\alpha(\mathbf{t}))^2}{A_\alpha(\mathbf{t})} \left(\sum_{j=1}^d \int_0^1 \dot{C}_{G_\alpha; j}(v^{t_1}, \dots, v^{t_d}) f(1, \dots, 1, v^{t_j}, 1, \dots, 1) dv - \int_0^1 f(v^{t_1}, \dots, v^{t_d}) dv \right); \quad (\text{A.34})$$

(iv) $\phi_{\text{MD}, \text{ML}} = A_\alpha \phi'_{\text{MD}} \circ g'_{0, \varphi_{\text{ML}}}$, where ϕ'_{MD} is a special case of the map ϕ' in (ii) defined via

$$(\phi'_{\text{MD}}(f))(\mathbf{t}) = \alpha^{-1}(\phi_{\text{MD}}(f))(\mathbf{t}) + K_{d+1}^{\text{MD}}(\mathbf{t})f(1, \dots, 1)$$

for every $f \in \ell^\infty([0, 1]^d)$ and $\mathbf{t} \in \mathcal{S}_d$, with ϕ_{MD} as in (A.34), K_{d+1}^{MD} given by

$$K_{d+1}^{\text{MD}}(\mathbf{t}) = K_\alpha(\mathbf{t}) - \frac{\{1 + A_\alpha(\mathbf{t})\}^2}{\alpha A_\alpha(\mathbf{t})} \left(\sum_{j=1}^d \int_0^1 \dot{C}_{G_\alpha; j}(v^{t_1}, \dots, v^{t_d}) dv - 1 \right)$$

and K_α is as in (A.31); $g'_{0, \varphi_{\text{ML}}}$ is as in (ii), for the particular choices of $\epsilon = 0$ and $\varphi = \varphi_{\text{ML}}$ in (A.41);

(v) $\phi_{\text{MD}, \text{GPWM}} : \ell^\infty([0, 1]^{d+1}) \mapsto \ell^\infty(\mathcal{S}_d)$ is defined, for all $f \in \ell^\infty([0, 1]^d)$ and $\mathbf{t} \in \mathcal{S}_d$, via

$$(\phi_{\text{MD}, \text{GPWM}}(f))(\mathbf{t}) = A_\alpha(\mathbf{t})(\phi_{\tau_{\text{GPWM}}}^{\text{MD}} \circ g_0(f))(\mathbf{t}) = A_\alpha(\mathbf{t})(\phi_{\tau_{\text{GPWM}}}^{\text{MD}}(f))(\mathbf{t}),$$

where $\phi_{\tau_{\text{GPWM}}}^{\text{MD}}$ is a special case of the map ϕ_τ in (A.30), with $\phi = \phi_{\text{MD}}$ in (A.34) and $\tau = \tau_{\text{GPWM}}$ in (A.43);

(vi) $\phi_\circ : \ell^\infty([0, 1]^d) \rightarrow \ell^\infty(\mathcal{S}_d)$ is a special case of the map ϕ defined by

$$(\phi_\circ(f))(\mathbf{t}) = \int_0^\infty f(e^{-vt_1}, \dots, e^{-vt_d}) \beta_\epsilon(e^{-v \max(\mathbf{t})}) dv - \sum_{j=1}^d \int_0^\infty \dot{C}_{G_\alpha; j}(v^{t_1}, \dots, v^{t_d}) f(1, \dots, 1, v^{t_j}, 1, \dots, 1) \beta_\epsilon(e^{-vt_j}) h_\circ(\mathbf{t}; v) dv, \quad (\text{A.35})$$

with $\max(\mathbf{t}) = \max(t_1, \dots, t_d)$, $\beta_\epsilon(v) = v^\epsilon(1-v)^\epsilon$ for $v \in (0, 1)$. Here h_\circ is either $h_p(\mathbf{t}; v) = -A_\alpha^{-1}(\mathbf{t})$ or $h_{\text{CFG}}(\mathbf{t}; v) = 1/v$ for $v > 0$, $\mathbf{t} \in \mathcal{S}_d$;

(vii) $\phi_{\circ, \text{ML}} = A_\alpha \phi'_\circ \circ g'_{\epsilon, \varphi_{\text{ML}}}$, where $g'_{\epsilon, \varphi_{\text{ML}}}$ is as in (ii), with $\varphi = \varphi_{\text{ML}}$ in (A.41), and ϕ'_\circ is a special case of the map ϕ' in (ii) defined via

$$(\phi'_\circ(f))(\mathbf{t}) = \alpha^{-1}(\phi_\circ(f))(\mathbf{t}) + K_{d+1}^\circ(\mathbf{t})f(1, \dots, 1),$$

for every $f \in \ell^\infty([0, 1])$ and $\mathbf{t} \in \mathcal{S}_d$, with ϕ_\circ as in (A.35) and

$$K_{d+1}^\circ(\mathbf{t}) = K_\alpha(\mathbf{t}) - \int_0^\infty \beta_\epsilon(e^{-v \max(\mathbf{t})}) \frac{h_\circ(\mathbf{t}; v)}{\alpha} dv + \sum_{j=1}^d \int_0^\infty \dot{C}_{G_\alpha; j}(v^{t_1}, \dots, v^{t_d}) \beta_\epsilon(e^{-vt_j}) \frac{h_\circ(\mathbf{t}; v)}{\alpha} dv;$$

(viii) $\phi_{\circ, \text{GPWM}} : \ell^\infty([0, 1]^{d+1}) \mapsto \ell^\infty(\mathcal{S}_d)$ is defined, for all $f \in \ell^\infty([0, 1]^d)$ and $\mathbf{t} \in \mathcal{S}_d$, via

$$(\phi_{\circ, \text{GPWM}}(f))(\mathbf{t}) = A_\alpha(\mathbf{t})(\phi_{\tau_{\text{GPWM}}}^\circ \circ g_\epsilon(f))(\mathbf{t})$$

where $\phi_{\tau_{\text{GPWM}}}^\circ$ is a special case of the map ϕ_τ in (A.30), with $\phi = \phi_\circ$ in (A.35) and $\tau = \tau_{\text{GPWM}}$ in (A.43).

A.8.2. General preliminary results

The results in this section rely on the following conditions on the estimators $\widehat{\alpha}_n$ of α .

Condition A.2. Let $\widehat{\alpha}_n$ be an estimator of α satisfying one of the following properties:

(i) There is a continuous linear map $\tau : \ell^\infty([0, 1]) \mapsto \mathbb{R}$ such that

$$\sqrt{n}(\widehat{\alpha}_n - \alpha) = \tau(\mathbb{B}_n) + o_p(1),$$

where \mathbb{B}_n is as in (A.22);

(ii) There is a measurable function $\zeta : (0, +\infty) \mapsto \mathbb{R}$ such that $\Phi_\alpha \zeta = 0$, $\Phi_\alpha \zeta^2 < \infty$ and

$$\sqrt{n}(\widehat{\alpha}_n - \alpha) = n^{-1/2}(\zeta(\xi_1) + \dots + \zeta(\xi_n)) + o_p(1).$$

When a general composite-estimator \widehat{A}_n^* is obtained combining an estimator $\widehat{A}_{\alpha,n}$ of A_α and $\widehat{\alpha}_n$ of α satisfying Conditions A.1(i) and A.2(i), respectively, the functional weak convergence of $\sqrt{n}(\widehat{A}_n^* - A^*)$ can be established by fairly direct arguments. The GPWM-based composite-estimators are examples of estimators for which such asymptotic results hold, see Appendix A.8.3 and Lemma A.11. When $\widehat{\alpha}_n$ complies with A.2(ii), to study the joint limit behaviour of $(\widehat{A}_{\alpha,n}, \widehat{\alpha}_n)$ and ultimately determining that of $\sqrt{n}(\widehat{A}_n^* - A^*)$ needs more complex asymptotic arguments. The composite-estimators based on the M-estimator (van der Vaart, 2000, Ch. 5) and the MLE are examples of estimators for which the second type of asymptotic results apply. The following propositions derive the required theory. Hereafter, the notations with the superscript “’” are specific to the second type of asymptotic results.

Proposition A.5. *Let \widehat{A}_n^* be the composite-estimator obtained by the composition of $\widehat{A}_{\alpha,n}$ and $\widehat{\alpha}_n$ satisfying Conditions A.1(i) and A.2(i), respectively. Let ϕ_τ be as in Definition A.4(i) and g_ϵ be as in (A.24). Then, in $\ell^\infty(\mathcal{S}_d)$*

$$\sqrt{n}\{\widehat{A}_n^*(\cdot) - A^*(\cdot)\} \rightsquigarrow A_\alpha(\cdot)(\phi_\tau \circ g_\epsilon(\mathbb{C}_Q))(\cdot), \quad n \rightarrow \infty. \quad (\text{A.36})$$

Proof. The claim in (A.36) relies on the following result.

Lemma A.6. *With $\|t\|_{1/h}$ being defined as in (3.4), the map*

$$g : (0, \infty) \mapsto \ell^\infty(\mathcal{S}_d) : h \mapsto \left(\ln \|t\|_{1/h}^{1/h} \right)_{t \in \mathcal{S}_d}$$

is Hadamard differentiable at α with derivative

$$\{(\dot{g}_\alpha(h))(t)\}_{t \in \mathcal{S}_d} = \left\{ -h\alpha^{-2} \|t\|_{1/\alpha}^{-1/\alpha} \sum_{1 \leq j \leq d: t_j > 0} t_j^{1/\alpha} \ln t_j \right\}_{t \in \mathcal{S}_d}, \quad 0 < h < \infty.$$

For the proof see Section 4 of the supplementary material. For simplicity we focus on $\ln A^*$ and $\ln \widehat{A}_n^*$. For $t \in \mathcal{S}_d$, we have that

$$\begin{aligned} \sqrt{n}\{\ln \widehat{A}_n^*(t) - \ln A^*(t)\} &= \sqrt{n}\{\widehat{\alpha}_n^{-1} \ln \widehat{A}_{\alpha,n}(t) - \alpha^{-1} \ln A_\alpha(t)\} - \sqrt{n}(\ln \|t\|_{1/\widehat{\alpha}_n}^{1/\widehat{\alpha}_n} - \ln \|t\|_{1/\alpha}^{1/\alpha}) \\ &\equiv T_{1,n}(t) + T_{2,n}(t). \end{aligned}$$

By Condition A.2(i), the functional version of Slutsky's lemma (van der Vaart and Wellner, 1996, p. 32) and the delta method (van der Vaart, 2000, Theorem 3.1) it follows that

$$T_{1,n}(\cdot) = \alpha^{-1} \sqrt{n}\{\ln \widehat{A}_{\alpha,n}(\cdot) - \ln A_\alpha(\cdot)\} - \alpha^{-2} \ln A_\alpha(\cdot) \sqrt{n}(\widehat{\alpha}_n - \alpha) + o_p(1).$$

By Lemma A.6 and the functional delta method (van der Vaart, 2000, Ch. 20) it follows that

$$T_{2,n}(\cdot) = \{K_\alpha(\cdot) + \alpha^{-2} \ln A_\alpha(\cdot)\} \sqrt{n}(\widehat{\alpha}_n - \alpha) + o_p(1),$$

where K_α is as in (A.31). Then, by Conditions A.1(i) and A.2(i) and the identity $\tau \circ \pi_{d+1}(\mathbb{C}_{Q,n}) = \tau \circ \pi_{d+1}(w_\epsilon g_\epsilon(\mathbb{C}_{Q,n}))$, with π_{d+1} as in (A.25) and w_ϵ as in (A.24), we obtain

$$\begin{aligned} T_{1,n}(\cdot) + T_{2,n}(\cdot) &= \alpha^{-1}(\phi \circ g_\epsilon(\mathbb{C}_{G_{\alpha,n}}))(\cdot) + K_\alpha(\cdot) \sqrt{n}(\widehat{\alpha}_n - \alpha) + o_p(1) \\ &= \alpha^{-1}(\phi \circ g_\epsilon(\mathbb{C}_{G_{\alpha,n}}))(\cdot) + K_\alpha(\cdot) \tau(\mathbb{B}_n) + o_p(1) \\ &= (\phi_\tau \circ g_\epsilon(\mathbb{C}_{Q,n}))(\cdot) + o_p(1), \end{aligned} \quad (\text{A.37})$$

Thus, the result in (A.36) follows from (A.28), by applying the continuous mapping theorem and the functional delta method in the last line of (A.37). \square

Proposition A.7. Let \widehat{A}^*_n be the composite-estimator obtained by the composition of $\widehat{A}_{\alpha,n}$ and $\widehat{\alpha}_n$ complying with Condition A.1(i)-(ii) and Condition A.2(ii), respectively, with $\varphi = \zeta \circ \Phi_\alpha^-$ satisfying

$$-\infty < E(\omega'_{\epsilon,u}(U)\varphi(V)) < \infty, \quad \forall u \in (0, 1]^d \setminus \{\mathbf{1}\}, \quad (\text{A.38})$$

where $\omega'_{\epsilon,u}$ is given in (A.27). Let ϕ' and $g'_{\epsilon,\varphi}$ be as in Definition A.4(ii). Then, in $\ell^\infty(S_d)$ as $n \rightarrow \infty$

$$\sqrt{n}\{\widehat{A}^*_n(\cdot) - A^*(\cdot)\} \rightsquigarrow A_\alpha(\cdot)(\phi' \circ g'_{\epsilon,\varphi}(\mathbb{C}_Q))(\cdot),$$

where $g'_{\epsilon,\varphi}(\mathbb{C}_Q)$ is a zero-mean Gaussian process with covariance function defined in (A.39).

Proof. For any $u_i \in [0, 1]^d$, $i \leq k$, $k \in \mathbb{N}_+$, the random vectors $(\omega'_{\epsilon,u_1}(U_i), \dots, \omega'_{\epsilon,u_k}(U_i), \varphi(V_i))$, $i = 1, \dots, n$, are iid with zero-mean and finite pairwise covariances, by arguments in Genest and Segers (2009), Gudendorf and Segers (2012), Condition A.2(ii) and (A.38). Let

$$(g'_\epsilon(\mathbb{C}_{Q,n}))(u) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \omega'_{\epsilon,u}(U_i), \quad \bar{\varphi}_n(u) = \delta(u) \frac{1}{\sqrt{n}} \sum_{i=1}^n \varphi(V_i), \quad u \in [0, 1]^d,$$

where $\delta(u) = 1$ for all $u \in [0, 1]^d$. Note that $(g'_\epsilon(\mathbb{C}_{Q,n}))(u) = (g_\epsilon(\mathbb{C}_{Q,n}))(u)$, with $u \in [0, 1]^d$. Then, $g'_\epsilon(\mathbb{C}_{Q,n})$ and $\bar{\varphi}_n$ are asymptotically tight (van der Vaart and Wellner, 1996, Definition 1.3.7) and, by the central limit theorem,

$$((g'_\epsilon(\mathbb{C}_{Q,n}))(u_1), \dots, (g'_\epsilon(\mathbb{C}_{Q,n}))(u_k), \bar{\varphi}_n(u_1), \dots, \bar{\varphi}_n(u_k)) \rightsquigarrow N(\mathbf{0}, \Sigma)$$

as $n \rightarrow \infty$. By arguments in van der Vaart and Wellner (1996, p. 42 point 3), these facts are sufficient to claim that the class of functions $\mathcal{Y}_{\epsilon,\varphi} := \{(a, b) \mapsto \omega'_{\epsilon,u}(a) + \delta(u)\varphi(b) : u \in [0, 1]^d\}$ is C_Q -Donsker (van der Vaart and Wellner, 1996, pp. 80-82). Indeed, introducing the map

$$g'_{\epsilon,\varphi} : M \mapsto \{Mf : f \in \mathcal{Y}_{\epsilon,\varphi}\}$$

defined on the space of signed measure M on $[0, 1]^{d+1}$, we have that $(g'_{\epsilon,\varphi}(\mathbb{C}_{Q,n}))(\omega'_{\epsilon,u} + \delta(u)\varphi) = (g'_\epsilon(\mathbb{C}_{Q,n}))(u) + \bar{\varphi}_n(u)$, $\forall u \in [0, 1]^d$. Then, as $n \rightarrow \infty$, $g'_{\epsilon,\varphi}(\mathbb{C}_{Q,n}) \rightsquigarrow g'_{\epsilon,\varphi}(\mathbb{C}_Q)$ in $\ell^\infty(\mathcal{Y}_{\epsilon,\varphi})$, where $g'_{\epsilon,\varphi}(\mathbb{C}_Q)$ is a zero-mean Gaussian process with covariance function

$$\begin{aligned} \text{Cov} \{ (g'_{\epsilon,\varphi}(\mathbb{C}_Q))(\omega'_{\epsilon,u} + \delta(u)\varphi), (g'_{\epsilon,\varphi}(\mathbb{C}_Q))(\omega'_{\epsilon,v} + \delta(v)\varphi) \} \\ = \begin{cases} E(\{\omega_{\epsilon,u}(U) + \varphi(V)\}\{\omega_{\epsilon,v}(U) + \varphi(V)\}), & u, v \in \mathcal{V} \\ E(\{\omega_{\epsilon,u}(U) + \varphi(V)\}\varphi(V)), & u \in \mathcal{V}, v \in \mathcal{V}^c \\ E(\varphi^2(V)), & u, v \in \mathcal{V}^c \end{cases} \end{aligned} \quad (\text{A.39})$$

where $\mathcal{V} = (0, 1]^d \setminus \{\mathbf{1}\}$. Since each element of $\mathcal{Y}_{\epsilon,\varphi}$ corresponds to a unique $u \in [0, 1]^d$, we can equivalently think of the codomain of $g'_{\epsilon,\varphi}$ as $\ell^\infty([0, 1]^d)$ (as done in (A.33)) and consider the processes $g'_{\epsilon,\varphi}(\mathbb{C}_{Q,n})$, $g'_{\epsilon,\varphi}(\mathbb{C}_Q)$ as indexed on $[0, 1]^d$. From (A.29), Condition A.2(ii) and the first line of (A.37), it follows that

$$\sqrt{n}\{\ln \widehat{A}^*_n(\cdot) - \ln A^*(\cdot)\} = (\phi' \circ g'_{\epsilon,\varphi}(\mathbb{C}_{Q,n}))(\cdot) + o_p(1).$$

The final results are obtained by applying the continuous mapping theorem and the functional delta method to the above expression. \square

A.8.3. Main body of the proof

In what follows, the asymptotic results concerning the ML- and GPWM-based composite-estimators are established by verifying the assumptions of Proposition A.7 and Proposition A.5, respectively. Firstly, we address the madogram-based case; then, we simultaneously examine P- and CFG-based estimators, due to their similar traits.

We start analyzing the case when α is estimated with the ML estimator in (4.3) and A_α with the MD estimator in (4.7). We recall that the estimator in (4.3) is the unique solution of log-likelihood equation $n^{-1} \sum_{i=1}^n \dot{\mathcal{L}}_{\tilde{\alpha}}(\xi_i) = 0$, where

$$\dot{\mathcal{L}}_{\tilde{\alpha}}(x) = \partial/\partial \tilde{\alpha} \ln \Phi_{\tilde{\alpha}}(x) = 1/\tilde{\alpha} + \ln x(x^{-\tilde{\alpha}} - 1), \quad x > 0.$$

Noting that ξ^{-1} is a Weibull random variable, then by using similar arguments to van der Vaart (2000, Theorem 5.41 and 5.42) it follows that $\widehat{\alpha}_n^{\text{ML}} \xrightarrow{P} \alpha$ as $n \rightarrow \infty$ and

$$\sqrt{n}(\widehat{\alpha}_n^{\text{ML}} - \alpha) = i_\alpha^{-1} n^{-1/2} \{\dot{\mathcal{L}}_\alpha(\xi_1) + \cdots + \dot{\mathcal{L}}_\alpha(\xi_n)\} + o_p(1), \quad (\text{A.40})$$

where $i_\alpha = \alpha^{-2}\{(1 - \zeta^2) + \pi^2/6\}$ is the Fisher information and ζ is Euler's constant.

Assuming that Condition 4.1(i) holds true, then, by Lemma A.8, $\widehat{A}_{\alpha,n}^{\text{MD}}$ satisfies Condition A.1(i)-(ii) with $\phi = \phi_{\text{MD}}$ in (A.34), and $\epsilon = 0$. Furthermore, $\widehat{\alpha}_n^{\text{ML}}$ satisfies Condition A.2(ii) by (A.40) with $\zeta = \zeta_{\text{ML}} = i_\alpha^{-1} \dot{\mathcal{L}}_\alpha$. Define

$$\varphi_{\text{ML}}(v) := \zeta_{\text{ML}} \circ \Phi_\alpha^\leftarrow(v) = i_\alpha^{-1} \alpha^{-1} \{1 + (1 + \ln v) \ln(-\ln v)\}, \quad v \in (0, 1), \quad (\text{A.41})$$

then (A.38) is satisfied with $\varphi = \varphi_{\text{ML}}$, by Lemma A.9. Therefore, from Proposition A.7 it follows that, in $\ell^\infty(\mathcal{S}_d)$,

$$\sqrt{n} \left\{ \widehat{A}_n^{\star, \text{MD, ML}}(\cdot) - A^\star(\cdot) \right\} \rightsquigarrow (\phi_{\text{MD, ML}}(\mathbb{C}_Q))(\cdot), \quad n \rightarrow \infty,$$

where $\phi_{\text{MD, ML}} = A_\alpha \phi'_{\text{MD}} \circ g'_{0, \varphi_{\text{ML}}}$, with details given in Definition A.4(iv), and where $g'_{0, \varphi_{\text{ML}}}(\mathbb{C}_Q)$ is a zero-mean Gaussian process with covariance function

$$\text{Cov}\{g'_{0, \varphi_{\text{ML}}}(\mathbb{C}_Q)(\mathbf{u}), g'_{0, \varphi_{\text{ML}}}(\mathbb{C}_Q)(\mathbf{v})\} = C_{G_\alpha}(\min(\mathbf{u}, \mathbf{v})) - C_{G_\alpha}(\mathbf{u})C_{G_\alpha}(\mathbf{v}) + T_\alpha(\mathbf{u}) + T_\alpha(\mathbf{v}) + 1,$$

for every $\mathbf{u}, \mathbf{v} \in [0, 1]^d$, with

$$T_\alpha(\cdot) = \frac{1}{i_\alpha \alpha} \left(C_{G_\alpha}(\mathbf{u}) - \int_0^1 \frac{\partial}{\partial v} C_Q(\mathbf{u}, v) (1 + \ln v) \ln(-\ln v) dv \right).$$

Finally, from the weak convergence result and the functional version of Slutsky's lemma it follows that $\|\widehat{A}_n^{\star, \text{MD, ML}} - A^\star\|_\infty \xrightarrow{P} 0$ as $n \rightarrow \infty$.

Next, we study the case when α is estimated with the GPWM estimator in (4.1) and A_α with the MD estimator in (4.7). Here, we additionally assume that $\alpha > 1/(k-1)$. By Lemma A.11, the estimator $\widehat{\alpha}_n^{\text{GPWM}}$ satisfies Condition A.2(i) with $\tau = \tau_{\text{GPWM}}$ given in (A.43). Then, by Proposition A.5 it follows that, in $\ell^\infty(\mathcal{S}_d)$,

$$\sqrt{n} \left\{ \widehat{A}_n^{\star, \text{MD, GPWM}}(\cdot) - A^\star(\cdot) \right\} \rightsquigarrow (\phi_{\text{MD, GPWM}}(\mathbb{C}_Q))(\cdot), \quad n \rightarrow \infty,$$

where $\phi_{\text{MD, GPWM}}(f)$, $f \in \ell^\infty([0, 1]^{d+1})$, is given in Definition A.4(v). Furthermore, given the result in Lemma A.11 and since $\|n^{-1/2} \mathbb{B}_n\|_\infty \xrightarrow{\text{as}} 0$, then we have that $\widehat{\alpha}_n^{\text{GPWM}} \xrightarrow{\text{as}} \alpha$ as $n \rightarrow \infty$. Consequently, by Lemma A.10,

$$\left\| \widehat{A}_n^{\star, \text{MD, GPWM}} - A^\star \right\|_\infty \xrightarrow{\text{as}} 0, \quad n \rightarrow \infty.$$

Now, we study the case when α is estimated with the ML estimator in (4.3) and A_α with the P and CFG estimator in (4.4) and (4.6), respectively. Assuming that Conditions 4.1(i) and 4.1(ii) hold true, then by Segers (2012, Proposition 3.1 and 4.2) we have

$$\widehat{\mathbb{C}}_{G_\alpha, n}(\mathbf{u}) = \mathbb{C}_{G_\alpha, n}(\mathbf{u}) - \sum_{j=1}^d \dot{C}_{G_\alpha; j} \mathbb{C}_{G_\alpha, n}(1, \dots, 1, u_j, 1, \dots, 1) + R_n(\mathbf{u}),$$

where $\widehat{\mathbb{C}}_{G_\alpha, n}$ is as in (A.23) and almost surely $\|R_n\|_\infty = O(n^{-1/4}(\ln n)^{1/2}(\ln \ln n)^{1/4})$, $n \rightarrow \infty$. Then, using similar arguments to those in Gudendorf and Segers (2012, pp. 3082–3083) (and the functional delta method for the P

estimator) we have that $\widehat{A}_{\alpha,n}^P$ and $\widehat{A}_{\alpha,n}^{CFG}$ satisfy Condition A.1(i)-(ii) with $\phi = \phi_P$ and $\phi = \phi_{CFG}$. Precisely, for any fixed $\epsilon \in (0, 1/2)$

$$\sqrt{n}\{\ln \widehat{A}_{\alpha,n}^\circ(\cdot) - \ln A_\alpha(\cdot)\} = (\phi_\circ \circ g_\epsilon(\mathbb{C}_{G_{\alpha,n}}))(\cdot) + o_p(1),$$

where ϕ_\circ is given in A.4(vi) and satisfies the representation in (A.29). By Lemma A.9, the expectation in (A.38) is finite for $\varphi = \varphi_{ML}$ and any given $\epsilon \in (0, 1/2)$. Then, by Proposition A.7,

$$\sqrt{n}\{\widehat{A}_n^{\star, ML}(\cdot) - A^\star(\cdot)\} \rightsquigarrow (\phi_{\circ, ML}(\mathbb{C}_Q))(\cdot), \quad n \rightarrow \infty$$

in $\ell^\infty(\mathcal{S}_d)$, where $\phi_{\circ, ML} = A_\alpha \phi'_\circ \circ g'_{\epsilon, \varphi_{ML}}$, with details given in Definition A.4(vii), and where $g'_{\epsilon, \varphi_{ML}}(\mathbb{C}_Q)$ is a zero-mean Gaussian process with covariance function as in (A.39), for $\varphi = \varphi_{ML}$. Finally, from this result and the functional version of Slutsky's lemma, it follows that $\|\widehat{A}_n^{\star, P, ML} - A^\star\|_\infty \xrightarrow{P} 0$ and $\|\widehat{A}_n^{\star, CFG, ML} - A^\star\|_\infty \xrightarrow{P} 0$ as $n \rightarrow \infty$.

Concluding, we study the case when α is estimated with the GPWM estimator in (4.1) and A_α with the P and CFG estimators in (4.4) and (4.6). Assuming in addition to the previous case that $\alpha > 1/(k-1)$, then by Proposition A.5 we have that in $\ell^\infty(\mathcal{S}_d)$

$$\sqrt{n}\{\widehat{A}_n^{\star, \circ, GPWM}(\cdot) - A^\star(\cdot)\} \rightsquigarrow (\phi_{\circ, GPWM}(\mathbb{C}_Q))(\cdot), \quad n \rightarrow \infty,$$

where $\phi_{\circ, GPWM}(f)$, $f \in \ell^\infty([0, 1]^{d+1})$, is given in Definition A.4(viii). Ultimately, from this result and the functional version of Slutsky's lemma, it follows that

$$\|\widehat{A}_n^{\star, P, GPWM} - A^\star\|_\infty \xrightarrow{P} 0, \quad \|\widehat{A}_n^{\star, CFG, GPWM} - A^\star\|_\infty \xrightarrow{P} 0, \quad n \rightarrow \infty.$$

A.8.4. Auxiliary results

Lemma A.8. Under Condition 4.1(i) we have

$$\sqrt{n}\{\ln \widehat{A}_{\alpha,n}^{MD}(\cdot) - \ln A_\alpha(\cdot)\} = (\phi_{MD}(\mathbb{C}_{G_{\alpha,n}}))(\cdot) + o_p(1),$$

where ϕ_{MD} is given in Definition A.4(iii).

Lemma A.9. Let (U, V) be defined as in (A.20). Let $\omega'_{\epsilon, u}$ be the function defined in (A.27) and φ_{ML} in (A.41). Then, for every $\epsilon \in [0, 1/2)$ and $u \in (0, 1]^d \setminus \{1\}$, we have $E(\omega'_{\epsilon, u}(U)\varphi_{ML}(V)) \in \mathbb{R}$, i.e. the expectation is finite.

Lemma A.10. Let $\widehat{\alpha}_n$ be an estimator of α satisfying $\widehat{\alpha}_n \xrightarrow{as} \alpha$ as $n \rightarrow \infty$ and $\widehat{A}_{\alpha,n}^{MD}$ be the estimator of A_α given in (4.7). Let

$$\widehat{A}_n^\star(t) = (\widehat{A}_{\alpha,n}^{MD}(t)/\|t\|_{1/\widehat{\alpha}_n})^{1/\widehat{\alpha}_n}, \quad t \in \mathcal{S}_d.$$

Then,

$$\|\widehat{A}_n^\star - A^\star\|_\infty \xrightarrow{as} 0, \quad n \rightarrow \infty. \quad (\text{A.42})$$

Lemma A.11. Assume that $\alpha > 1/(k-1)$. Then, almost surely as $n \rightarrow \infty$

$$\sqrt{n}(\widehat{\alpha}_n^{GPWM} - \alpha) = \tau_{GPWM}(\mathbb{B}_n) + o(1),$$

where $\tau_{GPWM} : \ell^\infty([0, 1]) \rightarrow \mathbb{R}$ is defined as

$$\begin{aligned} \tau_{GPWM}(f) &= -2 \int_0^1 f(v) \frac{v(-\ln v)^k \{\mu_{1,k-1} - \mu_{1,k}/(-\ln v)\}}{\Phi_\alpha(\Phi_\alpha^\leftarrow(v))(k\mu_{1,k-1} - 2\mu_{1,k})^2} dv, \\ \mu_{a,b} &= \int_0^1 \Phi_\alpha^\leftarrow(v) v^a (-\ln v)^b dv, \quad a, b \in \mathbb{N} \end{aligned} \quad (\text{A.43})$$

and $\Phi_\alpha(v) = \partial/\partial v \Phi_\alpha(v)$, $v \in (0, 1)$.

For the proofs see Section 4 of the supplementary material.

Acknowledgements

S. A. Padoan is supported by the Bocconi Institute for Data Science and Analytics (BIDSA). E. Hashorva is supported by SNSF Grant 200021-175752/1.

References

- Anderson, C. W. and Turkman, K. F. (1995). *Sums and Maxima of Stationary Sequences with Heavy Tailed Distributions.*, *Sankhyā: The Indian Journal of Statistics, Series A (1961-2002)* **57**, 1-10.
- Barakat, C., Thiran, P., Iannaccone, G., Diot, C. and Owezarski, P. (2002). *A flow-based model for internet backbone traffic*, in Proc. SIGCOMM Internet Meas. Workshop, New York, 35–47.
- Barakat, H. M. and El-Shandidy, M. A. (1990). On the limit distribution of the extremes of a random number of independent random variables. *J. Statist. Plann. Inference*, **26**, 353–361. [MR1086106](#)
- Berghaus, B., Bücher, A., and Dette, H. (2013). Minimum distance estimators of the Pickands dependence function and related tests of multivariate extreme-value dependence. *J. SFS*, **154**, 116–137. [MR3089619](#)
- Bingham, N., Goldie, C. and Teugels, J. (1989). *Regular variation*, Encyclopedia of Mathematics and its Applications, 27. Cambridge University Press, Cambridge. [MR1015093](#)
- Bonald, T., Proutire, A., Rgni, G. and Roberts, J.W. (2001). Insensitivity results in statistical bandwidth sharing. *Teletr. Scie. and Engin.*, **4**, 125–136.
- Capéraà, P., Fougères, A.-L. and Genest, C. (1997). A nonparametric estimation procedure for bivariate extreme value copulas. *Biometrika*, **84** 567–577. [MR1603985](#)
- Cormier, E., Genest, C. and Neslehova, J. G. (2014). Using B-splines for nonparametric inference on bivariate extreme-value copulas. *Extremes*, **17**, 633–659. [MR3282690](#)
- de Haan, L. (1984). A spectral representation for max-stable processes. *Ann. Probab.*, **12**, 1194–1204. [MR0757776](#)
- Falk, M., Hüsler, J. and Reiss, R-D. (2011). *Laws of small numbers: extremes and rare events*, extended edn. Birkhäuser/Springer Basel AG, Basel. [MR2732365](#)
- Fougères, A.-L., Nolan, P. N. and Rootzén, H. (2009). Models for dependent extremes using stable mixtures. *Scand. J. Stat.*, **36**, 42–59. [MR2508330](#)
- Fougères, A.-L., Mercadier, M. and Nolan, P. N. (2013). Dense classes of multivariate extreme value distributions. *J. Multivar. Anal.*, **116**, 109–129. [MR3049895](#)
- Embrechts, P., Klüppelberg, C., and Mikosch, T. (1997). *Modelling extremal events, for insurance and finance*, Springer-Verlag, Berlin. [MR1458613](#)
- Genest, C. and Segers, J. (2009). Rank-based inference for bivariate extreme-value copulas. *Ann. Statist.*, **37**, 2990–3022. [MR2541453](#)
- Gentle, J. E. (2009). *Computational statistics*, Statistics and Computing. Springer, New York. [MR2537506](#)
- Gudendorf, G. and Segers, J. (2012). Nonparametric estimation of multivariate extreme-value copulas. *J. Statist. Plann. Inference*, **142**, 3073–3085. [MR2956794](#)
- Guillou, A., Naveau, P. and Schorgen, A. (2014). Madogram and asymptotic independence among maxima. *REV-STAT*, **12** 119–134. [MR3228763](#)
- Hashorva, E., Ratovomirija, G. and Tamraz, M. (2017). On some new dependence models derived from multivariate collective models in insurance applications. *Scand. Actuar. J.*, **8**, 730–750. [MR3692985](#)
- Joe, H. (2015). *Dependence modeling with copulas*, CRC Press. [MR3328438](#)
- Klüppelberg, C. and May, A. (2006). Bivariate extreme value distributions based on polynomial dependence functions *Math. Meth. Appl. Sci.* **29** 1467–1480. [MR2247312](#)
- Kojadinovic, I., and Yan, J. (2010). Modeling multivariate distributions with continuous margins using the copula R package *Journal of Statistical Software*, **34**(9), 1–20.
- Loiseau, P., Goncalves, P., Dewaele, G., Borgnat, P., Abry, P. and Primet, P. V. B. (2010). Investigating self-similarity and heavy-tailed distributions on a large-scale experimental facility. *IEEE/ACM Trans. Netw.*, **18**, 1261–1274.
- Marcon, G., Padoan, S. A., Naveau, P., Muliere, P., and Segers, J. (2017). Multivariate nonparametric estimation of the Pickands dependence function using Bernstein polynomials. *J. Statist. Plann. Inference*, **183**, 1–17. [MR3589492](#)

- Markovich, N. (2007). *Nonparametric analysis of univariate heavytailed data: research and practice*, John Wiley & Sons, Ltd. [MR2364666](#)
- Maulik, K. and Resnick, S. (2004). Characterizations and examples of hidden regular variation *Extremes* **7** 31–67. [MR2201191](#)
- Mitra, A. and Resnick, S. (2011). Hidden regular variation and detection of hidden risks *Stochastic Models* **27** 591–614.
- Pickands III, J. (1981). Multivariate extreme value distributions, with a discussion *Bull. Inst. Internat. Statist.*, **49**, 859–878, 894–902. [MR0820979](#)
- R Core Team (2014). R: A Language and Environment for Statistical Computing.
- Resnick, S. I. (2007). *Extreme Values, Regular Variation, and Point Processes*, reprint of the 1987 original. Springer Series in Operations Research and Financial Engineering. Springer, New York. [MR2364939](#)
- Resnick, S. (2002). Hidden regular variation, second order regular variation and asymptotic independence *Extremes* **5** 303–336. [MR2002121](#)
- Robert, C. (2013). Some new classes of stationary max-stable random fields. *Statist. Probab. Lett.*, **83**, 1496–1503. [MR3048315](#)
- Robert, C. Y. and Segers J. (2008). Tails of random sums of a heavy-tailed number of light-tailed terms. *Insurance Math. Econom.*, **43**, 85–92. [MR2442033](#)
- Segers, J. (2012). Asymptotics of empirical copula processes under non-restrictive smoothness assumptions *Bernoulli*, **18**, 764–782. [MR2948900](#)
- Silvestrov, D. S. and Teugels, J. L. (1998). Limit theorems for extremes with random sample sizes. *Adv. in Appl. Probab.*, **30**, 777–806. [MR1663553](#)
- Stephenson, A. G. (2002). evd: Extreme Value Distributions *R News*, **2**.
- Taqqu, M. S., Willinger, W. and Sherman, R. (1997). Proof of a fundamental result in self-similar traffic modeling. *Comput. Commun. Rev.*, **27**, 5–23.
- Uchaikin, V. V. and Zolotarev, V. M. (2011). *Chance and stability. Stable distributions and their applications*, with a foreword by V. Yu. Korolev and Zolotarev. Modern Probability and Statistics. VSP, Utrecht. [MR1745764](#)
- van de Meent, R. (2006). *Network Link Dimensioning – A Measurement and Modeling based Approach*, Ph.D. thesis, Univ. of Twente. <http://purl.org/utwente/56434>
- van der Vaart, A. W. (2000). *Asymptotic Statistics*, Cambridge University Press. [MR1652247](#)
- van der Vaart, A. W. and Wellner, J. A. (1996). *Weak Convergence and Empirical Processes. With Applications to Statistics*, Springer Verlag, New York. [MR1385671](#)
- Willinger, W., Taqqu, M. S., Sherman, R. and Wilson, D. V. (1997). Self-similarity through high-variability: Statistical analysis of ethernet lan traffic at the source level. *IEEE/ACM Trans. Netw.*, **5**, 71–86.
- Zhang, D., Wells, M. T. and Peng, L. (2008). Nonparametric estimation of the dependence function for a multivariate extreme value distribution *J. Multivariate Anal.* **99** 577–588. [MR2406072](#)