

OPTIMAL ESTIMATION OF SIMULTANEOUS SIGNALS USING ABSOLUTE INNER PRODUCT WITH APPLICATIONS TO INTEGRATIVE GENOMICS

Rong Ma¹, T. Tony Cai² and Hongzhe Li¹

Department of Biostatistics, Epidemiology and Informatics¹

Department of Statistics²

University of Pennsylvania

Philadelphia, PA 19104

Abstract

Integrating the summary statistics from genome-wide association study (GWAS) and expression quantitative trait loci (eQTL) data provides a powerful way of identifying the genes whose expression levels are potentially associated with complex diseases. A parameter called T -score that quantifies the genetic overlap between a gene and the disease phenotype based on the summary statistics is introduced based on the mean values of two Gaussian sequences. Specifically, given two independent samples $\mathbf{x}_n \sim N(\theta, \Sigma_1)$ and $\mathbf{y}_n \sim N(\mu, \Sigma_2)$, the T -score is defined as $\sum_{i=1}^n |\theta_i \mu_i|$, a non-smooth functional, which characterizes the amount of shared signals between two absolute normal mean vectors $|\theta|$ and $|\mu|$. Using approximation theory, estimators are constructed and shown to be minimax rate-optimal and adaptive over various parameter spaces. Simulation studies demonstrate the superiority of the proposed estimators over existing methods. The method is applied to an integrative analysis of heart failure genomics datasets and we identify several genes and biological pathways that are potentially causal to human heart failure.

KEY WORDS: eQTL; Genetic association; Minimax lower bound; Non-smooth functional.

1 Introduction

1.1 Integrating summary data from GWAS and eQTL studies

Integrative genomics aims to integrate various biological data sets for systematic discovery of genetic basis that underlies and modifies human disease (Giallourakis et al., 2005). To realize its full potential in genomic research, methods of both computational efficiency and theoretical

guarantee for such integrative analyses are needed in various applications. This paper proposes a method that combines datasets from genome-wide association studies (GWAS) and expression quantitative trait loci (eQTL) studies in order to identify genetically regulated disease genes and to provide an integrative view of the underlying biological mechanism of complex diseases such as heart failure. Results from GWAS have revealed that the majority of disease-associated single nucleotide polymorphisms (SNPs) lie in non-coding regions of the genome (Hindorff et al., 2009). These SNPs likely regulate the expression of a set of downstream genes that may have effects on diseases (Nicolae et al., 2010). On the other hand, eQTL studies measure the association between both cis- and trans- SNPs and the expression levels of genes, which characterizes how genetic variants regulate transcriptions. A key next step in human genetic research is to explore whether these intermediate cellular level eQTL signals are located in the same loci (“colocalize”) as GWAS signals and potentially mediate the genetic effects on disease, and to find disease genes whose eQTL overlap significantly with the set of loci associated with the disease (He et al., 2013).

This paper focuses on the integrative analysis of the summary statistics of GWAS and eQTL studies performed on possibly different set of subjects. Due to the privacy and confidentiality concerns of GWAS/eQTL participants, the raw genotype data are often not available, instead most of the published papers provide summary statistics that include single SNP analysis results such as the estimated effect size, its p-value and the minor allele frequency. Based on these summary statistics, we propose a method that identifies potential disease genes by measuring their genetic overlaps to the disease. In particular, we propose a gene-specific measure, T -score, that characterizes the total amount of simultaneous SNP signals that share the same loci in both GWAS and eQTL study of a relevant normal tissues. Such a measure enables us to prioritize genes whose expression levels may underlie and modify human disease (Zhao et al., 2017).

Treating SNP-specific GWAS and eQTL summary z -score statistics (as obtained for linear or logistic regression coefficients) as two independent sequences of Gaussian random variables, we define the parameter T -score as the sum of the product of the absolute values of two normal means over a given set of n SNPs. Specifically, for any individual gene g , we denote \mathbf{x}_n^g the vector of z -scores from eQTL study, and \mathbf{y}_n the vector of z -scores from GWAS. We assume $\mathbf{x}_n^g \sim N(\theta^g, \Sigma_1)$ and $\mathbf{y}_n \sim N(\mu, \Sigma_2)$ for some $\theta^g, \mu \in \mathbb{R}^n$ and covariance matrices $\Sigma_1, \Sigma_2 \in \mathbb{R}^{n \times n}$ with unit diagonals. The T -score for gene g is then defined as

$$T\text{-score}(g) = \sum_{i=1}^n |\theta_i^g \mu_i|, \quad (1)$$

where the summation is over a given set of n SNPs. The T -score quantifies the amount of simultaneous signals contained in two Gaussian mean vectors, regardless of the directions of the signals. Intuitively, a large T -score would possibly result from a large number of contributing components i 's whose means θ_i^g and μ_i are simultaneously large in absolute values. The supports (nonzero

coordinates) of the mean vectors θ (hereafter we omit its dependence on g for simplicity) and μ are assumed to have sparse overlaps since it has been observed that, for a relatively large set of SNPs, only a small subset of SNPs are associated with both disease and gene expression (He et al., 2013). By estimating the T -scores for all the genes using summary statistics, we would be able to, after proper normalizations that accounts for study sample sizes, the number of SNPs and effect sizes (see Section 2.5), identify and prioritize those genetically regulated candidate disease genes. Besides, the T -scores can also be used in the Gene Set Enrichment Analysis to identify the disease associated gene sets and pathways, or to quantify the genetic sharing among different complex traits using the GWAS summary statistics (Bulik-Sullivan et al., 2015).

1.2 Related works

Statistically, estimation of T -score involves estimating a non-smooth functional – the absolute value function – of Gaussian random variables. Unlike the problems of estimating smooth functionals such as the linear or quadratic functionals (Ibragimov and Khas’minskii, 1985; Donoho and Nussbaum, 1990; Fan, 1991; Efromovich and Low, 1994; Cai and Low, 2006) where some natural unbiased estimators are available, much less is known for estimating the non-smooth functionals. Using approximation theory, Cai and Low (2011) established the minimax risk and constructed a minimax optimal procedure for estimating a non-smooth functional. More recently, this idea has been adapted to statistical information theory that also considered estimation of non-smooth functionals such as the Rényi entropy, support size, and L_1 distance (Jiao et al., 2015, 2016; Wu and Yang, 2016, 2019; Acharya et al., 2016). Nonetheless, how to estimate the absolute inner product of two Gaussian mean vectors (T -score) remains unknown.

In the statistical genetics and genomics literature, several approaches have been proposed for integrating GWAS and eQTL data sets. Under the colocalization framework, methods such as Nica et al. (2010) and Giambartolomei et al. (2014) were developed to detect colocalised SNPs. However, these methods do not directly identify the potential causal genes. Under the transcriptome-wise association study (TWAS) framework, Zhu et al. (2016) proposed a summary data-based Mendelian randomization method for causal gene identification, by posing some structural causality assumptions. Gamazon et al. (2015) developed a gene-based association method called PrediXcan that directly tests the molecular mechanisms through which genetic variation affects phenotype. Nevertheless, there is still a need for a quantitative measure of the genetic sharing between genes and the disease that can be estimated from the GWAS/eQTL summary statistics.

As a related but different quantity, the genetic covariance ρ , proposed by Bulik-Sullivan et al. (2015) as a measure of the genetic sharing between two traits, can be expressed using our notation as $\rho = \sum_{i=1}^n \theta_i \mu_i$. In addition to the difference due to the absolute value function, in the original definition of genetic covariance ρ , the mean vectors θ and μ represent the conditional effect sizes (i.e., conditional on all other SNPs in the genome), whereas the mean vectors in our T -score correspond

to the marginal effect sizes, so as to be directly applicable to the standard GWAS/eQTL summary statistics. In addition, unlike the LD-score regression approach considered in Bulik-Sullivan et al. (2015), our proposed method takes advantage of the fact that the support overlap between θ and μ are expected to be very sparse.

1.3 Main contributions

In this paper, we propose an estimator of the T -score, based on the idea of thresholding and truncating the best polynomial approximation estimator. To the best of our knowledge, this is the first result concerning estimation of such absolute inner product of two Gaussian mean vectors. Under the framework of statistical decision theory, the minimax lower bounds are obtained and we show that our proposed estimators are minimax rate-optimal over various parameter spaces. In addition, our results indicate that the proposed estimators are locally adaptive to the unknown sparsity level and the signal strength (Section 2). Our simulation study shows the strong empirical performance and robustness of the proposed estimators in various settings, and provides guidelines for using our proposed estimators in practice (Section 3). Analysis of GWAS and eQTL data sets of heart failure using the proposed method identifies several important genes that are functionally relevant to the etiology of human heart failure (Section 4).

2 Minimax Optimal Estimation of T-score

2.1 Minimax Lower Bounds

We start with establishing the minimax lower bounds for estimating T -score over various parameter spaces. Throughout, we denote $T(\theta, \mu) = \sum_{i=1}^n |\theta_i \mu_i|$. For a vector $a = (a_1, \dots, a_n)^\top \in \mathbb{R}^n$, we define the ℓ_∞ norm $\|a\|_\infty = \max_{1 \leq j \leq n} |a_j|$. For sequences $\{a_n\}$ and $\{b_n\}$, we write $a_n \lesssim b_n$ or $b_n \gtrsim a_n$ if there exists a constant C such that $a_n \leq C b_n$ for all n . We write $a_n \asymp b_n$ if $a_n \lesssim b_n$ and $a_n \gtrsim b_n$.

As of both practical and theoretical interest, we focus on the class of mean vector pairs (θ, μ) with only a small fraction of support overlaps. Specifically, for any $s < n$, we define the parameter space for (θ, μ) as $D(s) = \{(\theta, \mu) \in \mathbb{R}^n \times \mathbb{R}^n : |\text{supp}(\theta) \cap \text{supp}(\mu)| \leq s\}$. Intuitively, in addition to the sparsity s , the difficulty of estimating T -score $T(\theta, \mu)$ should also rely on the magnitudes of the mean vectors θ and μ , and the covariance matrices Σ_1 and Σ_2 . Towards this end, we define the parameter space for $(\theta, \mu, \Sigma_1, \Sigma_2)$ as

$$D^\infty(s, L_n) = \{(\theta, \mu, \Sigma_1, \Sigma_2) : (\theta, \mu) \in D(s), \max(\|\theta\|_\infty, \|\mu\|_\infty) \leq L_n, \Sigma_1 = \Sigma_2 = \mathbf{I}_n\},$$

where both s and L_n can growth with n . In particular, we calibrate the sparsity $s \asymp n^\beta$ for some $0 < \beta < 1$. Throughout, we consider the normalized loss function as the squared distance scaled

by n^{-2} and define the estimation risk for some estimator \hat{T} as

$$\mathcal{R}(\hat{T}) = \frac{1}{n^2} \mathbb{E}(\hat{T} - T(\theta, \mu))^2.$$

To simplify our statement, we define the rate function

$$\psi(s, n) = \log \left(1 + \frac{n}{s^2} \right) + \frac{1}{\log s}.$$

The following theorem establishes the minimax lower bound over the parameter space $D^\infty(s, L_n)$.

Theorem 1. *Let $\mathbf{x}_n \sim N(\theta, \Sigma_1)$ and $\mathbf{y}_n \sim N(\mu, \Sigma_2)$ be multivariate Gaussian random vectors where $(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)$ and $L_n \gtrsim \sqrt{\log n}$. Then*

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)} \mathcal{R}(\hat{T}) \gtrsim \frac{L_n^2 s^2 \psi(s, n)}{n^2} \quad (2)$$

where \hat{T} is any estimator based on $(\mathbf{x}_n, \mathbf{y}_n)$.

From the above theorem and the definition of the rate function $\psi(s, n)$, apparently, there is a discrepancy between the cases when $\beta \in (0, 1/2)$ and when $\beta \in [1/2, 1)$. Specifically, when $\beta \in (0, 1/2)$, the minimax lower bound (2) becomes

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)} \mathcal{R}(\hat{T}) \gtrsim \frac{L_n^2 s^2 \log n}{n^2}, \quad (3)$$

whereas when $\beta \in [1/2, 1)$, we have

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)} \mathcal{R}(\hat{T}) \gtrsim \frac{L_n^2 s^2}{n^2 \log n}. \quad (4)$$

These results suggest that the minimax optimal estimators of $T(\theta, \mu)$ might not be the same across different regions of β .

2.2 Optimal estimators of T-score via polynomial approximation

In general, the proposed estimators are based on the idea of optimal estimation of the absolute value of normal means studied by Cai and Low (2011). Therein, the best polynomial approximation of the absolute value function was applied to obtain the rate-optimal estimator, as well as the minimax lower bound. Specifically, it was shown that, if we define the $2K$ -degree polynomial

$$G_K(x) = \frac{2}{\pi} T_0(x) + \frac{4}{\pi} \sum_{k=1}^K (-1)^{k+1} \frac{T_{2k}(x)}{4k^2 - 1} \equiv \sum_{k=0}^K g_{2k} x^{2k}, \quad (5)$$

where $T_k(x) = \sum_{j=0}^{\lfloor k/2 \rfloor} (-1)^j \frac{k!}{k-j!} \binom{k-j}{j} 2^{k-2j-1} x^{k-2j}$ are Chebyshev polynomials, then for any $X \sim N(\theta, 1)$, if H_k are Hermite polynomials with respect to the standard normal density ϕ such that

$$\frac{d^k}{dy^k} \phi(y) = (-1)^k H_k(y) \phi(y), \quad (6)$$

the estimator

$$\tilde{S}_K(X) \equiv \sum_{k=0}^K g_{2k} M_n^{-2k+1} H_{2k}(X) \quad (7)$$

for some properly chosen K and M_n has some optimality properties for estimating $|\theta|$. This important result motivates our construction of the optimal estimators of T -score.

We begin by considering the setting where $\mathbf{x}_n = (x_1, \dots, x_n)^\top \sim N(\theta, \mathbf{I}_n)$ and $\mathbf{y}_n = (y_1, \dots, y_n)^\top \sim N(\mu, \mathbf{I}_n)$. To estimate $T(\theta, \mu)$, we first split each sample into two copies, one is used for testing, and the other is used for estimation. Specifically, for $x_i \sim N(\theta_i, 1)$, one can generate x_{i1} and x_{i2} from x_i by letting $z_i \sim N(0, 1)$ and setting $x'_{i1} = x_i + z_i$ and $x'_{i2} = x_i - z_i$. Let $x_{il} = x'_{il}/\sqrt{2}$ for $l = 1, 2$, then $x_{il} \sim N(\theta'_i, 1)$ for $l = 1, 2$ and $i = 1, \dots, n$ with $\theta'_i = \theta_i/\sqrt{2}$. Similarly, one can construct $y_{il} \sim N(\mu'_i, 1)$ for $l = 1, 2$ and $i = 1, \dots, n$ with $\mu'_i = \mu_i/\sqrt{2}$. Since $T(\theta, \mu) = 2T(\theta', \mu')$, estimating $T(\theta, \mu)$ with $\{x_i, y_i\}_{i=1}^n$ is then equivalent to estimating $T(\theta', \mu')$ with $\{x_{il}, y_{il}\}_{i=1}^n, l = 1, 2$.

In light of the estimator defined in (7), we consider a slightly adjusted statistic $S_K(X) = \sum_{k=1}^K g_{2k} M_n^{-2k+1} H_{2k}(X)$ and define its truncated version

$$\delta_K(X) = \min\{S_K(X), n^2\},$$

with $M_n = 8\sqrt{\log n}$ and $K \geq 1$ to be specified later. It will be seen that the above truncation is important in reducing the variance of $\delta_K(X)$. Following the sample splitting procedure, we construct an estimator of $|\theta'_i|$ as

$$\hat{V}_{i,K}(x_i) = \delta_K(x_{i1})I(|x_{i2}| \leq 2\sqrt{2\log n}) + |x_{i1}|I(|x_{i2}| > 2\sqrt{2\log n}),$$

and a similar estimator of $|\mu'_i|$ as $\hat{V}_{i,K}(y_i)$. To further exploit the sparse structure, we also consider their thresholded version

$$\hat{V}_{i,K}^S(x_i) = \delta_K(x_{i1})I(\sqrt{2\log n} < |x_{i2}| \leq 2\sqrt{2\log n}) + |x_{i1}|I(|x_{i2}| > 2\sqrt{2\log n})$$

as an estimator of $|\theta'_i|$ and similarly $\hat{V}_{i,K}^S(y_i)$ for $|\mu'_i|$. Intuitively, both $\hat{V}_{i,K}(x_i)$ and $\hat{V}_{i,K}^S(x_i)$ are hybrid estimators: $\hat{V}_{i,K}(x_i)$ is a composition of an estimator based on polynomial approximation designed for small to moderate observations (less than $2\sqrt{2\log n}$ in absolute value) and the simple absolute value estimator applied to large observations (larger than $2\sqrt{2\log n}$ in absolute value), whereas $\hat{V}_{i,K}^S(x_i)$ has an additional thresholding procedure for small observations (less than $\sqrt{2\log n}$

in absolute value). Consequently, we propose two estimators of $T(\theta, \mu)$, namely,

$$\widehat{T_K(\theta, \mu)} = 2 \sum_{i=1}^n \widehat{V}_{i,K}(x_i) \widehat{V}_{i,K}(y_i), \quad (8)$$

as the hybrid non-thresholding estimator and

$$\widehat{T_K^S(\theta, \mu)} = 2 \sum_{i=1}^n \widehat{V}_{i,K}^S(x_i) \widehat{V}_{i,K}^S(y_i), \quad (9)$$

as the hybrid thresholding estimator. Both estimators rely on K , which is the tuning parameter to be specified later.

2.3 Theoretical properties and minimax optimality

In this section, we formally study the theoretical properties of $\widehat{T_K(\theta, \mu)}$ and $\widehat{T_K^S(\theta, \mu)}$. The following theorem provides the risk upper bounds and therefore establishes the minimax optimality of $\widehat{T_K(\theta, \mu)}$ and $\widehat{T_K^S(\theta, \mu)}$ over $D^\infty(s, L_n)$ with different sparsity levels.

Theorem 2. *Let $\mathbf{x}_n \sim N(\theta, \boldsymbol{\Sigma}_1)$ and $\mathbf{y}_n \sim N(\mu, \boldsymbol{\Sigma}_2)$ be multivariate Gaussian random vectors with $(\theta, \mu, \boldsymbol{\Sigma}_1, \boldsymbol{\Sigma}_2) \in D^\infty(s, L_n)$, $L_n \gtrsim \sqrt{\log n}$ and $s \asymp n^\beta$. Then*

1. *for any $\beta \in (0, 1)$, let K be any finite positive integer, we have,*

$$\sup_{(\theta, \mu, \boldsymbol{\Sigma}_1, \boldsymbol{\Sigma}_2) \in D^\infty(s, L_n)} \mathcal{R}(\widehat{T_K^S(\theta, \mu)}) \lesssim \frac{L_n^2 s^2 \log n}{n^2}; \quad (10)$$

2. *for any $\beta \in (1/2, 1)$, let $K = r \log n$ for some $0 < r < \frac{2\beta-1}{12}$, we have,*

$$\sup_{(\theta, \mu, \boldsymbol{\Sigma}_1, \boldsymbol{\Sigma}_2) \in D^\infty(s, L_n)} \mathcal{R}(\widehat{T_K(\theta, \mu)}) \lesssim \frac{L_n^2 s^2}{n^2 \log n}. \quad (11)$$

From the above theorem, if $0 < \beta < 1/2$, the risk upper bound (10) along with the minimax lower bound (3) implies

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \boldsymbol{\Sigma}_1, \boldsymbol{\Sigma}_2) \in D^\infty(s, L_n)} \mathcal{R}(\hat{T}) \asymp \frac{L_n^2 s^2 \log n}{n^2}, \quad (12)$$

where the optimal rate can be attained by the thresholding estimator $\widehat{T_K^S(\theta, \mu)}$ with K being any finite positive integer. If $1/2 < \beta < 1$, then the risk upper bound (11) along with the minimax

lower bound (4) implies

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)} \mathcal{R}(\hat{T}) \asymp \frac{L_n^2 s^2}{n^2 \log n}, \quad (13)$$

where the optimal rate can be attained by the non-thresholding estimator $\widehat{T_K(\theta, \mu)}$ with $K = r \log n$ for some small r . In both cases, the tuning parameter K plays an important role in controlling the bias-variance trade-off. An important consequence of our results concerns the *local adaptivity* of $\widehat{T_K(\theta, \mu)}$ and $\widehat{T_K^S(\theta, \mu)}$ with respect to s and L_n . Specifically, the estimator $\widehat{T_K(\theta, \mu)}$ with $K = r \log n$ for some $0 < r < \frac{2\beta-1}{12}$ is simultaneously rate-optimal for any L_n and any $\beta' \in (1/2, 1)$ with $\beta' > \beta$, whereas the estimator $\widehat{T_K(\theta, \mu)}$ with K being any finite positive integer is simultaneously rate-optimal for any L_n and $\beta \in (0, 1/2)$.

Unfortunately, even with an appropriate choice of K , neither $\widehat{T_K^S(\theta, \mu)}$ nor $\widehat{T_K(\theta, \mu)}$ is simultaneously optimal across all $\beta \in (0, 1)$. However, since the difference in the optimal rates of convergence between (12) and (13) is only of a factor of $\log n$, we argue that in practice, even when $\beta \in (1/2, 1)$, the nearly optimal thresholding estimator $\widehat{T_K^S(\theta, \mu)}$ would perform just as well as the non-thresholding minimax optimal estimator $\widehat{T_K(\theta, \mu)}$. Moreover, as for the optimal choice of K , since the difference between $K \asymp \log n$ and $K \asymp 1$ is minor for most applications with small to moderately large sample sizes (e.g., with $n \sim 10^5$, we have $\log n \approx 11.51$), in practice it suffices to choose K as some fixed moderate constant.

2.4 Sparse estimation via simple thresholding

According to our previous analysis, if the parameter space is very sparse, i.e., $0 < \beta < 1/2$, the proposed estimator $\widehat{T_K^S(\theta, \mu)}$ is minimax optimal if we choose K as any constant positive integer. In other words, any constant degree polynomial approximation suffices for the optimal estimation of $T(\theta, \mu)$, including the constant function. It means that in this case the polynomial approximation is essentially redundant for our purpose.

In light of the above observation, we consider the following simple thresholding estimator

$$\widetilde{T(\theta, \mu)} = 2 \sum_{i=1}^n \hat{U}_i(x_i) \hat{U}_i(y_i), \quad \text{where} \quad \hat{U}_i(x_i) = |x_{i1}| I(|x_{i2}| > 2\sqrt{2 \log n}).$$

In the following, we show that $\widetilde{T(\theta, \mu)}$ has the same rates of convergence as $\widehat{T_K^S(\theta, \mu)}$ over the parameter spaces $D^\infty(n^\beta, L_n)$, and, in light of Theorem 1, is minimax optimal and adaptive over any sparsity level $\beta \in (0, 1/2)$.

Theorem 3. *Let $\mathbf{x}_n \sim N(\theta, \Sigma_1)$ and $\mathbf{y}_n \sim N(\mu, \Sigma_2)$ be multivariate Gaussian random vectors*

with $(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)$ and $L_n \gtrsim \sqrt{\log n}$. Then

$$\sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)} \mathcal{R}(\widetilde{T(\theta, \mu)}) \lesssim \frac{L_n^2 s^2 \log n}{n^2}. \quad (14)$$

Since our simple thresholding estimator $\widetilde{T(\theta, \mu)}$ completely drops the polynomial components in $T_K^S(\theta, \mu)$, its variance is significantly reduced. As a consequence, we find that as long as $\max(\|\theta\|_\infty, \|\mu\|_\infty) \leq \sqrt{n}$, the isotopic Gaussian assumption $\Sigma_1 = \Sigma_2 = \mathbf{I}_n$ can be removed without changing the rate of convergence. Towards this end, we define the enlarged parameter space

$$D_0^\infty(s, L_n) = \left\{ (\theta, \mu, \Sigma_1, \Sigma_2) : \begin{array}{l} (\theta, \mu) \in D(s), \max(\|\theta\|_\infty, \|\mu\|_\infty) \leq L_n, \\ \Sigma_1, \Sigma_2 \succeq 0, \Sigma_1 \text{ and } \Sigma_2 \text{ have unit diagonals.} \end{array} \right\}.$$

In particular, as Σ_1 and Σ_2 have unit diagonals, the sample splitting procedure (Section 2.1) still applies, which only leads to a 1/2-scaling of the off-diagonal entries of the covariance matrices.

Theorem 4. *Let $\mathbf{x}_n \sim N(\theta, \Sigma_1)$ and $\mathbf{y}_n \sim N(\mu, \Sigma_2)$ where $(\theta, \mu, \Sigma_1, \Sigma_2) \in D_0^\infty(s, \sqrt{n})$. Then we have*

$$\sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D_0^\infty(s, \sqrt{n})} \mathcal{R}(\widetilde{T(\theta, \mu)}) \lesssim \frac{s^2 \log^2 n}{n^2}. \quad (15)$$

By definition, we have $D^\infty(s, L_n) \subset D_0^\infty(s, L_n)$. It then follows from Theorem 1 that for any $s \lesssim \sqrt{n}$,

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D_0^\infty(s, \sqrt{n})} \mathcal{R}(\hat{T}) \asymp \frac{s^2 \log^2 n}{n^2}, \quad (16)$$

where the minimax optimal rate can be attained by $\widetilde{T(\theta, \mu)}$. This establishes the minimax optimality and adaptivity of $\widetilde{T(\theta, \mu)}$ over $D_0^\infty(n^\beta, \sqrt{n})$ for any $\beta \in (0, 1/2)$. The result confirms an important advantage of $\widetilde{T(\theta, \mu)}$ over $T_K^S(\theta, \mu)$, namely, its guaranteed theoretical performance over arbitrary correlation structures, which complies with the fact that in many applications the observations are not necessarily independent.

2.5 Normalization and linkage disequilibrium

Dealing with linkage disequilibrium (LD) among the SNPs (Reich et al., 2001; Daly et al., 2001; Pritchard and Przeworski, 2001) is essential in any genetic studies. In this paper, we follow the idea of Bulik-Sullivan et al. (2015) and propose to use the normalized T -score

$$\text{Normalized } T\text{-score}(g) = \frac{\sum_{i=1}^n |\theta_i^g \mu_i|}{\|\theta^g\|_2 \|\mu\|_2}$$

as a measure of genetic overlap between gene g and the outcome disease. In particular, the estimation of the ℓ_2 norms $\|\theta^g\|_2$ and $\|\mu\|_2$, or in our context, the SNP-heritability of the traits (Yang

et al., 2010), can be easily accomplished using summary statistics. As a result, every normalized T -score lies in between 0 and 1, which is scale-invariant (e.g., invariance to study sample sizes and SNP effect sizes) and comparable across many different genes or studies. In addition, as similarly argued in Bulik-Sullivan et al. (2015), the normalized T -score is less sensitive to the choice of the n -SNP sets used in the analysis.

Moreover, in Theorem 4, we show that the simple thresholding estimator $\widetilde{T}(\theta, \mu)$ does not require the independence of the z -scores, which theoretically guarantees its applicability in the presence of arbitrary LD structure among the SNPs. However, our theoretical results concerning $\widehat{T}_K(\theta, \mu)$ and $\widehat{T}_K^S(\theta, \mu)$ rely on such an independence assumption. In our simulation studies, we found that the empirical performance (including optimality) of $\widehat{T}_K(\theta, \mu)$ and $\widehat{T}_K^S(\theta, \mu)$ is not likely affected by the dependence due to the LD structure. As a result, our proposed estimation method, although partially analysed under the independence assumption, can be directly applied to the summary statistics without specifying the underlying LD or covariance structure.

3 Simulation Studies

In this section, we demonstrate and compare the empirical performance of our proposed estimators and some alternative estimators under various settings. Since our goal is to estimate the T -score for each gene and to identify important genes that have strong association with disease, we generate data that mimic genetic association data, including block-wise and sparse signals. In particular, the block-wise correlated structure of the SNPs and therefore also the corresponding z -scores is widely observed in large-scale genetic studies due to LD among the SNPs. In the following, we consider the block-wise signals with both the standard independent errors or errors with local dependence to show the robustness of our proposed estimator.

We generate a pair of n -dimensional vectors, denoted as \mathbf{x}_n and \mathbf{y}_n , with $n = 1.5 \times 10^5, 3 \times 10^5$ and 5×10^5 , from a multivariate normal distribution $N(\theta, \Sigma)$ and $N(\mu, \Sigma)$, respectively. The mean vectors (θ, μ) , and the covariance matrix Σ are generated as follows:

- Block-wise signals: we first partition the n coordinates of θ and μ into blocks with each block consisting of 10 neighboring coordinates. For any sparsity level $s = 50, 100, 200, 300$ or 400 , we randomly pick $s/10$ blocks as the support. In each of these blocks, we assign coordinates of θ and μ with symmetric triangle-shaped values, whose maximal value is generated from a uniform distribution over $[3, 5]$. For the rest of the $(n - s)/10$ blocks, we set them as 0.
- Global covariance: $\Sigma = \mathbf{I}$.
- Block-wise covariances: Σ is block diagonal where each block is either a 10×10 Toeplitz matrix (see Supplementary Material for details) or a 1000×1000 exchangeable covariance matrix whose off-diagonal elements are 0.5. These two covariance matrices are denoted as Σ_1 and Σ_2 , respectively.

In particular, for the correlated cases, the correlation exists over either a small block of 10 elements whose correlation decays in distance, or over a large interval of 1000 elements with a constant correlation.

We compare the following five estimators of $T(\theta, \mu)$: (1) the hybrid thresholding estimator $\widehat{T_K^S(\theta, \mu)}$; (2) the hybrid thresholding estimator without sample splitting procedure, denoted as $\widehat{T_K^{S^*}(\theta, \mu)}$; (3) the proposed simple thresholding estimator $\widehat{T(\theta, \mu)}$; (4) the hybrid non-thresholding estimator $\widehat{T_K(\theta, \mu)}$; and (5) the estimator that simply calculates the absolute inner product of observed vectors, denoted as Naive.

For $\widehat{T_K^S(\theta, \mu)}$, $\widehat{T_K(\theta, \mu)}$ and $\widehat{T_K^{S^*}(\theta, \mu)}$, we choose $K = 8$ (see the last paragraph of Section 2.3). Each setting was repeated 100 times. We compare the estimators using the empirical version of the following rescaled mean square error:

$$\text{RMSE}(\hat{T}) = \frac{1}{s} \sqrt{\mathbb{E}(\hat{T} - T)^2}.$$

In particular, as the empirical RMSEs are subject to a factor of s , small numerical differences in RMSE actually indicate large differences in overall MSE. Tables 1-3 give the empirical RMSE of the five estimators under the settings with either independent or dependent observations. In general, $\widehat{T_K^{S^*}(\theta, \mu)}$ outperforms all the other estimators in all the settings. The performances of $\widehat{T_K^S(\theta, \mu)}$, $\widehat{T(\theta, \mu)}$ and $\widehat{T_K(\theta, \mu)}$ are roughly the same, with $\widehat{T_K^S(\theta, \mu)}$ and $\widehat{T(\theta, \mu)}$ being slightly better over the more sparse cases. The Naive estimator performs poorly in all the settings. The superiority of $\widehat{T_K^{S^*}(\theta, \mu)}$ over $\widehat{T_K^S(\theta, \mu)}$ might be explained by the fact that, by using the sample splitting procedure, we introduced extra variability to our estimator. Since the sample splitting is used only to facilitate our theoretical analysis, in applications we suggest to use $\widehat{T_K^{S^*}(\theta, \mu)}$ for its better performance.

As for the robustness of our estimators to the presence of correlated observations, we see that all of the proposed estimators have roughly the same performance as the independent cases. In real applications, we expect that the observed data are more likely to be “block-wise signal with block-wise correlation.” Our simulation shows that $\widehat{T_K^{S^*}(\theta, \mu)}$ provides the best estimate of the T-scores.

4 Integrative data analysis of human heart failure

Finally, we apply our proposed estimation procedure to identify genes whose expressions are possibly causally linked to heart failure by integrating GWAS and eQTL data. GWAS results were obtained from a heart failure genetic association study at the University of Pennsylvania, a prospective study of patients recruited from the University of Pennsylvania, Case Western Reserve University, and the University of Wisconsin, where genotype data were collected from 4,523 controls and 2,206 cases using the Illumina OmniExpress Plus array. GWAS summary statistics were calculated controlling

Table 1: Empirical RMSE under covariance $\Sigma = \mathbf{I}_n$. $T_K^{S^*}(\theta, \mu)$: the hybrid thresholding estimator without sample splitting procedure; $T_K^S(\theta, \mu)$: the hybrid thresholding estimator; $T(\theta, \mu)$: the proposed simple thresholding estimator; $T_K(\theta, \mu)$: the hybrid non-thresholding estimator; Naive: the estimator that simply calculates the absolute inner product of observed vectors.

n	s	$T_K^{S^*}(\theta, \mu)$	$T_K^S(\theta, \mu)$	$T(\theta, \mu)$	$T_K(\theta, \mu)$	Naive
150,000	100	4.091	5.818	5.764	5.847	954.4
	250	4.058	6.348	6.157	6.382	381.3
	400	3.949	6.274	6.042	238.4	6.307
	550	4.024	6.272	6.069	6.304	173.2
	700	3.973	5.84	5.702	5.870	135.9
300,000	100	4.271	6.579	6.385	6.613	1909.1
	250	4.222	6.244	6.118	6.276	763.3
	400	4.143	6.015	5.882	6.046	476.8
	550	4.214	5.991	5.874	6.022	346.6
	700	4.242	6.144	6.026	6.175	272.4
500,000	100	4.380	6.139	6.031	6.170	3182.7
	250	4.428	6.251	6.157	6.283	1272.6
	400	4.374	6.109	5.997	6.140	795.3
	550	4.395	6.149	6.056	6.180	578.3
	700	4.363	6.221	6.117	6.253	454.2

Table 2: Empirical RMSE under covariance Σ_1 .

n	s	$T_K^{S^*}(\theta, \mu)$	$T_K^S(\theta, \mu)$	$T(\theta, \mu)$	$T_K(\theta, \mu)$	Naive
150,000	100	4.090	6.704	6.502	7.042	953.5
	250	4.134	5.816	5.702	6.104	381.9
	400	4.068	6.087	5.896	6.396	237.9
	550	3.947	5.991	5.826	6.293	172.9
	700	4.163	6.011	5.913	6.310	135.8
300,000	100	3.868	6.088	6.009	6.403	1912.2
	250	4.196	5.963	5.797	6.262	764.7
	400	4.215	6.171	6.09	6.478	476.8
	550	4.238	6.005	5.888	6.306	345.7
	700	4.409	6.056	5.965	6.136	272.6
500,000	100	4.288	5.385	5.345	5.651	3184.8
	250	4.353	6.227	6.027	6.531	1273.0
	400	4.478	5.878	5.877	6.170	795.3
	550	4.370	5.929	5.855	6.222	577.7
	700	4.464	6.135	6.083	6.438	455.1

for age, gender, and the first two principal components of the genotypes.

Heart failure eQTL data were obtained from the MAGNet eQTL study (<https://www.med>).

Table 3: Empirical RMSE under covariance Σ_2 .

n	s	$\widehat{T_K^{S^*}(\theta, \mu)}$	$\widehat{T_K^S(\theta, \mu)}$	$\widehat{T(\theta, \mu)}$	$\widehat{T_K(\theta, \mu)}$	Naive
150,000	100	4.336	5.88	5.807	5.906	485.3
	250	4.645	5.857	5.828	5.885	189.7
	400	4.770	6.543	6.438	6.575	120.1
	550	4.566	6.035	5.985	6.065	87.12
	700	4.523	5.983	5.938	6.012	68.89
300,000	100	4.484	6.005	5.980	6.029	961.0
	250	4.718	5.86	5.865	5.887	380.6
	400	4.753	6.26	6.241	6.291	241.8
	550	4.729	5.985	5.949	6.015	174.8
	700	4.84	6.334	6.289	6.365	135.7
500,000	100	4.836	6.292	6.279	6.313	1593.5
	250	4.749	6.629	6.569	6.658	633.4
	400	4.889	6.409	6.401	6.438	398.1
	550	4.867	6.343	6.309	6.373	288.8
	700	4.887	6.131	6.112	6.159	227.7

upenn.edu/magnet/index.shtml), where the left ventricular free-wall tissue was collected from 136 donor hearts without heart failure. Genotype data were collected using Affymetrix genome-wide SNP array 6.0 and only markers in Hardy-Weinberg equilibrium with minor allele frequencies above 5% were considered. Gene expression data were collected using Affymetrix GeneChip ST1.1 arrays, normalized using RMA (Irizarry et al., 2003) and batch-corrected using ComBat (Johnson et al., 2007). To obtain a common set of SNPs, SNPs were imputed using 1000 Genomes Project data. Summary statistics for the MAGNet eQTL data were obtained using the fast marginal regression algorithm of Sikorska et al. (2013) controlling for age and gender.

4.1 Ranking of Potential Heart Failure Causal Genes

After matching the SNPs of both eQTL and GWAS data, we had a total of 347,019 SNPs and 19,081 genes with expression data available. For each gene, the vector of z -scores from eQTL analysis was obtained and the value of its absolute inner product with the vector of z -scores from GWAS data was estimated by our proposed estimator $\widehat{T_K^{S^*}(\theta, \mu)}$. We chose $K = 8$ as in our simulation study. After obtaining the estimates of T -scores for all the genes, and the corresponding SNP-heritability, we rank these genes in the order of their normalized T -scores as defined in Section 2.5. As a result, genes with the highest ranks are considered important in gaining insights into the biological mechanisms of heart failure.

To assess that the top scored genes indeed represent true biological signals, we calculated the T -scores for two “null datasets” that are created using permutations. For the first dataset, we

Table 4: Top eight heart failure associated genes based on the estimated normalized T -scores and their functional annotations.

Gene Name	Annotations
TMEM37	voltage-gated ion channel activity (Chen et al., 2007)
ADCY7	adenylate cyclase activity; fibrotic myocardial degeneration (Nojiri et al., 2006)
C1QC	Wnt signaling activity; associated with heart failure (Naito et al., 2012)
FAM98A	associated with ventricular septal defect (Liu et al., 2018)
BMP2	associated with heart-valve development (Rivera-Feliciano and Tabin, 2006)
SLCO2B1	organic anion transporter; associated with cardiac glycoside (Mikkaichi et al., 2004)
C1QA	Wnt signaling activity; associated with heart failure (Naito et al., 2012)
FCGR2B	intracellular signaling activity; associated with vascular disease pathogenesis (Tanigaki et al., 2015)

randomly permuted the labels of the SNPs of the GWAS z -scores by sampling without replacement before estimating the normalized T -scores with eQTL z -scores. For the second dataset, we permuted the labels of the SNPs of the GWAS z -scores in a circular manner similar to Cabrera et al. (2012). Specifically, for each chromosome, we randomly chose one SNP as start of the chromosome and move the SNPs on the fragment before this SNP to the end. Such a cyclic permutation preserves the local dependence of the z -scores. By permuting the data from one phenotype, we break the original matching of the z -scores between the two phenotypes. The permutation was performed 50 times and the null distribution of T -scores based on the permuted data was obtained. Figure 1 shows the ranked normalized T -scores based on the original data and the box plots of the ranked z -scores based on 50 permutations of the z -scores. We find that all the top ranked genes have larger T -scores than the ones based on permutations. In addition, about 30 top ranked genes in the top plot and about 10 top ranked genes in the bottom plot have true T -scores larger than all the T -scores from the permuted datasets. This further confirms that the top ranked genes based on their estimated normalized T -scores are not due to random noise and indeed represent certain sharing of genetic variants between heart failure and gene expression levels.

Table 4 lists the top eight highest ranked genes along with their biological annotations. All of the genes are either directly or indirectly associated with human heart failure, including those related to fibrotic myocardial degeneration, Wnt signalling activity and heart-valve development. It is interesting that our proposed methods can identify these relevant genes using only the gene expression data measured on normal heart tissues.




Figure 1: Estimated score for the top 50 genes and the box plots of the top scores based on 50 permutations. Top: random permutation of the GWAS scores; bottom: cyclic permutations of the GWAS scores.

4.2 Gene Set Enrichment Analysis

To complete our analysis, we finish this section with the gene set enrichment analysis (GSEA) (Subramanian et al., 2005) using the normalized T -scores in order to identify the heart failure associated biological processes. In the following analysis, we removed genes with low expression and small variability across the samples, which resulted in a total of 6,355 genes. For the normalized T -scores T_j , $1 \leq j \leq J$, given a gene set S , the Kolmogorov-Smirnov statistic defined as

$$\sup_t \left| \frac{1}{k} \sum_{j \in S} I(T_j \leq t) - \frac{1}{k'} \sum_{j \in S^c} I(T_j \leq t) \right|,$$

is calculated, where k and k' are the number of genes in S and S^c , respectively. This tests whether the distribution of T_j differs between genes in S and genes in S^c . For a given gene set, significance of this test implies that this gene set is enriched by genes that shared similar genetic variants as those for heart failure, suggesting their relevance to the etiology of heart failure.

Table 5: Top six GO biological processes that are associated with heart failure based on the gene set enrichment analysis

GO term	p-value
<i>Biological Process</i>	
regulation of skeletal muscle contraction by regulation of release of sequestered calcium ion	7.9×10^{-7}
linoleic acid metabolic process	1.0×10^{-6}
regulation of skeletal muscle contraction by calcium ion signaling	3.4×10^{-6}
positive regulation of sequestering of calcium ion	3.4×10^{-6}
cellular response to caffeine	1.0×10^{-5}
cellular response to purine-containing compound	1.0×10^{-5}

This analysis was applied to the gene sets from Gene Ontology (GO) (Botstein et al. 2000) that contain at least 10 genes. Specifically, 5,023 biological processes were tested. Figure 2 in our supplementary material presents the directed acyclic graphs of the GO biological processes that linked to the most significant GO terms from the simultaneous signal GSEA analysis. Table 5 shows the top six GO biological processes identified from the GSEA analysis. Among these processes, regulation of skeletal muscle contraction, linoleic acid metabolic process and calcium ion regulation are strongly implicated in human heart failure. Murphy et al. (2011) showed that skeletal muscle reflexes are essential to the initiation and regulation of the cardiovascular response to exercise, and alteration of this reflex mechanism can happen in disease states such as hypertension and heart failure. In Farvid et al. (2014), a systematic review of literature and meta-analysis was carried out, which supports a significant inverse association between dietary linoleic acid intake, when replacing either carbohydrates or saturated fat, and risk of coronary heart disease. Moreover, the importance of calcium-dependent signaling in the heart failure was reported in Marks (2003), who suggested that impaired calcium release causes decreased muscle contraction (systolic dysfunction) and defective calcium removal hampers relaxation (diastolic dysfunction).

5 Discussion

This paper considers the optimal estimation over sparse parameter spaces. However, in Section 2, the minimax rates of convergence were only established for parameter spaces $D^\infty(n^\beta, L_n)$ with $\beta \in (0, 1/2) \cup (1/2, 1)$. Apparently, there was a missing gap at $\beta = 1/2$. Our theoretical analysis suggests a lower bound (2) with the rate function $\psi(s, n) \asymp \log^{-1} n$. However, none of our proposed estimators attain such rates, leaving it an open question how to optimality estimate $T(\theta, \mu)$ when $s \asymp \sqrt{n}$.

In some applications, we may need to consider non-sparse parameter spaces. In this case, our theoretical analysis shows that the estimator $T_K(\widehat{\theta}, \widehat{\mu})$ with $K = r \log n$ for some small constant

$r > 0$ can still be applied. Specifically, from our proof of Theorem 1 and Theorem 2, it follows that, if we define the non-sparse parameter space as $\mathcal{D}_{\hat{U}}^{\infty}(L_n) = \{(\theta, \mu, \Sigma_1, \Sigma_2) : (\theta, \mu) \in \mathbb{R}^n \times \mathbb{R}^n, \max(\|\theta\|_{\infty}, \|\mu\|_{\infty}) \leq L_n, \Sigma_1 = \Sigma_2 = \mathbf{I}_n\}$ with $L_n \gtrsim \sqrt{\log n}$, then for $\mathbf{x}_n \sim N(\theta, \Sigma_1)$ and $\mathbf{y}_n \sim N(\mu, \Sigma_2)$, the minimax optimal rate for estimating $T(\theta, \mu)$ over $\mathcal{D}_{\hat{U}}^{\infty}(L_n)$ is

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in \mathcal{D}_{\hat{U}}^{\infty}(L_n)} \mathcal{R}(\hat{T}) \asymp \frac{L_n}{\log n}, \quad (17)$$

which can be attained by the above $T_K(\widehat{\theta}, \widehat{\mu})$.

In addition to the estimation problem, it is also of interest to conduct hypothesis testing or construct confidence intervals for T -score. These problems can be technically challenging due to the non-smooth functional. Moreover, the polynomial approximation technique adopted in the current work may introduce extra complexities to the higher-order asymptotic analysis. We leave these problems for future investigations.

Supplementary Materials

Our supplementary material includes proofs of the main theorems and the technical lemmas.

References

- Acharya, J., H. Das, A. Orlytsky, and A. T. Suresh (2016). A unified maximum likelihood approach for optimal distribution property estimation. *arXiv preprint arXiv:1611.02960*.
- Botstein, D., J. M. Cherry, M. Ashburner, C. Ball, J. Blake, H. Butler, A. Davis, K. Dolinski, S. Dwight, J. Eppig, et al. (2000). Gene ontology: tool for the unification of biology. *Nature Genetics* 25(1), 25–9.
- Brown, L. D. and M. G. Low (1996). A constrained risk inequality with applications to nonparametric functional estimation. *The Annals of Statistics* 24(6), 2524–2535.
- Bulik-Sullivan, B., H. K. Finucane, V. Anttila, A. Gusev, F. R. Day, P.-R. Loh, L. Duncan, J. R. Perry, N. Patterson, E. B. Robinson, et al. (2015). An atlas of genetic correlations across human diseases and traits. *Nature Genetics* 47(11), 1236.
- Cabrera, C. P., P. Navarro, J. E. Huffman, A. F. Wright, C. Hayward, H. Campbell, J. F. Wilson, I. Rudan, N. D. Hastie, V. Vitart, et al. (2012). Uncovering networks from genome-wide association studies via circular genomic permutation. *G3: Genes, Genomes, Genetics* 2(9), 1067–1075.
- Cai, T. T. and M. G. Low (2006). Optimal adaptive estimation of a quadratic functional. *The Annals of Statistics* 34(5), 2298–2325.

- Cai, T. T. and M. G. Low (2011). Testing composite hypotheses, hermite polynomials and optimal estimation of a nonsmooth functional. *The Annals of Statistics* 39(2), 1012–1041.
- Chen, R.-S., T.-C. Deng, T. Garcia, Z. M. Sellers, and P. M. Best (2007). Calcium channel γ subunits: a functionally diverse protein family. *Cell Biochemistry and Biophysics* 47(2), 178–186.
- Daly, M. J., J. D. Rioux, S. F. Schaffner, T. J. Hudson, and E. S. Lander (2001). High-resolution haplotype structure in the human genome. *Nature Genetics* 29(2), 229–232.
- Donoho, D. L. and M. Nussbaum (1990). Minimax quadratic estimation of a quadratic functional. *Journal of Complexity* 6(3), 290–323.
- Efromovich, S. and M. G. Low (1994). Adaptive estimates of linear functionals. *Probability Theory and Related Fields* 98(2), 261–275.
- Fan, J. (1991). On the estimation of quadratic functionals. *The Annals of Statistics*, 1273–1294.
- Farvid, M. S., M. Ding, A. Pan, Q. Sun, S. E. Chiuve, L. M. Steffen, W. C. Willett, and F. B. Hu (2014). Dietary linoleic acid and risk of coronary heart disease: a systematic review and meta-analysis of prospective cohort studies. *Circulation* 130(18), 1568–1578.
- Gamazon, E. R., H. E. Wheeler, K. P. Shah, S. V. Mozaffari, K. Aquino-Michaels, R. J. Carroll, A. E. Eyler, J. C. Denny, D. L. Nicolae, N. J. Cox, et al. (2015). A gene-based association method for mapping traits using reference transcriptome data. *Nature Genetics* 47(9), 1091.
- Giallourakis, C., C. Henson, M. Reich, X. Xie, and V. K. Mootha (2005). Disease gene discovery through integrative genomics. *Annu. Rev. Genomics Hum. Genet.* 6, 381–406.
- Giambartolomei, C., D. Vukcevic, E. E. Schadt, L. Franke, A. D. Hingorani, C. Wallace, and V. Plagnol (2014). Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genetics* 10(5), e1004383.
- He, X., C. K. Fuller, Y. Song, Q. Meng, B. Zhang, X. Yang, and H. Li (2013). Sherlock: detecting gene-disease associations by matching patterns of expression QTL and GWAS. *The American Journal of Human Genetics* 92(5), 667–680.
- Hindorf, L. A., P. Sethupathy, H. A. Junkins, E. M. Ramos, J. P. Mehta, F. S. Collins, and T. A. Manolio (2009). Potential etiologic and functional implications of genome-wide association loci for human diseases and traits. *Proceedings of the National Academy of Sciences* 106(23), 9362–9367.
- Ibragimov, I. A. and R. Z. Khas'minskii (1985). On nonparametric estimation of the value of a linear functional in gaussian white noise. *Theory of Probability & Its Applications* 29(1), 18–32.

- Irizarry, R. A., B. Hobbs, F. Collin, Y. D. Beazer-Barclay, K. J. Antonellis, U. Scherf, and T. P. Speed (2003). Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4(2), 249–264.
- Jiao, J., Y. Han, and T. Weissman (2016). Minimax estimation of the l_1 distance. In *Information Theory (ISIT), 2016 IEEE International Symposium on*, pp. 750–754. IEEE.
- Jiao, J., K. Venkat, Y. Han, and T. Weissman (2015). Minimax estimation of functionals of discrete distributions. *IEEE Transactions on Information Theory* 61(5), 2835–2885.
- Johnson, W. E., C. Li, and A. Rabinovic (2007). Adjusting batch effects in microarray expression data using empirical bayes methods. *Biostatistics* 8(1), 118–127.
- Liu, H., Y. Hu, B. Zhuang, J. Yin, X.-H. Chen, J. Wang, M.-M. Li, J. Xu, X.-Y. Wang, Z.-B. Yu, et al. (2018). Differential expression of circrnas in embryonic heart tissue associated with ventricular septal defect. *International Journal of Medical Sciences* 15(7), 703.
- Marks, A. R. (2003). Calcium and the heart: a question of life and death. *The Journal of Clinical Investigation* 111(5), 597–600.
- Mikkaichi, T., T. Suzuki, M. Tanemoto, S. Ito, and T. Abe (2004). The organic anion transporter (oatp) family. *Drug Metabolism and Pharmacokinetics* 19(3), 171–179.
- Murphy, M. N., M. Mizuno, J. H. Mitchell, and S. A. Smith (2011). Cardiovascular regulation by skeletal muscle reflexes in health and disease. *American Journal of Physiology-Heart and Circulatory Physiology*.
- Naito, A. T., T. Sumida, S. Nomura, M.-L. Liu, T. Higo, A. Nakagawa, K. Okada, T. Sakai, A. Hashimoto, Y. Hara, et al. (2012). Complement c1q activates canonical wnt signaling and promotes aging-related phenotypes. *Cell* 149(6), 1298–1313.
- Nica, A. C., S. B. Montgomery, A. S. Dimas, B. E. Stranger, C. Beazley, I. Barroso, and E. T. Dermitzakis (2010). Candidate causal regulatory effects by integration of expression QTLs with complex trait genetic associations. *PLoS Genetics* 6(4), e1000895.
- Nicolae, D. L., E. Gamazon, W. Zhang, S. Duan, M. E. Dolan, and N. J. Cox (2010). Trait-associated SNPs are more likely to be eQTLs: annotation to enhance discovery from GWAS. *PLoS Genetics* 6(4), e1000888.
- Nojiri, H., T. Shimizu, M. Funakoshi, O. Yamaguchi, H. Zhou, S. Kawakami, Y. Ohta, M. Sami, T. Tachibana, H. Ishikawa, et al. (2006). Oxidative stress causes heart failure with impaired mitochondrial respiration. *Journal of Biological Chemistry* 281(44), 33789–33801.

- Pritchard, J. K. and M. Przeworski (2001). Linkage disequilibrium in humans: models and data. *The American Journal of Human Genetics* 69(1), 1–14.
- Reich, D. E., M. Cargill, S. Bolk, and J. Ireland (2001). Linkage disequilibrium in the human genome. *Nature* 411(6834), 199.
- Rivera-Feliciano, J. and C. J. Tabin (2006). Bmp2 instructs cardiac progenitors to form the heart-valve-inducing field. *Developmental Biology* 295(2), 580–588.
- Sikorska, K., E. Lesaffre, P. F. Groenen, and P. H. Eilers (2013). GWAS on your notebook: fast semi-parallel linear and logistic regression for genome-wide association studies. *BMC bioinformatics* 14(1), 166.
- Subramanian, A., P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette, A. Paulovich, S. L. Pomeroy, T. R. Golub, E. S. Lander, et al. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences* 102(43), 15545–15550.
- Tanigaki, K., N. Sundgren, A. Khera, W. Vongpatanasin, C. Mineo, and P. W. Shaul (2015). Fc γ receptors and ligands and cardiovascular disease. *Circulation Research* 116(2), 368–384.
- Wu, Y. and P. Yang (2016). Minimax rates of entropy estimation on large alphabets via best polynomial approximation. *IEEE Transactions on Information Theory* 62(6), 3702–3720.
- Wu, Y. and P. Yang (2019). Chebyshev polynomials, moment matching, and optimal estimation of the unseen. *The Annals of Statistics* 47(2), 857–883.
- Yang, J., B. Benyamin, B. P. McEvoy, S. Gordon, A. K. Henders, D. R. Nyholt, P. A. Madden, A. C. Heath, N. G. Martin, G. W. Montgomery, et al. (2010). Common SNPs explain a large proportion of the heritability for human height. *Nature Genetics* 42(7), 565.
- Zhao, S. D., T. T. Cai, T. P. Cappola, K. B. Margulies, and H. Li (2017). Sparse simultaneous signal detection for identifying genetically controlled disease genes. *Journal of the American Statistical Association* 112(519), 1032–1046.
- Zhu, Z., F. Zhang, H. Hu, A. Bakshi, M. R. Robinson, J. E. Powell, G. W. Montgomery, M. E. Goddard, N. R. Wray, P. M. Visscher, et al. (2016). Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nature Genetics* 48(5), 481.

Supplement to “Optimal Estimation of Simultaneous Signals Using Absolute Inner Product with Applications to Integrative Genomics”

Rong Ma¹, T. Tony Cai² and Hongzhe Li¹

Department of Biostatistics, Epidemiology and Informatics¹

Department of Statistics²

University of Pennsylvania

Philadelphia, PA 19104

Abstract

This supplementary material includes proofs of the main theorems and the technical lemmas of the paper “Optimal Estimation of Simultaneous Signals Using Absolute Inner Product with Applications to Integrative Genomics.” Some supplementary figures and tables are also included.

1 Proofs of the Minimax Lower Bounds

1.1 Proof of Theorem 1

By the definition of the rate function $\psi(s, n)$, it suffices to show the following two statements.

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)} \mathbb{E}(\hat{T} - T(\theta, \mu))^2 \gtrsim L_n^2 s^2 \log \left(1 + \frac{n}{s^2} \right), \quad (18)$$

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)} \mathbb{E}(\hat{T} - T(\theta, \mu))^2 \gtrsim \frac{L_n^2 s^2}{\log s}. \quad (19)$$

Proof of (18). Let $l(s, n)$ be the class of all subsets of $\{1, \dots, n\}$ of s elements. For $I \in l(s, n)$, we denote $\theta_I = \{\theta \in \mathbb{R}^n : \theta_i = 0, \forall i \notin I, \text{ and } \theta_i = \rho, \forall i \in I\}$. Suppose that μ is fixed with $\mu = \mu^*$ where $\mu_i^* = L_n$ for all $1 \leq i \leq n$. Denote

$$g_I(x_1, \dots, x_n, y_1, \dots, y_n) = \prod_{i=1}^n \psi_{\theta_{I,i}}(x_i) \prod_{i=1}^n \psi_{\mu_i^*}(y_i),$$

where ψ_{θ_i} denotes the density function of $N(\theta_i, 1)$ and $\theta_{I,i}$ is the i -th component of θ_I . In this way, we are considering the class of probability measures for $\{x_i, y_i\}_{1 \leq i \leq n}$ where the mean vector

for $\{x_i\}$ is the s -sparse vector θ_I whereas the mean vector for $\{y_i\}$ is μ^* . By averaging over all the possible $I \in l(s, n)$, we have the mixture probability measure

$$g = \frac{1}{\binom{n}{s}} \sum_{I \in l(s, n)} g_I.$$

On the other hand, we consider the probability measure

$$f = \prod_{i=1}^n \phi(x_i) \prod_{i=1}^n \psi_{\mu^*}(y_i)$$

where ϕ is the normal density of $N(0, 1)$. From the above construction, we consider $D_s^\theta(\rho) = \{(\theta, \mu, \Sigma_1, \Sigma_2) : \theta = \theta_I, I \in l(s, n), \mu = \mu^*, \Sigma_1 = \Sigma_2 = I\} \cup \{(\theta, \mu, \Sigma_1, \Sigma_2) : \theta = 0, \mu = \mu^*, \Sigma_1 = \Sigma_2 = I\}$. Apparently, for $\rho \leq L_n$, $D_s^\theta(\rho) \subset D^\infty(s, L_n)$. In the following, we will consider the χ^2 -divergence between g and f and obtain the minimax lower bound over $D_s(\rho)$ using the constrained risk inequality of Brown and Low (1996). Note that

$$\int \frac{g^2}{f} = \frac{1}{\binom{n}{s}^2} \sum_{I \in l(s, n)} \sum_{I' \in l(s, n)} \int \frac{g_I g_{I'}}{f}$$

and for any I and I' ,

$$\begin{aligned} \int \frac{g_I g_{I'}}{f} &= \frac{1}{(2\pi)^{n/2}} \int \exp \left\{ - \frac{\sum_i (x_i - \theta_{I,i})^2 + \sum_i (x_i - \theta_{I',i})^2 - \sum_i x_i^2}{2} \right\} \\ &\quad \times \frac{1}{(2\pi)^{n/2}} \int \exp \left\{ - \frac{\sum_i (y_i - \mu_i^*)^2}{2} \right\} \\ &= \frac{1}{(2\pi)^{n/2}} \int \exp \left\{ - \frac{\sum_i (x_i - \theta_{I,i} - \theta_{I',i})^2 - 2 \sum_{i=1}^n \theta_{I,i} \theta_{I',i}}{2} \right\} \\ &= \exp(\rho^2 j) \end{aligned}$$

where j is the number of points in the set $I \cap I'$. Following the similar arguments, we have, by taking $\rho = \sqrt{\log(1 + \frac{n}{s^2})}$,

$$\int \frac{g^2}{f} = \mathbb{E} \exp(J\rho^2) \leq e. \quad (20)$$

Then if some estimator δ satisfies

$$\mathbb{E}_f(\delta - 0)^2 \leq Cs^2 \|\mu^*\|_\infty^2 \log \left(1 + \frac{n}{s^2} \right) \quad (21)$$

then by the constrained risk inequality

$$\begin{aligned}\mathbb{E}_g\left(\delta - s\rho\|\mu^*\|_\infty\right)^2 &\geq s^2\rho^2\|\mu^*\|_\infty^2 - 2\rho s\|\mu^*\|_\infty C^{1/2}s\|\mu^*\|_\infty \log^{1/2}\left(1 + \frac{n}{s^2}\right) \\ &= s^2\|\mu^*\|_\infty^2 \log\left(1 + \frac{n}{s^2}\right) - \sqrt{2C}s^2\|\mu^*\|_\infty^2 \log\left(1 + \frac{n}{s^2}\right),\end{aligned}$$

for any such estimator δ . Recall that $\|\mu^*\|_\infty = L_n$. By choosing C sufficiently small, we conclude that there exists some $I \in l(s, n)$ such that

$$\mathbb{E}_{g_I}\left(\delta - s\rho\|\mu^*\|_\infty\right)^2 \geq Cs^2L_n^2 \log\left(1 + \frac{n}{s^2}\right) \quad (22)$$

for all δ . Therefore we have

$$\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D_s^\theta(\rho)} \mathbb{E}(\hat{T} - T(\theta, \mu))^2 \geq Cs^2L_n^2 \log\left(1 + \frac{n}{s^2}\right). \quad (23)$$

The lower bound (18) then follows from the fact that $L_n \gtrsim \sqrt{\log n}$.

Proof of (19). Now we prove the second part of the theorem. It follows by Lemma 1 in Cai and Low (2011) that there exist measures ν_i on $[-M_n, M_n]$ for $i = 0, 1$, such that:

1. ν_0 and ν_1 are symmetric around 0;
2. $\int t^l \nu_1(dt) = \int t^l \nu_0(dt)$, for $l = 0, 1, \dots, k_n$;
3. $\int |t| \nu_1(dt) - \int |t| \nu_0(dt) = 2M_n \delta_{k_n}$.
4. $\int |t| \nu_0(dt) > 0$.

where δ_{k_n} is the distance in the uniform norm on $[-1, 1]$ from the absolute value function $f(x) = |x|$ to the space of polynomials of no more than degree k_n . In addition, $\delta_{k_n} = \beta_* k_n^{-1}(1 + o(1))$ as $k_n \rightarrow \infty$. Now we consider product priors on the n -vector θ , which are supported on the first $s \leq n$ components. Let

$$\nu_{i1}^n = \Pi^{\otimes s} \nu_i \cdot \Pi^{\otimes n-s} 1_{\{0\}}, \quad \nu_{i2}^n = \Pi^{\otimes n} 1_{\{\mu^*\}}$$

for $i = 0$ and 1 . In other words, we put independent priors ν_i for the first s components of the vector θ , while keeping the other coordinates as 0, and we fix $\mu = \mu^*$.

Following the above construction, we have

$$\mathbb{E}_{\nu_{11}^n} \frac{1}{n} \sum_{i=1}^n |\theta_i| - \mathbb{E}_{\nu_{01}^n} \frac{1}{n} \sum_{i=1}^n |\theta_i| = \frac{s}{n} \left[\mathbb{E}_{\nu_{11}} |\theta_i| - \mathbb{E}_{\nu_{01}} |\theta_i| \right] = 2sM\delta_{k_n}/n,$$

and

$$\mathbb{E}_{\nu_{12}^n} \frac{1}{n} \sum_{i=1}^n |\mu_i| - \mathbb{E}_{\nu_{02}^n} \frac{1}{n} \sum_{i=1}^n |\mu_i| = 0$$

Then we have

$$\begin{aligned} \mathbb{E}_{\nu_{11}^n \nu_{12}^n} \frac{1}{n} T(\theta, \mu) - \mathbb{E}_{\nu_{01}^n \nu_{02}^n} \frac{1}{n} T(\theta, \mu) &= \frac{s}{n} (\mathbb{E}_{\nu_{11}} |\theta_i| \mathbb{E}_{\nu_{12}} |\mu_i| - \mathbb{E}_{\nu_{01}} |\theta_i| \mathbb{E}_{\nu_{02}} |\mu_i|) \\ &= \frac{s}{n} (\mathbb{E}_{\nu_{11}} |\theta_i| \mathbb{E}_{\nu_{12}} |\mu_i| - \mathbb{E}_{\nu_{01}} |\theta_i| \mathbb{E}_{\nu_{12}} |\mu_i|) \\ &\quad + \frac{s}{n} (\mathbb{E}_{\nu_{01}} |\theta_i| \mathbb{E}_{\nu_{12}} |\mu_i| - \mathbb{E}_{\nu_{01}} |\theta_i| \mathbb{E}_{\nu_{02}} |\mu_i|) \\ &= \mathbb{E}_{\nu_{12}} |\mu_i| \frac{s}{n} (\mathbb{E}_{\nu_{11}} |\theta_i| - \mathbb{E}_{\nu_{01}} |\theta_i|) + \mathbb{E}_{\nu_{01}} |\theta_i| \frac{s}{n} (\mathbb{E}_{\nu_{12}} |\mu_i| - \mathbb{E}_{\nu_{02}} |\mu_i|) \\ &= \frac{2sM_n \delta_{k_n}}{n} \mathbb{E}_{\nu_{12}} |\mu_i| \\ &= \frac{2sM_n L_n \delta_{k_n}^2}{n} \end{aligned} \tag{24}$$

We further have

$$V_0^2 \equiv \frac{1}{n^2} \mathbb{E}_{\nu_{01}^n \nu_{02}^n} (T(\theta, \mu) - \mathbb{E}_{\nu_{01}^n \nu_{02}^n} T(\theta, \mu))^2 \leq \frac{sM_n^2 L_n^2}{n^2} \tag{25}$$

Set $f_{0,M_n}(y) = \int \phi(y-t) \nu_0(dt)$ and $f_{1,M_n}(y) = \int \phi(y-t) \nu_1(dt)$. Note that since $g(x) = \exp(-x)$ is a convex function of x , and ν_0 is symmetric,

$$\begin{aligned} f_{0,M_n}(y) &\geq \frac{1}{\sqrt{2\pi}} \exp\left(-\int \frac{(y-t)^2}{2} \nu_0(dt)\right) \\ &= \phi(y) \exp\left(-\frac{1}{2} M_n^2 \int t^2 \nu_0'(dt)\right) \\ &\geq \phi(y) \exp\left(-\frac{1}{2} M_n^2\right). \end{aligned}$$

Let H_r be the Hermite polynomial defined by

$$\frac{d^r}{dy^r} \phi(y) = (-1)^r H_r(y) \phi(y) \tag{26}$$

which satisfy

$$\int H_r^2(y) \phi(y) dy = r! \quad \text{and} \quad \int H_r(y) H_l(y) \phi(y) dy = 0 \tag{27}$$

when $r \neq l$. Then

$$\phi(y-t) = \sum_{k=0}^{\infty} H_k(y) \phi(y) \frac{t^k}{k!}$$

and it follows that

$$\begin{aligned}
\int \frac{(f_{1,M_n}(y) - f_{0,M_n}(y))^2}{f_{0,M_n}(y)} dy &\leq \int (f_{1,M_n}(y) - f_{0,M_n}(y))^2 e^{M_n^2/2} / \phi(y) dy \\
&= e^{M_n^2/2} \int \left\{ \sum_{k=0}^{\infty} H_k(y) \frac{\phi(y)}{k!} \left[\int t^k \nu_1(dt) - \int t^k \nu_0(dt) \right] \right\}^2 / \phi(y) dy \\
&= e^{M_n^2/2} \int \sum_{k=k_n+1}^{\infty} H_k^2(y) \phi(y) \frac{M_n^{2k}}{(k!)^2} \left[\int t^k \nu_1(dt) - \int t^k \nu_0(dt) \right]^2 dy \\
&= e^{M_n^2/2} \sum_{k=k_n+1}^{\infty} \frac{M_n^{2k}}{k!} \left[\int t^k \nu_1(dt) - \int t^k \nu_0(dt) \right]^2 \\
&\leq e^{M_n^2/2} \sum_{k=k_n+1}^{\infty} \frac{M_n^{2k}}{k!}.
\end{aligned}$$

It then follows

$$I_n^2 = \prod_{i=1}^s \int \frac{(f_{1,M_n}(x_i))^2}{f_{0,M_n}(x_i)} dx_i - 1 \leq \left(1 + e^{M_n^2/2} \sum_{k=k_n+1}^{\infty} \frac{1}{k!} M_n^{2k} \right)^s - 1 \leq \left(1 + e^{M_n^2/2} D \frac{1}{k_n!} M_n^{2k_n} \right)^s - 1,$$

for some $D > 0$. Since $k! > (k/e)^k$, we also have

$$I_n^2 \leq \left(1 + e^{M_n^2/2} D \left(\frac{eM_n^2}{k_n} \right)^{k_n} \right)^s - 1 \quad (28)$$

Now let $M_n \asymp \sqrt{\log s}$ and $k_n \asymp \log s$. It then can be checked that $I_n < c$ for some constant $c > 0$. Therefore, Corollary 1 in Cai and Low (2011) along with the fact that $L_n \gtrsim \sqrt{\log n}$ yields

$$\begin{aligned}
\inf_{\hat{T}} \sup_{(\theta, \mu, \Sigma_1, \Sigma_2) \in D^\infty(s, L_n)} \frac{1}{n^2} \mathbb{E}(\hat{T} - T(\theta, \mu))^2 &\geq \frac{(2sM_n \delta_{k_n} L_n/n - (s^{1/2} M_n L_n/n) I_n)^2}{(I_n + 2)^2} \\
&\geq \frac{Cs^2 L_n^2}{n^2 \log s}.
\end{aligned} \quad (29)$$

2 Proofs of the Risk Upper Bounds

2.1 Proof of Theorem 2

For the hybrid estimators defined in Section 2.2 of the main paper, the key is to study the bias and variance of a single component. Let $x_1, x_2 \sim N(\theta, 1)$, $y_1, y_2 \sim N(\mu, 1)$. Denote $\delta(x) = \min\{S_K(x), n^2\}$. In the following, we analyse the two hybrid estimators separately.

Part I: Analysis of $T_K^S(\widehat{\theta}, \mu)$. Let

$$\begin{aligned} \xi = \xi(x_1, x_2, y_1, y_2) &= [\delta(x_1)I(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + |x_1|I(|x_2| > 2\sqrt{2\log n})] \\ &\quad \times [\delta(y_1)I(\sqrt{2\log n} < |y_2| \leq 2\sqrt{2\log n}) + |y_1|I(|y_2| > 2\sqrt{2\log n})] \end{aligned} \quad (30)$$

Note that

$$\begin{aligned} \mathbb{E}(\xi) &= [\mathbb{E}\delta(x_1)P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n})] \\ &\quad \times [\mathbb{E}\delta(y_1)P(\sqrt{2\log n} < |y_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|y_1|P(|y_2| > 2\sqrt{2\log n})] \end{aligned} \quad (31)$$

We denote

$$\tilde{\sigma}_x^2 = \text{Var}(\delta(x_1)I(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + |x_1|I(|x_2| > 2\sqrt{2\log n})),$$

$$\tilde{\sigma}_y^2 = \text{Var}(\delta(y_1)I(\sqrt{2\log n} < |y_2| \leq 2\sqrt{2\log n}) + |y_1|I(|y_2| > 2\sqrt{2\log n})),$$

and

$$\tilde{\theta}_x = \mathbb{E}\delta(x_1)P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}),$$

$$\tilde{\mu}_y = \mathbb{E}\delta(y_1)P(\sqrt{2\log n} < |y_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|y_1|P(|y_2| > 2\sqrt{2\log n}).$$

Then we have

$$\text{Var}(\xi) = \tilde{\sigma}_x^2\tilde{\sigma}_y^2 + \tilde{\sigma}_x^2\tilde{\mu}_y^2 + \tilde{\sigma}_y^2\tilde{\theta}_x^2$$

The following proposition is key to our calculation of the estimation risk.

Proposition 1. *For all $\theta, \mu \in \mathbb{R}$, we have*

$$|B_x| \equiv |\tilde{\theta}_x - |\theta|| \lesssim \sqrt{\log n}, \quad |B_y| \equiv |\tilde{\mu}_y - |\mu|| \lesssim \sqrt{\log n},$$

and

$$\tilde{\sigma}_x^2 \lesssim \log n, \quad \tilde{\sigma}_y^2 \lesssim \log n.$$

In particular, when $\theta = 0$, we have $|B_x| \leq n^{-2}\log n$ and $\tilde{\sigma}_x^2 \lesssim n^{-1}\log n$, whereas when $\mu = 0$, we have $|B_y| \leq n^{-2}\log n$, and $\tilde{\sigma}_y^2 \lesssim n^{-1}\log n$.

Now the bias of the estimator ξ satisfies

$$\begin{aligned} B(\xi) &= \mathbb{E}\xi - |\theta||\mu| = \tilde{\theta}_x\tilde{\mu}_y - |\theta||\mu| \\ &\leq |\theta| \cdot |\tilde{\mu}_y - |\mu|| + |\mu| \cdot |\tilde{\theta}_x - |\theta|| + |\tilde{\mu}_y - |\mu|| \cdot |\tilde{\theta}_x - |\theta|| \\ &\leq |\mu||B_y| + |\theta||B_x| + |B_x||B_y|. \end{aligned} \quad (32)$$

and the variance

$$\begin{aligned}\text{Var}(\xi) &= \tilde{\sigma}_x^2 \tilde{\sigma}_y^2 + \tilde{\sigma}_x^2 \tilde{\mu}_y^2 + \tilde{\sigma}_y^2 \tilde{\theta}_x^2 \\ &\lesssim \tilde{\sigma}_x^2 \tilde{\sigma}_y^2 + |\mu|^2 \tilde{\sigma}_x^2 + |\theta|^2 \tilde{\sigma}_y^2 + B_x^2 \tilde{\sigma}_y^2 + B_y^2 \tilde{\sigma}_x^2.\end{aligned}\quad (33)$$

Now let $(x_{1\ell}, x_{2\ell}, \dots, x_{n\ell}) \sim N(\theta, \mathbf{I}_n)$ and $(y_{1\ell}, y_{2\ell}, \dots, y_{n\ell}) \sim N(\mu, \mathbf{I}_n)$ for $\ell = 1, 2$, and let

$$\widehat{T^S(\theta, \mu)} = \sum_{i=1}^n \xi(x_{i1}, x_{i2}, y_{i1}, y_{i2}).$$

It follows from (32) and Cauchy-Schwartz inequality that the bias of $\widehat{T^S(\theta, \mu)}$ is bounded by

$$|B(\widehat{T^S(\theta, \mu)})| \lesssim (\|\theta\|_\infty + \|\mu\|_\infty) s \sqrt{\log n} + s \log n$$

From (33) we have the variance of $\widehat{T^S(\theta, \mu)}$ is bounded by

$$\begin{aligned}\text{Var}(\widehat{T^S(\theta, \mu)}) &\leq \sum_{i=1}^n \text{Var}(\xi(x_{i1}, x_{i2}, y_{i1}, y_{i2})) \\ &\lesssim s \log^2 n + (\|\theta\|_\infty^2 + \|\mu\|_\infty^2) s \log n\end{aligned}\quad (34)$$

Therefore the mean squared error of $\widehat{T^S(\theta, \mu)}$ satisfies

$$\begin{aligned}\mathbb{E}(\widehat{T^S(\theta, \mu)} - T(\theta, \mu))^2 &\leq B^2(\widehat{T^S(\theta, \mu)}) + \text{Var}(\widehat{T^S(\theta, \mu)}) \\ &\lesssim s^2 \log n (\|\theta\|_\infty^2 + \|\mu\|_\infty^2 + \log n).\end{aligned}$$

The final result follows from the fact that $\max(\|\theta\|_\infty, \|\mu\|_\infty) \leq L_n$ and $L_n \gtrsim \log n$.

Part II: Analysis of $\widehat{T_K(\theta, \mu)}$. With a slight abuse of notation, we denote

$$\begin{aligned}\xi &= \xi(x_1, x_2, y_1, y_2) = [\delta(x_1)I(|x_2| \leq 2\sqrt{2\log n}) + |x_1|I(|x_2| > 2\sqrt{2\log n})] \\ &\quad \times [\delta(y_1)I(|y_2| \leq 2\sqrt{2\log n}) + |y_1|I(|y_2| > 2\sqrt{2\log n})]\end{aligned}\quad (35)$$

Note that

$$\begin{aligned}\mathbb{E}(\xi) &= [\mathbb{E}\delta(x_1)P(|x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n})] \\ &\quad \times [\mathbb{E}\delta(y_1)P(|y_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|y_1|P(|y_2| > 2\sqrt{2\log n})]\end{aligned}\quad (36)$$

We denote

$$\tilde{\sigma}_x^2 = \text{Var}(\delta(x_1)I(|x_2| \leq 2\sqrt{2\log n}) + |x_1|I(|x_2| > 2\sqrt{2\log n})),$$

$$\tilde{\sigma}_y^2 = \text{Var}(\delta(y_1)I(|y_2| \leq 2\sqrt{2\log n}) + |y_1|I(|y_2| > 2\sqrt{2\log n})),$$

and

$$\tilde{\theta}_x = \mathbb{E}\delta(x_1)P(|x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}),$$

$$\tilde{\mu}_y = \mathbb{E}\delta(y_1)P(|y_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|y_1|P(|y_2| > 2\sqrt{2\log n}).$$

Proposition 2. *Let $B_x = \tilde{\theta}_x - |\theta|$, $B_y = \tilde{\mu}_y - |\mu|$. For all $\theta, \mu \in \mathbb{R}$ and $K = r \log n$ for some $0 < r < 1/3$, we have*

$$|B_x| \lesssim \frac{1}{\sqrt{\log n}}, \quad |B_y| \lesssim \frac{1}{\sqrt{\log n}},$$

and

$$\tilde{\sigma}_x^2 = O(n^{6r} \log^3 n), \quad \tilde{\sigma}_y^2 = O(n^{6r} \log^3 n).$$

In particular, when $\theta = 0$, we have $|B_x| \leq n^{6r-2} \log^3 n$ and $\tilde{\sigma}_x^2 \lesssim n^{6r} \log^3 n$, whereas when $\mu = 0$, we have $|B_y| \leq n^{6r-2} \log^3 n$, and $\tilde{\sigma}_y^2 \lesssim n^{6r} \log^3 n$.

Again, let $(x_{1\ell}, x_{2\ell}, \dots, x_{n\ell}) \sim N(\theta, \mathbf{I}_n)$ and $(y_{1\ell}, y_{2\ell}, \dots, y_{n\ell}) \sim N(\mu, \mathbf{I}_n)$ for $\ell = 1, 2$, and let $\widehat{T(\theta, \mu)} = \sum_{i=1}^n \xi(x_{i1}, x_{i2}, y_{i1}, y_{i2})$. It follows that the bias can be bounded by

$$|B(\widehat{T(\theta, \mu)})| \lesssim \frac{\|\theta\|_\infty s}{\sqrt{\log n}} + \frac{\|\mu\|_\infty s}{\sqrt{\log n}} + \frac{s}{\log n}.$$

On the other hand, the variance of $\widehat{T(\theta, \mu)}$ is bounded by

$$\begin{aligned} \text{Var}(\widehat{T(\theta, \mu)}) &\leq \sum_{i=1}^n \text{Var}(\xi(x_{i1}, x_{i2}, y_{i1}, y_{i2})) \\ &\lesssim n^{12r+1} \log^6 n + n^{6r+1} \log^3 n \cdot (\|\theta\|_\infty^2 + \|\mu\|_\infty^2) \end{aligned} \quad (37)$$

$$\lesssim |B(\widehat{T(\theta, \mu)})|^2, \quad (38)$$

as long as $0 < r < \frac{2\beta-1}{12}$. In this case, we have

$$\mathbb{E}(\widehat{T(\theta, \mu)} - T(\theta, \mu))^2 \lesssim \frac{s^2}{\log^2 n} + \frac{s^2 \|\theta\|_\infty^2}{\log n} + \frac{s^2 \|\mu\|_\infty^2}{\log n}.$$

The final result follows from the fact that $\max(\|\theta\|_\infty, \|\mu\|_\infty) \leq L_n$ and $L_n \gtrsim \sqrt{\log n}$.

2.2 Proof of Theorem 3

Let $x_1, x_2 \sim N(\theta, 1)$ and $y_1, y_2 \sim N(\mu, 1)$. Define

$$\xi = \xi(x_1, x_2, y_1, y_2) = [|x_1|I(|x_2| > 2\sqrt{2\log n})] \times [|y_1|I(|y_2| > 2\sqrt{2\log n})]. \quad (39)$$

Note that

$$\mathbb{E}(\xi) = \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}) \times \mathbb{E}|y_1|P(|y_2| > 2\sqrt{2\log n}) \quad (40)$$

Denote

$$\tilde{\sigma}_x^2 = \text{Var}(|x_1|I(|x_2| > 2\sqrt{2\log n})), \quad \tilde{\sigma}_y^2 = \text{Var}(|y_1|I(|y_2| > 2\sqrt{2\log n})),$$

and

$$\tilde{\theta}_x = \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}), \quad \tilde{\mu}_y = \mathbb{E}|y_1|P(|y_2| > 2\sqrt{2\log n}).$$

Then we have

$$\text{Var}(\xi) = \tilde{\sigma}_x^2\tilde{\sigma}_y^2 + \tilde{\sigma}_x^2\tilde{\mu}_y^2 + \tilde{\sigma}_y^2\tilde{\theta}_x^2$$

Proposition 3. *Let $B_x = \tilde{\theta}_x - |\theta|$ and $B_y = \tilde{\mu}_y - |\mu|$. For all $\theta, \mu \in \mathbb{R}$, we have*

$$|B_x| \lesssim \sqrt{\log n}, \quad |B_y| \lesssim \sqrt{\log n},$$

and

$$\tilde{\sigma}_x^2 \lesssim \log n, \quad \tilde{\sigma}_y^2 \lesssim \log n.$$

In particular, when $\theta = 0$ we have $|B_x| \lesssim n^{-4}$ and $\tilde{\sigma}_x^2 \lesssim n^{-4}$, whereas when $\mu = 0$ we have $|B_y| \lesssim n^{-4}$ and $\tilde{\sigma}_y^2 \lesssim n^{-4}$.

Now the bias of the estimator ξ is

$$\begin{aligned} B(\xi) &= \mathbb{E}\xi - |\theta||\mu| = \tilde{\theta}_x\tilde{\mu}_y - |\theta||\mu| \\ &\leq |\theta||B_y| + |\mu||B_x| + |B_x||B_y|, \end{aligned} \quad (41)$$

whereas the variance is bounded by

$$\text{Var}(\xi) = \tilde{\sigma}_x^2\tilde{\sigma}_y^2 + \tilde{\sigma}_x^2\tilde{\mu}_y^2 + \tilde{\sigma}_y^2\tilde{\theta}_x^2 \lesssim \tilde{\sigma}_x^2\tilde{\sigma}_y^2 + |\mu|^2\tilde{\sigma}_x^2 + |\theta|^2\tilde{\sigma}_y^2 + \log n(\tilde{\sigma}_y^2 + \tilde{\sigma}_x^2). \quad (42)$$

Now let $(x_{1\ell}, x_{2\ell}, \dots, x_{n\ell}) \sim N(\theta, \mathbf{I}_n)$ and $(y_{1\ell}, y_{2\ell}, \dots, y_{n\ell}) \sim N(\mu, \mathbf{I}_n)$ for $\ell = 1, 2$, and let

$$\widehat{T(\theta, \mu)} = \sum_{i=1}^n \xi(x_{i1}, x_{i2}, y_{i1}, y_{i2}).$$

It then follows that

$$|B(\widehat{T(\theta, \mu)})| \leq s \log n + (\|\theta\|_\infty + \|\mu\|_\infty)s\sqrt{\log n},$$

and

$$\text{Var}(\widehat{T(\theta, \mu)}) \leq s \log^2 n + (\|\theta\|_\infty^2 + \|\mu\|_\infty^2) s \log n. \quad (43)$$

Therefore the mean squared error of $\widehat{T(\theta, \mu)}$ satisfies

$$\begin{aligned} \mathbb{E}(\widehat{T(\theta, \mu)} - T(\theta, \mu))^2 &\leq B^2(\widehat{T(\theta, \mu)}) + \text{Var}(\widehat{T(\theta, \mu)}) \\ &\lesssim s^2 \log n (\|\theta\|_\infty^2 + \|\mu\|_\infty^2 + \log n). \end{aligned}$$

The final result follows from the fact that $\max(\|\theta\|_\infty, \|\mu\|_\infty) \leq L_n$ and $L_n \gtrsim \sqrt{\log n}$.

2.3 Proof of Theorem 4

By our sample splitting argument, it suffices to obtain the mean squared risk bound for the estimator $\widehat{T(\theta, \mu)} = \sum_{i=1}^n \hat{U}_i(x_i) \hat{U}_i(y_i)$ of $T(\theta, \mu) = \sum_{i=1}^n |\theta_i| |\mu_i|$ where $\hat{U}_i(x_i) = |x_{i1}| I(|x_{i2}| > 2\sqrt{2 \log n})$, $\hat{U}_i(y_i) = |y_{i1}| I(|y_{i2}| > 2\sqrt{2 \log n})$ and $(x_{1\ell}, x_{2\ell}, \dots, x_{n\ell}) \sim N(\theta, \Sigma_1)$ and $(y_{1\ell}, y_{2\ell}, \dots, y_{n\ell}) \sim N(\mu, \Sigma_2)$ for $\ell = 1, 2$. By Proposition 3, the bias of the estimator $\xi_i = \hat{U}_i(x_i) \hat{U}_i(y_i)$ satisfies

$$B(\xi_i) = \mathbb{E}\xi_i - |\theta_i| |\mu_i| \leq |\mu_i| |B_{ix}| + |\theta_i| |B_{iy}| + |B_{ix}| |B_{iy}|, \quad (44)$$

where $B_{ix} = \hat{U}_i(x_i) - |\theta_i|$ and $B_{iy} = \hat{U}_i(y_i) - |\mu_i|$. The variance satisfies

$$\text{Var}(\xi_i) \lesssim \tilde{\sigma}_{ix}^2 \tilde{\sigma}_{iy}^2 + |\mu_i|^2 \tilde{\sigma}_{ix}^2 + |\theta_i|^2 \tilde{\sigma}_{iy}^2 + \log n (\tilde{\sigma}_{iy}^2 + \tilde{\sigma}_{ix}^2), \quad (45)$$

where $\tilde{\sigma}_{ix}^2 = \text{Var}(\hat{U}_i(x_i))$ and $\tilde{\sigma}_{iy}^2 = \text{Var}(\hat{U}_i(y_i))$. The covariance between two copies

$$\begin{aligned} \text{Cov}(\xi_i, \xi_j) &= \mathbb{E}\xi_i \xi_j - \mathbb{E}\xi_i \mathbb{E}\xi_j \\ &= \mathbb{E}\hat{U}_i(x_i) \hat{U}_i(y_i) \hat{U}_j(x_j) \hat{U}_j(y_j) - \mathbb{E}\hat{U}_i(x_i) \hat{U}_i(y_i) \mathbb{E}\hat{U}_j(x_j) \hat{U}_j(y_j) \\ &= \mathbb{E}\hat{U}_i(x_i) \hat{U}_j(x_j) \mathbb{E}\hat{U}_i(y_i) \hat{U}_j(y_j) - \mathbb{E}\hat{U}_i(x_i) \mathbb{E}\hat{U}_j(x_j) \mathbb{E}\hat{U}_i(y_i) \mathbb{E}\hat{U}_j(y_j) \end{aligned}$$

Thus, we have

$$\begin{aligned} |\text{Cov}(\xi_i, \xi_j)| &\leq |\text{Cov}(\hat{U}_i(x_i), \hat{U}_j(x_j))| \cdot \tilde{\mu}_i \tilde{\mu}_j + |\text{Cov}(\hat{U}_i(y_i), \hat{U}_j(y_j))| \cdot \tilde{\theta}_i \tilde{\theta}_j \\ &\quad + |\text{Cov}(\hat{U}_i(x_i), \hat{U}_j(x_j)) \text{Cov}(\hat{U}_i(y_i), \hat{U}_j(y_j))|, \end{aligned}$$

where $\tilde{\theta}_j = \mathbb{E}\hat{U}_j(x_j)$ and $\tilde{\mu}_j = \mathbb{E}\hat{U}_j(y_j)$. Note that

$$|\text{Cov}(\hat{U}_i(x_i), \hat{U}_j(x_j))| \leq |\mathbb{E}\hat{U}_i(x_i) \hat{U}_j(x_j)| + |\tilde{\theta}_i \tilde{\theta}_j|$$

where

$$\mathbb{E}\hat{U}_i(x_i)\hat{U}_j(x_j) = \mathbb{E}|x_{i2}||x_{j2}|P(|x_{i2}| > 2\sqrt{2\log n}, |x_{j2}| > 2\sqrt{2\log n}).$$

Suppose one of θ_i and θ_j is 0, and the other bounded by $C_n \gtrsim \sqrt{\log n}$. Then by the proof of Proposition 3, we have

$$\mathbb{E}\hat{U}_i(x_i)\hat{U}_j(x_j) = O(n^{-4}C_n^2)$$

and

$$|\tilde{\theta}_i\tilde{\theta}_j| = O(n^{-4}C_n).$$

So

$$|\text{Cov}(\hat{U}_i(x_i), \hat{U}_j(x_j))| = O(n^{-4}C_n^2).$$

As a result, since $\mu_i, \mu_j \lesssim C_n$, we have

$$|\text{Cov}(\xi_i, \xi_j)| \leq O(n^{-4}C_n^4).$$

Hence, summation over $O(n^2)$ terms will be bounded by $O(n^{-2}C_n^4)$. On the other hand, if neither θ_i or θ_j is zero, we have the trivial bound from Proposition 3

$$|\text{Cov}(\xi_i, \xi_j)| \leq \log^2 n,$$

and the summation over $O(s^2)$ terms will be bounded by $s^2 \log^2 n$. Thus, as long as $C_n \lesssim \sqrt{n}$, we have

$$\text{Var}(\widetilde{T(\theta, \mu)}) \leq \sum_{i=1}^n \text{Var}(\xi(x_{i1}, x_{i2}, y_{i1}, y_{i2})) + O(n^{-2}C_n^4) + O(s^2 \log^2 n) \lesssim s^2 \log^2 n. \quad (46)$$

Now note that $|B(\widetilde{T(\theta, \mu)})| \lesssim s \log n$, it follows that

$$\mathbb{E}(\widetilde{T(\theta, \mu)} - T(\theta, \mu))^2 \lesssim s^2 \log^2 n.$$

3 Proof of Propositions 1-3

3.1 Proof of Proposition 1

In the following, we divide into four cases according to the value of $|\theta|$. When $|\theta| = 0$, we show that we are actually estimating $|\theta|$ by 0. When $0 \leq |\theta| \leq \sqrt{2\log n}$, we show that the estimator ξ' behaves essentially like δ , which is a good estimator when $|\theta|$ is small. When $\sqrt{2\log n} < |\theta| \leq 4\sqrt{2\log n}$, we show that the hybrid estimator ξ uses either $\delta(x_1)$ or $|x_1|$ and in this case both are good estimators of $|\theta|$. When $|\theta|$ is large, the hybrid estimator is essentially the same as $|x_1|$. We need the following

lemmas to facilitate our proof.

Lemma 1. Consider $G_K(x)$ defined in the main paper. The constant term of $\tilde{G}_K(x) = \sum_{l=0}^K \tilde{g}_{2l} x^{2l}$, with $\tilde{g}_{2l} = M_n^{-2l+1} g_{2l}$, satisfies

$$\tilde{g}_0 = M_n g_0 \leq \frac{2M_n}{\pi(2K+1)}. \quad (47)$$

Lemma 2. Let $X \sim N(\theta, 1)$ and $S_K(x) = \sum_{k=1}^K g_{2k} M_n^{-2k+1} H_{2k}(x)$ with $M_n = 8\sqrt{\log n}$ and $K = O(1)$. Then for all $|\theta| \leq 4\sqrt{2\log n}$,

$$\left| \mathbb{E}S_K(X) - |\theta| \right| \leq \frac{4M_n}{\pi(2K+1)} \lesssim \sqrt{\log n},$$

and $\mathbb{E}S_K^2(X) \lesssim \log n$.

Lemma 3. Suppose $I(A)$ and $I(B)$ are indicator random variables independent of X and Y , with $A \cap B = \emptyset$ then

$$\begin{aligned} \text{Var}(XI(A) + YI(B)) &= \text{Var}(X)P(A) + \text{Var}(Y)P(B) + (\mathbb{E}X)^2P(A)P(A^c) \\ &\quad + (\mathbb{E}Y)^2P(B)P(B^c) - 2\mathbb{E}X\mathbb{E}YP(A)P(B). \end{aligned} \quad (48)$$

In particular, if $A^c = B$, then we have

$$\text{Var}(XI(A) + YI(A^c)) = \text{Var}(X)P(A) + \text{Var}(Y)P(A^c) + (\mathbb{E}X - \mathbb{E}Y)^2P(A)P(A^c). \quad (49)$$

Applying Lemma 3, we have

$$\begin{aligned} \tilde{\sigma}_x^2 &= \text{Var}(\delta(x_1))P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \text{Var}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) \\ &\quad + (\mathbb{E}\delta(x_1))^2P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n})(1 - P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n})) \\ &\quad + (\mathbb{E}|x_1|)^2P(|x_2| > 2\sqrt{2\log n})(1 - P(|x_2| > 2\sqrt{2\log n})) \\ &\quad - 2\mathbb{E}\delta(x_1)\mathbb{E}|x_1|P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n})P(|x_2| > 2\sqrt{2\log n}). \end{aligned}$$

$$\begin{aligned} \tilde{\sigma}_y^2 &= \text{Var}(\delta(y_1))P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \text{Var}(|y_1|)P(|y_2| > 2\sqrt{2\log n}) \\ &\quad + (\mathbb{E}\delta(y_1))^2P(\sqrt{2\log n} < |y_2| \leq 2\sqrt{2\log n})(1 - P(\sqrt{2\log n} < |y_2| \leq 2\sqrt{2\log n})) \\ &\quad + (\mathbb{E}|y_1|)^2P(|y_2| > 2\sqrt{2\log n})(1 - P(|y_2| > 2\sqrt{2\log n})) \\ &\quad - 2\mathbb{E}\delta(y_1)\mathbb{E}|y_1|P(\sqrt{2\log n} < |y_2| \leq 2\sqrt{2\log n})P(|y_2| > 2\sqrt{2\log n}). \end{aligned}$$

Case 1. $\theta = 0$. Note that $\delta(x_1)$ can be written as

$$\delta(x_1) = S_K(x_1) - (S_K(x_1) - n^2)I(S_K(x_1) \geq n^2).$$

Consequently,

$$\begin{aligned}
|B_x| &= |\mathbb{E}(\delta(x_1))P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n})| \\
&\leq |\mathbb{E}S_K(x_1)| + \mathbb{E}\{(S_K(x_1) - n^2)I(S_K(x_1) \geq n^2)\} + \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}) \\
&\equiv B_1 + B_2 + B_3.
\end{aligned}$$

By definition of $S_K(x_1)$ we have

$$B_1 = 0. \quad (50)$$

It follows from the standard bound for normal tail probability $\Phi(-z) \leq z^{-1}\phi(z)$ for $z > 0$ that

$$P(|x_2| > 2\sqrt{2\log n}) = 2\Phi(-2\sqrt{2\log n}) \leq \frac{1}{2\sqrt{\pi\log n}}n^{-4}. \quad (51)$$

And in this case

$$\mathbb{E}|x_1| = 2\phi(0). \quad (52)$$

It then follows that

$$B_3 \leq 2\phi(0) \cdot \frac{1}{2\sqrt{\pi\log n}}n^{-4} = \frac{1}{\pi\sqrt{2\log n}}n^{-4}. \quad (53)$$

Now consider B_2 . Note that for any random variable X and any constant $\lambda > 0$,

$$\mathbb{E}(XI(X \geq \lambda)) \leq \lambda^{-1}\mathbb{E}(X^2I(X \geq \lambda)) \leq \lambda^{-1}\mathbb{E}X^2.$$

This together with Lemma 2 yields that

$$B_2 \leq \mathbb{E}\{(S_K(x_1)I(S_K(x_1) \geq n^2))\} \leq n^{-2}\mathbb{E}(S_K^2(x_1)) \lesssim n^{-2}\log n. \quad (54)$$

Combining the three pieces together, we have

$$|B_x| \leq B_1 + B_2 + B_3 \lesssim \frac{\log n}{n^2}.$$

We now consider the variance. It follows that

$$\begin{aligned}
\tilde{\sigma}_x^2 &\leq \text{Var}(S_K(x_1))P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \text{Var}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) \\
&\quad + (\mathbb{E}\delta(x_1))^2 + (\mathbb{E}|x_1|)^2P(|x_2| > 2\sqrt{2\log n}) - 2\mathbb{E}\delta(x_1)\mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}) \\
&\leq \mathbb{E}S_K^2(x_1)n^{-1} + (\mathbb{E}\delta(x_1))^2 + [\mathbb{E}x_1^2 + (\mathbb{E}|x_1|)^2 - 2\mathbb{E}\delta(x_1)\mathbb{E}|x_1|] \cdot \frac{1}{2n^4\sqrt{\pi\log n}} \\
&\lesssim n^{-1}\log n + n^{-4}\log^4 n + \frac{1}{n^4\sqrt{\log n}} \\
&\lesssim n^{-1}\log n
\end{aligned}$$

where we use the fact that

$$P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) \leq P(\sqrt{2\log n} \leq |x|) \leq \Phi(-\sqrt{2\log n}) \leq n^{-1}.$$

Case 2. $0 < |\theta| \leq \sqrt{2\log n}$. In this case

$$\begin{aligned} |B_x| &= |\mathbb{E}(\delta(x_1))P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) - |\theta| \\ &\leq |\mathbb{E}S_K(x_1) - |\theta|| + \mathbb{E}\{(S_K(x_1) - n^2)I(S_K(x_1) \geq n^2)\} \\ &\quad + (\mathbb{E}|x_1|)P(|x_2| > 2\sqrt{2\log n}) + |\theta|(1 - P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n})) \\ &\equiv B_1 + B_2 + B_3 + B_4 \end{aligned}$$

From Lemma 2 we have

$$B_1 \lesssim \sqrt{\log n}.$$

Again, the standard bound for normal tail probability yields

$$P(|x_2| > 2\sqrt{2\log n}) \leq 2\Phi(-\sqrt{2\log n}) \leq \frac{1}{\sqrt{\pi \log n}} n^{-1}$$

Note that

$$\mathbb{E}|x_1| = |\theta| + 2\phi(\theta) - 2|\theta|\Phi(-|\theta|) \leq |\theta| + 1 \leq \sqrt{2\log n} + 1.$$

Then we have

$$B_3 \leq \left(\sqrt{2\log n} + 1 \right) \cdot \frac{1}{\sqrt{\pi \log n}} n^{-1} \leq 3n^{-1},$$

and

$$B_4 \leq |\theta| \leq \sqrt{2\log n}.$$

Note that B_2 follows (54), and we have

$$|B_x| \leq B_1 + B_2 + B_3 + B_4 \lesssim \sqrt{\log n}.$$

For the variance, note that

$$\begin{aligned} (\mathbb{E}\delta(x_1))^2 &\leq \mathbb{E}\delta^2(x_1) \\ &= \mathbb{E}(\min\{S_K^2(x_1), n^4\}) \\ &= \mathbb{E}[S_K^2(x_1) - (S_K^2(x_1) - n^4)I(S_K^2(x_1) > n^4)] \\ &\leq \mathbb{E}S_K^2(x_1) \\ &\lesssim \log n, \end{aligned}$$

and

$$\mathbb{E}x_1^2 + (\mathbb{E}|x_1|)^2 - 2\mathbb{E}\delta(x_1)\mathbb{E}|x_1| \leq \text{Var}(x_1) + (\mathbb{E}|x_1|)^2 + (\sqrt{2\log n})^2 \lesssim \log n$$

Then we have

$$\begin{aligned} \tilde{\sigma}_x^2 &\leq \text{Var}(S_K(x_1)) + \text{Var}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) + (\mathbb{E}\delta(x_1))^2 \\ &\quad + (\mathbb{E}|x_1|)^2P(|x_2| > 2\sqrt{2\log n}) + 2|\mathbb{E}\delta(x_1)| \cdot \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}) \\ &\leq \mathbb{E}S_K^2(x_1) + (\mathbb{E}\delta(x_1))^2 + [\mathbb{E}x_1^2 + (\mathbb{E}|x_1|)^2 + 2|\mathbb{E}\delta(x_1)| \cdot \mathbb{E}|x_1|] \cdot \frac{1}{n\sqrt{\pi\log n}} \\ &\leq 2\log n + \frac{5\sqrt{\log n}}{\sqrt{\pi n}} \\ &\lesssim \log n. \end{aligned}$$

Case 3 $\sqrt{2\log n} \leq |\theta| \leq 4\sqrt{2\log n}$. In this case,

$$\begin{aligned} |B_x| &= |\mathbb{E}(\delta(x_1))P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) - |\theta|| \\ &\leq |\mathbb{E}(\delta(x_1)) - |\theta|| + |\mathbb{E}|x_1| - |\theta|| + |\theta|P(|x_2| \leq 2\sqrt{2\log n}) \\ &\leq |\mathbb{E}S_K(x_1) - |\theta|| + \mathbb{E}\{(S_K(x_1) - n^2)I(S_K(x_1) \geq n^2)\} + 2\phi(\theta) + 4\sqrt{2\log n} \\ &\lesssim \sqrt{\log n} + n^{-2}\log n + n^{-1} \\ &\lesssim \sqrt{\log n}. \end{aligned}$$

For the variance, similarly since

$$\begin{aligned} \mathbb{E}x_1^2 + (\mathbb{E}|x_1|)^2 + 2|\mathbb{E}\delta(x_1)| \cdot \mathbb{E}|x_1| &\leq \text{Var}(x_1) + (\mathbb{E}|x_1|)^2 + (4\sqrt{2\log n})^2 \\ &\quad + \left(4\sqrt{2\log n} + \frac{2M_n}{\pi K}\right)4\sqrt{2\log n} \\ &\leq 1 + (32 + 32 + 32)\log n + \frac{64\sqrt{2}\log n}{\pi K} \\ &\lesssim \log n, \end{aligned}$$

and then

$$\begin{aligned} \tilde{\sigma}_x^2 &\leq \text{Var}(S_K(x_1)) + \text{Var}(|x_1|) + (\mathbb{E}\delta(x_1))^2 + (\mathbb{E}|x_1|)^2 + 2|\mathbb{E}\delta(x_1)| \cdot \mathbb{E}|x_1| \\ &\leq \mathbb{E}S_K^2(x_1) + (\mathbb{E}\delta(x_1))^2 + [\mathbb{E}x_1^2 + (\mathbb{E}|x_1|)^2 + 2\mathbb{E}\delta(x_1)\mathbb{E}|x_1|] \\ &\lesssim \log n. \end{aligned}$$

Case 4. $|\theta| > 4\sqrt{2\log n}$. In this case, the standard bound for normal tail probability yields that

$$P(|x_2| \leq 2\sqrt{2\log n}) \leq 2\Phi(-(|\theta|/ - 2\sqrt{2\log n})) \leq 2\Phi\left(-\frac{|\theta|}{2}\right) \leq \frac{4}{|\theta|}\phi\left(\frac{|\theta|}{2}\right).$$

In particular,

$$P(|x_2| \leq 2\sqrt{2\log n}) \leq 2\Phi(-2\sqrt{2\log n}) \leq \frac{1}{2\sqrt{\pi\log n}}n^{-4}.$$

Also note that

$$\begin{aligned}\mathbb{E}\delta(x_1) &= \mathbb{E}\min\{S_K(x_1), n^2\} \\ &= \mathbb{E}(S_K(x_1)1\{S_K(x_1) \leq n^2\} + n^21\{S_K(x_1) > n^2\}) \\ &\leq n^2\end{aligned}$$

Hence,

$$\begin{aligned}|B_x| &\leq |\mathbb{E}(\delta(x_1))P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) - |\theta|| \\ &\leq |\mathbb{E}(\delta(x_1))|P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}|x_1| - |\theta| + \mathbb{E}|x_1|P(|x_2| \leq 2\sqrt{2\log n}) \\ &\leq |\mathbb{E}|x_1| - |\theta| + (|\mathbb{E}\delta(x_1)| + \mathbb{E}|x_1|)P(|x_2| \leq 2\sqrt{2\log n}) \\ &\leq 2\phi(\theta) + (n^2 + |\theta| + 1)P(|x_2| \leq 2\sqrt{2\log n}) \\ &\leq 2\phi(\theta) + 4\phi\left(\frac{|\theta|}{2}\right) + \frac{1}{2}n^{-2} \\ &\leq 6\phi\left(\frac{\theta}{2}\right) + \frac{1}{2n^2} \\ &\leq \frac{1}{n^2}.\end{aligned}$$

For the variance, similarly we have

$$\begin{aligned}\tilde{\sigma}_x^2 &\leq \text{Var}(\delta(x_1))P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) + \text{Var}(|x_1|) + (\mathbb{E}|x_1|)^2P(|x_2| \leq 2\sqrt{2\log n}) \\ &\quad + (\mathbb{E}\delta(x_1))^2P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) \\ &\quad + 2|\mathbb{E}\delta(x_1)| \cdot \mathbb{E}|x_1|P(\sqrt{2\log n} < |x_2| \leq 2\sqrt{2\log n}) \\ &\leq 1 + [\text{Var}(\delta(x_1)) + (\mathbb{E}\delta(x_1))^2 + (\mathbb{E}|x_1|)^2 + 2|\mathbb{E}\delta(x_1)| \cdot \mathbb{E}|x_1|]P(|x_2| \leq 2\sqrt{2\log n}) \\ &\leq 1 + [\log n + (n^2 + |\theta| + 1)^2]P(|x_2| \leq 2\sqrt{2\log n}) \\ &= 1 + o(1).\end{aligned}$$

Obviously, the same argument holds for y_1 , y_2 and $|\mu|$.

3.2 Proof of Proposition 2

We only prove the proposition for θ . The argument for μ is the same. We need the following lemma for the proof.

Lemma 4. *Let $X \sim N(\theta, 1)$ and $S_K(x) = \sum_{k=1}^K g_{2k} M_n^{-2k+1} H_{2k}(x)$ with $M_n = 8\sqrt{\log n}$ and $K = r \log n$ for some $r > 0$. Then for all $|\theta| \leq 4\sqrt{2 \log n}$,*

$$\left| \mathbb{E} S_K(X) - |\theta| \right| \leq \frac{4M_n}{\pi(2K+1)} \lesssim \frac{1}{\sqrt{\log n}},$$

and $\mathbb{E} S_K^2(X) \lesssim n^{6r} \log^3 n$.

Case 1. $\theta = 0$. Note that $\delta(x_1)$ can be written as

$$\delta(x_1) = S_K(x_1) - (S_K(x_1) - n^2)I(S_K(x_1) \geq n^2).$$

Consequently,

$$\begin{aligned} |B_x| &\leq |\mathbb{E} S_K(x_1)| + \mathbb{E}\{(S_K(x_1) - n^2)I(S_K(x_1) \geq n^2)\} \\ &\quad + (|\mathbb{E} S_K(x_1)| + \mathbb{E}|x_1|)\mathbb{P}(|x_2| > 2\sqrt{2 \log n}) \\ &\equiv B_1 + B_2 + B_3. \end{aligned}$$

By definition of $S_K(x_1)$ we have

$$B_1 = 0. \tag{55}$$

It follows from the standard bound for normal tail probability $\Phi(-z) \leq z^{-1}\phi(z)$ for $z > 0$ that

$$\mathbb{P}(|x_2| > 2\sqrt{2 \log n}) = 2\Phi(-2\sqrt{2 \log n}) \leq \frac{1}{2\sqrt{\pi \log n}} n^{-4}. \tag{56}$$

And in this case

$$\mathbb{E}|x_1| = 2\phi(0). \tag{57}$$

It then follows that

$$B_3 \leq 2\phi(0) \cdot \frac{1}{2\sqrt{\pi \log n}} n^{-4} = \frac{1}{\pi\sqrt{2 \log n}} n^{-4}. \tag{58}$$

Now consider B_2 . Note that for any random variable X and any constant $\lambda > 0$,

$$\mathbb{E}(XI(X \geq \lambda)) \leq \lambda^{-1}\mathbb{E}(X^2I(X \geq \lambda)) \leq \lambda^{-1}\mathbb{E}X^2.$$

This together with Lemma 4 yields that

$$B_2 \leq \mathbb{E}\{(S_K(x_1)I(S_K(x_1) \geq n^2))\} \leq n^{-2}\mathbb{E}(S_K^2(x_1)) \lesssim n^{6r-2} \log^3 n. \quad (59)$$

Combining the three pieces together, we have

$$|B_x| \leq B_1 + B_2 + B_3 \lesssim n^{6r-2} \log^3 n.$$

We now consider the variance. It follows from Lemma 3 and Lemma 4 that

$$\begin{aligned} \tilde{\sigma}_x^2 &\leq \text{Var}(S_K(x_1))P(|x_2| \leq 2\sqrt{2\log n}) + \text{Var}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) \\ &\quad + [(\mathbb{E}\delta(x_1))^2 + (\mathbb{E}|x_1|)^2]P(|x_2| > 2\sqrt{2\log n}) - 2\mathbb{E}\delta(x_1)\mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}) \\ &\leq \mathbb{E}S_K^2(x_1) + [(\mathbb{E}\delta(x_1))^2 + (\mathbb{E}|x_1|)^2 - 2\mathbb{E}\delta(x_1)\mathbb{E}|x_1|] \cdot \frac{1}{2n^4\sqrt{\pi\log n}} \\ &\lesssim n^{6r} \log^3 n \end{aligned}$$

as long as $r < 3/4$.

Case 2. $0 < |\theta| \leq \sqrt{2\log n}$. In this case

$$\begin{aligned} |B_x| &= |\mathbb{E}(\delta(x_1))P(|x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) - |\theta|| \\ &\leq |\mathbb{E}(\delta(x_1)) - |\theta|| + |\mathbb{E}|x_1| - |\theta||P(|x_2| > 2\sqrt{2\log n}) \\ &\lesssim |\mathbb{E}S_K(x_1) - |\theta|| + \mathbb{E}\{(S_K(x_1) - n^2)I(S_K(x_1) \geq n^2)\} + n^{-4} \\ &\lesssim 1/\sqrt{\log n} + n^{6r-2} \log^3 n + n^{-4} \\ &\lesssim 1/\sqrt{\log n} \end{aligned}$$

as long as $r < 1/3$. Similarly, note that

$$\mathbb{E}\delta(x_1) = \mathbb{E}S_K(x_1) + B_2$$

then

$$\begin{aligned} (\mathbb{E}\delta(x_1) - \mathbb{E}|x_1|)^2 &\leq 2(\mathbb{E}\delta(x_1))^2 + 2(\mathbb{E}|x_1|)^2 \\ &\leq 2(\mathbb{E}S_K(x_1) + B_2)^2 + 2(|\theta| + 1)^2 \\ &\lesssim n^{6r} \log^3 n. \end{aligned}$$

Hence the variance can be bounded as follows.

$$\begin{aligned}\tilde{\sigma}_x^2 &\leq \mathbb{E}S_K^2(x_1) + \mathbb{E}x_1^2\mathbb{P}(|x_2| > 2\sqrt{2\log n}) \\ &\quad + (\mathbb{E}\delta(x_1) - \mathbb{E}|x_1|)^2\mathbb{P}(|x_2| > 2\sqrt{2\log n}) \\ &\lesssim n^{6r}\log^3 n.\end{aligned}$$

Case 3. $\sqrt{2\log n} < |\theta| \leq 4\sqrt{2\log n}$. In this case

$$\begin{aligned}|B_x| &= |\mathbb{E}(\delta(x_1))P(|x_2| \leq 2\sqrt{2\log n}) + \mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) - |\theta|| \\ &\leq |\mathbb{E}(\delta(x_1)) - |\theta|| + |\mathbb{E}|x_1| - |\theta|| \\ &\lesssim |\mathbb{E}S_K(x_1) - |\theta|| + \mathbb{E}\{(S_K(x_1) - n^2)I(S_K(x_1) \geq n^2)\} + 2\phi(\theta) \\ &\lesssim 1/\sqrt{\log n} + n^{6r-2}\log^3 n + n^{-1} \\ &\lesssim 1/\sqrt{\log n}\end{aligned}$$

as long as $r < 1/3$. The variance can be bounded similar to the Case 2.

Case 4. $|\theta| > 4\sqrt{2\log n}$. In this case, same argument follows from the proof of Case 4 in Proposition 1.

Proof of Proposition 3. Similar to the proofs of previous propositions, we only prove the statements about θ . The argument for μ is the same.

Case 1. $\theta = 0$. Note that $|B_x| = \mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n})$. It follows from the standard bound for normal tail probability $\Phi(-z) \leq z^{-1}\phi(z)$ for $z > 0$ that

$$P(|x_2| > 2\sqrt{2\log n}) = 2\Phi(-2\sqrt{2\log n}) \leq \frac{1}{2\sqrt{\pi\log n}}n^{-4}. \quad (60)$$

And in this case $E|x_1| = 2\phi(0)$. It then follows that

$$|B_x| \leq 2\phi(0) \cdot \frac{1}{2\sqrt{\pi\log n}}n^{-4} = \frac{1}{\pi\sqrt{2\log n}}n^{-4}. \quad (61)$$

We now consider the variance. It follows that

$$\tilde{\sigma}_x^2 \leq \text{Var}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) \leq \mathbb{E}x_1^2 \cdot \frac{1}{2n^4\sqrt{\pi\log n}} \lesssim \frac{1}{n^4\sqrt{\log n}}.$$

Case 2. $0 < |\theta| \leq \sqrt{2\log n}$. In this case

$$|B_x| = |\mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}) - \theta| \leq \mathbb{E}|x_1|P(|x_2| > 2\sqrt{2\log n}) + |\theta|.$$

The standard bound for normal tail probability yields

$$P(|x_2| > 2\sqrt{2\log n}) \leq 2\Phi(-\sqrt{2\log n}) \leq \frac{1}{\sqrt{\pi \log n}} n^{-1}$$

Note that

$$\mathbb{E}|x_1| = |\theta| + 2\phi(\theta) - 2|\theta|\Phi(-|\theta|) \leq |\theta| + 1 \leq \sqrt{2\log n} + 1.$$

Then we have

$$|B_x| \leq \left(\sqrt{2\log n} + 1 \right) \cdot \frac{1}{\sqrt{\pi \log n}} n^{-1} + |\theta| \leq 3n^{-1} + \sqrt{2\log n}.$$

On the other hand, since

$$\mathbb{E}x_1^2 \leq \text{Var}(x_1) + (\mathbb{E}|x_1|)^2 \leq 1 + 2\log n.$$

we have

$$\tilde{\sigma}_x^2 \leq \text{Var}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) \leq \mathbb{E}x_1^2 \cdot \frac{1}{n\sqrt{\pi \log n}} \lesssim \frac{\sqrt{\log n}}{n}.$$

Case 3 $\sqrt{2\log n} \leq |\theta| \leq 4\sqrt{2\log n}$. In this case 2,

$$\begin{aligned} |B_x| &= |\mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) - |\theta|| \\ &\leq |\mathbb{E}|x_1| - |\theta|| + |\theta|P(|x_2| \leq 2\sqrt{2\log n}) \\ &\lesssim \sqrt{\log n}. \end{aligned}$$

For the variance, similarly we have

$$\tilde{\sigma}_x^2 \leq \mathbb{E}x_1^2 \lesssim \log n.$$

Case 4. $|\theta| > 4\sqrt{2\log n}$. In this case, the standard bound for normal tail probability yields that

$$P(|x_2| \leq 2\sqrt{2\log n}) \leq 2\Phi(-(|\theta| - 2\sqrt{2\log n})) \leq 2\Phi\left(-\frac{|\theta|}{2}\right) \leq \frac{4}{|\theta|} \phi\left(\frac{|\theta|}{2}\right).$$

In particular,

$$P(|x_2| \leq 2\sqrt{2\log n}) \leq 2\Phi(-2\sqrt{2\log n}) \leq \frac{1}{2\sqrt{\pi \log n}} n^{-4}.$$

Hence,

$$\begin{aligned} |B_x| &\leq |\mathbb{E}(|x_1|)P(|x_2| > 2\sqrt{2\log n}) - |\theta|| \leq |\mathbb{E}|x_1| - |\theta|| + \mathbb{E}|x_1|P(|x_2| \leq 2\sqrt{2\log n}) \\ &\leq 2\phi(\theta) + (|\theta| + 1)P(|x_2| \leq 2\sqrt{2\log n}) \leq 2\phi(\theta) + 4\phi\left(\frac{|\theta|}{2}\right) \lesssim \frac{1}{n^2}. \end{aligned}$$

For the variance, similarly we have

$$\begin{aligned} \tilde{\sigma}_x^2 &\leq \text{Var}(|x_1|) + (\mathbb{E}|x_1|)^2 P(|x_2| \leq 2\sqrt{2\log n}) \\ &= 1 + o(1). \end{aligned}$$

4 Proofs of Technical Lemmas

Proof of Lemma 1. By Lemma 2 of Cai and Low (2011), for $x \in [-1, 1]$, we have

$$\max_{x \in [-1, 1]} |G_K(x) - |x|| \leq \frac{2}{\pi(2K + 1)}.$$

Then for $x' \in [-M_n, M_n]$, we have $x' = M_n x$, and thus

$$\max_{x'} |\tilde{G}_K(x') - |x'|| \leq \frac{2M_n}{\pi(2K + 1)}. \quad (62)$$

Set $x' = 0$, we obtain the statement.

Proof of Lemma 2. The first statement follows from Lemma 2 in Cai and Low (2011) and that

$$|\mathbb{E}S_K(X) - |\theta|| \leq \left| \sum_{k=0}^K g_{2k} M_n^{-2k+1} \theta^{2k} - |\theta| \right| + \frac{2M_n}{\pi(2K + 1)} \quad (63)$$

$$\leq \frac{4M_n}{\pi(2K + 1)}. \quad (64)$$

To bound $\mathbb{E}S_K^2(X)$, it follows from Lemma 3 in Cai and Low (2011) and that

$$\mathbb{E}S_K^2(X) \leq \left(\sum_{k=1}^K |g_{2k}| M_n^{-2k+1} (\mathbb{E}H_{2k}^2(X))^{1/2} \right)^2 \quad (65)$$

$$\leq 2^{6K} \left(\sum_{k=1}^K M_n^{-2k+1} (2M_n^2)^k \right)^2 \quad (66)$$

$$\leq 2^{8K} M_n^2 K^2 \quad (67)$$

$$\lesssim \log n. \quad (68)$$

Proof of Lemma 3. For equation (48), since events A and B are independent of random variables X and Y , we have

$$\begin{aligned} \text{Var}(XI(A) + YI(B)) &= \mathbb{E}[X^2I(A) + Y^2I(B) + 2XYI(A)I(B)] - (\mathbb{E}XP(A) + \mathbb{E}YP(B))^2 \\ &= \mathbb{E}X^2P(A) + \mathbb{E}Y^2P(B) - (\mathbb{E}X)^2P^2(A) - (\mathbb{E}Y)^2P^2(B) \\ &\quad - 2\mathbb{E}X\mathbb{E}YP(A)P(B) \\ &= \text{Var}(X)P(A) + \text{Var}(Y)P(B) + (\mathbb{E}X)^2(P(A) - P^2(A)) \\ &\quad + (\mathbb{E}Y)^2(P(B) - P^2(B)) - 2\mathbb{E}X\mathbb{E}YP(A)P(B) \end{aligned}$$

Equation (49) follows directly from the above derivation.

Proof of Lemma 4. It follows from Lemma 2 in Cai and Low (2011) that

$$\left| \mathbb{E}S_K(X) - |\theta| \right| = \left| \sum_{k=0}^K g_{2k} M_n^{-2k+1} \theta^{2k} - |\theta| \right| + \frac{2M_n}{\pi(2K+1)} \quad (69)$$

$$\leq \frac{4M_n}{\pi(2K+1)}. \quad (70)$$

To bound $\mathbb{E}S_K^2(X)$, it follows that when $K = r \log n$ for some $r > 0$,

$$\mathbb{E}S_K^2(X) \leq \left(\sum_{k=1}^K |g_{2k}| M_n^{-2k+1} (EH_{2k}^2(X))^{1/2} \right)^2 \quad (71)$$

$$\leq 2^{6K} \left(\sum_{k=1}^K M_n^{-2k+1} (2M_n^2)^k \right)^2 \quad (72)$$

$$\leq 2^{8K} M_n^2 K^2 \quad (73)$$

$$\lesssim n^{6r} \log^3 n. \quad (74)$$

A Supplementary Figures and Tables

Simulations. In our Section 3, to generate dependent observations from a non-identity covariance matrix, we considered the block-wise covariances where Σ is block diagonal where each block is either a 10×10 Toeplitz matrix or a 1000×1000 exchangeable covariance matrix whose off-diagonal elements are 0.5. In particular, the 10×10 Toeplitz matrix is given as follows

$$\begin{bmatrix} 1 & 0.9 & 0.8 & 0.7 & 0.6 & 0.5 & \dots & 0.1 \\ 0.9 & 1 & 0.9 & 0.8 & 0.7 & 0.6 & \dots & 0.2 \\ 0.8 & 0.9 & 1 & 0.9 & 0.8 & 0.7 & \dots & 0.3 \\ \dots & & & & & & & \\ 0.1 & 0.2 & 0.3 & 0.4 & 0.5 & 0.6 & \dots & 1 \end{bmatrix}.$$

Real data analysis. In our gene set enrichment analysis, 5,023 biological processes from Gene Ontology (GO) (Botstein et al. 2000) that contain at least 10 genes were tested. Figure 2 presents the directed acyclic graphs of the GO biological processes that linked to the most significant GO terms from the simultaneous signal GSEA analysis.




Figure 2: Directed acyclic graph of GO biological processes connected by some path to the most significant processes from the GSEA analysis. Yellow: least significant; Red: most significant; Rectangles: top GSEA results.