

On the chordality of polynomial sets in triangular decomposition in top-down style*

Chenqi Mou and Yang Bai

LMIB – School of Mathematics and Systems Science /
Beijing Advanced Innovation Center for Big Data and Brain
Computing
Beihang University, Beijing 100191, China
{chenqi.mou, yangbai}@buaa.edu.cn

Abstract

In this paper the chordal graph structures of polynomial sets appearing in triangular decomposition in top-down style are studied when the input polynomial set to decompose has a chordal associated graph. In particular, we prove that the associated graph of one specific triangular set computed in any algorithm for triangular decomposition in top-down style is a subgraph of the chordal graph of the input polynomial set and that all the polynomial sets including all the computed triangular sets appearing in one specific simply-structured algorithm for triangular decomposition in top-down style (Wang’s method) have associated graphs which are subgraphs of the the chordal graph of the input polynomial set. These subgraph structures in triangular decomposition in top-down style are multivariate generalization of existing results for Gaussian elimination and may lead to specialized efficient algorithms and refined complexity analyses for triangular decomposition of chordal polynomial sets.

Key words: Chordal graph, triangular decomposition, top-down style, Wang’s method

1 Introduction

This paper is inspired by the pioneering work on the connections between chordal graphs and triangular sets [10], where the authors introduced the concept of chordal networks and proposed an algorithm for constructing

*This work was partially supported by the National Natural Science Foundation of China (NSFC 11401018 and 11771034)

chordal networks for polynomial sets based on the computation of triangular decomposition. In particular, the authors found that for input polynomial sets with chordal associated graphs, the elimination methods due to Wang become more efficient. It is worth mentioning that the authors also studied the connections between chordal graphs and Gröbner bases [9], and found that the chordal structures of polynomial sets are destroyed in the computation of Gröbner bases.

The chordal graph structures have been studied and applied in the prediction of structures of matrices appearing in solving (especially sparse) linear systems. It is shown that the sparsity of the matrices handled by Gaussian elimination can be controlled if the associated graph of this input matrix is chordal [23, 25, 18].

Like the Gröbner basis which has been greatly developed in its theory, methods, implementations, and applications [6, 12, 13, 11], the triangular set is also a powerful algebraic tool in the study on and computation of polynomials symbolically, especially for elimination theory and polynomial system solving [31, 16, 21, 27, 2, 30, 22, 8], with diverse applications [32, 7]. The process of decomposing a polynomial set into finitely many triangular sets or systems (probably with additional properties like being regular or normal, etc.) with associated zero and ideal relationships is called triangular decomposition of the input polynomial set. Triangular decomposition of polynomial sets can be regarded as multivariate generalization of Gaussian elimination for solving linear equations.

The top-down strategy in triangular decomposition means that the variables appearing in the input polynomial set are handled in a strictly decreasing order, and it is a widely-used strategy in the design and implementations of algorithms for triangular decomposition. In particular, most algorithms for triangular decomposition due to Wang are in the top-down style [27, 28, 29]. A Boolean algorithm for triangular decomposition in top-down style with refinement in the Boolean settings has also been proposed [17]. The fact that elimination in it is performed in a strictly decreasing order makes triangular decomposition in top-down style the closest among all kinds of triangular decomposition to Gaussian elimination, in which the elimination of entries in different columns of the matrix is also performed in a strict order.

The study in this paper arises naturally after summing up what are stated above: we study the chordal structures of polynomial sets appearing in the algorithms for triangular decomposition in top-down style, in particular the graph structures of the triangular sets computed by such algorithms. This is multivariate generalization of the study on the roles chordal structures play in Gaussian elimination, and it is highly non-trivial because in this polynomial (and thus nonlinear) case splitting occurs in the triangular decomposition which results in a complicated decomposition process.

With the introduction of associated graphs of polynomial sets in Sec-

tion 2, we define a polynomial set to be chordal if its associated graph is chordal. A chordal graph implies a perfect elimination ordering of the vertexes, and we assume that the polynomial set \mathcal{F} we want to decompose is chordal with the variables ordered as one perfect elimination ordering of its chordal associated graph $G(\mathcal{F})$. Under such assumptions, in this paper the graph structures of polynomial sets after reduction of one variable and all the variables in triangular decomposition in top-down style are exploited in Section 3. In particular, it is proved that the associated graph of one specific triangular set computed in an arbitrary algorithm for triangular decomposition in top-down style will be a subgraph of $G(\mathcal{F})$. Then in Section 4 for a specific simply-structured algorithm for triangular decomposition in top-down style, namely Wang's method, after reformulation of the underlying structures of its decomposition tree, we prove that each polynomial set occurring in the decomposition process has an associated graph being subgraph of $G(\mathcal{F})$, which directly implies that all the triangular sets computed by Wang's method have associated graphs which are subgraphs of $G(\mathcal{F})$. This paper ends with brief discussions on the applications of graph structures of polynomial sets on expressing the variable sparsity of polynomial sets and potential refined complexity analyses on triangular decomposition in top-down style in Section 5.

2 Preliminaries

Let \mathbb{K} be a field, and $\mathbb{K}[x_1, \dots, x_n]$ be the multivariate polynomial ring over \mathbb{K} in the variables x_1, \dots, x_n . For the sake of simplicity, we write (x_1, \dots, x_n) as \mathbf{x} and $\mathbb{K}[x_1, \dots, x_n]$ as $\mathbb{K}[\mathbf{x}]$.

2.1 Polynomial sets and associated graphs

For a polynomial $F \in \mathbb{K}[\mathbf{x}]$, define the (variable) *support* of F , denoted by $\text{supp}(F)$, to be the set of variables in x_1, \dots, x_n which effectively appear in F . For a polynomial set $\mathcal{F} \subset \mathbb{K}[\mathbf{x}]$, $\text{supp}(\mathcal{F}) := \cup_{F \in \mathcal{F}} \text{supp}(F)$.

The *associated graph* $G(\mathcal{F})$ of a polynomial set $\mathcal{F} \subset \mathbb{K}[\mathbf{x}]$ is an undirected graph constructed in the following way:

- (a) The vertexes of $G(\mathcal{F})$ are the variables in $\text{supp}(\mathcal{F})$.
- (b) There exists an edge connecting two vertexes x_i and x_j in $G(\mathcal{F})$ for $1 \leq i \neq j \leq n$ if there exists one polynomial $F \in \mathcal{F}$ such that $x_i, x_j \in \text{supp}(F)$.

Example 2.1 The associated graphs of

$$\begin{aligned}\mathcal{P} &= \{x_2 + x_1, x_3 + x_1, x_4^2 + x_2, x_4^3 + x_3, x_5 + x_2, x_5 + x_3 + x_2\} \\ \mathcal{Q} &= \{x_2 + x_1, x_3 + x_1, x_3, x_4^2 + x_2, x_4^3 + x_3, x_5 + x_2\}\end{aligned}$$




Figure 1: The associated graphs $G(\mathcal{P})$ (left) and $G(\mathcal{Q})$ (right) in Example 2.1

are shown in Figure 1.

Definition 2.1 Let $G = (V, E)$ be a graph with $V = \{x_1, \dots, x_n\}$. Then an ordering $x_{i_1} < x_{i_2} < \dots < x_{i_n}$ of the vertexes is called a *perfect elimination ordering* of G if for each $j = i_1, \dots, i_n$, the restriction of G on the following set

$$X_j = \{x_j\} \cup \{x_k : x_k < x_j \text{ and } (x_k, x_j) \in E\} \quad (1)$$

is a clique. A graph G is said to be *chordal* if there exists a perfect elimination ordering of it.

An equivalent condition for a graph $G = (V, E)$ to be *chordal* is the following: for any cycle C contained in G of four or more vertexes, there is an edge $e \in E \setminus C$ connecting two vertexes in C . A chordal graph is also called a triangulated one.

Definition 2.2 A polynomial set $\mathcal{F} \subset \mathbb{K}[\mathbf{x}]$ is said to be *chordal* if its associated graph $G(\mathcal{F})$ is chordal.

Example 2.2 In Example 2.1 and Figure 1, the associated graph $G(\mathcal{P})$ is chordal by definition and thus \mathcal{P} is chordal, while $G(\mathcal{Q})$ is not.

2.2 Triangular sets and triangular decomposition

Throughout this subsection the variables are ordered as $x_1 < \dots < x_n$. For an arbitrary polynomial $F \in \mathbb{K}[\mathbf{x}]$, denote the greatest variable appearing in F by $\text{lv}(F)$. Let $\text{lv}(F) = x_k$. Write $F = Ix_k^d + R$ with $I \in \mathbb{K}[x_1, \dots, x_{k-1}]$, $R \in \mathbb{K}[x_1, \dots, x_k]$, and $\deg(R, x_k) < d$. Then the polynomial I and R are called the *initial* and *tail* of F and denoted by $\text{ini}(F)$ and $\text{tail}(F)$ respectively. For two polynomial sets $\mathcal{F}, \mathcal{G} \subset \mathbb{K}[\mathbf{x}]$, the set of common zeros of \mathcal{F} is denoted by $Z(\mathcal{F})$, and $Z(\mathcal{F}/\mathcal{G}) := Z(\mathcal{F}) \setminus Z(\prod_{G \in \mathcal{G}} G)$.

Definition 2.3 An ordered set of non-constant polynomials $\mathcal{T} = [T_1, \dots, T_r] \subset \mathbb{K}[\mathbf{x}]$ is called a *triangular set* if $\text{lv}(T_1) < \dots < \text{lv}(T_r)$. A tuple $(\mathcal{T}, \mathcal{U})$ with $\mathcal{T}, \mathcal{U} \subset \mathbb{K}[\mathbf{x}]$ is called a *triangular system* if \mathcal{T} is a triangular set.

Definition 2.4 Let $\mathcal{F} \subset \mathbb{K}[\mathbf{x}]$ be a polynomial set. Then a finite number of triangular sets $\mathcal{T}_1, \dots, \mathcal{T}_r \subset \mathbb{K}[\mathbf{x}]$ are called a *decomposition* of \mathcal{F} into *triangular sets* if the zero relationship $Z(\mathcal{F}) = \cup_{i=1}^r Z(\mathcal{T}_i / \text{ini}(\mathcal{T}_i))$ holds, where $\text{ini}(\mathcal{T}_i) := \{\text{ini}(T) : T \in \mathcal{T}_i\}$.

Definition 2.5 Let $\mathcal{F} \subset \mathbb{K}[\mathbf{x}]$ be a polynomial set. Then a finite number of triangular systems $(\mathcal{T}_1, \mathcal{U}_1), \dots, (\mathcal{T}_r, \mathcal{U}_r)$ are called a *decomposition* of \mathcal{F} into *triangular systems* if the zero relationship $Z(\mathcal{F}) = \cup_{i=1}^r Z(\mathcal{T}_i / \mathcal{U}_i)$ holds.

As shown in Definitions 2.3, 2.4, and 2.5, triangular systems are generalization of triangular sets. For a triangular system $(\mathcal{T}, \mathcal{U})$, \mathcal{T} is a triangular set which represents the equations $\mathcal{T} = 0$ while \mathcal{U} is a polynomial set which represents the inequations $\mathcal{U} \neq 0$.

In general, the process of computing a decomposition of a polynomial set into triangular sets or triangular systems is called *triangular decomposition*. There exist many algorithms for decomposing polynomial sets into triangular sets or systems with different properties. One of the main strategies for designing such algorithms for triangular decomposition is to carry out reduction on polynomials containing the largest (unprocessed) variable until there is only one such polynomial with producing new polynomials whose leading variables are strictly smaller than the currently processed variable.

For any polynomial set $\mathcal{P} \subset \mathbb{K}[\mathbf{x}]$ and an integer i ($1 \leq i \leq n$), we denote $\mathcal{P}^{(i)} := \{P \in \mathcal{P} : \text{lv}(P) = x_i\}$. The smallest integer i ($1 \leq i \leq n$) such that $\#(\mathcal{P}^{(j)}) = 0$ or 1 for each $j = i+1, \dots, n$ is called the *level* of \mathcal{P} and denoted by $\text{level}(\mathcal{P})$. For two polynomial sets \mathcal{P} and \mathcal{Q} in $\mathbb{K}[\mathbf{x}]$, \mathcal{P} is said to be of *lower rank* than \mathcal{Q} if either $\text{level}(\mathcal{P}) < \text{level}(\mathcal{Q})$ or $\text{level}(\mathcal{P}) = \text{level}(\mathcal{Q})$ but the minimal degree in x_k of polynomials in $\mathcal{P}^{(k)}$ is strictly smaller than that in $\mathcal{Q}^{(k)}$, where $k = \text{level}(\mathcal{P}) = \text{level}(\mathcal{Q})$.

Let \mathcal{F} be a polynomial set in $\mathbb{K}[\mathbf{x}]$ and Φ be a set of polynomial sets, initialized with $\{\mathcal{F}\}$. Then an algorithm \mathcal{A} for computing triangular decomposition of \mathcal{F} is said to be in *top-down style* if for each polynomial set $\mathcal{P} \in \Phi$ with $\text{level}(\mathcal{P}) = k$, the algorithm \mathcal{A} computes one polynomial set \mathcal{P}' such that $\#(\mathcal{P}'^{(k)}) = 1$ and $\mathcal{P}'^{(i)} = \mathcal{P}^{(i)}$ for $i = k+1, \dots, n$ and finitely many $\mathcal{Q}_1, \dots, \mathcal{Q}_s$ which are all of lower ranks than \mathcal{P} and are put into Φ for later computation with \mathcal{A} such that $Z(\mathcal{P}) = Z(\mathcal{P}') \cup Z(\mathcal{Q}_1) \cup \dots \cup Z(\mathcal{Q}_s)$.

Remark 2.1 As mentioned in the introduction, algorithms for triangular decomposition in top-down style are polynomial generalization of Gaussian elimination for transforming a non-singular matrix into echelon form. The requirements on $\mathcal{Q}_1, \dots, \mathcal{Q}_s$ above to have lower ranks than \mathcal{P} guarantee termination of the algorithm \mathcal{A} .

3 Chordality of polynomial sets in general triangular decomposition in top-down style

In this section, the graph structures of polynomial sets in an arbitrary algorithm for triangular decomposition in top-down style are studied when the input polynomial set is chordal. We start this section with the connections between the associated graphs of a triangular set reduced from a chordal polynomial set and the chordal associated graph of the input polynomial set.

Proposition 3.1 *Let $\mathcal{P} \subset \mathbb{K}[\mathbf{x}]$ be a chordal polynomial set with $x_1 < \dots < x_n$ as one perfect elimination ordering, and for $i = 1, \dots, n$, let $T_i \in \mathbb{K}[\mathbf{x}]$ be a polynomial such that $\text{lv}(T_i) = x_i$ and $\text{supp}(T_i) \subset \text{supp}(\mathcal{P}^{(i)})$ (T_i is set null if $\mathcal{P}^{(i)} = \emptyset$). Then $\mathcal{T} = [T_1, \dots, T_n]$ is a triangular set, and $G(\mathcal{T}) \subset G(\mathcal{P})$. In particular, if $\text{supp}(T_i) = \text{supp}(\mathcal{P}^{(i)})$ for $i = 1, \dots, n$, then $G(\mathcal{T}) = G(\mathcal{P})$.*

Proof It is straightforward that \mathcal{T} is a triangular set with $\text{lv}(T_i) = x_i$ if $\mathcal{P}^{(i)} \neq \emptyset$ for $i = 1, \dots, n$.

For any edge $(x_i, x_j) \in G(\mathcal{T})$, there exists an integer k ($i, j \leq k \leq n$) such that $x_i, x_j \in \text{supp}(T_k)$. Then $x_i, x_j \in \text{supp}(\mathcal{P}^{(k)})$, and thus $(x_i, x_k) \in G(\mathcal{P})$ and $(x_j, x_k) \in G(\mathcal{P})$. Since $G(\mathcal{P})$ is chordal with $x_1 < \dots < x_n$ as a perfect elimination ordering and $x_i < x_k, x_j < x_k$, we know that $(x_i, x_j) \in G(\mathcal{P})$ by Definition 2.1. This proves the inclusion $G(\mathcal{T}) \subset G(\mathcal{P})$.

In particular, when $\text{supp}(T_i) = \text{supp}(\mathcal{P}^{(i)})$ for $i = 1, \dots, n$, next we show the inclusion $G(\mathcal{T}) \supset G(\mathcal{P})$, which implies the equality $G(\mathcal{T}) = G(\mathcal{P})$. For any $(x_i, x_j) \in G(\mathcal{P})$, there exists an integer k and a polynomial P such that $x_i, x_j \in \text{supp}(P)$ with $P \in \mathcal{P}^{(k)}$. Since $\text{supp}(P) \subseteq \text{supp}(\mathcal{T}_k)$, $x_i, x_j \in \text{supp}(T_k)$, and thus $(x_i, x_j) \in G(\mathcal{T})$. \square

Example 3.1 Proposition 3.1 does not necessarily hold in general if the polynomial set \mathcal{P} is not chordal. Consider the same \mathcal{Q} as in Example 2.1 whose associated graph $G(\mathcal{Q})$ is not chordal. Let

$$\mathcal{T} = [x_2 + x_1, x_3 + x_1, -x_2x_4 + x_3, x_5 + x_2].$$

Then one can check that for $i = 2, \dots, 5$, $\text{supp}(\mathcal{T}^{(i)}) = \text{supp}(\mathcal{Q}^{(i)})$, but the associated graph $G(\mathcal{T})$, as shown in Figure 2, is not a subgraph of $G(\mathcal{Q})$.

Proposition 3.1 above relates the associated graph of a triangular set and that of a chordal polynomial set when their variables satisfy certain conditions. The following theorem is for relating the associated graph of a chordal polynomial set before and after one kind of commonly used reduction in triangular decomposition.

Theorem 3.2 *Let $\mathcal{P} \subset \mathbb{K}[\mathbf{x}]$ be a chordal polynomial set such that $\mathcal{P}^{(n)} \neq \emptyset$ and $x_1 < \dots < x_n$ is one perfect elimination ordering. Let $T \in \mathbb{K}[\mathbf{x}]$ be a*




Figure 2: The associated graph $G(\mathcal{T})$ in Example 3.1

polynomial such that $\text{lv}(T) = x_n$ and $\text{supp}(T) \subset \text{supp}(\mathcal{P}^{(n)})$, and $\mathcal{R} \subset \mathbb{K}[\mathbf{x}]$ be a polynomial set such that $\text{supp}(\mathcal{R}) \subset \text{supp}(\mathcal{P}^{(n)}) \setminus \{x_n\}$. Then for the polynomial set $\tilde{\mathcal{P}} = \{\tilde{\mathcal{P}}^{(1)}, \dots, \tilde{\mathcal{P}}^{(n-1)}, T\}$, where $\tilde{\mathcal{P}}^{(k)} = \mathcal{P}^{(k)} \cup \mathcal{R}^{(k)}$ for $k = 1, \dots, n-1$, we have $G(\tilde{\mathcal{P}}) \subset G(\mathcal{P})$. In particular, if $\text{supp}(T) = \text{supp}(\mathcal{P}^{(n)})$, then $G(\tilde{\mathcal{P}}) = G(\mathcal{P})$.

Proof To prove the inclusion $G(\tilde{\mathcal{P}}) \subset G(\mathcal{P})$, it suffices to show that for each edge $(x_i, x_j) \in G(\tilde{\mathcal{P}})$, we have $(x_i, x_j) \in G(\mathcal{P})$. For an arbitrary edge $(x_i, x_j) \in G(\tilde{\mathcal{P}})$, there exists a polynomial $P \in \tilde{\mathcal{P}}$ and an integer k ($i, j \leq k \leq n$) such that $x_i, x_j \in \text{supp}(P)$ and $P \in \tilde{\mathcal{P}}^{(k)}$.

If $k = n$, then $x_i, x_j \in \text{supp}(T)$, and by $\text{supp}(T) \subset \text{supp}(\mathcal{P}^{(n)})$ we have $x_i, x_j \in \text{supp}(\mathcal{P}^{(n)})$. This implies that $(x_i, x_n), (x_j, x_n) \in G(\mathcal{P}^{(n)}) \subset G(\mathcal{P})$ and by the chordality of $G(\mathcal{P})$ we have $(x_i, x_j) \in G(\mathcal{P})$.

Else if $k < n$, then by $\tilde{\mathcal{P}}^{(k)} = \mathcal{P}^{(k)} \cup \mathcal{R}^{(k)}$ there are two cases for P accordingly. When $P \in \mathcal{P}^{(k)} \subset \mathcal{P}$, clearly $(x_i, x_j) \in G(\mathcal{P})$; when $P \in \mathcal{R}^{(k)}$, we have $x_i, x_j \in \text{supp}(\mathcal{R}^{(k)}) \subset \text{supp}(\mathcal{P}^{(n)})$, and thus $(x_i, x_n), (x_j, x_n) \in G(\mathcal{P}^{(n)}) \subset G(\mathcal{P})$, and the chordality $G(\mathcal{P})$ implies $(x_i, x_j) \in G(\mathcal{P})$.

In particular, if $\text{supp}(T) = \text{supp}(\mathcal{P}^{(n)})$, then by $G(\mathcal{P}^{(k)}) \subset G(\tilde{\mathcal{P}}^{(k)})$ for $k = 1, \dots, n-1$ and $G(\mathcal{P}^{(n)}) \subset G(T)$, we have $G(\mathcal{P}) \subset G(\tilde{\mathcal{P}})$. This proves the equality $G(\tilde{\mathcal{P}}) = G(\mathcal{P})$. \square

Example 3.2 Let \mathcal{P} be the chordal polynomial set as in Example 2.1. Then $\mathcal{P}^{(5)} = \{x_5 + x_2, x_5 + x_3 + x_2\}$. If we take $T = x_5 + x_2$, and $\mathcal{R} = \{\text{prem}(x_5 + x_3 + x_2, x_5 + x_2)\} = \{x_3\}$, then $\tilde{\mathcal{P}}$ equals \mathcal{Q} in Example 2.1, and $G(\tilde{\mathcal{P}})$ is a (strict) subgraph of $G(\mathcal{P})$; If we take $T = x_5 + x_3 + x_2$, and $\mathcal{R} = \{\text{prem}(x_5 + x_2, x_5 + x_3 + x_2)\} = \{-x_3\}$, then $\text{supp}(T) = \text{supp}(\mathcal{P}^{(5)})$ and thus $G(\tilde{\mathcal{P}}) = G(\mathcal{P})$.

Next we introduce some notations to formulate the reduction process in Theorem 3.2. Denote the power set of a set S by 2^S . For an integer i ($1 \leq i \leq n$), let f_i be a mapping

$$\begin{aligned} f_i : 2^{\mathbb{K}[\mathbf{x}_i] \setminus \mathbb{K}[\mathbf{x}_{i-1}]} &\rightarrow (\mathbb{K}[\mathbf{x}_i] \setminus \mathbb{K}[\mathbf{x}_{i-1}]) \times 2^{\mathbb{K}[\mathbf{x}_{i-1}]} \\ \mathcal{P} &\mapsto (T, \mathcal{R}) \end{aligned} \quad (2)$$

such that $\text{supp}(T) \subset \text{supp}(\mathcal{P})$ and $\text{supp}(\mathcal{R}) \subset \text{supp}(\mathcal{P})$, where $\mathbb{K}[\mathbf{x}_0]$ is understood as \mathbb{K} . For a polynomial set $\mathcal{P} \subset \mathbb{K}[\mathbf{x}]$ and a fixed integer i ($1 \leq i \leq n$), suppose that $(T_i, \mathcal{R}_i) = f_i(\mathcal{P}^{(i)})$ for some f_i as stated above. For $j = 1, \dots, n$, define the polynomial set

$$\text{red}_i(\mathcal{P}^{(j)}) := \begin{cases} \mathcal{P}^{(j)}, & \text{if } j > i \\ \{T_i\}, & \text{if } j = i \\ \mathcal{P}^{(j)} \cup \mathcal{R}_i^{(j)}, & \text{if } j < i \end{cases}$$

and $\text{red}_i(\mathcal{P}) := \cup_{j=1}^n \text{red}_i(\mathcal{P}^{(j)})$. In particular, write

$$\overline{\text{red}}_i(\mathcal{P}) := \text{red}_i(\text{red}_{i+1}(\dots(\text{red}_n(\mathcal{P}))\dots)) \quad (3)$$

for simplicity.

Here $\text{red}_i(\cdot)$ denotes the result of reduction with respect to x_i and $\overline{\text{red}}_i(\cdot)$ denotes the result of successive reduction with respect to x_n, x_{n-1}, \dots, x_i . Following the above terminologies, Theorem 3.2 can be reformulated as $G(\text{red}_n(\mathcal{P})) \subset G(\mathcal{P})$, and the equality holds if $\text{supp}(T_n) = \text{supp}(\mathcal{P}^{(n)})$.

Proposition 3.3 *Let $\mathcal{P} \subset \mathbb{K}[\mathbf{x}]$ be a chordal polynomial set with $x_1 < \dots < x_n$ as one perfect elimination ordering. For each i ($1 \leq i \leq n$), suppose that $(T_i, \mathcal{R}_i) = f_i(\overline{\text{red}}_{i+1}(\mathcal{P})^{(i)})$ for some f_i as in (2) and $\text{supp}(T_i) = \text{supp}(\overline{\text{red}}_{i+1}(\mathcal{P})^{(i)})$, where $\overline{\text{red}}_{n+1}(\mathcal{P})$ is understood as \mathcal{P} . Then $G(\overline{\text{red}}_1(\mathcal{P})) = G(\mathcal{P})$.*

Proof Repeated use of Theorem 3.2 implies

$$G(\mathcal{P}) = G(\text{red}_n(\mathcal{P})) = G(\overline{\text{red}}_{n-1}(\mathcal{P})) = \dots = G(\overline{\text{red}}_1(\mathcal{P})),$$

and the conclusion follows. \square

Remark 3.1 Note that $\overline{\text{red}}_1(\mathcal{P})$ forms a triangular set after reordering if $\overline{\text{red}}_1(\mathcal{P})$ does not contain any non-zero constant. Indeed, the reduction process to compute this triangular set is commonly used in algorithms for triangular decomposition in top-down style, and the mapping f_i in (2) is abstraction of specific reductions used in different kinds of algorithms for triangular decomposition [20]. For example, one specific kind of such reduction is performed by using pseudo-divisions, and in this case \mathcal{R} in (2) consists of pseudo-remainders which do not contain x_i .

Proposition 3.3 holds because after every reduction $G(\overline{\text{red}}_i(\mathcal{P}))$ remains the same as the chordal graph $G(\mathcal{P})$, and thus the hypotheses of Theorem 3.2 remain satisfied. If we loosen the condition $\text{supp}(T_i) = \text{supp}(\overline{\text{red}}_{i+1}(\mathcal{P})^{(i)})$ in Proposition 3.3 to $\text{supp}(T_i) \subset \text{supp}(\overline{\text{red}}_{i+1}(\mathcal{P})^{(i)})$, then in general we will not have

$$G(\overline{\text{red}}_1(\mathcal{P})) \subset \dots \subset G(\overline{\text{red}}_{n-1}(\mathcal{P})) \subset G(\text{red}_n(\mathcal{P})) \subset G(\mathcal{P}),$$

as shown by the following example (though the last inclusion always holds because $G(\mathcal{P})$ is chordal).

Example 3.3 Let us continue with Example 3.2 with \mathcal{P} and $\mathcal{Q} = \text{red}_5(\mathcal{P})$, where $G(\mathcal{Q}) \subsetneq G(\mathcal{P})$. Take

$$\begin{aligned} T_4 &= \text{prem}(x_4^3 + x_3, x_4^2 + x_2) = -x_2x_4 + x_3, \\ \mathcal{R}_4 &= \{\text{prem}(x_4^2 + x_2, -x_2x_4 + x_3)\} = \{x_3^2 - x_2^3\}, \end{aligned}$$

then

$$\mathcal{Q}' := \overline{\text{red}}_4(\mathcal{P}) = \{x_2 + x_1, x_3 + x_1, x_3^2 - x_2^3, x_3, -x_2x_4 + x_3, x_5 + x_2\}.$$

The associated graph $G(\mathcal{Q}')$ is shown below. Note that $G(\mathcal{Q}') \not\subset G(\mathcal{Q})$ but $G(\mathcal{Q}') \subset G(\mathcal{P})$.




Figure 3: The associated graph $G(\mathcal{Q}')$ in Example 3.3

Despite of this example where successive inclusions of the associated graphs in the reduction chain does not hold, it can be proved that for each $i = n, \dots, 1$, $G(\overline{\text{red}}_i(\mathcal{P}))$ is a subgraph of the original graph $G(\mathcal{P})$.

Lemma 3.4 *Let $\mathcal{P} \subset \mathbb{K}[x]$ be a chordal polynomial set with $x_1 < \dots < x_n$ as one perfect elimination ordering and $\overline{\text{red}}_i(\mathcal{P})$ be defined in (3) for $i = n, \dots, 1$. Then for each $i = n, \dots, 1$ and any two variables x_p and x_q , if there exists an integer k such that $x_p, x_q \in \text{supp}(\overline{\text{red}}_i(\mathcal{P})^{(k)})$, then $(x_p, x_q) \in G(\mathcal{P})$.*

Proof We induce on the integer i . In the case $i = n$, from the proof of Theorem 3.2 one can easily find that the proposition is true. Now suppose that the proposition holds for $i = j$ ($< n$), and next we prove that it also holds for $i = j - 1$, namely for any x_p and x_q , if there exists k ($\geq p, q$) such that $x_p, x_q \in \text{supp}(\overline{\text{red}}_{j-1}(\mathcal{P})^{(k)})$, then $(x_p, x_q) \in G(\mathcal{P})$.

First by $\overline{\text{red}}_{j-1}(\mathcal{P}) = \text{red}_{j-1}(\overline{\text{red}}_j(\mathcal{P}))$ we know that there exists a polynomial set $\tilde{\mathcal{R}}$ such that

$$\overline{\text{red}}_{j-1}(\mathcal{P})^{(k)} = \overline{\text{red}}_j(\mathcal{P})^{(k)} \cup \tilde{\mathcal{R}}^{(k)}$$

and $\text{supp}(\tilde{\mathcal{R}}) \subset \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(j-1)}) \setminus \{x_{j-1}\}$.

(a) If $\tilde{\mathcal{R}} = \emptyset$, then $x_p, x_q \in \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(k)})$, and by the induction assumption we know that $(x_p, x_q) \in G(\mathcal{P})$.

(b) If $\tilde{\mathcal{R}} \neq \emptyset$, then $x_k \in \text{supp}(\tilde{\mathcal{R}}^{(k)}) \subset \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(j-1)})$. Next we consider the following three cases.

Case (1): $x_p, x_q \in \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(k)})$: with the same argument as in (a) we know that $(x_p, x_q) \in G(\mathcal{P})$.

Case (2): $x_p, x_q \in \text{supp}(\tilde{\mathcal{R}}^{(k)}) \subset \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(j-1)})$: by the induction assumption we know that $(x_p, x_q) \in G(\mathcal{P})$.

Case (3): $x_p \in \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(k)})$ and $x_q \in \text{supp}(\tilde{\mathcal{R}}^{(k)}) \subset \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(j-1)})$: Since $x_p, x_k \in \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(k)})$, by the induction assumption we have $(x_p, x_k) \in G(\mathcal{P})$; since $x_q, x_k \in \text{supp}(\overline{\text{red}}_j(\mathcal{P})^{(j-1)})$, by the induction assumption we have $(x_q, x_k) \in G(\mathcal{P})$. Then by the chordality of \mathcal{P} , $(x_p, x_k) \in G(\mathcal{P})$ and $(x_q, x_k) \in G(\mathcal{P})$ imply that $(x_p, x_q) \in G(\mathcal{P})$.

This ends the proof of this proposition with induction on i . \square

Theorem 3.5 *Let $\mathcal{P} \subset \mathbb{K}[\mathbf{x}]$ be a chordal polynomial set with $x_1 < \dots < x_n$ as one perfect elimination ordering and $\overline{\text{red}}_i(\mathcal{P})$ be defined in (3) for $i = n, \dots, 1$. Then for each $i = n, \dots, 1$, $G(\overline{\text{red}}_i(\mathcal{P})) \subset G(\mathcal{P})$.*

Proof By the construction of $\overline{\text{red}}_i(\mathcal{P})$, we know that all the vertexes of $G(\overline{\text{red}}_i(\mathcal{P}))$ are also vertexes of $G(\mathcal{P})$. For each edge $(x_p, x_q) \in G(\overline{\text{red}}_i(\mathcal{P}))$, there exists an integer k ($p, q \leq k \leq n$) and a polynomial P such that $x_p, x_q \in \text{supp}(P)$ and $P \in \overline{\text{red}}_i(\mathcal{P})^{(k)}$. Then by Lemma 3.4, we know that $(x_p, x_q) \in G(\mathcal{P})$, and thus $G(\overline{\text{red}}_i(\mathcal{P})) \subset G(\mathcal{P})$. \square

Corollary 3.6 *Let $\mathcal{P} \subset \mathbb{K}[\mathbf{x}]$ be a chordal polynomial set with $x_1 < \dots < x_n$ as one perfect elimination ordering and $\overline{\text{red}}_i(\mathcal{P})$ be defined in (3) for $i = n, \dots, 1$. If $\mathcal{T} := \overline{\text{red}}_1(\mathcal{P})$ does not contain any nonzero constant, then \mathcal{T} forms a triangular set such that $G(\mathcal{T}) \subset G(\mathcal{P})$.*

Remark 3.2 Corollary 3.6 tells us that under the conditions that the input polynomial set is chordal and the variable ordering is one perfect elimination ordering, the associated graph of one specific triangular set computed in any algorithm for triangular decomposition in top-down style is a subgraph of the associated graph of the input polynomial set. In fact, this triangular set is the “main branch” in the triangular decomposition in the sense that other branches are obtained by adding additional constraints in the splitting in the process of triangular decomposition.

Note that in the case when the input polynomial set \mathcal{P} is not chordal, a process of chordal completion can be carried out on $G(\mathcal{P})$ to make it a chordal graph (in the worst case this chordal completion results in a complete graph which is trivially chordal). Then the conditions of Corollary 3.6 will be satisfied after this chordal completion.

The chordality of any triangular sets other than the specific one above in a triangular decomposition computed by an algorithm in top-down style is dependent on the strategy how splitting occurs in the algorithm. Therefore

in the next section we focus on Wang's method, one specific algorithm for triangular decomposition in top-down style, and prove that the associated graphs of all the triangular sets computed by Wang's method are subgraphs of the associated graph of a chordal input polynomial set.

4 Chordality of polynomial sets computed by Wang's method

A simply-structured algorithm was proposed by Wang for triangular decomposition in top-down style in 1993 [27], which is referred to as Wang's method in the literatures (see. e.g., [3]). In this section the chordality of polynomial sets in the decomposition process of Wang's method is studied.

4.1 Restatement of Wang's method

For the self-containness of this paper, Wang's method for triangular decomposition is outlined in Algorithm 1 below. In this algorithm, the data structure $(\mathcal{P}, \mathcal{Q}, i)$ is used to represent two polynomial sets \mathcal{P} and \mathcal{Q} such that $\#(\mathcal{P}^{(j)}) = 1$ or 0 for $j = i+1, \dots, n$, and the subroutine $\text{pop}(\Phi)$ returns an element in Φ and then remove it from Φ .

As shown in Algorithm 1, for each $(\mathcal{P}, \mathcal{Q}, i)$ picked from Φ , if \mathcal{P} is already a triangular set (namely $i = 0$, which means $\#(\mathcal{P}^{(j)})$ is either 0 or 1 for $j = 1, \dots, n$ and \mathcal{P} contains no non-zero constant), then $(\mathcal{P}, \mathcal{Q})$ is included in the output Ψ . Note that \mathcal{Q} in Algorithm 1 is for collecting the inequations $\text{ini}(T) \neq 0$ for T in Line 8 of Algorithm 1 and it does not impose any influence on the graph structures of the triangular sets computed by Wang's method.

The decomposition process in Wang's method (Algorithm 1) applied to \mathcal{F} can be viewed as a binary tree with its root as $(\mathcal{F}, \emptyset, n)$. The nodes of this binary tree are all the tuples $(\mathcal{P}, \mathcal{Q}, i)$ picked from Φ , and each node $(\mathcal{P}, \mathcal{Q}, i)$ has two children $(\mathcal{P}', \mathcal{Q}', i)$ and $(\mathcal{P}'', \mathcal{Q}'', i)$ where

$$\begin{aligned}\mathcal{P}' &:= \mathcal{P} \setminus \mathcal{P}^{(i)} \cup \{T\} \cup \{\text{prem}(P, T) : P \in \mathcal{P}\}, & \mathcal{Q}' &:= \mathcal{Q} \cup \{\text{ini}(T)\}, \\ \mathcal{P}'' &:= \mathcal{P} \setminus \{T\} \cup \{\text{ini}(T), \text{tail}(T)\}, & \mathcal{Q}'' &:= \mathcal{Q},\end{aligned}$$

with T as a polynomial in $\mathcal{P}^{(i)}$ with minimal degree in x_i . In fact, the left child node $(\mathcal{P}', \mathcal{Q}', i)$ corresponds to the case when $\text{ini}(T) \neq 0$ and thus reduction of $\mathcal{P}^{(i)}$ are performed with respect to T ; while the right child node $(\mathcal{P}'', \mathcal{Q}'', i)$ corresponds to the case $\text{ini}(T) = 0$ and thus T is replaced by $\text{ini}(T)$ and $\text{tail}(T)$ (since $T = 0$ and $\text{ini}(T) = 0$ imply $\text{tail}(T) = 0$).

The binary decomposition tree for Wang's method and the splitting at one node are illustrated in Figures 4 and 5 respectively.

Algorithm 1: Wang's method for triangular decomposition $\Psi := \text{TriDecWang}(\mathcal{F})$

Input: \mathcal{F} , a polynomial set in $\mathbb{K}[\mathbf{x}]$

Output: Ψ , a set of finitely many triangular systems which form a triangular decomposition of \mathcal{F}

```

1   $\Phi := \{(\mathcal{F}, \emptyset, n)\};$ 
2  while  $\Phi \neq \emptyset$  do
3       $(\mathcal{P}, \mathcal{Q}, i) := \text{pop}(\Phi);$ 
4      if  $i = 0$  then
5           $\Psi := \Psi \cup \{(\mathcal{P}, \mathcal{Q})\};$ 
6          Break;
7      while  $\#(\mathcal{P}^{(i)}) > 1$  do
8           $T :=$  a polynomial in  $\mathcal{P}^{(i)}$  with minimal degree in  $x_i$ ;
9           $\Phi := \Phi \cup \{(\mathcal{P} \setminus \{T\} \cup \{\text{ini}(T), \text{tail}(T)\}, \mathcal{Q}, i)\};$ 
10          $\overline{\mathcal{P}} := \mathcal{P}^{(i)} \setminus \{T\};$ 
11          $\mathcal{P} := \mathcal{P} \setminus \overline{\mathcal{P}};$ 
12         for  $P \in \mathcal{P}^{(i)}$  do
13              $\mathcal{P} := \mathcal{P} \cup \{\text{prem}(P, T)\};$ 
14              $\mathcal{Q} := \mathcal{Q} \cup \{\text{ini}(T)\};$ 
15          $\Phi := \Phi \cup \{(\mathcal{P}, \mathcal{Q}, i - 1)\};$ 
16 for  $(\mathcal{P}, \mathcal{Q}) \in \Psi$  do
17     if  $\mathcal{P}$  contains a non-zero constant then
18          $\Psi := \Psi \setminus \{(\mathcal{P}, \mathcal{Q})\}$ 
19 return  $\Psi$ 

```

4.2 Chordality of polynomial sets in Wang's method

With a chordal polynomial set as input for Wang's method with respect to a perfect elimination ordering, the relationships are first clarified in the following propositions between the associated graphs of the polynomials in the left nodes and the associated graph of the input polynomial set and between the associated graphs of the polynomials in the right child nodes and those of the polynomials in the parent node.

Proposition 4.1 *Let $\mathcal{F} \subset \mathbb{K}[\mathbf{x}]$ be a chordal polynomial set with $x_1 < \dots < x_n$ as one perfect elimination ordering and $(\mathcal{P}, \mathcal{Q}, i)$ be any node in the binary decomposition tree for Wang's method applied to \mathcal{F} such that $G(\mathcal{P}) \subset G(\mathcal{F})$, T be a polynomial in \mathcal{P} with minimal degree in x_i . Denote $\mathcal{P}' = \mathcal{P} \setminus \mathcal{P}^{(i)} \cup \{T\} \cup \{\text{prem}(P, T) : P \in \mathcal{P}^{(i)}\}$. Then $G(\mathcal{P}') \subset G(\mathcal{F})$.*

Proof Clearly the vertexes of $G(\mathcal{P}')$ are also those of $G(\mathcal{P})$ and thus those




Figure 4: Binary decomposition tree for Wang's method


of $G(\mathcal{F})$, and it suffices to prove that for any edge $(x_p, x_q) \in G(\mathcal{P}')$, we have $(x_p, x_q) \in G(\mathcal{F})$.

Denote $\mathcal{R} := \{\text{prem}(P, T) : P \in \mathcal{P}^{(i)}\}$. Then by definition $\text{supp}(\mathcal{R}^{(l)}) \subset \text{supp}(\mathcal{P}^{(i)})$ for any $l = 1, \dots, i$. Furthermore, the following relationships hold: $\mathcal{P}'^{(l)} = \mathcal{P}^{(l)} \cup \mathcal{R}^{(l)}$ for $l = 1, \dots, i-1$ and $\mathcal{P}'^{(i)} = \{T\} \cup \mathcal{R}^{(i)}$. For any edge $(x_p, x_q) \in G(\mathcal{P}')$, there exist an integer k ($p, q \leq k \leq i$) and a polynomial $P \in \mathcal{P}'^{(k)}$ such that $x_p, x_q \in \text{supp}(P)$.

In the case when $k = i$, we have $(x_p, x_q) \in G(\mathcal{P}'^{(i)})$. Then $x_p, x_q \in \text{supp}(T) \cup \text{supp}(\mathcal{R}^{(i)}) \subset \text{supp}(\mathcal{P}^{(i)})$ and thus $(x_p, x_i), (x_q, x_i) \in G(\mathcal{P}^{(i)}) \subset G(\mathcal{F})$. By the chordality of \mathcal{F} , we have $(x_p, x_q) \in G(\mathcal{F})$.

In the case when $k < i$, we have $(x_p, x_q) \in G(\mathcal{P}'^{(k)})$ with $\mathcal{P}'^{(k)} = \mathcal{P}^{(k)} \cup \mathcal{R}^{(k)}$. If $P \in \mathcal{P}^{(k)}$, then it is obvious that $(x_p, x_q) \in G(\mathcal{P}^{(k)}) \subset G(\mathcal{F})$; otherwise if $P \in \mathcal{R}^{(k)}$, then $x_p, x_q \in \text{supp}(R^{(k)}) \subset \text{supp}(\mathcal{P}^{(i)})$ and thus $(x_p, x_i), (x_q, x_i) \in G(\mathcal{P}) \subset G(\mathcal{F})$, then by the chordality of \mathcal{F} , we have $(x_p, x_q) \in G(\mathcal{F})$. \square

Proposition 4.2 *Let $(\mathcal{P}, \mathcal{Q}, i)$ be any node in the binary decomposition tree for Wang's method and T be a polynomial in $\mathcal{P}^{(i)}$ with minimal degree in x_i . Denote $\mathcal{P}'' = \mathcal{P} \setminus \{T\} \cup \{\text{ini}(T), \text{tail}(T)\}$. Then $G(\mathcal{P}'') \subset G(\mathcal{P})$. In particular, if $\text{supp}(\text{tail}(T)) = \text{supp}(T)$, then $G(\mathcal{P}'') = G(\mathcal{P})$.*



$$\begin{aligned}\mathcal{P}' &= \mathcal{P} \setminus \mathcal{P}^{(i)} \cup \{T\} \cup \{\text{prem}(P, T) : P \in \mathcal{P}^{(i)}\}, & \mathcal{Q}' &= \mathcal{Q} \cup \{\text{ini}(T)\} \\ \mathcal{P}'' &= \mathcal{P} \setminus \{T\} \cup \{\text{ini}(T), \text{tail}(T)\}, & \mathcal{Q}'' &= \mathcal{Q}\end{aligned}$$

Figure 5: Splitting at one node $(\mathcal{P}, \mathcal{Q}, i)$ in the binary decomposition tree

Proof Since \mathcal{P}'' is constructed by replacing T in \mathcal{P} with $\text{ini}(T)$ and $\text{tail}(T)$, we only need to study the differences between $G(\mathcal{P})$ and $G(\mathcal{P}'')$ caused by this replacement. First, by $\text{supp}(\text{ini}(T)) \cup \text{supp}(\text{tail}(T)) \subset \text{supp}(T)$ we have $\text{supp}(\mathcal{P}'') \subset \text{supp}(\mathcal{P})$. Second, for any edge (x_p, x_q) in $G(\text{ini}(T))$ or in $G(\text{tail}(T))$, we have $(x_p, x_q) \in G(T)$, which means that all the edges of $G(\mathcal{P}'')$ are also edges of $G(\mathcal{P})$. Therefore, $G(\mathcal{P}'') \subset G(\mathcal{P})$.

In particular, if $\text{supp}(\text{tail}(T)) = \text{supp}(T)$, then $\text{supp}(\text{ini}(T)) \cup \text{supp}(\text{tail}(T)) = \text{supp}(T)$ and any edge $(x_p, x_q) \in \text{supp}(T)$ is also contained in $G(\text{tail}(T))$, and thus $G(\mathcal{P}'') = G(\mathcal{P})$. \square

Example 4.1 Let

$$\begin{aligned}\mathcal{P}_1 &= [x_1 + x_2, x_1 + x_3, x_2 + x_3, x_4^3 + x_1, x_3x_4^2 + x_3 + x_4], \\ \mathcal{P}_2 &= [x_1 + x_2, x_1 + x_3, x_2 + x_3, x_4^3 + x_1, x_3x_4^2 + x_4].\end{aligned}$$

Then $G(\mathcal{P}_1) = G(\mathcal{P}_2)$ is shown in Figure 6 below (left). Let \mathcal{P}_1'' and \mathcal{P}_2'' be constructed from \mathcal{P}_1 and \mathcal{P}_2 with respect to x_4 respectively. Then $x_3x_4^2 + x_3 + x_4$ and $x_3x_4^2 + x_4$ are chosen as T respectively and

$$\begin{aligned}\mathcal{P}_1'' &= [x_1 + x_2, x_1 + x_3, x_2 + x_3, x_3, x_4^3 + x_1, x_3 + x_4], \\ \mathcal{P}_2'' &= [x_1 + x_2, x_1 + x_3, x_2 + x_3, x_3, x_4^3 + x_1, x_4].\end{aligned}$$

One may check that $G(\mathcal{P}_1'') = G(\mathcal{P}_1)$ while $G(\mathcal{P}_2'') \neq G(\mathcal{P}_2)$, with $G(\mathcal{P}_2'')$ shown in Figure 6 below (right).

Next we prove that with a chordal input polynomial set, all the polynomials in the nodes of the decomposition tree of Wang's method, and thus all the computed triangular sets, have associated graphs which are subgraphs of that of the input polynomial set.

Theorem 4.3 *Let $\mathcal{F} \subset \mathbb{K}[x]$ be a chordal polynomial set with $x_1 < \dots < x_n$ as one perfect elimination ordering and $(\mathcal{P}, \mathcal{Q}, i)$ be any node in the binary decomposition tree for Wang's method applied to \mathcal{F} . Then $G(\mathcal{P}) \subset G(\mathcal{F})$.*




Figure 6: The associated graphs $G(\mathcal{P}_1) = G(\mathcal{P}_2) = G(\mathcal{P}_1'')$ (left) and $G(\mathcal{P}_2'')$ (right) in Example 4.1

Proof We induce on the depth d of $(\mathcal{P}, \mathcal{Q}, i)$ in the binary decomposition tree. When $d = 1$, then $(\mathcal{P}, \mathcal{Q}, i)$ is a child node of $(\mathcal{F}, \emptyset, n)$, and $G(\mathcal{P}) \subset G(\mathcal{F})$ by Proposition 4.1 if it is a left child node or by Proposition 4.2 otherwise. Now assume that the first polynomial in any node of depth d in the decomposition tree has an associated graph which is a subgraph of $G(\mathcal{F})$. Let $(\mathcal{P}, \mathcal{Q}, i)$ be of depth $d + 1$ and $(\tilde{\mathcal{P}}, \tilde{\mathcal{Q}}, i)$ be its parent of depth d in the decomposition. Then $G(\mathcal{P}) \subset G(\mathcal{F})$ by Proposition 4.1 if $(\mathcal{P}, \mathcal{Q}, i)$ is a left child node or $G(\mathcal{P}) \subset G(\tilde{\mathcal{P}}) \subset G(\mathcal{F})$ by Proposition 4.2 otherwise. This ends the inductive proof. \square

Corollary 4.4 *Let $\mathcal{F} \subset \mathbb{K}[\mathbf{x}]$ be a chordal polynomial set with $x_1 < \dots < x_n$ as one perfect elimination ordering and $\mathcal{T}_1, \dots, \mathcal{T}_r$ be the triangular sets computed by Wang's method applied to \mathcal{F} . Then $G(\mathcal{T}_i) \subset G(\mathcal{F})$ for $i = 1, \dots, r$.*

Proof Straightforward from Theorem 4.3 with the fact that each triangular set \mathcal{T}_j for some i ($1 \leq i \leq r$) is from a node $(\mathcal{T}_i, \mathcal{Q}_i, 0)$ in the decomposition tree such that \mathcal{T}_i contains no non-zero constant. \square

4.3 An illustrative example

Here we illustrate the changes of chordality of polynomials computed in the triangular decomposition via Wang's method applied to

$$\mathcal{F} = \{x_2 + x_1 + 2, (x_2 + 2)x_3 + x_1, (x_3 + x_2)x_4 + x_3 - 1, x_4 + x_2\} \quad (4)$$

in $\mathbb{Q}[x_1, x_2, x_3, x_4]$ for the variable ordering $x_1 < x_2 < x_3 < x_4$. The associated graph $G(\mathcal{F})$ is shown in Figure 7, and one can check that $G(\mathcal{F})$ is chordal with $x_1 < x_2 < x_3 < x_4$ as one perfect elimination ordering.

First $T = (x_3 + x_2)x_4 + x_3 - 1$ is chosen as the polynomial in $\mathcal{F}^{(4)}$ with minimal degree in x_4 , then a new polynomial set $\mathcal{F}' = \{x_2 + x_1 + 2, (x_2 + 2)x_3 + x_1, (x_3 + x_2), x_3 - 1, x_4 + x_2\}$, which corresponds to the right child node for the case $\text{ini}(T) = x_3 + x_2 = 0$ in the binary decomposition tree, is




Figure 7: The associated graph $G(\mathcal{F})$ with \mathcal{F} from (4)

added to Φ for further computation. Psuedo division of $x_4 + x_2$ over T with respect to x_4 is performed to result in

$$\mathcal{P} = \{x_2 + x_1 + 2, (x_2 + 2)x_3 + x_1, (x_2 - 1)x_3 + x_2^2 + 1, (x_3 + x_2)x_4 + x_3 - 1\},$$

and thus the left child node is $(\mathcal{P}, \{x_3 + x_2\}, 3)$ in the binary tree.

Next $T' = ((x_2 + 2)x_3 + x_1)$ is chosen as the polynomial in $\mathcal{P}^{(3)}$ with minimal degree in x_3 , then a new polynomial set $\mathcal{F}'' = \{x_1, x_2 + x_1 + 2, x_2 + 2, (x_2 - 1)x_3 + x_2^2 + 1, (x_3 + x_2)x_4 + x_3 - 1\}$ is added to Φ , and the pseudo division of $(x_2 - 1)x_3 + x_2^2 + 1$ over $(x_2 + 2)x_3 + x_1$ results in

$$\mathcal{P}' = \{x_2 + x_1 + 2, x_2^3 + 2x_2^2 - (x_1 - 1)x_2 + x_1 + 2, (x_2 + 2)x_3 + x_1, (x_3 + x_2)x_4 + x_3 - 1\}$$

and the left node is $(\mathcal{P}', \{x_x + x_2, x_2 + 2\}, 2)$.

At this step $T'' = x_2 + x_1 + 2$ is chosen as the polynomial in $\mathcal{P}'^{(2)}$ with minimal degree in x_2 , then no polynomial set is added to Φ since $\text{ini}(T'') = 1$ and the pseudo-division of $x_2^3 + 2x_2^2 - (x_1 - 1)x_2 + x_1 + 2$ over T'' with respect to x_2 results in the first triangular set

$$\mathcal{T}_1 = [-x_1^3 + x_1^2 + 14x_1 + 16, x_2 + x_1 + 2, (x_2 + 2)x_3 + x_1, (x_3 + x_2)x_4 + x_3 - 1]. \quad (5)$$

With similar treatments on \mathcal{F}' and \mathcal{F}'' in Φ , the other two triangular sets

$$\begin{aligned} \mathcal{T}_2 &= [x_1 + 1, x_2 + 1, x_3 - 1, x_4 + x_2], \\ \mathcal{T}_3 &= [x_1, x_2 + 2, (x_2 - 1)x_3 + x_2^2 + 1, (x_3 + x_2)x_4 + x_3 - 1] \end{aligned} \quad (6)$$

are computed.

The associated graphs of all these three computed triangular sets are shown in Figure 8. One can find that the associated graphs $G(\mathcal{F})$ and $G(\mathcal{T}_1)$ are the same, while $G(\mathcal{T}_2)$ and $G(\mathcal{T}_3)$ are strict subgraphs of $G(\mathcal{F})$.

5 Further remarks on the applications of graphs structures of polynomial sets

5.1 Variable sparsity of polynomial sets

When referring to a polynomial set $\mathcal{F} \subset \mathbb{K}[\mathbf{x}]$ to be sparse, one usually means that the percentage of terms effectively appearing in \mathcal{F} in all the possible




Figure 8: The associated graphs $G(\mathcal{T}_1)$ (left), $G(\mathcal{T}_2)$ (middle), and $G(\mathcal{T}_3)$ (right) with \mathcal{T}_1 from (5) and $\mathcal{T}_2, \mathcal{T}_3$ from (6)

terms in the variables x_1, \dots, x_n up to a certain degree is low. This kind of sparsity for polynomial sets is convenient for the computation of Gröbner bases which is essentially based on reduction with respect to terms. In fact, efficient algorithms for computing Gröbner bases for sparse polynomial sets defined in this way have been proposed, implemented, and analyzed [15].

Instead of terms of polynomials, triangular sets focus on the variables of polynomials. As exploited in [10], sparsity of the polynomial sets with respect to their variables are partially reflected in their associated graphs. To make it precise, let $G(\mathcal{F}) = (V, E)$ be the associated graph of a polynomial set $\mathcal{F} = \{F_1, \dots, F_r\} \subset \mathbb{K}[\mathbf{x}]$. Then the *variable sparsity* $s_v(\mathcal{F})$ of \mathcal{F} can be defined as

$$s_v(\mathcal{F}) = |E| / \binom{2}{|V|},$$

where the denominator is the number of edges of a complete graph composed of $|V|$ vertexes.

Furthermore, the associated graph $G(\mathcal{F})$ can be extended to a weighted one $G^w(\mathcal{F})$ by associating the number $\#\{F \in \mathcal{F} : x_i, x_j \in \text{supp}(F)\}$ to each edge (x_i, x_j) of $G(\mathcal{F})$. Let $G^w(\mathcal{F}) = (V, E)$, with the weight w_e for each $e \in E$, be the weighted associated graph of \mathcal{F} . Then the *weighted variable sparsity* $s_v^w(\mathcal{F})$ of \mathcal{F} can be defined as

$$s_v^w(\mathcal{F}) = \frac{\sum_{e \in E} w_e}{r \cdot \binom{2}{|V|}},$$

where r is the number of polynomials in \mathcal{F} .

5.2 Complexity analysis for triangular decomposition in top-down style

In general, due to the complicated behaviors in the decomposition process, the complexity of triangular decomposition is not as clearly known as that of computation of Gröbner bases [26, 24, 19, 4, 5, 14].

For a graph G , another graph G' is called a *chordal completion* if G' is chordal with G as its subgraph. The *treewidth* of a graph G is defined

to be the minimum of the sizes of the largest cliques in all the possible chordal completions of G . It has been shown that many NP-complete problems related to graphs can be solved efficiently if the graphs have bounded treewidth [1].

As shown by Theorem 4.3, when the input polynomial set is chordal, the associated graphs of all the polynomials in the decomposition process of Wang’s method are subgraphs of the chordal associated graph of the input polynomial set. In other words, the input chordal associated graph imposes some kind of upper bound for all the polynomials in the decomposition process. Furthermore, the complexity of computing Gröbner bases has been analyzed for polynomial systems by using the treewidth of their associated graphs [9]. These comments lead to the hope of refined complexity analysis of triangular decomposition in top-down style, especially on Wang’s method, from the viewpoint of chordal graphs and their treewidth.

Acknowledgements The authors would like to thank Dongming Wang and Diego Cifuentes for helpful discussions on Wang’s methods for triangular decomposition and on chordal graph structures of polynomial sets in triangular decomposition respectively.

References

- [1] Stefan Arnborg and Andrzej Proskurowski. Linear time algorithms for NP-hard problems restricted to partial k -trees. *Discrete Appl. Math.*, 23(1):11–24, 1989.
- [2] Philippe Aubry, Daniel Lazard, and Marc Moreno Maza. On the theories of triangular sets. *J. Symbolic Comput.*, 28(1–2):105–124, 1999.
- [3] Philippe Aubry and Marc Moreno Maza. Triangular sets for solving polynomial systems: A comparative implementation of four methods. *J. Symbolic Comput.*, 28(1–2):125–154, 1999.
- [4] Magali Bardet, Jean-Charles Faugère, and Bruno Salvy. On the complexity of Gröbner basis computation of semi-regular overdetermined algebraic equations. In *International Conference on Polynomial System Solving - ICPSS*, pages 71 –75, 2004.
- [5] Magali Bardet, Jean-Charles Faugère, Bruno Salvy, and Pierre-Jean Spaenlehauer. On the complexity of solving quadratic Boolean systems. *J. Complexity*, 29(1):53–75, 2013.
- [6] Bruno Buchberger. *Ein Algorithmus zum Auffinden der Basiselemente des Restklassenrings nach einem nulldimensionalen Polynomideal*. PhD thesis, Universität Innsbruck, Austria, 1965.

- [7] Fengjuan Chai, Xiao-Shan Gao, and Chunming Yuan. A characteristic set method for solving Boolean equations and applications in cryptanalysis of stream ciphers. *J. Syst. Sci. Complex.*, 21(2):191–208, 2008.
- [8] Changbo Chen and Marc Moreno Maza. Algorithms for computing triangular decompositions of polynomial systems. *J. Symbolic Comput.*, 47(6):610–642, 2012.
- [9] Diego Cifuentes and Pablo A Parrilo. Exploiting chordal structure in polynomial ideals: A Gröbner bases approach. *SIAM J. Discrete Math.*, 30(3):1534–1570, 2016.
- [10] Diego Cifuentes and Pablo A Parrilo. Chordal networks of polynomial ideals. *SIAM J. Appl. Algebra Geom.*, 1(1):73–110, 2017.
- [11] David A. Cox, John B. Little, and Donal O’Shea. *Using Algebraic Geometry*. Springer Verlag, 1998.
- [12] Jean-Charles Faugère. A new efficient algorithm for computing Gröbner bases (F_4). *J. Pure Appl. Algebra*, 139(1–3):61–88, 1999.
- [13] Jean-Charles Faugère and Antoine Joux. Algebraic cryptanalysis of hidden field equation (HFE) cryptosystems using Gröbner bases. In Dan Boneh, editor, *Advances in Cryptology – CRYPTO 2003*, pages 44–60. Springer, 2003.
- [14] Jean-Charles Faugère and Chenqi Mou. Sparse fglm algorithms. *J. Symbolic Comput.*, 80(3):538–569, 2017.
- [15] Jean-Charles Faugère, Pierre-Jean Spaenlehauer, and Jules Svartz. Sparse Gröbner bases: The unmixed case. In Katsusuke Nabeshima and Kosaku Nagasaka, editors, *Proceedings of ISSAC 2014*, pages 178–185. ACM, 2014.
- [16] Xiao-Shan Gao and Shang-Ching Chou. Solving parametric algebraic systems. In Paul Wang, editor, *Proceedings of ISSAC 1992*, pages 335–341. ACM, 1992.
- [17] Xiao-Shan Gao and Zhenyu Huang. Characteristic set algorithms for equation solving in finite fields. *J. Symbolic Comput.*, 47(6):655–679, 2012.
- [18] John R. Gilbert. Predicting structure in sparse matrix computations. *SIAM J. Matrix Anal. Appl.*, 15(1):62–79, 1994.
- [19] Zhenyu Huang, Yao Sun, and Dongdai Lin. On the efficiency of solving Boolean polynomial systems with the characteristic set method. *arXiv preprint arXiv:1405.4596*, 2014.

- [20] Meng Jin, Xiaoliang Li, and Dongming Wang. A new algorithmic scheme for computing characteristic sets. *J. Symbolic Comput.*, 50:431–449, 2013.
- [21] Michael Kalkbrener. A generalized Euclidean algorithm for computing triangular representations of algebraic varieties. *J. Symbolic Comput.*, 15(2):143–167, 1993.
- [22] Xiaoliang Li, Chenqi Mou, and Dongming Wang. Decomposing polynomial sets into simple sets over finite fields: The zero-dimensional case. *Comput. Math. Appl.*, 60(11):2983–2997, 2010.
- [23] Seymour Parter. The use of linear graphs in Gauss elimination. *SIAM Rev.*, 3(2):119–130, 1961.
- [24] Adrien Poteaux and Éric Schost. On the complexity of computing with zero-dimensional triangular sets. *J. Symbolic Comput.*, 50:110–138, 2013.
- [25] Donald J. Rose. Triangulated graphs and the elimination process. *J. Math. Anal. Appl.*, 32(3):597–609, 1970.
- [26] Agnes Szanto. *Computation with Polynomial Systems*. PhD thesis, Cornell University, USA, 1999.
- [27] Dongming Wang. An elimination method for polynomial systems. *J. Symbolic Comput.*, 16(2):83–114, 1993.
- [28] Dongming Wang. Decomposing polynomial systems into simple systems. *J. Symbolic Comput.*, 25(3):295–314, 1998.
- [29] Dongming Wang. Computing triangular systems and regular systems. *J. Symbolic Comput.*, 30(2):221–236, 2000.
- [30] Dongming Wang. *Elimination Methods*. Springer-Verlag, Wien, 2001.
- [31] Wen-Tsun Wu. On zeros of algebraic equations: An application of Ritt principle. *Kexue Tongbao*, 31(1):1–5, 1986.
- [32] Wen-Tsun Wu. *Mechanical Theorem Proving in Geometries: Basic Principles*. Springer-Verlag, Wien [Translated from the Chinese by X. Jin and D. Wang], 1994.