

# Comment on “Phase Control of Directed Diffusion in a Symmetric Optical Lattice”

Ricardo Chacón

*Departamento de Física Aplicada, E.I.I., Universidad de Extremadura, Apartado Postal 382,  
E-06006 Badajoz, Spain, and Instituto de Computación Científica Avanzada (ICCAEEx),  
Universidad de Extremadura, E-06006 Badajoz, Spain*

(Dated: March 15, 2022)

PACS numbers:

Motivated by investigation into the mechanisms that yield directed diffusion in a symmetric periodic potential, Schiavoni *et al.* [1] have studied experimentally and numerically cold atoms in a one-dimensional dissipative optical lattice where directed motion appears as a result of the breaking of the system’s temporal symmetry after applying a biharmonic phase modulation to one of the lattice beams. In the accelerated reference frame, the atoms experience a stationary optical potential together with an inertial force

$$F(t) = \gamma [A \cos(\omega t) + B \cos(2\omega t - \phi)], \quad (1)$$

where  $\gamma$  is an amplitude factor,  $A = 1 - B$ , and the parameters  $B \in [0, 1]$  and  $\phi \in [0, 2\pi]$  account for the relative amplitude and initial phase difference of the two harmonics, respectively. Commenting on their experimental results, the authors claim that: “By increasing  $B$  from the zero value the atoms are set into directed motion, and a maximum for the c.m. velocity is reached for  $B \simeq 0.5$ , i.e., for about equal amplitudes of the even and odd harmonics.” This statement has had the unfortunate consequence that most of the subsequent published papers citing Ref. [1] have considered  $B = 1/2$  as the condition that maximizes the ratchet transport in systems subjected to a biharmonic temporal force.

This Comment will question the above statement. I shall argue that the maximum c.m. velocity is reached for  $B = 1/3$  as predicted by the theory of ratchet universality [2,3]. Indeed, it has been demonstrated for temporal and spatial biharmonic forces that optimal enhancement of directed ratchet transport is achieved when maximally effective (i.e., critical) symmetry breaking occurs, which implies the existence of a particular universal waveform [2,3]. Specifically, the optimal value of the relative amplitude  $B$  comes from the condition that the amplitude of the odd harmonic must be twice that of the even harmonic in Eq. (1), i.e.,  $1 - B_{\text{opt}} = 2B_{\text{opt}} \implies B_{\text{opt}} = 1/3$ . This condition has been experimentally confirmed in the context of a Bose-Einstein condensate exposed to a

sawtooth-like optical lattice potential [4]. Notice that this means that the contributions of the amplitudes of the two harmonics to the directed motion of the atoms are *not* independent, which is solely taken into account in the estimate of the optical pumping rate (escape rate)  $\Gamma' \propto \sin^2 k\Delta z$ , with  $\Delta z \propto A^2 B$  being the displacement of the centre of oscillation of the atoms in a potential well from the well centre [1], *after* the substitution  $A = 1 - B$ . In such a case, one obtains that  $\Gamma' = \Gamma'(B)$  presents a single maximum at  $B = 1/3$  for which the asymmetry between the escape rates towards the left and right wells is maximal, and hence a maximal nonzero current of atoms is expected again for  $B = 1/3$ , as is indeed confirmed by the experimental results [1] (see Fig. 1). Ratchet universality predicts that the strength of the nonzero current has the functional dependence  $\sim S(B)p(\phi)$  [3], where  $S(B)$  accounts for the degree of breakage of the shift symmetry  $F(t + T/2) = -F(t)$ , while the  $2\pi$ -periodic function  $p(\phi)$  accounts for the degree of breakage of the time-reversal symmetry  $F(-t) = F(t)$  and presents two extrema at the optimal values  $\phi_{\text{opt}} = \{\pi/2, 3\pi/2\}$ . This dependence on  $\phi$  is indeed confirmed by the experimental results shown in Fig. 2 of Ref. [1]. Also,  $S(B)$  presents features similar to those of the impulse,  $I[F] \equiv \left| \int_{T/2} F(t) dt \right|$ , transmitted by the normalized version of the biharmonic force (1),  $F^*(t)$ , for *any* value of  $\phi$  (see Ref. [3] for additional details). Figure 1 shows plots of the function  $S(B)$  for two limiting cases of the initial phase difference: one of the optimal values ( $\phi = \pi/2$ ) and one of the least favourable values ( $\phi = 0$ ) [5]. These curves fit the experimental data reasonably well, and present a single maximum at  $B = 1/3$ , as expected [2,3]. The importance of the results of Ref. [1] is that they provide a first experimental proof of ratchet universality in the context of cold atoms in optical lattices.

The author thanks F. Renzoni and P. J. Martínez for their help with the experimental data recovery and interchanges about this issue.

- 
- [1] M. Schiavoni, L. Sánchez-Palencia, F. Renzoni, and G. Grynberg, Phys. Rev. Lett. **90**, 094101 (2003).  
[2] R. Chacón, J. Phys. A: Math. Theor. **40**, F413 (2007).  
[3] R. Chacón, J. Phys. A: Math. Theor. **43**, 322001 (2010).

- [4] T. Salger *et al.*, Science **326**, 1241 (2009).  
[5] The corresponding analytical expressions for  $S_{\phi=0, \pi/2}$  can be obtained straightforwardly from the respective expressions in Ref. [3] after the substitution  $\eta \rightarrow 1 - B$ .

### A. Figure Captions

Fig. 1 (colour online). Velocity of the centre of mass of the atomic cloud for  $\phi = \pi/2$  (experimental data from Fig. 3 in Ref. [1]; dots), and the curves

$S_{\phi=0} = 1.95I \left[ F_{\phi=0}^* \right]$  (dashed line) and  $S_{\phi=\pi/2} = 6.6 \left( I \left[ F_{\phi=\pi/2}^* \right] - 1 \right)$  (solid line) as functions of the relative amplitude  $B$  [see the text; Eq. (1)].

