

DISJOINTNESS OF CONTINUOUS G-FRAMES AND RIESZ-TYPE CONTINUOUS G-FRAMES

Y. KHEDMATI AND M. R. ABDOLLAHPOUR*

ABSTRACT. In this paper we introduce concepts of disjoint, strongly disjoint and weakly disjoint continuous g -frames in Hilbert spaces and we get some equivalent conditions to these notions. We also construct a continuous g -frame by disjoint continuous g -frames. Furthermore, we provide some results related to the Riesz-type continuous g -frames.

1. INTRODUCTION

In 1952, the concept of frames for Hilbert spaces was defined by Duffin and Schaeffer [6]. Frames are important tools in the signal processing, image processing, data compression, etc. Let \mathcal{H} be a separable Hilbert space. We call a sequence $F = \{f_i\}_{i \in I} \subseteq \mathcal{H}$ a frame for \mathcal{H} if there exist two constant $A_F, B_F > 0$ such that

$$(1.1) \quad A_F \|f\|^2 \leq \sum_{i \in I} |\langle f, f_i \rangle|^2 \leq B_F \|f\|^2, \quad f \in \mathcal{H}.$$

If in (1.1), $A_F = B_F = 1$ we say that $F = \{f_i\}_{i \in I}$ is a Parseval frame for \mathcal{H} . Let $F = \{f_i\}_{i \in I}$ be a frame for \mathcal{H} , then the operator

$$T_F : l_2(I) \rightarrow \mathcal{H}, \quad T_F(\{c_i\}_{i \in I}) = \sum_{i \in I} c_i f_i,$$

is well define and onto, also its adjoint is

$$T_F^* : \mathcal{H} \rightarrow l_2(I), \quad T_F^* f = \{\langle f, f_i \rangle\}_{i \in I}.$$

The operators T_F and T_F^* are called the synthesis and analysis operators of frame F . The concepts of disjoint frames and strongly disjoint frames introduced by Han and Larson [10].

Definition 1.1. Let $F = \{f_i\}_{i \in I}$ and $G = \{g_i\}_{i \in I}$ be frames for Hilbert spaces \mathcal{H} and \mathcal{K} , respectively. We say that F and G are

- (i) Disjoint, if $\{f_i \oplus g_i\}_{i \in I}$ is a frame for $\mathcal{H} \oplus \mathcal{K}$.

MSC(2010): Primary 41A58, 42C15.

Keywords: continuous frame, continuous g -frame, Riesz-type continuous g -frame.

*Corresponding author .

- (ii) Strongly disjoint, if there are invertible operator $L_1 \in B(\mathcal{H})$ and $L_2 \in B(\mathcal{K})$ such that $\{L_1 f_i\}_{i \in I}$, $\{L_2 g_i\}_{i \in I}$ and $\{L_1 f_i \oplus L_2 g_i\}_{i \in I}$ are respective Parseval frames for \mathcal{H} , \mathcal{K} and $\mathcal{H} \oplus \mathcal{K}$.

Proposition 1.2. [10] *Let $F = \{f_i\}_{i \in I}$ and $G = \{g_i\}_{i \in I}$ be frames for Hilbert spaces \mathcal{H} and \mathcal{K} , respectively. Then*

- (i) *F and G are disjoint if and only if $\text{Range}T_F^* \cap \text{Range}T_G^* = \{0\}$ and $\text{Range}T_F^* + \text{Range}T_G^*$ is closed subspace of $l_2(I)$.*
- (ii) *F and G are strongly disjoint if and only if $\text{Range}T_F^*$ and $\text{Range}T_G^*$ are orthogonal.*

In 1993, Ali, Antoine and Gazeau developed the notion of ordinary frame to a family indexed by a measurable space which are known as continuous frames [4].

Definition 1.3. Let \mathcal{H} be a complex Hilbert space and (Ω, μ) be a measure space. The mapping $F : \Omega \rightarrow \mathcal{H}$ is called a continuous frame if

- (i) F is weakly-measurable, i.e., for all $f \in \mathcal{H}$, $\omega \rightarrow \langle f, F(\omega) \rangle$ is a measurable function on Ω ,
- (ii) there exist constants $A_F, B_F > 0$ such that

$$A_F \|f\|^2 \leq \int_{\Omega} |\langle f, F(\omega) \rangle|^2 d\mu(\omega) \leq B_F \|f\|^2, \quad f \in \mathcal{H}.$$

If $F : \Omega \rightarrow \mathcal{H}$ is a continuous frame then the operator $S_F : \mathcal{H} \rightarrow \mathcal{H}$ defined by

$$\langle S_F f, g \rangle = \int_{\Omega} \langle f, F(\omega) \rangle \langle F(\omega), g \rangle d\mu(\omega), \quad f, g \in \mathcal{H},$$

is positive and invertible. S_F is called the continuous frame operator of F .

In 2006, g -frames or generalized frames introduced by Sun [11]. Abdollahpour and Faroughi introduced and investigated continuous g -frames and Riesz-type continuous g -frames [2]. Disjointness notions were developed to continuous frames by Gabardo and Han [7] and to g -frames by Abdollahpour [1]. In the rest of this paper we assume that \mathcal{H} and \mathcal{K} are complex Hilbert spaces and (Ω, μ) is a measure space with positive measure μ and $\{\mathcal{K}_{\omega} : \omega \in \Omega\}$ is a family of Hilbert spaces. Now, we summarize some facts about continuous g -frames from [2].

We say that $F \in \prod_{\omega \in \Omega} \mathcal{K}_{\omega}$ is strongly measurable if F as a mapping of Ω to $\bigoplus_{\omega \in \Omega} \mathcal{K}_{\omega}$ is measurable, where

$$\prod_{\omega \in \Omega} \mathcal{K}_{\omega} = \left\{ f : \Omega \rightarrow \bigcup_{\omega \in \Omega} \mathcal{K}_{\omega} : f(\omega) \in \mathcal{K}_{\omega} \right\}.$$

Definition 1.4. We say that $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a continuous g -frame for \mathcal{H} with respect to $\{\mathcal{K}_\omega : \omega \in \Omega\}$ if

- (i) for each $f \in \mathcal{H}$, $\{\Lambda_\omega f : \omega \in \Omega\}$ is strongly measurable,
- (ii) there are two constants $0 < A_\Lambda \leq B_\Lambda < \infty$ such that

$$(1.2) \quad A_\Lambda \|f\|^2 \leq \int_\Omega \|\Lambda_\omega f\|^2 d\mu(\omega) \leq B_\Lambda \|f\|^2, \quad f \in \mathcal{H}.$$

We call A_Λ, B_Λ the lower and upper continuous g -frame bounds, respectively. Λ is called a tight continuous g -frame if $A_\Lambda = B_\Lambda$, and a Parseval continuous g -frame if $A_\Lambda = B_\Lambda = 1$. If for each $\omega \in \Omega$, $\mathcal{K} = \mathcal{K}_\omega$, then Λ is called a continuous g -frame with respect to \mathcal{K} . $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is called a continuous g -Bessel family if the right hand inequality in (1.2) holds for all $f \in \mathcal{H}$. In this case, B_Λ is called the Bessel constant.

If there is no confusion, we use continuous g -frame (continuous g -Bessel family) instead of continuous g -frame for \mathcal{H} with respect to $\{\mathcal{K}_\omega : \omega \in \Omega\}$ (continuous g -Bessel family for \mathcal{H} with respect to $\{\mathcal{K}_\omega : \omega \in \Omega\}$).

Proposition 1.5. [2] *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be a continuous g -frame. Then there exists a unique positive and invertible operator $S_\Lambda : \mathcal{H} \rightarrow \mathcal{H}$ such that*

$$\langle S_\Lambda f, g \rangle = \int_\Omega \langle f, \Lambda_\omega^* \Lambda_\omega g \rangle d\mu(\omega), \quad f, g \in \mathcal{H},$$

and $A_\Lambda I \leq S_\Lambda \leq B_\Lambda I$.

The operator S_Λ in Proposition 1.5 is called the continuous g -frame operator of Λ . Also, we have

$$(1.3) \quad \langle f, g \rangle = \int_\Omega \langle S_\Lambda^{-1} f, \Lambda_\omega^* \Lambda_\omega g \rangle d\mu(\omega) = \int_\Omega \langle f, \Lambda_\omega^* \Lambda_\omega S_\Lambda^{-1} g \rangle d\mu(\omega),$$

for all $f, g \in \mathcal{H}$.

We consider the space

$$\widehat{\mathcal{K}} = \left\{ F \in \prod_{\omega \in \Omega} \mathcal{K}_\omega : F \text{ is strongly measurable, } \int_\Omega \|F(\omega)\|^2 d\mu(\omega) < \infty \right\}.$$

It is clear that $\widehat{\mathcal{K}}$ is a Hilbert space with point wise operations and with the inner product given by

$$\langle F, G \rangle = \int_\Omega \langle F(\omega), G(\omega) \rangle d\mu(\omega), \quad F, G \in \widehat{\mathcal{K}}.$$

Proposition 1.6. [2] *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be a continuous g -Bessel family. Then the mapping $T_\Lambda : \widehat{\mathcal{K}} \rightarrow \mathcal{H}$ defined by*

$$(1.4) \quad \langle T_\Lambda F, g \rangle = \int_\Omega \langle \Lambda_\omega^* F(\omega), g \rangle d\mu(\omega), \quad F \in \widehat{\mathcal{K}}, \quad g \in \mathcal{H},$$

is linear and bounded with $\|T_\Lambda\| \leq \sqrt{B_\Lambda}$. Also, for each $g \in \mathcal{H}$ and $\omega \in \Omega$,

$$(T_\Lambda^* g)(\omega) = \Lambda_\omega g.$$

Theorem 1.7. [2] *Let (Ω, μ) be a measure space, where μ is σ -finite. Suppose that $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a family of operators such $\{\Lambda_\omega f : \omega \in \Omega\}$ is strongly measurable, for each $f \in \mathcal{H}$. Then Λ is a continuous g -frame if and only if the operator $T_\Lambda : \widehat{\mathcal{K}} \rightarrow \mathcal{H}$ defined by (1.4) is bounded and onto.*

The operators T_Λ and T_Λ^* in Theorem 1.7 are called the synthesis and analysis operators of Λ , respectively.

Definition 1.8. Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two continuous g -frames such that

$$\langle f, g \rangle = \int_\Omega \langle f, \Theta_\omega^* \Lambda_\omega g \rangle d\mu(\omega), \quad f, g \in \mathcal{H},$$

then Θ is called a dual continuous g -frame of Λ .

Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be a continuous g -frame. Then $\tilde{\Lambda} = \{\Lambda_\omega S_\Lambda^{-1} \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a continuous g -frame and by (1.3), $\tilde{\Lambda}$ is a dual of Λ and we call $\tilde{\Lambda}$ the canonical dual of Λ . One can always get a tight continuous g -frame from any continuous g -frame, in fact, if $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a continuous g -frame then $\{\Lambda_\omega S_\Lambda^{-1/2} \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a Parseval continuous g -frame.

Two continuous g -Bessel families $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ are weakly equal, if for all $f \in \mathcal{H}$,

$$\Lambda_\omega f = \Theta_\omega f, \quad a.e. \quad \omega \in \Omega.$$

If the continuous g -frame $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ have only one dual (weakly), i.e., every dual of Λ is weakly equal to the canonical dual of Λ , then Λ is called a Riesz-type continuous g -frame.

Theorem 1.9. [2] *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be a continuous g -frame. Then Λ is a Riesz-type continuous g -frame if and only if $\text{Range } T_\Lambda^* = \widehat{\mathcal{K}}$.*

We mention that the authors of this paper studied some properties of continuous g -frames and Riesz-type continuous g -frames in [3].

2. DISJOINTNESS OF CONTINUOUS g -FRAMES

In this section we study disjointness, strongly disjointness, weakly disjointness for continuous g -frames. We prove some results concern with these concepts and we construct a continuous g -frame by disjoint and strongly disjoint continuous g -frames.

Definition 2.1. Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two continuous g -frames. Then Λ and Θ are called:

- (i) Strongly disjoint, if $\text{Range}T_\Lambda^* \perp \text{Range}T_\Theta^*$.
- (ii) Disjoint, if $\text{Range}T_\Lambda^* \cap \text{Range}T_\Theta^* = \{0\}$ and $\text{Range}T_\Lambda^* + \text{Range}T_\Theta^*$ is a closed subspace of $\widehat{\mathcal{K}}$.
- (iii) Complementary pair, if $\text{Range}T_\Lambda^* \cap \text{Range}T_\Theta^* = \{0\}$ and $\text{Range}T_\Lambda^* + \text{Range}T_\Theta^* = \widehat{\mathcal{K}}$.
- (iv) Strongly complementary pair, if $\text{Range}T_\Lambda^* \oplus \text{Range}T_\Theta^* = \widehat{\mathcal{K}}$.
- (v) Weakly disjoint, if $\text{Range}T_\Lambda^* \cap \text{Range}T_\Theta^* = \{0\}$.

Theorem 2.2. Let (Ω, μ) be a measure space. Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two continuous g -frames. Consider $\Gamma = \{\Gamma_\omega \in B(\mathcal{H} \oplus \mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ where

$$\Gamma_\omega(h \oplus k) = \Lambda_\omega h + \Theta_\omega k, \quad \omega \in \Omega, h \in \mathcal{H}, k \in \mathcal{K}.$$

Then Λ and Θ are

- (i) Strongly disjoint if and only if there exist invertible operators $L_1 \in B(\mathcal{H})$ and $L_2 \in B(\mathcal{K})$ such that $\{\Lambda_\omega L_1 \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$, $\{\Theta_\omega L_2 \in B(\mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\{\Delta_\omega \in B(\mathcal{H} \oplus \mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ are Parseval continuous g -frames, where

$$\Delta_\omega(h \oplus k) = \Lambda_\omega L_1 h + \Theta_\omega L_2 k, \quad \omega \in \Omega, h \in \mathcal{H}, k \in \mathcal{K}.$$

- (ii) Disjoint if and only if Γ is a continuous g -frame.
- (iii) Complementary pair if and only if Γ is a Riesz-type continuous g -frame.
- (iv) Strongly complementary pair if and only if they are strongly disjoint and Γ is a Riesz-type continuous g -frame.
- (v) Weakly disjoint if and only if

$$\{f \oplus g : \Gamma_\omega(f \oplus g) = 0, \omega \in \Omega\} = \{0\}.$$

Proof. (i) For any $h \in \mathcal{H}$ and $k \in \mathcal{K}$ we have

$$\begin{aligned}
& \int_{\Omega} \|\Lambda_{\omega} S_{\Lambda}^{-1/2} h + \Theta_{\omega} S_{\Theta}^{-1/2} k\|^2 d\mu(\omega) \\
&= \int_{\Omega} \|\Lambda_{\omega} S_{\Lambda}^{-1/2} h\|^2 d\mu(\omega) + \int_{\Omega} \|\Theta_{\omega} S_{\Theta}^{-1/2} k\|^2 d\mu(\omega) \\
&+ 2\operatorname{Re} \int_{\Omega} \langle \Lambda_{\omega} S_{\Lambda}^{-1/2} h, \Theta_{\omega} S_{\Theta}^{-1/2} k \rangle d\mu(\omega) \\
&= \int_{\Omega} \|\Lambda_{\omega} S_{\Lambda}^{-1/2} h\|^2 d\mu(\omega) + \int_{\Omega} \|\Theta_{\omega} S_{\Theta}^{-1/2} k\|^2 d\mu(\omega) \\
&= \|h\|^2 + \|k\|^2 = \|h \oplus k\|^2.
\end{aligned}$$

It is sufficient to take $L_1 = S_{\Lambda}^{-1/2}$ and $L_2 = S_{\Theta}^{-1/2}$.

Conversly, for every $h \in \mathcal{H}$ and $k \in \mathcal{K}$ we have

$$\begin{aligned}
\|h \oplus k\|^2 &= \int_{\Omega} \|\Delta_{\omega}(h \oplus k)\|^2 d\mu(\omega) \\
&= \int_{\Omega} \|\Lambda_{\omega} L_1 h\|^2 d\mu(\omega) + \int_{\Omega} \|\Theta_{\omega} L_2 k\|^2 d\mu(\omega) \\
&+ 2\operatorname{Re} \int_{\Omega} \langle \Lambda_{\omega} L_1 h, \Theta_{\omega} L_2 k \rangle d\mu(\omega) \\
&= \|h\|^2 + \|k\|^2 + 2\operatorname{Re} \int_{\Omega} \langle \Lambda_{\omega} L_1 h, \Theta_{\omega} L_2 k \rangle d\mu(\omega).
\end{aligned}$$

Therefore

$$\operatorname{Re} \int_{\Omega} \langle \Lambda_{\omega} L_1 h, \Theta_{\omega} L_2 k \rangle d\mu(\omega) = 0, \quad h \in \mathcal{H}, k \in \mathcal{K}.$$

On the other hand

$$\operatorname{Im} \int_{\Omega} \langle \Lambda_{\omega} L_1 h, \Theta_{\omega} L_2 k \rangle d\mu(\omega) = -\operatorname{Re} \int_{\Omega} \langle \Lambda_{\omega} L_1(ih), \Theta_{\omega} L_2 k \rangle d\mu(\omega) = 0.$$

Thus

$$\int_{\Omega} \langle \Lambda_{\omega} L_1 h, \Theta_{\omega} L_2 k \rangle d\mu(\omega) = 0, \quad h \in \mathcal{H}, k \in \mathcal{K}.$$

Now, since L_1 and L_2 are invertible operators, $\operatorname{Range} T_{\Lambda}^* \perp \operatorname{Range} T_{\Theta}^*$.

(ii) Let Γ be a continuous g -frame for $\mathcal{H} \oplus \mathcal{K}$. Then for any $h \in \mathcal{H}, k \in \mathcal{K}$ we have

$$(2.1) \quad A_{\Gamma}(\|h\|^2 + \|k\|^2) \leq \|T_{\Lambda}^* h + T_{\Theta}^* k\|^2 \leq B_{\Gamma}(\|h\|^2 + \|k\|^2).$$

Let there exist $h_1 \in \mathcal{H}$ and $k_1 \in \mathcal{K}$ such that $T_{\Lambda}^* h_1 = T_{\Theta}^* k_1$. By the left hand inequality (2.1), $h_1 = k_1 = 0$ and so, $T_{\Lambda}^* h_1 = T_{\Theta}^* k_1 = 0$. Consequently, $\operatorname{Range} T_{\Lambda}^* \cap \operatorname{Range} T_{\Theta}^* = \{0\}$. Also, $\operatorname{Range} T_{\Lambda}^* + \operatorname{Range} T_{\Theta}^* =$

$RangeT_\Gamma^*$ is a closed subspace of $\widehat{\mathcal{K}}$.
Conversely, the operator

$$\begin{aligned} L : RangeT_\Lambda^* \oplus RangeT_\Theta^* &\rightarrow RangeT_\Lambda^* + RangeT_\Theta^*, \\ L(F \oplus G) &= F + G \end{aligned}$$

is a bijective bounded operator, In fact

$$\begin{aligned} \|L(F \oplus G)\|^2 &= \|F + G\|^2 \leq (\|F\| + \|G\|)^2 \\ &\leq 2(\|F\|^2 + \|G\|^2) = 2\|F \oplus G\|^2. \end{aligned}$$

Then for any $h \in \mathcal{H}$, $k \in \mathcal{K}$ we have

$$\begin{aligned} \|L^{-1}\|^{-2} \cdot \min\{A_\Lambda, A_\Theta\} \cdot \|h \oplus k\|^2 &\leq \int_\Omega \|\Gamma_\omega(h \oplus k)\|^2 d\mu(\omega) \\ &= \|L(T_\Lambda^*h \oplus T_\Theta^*k)\|^2 \\ &\leq \|L\|^2 \cdot \max\{B_\Lambda, B_\Theta\} \cdot \|h \oplus k\|^2. \end{aligned}$$

(iii) Let Γ be a Riesz-type continuous g -frame for $\mathcal{H} \oplus \mathcal{K}$. Then by Theorem 1.9, we have

$$RangeT_\Lambda^* + RangeT_\Theta^* = RangeT_\Gamma^* = \widehat{\mathcal{K}}.$$

On the other hand, let $\phi \in RangeT_\Lambda^* \cap RangeT_\Theta^*$. Then there exist $h \in \mathcal{H}$ and $k \in \mathcal{K}$ such that $T_\Lambda^*h = T_\Theta^*k = \phi$. Therefore

$$T_\Gamma^*(h \oplus 0) = \phi = T_\Gamma^*(0 \oplus k),$$

since T_Γ^* is one-to-one, $h = k = 0$ and so $\phi = 0$.

Conversely, by the part (ii), Γ is a continuous g -frame and

$$RangeT_\Gamma^* = RangeT_\Lambda^* + RangeT_\Theta^* = \widehat{\mathcal{K}}.$$

So, by Theorem 1.9, Γ is a Riesz-type continuous g -frame.

(iv) By applying the definition of strongly disjoint and (iii), the poof is completed.

(v) Let Λ and Θ be weakly disjoint and $h \in \mathcal{H}, k \in \mathcal{K}$ such that $\Gamma_\omega(h \oplus k) = 0$ for all $\omega \in \Omega$, then

$$T_\Lambda^*h = T_\Theta^*(-k) \in RangeT_\Lambda^* \cap RangeT_\Theta^* = \{0\},$$

and so, $h = k = 0$. Conversely, let $\phi \in RangeT_\Lambda^* \cap RangeT_\Theta^*$. Then there exist $h \in \mathcal{H}, k \in \mathcal{K}$ such that $T_\Lambda^*h = T_\Theta^*k = \phi$, therefore

$$T_\Gamma^*(h \oplus (-k)) = T_\Lambda^*h - T_\Theta^*k = 0.$$

Hence $h = k = 0$, consequently, $\phi = 0$. □

Lemma 2.3. [5] *Suppose that $T : \mathcal{K} \rightarrow \mathcal{H}$ is a linear bounded, surjective operator. Then there exists a linear bounded operator (called the pseudo-inverse of T) $T^\dagger : \mathcal{H} \rightarrow \mathcal{K}$ for which $TT^\dagger f = f$, for any $f \in \mathcal{H}$.*

By generalizing a result from [8] we get a following proposition to construct a continuous g -frame from disjoint continuous g -frames.

Proposition 2.4. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two disjoint continuous g -frames and $L_1, L_2 \in B(\mathcal{H})$. If L_1 or L_2 is surjective, then $\Lambda L_1^* + \Theta L_2^* = \{\Lambda_\omega L_1^* + \Theta_\omega L_2^* \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a continuous g -frame.*

Proof. Let L_1 is surjective, then by Lemma 2.3, there exist $L_1^\dagger \in B(\mathcal{H})$ such that $L_1 L_1^\dagger = I$, so $(L_1^\dagger)^* L_1^* = I$. Hence for any $h \in \mathcal{H}$ we have

$$\|h\| = \|(L_1^\dagger)^* L_1^* h\| \leq \|L_1^\dagger\| \|L_1^* h\|,$$

thus

$$\|L_1^* h\| \geq \frac{\|h\|}{\|L_1^\dagger\|}.$$

Suppose $\Gamma_\omega(h \oplus k) = \Lambda_\omega h + \Theta_\omega k$, for all $h, k \in \mathcal{H}$ and for all $\omega \in \Omega$. By part (ii) of Theorem 2.2, $\Gamma = \{\Gamma_\omega \in B(\mathcal{H} \oplus \mathcal{H}) : \omega \in \Omega\}$ is a continuous g -frame. Therefore, for any $h \in \mathcal{H}$ we have

$$\begin{aligned} \frac{A_\Gamma}{\|L_1^\dagger\|^2} \|h\|^2 &\leq A_\Gamma \|L_1^* h\|^2 \leq A_\Gamma (\|L_1^* h\|^2 + \|L_2^* h\|^2) \\ &= A_\Gamma \|L_1^* h \oplus L_2^* h\|^2 \\ &\leq \int_\Omega \|\Gamma_\omega(L_1^* h \oplus L_2^* h)\|^2 d\mu(\omega) \\ &\leq B_\Gamma \|L_1^* h \oplus L_2^* h\|^2 \\ &\leq B_\Gamma (\|L_1^* h\|^2 + \|L_2^* h\|^2) \\ &\leq 2B_\Gamma \cdot \max\{\|L_1\|^2, \|L_2\|^2\} \|h\|^2. \end{aligned}$$

Therefore we have

$$\begin{aligned} \frac{A_\Gamma}{\|L_1^\dagger\|^2} \|h\|^2 &\leq \int_\Omega \|(\Lambda_\omega L_1 + \Theta_\omega L_2)h\|^2 d\mu(\omega) \\ &\leq 2B_\Gamma \cdot \max\{\|L_1\|^2, \|L_2\|^2\} \|h\|^2, \quad h \in \mathcal{H}. \end{aligned}$$

The proof is similar whenever L_2 is surjective. \square

Corollary 2.5. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two disjoint continuous g -frames. Then $\Lambda + \Theta = \{\Lambda_\omega + \Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a continuous g -frame.*

Proof. By considering $L_1 = L_2 = I$ in Proposition 2.4 the proof is completed. \square

In the following results, we construct a continuous g -frame by strongly disjoint continuous g -frames by generalizing a result from [10].

Proposition 2.6. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be strongly disjoint continuous g -frames and let $L_1, L_2 \in B(\mathcal{H})$ such that $L_1^* L_1 + L_2^* L_2 = AI$ for some $A > 0$. Then $\Lambda L_1 + \Theta L_2 = \{\Lambda_\omega L_1 + \Theta_\omega L_2 \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a continuous g -frame. In particular, $\alpha\Lambda + \beta\Theta = \{\alpha\Lambda_\omega + \beta\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a continuous g -frame for $\alpha, \beta \in \mathbb{C}$ with $|\alpha|^2 + |\beta|^2 > 0$.*

Proof. For any $h \in \mathcal{H}$ we have

$$\begin{aligned} \int_{\Omega} \|(\Lambda_\omega L_1 + \Theta_\omega L_2)h\|^2 d\mu(\omega) &= \|T_\Lambda^* L_1 h + T_\Theta^* L_2 h\|^2 \\ &= \|T_\Lambda^* L_1 h\|^2 + \|T_\Theta^* L_2 h\|^2, \end{aligned}$$

and

$$\begin{aligned} (B_\Lambda \|L_1\|^2 + B_\Theta \|L_2\|^2) \|h\|^2 &\geq \|T_\Lambda^* L_1 h\|^2 + \|T_\Theta^* L_2 h\|^2 \\ &\geq \min\{A_\Lambda, A_\Theta\} (\|L_1 h\|^2 + \|L_2 h\|^2) \\ &= \min\{A_\Lambda, A_\Theta\} \langle (L_1^* L_1 + L_2^* L_2) h, h \rangle \\ &= A \cdot \min\{A_\Lambda, A_\Theta\} \|h\|^2. \end{aligned}$$

By taking $L_1 = \alpha I$ and $L_2 = \beta I$ the particular case is obvious. \square

Proposition 2.7. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be strongly disjoint Parseval continuous g -frames and let $L_1, L_2 \in B(\mathcal{H})$. Then $L_1^* L_1 + L_2^* L_2 = AI$ for some $A > 0$ if and only if $\Lambda L_1 + \Theta L_2 = \{\Lambda_\omega L_1 + \Theta_\omega L_2 \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a tight continuous g -frame with bound A . In particular, $\alpha\Lambda + \beta\Theta = \{\alpha\Lambda_\omega + \beta\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a tight continuous g -frame if and only if $|\alpha|^2 + |\beta|^2 > 0$ for $\alpha, \beta \in \mathbb{C}$.*

Proof. For any $h \in \mathcal{H}$ we have

$$\begin{aligned} \int_{\Omega} \|(\Lambda_\omega L_1 + \Theta_\omega L_2)h\|^2 d\mu(\omega) &= \|T_\Lambda^* L_1 h + T_\Theta^* L_2 h\|^2 \\ &= \|T_\Lambda^* L_1 h\|^2 + \|T_\Theta^* L_2 h\|^2 \\ &= \|L_1 h\|^2 + \|L_2 h\|^2 \\ &= \langle (L_1^* L_1 + L_2^* L_2) h, h \rangle = A \|h\|^2. \end{aligned}$$

In particular, it is sufficient to take $L_1 = \alpha I$ and $L_2 = \beta I$. \square

Corollary 2.8. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be strongly disjoint Parseval continuous g -frames and let $L_1, L_2 \in B(\mathcal{H})$. Then $L_1^* L_1 + L_2^* L_2 = I$ if and only if $\Lambda L_1 + \Theta L_2 = \{\Lambda_\omega L_1 + \Theta_\omega L_2 \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a Parseval continuous g -frame. In particular, $\alpha\Lambda + \beta\Theta = \{\alpha\Lambda_\omega + \beta\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a Parseval continuous g -frame if and only if $|\alpha|^2 + |\beta|^2 = 1$ for $\alpha, \beta \in \mathbb{C}$.*

Now, to get a dual continuous g -frames by strongly disjoint continuous g -frames we generalize results of [1] and [9].

Proposition 2.9. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Psi = \{\Psi_\omega \in B(\mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be duals of continuous g -frames $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Phi = \{\Phi_\omega \in B(\mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$, respectively. If Λ, Φ and Θ, Ψ are strongly disjoint. Then $\Gamma = \{\Gamma_\omega \in B(\mathcal{H} \oplus \mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Delta = \{\Delta_\omega \in B(\mathcal{H} \oplus \mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ are dual continuous g -frames, where*

$$\Gamma_\omega(h \oplus k) = \Lambda_\omega h + \Psi_\omega k, \quad \Delta_\omega(h \oplus k) = \Theta_\omega h + \Phi_\omega k,$$

for all $\omega \in \Omega$ and for all $h \in \mathcal{H}, k \in \mathcal{K}$.

Proof. It is clear that Γ and Δ are continuous g -Bessel families for $\mathcal{H} \oplus \mathcal{K}$. For any $h_1, h_2 \in \mathcal{H}$ and $k_1, k_2 \in \mathcal{K}$, we have

$$\begin{aligned} & \int_{\Omega} \langle \Gamma_\omega(h_1 \oplus k_1), \Delta_\omega(h_2 \oplus k_2) \rangle d\mu(\omega) \\ &= \int_{\Omega} \langle \Lambda_\omega h_1, \Theta_\omega h_2 \rangle d\mu(\omega) + \int_{\Omega} \langle \Lambda_\omega h_1, \Phi_\omega k_2 \rangle d\mu(\omega) \\ &+ \int_{\Omega} \langle \Psi_\omega k_1, \Theta_\omega h_2 \rangle d\mu(\omega) + \int_{\Omega} \langle \Psi_\omega k_1, \Phi_\omega k_2 \rangle d\mu(\omega) \\ &= \langle h_1, h_2 \rangle + \langle k_1, k_2 \rangle \\ &= \langle h_1 \oplus k_1, h_2 \oplus k_2 \rangle. \end{aligned}$$

Thus by Proposition 3.2 of [2], Γ and Δ are dual continuous g -frames for $\mathcal{H} \oplus \mathcal{K}$. \square

Example 2.10. Let $F : \Omega \rightarrow \mathcal{H}$ and $G : \Omega \rightarrow \mathcal{K}$ be two continuous frames. We define two families of bounded operators $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathbb{C}^2) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathbb{C}^2) : \omega \in \Omega\}$ where

$$\Lambda_\omega f = (\langle f, F(\omega) \rangle_{\mathcal{H}}, 0), \quad \Theta_\omega f = (\langle S_F^{-1} f, F(\omega) \rangle_{\mathcal{H}}, 0),$$

for all $f \in \mathcal{H}, \omega \in \Omega$. It is obvious that Λ and Θ are continuous g -frames. Also for any $f, g \in \mathcal{H}$

$$\int_{\Omega} \langle \Theta_\omega f, \Lambda_\omega g \rangle d\mu(\omega) = \int_{\Omega} \langle f, S_F^{-1} F(\omega) \rangle \langle F(\omega), g \rangle d\mu(\omega) = \langle f, g \rangle.$$

So, Λ and Θ are dual continuous g -frames. We also define two families of bounded operators $\Phi = \{\Phi_\omega \in B(\mathcal{K}, \mathbb{C}^2) : \omega \in \Omega\}$ and $\Psi = \{\Psi_\omega \in B(\mathcal{K}, \mathbb{C}^2) : \omega \in \Omega\}$ where

$$\Phi_\omega g = (0, \langle S_G^{-1}g, G(\omega) \rangle_\mathcal{K}), \quad \Psi_\omega g = (0, \langle g, G(\omega) \rangle_\mathcal{K}),$$

for all $g \in \mathcal{K}$, $\omega \in \Omega$. Similarly, Ψ and Φ are dual continuous g -frames. On the other hand, for any $f \in \mathcal{H}$ and $g \in \mathcal{K}$

$$\langle T_\Lambda^* f, T_\Phi^* g \rangle = \int_\Omega \langle \Lambda_\omega f, \Phi_\omega g \rangle d\mu(\omega) = 0,$$

so Λ and Φ are strongly disjoint. Also, Θ and Ψ are strongly disjoint. Let us consider

$$\Gamma_\omega : \mathcal{H} \oplus \mathcal{K} \rightarrow \mathbb{C}^2, \quad \Gamma_\omega(f \oplus g) = (\langle f, F(\omega) \rangle_\mathcal{H}, \langle g, G(\omega) \rangle_\mathcal{K}),$$

and

$$\Delta_\omega : \mathcal{H} \oplus \mathcal{K} \rightarrow \mathbb{C}^2, \quad \Delta_\omega(f \oplus g) = (\langle S_F^{-1}f, F(\omega) \rangle_\mathcal{H}, \langle S_G^{-1}g, G(\omega) \rangle_\mathcal{K}),$$

for all $f \in \mathcal{H}$ and $g \in \mathcal{K}$. Then

$$\begin{aligned} \int_\Omega \langle \Gamma_\omega(h_1 \oplus k_1), \Delta_\omega(h_2 \oplus k_2) \rangle d\mu(\omega) \\ &= \int_\Omega \langle h_1, F(\omega) \rangle_\mathcal{H} \langle S_F^{-1}F(\omega), h_2 \rangle_\mathcal{H} d\mu(\omega) \\ &\quad + \int_\Omega \langle k_1, G(\omega) \rangle_\mathcal{K} \langle S_G^{-1}G(\omega), k_2 \rangle_\mathcal{K} d\mu(\omega) \\ &= \langle h_1, h_2 \rangle + \langle k_1, k_2 \rangle \\ &= \langle h_1 \oplus k_1, h_2 \oplus k_2 \rangle. \end{aligned}$$

Which Proposition 2.9 confirm this result.

Proposition 2.11. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two strongly disjoint continuous g -frames and $L_1, L_2 \in B(\mathcal{H})$. If L_1 is surjective, then $\tilde{\Lambda}L_1^\dagger = \{\Lambda_\omega S_\Lambda^{-1}L_1^\dagger \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a dual for both $\Lambda L_1^* = \{\Lambda_\omega L_1^* \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Lambda L_1^* + \Theta L_2^* = \{\Lambda_\omega L_1^* + \Theta_\omega L_2^* \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$.*

Proof. It is obvious that $\tilde{\Lambda}L_1^\dagger$, ΛL_1^* and $\Lambda L_1^* + \Theta L_2^*$ are continuous g -Bessel families. Since L_1 is surjective, then by Lemma 2.3 there exist

$L_1^t \in B(\mathcal{H}, \mathcal{K}_\omega)$, such that $L_1 L_1^t = I$. For any $h, k \in \mathcal{H}$ we have

$$\begin{aligned}
& \int_{\Omega} \langle (\Lambda_\omega L_1^* + \Theta_\omega L_2^*)h, \Lambda_\omega S_\Lambda^{-1} L_1^\dagger k \rangle d\mu(\omega) \\
&= \int_{\Omega} \langle \Lambda_\omega L_1^* h, \Lambda_\omega S_\Lambda^{-1} L_1^\dagger k \rangle d\mu(\omega) \\
&+ \int_{\Omega} \langle \Theta_\omega L_2^* h, \Lambda_\omega S_\Lambda^{-1} L_1^\dagger k \rangle d\mu(\omega) \\
&= \langle L_1^* h, L_1^\dagger k \rangle + \langle T_\Theta^* L_2^* h, T_\Lambda^* S_\Lambda^{-1} L_1^\dagger k \rangle \\
&= \langle h, L_1 L_1^\dagger k \rangle + 0 = \langle h, k \rangle + 0 = \langle h, k \rangle.
\end{aligned}$$

And also we have

$$\int_{\Omega} \langle \Lambda_\omega L_1^* f, \Lambda_\omega S_\Lambda^{-1} L_1^\dagger g \rangle d\mu(\omega) = \langle L_1^* f, L_1^\dagger g \rangle = \langle f, L_1 L_1^\dagger g \rangle = \langle f, g \rangle.$$

Therefore the proof is completed. \square

Corollary 2.12. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two strongly disjoint continuous g -frames. Then $\tilde{\Lambda} = \{\Lambda_\omega S_\Lambda^{-1} \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a dual for both $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Lambda + \Theta = \{\Lambda_\omega + \Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$.*

Proof. By considering $L_1 = L_2 = I$ in Proposition 2.11 the proof is completed. \square

3. SOME RESULTS RELATED TO RIESZ-TYPE CONTINUOUS g -FRAMES

In this section by generalizing some results of [12], we get some equivalent conditions for Riesz-type continuous g -frames.

Theorem 3.1. *Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be a continuous g -frame. Then the following are equivalent:*

- (i) Λ is a Riesz-type continuous g -frame.
- (ii) There exist constants $A, B > 0$ such that

$$(3.1) \quad A\|\phi\|^2 \leq \|T_\Lambda \phi\|^2 \leq B\|\phi\|^2, \quad \phi \in \widehat{\mathcal{K}}.$$

- (iii) If

$$\int_{\Omega} \langle \Lambda_\omega^* \phi(\omega), f \rangle d\mu(\omega) = 0$$

for some $\phi \in \widehat{\mathcal{K}}$ and for any $f \in \mathcal{H}$, then $\phi = 0$.

Proof. (i) \Rightarrow (ii) By Proposition 1.6, it remains to prove the left-hand inequality in (3.1). By Theorem 1.9, for any $\phi \in \widehat{\mathcal{K}}$, there exist $f \in \mathcal{H}$ such that $T_\Lambda^* f = \phi$. Then

$$\begin{aligned} \|\phi\|^4 &= \left(\int_\Omega \|\Lambda_\omega(f)\|^2 d\mu(\omega) \right)^2 = |\langle S_\Lambda f, f \rangle|^2 \\ &\leq \|S_\Lambda f\|^2 \|f\|^2 \\ &\leq \frac{1}{A_\Lambda} \|S_\Lambda f\|^2 \int_\Omega \|\Lambda_\omega f\|^2 d\mu(\omega), \end{aligned}$$

and hence

$$A_\Lambda \|\phi\|^2 \leq \|S_\Lambda f\|^2 = \|T_\Lambda T_\Lambda^* f\|^2 = \|T_\Lambda \phi\|^2.$$

(ii) \Rightarrow (iii) Let for some $\phi \in \widehat{\mathcal{K}}$ and any $f \in \mathcal{H}$, we have

$$\langle T_\Lambda \phi, f \rangle = \int_\omega \langle \Lambda_\omega^* \phi(\omega), f \rangle d\mu(\omega) = 0.$$

Then $T_\Lambda \phi = 0$ and by inequality (3.1), $\phi = 0$.

(iii) \Rightarrow (i) Since Λ is a continuous g -frame, T_Λ is onto and by (iii) T_Λ is one to one, so T_Λ is invertible. Consequently, T_Λ^* is invertible. Therefore, by Theorem 1.9, the proof is completed. \square

Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{K}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two continuous g -Bessel families. Consider the well defined operator $S_{\Theta\Lambda} : \mathcal{H} \rightarrow \mathcal{K}$, $S_{\Theta\Lambda} = T_\Theta T_\Lambda^*$. Then

$$\langle S_{\Theta\Lambda} f, g \rangle = \int_\Omega \langle \Lambda_\omega f, \Theta_\omega g \rangle d\mu(\omega), \quad f \in \mathcal{H}, g \in \mathcal{K},$$

and $S_{\Theta\Lambda}^* = S_{\Lambda\Theta}$.

Theorem 3.2. Let $\Lambda = \{\Lambda_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ and $\Theta = \{\Theta_\omega \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ be two continuous g -Bessel family such that $S_{\Lambda\Theta} = I_{\mathcal{H}}$. Assume that $L_1, L_2 : \mathcal{H} \rightarrow \mathcal{H}$ are bounded linear operators so that $L_1^* L_2 = I$. Then the following statements are equivalent:

- (i) $\Gamma = \{\Lambda_\omega L_1 + \Theta_\omega L_2 \in B(\mathcal{H}, \mathcal{K}_\omega) : \omega \in \Omega\}$ is a Riesz-type continuous g -frame.
- (ii) The operator $T_\Lambda^* L_1 + T_\Theta^* L_2$ is surjective.
- (iii) There exists a constant $M > 0$ such that

$$M \|\phi\|^2 \leq \|(L_1^* T_\Lambda + L_2^* T_\Theta) \phi\|^2, \quad \phi \in \widehat{\mathcal{K}}.$$

Proof. For any $f \in \mathcal{H}$ we have

$$\begin{aligned}
& \int_{\Omega} \|(\Lambda_{\omega}L_1 + \Theta_{\omega}L_2)f\|^2 d\mu(\omega) \\
&= \int_{\Omega} \|\Lambda_{\omega}L_1f\|^2 d\mu(\omega) + \int_{\Omega} \langle \Lambda_{\omega}L_1f, \Theta_{\omega}L_2f \rangle d\mu(\omega) \\
&+ \int_{\Omega} \langle \Theta_{\omega}L_2f, \Lambda_{\omega}L_1f \rangle d\mu(\omega) + \int_{\Omega} \|\Theta_{\omega}L_2f\|^2 d\mu(\omega) \\
&= \int_{\Omega} \|\Lambda_{\omega}L_1f\|^2 d\mu(\omega) + 2\|f\|^2 + \int_{\Omega} \|\Theta_{\omega}L_2f\|^2 d\mu(\omega).
\end{aligned}$$

So

$$\begin{aligned}
2\|f\|^2 &\leq \int_{\Omega} \|(\Lambda_{\omega}L_1 + \Theta_{\omega}L_2)f\|^2 d\mu(\omega) \\
&\leq (B_{\Lambda}\|L_1\|^2 + 2 + B_{\Theta}\|L_2\|^2)\|f\|^2.
\end{aligned}$$

Hence Γ is a continuous g -frame. On the other hand

$$T_{\Gamma}^* = T_{\Lambda}^*L_1 + T_{\Theta}^*L_2.$$

By Theorem 1.9, (i) and (ii) are equivalent.

(i) \Leftrightarrow (iii) It is concluded by Theorem 3.1 and $T_{\Gamma} = L_1^*T_{\Lambda} + L_2^*T_{\Theta}$. \square

Proposition 3.3. *Let (Ω, μ) be a measure space, that μ is σ -finite and $\Lambda = \{\Lambda_{\omega} \in B(\mathcal{H}, \mathcal{K}_{\omega}) : \omega \in \Omega\}$ is a continuous g -frame. Suppose that $\Theta = \{\Theta_{\omega} \in B(\mathcal{K}, \mathcal{K}_{\omega}) : \omega \in \Omega\}$ is a continuous g -Bessel family. If $S_{\Theta\Lambda}$ is surjective, then Θ is a continuous g -frame. If Θ is a continuous g -frame and Λ is a Riesz-type continuous g -frame then $S_{\Theta\Lambda}$ is surjective.*

Proof. Since $S_{\Theta\Lambda}$ is surjective, it follows that T_{Θ} is surjective. On the other hand, by Proposition 1.6, T_{Θ} is bounded. Hence by Theorem 1.7, Θ is a continuous g -frame.

If Λ is a Riesz-type continuous g -frame and Θ is a continuous g -frame then by Theorems 1.9 and 1.7, T_{Λ}^* and T_{Θ} are surjective. So, $S_{\Theta\Lambda}$ is surjective. \square

Theorem 3.4. *Let $\Lambda = \{\Lambda_{\omega} \in B(\mathcal{H}, \mathcal{K}_{\omega}) : \omega \in \Omega\}$ be a continuous g -frame and $\Theta = \{\Theta_{\omega} \in B(\mathcal{K}, \mathcal{K}_{\omega}) : \omega \in \Omega\}$ be a continuous g -Bessel family. Suppose that there exists a number λ with $0 < \lambda < A_{\Lambda}$ such that*

$$(3.2) \quad \|S_{\Theta\Lambda}f - S_{\Lambda}f\| \leq \lambda\|f\|, \quad f \in \mathcal{H}.$$

Then Λ is a Riesz-type continuous g -frame if and only if Θ is a Riesz-type continuous g -frame.

Proof. For all $f \in \mathcal{H}$ we have

$$\begin{aligned}\|S_{\Theta\Lambda}f\| &= \|S_{\Theta\Lambda}f - S_{\Lambda}f + S_{\Lambda}f\| \geq \|S_{\Lambda}f\| - \|S_{\Theta\Lambda}f - S_{\Lambda}f\| \\ &\geq (A_{\Lambda} - \lambda)\|f\|.\end{aligned}$$

Then $S_{\Theta\Lambda}$ is injective with closed range. On the other hand,

$$\|S_{\Lambda\Theta}f - S_{\Lambda}f\| \leq \|(S_{\Theta\Lambda} - S_{\Lambda})^*\| \|f\| \leq \lambda \|f\|.$$

So $S_{\Lambda\Theta}$ is also injective with closed range. Therefore

$$\text{Range } S_{\Theta\Lambda} = \ker(S_{\Lambda\Theta})^{\perp} = \mathcal{H}$$

and $\text{Range } S_{\Lambda\Theta} = \mathcal{H}$. Thus, $S_{\Theta\Lambda}$ and $S_{\Lambda\Theta}$ are invertible. Hence, T_{Λ}^* is invertible if and only if T_{Θ}^* is invertible. Then by Theorem 1.9 the proof is completed. \square

Acknowledgment:

REFERENCES

- [1] M. R. Abdollahpour, *Dilation of dual g-frames to dual g-Riesz bases*, Banach Journal of Mathematical Analysis **9(1)** (2015), 54-66.
- [2] M. R. Abdollahpour and M. H. Faroughi, *Continuous g-frames in Hilbert spaces*, Southeast Asian Bulletin of Mathematics **32(1)** (2008), 1-19.
- [3] M. R. Abdollahpour and Y. Khedmati, *On some properties of continuous g-frames and Riesz-type continuous g-frames*, Indian Journal of Pure and Applied Mathematics **48(1)** (2017), 59-74.
- [4] S. T. Ali, J. P. Antoine and J. P. Gazeau, *Continuous frames in Hilbert space*, Annals of Physics **222(1)** (1993), 1-37.
- [5] O. Christensen and K. Jensen Torben, *An introduction to the theory of bases, frames, and wavelets*, (1999).
- [6] R. J. Duffin and A. C. Schaeffer, *A class of nonharmonic Fourier series*, Transactions of the American Mathematical Society **72(2)** (1952), 341-366.
- [7] J. P. Gabardo and D. Han, *Frames associated with measurable space*, Advances in Computational Mathematics **18(2)** (2003), 127-147.
- [8] X. Guo, *Constructions of frames by disjoint frames*, Numerical Functional Analysis and Optimization **35(5)** (2014), 576-587.
- [9] X. Guo, *Characterizations of disjointness of g-frames and constructions of g-frames in Hilbert spaces*, Complex Analysis and Operator Theory **8(7)** (2014), 1547-1563.
- [10] D. Han and D. Larson, *Frames, bases and group representations*, Memoirs of the American Mathematical Society **697** (2000), 149-182.
- [11] W. Sun, *G-frames and g-Riesz bases*, Journal of Mathematical Analysis and Applications **322(1)** (2006), 437-452.
- [12] Z. Q. Xiang, *New characterizations of Riesz-type frames and stability of alternate duals of continuous frames*, Advances in Mathematical Physics (2013).

YAVAR KHEDMATI AND MOHAMMAD REZA ABDOLLAHPOUR

DEPARTMENT OF MATHEMATICS

FACULTY OF SCIENCES

UNIVERSITY OF MOHAGHEGH ARDABILI

ARDABIL 56199-11367

IRAN

E-mail address: khedmati.y@uma.ac.ir, khedmatiy.y@gmail.com

E-mail address: m.abdollah@uma.ac.ir, mrabdollahpour@yahoo.com