arXiv:1802.04077v1 [math.FA] 12 Feb 2018

COMPACT OPERATORS ON THE SETS OF FRACTIONAL DIFFERENCE SEQUENCES

FARUK ÖZGER

Abstract. Fractional difference sequence spaces have been studied in the literature recently. In this work, some identities or estimates for the operator norms and the Hausdorff measures of noncompactness of certain operators on some difference sequence spaces of fractional orders are established. Some classes of compact operators on those spaces are characterized. The results of this work are more general and comprehensive then many other studies in literature.

1. Introduction

The most common type of sets of sequences are probably the sets of difference sequences among the sequence spaces studied. The difference sequence spaces first introduced in Kızmaz's study [15]. Many authors have made efforts to investigate the topological structures of these spaces during the past decade (see [4],[6],[12],[13],[18],[19],[23]). Compact operators on the sets of difference sequences have been characterized in ([7],[8],[10],[20]). We also refer to ([1],[2],[3],[5],[9],[11],[21]) for further studies in theory of FK-spaces and its applications. More recently, certain difference sequence spaces of fractional orders have been introduced by Baliarsingh [22]. Certain Euler difference sequence spaces of fractional order and related dual properties have been studied by Kadak and Baliarsingh [24]. Topological properties of certain sequence spaces that are combined by the mean operator and the fractional difference operator are investigated by Furkan [14].

The rest of the paper is organized as follows. In the rest of this section, we consider fractional operators, their properties and fractional sets of sequences $c_0(\Delta^{(\widetilde{\alpha})})$, $c(\Delta^{(\widetilde{\alpha})})$ and $\ell_\infty(\Delta^{(\widetilde{\alpha})})$. In section 2, we will determine the β duals of fractional sets of sequences and characterize matrix transformations on them. We also examine operator norms of our spaces. In section 3, we will study on characterizations of some compact operators by applying Hausdorff measure of noncompactness.

1.1. **Fractional Difference Operators.** The gamma function of a real number x (except zero and the negative integers) is defined by an improper integral:

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt.$$

2000 *Mathematics Subject Classification.* Primary: 40H05; Secondary: 46H05.

Key words and phrases. Fractional difference sequence spaces, compact operators, Hausdorff measure of noncompactness

This work is a part of the research project supported by Izmir Katip Celebi University Scientific Research Project Coordination Unit.

It is known that for any natural number n, $\Gamma(n+1)=n!$ and $\Gamma(n+1)=n\Gamma(n)$ holds for any real number $n\notin\{0,-1,-2,...\}$.

The fractional difference operator for a fraction $\tilde{\alpha}$ have been defined in [22] as

$$\Delta^{(\tilde{\alpha})}(x_k)=\sum_{i=0}^{\infty}(-1)^i\frac{\Gamma(\tilde{\alpha}+1)}{\Gamma(\tilde{\alpha}-i+1)}x_{k-i}. \tag{1.1}$$

It is assumed that the series defined in (1.1) is convergent for $x\in\omega$.

Let m be a positive integer then recall the difference operators $\Delta^{(1)}$ and $\Delta^{(m)}$ by :

$$(\Delta^{(1)}x)_k=\Delta^{(1)}x_k=x_k-x_{k-1}$$

and

$$(\Delta^{(m)}x)_k=\sum_{i=0}^{m}(-1)^i\binom{m}{i}x_{k-i}.$$

We write Δ and $\Delta^{(m)}$ for the matrices with $\Delta_{nk}=(\Delta^{(1)}e^{(k)})_n$ and $\Delta^{(m)}_{nk}=(\Delta^{(m)}e^{(k)})_n$ for all n and k.

We may write the fractional difference operator as an infinite matrix:

$$\Delta^{(\tilde{\alpha})}_{nk}=\begin{cases}(-1)^{n-k}\frac{\Gamma(\tilde{\alpha}+1)}{(n-k)!\Gamma(\tilde{\alpha}-n+k+1)} & (0\leq k\leq n)\\ 0 & (k>n).\end{cases}$$

Remark 1.1. *The inverse of fractional difference matrix is given by*

$$\Delta^{(-\tilde{\alpha})}_{nk}=\begin{cases}(-1)^{n-k}\frac{\Gamma(-\tilde{\alpha}+1)}{(n-k)!\Gamma(-\tilde{\alpha}-n+k+1)} & (0\leq k\leq n)\\ 0 & (k>n).\end{cases}$$

For some values of $\tilde{\alpha}$, we have

$$\begin{aligned}\Delta^{1/2}x_k &= x_k-\frac{1}{2}x_{k-1}-\frac{1}{8}x_{k-2}-\frac{1}{16}x_{k-3}-\frac{5}{128}x_{k-4}-...\\ \Delta^{-1/2}x_k &= x_k+\frac{1}{2}x_{k-1}+\frac{3}{8}x_{k-2}+\frac{5}{16}x_{k-3}+\frac{35}{128}x_{k-4}+...\\ \Delta^{2/3}x_k &= x_k-\frac{2}{3}x_{k-1}-\frac{1}{9}x_{k-2}-\frac{4}{81}x_{k-3}-\frac{7}{243}x_{k-4}-...\end{aligned}$$

Theorem 1.2. *The following results hold:*

(i) $\Delta^{(\tilde{\alpha})}\circ\Delta^{(-\tilde{\alpha})}=I$, *where* I *is identity on* $x\in\omega$.
(ii) $\Delta^{(\tilde{\alpha})}\Delta^{(\tilde{\beta})}=\Delta^{(\tilde{\alpha}+\tilde{\beta})}$.
(iii) $\Delta^{(\tilde{\alpha})}(\Delta^{(-\tilde{\alpha})}x_k)=x_k$.

Proof. Since the proofs of Parts (i) and (ii) can similarly be obtained, we only prove Part (iii).

$$\begin{aligned}\Delta^{(\tilde{\alpha})}(\Delta^{(-\tilde{\alpha})}x_k) &= \Delta^{(\tilde{\alpha})}\left\{x_k + x_{k-1}\alpha + x_{k-2}\tfrac{\alpha(\alpha+1)}{2!} + x_{k-3}\tfrac{\alpha(\alpha+1)(\alpha+2)}{3!} + x_{k-4}\tfrac{\alpha(\alpha+1)(\alpha+2)(\alpha+3)}{4!} + \cdots\right\}\\ &= x_k + x_{k-1}\{-\alpha+\alpha\} + x_{k-2}\left\{\tfrac{\alpha(\alpha-1)}{2!} + \alpha^2 + \tfrac{\alpha(\alpha+1)}{2!}\right\} +\\ &\quad x_{k-3}\left\{-\tfrac{\alpha(\alpha-1)(\alpha-2)}{3!} + \tfrac{\alpha^2(\alpha-1)}{2!} - \tfrac{\alpha^2(\alpha+1)}{2!} + \tfrac{\alpha(\alpha+1)(\alpha+2)}{3!}\right\} +\\ &\quad x_{k-4}\left\{\tfrac{\alpha(\alpha-1)(\alpha-2)(\alpha-3)}{4!} - \tfrac{\alpha^2(\alpha-1)(\alpha-2)}{3!} + \tfrac{\alpha^2(\alpha+1)(\alpha-1)}{2!2!} - \tfrac{\alpha^2(\alpha+1)(\alpha+2)}{3!}\right\} + \cdots\\ &= x_k.\end{aligned}$$

□

We refer to [22] for more properties of the fractional difference operators.

Note that the results of this work are more general and comprehensive then many other studies related to difference sequence spaces in literature.

1.2. **Preliminaries Results and Notations.** For the readers convenience, we state the known results that are used here and in the sequel.

Let ω denote the set of all complex sequences $x = (x_k)_{k=0}^{\infty}$. We write ℓ_∞, c, c_0 and ϕ for the sets of all bounded, convergent, null and finite sequences, respectively; also bs, cs and ℓ_1 denote the sets of all bounded, convergent and absolutely convergent series.

A subspace X of ω is said to be a BK space if it is a Banach space with continuous coordinates $P_n : X \to \mathbb{C}$ $(n = 0, 1, \dots)$, where $P_n(x) = x_n$ for all $x \in X$. A BK space $X \supset \phi$ is said to have AK if every sequence $x = (x_k)_{k=0}^{\infty} \in X$ has a unique representation $x = \lim_{m\to\infty} x^{[m]}$, where $x^{[m]} = \sum_{n=0}^{m} x_n e^{(n)}$ is the m section of the sequence x. Let X be a normed space. Then $S_X = \{x \in X : \|x\| = 1\}$ and $\bar{B}_X = \{x \in X : \|x\| \leq 1\}$ denote the unit sphere and closed unit ball in X, where X is a normed space. By N_r we denote any subset of $\mathbb{N}_0$ with elements greater or equal to r.

Given any infinite matrix $A = (a_{nk})_{n,k=0}^{\infty}$ of complex numbers and any sequence x, we write $A_n = (a_{nk})_{k=0}^{\infty}$ for the sequence in the n^{th} row of A, $A_n x = \sum_{k=0}^{\infty} a_{nk}x_k$ $(n = 0, 1, \dots)$ and $Ax = (A_n x)_{n=0}^{\infty}$, provided $A_n \in x^\beta$ for all n. If X and Y are subsets of ω, then

$$X_A = \{x \in \omega : Ax \in X\}$$

denotes the matrix domain of A in X and (X, Y) is the class of all infinite matrices that map X into Y; so $A \in (X, Y)$ if and only if $X \subset Y_A$.

Given $a \in \omega$, we write

$$\|a\|_X^* = \sup_{x\in S_X}\left|\sum_{k=1}^{\infty} a_k x_k\right|$$

provided the expression on the write hand side is defined and finite which is the case whenever X is a BK space and $a \in X^\beta$ ([2], Theorem 7.2.9, p. 107).

An infinite matrix $T = (t_{nk})_{n,k=0}^{\infty}$ is said to be a triangle if $t_{nk} = 0$ $(k > n)$ and $t_{nn} \neq 0$ for all n.

A sequence $(b_n)_{n=0}^{\infty}$ in a linear metric space X is called a Schauder basis if for each $x \in X$ there exists a unique sequence $(\lambda_n)_{n=0}^{\infty}$ of scalars such that $x = \sum_{n=0}^{\infty} \lambda_n b_n$.

1.3. **The Difference Sequence Spaces of Fractional Order.** Consider now the following fractional difference sequence spaces.

$$\begin{aligned}
c_0(\Delta^{(\tilde{\alpha})}) &:= \left\{ x=(x_k)\in\omega : \lim_k \sum_{i=0}^{\infty}(-1)^i \frac{\Gamma(\tilde{\alpha}+1)}{i!\Gamma(\tilde{\alpha}-i+1)}x_{k-i}=0 \right\},\\
c(\Delta^{(\tilde{\alpha})}) &:= \left\{ x=(x_k)\in\omega : \lim_k \sum_{i=0}^{\infty}(-1)^i \frac{\Gamma(\tilde{\alpha}+1)}{i!\Gamma(\tilde{\alpha}-i+1)}x_{k-i} \text{ exists} \right\},\\
\ell_\infty(\Delta^{(\tilde{\alpha})}) &:= \left\{ x=(x_k)\in\omega : \sup_k \left|\sum_{i=0}^{\infty}(-1)^i \frac{\Gamma(\tilde{\alpha}+1)}{i!\Gamma(\tilde{\alpha}-i+1)}x_{k-i}\right|<\infty \right\}.
\end{aligned}$$

Let us now define the sequence $y=(y_k)$ which will be used, by the $\Delta^{(\tilde{\alpha})}$-transform of a sequence $x=(x_k)$, that is,

$$\begin{aligned}
y_k &= x_k - \tilde{\alpha}x_{k-1} + \frac{\tilde{\alpha}(\tilde{\alpha}-1)}{2!}x_{k-2} - \frac{\tilde{\alpha}(\tilde{\alpha}-1)(\tilde{\alpha}-2)}{3!}x_{k-3}+\cdots\\
&= \sum_{i=0}^{\infty}(-1)^i \frac{\Gamma(\tilde{\alpha}+1)}{i!\Gamma(\tilde{\alpha}-i+1)}x_{k-i}.
\end{aligned}$$

Hence, those spaces can be considered as the matrix domains of the triangle $\Delta^{(\tilde{\alpha})}$ in the classical sequence spaces c_0, c, ℓ_∞. We also have the following relation between the sequences $x=(x_k)$ and $y=(y_k)$:

$$x_k = \sum_{i=0}^{\infty}(-1)^i \frac{\Gamma(-\tilde{\alpha}+1)}{i!\Gamma(-\tilde{\alpha}-i+1)}y_{k-i}.$$

Lemma 1.3. *([2], Theorem 4.3.12, p. 63) Let $(X,\|.\|)$ be a BK space. Then X_T is a BK space with $\|.\|_T = \|T(.)\|$.*

By Lemma 1.3, defined fractional difference sequence spaces are complete, linear BK–spaces with the following norm:

$$\|x\| = \sup_n \left|\sum_{i=0}^{\infty}(-1)^i \frac{\Gamma(\tilde{\alpha}+1)}{i!\Gamma(\tilde{\alpha}-i+1)}x_{n-i}\right|.$$

Remark 1.4. *([1], Remark 2.4) The matrix domain X_T of a linear metric sequence space X has a basis if and only if X has a basis.*

Theorem 1.5. *Let us define the sequences $c^{(n)}$ for $n=0,1,\ldots$ and $c^{(-1)}$ by*

$$c_k^{(n)} = \begin{cases} 0 & (0\le k<n)\\ (-1)^{k-n}\frac{\Gamma(-\tilde{\alpha}+1)}{(k-n)!\Gamma(-\tilde{\alpha}+n-k+1)} & (k\ge n)\end{cases}$$

and

$$c_k^{(-1)} = \sum_{n=0}^{k} (-1)^{k-n} \frac{\Gamma(-\tilde{\alpha}+1)}{(k-n)!\Gamma(-\tilde{\alpha}+n-k+1)} \quad \text{for} \quad k = 0, 1, \ldots$$

(i) *Then* $\left(c^{(n)}\right)_{n=0}^{\infty}$ *is a Schauder basis for* $c_0(\Delta^{(\tilde{\alpha})})$ *and every sequence* $x = (x_n)_{n=0}^{\infty} \in c_0(\Delta^{(\tilde{\alpha})})$ *has a unique representation*

$$x = \sum_n (\Delta_n^{(\tilde{\alpha})} x) c^{(n)} \ \forall n.$$

(ii) *Then* $\left(c^{(n)}\right)_{n=-1}^{\infty}$ *is a Schauder basis for* $c(\Delta^{(\tilde{\alpha})})$*, and every sequence* $x = (x_n)_{n=0}^{\infty} \in c(\Delta^{(\tilde{\alpha})})$ *has a unique representation*

$$x = \xi c^{(-1)} + \sum_n (y_n - \xi)\, c^{(n)},$$

where $\xi = \lim_{n\to\infty} y_n$.

(iii) *The set* $\ell_\infty(\Delta^{(\tilde{\alpha})})$ *has no Schauder basis.*

2. The β Duals and Operator Norms of Fractional Spaces

If x and y are sequences and X and Y are subsets of ω, then we write $x \cdot y = (x_k y_k)_{k=0}^{\infty}$,

$$x^{-1} * Y = \{a \in \omega : a \cdot x \in Y\}$$

and

$$M(X, Y) = \bigcap_{x \in X} x^{-1} * Y = \{a \in \omega : a \cdot x \in Y \text{ for all } x \in X\}$$

for the multiplier space of X and Y; in particular, we use the notation $x^{\beta} = x^{-1} * cs$ $X^{\beta} = M(X, cs)$ for the β dual of X.

Lemma 2.1. *Let* T *be a triangle and* S *be its inverse and* $R = S^t$*, the transpose of* S.

(i) *Let* X *be a BK space with AK or* $X = \ell_\infty$*. Then* $a \in (X_T)^{\beta}$ *if and only if* $a \in \left(X^{\beta}\right)_R$ *and* $W \in (X, c_0)$ *where the triangle* W *is defined for* $n = 0, 1, 2, \ldots$ *by*

$$w_{nk} = \begin{cases} 0 & (k > n) \\ \sum_{j=n}^{\infty} a_j s_{jk} & (0 \le k \le n). \end{cases} \tag{2.1}$$

Moreover, if $a \in (X_T)^{\beta}$ *then*

$$\sum_k a_k z_k = \sum_k (R_k a)(T_k z) \ \forall z \in X_T. \tag{2.2}$$

(ii) *We have* $a \in (c_T)^{\beta}$ *if and only if* $a \in (\ell_1)_R$ *and* $W \in (c, c)$*. Moreover, if* $a \in (c_T)^{\beta}$ *then we have*

$$\sum_k a_k z_k = \sum_k (R_k a)(T_k z) - \lim_k T_k z \lim_n \sum_{k=0}^{n} w_{nk} \ \forall z \in c_T. \tag{2.3}$$

Remark 2.2. *We have the following results:*

(i) *If X be a BK space with AK then the condition $W \in (X, c_0)$ in Lemma 2.1(i) can be replaced by*

$$W \in (X, \ell_\infty). \tag{2.4}$$

(ii) *The condition $W \in (c, c)$ in Lemma 2.1(ii) can be replaced by the conditions*

$$W \in (c_0, \ell_\infty) \ \textit{and} \tag{2.5}$$

$$\lim_n W_n e = \gamma \ \textit{exists}. \tag{2.6}$$

Proof. By definition of the triangle W we have

$$\lim_n W_n e^{(k)} = \lim_n \sum_{j=n}^{\infty} = 0 \ \forall k. \tag{2.7}$$

(i) The conditions (2.4) and (2.7) imply $W \in (X, c_0)$ because X has AK and c_0 is a closed subspace of ℓ_∞. On the other hand clearly $W \in (X, c_0)$ implies (2.4).
(ii) Since c_0 has AK and c is a closed subspace of ℓ_∞, the conditions (2.5) and (2.7) imply $W \in (c_0, c_0)$. Then $W \in (c_0, c)$ and the condition (2.6) imply $W \in (c, c)$. On the other hand $W \in (c, c)$ implies (2.5) and $We \in c$, that is the condition (2.6) holds.

□

Theorem 2.3. *We have*

(i) *$a \in \left(c_0(\Delta^{(\tilde{\alpha})})\right)^\beta$ if and only if*

$$\sum_k \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_j \right| < \infty \tag{2.8}$$

and

$$\sup_n \left(\sum_{k=0}^{n} \left| \sum_{j=n}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_j \right| \right) < \infty; \tag{2.9}$$

moreover, if $a \in \left(c_0(\Delta^{(\tilde{\alpha})})\right)^\beta$ then $\forall x \in c_0(\Delta^{(\tilde{\alpha})})$

$$\sum_k a_k x_k = \sum_k \left(\sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_j \right) y_k. \tag{2.10}$$

(ii) $a \in \left(c(\Delta^{(\tilde{\alpha})})\right)^{\beta}$ *if and only if (2.8), (2.9) and*

$$\lim_n \sum_{k=0}^{n} \left(\sum_{j=n}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_j \right) = \rho; \tag{2.11}$$

moreover, if $a \in \left(c(\Delta^{(\tilde{\alpha})})\right)^{\beta}$ *then* $\forall x \in c(\Delta^{(\tilde{\alpha})})$

$$\sum_k a_k x_k = \sum_k \left(\sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_j \right) y_k - \rho \lim_k y_k.$$

(iii) $a \in \left(\ell_\infty(\Delta^{(\tilde{\alpha})})\right)^{\beta}$ *if and only if (2.8) and*

$$\lim_n \sum_{k=0}^{n} \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_j \right| = 0; \tag{2.12}$$

moreover, if $a \in \left(\ell_\infty(\Delta^{(\tilde{\alpha})})\right)^{\beta}$ *then (2.10) holds* $\forall x \in \ell_\infty(\Delta^{(\tilde{\alpha})})$.

Proof. We apply Lemma 2.1 and Remark 2.2.

The triangles R and W defined for $n = 0, 1, \ldots$ by

$$R_k a = \sum_{j=k}^{\infty} s_{jk} a_j = \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_j \quad (k = 0, 1, \ldots) \tag{2.13}$$

and

$$w_{nk} = \begin{cases} \sum\limits_{j=n}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_j & (0 \le k \le n) \\ 0 & (k > n). \end{cases} \tag{2.14}$$

The condition $Ra \in \ell_1$ of Lemma 2.1 holds in each part because $c_0^{\beta} = c^{\beta} = \ell_\infty^{\beta}$.

By Lemma 2.1 and Remark 2.2(ii), we must add the condition $W \in (c_0, \ell_\infty)$ which is equivalent to

$$\sup_n \sum_{k=0}^{n} |w_{nk}| < \infty$$

and this is the condition in (2.6) to (2.9), and (2.9) is the condition in (2.11).

By Lemma 2.1, we must add the condition $W \in (\ell_\infty, c_0)$ which is equivalent to

$$\lim_n \sum_{k=0}^{n} |w_{nk}| = 0.$$

Note that, (2.13) in Parts (i) and (iii) and (2.14) come from (2.2) and (2.3), respectively. □

2.1. **Operators Norms and Matrix Transformations of Fractional Sequence Spaces.** Let us now establish identities and inequalities of operator norms for fractional sequence spaces and then characterize some classes of matrix mappings on them. The following notations and results are needed to characterize certain classes of matrix mappings on the sets of fractional sequences and for determination of the operator norms of our spaces.

Lemma 2.4. *Let X and Y be BK spaces*

(i) *Then we have $(X,Y) \subset \mathcal{B}(X,Y)$, that is, every $A \in (X,Y)$ defines an operator $L_A \in \mathcal{B}(X,Y)$, where $L_A(x) = Ax$ for all $x \in X$ (see* [5]*, Theorem 1.23).*

(ii) *If X has AK then we have $\mathcal{B}(X,Y) \subset (X,Y)$, that is, every operator $L \in \mathcal{B}(X,Y)$ is given by a matrix $A \in (X,Y)$ such that $L(x) = Ax$ for all $x \in X$ (see* [1]*, Theorem 1.9).*

Lemma 2.5. *Let Y be an arbitrary subset of ω.*

(i) *Let X be a BK space with AK or $X = \ell_\infty$, and $R = S^t$. Then $A \in (X_T, Y)$ if and only if $\hat{A} \in (X,Y)$ and $W^{(A_n)} \in (X, c_0)$ for all $n = 0, 1, \ldots$. Here $\hat{A}$ is the matrix with rows $\hat{A}_n = RA_n$ for $n = 0, 1, \ldots$, and the triangles $W^{(A_n)}$ $(n = 0, 1, \ldots)$ are defined as in 2.16 with a_j replaced by a_{nj}.*

Moreover, if $A \in (X,Y)$ then we have $Az = \hat{A}(Tz)$ for all $z \in Z = X_T$ (see [11]*, Theorem 3.4, Remark 3.5(a)).*

(ii) *We have $A \in (c_T, Y)$ if and only if $\hat{A} \in (c_0, Y)$ and $W^{(A_n)} \in (c,c)$ for all $n = 0, 1, \ldots$ and $\hat{A}e - (\gamma_n)_{n=0}^{\infty} \in Y$, where*

$$\gamma_n = \lim_m \sum_{k=0}^{m} w_{mk}^{(A_n)} \text{for } n = 0, 1, \ldots$$

Moreover, if $A \in (c_T, Y)$ then we have $Az = \hat{A}(Tz) - \eta(\gamma_n)_{n=0}^{\infty}$ for all $z \in c_T$, where $\eta = \lim_k T_k z$ ([11]*, Theorem 1.23).*

Theorem 2.6. *Let $X = c_0(\Delta^{(\tilde{\alpha})})$ or $X = \ell_\infty(\Delta^{(\tilde{\alpha})})$.*

(i) *Let $Y = c_0, c, \ell_\infty$. If $A \in (X_T, Y)$ then, putting*

$$\|A\|_{(X_T,\infty)} = \sup_n \left\| \hat{A}_n \right\|_1 = \sup_n \sum_k \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_{nj} \right|,$$

we have $\|L_A\| = \|A\|_{(X_T,\infty)}$.

(ii) *Let $Y = \ell_1$. If $A \in (X_T, \ell_1)$. Then we have*

$$\|A\|_{(X_T,1)} = \sup_{\substack{N \subset \mathbb{N} \\ N \text{ finite}}} \left(\sum_k \left| \sum_{n \in N} \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_{nj} \right| \right) \leq \|L_A\| \leq 4 \|A\|_{(X_T,1)}.$$

Proof. The proof is based on the results in ([16], Theorem 2.8) □

Theorem 2.7. *The operator norm of the set $c(\Delta^{(\tilde{\alpha})})$ is given.*

(i) *Let* $A \in \left(c(\Delta^{(\widetilde{\alpha})}), Y\right)$*, where* Y *is any of the spaces* c_0, c *or* ℓ_∞*. Then we have*

$$\|L_A\| = \|A\|_{\left(c(\Delta^{(\widetilde{\alpha})}),\infty\right)} = \sup_n \left(\sum_k \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_{nj} \right| + |\gamma_n| \right),$$

where $\gamma_n = \lim_m \sum_{k=0}^{m} w_{mk}^{(A_n)}$ *for* $n = 0, 1, \ldots$

(ii) *Let* $A \in \left(c(\Delta^{(\widetilde{\alpha})}), \ell_1\right)$*, then, putting,*

$$\|A\|_{\left(c(\Delta^{(\widetilde{\alpha})}),1\right)} = \sup_{\substack{N \subset \mathbb{N} \\ N \text{ finite}}} \left(\sum_k \left| \sum_{n \in N} \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)} a_{nj} \right| + \left| \sum_{n \in N} \gamma_n \right| \right)$$

we have

$$\|A\|_{\left(c(\Delta^{(\widetilde{\alpha})}),1\right)} \leq \|L_A\| \leq 4 \|A\|_{\left(c(\Delta^{(\widetilde{\alpha})}),1\right)}.$$

Proof. The proof is based on the results in ([16], Theorem 2.9) □

Theorem 2.8. *The necessary and sufficient conditions for* $A \in (\ell_\infty(\Delta^{(\widetilde{\alpha})}), Y)$ $A \in (c_0(\Delta^{(\widetilde{\alpha})}), Y)$ *and* $A \in (c(\Delta^{(\widetilde{\alpha})}), Y)$*, where* $Y \in \{\ell_\infty, c_0, c, \ell_1\}$ *can be read from the following table:*

From / *To*	$\ell_\infty(\Delta^{(\widetilde{\alpha})})$	$c_0(\Delta^{(\widetilde{\alpha})})$	$c(\Delta^{(\widetilde{\alpha})})$
ℓ_∞	**1.**	**1.**	**2.**
c_0	**3.**	**3.**	**4.**
c	**5.**	**5.**	**6.**
ℓ_1	**7.**	**7.**	**8.**

[**1**] 1A and 1B where

(1A) $\|A\|_{(c(\Delta^{(\widetilde{\alpha})}),\infty)} = \sup_n \sum_{k=0}^{\infty} |\hat{a}_{nk}| < \infty$,

(1B) $\|W^{(A_n)}\|_{(\ell_\infty,c_0)} = \lim_{m\to\infty} \sum_{k=0}^{m} \left| w_{mk}^{(A_n)} \right| = 0$ for all n.

[**2**] 1A and 2A where

(2A) $\|W^{(A_n)}\|_{(\ell_\infty,\ell_\infty)} = \sup_m \sum_{k=0}^{m} \left| w_{mk}^{(A_n)} \right| < \infty$ for all n.

[**3**] 1A, 2A, 3A and 3B where

(3A) $\lim_{m\to\infty} \sum_{k=0}^{m} w_{mk}^{(n)} = \gamma_n$ exists for each n,

(3B) $\sup_n \left| \sum_{k=0}^{\infty} \hat{a}_{nk} - \gamma_n \right| = 0$.

[**4**] 1B and 4A where

(4A) $\lim_{m\to\infty} \sum_{k=0}^{m} |\hat{a}_{nk}| = 0$.

[**5**] 1A, 2A and 5A where

(5A) $\lim_{n\to\infty} \hat{a}_{nk} = 0$ for each k.

[**6**] 1A, 2A, 3A, 5A and 6A where

(6A) $\lim_{n\to\infty}\left(\sum_{k=0}^{\infty}\hat{a}_{nk}-\gamma_n\right)=0.$

[**7**] 1B, 7A, 7B and 7C where

(7A) $\lim_{n\to\infty}\hat{a}_{nk}=\hat{\alpha}_k$ exists for each n,

(7B) $\sum_{k=0}^{\infty}|\hat{a}_{nk}|\,,\sum_{k=0}^{\infty}|\hat{\alpha}_k|<\infty$ for all n,

(7C) $\lim_{n\to\infty}\left(\sum_{k=0}^{\infty}\hat{a}_{nk}-\hat{\alpha}_k\right)=0.$

[**8**] 1A, 2B and 7A.

[**9**] 1A, 2B, 3A, 7A and 9A where

(9A) $\lim_{n\to\infty}\left(\sum_{k=0}^{\infty}\hat{a}_{nk}-\gamma_n\right)=\delta$ exists.

[**10**] 1B and 10A where

(10A) $\sup_{K\subset\mathbb{N}}\sum_{n=0}^{\infty}\left|\sum_{k\in K}\hat{a}_{nk}\right|<\infty.$

[**11**] 2A and 10A.

[**12**] 2A, 3A, 10A and 12A where

(12A) $\sum_{n=0}^{\infty}\left|\sum_{k=0}^{\infty}\hat{a}_{nk}-\gamma_n\right|<\infty,$

where for given a matrix $A=(a_{nk})_{n,k=0}^{\infty}$, we define the matrices $\hat{A}=(\hat{a}_{nk})_{n,k=0}^{\infty}$ and $W^{(A_n)}=(w_{mk}^{(A_n)})_{m,k=0}^{\infty}$ by

$$\hat{a}_{nk}=\sum_{j=k}^{\infty}(-1)^{j-k}\frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)}a_{nj}\text{ for all }n,k\in\mathbb{N}_0 \tag{2.15}$$

and

$$w_{mk}^{(A_n)}=\begin{cases}\sum_{j=m}^{\infty}(-1)^{j-k}\frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}+j-k+1)}a_{nj} & (0\le k\le m)\\ 0 & (k>m)\end{cases} \tag{2.16}$$

for $n,m\in\mathbb{N}_0$,

Proof. Note that the entries of the triangles $\hat{A}$ and $W^{(A_n)}$ are given above and assume that $Y\in\{c_0,c,\ell_\infty,\ell_1\}$.

[1], [4], [7], [10] Taking into account Lemma 2.5(i) we have $A\in(\ell_\infty(\Delta^{(\tilde{\alpha})}),Y)$ if and only if $\hat{A}\in(\ell_\infty,Y)$ and $W^{(A_n)}\in(\ell_\infty,c_0)$ for each n. First, $\hat{A}\in(\ell_\infty,Y)$ satisfies (1A) in [1], by [[2], Theorem 1.3.3], (4A) in [4] by [[21], 21. (21.1)], (7A) , (7B) and (7C) in [7] by [[2], Theorem 1.7.18 (ii)], and (10A) in [10] by [[2], 8.4.9A]. Also, $W^{(A_n)}\in(\ell_\infty,c_0)$ for all n satisfies (2A) in [1], [4], [7], [10] by [[2], Theorem 1.3.3].

[2], [5], [8], [11] Remark 2.2(i) and Lemma 2.5(i) satisfy $A\in(\ell_\infty(\Delta^{(\tilde{\alpha})}),Y)$ if and only if $\hat{A}\in(c_0,Y)$ and $W^{(A_n)}\in(c_0,c_0)$ for each $n=0,1,\ldots$. First $\hat{A}\in(c_0,c_0)$ satisfies (1A) in [2] by [[2], Theorem 1.3.3], (1A) and (5A) in [5] by [[2], 8.4.5A], (1A) and (7A) in [8] by [[2],

8.4.5A] and (10A) in [11] by [15, 8.4.3B]. Also by [[2], Theorem 1.3.3] $W^{(A_n)} \in (c_0, c_0)$ for all n satisfies (2A) in [2], [5], [8], [11].

[3], [6], [9], [12] Remark 2.2(i) and Lemma 2.5(i) satisfy $A \in (c_0(\Delta^{(\tilde{\alpha})}), Y)$ if and only if $\hat{A} \in (c_0, Y)$ and $W^{(A_n)} \in (c_0, \ell_\infty)$ for each n, and

$$\lim_{n\to\infty} W_m^{(A_n)} = \lim_m \sum_{k=0}^{m} w_{mk}^{(A_n)} \text{ for } n = 0, 1, \dots$$

and for each n

$$\left\{ \hat{A} - \left(\lim_m \sum_{k=0}^{m} w_{mk}^{(A_n)} \right) \right\} \in Y.$$

It means, we have to add the last two conditions to those for $A \in (c_0(\Delta^{(\tilde{\alpha})}), Y)$, that is, (3A) and (3B) in [3] to those in [2], (3A) and (6A) in 6. to those in 5., (3A) and (9A) in [9] to those in [8] and (3A) and (12A) in [12] to those in [11]. □

3. Compact Operators on Fractional Spaces $c_0(\Delta^{(\tilde{\alpha})}), c(\Delta^{(\tilde{\alpha})})$ and $\ell_\infty(\Delta^{(\tilde{\alpha})})$

In this section, we give our main results related to compact operators on fractional sequence spaces. We recall the definition of the Hausdorff measure of noncompactness of bounded subsets of a metric space, and the Hausdorff measure of noncompactness of operators between Banach spaces.

If X and Y are infinite–dimensional complex Banach spaces then a linear operator $L : X \to Y$ is said to be compact if the domain of L is all of X, and, for every bounded sequence (x_n) in X, the sequence $(L(x_n))$ has a convergent subsequence. We denote the class of such operators by $\mathcal{C}(X, Y)$.

Let (X, d) be a metric space, $B(x_0, \delta) = \{x \in X : d(x, x_0) < \delta\}$ denote the open ball of radius $\delta > 0$ and center in $x_0 \in X$, and $\mathcal{M}_X$ be the collection of bounded sets in X. The Hausdorff measure of noncompactness of $Q \in \mathcal{M}_X$ is

$$\chi(Q) = \inf\{\epsilon > 0 : Q \subset \bigcup_{k=1}^{n} B(x_k, \delta_k) : x_k \in X,\ \delta_k < \epsilon,\ 1 \le k \le n,\ n \in \mathbb{N}\}.$$

Let X and Y be Banach spaces and χ_1 and χ_2 be measures of noncompactness on X and Y. Then the operator $L : X \to Y$ is called (χ_1, χ_2)–bounded if $L(Q) \in \mathcal{M}_Y$ for every $Q \in \mathcal{M}_X$ and there exists a positive constant C such that

$$\chi_2(L(Q)) \le C\chi_1(Q) \text{ for every } Q \in \mathcal{M}_X. \tag{3.1}$$

If an operator L is (χ_1, χ_2)–bounded then the number

$$\|L\|_{(\chi_1,\chi_2)} = \inf\{C \ge 0 : (3.1) \text{ holds for all } Q \in \mathcal{M}_X\}$$

is called the (χ_1, χ_2)–measure of noncompactness of L. In particular, if $\chi_1 = \chi_2 = \chi$, then we write $\|L\|_\chi$ instead of $\|L\|_{(\chi,\chi)}$.

Lemma 3.1. *([5], Theorem 2.25 and Corollary 2.26) Let X and Y are Banach spaces and $L \in \mathcal{B}(X, Y)$. Then we have*

$$\|L\|_\chi = \chi\left(L(\bar{B}_X)\right) = \chi\left(L(S_X)\right), \tag{3.2}$$

$$L \in \mathcal{C}(X,Y) \text{ if and only if } \|L\|_\chi = 0. \tag{3.3}$$

Lemma 3.2. *(Goldenstein, Gohberg, Markus* [5]*, Theorem 2.23) Let X be a Banach space with Schauder basis $(b_n)_{n=0}^{\infty}$, $Q \in \mathcal{M}_X$, $P_n : X \to X$ be the projector onto the linear span of $\{b_0, b_1, \dots b_n\}$. I be the identity map on X and $R_n = I - P_n$ $(n = 0, 1, \dots)$. Then we have*

$$\frac{1}{a} \cdot \limsup_{n\to\infty} \left(\sup_{x\in Q} \|R_n(x)\| \right) \leq \chi(Q) \leq \limsup_{n\to\infty} \left(\sup_{x\in Q} \|R_n(x)\| \right), \tag{3.4}$$

where $a = \limsup_{n\to\infty} \|R_n\|$.

Lemma 3.3. *(*[5]*, Theorem 2.8) Let Q be a bounded subset of the normed space X, where X is ℓ_p for $1 \leq p < \infty$ or c_0. If $P_n : X \to X$ is the operator defined by $P_n(x) = x^{[n]}$ for $x = (x_k)_{k=0}^{\infty} \in X$, then we have*

$$\chi(Q) = \lim_n \left(\sup_{x\in Q} \|R_n(x)\| \right).$$

The final results of this section give the estimates of the Hausdorff measure of noncompactness of L_A when $A \in (X_T, c)$ for $X = c_0, \ell_\infty, c$.

Lemma 3.4. *(* [17]*, Corollary 5.13) If $A \in ((c_0)_T, c)$ or $A \in ((\ell_\infty)_T, c)$ then we have*

$$\frac{1}{2} \cdot \lim_{r\to\infty} \left(\sup_{n\geq r} \left\| \hat{A}_n - \hat{\alpha} \right\|_1 \right) \leq \|L_A\|_\chi \leq \lim_{r\to\infty} \left(\sup_{n\geq r} \left\| \hat{A}_n - \hat{\alpha} \right\|_1 \right), \tag{3.5}$$

where $\hat{\alpha} = (\alpha_k)_{k=0}^{\infty}$ with $\hat{\alpha}_k = \lim_{n\to\infty} \hat{a}_{nk}$ for $k = 0, 1, \dots$

Lemma 3.5. *(* [17]*, Corollary 5.14) If $A \in (c_T, c)$ then we have*

$$\begin{aligned} &\frac{1}{2} \cdot \lim_{r\to\infty} \left(\sup_{n\geq r} \left(\left| \beta - \delta_n - \sum_{k=0}^{\infty} \alpha_k \right| + \sum_{k=0}^{\infty} |\hat{a}_{nk} - \alpha_k| \right) \right) \\ \leq\ &\|L_A\|_\chi \leq \lim_{r\to\infty} \left(\sup_{n\geq r} \left(\left| \beta - \delta_n - \sum_{k=0}^{\infty} \alpha_k \right| + \sum_{k=0}^{\infty} |\hat{a}_{nk} - \alpha_k| \right) \right), \end{aligned} \tag{3.6}$$

where $\gamma_n = \lim_{m\to\infty} w_{mk}^{(A_n)}$ for $n = 0, 1, \dots$, $\beta = \lim_{n\to\infty} (\sum_{k=0}^{\infty} \hat{a}_{nk} - \gamma_n)$ and $\hat{\alpha} = (\alpha_k)_{k=0}^{\infty}$ with $\hat{\alpha}_k = \lim_{n\to\infty} \hat{a}_{nk}$ for $k = 0, 1, \dots$.

We now establish necessary and sufficient conditions for a matrix operator to be a compact operator from fractional difference sequence spaces into Y, where $Y \in \{c_0, c, \ell_\infty, \ell_1\}$. This is achieved applying the results given above about Hausdorff measure of noncompactness.

Theorem 3.6. *The identities or estimates for L_A when $A \in (\ell_\infty(\Delta^{(\widetilde{\alpha})}), Y)$, $A \in (c_0(\Delta^{(\tilde{\alpha})}), Y)$ and $A \in (c(\Delta^{(\widetilde{\alpha})}), Y)$, where $Y \in \{\ell_\infty, c_0, c, \ell_1\}$ can be read from the following table:*

From / *To*	$\ell_\infty(\Delta^{(\widetilde{\alpha})})$	$c_0(\Delta^{(\tilde{\alpha})})$	$c(\Delta^{(\widetilde{\alpha})})$
ℓ_∞	**1.**	**1.**	**2.**
c_0	**3.**	**3.**	**4.**
c	**5.**	**5.**	**6.**
ℓ_1	**7.**	**7.**	**8.**

Here

1. $0 \le \|L_A\|_\chi \le \lim_{r\to\infty}\left(\sup_{n\ge r}\sum_{k=0}^{\infty}|\hat{a}_{nk}|\right)$;

2. $0 \le \|L_A\|_\chi \le \lim_{r\to\infty}\left(\sup_{n\ge r}\sum_{k=0}^{\infty}|\hat{a}_{nk}|+|\gamma_n|\right)$;

3. $\|L_A\|_\chi = \lim_{r\to\infty}\left\|\hat{A}^{[r]}\right\|_{(\ell_\infty,\ell_\infty)}$;

4. $\|L_A\|_\chi = \lim_{r\to\infty}\left(\sup_{n\ge r}\sum_{k=0}^{\infty}|\hat{a}_{nk}|+|\gamma_n|\right)$;

5. $\frac{1}{2}\cdot\lim_{r\to\infty}\left\|\hat{B}^{[r]}\right\|_{((\ell_\infty,\ell_\infty)} \le \|L_A\|_\chi \le \lim_{r\to\infty}\left\|\hat{B}^{[r]}\right\|_{((\ell_\infty,\ell_\infty)}$;

6. $\frac{1}{2}\cdot\lim_{r\to\infty}\left(\sup_{n\ge r}\sum_{k=0}^{\infty}\left|\hat{b}_{nk}\right|+|\delta_n|\right) \le \|L_A\|_\chi \le \lim_{r\to\infty}\left(\sup_{n\ge r}\sum_{k=0}^{\infty}\left|\hat{b}_{nk}\right|+|\delta_n|\right)$;

7. $\lim_{r\to\infty}\sup_{\substack{N\subset\mathbb{N}_0\\ finite}}\left\|\sum_{n\in\mathbb{N}}\hat{A}_n^{[r]}\right\|_1 \le \|L_A\|_\chi \le 4\lim_{r\to\infty}\sup_{\substack{N\subset\mathbb{N}_0\\ finite}}\left\|\sum_{n\in\mathbb{N}}\hat{A}_n^{[r]}\right\|_1$;

8. $\lim_{r\to\infty}\sup_{\substack{N\subset\mathbb{N}_0\\ finite}}\left(\left\|\sum_{n\in\mathbb{N}}\hat{A}_n^{[r]}\right\|_1+\left|\sum_{n\in N}\gamma_n\right|\right) \le \|L_A\|_\chi \le 4\lim_{r\to\infty}\sup_{\substack{N\subset\mathbb{N}_0\\ finite}}\left(\left\|\sum_{n\in\mathbb{N}}\hat{A}_n^{[r]}\right\|_1+\left|\sum_{n\in N}\gamma_n\right|\right)$,

where the notations used in the theorem are defined as follows:

Let $A=(a_{nk})_{n,k=0}^{\infty}$ *be an infinite matrix and* $r\in\mathbb{N}_0$. *Then* $A^{[r]}$ *denotes the matrix with rows* $A_n^{[r]}=0$ *for* $0\le n\le r$ *and* $A_n^{[r]}=A_n$ *for* $n\ge r+1$.

We write $\hat{A}$ *for the matrix with*

$$\hat{a}_{nk}=\sum_{j=k}^{\infty}(-1)^{j-k}\frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)}a_{nj} \text{ for all } n,k\in\mathbb{N}_0;$$

and $\hat{\alpha}=(\hat{\alpha}_k)_{k=0}^{\infty}$ *and* $\gamma=(\gamma_n)_{n=0}^{\infty}$ *for the sequences with* $\hat{\alpha}_k=\lim_{n\to\infty}\hat{a}_{nk}$ *for* $k=0,1,\ldots$ *and*

$$\gamma_n=\lim_m\sum_{k=0}^{m}w_{mk}^{(A_n)}=\lim_m\sum_{k=0}^{m}\sum_{j=m}^{\infty}(-1)^{j-m}\frac{\Gamma(-\tilde{\alpha}+1)}{(j-m)!\Gamma(-\tilde{\alpha}-j+m+1)}a_{nj};$$

also $\beta=\lim_{n\to\infty}(\sum_{k=0}^{\infty}\hat{a}_{nk}-\gamma_n)$. *We also write* $\hat{B}=(\hat{b}_{nk})_{n,k=0}^{\infty}$ *for the matrix with* $\hat{b}_{nk}=\hat{a}_{nk}-\hat{\alpha}_k$ *for each* $n,k\in\mathbb{N}_0$ *and* $\delta=(\delta_n)_{n=0}^{\infty}$ *for the sequence with* $\delta_n=\sum_{k=0}^{\infty}\hat{\alpha}_k-\gamma_n+\beta$ $(n=0,1,\ldots)$

Proof. The conditions in **1.** and **2.** are immediate consequence of ([16], Corollary 3.6(a)). We define $P_r:\ell_\infty\to\ell_\infty$ by $P_r(x)=x^{[r]}$ for all $x\in\ell_\infty$ and $r=0,1,\ldots$, $R_r=I-P_r$, and write $L=L_A$ and $\bar{B}=\bar{B}_{\ell_\infty}$ for short. Then it follows from (3.1), ([5], Theorem 2.12) and Lemma 2.6(i) that

$$\begin{aligned}0 &\le \|L\|_\chi=\chi(L(\bar{B}))\\ &\le \chi(P_r(L(\bar{B})))+\chi(R_r(L(\bar{B})))\\ &= \chi(R_r(L(\bar{B})))\le\sup_{x\in\bar{B}}\|R_r(L(x))\|_\infty=\left\|\hat{A}^{[r]}\right\|_{(X,\infty)}.\end{aligned}$$

Then, **3.** holds.

The conditions in **4.** and **6.** are immediate consequence of ([16], Theorem 3.7 (b), (a)). Part **5.** follows by a similar argument as part **3.**; we use Lemma 2.7(i) instead of Lemma 2.6(i). □

Corollary 3.7. *Let X be one of the spaces $c_0(\Delta^{(\widetilde{\alpha})})$ or $\ell_\infty(\Delta^{(\widetilde{\alpha})})$. We obtain as an immediate consequence of (3.3) and Theorem 3.6 ((1.1), (3.1) and (5.1)).*

(i) *If $A \in (X, c_0)$, then L_A is compact if and only if*

$$\lim_{r\to\infty}\left(\sup_{n\geq r}\left(\sum_{k=0}^{\infty}|\hat{a}_{nk}|\right)\right)=0. \tag{3.7}$$

(ii) *If $A \in (X, c)$, then L_A is compact if and only if*

$$\lim_{r\to\infty}\sup_{n\geq r}\left(\sum_{k=0}^{\infty}|\hat{a}_{nk}-\hat{\alpha}_k|\right)=0.$$

(iii) *If $A \in (X, \ell_\infty)$, then L_A is compact if the condition (3.7) holds.*

Corollary 3.8. *We obtain as an immediate consequence of (3.3) and Theorem 3.6 ((2.1), (4.1) and (6.1)).*

(i) *If $A \in (c(\Delta^{(\widetilde{\alpha})}), c_0)$, then L_A is compact if and only if*

$$\lim_{r\to\infty}\left(\sup_{n\geq r}\left(\sum_{k=0}^{\infty}|\hat{a}_{nk}|+|\gamma_n|\right)\right)=0. \tag{3.8}$$

(ii) *If $A \in (c(\Delta^{(\widetilde{\alpha})}), c)$, then L_A is compact if and only if*

$$\lim_{r\to\infty}\sup_{n\geq r}\left(\sum_{k=0}^{\infty}|\hat{a}_{nk}-\hat{\alpha}_k|+\left|\sum_{k=0}^{\infty}\hat{\alpha}_k-\gamma_n-\beta\right|\right)=0.$$

(iii) *If $A \in (c(\Delta^{(\widetilde{\alpha})}), \ell_\infty)$, then L_A is compact if (3.8) holds.*

Corollary 3.9. *We obtain as an immediate consequence of (3.3), Theorem 3.6 ((7.1), (8.1)).*

(i) *If $A \in (c_0(\Delta^{(\tilde{\alpha})}), \ell_1)$ or $A \in (\ell_\infty(\Delta^{(\tilde{\alpha})}), \ell_1)$, then L_A is compact if and only if*

$$\lim_{r\to\infty}\left(\sup_{\substack{N\subset\mathbb{N}_0\\ finite}}\left\|\sum_{n\in N_r}\sum_{j=k}^{\infty}(-1)^{j-k}\frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)}a_{nj}\right\|_1\right)=0.$$

(ii) *If $A \in (c(\Delta^{(\tilde{\alpha})}), \ell_1)$, then L_A is compact if and only if*

$$\lim_{r\to\infty}\sup_{\substack{N\subset\mathbb{N}_0\\ finite}}\left(\sum_{k=0}^{\infty}\left|\sum_{n\in N_r}\sum_{j=k}^{\infty}(-1)^{j-k}\frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)!\Gamma(-\tilde{\alpha}-j+k+1)}a_{nj}\right|+\left|\sum_{n\in N_r}\gamma_n\right|\right)=0.$$

Compliance with ethical standards

Conflict of interest. The authors declare that they have no conflict of interest.

Ethical approval. This article does not contain any studies with human participants or animals performed by any of the authors.

References

[1] A.M. Jarrah, E. Malkowsky, Ordinary, absolute and strong summability and matrix transformations, Filomat **17** (2003), 59–78.

[2] A. Wilansky, Summability through Functional Analysis, North–Holland Mathematics Studies 85, Amsterdam, New York, Oxford, 1984.

[3] B. Altay and F. Başar, Certain topological properties and duals of the matrix domain of a triangle matrix in a sequence space, J. Math. Anal. Appl. **336**(1) (2007), 632–645.

[4] C. Aydn and F. Başar, Some new difference sequence spaces. Appl. Math. Comput. **157**(3) (2004), 677693.

[5] E. Malkowsky and V. Rakočević, An introduction into the theory of sequence spaces and measures of noncompactness, Zb. Rad. (Beogr.) **9**(17) (2000), 143–234.

[6] E. Malkowsky and F. Özger, A note on some sequence spaces of weighted means, Filomat **26**(3) (2012), 511-518.

[7] E. Malkowsky and F. Özger, Compact Operators on Spaces of Sequences of Weighted Means, AIP Conf. Proc. **1470** (2012), 179–182.

[8] E. Malkowsky, F. Özger and A. Alotatibi, Some Notes on Matrix Mappings and their Hausdorff Measure of Noncompactness, Filomat **28**5 (2014), 10591072.

[9] E. Malkowsky, F. Ozger and V. Veličković, Some Spaces Related to Cesaro Sequence Spaces and an Application to Crystallography, MATCH Commun. Math. Comput. Chem. **70**(3) (2013), 867–884.

[10] E. Malkowsky and V. Rakočević, The measure of noncompactness of linear operators between spaces of mth-order difference sequences, Stud. Sci. Math. Hungar. **33** (1999) 381391.

[11] E. Malkowsky and V. Rakočević, On matrix domains of triangles, Appl. Math. Comput. **189** (2007), 1148–1163.

[12] F. Özger and F. Başar, Domain of the double sequential band matrix $B(\widetilde{r},\widetilde{s})$ on some Maddox's spaces, AIP Conf. Proc. **1470** (2012), 152–155.

[13] F. Özger and F. Başar, Domain of the double sequential band matrix $B(\widetilde{r},\widetilde{s})$ on some Maddox's spaces, Acta Mathematica Scientia **34**(2) (2014), 394-408.

[14] H. Furkan, On some λ difference sequence spaces of fractional order, J. Egypt. Math. Soc. **25** (2017), 37–42.

[15] H. Kızmaz, *On certain sequence spaces*, Canad. Math. Bull. **24**(2) (1981), 169–176.

[16] I. Djolović and E. Malkowsky, A note on compact operators on matrix domains, J. Math. Anal. Appl. **340** (2008), 291–303.

[17] I. Djolović and E. Malkowsky, Matrix transformations and compact operators on some new mth order difference sequences, Appl. Math. Comput. **198**(2) (2008), 700–714.

[18] M. Candan, Domain of the double sequential band matrix in the classical sequence spaces, J. Inequal. Appl. **2012**(281) (2012), doi:10.1186/1029-242X-2012-281.

[19] M. Kirişçi and F. Başar, Some new sequence spaces derived by the domain of generalized difference matrix, Comput. Math. Appl. **60**(5) (2010), 1299–1309.

[20] M. Mursaleen, V. Karakaya, H. Polat and N. Simek, Measure of noncompactness of matrix operators on some difference sequence spaces of weighted means, Comput. Math. Appl. **62**(2) (2011), 814–820.

[21] M. Stieglitz and H. Tietz, Matrixtransformationen von Folgenräumeneine Ergebnisübersicht, Math. Z. **154** (1977), 1–16.

[22] P. Baliarsingh and S. Dutta, On the classes of fractional order of difference sequence spaces and matrix transformations, Appl. Math. Comput. **250** (2015), 665–674.

[23] R. Çolak and M. Et, On some generalized difference sequence spaces and related matrix transformations. Hokkaido Math J. **26**(3) (1997), 483-492.

[24] U. Kadak and P. Baliarsingh, On certain Euler difference sequence spaces of fractional order and related dual properties, J. Nonlinear Sci. Appl. **8** (2015), 997–1004.

Department of Engineering Sciences, Izmir Katip Celebi University, Izmir, Turkey
E-mail address, Özger: faruk.ozger@ikc.edu.tr; farukozger@gmail.com