

CHARACTERIZATIONS OF COMPACT OPERATORS ON ℓ_p TYPE FRACTIONAL SETS OF SEQUENCES

FARUK ÖZGER

ABSTRACT. Among the sets of sequences studied, difference sets of sequences are probably the most common type of sets. This paper considers some ℓ_p type fractional difference sequence spaces via Euler gamma function. Although we characterize compactness conditions on those spaces using the main tools of Hausdorff measure of noncompactness, we can only obtain sufficient conditions when the final space is ℓ_∞ . However, we use some recent results to exactly characterize the classes of compact matrix operators when the final space is the set of bounded sequences.

1. INTRODUCTION

The Euler gamma function of a real number x (except zero and the negative integers) is defined by an improper integral:

$$\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt.$$

It is known that for any natural number n , $\Gamma(n+1) = n!$, and $\Gamma(n+1) = n\Gamma(n)$ holds for any real number $n \notin \{0, -1, -2, \dots\}$.

The fractional difference operator for a fraction $\tilde{\alpha}$ have been defined in [31] as

$$(1.1) \quad \Delta^{(\tilde{\alpha})}(x_k) = \sum_{i=0}^{\infty} (-1)^i \frac{\Gamma(\tilde{\alpha}+1)}{\Gamma(\tilde{\alpha}-i+1)} x_{k-i}.$$

It is assumed that the series defined in (1.1) is convergent for $x \in \omega$.

Let m be a positive integer, then recall the difference operators $\Delta^{(1)}$ and $\Delta^{(m)}$ by:

$$(\Delta^{(1)}x)_k = \Delta^{(1)}x_k = x_k - x_{k-1}$$

and

$$(\Delta^{(m)}x)_k = \sum_{i=0}^m (-1)^i \binom{m}{i} x_{k-i}.$$

We write Δ and $\Delta^{(m)}$ for the matrices with $\Delta_{nk} = (\Delta^{(1)}e^{(k)})_n$ and $\Delta_{nk}^{(m)} = (\Delta^{(m)}e^{(k)})_n$ for all n and k . The topological properties of some spaces that are constructed by the matrix

2010 *Mathematics Subject Classification.* 46B45, 47B37.

Key words and phrases. Euler gamma function, fractional operator, compact operator, Hausdorff measure of noncompactness .

operator $\Delta^{(m)}$ were studied in the paper [32]. In [23], some identities and estimates for the Hausdorff measure of noncompactness of matrix operators from $\Delta^{(m)}$ type spaces into the sets of bounded, convergent, null sequences and also absolutely convergent series were established.

We also write fractional difference operator as an infinite matrix:

$$\Delta_{nk}^{(\tilde{\alpha})} = \begin{cases} (-1)^{n-k} \frac{\Gamma(\tilde{\alpha}+1)}{(n-k)! \Gamma(\tilde{\alpha}-n+k+1)} & (0 \leq k \leq n) \\ 0 & (k > n). \end{cases}$$

Remark 1.1. The inverse of fractional difference matrix is given by

$$\Delta_{nk}^{(-\tilde{\alpha})} = \begin{cases} (-1)^{n-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(n-k)! \Gamma(-\tilde{\alpha}-n+k+1)} & (0 \leq k \leq n) \\ 0 & (k > n). \end{cases}$$

For some values of $\tilde{\alpha}$, we have

$$\begin{aligned} \Delta^{1/2} x_k &= x_k - \frac{1}{2} x_{k-1} - \frac{1}{8} x_{k-2} - \frac{1}{16} x_{k-3} - \frac{5}{128} x_{k-4} - \dots \\ \Delta^{-1/2} x_k &= x_k + \frac{1}{2} x_{k-1} + \frac{3}{8} x_{k-2} + \frac{5}{16} x_{k-3} + \frac{35}{128} x_{k-4} + \dots \\ \Delta^{2/3} x_k &= x_k - \frac{2}{3} x_{k-1} - \frac{1}{9} x_{k-2} - \frac{4}{81} x_{k-3} - \frac{7}{243} x_{k-4} - \dots \end{aligned}$$

The idea of constructing new sequence spaces via infinite matrices started with Kızmaz's study [20] and then it has been developed by numerous researchers using different triangles [2], [4], [6], [7], [18].

In the studies [3], [5], [16], [17], [21], [25], [26] [27] and [30] different difference sequence spaces have been studied based on some newly defined infinite matrices. Some new results on the visualization and animations of the topologies of certain sequence spaces have been illustrated in [9], [11], [12]. The authors applied their software package for this purpose. Note that, those results have an interesting and important application in crystallography.

Many authors have made efforts to apply Hausdorff measure of noncompactness to find compactness conditions of certain sets of sequences during the past decade [8], [9], [15] and [28].

Fractional difference sequence spaces have been studied in the literature recently [19], [31], [33]. The authors of those papers especially studied on the properties of fractional operators in their researches in addition to focusing on certain fractional sequence spaces.

In this work, we consider the fractional sequence spaces $\ell_p(\Delta^{(\tilde{\alpha})})$ for $1 \leq p < \infty$ and determine norm operators of our spaces. We establish some identities or estimates for the Hausdorff measures of noncompactness of certain operators on difference sequence spaces $\ell_p(\Delta^{(\tilde{\alpha})})$ of fractional orders. We characterize some classes of compact operators on those spaces. Note that, we can only obtain sufficient conditions when the final space is ℓ_∞ . However, we use the results in [24] and [34] to obtain necessary and sufficient conditions for the classes of compact matrix operators from $\ell_p(\Delta^{(\tilde{\alpha})})$ spaces into the sets of bounded sequences and for the classes $(\ell_1(\Delta^{(\tilde{\alpha})}), \ell_\infty)$ and $(\ell_\infty(\Delta^{(\tilde{\alpha})}), \ell_\infty)$.

2. PRELIMINARIES

The β dual of a set X is defined by

$$X^\beta = \{a \in \omega : a \cdot x \in cs \text{ for all } x \in X\}.$$

Note that $c_0^\beta = c^\beta = \ell_\infty^\beta = \ell_1$ and $\ell_p^\beta = \ell_q$.

Given any infinite matrix $A = (a_{nk})_{n,k=0}^\infty$ of complex numbers and any sequence x , we write $A_n = (a_{nk})_{k=0}^\infty$ for the sequence in the n^{th} row of A , $A_n x = \sum_{k=0}^\infty a_{nk} x_k$ ($n = 0, 1, \dots$) and $Ax = (A_n x)_{n=0}^\infty$, provided $A_n \in X^\beta$ for all n .

If X and Y are subsets of ω , then

$$X_A = \{x \in \omega : Ax \in X\}$$

denotes the matrix domain of A in X and (X, Y) is the class of all infinite matrices that map X into Y ; so $A \in (X, Y)$ if and only if $X \subset Y_A$.

Consider now the following fractional difference sequence spaces for $1 \leq p < \infty$:

$$\ell_p(\Delta^{(\tilde{\alpha})}) := \left\{ x = (x_k) \in \omega : \sum_{n=0}^\infty \left| \sum_{i=0}^\infty (-1)^i \frac{\Gamma(\tilde{\alpha} + 1)}{i! \Gamma(\tilde{\alpha} - i + 1)} x_{n-i} \right|^p < \infty \right\}.$$

Now let us define the sequence $y = (y_k)$ which will be used, by the $\Delta^{(\tilde{\alpha})}$ -transform of a sequence $x = (x_k)$, that is,

$$\begin{aligned} y_k &= x_k - \tilde{\alpha} x_{k-1} + \frac{\tilde{\alpha}(\tilde{\alpha} - 1)}{2!} x_{k-2} - \frac{\tilde{\alpha}(\tilde{\alpha} - 1)(\tilde{\alpha} - 2)}{3!} x_{k-3} + \dots \\ &= \sum_{i=0}^\infty (-1)^i \frac{\Gamma(\tilde{\alpha} + 1)}{i! \Gamma(\tilde{\alpha} - i + 1)} x_{k-i}. \end{aligned}$$

Hence, those spaces can be considered as the matrix domains of the triangle $\Delta^{(\tilde{\alpha})}$ in the classical sequence spaces ℓ_p , where $1 \leq p < \infty$. We also have the following relation between the sequences $x = (x_k)$ and $y = (y_k)$:

$$x_k = \sum_{i=0}^\infty (-1)^i \frac{\Gamma(-\tilde{\alpha} + 1)}{i! \Gamma(-\tilde{\alpha} - i + 1)} y_{k-i}.$$

Lemma 2.1. [1, Theorem 4.3.12, p. 63] Let $(X, \|\cdot\|)$ be a BK space. Then X_T is a BK space with $\|\cdot\|_T = \|T(\cdot)\|$.

By Lemma 2.1, defined fractional difference sequence space is a complete, linear, BK space with the following norm:

$$\|x\| = \left(\sum_n \left| \sum_{i=0}^\infty (-1)^i \frac{\Gamma(\tilde{\alpha} + 1)}{i! \Gamma(\tilde{\alpha} - i + 1)} x_{n-i} \right|^p \right)^{\frac{1}{p}}.$$

Let X be a normed space. Then $S_X = \{x \in X : \|x\| = 1\}$ and $\bar{B}_X = \{x \in X : \|x\| \leq 1\}$ denote the unit sphere and closed unit ball in X , where X is a normed space.

By \mathcal{F}_r ($r = 0, 1, \dots$), we denote the subcollection of \mathcal{F} consisting of all nonempty and finite subsets of \mathbb{N} with terms that are greater than r , that is

$$\mathcal{F}_r = \{N \in \mathcal{F} : n > r \text{ for all } n \in N\} \quad (r = 0, 1, \dots).$$

Given $a \in \omega$, we write

$$\|a\|_X^* = \sup_{x \in S_X} \left| \sum_{k=1}^{\infty} a_k x_k \right|$$

provided the expression on the write hand side is defined and finite which is the case whenever X is a BK space and $a \in X^\beta$.

Lemma 2.2. *Let X and Y be BK spaces.*

- (i) *Then we have $(X, Y) \subset \mathcal{B}(X, Y)$, that is, every $A \in (X, Y)$ defines an operator $L_A \in \mathcal{B}(X, Y)$, where $L_A(x) = Ax$ for all $x \in X$ (see [14, Theorem 1.23]).*
- (ii) *We have $\|x\|_X^* = \|x\|_{X^\beta}$ for all $x \in X^\beta$, where $\|\cdot\|_{X^\beta}$ is the natural norm on the dual set X^β (see [13, Theorem 3.2]).*

Lemma 2.3. *Let Y be an arbitrary subset of ω and X be a BK space with AK or $X = \ell_\infty$, and $R = S^t$. Then $A \in (X_T, Y)$ if and only if $\hat{A} \in (X, Y)$ and $W^{(n)} \in (X, c_0)$ for all $n = 0, 1, \dots$. Here \hat{A} is the matrix with rows $\hat{A}_n = RA_n$ for $n = 0, 1, \dots$, and the triangles $W^{(n)}$ ($n = 0, 1, \dots$) are defined by $w_{mk}^{(n)} = \sum_{j=m}^{\infty} a_{nj} s_{jk}$.*

Moreover, if $A \in (X, Y)$ then we have $Az = \hat{A}(Tz)$ for all $z \in Z = X_T$ (see [13], Theorem 3.4, Remark 3.5(a)).

We have the following results the for operator norms of bounded operators by [22, Theorem 2.8].

Lemma 2.4. *Let X be a BK space.*

- (i) *If $A \in (X, \ell_1)$, then*

$$\|A\|_{(X, \ell_1)} \leq \|L_A\| \leq 4 \cdot \|A\|_{(X, \ell_1)},$$

where

$$\|A\|_{(X, \ell_1)} = \sup_{N \in \mathcal{F}} \left\| \sum_{n \in N} A_n \right\|_X^* < \infty.$$

- (ii) *If $A \in (X, Y)$, then*

$$\|L_A\| = \|A\|_{(X, \ell_\infty)} = \sup_n \|A_n\|_X^* < \infty,$$

where Y is any of the spaces c_0, c, ℓ_∞ .

We obtain the following result as an immediate consequence of Lemma 2.2(ii).

Lemma 2.5. *Let $1 \leq p < \infty$, then we have $\|x\|_{\ell_p}^* = \|x\|_{\ell_q}$ for all $x = (x_k) \in \ell_q$.*

Remark 2.6. *Let $1 \leq p < \infty$, A be an infinite matrix and X be BK space. If $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), Y)$, then $\hat{A} \in (\ell_p, Y)$ such that $Ax = \hat{A}y$ for all $x \in \ell_p(\Delta^{(\tilde{\alpha})})$ and $y \in \ell_p$. Here x*

and y connected by (2.1) and $\hat{A} = (\hat{a}_{nk})$ is defined by

$$(2.1) \quad \hat{a}_{nk} = \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha} + 1)}{(j-k)! \Gamma(-\tilde{\alpha} + j - k + 1)} a_{nj} \text{ for all } n, k \in \mathbb{N}_0;$$

and for $k = 0, 1, \dots$ we also define $\hat{\alpha} = (\hat{\alpha}_k)_{k=0}^{\infty}$ by

$$(2.2) \quad \hat{\alpha}_k = \lim_{n \rightarrow \infty} \hat{a}_{nk}.$$

Proof. Let $1 \leq p < \infty$, $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), Y)$ and $x \in \ell_p(\Delta^{(\tilde{\alpha})})$. So, $A_n \in (\ell_p(\Delta^{(\tilde{\alpha})}))^\beta$ for all $n = 0, 1, \dots$. Then, we have $\hat{A}_n \in \ell_p^\beta = \ell_q$ for all $n = 0, 1, \dots$ and $Ax = \hat{A}y$ satisfies. Therefore, $\hat{A}y \in Y$ and this means $\hat{A} \in (\ell_p, Y)$. \square

Lemma 2.7. *If $a = (a_k) \in (\ell_p(\Delta^{(\tilde{\alpha})}))^\beta$, then $\bar{a} = (\bar{a}_k) \in \ell_q$ and the equality*

$$(2.3) \quad \sum_{k=0}^{\infty} a_k x_k = \sum_{k=0}^{\infty} \bar{a}_k y_k.$$

satisfies for every $x = (x_k) \in \ell_p(\Delta^{(\tilde{\alpha})})$, where

$$(2.4) \quad \bar{a}_k = \sum_{i=k}^{\infty} (-1)^{i-k} \frac{\Gamma(-\tilde{\alpha} + 1)}{(i-k)! \Gamma(-\tilde{\alpha} - i + k + 1)} a_i.$$

Let us now write $S = S_{\ell_p}$ and $\hat{S} = S_{\ell_p(\Delta^{(\tilde{\alpha})})}$, for short.

Lemma 2.8. *Let $1 \leq p < \infty$ and $\bar{a} = (\bar{a}_k)$ be defined in (2.4), then we have*

$$\|a\|_{\ell_p(\Delta^{(\tilde{\alpha})})}^* = \|\bar{a}\|_{\ell_q} = \begin{cases} \left(\sum_{k=0}^{\infty} |\bar{a}_k|^q \right)^{\frac{1}{q}} & (1 < p \leq \infty) \\ \sup_k |\bar{a}_k| & (p = 1). \end{cases}$$

for all $a = (a_k) \in (\ell_p(\Delta^{(\tilde{\alpha})}))^\beta$.

Proof. Assume that $\bar{a} \in (\ell_p(\Delta^{(\tilde{\alpha})}))^\beta$. So, we have by Lemma 2.7 that $\bar{a} \in \ell_q$ and the equality (2.3) yields for all sequences $x = (x_k) \in (\ell_p(\Delta^{(\tilde{\alpha})}))$ and $y = (y_k) \in \ell_p$ which are connected by (2.1). Additionally, we have by Lemma 2.5 that $x \in \hat{S}$ if and only if $y \in S$. Hence, we have by (2.3) that

$$(2.5) \quad \|a\|_{\ell_p(\Delta^{(\tilde{\alpha})})}^* = \sup_{x \in \hat{S}} \left| \sum_{k=0}^{\infty} a_k x_k \right| = \sup_{y \in S} \left| \sum_{k=0}^{\infty} \bar{a}_k y_k \right| = \|\bar{a}\|_{\ell_q}^* < \infty.$$

Moreover, we obtain by (2.5) and Lemma 2.5 that

$$\|a\|_{\ell_p(\Delta^{(\tilde{\alpha})})}^* = \|\bar{a}\|_{\ell_q}^* = \|\bar{a}\|_{\ell_q}$$

because $\bar{a} \in \ell_q$. This completes the proof. \square

We obtain estimates or identities for the norms of the matrix operators for the classes $(\ell_p(\Delta^{(\tilde{\alpha})}), c_0)$, $(\ell_p(\Delta^{(\tilde{\alpha})}), c)$, $(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)$ and $(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_1)$.

Theorem 2.9. *Let A be in any of the classes $(\ell_p(\Delta^{(\tilde{\alpha})}), c_0)$, $(\ell_p(\Delta^{(\tilde{\alpha})}), c)$ or $(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)$, then*

$$\|L_A\| = \|A\|_{(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)},$$

where

$$\|A\|_{(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)} = \sup_n \left(\sum_{k=0}^{\infty} |\hat{a}_{nk}|^q \right)^{\frac{1}{q}} < \infty$$

Proof. This is an immediate consequence of Lemma 2.2 and Lemma 2.4. \square

Theorem 2.10. *Let $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), \ell_1)$, then*

$$\|A\|_{(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_1)} \leq \|L_A\| \leq 4 \cdot \|A\|_{(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_1)},$$

where

$$\|A\|_{(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_1)} = \sup_{N \in \mathcal{F}} \left(\sum_{k=0}^{\infty} \left| \sum_{n \in N} \hat{a}_{nk} \right| \right) < \infty.$$

Proof. This is an immediate consequence of Lemma 2.2 and Lemma 2.4. \square

3. MAIN RESULTS RELATED TO COMPACT OPERATORS

The following notations are needed to establish estimates and identities for the Hausdorff measure of noncompactness of matrix operators and characterize the classes of compact operators. We also use the results in Katarina's paper [24] to prove our results.

We recall the definition of the Hausdorff measure of noncompactness of bounded subsets of a metric space, and the Hausdorff measure of noncompactness of operators between Banach spaces.

If X and Y are infinite-dimensional complex Banach spaces then a linear operator $L : X \rightarrow Y$ is said to be compact if the domain of L is all of X , and, for every bounded sequence (x_n) in X , the sequence $(L(x_n))$ has a convergent subsequence. We denote the class of such operators by $\mathcal{C}(X, Y)$.

Definition 3.1. Let (X, d) be a metric space, $B(x_0, \delta) = \{x \in X : d(x, x_0) < \delta\}$ denote the open ball of radius $\delta > 0$ and center in $x_0 \in X$, and \mathcal{M}_X be the collection of bounded sets in X . The Hausdorff measure of noncompactness of $Q \in \mathcal{M}_X$ is

$$\chi(Q) = \inf\{\epsilon > 0 : Q \subset \bigcup_{k=1}^n B(x_k, \delta_k) : x_k \in X, \delta_k < \epsilon, 1 \leq k \leq n, n \in \mathbb{N}\}.$$

Let X and Y be Banach spaces and χ_1 and χ_2 be measures of noncompactness on X and Y . Then the operator $L : X \rightarrow Y$ is called (χ_1, χ_2) -bounded if $L(Q) \in \mathcal{M}_Y$ for every $Q \in \mathcal{M}_X$ and there exists a positive constant C such that

$$(3.1) \quad \chi_2(L(Q)) \leq C\chi_1(Q) \text{ for every } Q \in \mathcal{M}_X.$$

If an operator L is (χ_1, χ_2) -bounded then the number

$$\|L\|_{(\chi_1, \chi_2)} = \inf \{C \geq 0 : (3.1) \text{ holds for all } Q \in \mathcal{M}_X\}$$

is called the (χ_1, χ_2) -measure of noncompactness of L . In particular, if $\chi_1 = \chi_2 = \chi$, then we write $\|L\|_\chi$ instead of $\|L\|_{(\chi, \chi)}$.

Lemma 3.2. [14, Corollary 2.26.] *Let X and Y be Banach spaces and $L \in \mathcal{B}(X, Y)$. Then we have*

$$(3.2) \quad \|L\|_\chi = \chi(L(\bar{B}_X)) = \chi(L(S_X)),$$

$$(3.3) \quad L \in \mathcal{C}(X, Y) \text{ if and only if } \|L\|_\chi = 0.$$

Lemma 3.3. [14, Theorem 2.23.] *Let X be a Banach space with Schauder basis $(b_n)_{n=0}^\infty$, $Q \in \mathcal{M}_X$, $P_n : X \rightarrow X$ be the projector onto the linear span of $\{b_0, b_1, \dots, b_n\}$. I be the identity map on X and $R_n = I - P_n$ ($n = 0, 1, \dots$). Then we have*

$$(3.4) \quad \frac{1}{a} \cdot \limsup_{n \rightarrow \infty} \left(\sup_{x \in Q} \|R_n(x)\| \right) \leq \chi(Q) \leq \limsup_{n \rightarrow \infty} \left(\sup_{x \in Q} \|R_n(x)\| \right),$$

where $a = \limsup_{n \rightarrow \infty} \|R_n\|$.

Lemma 3.4. [14, Theorem 2.8.] *Let Q be a bounded subset of the normed space X , where X is ℓ_p for $1 \leq p < \infty$ or c_0 . If $P_n : X \rightarrow X$ is the operator defined by $P_n(x) = x^{[n]}$ for $x = (x_k)_{k=0}^\infty \in X$, then we have*

$$\chi(Q) = \lim_n \left(\sup_{x \in Q} \|R_n(x)\| \right).$$

When the first space is ℓ_1 , a problem takes place since β dual of ℓ_1 is ℓ_∞ , which has no AK . In this case, the following study of Sargent [34] is used in order to characterize compact operators.

Lemma 3.5. [34, Theorem 5.] *If $L \in B(\ell_1, \ell_\infty)$, then L is compact if and only if*

$$\lim_{m \rightarrow \infty} \sup_{1 \leq n \leq m} |a_{n, k_1} - a_{n, k_2}| = \sup_n |a_{n, k_1} - a_{n, k_2}|$$

uniformly in k_1 and k_2 , ($1 \leq k_1, k_2 < \infty$).

The following results (by [22, Corollary 3.6] and [28, Theorems 3.7 and 3.11]) are needed to determine estimates for the norms of continuous linear operators L_A on our spaces and establish necessary and sufficient conditions for a matrix operator to be a compact operator.

Lemma 3.6. *Then we have:*

(i) *If $A \in (X, c_0)$, then we have*

$$\|L_A\|_\chi = \limsup_{n \rightarrow \infty} \|A_n\|_X^*.$$

(ii) *If $A \in (X, c)$, then we have*

$$\frac{1}{2} \cdot \limsup_{n \rightarrow \infty} \|A_n - \alpha\|_X^* \leq \|L_A\|_\chi \leq \limsup_{n \rightarrow \infty} \|A_n - \alpha\|_X^*.$$

(iii) If $A \in (X, \ell_1)$, then we have

$$\lim_{r \rightarrow \infty} \left(\sup_{N \in \mathcal{F}_r} \left\| \sum_{n \in N} A_n \right\|_X^* \right) \leq \|L_A\|_\chi \leq 4 \cdot \lim_{r \rightarrow \infty} \left(\sup_{N \in \mathcal{F}_r} \left\| \sum_{n \in N} A_n \right\|_X^* \right).$$

We are now ready to state the main results related to compact operators. We start with establishing some estimates for the norms of bounded linear operators L_A on the given fractional sequence spaces $\ell_p(\Delta^{(\tilde{\alpha})})$.

Theorem 3.7. Let $1 < p < \infty$ and $q = \frac{p}{p-1}$.

(i) If $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), c_0)$, then we have

$$(3.5) \quad \|L_A\|_\chi = \lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sum_k |\hat{a}_{nk}|^q \right)^{\frac{1}{q}}.$$

(ii) If $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), c)$, then we have

$$(3.6) \quad \frac{1}{2} \cdot \lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sum_k |\hat{a}_{nk} - \hat{\alpha}_k|^q \right)^{\frac{1}{q}} \leq \|L_A\|_\chi \leq \lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sum_k |\hat{a}_{nk} - \hat{\alpha}_k|^q \right)^{\frac{1}{q}}.$$

(iii) If $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), \ell_1)$, then we have

$$(3.7) \quad \lim_{r \rightarrow \infty} \left(\sup_{N \in \mathcal{F}_r} \left(\sum_k \left| \sum_{n \in N} \hat{a}_{nk} \right|^q \right)^{\frac{1}{q}} \right) \leq \|L_A\|_\chi \leq 4 \cdot \lim_{r \rightarrow \infty} \left(\sup_{N \in \mathcal{F}_r} \left(\sum_k \left| \sum_{n \in N} \hat{a}_{nk} \right|^q \right)^{\frac{1}{q}} \right).$$

Proof. Let $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), c_0)$. By Lemma 2.8, we have

$$\|A\|_{\ell_p(\Delta^{(\tilde{\alpha})})}^* = \|\hat{A}_n\|_{\ell_q} = \left(\sum_{k=0}^{\infty} |\hat{a}_{nk}|^q \right)^{\frac{1}{q}}.$$

for all $n = 0, 1, \dots$ because $A_n \in (\ell_p(\Delta^{(\tilde{\alpha})}))^\beta$ for all $n = 0, 1, \dots$. Taking into account this result and Lemma 3.6(i) we obtain (3.5).

In order to prove the second part we start with $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), c)$. By Remark 2.1 $\hat{A} \in (\ell_p, c)$. So, we have by Lemma 3.6(ii) that

$$\frac{1}{2} \cdot \limsup_{n \rightarrow \infty} \|\hat{A}_n - \hat{\alpha}\|_{\ell_q} \leq \|L_A\|_\chi \leq \limsup_{n \rightarrow \infty} \|\hat{A}_n - \hat{\alpha}\|_{\ell_q},$$

where $\hat{\alpha}$ defined in (2.2). Hence, by Lemma 3.2 we get

$$(3.8) \quad \|L_A\|_\chi = \chi(L_A(\hat{S})) = \chi(A\hat{S}),$$

and

$$(3.9) \quad \|L_{\hat{A}}\|_\chi = \chi(L_{\hat{A}}(S)) = \chi(\hat{A}S).$$

Moreover, $x \in \hat{S}$ if and only if $y \in S$. We have $A\hat{S} = \hat{A}S$ because $Ax = \hat{A}y$ by Remark 2.1. Taking into account the equalities (3.8) and (3.9) we conclude that $\|L_A\|_\chi = \|\hat{L}_{\hat{A}}\|_\chi$ which completes the second part of the proof.

Finally, let $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), \ell_1)$. We derive from Lemma 2.5 that

$$(3.10) \quad \left\| \sum_{n \in N} A_n \right\|_{\ell_p(\Delta^{(\tilde{\alpha})})}^* = \left\| \sum_{n \in N} \hat{A}_n \right\|_{\ell_q}$$

because $A_n \in (\ell_p(\Delta^{(\tilde{\alpha})}))^\beta$. Hence, we obtain the condition in (3.7) by (3.10) and Lemma 3.6(iii). \square

Theorem 3.8. *Let $1 \leq p < \infty$.*

(i) *If $A \in (\ell_1(\Delta^{(\tilde{\alpha})}), c_0)$ then we have*

$$(3.11) \quad \|L_A\|_\chi = \lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sup_k |\hat{a}_{nk}| \right).$$

(ii) *If $A \in (\ell_1(\Delta^{(\tilde{\alpha})}), c)$ then we have*

$$\frac{1}{2} \cdot \lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sup_k |\hat{a}_{nk} - \hat{\alpha}_k| \right) \leq \|L_A\|_\chi \leq \lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sup_k |\hat{a}_{nk} - \hat{\alpha}_k| \right).$$

Proof. This is an immediate consequence of Theorem 2.9, Lemma 3.6 and Theorem 3.7. \square

We now use the results in [24] and [34] to obtain necessary and sufficient conditions for the classes of compact matrix operators in $(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)$, $(\ell_1(\Delta^{(\tilde{\alpha})}), \ell_\infty)$ and $(\ell_\infty(\Delta^{(\tilde{\alpha})}), \ell_\infty)$.

Theorem 3.9. *Let $1 < p < \infty$ and $q = \frac{p}{p-1}$.*

(i) *If $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)$, then L_A is compact if and only if*

$$\lim_{r \rightarrow \infty} \sup_n \left(\sum_{k=r+1}^{\infty} |\hat{a}_{nk}|^q \right)^{\frac{1}{q}} = 0.$$

(ii) *If $A \in (\ell_1(\Delta^{(\tilde{\alpha})}), \ell_\infty)$, then L_A is compact if and only if*

$$\lim_{m \rightarrow \infty} \sup_{1 \leq n \leq m} |\hat{a}_{n,k_1} - \hat{a}_{n,k_2}| = \sup_n |\hat{a}_{n,k_1} - \hat{a}_{n,k_2}|$$

uniformly in k_1 and k_2 , $(1 \leq k_1, k_2 < \infty)$.

Proof. Assume $A \in (\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)$. By Lemma 2.3, we obtain $L_A \in \mathcal{B}(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)$ and $\hat{A} \in (\ell_p, \ell_\infty)$ such that $Ax = \hat{A}(\Delta^{(-\tilde{\alpha})}x)$ for all $x \in \ell_p(\Delta^{(\tilde{\alpha})})$, where $\Delta^{(-\tilde{\alpha})}$ is the inverse of the triangle $\Delta^{(\tilde{\alpha})}$ and \hat{a}_{nk} defined in (2.1). We have $L_{\hat{A}} \in \mathcal{B}(\ell_p, \ell_\infty)$ and $\hat{A}x = L_{\hat{A}}(x)$ for all $x \in \ell_p$ because ℓ_p is a BK space. If we write $(\hat{A}_n)_r$ for the r th row of the matrix \hat{A} with \hat{a}_{nk} replaced by 0 for $k > r$,

$$(\hat{A}_n)_r = \sum_{k=0}^r \hat{a}_{nk} e^{(k)}$$

and apply [34, (b), p.85] we obtain $L_{\hat{A}}$ is compact if and only if

$$\sup_n \|\hat{A}_n\|_q = \sup_n \left(\sum_k |\hat{a}_{nk}|^q \right)^{\frac{1}{q}} < \infty$$

and

$$\lim_{r \rightarrow \infty} \sup_n \|\hat{A}_n - (\hat{A}_n)_r\|_q = \lim_{r \rightarrow \infty} \sup_n \left(\sum_{k=r+1}^{\infty} |\hat{a}_{nk}|^q \right)^{\frac{1}{q}} = 0.$$

The first condition yields from the characterizations of matrix transformations, [1, Example 8.4.5D and Example 8.4.6D]. Hence $L_{\hat{A}}$ is compact if and only if the condition

$$\lim_{r \rightarrow \infty} \sup_n \left(\sum_{k=r+1}^{\infty} |\hat{a}_{nk}|^q \right)^{\frac{1}{q}} = 0$$

yields. Taking into account the equalities (3.8) and (3.9) we have L_A is compact if and only if $L_{\hat{A}}$ is compact. This completes the first part of the proof.

We know that L_A is compact if and only if $L_{\hat{A}}$ is compact with the equalities (3.8) and (3.9). Also $A \in (\ell_1(\Delta^{(\hat{\alpha})}), \ell_\infty)$ implies $\hat{A} \in (\ell_1, \ell_\infty)$. As an immediate consequence of Lemma 3.5 we prove the second part of the theorem. \square

Theorem 3.10. *Let $A \in (\ell_\infty(\Delta^{(\hat{\alpha})}), \ell_\infty)$. Then L_A is compact if and only if*

$$(3.12) \quad \lim_{r \rightarrow \infty} \sup_n \sum_{k=r+1}^{\infty} |\hat{a}_{nk}| = 0.$$

Proof. By the help of Lemma 2.3 we have $\hat{A} \in (\ell_\infty, \ell_\infty)$ because ℓ_∞ is a BK space and also by Lemma 2.2 the associated bounded linear operator $L_{\hat{A}}$. Since $(\ell_\infty)^\beta = \ell_1$, therefore by the condition (b) in [34, p. 85] we have: The spaces ℓ_∞ and c_0 have the same dual space ℓ_1 , and therefore the conditions [34, (b) and (f), p.85] are the same. It follows that $L_{\hat{A}}$ is compact if and only if

$$\sup_n \|\hat{A}_n\|_1 = \sup_n \sum_{k=0}^{\infty} |\hat{a}_{nk}| < \infty$$

and

$$\lim_{r \rightarrow \infty} \sup_n \|\hat{A}_n - (\hat{A}_n)_r\|_1 = \lim_{r \rightarrow \infty} \sup_n \sum_{k=r+1}^{\infty} |\hat{a}_{nk}| = 0.$$

So the first condition is satisfied [1, Example 8.4.5A], and it follows that $L_{\hat{A}}$ is compact if and only if the condition (3.12) yields. This completes the proof because L_A is compact if and only if $L_{\hat{A}}$ is compact by Lemma 2.2 and the equalities (3.8) and (3.9). \square

4. CONCLUSION

The main aim of this paper is to consider fractional sequence spaces $\ell_p(\Delta^{(\tilde{\alpha})})$ for $1 \leq p < \infty$ via Euler gamma function. We derive some estimates and identities for the norms of bounded linear operators on fractional sequence spaces $\ell_p(\Delta^{(\tilde{\alpha})})$. In general, Hausdorff measure of noncompactness is used to find necessary and sufficient conditions for a matrix operator on a given sequence space to be a compact operator. However we can only obtain sufficient conditions when the final space is ℓ_∞ , that is for a matrix class (X, ℓ_∞) . This is why we use the results in [24] and [34] to exactly obtain necessary and sufficient conditions for the classes of compact matrix operators in $(\ell_p(\Delta^{(\tilde{\alpha})}), \ell_\infty)$ and $(\ell_\infty(\Delta^{(\tilde{\alpha})}), \ell_\infty)$. In addition to these characterizations, we apply Hausdorff measure of noncompactness to establish necessary and sufficient conditions for a matrix operator to be a compact operator from fractional sequence spaces $\ell_p(\Delta^{(\tilde{\alpha})})$ and $\ell_1(\Delta^{(\tilde{\alpha})})$ into Y , where Y is any of the spaces of convergent to zero, convergent and bounded sequences c_0, c, ℓ_1 .

All these results are given with the following table:

To	From	$\ell_p(\Delta^{(\tilde{\alpha})})$	$\ell_1(\Delta^{(\tilde{\alpha})})$
c_0		1.	5.
c		2.	6.
ℓ_∞		3.	7.
ℓ_1		4.	-

Table. Necessary and sufficient conditions for an operator to be compact

- [1.] $\lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sum_{k=0}^{\infty} \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)! \Gamma(-\tilde{\alpha}+j-k+1)} a_{nj} \right|^q \right)^{\frac{1}{q}} = 0.$
- [2.] $\lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sum_{k=0}^{\infty} \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)! \Gamma(-\tilde{\alpha}+j-k+1)} a_{nj} - \lim_{n \rightarrow \infty} \hat{a}_{nk} \right|^q \right)^{\frac{1}{q}} = 0.$
- [3.] $\lim_{r \rightarrow \infty} \sup_n \left(\sum_{k=r+1}^{\infty} \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)! \Gamma(-\tilde{\alpha}+j-k+1)} a_{nj} \right|^q \right)^{\frac{1}{q}} = 0.$
- [4.] $\lim_{r \rightarrow \infty} \sup_{N \in \mathcal{F}_r} \left(\sum_{k=0}^{\infty} \left| \sum_{n \in N} \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)! \Gamma(-\tilde{\alpha}+j-k+1)} a_{nj} \right|^q \right)^{\frac{1}{q}} = 0.$
- [5.] $\lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sup_k \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)! \Gamma(-\tilde{\alpha}+j-k+1)} a_{nj} \right| \right) = 0.$
- [6.] $\lim_{r \rightarrow \infty} \sup_{n \geq r} \left(\sup_k \left| \sum_{j=k}^{\infty} (-1)^{j-k} \frac{\Gamma(-\tilde{\alpha}+1)}{(j-k)! \Gamma(-\tilde{\alpha}+j-k+1)} a_{nj} - \lim_{n \rightarrow \infty} \hat{a}_{nk} \right| \right) = 0.$
- [7.] $\lim_{m \rightarrow \infty} \sup_{1 \leq n \leq m} |\hat{a}_{n,k_1} - \hat{a}_{n,k_2}| = \sup_n |\hat{a}_{n,k_1} - \hat{a}_{n,k_2}|,$
uniformly in k_1 and k_2 , ($1 \leq k_1, k_2 < \infty$).

COMPLIANCE WITH ETHICAL STANDARDS

Conflict of interest. The authors declare that they have no conflict of interest.

Ethical approval. This article does not contain any studies with human participants or animals performed by any of the authors.

REFERENCES

- [1] A. Wilansky, Summability through Functional Analysis, North-Holland Mathematics Studies 85, Amsterdam, New York, Oxford, 1984.
- [2] A. Karaisa and F. Özger, Almost difference sequence space derived by using a generalized weighted mean. *J. Comput. Anal. Appl.* **19**(1), (2015) 27–38.
- [3] A. Karaisa and F. Özger, On almost convergence and difference sequence spaces of order m with core theorems, *Gen. Math. Notes*, **26**(1), (2015), 102–125.
- [4] B. Altay and F. Başar, Certain topological properties and duals of the matrix domain of a triangle matrix in a sequence space, *J. Math. Anal. Appl.* **336**(1) (2007), 632–645.
- [5] C. Aydın and F. Başar, Some new difference sequence spaces. *Appl. Math. Comput.* **157**(3) (2004), 677693.
- [6] E. Malkowsky and F. Özger, Some mixed paranorm spaces, *Filomat* **31**(4) (2017), 1079–1098.
- [7] E. Malkowsky and F. Özger, A note on some sequence spaces of weighted means, *Filomat* **26**(3) (2012), 511–518.
- [8] E. Malkowsky and F. Özger, Compact operators on spaces of sequences of weighted means, *AIP Conf. Proc.* **1470** (2012), 179–182.
- [9] E. Malkowsky, F. Özger and A. Alotatibi, Some notes on matrix mappings and their Hausdorff measure of noncompactness, *Filomat* **285** (2014), 1059–1072.
- [10] E. Malkowsky, F. Özger and V. Veličković, Some spaces related to Cesaro sequence spaces and an application to crystallography, *MATCH Commun. Math. Comput. Chem.* **70**(3) (2013), 867–884.
- [11] E. Malkowsky, F. Özger and V. Veličković, Matrix transformations on mixed paranorm spaces, *Filomat* **31**(10) (2017), 2957–2966.
- [12] E. Malkowsky, F. Özger and V. Veličković, Visualization of the spaces $W(u, v; \ell_p)$ and their duals, *AIP Conf. Proc.* **1759** (2016), doi: 10.1063/1.4959634.
- [13] E. Malkowsky and V. Rakočević, On matrix domains of triangles, *Appl. Math. Comput.* **189** (2007), 1148–1163.
- [14] E. Malkowsky and V. Rakočević, An introduction into the theory of sequence spaces and measures of noncompactness, *Zb. Rad. (Beogr.)* **9**(17) (2000), 143–234.
- [15] E.E. Kara, M. Başarır and M. Mursaleen, Compactness of matrix operators on some sequence spaces derived by Fibonacci Numbers, *Kragujevac Journal of Math.* **39**(2) (2015) 217–230.
- [16] F. Özger and F. Başar, Domain of the double sequential band matrix $B(\tilde{r}, \tilde{s})$ on some Maddox’s spaces, *AIP Conf. Proc.* **1470** (2012), 152–155.
- [17] F. Özger and F. Başar, Domain of the double sequential band matrix $B(\tilde{r}, \tilde{s})$ on some Maddox’s spaces, *Acta Mathematica Scientia* **34**(2) (2014), 394–408.
- [18] F. Nuray, U. Ulusu and E. Dndar, Lacunary statistical convergence of double sequences of sets. *Soft Comput.* **20** (2016), 2883–2888.
- [19] H. Furkan, On some λ difference sequence spaces of fractional order, *J. Egypt. Math. Soc.* **25** (2017), 37–42.
- [20] H. Kızmaz, *On certain sequence spaces*, *Canad. Math. Bull.* **24**(2) (1981), 169–176.
- [21] H. Polat and B. Altay, On some new Euler difference sequence spaces. *Southeast Asian Bull Math.* **30** (2006), 209–220.
- [22] I. Djolović and E. Malkowsky, A note on compact operators on matrix domains, *J. Math. Anal. Appl.* **340** (2008), 291–303.
- [23] I. Djolović and E. Malkowsky, Matrix transformations and compact operators on some new m th order difference sequences, *Appl. Math. Comput.* **198**(2) (2008), 700–714.
- [24] K. Petković, Some new results related to compact matrix operators in the classes $A \in ((\ell_p)_T, \ell_\infty)$, *Acta Mathematica Sinica* **31**(8), (2015), 1339–1347.

- [25] M. Candan, Domain of the double sequential band matrix in the classical sequence spaces, *J. Inequal. Appl.* **2012**(281) (2012), doi:10.1186/1029-242X-2012-281.
- [26] M. Kirişçi and F. Başar, Some new sequence spaces derived by the domain of generalized difference matrix, *Comput. Math. Appl.* **60**(5) (2010), 1299–1309.
- [27] M. Mursaleen, V. Karakaya, H. Polat and N. Simek, Measure of noncompactness of matrix operators on some difference sequence spaces of weighted means, *Comput. Math. Appl.* **62**(2) (2011), 814–820.
- [28] M. Mursaleen, A.K. Noman, Compactness by Hausdorff the measures of noncompactness, *Nonlinear Analysis*, 73, (2010) 2541-2557.
- [29] M. Stieglitz and H. Tietz, Matrixtransformationen von Folgenräumen eine Ergebnisübersicht, *Math. Z.* **154** (1977), 1–16.
- [30] E. Yavuz and Ö. Talo, Abel summability of sequences of fuzzy numbers, *Soft Comput.* **20** (2016), 1041-1046.
- [31] P. Baliarsingh and S. Dutta, On the classes of fractional order of difference sequence spaces and matrix transformations, *Appl. Math. Comput.* **250** (2015), 665–674.
- [32] R. Çolak and M. Et, On some generalized difference sequence spaces and related matrix transformations. *Hokkaido Math J.* **26**(3) (1997), 483-492.
- [33] U. Kadak and P. Baliarsingh, On certain Euler difference sequence spaces of fractional order and related dual properties, *J. Nonlinear Sci. Appl.* **8** (2015), 997–1004.
- [34] W. L. C. Sargent, On compact matrix transformations between sectionally bounded *BK* spaces, *J. London Math. Soc.* 41, (1966) 79-87.

(Özger) DEPARTMENT OF ENGINEERING SCIENCES, İZMİR KATIP ÇELEBİ UNIVERSITY, İZMİR, TURKEY
E-mail address, Özger: farukozger@gmail.com