

MEASURE-GEOMETRIC LAPLACIANS ON THE REAL LINE

M. KESSEBÖHMER, T. SAMUEL, AND H. WEYER

ABSTRACT. Motivated by the fundamental theorem of calculus, and based on the works of Feller as well as Kac and Kreĭn, given an atomless Borel probability measure η supported on a compact subset of \mathbb{R} , Freiberg and Zähle introduced a measure-geometric approach to define a first order differential operator ∇_η and a second order differential operator Δ_η , with respect to η . We generalise this approach to measures of the form $\eta := \nu + \delta$, where ν is continuous and δ is finitely supported. We determine analytic properties of ∇_η and Δ_η and show that Δ_η is a densely defined, unbounded, linear, self-adjoint operator with compact resolvent. Moreover, we give a systematic way to calculate the eigenvalues and eigenfunctions of Δ_η . For two leading examples, we determine the eigenvalues and the eigenfunctions, as well as the asymptotic growth rates of the eigenvalue counting function.

1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

Kac posed the following famous question in [16]. “Can one hear the shape of a drum?” Namely, can one reconstruct the geometry of a n -dimensional manifold from the eigenvalues of the associated Laplacian. In 1964 Milnor [26] showed the existence of a pair of 16-dimensional tori whose associated Laplacians have the same eigenvalues but which have different shapes. Subsequently, for a given $n \geq 4$, Urakawa [29] produced the first examples of domains in \mathbb{R}^n with this property. The problem in two dimensions remained open until 1992, when Gordon, Webb, and Wolpert [12] constructed a pair of regions in the plane that have different shapes but whose associated Laplacians have identical eigenvalues. Nevertheless, as observed by Weyl [30], Berry [3, 4], Lapidus *et al.* [21, 22, 23, 24, 25], Beals and Greiner [2] and many others, the spectrum of a Laplacian still tells us a lot about the shape of the underlying geometric structure.

As a special case of the generalised Kreĭn-Feller operator [6, 15], in [10] Freiberg and Zähle introduced a measure-geometric approach to define a first order differential operator ∇_η and a second order differential operator $\Delta_{\eta,\eta} := \nabla_\eta \circ \nabla_\eta$, with respect to an atomless Borel probability measure η supported on a compact subset of \mathbb{R} . In the case that η is the Lebesgue measure, it was shown that ∇_η coincides with the weak derivative. Moreover, a harmonic calculus for $\Delta_{\eta,\eta}$ was developed and, when η is a self-similar measure supported on a Cantor set, Freiberg and Zähle proved that the eigenvalue counting function of $\Delta_{\eta,\eta}$ is comparable to the square-root function. In [17] the exact eigenvalues and eigenfunctions of $\Delta_{\eta,\eta}$ were obtained and it was shown that the eigenvalues do not depend on the given measure. Arzt [1], Freiberg [7, 8, 9], Fujita [11], and Kotani and Watanabe [20] have also considered the Kreĭn-Feller operator $\Delta_{\eta,\Lambda} := \nabla_\eta \circ \nabla_\Lambda$, where η denotes a continuous Borel probability measure and Λ denotes the Lebesgue measure. In the case that η is a purely atomic measure, it has been shown in [2] that the eigenvalues of $\Delta_{\eta,\Lambda}$ highly depend on the position and weights of the atoms. Interestingly, the operators $\Delta_{\eta,\Lambda}$ and $\Delta_{\eta,\eta}$, in the case that η is continuous, also appear as the infinitesimal generator of Liouville Brownian motion [14] and Liouville quantum gravity [28].

In [18] it was shown that the framework of Freiberg and Zähle can be extended to include purely atomic measures η . Unlike in the case when one has a measure with a continuous distribution function, it was proven that the operators ∇_η and $\Delta_{\eta,\eta}$ are no longer symmetric. To circumvent this problem, the η -Laplacian was defined by $\Delta_\eta = -\nabla_\eta^* \circ \nabla_\eta$, where ∇_η^* denotes the adjoint of ∇_η . Further, a matrix representation for these operators was given and shown to coincide with the normalised graph Laplacian of a cycle graph [5]. Moreover, the eigenvalues of Δ_η depend only on the weights of the atoms and are independent of the positions of the atoms.

In this article, we continue to develop the program of Freiberg and Zähle for measures of the form $\eta = \nu + \delta$, where ν is continuous and δ is a finite sum of weighted Dirac point masses. Indeed, we show, for such an η , one can define a first and a second order measure geometric differential operator, ∇_η and $\Delta_\eta = -\nabla_\eta^* \circ \nabla_\eta$ respectively. Moreover, we determine properties of both operators; in particular, we show the following.

(M. Kesseböhmer and H. Weyer) FB 3 – MATHEMATIK, UNIVERSITÄT BREMEN, BIBLIOTHEKSTR. 1, 28359 BREMEN, GERMANY

(T. Samuel) SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, EDGBASTON, BIRMINGHAM, B15 2TT, UK

2010 *Mathematics Subject Classification.* 47G30; 42B35; 35P20.

Key words and phrases. measure-geometric Laplacians; spectral asymptotics; harmonic analysis.

Theorem 1.1. *The operator Δ_η is densely defined on the space L_η^2 of square- η -integrable functions. Further, it is linear, self-adjoint, non-positive and has compact resolvent.*

For ease of exposition in the following result we assume that ν is a probability measure on $(0, 1]$ with distribution function F_ν and let $\delta = \sum_{i=1}^N \alpha_i \delta_{z_i}$ with $0 < z_1 < \dots < z_N = 1$.

Theorem 1.2. *A square- η -integrable function f is an eigenfunction of Δ_η with corresponding eigenvalue λ if and only if it is of the form*

$$f(x) = \begin{cases} a_1 \sin(bF_\nu(x) + \gamma_1) & \text{if } x \in (0, z_1], \\ \vdots & \vdots \\ a_N \sin(bF_\nu(x) + \gamma_N) & \text{if } x \in (z_{N-1}, 1], \end{cases}$$

and $\lambda = -b^2$, where $b, a_1, \dots, a_N, \gamma_1, \dots, \gamma_N \in \mathbb{R}$ satisfy the following system of equations:

$$\alpha_j b a_{j+1} \cos(bF_\nu(z_j) + \gamma_{j+1}) = a_{j+1} \sin(bF_\nu(z_j) + \gamma_{j+1}) - a_j \sin(bF_\nu(z_j) + \gamma_j),$$

$$\alpha_j b^2 a_j \sin(bF_\nu(z_j) + \gamma_j) = a_j b \cos(bF_\nu(z_j) + \gamma_j) - a_{j+1} b \cos(bF_\nu(z_j) + \gamma_{j+1}),$$

for $j \in \{1, \dots, N-1\}$, and

$$\alpha_N b a_1 \cos(\gamma_1) = a_1 \sin(\gamma_1) - a_N \sin(b + \gamma_N),$$

$$\alpha_N b^2 a_N \sin(b + \gamma_N) = a_N b \cos(b + \gamma_N) - a_1 b \cos(\gamma_1).$$

Remark 1.3. To prove Theorem 1.1, it is sufficient to consider the case when the continuous part ν of η restricted to the interval between two consecutive atoms is either zero or Lebesgue, and to prove Theorem 1.2, it is sufficient to consider the case when $\nu = \Lambda$. Indeed, the general case follows by appropriately composing the operator with the distribution function F_ν as in [17].

Theorem 1.2 shows that the eigenvalues depend on the weights of the Dirac point masses and their positions relative to ν , but that they are independent of the distribution of ν ; this condition is different than that given for the Krein-Feller operator $\Delta_{\eta, \Lambda}$, where η is a purely atomic measure, compare with [2].

Further, we investigate two leading examples in detail and determine their eigenvalues and eigenfunctions explicitly. In contrast to the classical theory and to the case of atomless measures we see that all eigenspaces are one-dimensional; however, the asymptotic growth rate of the eigenvalue counting function N_η behaves as in the classical situation, namely

$$\lim_{x \rightarrow \infty} \frac{\pi N_\eta(x)}{\sqrt{x}} = 1.$$

This article is structured as follows. In Section 2 we define the operators ∇_η , ∇_η^* and Δ_η and prove that Δ_η is a densely defined self-adjoint operator on L_μ^2 . Further, we show that ∇_η and ∇_η^* are closed, give an explicit description of their domains and ranges, and prove that Δ_η has compact resolvent. From this and Remark 1.3 we conclude the proof of Theorem 1.1. In Section 3 we determine spectral properties of Δ_η . We divide this section into three parts. In the first part (Section 3.1), we give a system of equations which allows one to obtain the eigenvalues and find a general form of the eigenfunctions, hence proving Theorem 1.2. In the second part (Section 3.2), we solve the system of equations given in Theorem 1.2 for the case that $N = 1$ and illustrate the results in an example. The third part (Section 3.3) deals with the case when $N = 2$ and when the Dirac point masses are uniformly distributed and equally weighted. We end this final section with an example which illustrates our results in this latter setting.

2. THE OPERATORS ∇_η AND Δ_η

Let η denote a finite Borel measure on \mathbb{R} and let $a, b \in \mathbb{R}$ be such that the convex hull of $\text{supp}(\eta)$ is equal to $[a, b]$. Here $\text{supp}(\eta)$ denotes the support of η , that is, the smallest compact set with full measure. We assume that $\eta(\{a\}) = 0$ and set $M = (a, b]$. For $K \subseteq M$, we let $\eta|_K$ be the restriction of η to the set K , that is, $\eta|_K(A) := \eta(A \cap K)$ for all Borel sets $A \subseteq \mathbb{R}$; the same notation is used for functions. When it is clear from context, we write η for $\eta|_K$. We denote the set of real-valued square- η -integrable functions with domain equal to M by \mathcal{Q}_η^2 , define \mathcal{N}_η to be the set of \mathcal{Q}_η^2 -functions which are constant zero η -almost everywhere, and let $L_\eta^2 := \mathcal{Q}_\eta^2 / \mathcal{N}_\eta$. Following convention, when we write $f \in L_\eta^2$, we mean that there exists an equivalence class of L_η^2 to which f belongs. When it is clear

from context, we will use the same notation for a function $f \in \mathfrak{L}_\eta^2$ and for the equivalence class in L_η^2 to which it belongs. We equip L_η^2 with the inner product given by

$$\langle f, g \rangle_\eta := \int f g \, d\eta.$$

We denote by $\|\cdot\|_\eta$ the associate L_η^2 -norm. For $d \in \mathbb{R}$ we set $d\mathbb{Z} := \{dk : k \in \mathbb{Z}\}$ and let $\delta_{d\mathbb{Z}}$ denote the Dirac comb $\delta_{d\mathbb{Z}} := \sum_{k \in \mathbb{Z}} \delta_{dk}$. Here, for $z \in \mathbb{R}$, we write δ_z for the Dirac point mass at z . For a function $f : M \rightarrow \mathbb{R}$ we let $\mathbf{f} : \mathbb{R} \rightarrow \mathbb{R}$ be the periodic extension of f , that is $\mathbf{f}(x) = f(x)$ for all $x \in M$ and $\mathbf{f}(x) = \mathbf{f}(x + (b-a)k)$ for all $x \in \mathbb{R}$ and $k \in \mathbb{Z}$. We define the set \mathcal{D}_η^1 of η -differentiable functions by

$$\mathcal{D}_\eta^1 := \left\{ f \in \mathfrak{L}_\eta^2 : \text{there exists } f' \in L_\eta^2 \text{ with } \mathbf{f}(x) = \mathbf{f}(y) + \int_{[y,x]} \mathbf{f}' \, d\eta * \delta_{(b-a)\mathbb{Z}} \text{ for all } x, y \in \mathbb{R} \text{ with } y < x \right\}. \quad (1)$$

Here $\eta * \delta_{(b-a)\mathbb{Z}}$ denotes the convolution of the measure η and the Dirac comb $\delta_{(b-a)\mathbb{Z}}$; see, for instance, [13] for the definition of and results on convolutions of measures.

If $f \in \mathcal{D}_\eta^1$, then f is left-continuous with discontinuities occurring only in a subset of $\{z_1, \dots, z_N\}$. As the function f' defined in (1) is unique in L_η^2 , the operator $\nabla_\eta : \mathcal{D}_\eta^1 \rightarrow L_\eta^2$ given by $\nabla_\eta f := f'$ is well-defined and called the η -derivative. By the linearity of the integral equation in (1), it follows that ∇_η is linear. Additionally, if $f, g \in \mathcal{D}_\eta^1$ with $f \neq g$, then $\|f - g\|_\eta \neq 0$. Thus, we may view \mathcal{D}_η^1 as a collection of real-valued square- η -integrable functions, or as a collection of equivalence classes of L_η^2 and in the latter setting, we define ∇_η accordingly; namely, if $f \in \mathcal{D}_\eta^1$, then ∇_η maps the equivalence class of f to the equivalence class of $\nabla_\eta f$.

Remark 2.1. Let $C(M)$ denote the set of continuous functions $f : M \rightarrow \mathbb{R}$ and let $C^1(M)$ denote the set of $f \in C(M)$ such that f is differentiable on (a, b) and left-differentiable at b . In the case that η is a continuous Borel measure, the set \mathcal{D}_η^1 given in (1) is contained in $C(M)$ and equal to the set \mathcal{D}_η^1 given in [10], which, if $\eta = \Lambda$, is in turn equal to the Sobolev space W_2^1 . Moreover, an application of the fundamental theorem of calculus yields that $C^1(M)$ is contained in \mathcal{D}_Λ^1 . Thus, we have $C^1(M) \subseteq \mathcal{D}_\Lambda^1 \subseteq C(M) \subseteq L_\Lambda^2$, where each set is dense with respect to $\|\cdot\|_\Lambda$, in the succeeding one.

Remark 2.2. By Remark 1.3, and a rescaling and translation argument, it is sufficient to prove Theorem 1.1 under the following assumptions. There exists $N \in \mathbb{N}$ and $\underline{c} = (c_1, \dots, c_N) \in \{0, 1\}^N$ with $\underline{c} \neq \underline{0}$, such that

- (i) $a = 0$ and $b = 1$, in which case $M = \mathcal{I} := (0, 1]$, and
- (ii) $\eta = \Gamma + \sum_{i=1}^N \alpha_i \delta_{z_i}$, where $0 < z_1 < \dots < z_N = 1$ and $d\Gamma = (\sum_{i=1}^N c_i \mathbb{1}_{(z_{i-1}, z_i]}) d\Lambda$.

Thus, throughout this section we assume that η has this form.

For convenience, we set $z_{N+1} := 1 + z_1$ and $A_i := (z_{i-1}, z_i]$, for $i \in \{1, \dots, N\}$. Given a bounded interval J , following convention, we let J° denote the interior of J . Further, we let $\mathbb{1}_J$ denote the characteristic function on J , and in the case that $J = \mathcal{I}$, we write $\mathbb{1}$ for $\mathbb{1}_J$.

As with the classical weak Laplacian, $\nabla_\eta f$ reflects local properties of f . Indeed, we have, for $i \in \{1, \dots, N\}$,

$$\nabla_\eta f(z_i) = \lim_{\varepsilon \searrow 0} \frac{\mathbf{f}(z_i + \varepsilon) - \mathbf{f}(z_i)}{\alpha_i} \quad \text{and} \quad \nabla_\eta f(x) = f'_\Gamma(x), \quad (2)$$

where $x \in A_i^\circ$ and $f'_\Gamma \in L_\Gamma^2$ is such that

$$\mathbf{f}(x) = \lim_{\varepsilon \searrow 0} \mathbf{f}(z_i + \varepsilon) + \int_{[z_i, x]} \mathbf{f}'_\Gamma \, d\Gamma.$$

If $c_i = 1$, then $f'_\Gamma|_{A_i^\circ}$ coincides with the weak derivative, and if $c_i = 0$, then $f'_\Gamma|_{A_i^\circ}$ can be chosen arbitrarily.

Proposition 2.3. *The set \mathcal{D}_η^1 is dense in L_η^2 with respect to $\|\cdot\|_\eta$.*

Proof. For $i \in \{1, \dots, N\}$ we show, for given $f \in L_\Lambda^2$ and $\varepsilon > 0$, that there exists $g \in \mathcal{D}_\eta^1$ with $\|f - g\|_\eta < \varepsilon$. Following this, for $i \in \{1, \dots, N\}$, we construct a sequence $(h_n)_{n \in \mathbb{N}}$ in \mathcal{D}_η^1 with $\lim_{n \rightarrow \infty} \|h_n - \mathbb{1}_{[z_i]}\|_\eta = 0$.

Let us begin by showing the first statement for a fixed $i \in \{1, \dots, N-1\}$; the case $i = N$ follows by a similar argument and is left to the reader. For $0 < \varepsilon < (z_{i+1} - z_i)/2$, $n \in \mathbb{N}_0$ and $x \in (0, 1]$ set $c_{\varepsilon, n} := \pi n / (z_{i+1} - z_i - 2\varepsilon)$ and

$f_{\varepsilon,n}(x) := \int_0^x f'_{\varepsilon,n} d\Lambda$, where

$$f'_{\varepsilon,n}(x) := \begin{cases} -(32c_{\varepsilon,n}/\varepsilon)(x - (z_i + \varepsilon/4)) & \text{if } x \in (z_i + \varepsilon/4, z_i + 3\varepsilon/8], \\ (32c_{\varepsilon,n}/\varepsilon)(x - (z_i + \varepsilon/2)) & \text{if } x \in (z_i + 3\varepsilon/8, z_i + \varepsilon/2], \\ -c_{\varepsilon,n} \cos((2\pi/\varepsilon)(x - (z_i + \varepsilon/2))) + c_{\varepsilon,n} & \text{if } x \in (z_i + \varepsilon/2, z_i + \varepsilon], \\ 2c_{\varepsilon,n} \cos(2c_{\varepsilon,n}(x - (z_i + \varepsilon))) & \text{if } x \in (z_i + \varepsilon, z_{i+1} - \varepsilon], \\ c_{\varepsilon,n} \cos((2\pi/\varepsilon)(x - (z_{i+1} - \varepsilon))) + c_{\varepsilon,n} & \text{if } x \in (z_{i+1} - \varepsilon, z_{i+1} - \varepsilon/2], \\ -(32c_{\varepsilon,n}/\varepsilon)(x - (z_{i+1} - \varepsilon/2)) & \text{if } x \in (z_{i+1} - \varepsilon/2, z_{i+1} - 3\varepsilon/8], \\ (32c_{\varepsilon,n}/\varepsilon)(x - (z_{i+1} - \varepsilon/4)) & \text{if } x \in (z_{i+1} - 3\varepsilon/8, z_{i+1} - \varepsilon/4], \\ 0 & \text{otherwise.} \end{cases}$$

By definition, $f_{\varepsilon,n} \in \mathcal{D}_\eta^1$, $f_{\varepsilon,n}(x) = 0$, for $x \in I \setminus A_i^o$, and $f_{\varepsilon,n}(x) = \sin(2c_{\varepsilon,n}(x - (z_i + \varepsilon)))$, for $x \in (z_i + \varepsilon, z_{i+1} - \varepsilon)$. Further, set

$$g_{\varepsilon,n}(x) := \begin{cases} -(1/2) \cos((2\pi/\varepsilon)(x - (z_i + \varepsilon/2))) + 1/2 & \text{if } x \in (z_i + \varepsilon/2, z_i + \varepsilon], \\ \cos(2c_{\varepsilon,n}(x - (z_i + \varepsilon))) & \text{if } x \in (z_i + \varepsilon, z_{i+1} - \varepsilon], \\ (1/2) \cos((2\pi/\varepsilon)(x - (z_{i+1} - \varepsilon))) + 1/2 & \text{if } x \in (z_{i+1} - \varepsilon, z_{i+1} - \varepsilon/2], \\ 0 & \text{otherwise.} \end{cases}$$

Notice, $g_{\varepsilon,n} \in \mathcal{D}_\eta^1$, $g_{\varepsilon,n}(x) = 0$, for $x \in I \setminus A_i^o$, and $g_{\varepsilon,n}(x) = \cos(2c_{\varepsilon,n}(x - (z_i + \varepsilon)))$, for $x \in (z_i + \varepsilon, z_{i+1} - \varepsilon)$.

The functions $f_{\varepsilon,n}$ and $g_{\varepsilon,n}$ are differentiable (in the classical sense) and hence, by Remark 2.1, lie in $\mathcal{D}_{\eta|_{A_i}}^1$. For $i \in \{1, \dots, N\}$, define $\mathcal{F}_i := \{f_{\varepsilon,n}, g_{\varepsilon,n} : 0 < \varepsilon < (z_{i+1} - z_i)/2 \text{ and } n \in \mathbb{N}_0\}$. From the fact that the set $\{x \mapsto \sin(2\pi nx/\Lambda(A_i)) : n \in \mathbb{N}\} \cup \{x \mapsto \cos(2\pi nx/\Lambda(A_i)) : n \in \mathbb{N}_0\}$ forms a basis of $L_{\Lambda|_{A_i}}^2$, it follows that the span of $\bigcup_{i=1}^N \mathcal{F}_i \subseteq \mathcal{D}_\eta^1$ is a dense subset of L_Λ^2 with respect to $\|\cdot\|_\Lambda$.

For the second statement, again let $i \in \{1, \dots, N-1\}$ be fixed; the case $i = N$ follows by a similar argument and is left to the reader. Let K be the smallest natural number with $1/K < \min\{z_i - z_{i-1} : i \in \{1, \dots, N\}\}$. We divide the proof into four cases, namely when c_i and c_{i+1} are zero or one.

In the case that the atom is isolated, that is $c_i = c_{i+1} = 0$, the function $h := \mathbb{1}_{A_i}$ lies in \mathcal{D}_η^1 and $\|h - \mathbb{1}_{\{z_i\}}\|_\eta = 0$. For the cases when $c_{i+1} = 1$, we set, for $n \in \mathbb{N}$ with $n \geq K$,

$$h_n(x) := \begin{cases} (1 - \cos(n\pi(x - (z_i - 1/n))))/2 & \text{if } c_i = 1 \text{ and } x \in (z_i - 1/n, z_i + 1/n), \text{ or} \\ & \text{if } c_i = 0 \text{ and } x \in (z_i, z_i + 1/n), \\ 1 & \text{if } c_i = 0 \text{ and } x \in (z_{i-1}, z_i], \\ 0 & \text{otherwise.} \end{cases}$$

Observe that $\lim_{n \rightarrow \infty} \|h_n - \mathbb{1}_{\{z_i\}}\|_\eta = 0$ and moreover, that $h_n \in \mathcal{D}_\eta^1$.

Finally, we consider the case $c_i = 1$ and $c_{i+1} = 0$. For this let $j \in \{i+2, \dots, N\}$ be the smallest such integer with $c_j = 1$. If no such j exists then let $j \in \{1, \dots, i\}$ be the smallest such integer with $c_j = 1$. Observe that in the first case it is sufficient to approximate $\mathbb{1}_{\{z_i, \dots, z_j\}}$ and, in the second case, $\mathbb{1}_{\{z_1, \dots, z_j, z_i, \dots, z_N\}}$. Here we prove the former of these two cases as the latter follows analogously. For $n \in \mathbb{N}$ with $n \geq K$, set

$$h_n(x) := \begin{cases} (1 - \cos(n\pi(x - (z_i - 1/n))))/2 & \text{if } x \in (z_i - 1/n, z_i], \\ (1 - \cos(n\pi(x - (z_j + 1/n))))/2 & \text{if } x \in (z_j, z_j + 1/n], \\ 1 & \text{if } x \in (z_i, z_j], \\ 0 & \text{otherwise.} \end{cases}$$

By definition, we have $\lim_{n \rightarrow \infty} \|h_n - \mathbb{1}_{\{z_i, \dots, z_j\}}\|_\eta = 0$ and $h_n \in \mathcal{D}_\eta^1$. □

From (1), if $f \in \mathcal{D}_\eta^1$, then

$$\langle \nabla_\eta f, \mathbb{1} \rangle_\eta = \int \nabla_\eta f d\eta = 0, \tag{3}$$

which implies that the zero function is the only constant function which can occur as an η -derivative.

Letting ϱ denote the (natural) quotient map from L_η^2 to $L_\eta^2/\{c\mathbb{1} : c \in \mathbb{R}\}$, in the following proposition we show that the image of the range of ∇_η under ϱ is dense in $L_\eta^2/\{c\mathbb{1} : c \in \mathbb{R}\}$. By the continuity of the inner product $\langle \cdot, \cdot \rangle_\eta$, this implies that the orthogonal complement of the range of ∇_η is equal to the set of constant functions on I .

Proposition 2.4. *The image under ϱ of the range of ∇_η is dense in the quotient space $L_\eta^2/\{c\mathbb{1} : c \in \mathbb{R}\}$.*

Proof. The set of bounded continuous functions $C_B(I)$ with domain equal to I is dense in L_η^2 , and so the image of $E := \{g \in C_B(I) : \langle g, \mathbb{1} \rangle_\eta = 0\}$ under ϱ is dense in $L_\eta^2/\{c\mathbb{1} : c \in \mathbb{R}\}$. For $g \in E$, setting $f(x) = \langle g, \mathbb{1}_{[0,x)} \rangle_\eta$ for $x \in I$, observe that f is left-continuous and bounded, and so $f \in L_\eta^2$. Since $g \in E$, by (3), we have $f \in \mathcal{D}_\eta^1$, where $\nabla_\eta f = g$. In other words, E is contained in the range of ∇_η and hence the image of the range of ∇_η under ϱ is dense in $L_\eta^2/\{c\mathbb{1} : c \in \mathbb{R}\}$. \square

As in [18, 19], we use a Dirichlet form \mathcal{E}_η to define the measure geometric Laplacian Δ_η . For this we use the following properties of unbounded operators; see for instance [27] for further details. The *graph of a densely defined linear operator* T on L_η^2 is $\Gamma(T) := \{(f, T(f)) \in L_\eta^2 \times L_\eta^2 : f \in \text{Dom}(T)\}$, where $\text{Dom}(T)$ denotes the domain of T . In the case that $\Gamma(T)$ is closed in $L_\eta^2 \times L_\eta^2$, we say that T is *closed*. If T_1 is a densely defined operator on L_η^2 and if $\Gamma(T_1) \supseteq \Gamma(T)$, then T_1 is called an *extension* of T . When T has a closed extension, T is said to be *closable*. The smallest closed extension of T , denoted by \overline{T} , is the *closure* of T .

For a densely defined operator T on L_η^2 we let $\text{Dom}(T^*)$ be the set of $f \in L_\eta^2$ for which there exists $h \in L_\eta^2$ with $\langle T(g), f \rangle_\eta = \langle g, h \rangle_\eta$ for all $g \in \text{Dom}(T)$. For each such $f \in \text{Dom}(T^*)$, we define $T^*(f) := h$. We refer to T^* as the *adjoint* of T . We call T *symmetric* if $\text{Dom}(T) \subseteq \text{Dom}(T^*)$ and $T(f) = T^*(f)$, for all $f \in \text{Dom}(T)$. Equivalently, T is symmetric if and only if $\langle T(f), g \rangle_\eta = \langle f, T(g) \rangle_\eta$ for all $f, g \in \text{Dom}(T)$. If in addition to T being symmetric, we have that $\text{Dom}(T) = \text{Dom}(T^*)$, then we say that T is *self-adjoint*.

Theorem 2.5 ([27]). *If T is an unbounded, densely defined operator on L_η^2 , then the following holds.*

- (i) *The operator T^* is closed.*
- (ii) *The operator T is closable if and only if $\text{Dom}(T^*)$ is dense in L_η^2 in which case $\overline{T} = T^{**}$.*
- (iii) *If T is closable, then $(\overline{T})^* = T^*$.*

In the following proposition we show that the domain of ∇_η^* is equal to

$$\mathcal{D}_\eta^{1*} := \left\{ f \in \mathcal{Q}_\eta^2 : \text{there exists } f^* \in L_\eta^2 \text{ with } \mathbf{f}(x) = \mathbf{f}(y) + \int \mathbb{1}_{(x,y]} \mathbf{f}^* d\eta * \delta_{\mathbb{Z}} \text{ for all } x, y \in \mathbb{R} \text{ with } x < y \right\}. \quad (4)$$

Notice, if $f \in \mathcal{D}_\eta^{1*}$, then f is right-continuous with discontinuities occurring only at points in a subset of $\{z_1, \dots, z_N\}$. Moreover, if $f, g \in \mathcal{D}_\eta^{1*}$ with $f \neq g$, then $\|f - g\|_\eta \neq 0$. Thus, as with \mathcal{D}_η^1 , we may view \mathcal{D}_η^{1*} as a collection of real-valued square- η -integrable functions, or as a collection of equivalence classes of L_η^2 .

Proposition 2.6. *The domain of ∇_η^* is equal to \mathcal{D}_η^{1*} .*

Proof. Let $f \in \text{Dom}(\nabla_\eta^*)$. Using the fact that $\langle \nabla_\eta^* f, \mathbb{1} \rangle_\eta = \langle f, \nabla_\eta \mathbb{1} \rangle_\eta = \langle f, 0 \rangle_\eta = 0$ and Fubini's Theorem, we obtain, for $g \in \mathcal{D}_\eta^1$,

$$\begin{aligned} \int \mathbb{1}_{(0,1]} f(\nabla_\eta g) d\eta &= \int \mathbb{1}_{(0,1]} (\nabla_\eta^* f) g d\eta \\ &= \int \mathbb{1}_{(0,1]}(x) (\nabla_\eta^* f)(x) \left(\mathbf{g}(0) + \int \mathbb{1}_{[0,x)}(y) (\nabla_\eta g)(y) d\eta(y) \right) d\eta(x) \\ &= \int \mathbb{1}_{[0,1)}(y) \nabla_\eta g(y) \int \mathbb{1}_{(y,1]}(x) (\nabla_\eta^* f)(x) d\eta(x) d\eta(y). \end{aligned}$$

This, together with Proposition 2.4 and the fact that $\langle \mathbb{1}, \nabla_\eta g \rangle_\eta = 0$, implies $\varrho(f - \int \mathbb{1}_{(\cdot,1]} \nabla_\eta^* f d\eta) = 0$. Hence, there exists $c \in \mathbb{R}$ so that, for η -almost all $y \in (0, 1]$,

$$f(y) - \int \mathbb{1}_{(y,1]} \nabla_\eta^* f d\eta = c.$$

This yields that f is a right-continuous function with $c = \mathbf{f}(0)$. When setting $f^* := \nabla_\eta^* f$, since $\mathbf{f}(0) = f(1)$ and $\langle \nabla_\eta^* f, \mathbb{1} \rangle_\eta = 0$, we have that \mathbf{f}^* fulfils the integral equation in (4) and therefore, $\text{Dom}(\nabla_\eta^*) \subseteq \mathcal{D}_\eta^{1*}$.

For the reverse inclusion, let $g \in \text{Dom}(\nabla_\eta)$, $f \in \mathcal{D}_\eta^{1*}$ and f^* be as in (4). By Fubini's Theorem and the fact that $\langle f^*, \mathbb{1} \rangle_\eta = \langle \nabla_\eta g, \mathbb{1} \rangle_\eta = 0$, we have the following chain of equalities, which yields the result.

$$\begin{aligned} \int (\nabla_\eta g) f d\eta &= \int \nabla_\eta g(y) \left(f(1) + \int \mathbb{1}_{(y,1]}(x) f^*(x) d\eta(x) \right) d\eta(y) \\ &= \int f^*(x) \int \nabla_\eta g(y) \mathbb{1}_{[0,x)}(y) d\eta(y) d\eta(x) = \int f^*(x) g(x) d\eta(x) \end{aligned} \quad \square$$

Corollary 2.7. For $f \in \text{Dom}(\nabla_\eta^*)$, we have that $\nabla_\eta^* f = f^*$, where f^* is defined as in (4).

The function $\nabla_\eta^* f$, when it exists, reflects local properties of f . Indeed, we have

$$\nabla_\eta^* f(z_i) = \lim_{\varepsilon \searrow 0} \frac{\mathbf{f}(z_i) - \mathbf{f}(z_i - \varepsilon)}{\alpha_i} \quad \text{and} \quad \nabla_\eta^* f(x) = f_\Lambda^*(x), \quad (5)$$

for $i \in \{1, \dots, N\}$ and $x \in A_i^\circ \cap \mathcal{I}$, where $f_\Lambda^* \in L_\Lambda^2$ is equal to the negative of the weak derivative of f on A_i° , if $c_i = 1$ and otherwise can be chosen arbitrarily on A_i° .

Proposition 2.8. The operator ∇_η is densely defined, unbounded and closed.

Proof. By Proposition 2.3, the operator ∇_η is densely defined. Set $K := \Gamma(\mathcal{I})$ and define for $m \in \mathbb{N}$ and $x \in \mathcal{I}$ $g_m(x) = \sin(2\pi m F_\Gamma(x)/K) \in \mathcal{D}_\eta^1$, in which case

$$\nabla_\eta g_m(x) = \begin{cases} (2\pi m/K) \cos(2\pi m F_\Gamma(x)/K) & \text{if } x \in \mathcal{I} \setminus \{z_1, \dots, z_N\}, \\ 0 & \text{otherwise.} \end{cases}$$

This implies that ∇_η is an unbounded operator as $\|g_m\|_\eta \leq 1 + \sum_{i=1}^N \alpha_i$ and $\|\nabla_\eta g_m\|_\eta = \sqrt{2}\pi m$. Theorem 2.5 gives that ∇_η^{**} is closed. To complete the proof, it suffices to show that $\text{Dom}(\nabla_\eta) = \text{Dom}(\nabla_\eta^{**})$. However, this follows by an analogously argument to that given in the proof of Proposition 2.6; noting, by a similar proof to that of Proposition 2.3, we have \mathcal{D}_η^{1*} is dense in L_η^2 with respect to $\|\cdot\|_\eta$. \square

The non-negative symmetric bilinear form $\mathcal{E}: \mathcal{D}_\eta^1 \times \mathcal{D}_\eta^1 \rightarrow \mathbb{R}$ defined by $\mathcal{E}(f, g) = \mathcal{E}_\eta(f, g) := \langle \nabla_\eta f, \nabla_\eta g \rangle_\eta$ is called the η -energy form. In our next result, we show that \mathcal{E} is a Dirichlet form. With this at hand, we may then define the η -Laplacian Δ_η .

Proposition 2.9. The η -energy form \mathcal{E} is a Dirichlet form with domain \mathcal{D}_η^1 .

Proof. Using the properties of the inner product $\langle \cdot, \cdot \rangle_\eta$ and the operator ∇_η it follows that \mathcal{E} is bilinear, symmetric and that $\mathcal{E}(u, u) \geq 0$, for all $u \in \mathcal{D}_\eta^1$. Moreover, this yields that \mathcal{D}_η^1 equipped with $\langle \cdot, \cdot \rangle_\mathcal{E} := \langle \cdot, \cdot \rangle_\eta + \mathcal{E}(\cdot, \cdot)$ is an inner product space. All that remains is to show that \mathcal{D}_η^1 is complete with respect to the norm induced by $\langle \cdot, \cdot \rangle_\mathcal{E}$ and that the Markov property holds.

If $(f_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in $(\mathcal{D}_\eta^1, \langle \cdot, \cdot \rangle_\mathcal{E})$, then both $(f_n)_{n \in \mathbb{N}}$ and $(\nabla_\eta f_n)_{n \in \mathbb{N}}$ are Cauchy-sequences in L_η^2 . Hence, there exist $\tilde{f}_0, \tilde{f}_1 \in L_\eta^2$ with $\lim_{n \rightarrow \infty} \|f_n - \tilde{f}_0\|_\eta = 0$ and $\lim_{n \rightarrow \infty} \|\nabla_\eta f_n - \tilde{f}_1\|_\eta = 0$. Proposition 2.8 implies that $\nabla_\eta \tilde{f}_0 = \tilde{f}_1$. Thus, (f_n) converges to $\tilde{f}_0 \in \mathcal{D}_\eta^1$ with respect to the norm induced by $\langle \cdot, \cdot \rangle_\mathcal{E}$.

For the Markov property it is sufficient to show, for $u \in \mathcal{D}_\eta^1$, that $u_+ \in \mathcal{D}_\eta^1$ and $|\nabla_\eta u_+(x)| \leq |\nabla_\eta u(x)|$ for all $x \in \mathcal{I}$. Here $u_+ := \min(\max(u, 0), 1)$. Define, for $x \in \mathcal{I} \setminus \{z_1, \dots, z_N\}$,

$$u'_+(x) := \begin{cases} \nabla_\eta u(x) & \text{if } u(x) \in [0, 1], \\ 0 & \text{otherwise.} \end{cases}$$

and for $i \in \{1, \dots, N\}$ set $u'_+(z_i) := \lim_{\varepsilon \searrow 0} (u_+(x + \varepsilon) - u_+(x))/\alpha_i$. A direct calculation shows, for $x, y \in \mathbb{R}$ with $x < y$, that this function fulfils $u_+(x) = u_+(y) + \int \mathbf{1}_{[y, x]} u'_+ d\eta * \delta_\mathbb{Z}$. Hence, $u_+ \in \mathcal{D}_\eta^1$ with $\nabla_\eta u_+ = u'_+$. By definition, $|\nabla_\eta u_+(x)| \leq |\nabla_\eta u(x)|$ for all $x \in \mathcal{I} \setminus \{z_1, \dots, z_N\}$. In the case that $x \in \{z_1, \dots, z_N\}$, we have

$$\lim_{\varepsilon \searrow 0} |u_+(x + \varepsilon) - u_+(x)| \leq \lim_{\varepsilon \searrow 0} |u(x + \varepsilon) - u(x)|,$$

and so, by (2), it follows that $|\nabla_\eta u_+(z_i)| \leq |\nabla_\eta u(z_i)|$ for $i \in \{1, \dots, N\}$. \square

We write $f \in \mathcal{D}_\eta^2$ if $f \in \mathcal{D}_\eta^1$ and if there exists $h \in L_\eta^2$ such that $\mathcal{E}(f, g) = -\langle h, g \rangle_\eta$, for all $g \in \mathcal{D}_\eta^1$. We call the operator $\Delta_\eta: \mathcal{D}_\eta^2 \rightarrow L_\eta^2$ defined by $\Delta_\eta f := h$ the η -Laplacian. Notice, for an arbitrary $g \in \mathcal{D}_\eta^1$, that $\langle \nabla_\eta f, \nabla_\eta g \rangle_\eta = -\langle \Delta_\eta f, g \rangle_\eta$, and thus $\Delta_\eta = -\nabla_\eta^* \circ \nabla_\eta$.

Theorem 2.10. The operator Δ_η is densely defined on L_η^2 , linear, self-adjoint and non-positive.

Proof. That the operator is densely defined follows from the observation that the functions $f_{\varepsilon, n}, g_{\varepsilon, n}, h$ and h_n , as defined in Proposition 2.3, lie in \mathcal{D}_η^2 . Linearity follows from the linearity of ∇_η and the bilinearity of the inner product. The fact that Δ_η is self-adjoint is a consequence of Theorem 2.5 and Proposition 2.8. Further, since $\langle \Delta_\eta f, f \rangle_\eta = -\langle \nabla_\eta f, \nabla_\eta f \rangle_\eta \leq 0$ we have the operator Δ_η is non-positive. \square

We conclude this section with the following theorem, in which we show that Δ_η has compact resolvent. For this we use the following notation. We denote the closed unit ball in a normed space $(X, \|\cdot\|)$ by $B(X, \|\cdot\|)$ and for $i \in \{1, \dots, N\}$, let $(W_i^{1,2}, \|\cdot\|_i)$ denote the Sobolev space $(W^{1,2}(A_i^\circ), \|\cdot\|_{1,2})$.

Theorem 2.11. *The operator Δ_η has compact resolvent.*

Proof. Let λ denote a fixed element of the resolvent set. We show the embedding $\pi: \mathcal{D}_\eta^1 \rightarrow L_\eta^2$ is compact, and that $(\lambda \text{Id} - \Delta_\eta)^{-1}: L_\eta^2 \rightarrow \mathcal{D}_\eta^1$ is continuous. This is sufficient to prove the result since the composition of a compact operator and a continuous operator is compact.

Let $(f_n)_{n \in \mathbb{N}}$ be a sequence in $B(\mathcal{D}_\eta^1, \|\cdot\|_\mathcal{E})$. To show π is compact, it is sufficient to show that $(f_n)_{n \in \mathbb{N}}$ has a convergent subsequence with respect to $\|\cdot\|_\eta$. It is known, for $i \in \{1, \dots, N\}$, that the unit ball $B(W_i^{1,2}, \|\cdot\|_i)$ is compact in L_Λ^2 . By (2), if $f \in B(\mathcal{D}_\eta^1, \|\cdot\|_\mathcal{E})$, then $f|_{A_i^\circ} \in B(W_i^{1,2}, \|\cdot\|_i)$. This yields the existence of a subsequence $(n_k)_{k \in \mathbb{N}}$ such that for $i \in \{1, \dots, N\}$ with $c_i = 1$, the sequence $(f_{n_k}|_{A_i^\circ})_{k \in \mathbb{N}}$ converges in $W_i^{1,2}$. Combined with the Bolzano-Weierstrass Theorem and the left continuity of elements in \mathcal{D}_η^1 , this yields the existence of a subsequence $(n_l)_{l \in \mathbb{N}}$, such that $(f_{n_l})_{l \in \mathbb{N}}$ converges with respect to $\|\cdot\|_{\eta|_{A_i}}$ for all $i \in \{1, \dots, N\}$, and hence, with respect to $\|\cdot\|_\eta$.

To conclude the proof it is sufficient to find, for each λ belonging to the resolvent set, a constant $C \in \mathbb{R}$ such that $\|(\Delta_\eta - \lambda \text{Id})^{-1}f\|_\mathcal{E} \leq C\|f\|_\eta$ for all $f \in L_\eta^2$. This is done in the following sequence of inequalities, in which we use that, since $(\Delta_\eta - \lambda \text{Id})^{-1}$ is a bounded linear operator on L_η^2 , there exists a $K \in \mathbb{R}$ with $\|(\Delta_\eta - \lambda \text{Id})^{-1}f\|_\eta \leq K\|f\|_\eta$.

$$\begin{aligned} \|(\Delta_\eta - \lambda \text{Id})^{-1}f\|_\mathcal{E}^2 &= \langle (\Delta_\eta - \lambda \text{Id})^{-1}f, (\Delta_\eta - \lambda \text{Id})^{-1}f \rangle_\eta + \langle \nabla_\eta (\Delta_\eta - \lambda \text{Id})^{-1}f, \nabla_\eta (\Delta_\eta - \lambda \text{Id})^{-1}f \rangle_\eta \\ &\leq (1 + |\lambda|) \|(\Delta_\eta - \lambda \text{Id})^{-1}f\|_\eta^2 + \langle f, (\Delta_\eta - \lambda \text{Id})^{-1}f \rangle_\eta \\ &\leq ((1 + |\lambda|)K^2 + K) \|f\|_\eta^2 \end{aligned} \quad \square$$

Since every eigenfunction of the resolvent operator is also an eigenfunction of Δ_η , the spectral theorem for compact operators together with the fact that Δ_η is non-positive imply that the eigenfunctions of Δ_η form a basis of L_η^2 and that the eigenvalues of Δ_η are non-positive and form a countable unbounded monotonic sequence. Moreover, all eigenvalues of Δ_η have finite multiplicity.

Proof of Theorem 1.1. This is a direct consequence of Remark 2.2 and Theorems 2.10 and 2.11. \square

3. EIGENVALUES AND EIGENFUNCTIONS OF Δ_η

3.1. Proof of Theorem 1.2 and general observations. Let $N \in \mathbb{N}$ denote a positive integer and for $i \in \{1, \dots, N\}$ let $\alpha_i > 0$ and $z_i \in \mathcal{I}$ with $0 < z_1 < \dots < z_N \leq 1$. In this section we determine a systematic way to compute the eigenvalues and eigenfunctions of the η -Laplacian, where $\eta = \Lambda + \sum_{i=1}^N \alpha_i \delta_{z_i}$. Without loss of generality, the assumption $z_N = 1$ can be made, since we can obtain the eigenfunctions of measures not having this property by applying an appropriate translation argument. To be precise, set $\hat{z}_i := z_i + 1 - z_N$ for $i \in \{1, \dots, N\}$ and define the measure $\hat{\eta} = \Lambda + \sum_{i=1}^N \alpha_i \delta_{\hat{z}_i}$. By construction, we have that f is an eigenfunction of $\Delta_{\hat{\eta}}$ if and only if

$$x \mapsto \begin{cases} f(x - z_N + 1) & x \in (0, z_N], \\ f(x - z_N) & x \in (z_N, 1], \end{cases}$$

is an eigenfunction of Δ_η .

Proof of Theorem 1.2. Combining (2), (5) and the fact that elements of $\text{Dom}(\nabla_\eta)$ are left-continuous, together with Picard-Lindelöf's Theorem, if $\Delta_\eta f = \lambda f$, for some $\lambda \in \mathbb{R}$ and $f \in \mathcal{D}_\eta^2$, then there exist $b, a_1, \dots, a_N, \gamma_1, \dots, \gamma_N \in \mathbb{R}$ with $\gamma_1 \in (-\pi/2, \pi/2]$ and $\lambda = -b^2$, such that

$$f(x) = \begin{cases} a_1 \sin(bx + \gamma_1) & \text{if } x \in (0, z_1], \\ \vdots & \vdots \\ a_N \sin(bx + \gamma_N) & \text{if } x \in (z_{N-1}, 1]. \end{cases} \quad (6)$$

Since $\nabla_\eta f$ is right-continuous and $\Delta_\eta f$ is left-continuous, by (2) and (5), we have f is an eigenfunction of Δ_η if and only if $b, a_1, \dots, a_N, \gamma_1, \dots, \gamma_N$ satisfy the following system of equations:

$$\begin{aligned} \alpha_j b a_{j+1} \cos(bz_j + \gamma_{j+1}) &= a_{j+1} \sin(bz_j + \gamma_{j+1}) - a_j \sin(bz_j + \gamma_j), \\ \alpha_j b^2 a_j \sin(bz_j + \gamma_j) &= a_j b \cos(bz_j + \gamma_j) - a_{j+1} b \cos(bz_j + \gamma_{j+1}), \end{aligned} \quad (7)$$

for $j \in \{1, \dots, N-1\}$, and

$$\begin{aligned} a_N b a_1 \cos(\gamma_1) &= a_1 \sin(\gamma_1) - a_N \sin(b + \gamma_N), \\ a_N b^2 a_N \sin(b + \gamma_N) &= a_N b \cos(b + \gamma_N) - a_1 b \cos(\gamma_1). \end{aligned} \quad (8)$$

This concludes the proof. \square

Let f be an eigenfunction of Δ_η with eigenvalue λ , and let b, a_1, \dots, a_N be as in (6). By the fact that $\lambda = -b^2$, if $b = 0$, then f is a step function, and so, (7) and (8) imply that f is a constant function. If $a_i = 0$ for some $i \in \{1, \dots, N-1\}$ and if $b \neq 0$, then (7) yields that $a_{j+1} \cos(bz_j + \gamma_{j+1}) = a_{j+1} \sin(bz_j + \gamma_{j+1}) = 0$, and hence, $a_{j+1} = 0$. An analogue results holds when one assumes that $a_N = 0$. This implies, if $a_i = 0$ for some $i \in \{1, \dots, N\}$, then $f \equiv 0$.

Corollary 3.1. *Every constant function is an eigenfunction with corresponding eigenvalue equal to zero. Moreover, the eigenspace $E_0 := \{f \in \mathcal{D}_\eta^2 : \Delta_\eta f \equiv 0\}$ is one-dimensional. Further, if $f \notin E_0$ is an eigenfunction of Δ_η , then $b \neq 0$ and $a_i \neq 0$ for all $i \in \{1, \dots, N\}$, where b, a_1, \dots, a_N are as in (6).*

If the atoms are equally distributed, namely $z_i - z_{i-1} = 1/N$ for $i \in \{1, \dots, N\}$, two properties which follow directly from (7) and (8) are the following. If f is an eigenfunction of Δ_η , then, for $r \in \{2, \dots, N\}$,

$$f_r(x) = \begin{cases} a_{N-r+2} \sin(b(x + (N-r+1)/N) + \gamma_{N-r+2}) & \text{if } x \in (0, 1/N], \\ \vdots & \vdots \\ a_N \sin(b(x + (N-r+1)/N) + \gamma_N) & \text{if } x \in ((r-2)/N, (r-1)/N], \\ a_1 \sin(b(x - (r-1)/N) + \gamma_1) & \text{if } x \in ((r-1)/N, r/N], \\ \vdots & \vdots \\ a_{N-r+1} \sin(b(x - (r-1)/N) + \gamma_{N-r+1}) & \text{if } x \in ((N-1)/N, 1], \end{cases} \quad (9)$$

is an eigenfunction of Δ_{η_r} , where $\eta_r := \Lambda + \sum_{i=1}^{r-1} \alpha_{i+N-r+1} \delta_{z_i} + \sum_{i=r}^N \alpha_{i-r+1} \delta_{z_i}$. Note, if $\alpha_1 = \dots = \alpha_N$, then $\eta_r = \eta$.

Corollary 3.2. *Let $N = pk$ with $p, k \in \mathbb{N}$ and suppose that $\alpha_{i+p} = \alpha_i$ for all $i \in \{1, \dots, N-p\}$. Set*

$$\eta^{(p)} := \Lambda + \sum_{i=1}^p \alpha_i \delta_{i/p} \quad \text{and} \quad \eta^{(N)} := \Lambda + \sum_{i=1}^N \alpha_i / k \delta_{i/N}.$$

If $f^{(p)}$ is an eigenfunction of $\Delta_{\eta^{(p)}}$ with eigenvalue $\lambda^{(p)}$, then $f^{(N)}$ is an eigenfunction of $\Delta_{\eta^{(N)}}$ with eigenvalue $\lambda^{(n)} := k^2 \lambda^{(p)}$, where $f^{(N)}(x) := \mathbf{f}^{(p)}(kx)$ for $x \in \mathcal{I}$.

3.2. $N = 1$. Here we consider the case $N = 1$, namely when $\eta = \Lambda + \alpha \delta_z$, for some $\alpha > 0$ and $z \in \mathcal{I}$. As in Section 3.1, without loss of generality we may assume that $z = 1$. The main results of this section are Theorem 3.4 and Corollary 3.5, in which we explicitly compute the eigenvalues and eigenfunctions of Δ_η . For the proofs of these results we will require the following preliminaries. For $\beta > 0$ and $k \in \mathbb{Z}$, let $c_k^\pm = c_k^\pm(\beta)$ denote the unique solution in the interval $(-\pi/2, \pi/2)$ to the equation

$$\tan(c_k^\pm) = -2c_k^\pm \beta \pm \beta \pi/2 + 2\pi \beta k \pm 1 \quad (10)$$

and set

$$\xi_k^\pm = \xi_k^\pm(\beta) := (\tan(c_k^\pm) \mp 1)/\beta. \quad (11)$$

Analogously, for $\beta < 0$ we denote by $C_k^\pm = C_k^\pm(\beta)$ the set of solutions to (10). Note, the cardinality $|C_k^\pm|$ of C_k^\pm is less than or equal to three. We denote the elements of C_k^\pm by $c_{k,i}^\pm = c_{k,i}^\pm(\beta)$, where $i \in \{1, \dots, |C_k^\pm|\}$. For every $c_{k,i}^\pm$ define the values $\xi_{k,i}^\pm = \xi_{k,i}^\pm(\beta)$ similar to (11) and denote by $\Xi_k^\pm = \Xi_k^\pm(\beta)$ the set $\{\xi_{k,i}^\pm : i \in \{1, \dots, |C_k^\pm|\}\}$.

Lemma 3.3. *Let $\beta \in \mathbb{R} \setminus \{0\}$. The pair $(\xi, c) \in \mathbb{R} \setminus \{0\} \times \mathbb{R}$ is a solution to the system of equations*

$$\begin{aligned} \beta \xi \cos(c) &= \sin(c) - \sin(\xi + c) \\ \beta \xi^2 \sin(\xi + c) &= \xi \cos(\xi + c) - \xi \cos(c) \end{aligned} \quad (12)$$

if and only if

$$(\xi, c) \in \begin{cases} \{(\xi_k^\pm, c_k^\pm) : k \in \mathbb{Z}\} & \text{if } \beta > 0, \\ \{(\xi_{k,i}^\pm, c_{k,i}^\pm) : k \in \mathbb{Z} \text{ and } i \in \{1, \dots, |C_k^\pm|\}\} & \text{if } \beta < 0. \end{cases}$$

The system of equations given in (12) is also solved by $(0, c)$ for all $c \in \mathbb{R}$. Further, if $c = \pi/2 + \pi k$, for some $k \in \mathbb{Z}$, then the only solution to (12) is given when $\xi = 0$.

Proof. The backwards implication follows by substituting the given values for ξ and c directly into (12). We now show the forwards implication. Substituting the first equation of (12) into the second equation of (12), and using the identity $\cos^2(\arcsin(x)) = 1 - x^2$, for $x \in [-1, 1]$, we obtain

$$(\beta\xi \sin(c) - \beta^2\xi^2 \cos(c) + \cos(c))^2 = 1 - (\sin(c) - \beta\xi \cos(c))^2.$$

From this it follows that $\cos(c) \neq 0$ and hence,

$$\beta^2\xi^2 - 2 \tan(c)\beta\xi + (\tan^2(c) - 1) = 0.$$

Thus, we have that either $\xi = 0$ and $c = \pm\pi/4 + \pi k$, for some $k \in \mathbb{Z}$, or that $\beta\xi = \tan(c) \pm 1$. In the case $\xi \neq 0$, substituting this value into the first equation of (12) yields $(\tan(c) \pm 1) \cos(c) = \sin(c) - \sin((\tan(c) \pm 1)/\beta + c)$, or equivalently,

$$\cos(c) = \sin\left(\frac{\mp \tan(c) - 1}{\beta} \mp c\right).$$

This leads to the following four cases:

- (i) $\beta\xi = \tan(c) + 1$ and $\tan(c) = -2c\beta - \beta\pi/2 + 2\pi\beta k - 1$, for $k \in \mathbb{Z}$,
- (ii) $\beta\xi = \tan(c) + 1$ and $\tan(c) = -\beta\pi/2 + 2\pi\beta k - 1$, for $k \in \mathbb{Z}$,
- (iii) $\beta\xi = \tan(c) - 1$ and $\tan(c) = -2c\beta + \beta\pi/2 + 2\pi\beta k + 1$, for $k \in \mathbb{Z}$, or
- (iv) $\beta\xi = \tan(c) - 1$ and $\tan(c) = \beta\pi/2 + 2\pi\beta k + 1$, for $k \in \mathbb{Z}$.

By substituting these values into (12), one sees that Cases (i) and (iii) yield solutions to (12). Cases (ii) and (iv) do not yield solutions, except when $c = 0$, but this is the same solution given by Cases (i) and (iii) when $c = 0$.

The last statement follows by substituting the given values for ξ and c directly into (12). \square

By (6), if $\Delta_\eta f = \lambda f$, for some $\lambda \in \mathbb{R}$, then there exist $a, b \in \mathbb{R}$ and $\gamma \in (-\pi/2, \pi/2]$ such that $\lambda = -b^2$ and $f(x) = a \sin(bx + \gamma)$ for $x \in \mathcal{I}$. By linearity, without loss of generality we may assume $a = 1$. Further, (7) and (8) imply that f is an eigenfunction of Δ_η if and only if b and γ satisfy the following system of equations.

$$\begin{aligned} \alpha b \cos(\gamma) &= \sin(\gamma) - \sin(b + \gamma) \\ \alpha b^2 \sin(b + \gamma) &= b \cos(b + \gamma) - b \cos(\gamma) \end{aligned} \tag{13}$$

Thus, if f is non-constant, then $\gamma \neq \pi/2$. Suppose, by way of contradiction, that $\gamma = \pi/2$. In this case (13) implies that $0 = 1 - \sin(b + \pi/2)$ and $\alpha b^2 \sin(b + \pi/2) = b \cos(b + \pi/2)$. The first yields that $b = 2\pi n$, for some $n \in \mathbb{Z}$. Substituting this value for b into the second equation yields $\alpha(2\pi n)^2 = 0$, and so $n = 0$. Hence, $b = 0$, in which case $f = 1$.

For $k \in \mathbb{Z}$, let $\gamma = \gamma^{(k,1)}(\alpha)$ denote the unique solution in the interval $(-\pi/2, \pi/2)$ to $\tan(\gamma) = -2\gamma\alpha + \alpha\pi/2 + 2\pi\alpha k + 1$, and set $b^{(k,1)}(\alpha) := -2\gamma^{(k,1)}(\alpha) + \pi/2 + 2\pi k$. As we will shortly see, $b^{(k,1)}$ and $\gamma^{(k,1)}$ completely determine the eigenfunctions and eigenvalues of Δ_η . We have introduced the extra index 1 to indicate that they give rise to solutions to the eigenvalue problem when η has a single atom; this will become important in the subsequent section where we consider measures with two atoms.

Notice, if $k = 0$, then $\gamma^{(k,1)} = \pi/4$ and $b^{(k,1)} = 0$; if $b^{(k,1)} = 0$, then $\gamma^{(k,1)} = \pi/4$ and $k = 0$; if $\gamma^{(k,1)} = \pi/4$, then $k = 0$ and $b^{(k,1)} = 0$.

Theorem 3.4. *The eigenvalues of Δ_η are $\lambda^{(k,1)} = -(b^{(k,1)}(\alpha))^2$ for $k \in \mathbb{Z}$, with corresponding eigenfunctions $f^{(k,1)}(x) := \sin(b^{(k,1)}(\alpha)x + \gamma^{(k,1)}(\alpha))$. Further, each eigenvalue has multiplicity one.*

Proof. This is a direct consequence of Lemma 3.3 with $\beta = \alpha$, $c_k^+ = \gamma^{(k,1)}$ and $\xi_k^+ = b^{(k,1)}$ and the observation that $-c_{-k}^- = c_k^+$ and $-\xi_{-k}^- = \xi_k^+$. \square

Note, the only eigenfunction $f^{(k,1)}$ with $\mathbf{f}^{(k,1)}$ continuous is the constant function $f^{(0,1)}$. Indeed, if there exists $k \in \mathbb{Z} \setminus \{0\}$ with $\mathbf{f}^{(k,1)}$ continuous, then $\sin(b^{(k,1)} + \gamma^{(k,1)}) = \sin(\gamma^{(k,1)})$, and so, by (13), we would have $\cos(\gamma^{(k,1)}) = 0$ as $b^{(k,1)} \neq 0$, contradicting the fact that $\gamma = \gamma^{(k,1)}(\alpha) \in (-\pi/2, \pi/2)$. Further, for $k_1, k_2 \in \mathbb{Z}$ with $k_1 \neq k_2$, by definition $\gamma^{(k_1,1)} \neq \gamma^{(k_2,1)}$ and, since $\gamma^{(k,1)} \in (-\pi/2, \pi/2)$ for $k \in \mathbb{Z} \setminus \{0\}$, we have $(b^{(k_1,1)})^2 \neq (b^{(k_2,1)})^2$. Hence, all eigenvalues have multiplicity one.

Corollary 3.5. *Letting $N_\eta: \mathbb{R}^+ \rightarrow \mathbb{R}$ denote the eigenvalue counting function of $-\Delta_\eta$, we have $\lim_{x \rightarrow \infty} \frac{\pi N_\eta(x)}{\sqrt{x}} = 1$.*

In contrast to the case when η is atomless, the eigenvalues of Δ_η do not occur in pairs. Indeed, let λ_k denote the k -th largest eigenvalue of Δ_η , then




FIGURE 1. Graphs of the eigenfunctions $f^{(k,1)}$ of Δ_η for $k \in \{1, 2, 3\}$, where $\eta = \Lambda + \pi^{-1}\delta_1$.

- (i) $\lim_{k \rightarrow \infty} \lambda_k / (k\pi + \pi/2)^2 = -1$,
- (ii) $\lim_{k \rightarrow +\infty} \gamma^{(k,1)} = -\lim_{k \rightarrow -\infty} \gamma^{(k,1)} = \pi/2$, and
- (iii) $\lim_{k \rightarrow +\infty} b^{(k,1)} / (2\pi k) = \lim_{k \rightarrow -\infty} b^{(k,1)} / (2\pi k) = 1$.

Example 3.6. For $\eta = \Lambda + \pi^{-1}\delta_1$ we have $\lambda^{(1,1)} \approx -29.3$, $\lambda^{(2,1)} \approx -130.4$ and $\lambda^{(3,1)} \approx -309.1$; see Figure 1 for a graphical representation of $f^{(1,1)}$, $f^{(2,2)}$ and $f^{(3,1)}$.

3.3. N = 2: Uniformly distributed Dirac point masses. Let α denote a positive real number, let $z_1, z_2 \in (0, 1]$ be such that $z_2 - z_1 = 1/2$ and let $\eta = \Lambda + \sum_{i=1}^2 \alpha \delta_{i/2}$. As in Section 3.2, without loss of generality, we may assume that $z_2 = 1$, and hence that $z_1 = 1/2$. The main results of this section are Theorem 3.8 and Corollary 3.9, in which we determine the eigenvalues and eigenfunctions of Δ_μ .

By (6), if $\Delta_\eta f = \lambda f$, for some $\lambda \in \mathbb{R}$, then there exist $b, a_1, a_2, \gamma_1, \gamma_2 \in \mathbb{R}$ with $\gamma_1 \in (-\pi/2, \pi/2]$ such that

$$f(x) = \begin{cases} a_1 \sin(bx + \gamma_1) & \text{if } x \in (0, 1/2], \\ a_2 \sin(bx + \gamma_2) & \text{if } x \in (1/2, 1]. \end{cases}$$

Corollary 3.1 implies the constant function $\mathbb{1}$ is an eigenfunction of Δ_η and the eigenspace $E_0 = \{f \in \mathcal{D}_\eta^2 : \Delta_\eta f \equiv 0\}$ is one-dimensional. In other words, if $b = 0$, then $f \in E_0$.

With this at hand we may assume that $b \neq 0$. From the system of equations given in (7) and (8) it follows that if f is an eigenfunction, then $a_1, a_2, b, \gamma_1, \gamma_2$ fulfil the following equations.

$$\begin{aligned} aba_1 \cos(\gamma_1) &= a_1 \sin(\gamma_1) - a_2 \sin(b + \gamma_2) \\ aba_2 \sin(b + \gamma_2) &= a_2 \cos(b + \gamma_2) - a_1 \cos(\gamma_1) \\ aba_2 \cos(b/2 + \gamma_2) &= a_2 \sin(b/2 + \gamma_2) - a_1 \sin(b/2 + \gamma_1) \\ aba_1 \sin(b/2 + \gamma_1) &= a_1 \cos(b/2 + \gamma_1) - a_2 \cos(b/2 + \gamma_2) \end{aligned} \tag{14}$$

As discussed directly above Corollary 3.1, we have $a_1, a_2 \neq 0$, since otherwise (14) yields $f \equiv 0$.

By (9), we have that

$$f_2(x) = \begin{cases} a_2 \sin(b(x + 1/2) + \gamma_2) & \text{if } x \in (0, 1/2], \\ a_1 \sin(b(x - 1/2) + \gamma_1) & \text{if } x \in (1/2, 1], \end{cases} \tag{15}$$

is also an eigenfunction of Δ_η . Hence, without loss of generality, we may assume $a_1 = 1$ and $|a_2| \leq 1$. Our aim is to find all tuples $(b, a_2, \gamma_1, \gamma_2) \in \mathbb{R}^4$ such that f is a non-constant eigenfunction. We start with the special cases that $a_2 = 1$ and $b = \pm 1/\alpha$. In the second step we discuss the case $a_2 = 1$ and $b \neq \pm 1/\alpha$. Noting that $(b, a_2, \gamma_1, \gamma_2)$ leads to an eigenfunction if and only if $(b, -a_2, \gamma_1, \gamma_2 + \pi)$ does, solves the case $a_2 = -1$. We then show that f is not an eigenfunction if $|a_2| < 1$.

Suppose that $a_2 = 1$ and $b = -1/\alpha$. The first two equations in (14) implies that $\sin(\gamma_1) = -\cos(b + \gamma_2)$. This yields that $b + \gamma_2 = \gamma_1 + \pi/2 + 2\pi k$ for some $k \in \mathbb{Z}$. Substituting this into the first equation of (14) implies that $\gamma_1 = 0$ and hence, $\gamma_2 = -b + \pi/2 + 2\pi k$ for some $k \in \mathbb{Z}$. Subsequently, the third and fourth equations of (14) yields that f is an eigenfunction if and only if $\alpha = 1/(\pi + 2\pi m)$ for some $m \in \mathbb{Z}$, in which case $\gamma_2 \bmod 2\pi = 3\pi/2$.

One can show in a similar manner, that if $a_2 = 1$ and $b = 1/\alpha$, then f is an eigenfunction of Δ_η if and only if $\alpha = 1/(2 \arctan(1/2) - 2 \arctan(2) + 2\pi m)$ for some $m \in \mathbb{Z}$; in which case

$$\gamma_1 = \arctan(2) \quad \text{and} \quad \gamma_2 = \arctan(2) - 2 \arctan(1/2) + \pi/2 \bmod 2\pi.$$

Corollary 3.7. *If there exists $m \in \mathbb{N}_0$ with $\alpha = \alpha' := 1/(\pi + 2\pi m)$ or $\alpha = \alpha'' := 1/(2 \arctan(1/2) - 2 \arctan(2) + 2\pi m)$, then $\lambda = -1/\alpha^2$ is an eigenvalue of Δ_η with multiplicity one. The corresponding eigenfunction is*

$$f(x) = \begin{cases} \sin(-(1/\alpha)x) & \text{if } x \in (0, 1/2] \text{ and } \alpha = \alpha', \\ \sin(-(1/\alpha)x + 3\pi/2) & \text{if } x \in (1/2, 1] \text{ and } \alpha = \alpha', \\ \sin((1/\alpha)x + \arctan(2)) & \text{if } x \in (0, 1/2] \text{ and } \alpha = \alpha'', \\ \sin((1/\alpha)x + \arctan(2) - 2 \arctan(1/2) + \pi/2) & \text{if } x \in (1/2, 1] \text{ and } \alpha = \alpha''. \end{cases}$$

We now consider the case $a_2 = 1$ and $b \in \mathbb{R} \setminus \{0, \pm 1/\alpha\}$. From (14) it follows that $\cos(\gamma_1) \neq 0$ and $\cos(b/2 + \gamma_2) \neq 0$. This implies f is discontinuous at the atoms and hence that $b + \gamma_2 - \gamma_1 \neq 0$ and $\gamma_1 - \gamma_2 \neq 0$; thus $\gamma_1 \neq \pi/2$. Define $g: \mathcal{I} \rightarrow \mathbb{R}$ by $g(x) := \sin((b + \gamma_2 - \gamma_1)x + \gamma_1)$ and set $\beta_1 := \alpha b/(b + \gamma_2 - \gamma_1)$. Setting $\beta = \beta_1$, $c = \gamma_1$ and $\xi = b + \gamma_2 - \gamma_1$, the equalities in (14) imply those of (12), and so there exists a $k \in \mathbb{Z}$ with $b + \gamma_2 = -\gamma_1 \pm \pi/2 + 2\pi k$. If $b + \gamma_2 = -\gamma_1 + \pi/2 + 2\pi k$, then $\sin(b + \gamma_2) = \cos(\gamma_1)$ and $\sin(b/2 + \gamma_1) = \cos(b/2 + \gamma_2)$. Combining this with (14) yields

$$\begin{aligned} \tan(\gamma_1) &= 1 + \alpha b, & \tan(b + \gamma_2) &= 1/(1 + \alpha b), \\ \tan(b/2 + \gamma_2) &= 1 + \alpha b, & \tan(b/2 + \gamma_1) &= 1/(1 + \alpha b). \end{aligned} \tag{16}$$

For $k \in \mathbb{Z}$, let $\gamma_1^{(k,2)}$ denote the unique solution of $\tan(\gamma_1) = 1 - 4\alpha\gamma_1 + \alpha\pi + 2\pi\alpha k$. Substituting the first equation of (16) into the last, we observe that $\gamma_1 = \gamma_1^{(k,2)}$, $b = b^{(k,2)} = -4\gamma_1^{(k,2)} + \pi + 2\pi k$ and $\gamma_2 = \gamma_2^{(k,2)} = -b^{(k,2)} - \gamma_1^{(k,2)} + \pi/2$, for some $k \in \mathbb{Z}$. Observe that $\gamma_1^{(0,2)} = \pi/4$, and hence that $b^{(0,2)} = 0$, contradicting our initial assumption that f is a non-constant eigenfunction.

Similar to the case when $N = 1$, if $b + \gamma_2 = -\gamma_1 - \pi/2 + 2\pi k$, then analogue calculations yield the same eigenfunctions, namely that $b = -b^{(k,2)}$, $\gamma_1 = -\gamma_1^{(k,2)}$ and $\gamma_2 = -\gamma_2^{(k,2)}$.

A direct calculation shows, for $x \in (0, 1/2]$, that $\sin(b^{(k,2)}(x + 1/2) + \gamma_2^{(k,2)}) = \sin(b^{(k,2)}x + \gamma_1^{(k,2)} - \pi k)$. This means that if k is even, the eigenfunctions are periodic with period $1/2$, namely $f = f_2$ where f_2 is defined as in (15). This means, as discussed in Corollary 3.2, that these are concatenations of rescaled solutions for a measure with one atom, namely if $k = 2m$, then $\gamma_1^{(k,2)}(\alpha) = \gamma^{(m,1)}(2\alpha)$ and $b^{(k,2)}(\alpha) = 2b^{(m,1)}(2\alpha)$. On the other hand if k is odd, then $f = -f_2$. Summarising, we have, for $k \in \mathbb{Z} \setminus \{0\}$, that

$$f^{(k,2)}(x) = \begin{cases} \sin(b^{(k,2)}x + \gamma_1^{(k,2)}) & \text{if } x \in (0, 1/2], \\ \sin(b^{(k,2)}x + \gamma_2^{(k,2)}) & \text{if } x \in (1/2, 1] \end{cases}$$

is an eigenfunction of Δ_η with corresponding eigenvalue $\lambda^{(k,2)} = -(b^{(k,2)})^2$, and $f^{(0,2)} := \mathbb{1}$ is an eigenfunction of Δ_η with corresponding eigenvalue $\lambda^{(0,2)} := 0$.

If α is of the form discussed in Corollary 3.7, then the eigenfunction f given there belongs to $\{f^{(k,2)} : k \in \mathbb{Z}\}$. Indeed, if there exists $m \in \mathbb{N}_0$ with $\alpha = 1/(\pi + 2\pi m)$, then $f = f^{(-m-1,2)}$, and if there exists $m \in \mathbb{N}$ with $\alpha = 1/(2 \arctan(1/2) - 2 \arctan(2) + 2\pi m)$, then $f = f^{(m,2)}$.

To conclude, by way of contradiction, we show that there does not exist any other eigenfunction of Δ_η other than those discussed above. To this end, assume that h is an eigenfunction of Δ_η in the orthogonal complement of $\text{span}\{f^{(k,2)} : k \in \mathbb{Z}\}$. By the discussion following (9), we have

$$h(x) = \begin{cases} \sin(bx + \gamma_1) & \text{if } x \in (0, 1/2], \\ a_2 \sin(bx + \gamma_2) & \text{if } x \in (1/2, 1], \end{cases}$$

for some $b \in \mathbb{R}$ and $a_2, \gamma_1, \gamma_2 \in \mathbb{R}$ with $|a_2| < 1$. Letting $\psi(x) = x/2$ for all $x \in \mathbb{R}$, by definition and Theorems 2.11 and 3.4, we have $\{f^{(2m,2)} \circ \psi : m \in \mathbb{Z}\}$ forms an orthonormal basis for $L^2_{\Lambda+2\alpha\delta_1}$, and thus, $\{f^{(2m,2)}|_{(0,1/2]} : m \in \mathbb{Z}\}$ is an orthonormal basis for $L^2_{\Lambda|_{(0,1/2]} + \alpha\delta_{1/2}}$. This implies that $\langle h, f \rangle_\eta = 0$ for all $f \in \text{span}\{f^{(2m,2)} : m \in \mathbb{Z}\}$. Using this, we obtain, for all Borel sets $A \subset (0, 1/2]$, that

$$\int \mathbb{1}_A(x)(h(x) + h(x + 1/2)) d\eta(x) = 0.$$




FIGURE 2. Graphs of the eigenfunctions $f^{(k,2)}$ of Δ_η , for $k \in \{-2, -1, 1, 2, 3, 4\}$, where $\eta = \Lambda + \pi^{-1}\delta_{1/2} + \pi^{-1}\delta_1$.

This yields $\eta(\{x \in (0, 1/2]: h(x) > -h(x + 1/2)\}) = \eta(\{x \in (0, 1/2]: h(x) < -h(x + 1/2)\}) = 0$. Hence, we have $h(x) = -h(x + 1/2)$ for η -almost all $x \in (0, 1/2]$. Since $h \in \mathcal{D}_\eta^2$, it is left-continuous, which implies that $h(x) = -h(x + 1/2)$ for all $x \in (0, 1/2]$. Therefore,

$$h(x) = \begin{cases} \sin(bx + \gamma_1) & \text{if } x \in (0, 1/2], \\ -\sin(b(x - 1/2) + \gamma_1) & \text{if } x \in (1/2, 1], \end{cases}$$

contradicting our initial assumption that $|a_2| < 1$.

For $k_1, k_2 \in \mathbb{Z}$ with $k_1 \neq k_2$ we observe that $\gamma^{(k_1,1)} \neq \gamma^{(k_2,1)}$. Further, an elementary calculation shows that $(b^{(k_1,1)})^2 \neq (b^{(k_2,1)})^2$, which implies all eigenvalues of Δ_η have multiplicity one. Combining the above we obtain the following.

Theorem 3.8. *The eigenvalues of Δ_η are $\lambda^{(k,2)} := -(b^{(k,2)})^2$ for $k \in \mathbb{Z}$, with corresponding eigenfunctions $f^{(k,2)}$. Further, each eigenvalue has multiplicity one.*

Notice the only eigenfunction $f^{(k,2)}$ with $\mathbf{f}^{(k,2)}$ continuous at an atom is the constant function $f^{(0,2)}$. Indeed, if there exists a $k \in \mathbb{Z} \setminus \{0\}$ such that $\mathbf{f}^{(k,2)}$ is continuous at an atom, we would have $\cos(\gamma_1^{(k,2)}) = 0$ or $\cos(b^{(k,2)}/2 + \gamma_2^{(k,2)}) = 0$. Substituting the defining equations for $b^{(k,2)}$ and $\gamma_2^{(k,2)}$ into the latter, we obtain, in both cases, that $\gamma_1^{(k,2)} = \pi/2$, which contradicts the fact that $\gamma_1^{(k,2)} \in (-\pi/2, \pi/2)$. We also note the following.

- (i) $\lim_{k \rightarrow +\infty} \gamma_1^{(k,2)} = -\lim_{k \rightarrow -\infty} \gamma_1^{(k,2)} = \pi/2$
- (ii) $\lim_{k \rightarrow +\infty} \gamma_2^{(k,2)} \bmod 2\pi = \pi$ and $\lim_{k \rightarrow -\infty} \gamma_2^{(k,2)} \bmod 2\pi = 0$
- (iii) $\lim_{k \rightarrow +\infty} b^{(k,2)}/(2\pi k) = \lim_{k \rightarrow -\infty} b^{(k,2)}/(2\pi k) = 1$

Corollary 3.9. *Letting $N_\eta: \mathbb{R}^+ \rightarrow \mathbb{R}$ denote the eigenvalue counting function of $-\Delta_\eta$, we have $\lim_{x \rightarrow \infty} \frac{\pi N_\eta(x)}{\sqrt{x}} = 1$.*

In contrast to the case when η is atomless and the case when $N = 1$, the eigenvalues of Δ_η do not occur in pairs. However, we have that $\lim_{k \rightarrow \infty} -b^{(k,2)}/b^{(-k-1,2)} = 1$.

Example 3.10. For $\eta = \Lambda + \pi^{-1}\delta_{\frac{1}{2}} + \pi^{-1}\delta_1$, we have that $\lambda^{(-2,2)} = 4\pi^2$, $\lambda^{(-1,2)} = \pi^2$, $\lambda^{(1,2)} \approx 21.8$, $\lambda^{(2,2)} \approx 106.9$, $\lambda^{(3,2)} \approx 267.2$ and $\lambda^{(4,2)} \approx 505.3$; see Figure 2 for a graphical representation of $f^{(k,2)}$ for $k \in \{-2, -1, 1, 2, 3, 4\}$.

ACKNOWLEDGEMENTS

The authors thank Hendrik Vogt for several enlightening discussions. They acknowledge the support of the Deutsche Forschungsgemeinschaft (DFG grant Ke 1440/3-1). Part of this work was completed while the first and second authors were visiting Institut Mittag-Leffler as part of the research program *Fractal Geometry and Dynamics*. They are extremely grateful to the organisers and staff for their very kind hospitality, financial support and a stimulating atmosphere. The third author acknowledges the support from the BremenIDEA program at Universität Bremen.

REFERENCES

- [1] P. Arzt. Measure theoretic trigonometric functions. *J. Fractal Geom.*, 2(2):115–169, 2015.
- [2] R. Beals and P. C. Greiner. Strings, waves, drums: spectra and inverse problems. *Anal. Appl. (Singap.)*, 7(2):131–183, 2009.
- [3] M. V. Berry. Distribution of modes in fractal resonators. In *Structural stability in physics (Proc. Internat. Symposia Appl. Catastrophe Theory and Topological Concepts in Phys., Inst. Inform. Sci., Univ. Tübingen, Tübingen, 1978)*, volume 4 of *Springer Ser. Synergetics*, pages 51–53. Springer, Berlin, 1979.
- [4] M. V. Berry. Some geometric aspects of wave motion: wavefront dislocations, diffraction catastrophes, diffractals. In *Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979)*, Proc. Sympos. Pure Math., XXXVI, pages 13–28. Amer. Math. Soc., Providence, R.I., 1980.
- [5] N. Biggs. *Algebraic graph theory*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 1993.
- [6] W. Feller. Generalized second order differential operators and their lateral conditions. *Illinois J. Math.*, 1:459–504, 1957.
- [7] U. Freiberg. Analytical properties of measure geometric Krein-Feller-operators on the real line. *Math. Nachr.*, 260:34–47, 2003.
- [8] U. Freiberg. A survey on measure geometric Laplacians on Cantor like sets. *Arab. J. Sci. Eng. Sect. C Theme Issues*, 28(1):189–198, 2003. Wavelet and fractal methods in science and engineering, Part I.
- [9] U. Freiberg. Spectral asymptotics of generalized measure geometric Laplacians on Cantor like sets. *Forum Math.*, 17(1):87–104, 2005.
- [10] U. Freiberg and M. Zähle. Harmonic calculus on fractals—a measure geometric approach. I. *Potential Anal.*, 16(3):265–277, 2002.
- [11] T. Fujita. A fractional dimension, self-similarity and a generalized diffusion operator. In *Probabilistic methods in mathematical physics (Katata/Kyoto, 1985)*, pages 83–90. Academic Press, Boston, MA, 1987.
- [12] C. Gordon, D. Webb, and S. Wolpert. Isospectral plane domains and surfaces via Riemannian orbifolds. *Invent. Math.*, 110(1):1–22, 1992.
- [13] P. R. Halmos. *Measure Theory*. D. Van Nostrand Company, Inc., New York, N. Y., 1950.
- [14] X. Jin. Spectral representation of one-dimensional Liouville brownian motion and Liouville brownian excursion, 2017.
- [15] I. S. Kac and M. G. Kreĭn. Criteria for the discreteness of the spectrum of a singular string. *Izv. Vysš. Učebn. Zaved. Matematika*, 1958(2 (3)):136–153, 1958.
- [16] M. Kac. Can one hear the shape of a drum? *Amer. Math. Monthly*, 73(4, part II):1–23, 1966.
- [17] M. Kesseböhmer, T. Samuel, and H. Weyer. A note on measure-geometric Laplacians. *Monatsh. Math.*, 181(3):643–655, 2016.
- [18] M. Kesseböhmer, T. Samuel, and H. Weyer. Measure-geometric Laplacians for discrete distributions. In To appear in *Horizons of Fractal Geometry and Complex Dimensions*. Contemp. Math., 2017.
- [19] J. Kigami and M. L. Lapidus. Self-similarity of volume measures for Laplacians on p.c.f. self-similar fractals. *Comm. Math. Phys.*, 217(1):165–180, 2001.
- [20] S. Kotani and S. Watanabe. Kreĭn’s spectral theory of strings and generalized diffusion processes. In *Functional analysis in Markov processes (Katata/Kyoto, 1981)*, volume 923 of *Lecture Notes in Math.*, pages 235–259. Springer, Berlin-New York, 1982.
- [21] M. L. Lapidus. Fractal drum, inverse spectral problems for elliptic operators and a partial resolution of the Weyl-Berry conjecture. *Trans. Amer. Math. Soc.*, 325(2):465–529, 1991.
- [22] M. L. Lapidus. Vibrations of fractal drums, the Riemann hypothesis, waves in fractal media and the Weyl-Berry conjecture. In *Ordinary and partial differential equations, Vol. IV (Dundee, 1992)*, volume 289 of *Pitman Res. Notes Math. Ser.*, pages 126–209. Longman Sci. Tech., Harlow, 1993.
- [23] M. L. Lapidus and C. Pomerance. Fonction zêta de Riemann et conjecture de Weyl-Berry pour les tambours fractals. *C. R. Acad. Sci. Paris Sér. I Math.*, 310(6):343–348, 1990.
- [24] M. L. Lapidus and C. Pomerance. The Riemann zeta-function and the one-dimensional Weyl-Berry conjecture for fractal drums. *Proc. London Math. Soc. (3)*, 66(1):41–69, 1993.
- [25] M. L. Lapidus and C. Pomerance. Counterexamples to the modified Weyl-Berry conjecture on fractal drums. *Math. Proc. Cambridge Philos. Soc.*, 119(1):167–178, 1996.
- [26] J. Milnor. Eigenvalues of the Laplace operator on certain manifolds. *Proc. Nat. Acad. Sci. U.S.A.*, 51:542, 1964.
- [27] M. Reed and B. Simon. *Methods of modern mathematical physics. I*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, second edition, 1980. Functional analysis.
- [28] R. Rhodes and V. Vargas. Spectral dimension of Liouville quantum gravity. *Ann. Henri Poincaré*, 15(12):2281–2298, 2014.
- [29] H. Urakawa. Bounded domains which are isospectral but not congruent. *Ann. Sci. École Norm. Sup. (4)*, 15(3):441–456, 1982.
- [30] H. Weyl. Über die Abhängigkeit der Eigenschwingungen einer Membran von deren Begrenzung. *J. Angew. Math.*, 141:1–11, 1912.