

EXPLICIT SOLUTIONS OF RICCI FLOW EQUATION FOR LOCALLY HOMOGENEOUS \mathbb{S}^1 -TRIPLES ON COMPACT RIEMANN SURFACES

ARASH BAZDAR

ABSTRACT. Let (M, g) be a compact Riemann surfaces, and $\pi : P \rightarrow M$ be a principal \mathbb{S}^1 -bundle over M endowed with a connection A . Fixing an inner product on the Lie algebra of \mathbb{S}^1 , the connection A and metric g define a Riemannian metric g_A on P . In this article, we show that the Ricci flow equation of metric g_A is equivalent to a system of differential equations. We will give an explicit solution of normalized Ricci flow equation of metric g_A in the case where the base manifold is of constant curvature and the initial connection A is Yang-Mills. Finally, we will describe some asymptotic behaviors of these flows.

Keywords: Ricci flow, theory of connections, geometric structure

1. LOCALLY HOMOGENEOUS TRIPLES WITH STRUCTURE GROUP $K = \mathbb{S}^1$ OVER RIEMANN SURFACES

1.1. Metric connection on principal K -bundles. A Riemannian metric g on a differentiable manifold M is defined to be locally homogeneous if, for every two points $x, x' \in M$ there exist open neighborhoods $U \ni x, U' \ni x'$ and an isometry $\varphi : U \rightarrow U'$ such that $\varphi(x) = x'$. Inspiring from this definition we consider the following geometric objects:

Definition 1.1. Let M be a smooth manifold and K be a compact Lie group. A locally homogeneous triple with structure group K on M is a triple $(g, P \xrightarrow{\pi} M, A)$, where $\pi : P \rightarrow M$ is a principal K -bundle on M , g is Riemannian metric on M , and A is connection on P such that the following locally homogeneity condition is satisfied: for every two points $x, x' \in M$ there exists an isometry $\varphi : U \rightarrow U'$ between open neighborhoods $U \ni x, U' \ni x'$ with $\varphi(x) = x'$, and a φ -covering bundle isomorphism $\Phi : P_U \rightarrow P_{U'}$ such that $\Phi^*(A_{U'}) = A_U$.

In these formulae, for an open set $U \subset M$, the subscript U is used to denote the restriction of the indicated objects to U . For connections on principal bundles we adopt the conventions of [4], so in Definition 1.1 the symbol A stands for a K -invariant horizontal distribution of P . Fixing an ad-invariant inner product on the Lie algebra \mathfrak{k} of K , one can endow P with a locally homogeneous Riemannian metric g_A called the connection metric. Let $\langle\langle \cdot, \cdot \rangle\rangle$ be an ad-invariant inner product on \mathfrak{k} and define a Riemannian metric g_A on P characterized by the following condition

1. The canonical bundle isomorphism $V_P \simeq P \times \mathfrak{k}$ is an orthogonal bundle isomorphism with respect to the inner products defined by g_A and $\langle\langle \cdot, \cdot \rangle\rangle$.
2. The restriction of π_* to the horizontal subbundle $A \subset T_P$ gives an orthogonal bundle isomorphism $A \rightarrow \pi^*(T_M)$.

3. The direct sum decomposition $T_P \simeq A \oplus V_P$ of the tangent bundle T_P is g_A -orthogonal.

Remark 1. Let $(g, P \xrightarrow{\pi} M, A)$ be a locally homogeneous K -triple over M , and let $\langle\langle \cdot, \cdot \rangle\rangle$ be an ad-invariant inner product on \mathfrak{k} . The connection metric g_A on P is locally homogeneous.

A very well-known theorem of Singer [6] assert that any locally homogeneous, complete, simply connected Riemannian manifold is homogeneous. Let $\pi : \tilde{M} \rightarrow M$ be the universal cover of M . The metric $\tilde{g} := \pi^*g$ defines a complete Riemannian metric on the simply connected manifold \tilde{M} . Since (\tilde{M}, \tilde{g}) is locally homogeneous, by theorem of Singer, it is also homogeneous. In particular, \tilde{M} admits a connected transitive group of isometries denoted by G . Therefore, For a compact, locally homogeneous Riemannian manifold (M, g) , there exists a homogeneous model geometry given by (\tilde{M}, G) . Generalizing the method of Singer, one can prove a similar classification theorem for locally homogeneous triples [1], [2].

A model geometry is a simply connected manifold X together with a transitive action of a Lie group G on X with compact stabilizers. In dimension three there exists (up to equivalence) eight maximal model geometries which admit compact quotients: \mathbf{E}^3 , \mathbf{S}^3 , \mathbf{H}^3 , $\mathbf{S}^2 \times \mathbb{R}$, $\mathbf{H}^2 \times \mathbb{R}$, $\widetilde{\mathbf{SL}_2(\mathbb{R})}$, \mathbf{Nil} and \mathbf{Sol} . We refer to [8], [7] for the explicit description of the pair (X, G) corresponding to each symbol above. A geometric structure on a manifold M is a diffeomorphism from M to X/Γ for some model geometry (X, G) , where $\Gamma \subset G$ is a discrete subgroup of G acting freely on X . Excepting \mathbf{H}^3 and \mathbf{Sol} , for any model geometry of Thurston geometries there exist compact 3-manifolds with a geometric structure which are associated with locally homogeneous triples with structure group \mathbb{S}^1 . For instance any non-trivial \mathbb{S}^1 -bundle on a Riemann surface of genus g has a geometric (X, G) -structure, where

1. $(X, G) = \mathbf{S}^3$ when $g = 0$,
2. $(X, G) = \mathbf{Nil}$ when $g = 1$,
3. $(X, G) = \widetilde{\mathbf{SL}_2(\mathbb{R})}$ when $g \geq 2$.

The trivial \mathbb{S}^1 -bundles over Riemann surfaces have geometric structures with model geometry $\mathbf{S}^2 \times \mathbb{R}$, \mathbf{E}^3 , or $\mathbf{H}^2 \times \mathbb{R}$. Moreover any \mathbb{S}^1 -bundle over a Riemann surface has a geometric metric which is associated with a locally homogeneous triple in the sense of Definition 1.1. These examples shows the abundance of geometric principal \mathbb{S}^1 -bundles in the class of geometric three manifolds.

Suppose M is a compact manifold, and K is a compact Lie group, then any locally homogeneous triple $(g, P \xrightarrow{\pi} M, A)$ with structure group K on M can be identified with a Γ -quotient of the homogeneous triple $(\tilde{g} := \pi^*g, Q := \pi^*P \rightarrow \tilde{M}, B)$ on the universal cover \tilde{M} [2]. Thanks to this result, instead of studying the Ricci flow of locally homogeneous triples over compact Riemann surfaces, it is enough to study the Ricci flow on compact Riemann surfaces with metric of constant curvature. We will study the (normalized) Ricci flow equation of locally homogeneous \mathbb{S}^1 -triples over compact surfaces M with a metric of constant curvature. We will give an explicit solution of the (normalized) Ricci flow equations for the metric g_A in the case where the initial connection A is chosen to be Yang-Mills. Note that this is justified because it is proved that an \mathbb{S}^1 -triple $(g, P \xrightarrow{\pi} M, A)$ is locally homogeneous if and only if the connection A is Yang-Mills [2]. Finally, we will study some asymptotic behaviors of these flows in several interesting cases as it is down in [3].

1.2. Levi-Civita connection associated with connection metric on circle bundles over Riemann surfaces. Let $\pi : P \rightarrow M$ be a principal \mathbb{S}^1 -bundle over a Riemann surface M , g a Riemannian metric on M and A a connection on P . We can endow P with the connection metric g_A . We summarize these information by saying that $(g, P \xrightarrow{\pi} M, A)$ is a \mathbb{S}^1 -triple. Identify the Lie algebra of \mathbb{S}^1 with \mathbb{R} and denote the connection 1-form associated with A by $\omega_A \in A^1(P, \mathbb{R})$. Let $\varphi = (\varphi_1, \varphi_2)$ be an orthonormal frame of Riemann surface (M, g) over an open neighborhood $U \subset M$ and let $(e_1 := \tilde{\varphi}_1, e_2 := \tilde{\varphi}_2)$ be the A -horizontal lifts of φ defined on $P_U := \pi^{-1}(U)$. Therefore, we have

$$\omega(e_i) = 0, \pi_*(e_i) = \varphi_i \quad \text{for } i = 1, 2.$$

Let ξ be an element of the Lie algebra of \mathbb{S}^1 and let $\xi^\# : P \rightarrow T_P$ be the fundamental vector field defined on P trivializing the vertical bundle $V_P \simeq P \times \mathfrak{k}$. The vertical section $\xi^\# \in \mathcal{X}^v(P)$ at a point $y \in P$ is defined by

$$\xi_y^\# := \left. \frac{d}{dt} \right|_{t=0} (y \exp(t\xi)).$$

Note that we can define a smooth map $f : P \rightarrow \mathbb{R}$ at a point $y \in M$ by

$$f(y) := \sqrt{g_A(\xi_y^\#, \xi_y^\#)} \quad \text{where } y \in P_x$$

For all $y \in P_U$ set $e_3(y) := f(y)^{-1} \xi_y^\#$ and note that (e_1, e_2, e_3) is an orthonormal frame on P_U with respect to metric g_A . Use (η^1, η^2) to denote the dual frame associated with (φ_1, φ_2) over $U \subset M$. By a well-known theorem of differential geometry [5], there exists a unique skew-symmetric matrix (η_j^i) of real-valued 1-forms $\eta_j^i \in A^1(U, \mathbb{R})$ such that

$$(1) \quad \begin{aligned} d\eta^1 &= -\eta_2^1 \wedge \eta^2, \\ d\eta^2 &= -\eta_1^2 \wedge \eta^1. \end{aligned}$$

Putting $\theta^1 = \pi^* \eta^1$, $\theta^2 = \pi^* \eta^2$ and $\theta^3 = f\omega$ we will obtain a co frame $(\theta^1, \theta^2, \theta^3)$ dual to (e_1, e_2, e_3) on P_U . Taking pullback of both sides of equations (1) we obtain

$$(2) \quad \begin{aligned} d\theta^1 &= -\pi^* \eta_2^1 \wedge \theta^2, \\ d\theta^2 &= -\pi^* \eta_1^2 \wedge \theta^1. \end{aligned}$$

The curvature 2-form of connection A is denoted by $F_\omega = d\omega \in A^2(P, \mathbb{R})$. Since \mathbb{S}^1 is abelian F_ω is \mathbb{S}^1 -invariant and descend to a closed 2-form on M . Using the dual frame $(\theta^1, \theta^2, \theta^3)$ on P_U , the curvature 2-form F_ω can be written as follow

$$(3) \quad F_\omega = c \theta^1 \wedge \theta^2 \quad \text{where } c \in \mathcal{C}^\infty(P_U, \mathbb{R}).$$

The map $f \in \mathcal{C}^\infty(P)$ is \mathbb{S}^1 -invariant and it descends to a smooth map $f : M \rightarrow \mathbb{R}$, so there exists the maps $f_1, f_2 \in \mathcal{C}^\infty(P_U)$ such that $df = f_1 \theta^1 + f_2 \theta^2$. Using this we deduce that

$$(4) \quad d\theta^3 = d(f\omega) = fd\omega + df \wedge \omega = fc \theta^1 \wedge \theta^2 + \frac{f_1}{f} \theta^1 \wedge \theta^3 + \frac{f_2}{f} \theta^2 \wedge \theta^3.$$

The equations (2) and (4) lead to the following system of differential equations

$$(5) \quad \begin{aligned} d\theta^1 &= -\pi^* \eta_2^1 \wedge \theta^2, \\ d\theta^2 &= -\pi^* \eta_1^2 \wedge \theta^1, \\ d\theta^3 &= fc \theta^1 \wedge \theta^2 + \frac{f_1}{f} \theta^1 \wedge \theta^3 + \frac{f_2}{f} \theta^2 \wedge \theta^3. \end{aligned}$$

The equations (5) can be used to find the components of Levi-Civita connection for metric g_A . In fact, we search a unique skew-symmetric matrix (θ_j^i) of real-valued 1-forms $\theta_j^i \in A^1(P_U, \mathbb{R})$ satisfying the following differential equations

$$(6) \quad d\theta^i = -\sum_j \theta_j^i \wedge \theta^j.$$

The solutions of these equations are the components of the Levi-Civita connection 1-form associated to the metric g_A . These solutions can be find in two steps:

Step 1. First of all, we try to find the real-valued 1-forms μ_j^i on P_U which satisfy in (6). Unfortunately, there exists no reason to be sure that (μ_j^i) is skew-symmetric. Fortunately, we can modify this solution (μ_j^i) in an appropriate way such that the modified solution is skew-symmetric and still satisfies in (6).

Step 2. Put $\theta_j^i := \mu_j^i + \sum_k S_{jk}^i \theta^k$ and search a unique tensor S such that:

1. $S_{jk}^i = S_{kj}^i$ for all i, j, k .
2. (θ_j^i) is skew-symmetric.

It is easy to verify that the components of skew-symmetric matrix (θ_j^i) defined by $\theta_j^i := \mu_j^i + \sum_k S_{jk}^i \theta^k$ satisfy in (6). Therefore, we just need to determined the unique tensor S as above. These two steps give an algorithm to find the skew-symmetric matrix (θ_j^i) of real valued 1-forms which defines the Levi-Civita connection of metric g_A . First step is to find $\theta_j^i \in A^1(P_U, \mathbb{R})$ satisfying the equation (6). Thanks to the equations (5) we conclude that the following matrix of 1-forms has this property.

$$\mu_j^i := \begin{pmatrix} 0 & \pi^* \eta_2^1 & 0 \\ \pi^* \eta_1^2 & 0 & 0 \\ f_c \theta^2 + \frac{f_1}{f} \theta^3 & \frac{f_2}{f} \theta^3 & 0 \end{pmatrix}.$$

Second step is to modify μ_j^i by adding a tensor field S_{jk}^i such that $S_{jk}^i = S_{kj}^i$ for all i, j, k . Therefore, we have to solve the following system of equations

$$(7) \quad \begin{aligned} \theta_j^i &= \mu_j^i + \sum_k S_{jk}^i \theta^k \\ \theta_j^i &= -\theta_i^j \\ S_{jk}^i &= S_{kj}^i \end{aligned}.$$

If $i = j$, then by the skew-symmetry property of θ_i^i we have

$$(8) \quad 0 = \theta_i^i = \mu_i^i + \sum_k S_{ik}^i \theta^k = \sum_k S_{ik}^i \theta^k.$$

So, for all i and k we have $S_{ik}^i = S_{ki}^i = 0$. By the first equation of (7) we have

$$(9) \quad \theta_2^1 = \mu_2^1 + \sum_k S_{2k}^1 \theta^k = \pi^* \eta_2^1 + \sum_k S_{2k}^1 \theta^k,$$

$$(10) \quad \theta_1^2 = \mu_1^2 + \sum_k S_{1k}^2 \theta^k = \pi^* \eta_1^2 + \sum_k S_{1k}^2 \theta^k.$$

Using (9), (10) and $\theta_2^1 = -\theta_1^2$ we get $S_{2k}^1 = -S_{1k}^2$. In the same way, the first equation of (7) implies

$$(11) \quad \theta_3^1 = \mu_3^1 + \sum_k S_{3k}^1 \theta^k = \sum_k S_{3k}^1 \theta^k,$$

$$(12) \quad \theta_1^3 = \mu_1^3 + \sum_k S_{1k}^3 \theta^k = f_c \theta^2 + \frac{f_1}{f} \theta^3 + \sum_k S_{1k}^3 \theta^k.$$

Using (11), (12) and $\theta_3^1 = -\theta_1^3$ we obtain

$$S_{11}^3 = 0, \quad S_{12}^3 + S_{32}^1 = -fc, \quad S_{33}^1 = -\frac{f_1}{f}.$$

In the same way, we have the following equations

$$(13) \quad \theta_3^2 = \mu_3^2 + \sum_k S_{3k}^2 \theta^k = \sum_k S_{3k}^2 \theta^k,$$

$$(14) \quad \theta_2^3 = \mu_2^3 + \sum_k S_{2k}^3 \theta^k = \frac{f_2}{f} \theta^3 + \sum_k S_{2k}^3 \theta^k,$$

Using (13), (14) and $\theta_3^2 = -\theta_2^3$ we obtain

$$S_{22}^3 = 0, \quad S_{33}^2 = -\frac{f_2}{f}, \quad S_{21}^3 + S_{31}^2 = 0.$$

Using the above equations, we deduce that $S_{12}^3 = S_{32}^1 = -\frac{1}{2}fc$ and $S_{13}^2 = \frac{1}{2}fc$. So, we have proved the next proposition.

Proposition 1.2. *Let $\pi : P \rightarrow M$ be a principal \mathbb{S}^1 -bundle over a Riemann surface (M, g) endowed with a connection A . Using the same notation as above, the Levi-Civita connection (θ_j^i) of connection metric g_A is given by*

- (1) $\theta_2^1 = \pi^*(\eta_2^1) - \frac{1}{2}fc\theta^3$.
- (2) $\theta_3^1 = -\frac{1}{2}fc\theta^2 - \frac{f_1}{f}\theta^3$.
- (3) $\theta_3^2 = \frac{1}{2}fc\theta^1 - \frac{f_2}{f}\theta^3$.

1.3. The curvature tensors of connection metric on \mathbb{S}^1 -bundles over Riemann surfaces. Let $\pi : P \rightarrow M$ be a principal \mathbb{S}^1 -bundle over a Riemann surface (M, g) endowed with connection metric g_A . Let (θ_j^i) be the skew-symmetric matrix of 1-forms which defines the Levi-Civita connection of g_A as in Proposition 1.2. Using the co frame $(\theta^1, \theta^2, \theta^3)$ defined on P_U , the curvature 2-form F_ω can be written as $F_\omega = c\theta^1 \wedge \theta^2$, where c is a smooth function on P_U . We shall use the following decomposition in next calculations.

$$(15) \quad dc = \alpha\theta^1 + \beta\theta^2, \quad df = f_1\theta^1 + f_2\theta^2, \quad df_i = f_{i1}\theta^1 + f_{i2}\theta^2 \quad \text{for } i = 1, 2.$$

If $\Omega^\theta = (\Omega_j^i)$ denotes the curvature 2-form of Levi-Civita connection $\theta = (\theta_j^i)$, then we will use the structure equation $\Omega_j^i = d\theta_j^i + \sum_{k=1}^3 \theta_k^i \wedge \theta_j^k$ to determine the components of Ω^θ [4]. In order to calculate $\Omega_2^1 = d\theta_2^1 + \theta_3^1 \wedge \theta_2^3$, we need the following

$$(16) \quad d\theta_2^1 = \pi^*d\eta_2^1 - \frac{1}{2}d(fc\theta^3).$$

we also know that

$$d\theta^3 = d(f\omega) = fd\omega + df \wedge \omega = fc\theta^1 \wedge \theta^2 + \frac{f_1}{f}\theta^1 \wedge \theta^3 + \frac{f_2}{f}\theta^2 \wedge \theta^3.$$

Using this equation, it follows that

$$(17) \quad \begin{aligned} d(fc\theta^3) &= f(dc \wedge \theta^3) + c(df \wedge \theta^3) + fc d\theta^3 \\ &= f\alpha\theta^1 \wedge \theta^3 + f\beta\theta^2 \wedge \theta^3 + cf_1\theta^1 \wedge \theta^3 \\ &\quad + cf_2\theta^2 \wedge \theta^3 + f^2c^2\theta^1 \wedge \theta^2 + cf_1\theta^1 \wedge \theta^3 + cf_2\theta^2 \wedge \theta^3. \end{aligned}$$

If K_g denotes the Gauss curvature of surface (M, g) , then $d\eta_2^1 = K(\eta^1 \wedge \eta^2)$ and hence $d\theta_2^1 = \pi^* K_g(\theta^1 \wedge \theta^2)$. Putting $\tilde{K} = \pi^* K_g$ to save on notation and using the equations (16), (17) we get

$$(18) \quad d\theta_2^1 = (\tilde{K} - \frac{1}{2}f^2c^2)\theta^1 \wedge \theta^2 - (\frac{1}{2}f\alpha + cf_1)\theta^1 \wedge \theta^3 - (\frac{1}{2}f\beta + cf_2)\theta^2 \wedge \theta^3.$$

We have also

$$(19) \quad \begin{aligned} \theta_3^1 \wedge \theta_2^3 &= (-\frac{1}{2}fc\theta^2 - \frac{f_1}{f}\theta^3) \wedge (-\frac{1}{2}fc\theta^1 + \frac{f_2}{f}\theta^3) \\ &= -\frac{1}{4}f^2c^2\theta^1 \wedge \theta^2 - \frac{1}{2}cf_2\theta^2 \wedge \theta^3 - \frac{1}{2}cf_1\theta^1 \wedge \theta^3. \end{aligned}$$

Therefore, (18) and (19) imply

$$\Omega_2^1 = d\theta_2^1 + \theta_3^1 \wedge \theta_2^3 = (\tilde{K} - \frac{3}{4}f^2c^2)\theta^1 \wedge \theta^2 - (\frac{1}{2}f\alpha + \frac{3}{2}cf_1)\theta^1 \wedge \theta^3 - (\frac{1}{2}f\beta + \frac{3}{2}cf_2)\theta^2 \wedge \theta^3.$$

To determine Ω_3^1 , we need to calculate $d\theta_3^1 = -\frac{1}{2}d(fc\theta^2) - d(f_1\omega)$. But

$$(20) \quad d(fc\theta^2) = fdc\wedge\theta^2 + cdf\wedge\theta^2 + fc d\theta^2 = f\alpha\theta^1\wedge\theta^2 + cf_1\theta^1\wedge\theta^2 + fc(-\pi^*\eta_1^2\wedge\theta^1).$$

and

$$(21) \quad \begin{aligned} d(f_1\omega) &= f_1d\omega + df_1\wedge\omega = cf_1\theta^1\wedge\theta^2 + (f_{11}\theta^1 + f_{12}\theta^2)\wedge\frac{1}{f}\theta^3 \\ &= cf_1\theta^1\wedge\theta^2 + \frac{f_{11}}{f}\theta^1\wedge\theta^3 + \frac{f_{12}}{f}\theta^2\wedge\theta^3. \end{aligned}$$

Therefore, (20) and (21) give

$$(22) \quad d\theta_3^1 = -(\frac{1}{2}f\alpha + \frac{3}{2}cf_1)\theta^1\wedge\theta^2 - \frac{f_{11}}{f}\theta^1\wedge\theta^3 - \frac{f_{12}}{f}\theta^2\wedge\theta^3 - \frac{1}{2}fc(\pi^*\eta_2^1)\wedge\theta^1.$$

We have also

$$(23) \quad \begin{aligned} \theta_2^1 \wedge \theta_3^2 &= (\pi^*\eta_2^1 - \frac{1}{2}fc\theta^3) \wedge (\frac{1}{2}fc\theta^1 - \frac{f_2}{f}\theta^3) \\ &= \frac{1}{2}fc(\pi^*\eta_2^1)\wedge\theta^1 - \frac{f_2}{f}(\pi^*\eta_2^1)\wedge\theta^3 + \frac{1}{4}f^2c^2\theta^1\wedge\theta^3. \end{aligned}$$

Using the decompositions $\pi^*\eta_2^1 = \eta_{2,1}^1\theta^1 + \eta_{2,2}^1\theta^2$, and the equations (22) and (23) we obtain

$$\Omega_3^1 = -(\frac{1}{2}f\alpha + \frac{3}{2}cf_1)\theta^1\wedge\theta^2 + (\frac{1}{4}f^2c^2 - \frac{f_{11}}{f} - \frac{f_2}{f}\eta_{2,1}^1)\theta^1\wedge\theta^3 - (\frac{f_2}{f}\eta_{2,2}^1 + \frac{f_{12}}{f})\theta^2\wedge\theta^3.$$

In order to determine Ω_3^2 , we need first to calculate

$$(24) \quad d\theta_3^2 = \frac{1}{2}d(fc\theta^1) - d(f_2\omega)$$

But, we have

$$(25) \quad \begin{aligned} d(fc\theta^1) &= fc d\theta^1 + fdc\wedge\theta^1 + cdf\wedge\theta^1 \\ &= -fc(\pi^*\eta_2^1)\wedge\theta^2 - f\beta\theta^1\wedge\theta^2 - cf_2\theta^1\wedge\theta^2. \end{aligned}$$

and

$$(26) \quad \begin{aligned} d(f_2\omega) &= f_2d\omega + df_2\wedge\omega = cf_2\theta^1\wedge\theta^2 + (f_{21}\theta^1 + f_{22}\theta^2)\wedge\frac{1}{f}\theta^3 \\ &= cf_2\theta^1\wedge\theta^2 + \frac{f_{21}}{f}\theta^1\wedge\theta^3 + \frac{f_{22}}{f}\theta^2\wedge\theta^3. \end{aligned}$$

Secondly, we need the following calculation

$$\begin{aligned}
 \theta_1^2 \wedge \theta_3^1 &= \left(-\pi^* \eta_2^1 + \frac{1}{2} f c \theta^3 \right) \wedge \left(-\frac{1}{2} f c \theta^2 - \frac{f_1}{f} \theta^3 \right) \\
 &= \frac{1}{2} f c (\pi^* \eta_2^1) \wedge \theta^2 + \frac{f_1}{f} (\pi^* \eta_2^1) \wedge \theta^3 + \frac{1}{4} f^2 c^2 \theta^2 \wedge \theta^3.
 \end{aligned}
 \tag{27}$$

Finally, the equations (24), (25), (26) and (27) imply that

$$\begin{aligned}
 \Omega_3^2 &= d\theta_3^2 + \theta_1^2 \wedge \theta_3^1 = -\left(\frac{1}{2} f \beta + \frac{3}{2} c f_2\right) \theta^1 \wedge \theta^2 + \left(\frac{f_1}{f} \eta_{2,1}^1 - \frac{f_{21}}{f}\right) \theta^1 \wedge \theta^3 \\
 &\quad + \left(\frac{f_1}{f} \eta_{2,2}^1 - \frac{f_{22}}{f} + \frac{1}{4} f^2 c^2\right) \theta^2 \wedge \theta^3.
 \end{aligned}$$

Using these notations, we have proved the next proposition.

Proposition 1.3. *Let $\pi : P \rightarrow M$ be a principal \mathbb{S}^1 -bundle over a Riemann surface (M, g) endowed with a connection A . Let $\theta = (\theta_j^i)$ be the connection 1-form of the Levi-Civita connection of g_A . Using the same notation as above, the components of curvature 2-form $\Omega^\theta = (\Omega_j^i)$ corresponding to θ is given by*

$$\begin{aligned}
 (1) \quad \Omega_2^1 &= (\tilde{K} - \frac{3}{4} f^2 c^2) \theta^1 \wedge \theta^2 + (-\frac{1}{2} f \alpha - \frac{3}{2} c f_1) \theta^1 \wedge \theta^3 + (-\frac{1}{2} f \beta - \frac{3}{2} c f_2) \theta^2 \wedge \theta^3 \\
 (2) \quad \Omega_3^1 &= (-\frac{1}{2} f \alpha - \frac{3}{2} c f_1) \theta^1 \wedge \theta^2 + (\frac{1}{4} f^2 c^2 - \frac{f_{11}}{f} - \frac{f_2}{f} \eta_{2,1}^1) \theta^1 \wedge \theta^3 - (\frac{f_2}{f} \eta_{2,2}^1 + \frac{f_{12}}{f}) \theta^2 \wedge \theta^3 \\
 (3) \quad \Omega_3^2 &= (-\frac{1}{2} f \beta - \frac{3}{2} c f_2) \theta^1 \wedge \theta^2 + (\frac{f_1}{f} \eta_{2,1}^1 - \frac{f_{21}}{f}) \theta^1 \wedge \theta^3 + (\frac{f_1}{f} \eta_{2,2}^1 - \frac{f_{22}}{f} + \frac{1}{4} f^2 c^2) \theta^2 \wedge \theta^3
 \end{aligned}$$

1.4. The Ricci curvature of metric connection on principal \mathbb{S}^1 -bundles over Riemann surfaces. Let $(g, P \xrightarrow{\pi} M, A)$ be a \mathbb{S}^1 -triple over M and, let g_A be the connection metric on P . Let (e_1, e_2, e_3) be the g_A -orthonormal frame on P_U chosen as previous. For an arbitrary point $y \in P_U$ and two vectors $v, w \in T_y P$ the Ricci curvature tensor is given by $\text{Ric}(v, w) = \sum_{m=1}^3 g_A(R^\theta(v, e_m)e_m, w)$, where $R^\theta \in A^2(P, \text{End}(T_P))$ is the Riemann curvature tensor associated to Levi-Civita connection ∇^θ on tangent bundle T_P . The components of Ricci curvature tensor in g_A -orthonormal frame (e_1, e_2, e_3) is given by

$$R_{lk} = \text{Ric}(e_l, e_k) = \sum_m g_A(R^\theta(e_l, e_m)e_m, e_k).$$

The Riemann curvature tensor R^θ is related to curvature 2-form $\Omega^\theta = (\Omega_j^i)$ by

$$R^\theta(e_l, e_m)e_m = \sum_i \Omega_m^i(e_l, e_m)e_i.$$

Since (e_1, e_2, e_3) is g_A -orthonormal, we have

$$(28) \quad R_{lk} = \sum_{m,i} g_A(\Omega_m^i(e_l, e_m)e_i, e_k) = \sum_m \Omega_m^k(e_l, e_m).$$

Using Proposition 1.2 and Proposition 1.3 and (28) the components of Ricci tensor of g_A can be computed. If in (28) we substitute $l = 1$, then $R_{1k} = \Omega_2^k(e_1, e_2) + \Omega_3^k(e_1, e_3)$, calculating this for $k = 1, 2, 3$ and using Proposition 1.3 we obtain

$$R_{11} = \tilde{K} - \frac{1}{2} f^2 c^2 - \frac{f_{11}}{f} - \frac{f_2}{f} \eta_{2,1}^1, \quad R_{12} = \frac{f_1}{f} \eta_{2,1}^1 - \frac{f_{21}}{f}, \quad R_{13} = \frac{1}{2} f \beta + \frac{3}{2} c f_2.$$

If we substitute $l = 2$ and $l = 3$ in (28), then in a similar way we obtain

$$R_{21} = -\frac{f_2}{f} \eta_{2,2}^1 - \frac{f_{12}}{f}, \quad R_{22} = \tilde{K} - \frac{1}{2} f^2 c^2 + \frac{f_1}{f} \eta_{2,2}^1 - \frac{f_{22}}{f}, \quad R_{23} = -\frac{1}{2} f \alpha - \frac{3}{2} c f_1.$$

and

$$R_{31} = \frac{1}{2}f\beta + \frac{3}{2}cf_2, \quad R_{32} = -\frac{1}{2}f\alpha - \frac{3}{2}cf_1, \quad R_{33} = \frac{1}{2}f^2c^2 - \frac{f_{11}}{f} - \frac{f_2}{f}\eta_{2,1}^1 + \frac{f_1}{f}\eta_{2,2}^1 - \frac{f_{22}}{f}.$$

With the same notation as previous, we have $\Delta f = -f_{11} - f_{22} - f_2\eta_{2,1}^1 + f_1\eta_{2,2}^1$, where $\Delta f = -*d*df$ is the laplacian of function $f : P \rightarrow \mathbb{R}$. Using $df = f_1\theta^1 + f_2\theta^2$ we obtain

$$\Delta f = -*d*(f_1\theta^1 + f_2\theta^2) = -f_{11} - f_{22} - f_2\eta_{2,1}^1 + f_1\eta_{2,2}^1.$$

So, we can write $R_{33} = \frac{1}{2}f^2c^2 + \frac{\Delta f}{f}$. The Ricci tensor of metric g_A with respect the frame (e_1, e_2, e_3) has the following representation

$$\text{Ric}(g_A) = \begin{pmatrix} \mathcal{R} & \mathcal{S} \\ \mathcal{S}^t & \varphi \end{pmatrix}.$$

where

$$\mathcal{R} = \begin{pmatrix} \tilde{K} - \frac{1}{2}f^2c^2 & 0 \\ 0 & \tilde{K} - \frac{1}{2}f^2c^2 \end{pmatrix} - \frac{1}{f} \begin{pmatrix} f_{11} + f_2\eta_{2,1}^1 & f_{21} - f_1\eta_{2,1}^1 \\ f_{12} + f_2\eta_{2,2}^1 & f_{22} - f_1\eta_{2,2}^1 \end{pmatrix},$$

$$\mathcal{S} = \begin{pmatrix} \frac{1}{2}f\beta + \frac{3}{2}cf_2 \\ -\frac{1}{2}f\alpha - \frac{3}{2}cf_1 \end{pmatrix} \quad \text{and} \quad \varphi = \frac{1}{2}f^2c^2 + \frac{\Delta f}{f}.$$

Remark 2. The equation $d^2f = 0$ implies $f_{21} - f_1\eta_{2,1}^1 = f_{12} + f_2\eta_{2,2}^1$ which is equivalent to $R_{12} = R_{21}$.

Remark 3. Let ∇ denote the Linear connection on tangent bundle T_P induced by the Levi-Civita connection θ . Since, $df = f_1\theta^1 + f_2\theta^2$ one has the next formula

$$\begin{aligned} \nabla df &= \nabla(f_1\theta^1 + f_2\theta^2) = df_1 \otimes \theta^1 + f_1\nabla\theta^1 + df_2 \otimes \theta^2 + f_2\nabla\theta^2 = f_{11}\theta^1 \otimes \theta^1 \\ &+ f_{12}\theta^2 \otimes \theta^1 + f_1(-\pi^*\eta_2^1 \otimes \theta^2) + f_{21}\theta^1 \otimes \theta^2 + f_{22}\theta^2 \otimes \theta^2 + f_2(\pi^*\eta_2^1 \otimes \theta^1). \end{aligned}$$

Thus, the matrix representation of ∇df with respect to $(\theta^1, \theta^2, \theta^3)$ is given by

$$\nabla df = \begin{pmatrix} f_{11} + f_2\eta_{2,1}^1 & f_{21} - f_1\eta_{2,1}^1 \\ f_{12} + f_2\eta_{2,2}^1 & f_{22} - f_1\eta_{2,2}^1 \end{pmatrix}.$$

Using these remarks and the fact that $|F_\omega|_g^2 = c^2$ we can rewrite \mathcal{R} in the following way:

$$\mathcal{R} = (\tilde{K} - \frac{1}{2}f^2|F_\omega|_g^2)\pi^*g - \frac{1}{f}\nabla df = \pi^*\text{Ric}(g) - \frac{1}{2}f^2|F_\omega|_g^2\pi^*g - \frac{1}{f}\nabla df.$$

We can also rewrite \mathcal{S} in the invariant way as

$$\mathcal{S} = \frac{1}{2}(f\beta + 3cf_2)\theta^1 \otimes \theta^3 - \frac{1}{2}(f\alpha + 3cf_1)\theta^2 \otimes \theta^3 = -\frac{1}{2}f^2(d^*F_\omega) \otimes \omega - \frac{3}{2f}(*F_\omega)(*df) \otimes \omega.$$

Since these formula are written in an invariant way they are independent of chosen frame (e_1, e_2, e_3) and the next proposition is proved.

Proposition 1.4. *Let $(g, P \xrightarrow{\pi} M, A)$ be a \mathbb{S}^1 -triple on M . If ω and F_ω denote, respectively, the connection 1-form and the curvature 2-form associated with A , then the matrix representation of Ricci curvature tensor of g_A with respect to orthonormal frame $(\theta^1, \theta^2, \theta^3)$ is given by*

$$\text{Ric}(g_A) = \begin{pmatrix} \mathcal{R} & \mathcal{S} \\ \mathcal{S}^t & \varphi \end{pmatrix}.$$

where, $\mathcal{R} = (\pi^* K_g - \frac{1}{2} f^2 |F_\omega|_g^2) \pi^* g - \frac{1}{f} \nabla df$, $\mathcal{S} = -\frac{1}{2} f^2 (d^* F_\omega) \otimes \omega - \frac{3}{2f} (*F_\omega)(*df) \otimes \omega$ and $\varphi = \frac{1}{2} f^2 |F_\omega|_g^2 + \frac{\Delta f}{f}$.

Corollary 1.5. *Suppose that $f : M \rightarrow \mathbb{R}$ is a constant function on M and A is a Yang-Mills connection. Then*

- (1) *If $R_g = 2K_g$ denotes the scalar curvature of g , then the scalar curvature of g_A is given by*

$$R_{g_A} = \pi^* R_g - \frac{1}{2} f^2 |F_\omega|_g^2.$$

- (2) *The horizontal and vertical subspaces are orthogonal relative to the Ricci tensor of metric g_A and we have*

$$\text{Ric}(g_A) = (\pi^* K_g - \frac{1}{2} f^2 |F_\omega|_g^2) \pi^* g + \frac{1}{2} f^4 |F_\omega|_g^2 \omega \otimes \omega.$$

2. THE RICCI FLOW EQUATION OF METRIC CONNECTION ON PRINCIPAL \mathbb{S}^1 -BUNDLES OVER COMPACT RIEMANN SURFACES

2.1. Ricci flow equation on principal \mathbb{S}^1 -bundles over compact Riemann surfaces. Let (M, g) be a connected, compact, oriented Riemann surface and, let $(g, P \xrightarrow{\pi} M, A)$ be a \mathbb{S}^1 -triple over M . The connection metric on P is defined by $g_A = \pi^* g + f^2 \omega \otimes \omega$, where $\omega \in A^1(P, \mathbb{R})$ and $f \in C^\infty(P)$ is the smooth \mathbb{S}^1 -invariant map defined as in Section 1.2. The variation of connection metric $g_A(t)$ with time is as follow

$$\dot{g}_A(t) = \pi^* \dot{g}_t + f_t^2 \dot{\omega}_t \otimes \omega_t + f_t^2 \omega_t \otimes \dot{\omega}_t + 2f_t \dot{f}_t \omega_t \otimes \omega_t.$$

The Ricci flow equation of metric g_A is the following evolution equation

$$\dot{g}_A(t) = -2\text{Ric}(g_A(t)).$$

Using Proposition 1.4 the Ricci flow equation of g_A is equivalent to the following system of differential equations:

$$(29) \quad \begin{aligned} \dot{g}_t &= -2\text{Ric}(g_t) + f_t^2 |F_{\omega_t}|_{g_t}^2 g_t + \frac{2}{f_t} \nabla df_t. \\ \dot{\omega}_t &= d^* F_{\omega_t} + 3f_t^{-3} (*F_{\omega_t})(*df_t). \\ \dot{f}_t &= -\frac{1}{2} f_t^3 |F_{\omega_t}|_{g_t}^2 - \Delta f_t. \end{aligned}$$

2.2. Ricci flow equation of locally homogeneous \mathbb{S}^1 -triples on compact Riemann surfaces with constant curvature. Let (M, g_0) be a connected, oriented, compact Riemann surface with constant Gauss curvature K_0 , and let $\pi : P \rightarrow M$ be a principal \mathbb{S}^1 -bundle endowed with a Yang-Mills connection A_0 . Suppose that $\omega_0 \in A^1(P, \mathbb{R})$ (resp. $F_{\omega_0} = d\omega_0 \in A^2(P, \mathbb{R})$), denotes the connection 1-form (resp. the curvature 2-form) associated with connection A_0 . Let $g_t = \lambda_t g_0$ be a one-parameter family of metric on M . If f_t denotes the size of the fiber at time $t \geq 0$ and $\omega_t = \omega_0$, then we search a solution of equations (29) with the initial conditions $\lambda(0) = f(0) = 1$. Moreover, we suppose that $f_t : M \rightarrow \mathbb{R}$ is constant with respect to M and hence $df_t = \Delta f_t = 0$. Since A_0 is Yang-Mills and $\omega_t = \omega_0$ we have

$$0 = \dot{\omega}_t = d^* F_{\omega_t} = d^* F_{\omega_0} = 0.$$

This means that the second equation of Ricci flow equations (29) is automatically verified and they are equivalent to

$$(30) \quad \begin{aligned} \dot{g}_t &= -2\text{Ric}(g_t) + f_t^2 |F_{\omega_0}|_{g_t}^2 g_t \cdot \\ \dot{f}_t &= -\frac{1}{2} f_t^3 |F_{\omega_0}|_{g_t}^2 \cdot \end{aligned}$$

To save on notation denote the norm square of curvature 2-form $F_{\omega_0} \in A^2(M, \mathbb{R})$ by $|F_{\omega_0}|_{g_0}^2 =: F_0$ and note that

$$(31) \quad |F_{\omega_0}|_{g_t}^2 = g_t(F_{\omega_0}, F_{\omega_0}) = \lambda_t^{-2} g_0(F_{\omega_0}, F_{\omega_0}) = \lambda_t^{-2} |F_{\omega_0}|_{g_0}^2 = \lambda_t^{-2} F_0 \cdot$$

The Ricci tensor is scale invariant, i.e. $\text{Ric}(g_t) = \text{Ric}(\lambda(t)g_0) = \text{Ric}(g_0) = K_0 g_0$. Hence, the equations (30) take the form

$$(32) \quad \begin{aligned} \dot{\lambda}_t &= -2K_0 + F_0 f_t^2 \lambda_t^{-1}, \\ \dot{f}_t &= -\frac{1}{2} F_0 f_t^3 \lambda_t^{-2}. \end{aligned}$$

Lemma 2.1. *Let $\pi : P \rightarrow M$ be a \mathbb{S}^1 -bundle on a compact Riemann surface (M, g) and A be a Yang-Mills connection on P . The curvature 2-form $F_A \in A^2(M, \mathbb{R})$ is a constant multiple of the volume form of (M, g) . More precisely, there exists a constant $c \in \mathbb{R}$ such that*

$$F_A = c \text{Vol}_g \cdot$$

Proof: Since the curvature F_A is a 2-form on M , there exists a smooth function $f_A \in C^\infty(M)$ such that $F_A = f_A \text{Vol}_g$. Since A is Yang-Mills $d^* F_A = 0$, we know also that $dF_A = 0$. Therefore

$$0 = (dd^* + d^*d) F_A = (dd^* + d^*d) f_A \text{Vol}_g = (\Delta f_A) \text{Vol}_g$$

This means $\Delta f_A = 0$ on M and as M is compact, there exists a constant $c \in \mathbb{R}$ such that $f_A \equiv c$. This constant can be determined using $c \int_M \text{Vol}_g = 2\pi c_1(P)$.

Example. Suppose (M, g_0) is a Riemann surface of constant positive curvature $K_0 > 0$ and A_0 is a non flat Yang-Mills connection on \mathbb{S}^1 -bundle $P \rightarrow M$ such that $F_0 = K_0$. Note that there exists always such a connection A_0 , in fact using Lemma 2.1 we can scale an arbitrary non flat Yang-Mills connection in an appropriately way to obtain a Yang-Mills connection A_0 for which $F_0 = K_0$. The initial metric $g_{A_0} = \pi^* g + \omega_0 \otimes \omega_0$ is an Einstein metric of positive scalar curvature. Putting $\varphi_t := f_t^2$, the Ricci flow equation is equivalent to

$$(33) \quad \begin{aligned} \dot{\lambda}_t &= -2K_0 + K_0 \varphi_t \lambda_t^{-1} \\ \dot{\varphi}_t &= -K_0 \varphi_t^2 \lambda_t^{-2} \end{aligned}$$

Supposing that $\varphi_t \lambda_t^{-1} = 1$ and integrating (33) we obtain $\varphi_t = \lambda_t = 1 - K_0 t$. Hence, the Ricci flow solution exists for all finite time and

$$g_A(t) = (1 - K_0 t) g_{A_0} \quad \text{for all } 0 \leq t < 1/K_0.$$

Therefore, the metric shrinks to a point as t approaches to $1/K_0$, while the curvatures becomes infinite. For instance, let $\pi : \mathbb{S}^3 \rightarrow \mathbb{S}^2$ be the famous \mathbb{S}^1 -bundle of Hopf fibration endowed with a Yang-Mills connection A_0 . Suppose $g_{\mathbb{S}^2}$ is the standard metric on 2-sphere induced by the embedding $\mathbb{S}^2 \subset \mathbb{R}^3$ and put $g_0 := \frac{1}{4} g_{\mathbb{S}^2}$. Using Lemma 2.1 the curvature F_{A_0} of Yang-Mills connection A_0 satisfies $F_{A_0} = 2 \text{Vol}_{g_0}$. The metric $g_{A_0} := \pi^* g_0 + \omega_0 \otimes \omega_0$ induced by the connection A_0 and metric g_0 coincides with the standard metric $g_{\mathbb{S}^3}$ on 3-sphere \mathbb{S}^3 . Since $F_0 = K_0 = 4$, the

Ricci-flow solution is given by $g_A(t) = (1 - 4t)g_{\mathbb{S}^3}$. The sphere \mathbb{S}^3 is collapsing to a point and the curvatures becomes infinite as $t \rightarrow 1/4$.

2.3. Explicit solutions of normalized Ricci flow equation for locally homogeneous \mathbb{S}^1 -triples on compact Riemann surfaces. Let (M, g) be a connected, oriented, compact Riemann surface endowed with a metric of constant curvature. An \mathbb{S}^1 -triple $(g, P \xrightarrow{\pi} M, A)$ is locally homogeneous if and only if the connection A is Yang-Mills [2]. Suppose that $(g, P \xrightarrow{\pi} M, A)$ is a locally homogeneous \mathbb{S}^1 -triple over M , then one can endow the total space P with a locally homogeneous metric g_A . The total space P is a compact manifold of dimension three and the normalized Ricci-flow equation of metric g_A on P is the next evolution equation [3]

$$(34) \quad \frac{\partial}{\partial t} g_A(t) = -2\text{Ric } g_A(t) + \frac{2}{3}r(t)g_A(t).$$

where $r(t) := \int_P R(t)d\mu_t / \int_P d\mu_t$ is the average of scalar curvature $R(t)$ of $g_A(t)$ and μ_t denotes its volume form. The factor $r(t)$ is used to normalize the equation so that the volume is constant. Let $\omega_0 \in A^1(P, \mathbb{R})$ (resp. $F_{\omega_0} = d\omega_0 \in A^2(P, \mathbb{R})$), be the connection 1-form (resp. the curvature 2-form) associated with connection A_0 . Consider the one-parameter family of metrics $g_t = \lambda_t g_0$ on M and, let f_t be the size of the fiber at time t . Let $\omega_t = \omega_0$ be the initial Yang-Mills connection, for all $t \geq 0$. We search a solution of the normalized Ricci-flow equation (34) with the initial conditions $\lambda(0) = f(0) = 1$. Since A is Yang-Mills, Corollary 1.5 implies that $R(t) = 2K_{g_t} - \frac{1}{2}f_t^2|F_{\omega_t}|_{g_t}^2 = 2K_0\lambda_t^{-1} - \frac{1}{2}F_0f_t^2\lambda_t^{-2}$. Therefore, the scalar curvature $R(t)$ is constant on P and we have $r(t) = R(t)$. The normalized Ricci flow equation (34) take the form

$$\begin{aligned} \dot{g}_t &= -2\text{Ric}(g_t) + f_t^2|F_{\omega_t}|_{g_t}^2 g_t + \frac{2}{3}(2K_{g_t} - \frac{1}{2}f_t^2|F_{\omega_t}|_{g_t}^2)g_t, \\ 2\dot{f}_t &= -2(\frac{1}{2}f_t^2|F_{\omega_t}|_{g_t}^2) + \frac{2}{3}(2K_{g_t} - \frac{1}{2}f_t^2|F_{\omega_t}|_{g_t}^2). \end{aligned}$$

Or, equivalently

$$(35) \quad \begin{aligned} \dot{g}_t &= -\frac{2}{3}\text{Ric}(g_t) + \frac{2}{3}f_t^2|F_{\omega_t}|_{g_t}^2 g_t, \\ \dot{f}_t &= \frac{2}{3}K_{g_t}f_t - \frac{2}{3}f_t^3|F_{\omega_t}|_{g_t}^2. \end{aligned}$$

Since $g_t = \lambda_t g_0$ we deduce $K_{g_t} = \lambda_t^{-1}K_0$, $\text{Ric}(g_t) = \text{Ric}(g_0) = K_0 g_0$. By (31) we have $|F_{\omega_t}|_{g_t}^2 = \lambda_t^{-2}F_0$. Make use of these formulae in (35) leads to the following system of ordinary differential equations

$$(36) \quad \begin{aligned} \dot{\lambda}_t &= -\frac{2}{3}K_0 + \frac{2}{3}F_0f_t^2\lambda_t^{-1}, \\ \dot{f}_t &= \frac{2}{3}K_0f_t\lambda_t^{-1} - \frac{2}{3}F_0f_t^3\lambda_t^{-2}. \end{aligned}$$

In the remaining part of the paper, we give an explicit solutions of these differential equations and, we describe some asymptotical behaviors of their solutions. A simple observation is that

$$\dot{\lambda}_t f_t + \lambda_t \dot{f}_t = -\frac{2}{3}K_0 f_t + \frac{2}{3}F_0 f_t^3 \lambda_t^{-1} + \frac{2}{3}K_0 f_t - \frac{2}{3}F_0 f_t^3 \lambda_t^{-1} = 0.$$

The initial conditions $f(0) = \lambda(0) = 1$ imply that $f(t)\lambda(t) = 1$, for all $t \geq 0$. So, to solve the equations (36) it is enough to solve the system below

$$(37) \quad \dot{\lambda}_t = -\frac{2}{3}K_0 + \frac{2}{3}F_0\lambda_t^{-3}, \quad f_t = \lambda_t^{-1}.$$

Thanks to Corollary 1.5 the scalar curvature of $g_{A_0}(t)$ can be computed by

$$(38) \quad R_t = 2K_{g_t} - \frac{1}{2}f_t^2|F_{\omega_t}|_{g_t}^2 = 2K_0\lambda_t^{-1} - \frac{1}{2}F_0\lambda_t^{-4}.$$

To study the equations (37), we will consider two situations. Firstly, we suppose that $K_0 \neq 0$ and we set $a_0 := \sqrt[3]{\frac{F_0}{K_0}}$. The equations (37) is equivalent to

$$\dot{\lambda}_t = -\frac{2}{3}K_0 + \frac{2}{3}F_0\lambda_t^{-3} = -\frac{2}{3}K_0(1 - \frac{F_0}{K_0}\lambda_t^{-3}).$$

Solving this equation leads to the following explicit solution

$$a_0 \ln \frac{\sqrt[3]{|\lambda_t - a_0|}}{\sqrt[6]{\lambda_t^2 + a_0\lambda_t + a_0^2}} - \frac{a_0}{\sqrt{3}} \tan^{-1}\left(\frac{2\lambda_t + a_0}{\sqrt{3}a_0}\right) + \lambda_t = -\frac{2}{3}K_0t + c_0.$$

Here, $c_0 \in \mathbb{R}$ is a real constant which can be determined by the initial condition $\lambda(0) = 1$. The rest of this work is dedicated to describe the asymptotical behavior of (normalized) Ricci flow in some interesting cases:

Spherical geometry: if $K_0 > 0$ and A_0 is non flat, then $\lambda_t \rightarrow a_0$ and $f_t \rightarrow a_0^{-1}$ as t goes to infinity and the Ricci flow converge to some Einstein metric. The scalar curvature of metric $g_{A_0}(t)$ is computed due to equations (38) and $R_t \rightarrow \frac{3}{2}K_0a_0^{-1}$ as t goes to infinity. In this case, the circle bundle $\pi : P \rightarrow M$ has a geometric structure with model geometry \mathbf{S}^3 .

Geometry of the universal cover of $\mathrm{SL}(2, \mathbb{R})$: if $K_0 < 0$ and A_0 is non flat, then from the first equation of (36) we have $\dot{\lambda}_t = -\frac{2}{3}K_0 + \frac{2}{3}F_0f_t^2\lambda_t^{-1}$ which implies that $\dot{\lambda}_t \geq \frac{2}{3}F_0f_t^2\lambda_t^{-1}$. Use $f_t\lambda_t = 1$ and integrate this to obtain $\lambda_t^4 \geq \frac{2}{3}F_0t + C_0$. Therefore, $\lambda_t \rightarrow \infty$ and $f_t \rightarrow 0$ as t goes to infinity. This means that the metric on the base expand without bound while the fiber of the bundle shrinks to zero. Using the equation (38), the scalar curvature falls off to zero as t goes to infinity. In this case, the circle bundle $\pi : P \rightarrow M$ has a geometric structure with model geometry $\widetilde{\mathrm{SL}}_2(\mathbb{R})$.

Euclidean geometry: If $K_0 = 0$ and A_0 is flat ($F_{A_0} = 0$), then for all $t \geq 0$ we have $\lambda_t = f_t = 1$. Hence, the Ricci flow solution is trivial, $g_A(t) = g_A(0)$ for all $t \geq 0$. In this case, the circle bundle $\pi : P \rightarrow M$ has a geometric structure with model geometry \mathbf{E}^3 .

Now, suppose that $K_0 \neq 0$ and A_0 is flat, then the Ricci flow equations are equivalent to

$$(39) \quad \dot{\lambda}_t = -\frac{2}{3}K_0, \quad \dot{f}_t = \frac{2}{3}K_0f_t\lambda_t^{-1}$$

Integrating this, it follows that $\lambda_t = 1 - \frac{2}{3}K_0t$ and $f_t = (1 - \frac{2}{3}K_0t)^{-1}$. Therefore, the solution of Ricci flow equation is given by

$$g_A(t) = (1 - \frac{2}{3}K_0t)\pi^*g_0 + (1 - \frac{2}{3}K_0t)^{-2}\omega_0 \otimes \omega_0.$$

The scalar curvature of this metric is given by $R_t = 6K_0(3 - 2K_0t)^{-1}$. Now, we can consider two cases:

Geometry of $\mathbf{S}^2 \times \mathbf{R}$: if $K_0 > 0$ (and A_0 is flat), then at time $t = \frac{3}{2K_0}$ the metric vanishes and a curvature singularity appear. Also, we have $f_t \rightarrow \infty$, which means that the size of the fiber expands. In this case, the circle bundle P has a

geometric structure with model geometry $\mathbf{S}^2 \times \mathbf{R}$. This is the only class of locally homogeneous geometries whose Ricci flows do achieve curvature singularities [3].

Geometry of $\mathbf{H}^2 \times \mathbf{R}$: if $K_0 < 0$ (and A_0 is flat), then the Ricci flow solution exists for any time $t \geq 0$. If t goes to infinity, then $\lambda_t \rightarrow \infty$ and $f_t \rightarrow 0$, which means that g_t is nonsingular and the fiber collapses. The scalar curvature falls off to zero when t goes to infinity. In this case, the circle bundle $\pi : P \rightarrow M$ has a geometric structure with model geometry $\mathbf{H}^2 \times \mathbf{R}$.

Geometry of the Heisenberg group: If $K_0 = 0$ and A_0 is non flat ($F_{A_0} \neq 0$), then the Ricci flow equations take the form

$$\dot{\lambda}_t = \frac{2}{3}F_0\lambda_t^{-3}, \quad f_t = \lambda_t^{-1}.$$

The solutions are given by $\lambda(t) = (1 + \frac{8}{3}F_0t)^{1/4}$ and $f(t) = (1 + \frac{8}{3}F_0t)^{-1/4}$. Hence, the Ricci flow solution is given by

$$g_A(t) = (1 + \frac{8}{3}F_0t)^{1/4}\pi^*g_0 + (1 + \frac{8}{3}F_0t)^{-1/2}\omega_0 \otimes \omega_0.$$

If t goes to infinity, then $\lambda_t \rightarrow \infty$ and $f_t \rightarrow 0$, which means that the metric g_t on the base manifold M expands and the fiber collapses. Using remark 1.5 the scalar curvature of $g_A(t)$ is given by

$$R_t = -\frac{1}{2}f_t^2|F_{\omega_t}|_{g_t}^2 = -\frac{1}{2}F_0f_t^2\lambda_t^{-2} = -\frac{1}{2}F_0f_t^4 = -\frac{1}{2}F_0(1 + \frac{8}{3}F_0t)^{-1}.$$

Hence, the scalar curvature falls off to zero as t goes to infinity. In this case, the circle bundle $\pi : P \rightarrow M$ over surface M has a geometric structure with model geometry \mathbf{Nil} .

REFERENCES

- [1] A. Bazdar, Locally homogeneous triples: Extension theorems for parallel sections and parallel bundle isomorphisms. *Mediterr. J. Math.* (2017).
- [2] A. Bazdar, Geometric principal bundles over geometric Riemannian manifolds., Ph.D. thesis, Aix-Marseille University. (2017).
- [3] J. Isenberg, Ricci flow of locally homogeneous geometries on closed manifolds. *J. Differ. Geom.* 35 (3), (1992) 723-741.
- [4] S. Kobayashi, K. Nomizu, *Foundations of Differential Geometry I*, Interscience Publ., New York, Vol. I (1963).
- [5] S. Morita, *Geometry of Differential Forms*. Translations of Mathematical Monographs, vol. 201. AMS, Providence, (2001).
- [6] I.M. Singer, Infinitesimally homogeneous spaces, *Comm. Pure Appl. Math.*,13, (1960) 685-697.
- [7] P. Scott, The geometries of 3-manifolds, *Bull. London Math. Soc.* 15, no. 5, (1983) 401-487.
- [8] W.P. Thurston, Three-dimensional manifolds, Kleinian groups and hyperbolic geometry, *AMS, Bulletin. New Series*, vol. 6 (1982) 357-381.

AIX MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M, UMR 7373, 13453 MARSEILLE, FRANCE

E-mail address: arash.bazdar@univ-amu.fr