

REMARKS ON HYPERGEOMETRIC CAUCHY NUMBERS

MIHO AOKI AND TAKAO KOMATSU

ABSTRACT. For a positive integer N , hypergeometric Cauchy numbers $c_{N,n}$ are defined by

$$\frac{1}{{}_2F_1(1, N; N+1; -x)} = \frac{(-1)^{N-1} x^N / N}{\log(1+x) - \sum_{n=1}^{N-1} (-1)^{n-1} x^n / n} = \sum_{n=0}^{\infty} c_{N,n} \frac{x^n}{n!},$$

where ${}_2F_1(a, b; c; z)$ is the Gauss hypergeometric function. When $N = 1$, $c_n = c_{1,n}$ are the classical Cauchy numbers. In 1875, Glaisher gave several interesting determinant expressions of numbers, including Bernoulli, Cauchy and Euler numbers (see (4), (6) and (3) in the text). Hypergeometric numbers can be recognized as one of the most natural extensions of the classical Cauchy numbers in terms of determinants (see Section 2), though many kinds of generalizations of the Cauchy numbers have been considered by many authors. In addition, there are some relations between the hypergeometric Cauchy numbers and the classical Cauchy numbers. In this paper, we give the determinant expressions of hypergeometric Cauchy numbers and their generalizations, and show some interesting expressions of hypergeometric Cauchy numbers. As applications, we can get the inversion relations such that hypergeometric Cauchy numbers as $c_{N,n}/n!$ and the numbers $N/(N+n)$ are interchanged in terms of determinants of the so-called Hassenberg matrices.

AMS Subject Classifications: 11B75, 11B37, 11C20, 15A15, 33C05.

Keywords: Cauchy numbers, hypergeometric Cauchy numbers, hypergeometric functions, determinants, recurrence relations.

Abbreviated title:

REMARKS ON HYPERGEOMETRIC CAUCHY NUMBERS

1. INTRODUCTION

Denote ${}_2F_1(a, b; c; z)$ be the Gauss hypergeometric function defined by

$${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)^{(n)}(b)^{(n)}}{(c)^{(n)}} \frac{z^n}{n!}$$

The first author was supported by JSPS KAKENHI Grant Numbers 26400015.

with the rising factorial $(x)^{(n)} = x(x+1) \dots (x+n-1)$ ($n \geq 1$) and $(x)^{(0)} = 1$. For $N \geq 1$, define the hypergeometric Cauchy numbers $c_{N,n}$ ([19]) by

$$(1) \quad \frac{1}{{}_2F_1(1, N; N+1; -x)} = \frac{(-1)^{N-1} x^N / N}{\log(1+x) - \sum_{n=1}^{N-1} (-1)^{n-1} x^n / n} = \sum_{n=0}^{\infty} c_{N,n} \frac{x^n}{n!}.$$

When $N = 1$, $c_n = c_{1,n}$ are classical Cauchy numbers ([5]) defined by

$$\frac{x}{\log(1+x)} = \sum_{n=0}^{\infty} c_n \frac{x^n}{n!}.$$

Notice that $b_n = c_n/n!$ are sometimes called the Bernoulli numbers of the second kind. In addition, the hypergeometric Cauchy polynomials $c_{M,N,n}(z)$ ([19]) are defined by

$$\begin{aligned} \frac{1}{(1+x)^z} \frac{1}{{}_2F_1(M, N; N+1; -x)} &= \frac{1}{(1+x)^z} \sum_{n=0}^{\infty} c_{M,N,n} \frac{x^n}{n!} \\ &= \sum_{n=0}^{\infty} c_{M,N,n}(z) \frac{x^n}{n!}. \end{aligned}$$

so that $c_{1,N,n}(0) = c_{N,n}$.

Similar hypergeometric numbers are hypergeometric Bernoulli numbers $B_{N,n}$ and hypergeometric Euler numbers. For $N \geq 1$, define hypergeometric Bernoulli numbers $B_{N,n}$ ([9, 10, 11, 12, 14, 32]) by

$$(2) \quad \frac{1}{{}_1F_1(1; N+1; x)} = \frac{x^N / N!}{e^x - \sum_{n=0}^{N-1} x^n / n!} = \sum_{n=0}^{\infty} B_{N,n} \frac{x^n}{n!},$$

where ${}_1F_1(a; b; z)$ is the confluent hypergeometric function defined by

$${}_1F_1(a; b; z) = \sum_{n=0}^{\infty} \frac{(a)^{(n)}}{(b)^{(n)}} \frac{z^n}{n!}.$$

When $N = 1$, $B_{1,n} = B_n$ are classical Bernoulli numbers, defined by

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}.$$

The hypergeometric Euler numbers $E_{N,n}$ ([29]) are defined by

$$\frac{1}{{}_1F_2(1; N+1, (2N+1)/2; x^2/4)} = \sum_{n=0}^{\infty} E_{N,n} \frac{x^n}{n!},$$

where ${}_1F_2(a; b, c; z)$ is the hypergeometric function defined by

$${}_1F_2(a; b, c; z) = \sum_{n=0}^{\infty} \frac{(a)^{(n)}}{(b)^{(n)}(c)^{(n)}} \frac{z^n}{n!}.$$

When $N = 0$, then $E_n = E_{0,n}$ are classical Euler numbers defined by

$$\frac{1}{\cosh x} = \sum_{n=0}^{\infty} E_n \frac{x^n}{n!}.$$

Several kinds of generalizations of the Cauchy numbers (or the Bernoulli numbers of the second kind) have been considered by many authors. For example, poly-Cauchy numbers [17], multiple Cauchy numbers, shifted Cauchy numbers [26], generalized Cauchy numbers [24], incomplete Cauchy numbers [20, 22, 25], various types of q -Cauchy numbers [4, 18, 21, 27], Cauchy Carlitz numbers [15, 16]. The situations are similar and even more for Bernoulli numbers and Euler numbers. Hypergeometric numbers can be recognized as one of the most natural extensions of the classical numbers in terms of determinants, though many kinds of generalizations of the classical numbers have been considered by many authors. In [23, 29], the hypergeometric Euler numbers $E_{N,2n}$ can be expressed as

$$E_{N,2n} = (-1)^n (2n)! \begin{vmatrix} \frac{(2N)!}{(2N+2)!} & 1 & & & \\ \frac{(2N)!}{(2N+4)!} & & \ddots & & \\ \vdots & & & \ddots & \\ \frac{(2N)!}{(2N+2n)!} & \cdots & \frac{(2N)!}{(2N+4)!} & \frac{(2N)!}{(2N+2)!} & 1 \end{vmatrix} \quad (N \geq 0, n \geq 1).$$

When $N = 0$, this is reduced to a famous determinant expression of Euler numbers (*cf.* [6, p.52]):

$$(3) \quad E_{2n} = (-1)^n (2n)! \begin{vmatrix} \frac{1}{2!} & & & & \\ \frac{1}{4!} & \frac{1}{2!} & & & \\ \vdots & & \ddots & & \\ \frac{1}{(2n-2)!} & \frac{1}{(2n-4)!} & & \frac{1}{2!} & 1 \\ \frac{1}{(2n)!} & \frac{1}{(2n-2)!} & \cdots & \frac{1}{4!} & \frac{1}{2!} \end{vmatrix}.$$

In addition, When $N = 1$, $E_{1,n}$ can be expressed by Bernoulli numbers as $E_{1,n} = -(n-1)B_n$ ([29]).

In [1], the hypergeometric Bernoulli numbers $B_{N,n}$ can be expressed as

$$B_{N,n} = (-1)^n n! \begin{vmatrix} \frac{N!}{(N+1)!} & 1 & & & \\ \frac{N!}{(N+2)!} & \frac{N!}{(N+1)!} & & & \\ \vdots & \vdots & \ddots & & \\ \frac{N!}{(N+n-1)!} & \frac{N!}{(N+n-2)!} & \cdots & \frac{N!}{(N+1)!} & 1 \\ \frac{N!}{(N+n)!} & \frac{N!}{(N+n-1)!} & \cdots & \frac{N!}{(N+2)!} & \frac{N!}{(N+1)!} \end{vmatrix} \quad (N \geq 1, n \geq 1).$$

When $N = 1$, we have a determinant expression of Bernoulli numbers ([6, p.53]):

$$(4) \quad B_n = (-1)^n n! \begin{vmatrix} \frac{1}{2!} & \frac{1}{3!} & & & \\ \vdots & \vdots & \ddots & 1 & \\ \frac{1}{n!} & \frac{1}{(n-1)!} & \cdots & \frac{1}{2!} & 1 \\ \frac{1}{(n+1)!} & \frac{1}{n!} & \cdots & \frac{1}{3!} & \frac{1}{2!} \end{vmatrix}.$$

In addition, relations between $B_{N,n}$ and $B_{N-1,n}$ are shown in [1].

In this paper, we shall give similar determinant expression of hypergeometric Cauchy numbers and their generalizations. We also study some interesting relations between the hypergeometric Cauchy numbers and the classical Cauchy numbers. As applications, we can get the inversion relations such that hypergeometric Cauchy numbers as $c_{N,n}/n!$ and the numbers $N/(N+n)$ are interchanged in terms of determinants of the so-called Hassenberg matrices.

2. A DETERMINANT EXPRESSION OF THE HYPERGEOMETRIC CAUCHY NUMBERS

From the definition (1), we have

$$(5) \quad \sum_{i=0}^n \frac{(-1)^i c_{N,i}}{(N+n-i)!} = 0 \quad (n \geq 1)$$

with $c_{N,0} = 1$ ([19, Proposition 1]).

By using this expression, the first few values of $c_{N,n}$ are given:

$$\begin{aligned} c_{N,0} &= 1, \\ c_{N,1} &= \frac{N}{N+1}, \\ c_{N,2} &= -\frac{2N}{(N+1)^2(N+2)}, \\ c_{N,3} &= \frac{6N(N^2+N+2)}{(N+1)^3(N+2)(N+3)}, \\ c_{N,4} &= -\frac{4!N(N^5+5N^4+14N^3+24N^2+20N+12)}{(N+1)^4(N+2)^2(N+3)(N+4)}, \\ c_{N,5} &= \frac{5!N(N^7+8N^6+35N^5+96N^4+160N^3+184N^2+116N+48)}{(N+1)^5(N+2)^2(N+3)(N+4)(N+5)}. \end{aligned}$$

In [19, Theorem 1], an explicit expression of hypergeometric Cauchy numbers is given.

LEMMA 1. For $N, n \geq 1$, we have

$$c_{N,n} = (-1)^n n! \sum_{r=1}^n (-N)^r \sum_{\substack{i_1 + \dots + i_r = n \\ i_1, \dots, i_r \geq 1}} \frac{1}{(N + i_1) \cdots (N + i_r)}.$$

Such values of $c_{N,n}$ can be expressed in terms of the determinant.

THEOREM 1. For $N, n \geq 1$, we have

$$c_{N,n} = n! \begin{vmatrix} \frac{N}{N+1} & 1 & & & \\ \frac{N}{N+2} & \frac{N}{N+1} & & & \\ \vdots & \vdots & \ddots & 1 & \\ \frac{N}{N+n-1} & \frac{N}{N+n-2} & \cdots & \frac{N}{N+1} & 1 \\ \frac{N}{N+n} & \frac{N}{N+n-1} & \cdots & \frac{N}{N+2} & \frac{N}{N+1} \end{vmatrix}.$$

Remark. When $N = 1$, we have a determinant expression of Cauchy numbers ([6, p.50]):

$$(6) \quad c_n = n! \begin{vmatrix} \frac{1}{2} & 1 & & & \\ \frac{1}{3} & \frac{1}{2} & & & \\ \vdots & \vdots & \ddots & 1 & \\ \frac{1}{n} & \frac{1}{n-1} & \cdots & \frac{1}{2} & 1 \\ \frac{1}{n+1} & \frac{1}{n} & \cdots & \frac{1}{3} & \frac{1}{2} \end{vmatrix}.$$

The value of this determinant, that is, $b_n = c_n/n!$ are called *Bernoulli numbers of the second kind*.

Proof of Theorem 1. Put $b_{N,n} = c_{N,n}/n!$. Then, we shall prove that

$$(7) \quad b_{N,n} = \begin{vmatrix} \frac{N}{N+1} & 1 & & & \\ \frac{N}{N+2} & \frac{N}{N+1} & & & \\ \vdots & \vdots & \ddots & 1 & \\ \frac{N}{N+n-1} & \frac{N}{N+n-2} & \cdots & \frac{N}{N+1} & 1 \\ \frac{N}{N+n} & \frac{N}{N+n-1} & \cdots & \frac{N}{N+2} & \frac{N}{N+1} \end{vmatrix}.$$

Since by (5)

$$c_{N,n} = \sum_{i=0}^{n-1} (-1)^{n-i-1} \frac{n!}{i!} \frac{N}{N+n-i} c_{N,i},$$

we have

$$b_{N,n} = N \sum_{i=0}^{n-1} \frac{(-1)^{n-i-1}}{N+n-i} b_{N,i}.$$

When $n = 1$, (7) is true as $b_{N,1} = N/(N+1)$. Assume that the result (7) is true up to $n-1$. Then, by expanding the last column of the right-hand side

of (7), it is equal to

$$\begin{aligned}
& \frac{N}{N+1} b_{N,n-1} - \left| \begin{array}{ccccc} \frac{N}{N+1} & 1 & & & \\ \frac{N}{N+2} & \frac{N}{N+1} & & & \\ \vdots & \vdots & \ddots & 1 & \\ \frac{N}{N+n-2} & \frac{N}{N+n-3} & \cdots & \frac{N}{N+1} & 1 \\ \frac{N}{N+n} & \frac{N}{N+n-1} & \cdots & \frac{N}{N+3} & \frac{N}{N+2} \end{array} \right| \\
&= \frac{N}{N+1} b_{N,n-1} - \frac{N}{N+2} b_{N,n-2} + \left| \begin{array}{ccccc} \frac{N}{N+1} & 1 & & & \\ \frac{N}{N+2} & \frac{N}{N+1} & & & \\ \vdots & \vdots & \ddots & 1 & \\ \frac{N}{N+n-3} & \frac{N}{N+n-4} & \cdots & \frac{N}{N+1} & 1 \\ \frac{N}{N+n} & \frac{N}{N+n-1} & \cdots & \frac{N}{N+4} & \frac{N}{N+3} \end{array} \right| \\
&= \frac{N}{N+1} b_{N,n-1} - \frac{N}{N+2} b_{N,n-2} + \frac{N}{N+3} b_{N,n-3} \\
&\quad - \cdots + (-1)^{n-2} \left| \begin{array}{cc} \frac{N}{N+1} & 1 \\ \frac{N}{N+n} & \frac{N}{N+n-1} \end{array} \right| \\
&= N \sum_{i=0}^{n-1} \frac{(-1)^{n-i-1}}{N+n-i} b_{N,i} = b_{N,n}.
\end{aligned}$$

Note that

$$b_{N,1} = \frac{N}{N+1} \quad \text{and} \quad b_{N,0} = 1.$$

□

3. A RELATION BETWEEN $c_{N,n}$ AND $c_{N-1,n}$

In this section, we show the following relation between $c_{N,n}$ and $c_{N-1,n}$.

PROPOSITION 1. *For $N \geq 2$ and $n \geq 1$, we have*

$$c_{N,n} = \sum_{m=0}^n \left(\frac{N}{1-N} \right)^m \sum_{0 \leq i_m < \cdots < i_1 < i_0 = n} \frac{n!}{i_m!} c_{N-1,i_m} \prod_{k=1}^m \frac{c_{N-1,i_{k-1}-i_k+1}}{(i_{k-1}-i_k+1)!},$$

where $i_0 = n$.

Examples

$$\begin{aligned}
\text{(i)} \quad c_{N,1} &= c_{N-1,1} + \frac{N}{1-N} \times \frac{c_{N-1,0} c_{N-1,2}}{2} \\
\text{(ii)} \quad c_{N,2} &= c_{N-1,2} + \frac{N}{1-N} \left(\frac{c_{N-1,3}}{3} + c_{N-1,1} c_{N-1,2} \right) + \left(\frac{N}{1-N} \right)^2 \times \frac{c_{N-1,2}^2}{2}
\end{aligned}$$

LEMMA 2. *For $N \geq 2$ and $n \geq 0$, we have*

$$c_{N,n} = c_{N-1,n} - \frac{N}{(n+1)(N-1)} \sum_{m=0}^{n-1} \binom{n+1}{m} c_{N,m} c_{N-1,n-m+1}.$$

Proof. From (1), we have

$$(-1)^{N-1} \times \frac{x^N}{N} = \left(\sum_{n=0}^{\infty} c_{N,n} \frac{x^n}{n!} \right) \left\{ \log(1+x) - \sum_{n=1}^{N-2} (-1)^{n-1} \frac{x^n}{n} - (-1)^{N-2} \frac{x^{N-1}}{N-1} \right\}.$$

By dividing $\log(1+x) - \sum_{n=1}^{N-2} (-1)^{n-1} x^n/n$, we have

$$\frac{1-N}{N} x \times \sum_{n=0}^{\infty} c_{N-1,n} \frac{x^n}{n!} = \left(\sum_{n=0}^{\infty} c_{N,n} \frac{x^n}{n!} \right) \left(1 - \sum_{n=0}^{\infty} c_{N-1,n} \frac{x^n}{n!} \right),$$

and hence

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{(1-N)n}{N} c_{N-1,n-1} \frac{x^n}{n!} &= \sum_{n=0}^{\infty} c_{N,n} \frac{x^n}{n!} - \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} c_{N,m} c_{N-1,k} \frac{x^{m+k}}{m!k!} \\ &= \sum_{n=0}^{\infty} c_{N,n} \frac{x^n}{n!} - \sum_{n=0}^{\infty} \sum_{m=0}^n c_{N,m} c_{N-1,n-m} \frac{x^n}{m!(n-m)!} \\ &= \sum_{n=0}^{\infty} \left\{ c_{N,n} - \sum_{m=0}^n \binom{n}{m} c_{N,m} c_{N-1,n-m} \right\} \frac{x^n}{n!}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} c_{N-1,n-1} &= \frac{N}{(1-N)n} \left\{ c_{N,n} - \sum_{m=0}^n \binom{n}{m} c_{N,m} c_{N-1,n-m} \right\} \\ &= \frac{N}{(1-N)n} \left\{ -\binom{n}{n-1} c_{N,n-1} c_{N-1,1} - \sum_{m=0}^{n-2} \binom{n}{m} c_{N,m} c_{N-1,n-m} \right\} \\ &= \frac{N}{(1-N)n} \left\{ \frac{(1-N)n}{N} c_{N,n-1} - \sum_{m=0}^{n-2} \binom{n}{m} c_{N,m} c_{N-1,n-m} \right\} \\ &= c_{N,n-1} - \frac{N}{(1-N)n} \sum_{m=0}^{n-2} \binom{n}{m} c_{N,m} c_{N-1,n-m}, \end{aligned}$$

for $n \geq 1$, and the proof is complete. \square

Proof of Proposition 1. We give the proof by induction for n . In the case $n = 1$, the assertion means

$$c_{N,1} = c_{N-1,1} + \frac{N}{1-N} \times \frac{c_{N-1,0} c_{N-1,2}}{2},$$

and this equality follows from $c_{N,1} = N/(N+1)$, $c_{N-1,1} = (N-1)/N$, $c_{N-1,0} = 1$ and $c_{N-1,2} = -2(N-1)/N^2(N+1)$. Assume that the assertion holds up to $n-1$. By Lemma 2, we have

$$\begin{aligned}
c_{N,n} &= c_{N-1,n} - \frac{N}{(n+1)(N-1)} \sum_{i_1=0}^{n-1} \binom{n+1}{i_1} c_{N,i_1} c_{N-1,n-i_1+1} \\
&= c_{N-1,n} - \frac{N}{(n+1)(N-1)} \sum_{i_1=0}^{n-1} \binom{n+1}{i_1} c_{N-1,n-i_1+1} \\
&\quad \times \sum_{m=1}^{i_1} \left(\frac{N}{1-N} \right)^{m-1} \sum_{0 \leq i_m < \dots < i_1} \frac{i_1!}{i_m!} c_{N-1,i_m} \prod_{k=2}^m \frac{c_{N-1,i_{k-1}-i_k+1}}{(i_{k-1}-i_k+1)!} \\
&= c_{N-1,n} + \sum_{i_1=0}^{n-1} \sum_{m=1}^{i_1} \sum_{0 \leq i_m < \dots < i_1} \left(\frac{N}{1-N} \right)^m \frac{n!}{i_m!} c_{N-1,i_m} \prod_{k=1}^m \frac{c_{N-1,i_{k-1}-i_k+1}}{(i_{k-1}-i_k+1)!} \\
&= c_{N-1,n} + \sum_{m=1}^n \left(\frac{N}{1-N} \right)^m \sum_{0 \leq i_m < \dots < i_1 < i_0=n} \frac{n!}{i_m!} c_{N-1,i_m} \prod_{k=1}^m \frac{c_{N-1,i_{k-1}-i_k+1}}{(i_{k-1}-i_k+1)!} \\
&= \sum_{m=0}^n \left(\frac{N}{1-N} \right)^m \sum_{0 \leq i_m < \dots < i_1 < i_0=n} \frac{n!}{i_m!} c_{N-1,i_m} \prod_{k=1}^m \frac{c_{N-1,i_{k-1}-i_k+1}}{(i_{k-1}-i_k+1)!}.
\end{aligned}$$

□

4. MULTIPLE HYPERGEOMETRIC CAUCHY NUMBERS

For positive integers N and r , define the *hypergeometric Cauchy numbers* $c_{N,n}^{(r)}$ by the generating function

$$\begin{aligned}
\frac{1}{({}_2F_1(1, N; N+1; -x))^r} &= \left(\frac{(-1)^{N-1} x^N / N}{\log(1+x) - \sum_{n=1}^{N-1} (-1)^{n-1} x^n / n} \right)^r \\
(8) \quad &= \sum_{n=0}^{\infty} c_{N,n}^{(r)} \frac{x^n}{n!}.
\end{aligned}$$

From the definition (8),

$$\begin{aligned}
&\left(\frac{(-1)^{N-1} x^N}{N} \right)^r \\
&= \left(\sum_{i=0}^{\infty} \frac{(-1)^{i+N-1} x^{N+i}}{N+i} \right)^r \left(\sum_{n=0}^{\infty} c_{N,n}^{(r)} \frac{x^n}{n!} \right) \\
&= x^{rN} (-1)^{r(N-1)} \left(\sum_{l=0}^{\infty} \sum_{\substack{i_1+\dots+i_r=l \\ i_1, \dots, i_r \geq 0}} \frac{(-1)^l l!}{(N+i_1) \cdots (N+i_r)} \frac{x^l}{l!} \right) \left(\sum_{n=0}^{\infty} c_{N,n}^{(r)} \frac{x^n}{n!} \right) \\
&= x^{rN} (-1)^{r(N-1)} \sum_{n=0}^{\infty} \sum_{m=0}^n \sum_{\substack{i_1+\dots+i_r=n-m \\ i_1, \dots, i_r \geq 0}} \binom{n}{m} \frac{(-1)^{n-m} (n-m)!}{(N+i_1) \cdots (N+i_r)} c_{N,m}^{(r)} \frac{x^n}{n!}.
\end{aligned}$$

Hence, as a generalization of Proposition (5), for $n \geq 1$, we have the following.

PROPOSITION 2.

$$\sum_{m=0}^n \sum_{\substack{i_1+\dots+i_r=n-m \\ i_1, \dots, i_r \geq 0}} \frac{(-1)^{n-m} c_{N,m}^{(r)}}{m!(N+i_1) \cdots (N+i_r)} = 0.$$

By using Proposition 2 or

$$(9) \quad c_{N,n}^{(r)} = -n!N^r \sum_{m=0}^{n-1} \sum_{\substack{i_1+\dots+i_r=n-m \\ i_1, \dots, i_r \geq 0}} \frac{(-1)^{n-m} c_{N,m}^{(r)}}{m!(N+i_1) \cdots (N+i_r)}$$

with $c_{N,0}^{(r)} = 1$ ($N \geq 1$), some values of $c_{N,n}^{(r)}$ ($0 \leq n \leq 4$) are explicitly given by the following.

$$\begin{aligned} c_{N,0}^{(r)} &= 1, \\ c_{N,1}^{(r)} &= \frac{rN}{N+1}, \\ c_{N,2}^{(r)} &= \frac{r(r+1)N^2}{(N+1)^2} - \frac{2rN}{N+2}, \\ c_{N,3}^{(r)} &= \frac{r(r+1)(r+2)N^3}{(N+1)^3} - \frac{6r(r+1)N^2}{(N+1)(N+2)} + \frac{6rN}{N+3}, \\ c_{N,4}^{(r)} &= \frac{r(r+1)(r+2)(r+3)N^4}{(N+1)^4} - \frac{12r(r+1)(r+2)N^3}{(N+1)^2(N+2)} \\ &\quad + \frac{24r(r+1)N^2}{(N+1)(N+3)} + \frac{12r(r+1)N^2}{(N+2)^2} - \frac{24rN}{N+4}. \end{aligned}$$

As a generalization of Lemma 1, we have an explicit expression of $c_{N,n}^{(r)}$.

PROPOSITION 3. *For $N, n \geq 1$, we have*

$$c_{N,n}^{(r)} = n! \sum_{k=1}^n (-1)^{n-k} \sum_{\substack{e_1+\dots+e_k=n \\ e_1, \dots, e_k \geq 1}} D_r(e_1) \cdots D_r(e_k),$$

where

$$(10) \quad D_r(e) = \sum_{\substack{i_1+\dots+i_r=e \\ i_1, \dots, i_r \geq 0}} \frac{N^r}{(N+i_1) \cdots (N+i_r)}.$$

The first few values of $D_r(e)$ are given by the following.

$$\begin{aligned} D_r(1) &= \frac{rN}{N+1}, \\ D_r(2) &= \frac{rN}{N+2} + \frac{r(r-1)N^2}{2(N+1)^2}, \end{aligned}$$

$$\begin{aligned}
D_r(3) &= \frac{rN}{N+3} + \frac{r(r-1)N^2}{(N+1)(N+2)} + \binom{r}{3} \frac{N^3}{(N+1)^3}, \\
D_r(4) &= \frac{rN}{N+4} + \frac{r(r-1)N^2}{(N+1)(N+3)} + \binom{r}{2} \frac{N^2}{(N+1)^2} \\
&\quad + r \binom{r-1}{2} \frac{N^3}{(N+1)^2(N+2)} + \binom{r}{4} \frac{N^4}{(N+1)^4}.
\end{aligned}$$

We shall introduce the Hasse-Teichmüller derivative in order to prove Proposition 3 easily. Let \mathbb{F} be a field of any characteristic, $\mathbb{F}[[z]]$ the ring of formal power series in one variable z , and $\mathbb{F}((z))$ the field of Laurent series in z . Let n be a nonnegative integer. We define the Hasse-Teichmüller derivative $H^{(n)}$ of order n by

$$H^{(n)} \left(\sum_{m=R}^{\infty} c_m z^m \right) = \sum_{m=R}^{\infty} c_m \binom{m}{n} z^{m-n}$$

for $\sum_{m=R}^{\infty} c_m z^m \in \mathbb{F}((z))$, where R is an integer and $c_m \in \mathbb{F}$ for any $m \geq R$. Note that $\binom{m}{n} = 0$ if $m < n$.

The Hasse-Teichmüller derivatives satisfy the product rule [34], the quotient rule [7] and the chain rule [8]. One of the product rules can be described as follows.

LEMMA 3. *For $f_i \in \mathbb{F}[[z]]$ ($i = 1, \dots, k$) with $k \geq 2$ and for $n \geq 1$, we have*

$$H^{(n)}(f_1 \cdots f_k) = \sum_{\substack{i_1 + \cdots + i_k = n \\ i_1, \dots, i_k \geq 0}} H^{(i_1)}(f_1) \cdots H^{(i_k)}(f_k).$$

The quotient rules can be described as follows.

LEMMA 4. *For $f \in \mathbb{F}[[z]] \setminus \{0\}$ and $n \geq 1$, we have*

$$(11) \quad H^{(n)} \left(\frac{1}{f} \right) = \sum_{k=1}^n \frac{(-1)^k}{f^{k+1}} \sum_{\substack{i_1 + \cdots + i_k = n \\ i_1, \dots, i_k \geq 1}} H^{(i_1)}(f) \cdots H^{(i_k)}(f)$$

$$(12) \quad = \sum_{k=1}^n \binom{n+1}{k+1} \frac{(-1)^k}{f^{k+1}} \sum_{\substack{i_1 + \cdots + i_k = n \\ i_1, \dots, i_k \geq 0}} H^{(i_1)}(f) \cdots H^{(i_k)}(f).$$

Proof of Proposition 3. Put $h(x) = (f(x))^r$, where

$$f(x) = \frac{\sum_{i=N}^{\infty} (-1)^{i-1} \frac{x^i}{i}}{(-1)^{N-1} \frac{x^N}{N}} = \sum_{j=0}^{\infty} \frac{(-1)^j N}{N+j} x^j.$$

Since

$$\begin{aligned} H^{(i)}(f) \Big|_{x=0} &= \sum_{j=i}^{\infty} \frac{(-1)^j N}{N+j} \binom{j}{i} x^{j-i} \Big|_{x=0} \\ &= \frac{(-1)^i N}{N+i} \end{aligned}$$

by the product rule of the Hasse-Teichmüller derivative in Lemma 3, we get

$$\begin{aligned} H^{(e)}(h) \Big|_{x=0} &= \sum_{\substack{i_1+\dots+i_r=e \\ i_1, \dots, i_r \geq 0}} H^{(i_1)}(f) \Big|_{x=0} \cdots H^{(i_r)}(f) \Big|_{x=0} \\ &= \sum_{\substack{i_1+\dots+i_r=e \\ i_1, \dots, i_r \geq 0}} \frac{(-1)^{i_1} N}{N+i_1} \cdots \frac{(-1)^{i_r} N}{N+i_r} \\ &= (-1)^e \sum_{\substack{i_1+\dots+i_r=e \\ i_1, \dots, i_r \geq 0}} \frac{N^r}{(N+i_1) \cdots (N+i_r)} := (-1)^e D_r(e). \end{aligned}$$

Hence, by the quotient rule of the Hasse-Teichmüller derivative in Lemma 4 (11), we have

$$\begin{aligned} \frac{c_{N,n}^{(r)}}{n!} &= \sum_{k=1}^n \frac{(-1)^k}{h^{k+1}} \Big|_{x=0} \sum_{\substack{e_1+\dots+e_k=n \\ e_1, \dots, e_k \geq 1}} H^{(e_1)}(h) \Big|_{x=0} \cdots H^{(e_k)}(h) \Big|_{x=0} \\ &= \sum_{k=1}^n (-1)^k \sum_{\substack{e_1+\dots+e_k=n \\ e_1, \dots, e_k \geq 1}} (-1)^n D_r(e_1) \cdots D_r(e_k). \end{aligned}$$

□

Now, we can also show a determinant expression of $c_{N,n}^{(r)}$.

THEOREM 2. *For $N, n \geq 1$, we have*

$$c_{N,n}^{(r)} = n! \begin{vmatrix} D_r(1) & 1 & & & \\ D_r(2) & D_r(1) & & & \\ \vdots & \vdots & \ddots & & \\ D_r(n-1) & D_r(n-2) & \cdots & D_r(1) & 1 \\ D_r(n) & D_r(n-1) & \cdots & D_r(2) & D_r(1) \end{vmatrix}.$$

where $D_r(e)$ are given in (10).

Remark. When $r = 1$ in Theorem 2, we have the result in Theorem 1.

Proof. For simplicity, put $b_{N,n}^{(r)} = c_{N,n}^{(r)}/n!$. Then, we shall prove that for any $n \geq 1$

$$(13) \quad b_{N,n}^{(r)} = \begin{vmatrix} D_r(1) & 1 & & & \\ D_r(2) & D_r(1) & & & \\ \vdots & \vdots & \ddots & 1 & \\ D_r(n-1) & D_r(n-2) & \cdots & D_r(1) & 1 \\ D_r(n) & D_r(n-1) & \cdots & D_r(2) & D_r(1) \end{vmatrix}.$$

When $n = 1$, (13) is valid because

$$D_r(1) = \frac{rN^r}{N^{r-1}(N+1)} = \frac{rN}{N+1} = b_{N,1}^{(r)}.$$

Assume that (13) is valid up to $n-1$. Notice that by (9), we have

$$b_{N,n}^{(r)} = \sum_{l=1}^n (-1)^{l-1} b_{N,n-l}^{(r)} D_r(l).$$

Thus, by expanding the first row of the right-hand side (13), it is equal to

$$\begin{aligned} & D_r(1) b_{N,n-1}^{(r)} - \begin{vmatrix} D_r(2) & 1 & & & \\ D_r(3) & D_r(1) & & & \\ \vdots & \vdots & \ddots & 1 & \\ D_r(n-1) & D_r(n-3) & \cdots & D_r(1) & 1 \\ D_r(n) & D_r(n-2) & \cdots & D_r(2) & D_r(1) \end{vmatrix} \\ &= D_r(1) b_{N,n-1}^{(r)} - D_r(2) b_{N,n-2}^{(r)} \\ & \quad + \begin{vmatrix} D_r(3) & 1 & & & \\ D_r(4) & D_r(1) & & & \\ \vdots & \vdots & \ddots & 1 & \\ D_r(n-1) & D_r(n-4) & \cdots & D_r(1) & 1 \\ D_r(n) & D_r(n-3) & \cdots & D_r(2) & D_r(1) \end{vmatrix} \\ &= D_r(1) b_{N,n-1}^{(r)} - D_r(2) b_{N,n-2}^{(r)} + \cdots + (-1)^{n-2} \begin{vmatrix} D_r(n-1) & 1 \\ D_r(n) & D_r(1) \end{vmatrix} \\ &= \sum_{l=1}^n (-1)^{l-1} D_r(l) b_{N,n-l}^{(r)} = b_{N,n}^{(r)}. \end{aligned}$$

Note that $b_{N,1}^{(r)} = D_r(1)$ and $b_{N,0}^{(r)} = 1$. □

5. A RELATION BETWEEN $c_{N,n}^{(r)}$ AND $c_{N,n}$

In this section, we show the following relation between $c_{N,n}^{(r)}$ and $c_{N,n}$.

LEMMA 5. For $r \geq 2$ and $N, n \geq 0$, we have

$$c_{N,n}^{(r)} = \sum_{\substack{n_1, \dots, n_r \geq 0 \\ n_1 + \dots + n_r = n}} \frac{n!}{n_1! \dots n_r!} C_{N,n_1} \dots C_{N,n_r},$$

Proof. From the definition (8), we have

$$\begin{aligned} \sum_{n=0}^{\infty} C_{N,n}^{(r)} \frac{n!}{x^n} &= \left(\sum_{n=0}^{\infty} C_{N,n} \frac{x^n}{n!} \right)^r \\ &= \sum_{n=0}^{\infty} \sum_{\substack{n_1, \dots, n_r \geq 0 \\ n_1 + \dots + n_r = n}} \frac{n!}{n_1! \dots n_r!} C_{N,n_1} \dots C_{N,n_r} \frac{x^n}{n!}, \end{aligned}$$

and we get the assertion. \square

Examples

- (i) $c_{N,0}^{(r)} = c_{N,0}^{r+1}$
- (ii) $c_{N,1}^{(r)} = r c_{N,1}$
- (iii) $c_{N,2}^{(r)} = r c_{N,2} c_{N,0}^{N-1} + r(r-1) c_{N,1}^2 c_{N,0}^{r-2}$

6. APPLICATIONS BY THE TRUDI'S FORMULA AND INVERSION EXPRESSIONS

The expressions in Theorem 1 and Theorem 2 are useful for several applications too. We can obtain different explicit expressions for the numbers $c_{N,n}^{(r)}$, $c_{N,n}$ and c_n by using the Trudi's formula. We also show some inversion formulas. The following relation is known as Trudi's formula [30, Vol.3, p.214], [35] and the case $a_0 = 1$ of this formula is known as Brioschi's formula [2], [30, Vol.3, pp.208–209].

LEMMA 6. For a positive integer m , we have

$$\begin{aligned} & \begin{vmatrix} a_1 & a_2 & \cdots & \cdots & a_m \\ a_0 & a_1 & \ddots & & \vdots \\ & \ddots & \ddots & \ddots & \vdots \\ & & \ddots & a_1 & a_2 \\ & & & a_0 & a_1 \end{vmatrix} \\ &= \sum_{t_1 + 2t_2 + \dots + mt_m = m} \binom{t_1 + \dots + t_m}{t_1, \dots, t_m} (-a_0)^{m-t_1-\dots-t_m} a_1^{t_1} a_2^{t_2} \dots a_m^{t_m}, \end{aligned}$$

where $\binom{t_1 + \dots + t_m}{t_1, \dots, t_m} = \frac{(t_1 + \dots + t_m)!}{t_1! \dots t_m!}$ are the multinomial coefficients.

In addition, there exists the following inversion formula, which is based upon the relation:

$$\sum_{k=0}^n (-1)^{n-k} \alpha_k R(n-k) = 0 \quad (n \geq 1).$$

LEMMA 7. *If $\{\alpha_n\}_{n \geq 0}$ is a sequence defined by $\alpha_0 = 1$ and*

$$\alpha_n = \begin{vmatrix} R(1) & 1 & & \\ R(2) & \ddots & \ddots & \\ \vdots & \ddots & \ddots & 1 \\ R(n) & \cdots & R(2) & R(1) \end{vmatrix}, \text{ then } R(n) = \begin{vmatrix} \alpha_1 & 1 & & \\ \alpha_2 & \ddots & \ddots & \\ \vdots & \ddots & \ddots & 1 \\ \alpha_n & \cdots & \alpha_2 & \alpha_1 \end{vmatrix}.$$

Moreover, if

$$A = \begin{pmatrix} 1 & & & \\ \alpha_1 & 1 & & \\ \vdots & \ddots & \ddots & \\ \alpha_n & \cdots & \alpha_1 & 1 \end{pmatrix}, \text{ then } A^{-1} = \begin{pmatrix} 1 & & & \\ R(1) & 1 & & \\ \vdots & \ddots & \ddots & \\ R(n) & \cdots & R(1) & 1 \end{pmatrix}.$$

From Trudi's formula, it is possible to give the combinatorial expression

$$\alpha_n = \sum_{t_1+2t_2+\cdots+nt_n=n} \binom{t_1+\cdots+t_n}{t_1, \dots, t_n} (-1)^{n-t_1-\cdots-t_n} R(1)^{t_1} R(2)^{t_2} \cdots R(n)^{t_n}.$$

By applying these lemmata to Theorem 2, we obtain an explicit expression for the generalized hypergeometric Cauchy numbers $c_{N,n}^{(r)}$. A different version can be seen in [28].

THEOREM 3. *For $n \geq 1$*

$$\begin{aligned} & c_{N,n}^{(r)} \\ &= n! \sum_{t_1+2t_2+\cdots+nt_n=n} \binom{t_1+\cdots+t_n}{t_1, \dots, t_n} (-1)^{n-t_1-\cdots-t_n} D_r(1)^{t_1} D_r(2)^{t_2} \cdots D_r(n)^{t_n}, \end{aligned}$$

where $D_r(e)$ are given in (10). Moreover,

$$D_r(n) = \begin{vmatrix} \frac{c_{N,1}^{(r)}}{1!} & 1 & & \\ \frac{c_{N,2}^{(r)}}{2!} & \ddots & \ddots & \\ \vdots & \ddots & \ddots & 1 \\ \frac{c_{N,n}^{(r)}}{n!} & \cdots & \frac{c_{N,2}^{(r)}}{2!} & \frac{c_{N,1}^{(r)}}{1!} \end{vmatrix},$$

and

$$\left(\begin{array}{cccc} 1 & & & \\ \frac{c_{N,1}^{(r)}}{1!} & 1 & & \\ \frac{c_{N,2}^{(r)}}{2!} & \frac{c_{N,1}^{(r)}}{1!} & 1 & \\ \vdots & & \ddots & \ddots \\ \frac{c_{N,n}^{(r)}}{n!} & \dots & \frac{c_{N,2}^{(r)}}{2!} & \frac{c_{N,1}^{(r)}}{1!} & 1 \end{array} \right)^{-1} = \left(\begin{array}{ccccc} 1 & & & & \\ D_r(1) & 1 & & & \\ D_r(2) & D_r(1) & 1 & & \\ \vdots & & \ddots & \ddots & \\ D_r(n) & \dots & D_r(2) & D_r(1) & 1 \end{array} \right).$$

When $r = 1$ in Theorem 3, we have an explicit expression for the numbers $c_{N,n}$.

COROLLARY 1. *For $n \geq 1$*

$$c_{N,n} = n! \sum_{t_1+2t_2+\dots+nt_n=n} \binom{n-t_1-\dots-t_n}{t_1, \dots, t_n} (-1)^{t_1+\dots+t_n} \times \left(\frac{N}{N+1} \right)^{t_1} \left(\frac{N}{N+2} \right)^{t_2} \dots \left(\frac{N}{N+n} \right)^{t_n}$$

and

$$\frac{N}{N+n} = \left| \begin{array}{cccc} \frac{c_{N,1}}{1!} & 1 & & \\ \frac{c_{N,2}}{2!} & \ddots & \ddots & \\ \vdots & \ddots & \ddots & 1 \\ \frac{c_{N,n}}{n!} & \dots & \frac{c_{N,2}}{2!} & \frac{c_{N,1}}{1!} \end{array} \right|,$$

When $r = N = 1$ in Theorem 3, we have a different expression of the classical Cauchy numbers.

COROLLARY 2. *We have for $n \geq 1$*

$$c_n = n! \sum_{t_1+2t_2+\dots+nt_n=n} \binom{t_1+\dots+t_n}{t_1, \dots, t_n} (-1)^{n-t_1-\dots-t_n} \times \left(\frac{1}{2} \right)^{t_1} \left(\frac{1}{3} \right)^{t_2} \dots \left(\frac{1}{n+1} \right)^{t_n}$$

and

$$\frac{1}{n+1} = \left| \begin{array}{cccc} \frac{c_1}{1!} & 1 & & \\ \frac{c_2}{2!} & \ddots & \ddots & \\ \vdots & \ddots & \ddots & 1 \\ \frac{c_n}{n!} & \dots & \frac{c_2}{2!} & \frac{c_1}{1!} \end{array} \right|.$$

7. ADDITIONAL COMMENTS

Hypergeometric Cauchy numbers are not integers, but fractions. Hence, combinatorial interpretations of the above results or congruent relations seem to be difficult to obtain. Nevertheless, definition (1) is not obvious or artificial, but has motivations from Combinatorics, in particular, graph theory. In 1989, Cameron [3] considered the operator A defined on the set of sequences of non-negative integers as follows. For $x = \{x_n\}_{n \geq 1}$ and $z = \{z_n\}_{n \geq 1}$, let $Ax = z$, where

$$(14) \quad 1 + \sum_{n=1}^{\infty} z_n t^n = \left(1 - \sum_{n=1}^{\infty} x_n t^n \right)^{-1}.$$

For hypergeometric Cauchy numbers, we have

$$x_n = \frac{c_{N,n}}{n!} \quad \text{and} \quad z_n = \frac{(-1)^{n-1} N}{N+n}$$

and vice versa. If x enumerates a class C , then Ax enumerates the class of disjoint unions of members. More concrete examples can be seen in [3].

REFERENCES

- [1] M. Aoki and T. Komatsu, *Remarks on hypergeometric Bernoulli numbers*, preprint.
- [2] F. Brioschi, *Sulle funzioni Bernoulliane ed Euleriane*, Annali de Mat., i. (1858) 260–263; Opere Mat., i. pp. 343–347.
- [3] P. J. Cameron, *Some sequences of integers*, Discrete Math. **75** (1989), 89–102.
- [4] M. Cenkci and T. Komatsu, *Poly-Bernoulli numbers and polynomials with a q parameter*, J. Number Theory **152** (2015), 38–54.
- [5] L. Comtet, *Advanced Combinatorics*, Reidel, Dordrecht, 1974.
- [6] J. W. L. Glaisher, *Expressions for Laplace's coefficients, Bernoullian and Eulerian numbers etc. as determinants*, Messenger (2) **6** (1875), 49–63.
- [7] R. Gottfert, H. Niederreiter, *Hasse-Teichmüller derivatives and products of linear recurring sequences*, Finite Fields: Theory, Applications, and Algorithms (Las Vegas, NV, 1993), Contemporary Mathematics, vol. 168, American Mathematical Society, Providence, RI, 1994, pp.117–125.
- [8] H. Hasse, *Theorie der höheren Differentiale in einem algebraischen Funktionenkörper mit Vollkommenem Konstantenkörper bei beliebiger Charakteristik*, J. Reine Angew. Math. **175** (1936), 50–54.
- [9] A. Hassen and H. D. Nguyen, *Hypergeometric Bernoulli polynomials and Appell sequences*, Int. J. Number Theory **4** (2008), 767–774.
- [10] A. Hassen and H. D. Nguyen, *Hypergeometric zeta functions*, Int. J. Number Theory **6** (2010), 99–126.
- [11] F. T. Howard, *A sequence of numbers related to the exponential function*, Duke Math. J. **34** (1967), 599–615.
- [12] F. T. Howard, *Some sequences of rational numbers related to the exponential function*, Duke Math. J. **34** (1967), 701–716.
- [13] S. Hu and M.-S. Kim, *On hypergeometric Bernoulli numbers and polynomials*, Acta Math. Hungar. **154** (2018), 134–146.
- [14] K. Kamano, *Sums of products of hypergeometric Bernoulli numbers*, J. Number Theory **130** (2010), 2259–2271.
- [15] M. Kaneko and T. Komatsu, *Cauchy-Carlitz numbers*, J. Number Theory **163** (2016), 238–254.

- [16] M. Kaneko and T. Komatsu, *Bernoulli-Carlitz and Cauchy-Carlitz numbers with Stirling-Carlitz numbers*, RIMS Kokyuroku Bessatsu (to appear). arXiv:1704.07135
- [17] T. Komatsu, *Poly-Cauchy numbers*, Kyushu J. Math. **67** (2013), 143–153.
- [18] T. Komatsu, *Poly-Cauchy numbers with a q parameter*, Ramanujan J. **31** (2013), 353–371.
- [19] T. Komatsu, *Hypergeometric Cauchy numbers*, Int. J. Number Theory **9** (2013), 545–560.
- [20] T. Komatsu, *Incomplete poly-Cauchy numbers*, Monatsh. Math. **180** (2016), 271–288.
- [21] T. Komatsu, *q -poly-Bernoulli numbers and q -poly-Cauchy numbers with a parameter by Jackson's integrals*, Indag. Math. **27** (2016), 100–111.
- [22] T. Komatsu, *Generalized incomplete poly-Bernoulli and poly-Cauchy numbers*, Period. Math. Hungar. **75** (2017), 96–113.
- [23] T. Komatsu, *Complementary Euler numbers*, Period. Math. Hungar. **75** (2017), 302–314.
- [24] T. Komatsu, V. Laohakosol and K. Liptai, *A generalization of poly-Cauchy numbers and their properties*, Abstr. Appl. Anal. **2013** (2013), Article ID 179841, 8 pages.
- [25] T. Komatsu, I. Mező and L. Szalay, *Incomplete Cauchy numbers*, Acta Math. Hungar. **149** (2016), 306–323.
- [26] T. Komatsu and L. Szalay, *Shifted poly-Cauchy numbers*, Lithuanian Math. J. **54** (2014), 166–181.
- [27] T. Komatsu and L. Szalay, *q -multiparameter-Bernoulli polynomials and q -multiparameter-Cauchy polynomials by Jackson's integrals*, Integers **16** (2016), #A39, 11 pages.
- [28] T. Komatsu and P. Yuan, *Hypergeometric Cauchy numbers and polynomials*, Acta Math. Hungar. **153** (2017), 382–400.
- [29] T. Komatsu and H. Zhu, *Hypergeometric Euler numbers*, preprint. arXiv:1612.06210.
- [30] T. Muir, *The theory of determinants in the historical order of development*, Four volumes, Dover Publications, New York, 1960.
- [31] T. Muir, *A treatise on the theory of determinants*, Revised and enlarged by William H. Metzler, Dover Publications, Inc., New York 1960.
- [32] H. D. Nguyen, *Sums of products of hypergeometric Bernoulli polynomials*, MAA-NJ Section - Spring Meeting, Middle Country College, NJ, April 10, 2010.
- [33] H. D. Nguyen and L. G. Cheong, *New convolution identities for hypergeometric Bernoulli polynomials*, J. Number Theory **137** (2014), 201–221.
- [34] O. Teichmüller, *Differentialrechnung bei Charakteristik p* , J. Reine Angew. Math. **175** (1936), 89–99.
- [35] N. Trudi, *Intorno ad alcune formole di sviluppo*, Rendic. dell' Accad. Napoli, (1862), 135–143.

DEPARTMENT OF MATHEMATICS, SHIMANE UNIVERSITY, MATSUE, SHIMANE, JAPAN,
E-mail address: aoki@riko.shimane-u.ac.jp

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN 430072 CHINA,
E-mail address: komatsu@whu.edu.cn