

EXTREMALS FOR FRACTIONAL ORDER HARDY-SOBOLEV-MAZ'YA INEQUALITY

ARKA MALLICK

*TIFR Centre for Applicable Mathematics, Post Bag No. 6503 Sharadanagar,
Bangalore 560065, India.*

ABSTRACT. In this article, we derive the existence of positive solution of a semi-linear, non-local elliptic PDE, involving a singular perturbation of the fractional laplacian, coming from the fractional Hardy-Sobolev-Maz'ya inequality, derived in this paper. We also derive symmetry properties and a precise asymptotic behaviour of solutions.

MSC2010 Classification: *46E35, 35J70, 35J10, 35S05, 35B25*

Keywords: *Fractional Hardy-Sobolev-Maz'ya inequality, existence, cylindrical symmetry.*

1. INTRODUCTION

In this article, we study the following equation

$$\left. \begin{aligned} \frac{2}{c_{N,s}} (-\Delta)^s u - \beta \frac{u}{|x'|^{2s}} &= \frac{u^{2_t^*-1}}{|x'|^t}, \text{ in } \mathbb{R}^N, \\ u &\geq 0, \quad u \in \dot{\mathcal{H}}_\beta^s(\mathbb{R}^N), \end{aligned} \right\} \quad (1.1)$$

where $c_{N,s} = 2^{2s} \pi^{-N/2} \frac{\Gamma(\frac{N+2s}{2})}{|\Gamma(-s)|}$, $0 < s < 1$, $0 \leq t < 2s$, $2_t^* = \frac{2(N-t)}{N-2s}$, $\mathbb{R}^N = \mathbb{R}^k \times \mathbb{R}^{N-k}$, $2 \leq k \leq N-2$ and a point $x \in \mathbb{R}^N$ is denoted as $x = (x', x'') \in \mathbb{R}^k \times \mathbb{R}^{N-k}$. The space $\dot{\mathcal{H}}_\beta^s(\mathbb{R}^N)$ is defined in Section 2, where $0 < \beta \leq \mathcal{C}_{N,k,s}$ and $\mathcal{C}_{N,k,s}$ is the best constant of the following fractional version of Hardy-Maz'ya inequality

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy \geq \mathcal{C}_{N,k,s} \int_{\mathbb{R}^N} \frac{u^2(x)}{|x'|^{2s}} dx, \quad \forall u \in C_c^{0,1}(\mathbb{R}^k \setminus \{0\} \times \mathbb{R}^{N-k}). \quad (1.2)$$

An explicit expression of $\mathcal{C}_{N,k,s}$ and the proof of (1.2) can be found in Section 3. For $k = N$ the inequality (1.2) has been derived in [15], [16], [34]. Also, see [21], for similar inequalities.

The local counterpart (i.e. for the case $s = 1$) of (1.2) is derived by Maz'ya in [28] which can be state as follows

$$\int_{\mathbb{R}^N} |\nabla u(x)|^2 dx \geq \left(\frac{k-2}{2} \right)^2 \int_{\mathbb{R}^N} \frac{u^2(x)}{|x'|^2} dx, \quad \forall u \in C_c^\infty(\mathbb{R}^k \setminus \{0\} \times \mathbb{R}^{N-k}), \quad (1.3)$$

where the constant $\frac{(k-2)^2}{4}$ is the best possible. When $k = N$, (1.3) reduces to the usual Hardy inequality. Unlike the case of Hardy inequality, it was Maz'ya [28], who first observed that, the following Sobolev type improvement of (1.3) can be achieved when $k \geq 2$

$$C_\beta^t \left(\int_{\mathbb{R}^N} \frac{|u(x)|^q}{|x'|^t} dx \right)^{\frac{2}{q}} \leq \int_{\mathbb{R}^N} |\nabla u(x)|^2 dx - \beta \int_{\mathbb{R}^N} \frac{u^2(x)}{|x'|^2} dx, \quad (1.4)$$

E-mail address: arkamallick02@gmail.com, arka@math.tifrbng.res.in.

where $\beta \leq \frac{(k-2)^2}{4}$, $0 \leq t < 2$, $q = \frac{2(N-t)}{N-2}$ and $u \in C_c^\infty(\mathbb{R}^k \setminus \{0\} \times \mathbb{R}^{N-k})$. Note that, existence of nontrivial solution of (1.1) will follow for the case of $s = 1$, if we can show the existence of minimizers of (1.4). For $\beta = 0$, the existence of minimizers of (1.4) has been established in [2] by using concentration compactness principle due to P.L Lions [25], [24]. Whereas, for $0 < \beta < \frac{(k-2)^2}{4}$, the existence is proved in [29], by using blow up analysis for approximate solutions with a rescaling argument. On the other hand, since for $\beta = \frac{(k-2)^2}{4}$, the expected space in which the minimizers will belong is much bigger than the same for the case of $\beta < \frac{(k-2)^2}{4}$, one needs to employ a careful analysis. Using a penalty method, Tintarev and Tertikas proved the existence of minimizers for the case of $\beta = \frac{(k-2)^2}{4}$ in [33] and subsequently improved in [19].

The cylindrical symmetry of the local counterpart (i.e. $s = 1$) of (1.1), has been established in [26] by using moving plane method in the special case of $\beta = 0$. In fact, when $t = 1$, they have classified all the solutions by a careful asymptotic analysis. Subsequently, in the case of $0 \leq \beta < \frac{(k-2)^2}{4}$, $0 \leq t < 2$ and $s = 1$, cylindrical symmetry of solutions of (1.1) has been established in [18]. Finally, in a breakthrough paper, Sandeep and Mancini [27], established the uniqueness of positive extremals of (1.4). See also [8].

Thus we need the following fractional version of Hardy-Sobolev-Mazya inequality, to prove the existence of solution of (1.1).

$$C \left(\int_{\mathbb{R}^N} \frac{|u(x)|^{2_t^*}}{|x'|^t} dx \right)^{\frac{2}{2_t^*}} \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{u^2(x)}{|x'|^{2s}} dx, \quad (1.5)$$

where $0 \leq t < 2s$, $2_t^* := \frac{2(N-t)}{N-2s}$, $0 \leq \beta \leq \mathcal{C}_{N,k,s}$, $\mathcal{C}_{N,k,s}$ is the optimal constant of (1.2) and $C > 0$ is a constant independent of u . We will establish the inequality (1.5) in Section 3. In Section 4, we have proved the following theorems to establish the existence of a solution to the equation (1.1).

Theorem 1.1. *Let $0 < s < 1$, $0 \leq t < 2s$ and $0 \leq \beta \leq \mathcal{C}_{N,k,s}$. Define $S_k^t(\beta)$ in the following manner*

$$S_k^t(\beta) := \sup_{\substack{u \in C_c^{0,1}(\mathbb{R}^k \setminus \{0\} \times \mathbb{R}^{N-k}), \\ u \neq 0}} \frac{\int_{\mathbb{R}^N} \frac{|u|^{2_t^*}}{|x'|^t}}{\left(\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{u^2(x)}{|x'|^{2s}} dx \right)^{\frac{2_t^*}{2}}}. \quad (1.6)$$

Then, the supremum is achieved in $\dot{\mathcal{H}}_\beta^s(\mathbb{R}^N)$, if $0 < \beta < \mathcal{C}_{N,k,s}$, where the space $\dot{\mathcal{H}}_\beta^s(\mathbb{R}^N)$ is defined in Section 2.

Theorem 1.2. *The following infimum*

$$\kappa_k^t := \inf_{\substack{u \in \dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N), \\ u \neq 0}} = \frac{\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy}{\left(\int_{\mathbb{R}^N} \frac{|u|^{2_t^*}}{|x'|^{t+2_t^*\alpha}} dx \right)^{\frac{2}{2_t^*}}} \quad (1.7)$$

is achieved in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$. Here, $2_t^ = \frac{2(N-t)}{N-2s}$, $0 \leq t < 2s$, and $\alpha := (k-2s)/2$ and the space $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$ is defined in the Definition 2.1.*

We remark that, Theorem 1.1 settles the issue of existence of solution of (1.1), for $0 < \beta < \mathcal{C}_{N,k,s}$. Whereas, using a combination of Theorem 1.2 and Theorem 3.4, we can conclude that there exists a nontrivial solution of (1.1), for the case $\beta = \mathcal{C}_{N,k,s}$. To prove

Theorem 1.1, we have used an improved version of fractional Sobolev inequality derived in Proposition 2.3 originated in [31]. On the other hand, as pointed out in Section 4.2, the space $\dot{\mathcal{H}}_{\mathcal{C}_{N,k,s}}^s(\mathbb{R}^N)$ is much bigger than the space $\dot{\mathcal{H}}_{\beta}^s(\mathbb{R}^N)$, which is nothing but the usual homogeneous fractional Sobolev space $\dot{H}^s(\mathbb{R}^N)$, in the case of $0 < \beta < \mathcal{C}_{N,k,s}$. Because of this fact, we could not use Proposition 2.3. We also note that, in this case, we cannot use extension method, derived in [7], to convert (1.1) into a local one. Rather, by using Ekeland's variation principle, we were able to conclude that, the approximate solutions cannot concentrate near the singular set. However, since we are in a non local setup, so we faced a natural difficulty when we tried to cut off the approximate solutions. In this context, let us mention a paper by Ghoussoub, and Shakerian [20], where they used Ekeland's variation principle to get the existence of solution of a nonlocal equation, similar to (1.1). But, their arguments were based on extension technique. For a comprehensive study for the case $k = 1$, see [30].

Next, we prove some qualitative properties of solutions of (1.1) by means of following theorems.

Theorem 1.3. *Let u satisfies (1.1) weakly. Then, for any $0 < \beta \leq \mathcal{C}_{N,k,s}$, there exists a unique $\tilde{\alpha} \in (0, \frac{k-2s}{2}]$ given by (3.2), such that the following holds:*

$$0 \leq u(x) \leq \frac{C}{|x'|^{\tilde{\alpha}} (1 + |x|^{N-2s-2\tilde{\alpha}})}, \quad \forall x \in \mathbb{R}_k^N, \quad (1.8)$$

where $C > 0$ is constant, depends on u but independent of $x \in \mathbb{R}_k^N$ and $\mathbb{R}_k^N := (\mathbb{R}^k \setminus \{0\}) \times \mathbb{R}^{N-k}$. Moreover, if $0 < \beta < \mathcal{C}_{N,k,s}$, then $u \in C^\infty(\mathbb{R}_k^N)$.

Theorem 1.4. *For $0 < \beta \leq \mathcal{C}_{N,k,s}$ and $0 < s < 1$, any $u \in C_{loc}^{0,1}(\mathbb{R}_k^N)$ satisfying (1.1) is cylindrically symmetric i.e. radial with respect to $x' \in \mathbb{R}^k$ and there exist $x''_0 \in \mathbb{R}^{N-k}$ such that for any fixed $x' \neq 0$, $u(x', x'')$ is radial in the second variable with respect to x''_0 .*

We have used moving plane method to prove Theorem 1.4. Along with other hurdles, applying moving plane method in the non-local setup is inconvenient due to inadequacy of any direct small measure type lemma which was observed, in the local case, by Varadhan and successfully disseminated by Nirenberg and Berestycki [4]. As observed in [26], to prove such small measure type lemma, we could try to use the test function $w_\lambda = (u - u_\lambda)^+$, where u_λ denotes the usual reflexion of u along the hyperplane $T_\lambda = \{x_i = \lambda\}$. But since we are in non local set up, as noticed in [10], [14], the right test function should be an odd reflexion of w_λ along T_λ . However, in our set up we faced difficulty in showing w_λ belongs to right space simply because of the following reason. When $0 < \beta < \mathcal{C}_{N,k,s}$, to show the odd extension of w_λ belongs to $\dot{H}^s(\mathbb{R}^N)$, we need to show that

$$\int_{\{x_i < \lambda\}} \frac{w_\lambda^2(x)}{(\lambda - x_i)^{2s}} dx < \infty,$$

which follows by a fractional Hardy inequality proved in [5], when $s \neq \frac{1}{2}$ and $0 < s < 1$. But, when $s = \frac{1}{2}$, the best constant of the fractional Hardy inequality is zero. On the other hand, when $\beta = \mathcal{C}_{N,k,s}$, due to unavailability of right Hardy type inequality, the situation becomes more complex and it is not clear whether we can use the odd reflexion of w_λ as a test function or not. We have avoided this difficulty by approximating w_λ properly and using the precise bound on u , derived in Theorem 1.3.

Finally, let us describe the plan of this article. In Section 2, we will introduce the notations and all the function spaces used in this article. We will also recall some of the known results used in the proofs. Section 3 will be devoted to the proofs of inequality (1.2)

and (1.5). Section 4 contains the proofs of Theorem 1.1 and 1.2. Also, Section 5 and 6 contains the proofs of Theorem 1.3 and 1.4 respectively. Finally, in the Appendix we have proved the Lemma 2.1.

2. NOTATIONS AND PRELIMINARIES

Notations: We will denote the projection of a point $x \in \mathbb{R}^N$ to \mathbb{R}^k and \mathbb{R}^{N-k} by x' and x'' respectively.

For any $l \in \mathbb{N}$ and any $z \in \mathbb{R}^l$ we denote the l dimensional ball of radius $R > 0$ centered at z by $B_R^l(z)$.

For $2 \leq k \leq N - 2$, \mathbb{R}_k^N stands for the set $(\mathbb{R}^k \setminus \{0\}) \times \mathbb{R}^{N-k}$.

To avoid confusion, we clarify that $2^* := \frac{2N}{N-2s}$ and $\alpha := \frac{k-2s}{2}$.

2.1. Definitions and Different Notions Of Solution. In this section we will define different function spaces to be used. We will also define different notions of solution.

Definition 2.1.

- (i) We define $\dot{\mathcal{H}}_{\beta}^s(\mathbb{R}^N)$ as the completion of $C_c^{0,1}(\mathbb{R}^k \setminus \{0\} \times \mathbb{R}^{N-k})$ under the following norm

$$[u]_{s,\beta,\mathbb{R}^N}^2 := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{u^2(x)}{|x'|^{2s}} dx.$$

Here $0 \leq \beta \leq \mathcal{C}_{N,k,s}$ and $\mathcal{C}_{N,k,s}$ is the best constant of the Fractional Hardy-Sobolev-Maz'ya inequality.

- (ii) We also define the space $\dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$ as the completion of $C_c^{0,1}(\mathbb{R}^k \setminus \{0\} \times \mathbb{R}^{N-k})$ under the following norm

$$[[u]]_{s,\tilde{\alpha},\mathbb{R}^N}^2 := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy,$$

where $0 < \tilde{\alpha} \leq \alpha := (k - 2s)/2$.

- (iii) We recall, $\dot{H}^s(\mathbb{R}^N)$ is the completion of $C_c^\infty(\mathbb{R}^N)$ under the following norm

$$[u]_{s,\mathbb{R}^N}^2 := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s}} dx dy.$$

One can easily prove that the following characterization of $\dot{H}^s(\mathbb{R}^N)$. See [11]

$$\dot{H}^s(\mathbb{R}^N) := \{u \in L^{2^*}(\mathbb{R}^N) : [u]_{s,\mathbb{R}^N} < \infty\}.$$

- (iv) For any domain $\Omega \subset \mathbb{R}_+^{N+1}$ we recall

$$H^1(\Omega, 1 - 2s) := \{U \in L^2(\Omega, 1 - 2s) : |\nabla U| \in L^2(\Omega, 1 - 2s)\},$$

where, $L^2(\Omega, 1 - 2s) := \{U : \Omega \rightarrow \mathbb{R} \text{ measurable} \mid \int_{\Omega} t^{1-2s} U^2(x, t) dx dt < \infty\}$

- (v) We recall, $L_s(\mathbb{R}^N) := \left\{ f : \mathbb{R}^N \rightarrow \mathbb{R} \text{ measurable} \mid \int_{\mathbb{R}^N} \frac{|f|}{1+|x|^{N+2s}} dx < \infty \right\}$

Finally, let us state the following lemma regarding the precise representation of $\dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$.

Lemma 2.1. *For $0 < s < 1$, $k \geq 2$ and $0 < \tilde{\alpha} \leq \alpha$, we have the following representation of $\dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$.*

$$\dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N) = \left\{ u \in L^{2^*} \left(\mathbb{R}^N; \frac{1}{|x'|^{\tilde{\alpha}2^*}} \right) : \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2 dx dy}{|x - y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} < \infty \right\},$$

where $L^{2^*} \left(\mathbb{R}^N; \frac{1}{|x'|^{\alpha 2^*}} \right)$ consists of all measurable function f such that $\frac{f}{|x'|^{\alpha}} \in L^{2^*}(\mathbb{R}^N)$.

We have proved the lemma in the Appendix. We recall, for f belonging to the Schwartz class the fractional laplacian can be defined by the following integral representation. See [9]

$$(-\Delta)^s f(x) := c_{N,s} P.V. \int_{\mathbb{R}^N} \frac{f(x) - f(y)}{|x - y|^{N+2s}} dy, \text{ for } x \in \mathbb{R}^N,$$

where $c_{N,s} = 2^{2s} \pi^{-N/2} \frac{\Gamma(\frac{N+2s}{2})}{|\Gamma(-s)|}$. Next, we need the following two definitions.

Definition 2.2. For $0 < \beta \leq \mathcal{C}_{N,k,s}$, we say that, $u \in \dot{\mathcal{H}}_{\beta}^s(\mathbb{R}^N)$ is a weak solution of (1.1) if for every $\psi \in \dot{\mathcal{H}}_{\beta}^s(\mathbb{R}^N)$, we have

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))(\psi(x) - \psi(y))}{|x - y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{u(x)\psi(x)}{|x'|^{2s}} dx = \int_{\mathbb{R}^N} \frac{u^{2^*-1}(x)\psi(x)}{|x'|^t} dx$$

Definition 2.3. Let $\Omega \subset \mathbb{R}^N$ be any open set and $\mathcal{D}'(\Omega)$ denotes the space of all distributions over Ω . Assume that, $u \in L_s(\mathbb{R}^N)$ and $f \in \mathcal{D}'(\Omega)$. Then we say

$$(-\Delta)^s u(\geq) = (\leq) f, \text{ in the distributional sense,}$$

if for any nonnegative $\phi \in C_c^\infty(\Omega)$ we have

$$\int_{\mathbb{R}^N} u(x) (-\Delta)^s \phi(x) dx (\geq) = (\leq) \langle f, \phi \rangle_{\Omega},$$

$\langle f, \phi \rangle_{\Omega}$ denotes the action of f on ϕ .

2.2. Some Known Results. In this section, we will recall some known results.

Master Inequality: We recall the following integral version of Pólya-Szegő inequality. See [3].

Theorem 2.2. Let $f, g \in C(\mathbb{R}^N)$ be non negative vanishing at infinity i.e. they satisfy, $|\{f > t\}|_N < \infty, \forall t > \text{ess inf } f$ and $|\{g > t\}|_N < \infty, \forall t > \text{ess inf } g$, where for any measurable subset A of \mathbb{R}^N , $|A|_N$ denotes the N -dimensional Lebesgue measure of A . Then, for any fixed $\phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ increasing, convex and $K : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ decreasing, we have the following inequality

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \phi(|f^*(x) - g^*(y)|) K(|x - y|) dx dy \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \phi(|f(x) - g(y)|) K(|x - y|) dx dy, \quad (2.1)$$

where f^* and g^* denotes the Schwarz symmetrization of f, g respectively.

We need the following improved version of Sobolev inequality.

Proposition 2.3. Let $0 \leq t < 2s$ and $u \in \dot{H}(\mathbb{R}^N)$. Then, there exist constants $C, \theta_1, \theta_2 > 0$ independent of u such that

$$\int_{\mathbb{R}^N} \frac{|u|^{2^*}}{|x'|^t} dx \leq C [u]_{s, \mathbb{R}^N}^{\theta_1} \|u\|_{\mathcal{L}^{2, N-2s}}^{\theta_2}, \quad (2.2)$$

where

$$\|u\|_{\mathcal{L}^{2, N-2s}}^2 := \sup_{R>0, x \in \mathbb{R}^N} \frac{R^{N-2s}}{|B_R^N(x)|} \int_{B_R^N(x)} |u|^2 dy.$$

Proof. We divide the proof in the following two cases.

Case 1: $t = 0$ In this case, (2.2) was proved by Palatucci and Pisante in [31]. Their inequality states as follows: there exists a constant $C > 0$ such that for any $u \in \dot{H}^s(\mathbb{R}^N)$

$$\|u\|_{L^{2^*}(\mathbb{R}^N)} \leq C [u]_{s, \mathbb{R}^N}^\theta \|u\|_{\mathcal{L}^{2, N-2s}}^{1-\theta} \quad (2.3)$$

where $\frac{2}{2^*} \leq \theta < 1$.

Case 2: $0 < t < 2s$. Using Hölder inequality we get

$$\int_{\mathbb{R}^N} \frac{|u|^{2^*}}{|x'|^t} dx \leq \left(\int_{\mathbb{R}^N} |u|^{2^*} \right)^{\frac{2^*-2}{2^*}} \left(\int_{\mathbb{R}^N} \frac{|u|^{\frac{2 \cdot 2^*}{2^*-2^*+2}}}{|x'|^{\frac{2^* t}{2^*-2^*+2}}} \right)^{\frac{2^*-2^*+2}{2^*}}. \quad (2.4)$$

Now let $\xi := \frac{2^* t}{2^*-2^*+2}$. Then clearly $0 < \xi < 2s$ and $2_\eta^* = \frac{2^* 2}{2^*-2^*+2}$. Hence using (2.3), (2.4) and (1.5) we get the desired inequality (2.2) with $\theta_1 = (2_t^* - 2)\theta + \frac{2^*}{2} \left(1 - \frac{2_t^*-2}{2^*}\right)$ and $\theta_2 = (1 - \theta)(2_t^* - 2)$, where $\frac{2^*}{2} < \theta < 1$ same as in the inequality (2.3). \square

3. FRACTIONAL HARDY-MAZ'YA AND HARDY-SOBOLEV-MAZ'YA INEQUALITY

In this section, we will give two proofs of fractional Hardy-Maz'ya inequality (1.2). While in the first method, we get (1.2) with best possible constant, in the second method, we get the inequality with a rough constant. Also, as a consequence of the results, derived using both the methods we will be able to prove (1.5).

3.1. Ground State Representation and the Fractional Hardy-Maz'ya inequality.

In this subsection, we will derive an appropriate ground state representation. Similar representation was proved in [16] to derive the fractional Hardy inequality. In fact, we will use a few results derived in [16]. For reader's convenience, let us recall their result. Before that, we need to recall the following assumption.

Assumption 1: Let $\Omega \subset \mathbb{R}^N$ be any open set. We also assume that w is an almost everywhere positive measurable function in Ω and there exists a family of measurable function k_ϵ , $\epsilon > 0$ on $\Omega \times \Omega$ satisfying $k_\epsilon(x, y) = k_\epsilon(y, x)$, $0 \leq k_\epsilon(x, y) \leq k(x, y)$, and

$$\lim_{\epsilon \rightarrow 0} k_\epsilon(x, y) = k(x, y), \text{ for a.e. } x, y \in \Omega.$$

Moreover, the integrals

$$V_\epsilon(x) := \frac{2}{w(x)} \int_{\Omega} (w(x) - w(y)) k_\epsilon(x, y) dy$$

are absolutely convergent for a.e. $x \in \Omega$ and belong to $L^1_{loc}(\Omega)$. In addition to this, we assume that $V := \lim_{\epsilon \rightarrow 0} V_\epsilon$ exists weakly in $L^1_{loc}(\Omega)$.

Proposition 3.1 (Frank and Seiringer). *Under the Assumption 1, for any $u \in C_c^\infty(\Omega)$ we write $v := \frac{u}{w}$ and assume*

$$E[u] := \int_{\Omega} \int_{\Omega} |u(x) - u(y)|^2 k(x, y) dy dx < \infty,$$

$$E_w[v] := \int_{\Omega} \int_{\Omega} |v(x) - v(y)|^2 w(x)w(y)k(x, y) dy dx < \infty \text{ and}$$

$$\int_{\Omega} V^+ |u|^2 dx < \infty.$$

Then $E[u] - \int_{\mathbb{R}^N} V(x) |u(x)|^2 dx = E_w[v]$.

As a particular example, we take $\Omega = (\mathbb{R}^k \setminus \{0\}) \times \mathbb{R}^{N-k}$, $w(x) := 1/|x'|^{\tilde{\alpha}}$, for $0 < \tilde{\alpha} \leq \frac{k-2s}{2}$, $k(x, y) := \frac{1}{|x-y|^{N+2s}}$, $V(x) := \mathcal{C}_{N,k,s}(\tilde{\alpha}) \frac{1}{|x'|^{2s}}$, where $\mathcal{C}_{N,k,s}(\tilde{\alpha})$ is defined by $\mathcal{C}_{N,k,s}(\tilde{\alpha}) := \bar{\mathcal{C}}_{k,s}(\tilde{\alpha}) \int_{\mathbb{R}^{N-k}} \frac{dy''}{(1+|y''|^2)^{\frac{N+2s}{2}}}$ and the function $\bar{\mathcal{C}}_{k,s}(\tilde{\alpha})$ is defined in Lemma 3.3 below. We denote $\mathcal{C}_{N,k,s} := \mathcal{C}_{N,k,s}((k-2s)/2)$. We also take, $k_\epsilon(x, y) := \frac{1}{|x-y|^{N+2s}} \chi_{\{|x'-|y'|\|>\epsilon\}}$. Then we have the following lemma.

Lemma 3.2. *The following limit converges uniformly for x from compact sets in \mathbb{R}_k^N .*

$$2 \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^N} (w(x) - w(y)) k_\epsilon(x, y) dy = \mathcal{C}_{N,k,s}(\tilde{\alpha}) \frac{w(x)}{|x'|^{2s}}.$$

Proof. It is enough to notice that the following identity is true. The rest will follow from Lemma 3.3 below.

$$2 \int_{\mathbb{R}_k^N} (w(x) - w(y)) k_\epsilon(x, y) dy = I_{N,k,s} \int_{\|x'-|y'|\|>\epsilon} \frac{w(x') - w(y')}{|x' - y'|^{k+2s}} dy,$$

where $I_{N,k,s} := \int_{\mathbb{R}^{N-k}} \frac{dy''}{(1+|y''|^2)^{\frac{N+2s}{2}}} < \infty$. This proves the lemma. \square

Lemma 3.3. *One has uniformly for x from compact sets in $\mathbb{R}^k \setminus \{0\}$*

$$2 \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^k} \frac{(w(x') - w(y'))}{|x' - y'|^{k+2s}} dy = \bar{\mathcal{C}}_{k,s}(\tilde{\alpha}) \frac{w(x')}{|x'|^{2s}},$$

where

$$\begin{aligned} \bar{\mathcal{C}}_{k,s}(\tilde{\alpha}) &:= 2 \int_0^1 r^{2s-1} |1 - r\tilde{\alpha}|^2 \Phi_{k,s}(r) dr, \\ \Phi_{k,s}(r) &:= \left| S^{k-2} \right| \int_{-1}^1 \frac{(1-t^2)^{\frac{k-3}{2}} dt}{(1-2rt+r^2)^{\frac{k+2s}{2}}}, \quad 0 < \tilde{\alpha} \leq \frac{k-2s}{2}, \text{ and } k \geq 2. \end{aligned} \quad (3.1)$$

The above lemma was proved in [16] (Lemma 3.1), in the case of $\tilde{\alpha} = (k-2s)/2$. But, it is easy to see that the same proof will work even for $0 < \tilde{\alpha} < \frac{k-2s}{2}$. Notice that

$$\bar{\mathcal{C}}_{k,s}(0) = 0, \quad \bar{\mathcal{C}}_{k,s}\left(\frac{k-2s}{2}\right) = 2\pi^{k/2} \frac{\Gamma^2((k+2s)/4)}{\Gamma^2((k-2s)/4)} \frac{|\Gamma(-s)|}{\Gamma((k+2s)/2)}$$

i.e. the best constant of fractional Hardy inequality in dimension k . Also, $\bar{\mathcal{C}}_{k,s}(\tilde{\alpha})$ is strictly increasing and continuous in $[0, (k-2s)/2]$. So, for any $0 \leq \beta \leq \mathcal{C}_{N,k,s}$ there exists unique $\tilde{\alpha} \in [0, \frac{k-2s}{2}]$ such that

$$\beta = \mathcal{C}_{N,k,s}(\tilde{\alpha}) = \bar{\mathcal{C}}_{k,s}(\tilde{\alpha}) \int_{\mathbb{R}^{N-k}} \frac{dy''}{(1+|y''|^2)^{\frac{N+2s}{2}}}, \quad (3.2)$$

where $\bar{\mathcal{C}}_{k,s}(\tilde{\alpha})$ is defined in (3.1). Hence, summarizing the above discussion we have the following theorem.

Theorem 3.4 (Ground State Representation). *Let $u \in C_c^{0,1}(\mathbb{R}^k \setminus \{0\}) \times \mathbb{R}^{N-k}$, $k \geq 2$, and $0 < s < 1$. Then for any $0 < \beta \leq \mathcal{C}_{N,k,s}$ there exist a unique $0 < \tilde{\alpha} \leq (k-2s)/2$ such that*

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x-y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{u^2(x)}{|x'|^{2s}} dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|v(x) - v(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy. \quad (3.3)$$

Here, $v(x) = |x'|^{\tilde{\alpha}} u(x)$, $\mathcal{C}_{N,k,s} = \bar{\mathcal{C}}_{k,s} \int_{\mathbb{R}^{N-k}} \frac{dy''}{(1+|y''|^2)^{\frac{N+2s}{2}}}$ and $\bar{\mathcal{C}}_{k,s}$ is the best constant of the fractional Hardy inequality for dimension k . Moreover, $\tilde{\alpha} = (k-2s)/2$ whenever $\beta = \mathcal{C}_{N,k,s}$.

As a consequence of the above ground state representation, we have the fractional Hardy-Maz'ya inequality (1.2). Also, one can follow the same lines of Frank and Seiringer [17], to conclude that $\mathcal{C}_{N,k,s}$ is actually the best constant.

3.2. John Domain and Fractional Hardy-Sobolev-Maz'ya inequality. To prove the fractional Hardy-Sobolev-Maz'ya inequality, we will use the fact, \mathbb{R}_k^N is a John domain for $2 \leq k \leq N-2$ and a recent result by Dyda, Lehrbäck and Vähäkangas [12]. For reader's convenience we will state their result below. First, let us recall some definitions.

Definition 3.1 (Assouad dimension). For $D \subset \mathbb{R}^N$, the Assouad dimension denoted by $\dim_A(D)$ is the infimum of all exponent $\beta \geq 0$, for which there is a constant $C \geq 1$, such that for every $x \in D$ and every $0 < r < R$, the set $D \cap B_R^N(x)$ can be covered by at most $C \left(\frac{R}{r}\right)^\beta$ balls of radius r .

Definition 3.2 (John Domain). A domain $\Omega \subsetneq \mathbb{R}^N$, with $N \geq 2$, is a c -John domain, for $c \geq 1$, if each pair of points $x_1, x_2 \in \Omega$ can be joined by a rectifiable arc length parametrized curve $\gamma : [0, l] \rightarrow \Omega$ satisfying $\text{dist}(\gamma(t), \Omega) \geq \min\{t, l-t\}/c$ for every $t \in [0, l]$.

Theorem 3.5 (Dyda, Lehrbäck and Vähäkangas). Assume that $\Omega \subsetneq \mathbb{R}^N$ is an unbounded c -John Domain, $0 < s < 1$ and $1 < p \leq q \leq \frac{Np}{N-ps}$. Let $\beta \in \mathbb{R}$ be such that

$$\dim_A(\partial\Omega) < \min \left\{ \frac{q}{p} (N - sp + \beta), N - \frac{\beta}{p-1} \right\}.$$

Then for any $\tau \in (0, 1)$ there exists a constant $C = C(\beta, \tau, N, s, p, c) > 0$ such that for any $u \in \cup_{1 \leq r < \infty} L^r(\Omega)$ we have

$$\left(\int_{\Omega} |u(x)|^q \delta_{\partial\Omega}^{\frac{q}{p}(N-sp+\beta)-N}(x) dx \right)^{\frac{p}{q}} \leq C \int_{\Omega} \int_{B_{\tau\delta_{\partial\Omega}(x)}^N(x)} \frac{|u(x) - u(y)|^p}{|x-y|^{N+ps}} dy \delta_{\partial\Omega}^{\beta}(x) dx,$$

where $\delta_{\partial\Omega}^{\beta}(x) := \text{dist}(x, \partial\Omega)$.

Choosing $\beta = 2s - k$, $p = 2$, $q := \frac{2(N-t)}{N-2s}$, $0 \leq t < 2s$, $0 < s < 1$, $\Omega = \mathbb{R}^k \setminus \{0\} \times \mathbb{R}^{N-k}$, $2 \leq k \leq N-2$ in Theorem 3.5 and using (3.3) we get the inequality (1.5). Also notice that, choosing $\beta = 0$ and $q = p = 2$ we get the inequality (1.2) with a rough constant. For future reference, let us clarify that, as a consequence of the above discussion we get the following equivalent version of inequality (1.5).

$$C \left(\int_{\mathbb{R}^N} \frac{|u(x)|^{2_t^*}}{|x'|^{t+\tilde{\alpha}2_t^*}} dx \right)^{\frac{2}{2_t^*}} \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy, \quad \forall u \in C_c^{0,1}(\mathbb{R}_k^N), \quad (3.4)$$

where, $0 < \tilde{\alpha} \leq \frac{k-2s}{2}$, $0 \leq t < 2s$, $0 < s < 1$, $2 \leq k \leq N-2$, $2_t^* = \frac{2(N-t)}{N-2s}$ and $C > 0$ is constant depending only on N, s, k, t and $\tilde{\alpha}$.

4. EXISTENCE OF SOLUTION

In this section, we will prove Theorem 1.1 and Theorem 1.2.

4.1. Proof of Theorem 1.1.

Proof. It is evident that $\dot{\mathcal{H}}_\beta^s(\mathbb{R}^N) = \dot{H}^s(\mathbb{R}^N)$ for $\beta < \mathcal{C}_{N,k,s}$. In fact, $[u]_{s,\beta,\mathbb{R}^N}$ is an equivalent norm in $\dot{H}^s(\mathbb{R}^N)$. For $u, v \in \dot{H}^s(\mathbb{R}^N)$, let us define

$$E[u, v] := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{uv}{|x'|^{2s}} dx.$$

We will prove the theorem by dividing into two cases.

Case I : $t=0$. In this case we have

$$S_k(\beta) := S_k^0(\beta) = \sup_{u \in \dot{H}(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} |u|^{2^*}}{(E[u, u])^{\frac{2^*}{2}}}.$$

Let $\{u_m\}$ be any maximizing sequence. If necessary by normalizing we can assume $E[u_m, u_m] = 1$, $\forall m \in \mathbb{N}$. In addition to this, we could also assume that u_m 's are nonnegative and radially symmetric in the first variable. This follows from inequality (2.1) stated in Theorem 2.2.

For now our aim is to show that, there exist $R_m > 0$ and $x_m'' \in \mathbb{R}^{N-k}$ such that the sequence $\{v_m\}$ defined by

$$v_m(x', x'') := R_m^{\frac{N-2s}{2}} u_m(R_m x', R_m x'' + x_m'')$$

weakly converges to some non zero $v \in \dot{H}(\mathbb{R}^N)$. Using the fact that $\|u_m\|_{L^{2^*}(\mathbb{R}^N)}^2 \rightarrow S_k(\beta)$ and (2.3) we have that there exists $R_m > 0$ and $x_m \in \mathbb{R}^N$ such that

$$\frac{1}{R_m^{2s}} \int_{B_{R_m}^N(x_m)} |u_m(y)|^2 dy \geq C > 0,$$

for some constant C independent of m . Hence we have the following:

$$\frac{1}{R_m^{2s}} \int_{B_{R_m}^{N-k}(x_m'')} \int_{B_{R_m}^k(x_m'')} |u_m(y', y'')|^2 dy' dy'' \geq C > 0, \quad \forall m \in \mathbb{N}. \quad (4.1)$$

Now let us define

$$v_m(y', y'') := R_m^{\frac{N-2s}{2}} u_m(R_m y', R_m y'' + x_m'').$$

Then from (4.1) we have

$$\int_{B_1^{N-k}(0)} \int_{B_1^k(\bar{x}_m'')} |v_m(y)|^2 dy \geq C > 0, \quad \forall m \in \mathbb{N}, \quad (4.2)$$

where $\bar{x}_m' := \frac{x_m'}{R_m}$. Clearly, $E[v_m, v_m] = E[u_m, u_m] = 1$. Hence by compactness we can assume the following: upto a subsequence

- (i) $v_m \rightharpoonup v$ in $\dot{H}^s(\mathbb{R}^N)$,
- (ii) $v_m \rightarrow v$ in $L_{loc}^2(\mathbb{R}^N)$ and,
- (iii) $v_m \rightarrow v$ a.e.

We will prove that $v \neq 0$ by considering the following two cases.

Case I: Upto a subsequence $|\bar{x}_m'| \rightarrow \infty$.

Case II: $\{\bar{x}_m'\}$ is bounded sequence in \mathbb{R}^k .

In the first case, by using radial symmetry of v_m in the first variable we have

$$\int_{B_1^{N-k}(0)} \int_{B_2^k(0)} |v_m(y', y'')|^2 dy' dy'' \geq \int_{B_1^{N-k}(0)} \int_{B_1^k(0)} |v_m(y' + \bar{x}_m', y'')|^2 dy' dy''$$

$$= \int_{B_1^{N-k}(0)} \int_{B_1^k(\bar{x}'_m)} |v_m(y', y'')|^2 dy' dy'' \geq C > 0.$$

Hence by passing to the limit we get

$$\int_{B_1^{N-k}(0)} \int_{B_2^k(0)} |v(y', y'')|^2 dy' dy'' \geq C > 0,$$

which clearly shows that $v \neq 0$.

In the second case, upto a subsequence $\bar{x}'_m \rightarrow x'_0$. Then $B_1^k(\bar{x}'_m) \subset \Omega$, for sufficiently large open bounded set Ω in \mathbb{R}^k . So, using (4.2) we get

$$\int_{B_1^{N-k}} \int_{\Omega} |v_m|^2 dy \geq \int_{B_1^{N-k}} \int_{B_1^k(\bar{x}'_m)} |v_m|^2 dy \geq C > 0.$$

Then again as before, passing to the limit we conclude that $v \neq 0$. The rest of the proof is fairly standard. For the sake of the completeness, let us add the argument. By weak convergence of v_m to v we have

$$E[v, v] + E[v_m - v, v_m - v] = 1 + o(1),$$

as $m \rightarrow \infty$. Also by scale invariance and Brezis-Lieb lemma [6] we have

$$\begin{aligned} S_k(\beta) &= \lim_{m \rightarrow \infty} \int_{\mathbb{R}^N} |v_m|^{2^*} dy \\ &= \lim_{m \rightarrow \infty} \left[\int_{\mathbb{R}^N} |v|^{2^*} dy + \int_{\mathbb{R}^N} |v_m - v|^{2^*} dy \right] \\ &\leq S_k(\beta) (E[v, v])^{\frac{2^*}{2}} + S_k(\beta) \left(\lim_{m \rightarrow \infty} E[v_m - v, v_m - v] \right)^{\frac{2^*}{2}} \\ &\leq S_k(\beta) \left(E[v, v] + \lim_{m \rightarrow \infty} E[v_m - v, v_m - v] \right)^{\frac{2^*}{2}} \leq S_k(\beta). \end{aligned}$$

Hence we must have equality everywhere in the above estimate. So, $S_k(\beta) = \frac{\int_{\mathbb{R}^N} |v|^{2^*} dy}{(E[v, v])^{\frac{2^*}{2}}}$.

This proves the Case I, i.e. the existence of maximizer for $t = 0$.

Case II: $0 < t < 2s$. A careful inspection of the preceding proof yields that we only need the inequality (2.2), which was proved in Proposition 2.3, to conclude the proof. \square

Remark 4.1. In the proof of Theorem 1.1, we have essentially used the fact that $[\cdot]_{s, \mathbb{R}^N}$ is an equivalent norm in $\dot{\mathcal{H}}_{\beta}^s(\mathbb{R}^N)$ along with the improved Sobolev inequality (2.2). Hence we remark that, using the same arguments given there, we can conclude, $\bar{S}_k^t := (1/S_k^t(0))^{\frac{2}{2^*}}$ is attained by a nontrivial and nonnegative function $u_0 \in \dot{H}^s(\mathbb{R}^N)$. Hence, $\kappa_k^t < \bar{S}_k^t$.

4.2. Existence of Solution For $\beta = \mathcal{C}_{N, k, s}$. Firstly, we observe that if $\phi \in C_c^\infty(\mathbb{R}^N)$ with $\phi = 1$ in $B_1^N(0)$, then $\phi \in \dot{\mathcal{H}}^{s, \alpha}(\mathbb{R}^N)$ (See Lemma 2.1 of Appendix). Then, by ground state representation (3.3), we conclude that $\frac{\phi}{|x'|^\alpha} \in \dot{\mathcal{H}}_{\beta}^s(\mathbb{R}^N)$. Since, $\alpha = (k - 2s)/2$, so we can see that $\frac{\phi}{|x'|^\alpha} \notin \dot{H}^s(\mathbb{R}^N)$. This shows that, $\dot{\mathcal{H}}_{\beta}^s(\mathbb{R}^N)$ is much bigger than $\dot{H}^s(\mathbb{R}^N)$. Because of this fact, we were not able to follow the same arguments as in the case of $\beta < \mathcal{C}_{N, k, s}$. We have used Ekeland's Variation principle to prove the Theorem 1.2. We start by the following compactness result.

Lemma 4.2. *Let Ω be a bounded open set in \mathbb{R}^N . Then the following inclusion is compact.*

$$\dot{\mathcal{H}}^{s, \alpha}(\mathbb{R}^N) \hookrightarrow L^2 \left(\Omega; \frac{1}{|x'|^{2\alpha}} \right),$$

where $L^2\left(\Omega; \frac{1}{|x'|^{2\alpha}}\right)$ is the set of all measurable function $f : \mathbb{R}^N \rightarrow \mathbb{R}$ such that $\frac{f}{|x'|^\alpha} \in L^2(\Omega)$.

Proof. Let $u \in \dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$. Then by Hölder inequality we have

$$\begin{aligned} \int_{\Omega} \frac{u^2}{|x'|^{2\alpha}} &\leq |\Omega|^{\frac{2s}{N}} \left(\int_{\Omega} \frac{|u|^{2^*}}{|x'|^{\alpha 2^*}} \right)^{\frac{2}{2^*}} \\ &\leq |\Omega|^{\frac{2s}{N}} \frac{1}{\kappa_k^0} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy, \end{aligned} \quad (4.3)$$

where $|\Omega|$ denotes the N -dimensional Lebesgue measure of Ω . Hence, the inclusion, $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N) \hookrightarrow L^2\left(\Omega; \frac{1}{|x'|^{2\alpha}}\right)$ is continuous. To prove the compactness, it is enough to show that $u_m \rightarrow 0$ in $L^2\left(\Omega; \frac{1}{|x'|^{2\alpha}}\right)$, whenever $u_m \in \dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$ and u_m converges to zero weakly in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$, as well as in $L^2\left(\Omega; \frac{1}{|x'|^{2\alpha}}\right)$. To prove this, we consider, $\phi_\epsilon \in C^\infty(\mathbb{R}^k; [0, 1])$ such that

$$\phi_\epsilon = \begin{cases} 0, & \text{if } |x'| < \epsilon \\ 1, & \text{if } |x'| > 2\epsilon. \end{cases}$$

and define $\Omega_\epsilon := \Omega \cap B_\epsilon^k(0)$. Without loss of generality, we could assume that $\Omega \setminus \Omega_\epsilon$ is a domain with Lipschitz boundary. Also, let $M > 0$ be such that $[[u_m]]_{s,\alpha,\mathbb{R}^N}^2 < M$ and $\left\| \frac{u_m}{|x'|^\alpha} \right\|_{L^2(\Omega)} < M$. Now, note that,

$$\begin{aligned} \int_{\Omega \setminus \Omega_\epsilon} \int_{\Omega \setminus \Omega_\epsilon} \frac{|u_m(x) - u_m(y)|^2}{|x - y|^{N+2s}} dx dy &\leq C \int_{\Omega \setminus \Omega_\epsilon} \int_{\Omega \setminus \Omega_\epsilon} \frac{|u_m(x) - u_m(y)|^2}{|x - y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy \leq C \\ \text{and } \int_{\Omega \setminus \Omega_\epsilon} |u_m|^2 dx &\leq C \int_{\Omega} \frac{u_m^2}{|x'|^{2\alpha}} \leq C, \end{aligned}$$

where (and for the rest of the proof) $C > 0$ is a constant depending on N, s, k, Ω and M . So, u_m is bounded sequence in $H^s(\Omega \setminus \Omega_\epsilon)$. Hence, by compactness, (see [9]) $u_m \rightarrow u$ in $L^2(\Omega \setminus \Omega_\epsilon)$. By weak convergence of u_m to 0 in $L^2\left(\Omega; \frac{1}{|x'|^{2\alpha}}\right)$ we conclude that $u = 0$ and

$$\int_{\Omega} \frac{\phi_\epsilon^2 u_m^2}{|x'|^{2\alpha}} dx = \frac{1}{\epsilon^{2\alpha}} o(1), \text{ as } m \rightarrow \infty.$$

Also, by (4.3)

$$\int_{\Omega} |(1 - \phi_\epsilon) u_m|^2 \frac{dx}{|x'|^{2\alpha}} \leq C |\Omega_\epsilon|^{\frac{2s}{N}}.$$

Since,

$$\int_{\Omega} u_m^2 \frac{dx}{|x'|^{2\alpha}} \leq C \left[\int_{\Omega} \frac{\phi_\epsilon^2 u_m^2}{|x'|^{2\alpha}} dx + \int_{\Omega} |(1 - \phi_\epsilon) u_m|^2 \frac{dx}{|x'|^{2\alpha}} \right],$$

so first letting $m \rightarrow \infty$ and then letting $\epsilon \rightarrow 0$ in the last inequality, we see that $u_m \rightarrow 0$ in $L^2\left(\Omega; \frac{1}{|x'|^{2\alpha}}\right)$. This proves the lemma. \square

Next, we need the following lemma.

Lemma 4.3. *Let $\{u_m\}$ be a bounded sequence in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$ such that $u_m \rightharpoonup 0$. Then for any $\psi \in C_c^\infty(\mathbb{R}^N)$ with $\text{supp } \psi \subset B_{R_0}^N(0)$, for some $R_0 > 0$, we have*

$$\begin{aligned} & \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|(\psi u_m)(x) - (\psi u_m)(y)|^2}{|x - y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy \\ & \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{((\psi^2 u_m)(x) - (\psi^2 u_m)(y)) (u_m(x) - u_m(y))}{|x - y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy + \sigma_{R,m} + \tilde{\sigma}_R, \end{aligned} \quad (4.4)$$

for any $R > R_0 + 2$ and $m \in \mathbb{N}$. Here, for each R , $\lim_{m \rightarrow \infty} \sigma_{R,m} = 0$ and $\lim_{R \rightarrow \infty} \tilde{\sigma}_R = 0$.

Proof. Since, $R > R_0$, so, $\text{supp } \psi \subset B_{R-2}^N(0)$. Now, note that,

$$\begin{aligned} & ((\psi^2 u_m)(x) - (\psi^2 u_m)(y)) (u_m(x) - u_m(y)) \\ & = |(\psi u_m)(x) - (\psi u_m)(y)|^2 - u_m(x) u_m(y) |\psi(x) - \psi(y)|^2. \end{aligned}$$

So, using $2ab \leq (a^2 + b^2)$, with $a = \frac{|u_m(x)|}{|x'|^\alpha}$ and $b = \frac{|u_m(y)|}{|y'|^\alpha}$, we estimate

$$\begin{aligned} I_m & := \left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u_m(x) u_m(y) |\psi(x) - \psi(y)|^2}{|x - y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy \right| \\ & \leq \int_{\mathbb{R}^N} \frac{u_m^2(x)}{|x'|^{2\alpha}} \int_{\mathbb{R}^N} \frac{|\psi(x) - \psi(y)|^2}{|x - y|^{N+2s}} dy dx \leq I_{m,1} + I_{m,2} \end{aligned} \quad (4.5)$$

where

$$\begin{aligned} I_{m,1} & := \int_{|x| < R} \frac{u_m^2(x)}{|x'|^{2\alpha}} \int_{\mathbb{R}^N} \frac{|\psi(x) - \psi(x - y)|^2}{|y|^{N+2s}} dy dx \text{ and} \\ I_{m,2} & := \int_{|x| > R} \frac{u_m^2(x)}{|x'|^{2\alpha}} \int_{\mathbb{R}^N} \frac{|\psi(x) - \psi(x - y)|^2}{|y|^{N+2s}} dy dx. \end{aligned}$$

Now, since

$$\int_{\mathbb{R}^N} \frac{|\psi(x) - \psi(x - y)|^2}{|y|^{N+2s}} dy \leq C \left[\|\nabla \psi\|_{L^\infty(\mathbb{R}^N)} + \|\psi\|_{L^\infty(\mathbb{R}^N)} \right],$$

for some constant $C > 0$ depending only on N , so using Lemma (4.2) we conclude that, for each $R > 0$, $I_{m,1} = \sigma_{R,m}$, where $\lim_{m \rightarrow \infty} \sigma_{R,m} = 0$. To estimate $I_{m,2}$, we notice that, if $|x| > R$, $R > R_0 + 2$ and $|y| < 1$ then $\psi(x) = \psi(x - y) = 0$. Hence,

$$\begin{aligned} I_{m,2} & = \int_{|x| > R} \frac{u_m^2(x)}{|x'|^{2\alpha}} \int_{|y| > 1} \frac{|\psi(x - y)|^2}{|y|^{N+2s}} dy dx \\ & = \int_{|y| > 1} \frac{1}{|y|^{N+2s}} \int_{|x| > R} \frac{u_m^2(x)}{|x'|^{2\alpha}} |\psi(x - y)|^2 dx dy \end{aligned} \quad (4.6)$$

First using Höder inequality, then using (3.4) and boundedness of u_m in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$, we get from (4.6)

$$I_{m,2} \leq C \int_{|y| > 1} \frac{F_R(y)}{|y|^{N+2s}} dy,$$

where, $F_R(y) := \left(\int_{|x| > R} |\psi(x - y)|^{N/s} dx \right)^{2s/N}$. Then using Dominated Convergence Theorem we have $I_{m,2} \leq \tilde{\sigma}_R$, for all $m \in \mathbb{N}$ and $\lim_{R \rightarrow \infty} \tilde{\sigma}_R = 0$. Hence, using (4.5) we conclude the lemma. \square

Next, lemma shows that if the interior concentration happens, then it will happen away from the singular set.

Proposition 4.4. *Let $\{u_m\}$ be a nonnegative bounded sequence in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$ such that*

- (i) $\int_{\mathbb{R}^N} \frac{|u_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} dx = (\kappa_k^t)^{\frac{2_t^*}{2_t^*-2}}$
- (ii) $L_\alpha^s u_m = \frac{|u_m|^{2_t^*-2} u_m}{|x'|^{t+\alpha 2_t^*}} + f_m$, weakly in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$,

where $f_m \rightarrow 0$ in the dual of $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$. If for any, $R_m > 0$ and $\eta_m \in \mathbb{R}^{N-k}$, $\bar{u}_m(x', x'') := R_m^{(k-N)/2} u_m(x' R_m^{-1}, x'' R_m^{-1} + \eta_m)$ converges to zero weakly in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$, then there exist $R_m > 0$ and $\eta_m \in \mathbb{R}^{N-k}$ such that

$$\liminf_{m \rightarrow \infty} \int_{\Omega} \frac{|\bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} dx > 0,$$

where $\Omega := \{(x', x'') \in \mathbb{R}^N : \frac{1}{2} < |x'| < 1, |x''| < 1\}$.

Proof. We can choose $\eta_m \in \mathbb{R}^{N-k}$ and $R_m > 0$ such that

$$\begin{aligned} Q_m(1) &:= \sup_{\eta \in \mathbb{R}^{N-k}} \int_{B_1^k(0) \times B_1^{N-k}(\eta)} \frac{|\bar{u}_m|^{2_t^*}}{|x'|^{t+2_t^*}} dx = \int_{B_1^k(0) \times B_1^{N-k}(\eta)} \frac{|\bar{u}_m|^{2_t^*}}{|x'|^{t+2_t^*}} dx \\ &= \frac{(\kappa_k^t)^{\frac{2_t^*}{2_t^*-2}}}{2}. \end{aligned} \quad (4.7)$$

Then, clearly, $\{\bar{u}_m\}$ satisfies

$$L_\alpha^s \bar{u}_m = \frac{|\bar{u}_m|^{2_t^*-2} \bar{u}_m}{|x'|^{t+\alpha 2_t^*}} + \bar{f}_m, \text{ weakly in } \dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N), \quad (4.8)$$

where $\bar{f}_m \rightarrow 0$ in the dual of $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$. Now, there exist $N_0 \in \mathbb{N}$ and $\eta_1, \dots, \eta_{N_0} \in \mathbb{R}^{N-k}$ such that $B_1^{N-k}(0) \subset \cup_{j=1}^{N_0} B_{1/2}^{N-k}(\eta_j)$. Let $\psi_1, \dots, \psi_{N_0} \in C_c^\infty(\mathbb{R}^N; [0, 1])$ be such that

$$\psi_j = 1, \text{ on } B_{\frac{1}{2}}^k(0) \times B_{\frac{1}{2}}^{N-k}(\eta_j) \text{ and } \text{supp } \psi_j \subset B_1^k(0) \times B_1^{N-k}(\eta_j).$$

Now, using $\psi_j = 1$, on $B_{\frac{1}{2}}^k(0) \times B_{\frac{1}{2}}^{N-k}(\eta_j)$, translation invariance of norms in the second variable (i.e. x'') and employing the same calculations as in the Lemma 7.2, we can assert that $\psi_j^2 \bar{u}_m$ is a bounded sequence in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$. Hence using $\psi_j^2 \bar{u}_m$, as a test function in (4.8) we arrive at

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\left((\psi_j^2 \bar{u}_m)(x) - (\psi_j^2 \bar{u}_m)(y) \right) (\bar{u}_m(x) - \bar{u}_m(y))}{|x-y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy = \int_{\mathbb{R}^N} \frac{|\bar{u}_m|^{2_t^*-2} \bar{u}_m^2 \psi_j^2}{|x'|^{t+\alpha 2_t^*}} + \bar{\sigma}_m, \quad (4.9)$$

where $\bar{\sigma}_m \rightarrow 0$ as $m \rightarrow \infty$. Since $\bar{u}_m \rightharpoonup 0$ in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$, so, using Lemma 4.3, (4.9) and Hölder inequality we conclude that

$$\begin{aligned} &\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|(\psi_j \bar{u}_m)(x) - (\psi_j \bar{u}_m)(y)|^2}{|x-y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy \\ &\leq \left(\int_{B_1^k(0) \times B_1^{N-k}(\eta_j)} \frac{|\bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2_t^*-2}{2_t^*}} \left(\int_{\mathbb{R}^N} \frac{|\psi_j \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2}{2_t^*}} + \bar{\sigma}_m + \sigma_{R,m} + \tilde{\sigma}_R, \end{aligned} \quad (4.10)$$

where for any large R (depending only on j) $\lim_{m \rightarrow \infty} \sigma_{R,m} = 0$, $\lim_{R \rightarrow \infty} \tilde{\sigma}_R = 0$ and $\lim_{m \rightarrow \infty} \bar{\sigma}_m = 0$. Hence, using (3.4), (4.7) and (4.10) we have

$$\kappa_k^t \left(\int_{\mathbb{R}^N} \frac{|\psi_j \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2}{2_t^*}} \leq \frac{\kappa_k^t}{2^{\frac{2_t^*-2}{2_t^*}}} \left(\int_{\mathbb{R}^N} \frac{|\psi_j \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2}{2_t^*}} + \sigma_{R,m} + \tilde{\sigma}_R + \bar{\sigma}_m.$$

First letting, $m \rightarrow \infty$, then letting $R \rightarrow \infty$ we conclude that for any $j = 1, \dots, N_0$

$$\lim_{m \rightarrow \infty} \int_{\mathbb{R}^N} \frac{|\psi_j \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} dx = 0.$$

Hence

$$\int_{B_{\frac{1}{2}}^k(0) \times B_1^{N-k}(0)} \frac{|\bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} dx \leq \sum_{j=1}^{N_0} \int_{B_{\frac{1}{2}}^k(0) \times B_{\frac{1}{2}}^{N-k}(\eta_j)} \frac{|\bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} dx \rightarrow 0, \text{ as } m \rightarrow \infty.$$

Finally, using (4.7) we conclude the Proposition. \square

4.3. Proof of Theorem 1.2.

Proof. We can take a nonnegative minimizing sequence $\{u_m\}$ in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$ satisfying the following.

- (i) $\int_{\mathbb{R}^N} \frac{|u_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} dx = (\kappa_k^t)^{\frac{2_t^*}{2_t^*-2}},$
- (ii) $[[u_m]]_{s,\alpha,\mathbb{R}^N}^2 = (\kappa_k^t)^{\frac{2_t^*}{2_t^*-2}} + o(1), \text{ as } m \rightarrow \infty,$
- (iii) $L_\alpha^s u_m = \frac{|u_m|^{2_t^*-2} u_m}{|x'|^{t+\alpha 2_t^*}} + f_m \text{ weakly in } \dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N),$

where (iii) is a consequence of Ekeland's Variation Principle and $f_m \rightarrow 0$ in the dual of $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$. We claim that, there exist $R_m > 0$ and $\eta_m \in \mathbb{R}^{N-k}$, such that upto a subsequence, $\bar{u}_m \rightharpoonup u \neq 0$ in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$, where, \bar{u}_m is defined as in Proposition 4.4. Once the claim is established, we can argue similarly as in the proof of Theorem 1.1 to complete the Theorem. If possible, let us assume, that the claim is false. Then Proposition 4.4 guarantees the existence of $R_m > 0$ and $\eta_m \in \mathbb{R}^{N-k}$ such that $\bar{u}_m \rightharpoonup 0$ in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$,

$$\lim_{m \rightarrow \infty} \int_{\Omega} \frac{|\bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} dx > 0 \text{ and}$$

$$L_\alpha^s \bar{u}_m = \frac{|\bar{u}_m|^{2_t^*-2} \bar{u}_m}{|x'|^{t+\alpha 2_t^*}} + \bar{f}_m \text{ weakly in } \dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N),$$

where $\bar{f}_m \rightarrow 0$ in the dual of $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$ and $\Omega := \{(x', x'') \in \mathbb{R}^N : \frac{1}{2} < |x'| < 1, |x''| < 1\}$. Let $\psi \in C_c^\infty(\mathbb{R}_k^N, [0, 1])$, such that $\psi = 1$ in Ω . Since, ψ is supported away from $\{x' = 0\}$ so, we can perform a similar calculation, given in Lemma 7.2 to conclude $\{\psi^2 \bar{u}_m\}$ is a bounded sequence in $\dot{\mathcal{H}}^{s,\alpha}(\mathbb{R}^N)$. Then, proceeding similarly as in the proof of Proposition 4.4, we can derive

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|(\psi \bar{u}_m)(x) - (\psi \bar{u}_m)(y)|^2}{|x-y|^{N+2s} |x'|^\alpha |y'|^\alpha} dx dy$$

$$\leq \left(\int_{\mathbb{R}^N} \frac{|\bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2_t^*-2}{2_t^*}} \left(\int_{\mathbb{R}^N} \frac{|\psi \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2}{2_t^*}} + \bar{\sigma}_m + \sigma_{R,m} + \tilde{\sigma}_R,$$

$$\leq \kappa_k^t \left(\int_{\mathbb{R}^N} \frac{|\psi \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2}{2_t^*}} + \bar{\sigma}_m + \sigma_{R,m} + \tilde{\sigma}_R, \quad (4.11)$$

where for any large R (depending only on the support of ψ) $\lim_{m \rightarrow \infty} \sigma_{R,m} = 0$, $\lim_{R \rightarrow \infty} \tilde{\sigma}_R = 0$ and $\lim_{m \rightarrow \infty} \bar{\sigma}_m = 0$. Now, let us define $v_m := \frac{\psi \bar{u}_m}{|x'|^\alpha}$. Then by (3.3) and (4.11) we have

$$[v_m]_{s, \mathbb{R}^N}^2 - \mathcal{C}_{N,k,s} \int_{\mathbb{R}^N} \frac{v_m^2}{|x'|^{2s}} \leq \kappa_k^t \left(\int_{\mathbb{R}^N} \frac{|\psi \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2}{2_t^*}} + \bar{\sigma}_m + \sigma_{R,m} + \tilde{\sigma}_R.$$

Since, ψ is supported in \mathbb{R}_k^N , using Lemma 4.2 and inequality (1.5) for $\beta = 0$, we have

$$\bar{S}_k^t \left(\int_{\mathbb{R}^N} \frac{|\psi \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2}{2_t^*}} \leq \kappa_k^t \left(\int_{\mathbb{R}^N} \frac{|\psi \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} \right)^{\frac{2}{2_t^*}} + \bar{\sigma}_m + \sigma_{R,m} + \tilde{\sigma}_R. \quad (4.12)$$

As pointed out in Remark 4.1, $\kappa_k^t < \bar{S}_k^t$. Hence, from (4.12) we conclude that

$$\lim_{m \rightarrow \infty} \int_{\mathbb{R}^N} \frac{|\psi \bar{u}_m|^{2_t^*}}{|x'|^{t+\alpha 2_t^*}} = 0,$$

which contradicts Proposition 4.4. This proves the Theorem. \square

5. QUALITATIVE PROPERTIES OF SOLUTION

Main goal of this section is to prove Theorem 1.3. Following the ideas of [1], using (3.3), we could hide the singular weight in the operator. Therefore, working with the newly defined operator and using Moser iteration technique, we can prove the asymptotic estimate (1.8), given in Theorem 1.3. On the other hand, to prove the regularity of solution, we have used extension technique introduced by Caffarelli and Silvestre in [7].

The proof of inequality (1.8) is based on Moser iteration technique

5.1. L^∞ Estimates of Solutions. In this subsection, we will prove some L^∞ estimates of solution of (1.1), which seems inevitable to give an unified proof of the cylindrical symmetry of solutions. We start by introducing the operator L_α^s , which is defined on $\dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$ by the following inner product :

$$\langle L_\alpha^s v, \phi \rangle := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(v(x) - v(y))(\phi(x) - \phi(y))}{|x - y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy, \quad (5.1)$$

for all $\phi \in \dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$. Notice that as consequence of Theorem 3.3, if u satisfies (1.1) then $U := \mathcal{P}_\alpha[u] := |x'|^{\tilde{\alpha}} u \in \dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$ satisfies

$$L_\alpha^s U = \frac{U^{2_t^* - 1}}{|x'|^{t+2_t^* \tilde{\alpha}}} \quad (5.2)$$

weakly, where $0 < \tilde{\alpha} \leq \alpha$. First, we recall the following lemma, from [23], for convex functions, which will be used to derive a Kato type inequality for the newly defined operator L_α^s , in the proof of Proposition 5.2.

Lemma 5.1. *Let $I \subset \mathbb{R}$ be an interval, $a, b \in I$, $\theta_1, \theta_2 \geq 0$. If $f \in C^1(I)$ is convex, then*

$$(b - a) (\theta_1 f'(b) - \theta_2 f'(a)) \geq (f(b) - f(a)) (\theta_1 - \theta_2).$$

In particular, the following inequality holds:

$$(b - a) (f(b)f'(b) - f(a)f'(a)) \geq (f(b) - f(a))^2. \quad (5.3)$$

Proposition 5.2. *Let $V \in \dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$ be any non negative solution of (5.2), where $2 \leq k \leq N-2$, $0 < s < 1$, $0 \leq t < 2s$, $0 < \tilde{\alpha} \leq \alpha$ and $2_t^* = \frac{2(N-t)}{N-2s}$. Then $V \in L^\infty(\mathbb{R}^N)$.*

Proof. For $q \geq 1$ and $R > 0$ we define

$$\phi(r) := \phi_{q,R}(r) := \begin{cases} r^q, & \text{if } 0 \leq r \leq R \\ qR^{q-1}(r-R) + R^q, & \text{if } r > R. \end{cases}$$

Clearly, $\phi_{q,R}$ is Lipschitz and $\phi_{q,R}(0) = 0$ so, by Lemma 2.1, $\phi_{q,R}(V) \in \dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$. So, using (3.4) and Theorem 3.3 we have

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\phi(V(x)) - \phi(V(y))|^2}{|x-y|^{N+2s}} \frac{dx dy}{|x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} \geq C_{N,s,k} \left[\int_{\mathbb{R}^N} |\phi(V)|^{2_t^*} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \right]^{2/2_t^*}. \quad (5.4)$$

Notice that, $\phi \in C^1([0, \infty))$ so, using inequality 5.3 we have

$$\begin{aligned} & \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\phi(V(x)) - \phi(V(y))|^2}{|x-y|^{N+2s}} \frac{dx dy}{|x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} \\ & \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(\phi(V(x))\phi'(V(x)) - \phi(V(y))\phi'(V(y)))(V(x) - V(y))}{|x-y|^{N+2s}} \frac{dx dy}{|x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}}. \end{aligned} \quad (5.5)$$

Now, observe that the function g defined by $g := \phi\phi'$ is Lipschitz with $g(0) = 0$. Hence, $g(V) = \phi(V)\phi'(V) \in \dot{\mathcal{H}}^{s,\tilde{\alpha}}(\mathbb{R}^N)$. So, using $g(V)$ as test function in (5.2) and then employing (5.5) we have

$$\begin{aligned} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\phi(V(x)) - \phi(V(y))|^2}{|x-y|^{N+2s}} \frac{dx dy}{|x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} & \leq \int_{\mathbb{R}^N} \phi(V)\phi'(V) \frac{V^{2_t^*-1}}{|x'|^{t+\tilde{\alpha}2_t^*}} dx \\ & \leq 2q \int_{\mathbb{R}^N} \phi^2(V) V^{2_t^*-2} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}}, \end{aligned} \quad (5.6)$$

where in the last inequality we have used $r\phi'(r) \leq 2q\phi(r)$. Combining (5.4) and (5.6) we have

$$\left[\int_{\mathbb{R}^N} |\phi(V)|^{2_t^*} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \right]^{2/2_t^*} \leq 2qC_{N,s,k} \int_{\mathbb{R}^N} \phi^2(V) V^{2_t^*-2} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \quad (5.7)$$

We estimate the R.H.S. of (5.7) in the following manner. Consider $q = 2_t^*/2$ and $a > 0$ (to be chosen later). Then we have

$$\begin{aligned} 2qC_{N,s,k} \int_{\mathbb{R}^N} \phi^2(V) V^{2_t^*-2} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} & = 2qC_{N,s,k} \int_{\{V \leq a\}} \phi^2(V) V^{2_t^*-2} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \\ & \quad + 2qC_{N,s,k} \int_{\{V > a\}} \phi^2(V) V^{2_t^*-2} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}}. \end{aligned} \quad (5.8)$$

Now, first using Hölder inequality and then choosing $a > 0$ large enough we have

$$2qC_{N,s,k} \int_{\{V > a\}} \phi^2(V) V^{2_t^*-2} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \leq \frac{1}{2} \left[\int_{\mathbb{R}^N} |\phi(V)|^{2_t^*} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \right]^{\frac{2}{2_t^*}}$$

Hence incorporating this estimate in (5.8) and then using (5.7) we have

$$\begin{aligned} \left[\int_{\mathbb{R}^N} |\phi(V)|^{2_t^*} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \right]^{\frac{2}{2_t^*}} & \leq qa^{2_t^*-2} C_{N,s,k} \int_{\mathbb{R}^N} \frac{\phi^2(V) dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \\ & \leq qa^{2_t^*-2} C_{N,s,k} \int_{\mathbb{R}^N} \frac{|V|^{2_t^*}}{|x'|^{t+\tilde{\alpha}2_t^*}}, \end{aligned}$$

where in the last inequality we have used $\phi(r) \leq r^q$ and $q = 2^*/2$. Now letting $R \rightarrow \infty$ and using Fatou's lemma we conclude

$$\int_{\mathbb{R}^N} |V|^{2_t^* \frac{2_t^*}{2}} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} < \infty. \quad (5.9)$$

For $m \geq 1$ we define $\{q_m\}$ by

$$2q_{m+1} + 2_t^* - 2 = 2_t^* q_m, \quad q_1 = \frac{2_t^*}{2}. \quad (5.10)$$

Then using (5.7) and (5.10) we arrive at

$$\left[\int_{\mathbb{R}^N} |\phi_{q_{m+1},R}(V)|^{2_t^*} \frac{dx}{|x'|^{t+\tilde{\alpha}2_t^*}} \right]^{\frac{2_t^*}{2}} \leq q_{m+1} C_{N,s,k} \int_{\mathbb{R}^N} \frac{V^{q_m 2_t^*}}{|x'|^{t+\tilde{\alpha}2_t^*}} dx$$

Again letting $R \rightarrow 0$ and using Fatou's lemma we conclude

$$\left[\int_{\mathbb{R}^N} \frac{V^{q_{m+1} 2_t^*}}{|x'|^{t+\tilde{\alpha}2_t^*}} dx \right]^{\frac{1}{2_t^*(q_{m+1}-1)}} \leq (q_{m+1} C_{N,s,k})^{\frac{1}{2(q_{m+1}-1)}} \left[\int_{\mathbb{R}^N} \frac{V^{q_m 2_t^*}}{|x'|^{t+\tilde{\alpha}2_t^*}} dx \right]^{\frac{1}{2_t^*(q_m-1)}}. \quad (5.11)$$

For $m \geq 1$, set

$$I_m := \left[\int_{\mathbb{R}^N} \frac{V^{q_m 2_t^*}}{|x'|^{t+\tilde{\alpha}2_t^*}} dx \right]^{\frac{1}{2_t^*(q_m-1)}}, \quad D_m = (q_{m+1} C_{N,s,k})^{\frac{1}{2(q_{m+1}-1)}}.$$

Then (5.11) gives $I_{m+1} \leq D_m I_m, \forall m \geq 1$. Taking log in both side and then iterating we get

$$\log I_{m+1} \leq \sum_{j=1}^m \log D_j + \log I_1. \quad (5.12)$$

Since $q_1 > 1$, it is easy to see $\sum_{j=1}^{\infty} \log D_j < C_{N,s,k}$. Hence using (5.9), from (5.12) we get

$$I_{m+1} \leq C_{N,s,k}, \quad \forall m \geq 1.$$

So, for any $R > 0$ we have

$$\left[\int_{|x| \leq R} V^{q_m 2_t^*} dx \right]^{\frac{1}{2_t^*(q_m-1)}} \leq R^{\frac{t+\tilde{\alpha}2_t^*}{2_t^*(q_m-1)}} \left[\int_{|x| \leq R} \frac{V^{q_m 2_t^*}}{|x'|^{t+\tilde{\alpha}2_t^*}} dx \right]^{\frac{1}{2_t^*(q_m-1)}} \leq C_{N,s,k} R^{\frac{t+\tilde{\alpha}2_t^*}{2_t^*(q_m-1)}}.$$

Since $q_m \rightarrow \infty$ as $m \rightarrow \infty$, so $V \in L^\infty(B_R^N(0))$ and

$$\|V\|_{L^\infty(B_R^N(0))} \leq C_{N,s,k}, \quad \forall R > 0.$$

This proves the proposition. \square

Corollary 5.3. *As a consequence of the preceding Proposition and Theorem 3.3, we observe that if u solves (1.1) then $\mathcal{P}_{\tilde{\alpha}}[u] \in L^\infty(\mathbb{R}^N)$. Moreover, by Lemma 2.2 of [13], (1.1) is invariant under the Kelvin transformation i.e. $Ku(x) := \frac{1}{|x|^{N-2s}} u(x/|x|^2)$. Hence we have the following asymptotic estimate for any u solving (1.1)*

$$u(x) \leq \frac{C}{|x'|^{\tilde{\alpha}} (1 + |x|^{N-2s-2\tilde{\alpha}})}, \quad \forall x \in \mathbb{R}_k^N,$$

where $C > 0$ is constant, depends on u but independent of $x \in \mathbb{R}_k^N$.

5.2. Extension and Regularity. Throughout this section, we will consider, $\beta < \mathcal{C}_{k,s}$. Let $u \in \dot{H}^s(\mathbb{R}^N)$ be a positive solution of

$$(-\Delta)^s u - \beta \frac{u}{|x'|^{2s}} = \frac{u^{2^*_t-1}}{|x'|^t} \text{ in } \mathbb{R}^N.$$

We consider, the s -Harmonic extension U of u defined by

$$U(x, r) := d_{N,s} \int_{\mathbb{R}^N} \frac{r^{2s}}{(|x - \xi|^2 + r^2)^{\frac{N+2s}{2}}} u(\xi) d\xi, \text{ for } x \in \mathbb{R}^N, r \in (0, \infty). \quad (5.13)$$

where the constant $d_{N,s}$ is chosen so that $\int_{\mathbb{R}^N} \frac{dx}{(1+|x|^2)^{\frac{N+2s}{2}}} = \frac{2^{1-2s}\Gamma(1-s)}{d_{N,s}\Gamma(s)}$. Then, (See [7]) for any bounded domain $\Omega \subset \mathbb{R}_+^{N+1}$, $U \in H^1(\Omega, 1-2s)$ and satisfies (weakly)

$$\begin{cases} \operatorname{div}(r^{1-2s}\nabla U) = 0 & \text{in } \Omega, \\ -\lim_{r \rightarrow 0^+} r^{1-2s}\partial_r U(x, r) = a(x)U(x, 0) + b(x) & \text{on } \partial'\Omega, \end{cases} \quad (5.14)$$

where $\partial'\Omega$ is the interior of $\partial\Omega \cap \mathbb{R}^N$, $b(x) = 0$ and $a(x) := \left(\frac{\beta}{|x'|^{2s}} + \frac{U^{4s-2t}(x, 0)}{|x'|^t} \right)$. For

$R > 0$, we denote $Q_R := B_R^N(0) \times (0, R)$. We will use the following results from [22] to prove the smoothness of solution away from the set $\{x' = 0\}$.

Proposition 5.4. *Let $a \in L^{N/2s}(B_1^N)$, $b \in L^p(B_1^N)$ with $p > 2s$. Also, assume that $0 \leq \bar{U} \in H^1(Q_1, 1-2s)$ is weak solution of (5.14) in Q_1 . Then there exists $\delta > 0$ which depends only on n and s such that if $\|a^+\|_{L^{n/2s}(B_1^N)} < \delta$, then*

$$\|\bar{U}(\cdot, 0)\|_{L^q(\partial'Q_{1/2})} \leq C \left(\|\bar{U}\|_{L^2(Q_{1,1-2s})} + \|\nabla \bar{U}\|_{L^2(Q_{1,1-2s})} + \|b^+\|_{L^p(B_1^N)} \right),$$

where $C > 0$ depends only on N, p, s, δ and $q := \min\left\{\frac{2(N+1)}{N-2s}, \frac{N(p-1)}{(N-2s)p}, \frac{2N}{N-2s}\right\}$.

Proposition 5.5. *Let $\bar{U} \in H^1(Q_1, 1-2s)$ be a nonnegative weak solution of (5.14) and $a, b \in L^p(B_1)$ for some $p > N/2s$. Then we have the following Harnack inequality:*

$$\sup_{Q_{1/2}} \bar{U} \leq C \left(\inf_{Q_{1/2}} \bar{U} + \|b\|_{L^p(B_1)} \right),$$

where $C > 0$ depends only on N, p, s and $\|a\|_{L^p(B_1)}$. Consequently, there exists $\alpha \in (0, 1)$ depending only on N, p, s and $\|a\|_{L^p(B_1)}$ such that any weak solution \bar{U} of (5.14) is in $C^\alpha(\overline{Q_{1/2}})$.

5.3. Proof of Theorem 1.3.

Proof. Clearly, the first part of the theorem follows from Corollary 5.3. To prove the second part, we take $x_0 \in \mathbb{R}_k^N$ and $R_0 > 0$ be such that $\overline{B_{R_0}^N(x_0)} \subset \mathbb{R}_k^N$. Consider the s -harmonic extension U of u defined in (5.13). Clearly, U is nonnegative and satisfies (5.14) with $b = 0$, weakly in $B_{R_0}^N(x_0) \times (0, R_0)$. We define $V(x, r) := U(x_0 + R_0x, R_0r)$ for $x \in B_1^N(0)$ and $0 < r < 1$. Then, V satisfies (5.14) with $a(x) := R_0^{2s} \left(\frac{\beta}{|x'_0 + R_0x'|^{2s}} + \frac{u^{2^*_t-2}(x)}{|x'_0 + R_0x'|^t} \right)$ and $b(x) = 0$.

Since $\overline{B_{R_0}^N(x_0)} \subset \mathbb{R}_k^N$, $a \in L^{\frac{N}{2s-t}}(B_1^N)$. Using Proposition 5.4 whenever needed, we conclude $a \in L^p(B_{1/2}^N)$ with $p > N/2s$, for any $0 \leq t < 2s$. Hence by Proposition 5.5, $V \in C^\alpha(\overline{Q_{1/4}})$. So, by local Schauder estimates (See Theorem 2.11 of [22]) and bootstrapping argument we have $V \in C^\infty(\overline{Q_{1/4}})$ which in turn implies $U \in C^\infty(\overline{B_{R_0}^N(x_0)} \times [0, R])$. This implies $u \in C^\infty(\mathbb{R}_k^N)$. This proves the theorem. \square

6. CYLINDRICAL SYMMETRY OF POSITIVE SOLUTION

Our main goal of this section is to prove Theorem 1.4.

First, we will prove the following strong maximum principle for antisymmetric function. Notice that, here we are not assuming any lower semi continuity of u upto the boundary. To compensate this, we are assuming a global non negativity of u on the half plane. The proof is a suitable adaptation of the techniques introduced by Silvestre in [32]. For reader's convenience, we will add the proof.

Lemma 6.1. *Let $u \in L_s(\mathbb{R}^N)$ and $(-\Delta)^s u \geq 0$ in Ω , in the distributional sense, where $\Omega \subset \Omega_\lambda := \{x \in \mathbb{R}^N : x_1 < \lambda\}$ is open and bounded. Also, assume that $u \geq 0$ a.e. in Ω_λ and u is antisymmetric i.e. $u(x_\lambda) = -u(x)$ for a.e. $x \in \Omega_\lambda$, where $x_\lambda := (2\lambda - x_1, x_2, \dots, x_N)$. Then either $u > 0$ in Ω or $u \equiv 0$ a.e. in \mathbb{R}^N . Moreover, the above result still remains true if we replace x_1 by x_i for any $i \in \{1, 2, \dots, N\}$ in the definition of Ω_λ .*

Proof. Since $(-\Delta)^s u \geq 0$ in Ω and $u \in L_s(\mathbb{R}^N)$ so by Proposition 2.15 of [32] we have, u is lower semicontinuous in Ω and satisfies the following

$$u(x_0) \geq \int_{\mathbb{R}^N} u(x) \tau_\gamma(x - x_0) dx, \quad (6.1)$$

for any $x_0 \in \Omega$ and $\gamma < \text{dist}(x_0, \partial\Omega)$. Here, $\tau_\gamma(x) := (-\Delta)^s \Gamma_\gamma(x)$, $\Gamma_\gamma(x) := \frac{\Gamma(x/\gamma)}{\gamma^{N-2s}}$, and Γ is a $C^{1,1}$ regularization of $\Phi(x) := \frac{1}{|x|^{N-2s}}$ such that

$$\begin{aligned} \Gamma &\equiv \Phi \text{ in } \mathbb{R}^N \setminus B_1^N(0), \\ \Gamma &\leq \Phi \text{ in } B_1^N(0). \end{aligned}$$

Next, we claim that $\tau_\gamma(x - x_0) \geq \tau_\gamma(x_\lambda - x_0)$, $\forall x \in \Omega_\lambda$. To prove this, we notice that, for $x \in \Omega_\lambda \setminus B_\gamma^N(x_0)$

$$\begin{aligned} \tau_\gamma(x - x_0) &= P.V. \int_{\mathbb{R}^N} \frac{\Gamma_\gamma(x - x_0) - \Gamma_\gamma(y)}{|x - x_0 - y|^{N+2s}} dy \\ &= \int_{B_\gamma^N(x_0)} \frac{\Phi(y - x_0) - \Gamma_\gamma(y - x_0)}{|x - y|^{N+2s}} dy. \end{aligned}$$

To get the last equality we have used the fact that Φ is the fundamental solution of the fractional s -laplacian. Similarly, we have

$$\tau_\gamma(x_\lambda - x_0) = \int_{B_\gamma^N(x_0)} \frac{\Phi(y - x_0) - \Gamma_\gamma(y - x_0)}{|x_\lambda - y|^{N+2s}} dy.$$

Clearly, $|x - y| \leq |x_\lambda - y|$ if $x \in \Omega_\lambda \setminus B_\gamma^N(x_0)$ and $y \in B_\gamma^N(x_0)$. Hence, $\tau_\gamma(x - x_0) \geq \tau_\gamma(x_\lambda - x_0)$, $\forall x \in \Omega_\lambda \setminus B_\gamma^N(x_0)$. Now, let $x \in B_\gamma^N(x_0)$. Then, for γ small

$$\tau_\gamma(\tilde{x} - x_0) \leq \frac{\gamma^{2s}}{|\tilde{x} - x_0|^{N+2s}}, \text{ for } |\tilde{x} - x_0| \geq \frac{\text{dist}(x_0, \partial\Omega_\lambda)}{2},$$

where $C > 0$ is a generic constant. Hence, for γ sufficiently small $\tau_\gamma(x_\lambda - x_0) \leq C$, whereas $\tau_\gamma(x - x_0) = \frac{1}{\gamma^N} \tau_1\left(\frac{x - x_0}{\gamma}\right) \geq C$. This concludes our claim. Now, consider

$$\int_{\mathbb{R}^N} u(x) \tau_\gamma(x - x_0) dx = \int_{\Omega_\lambda} u(x) \tau_\gamma(x - x_0) dx + \int_{\mathbb{R}^N \setminus \Omega_\lambda} u(x) \tau_\gamma(x - x_0) dx.$$

Since, for sufficiently small γ , $\tau_\gamma(x - x_0) \geq \tau_\gamma(x_\lambda - x_0)$, $\forall x \in \Omega_\lambda$, we have

$$\int_{\Omega_\lambda} u(x) \tau_\gamma(x - x_0) dx \geq \int_{\Omega_\lambda} u(x) \tau_\gamma(x_\lambda - x_0) dx = - \int_{\Omega_\lambda} u(x_\lambda) \tau_\gamma(x_\lambda - x_0) dx$$

$$= - \int_{\mathbb{R}^N \setminus \Omega_\lambda} u(y) \tau_\gamma(y - x_0) dy.$$

Hence, $\int_{\mathbb{R}^N} u(x) \tau_\gamma(x - x_0) dx \geq 0$. Now, if possible, let us assume that there exists $x_0 \in \Omega$ such that $u(x_0) \leq 0$. Then, for $\gamma < \text{dist}(x_0, \partial\Omega)$ small enough we have from (6.1)

$$0 \geq u(x_0) \geq \int_{\mathbb{R}^N} u(x) \tau_\gamma(x - x_0) dx \geq 0.$$

So, $\int_{\mathbb{R}^N} u(x) \tau_\gamma(x - x_0) dx = 0$. Form here, using u is antisymmetric and non negative on Ω_λ one can easily conclude $u \equiv 0$ a.e. in Ω_λ . This proves the lemma. \square

6.1. Proof of Theorem 1.4.

Proof. For $u, v \in \dot{\mathcal{H}}_\beta^s(\mathbb{R}^N)$ we define

$$E[u, v] := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{u(x)v(x)}{|x'|^{2s}} dx.$$

Also, for $\lambda \in \mathbb{R}$ we define the following sets in \mathbb{R}^N

$$\Omega_\lambda := \{x_1 < \lambda\}, \text{ and } \Omega'_\lambda := \{x_{k+1} < \lambda\}.$$

Suppose u solves (1.1). We will show that, u is symmetric with respect to $\partial\Omega_0 = \{x_1 = 0\}$ and there exist $\lambda_0 \in \mathbb{R}$ such that for any fixed $x' \neq 0$, u is symmetric with respect to $\partial\Omega'_{\lambda_0}$. Once we show this, the rest of the proof will follow from standard arguments.

Step 1 : In this step we will show that u is symmetric with respect to $\partial\Omega_0$. Let $\lambda < 0$ and $w_\lambda := u - u_\lambda$, where $u_\lambda(x) := u(x_\lambda)$ and $x_\lambda := (2\lambda - x_1, x_2, \dots, x_N)$. We also define

$$v_\lambda(x) := \begin{cases} (u - u_\lambda)^+(x), & \text{if } x \in \Omega_\lambda \\ (u - u_\lambda)^-(x), & \text{if } x \in \mathbb{R}^N \setminus \Omega_\lambda, \end{cases}$$

$P_\lambda := \text{supp } v_\lambda \cap \Omega_\lambda$ and $Q_\lambda := \text{supp } v_\lambda \cap (\mathbb{R}^N \setminus \Omega_\lambda)$. Here, for any real number a , a^+ and a^- represents $\max\{a, 0\}$ and $\min\{a, 0\}$ respectively. Clearly, for each $\lambda < 0$ and $0 \leq t < 2s$

$$\int_{\mathbb{R}^N} \frac{|v_\lambda|^{2t}}{|x'|^t} \leq \int_{P_\lambda} \frac{u^{2t}}{|x'|^t} + \int_{Q_\lambda} \frac{u_\lambda^{2t}}{|x'|^t} = \int_{P_\lambda} \frac{u^{2t}}{|x'|^t} + \int_{P_\lambda} \frac{u^{2t}}{|x'_\lambda|^t} < \infty, \quad (6.2)$$

where to get the finiteness we have used (1.8). However, it is not clear whether v_λ belongs to $\dot{\mathcal{H}}_\beta^s(\mathbb{R}^N)$ or not. So, we will approximate v_λ in a proper way. Let $\eta \in C^\infty(\mathbb{R}^k)$ be such that $0 \leq \eta \leq 1$ and

$$\eta(x') = \begin{cases} 0, & \text{if } |x'| < 1 \\ 1, & \text{if } |x'| \geq 2. \end{cases}$$

For $\epsilon > 0$, we define

$$\eta_\epsilon(x') := \eta\left(\frac{x'}{\epsilon}\right) \text{ and } \eta_{\epsilon,\lambda}(x') := \eta\left(\frac{x'_\lambda}{\epsilon}\right).$$

For $h > 0$, we also define $\psi_h(x) := \psi(x/h)$ and $\psi_{h,\lambda}(x) := \psi(x_\lambda/h)$, where $\psi \in C_c^\infty(\mathbb{R}^N)$ such that $0 \leq \psi \leq 1$.

$$\psi(x) = \begin{cases} 1, & \text{if } |x| < 1 \\ 0, & \text{if } |x| \geq 2. \end{cases}$$

We further define, $\phi_{\epsilon,h,\lambda}(x', x'') := \eta_\epsilon(x') \eta_{\epsilon,\lambda}(x'') \psi_h(x) \psi_{h,\lambda}(x)$, $\Phi(x) := \phi_{\epsilon,h,\lambda}^2(x) v_\lambda(x)$ and $\tilde{\Phi}(x) := \phi_{\epsilon,h,\lambda}(x) v_\lambda(x)$. Then

$$\begin{aligned}
E[u_\lambda, \Phi] &= \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u_\lambda(x) - u_\lambda(y)) (\Phi(x) - \Phi(y))}{|x - y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{u_\lambda(x) \Phi(x)}{|x'|^{2s}} \\
&= \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(u(x) - u(y)) (\Phi_\lambda(x) - \Phi_\lambda(y))}{|x - y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \frac{u(x) \Phi_\lambda(x)}{|x'|^{2s}} \\
&\quad + \beta \int_{\mathbb{R}^N} u(x) \Phi_\lambda(x) \left[\frac{1}{|x'|^{2s}} - \frac{1}{|x'_\lambda|^{2s}} \right] dx,
\end{aligned}$$

where to get the last equality we have used the fact that $\Phi \in C_c^{0,1}(\mathbb{R}_k^N)$ supported away from $\partial\Omega_{2\lambda}$. Now, using Φ_λ as a test function in (1.1) we get

$$E[u_\lambda, \Phi] = \int_{\mathbb{R}^N} \frac{u^{2^*_t-1}(x) \Phi_\lambda(x)}{|x'|^t} + \beta \int_{\mathbb{R}^N} u(x) \Phi_\lambda(x) \left[\frac{1}{|x'|^{2s}} - \frac{1}{|x'_\lambda|^{2s}} \right] dx$$

Observing that v_λ is odd with respect to the reflection along $\partial\Omega_\lambda$ and

$$\left[\frac{1}{|x'|^t} - \frac{1}{|x'_\lambda|^t} \right] v_\lambda(x) \text{ and } \left[\frac{1}{|x'|^{2s}} - \frac{1}{|x'_\lambda|^{2s}} \right] v_\lambda(x)$$

both are nonpositive for all $x \in \mathbb{R}^N$, we arrive at the following inequality

$$\begin{aligned}
E[w_\lambda, \Phi] &\leq \int_{\mathbb{R}^N} \left(u^{2^*_t-1} - u_\lambda^{2^*_t-1} \right) (x) \Phi(x) \frac{dx}{|x'|^t} \\
&\leq C \left[\int_{P_\lambda} u^{2^*_t-2} \frac{\tilde{\Phi}^2}{|x'|^t} dx + \int_{Q_\lambda} u_\lambda^{2^*_t-2} \frac{\tilde{\Phi}^2}{|x'|^t} dx \right], \tag{6.3}
\end{aligned}$$

where $C > 0$ is a generic constant. Let $\xi = \frac{2^*_t}{2^*-2^*_t+2}$. Clearly, $0 \leq \xi < 2s$. Now, using Hölder inequality in (6.3) we get

$$\begin{aligned}
E[w_\lambda, \Phi] &\leq C \left(\int_{P_\lambda} u^{2^*} \right)^{\frac{2^*_t-2}{2^*}} \left(\int_{P_\lambda} \frac{|\tilde{\Phi}|^{2\xi}}{|x'|^\xi} \right)^{\frac{2}{2^*_t}} + C \left(\int_{Q_\lambda} u_\lambda^{2^*} \right)^{\frac{2^*_t-2}{2^*}} \left(\int_{Q_\lambda} \frac{|\tilde{\Phi}|^{2\xi}}{|x'|^\xi} \right)^{\frac{2}{2^*_t}} \\
&\leq C \left(\int_{P_\lambda} u^{2^*} \right)^{\frac{2^*_t-2}{2^*}} \left(\int_{\mathbb{R}^N} \frac{|\tilde{\Phi}|^{2\xi}}{|x'|^\xi} \right)^{\frac{2}{2^*_t}}. \tag{6.4}
\end{aligned}$$

Next, we will estimate $E[w_\lambda, \Phi]$ from below. For this we notice

$$\begin{aligned}
&(w_\lambda(x) - w_\lambda(y)) (\Phi(x) - \Phi(y)) \\
&= (w_\lambda(x) - w_\lambda(y)) (\phi_{\epsilon,h,\lambda}^2(x) v_\lambda(x) - \phi_{\epsilon,h,\lambda}^2(y) v_\lambda(y)) \\
&= \left| \tilde{\Phi}(x) - \tilde{\Phi}(y) \right|^2 \\
&\quad - [\phi_{\epsilon,h,\lambda}^2(x) v_\lambda(x) w_\lambda(y) + \phi_{\epsilon,h,\lambda}^2(y) v_\lambda(y) w_\lambda(x) - 2\phi_{\epsilon,h,\lambda}(x) \phi_{\epsilon,h,\lambda}(y) v_\lambda(x) v_\lambda(y)]
\end{aligned}$$

$$\geq \begin{cases} \left| \tilde{\Phi}(x) - \tilde{\Phi}(y) \right|^2 - \phi_{\epsilon,h,\lambda}^2(y)v_\lambda(y)w_\lambda(x), & \text{if } (x,y) \in P_\lambda^c \times (P_\lambda \cup Q_\lambda) \cup Q_\lambda^c \times (P_\lambda \cup Q_\lambda) \\ \left| \tilde{\Phi}(x) - \tilde{\Phi}(y) \right|^2 - \phi_{\epsilon,h,\lambda}^2(x)v_\lambda(x)w_\lambda(y), & \text{if } (x,y) \in (P_\lambda \cup Q_\lambda) \times P_\lambda^c \cup (P_\lambda \cup Q_\lambda) \times Q_\lambda^c \\ \left| \tilde{\Phi}(x) - \tilde{\Phi}(y) \right|^2 - |\phi_{\epsilon,h,\lambda}(x) - \phi_{\epsilon,h,\lambda}(y)|^2 v_\lambda(x)v_\lambda(y), & \text{otherwise,} \end{cases} \quad (6.5)$$

where $P_\lambda^c := \Omega_\lambda \setminus P_\lambda$ and $Q_\lambda^c := \Omega_\lambda^c \setminus Q_\lambda$. Since $\phi_{\epsilon,h,\lambda}$ symmetric, v_λ and w_λ antisymmetric with respect to the reflection along $\partial\Omega_\lambda$, so considering the sign of v_λ and w_λ in respective region we have

$$\begin{aligned} & \int_{P_\lambda^c} \int_{P_\lambda} \phi_{\epsilon,h,\lambda}^2(y)v_\lambda(y)w_\lambda(x) \frac{dydx}{|x-y|^{N+2s}} + \int_{P_\lambda^c} \int_{Q_\lambda} \phi_{\epsilon,h,\lambda}^2(y)v_\lambda(y)w_\lambda(x) \frac{dydx}{|x-y|^{N+2s}} \leq 0, \\ & \int_{Q_\lambda^c} \int_{P_\lambda} \phi_{\epsilon,h,\lambda}^2(y)v_\lambda(y)w_\lambda(x) \frac{dydx}{|x-y|^{N+2s}} + \int_{Q_\lambda^c} \int_{Q_\lambda} \phi_{\epsilon,h,\lambda}^2(y)v_\lambda(y)w_\lambda(x) \frac{dydx}{|x-y|^{N+2s}} \leq 0, \\ & \int_{P_\lambda} \int_{P_\lambda^c} \phi_{\epsilon,h,\lambda}^2(x)v_\lambda(x)w_\lambda(y) \frac{dydx}{|x-y|^{N+2s}} + \int_{Q_\lambda} \int_{P_\lambda^c} \phi_{\epsilon,h,\lambda}^2(x)v_\lambda(x)w_\lambda(y) \frac{dydx}{|x-y|^{N+2s}} \leq 0, \\ & \int_{P_\lambda} \int_{Q_\lambda^c} \phi_{\epsilon,h,\lambda}^2(x)v_\lambda(x)w_\lambda(y) \frac{dydx}{|x-y|^{N+2s}} + \int_{Q_\lambda} \int_{Q_\lambda^c} \phi_{\epsilon,h,\lambda}^2(x)v_\lambda(x)w_\lambda(y) \frac{dydx}{|x-y|^{N+2s}} \leq 0. \end{aligned}$$

Hence, integrating (6.5) we have

$$\begin{aligned} E[w_\lambda, \Phi] & \geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\tilde{\Phi}(x) - \tilde{\Phi}(y)|^2}{|x-y|^{N+2s}} dx dy - \beta \int_{\mathbb{R}^N} \tilde{\Phi}^2(x) \frac{dx}{|x|^{2s}} \\ & \quad - 2 \int_{P_\lambda} \int_{P_\lambda} \frac{v_\lambda(x)v_\lambda(y) |\phi_{\epsilon,h,\lambda}(x) - \phi_{\epsilon,h,\lambda}(y)|^2}{|x-y|^{N+2s}} dx dy \end{aligned} \quad (6.6)$$

Now, consider

$$\begin{aligned} I_{\epsilon,h,\lambda} & := \int_{P_\lambda} \int_{P_\lambda} \frac{v_\lambda(x)v_\lambda(y) |\phi_{\epsilon,h,\lambda}(x) - \phi_{\epsilon,h,\lambda}(y)|^2}{|x-y|^{N+2s}} dx dy \\ & \leq \int_{P_\lambda} \int_{P_\lambda} \frac{u^2(x) |\phi_{\epsilon,h,\lambda}(x) - \phi_{\epsilon,h,\lambda}(y)|^2}{|x-y|^{N+2s}} dx dy \\ & \leq 2 \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x-y|^{N+2s}} + 2 \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u_\lambda^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x-y|^{N+2s}} \\ & \quad + 2 \int_{P_\lambda} \int_{P_\lambda} \frac{u^2(x)\psi_{h,\lambda}^2(y) |\eta_\epsilon(x')\eta_{\epsilon,\lambda}(x') - \eta_\epsilon(y')\eta_{\epsilon,\lambda}(y')|^2}{|x-y|^{N+2s}} dx dy. \end{aligned} \quad (6.7)$$

By dividing \mathbb{R}^N in appropriate domain and using (1.8) (which essentially shows that $u \in L^2(\mathbb{R}^N)$, as $2 \leq k \leq N-2$) one can estimate the first two term of (1.4) to arrive at

$$\begin{aligned} & \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x-y|^{N+2s}} = \sigma_h \text{ and} \\ & \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u_\lambda^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x-y|^{N+2s}} = \sigma_{h,\lambda}, \end{aligned}$$

where for any $\lambda \in \mathbb{R}$ both σ_h and $\sigma_{h,\lambda}$ goes to zero as $h \rightarrow \infty$. To estimate the last term of (6.7) we use the fact that $u \in L^\infty(\Omega_\lambda)$. Hence, finally we arrive at the following inequality

$$I_{\epsilon,h,\lambda} \leq C_{N,s,t,\beta} [\sigma_h + \sigma_{h,\lambda}]$$

$$\begin{aligned}
& + C_{N,s,t,\beta} \|u\|_{L^\infty(\Omega_\lambda)} \int_{P_\lambda} \int_{P_\lambda} \frac{\psi_h^2(y) |\eta_\epsilon(x')\eta_{\epsilon,\lambda}(x') - \eta_\epsilon(y')\eta_{\epsilon,\lambda}(y')|^2}{|x-y|^{N+2s}} dx dy \\
& \leq C_{N,k,s,t,\beta} [\sigma_h + \sigma_{h,\lambda}] \\
& + h^{N-k} C_{N,k,s,t,\beta} \|u\|_{L^\infty(\Omega_\lambda)} \int_{\mathbb{R}^k} \int_{\mathbb{R}^k} \frac{|\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x'-y'|^{k+2s}} dx' dy' \\
& \leq C_{N,k,s,t,\beta} \left[\sigma_h + \sigma_{h,\lambda} + h^{N-k} \epsilon^{k-2s} \|u\|_{L^\infty(\Omega_\lambda)} \right]. \tag{6.8}
\end{aligned}$$

Hence, combining (6.4), (6.7), (6.8) and using (1.5) we conclude that there exist a constant $C_{N,k,s,t,\beta} > 0$ depending on N, k, s, t and β such that the following holds for any $h > 0$, $\epsilon > 0$ and $\lambda < 0$

$$\begin{aligned}
& \left(\int_{\mathbb{R}^N} \frac{|\tilde{\Phi}|^{2_\xi^*}}{|x'|^\xi} \right)^{\frac{2}{2_\xi^*}} - \left[\sigma_h + \sigma_{h,\lambda} + h^{N-k} \epsilon^{k-2s} \|u\|_{L^\infty(\Omega_\lambda)} \right] \\
& \leq C_{N,k,s,t,\beta} \left(\int_{P_\lambda} u^{2^*} \right)^{\frac{2_t^*-2}{2^*}} \left(\int_{\mathbb{R}^N} \frac{|\tilde{\Phi}|^{2_\xi^*}}{|x'|^\xi} \right)^{\frac{2}{2_\xi^*}}
\end{aligned}$$

First letting $\epsilon \rightarrow 0$ then letting $h \rightarrow \infty$ and using DCT (because (6.2) hold for each $\lambda < 0$) we arrive at

$$C_{N,k,s,t,\beta} \left(\int_{\mathbb{R}^N} \frac{|v_\lambda(x)|^{2_\xi^*}}{|x'|^\xi} \right)^{\frac{2}{2_\xi^*}} \leq \left(\int_{P_\lambda} u^{2^*} \right)^{\frac{2_t^*-2}{2^*}} \left(\int_{\mathbb{R}^N} \frac{|v_\lambda(x)|^{2_\xi^*}}{|x'|^\xi} \right)^{\frac{2}{2_\xi^*}} \tag{6.9}$$

If $v_\lambda \neq 0$ a.e., then using (6.9), we have that there exists a constant $C \equiv C(N, k, s, t, \beta)$ such that

$$0 < C(N, k, s, t, \beta) \leq \int_{P_\lambda} |u|^{2^*}, \text{ for any } \lambda < 0. \tag{6.10}$$

But, (6.8) yields a contradiction for large negative values of λ . So, for large negative values of λ we must have $v_\lambda \equiv 0$ a.e. in Ω_λ^i . Hence, the set defined by

$$A := \{\lambda \leq 0 \mid u \leq u_\lambda \text{ a.e. in } \Omega_\mu, \forall \mu \leq \lambda\}$$

is non empty. Let $\bar{\lambda} := \sup A$. For any $x' \in \mathbb{R}^k$ and $R > 0$, we also define the cylinder $C_R(x') := B_R^k(x') \times \mathbb{R}^{N-k}$. We claim that $\bar{\lambda} = 0$ and $u \leq u_{\bar{\lambda}}$ a.e. in $\Omega_{\bar{\lambda}}$. If possible, let us assume that $\bar{\lambda} < 0$. We define, $\tilde{w}_{\bar{\lambda}} := -w_{\bar{\lambda}} = u_{\bar{\lambda}} - u$. Let $\Omega \subset \Omega_{\bar{\lambda}}$. Since $w_{\bar{\lambda}} \in L_s(\mathbb{R}^N) \cap \dot{H}_\beta^s(\mathbb{R}^N)$ and $u_{\bar{\lambda}} \geq u$ a.e. in $\Omega_{\bar{\lambda}}$, $(-\Delta)^s \tilde{w}_{\bar{\lambda}} \geq 0$ in the distributional sense in Ω . So, $w_{\bar{\lambda}}$ is lower semicontinuous as well as antisymmetric and a.e. nonnegative (by continuity) in $\Omega_{\bar{\lambda}}$. Hence, by Lemma 6.1 we have either $w_{\bar{\lambda}} = 0$ a.e. in \mathbb{R}^N or $w_{\bar{\lambda}} > 0$ in Ω . We claim that, the second case can not occur. If it occurs, then $w_{\bar{\lambda}} > 0$ and lower semicontinuous in $\Omega_{\bar{\lambda}}$. Hence, by continuity of u , lower semicontinuity of $w_{\bar{\lambda}}$ and definition of $\bar{\lambda}$ we have for any $R_1 > 0$, $\delta > 0$ small and $R > 0$ large, there exists $\epsilon_0(R_1, R, \delta) > 0$ such that $\bar{\lambda} + \epsilon_0 < 0$ and

$$P_{\bar{\lambda}+\epsilon} \cap \Omega_{\bar{\lambda}-\delta} \cap C_{R_1}^c(0) \subset \mathbb{R}^N \setminus B_R^N(0), \forall 0 < \epsilon \leq \epsilon_0.$$

Now, since $\bar{\lambda}$ is the supremum so $v_{\bar{\lambda}+\epsilon}$ is not zero in a positive measure set, for any $0 < \epsilon \leq \epsilon_0$. Hence, using (6.10) we have

$$\begin{aligned}
0 < C(N, k, s, t, \beta) &\leq \int_{P_{\bar{\lambda}+\epsilon}} u^{2^*} \\
&\leq \int_{P_{\bar{\lambda}+\epsilon} \cap C_{R_1}(0)} u^{2^*} + \int_{P_{\bar{\lambda}+\epsilon} \cap C_{R_1}^c(0)} u^{2^*} \\
&\leq \int_{C_{R_1}(0)} u^{2^*} + \int_{P_{\bar{\lambda}+\epsilon} \cap \Omega_{\bar{\lambda}} \cap C_{R_1}^c(0)} u^{2^*} + \int_{P_{\bar{\lambda}+\epsilon} \cap \Omega_{\bar{\lambda}}^c \cap C_{R_1}^c(0)} u^{2^*} \\
&\leq \int_{C_{R_1}(0)} u^{2^*} + \int_{\mathbb{R}^N \setminus B_R^N(0)} u^{2^*} + \int_{\Omega_{\bar{\lambda}+\epsilon} \setminus \Omega_{\bar{\lambda}-\delta}} u^{2^*}.
\end{aligned}$$

Now, first choosing R_1 small, R large, δ small and then choosing ϵ small we can make the R.H.S. of the above inequality strictly less than $C(N, k, s, t, \beta)$, which gives a contradiction. So, either $\bar{\lambda} = 0$ or if $\bar{\lambda} < 0$ then $w_{\bar{\lambda}} = 0$ a.e. in \mathbb{R}^N . In the second case, $(-\Delta)^s w_{\bar{\lambda}} = 0$ in \mathbb{R}^N in the distributional sense. But, since $\bar{\lambda} < 0$ so, $(-\Delta)^s w_{\bar{\lambda}} > 0$ in Ω , in the distributional sense, for any open set $\Omega \Subset \Omega_{\bar{\lambda}}$. Which gives a contradiction. Hence, $\bar{\lambda} = 0$.

Repeating the same arguments for $\lambda > 0$, we can conclude that u is symmetric decreasing in x_1 direction.

Step 2 : In this step, we will show that there exist $\lambda_0 \in \mathbb{R}$ such that, for any $x' \in \mathbb{R}^k$ fixed, u is symmetric w.r.t the reflexion along $\partial\Omega'_{\lambda_0}$. We will only prove an analogous inequality of (6.10) derived in Step 1. Rest of the arguments will be similar to Step 1. We will exclude that part. We notice that in this case u may not be in $L^\infty(\Omega'_\lambda)$. Because of this we cannot use similar arguments of Step 1 to derive (6.10). We define $w_\lambda(x) = u(x) - u(x_\lambda)$, where $x_\lambda := (x', 2\lambda - x_{k+1}, x_{k+2}, \dots, x_N) = (x', x''_\lambda)$. Then clearly,

$$W_\lambda := \mathcal{P}_{\tilde{\alpha}}[w_\lambda] := |x|^{\tilde{\alpha}} w_\lambda = U - U_\lambda,$$

where we have denoted $\mathcal{P}_{\tilde{\alpha}}[u]$ by U and for any $0 < \beta \leq \mathcal{C}_{N,k,s}$, $\tilde{\alpha} \in [0, \tilde{\alpha} \leq \alpha]$ uniquely determined by (3.2). Since, u solves (1.1) so W_λ satisfies

$$L_{\tilde{\alpha}}^s W_\lambda = A_\lambda W_\lambda \text{ weakly in } \dot{\mathcal{H}}^{s, \tilde{\alpha}}(\mathbb{R}^N), \quad (6.11)$$

where

$$A_\lambda := \frac{1}{|x'|^{t+\tilde{\alpha}2_t^*}} \frac{U^{2_t^*-1} - U_\lambda^{2_t^*-1}}{U - U_\lambda}.$$

Also, define

$$v_\lambda(x) := \begin{cases} w_\lambda^+(x), & \text{if } x \in \Omega'_\lambda \\ w_\lambda^-(x), & \text{if } x \in (\Omega'_\lambda)^c. \end{cases}$$

Then

$$V_\lambda(x) := \mathcal{P}_{\tilde{\alpha}}[v_\lambda] = \begin{cases} W_\lambda^+(x), & \text{if } x \in \Omega'_\lambda \\ W_\lambda^-(x), & \text{if } x \in (\Omega'_\lambda)^c. \end{cases}$$

Similarly to the Step 1, we define

$$\begin{aligned}
P'_\lambda &:= \text{supp } v_\lambda \cap \Omega'_\lambda = \text{supp } V_\lambda \cap \Omega'_\lambda \\
Q'_\lambda &:= \text{supp } v_\lambda \cap (\Omega'_\lambda)^c = \text{supp } V_\lambda \cap (\Omega'_\lambda)^c.
\end{aligned}$$

We define, $\phi_{\epsilon,h,\lambda}(x', x'') := \eta_\epsilon(x')\psi_h(x)\psi_{h,\lambda}(x)$, $\Phi(x) := \phi_{\epsilon,h,\lambda}^2(x)V_\lambda(x)$ and $\tilde{\Phi}(x) := \phi_{\epsilon,h,\lambda}(x)V_\lambda(x)$, where ψ_h and $\psi_{h,\lambda}$ are same as defined in Step 1. Also, for $\epsilon < 1$, $\eta_\epsilon \in C^{0,1}(\mathbb{R}^k)$ and satisfies the following

$$\eta_\epsilon(x') = \begin{cases} 0, & \text{if } |x'| < \epsilon^2 \\ \frac{\ln\left(\frac{|x'|}{\epsilon^2}\right)}{|\ln \epsilon|}, & \text{if } \epsilon^2 \leq |x'| \leq \epsilon \\ 1, & \text{if } |x'| > \epsilon. \end{cases}$$

Then clearly, $\Phi \in C_c^{0,1}(\mathbb{R}_k^N)$ and so using Φ as a test function in (6.11) we get

$$\langle L_{\tilde{\alpha}}^s W_\lambda, \Phi \rangle = \int_{\mathbb{R}^N} \tilde{\Phi}^2 A_\lambda(x) dx. \quad (6.12)$$

In the next few paragraphs C will denote a positive constant possibly depending on N, s, k, t, β . We estimate

$$\begin{aligned} \int_{\mathbb{R}^N} \tilde{\Phi}^2 A_\lambda(x) dx &= \int_{\Omega'_\lambda} \tilde{\Phi}^2(x) A_\lambda(x) dx + \int_{\mathbb{R}^N \setminus \Omega'_\lambda} \tilde{\Phi}(x) A_\lambda(x) dx \\ &\leq C \int_{P'_\lambda} \frac{U^{2t^*-2}}{|x'|^{\tilde{\alpha}(2t^*-2)}} \frac{\tilde{\Phi}^2}{|x'|^{t+\tilde{\alpha}(2t^*-2)}} dx + C \int_{Q'_\lambda} \frac{U_\lambda^{2t^*-2}}{|x'|^{\tilde{\alpha}(2t^*-2)}} \frac{\tilde{\Phi}^2}{|x'|^{t+\tilde{\alpha}(2t^*-2)}} dx, \end{aligned}$$

Now using Hölder's inequality we have

$$\begin{aligned} \int_{\mathbb{R}^N} \tilde{\Phi}^2(x) A_\lambda(x) dx &\leq C \left(\int_{P'_\lambda} U^{2^*} \frac{dx}{|x'|^{\tilde{\alpha}2^*}} \right)^{\frac{2t^*-2}{2^*}} \left(\int_{\mathbb{R}^N} \frac{|\tilde{\Phi}|^{2\xi}}{|x'|^{\xi+\tilde{\alpha}2\xi}} \right)^{\frac{2}{2\xi}} \\ &\leq C \left(\int_{P'_\lambda} u^{2^*} \right)^{\frac{2t^*-2}{2^*}} \left(\int_{\mathbb{R}^N} \frac{|\tilde{\Phi}|^{2\xi}}{|x'|^{\xi+\tilde{\alpha}2\xi}} \right)^{\frac{2}{2\xi}}, \end{aligned} \quad (6.13)$$

where $\xi = \frac{t2^*}{2^*-2t^*+2}$. Next, we estimate $\langle L_{\tilde{\alpha}}^s W_\lambda, \Phi \rangle$. Note that, proceeding similarly as in Step 1 we will arrive at

$$\begin{aligned} \langle L_{\tilde{\alpha}}^s W_\lambda, \Phi \rangle &\geq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\tilde{\Phi}(x) - \tilde{\Phi}(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy \\ &\quad - 2 \int_{P_\lambda} \int_{P_\lambda} \frac{V_\lambda(x)V_\lambda(y) |\phi_{\epsilon,h,\lambda}(x) - \phi_{\epsilon,h,\lambda}(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy \end{aligned} \quad (6.14)$$

Using $2ab \leq (a^2 + b^2)$, with $a = \frac{V_\lambda(x)}{|x'|^{\tilde{\alpha}}}$ and $b = \frac{V_\lambda(y)}{|y'|^{\tilde{\alpha}}}$ whenever required, we estimate

$$\begin{aligned} I_{\epsilon,h,\lambda} &:= \int_{P_\lambda} \int_{P_\lambda} \frac{V_\lambda(x)V_\lambda(y) |\phi_{\epsilon,h,\lambda}(x) - \phi_{\epsilon,h,\lambda}(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy \\ &\leq 2 \int_{P_\lambda} \int_{P_\lambda} \frac{V_\lambda(x)V_\lambda(y) |\psi_{h,\lambda}(x) - \psi_{h,\lambda}(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy \\ &\quad + 2 \int_{P_\lambda} \int_{P_\lambda} \frac{V_\lambda(x)V_\lambda(y) \psi_{h,\lambda}^2(y) |\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy. \\ &\leq 2 \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{U^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x'|^{2\tilde{\alpha}} |x-y|^{N+2s}} + 2 \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{U_\lambda^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x'|^{2\tilde{\alpha}} |x-y|^{N+2s}} \end{aligned}$$

$$\begin{aligned}
& + 2 \int_{P_\lambda} \int_{P_\lambda} \frac{V_\lambda(x)V_\lambda(y)\psi_{h,\lambda}^2(y) |\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy. \\
& \leq 2 \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x-y|^{N+2s}} + 2 \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u_\lambda^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x-y|^{N+2s}} \\
& + 2 \|U\|_{L^\infty(\mathbb{R}^N)}^2 \int_{P_\lambda} \int_{P_\lambda} \frac{\psi_{h,\lambda}^2(y) |\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy, \tag{6.15}
\end{aligned}$$

where in the last inequality we have used Proposition 5.2. As remarked in the Step 1 we have

$$\begin{aligned}
& \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x-y|^{N+2s}} = \sigma_h \text{ and} \\
& \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u_\lambda^2(x) |\psi_h(x) - \psi_h(y)|^2}{|x-y|^{N+2s}} = \sigma_{h,\lambda},
\end{aligned}$$

where for any $\lambda \in \mathbb{R}$ both σ_h and $\sigma_{h,\lambda}$ goes to zero as $h \rightarrow \infty$. Also, by Lemma 7.1 we conclude that

$$\int_{P_\lambda} \int_{P_\lambda} \frac{\psi_{h,\lambda}^2(y) |\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy = \sigma_{\epsilon,h,\lambda},$$

where for any $h > 0$ and $\lambda \in \mathbb{R}$, $\sigma_{\epsilon,h,\lambda} \rightarrow 0$ as $\epsilon \rightarrow 0$. So, from (6.15) we have $I_{\epsilon,h,\lambda} \leq C(\sigma_h + \sigma_{h,\lambda} + \sigma_{\epsilon,h,\lambda})$.

Hence, using (3.4), (6.13) and (6.14) we have from (6.12) that there exist a constant $C_{N,k,s,t,\beta} > 0$ depending on the indexed variables such that

$$\begin{aligned}
C_{N,k,s,t,\beta} \left(\int_{\mathbb{R}^N} \frac{|\tilde{\Phi}(x)|^{2_\xi^*}}{|x'|^{\xi + \tilde{\alpha} 2_\xi^*}} dx \right)^{\frac{2}{2_\xi^*}} & - C(\sigma_h + \sigma_{h,\lambda} + \sigma_{\epsilon,h,\lambda}) \\
& \leq \left(\int_{P'_\lambda} u^{2^*} \right)^{\frac{2_t^* - 2}{2^*}} \left(\int_{\mathbb{R}^N} \frac{|\tilde{\Phi}(x)|^{2_\xi^*}}{|x'|^{\xi + \tilde{\alpha} 2_\xi^*}} dx \right)^{\frac{2}{2_\xi^*}}.
\end{aligned}$$

First letting $\epsilon \rightarrow 0$ then letting $h \rightarrow \infty$ and using DCT we arrive at

$$C_{N,k,s,t,\beta} \left(\int_{\mathbb{R}^N} \frac{|V_\lambda(x)|^{2_\xi^*}}{|x'|^{\xi + \tilde{\alpha} 2_\xi^*}} dx \right)^{\frac{2}{2_\xi^*}} \leq \left(\int_{P_\lambda} u^{2^*} \right)^{\frac{2_t^* - 2}{2^*}} \left(\int_{\mathbb{R}^N} \frac{|V_\lambda(x)|^{2_\xi^*}}{|x'|^{\xi + \tilde{\alpha} 2_\xi^*}} dx \right)^{\frac{2}{2_\xi^*}}$$

Hence, for $V_\lambda \neq 0$ a.e. we have

$$0 < C_{N,k,s,t,\beta} \leq \int_{P'_\lambda} |u|^{2^*}, \text{ for any } \lambda < 0,$$

which is the exact counterpart of inequality (6.10).

Combining Step 1 and Step 2 we conclude that u is cylindrically symmetric. \square

7. APPENDIX

7.1. A Density Property. Main aim of this section is to derive Lemma 2.1. The arguments are modifications of those in [11], where the Muckenhoupt A_1 properties of the

weights have been used crucially. We will sketch the proof by pointing out main steps. First, let us define

$$\mathcal{W} := \left\{ u \in L^{2^*} \left(\mathbb{R}^N; \frac{1}{|x'|^{\tilde{\alpha}2^*}} \right) : \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2 dx dy}{|x - y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} < \infty \right\},$$

endowed with the following norm

$$\|u\|_{\mathcal{W}} := [[u]]_{s, \tilde{\alpha}, \mathbb{R}^N} + \|u\|_{2^*, \tilde{\alpha}, \mathbb{R}^N}.$$

Here,

$$\|u\|_{2^*, \tilde{\alpha}, \mathbb{R}^N} := \left(\int_{\mathbb{R}^N} \frac{|u(x)|^{2^*}}{|x'|^{\tilde{\alpha}2^*}} dx \right)^{\frac{1}{2^*}},$$

and the semi-norm $[[u]]_{s, \tilde{\alpha}, \mathbb{R}^N}$ is same, as defined in Section 2. We also, define the following:

$$\begin{aligned} &\text{when } N' = 2N, w(z, z) = (z, z), \Theta(X) = |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}, X = (x, y); x, y, z \in \mathbb{R}^N, \\ &\text{and when } N' = N, w(z) = z, \Theta(X) = |x'|^{\tilde{\alpha}2^*}, X = x; x, y \in \mathbb{R}^N. \end{aligned} \quad (7.1)$$

Next, we will prove the following lemma.

Lemma 7.1. *Let $u \in C_c^\infty(\mathbb{R}^N)$. Also, we consider η_ϵ , defined by the following*

$$\eta_\epsilon(x') = \begin{cases} 0, & \text{if } |x'| < \epsilon^2 \\ \frac{\ln\left(\frac{|x'|}{\epsilon^2}\right)}{|\ln \epsilon|}, & \text{if } \epsilon^2 \leq |x'| \leq \epsilon \\ 1, & \text{if } |x'| > \epsilon. \end{cases}$$

Then, for any $0 < \tilde{\alpha} \leq (k - 2s)/2$, the following are true

$$\begin{aligned} &\text{(i) } \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy < \infty, \\ &\text{(ii) } \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u^2(x) |\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x - y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy = 0. \end{aligned}$$

In particular, $\eta_\epsilon u \in C_c^{0,1}(\mathbb{R}_k^N)$ converges to u under the semi norm $[[\cdot]]_{s, \tilde{\alpha}, \mathbb{R}^N}$, i.e $u \in \dot{\mathcal{H}}^{s, \tilde{\alpha}}(\mathbb{R}^N)$.

Proof. We will only prove (ii). One can easily check that (i) holds in fact for $u \in C_c^{0,1}(\mathbb{R}^N)$. Notice that

$$\begin{aligned} &\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u^2(x) |\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x - y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy \\ &= \int_{\mathbb{R}^k} \int_{\mathbb{R}^k} \frac{|\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx' dy' \int_{\mathbb{R}^{N-k}} u^2(x', x'') \int_{\mathbb{R}^{N-k}} \frac{dy''}{(|x' - y'|^2 + |x'' - y''|^2)^{\frac{N+2s}{2}}} dx'' \\ &\leq C \int_{\mathbb{R}^k} \int_{\mathbb{R}^k} \frac{|\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x' - y'|^{k+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx' dy', \end{aligned} \quad (7.2)$$

where (and for the rest of the proof) $C > 0$ is constant depending on $N, k, s, \tilde{\alpha}, \|u\|_{L^\infty(\mathbb{R}^N)}$ and $\text{supp } u$. We define

$$\begin{aligned} I_\epsilon(x', y') &:= \frac{|\eta_\epsilon(x') - \eta_\epsilon(y')|^2}{|x' - y'|^{k+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx' dy' \text{ and} \\ H_\epsilon &:= \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} I_\epsilon(x', y') dx' dy'. \end{aligned}$$

Then, in view of the (7.2), it is enough to show $H_\epsilon = o(1)$ as $\epsilon \rightarrow 0$. We define

$$\begin{aligned} H_{\epsilon,1} &:= \int_{|x'| < \epsilon^2} \int_{\epsilon^2 < |y'| < \epsilon} I_\epsilon, \quad H_{\epsilon,2} := \int_{|x'| > \epsilon} \int_{\epsilon^2 < |y'| < \epsilon} I_\epsilon, \\ H_{\epsilon,3} &:= \int_{|x'| < \epsilon^2} \int_{|y'| > \epsilon} I_\epsilon, \quad H_{\epsilon,4} := \int_{\epsilon^2 < |x'| < \epsilon} \int_{\epsilon^2 < |y'| < \epsilon} I_\epsilon. \end{aligned}$$

Then using the symmetry of I_ϵ we have

$$H_\epsilon := 2H_{\epsilon,1} + 2H_{\epsilon,2} + 2H_{\epsilon,3} + H_{\epsilon,4}.$$

We will show $H_{\epsilon,m} = o(1)$ as $\epsilon \rightarrow 0$ for $m = 1, 2, 3, 4$.

Step 1: In this step will estimate $H_{\epsilon,1}$. For this we define

$$\begin{aligned} F_{x'} &:= \{\epsilon^2 < |y'| < \epsilon\} \cap \{y' : |y' - x'| \geq \epsilon^2/2\}, \text{ and} \\ F'_{x'} &:= \{\epsilon^2 < |y'| < \epsilon\} \cap \{y' : |y' - x'| < \epsilon^2/2\}. \text{ Then} \\ H_{\epsilon,1} &= \int_{|x'| < \epsilon^2} \int_{F_{x'}} I_\epsilon + \int_{|x'| < \epsilon^2} \int_{F'_{x'}} I_\epsilon. \end{aligned}$$

We first consider

$$\begin{aligned} \int_{|x'| < \epsilon^2} \int_{F'_{x'}} I_\epsilon &\leq \frac{C}{|\ln \epsilon|^2} \int_{|x'| < \epsilon^2} \frac{1}{|x'|^{2\tilde{\alpha}}} \int_{F'_{x'}} \frac{|\ln |y'| - \ln |x'||^2}{|x' - y'|^{k+2s}} dy' dx' \\ &\leq \int_0^1 \frac{C}{|\ln \epsilon|^2} \int_{|x'| < \epsilon^2} \frac{1}{|x'|^{2\tilde{\alpha}}} \int_{F'_{x'}} \frac{dy'}{|x' - y'|^{k+2s-2} |y' + r(x' - y')|^2} dx' dr \\ &\leq \frac{C}{|\ln \epsilon|^2} \int_{|x'| < \epsilon^2} \frac{1}{|x'|^{2\tilde{\alpha}}} \int_{F'_{x'}} \frac{dy'}{|x' - y'|^{k+2s-2} |y'|^2} dx' \\ &\leq \frac{C}{\epsilon^4 |\ln \epsilon|^2} \int_{|x'| < \epsilon^2} \frac{1}{|x'|^{2\tilde{\alpha}}} \int_{\{|x' - y'| < \frac{\epsilon^2}{2}\}} \frac{dy'}{|x' - y'|^{k+2s-2}} dx' \\ &\leq \frac{C}{\epsilon^4 |\ln \epsilon|^2} \epsilon^{4s} \epsilon^{4-4s} = o(1), \text{ as } \epsilon \rightarrow 0, \end{aligned} \tag{7.3}$$

where in the last inequality we have used the fact that, for small $\epsilon > 0$, $\epsilon^{k-2\tilde{\alpha}} \leq \epsilon^{4s}$, for any $0 < 2\tilde{\alpha} \leq k - 2s$. Next, we consider

$$\begin{aligned} \int_{|x'| < \epsilon^2} \int_{F_{x'}} I_\epsilon &\leq \frac{C}{|\ln \epsilon|^2} \int_{|x'| \leq \epsilon^2} \frac{1}{|x'|^{\tilde{\alpha}}} \int_{F_{x'}} \frac{\ln^2(\frac{|y'|}{\epsilon^2})}{|x' - y'|^{k+2s} |y'|^{\tilde{\alpha}}} dy' dx' \\ &\leq \frac{C \epsilon^{k-2s-2\tilde{\alpha}}}{|\ln \epsilon|^2} \int_{|x'| \leq 1} \frac{1}{|x'|^{\tilde{\alpha}}} \int_{\substack{\{1 < |y'| < \frac{1}{\epsilon}\} \\ \cap \{|x' - y'| > \frac{1}{2}\}}} \frac{\ln^2 |y'|}{|x' - y'|^{k+2s} |y'|^{\tilde{\alpha}}} dy' dx' \\ &\leq o(1) + \frac{C}{|\ln \epsilon|^2} \int_{|x'| < 1} \int_{\substack{\{2 < |y'| < \frac{1}{\epsilon}\} \\ \cap \{|x' - y'| > \frac{1}{2}\}}} \frac{\ln^2 |y'|}{|x' - y'|^{k+2s} |y'|^{\tilde{\alpha}}} dy' dx' \\ &\leq o(1) + \frac{C}{|\ln \epsilon|^2} \int_{\{2 < |y'| < \epsilon^{-1}\}} \frac{\ln^2 |y'| dy'}{|y'|^{k+2s + \frac{k-2s}{2}}} \\ &\leq o(1) + \frac{C}{|\ln \epsilon|^2} \int_{\{2 < |y'| < \epsilon^{-1}\}} \frac{\ln^2 |y'| dy'}{|y'|^{k+2s}} = o(1), \text{ as } \epsilon \rightarrow 0. \end{aligned} \tag{7.4}$$

Hence, combining (7.3) and (7.4) we have $H_{\epsilon,1} = o(1)$, as $\epsilon \rightarrow 0$.

Step 2: In this step, we will show that $H_{\epsilon,m} = o(1)$, as $\epsilon \rightarrow 0$ for $m = 2, 3$. In fact, we will show this, only for the case $m = 2$. The assertion, for the case, $m = 3$, will follow similarly and much more easily.

By a change of variable we get

$$\begin{aligned} H_{\epsilon,2} &\leq \frac{1}{|\ln \epsilon|^2} \int_{\epsilon < |y'| < 1} \frac{1}{|y'|^{2\tilde{\alpha}}} \int_{|x'| > 1} \frac{\left| \ln \frac{|y'|}{\epsilon} - \ln \frac{1}{\epsilon} \right|^2}{|x' - y'|^{k+2s}} dy' dx' \\ &\leq H'_{\epsilon,2} + H''_{\epsilon,2}, \end{aligned} \quad (7.5)$$

where

$$\begin{aligned} H'_{\epsilon,2} &:= \frac{C}{|\ln \epsilon|^2} \int_0^1 \int_{\epsilon < |y'| < 1} \frac{1}{|y'|^{2\tilde{\alpha}}} \int_{\{|x'| > 1\} \cap \{|x' - y'| \leq \frac{1}{2}\}} \frac{dx'}{|x' - y'|^{k+2s-2} |x' + r(x' - y')|^2} dy' dr \\ &\leq \frac{C}{|\ln \epsilon|^2} \int_{\epsilon < |y'| < 1} \frac{1}{|y'|^{2\tilde{\alpha}}} \int_{\{|x' - y'| \leq \frac{1}{2}\}} \frac{dx'}{|x' - y'|^{k+2s-2}} dy' = o(1), \text{ as } \epsilon \rightarrow 0, \end{aligned}$$

and

$$\begin{aligned} H''_{\epsilon,2} &:= \frac{C}{|\ln \epsilon|^2} \int_{\epsilon < |y'| < 1} \frac{1}{|y'|^{2\tilde{\alpha}}} \int_{\{|x'| > 1\} \cap \{|x' - y'| \geq \frac{1}{2}\}} \frac{\left| \ln \frac{|y'|}{\epsilon} - \ln \frac{1}{\epsilon} \right|^2}{|x' - y'|^{k+2s}} dy' dx' \\ &\leq \frac{C}{|\ln \epsilon|^2} \int_{\epsilon < |y'| < 1} \frac{\ln^2 |y'|}{|y'|^{2\tilde{\alpha}}} dy' = \frac{C}{|\ln \epsilon|^2} \int_{\epsilon}^1 \frac{\ln^2 r}{r^{1-\alpha'}} dr \\ &= o(1), \text{ as } \epsilon \rightarrow 0, \end{aligned}$$

where $\alpha' = k - 2\tilde{\alpha} \geq 2s$. Hence, from (7.5) we have $H_{\epsilon,2} = o(1)$, as $\epsilon \rightarrow 0$.

Step 3: In this step, we will show that $H_{\epsilon,4} = o(1)$, as $\epsilon \rightarrow 0$. Similarly, considering different regions, we see that, it is enough to show the following:

$$H_{\epsilon,4,1} := \frac{1}{|\ln \epsilon|^2} \int \int_F \frac{|\ln |x'| - \ln |y'||^2}{|x' - y'|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx' dy' = o(1), \text{ as } \epsilon \rightarrow 0, \quad (7.6)$$

where F is defined as follows:

$$F := \{(x', y') : \epsilon^2 < |x'| \leq |y'| < \epsilon \text{ and } |y'| < 2|x'|\}.$$

Clearly, $F \subset \{(x', y') : \epsilon^2 < |x'| \leq |y'| < \epsilon \text{ and } |x' - y'| < 3|x'|\}$. So, using, $\ln r \leq r - 1$, for $r \geq 1$, we estimate

$$\begin{aligned} H_{\epsilon,4,1} &\leq \frac{1}{|\ln \epsilon|^2} \int_{\epsilon^2 < |x'| < \epsilon} \frac{1}{|x'|^{k-2s+2}} \int_{|x' - y'| < 3|x'|} \frac{dy'}{|x' - y'|^{k+2s-2}} dx' \\ &= \frac{C}{|\ln \epsilon|^2} \int_{\epsilon^2 < |x'| < \epsilon} \frac{dx'}{|x'|^k} = o(1), \text{ as } \epsilon \rightarrow 0. \end{aligned}$$

Combining Step 1, Step 2 and Step 3 we conclude the lemma. \square

In light of the Lemma 7.1, it is enough to prove that $C_c^\infty(\mathbb{R}^N)$ is dense in \mathcal{W} to conclude Lemma 2.1. The following Lemma shows that, we can approximate $u \in \mathcal{W}$ by a sequence of compactly supported functions lying in \mathcal{W} .

Lemma 7.2. *Let $u \in \mathcal{W}$, $0 < \tilde{\alpha} \leq \frac{k-2s}{2}$ and $\eta \in C_c^\infty(B_2^N(0); [0, 1])$ such that $\eta = 1$ in $B_1^N(0)$ and $\eta_j(x) = \eta(x/j)$. Then*

$$\lim_{j \rightarrow \infty} \left[\|u - \eta_j u\|_{2^*, \tilde{\alpha}, \mathbb{R}^N} + \|[u - \eta_j u]\|_{s, \tilde{\alpha}, \mathbb{R}^N} \right] = 0.$$

Proof. We define

$$I_j := \int_{\mathbb{R}^N} \int_{\mathbb{R}^N \setminus B_j^N(0)} \frac{|u(y)|^2 |\eta_j(x) - \eta_j(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dy dx.$$

Since, $\eta = 1$ on $B_1^N(0)$, so, to prove the Lemma, it is enough to prove, $\lim_{j \rightarrow \infty} I_j = 0$. We define the following sets

$$\begin{aligned} D_{j,0} &:= \{(x, y) \in \mathbb{R}^N \times (\mathbb{R}^N \setminus B_j^N(0)) : |x| \leq |y|/2\}, \\ D_{j,1} &:= \{(x, y) \in \mathbb{R}^N \times (\mathbb{R}^N \setminus B_j^N(0)) : |x| \geq |y|/2 \text{ and } |x-y| \geq j\}, \\ D_{j,2} &:= \{(x, y) \in \mathbb{R}^N \times (\mathbb{R}^N \setminus B_j^N(0)) : |x| \geq |y|/2 \text{ and } |x-y| \leq j\}. \end{aligned}$$

For $m = 0, 1, 2$, we write

$$I_{j,m} := \int \int_{D_{j,m}} \frac{|u(y)|^2 |\eta_j(x) - \eta_j(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dy dx.$$

Then

$$I_j = I_{j,0} + I_{j,1} + I_{j,2}. \quad (7.7)$$

Now, we break

$$\frac{|u(y)|^2 |\eta_j(x) - \eta_j(y)|^2}{|x-y|^{N+2s} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} = \frac{|\eta_j(x) - \eta_j(y)|^2}{|x-y|^{2s+2\sigma_m}} \frac{|u(y)|^2}{|x-y|^{N-2\sigma_m} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}},$$

where $\sigma_0 = s$, $\sigma_1 \in (0, s)$ and $s < \sigma_2 < 1$ such that $\frac{N(N-2\sigma_2)}{N-2s} > \max\{N-k, k\}$. We denote, $\sigma'_m := \frac{N-2\sigma_2}{N-2s}$. Then using Hölder inequality

$$I_{j,m} \leq \left(\int \int_{D_{j,m}} \frac{|\eta_j(x) - \eta_j(y)|^{\frac{N}{s}}}{|x-y|^{N+\sigma_m \frac{N}{s}}} dx dy \right)^{\frac{2s}{N}} \left(\int \int_{D_{j,m}} \frac{|u(y)|^{2^*}}{|x-y|^{N\sigma'_m} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy \right)^{\frac{N-2s}{N}}. \quad (7.8)$$

Clearly,

$$\int \int_{D_{j,m}} \frac{|\eta_j(x) - \eta_j(y)|^{\frac{N}{s}}}{|x-y|^{N+\sigma_m \frac{N}{s}}} dx dy \leq j^{(s-\sigma_m) \frac{N}{s}} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\eta_j(x) - \eta_j(y)|^{\frac{N}{s}}}{|x-y|^{N+\sigma_m \frac{N}{s}}} dx dy \leq C j^{\frac{(s-\sigma_m)N}{s}}. \quad (7.9)$$

Now, we consider

$$\begin{aligned} \int \int_{D_{j,0}} \frac{|u(y)|^{2^*}}{|x-y|^{N\sigma'_0} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy &\leq \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{\frac{2^*\tilde{\alpha}}{2}}} \int_{|x|<\frac{|y|}{2}} \frac{dx}{|x-y|^N |y'|^{\frac{2^*\tilde{\alpha}}{2}}} dy \\ &\leq C \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{\frac{2^*\tilde{\alpha}}{2}} |y|^{\frac{\tilde{\alpha} 2^*}{2}}} dy \leq C \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{\alpha 2^*}} dy. \end{aligned} \quad (7.10)$$

Since, $N\sigma'_1 > N > N-k$, we estimate

$$\begin{aligned} \int \int_{D_{j,1}} \frac{|u(y)|^{2^*}}{|x-y|^{N\sigma'_1} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy &\leq \int_{|y|>j} \int_{\substack{\{|x-y|>j\} \\ \cap \{|x'| \leq \frac{|y|}{2}\}}} \frac{|u(y)|^{2^*}}{|x-y|^{N\sigma'_1} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy \\ &+ \int_{|y|>j} \int_{\substack{\{|x-y|>j\} \\ \cap \{|x'| \geq \frac{|y|}{2}\}}} \frac{|u(y)|^{2^*}}{|x-y|^{N\sigma'_1} |x'|^{\tilde{\alpha}} |y'|^{\tilde{\alpha}}} dx dy \end{aligned}$$

$$\begin{aligned}
&\leq C \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{\frac{\tilde{\alpha}2^*}{2}}} \int_{|x'|<\frac{|y'|}{2}} \frac{dx'}{|x'-y'|^{N\sigma'_1-N+k}|x'|^{\frac{\tilde{\alpha}2^*}{2}}} dy' \\
&+ C \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{2^*\tilde{\alpha}}} \int_{|x-y|>j} \frac{dx}{|x-y|^{N\sigma'_1}} dy \\
&\leq C \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{\tilde{\alpha}2^*}} \frac{dy}{|y'|^{N\sigma'_1-N}} \\
&+ C \frac{1}{j^{N\sigma'_1-1}} \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{2^*\tilde{\alpha}}} dy \\
&\leq \frac{C}{j^{\frac{2N(s-\sigma_1)}{N-2s}}} \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{2^*\tilde{\alpha}}} dy. \tag{7.11}
\end{aligned}$$

Similarly, using $N > N\bar{\sigma}_2 > \max\{N-k, k\}$, we can derive

$$\int \int_{D_{j,2}} \frac{|u(y)|^{2^*}}{|x-y|^{N\sigma'_1}|x'|^{\tilde{\alpha}}|y'|^{\tilde{\alpha}}} dx dy \leq \frac{C}{j^{\frac{2N(s-\sigma_2)}{N-2s}}} \int_{|y|>j} \frac{|u(y)|^{2^*}}{|y'|^{2^*\tilde{\alpha}}} dy. \tag{7.12}$$

Hence, plugging (7.9), (7.10), (7.11) and (7.12) into (7.8) and then using (7.7) we get

$$I_j \leq C \|u\|_{L^{2^*}(\mathbb{R}^N \setminus B_j^N(0); \frac{1}{|x'|^{2\tilde{\alpha}}})}^2 \rightarrow 0, \text{ as } j \rightarrow \infty.$$

This proves the lemma. \square

The next proposition is a reminiscence of the fact, that Θ is in A_1 . Although, in this case, the proof is a direct consequence of Proposition 4.1 and 4.2 of [11].

Proposition 7.3. *There exists a constant $C > 0$ such that for every $X \in \mathbb{R}_k^N \times \mathbb{R}_k^N$, when $N' = 2N$ and $X \in \mathbb{R}_k^N$, when $N' = N$, the following inequality is true*

$$\sup_{r>0} \frac{1}{r^{N'}} \int_{B_r^N(0)} \frac{dz}{\Theta(X+w(z))} \leq \frac{C}{\Theta(X)}.$$

Using Proposition 7.3 and the fact, that the measure $\frac{dX}{\Theta(X)}$, is finite on compact sets of $\mathbb{R}^{N'}$, we can derive the following lemma which is related to the boundedness of the maximal operator.

Lemma 7.4. *Let $q > 1$ and $V : \mathbb{R}^{N'} \rightarrow \mathbb{R}$ be a measurable function. Then, for any $r > 0$,*

$$\int_{\mathbb{R}^{N'}} \left(\frac{1}{r^{n'}} \int_{B_r^{N'}(0)} |V(X-w(z))| dz \right)^q \frac{dX}{\Theta(X)} \leq C \int_{\mathbb{R}^{N'}} \frac{|V(X)|^q}{\Theta(X)},$$

for some constant $C > 0$.

Next, for $V : \mathbb{R}^{N'} \rightarrow \mathbb{R}$ measurable, we define the following operator

$$V \star \eta_0(X) := \int_{\mathbb{R}^N} V(X-w(z))\eta_0(z)dz,$$

where η_0 is a radially symmetric mollifier in \mathbb{R}^N , with $\eta_0 \geq 0$ and $\text{supp } \eta_0 \subset B_1^N(0)$. Notice that, when $N' = N$, $V \star \eta_0$ coincides with the usual convolution operator $V * \eta_0$. As a consequence of Lemma 7.4, we could control appropriate weighted L^p norm of $V \star \eta_0$. More precisely, we could derive the following proposition.

Proposition 7.5. *There exists a constant $C > 0$, such that for any measurable function $V : \mathbb{R}^{N'} \rightarrow \mathbb{R}$ we have*

$$\int_{\mathbb{R}^{N'}} |V \star \eta_0|^p \frac{dX}{\Theta(X)} \leq C \int_{\mathbb{R}^{N'}} |V(X)|^p \frac{dX}{\Theta(X)},$$

where $p = 2$, when $N' = 2N$ and $p = 2^*$, when $N' = N$.

7.2. Proof Of Lemma 2.1.

Proof. We define

$$L^p(\mathbb{R}^{N'}; \Theta) := \{V : \mathbb{R}^{N'} \rightarrow \mathbb{R} \text{ measurable} : \int_{\mathbb{R}^{N'}} |V(X)|^p \frac{dX}{\Theta(X)} < \infty\},$$

where p is defined in the Proposition 7.5. Then, since $\frac{dX}{\Theta(X)}$ is finite on compact sets of $\mathbb{R}^{N'}$, so using Lusin's theorem and Proposition 7.5, we can prove that $C_c^\infty(\mathbb{R}^{N'})$ is dense in $L^p(\mathbb{R}^{N'}; \Theta)$. As a consequence of this density and Proposition 7.2 and the fact, that for any $u \in \mathcal{W}$ and $\eta \in C_c^\infty(\mathbb{R}^N)$, $V^u \star \eta = V^{u \star \eta}$, we can prove that $C_c^\infty(\mathbb{R}^N)$ is dense in \mathcal{W} , where $V^u(x, y) := \frac{u(x) - u(y)}{|x - y|^{\frac{N}{2} + s}}$, for $x, y \in \mathbb{R}^N$. This proves that, $\dot{\mathcal{H}}^{s, \alpha}(\mathbb{R}^N) = \mathcal{W}$, which is exactly what we wanted to prove in Lemma 2.1. \square

ACKNOWLEDGMENTS

I would like to thank my Ph.D supervisor Prof. K. Sandeep for countless valuable discussions and suggestions.

REFERENCES

- [1] Boumediene Abdellaoui, María Medina, Ireneo Peral, and Ana Primo. The effect of the Hardy potential in some Calderón-Zygmund properties for the fractional Laplacian. *J. Differential Equations*, 260(11):8160–8206, 2016.
- [2] Marino Badiale and Gabriella Tarantello. A Sobolev-Hardy inequality with applications to a nonlinear elliptic equation arising in astrophysics. *Arch. Ration. Mech. Anal.*, 163(4):259–293, 2002.
- [3] Albert Baernstein, II. A unified approach to symmetrization. In *Partial differential equations of elliptic type (Cortona, 1992)*, Sympos. Math., XXXV, pages 47–91. Cambridge Univ. Press, Cambridge, 1994.
- [4] H. Berestycki and L. Nirenberg. On the method of moving planes and the sliding method. *Bol. Soc. Brasil. Mat. (N.S.)*, 22(1):1–37, 1991.
- [5] Krzysztof Bogdan and Bartłomiej Dyda. The best constant in a fractional Hardy inequality. *Math. Nachr.*, 284(5-6):629–638, 2011.
- [6] Haï m Brézis and Elliott and Lieb. A relation between pointwise convergence of functions and convergence of functionals. *Proc. Amer. Math. Soc.*, 88(3):486–490, 1983.
- [7] Luis Caffarelli and Luis Silvestre. An extension problem related to the fractional Laplacian. *Comm. Partial Differential Equations*, 32(7-9):1245–1260, 2007.
- [8] Daniele Castorina, Isabella Fabbri, Gianni Mancini, and Kunnath Sandeep. Hardy-Sobolev inequalities and hyperbolic symmetry. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 19(3):189–197, 2008.
- [9] Eleonora Di Nezza, Giampiero Palatucci, and Enrico Valdinoci. Hitchhiker's guide to the fractional Sobolev spaces. *Bull. Sci. Math.*, 136(5):521–573, 2012.
- [10] Serena Dipierro, Luigi Montoro, Ireneo Peral, and Berardino Sciunzi. Qualitative properties of positive solutions to nonlocal critical problems involving the Hardy-Leray potential. *Calc. Var. Partial Differential Equations*, 55(4):Art. 99, 29, 2016.
- [11] Serena Dipierro and Enrico Valdinoci. A density property for fractional weighted Sobolev spaces. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 26(4):397–422, 2015.
- [12] Bartłomiej Dyda, Juha Lehrbäck, and Antti V. Vähäkangas. Fractional Hardy-Sobolev type inequalities for half space and John domain. *arXiv: 1709.03296v1.*, 2017.
- [13] Mouhamed Moustapha Fall and Tobias Weth. Nonexistence results for a class of fractional elliptic boundary value problems. *J. Funct. Anal.*, 263(8):2205–2227, 2012.
- [14] PATRICIO FELMER and YING WANG. Radial symmetry of positive solutions to equations involving the fractional laplacian. *Communications in Contemporary Mathematics*, 16(01):1350023, 2014.

- [15] Rupert L. Frank, Elliott H. Lieb, and Robert Seiringer. Hardy-Lieb-Thirring inequalities for fractional Schrödinger operators. *J. Amer. Math. Soc.*, 21(4):925–950, 2008.
- [16] Rupert L. Frank and Robert Seiringer. Non-linear ground state representations and sharp Hardy inequalities. *J. Funct. Anal.*, 255(12):3407–3430, 2008.
- [17] Rupert L. Frank and Robert Seiringer. Sharp fractional Hardy inequalities in half-spaces. In *Around the research of Vladimir Maz'ya. I*, volume 11 of *Int. Math. Ser. (N. Y.)*, pages 161–167. Springer, New York, 2010.
- [18] Marita Gazzini and Roberta Musina. Hardy-Sobolev-Maz'ya inequalities: symmetry and breaking symmetry of extremal functions. *Commun. Contemp. Math.*, 11(6):993–1007, 2009.
- [19] Marita Gazzini and Roberta Musina. On a Sobolev-type inequality related to the weighted p -Laplace operator. *J. Math. Anal. Appl.*, 352(1):99–111, 2009.
- [20] Nassif Ghoussoub and Shaya Shakerian. Borderline variational problems involving fractional Laplacians and critical singularities. *Adv. Nonlinear Stud.*, 15(3):527–555, 2015.
- [21] V. P. Il'in. Some integral inequalities and their applications in the theory of differentiable functions of several variables *Mat. Sb. (N.S.)*, 54(96): 331–380,1961.
- [22] Tianling Jin, YanYan Li, and Jingang Xiong. On a fractional Nirenberg problem, part I: blow up analysis and compactness of solutions. *J. Eur. Math. Soc. (JEMS)*, 16(6):1111–1171, 2014.
- [23] Moritz Kassmann. A priori estimates for integro-differential operators with measurable kernels. *Calc. Var. Partial Differential Equations*, 34(1):1–21, 2009.
- [24] P.-L. Lions. The concentration-compactness principle in the calculus of variations. The locally compact case. I. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 1(2):109–145, 1984.
- [25] P.-L. Lions. The concentration-compactness principle in the calculus of variations. The locally compact case. II. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 1(4):223–283, 1984.
- [26] G. Mancini, I. Fabbri, and K. Sandeep. Classification of solutions of a critical Hardy-Sobolev operator. *J. Differential Equations*, 224(2):258–276, 2006.
- [27] Gianni Mancini and Kunnath Sandeep. On a semilinear elliptic equation in \mathbb{H}^n . *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 7(4):635–671, 2008.
- [28] Vladimir G. Maz'ja. *Sobolev spaces*. Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1985. Translated from the Russian by T. O. Shaposhnikova.
- [29] Roberta Musina. Ground state solutions of a critical problem involving cylindrical weights. *Nonlinear Anal.*, 68(12):3972–3986, 2008.
- [30] R. Musina and A. I. Nazarov. Fractional Hardy-Sobolev inequalities on half spaces. *ArXiv e-prints*, July 2017.
- [31] Giampiero Palatucci and Adriano Pisante. Improved Sobolev embeddings, profile decomposition, and concentration-compactness for fractional Sobolev spaces. *Calc. Var. Partial Differential Equations*, 50(3-4):799–829, 2014.
- [32] Luis Silvestre. Regularity of the obstacle problem for a fractional power of the Laplace operator. *Comm. Pure Appl. Math.*, 60(1):67–112, 2007.
- [33] A. Tertikas and K. Tintarev. On existence of minimizers for the Hardy-Sobolev-Maz'ya inequality. *Ann. Mat. Pura Appl. (4)*, 186(4):645–662, 2007.
- [34] Konstantinos Tzirakis. Sharp trace Hardy-Sobolev inequalities and fractional Hardy-Sobolev inequalities. *J. Funct. Anal.*, 270(12):4513–4539, 2016.