

# NNLO classical solution for Lipatov's effective action for reggeized gluons

S. Bondarenko, S. Pozdnyakov

*Physics Department, Ariel University, Ariel, Israel*

## Abstract

We consider the formalism of small- $x$  effective action for reggeized gluons, [1, 2, 3], and, following to the approach developed in [4, 5], calculate the classical gluon field to NNLO precision with fermion loops included. It is demonstrated, that the self-consistency of the equations of motion in each perturbative order in the approach is equivalent to the transversality conditions applied to the solutions of the equations in the lower orders, that allows to construct the solutions with the help of some recursive scheme. Applications of the obtained results are also discussed.

## 1 Introduction

In the framework of small- $x$  Lipatov's effective action, [2], the calculation of the amplitudes of high-energy scattering can be done by two different methods. The first one, which we can call as diagrammatic, see [6, 7, 8, 9], is based on the use of effective vertices of reggeons-gluons interactions obtained from the action and subsequent construction of the amplitudes with the use of the vertices. Another method, proposed as well in [2] and developed in [4, 5], is based on the formalism of the effective action. In this case, the field theory is constructed on the base of the fluctuations around the classical solutions of motion. Because of the dependence of the classical gluon fields on the reggeon fields, the result obtained is revealed as the Regge Field Theory (RFT), see [5, 10, 11], where the new RFT action is appearing as a functional of the reggeon fields only. In this form the action can be served as a generating functional of the reggeon-reggeon interactions and can be used also for the calculations of the reggeon loops arising in the RFT. This reformulation of the effective action for reggeized gluons as RFT can be considered as some generalization of Gribov's Regge calculus, [1], for the case of QCD degrees of freedom. The validity of the effective action approach was confirmed in different calculations, see [12, 13].

Therefore, the use of the approach in the form considered in [4] requires the knowledge of the classical solutions of the equations of motion for the gluon fields. In the perturbative scheme, used in [5], each term of the classical solution represents some bare vertex of reggeon-reggeon interactions, which will affect on the calculations of the gluons loops corrections as well. In the calculation of the NLO<sup>1</sup> propagator of reggeized gluons the only LO precision for the classical solutions is required, see [4]. Nevertheless, the calculation of the different vertices of reggeon-reggeon interactions with NLO and NNLO precision will require the NLO and NNLO precision for the classical gluon fields, this problem is considered in the present paper. We note also, that the classical solution with NNLO precision consists with fermion loops contribution as well, see [14].

The paper is organized as following. In the Section 2 we remind the main ideas of the formalism of the Lipatov's effective action, main results of [4] and discuss the self-consistency of the equations of motion and its relations to the transversality condition for a certain combination of fields. In Sections 3, 4 and 5 the calculations of the classical solutions for longitudinal and transverse components of the gluon field are represented. Section 6 is dedicated to the inclusion of the quark loop contribution in the approach. The last section is the Conclusion of the paper.

---

<sup>1</sup> In the RFT approach the LO contribution is represented by a bare action of the reggeized gluons, whereas the NLO contribution is given by RFT action with one gluon loops included, i.e. by the action with contribution of the terms quadratic on fluctuations in the Lipatov's effective action included.

## 2 Effective action for reggeized gluons with color field source

The Lipatov effective action, see [2, 15], is a non-linear gauge invariant action which describes clusters of particles in correspondence with their rapidities  $y = \frac{1}{2} \ln \left( \frac{p_+}{p_-} \right)$  whereas the interaction between the clusters with essentially different rapidities is realized by the reggeon fields exchange. The action has the following form

$$L = -\frac{1}{4} G_{\nu\mu}^a G_a^{\nu\mu} - v_+ J^+ - v_- J^- + L_{quark}, \quad (1)$$

where the quark part of the Lagrangian is given by

$$L_{quark} = \sum_{c=1}^{n_f} \bar{\psi}_c (i\gamma^\nu \partial_\nu - m + g\gamma^\nu T^a v_{a\nu}) \psi_c. \quad (2)$$

Under variation on the gluon fields these currents reproduce the Lipatov's induced currents

$$\delta(v_\pm J^\pm) = \delta(v_\pm) j^\pm, \quad (3)$$

where

$$j_a^\pm = -\text{tr}(f_a j^\pm) = \frac{1}{N} \text{tr}(f_a O(v_\pm) f_b O^T(v_\pm)) (\partial_i^2 A_\mp). \quad (4)$$

There are additional kinematical constraints for the reggeon fields

$$\partial_- A_+ = \partial_+ A_- = 0, \quad (5)$$

corresponding to the strong-ordering Sudakov components in the multi-Regge kinematics, see [2] and also [15].

The equations of motion in light-cone gauge ( $v_-^a = v_+^a = 0$ ) provided by Eq. (1) Lagrangian (without the quark term, see Section 6), have the following form:

$$(D_\mu G^{\mu\nu})_a = \partial_\mu G_a^{\mu\nu} + g f_{abc} v_\mu^b G^{c\mu\nu} = j_a^+ \delta^{\nu+} + j_a^- \delta^{\nu-}. \quad (6)$$

We solve these equations perturbatively, order by order, with the help of the usual perturbative expansion of the gluon fields:

$$v_+^a = \sum_{k=0}^{+\infty} g^k v_{+k}^a, \quad v^{ia} = \sum_{k=0}^{+\infty} g^k v_k^{ia}. \quad (7)$$

Here, among of four equations for three fields, one from the equations provides us with a transversality condition for a certain combination of fields. If this condition is satisfied, then the classical solutions for the fields exist and they have a very simple structure. Indeed, the same Eq. (6) we can rewrite in an another form. Due the covariant derivative presence, we can separate the terms in the equations and rewrite the equations as some recurrence relations between the different terms of the Eq. (7) series. Namely, the fields of  $k$  perturbative order will stay in the left-hand side of the equations and the terms which consists with the fields of perturbative order less than  $k$  but multiplied on the corresponding order of coupling constant  $g$  and denoted as  $\bar{j}_{k-1}^+$ ,  $\bar{j}_{k-1}^-$  and  $\bar{j}_{k-1}^i$  will stay on the right hand side of equations. Therefore, we obtain:

$$-\partial_- [\partial_i v_k^i + \partial_- v_{+k}] = \bar{j}_{k-1}^+, \quad (8)$$

$$\square v_k^j - \partial^j [\partial_i v_k^i + \partial_- v_{+k}] = \bar{j}_{k-1}^j, \quad (9)$$

$$\square v_{+k} - \partial_+ [\partial_i v_k^i + \partial_- v_{+k}] = \bar{j}_{k-1}^-. \quad (10)$$

Taking derivatives of the equations in correspondence to the indexes of the r.h.s of them and summing up the l.h.s. of the obtained expressions we obtain that the self-consistency of the solutions of the Eq. (8)-Eq. (10) is equivalent to the condition of the transversality being imposed on the r.h.s of the sum:

$$\partial_\mu \bar{j}_{k-1}^\mu = 0, \quad (11)$$

that means that the classical solutions of the lower perturbative orders and Eq. (4) induced current substituted in  $\bar{j}_{k-1}^\mu$  functions must have the form which satisfies the Eq. (11) condition. This conditions, therefore, provide a useful way for the constructions and check of the classical solutions of the equations of motion. Indeed, thus we have:

$$v_k^j = \square^{-1} [\bar{j}_{k-1}^j - \partial^j \partial_-^{-1} \bar{j}_{k-1}^+], \quad (12)$$

$$v_{+k} = \square^{-1} \left[ \bar{j}_{k-1}^- - \partial_+ \partial_-^{-1} \bar{j}_{k-1}^+ \right]. \quad (13)$$

Therefore, there are the LO and NLO expressions for the classical fields obtained in [4]:

$$v_{+0a} = A_{+a}, \quad (14)$$

$$v_{i0a} = \rho_i^b(x_\perp, x^-) U_{ab}, \quad (15)$$

$$v_{+1a} = -\frac{2}{g} \square^{-1} \left( (\partial_+ \partial^i U_{ab}) \rho_i^b \right), \quad (16)$$

$$v_{i1a} = -\square^{-1} \left[ \partial^j F_{jia} + \frac{1}{g} \partial_i \left( (\partial^j U_{ab}) \rho_j^b \right) - \partial_i \partial_-^{-1} j_{a1}^+ \right], \quad (17)$$

where

$$\rho_a^i = -\frac{1}{N} \partial_-^{-1} (\partial^i A_{-a}), \quad (18)$$

$$U^{ab}(v_+) = \text{tr} \left[ f^a \left( P e^{g \int_{-\infty}^{x^+} dx'^+ v_{+c}(x'^+, x^-, x_\perp) f^c} \right) f^b \left( P e^{g \int_{x^+}^{+\infty} dx'^+ v_{+d}(x'^+, x^-, x_\perp) f^d} \right) \right], \quad (19)$$

see Appendix A for details. To NNLO precision, we write our Eq. (7) ansatz in the following form:

$$v_+^a = A_+^a(x_\perp, x^+) + g v_{+1}^a(x_\perp, x^-, x^+) + g^2 v_{+2}^a(x_\perp, x^-, x^+), \quad (20)$$

$$v_i^a = v_{i0}^a(x_\perp, x^-, x^+) + g v_{i1}^a(x_\perp, x^-, x^+) + g^2 v_{i2}^a(x_\perp, x^-, x^+), \quad (21)$$

the calculations of these NNLO terms of the gluon fields are presented in the next Sections.

### 3 First equation of motion

In this section we consider the equation of motion arising after the variation of the Lagrangian with respect to  $v_{+a}$  field. We obtain:

$$\partial_\mu G_a^{\mu+} + g f_{abc} v_\mu^b G^{c\mu+} = j_a^+ \quad (22)$$

or

$$\partial_i G_a^{i+} + \partial_- G_a^{-+} + g f_{abc} v_i^b G^{ci+} = j_a^+, \quad (23)$$

where the current in the r.h.s. is given by Eq. (4). Using

$$G_a^{\mu+} = \partial^\mu v_a^+ - \partial^+ v_a^\mu + g f_{abc} v^{\mu b} v^{+c} = -\partial^+ v_a^\mu, \quad (24)$$

we obtain for Eq. (23) the following equation:

$$-\partial_i \partial^+ v_a^i - \partial_- \partial^+ v_a^- - g f_{abc} v_i^b \partial^+ v^{ic} = j_a^+. \quad (25)$$

Now we substitute inside of it the perturbative expressions of Eq. (14)-Eq. (17) obtaining the following terms.

1. The first term in the l.h.s of Eq. (25):

$$-\partial^i \partial^+ v_i^a = -\partial^i \partial_- v_{i0}^a - g \partial^i \partial_- v_{i1}^a - g^2 \partial^i \partial_- v_{i2}^a. \quad (26)$$

With the use of Eq. (15) expressions we obtain:

$$-\partial^i \partial_- v_{i0}^a = -(\partial^i \partial_- \rho_i^b) U_{ab} - g (\partial_- \rho_i^b) \left( \frac{1}{g} \partial^i U_{ab} \right) - g^2 (\partial^i \rho_i^b) \left( \frac{1}{g^2} \partial_- U_{ab} \right) - g^2 \rho_i^b \left( \frac{1}{g^2} \partial^i \partial_- U_{ab} \right). \quad (27)$$

The first term in the r.h.s of Eq. (27) is the same as the r.h.s of Eq. (25), they both have the  $g^0$  order precision as it must be for the this order classical solution.

2. The second term:

$$-\partial_- \partial^+ v_a^- = -\partial_- \partial^+ \left( A_{+a} + g v_{+1a} + g^2 v_{+2a} \right) = -\partial_- \partial^+ \left( g v_{+1a} + g^2 v_{+2a} \right) \quad (28)$$

and consists with the  $g$  and  $g^2$  order terms only.

3. The third term in Eq. (25) we write in the following form:

$$-gf_{abc}v_i^b\partial^+v^{ic} = gj_{a1}^+ + g^2L_{a2}^+(A_+, \rho) + O(g^3), \quad (29)$$

where

$$gj_{a1}^+ = -gf_{abc}(U^{bb'}\rho_{b'}^i)(U^{cc'}(\partial_-\rho_{ic'})) \quad (30)$$

and

$$g^2L_{a2}^+(A_+, \rho) = -g^2f_{abc}(v_{i0}^b\partial_-v_1^{ic} + v_{i1}^b\partial_-v_0^{ic}). \quad (31)$$

Summing up these three terms and rewriting the equation in the form of Eq. (8) we obtain:

$$-g^2\partial_- \left[ \partial^i v_{i2a} + \partial_- v_{+2a} \right] = g^2 \left[ (\partial^i \rho_i^b) \left( \frac{1}{g^2} \partial_- U_{ab} \right) - L_{a2}^+ - \square^{-1} \left[ \frac{2}{g} \partial_+ \partial_- j_{a1}^+ \right] \right] \quad (32)$$

or

$$\left[ \partial^i v_{i2a} + \partial_- v_{+2a} \right] = \partial_-^{-1} L_{a2}^+ + \square^{-1} \left[ \frac{2}{g} \partial_+ j_{a1}^+ \right] - \partial_-^{-1} \left( (\partial^i \rho_i^b) \left( \frac{1}{g^2} \partial_- U_{ab} \right) \right), \quad (33)$$

where the r.h.s. of the expression consists with the classical solutions of  $k = 0$  and  $k = 1$  orders.

## 4 The second and third equations of motion

The variation of the action with respect to  $v_{ia}$  gives

$$(D_+ G^{+i})_a + (D_- G^{-i})_a + (D_j G^{ji})_a = 0 \quad (34)$$

or

$$\square v_a^i - \partial^i \left[ \partial^j v_{ja} + \partial_- v_{+a} \right] + (\partial F)_a^i = 0, \quad (35)$$

where

$$(\partial F)_a^i = gf_{abc} \left( v_+^b \partial_- v^{ic} + \partial_- (v_-^b v^{ic}) + \partial_j (v^{jb} v^{ic}) + v^{jb} \partial_j v^{ic} - v_j^b \partial^i v^{jc} + g f^{cde} v^{jb} v_d^j v_e^i \right). \quad (36)$$

The Eq. (35) can be represented in the form of equation Eq. (9) and subsequently it can be written in the form of Eq. (12). Then, the only unknown functions in Eq. (35) are to the  $g^2$  order, we denote them as  $g^2 C_a^i$ . Calculation of this contribution is made in appendix B, finally the result for the classical solution of the transverse gluon fields to NNLO precision reads as

$$\begin{aligned} \square v_{2a}^i - \partial^i \left[ \partial_-^{-1} L_{a2}^+ - \partial_-^{-1} \left( (\partial^j \rho_j^b) \left( \frac{1}{g^2} \partial_- U_{ab} \right) \right) \right] + f_{abc} \left( v_{1+}^b \partial_- v_0^{ic} + v_{0+}^b \partial_- v_1^{ic} + \partial_- (v_{1+}^b v_0^{ic} + v_{0+}^b v_1^{ic}) \right. \\ \left. + \partial_j (v_0^{jb} v_1^{ic} + v_1^{jb} v_0^{ic}) + v_{0j}^b (\partial^j v_1^{ic} - \partial^i v_1^{jc}) + v_{1j}^b (\partial^j v_0^{ic} - \partial^i v_0^{jc}) + f^{cde} v_{j0}^b v_{0d}^j v_{0e}^i \right) + C_a^i = 0 \end{aligned} \quad (37)$$

or

$$\begin{aligned} v_{2a}^i &= \square^{-1} \left[ \partial_-^{-1} \partial^i \left( L_{a2}^+ - (\partial^j \rho_j^b) \left( \frac{1}{g^2} \partial_- U_{ab} \right) \right) - f_{abc} \left( \frac{1}{g} v_{j0}^b (\partial^j v_0^{ic} - \partial^i v_0^{jc}) + v_{0j}^b (\partial^j v_1^{ic} - \partial^i v_1^{jc}) \right. \right. \\ &\quad \left. \left. - v_0^{ic} \partial_- v_{1+}^b + 2A_+^b \partial_- v_1^{ic} + \partial_j (v_0^{jb} v_1^{ic} - v_0^{ib} v_1^{jc}) + f^{cde} v_{j0}^b v_{0d}^j v_{0e}^i \right) \right]. \end{aligned} \quad (38)$$

In turn, the substitution of Eq. (38) in Eq. (33) provides the classical solution for the longitudinal gluon field:

$$\begin{aligned} \partial_- v_{+2a} &= 2\square^{-1} \partial_+ \left[ L_{a2}^+ - (\partial^j \rho_j^b) \left( \frac{1}{g^2} \partial_- U_{ab} \right) + \frac{1}{g} j_{a1}^+ \right] + f_{abc} \square^{-1} \partial_i \left( \frac{1}{g} v_{j0}^b (\partial^j v_0^{ic} - \partial^i v_0^{jc}) \right. \\ &\quad \left. + v_{0j}^b (\partial^j v_1^{ic} - \partial^i v_1^{jc}) - v_0^{ic} \partial_- v_{1+}^b + 2A_+^b \partial_- v_1^{ic} + f^{cde} v_{j0}^b v_{0d}^j v_{0e}^i \right). \end{aligned} \quad (39)$$

This expression can be simplified if we are shortening the following expressions:

$$f^{ij} = -f^{ji}, \quad (\partial_i v_{j0}^b) f^{ij} = \frac{g}{2} \frac{(\partial_i v_{j0}^b - \partial_j v_{i0}^b)}{g} f^{ij} = O(g) \quad (40)$$

and

$$\frac{1}{g} v_{j0}^b \partial_i \left( \partial^j v_0^{ic} - \partial^i v_0^{jc} + g f^{cde} v_{0d}^j v_{0e}^i \right) = -v_{j0}^b \partial_i F_c^{ij}, \quad (41)$$

that gives:

$$v_{0j}^b \partial_i (\partial^j v_1^{ic} - \partial^i v_1^{jc}) = v_{j0}^b \square^{-1} \partial_k \partial^k \partial_i F_c^{ij}. \quad (42)$$

Then we obtain:

$$\begin{aligned} v_{+2a} &= \square^{-1} \partial_-^{-1} \left[ \frac{2}{g} \partial_+ j_{a1}^+ + f_{abc} \left( -v_{i0}^b \partial_- \left[ 2\partial_+ v_1^{ic} - \partial^i v_{1+}^c + 2\square^{-1} \partial_+ \partial_j F_c^{ji} \right] \right. \right. \\ &\quad \left. \left. + 2\partial_- v_{i0}^b \partial_+ v_1^{ic} + (\partial_i v_0^{ic}) \partial_- v_{1+}^b + 2\partial_i (A_+^b \partial_- v_1^{ic}) \right) \right]. \end{aligned} \quad (43)$$

Substituting Eq. (16) and Eq. (17) in Eq. (43) and preserving there only  $g^2$  order terms in we obtain finally:

$$v_{+2a} = \square^{-1} \partial_-^{-1} \left[ \frac{2}{g} \partial_+ j_{a1}^+ + f_{abc} \left( 2\partial_- v_{i0}^b \partial_+ v_1^{ic} + (\partial_i v_0^{ic}) \partial_- v_{1+}^b + 2\partial_i (A_+^b \partial_- v_1^{ic}) \right) \right], \quad (44)$$

where  $j_{a1}^+$  is defined by equation Eq. (30).

## 5 The fourth equation of motion

The variation of the action with respect to  $v_{-a}$  gives

$$\partial_\mu G_a^{\mu-} + g f_{abc} v_\mu^b G^{c\mu-} = j_a^-, \quad (45)$$

which can be rewritten as

$$\partial_i (\partial^i v_a^- - \partial^- v_a^i + g f_{abc} v_b^i v_c^-) + \partial_+ \partial^+ v_a^- + g f_{abc} v_i^b (\partial^i v^{-c} - \partial^- v^{ic} + g f^{cde} v_d^i v_e^-) + g f_{abc} v_+^b \partial^+ v^{-c} = \partial_i \partial^i A_{+a} \quad (46)$$

or

$$\square v_{+a} - \partial_+ \left[ \partial_i v_a^i + \partial_- v_{+a} \right] + g f_{abc} \left( \partial_i (v^{ib} v_+^c) + v_i^b (\partial^i v_+^c - \partial_+ v^{ic} + g f^{cde} v_d^i v_{+e}^-) + v_+^b \partial_- v_+^c \right) = \partial_i \partial^i A_{+a}. \quad (47)$$

Now we can verify that the equations of motion Eq. (25) and Eq. (47) give us the result Eq. (44) for  $v_{+2}$  found above. In the same time this calculation can serve as test of the fulfillment of the condition of the transversality introduced above. The reduction of the equation of motion (47) to the form of Eq. (10) is presented in the appendix C, we write here the result:

$$\begin{aligned} &\square v_{+2a} - \partial_+ \left[ \partial_i v_{2a}^i + \partial_- v_{+2a} \right] \\ &= -f_{abc} \left( -v_{i0}^b \partial_+ v_1^{ic} - 2\partial_i (A_+^b v_1^{ic}) + A_+^b \square^{-1} \partial_i \partial^i \partial_-^{-1} j_{1+}^c \right) + \square^{-1} \partial_i \partial^i \partial_-^{-1} \left( \frac{1}{g} \partial_+ j_{a1}^+ \right). \end{aligned} \quad (48)$$

Then, substituting expression from Eq. (33) and Eq. (31), we obtain

$$\begin{aligned} v_{+2a} &= \square^{-1} \partial_-^{-1} \left[ \left[ \frac{1}{g} \partial_+ j_{a1}^+ \right] - \partial_+ ((\partial^i \rho_i^b) (\frac{1}{g^2} \partial_- U_{ab})) \right. \\ &\quad \left. + f_{abc} \left( \partial_- v_{i0}^b \partial_+ v_1^{ic} + 2\partial_i (A_+^b \partial_- v_1^{ic}) - A_+^b \square^{-1} \partial_i \partial^i j_{1+}^c \right) \right]. \end{aligned} \quad (49)$$

It is easy to show that  $\partial_+ j_{a1}^+ = g f_{abc} f^{cde} A_+^b (v_{i0d} \partial_- v_{0e}^i) = -g f_{abc} A_+^b j_{1+}^{+c}$  up to the second order of  $g$ . We have:

$$\partial_+ j_{a1}^+ = g (f_{aec} f^{edb} + f_{abe} f^{edc}) v_+^b (v_{i0d} \partial_- v_{0c}^i). \quad (50)$$

Using

$$f_{abe} f^{edc} = f_{ade} f^{ebc} - f_{aec} f^{edb}, \quad (51)$$

we, therefore, obtain:

$$\partial_+ j_{a1}^+ = g f_{abe} f^{edc} v_+^b (v_{i0d} \partial_- v_{0c}^i) = -g f_{abc} v_+^b j_{1+}^{+c}, \quad (52)$$

that finally gives the same as Eq. (44) expression.

## 6 One-loop quark corrections

The full NNLO contribution to the classical solutions of the gluon equations of motion must include the contribution from the fermion loop as well, see Eq. (1). Expanding this part of the Lagrangian around  $\psi = \bar{\psi} = 0$ <sup>2</sup> classical solutions we have:

$$L_{quark}^\varepsilon = \sum_{c=1}^{n_f} \bar{\varepsilon}_{\psi c} (i\gamma^\nu \partial_\nu - m + g\gamma^\nu T^a v_{a\nu}) \varepsilon_{\psi c} = \sum_{c=1}^{n_f} \bar{\varepsilon}_{\psi c} (M_0 + M_1) \varepsilon_{\psi c}. \quad (53)$$

The bare quark Green's function of the problem is defined as usual:

$$M_{0x} G_q^0(x, y) = \delta^{(4)}(x - y) \quad (54)$$

and has the following momentum representation:

$$G_q^0(x, y) = (i\gamma_\nu \partial_x^\nu + m) \Delta_{xy}, \quad \Delta_{xy} = \int \frac{d^4 p}{(2\pi)^4} \frac{e^{ip(x-y)}}{p^2 - m^2 + i0}. \quad (55)$$

Below we will use the following properties of the  $\Delta$  function:

$$\partial_x^\nu \Delta_{xy} = -\partial_y^\nu \Delta_{yx}, \quad \partial_y^\mu \partial_x^\nu \Delta_{xy} = \partial_y^\nu \partial_x^\mu \Delta_{xy}. \quad (56)$$

The integration on the fluctuations

$$\begin{aligned} & \int \prod_{b=1}^{n_f} d\bar{\varepsilon}_{\psi b} d\varepsilon_{\psi b} \exp \left[ \sum_{c=1}^{n_f} i\bar{\varepsilon}_{\psi c} (M_0 + M_1) \varepsilon_{\psi c} \right] = \left( \det(M_0 + M_1) \right)^{n_f} = \exp \left[ n_f \text{tr} \ln (M_0 + M_1) \right] \\ & = \exp \left[ n_f \text{tr} \ln (M_0) \right] * \exp \left[ n_f \text{tr} \ln (1 + M_0^{-1} M_1) \right] \end{aligned} \quad (57)$$

results as the usual additional contribution to action:

$$\Gamma_q = -i n_f \text{tr} \left( \ln (1 + M_0^{-1} M_1) \right). \quad (58)$$

Expanding Eq. (58) we preserve the only  $g^2$  order contribution:

$$i n_f \frac{g^2}{4} \text{tr} \left( G_q^0(y, x) (\gamma^\nu v_{a\nu}(x)) G_q^0(x, y) (\gamma^\mu v_\mu^a(y)) \right), \quad (59)$$

that provides the following quark current to the equation of motion:

$$j_{quark\ a}^\rho(z) = i n_f \frac{g^2}{2} \text{tr} \left( \gamma^\rho G_q^0(z, y) \gamma^\mu v_{a\ \mu}(y) G_q^0(y, z) \right). \quad (60)$$

In order to construct additional parts to the classical gluon fields arising from this current we have to verify that the condition  $\partial_\rho j_{quark\ a}^\rho = 0$  is satisfied here. If it does, these additional parts can be calculated simply as

$$v_{q2a}^i = \square^{-1} \left[ j_{quark\ 2a}^i - \partial^i \partial_{-1}^{-1} j_{quark\ 2a}^+ \right] \quad (61)$$

and

$$v_{q+2a} = \square^{-1} \left[ j_{quark\ 2a}^- - \partial_+ \partial_{-1}^{-1} j_{quark\ 2a}^+ \right], \quad (62)$$

see Eq. (12)-Eq. (13). The check of  $\partial_\rho j_{quark\ a}^\rho = 0$  we begin rewriting Eq. (60) in the following form:

$$j_{quark\ a}^\rho(z) = -2i n_f g^2 \left( g^{\rho\nu} g^{\mu\sigma} + g^{\rho\sigma} g^{\mu\nu} - g^{\rho\mu} g^{\sigma\nu} \right) \int d^4 y (\partial_{\nu z} \Delta_{zy}) v_{\mu a}(y) (\partial_{\sigma y} \Delta_{yz}) + 2i n_f g^2 m^2 \int d^4 y \Delta_{zy} v_a^\rho(y) \Delta_{yz}. \quad (63)$$

Then we have

$$\begin{aligned} \partial_{\rho z} j_{quark\ a}^\rho(z) &= -2i n_f g^2 \int d^4 y \left( (\partial_{\rho z} \partial_z^\rho \Delta_{zy}) v_a^\nu(y) (\partial_{\nu y} \Delta_{yz}) + (\partial_{\nu y} \Delta_{yz}) v_a^\nu(y) (\partial_{\rho z} \partial_z^\rho \Delta_{zy}) \right. \\ &+ (\partial_z^\rho \Delta_{zy}) v_a^\nu(y) (\partial_{\nu z} \partial_{\rho y} \Delta_{yz}) + (\partial_{\rho z} \partial_{\nu z} \Delta_{zy}) v_a^\nu(y) (\partial_y^\rho \Delta_{yz}) \\ &\left. - \partial_{\rho z} \left[ (\partial_{\nu z} \Delta_{zy}) v_a^\rho(y) (\partial_y^\nu \Delta_{yz}) \right] - m^2 \partial_{\rho z} \left[ \Delta_{zy} v_a^\rho(y) \Delta_{yz} \right] \right). \end{aligned} \quad (64)$$

<sup>2</sup>In the approach we separate the classical solutions for the quark fields and fluctuations around these classical solution, in general the non-zero classical quark fields correspond to the production amplitudes in the approach.

The second and third lines of this expression are identically equal, and the difference between the first and fourth lines is given by an expression that is also zero:

$$\partial_{\rho z} j_{quark\ a}^{\rho}(z) = -2i n_f g^2 (v_a^{\nu}(z) (\partial_{\nu z} \Delta_{zy})_{y=z} + (\partial_{\nu z} \Delta_{zy})_{y=z} v_a^{\nu}(z)) = 0, \quad (65)$$

that completes the check and justifies the Eq. (61)-Eq. (62) expressions.

Inserting obtained classical gluon fields solutions in the Eq. (1) action, we will obtain the action which will depend only on the reggeon fields. The contribution of the quarks to the reggeon propagator can be calculated similarly to the done in [5] with  $g^2$  order of the contributions preserved. The components of the field strength tensor with LO precision reads as

$$G_{+-0}^a = 0, \quad G_{i+0}^a = \partial_i A_+^a, \quad G_{i-0}^a = -\partial_- v_{i0}^a, \quad G_{ij0} = 0, \quad (66)$$

and with quarks contributions included as

$$G_{+-q2}^a = 0, \quad G_{i+q2}^a = g^2 \partial_i v_{q+2}^a, \quad G_{i-q2}^a = -g^2 \partial_- v_{qi2}^a, \quad G_{ijq2} = 0. \quad (67)$$

It gives for the additional contribution to the Lagrangian of the reggeon fields:

$$\frac{1}{2} G_{\mu\nu 0} G_{q2}^{\mu\nu} = g^2 (\partial_- v_{i0}^a) (\partial_i v_{q+2}^a) + g^2 (\partial_- v_{qi2}^a) (\partial_i A_+^a). \quad (68)$$

Therefore, there is the additions contributions to the effective action to NNLO precision obtained from the inclusion of quarks in the approach:

$$S_{eff\ q2} = -\frac{g^2}{N} \int d^4x ((\partial_i v_{q+2}^a) (\partial_i A_-^b) + N (\partial_i A_+^a) (\partial_- v_{qi2}^b) + v_{q+2}^a (\partial_{\perp}^2 A_-^b)), \quad (69)$$

which can be written as

$$S_{eff\ q2} = -\frac{g^2}{N} \int d^4x \left[ \partial_i (v_{q+2}^a \partial_i A_-^b) + N (\partial_- (v_{qi2}^b \partial_i A_+^a) - v_{qi2}^b (\partial_i \partial_- A_+^a)) \right] = 0, \quad (70)$$

i.e. this additional part of the action does not contribute to the reggeon fields propagator. Nevertheless, there are additional contributions to the propagator arising from the modified classical solutions. We can calculate them following to the calculations of [5] and determining the additional contribution to the kernel of interacting of reggeized gluons:

$$(K_{xy}^{ab})_{q1}^{+-} = \left( \frac{\delta^2 \Gamma_q}{\delta A_{+x}^a \delta A_{-y}^b} \right)_{A_+, A_- = 0}. \quad (71)$$

Using Eq. (53)-Eq. (58), the new contribution to this kernel can be written as

$$-2i n_f g^2 \delta^{ab} \int d^4t d^4z \left[ \partial_{it} ((\partial_t^+ \Delta_{tz})(\partial_z^i \Delta_{zt}) + (\partial_t^i \Delta_{tz})(\partial_z^+ \Delta_{zt})) \right] \tilde{G}_{t^-y^-}^{-0} \delta_{t_{\perp} y_{\perp}}^2 \delta_{z_{\perp} x_{\perp}}^2 \delta_{z^+ x^+}. \quad (72)$$

In the momentum space this expression has the following form:

$$-2i n_f g^2 \delta^{ab} \int \frac{d^4p}{(2\pi)^3} \frac{d^4q}{(2\pi)^4} d^4t \frac{-ie^{-i(p-q)(t-x)} (p_i - q_i) (p_- q^i + q_- p^i)}{(p^2 + i0)(q^2 + i0)} \theta(t^- - y^-) \delta_{t_{\perp} y_{\perp}}^2 \delta_{p_- q_-}, \quad (73)$$

which after the integration on  $t^+$  and  $t_{\perp}$  reduces to the following integral:

$$-2i n_f g^2 \delta^{ab} \int \frac{d^4p}{(2\pi)^2} \frac{d^4q}{(2\pi)^4} dt^- \theta(t^- - y^-) \frac{ie^{-i(p-q)(y-x)} (p_i - q_i) (p_- q^i + q_- p^i)}{(p^2 + i0)(q^2 + i0)} \delta_{p_- q_-} \delta_{p_+ q_+}, \quad (74)$$

and which one after the integration on  $p_-$  and  $p_+$  variables reads as

$$-2i n_f g^2 \delta^{ab} \int \frac{d^2p_{\perp}}{(2\pi)^2} \frac{d^4q}{(2\pi)^4} \frac{ie^{-i(p-q)_i(y-x)^i} q_- (p_{\perp}^2 - q_{\perp}^2)}{(2q_+ q_- - p_{\perp}^2 + i0)(2q_+ q_- - q_{\perp}^2 + i0)} \int dt^- \theta(t^- - y^-), \quad (75)$$

where the integrand of  $q_+$  decreases at infinity as  $1/q_+^2$  and its poles are located on one side of the real axis that gives

$$(K_{xy}^{ab})_{q1}^{+-} = 0. \quad (76)$$

That means that the reggeon propagator does not change from taking the quark loop contribution into account, that is well known result, see [14].

## 7 Conclusion

In this paper we calculated classical solutions of the equations of motion for gluon field to NNLO precision basing on the Lipatov's effective action, which can be considered as extension of QCD for the case of high energy interactions, see [15]. In the light cone gauge, for the three unknown components of the gluon field, four equations of motion exist. It demonstrated, that in the perturbative scheme of the solution of the equations, the existing of the solutions is equivalent to the transversality condition Eq. (11) applied to the current built on found solutions of the lower order. Namely, beginning from the bare effective gluon current in Lagrangian Eq. (1) and LO solutions of the equations of motion, the contributions to the higher order of the classical solutions can be constructed with the help of Eq. (12)-Eq. (13) if condition Eq. (11) is satisfied for the constructed current, see also Eq. (61)-Eq. (43) where the fermion loop contribution to the NNLO classical solution is accounted.

The importance of the found solutions is then that they contribute to the construction of the QCD based Regge Field Theory (RFT). Considering the reggeized gluons  $A_+$  and  $A_-$  as the main degrees of freedom at high energy QCD interactions, the found solutions provide the NNLO structure of the effective action of RFT

$$\Gamma = \sum_{n,m=0} \left( A_+^{a_1} \cdots A_+^{a_n} K_{b_1 \cdots b_m}^{a_1 \cdots a_n} A_-^{b_1} \cdots A_-^{b_m} \right), \quad (77)$$

see calculations in [4, 5]. The action of the interacting reggeized gluons in this form allows to calculate the effective vertices of interacting reggeized gluons and correlation functions of the theory, that is important task from the point of view of unitarization of the amplitudes at high energy interactions. In this set-up, therefore, the NNLO classical solutions provides some complex bare reggeon-reggeon interactions vertices which can be used further for the calculations of the reggeon loops contribution into the amplitudes and calculations of the corrections to the simpler bare reggeon-reggeon interactions vertices.

Finally we conclude, that the classical solution for the gluon field calculated is the first step toward the calculations of the unitarization corrections in the framework of QCD RFT and we hope that it can be useful also in the calculations performing in the framework of the CGC approach, [16, 17], where this solution can provide some unitarization corrections as well.



## Appedix A: expressions for $\partial_- U^{ab}$ and $\partial_i U^{ab}$ functions

For the arbitrary representation of gauge field  $v_+ = \imath T^a v_+^a$  with  $D_+ = \partial_+ - g v_+$ , we can consider the following representation of  $O$  and  $O^T$  operators<sup>3</sup> introduced in [4]:

$$O_x = \delta^{ab} + g \int d^4 y G_{xy}^{+aa_1} (v_+(y))_{a_1 b} = 1 + g G_{xy}^+ v_{+y} \quad (\text{A.1})$$

and correspondingly

$$O_x^T = 1 + g v_{+y} G_{yx}^+, \quad (\text{A.2})$$

which is redefinition of the operator expansions used in [2] in terms of Green's function instead integral operators. The Green's function in above equations we understand as Green's function of the  $D_+$  operator and express it in the perturbative sense as

$$G_{xy}^+ = G_{xy}^{+0} + g G_{xz}^{+0} v_{+z} G_{zy}^+ \quad (\text{A.3})$$

and

$$G_{yx}^+ = G_{yx}^{+0} + g G_{yz}^{+0} v_{+z} G_{zx}^{+0} \quad (\text{A.4})$$

with the bare propagators defined as (there is no integration on  $x$  variable)

$$\partial_{+x} G_{xy}^{+0} = \delta_{xy}, \quad G_{yx}^{+0} \overleftarrow{\partial}_{+x} = -\delta_{xy}. \quad (\text{A.5})$$

Then we have

$$\partial_{kx} G_{xy}^+ = \partial_{kx} G_{xy}^{+0} + g G_{xz}^{+0} \partial_{kz} v_{+z} G_{zy}^+ \quad (\text{A.6})$$

for  $k = -, i$ , the difference is only  $\partial_- v_{+0} = 0$ , hence  $\partial_- v_0 = O(g)$ . Then we obtain

$$\partial_{kx} O_x = g \partial_{kx} G_{xy}^+ v_{+y} = -g G_{xy}^{+0} (\partial_{ky} v_{+y} + g \partial_{kz} (v_{+y} G_{yz}^+ v_{+z})) = -g G_{xy}^{+0} ((\partial_{ky} v_{+y}) O_y + v_{+y} (\partial_{ky} O_y)). \quad (\text{A.7})$$

Thus in NNLO we have

$$\partial_{-x} O_x = g^2 G_{xy}^{+0} \left( \frac{1}{g} \partial_{-y} v_{+y} \right) O_y, \quad O_x^T \overleftarrow{\partial}_{-x} = -g^2 O_y^T \left( \frac{1}{g} \partial_{-y} v_{+y} \right) G_{yx}^{+0}, \quad (\text{A.8})$$

this form is more convenient because  $\partial_- v_{+0} = 0$ . Correspondingly for transverse derivative

$$\partial_{ix} O_x = g G_{xy}^{+0} (\partial_{iy} v_{+y} O_y), \quad O_x^T \overleftarrow{\partial}_{ix} = -g (O_y^T v_{+y} \overleftarrow{\partial}_{iy}) G_{yx}^{+0}. \quad (\text{A.9})$$

We are interested in the order of  $g$  for the operators  $\partial_{-x} U_x^{ab}$  and  $\partial_{ix} U_x^{ab}$ . For  $U_x^{ab} = \text{tr} (f^a O_x f^b O_x^T)$ , see [4] for details, we can write

$$\partial_{-x} U_x^{ab} = g^2 \text{tr} \left( f^a G_{xy}^{+0} \left( \frac{1}{g} \partial_{-y} v_{+y} \right) O_y f^b O_x^T - \left( \frac{1}{g} \partial_{-y} v_{+y} \right) G_{yx}^+ f^a O_x f^b O_y^T \right), \quad (\text{A.10})$$

$$\partial_{ix} U_x^{ab} = g \text{tr} \left( f^a G_{xy}^{+0} (\partial_{iy} v_{+y} O_y) f^b O_x^T - G_{yx}^{+0} f^a O_x f^b (O_y^T v_{+y} \overleftarrow{\partial}_{iy}) \right). \quad (\text{A.11})$$

---

<sup>3</sup>Due the light cone gauge we consider here only  $O(x^+)$  operators. The construction of the representation of the  $O(x^-)$  operators can be done similarly.

## Appendix B: NNLO terms contribution in Eq. (34)

In this appendix we present a calculation of the NNLO terms, denoted as  $C_a^i$ , in Eq. (34) equation of motion. Substituting LO and NLO classical solutions into the Eq. (34) we obtain:

$$\square v_a^i - \partial^i \left[ \partial^j v_{j\ a} + \partial_- v_{+\ a} \right] + (\partial F)_a^i = 0, \quad (\text{B.1})$$

where

$$(\partial F)_a^i = g f_{abc} \left( v_+^b \partial_- v^{i\ c} + \partial_- (v_-^b v^{i\ c}) + \partial_j (v^{j\ b} v^{i\ c}) + v^{j\ b} \partial_j v^{i\ c} - v_j^b \partial^i v^{j\ c} + g f^{cde} v^j{}^b v_d^j v_e^i \right). \quad (\text{B.2})$$

The expression in the square brackets can be written as:

$$\left[ \partial^j v_{j\ a} + \partial_- v_{+\ a} \right] = \left[ \partial^j v_{j0\ a} + \partial_- A_{+\ a} \right] + g \left[ \partial^j v_{j1\ a} + \partial_- v_{+1\ a} \right] + g^2 \left[ \partial^j v_{j2\ a} + \partial_- v_{+2\ a} \right]. \quad (\text{B.3})$$

Using Eq. (33) we obtain:

$$-\partial^i \left[ \partial^j v_{j\ a} + \partial_- v_{+\ a} \right] = -\partial^i \partial_j v_0^j{}_{\ a} + g \partial^i \left[ \frac{1}{g} (\partial^j U^{ab}) \rho_j^b - \partial_-^{-1} j_{a1}^+ \right] - g^2 \partial^i \left[ \partial_-^{-1} L_{a2}^+ - \partial_-^{-1} ((\partial^i \rho_i^b) (\frac{1}{g^2} \partial_- U_{ab})) \right].$$

The LO terms are canceled in the equations due

$$\square v_0^i{}_{\ a} - \partial^i \partial_j v_0^j{}_{\ a} = 2 \partial_+ \partial_- v_0^i{}_{\ a} + \partial_j \left( \rho^{i\ b} (\partial^j U_{ab}(A_+)) - \rho^{j\ b} (\partial^i U_{ab}(A_+)) \right) \quad (\text{B.4})$$

with

$$\partial^j v_{0a}^i - \partial^i v_{0a}^j = \rho^{i\ b} (\partial^j U_{ab}(A_+)) - \rho^{j\ b} (\partial^i U_{ab}(A_+)). \quad (\text{B.5})$$

After the substituting of the NLO classical solutions NLO from Eq. (16) and Eq. (17) into the equations, there are the NNLO terms remain which we denote as  $g^2 C_a^i$ :

$$\begin{aligned} & g \square v_1^i{}_{\ a} + g \partial^i \left[ \frac{1}{g} (\partial^j U^{ab}) \rho_j^b - \partial_-^{-1} j_{a1}^+ \right] \\ & + 2 \partial_+ \partial_- v_0^i{}_{\ a} + \partial_j \left( \rho^{i\ b} (\partial^j U_{ab}(A_+)) - \rho^{j\ b} (\partial^i U_{ab}(A_+)) \right) \\ & + g f_{abc} \left( 2 A_+^b \partial_- v_0^i{}_{\ c} + \partial_j (v_0^j{}^b v_0^i{}_{\ c}) + v_{j0}^b (\partial^j v_0^i{}_{\ c} - \partial^i v_0^j{}_{\ c}) \right) = g^2 C_a^i. \end{aligned} \quad (\text{B.6})$$

The first line of the expression Eq. (B.6) can be simplified by substituting Eq. (17)

$$g \square v_1^i{}_{\ a} + g \partial^i \left[ \frac{1}{g} (\partial^j U^{ab}) \rho_j^b - \partial_-^{-1} j_{a1}^+ \right] = -g \partial_j F_a^{ji}. \quad (\text{B.7})$$

With the help of Eq. (15) and Eq. (19) expressions we can write:

$$\begin{aligned} \partial_+ U^{ab}(v_+) &= \text{tr} \left[ (f^a f^c - f^c f^a) \left( P e^{g \int_{-\infty}^{x^+} dx'^+ v_{+e}(x'^+, x^-, x_\perp) f^e} \right) f^b \left( P e^{g \int_{x^+}^{+\infty} dx'^+ v_{+d}(x'^+, x^-, x_\perp) f^d} \right) \right] g v_{+c} \\ &= -g f^{acf} v_{+c} U^{fb}(v_+) \end{aligned} \quad (\text{B.8})$$

and

$$2 \partial_+ \partial_- v_0^i{}_{\ a} = -2 g f_{abc} \partial_- (v_+^b v_0^i{}_{\ c}). \quad (\text{B.9})$$

Then, the Eq. (B.6) acquires the following form:

$$\begin{aligned} & -g \partial_j F_a^{ji} - 2 g^2 f_{abc} \partial_- (v_{+1}^b v_0^i{}_{\ c}) + \partial_j \left( \rho^{i\ b} (\partial^j U_{ab}(A_+)) - \rho^{j\ b} (\partial^i U_{ab}(A_+)) \right) \\ & + g f_{abc} \left( \partial_j (v_0^j{}^b v_0^i{}_{\ c}) + v_{j0}^b (\partial^j v_0^i{}_{\ c} - \partial^i v_0^j{}_{\ c}) \right) = g^2 C_a^i, \end{aligned} \quad (\text{B.10})$$

where

$$\partial_j F_a^{ji} = \partial_j \left[ \left( \rho^{i\ b} \left( \frac{1}{g} \partial^j U_{ab}(A_+) \right) - \rho^{j\ b} \left( \frac{1}{g} \partial^i U_{ab}(A_+) \right) \right) + f_{abc} v_0^j{}^b v_0^i{}_{\ c} \right], \quad (\text{B.11})$$

and correspondingly

$$C_a^i = f_{abc} \left[ v_{j0}^b \left( \rho^{i\ b} \left( \frac{1}{g} \partial^j U_{ab}(A_+) \right) - \rho^{j\ b} \left( \frac{1}{g} \partial^i U_{ab}(A_+) \right) \right) - 2 \partial_- (v_{+1}^b v_0^i{}_{\ c}) \right]. \quad (\text{B.12})$$

## Appendix C: NNLO terms contribution in Eq. (45)

In this appendix we present the fourth equation (47) of motion to the form of Eq. (10). We write Eq. (47) by analogy with the calculations from the appendix B in the form

$$\begin{aligned} & \square v_{+2a} - \partial_+ \left[ \partial_i v_{2a}^i + \partial_- v_{+2a} \right] + \bar{C}_a^i \\ & + f_{abc} \left( \partial_i (v_0^{ib} v_{+1}^c) + v_{i0}^b (\partial^i v_{+1}^c - \partial_+ v_1^{i\ c} + f^{cde} v_{0d}^i A_{+e}) + \partial_i (v_1^{ib} A_+^c) + v_{i1}^b \partial^i A_+^c + A_+^b \partial_- v_{+1}^c \right) = 0, \end{aligned} \quad (C.1)$$

where

$$g\bar{C}_a^i = \square v_{+1a} - \partial_+ \left[ \partial_i v_{1a}^i + \partial_- v_{+1a} \right] - \frac{1}{g} \partial_+ \left[ \partial_i v_{0a}^i \right] + f_{abc} \left( \partial_i (v_0^{ib} A_+^c) + v_{i0}^b (\partial^i A_+^c - \partial_+ v_0^{i\ c}) \right). \quad (C.2)$$

With the help of Eq. (17) and Eq. (16) we rewrite the first three terms in Eq. (C.2) and obtain:

$$g\bar{C}_a^i = -\frac{2}{g} \left( (\partial_+ \partial^i U_{ab}) \rho_i^b \right) - \square^{-1} \partial_i \partial^i \partial_+ \partial_-^{-1} j_{a1}^+ - \frac{1}{g} (\partial_i \rho^{ib}) \left[ \partial_+ U_{ab} \right] + f_{abc} \left( \partial_i (v_0^{ib} A_+^c) + v_{i0}^b (\partial^i A_+^c - \partial_+ v_0^{i\ c}) \right). \quad (C.3)$$

Applying Eq. (B.8) expression, we see that the following terms of  $g^2$  order are not canceled in Eq. (C.3):

$$\bar{C}_a^i = f_{abc} \left( A_+^b \left( \frac{1}{g} \partial^i U^{cd} \right) \rho_{id} + \partial^i (v_{+1}^b U^{cd}) \rho_{id} + \partial^i (v_{+1}^b v_{i0}^c) - v_{i0}^b \left( \frac{1}{g} \partial_+ v_0^{i\ c} \right) \right) - \square^{-1} \partial_i \partial^i \partial_-^{-1} \left( \frac{1}{g} \partial_+ j_{a1}^+ \right). \quad (C.4)$$

Using Eq. (B.8) and the antisymmetry property  $f^{abc}$ , we note that:

$$f_{abc} \left( \partial_i (v_0^{ib} v_{+1}^c) + \partial^i (v_{+1}^b v_{i0}^c) \right) = 0, \quad (C.5)$$

$$v_{i0}^b f^{cde} v_{0d}^i A_{+e} - v_{i0}^b \left( \frac{1}{g} \partial_+ v_0^{i\ c} \right) = O(g), \quad f_{abc} \left( v_{i0}^b \partial^i v_{+1}^c + \partial^i (v_{+1}^b U^{cd}) \rho_{id} \right) = O(g) \quad (C.6)$$

and

$$\begin{aligned} & f_{abc} \left( \partial_i (v_1^{ib} A_+^c) + v_{i1}^b (\partial^i A_+^c) + A_+^b \partial_- v_{+1}^c + A_+^b \left( \frac{1}{g} \partial^i U^{cd} \right) \rho_{id} \right) \\ & = f_{abc} \left( -2\partial_i (A_+^b v_1^{ic}) + A_+^b \left[ \partial_- v_{+1}^c + \partial^i v_{i1}^c \right] + A_+^b \left( \frac{1}{g} \partial^i U^{cd} \right) \rho_{id} \right) \\ & = f_{abc} \left( -2\partial_i (A_+^b v_1^{ic}) + A_+^b \square^{-1} \partial_i \partial^i \partial_-^{-1} j_{a1}^{+c} \right). \end{aligned} \quad (C.7)$$

Thus, with the use of Eq. (C.4)-Eq. (C.7) we rewrite Eq. (47) in the following form

$$\begin{aligned} & \square v_{+2a} - \partial_+ \left[ \partial_i v_{2a}^i + \partial_- v_{+2a} \right] \\ & = -f_{abc} \left( -v_{i0}^b \partial_+ v_1^{i\ c} - 2\partial_i (A_+^b v_1^{ic}) + A_+^b \square^{-1} \partial_i \partial^i \partial_-^{-1} j_{a1}^{+c} \right) + \square^{-1} \partial_i \partial^i \partial_-^{-1} \left( \frac{1}{g} \partial_+ j_{a1}^+ \right), \end{aligned} \quad (C.8)$$

where  $j_{a1}^+$  is defined by equation Eq. (30). The r.h.s. of the Eq. (C.8) consists with the classical solutions of  $k = 0$  and  $k = 1$  orders only.

## References

- [1] V. N. Gribov, Sov. Phys. JETP 26 (1968) 414.
- [2] L. N. Lipatov, Nucl. Phys. B **452**, 369 (1995); Phys. Rept. **286**, 131 (1997); Subnucl. Ser. **49**, 131 (2013); Int. J. Mod. Phys. Conf. Ser. **39**, 1560082 (2015); Int. J. Mod. Phys. A **31**, no. 28/29, 1645011 (2016); EPJ Web Conf. **125**, 01010 (2016).
- [3] L. N. Lipatov, Sov. J. Nucl. Phys. **23**, 338 (1976) [Yad. Fiz. **23** (1976) 642]; E. A. Kuraev, L. N. Lipatov and V. S. Fadin, Sov. Phys. JETP **45**, 199 (1977) [Zh. Eksp. Teor. Fiz. **72**, 377 (1977)]; I. I. Balitsky and L. N. Lipatov, Sov. J. Nucl. Phys. **28**, 822 (1978) [Yad. Fiz. **28**, 1597 (1978)].
- [4] S. Bondarenko, L. Lipatov and A. Prygarin, Eur. Phys. J. C **77** (2017) no.8, 527.
- [5] S. Bondarenko, L. Lipatov, S. Pozdnyakov, A. Prygarin, Eur. Phys. J. C **77** (2017) no.9, 630.
- [6] J. Bartels, Nucl. Phys. B 175 (1980) 365; J. Kwiecinski, M. Praszalowicz, Phys. Lett. B 94 (1980) 413.
- [7] E. N. Antonov, L. N. Lipatov, E. A. Kuraev and I. O. Cherednikov, Nucl. Phys. B **721** (2005) 111.
- [8] J. Bartels, V. S. Fadin, L. N. Lipatov and G. P. Vacca, Nucl. Phys. B **867**, 827 (2013).
- [9] A. van Hameren, P. Kotko and K. Kutak, JHEP **1301** (2013) 078; K. Kutak, A. Hameren and M. Serino, JHEP **1702** (2017) 009.
- [10] D. Amati, L. Caneschi and R. Jengo, Nucl. Phys. B **101** (1975) 397; R. Jengo, Nucl. Phys. B **108** (1976) 447; M. Ciafaloni, Nucl. Phys. B **146** (1978) 427.
- [11] E. Levin and A. Prygarin, Eur. Phys. J. C **53**, 385 (2008); S. Bondarenko, Eur. Phys. J. C **71**, 1587 (2011); S. Bondarenko, L. Horwitz, J. Levitan and A. Yahalom, Nucl. Phys. A **912**, 49 (2013).
- [12] L. N. Lipatov, Nucl. Phys. Proc. Suppl. **99A**, 175 (2001); M. A. Braun and M. I. Vyazovsky, Eur. Phys. J. C **51**, 103 (2007); M. A. Braun, M. Y. Salykin and M. I. Vyazovsky, Eur. Phys. J. C **65**, 385 (2010); M. A. Braun, L. N. Lipatov, M. Y. Salykin and M. I. Vyazovsky, Eur. Phys. J. C **71**, 1639 (2011); M. A. Braun, M. Y. Salykin and M. I. Vyazovsky, Eur. Phys. J. C **72**, 1864 (2012); M. Hentschinski and A. Sabio Vera, Phys. Rev. D **85**, 056006 (2012); M. A. Braun, M. Y. Salykin, S. S. Pozdnyakov and M. I. Vyazovsky, Eur. Phys. J. C **72**, 2223 (2012); J. Bartels, L. N. Lipatov and G. P. Vacca, Phys. Rev. D **86**, 105045 (2012); M. A. Braun, S. S. Pozdnyakov, M. Y. Salykin and M. I. Vyazovsky, Eur. Phys. J. C **73**, no. 9, 2572 (2013); G. Chachamis, M. Hentschinski, J. D. Madrigal Martnez and A. Sabio Vera, Phys. Part. Nucl. **45**, no. 4, 788 (2014); M. A. Braun, S. S. Pozdnyakov, M. Y. Salykin and M. I. Vyazovsky, Eur. Phys. J. C **74** (2014) no.8, 2989; M. A. Braun, S. S. Pozdnyakov, M. Y. Salykin and M. I. Vyazovsky, Eur. Phys. J. C **75** (2015) no.5, 222; M. A. Braun, Eur. Phys. J. C **75** (2015) no.7, 298; M. A. Braun and M. I. Vyazovsky, Phys. Rev. D **93** (2016) no.6, 065026; M. A. Braun, Eur. Phys. J. C **77** (2017) no.5, 279; M. A. Braun and M. Y. Salykin, Eur. Phys. J. C **77** (2017) no.7, 498; M. Hentschinski, arXiv:1802.06755 [hep-ph].
- [13] M. Nefedov and V. Saleev, Mod. Phys. Lett. A **32** (2017) no.40, 1750207.
- [14] V. S. Fadin and R. Fiore, Phys. Lett. B **294** (1992) 286; V. S. Fadin, R. Fiore and A. Quartarolo, Phys. Rev. D **50** (1994) 2265; V. S. Fadin, R. Fiore and A. Quartarolo, Phys. Rev. D **50** (1994) 5893; V. S. Fadin, R. Fiore, A. Flachi and M. I. Kotsky, Phys. Lett. B **422** (1998) 287.
- [15] S. Bondarenko and M. A. Zubkov, arXiv:1801.08066 [hep-ph].
- [16] L. McLerran and R. Venugopalan, Phys. Rev. D **49** (1994), 2233; **D49** (1994), 3352.
- [17] J. Jalilian-Marian, A. Kovner, L. McLerran and H. Weigert, Phys. Rev. D **55**, 5414 (1997); J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, Nucl. Phys. B **504**, 415 (1997); J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, Phys. Rev. D **59**, 014014 (1998); J. Jalilian-Marian, A. Kovner and H. Weigert, Phys. Rev. D **59**, 014015 (1998); E. Iancu, A. Leonidov and L. D. McLerran, Nucl. Phys. A **692**, 583 (2001); E. Iancu, A. Leonidov and L. D. McLerran, Phys. Lett. B **510**, 133 (2001); E. Ferreira, E. Iancu, A. Leonidov and L. McLerran, Nucl. Phys. A **703**, 489 (2002).