

Novel results on Hermite–Hadamard kind inequalities for η -convex functions by means of (k, r) -fractional integral operators*

Eze R. Nwaeze¹
enwaeze@tuskegee.edu

Delfim F. M. Torres^{2†}
delfim@ua.pt

¹Department of Mathematics, Tuskegee University,
Tuskegee, AL 36088, USA

²CIDMA, Department of Mathematics, University of Aveiro,
3810-193 Aveiro, Portugal

Abstract

We establish new integral inequalities of Hermite–Hadamard type for the recent class of η -convex functions. This is done via generalized (k, r) -Riemann–Liouville fractional integral operators. Our results generalize some known theorems in the literature. By choosing different values for the parameters k and r , one obtains interesting new results.

Keywords: Hermite–Hadamard inequalities, η -convexity, Riemann–Liouville integrals.

2010 Mathematics Subject Classification: 26A51, 26D15.

1 Introduction

Throughout this work, $I \subset \mathbb{R}$ shall denote an interval and I° the interior of I . We say that a function $g : I \rightarrow \mathbb{R}$ is convex if, for every $x, y \in I$ and $\beta \in [0, 1]$, one has

$$g(\beta x + (1 - \beta)y) \leq \beta g(x) + (1 - \beta)g(y). \quad (1)$$

Let $a, b \in I$. For a function g satisfying (1), the following inequalities hold:

$$g\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b g(x) dx \leq \frac{g(a) + g(b)}{2}. \quad (2)$$

Result (2) was proved by Hadamard in 1893 [6] and is celebrated in the literature as the Hermite–Hadamard integral inequality for convex functions [2]. Along the years, it has been extended to different classes of convex functions: see, e.g., [3, 8, 15] and references therein.

In 2016, the so called φ -convexity was introduced [5], subsequently denoted as η -convexity [4, 12]. Let us recall its definition here.

Definition 1 (See [5]). *A function $g : I \rightarrow \mathbb{R}$ is called convex with respect to η (for short, η -convex), if*

$$g(\beta x + (1 - \beta)y) \leq g(y) + \beta \eta(g(x), g(y))$$

for all $x, y \in I$ and $\beta \in [0, 1]$.

*This is a preprint of a paper whose final and definite form is a Springer chapter in the Book *Advances in Mathematical Inequalities and Applications*, published under the Birkhauser series *Trends in Mathematics*, ISSN: 2297-0215 [<http://www.springer.com/series/4961>].

[†]Corresponding author.

By taking $\eta(x, y) = x - y$, Definition 1 reduces to the classical notion (1) of convexity. It was further shown in [5] that for every convex function g there exists some η , different from $\eta(x, y) = x - y$, for which the function g is η -convex. The converse is, however, not necessarily true, that is, there are η -convex functions that are not convex.

Example 2. Consider function $g : \mathbb{R} \rightarrow \mathbb{R}$ defined piecewisely by

$$g(x) = \begin{cases} -x, & x \geq 0, \\ x, & x < 0, \end{cases}$$

and let $\eta : [-\infty, 0] \times [-\infty, 0] \rightarrow \mathbb{R}$ be given by $\eta(x, y) = -x - y$. Function g is clearly not convex but it is easy to see that it is η -convex. Indeed, in [12, Remark 4] it is noted that an η -convex function $g : [a, b] \rightarrow \mathbb{R}$ is integrable if η is bounded from above on $g([a, b]) \times g([a, b])$.

For the class of η -convex functions, the following theorem was obtained as an analogue of (2).

Theorem 3 (See [5]). Suppose that $g : I \rightarrow \mathbb{R}$ is an η -convex function such that η is bounded from above on $g(I) \times g(I)$. Then, for any $a, b \in I$ with $a < b$,

$$2g\left(\frac{a+b}{2}\right) - M_\eta \leq \frac{1}{b-a} \int_a^b g(x) dx \leq f(b) + \frac{\eta(g(a), g(b))}{2},$$

where M_η is an upper bound of η on $g([a, b]) \times g([a, b])$.

Recently, Rostamian Delavar and De La Sen obtained, among other results, the following theorem associated to η -convex functions [12].

Theorem 4 (See [12]). Suppose $g : [a, b] \rightarrow \mathbb{R}$ is a differentiable function and $|g'|$ is an η -convex function with η bounded from above on $[a, b]$. Then,

$$\left| \frac{g(a) + g(b)}{2} - \frac{1}{b-a} \int_a^b g(x) dx \right| \leq \frac{1}{8}(b-a)K,$$

where $K = \min \left\{ |g'(b)| + \frac{|\eta(g'(a), g'(b))|}{2}, |g'(a)| + \frac{|\eta(g'(b), g'(a))|}{2} \right\}$.

Still in the same spirit, Khan et al. established in 2017 the following result for η -convex functions via Riemann–Liouville fractional integral operators [9].

Theorem 5 (See [9]). Let $g : [a, b] \rightarrow \mathbb{R}$ be a differentiable function on (a, b) with $a < b$. If $|g'|$ is an η -convex function on $[a, b]$, then for $\alpha > 0$ the inequality

$$\left| \frac{g(a) + g(b)}{2} - \frac{\Gamma(\alpha + 1)}{2(b-a)^\alpha} \left[J_{a+}^\alpha g(b) + J_{b-}^\alpha g(a) \right] \right| \leq \frac{b-a}{2(\alpha+1)} \left(1 - \frac{1}{2^\alpha} \right) (2|g'(b)| + \eta(|g'(a)|, |g'(b)|))$$

holds, where

$$J_{a+}^\alpha g(x) = \frac{1}{\Gamma_1(\alpha)} \int_a^x (x-t)^{\alpha-1} g(t) dt$$

is the left Riemann–Liouville fractional integral and

$$J_{b-}^\alpha g(x) = \frac{1}{\Gamma_1(\alpha)} \int_x^b (t-x)^{\alpha-1} g(t) dt$$

is the right Riemann–Liouville fractional integral.

Fractional calculus is an area under strong development [11], and in [13] Sarikaya et al. proposed the following broader definition of the Riemann–Liouville fractional integral operators.

Definition 6 (See [13]). *The (k, r) -Riemann–Liouville fractional integral operators ${}_k^r \mathcal{J}_{a^+}^\alpha$ and ${}_k^r \mathcal{J}_{b^-}^\alpha$ of order $\alpha > 0$, for a real valued continuous function $g(x)$, are defined as*

$${}_k^r \mathcal{J}_{a^+}^\alpha g(x) = \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_a^x (x^{r+1} - t^{r+1})^{\frac{\alpha}{k}-1} t^r g(t) dt, \quad x > a, \quad (3)$$

and

$${}_k^r \mathcal{J}_{b^-}^\alpha g(x) = \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_x^b (t^{r+1} - x^{r+1})^{\frac{\alpha}{k}-1} t^r g(t) dt, \quad x < b, \quad (4)$$

where $k > 0$, $r \in \mathbb{R} \setminus \{-1\}$, and Γ_k is the k -gamma function given by

$$\Gamma_k(x) := \int_0^\infty t^{x-1} e^{-\frac{t^k}{k}} dt, \quad \operatorname{Re}(x) > 0,$$

with the properties $\Gamma_k(x+k) = x\Gamma_k(x)$ and $\Gamma_k(k) = 1$.

For some results related to the operators (3) and (4), we refer the interested readers to [7, 10, 14, 16]. Using these operators, Agarwal et al. established the following Hermite–Hadamard type result for convex functions [1].

Theorem 7 (See [1]). *Let $\alpha, k > 0$ and $r \in \mathbb{R} \setminus \{-1\}$. If g is a convex function on $[a, b]$, then*

$$g\left(\frac{a+b}{2}\right) \leq \frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha+k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k^r \mathcal{J}_{a^+}^\alpha G(b) + {}_k^r \mathcal{J}_{b^-}^\alpha G(a)] \leq \frac{g(a) + g(b)}{2},$$

where function G is defined by (5) below.

Inspired by the above works, it is our purpose to obtain here more general integral inequalities associated to η -convex functions via the (k, r) -Riemann–Liouville fractional operators. Theorems 8 and 12 generalize Theorems 7 and 5, respectively (see Remarks 9 and 13). In addition, two more fractional Hermite–Hadamard type inequalities are also established (see Theorems 14 and 15).

2 Main results

We establish four new results. For this, we start by making the following observations. Let g be a function defined on I with $[a, b] \subset I^\circ$ and define functions $G, \tilde{g} : [a, b] \rightarrow \mathbb{R}$ by

$$\tilde{g}(x) := g(a+b-x) \quad \text{and} \quad G(x) := g(x) + \tilde{g}(x). \quad (5)$$

For the fractional operators to be well defined, we shall assume $g \in L_\infty[a, b]$. By making use of the substitutions $w = \frac{t-a}{x-a}$ and $w = \frac{b-t}{b-x}$ in (3) and (4), respectively, one gets that

$${}_k^r \mathcal{J}_{a^+}^\alpha g(x) = (x-a) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_0^1 \frac{(wx + (1-w)a)^r g(wx + (1-w)a)}{[x^{r+1} - (wx + (1-w)a)^{r+1}]^{1-\frac{\alpha}{k}}} dw \quad (6)$$

and

$${}_k^r \mathcal{J}_{b^-}^\alpha g(x) = (b-x) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_0^1 \frac{(wx + (1-w)b)^r g(wx + (1-w)b)}{[(wx + (1-w)b)^{r+1} - x^{r+1}]^{1-\frac{\alpha}{k}}} dw. \quad (7)$$

Noting that $\tilde{g}((1-w)a + wb) = g(wa + (1-w)b)$, we also obtain

$${}_k^r \mathcal{J}_{a^+}^\alpha \tilde{g}(x) = (x-a) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_0^1 \frac{(wx + (1-w)a)^r g((1-w)x + wa)}{[x^{r+1} - (wx + (1-w)a)^{r+1}]^{1-\frac{\alpha}{k}}} dw \quad (8)$$

and

$${}_k^r \mathcal{J}_{b^-}^\alpha \tilde{g}(x) = (b-x) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_0^1 \frac{(wx + (1-w)b)^r g((1-w)x + wb)}{[(wx + (1-w)b)^{r+1} - x^{r+1}]^{1-\frac{\alpha}{k}}} dw. \quad (9)$$

We are now ready to formulate and prove our first result.

Theorem 8. Let $\alpha, k > 0$, $r \in \mathbb{R} \setminus \{-1\}$, and $g : I \rightarrow \mathbb{R}$ be a positive function on $[a, b] \subset I^\circ$ with $a < b$. If, in addition, g is η -convex on $[a, b]$ with η bounded on $g([a, b]) \times g([a, b])$, then the (k, r) -fractional integral inequality

$$\frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha+k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k^r \mathcal{J}_{a^+}^\alpha G(b) + {}_k^r \mathcal{J}_{b^-}^\alpha G(a)] \leq g(b) + \frac{\eta(g(a), g(b))}{2}$$

holds.

Proof. Function g is η -convex on $[a, b]$, which implies, by definition, the following inequalities for $t \in [0, 1]$:

$$g(ta + (1-t)b) \leq g(b) + t\eta(g(a), g(b)) \quad (10)$$

and

$$g((1-t)a + tb) \leq g(b) + (1-t)\eta(g(a), g(b)). \quad (11)$$

Adding inequalities (10) and (11), we get

$$g(ta + (1-t)b) + g((1-t)a + tb) \leq 2g(b) + \eta(g(a), g(b)). \quad (12)$$

Multiplying both sides of (12) by

$$(b-a) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \frac{(tb + (1-t)a)^r}{[b^{r+1} - (tb + (1-t)a)^{r+1}]^{1-\frac{\alpha}{k}}},$$

and integrating over $[0, 1]$ with respect to t , we get

$$\begin{aligned} & (b-a) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_0^1 \frac{(tb + (1-t)a)^r g((1-t)b + ta)}{[b^{r+1} - (tb + (1-t)a)^{r+1}]^{1-\frac{\alpha}{k}}} dt \\ & + (b-a) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_0^1 \frac{(tb + (1-t)a)^r g(tb + (1-t)a)}{[b^{r+1} - (tb + (1-t)a)^{r+1}]^{1-\frac{\alpha}{k}}} dt \\ & \leq [2g(b) + \eta(g(a), g(b))] (b-a) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \int_0^1 \frac{(tb + (1-t)a)^r}{[b^{r+1} - (tb + (1-t)a)^{r+1}]^{1-\frac{\alpha}{k}}} dt. \end{aligned}$$

Now, using (6) and (8) in the above inequality, we get

$${}_k^r \mathcal{J}_{a^+}^\alpha \tilde{g}(b) + {}_k^r \mathcal{J}_{a^+}^\alpha g(b) \leq \frac{(s+1)^{1-\frac{\alpha}{k}} (b^{s+1} - a^{s+1})^{\frac{\alpha}{k}}}{(s+1)\alpha\Gamma_k(\alpha)} [2g(b) + \eta(g(a), g(b))],$$

that is,

$${}_k^r \mathcal{J}_{a^+}^\alpha G(b) \leq \frac{(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}}{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha+k)} [2g(b) + \eta(g(a), g(b))]. \quad (13)$$

Similarly, multiplying again both sides of (12) by

$$(b-a) \frac{(r+1)^{1-\frac{\alpha}{k}}}{k\Gamma_k(\alpha)} \frac{(tb + (1-t)a)^r}{[(tb + (1-t)a)^{r+1} - a^{r+1}]^{1-\frac{\alpha}{k}}}$$

and integrating with respect to t over $[0, 1]$, we obtain that

$${}_k^r \mathcal{J}_{b^-}^\alpha G(a) \leq \frac{(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}}{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha+k)} [2g(b) + \eta(g(a), g(b))]. \quad (14)$$

Hence, the intended inequality follows by adding (13) and (14). \square

Remark 9. By taking $\eta(x, y) = x - y$ in our Theorem 8, we recover the right-hand side of the inequalities in Theorem 7.

For the rest of our results, we will need the following two lemmas.

Lemma 10 (See [1]). *Let $\alpha, k > 0$ and $r \in \mathbb{R} \setminus \{-1\}$. If $g : I \rightarrow \mathbb{R}$ is differentiable on I° and $a, b \in I^\circ$ such that $g' \in L[a, b]$ with $a < b$, then the following identity holds:*

$$\begin{aligned} \frac{g(a) + g(b)}{2} - \frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha + k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k^r \mathcal{J}_{a^+}^\alpha G(b) + {}_k^s \mathcal{J}_{b^-}^\alpha G(a)] \\ = \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \int_0^1 \Theta_{\alpha,r}(t) g'(ta + (1-t)b) dt, \end{aligned}$$

where $\Theta_{\alpha,r} : [0, 1] \rightarrow \mathbb{R}$ is defined by

$$\begin{aligned} \Theta_{\alpha,r}(t) := [(ta + (1-t)b)^{r+1} - a^{r+1}]^{\frac{\alpha}{k}} - [(tb + (1-t)a)^{r+1} - a^{r+1}]^{\frac{\alpha}{k}} \\ + [b^{r+1} - (tb + (1-t)a)^{r+1}]^{\frac{\alpha}{k}} - [b^{r+1} - (ta + (1-t)b)^{r+1}]^{\frac{\alpha}{k}}. \end{aligned}$$

Lemma 11. *Under the conditions of Lemma 10, we have that*

$$\int_0^1 |\Theta_{\alpha,r}(t)| dt = \frac{1}{b-a} (\mathfrak{R}_1 + \mathfrak{R}_2 + \mathfrak{R}_3 + \mathfrak{R}_4),$$

where

$$\begin{aligned} \mathfrak{R}_1 &= \int_{\frac{a+b}{2}}^b (w^{r+1} - a^{r+1})^{\frac{\alpha}{k}} dw - \int_a^{\frac{a+b}{2}} (w^{r+1} - a^{r+1})^{\frac{\alpha}{k}} dw, \\ \mathfrak{R}_2 &= \int_{\frac{a+b}{2}}^b [b^{r+1} - (b+a-w)^{r+1}]^{\frac{\alpha}{k}} dw - \int_a^{\frac{a+b}{2}} [b^{r+1} - (b+a-w)^{r+1}]^{\frac{\alpha}{k}} dw, \\ \mathfrak{R}_3 &= \int_a^{\frac{a+b}{2}} (b^{r+1} - w^{r+1})^{\frac{\alpha}{k}} dw - \int_{\frac{a+b}{2}}^b (b^{r+1} - w^{r+1})^{\frac{\alpha}{k}} dw, \end{aligned}$$

and

$$\mathfrak{R}_4 = \int_a^{\frac{a+b}{2}} [(b+a-w)^{r+1} - a^{r+1}]^{\frac{\alpha}{k}} dw - \int_{\frac{a+b}{2}}^b [(b+a-w)^{r+1} - a^{r+1}]^{\frac{\alpha}{k}} dw.$$

Proof. Using the substitution $w = ta + (1-t)b$, we get

$$\int_0^1 |\Theta_{\alpha,r}(t)| dt = \frac{1}{b-a} \int_a^b |\wp(w)| dw, \quad (15)$$

where

$$\begin{aligned} \wp(w) &= (w^{r+1} - a^{r+1})^{\frac{\alpha}{k}} - [(b+a-w)^{r+1} - a^{r+1}]^{\frac{\alpha}{k}} \\ &\quad + [b^{r+1} - (b+a-w)^{r+1}]^{\frac{\alpha}{k}} - (b^{r+1} - w^{r+1})^{\frac{\alpha}{k}}. \end{aligned}$$

The required result follows from (15) and by observing that \wp is a non-decreasing function on $[a, b]$, $\wp(a) = -2(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}} < 0$, $\wp(\frac{a+b}{2}) = 0$, and thus

$$\begin{cases} \wp(w) \leq 0 & \text{if } a \leq w \leq \frac{a+b}{2}, \\ \wp(w) > 0 & \text{if } \frac{a+b}{2} < w \leq b. \end{cases}$$

This concludes the proof. \square

Theorem 12. Let $\alpha, k > 0$, $r \in \mathbb{R} \setminus \{-1\}$, $g : I \rightarrow \mathbb{R}$ be a differentiable function on I° and $a, b \in I^\circ$ with $a < b$. Suppose $|g'|$ is η -convex on $[a, b]$ with η bounded on $|g'|([a, b]) \times |g'|([a, b])$. Then the following (k, r) -fractional integral inequality holds:

$$\left| \frac{g(a) + g(b)}{2} - \frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha + k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k\mathcal{J}_{a^+}^\alpha G(b) + {}_k\mathcal{J}_{b^-}^\alpha G(a)] \right| \leq \frac{1}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left[\Re |g'(b)| + \frac{\Xi}{b-a} \eta(|g'(a)|, |g'(b)|) \right],$$

where $\Re = \Re_1 + \Re_2 + \Re_3 + \Re_4$ (see Lemma 11) and $\Xi = \xi_1 + \xi_2 + \xi_3 + \xi_4$ with

$$\begin{aligned} \xi_1 &= \int_a^{\frac{a+b}{2}} (b-w)(b^{r+1} - w^{r+1})^{\frac{\alpha}{k}} dw - \int_{\frac{a+b}{2}}^b (b-w)(b^{r+1} - w^{r+1})^{\frac{\alpha}{k}} dw, \\ \xi_2 &= \int_{\frac{a+b}{2}}^b (b-w)(w^{r+1} - a^{r+1})^{\frac{\alpha}{k}} dw - \int_a^{\frac{a+b}{2}} (b-w)(w^{r+1} - a^{r+1})^{\frac{\alpha}{k}} dw, \\ \xi_3 &= \int_a^{\frac{a+b}{2}} (b-w)((b+a-w)^{r+1} - a^{r+1})^{\frac{\alpha}{k}} dw - \int_{\frac{a+b}{2}}^b (b-w)((b+a-w)^{r+1} - a^{r+1})^{\frac{\alpha}{k}} dw, \\ \xi_4 &= \int_{\frac{a+b}{2}}^b (b-w)(b^{r+1} - (b+a-w)^{r+1})^{\frac{\alpha}{k}} dw - \int_a^{\frac{a+b}{2}} (b-w)(b^{r+1} - (b+a-w)^{r+1})^{\frac{\alpha}{k}} dw. \end{aligned}$$

Proof. Since $|f'|$ is η -convex, it follows, by definition, that

$$|g'(ta + (1-t)b)| \leq |g'(b)| + t\eta(|g'(a)|, |g'(b)|) \quad (16)$$

for $t \in [0, 1]$. From [1, p. 9], we have

$$\int_0^1 t |\Theta_{\alpha, r}(t)| dt = \frac{\xi_1 + \xi_2 + \xi_3 + \xi_4}{(b-a)^2}. \quad (17)$$

Using Lemmas 10 and 11, inequality (16), identity (17), and properties of the modulus, we obtain

$$\begin{aligned} & \left| \frac{g(a) + g(b)}{2} - \frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha + k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k\mathcal{J}_{a^+}^\alpha G(b) + {}_k\mathcal{J}_{b^-}^\alpha G(a)] \right| \\ & \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \int_0^1 |\Theta_{\alpha, r}(t)| |g'(ta + (1-t)b)| dt \\ & \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \int_0^1 |\Theta_{\alpha, r}(t)| [|g'(b)| + t\eta(|g'(a)|, |g'(b)|)] dt \\ & = \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left(|g'(b)| \int_0^1 |\Theta_{\alpha, r}(t)| dt + \eta(|g'(a)|, |g'(b)|) \int_0^1 t |\Theta_{\alpha, r}(t)| dt \right) \\ & = \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left[|g'(b)| \frac{1}{b-a} (\Re_1 + \Re_2 + \Re_3 + \Re_4) + \eta(|g'(a)|, |g'(b)|) \frac{\xi_1 + \xi_2 + \xi_3 + \xi_4}{(b-a)^2} \right]. \end{aligned}$$

The desired result follows. \square

Remark 13. By taking $r = 0$ and $k = 1$ in Theorem 12, we recover Theorem 5. In this case,

$$\Re = \frac{4}{\alpha+1} (b-a)^{\alpha+1} \left(1 - \frac{1}{2^\alpha} \right)$$

and

$$\Xi = \frac{2}{\alpha+1} (b-a)^{\alpha+2} \left(1 - \frac{1}{2^\alpha} \right).$$

Theorem 14. Let g be differentiable on I° with $a, b \in I^\circ$. If $|g'|^q$ is η -convex on $[a, b]$ and $q > 1$ with η bounded on $|g'|^q([a, b]) \times |g'|^q([a, b])$, then the (k, r) -fractional integral inequality

$$\left| \frac{g(a) + g(b)}{2} - \frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha + k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k^r \mathcal{J}_{a^+}^\alpha G(b) + {}_k^r \mathcal{J}_{b^-}^\alpha G(a)] \right| \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left(|g'(b)|^q + \frac{\eta(|g'(a)|^q, |g'(b)|^q)}{2} \right)^{\frac{1}{q}} \|\Theta_{\alpha, r}\|_p$$

holds, where $\frac{1}{p} + \frac{1}{q} = 1$ and $\|\Theta_{\alpha, r}\|_p = \left(\int_0^1 |\Theta_{\alpha, r}(t)|^p dt \right)^{\frac{1}{p}}$.

Proof. Function $|g'|^q$ is η -convex, which implies

$$|g'(ta + (1-t)b)|^q \leq |g'(b)|^q + t\eta(|g'(a)|^q, |g'(b)|^q), \quad (18)$$

$t \in [0, 1]$. Using Lemma 10, inequality (18), Hölder's inequality and the properties of modulus, we get

$$\begin{aligned} & \left| \frac{g(a) + g(b)}{2} - \frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha + k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k^r \mathcal{J}_{a^+}^\alpha G(b) + {}_k^r \mathcal{J}_{b^-}^\alpha G(a)] \right| \\ & \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \int_0^1 |\Theta_{\alpha, r}(t)| |g'(ta + (1-t)b)| dt \\ & \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left(\int_0^1 |\Theta_{\alpha, r}(t)|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 |g'(ta + (1-t)b)|^q dt \right)^{\frac{1}{q}} \\ & \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left(\int_0^1 |\Theta_{\alpha, r}(t)|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 [|g'(b)|^q + t\eta(|g'(a)|^q, |g'(b)|^q)] dt \right)^{\frac{1}{q}} \\ & = \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left(\int_0^1 |\Theta_{\alpha, r}(t)|^p dt \right)^{\frac{1}{p}} \left(|g'(b)|^q + \frac{\eta(|g'(a)|^q, |g'(b)|^q)}{2} \right)^{\frac{1}{q}}. \end{aligned}$$

This completes the proof. \square

Theorem 15. Let g be differentiable on I° with $a, b \in I^\circ$. If $|g'|^q$ is η -convex on $[a, b]$ and $q > 1$ with η bounded on $|g'|^q([a, b]) \times |g'|^q([a, b])$, then the (k, r) -fractional integral inequality

$$\left| \frac{g(a) + g(b)}{2} - \frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha + k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k^r \mathcal{J}_{a^+}^\alpha G(b) + {}_k^r \mathcal{J}_{b^-}^\alpha G(a)] \right| \leq \frac{\Re^{\frac{1}{p}}}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left[|g'(b)|^q + \frac{\Xi}{b-a} \eta(|g'(a)|^q, |g'(b)|^q) \right]^{\frac{1}{q}}$$

holds, where $\frac{1}{p} + \frac{1}{q} = 1$ and \Re and Ξ are defined as in Theorem 12.

Proof. Following a similar approach as in the proof of Theorem 14, we have, by using Lemmas 10 and 11 combined with the power mean inequality plus inequality (18), that

$$\begin{aligned} & \left| \frac{g(a) + g(b)}{2} - \frac{(r+1)^{\frac{\alpha}{k}} \Gamma_k(\alpha + k)}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} [{}_k^r \mathcal{J}_{a^+}^\alpha G(b) + {}_k^r \mathcal{J}_{b^-}^\alpha G(a)] \right| \\ & \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \int_0^1 |\Theta_{\alpha, r}(t)| |g'(ta + (1-t)b)| dt \\ & \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left(\int_0^1 |\Theta_{\alpha, r}(t)| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |\Theta_{\alpha, r}(t)| |g'(ta + (1-t)b)|^q dt \right)^{\frac{1}{q}} \\ & \leq \frac{b-a}{4(b^{r+1} - a^{r+1})^{\frac{\alpha}{k}}} \left(\int_0^1 |\Theta_{\alpha, r}(t)| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |\Theta_{\alpha, r}(t)| [|g'(b)|^q + t\eta(|g'(a)|^q, |g'(b)|^q)] dt \right)^{\frac{1}{q}}. \end{aligned}$$

The required inequality follows. \square

Acknowledgements

This research was supported by FCT and CIDMA, project UID/MAT/04106/2013. The authors are grateful to the referees for their valuable comments and helpful suggestions.

References

- [1] P. Agarwal, M. Jleli and M. Tomar, Certain Hermite-Hadamard type inequalities via generalized k -fractional integrals, *J. Inequal. Appl.* **2017** (2017), Paper No. 55, 10 pp.
- [2] P. Cerone and S. S. Dragomir, *Mathematical inequalities*, CRC Press, Boca Raton, FL, 2011.
- [3] S. S. Dragomir, Generalization and reverses of the left Fejér inequality for convex functions, *J. Nonlinear Sci. Appl.* **10** (2017), no. 6, 3231–3244.
- [4] M. Eshaghi Gordji, S. S. Dragomir and M. Rostamian Delavar, An inequality related to η -convex functions (II), *Int. J. Nonlinear Anal. Appl.* **6** (2015), no. 2, 26–32.
- [5] M. Eshaghi Gordji, M. Rostamian Delavar and M. De La Sen, On ϕ -convex functions, *J. Math. Inequal.* **10** (2016), no. 1, 173–183.
- [6] J. Hadamard, Étude sur les propriétés des fonctions entières et en particulier d’une fonction considérée par Riemann, *J. Math. Pures Appl.* **9** (1893), 171–216.
- [7] M. Jleli, D. O’Regan and B. Samet, On Hermite-Hadamard type inequalities via generalized fractional integrals, *Turkish J. Math.* **40** (2016), no. 6, 1221–1230.
- [8] A. Kashuri and R. Liko, Hermite-Hadamard type inequalities for MT_m -preinvex functions, *Fasc. Math.* **58** (2017), 77–96.
- [9] M. A. Khan, Y. Khurshid and T. Ali, Hermite-Hadamard inequality for fractional integrals via η -convex functions, *Acta Math. Univ. Comenian. (N.S.)* **86** (2017), no. 1, 153–164.
- [10] S. Mubeen and G. M. Habibullah, k -fractional integrals and application, *Int. J. Contemp. Math. Sci.* **7** (2012), no. 1-4, 89–94.
- [11] E. R. Nwaeze and D. F. M. Torres, Chain rules and inequalities for the BHT fractional calculus on arbitrary timescales, *Arab. J. Math. (Springer)* **6** (2017), no. 1, 13–20. [arXiv:1611.09049](#)
- [12] M. Rostamian Delavar and M. De La Sen, Some generalizations of Hermite-Hadamard type inequalities, *SpringerPlus* **5** (2016), Paper No. 1661, 9 pp.
- [13] M. Z. Sarikaya, Z. Dahmani, M. E. Kiris and F. Ahmad, (k, s) -Riemann-Liouville fractional integral and applications, *Hacet. J. Math. Stat.* **45** (2016), no. 1, 77–89.
- [14] E. Set, M. Tomar and M. Z. Sarikaya, On generalized Grüss type inequalities for k -fractional integrals, *Appl. Math. Comput.* **269** (2015), 29–34.
- [15] Y. Shuang and F. Qi, Integral inequalities of the Hermite-Hadamard type for (α, m) -GA-convex functions, *J. Nonlinear Sci. Appl.* **10** (2017), no. 4, 1854–1860.
- [16] M. Tomar, S. Mubeen and J. Choi, Certain inequalities associated with Hadamard k -fractional integral operators, *J. Inequal. Appl.* **2016** (2016), Paper No. 234, 14 pp.