

February 2018

Maximum PBH Mass and Primordially

P.H. Frampton*

Dipartimento di Matematica e Fisica "Ennio De Giorgi"
Università del Salento and INFN Lecce, Via Arnesano 73100 Lecce, Italy.

and

9a South Parade, Oxford OX2 7JL, UK

Abstract

In order to avoid unacceptable μ -distortions inconsistent with observational data on the Cosmic Microwave Background, Primordial Black Holes (PBHs) must be less massive than $10^{12}M_{\odot}$, quite closely above the highest black hole mass yet observed. This comparableness leads us to posit that all supermassive black holes originate as PBHs.

*email: paul.h.frampton@gmail.com homepage: www.paulframpton.org

Introduction. Surely the most impressive prediction of general relativity (GR) theory is the existence of black holes in spacetime. The Schwarzschild solution of GR was discovered in 1916 [1] describing a static spherically-symmetric black hole. Because of the non-linearity of Einstein's equations, it took until 1963 before Kerr [2] discovered a solution of GR which described a rotating axially-symmetric black hole. There exist more general such solutions with electric charge but we shall assume that all the astrophysical black holes are electrically neutral so the Kerr solution is the most general classical solution needed.

One special property of Kerr black holes is commonly called the no-hair theorem [3] which states that they are completely characterised by mass and spin. This implies that there is no way of telling how a black hole was formed, whether primordially or by gravitational collapse of a pre-existing object. Nevertheless, the black holes formed by gravitational collapse can be formed only after the first stars are formed at about $t \sim 10^8$ yr (red shift $Z \sim 27$) while primordial black holes are formed before $t \sim 10^7$ seconds (red shift $Z \sim 2 \times 10^6$).

Stars have masses below $1,000M_\odot$ and therefore can collapse into black holes only lighter than this. Of course, heavier black holes can be produced by accretion and mergers of lighter black holes, although it seems extremely unlikely that a supermassive black hole like the one near the centre of gravity of the Milky Way called SagA* which has mass $M \sim 4 \times 10^6 M_\odot$ could have been so formed. Surely SagA* was rather seeded much earlier in the expansion era as we shall advocate in this Letter?

Let us begin by studying one thing which is certain, that the CMB spectrum is extraordinarily close to the black-body Planckian formula for frequency ν and temperature T :

$$F(\nu, T) = \left(\frac{2h\nu^3}{c^2} \right) [\exp(h\nu/kT) - 1]^{-1} \quad (1)$$

applicable when the electron-photon plasma is fully in thermal equilibrium, as happens when the Compton and Double Compton scatterings are faster than the cosmological expansion. Eq.(1) famously agrees with the measured CMB better than any terrestrially-measured black-body spectrum.

As a comparison to Eq.(1) absence of perfect thermal equilibrium in the electron-photon plasma can lead to a distorted CMB spectrum [4] which can be parametrised *e.g.* by

$$F(\nu, T) = \left(\frac{2h\nu^3}{c^2} \right) [\exp([h\nu/kT] + \mu) - 1]^{-1} \quad (2)$$

in which the chemical potential μ is strongly constrained, by the aforementioned accuracy of agreement with Eq.(1), to $\mu \leq 10^{-4}$ while planned experiments aim for $\mu \leq 10^{-9}$; a careful analysis is provided in [5].

Maximum PBH Mass. The mass of a PBH is tied to the horizon size at the time of PBH formation. At cosmological time t the PBH mass is given within an order of magnitude by

$$M_{PBH} \simeq 10^5 M_{\odot} \left(\frac{t}{1 \text{ second}} \right) \quad (3)$$

and therefore the maximum M_{PBH} depends on the maximum time of PBH formation.

In order that the CMB spectrum be given by Eq.(1) rather than by Eq.(2) with an unacceptably large μ , the PBH formation must take place while the electron-photon plasma remains in excellent thermal equilibrium. There is a cosmic time, called the thermalisation time t_{th} , after which the Compton and Double Compton scattering cannot keep up with the Hubble expansion so that thermal equilibrium becomes unacceptably inexact.

Assuming, as is justified *a posteriori*, that the thermalisation time occurs during the radiation-dominated era, its value is calculable with sufficient precision to cite a t_{th} accurate to an order of magnitude [5] so ignoring factors of order one we shall adopt here the value

$$t_{th} = 10^7 \text{ seconds} \quad (4)$$

which implies an upper limit on PBH mass:

$$M_{PBH} \leq 10^{12} M_{\odot} \quad (5)$$

Observed Supermassive Black Holes As already mentioned, the supermassive black hole (SMBH) at the centre of the Milky Way (MW), SagA*, has an unusually light mass for such a SMBH at a galactic centre:

$$M_{SMBH}^{(MW)} = M_{SagA*} \sim 4 \times 10^6 M_{\odot} \quad (6)$$

The most massive known SMBH at a galactic centre is in galaxy NGC1277 with

$$M_{SMBH}^{(NGC1277)} \sim 1.7 \times 10^{10} M_{\odot} \quad (7)$$

One cannot help noticing that the maximum PBH mass expressed in Eq.(5) is not much above the maximum so far observed SMBH mass given for NGC1277 in Eq.(7). We take this coincidence seriously and not as accidental.

Remarks on Primordiality. The black hole resulting from the gravitational collapse of a star cannot be more massive than the original star. If Population III stars once existed their masses are taken to be in a range up to a maximum of $\sim 1000 M_{\odot}$. These would be the most massive stars which have ever existed so a maximum value for gravity collapse black holes, at the time of their formation and ignoring all subsequent mergers and accretion is

$$\left\{ M_{BH}^{(collapse)} \right\}_{initial} \leq 10^3 M_{\odot} \quad (8)$$

Based on Eq.(8), the subsequent increase of mass of a gravity-collapsed black hole from $Z = 27$ to $Z = 0$ to reach the mass of sagA* in Eq.(6) would be by a factor of thousands which is impossible to underwrite by accretion and merger processes. In the case of NGC1277 the needed

increase of mass to reach Eq.(7) by accretion and merging would be by a factor of *ten million* which, without needing a calculation, is impossible. Our conclusion is that :
supermassive black holes are primordial or at least seeded by primordial black holes.

Let us conclude with some remarks about stellar mass black holes. Recently, the most exciting development was the discovery of gravitational waves from merger in a binary of stellar mass black holes at 410pc ($Z \sim 0.09$) from Earth with approximate masses:

$$36M_{\odot} + 29M_{\odot} \rightarrow 62M_{\odot} + (3M_{\odot} \text{ in gravitational waves}) \quad (9)$$

The energy emitted in gravitational waves is $3M_{\odot} \equiv 5 \times 10^{54}$ ergs. It was a surprise to many in the astrophysics community that the black hole masses in the first LIGO event, Eq.(9), were so large. It was not a surprise to physicists who had studied diligently the theory of dark matter suggested in [7] where large numbers of black holes with many solar masses were predicted before the LIGO announcement.

Because of the no-hair theorem, there is no way of knowing whether the initial black holes in Eq.(9) are primordial or the results of gravitational collapse. The LIGO discovery does offer support to our theory of dark matter but it is premature to take this support too seriously [8].

In order to confirm the theory of dark matter proposed in [7] the most promising method is by an extension of the microlensing observations reported in 2000 by the well-known MACHO Collaboration [9]. That remarkable experiment was completed at the end of the twentieth century when there was a prejudice that PBHs were mostly lighter than the Sun. Impressive light curves were measured for microlensing of stars in the Large Magellanic Cloud with durations ranging from two hours to almost one year corresponding to MACHO masses in the approximate range

$$10^{-5}M_{\odot} \leq M_{MACHO} \leq 25M_{\odot} \quad (10)$$

What is eagerly awaited therefore are the results from a search for longer duration light curves using the same strategy. This requires a suitable wide-angle Southern Hemisphere (to see the LMC and SMC) telescope. The best present choice (prior to LSST) is the Blanco 4m telescope which is fitted with a 520-megapixel camera (DECam). Such an experiment could identify the dark matter with certainty.

Acknowledgement

We acknowledge useful discussions with S. Sarkar and J. Silk. We thank INFN for support and the Physics Department at the University of Salento for hospitality.

References

- [1] K. Schwarzschild,
On the Gravitational Field of a Mass Point According to Einstein's Theory.
Sitzungsber.Preuss.Akad.Wiss.Berlin (Math.Phys.) **1916**, 189 (1916).
- [2] R.P. Kerr,
Gravitational Field of a Spinning Mass as an Example of Algebraically Special Metrics.
Phys. Rev. Lett. **11**, 237 (1963).
- [3] B. Carter,
Axisymmetric Black Hole Has Only Two Degrees of Freedom.
Phys. Rev. Lett. **26**, 331 (1971).
- [4] E. Pajer and M. Zaldarriaga,
A Hydrodynamical Approach to CMB μ -Distortion from Primordial Perturbations.
JCAP **1302**:036 (2013). [arXiv:1206.4479\[astro-ph.CO\]](#).
- [5] T. Nakata, B. Carr and J. Silk,
Limits on Primordial Black Holes from μ Distortions in Cosmic Microwave Background.
[ariv:1710.06945\[astro-ph.CO\]](#).
- [6] B.P. Abbott, *et al.* (LIGO Scientific and Virgo Collaborations),
Observation of Gravitational Waves from a Binary Black Hole Merger.
Phys. Rev.Lett. **116**, 061102 (2016). [arXiv 1602.03837\[gr-qc\]](#).
- [7] P.H. Frampton,
Searching for Dark Matter Constituents with Many Solar Masses.
Mod. Phys. Lett. **A31**, 1650093 (2016). [arXiv:1510.00400\[hep-ph\]](#).
- [8] S. Bird, I. Cholis, J.B. Muñoz, Y. Ali-Haïmoud, M. Kamionkowski, E.D. Kovetz,
A. Racanelli and A.G. Riess, *Did LIGO Detect Dark Matter?*
Phys. Rev. Lett. **116**, 201301 (2016). [arXiv:1603.00464\[astro-ph.CO\]](#).
- [9] C. Alcock, *et al.*, (The MACHO Collaboration),
The MACHO Project: Microlensing Results from 5.7 Years of LMC Observations.
Astrophys. J. **542**, 281 (2000). [arXiv:astro-ph/0001272](#).