# ON UNIQUENESS OF WEAK SOLUTIONS OF THE INCOMPRESSIBLE NAVIER-STOKES EQUATIONS IN 3-DIMENSIONAL CASE

KAMAL N. SOLTANOV

ABSTRACT. In this article we study the uniqueness of the weak solution of the incompressible Navier-Stokes Equation in the 3-dimensional case with use of different approach. Here the uniqueness of the obtained by Leray of the weak solution is proved in the case, when datums from spaces that are densely contained into spaces of datums for which was proved the existence of the weak solution. Moreover we investigate the solvability and uniqueness of the weak solutions of problems associated with investigation of the main problem.

## Contents

- 1. Introduction
- 2. Preliminary results
- 3. One conditional uniqueness theorem for problem  $(1.1^1)$  (1.3)
- 4. Restriction of problem  $(1.1^1) (1.3)$
- 5. Existence of Solution of Problem (3.3) (3.5)
  - 5.1. A priori estamations
  - 5.2. Boundedness of trilinear form
  - 5.3. Boundedness of u'
  - 5.4. Weakly compactness of operator B
  - 5.5. Realisation of initial condition
- 6. Uniqueness of Solution of Problem (3.3) (3.5)
- 7. Proof of Theorem 2
- 8. Conclusion
- 9. References

#### 1. INTRODUCTION

In this article we investigate Navier-Stokes equation in the incompressible case, i.e. we consider the following system of equations:

(1.1) 
$$\frac{\partial u_i}{\partial t} - \nu \Delta u_i + \sum_{j=1}^d u_j \frac{\partial u_i}{\partial x_j} + \frac{\partial p}{\partial x_i} = f_i, \quad i = \overline{1, d},$$

(1.2) 
$$\operatorname{div} u = \sum_{i=1}^{d} \frac{\partial u_i}{\partial x_i} = 0, \quad x \in \Omega \subset \mathbb{R}^d, t > 0$$

<sup>2010</sup> Mathematics Subject Classification. Primary 35K55, 35K61, 35D30, 35Q30; Secondary 76D03, 76N10.

Key words and phrases. 3D-Navier-Stokes Equations, Uniqueness, Solvability.

#### KAMAL N. SOLTANOV

(1.3) 
$$u(0,x) = u_0(x), \quad x \in \Omega; \quad u \mid_{(0,T) \times \partial \Omega} = 0$$

where  $\Omega \subset \mathbb{R}^d$  is a bounded domain with sufficiently smooth boundary  $\partial\Omega$ , T > 0 is a positive number. As it is well known Navier-Stokes equation describe the motion of a fluid in  $\mathbb{R}^d$  (d = 2 or 3). These equations are to be solved for an unknown velocity vector  $u(x,t) = \{u_i(x,t)\}_1^d \in \mathbb{R}^d$  and pressure  $p(x,t) \in \mathbb{R}$ , defined for position  $x \in \mathbb{R}^d$  and time  $t \ge 0$ ,  $f_i(x,t)$  are the components of a given, externally applied force (e.g. gravity),  $\nu$  is a positive coefficient (the viscosity),  $u_0(x) \in \mathbb{R}^d$  is a sufficiently smooth vector function (vector field).

As is well-known of [1] is shown (see, also, [2], [3], [4], [5], [7]) that the Navier– Stokes equations (1.1), (1.2), (1.3) in three dimensions have a weak solution (u, p) with suitable properties. But the uniqueness of weak solutions of the Navier–Stokes equation is not known in three space dimensions case. Uniqueness of weak solution in two space dimensions case were proved ([8], [7], see also [9]), and under complementary conditions on smoothnes of the solution three dimensions case was also studied (see, for example, [7], [28], [13], etc.). For the Euler equation, uniqueness of weak solutions is strikingly false (see, [10], [11]).

It is needed to note that the regularity of solutions in three dimensions case were investigated and partial regularity of the suitable weak solutions of the Navier– Stokes equation were obtained (see, [12], [14], [15], [7], [2]). There exist many works which study different properties of solutions of the Navier–Stokes equation (see, for example, [7], [2], [15], [5], [17], [18], [19], [20], [22], [24], [25], [26], [27], [28]), etc.) and also different modifications of Navier–Stokes equation (see, for example, [2], [7], [31], etc.).

It should be noted that under various complementary conditions of the type of certain smoothness of the weak solutions different results on the uniqueness of solution of the incompressible Navier-Stokes equation in 3D case earlier were obtained (see, e. g. [9], [7], [28], etc.). Here we would like to note the result of article [13] that possesses of some proximity to the result of this article. In this article the system of equations  $(1.1^1) - (1.3)$ , which is obtained from the incompressible Navier–Stokes system by using of the Hopf-Leray approach is examined (that below will be explained, e.g. as in [28]) in the following form

$$Nu = \mathbf{u} + \nu Au + B(u) = f, \quad \gamma_0 u = u_0,$$

where  $B(u) \equiv \sum_{j=1}^{3} u_j \frac{\partial u_i}{\partial x_j}$  and  $\gamma_0 u \equiv u(0)$ . In which the author shows that  $(N, \gamma_0)$ :  $Z \longrightarrow L^2(0, T : H^{-1/2}(\Omega)) \times H^{1/2}(\Omega)$  is the continuous operator under the condition that  $\Omega \subset R^3$  is a bounded region whose boundary  $\partial\Omega$  is a closed manifold of class  $C^{\infty}$ , where

$$Z = \left\{ \left. u \in L^2\left(0,T:H^{3/2}\left(\Omega\right)\right) \right| \; \stackrel{\bullet}{u} \in L^2\left(0,T:H^{-1/2}\left(\Omega\right)\right) \right\}.$$

Moreover, here the following result is proved: if to denote by  $F_{\gamma_0}$  the image:  $N(Z_{u_0}) = F_{\gamma_0}$  for  $u_0 \in H^{1/2}(\Omega)$  then for each  $f \in F_{\gamma_0}$  there exists only one solution  $u \in Z$  such that Nu = f and  $\gamma_0 u = u_0$ , here  $Z_{u_0} = \{u \in Z | \gamma_0 u = u_0\}$ . In this article also the density in  $L^2(0, T : H^{-1/2}(\Omega))$  of the defined above set  $F_{\gamma_0}$ in the topology of  $L^p(0, T : H^{-l}(\Omega))$  is shown under certain conditions on p, l. Here other interesting results for the operator N relatively of the dependence of the image of N from the selected domain of definition N are obtained. The proof given in [13] is similar to the proof of [7] and [28], but the result not follows from their results.

In this article we begin with the explanation why for the study of the posed question one must investigate the problem  $(1.1^1)$  - (1.3). For this we use the approach Hopf-Leray (with taking into account of the result of de Rham) for study the existence of the weak solution of the considered problem as usually all of the above mentioned authors.

Unlike above results here we study the question on the uniqueness in the case when the weak solution u of the problem  $(1.1^1) - (1.3)$  is contained of  $\mathcal{V}(Q^T)$ , and, as is well-known, the following condition is sufficiently for this: the functions  $u_0$ and f satisfy conditions

$$u_0 \in H(\Omega), \quad f \in L^2(0,T; V^*(\Omega)).$$

**Notation 1.** The result obtained for the problem  $(1.1^1)$  - (1.3) allows us to respond to the posed question, namely to prove the uniqueness of the vector velocity u.

So, in this article an investigation of the question on uniqueness of the weak solutions u in the sense of Hopf-Leray of the mixed problem with Dirichlet boundary condition for the incompressible Navier-Stokes system in the 3D case is given. For investigation we use an approach that is different from usual methods which are used for investigation of the question on the uniqueness of solution. The approach used here allows us to receive more general result on the posed question. More precisely with use of this approach more general uniqueness theorem of the weak solutions (of the vector velocity u) of the problem obtained from mixed problem for the incompressible Navier–Stokes equation, by using of the Hopf-Leray approach is proved. Moreover in order to carry out the proof of the main result in the beginning we study the auxiliary problems, more exactly we prove the existence and uniqueness of the weak solutions of auxiliary problems.

For study of the uniqueness of solution of the problem we use the variational formulation of the problem according to J. Leray [1] and E. Hopf [23] as above mentioned authors. As is well-known, on the existence of solution of problem  $(1.1^1)$  - (1.3) exist many results (see, [7], [28] and [6]). We will formulate here one of these general results from the book of [28]

**Theorem 1.** ([28]) Let  $\Omega$  be a Lipschitz open bounded set in  $\mathbb{R}^d$ ,  $d \leq 4$ . Let there be given f and  $u_0$  which

satisfy  $f \in L^2(0,T;V^*(\Omega))$  and  $u_0 \in H(\Omega)$ . Then there exists at least one function u which

satisfies  $u \in L^{2}(0,T;V(\Omega)), \frac{du}{dt} \in L^{1}(0,T;V^{*}(\Omega)), u(0) = u_{0}$  and the equation

(1.4) 
$$\frac{d}{dt}\langle u,v\rangle - \langle \nu\Delta u,v\rangle + \left\langle \sum_{j=1}^{d} u_j \frac{\partial u}{\partial x_j},v \right\rangle = \langle f,v\rangle$$

for any  $v \in V(\Omega)$ . Moreover,  $u \in L^{\infty}(0,T; H(\Omega))$  and u(t) is weakly continuous from [0,T] into  $H(\Omega)$  (i. e.  $\forall v \in H(\Omega), t \longrightarrow \langle u(t), v \rangle$  is a continuous scalar function, and consequently,  $\langle u(0), v \rangle = \langle u_0, v \rangle$ ).

"Moreover, in the case when d = 3 a weak solution u satisfy

$$u \in V(Q^T), \quad u' \equiv \frac{\partial u}{\partial t} \in L^{\frac{4}{3}}(0,T;V^*(\Omega)),$$

and also is almost everywhere equal to some continuous function from [0, T] into H, so that (1.3) is meaningful. with use of the obtained properties that any weak solution belong to the bounded subset of

$$\mathcal{V}\left(Q^{T}\right) \equiv V\left(Q^{T}\right) \cap W^{1,4/3}\left(0,T;V^{*}(\Omega)\right)$$

and satisfies the equation (1.4)."

In what follows we will base on the mentioned existence theorem of the solution of problem  $(1.1^1)$  - (1.3) and the added notation as principal result, since we as well investigate of the weak solution of the problem that is mentioned in Theorem 1, but by other way.

Then we can formulate the main result of this article in the following form.

**Theorem 2.** Let  $\Omega \subset \mathbb{R}^3$  be a domain of  $Lip_{loc}$  (will be defined below; see, Section 4), T > 0 be a number. If given functions  $u_0$ , f satisfy of conditions  $u_0 \in H^{1/2}(\Omega)$ ,  $f \in L^2(0,T; H^{1/2}(\Omega))$  then the weak solution  $u \in \mathcal{V}(Q^T)$  of the problem (1.1<sup>1</sup>) - (1.3) given by the above mentioned theorem is unique.

This article is organized as follow. In Section 2 we adduce some known results and the explanation of the relation between problems (1.1) - (1.3) and  $(1.1^1) - (1.3)$ . This section contains some necessary technical lemmas appropriate for the study of problem  $(1.1^1) - (1.3)$ . In Section 3 we prove one result on uniqueness of solution of problem  $(1.1^1) - (1.3)$  on some restriction on solution by use of the several modification of the well-known approach. In Section 4 we by use of the new approach transform problem  $(1.1^1) - (1.3)$  to auxiliary problems. In Section 5 we investigate the existence of the solution and, in Section 6 the uniqueness of solution of the auxiliary problem. In Section 7 we prove of the main result, i.e. Theorem 2.

### 2. Preliminary results

In this section, we briefly recall the background material, definitions of the appropriate spaces of Sobolev space type, deduce the necessary auxiliary results and introduce some of the notation that is needed for the results presented later in sections. Moreover, we recall the basic setup and results regarding of the weak solutions of the Navier–Stokes equations used throughout this paper. As is well known (see, e. g. [7], [28] and references therein) problem (1.1) - (1.3) possesses weak solution in the space  $\mathcal{V}(Q^T) \times L^2(Q^T), \mathcal{V}(Q^T)$  will be defined later on, for any  $u_{0i}(x)$ ,  $f_i(x,t)$  ( $i = \overline{1,3}$ ) which are contained in the suitable spaces (in the case d = 3, that we will investigate here, essentially).

**Definition 1.** Let  $\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz open set and  $Q^T \equiv (0,T) \times \Omega$ , T > 0 be a number. Let  $V(Q^T)$  be the space determined as

$$V\left(Q^{T}\right) \equiv L^{2}\left(0,T;V\left(\Omega\right)\right) \cap L^{\infty}\left(0,T;\left(H\left(\Omega\right)\right)^{d}\right),$$

where  $V(\Omega)$  is the closure in  $(H_0^1(\Omega))^d$  of

$$\left\{\varphi \mid \varphi \in \left(C_{0}^{\infty}\left(\Omega\right)\right)^{d}, \operatorname{div} \varphi = 0\right\}$$

the dual  $V(\Omega)$  determined as  $V^*(\Omega)$  and  $(H(\Omega))^d$  is the closure in  $(L^2(\Omega))^d$  of

$$\left\{\varphi \mid \varphi \in \left(C_0^{\infty}\left(\Omega\right)\right)^d, \operatorname{div} \varphi = 0\right\}.$$

Moreover we set also the space  $\mathcal{V}(Q^T) \equiv V(Q^T) \cap W^{1,4/3}(0,T;V^*(\Omega)).$ 

Here as is well-known  $L^{2}(\Omega)$  is the Lebesgue space and  $H^{1}(\Omega)$  is the Sobolev space, that are the Hilbert spaces and

$$H_{0}^{1}\left(\Omega\right)\equiv\left\{ v\mid v\in H^{1}\left(\Omega\right),\;v\mid_{\left.\partial\Omega\right.}=0\right\} .$$

In this case as is well-known (see, e.g. [28])  $H(\Omega)$  and  $V(\Omega)$  are the Hilbert spaces, also.

We assume that given functions  $u_0$  and f satisfy

$$u_0 \in (H(\Omega))^d$$
,  $f \in L^2(0,T; V^*(\Omega))$ 

where  $V^{*}(\Omega)$  is the dual space of  $V(\Omega)$ .

In order to adduce of the definition of the weak solution of the problem  $(1.1^1)$  - (1.3) we would like to note that we will investigate of the weak solutions of problem (1.1) - (1.3) in the sense of J. Leray [1] by use of his approach (see, also [7], [28]). This approach shows that for study of the posed problem it is sufficient to investigate of same question for the following problem by virtue of de Rham result (see, books [7], [28], etc. where sufficiently clearly explained this property of the posed problem):

(1.1<sup>1</sup>) 
$$\frac{\partial u_i}{\partial t} - \nu \Delta u_i + \sum_{j=1}^d u_j \frac{\partial u_i}{\partial x_j} = f(t, x)_i, \quad i = \overline{1, d}, \ \nu > 0$$

(1.2) 
$$\operatorname{div} u = \sum_{i=1}^{d} \frac{\partial u_i}{\partial x_i} = \sum_{i=1}^{d} D_i u_i = 0, \quad x \in \Omega \subset \mathbb{R}^d, \ t > 0,$$

(1.3) 
$$u(0,x) = u_0(x), \quad x \in \Omega; \quad u \mid_{(0,T) \times \partial \Omega} = 0$$

In order to explain that the investigation of the posed question for problem  $(1.1^1)$  - (1.3) is sufficient for our goal we represent here some results of the book [28] which have the immediate relation to this problem.

**Proposition 1.** ([28]) Let  $\Omega$  be a bounded Lipschitz open set in  $\mathbb{R}^d$  and  $f = (f_1, ..., f_n)$ ,  $f_i \in \mathcal{D}'(\Omega)$ ,  $1 \leq i \leq d$ . A necessary and sufficient condition that  $f = \operatorname{grad} p$  for some p in  $\mathcal{D}'(\Omega)$ , is that  $\langle f, v \rangle = 0 \ \forall v \in V(\Omega)$ .

**Proposition 2.** ([28]) Let  $\Omega$  be a bounded Lipschitz open set in  $\mathbb{R}^d$ .

(i) If a distribution p has all its first-order derivatives  $D_i p$ ,  $1 \le i \le d$  in  $L^2(\Omega)$ , then  $p \in L^2(\Omega)$  and

$$\|p\|_{L^2(\Omega)/R} \le c(\Omega) \|\operatorname{grad} p\|_{L^2(\Omega)};$$

(ii) If a distribution p has all its first derivatives  $D_i p$ ,  $1 \leq i \leq d$  in  $H^{-1}(\Omega)$ , then  $p \in L^2(\Omega)$  and

$$\|p\|_{L^2(\Omega)/R} \le c(\Omega) \|\operatorname{grad} p\|_{H^{-1}(\Omega)}.$$

In both cases, if  $\Omega$  is any open set in  $\mathbb{R}^d$ , then  $p \in L^2_{loc}(\Omega)$ .

Combining these results, one can note that if  $f \in H^{-1}(\Omega)$  (or  $f \in L^2(\Omega)$ ) and (f, v) = 0, then  $f = \operatorname{grad} p$  with  $p \in L^2(\Omega)$  (or  $p \in H^1(\Omega)$ ) if  $\Omega$  is a Lipschitz open bounded set.

**Theorem 3.** ([28]) Let  $\Omega$  be a Lipschitz open bounded set in  $\mathbb{R}^d$ . Then

$$H^{\perp} = \left\{ u \in L^{2}(\Omega) : u = \operatorname{grad} p, \ p \in H^{1}(\Omega) \right\};$$
$$H = \left\{ u \in L^{2}(\Omega) : \operatorname{div} u = 0, \ u \mid_{\partial \Omega} = 0 \right\}.$$

**Lemma 1.** ([28]) Let  $V, H, V^*$  be three Hilbert spaces, each space included in the following one  $V \subset H \equiv H^* \subset V^*$ ,  $V^*$  being the dual of V and all the injections are continuous. If a function u belongs to  $L^2(0,T;V)$  and its derivative u' belongs to  $L^2(0,T;V^*)$ , then u is almost everywhere equal to a function continuous from [0,T] into H and we have the following equality, which holds in the scalar distribution sense on (0,T):

$$\frac{d}{dt} \left\| u \right\|^2 = 2 \left\langle u', u \right\rangle.$$

Consequently, if one will seek of weak solution of the problem (1.1) - (1.3) by acording Hopf-Leray then one can get the following equation

(2.1) 
$$\frac{d}{dt}\langle u, v \rangle - \langle \nu \Delta u, v \rangle + \left\langle \sum_{j=1}^{d} u_j \frac{\partial u}{\partial x_j}, v \right\rangle = \langle f, v \rangle - \langle \nabla p, v \rangle,$$

where  $v \in V(\Omega)$  is arbitrary. Here if we consider of the last adding in the right side then at illumination of above results (1, 2 and 3) using integration by parts and taking into account that  $v \in V(\Omega)$ , i.e. that div v = 0 and  $v \mid_{(0,T) \times \partial \Omega} = 0$  we get the equation

(2.2) 
$$\langle \nabla p, v \rangle \equiv \int_{\Omega} \nabla p \cdot v \, dx = \int_{\Omega} p \operatorname{div} v \, dx = 0, \quad \forall v \in V(\Omega)$$

by virtue of de Rham result. Consequently taking into account (2.2) in (2.1) we obtain equation (1.4) that shows why for study of the posed question is enough to study problem  $(1.1^1) - (1.3)$ .

So we can continue our investigation of problem  $(1.1^1)$  - (1.3) in the case when d = 3.

Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain with the boundary  $\partial\Omega$  of the Lipschitz class. We will denote by  $\mathcal{H}^{1/2}(\Omega)$  the vector space defined by

$$\mathcal{H}^{1/2}\left(\Omega\right) \equiv \left\{w \mid w_{i} \in H^{1/2}\left(\Omega\right), \ i = 1, 2, 3\right\},\$$

where  $H^{1/2}(\Omega)$  is the Sobolev-Slobodeckij space  $W^{1/2,2}(\Omega)$  (see, [29], etc.). As is well-known (see, e.g. [29], [30] and references therein) the trace for the function of the space  $H^{1/2}(\Omega)$  is defined, which is necessary for application of our approach to the considered problem. We will prove the main theorem under this additional condition that is the sufficient condition for present investigation.

**Definition 2.** A  $u \in \mathcal{V}(Q^T)$  is called a solution of problem (1.1<sup>1</sup>) - (1.3) if u(t, x) satisfies the following equation

$$\frac{d}{dt}\langle u,v\rangle - \langle \nu\Delta u,v\rangle + \left\langle \sum_{j=1}^{d} u_j D_j u,v \right\rangle = \langle f,v\rangle$$

a. e. on (0,T) for any  $v \in V(\Omega)$  and u is weakly continuous from [0,T] into H, i. e.  $u(0,x) = u_0(x)$  holds. Consequently, in what follows we will use this definition together with the standard notation that is used usually. It should be noted that in the case when d = 3was proved, that the term  $\sum_{j=1}^{3} u_j D_j u \equiv B(u)$  belong to  $L^{4/3}(0,T; V^*(\Omega))$  (see, e. g. the books [7], [28]).

Let the posed problem have two different solutions  $u, v \in \mathcal{V}(Q^T)$ , then within the known approach we get the following problem for the function w(t, x) = u(t, x) - v(t, x)

(2.3) 
$$\frac{1}{2}\frac{\partial}{\partial t} \left\|w\right\|_{2}^{2} + \nu \left\|\nabla w\right\|_{2}^{2} + \sum_{j,k=1}^{3} \left\langle\frac{\partial v_{k}}{\partial x_{j}}w_{k}, w_{j}\right\rangle = 0,$$

(2.4)  $w(0,x) \equiv w_0(x) = 0, \quad x \in \Omega; \quad w \mid_{(0,T) \times \partial \Omega} = 0,$ 

where  $\langle g,h\rangle = \sum_{i=1\Omega}^{3} \int g_i h_i dx$  for any  $g,h \in (H(\Omega))^3$ , or  $g \in V(\Omega)$  and  $h \in V^*(\Omega)$ , respectively. So, for the proof of the uniqueness of solution it is necessary to show

that  $w \equiv 0$  in some sense. In the next section we will study the uniqueness by use of some modification of the above well-known approach, which gives we only the conditional result. But in sections 4-7 for study of the posed question we will pursue the basic approach of this paper, therefore further in this section we consider questions that are necessary for employing of this approach.

As our purpose is the investigation of the uniqueness of solution of problem  $(1.1^1) - (1.3)$  therefore we will go over to the discussion of this question. Beginning with mentioned explanations we will do some remarks about properties of solutions of problem  $(1.1^1) - (1.3)$ . As is known ([1], [2], [7]), problem  $(1.1^1) - (1.3)$  is solvable and possesses weak solution that is contained in the space  $\mathcal{V}(Q^T)$  denoted in Definition 1. Therefore we will conduct our study under the condition that problem  $(1.1^1) - (1.3)$  have weak solutions and they belong to  $\mathcal{V}(Q^T)$ . For the study of the uniqueness of the posed problem in the three dimensional case we will use the ordinary approach by assuming that problem  $(1.1^1) - (1.3)$  has, at least, two different solutions  $u, v \in \mathcal{V}(Q^T)$  but by employing a different procedure we will demonstrate that this is not possible.

Consequently, if we assume that problem  $(1.1^1) - (1.3)$  have two different solutions then they need to be different at least on some subdomain  $Q_1^T$  of  $Q^T$ . In other words there exist a subdomain  $\Omega_1$  of  $\Omega$  and an interval  $(t_1, t_2) \subseteq (0, T]$  such that

$$Q_1^T \subseteq (t_1, t_2) \times \Omega_1 \subseteq Q^T$$

with  $mes_4(Q_1^T) > 0$  for which the following is true

(2.5) 
$$mes_4(\{(t,x) \in Q^T \mid |u(t,x) - v(t,x)| > 0\}) = mes_4(Q_1^T) > 0$$

here we denote the measure of  $Q_1^T$  in  $\mathbb{R}^4$  as  $mes_4(Q_1^T)$  (Four dimensional Lebesgue measure). Whence follows, that for the subdomain  $\Omega_1$  takes place the inequation:  $mes_3(\Omega_1) > 0$ .

Even though we prove the following lemmas for d > 1, we will use them mostly for the case d = 4.

In the beginning we prove the following lemmas that we will use later on.

**Lemma 2.** Let  $G \subset \mathbb{R}^d$  be Lebesgue measurable subset then the following statements are equivalent:

1)  $\infty > mes_d(G) > 0;$ 

2) there exist a subset  $I \subset R^1$ ,  $mes_1(I) > 0$  and  $G_\beta \subset L_{\beta,d-1}$ ,  $mes_{d-1}(G_\beta) > 0$ such that  $G = \bigcup_{\beta \in I} G_\beta \cup N$ , where N is a set with  $mes_{d-1}(N) = 0$ , and  $L_{\beta,d-1}$  is the hyperplane of  $R^d$ , with  $co \dim_d L_{\beta,d-1} = 1$ , for any  $\beta \in I$ , which is generated by single vector  $y_0 \in R^d$  and defined in the following form

$$L_{\beta,d-1} \equiv \left\{ y \in R^d \mid \langle y_0, y \rangle = \beta \right\}, \quad \forall \beta \in I.$$

*Proof.* Let  $mes_d(G) > 0$  and consider the class of hyperplanes  $L_{\gamma,d-1}$  for which  $G \cap L_{\gamma,d-1} \neq \emptyset$  and  $\gamma \in I_1$ , here  $I_1 \subset R^1$  be some subset. It is clear that

$$G \equiv \bigcup_{\gamma \in I_1} \left\{ x \in G \cap L_{\gamma, d-1} \mid \gamma \in I_1 \right\}.$$

Then there exists a subclass of hyperplanes  $\{L_{\gamma,d-1} \mid \gamma \in I_1\}$  for which the inequality  $mes_{d-1}(G \cap L_{\gamma,d-1}) > 0$  is satisfied. The number of such type hyperplanes cannot be less than countable or equal it because  $mes_d(G) > 0$ , moreover this subclass of  $I_1$  must possess the  $R^1$  measure greater than 0 since  $mes_d(G) > 0$ . Indeed, let  $I_{1,0}$  be this subclass and  $mes_1(I_{1,0}) = 0$ . If we consider the set

$$\{(\gamma, y) \in I_{1,0} \times G \cap L_{\gamma,d-1} \mid \gamma \in I_{1,0}, y \in G \cap L_{\gamma,d-1}\} \subset \mathbb{R}^d$$

where  $mes_{d-1} (G \cap L_{\gamma,d-1}) > 0$  for all  $\gamma \in I_{1,0}$ , but  $mes_1 (I_{1,0}) = 0$ , then

$$mes_d(\{(\gamma, y) \in I_{1,0} \times G \cap L_{\gamma,d-1} \mid \gamma \in I_{1,0}\}) = 0.$$

On the other hand we have

$$0 = mes_d\left(\left\{(\gamma, y) \in I_1 \times G \cap L_{\gamma, d-1} \mid \gamma \in I_1\right\}\right) = mes_d\left(G\right)$$

as  $mes_{d-1}(G \cap L_{\gamma,d-1}) = 0$  for all  $\gamma \in I_1 - I_{1,0}$ . But this contradicts the condition  $mes_d(G) > 0$ . Consequently, the statement 2 holds.

Let the statement 2 holds. It is clear that the class of hyperplanes  $L_{\beta,d-1}$ defined by such way are paralell and also we can define the class of subsets of G as its cross-section with hyperplanes, i.e. in the form:  $G_{\beta} \equiv G \cap L_{\beta,d-1}$ ,  $\beta \in I$ . Then  $G_{\beta} \neq \emptyset$  and we can write  $G_{\beta} \equiv G \cap L_{\beta,d-1}$ ,  $\beta \in I$ , moreover  $G \equiv \bigcup_{\beta \in I} \{x \in G \cap L_{\beta,d-1} \mid \beta \in I\} \cup N$ . Whence we get

$$G \equiv \{(\beta, x) \in I \times G \cap L_{\beta, d-1} \mid \beta \in I, x \in G \cap L_{\beta, d-1}\} \cup N.$$

Consequently  $mes_d(G) > 0$  by virtue of conditions  $mes_1(I) > 0$  and  $mes_{d-1}(G_{\beta}) > 0$  for any  $\beta \in I$ .

From Lemma 2 it follows that for the study of the measure of some subset  $\Omega \subseteq \mathbb{R}^d$  it is enough to study its foliations by a class of suitable hyperplanes.

**Lemma 3.** Let problem  $(1.1^1) - (1.3)$  has, at least, two different solutions u, v that are contained in  $\mathcal{V}(Q^T)$  and assume that  $Q_1^T \subseteq Q^T$  is one of a subdomain of  $Q^T$ where u and v are different. Then there exists, at least, one class of parallel and different hyperplanes  $L_{\alpha}$ ,  $\alpha \in I \subseteq (\alpha_1, \alpha_2) \subset R^1$  ( $\alpha_2 > \alpha_1$ ) with codim<sub>R3</sub>  $L_{\alpha} = 1$ such, that  $u \neq v$  on  $Q_{L_{\alpha}}^T \equiv [(0, T) \times (\Omega \cap L_{\alpha})] \cap Q_1^T$ , and vice versa, here mes<sub>1</sub> (I) > 0 and  $L_{\alpha}$  are hyperplanes which are defined as follows: there is vector  $x_0 \in S_1^{R^3}(0)$ such that

$$L_{\alpha} \equiv \left\{ x \in \mathbb{R}^3 \mid \langle x_0, x \rangle = \alpha, \ \forall \alpha \in I \right\}.$$

*Proof.* Let problem  $(1.1^1)$  - (1.3) have two different solutions  $u, v \in \mathcal{V}(Q^T)$  then there exist a subdomain of  $Q^T$  on which these solutions are different. Then there are  $t_1, t_2 > 0$  such, that for any  $t \in J \subseteq [t_1, t_2] \subseteq [0, T)$  the following holds

(2.6) 
$$mes_{R^{3}}\left(\left\{x \in \Omega \mid |u(t,x) - v(t,x)| > 0\right\}\right) > 0$$

where  $mes_1(J) > 0$  by the virtue of the codition

$$mes_4(\{(t,x) \in Q^T \mid |u(t,x) - v(t,x)| > 0\}) > 0$$

and of Lemma 2. Hence follows, that there exist, at least, one class of parallel hyperplanes  $L_{\alpha}$ ,  $\alpha \in I \subseteq (\alpha_1, \alpha_2) \subset R^1$  with  $co \dim_{R^3} L_{\alpha} = 1$  such that

(2.7) 
$$mes_{R^{2}} \{ x \in \Omega \cap L_{\alpha} \mid |u(t,x) - v(t,x)| > 0 \} > 0, \ \forall \alpha \in I$$

for  $\forall t \in J$ , where the subset I is such that  $I \subseteq (\alpha_1, \alpha_2) \subset R^1$  with  $mes_1(I) > 0$ ,  $mes_1(J) > 0$  and (2.7) holds, by virtue of (2.6). This proves the "if" part of Lemma.

Now consider the converse assertion. Let there exist a class of hyperplanes  $L_{\alpha}$ ,  $\alpha \in I_1 \subseteq (\alpha_1, \alpha_2) \subset R^1$  with  $co \dim_{R^3} L_{\alpha} = 1$  that fulfills the condition of Lemma and  $I_1$  satisfies the same condition I. Then there exist, at least, one subset  $J_1$  of [0,T) such, that  $mes_1(J_1) > 0$  and the inequality  $u(t,x) \neq v(t,x)$  on  $Q_2^T$  with  $mes_4(Q_2^T) > 0$  defined as  $Q_2^T \equiv J_1 \times U_L$  takes place, where

(2.8) 
$$U_L \equiv \bigcup_{\alpha \in I_1} \{ x \in \Omega \cap L_\alpha \mid u(t, x) \neq v(t, x) \} \subset \Omega, \ t \in J_1$$

for which the inequality  $mes_{R^3}(U_L) > 0$  is satisfied by the condition and of Lemma 2.

So we get

$$u(t,x) \neq v(t,x)$$
 on  $Q_2^T \equiv J_1 \times U_L$ ,  $mes_4(Q_2^T) > 0$ .

Thus the fact that u(t,x) and v(t,x) are different functions in  $\mathcal{V}(Q^T)$  follows.  $\Box$ 

It is not difficult to see that result of Lemma 3 is independent of assumption:  $Q_1^T \subset Q^T$  or  $Q_1^T = Q^T$ .

May be one can prove more general lemmas of such type with the use of regularity properties of weak solutions of this problem (see, [10], [14], [15], etc.).

### 3. One conditional uniqueness theorem for problem $(1.1^1)$ - (1.3)

In the beginning we would like to show a what result one can receive relatively of the uniqueness of solution of the problem without of the complementary conditions. In other words we will show a what result one can obtain if to apply well-known approach for the investigation of the uniqueness of solution of problem  $(1.1^1)$  - (1.3).

Let the posed problem have two different solutions  $u, v \in \mathcal{V}(Q^T)$ , then within the known approach we get the following problem for the vector function w(t, x) = u(t, x) - v(t, x)

(3.1) 
$$\frac{1}{2}\frac{\partial}{\partial t} \|w\|_2^2 + \nu \|\nabla w\|_2^2 + \sum_{j,k=1}^3 \left\langle \frac{\partial v_k}{\partial x_j} w_k, w_j \right\rangle = 0,$$

(3.2) 
$$w(0,x) \equiv w_0(x) = 0, \quad x \in \Omega; \quad w \mid_{(0,T) \times \partial \Omega} = 0,$$

Here we will show a result when the solution of problem (3.1)-(3.2) only is zero, with use some approach that is based on the nature of the nonlinearity of this problem.

Consequently, for examination of the problem (3.1)-(3.2) in the beginning we need to study the following quadratic form ([21])

$$\sum_{j,k=1}^{d} \left( \frac{\partial v_k}{\partial x_j} w_k w_j \right) (t,x) \Longrightarrow F(t,x) \equiv \sum_{j,k=1}^{d} \left( a_{jk} w_k w_j \right) (t,x) \Longrightarrow$$
$$F(t,x) \equiv \sum_{j=1}^{d} \left( a_j \overline{w}_j^2 \right) (t,x), \quad a_j(t,x) \equiv G_j \left( D_1 v_1, \dots, D_1 v_d, \dots, D_d v_d \right)$$
here  $D_i v_k \equiv \frac{\partial v_k}{\partial x_i}, \quad i,k = \overline{1,d}$ 

i.e. the behavior of the surface generated by the quadratic polynomial function F(t,x), at the variable  $w_k$ ,  $k = \overline{1, d}$ , depende of the accelerations of the flow on the different directions.

Assume d = 3 then we have

$$a_1 = D_1 v_1; \ a_2 = D_2 v_2 - \frac{\left(D_1 v_2 + D_2 v_1\right)^2}{4a_1}; \ a_3 = \frac{\det \|D_i v_k\|_{i,k=1}^3}{\det \|D_i v_k\|_{i,k=1}^2},$$

therefore,

(3.3) 
$$F(t,x) \equiv \sum_{j,k=1}^{3} (a_{jk}w_k w_j)(t,x) \equiv \sum_{j=1}^{3} a_j(t,x) \cdot \overline{w}_j^2(t,x) =$$

for any  $(t, x) \in Q^T \equiv (0, T) \times \Omega$ , here

$$\|D_{i}v_{k}\|_{i,k=1}^{3} \equiv \left\| \begin{array}{ccc} D_{1}v_{1} & \frac{1}{2}\left(D_{1}v_{2}+D_{2}v_{1}\right) & \frac{1}{2}\left(D_{1}v_{3}+D_{3}v_{1}\right) \\ \frac{1}{2}\left(D_{1}v_{2}+D_{2}v_{1}\right) & D_{2}v_{2} & \frac{1}{2}\left(D_{2}v_{3}+D_{3}v_{2}\right) \\ \frac{1}{2}\left(D_{1}v_{3}+D_{3}v_{1}\right) & \frac{1}{2}\left(D_{2}v_{3}+D_{3}v_{2}\right) & D_{3}v_{3} \end{array} \right|$$

and

$$\|D_i v_k\|_{i,k=1}^2 \equiv \left\| \begin{array}{cc} D_1 v_1 & \frac{1}{2} \left( D_1 v_2 + D_2 v_1 \right) \\ \frac{1}{2} \left( D_1 v_2 + D_2 v_1 \right) & D_2 v_2 \end{array} \right\|$$

Thus we have

$$\begin{split} F\left(t,x\right) &\equiv \frac{1}{D_{1}v_{1}} \left[2D_{1}v_{1}w_{1} + \left(D_{1}v_{2} + D_{2}v_{1}\right)w_{2} + \left(D_{1}v_{3} + D_{3}v_{1}\right)w_{3}\right]^{2} + \\ &\frac{1}{\left(4D_{1}v_{1}\right)^{2}} \left(4D_{1}v_{1}D_{2}v_{2} - \left(D_{1}v_{2} + D_{2}v_{1}\right)^{2}\right) \times \\ &\left[\left(4D_{1}v_{1}D_{2}v_{2} - \left(D_{1}v_{2} + D_{2}v_{1}\right)^{2}\right)w_{2} + \\ \left(2D_{1}v_{1}\left(D_{2}v_{3} + D_{3}v_{2}\right) - \left(D_{1}v_{2} + D_{2}v_{1}\right)\left(D_{1}v_{3} + D_{3}v_{1}\right)\right)w_{3}\right]^{2} + \\ &\frac{1}{4} \left[4D_{1}v_{1}D_{2}v_{2}D_{3}v_{3} + \left(D_{1}v_{2} + D_{2}v_{1}\right)\left(D_{1}v_{3} + D_{3}v_{1}\right)\left(D_{2}v_{3} + D_{3}v_{2}\right) - \\ &D_{1}v_{1}\left(D_{2}v_{3} + D_{3}v_{2}\right)^{2} - D_{2}v_{2}\left(D_{1}v_{3} + D_{3}v_{1}\right)^{2} - D_{3}v_{3}\left(D_{1}v_{2} + D_{2}v_{1}\right)^{2}\right]w_{3}^{2} \end{split}$$

Hence, if one take account (3.3) in the equation (3.1) then we get

$$\frac{1}{2}\frac{\partial}{\partial t}\left\|w\right\|_{2}^{2}+\nu\left\|\nabla w\right\|_{2}^{2}+\sum_{j=1}^{3}\left\langle a_{j}\overline{w}_{j},\overline{w}_{j}\right\rangle=0,$$

and consequently,

(3.4) 
$$\frac{1}{2}\frac{\partial}{\partial t} \|w\|_2^2 = -\nu \|\nabla w\|_2^2 - \sum_{j=1}^3 \langle a_j \overline{w}_j, \overline{w}_j \rangle, \quad \|w_0\|_2 = 0.$$

This shows that if the quadratic form function  $F(t, x, w_i, w_j) = \sum_{j=1}^{3} a_j(t, x) \cdot \sum_{j=1}^{3} a_j(t, x)$ 

 $\overline{w}_i^2(t,x)$  for a.e.  $(t,x) \in Q^T$  is the quadratic polynomial that describe an ellipsoid in  $R^3$  at the variables  $(\overline{w}_i, \overline{w}_i)$ , i.e.  $a_i(t, x) \geq 0$ , then the posed problem have unique solution. It is needed to note that images of functions  $a_i(t, x)$  are contained a.e. in the bounded subset of same space where are contained images of functions  $D_i v_k$ . So is remains to investigate the cases when the above condition not is fulfilled. Here the following variants are possible:

1. Let  $F(t, x, w_i, w_i)$  describe some other surface in  $\mathbb{R}^3$ , but

$$\sum_{j=1}^{3} \langle a_j \overline{w}_j, \overline{w}_j \rangle \equiv \sum_{j=1}^{3} \int_{\Omega} a_j \overline{w}_j^2 dx \ge 0.$$

In this case we can conclude that problem have unique solution (and a solution is stable).

2. Let 
$$\sum_{j=1}^{3} \int a_j \overline{w}_j^2 dx < 0$$
 then we will study  $F(t,x) \equiv \sum_{j,k=1}^{3} (D_i v_k w_k w_j)(t,x).$ 

In this case the problem (3.4) is possible to investigate by following way. For the multilinear form in the equation it is necessary to derive suitable estimations. Thus

$$\left|\frac{1}{2}\frac{\partial}{\partial t}\left\|w\right\|_{2}^{2}=-\nu\left\|\nabla w\right\|_{2}^{2}+\left|\sum_{j,k=1}^{3}\left\langle D_{i}v_{k}w_{k},w_{j}\right\rangle\right|\leq$$

(3.5) 
$$-\sum_{j=1}^{3} \int_{\Omega} \nu |\nabla w_j(t,x)|^2 dx + \sum_{j,k=1}^{3} \int_{\Omega} |(D_i v_k w_k w_j)(t,x)| dx$$

as in this case the second adding in the right part is negative. So, in order that to continue the inequation (3.5), we will use of the corresponding estimations. Here using Hölder inequality to multilinear term we obtain the estimation  $^{1}$ 

$$|\langle D_i v_j w_i, w_j \rangle| \le ||D_i v_j||_2 ||w_i||_{p_1} ||w_j||_{p_2},$$

where  $p_1^{-1} + p_2^{-1} = 2^{-1}$ , for us sufficiently to choose,  $p_1 = p_2 = 4$ . Consequently, for  $F(t, x, w_i, w_i)$  takes place the estimation

$$\int_{\Omega} |F(t,x)| \, dx \le \sum_{i,j=1}^{3} \|D_j v_i\|_2 \, \|w_i\|_4 \, \|w_j\|_4 \, .$$

<sup>1</sup>It is known that ([2], [7])  $|\langle u_k D_i v_j, w_l \rangle| \le ||u_k||_q ||D_i v_j||_2 ||w_l||_n$ ,  $n \ge 3$ ;  $||v_j||_4 \le C (mes \ \Omega) ||Dv_j||_2^{\frac{1}{2}} ||v_j||_2^{\frac{1}{2}}$ , n = 2

Here we will use the known Gagliardo-Nirenberg-Sobolev inequation, that can be formulated as follows (see, for example, [30])

(3.6) 
$$||u||_{p_0,s} \le C(p_0, p_1, p_2, s, m) \left(\sum_{|\alpha|=m} ||D^{\alpha}u||_{p_1}\right)^{\sigma} \cdot ||u||_{p_2}^{1-\sigma}$$

inequation holds if  $1 \le p_1, p_2, p_0 \le \infty, 0 \le s < m$ , where  $C(p_0, p_1, p_2, s, m) > 0$  is constant,

$$\frac{d}{p_0} - s = \sigma \left(\frac{d}{p_1} - m\right) + (1 - \sigma) \frac{d}{p_2}, \quad \frac{s}{m} \le \sigma \le 1$$

with the following exclusions:

a) if  $s = 0, s < \frac{d}{p_1}, p_2 = \infty$  then (3.6) holds under complementary condition: or  $\lim_{x \to \infty} u(x) = 0$ , or  $u \in L^q$  for some q > 0;

b) if  $1 \le p_1 < \infty, m - s - \frac{n}{p_1} = 0, p_0 = \infty$  then (3.6) not holds in the case  $\sigma = 1$ . Hence use G-N-S inequation we get

$$||w_j||_4 \le c ||w_j||_2^{1-\sigma} ||\nabla w_j||_2^{\sigma}, \quad \sigma = \frac{3}{4},$$

here  $c \equiv C(4, 2, 2, 0, 1)$  for this case, or

$$\|w_j\|_4 \le c \|w_j\|_2^{\frac{1}{4}} \|\nabla w_j\|_2^{\frac{3}{4}} \Longrightarrow \|w_j\|_4^2 \le c^2 \|w_j\|_2^{\frac{1}{2}} \|\nabla w_j\|_2^{\frac{3}{2}}.$$

Thus

$$\int_{\Omega} |F(t,x)| \, dx \le c^2 \sum_{i,j=1}^3 \|D_j v_i\|_2 \, \|w_i\|_2^{\frac{1}{4}} \, \|\nabla w_i\|_2^{\frac{3}{4}} \, \|w_j\|_2^{\frac{1}{4}} \, \|\nabla w_j\|_2^{\frac{3}{4}}$$

if one take into account the above estimation in (3.6) then

$$\frac{1}{2}\frac{\partial}{\partial t}\|w(t)\|_{2}^{2} \leq -\sum_{j=1}^{3}\nu\|\nabla w_{j}(t)\|_{2}^{2} + c^{2}\sum_{i,j=1}^{3}\|D_{j}v_{i}(t)\|_{2}\|w_{i}(t)\|_{2}^{\frac{1}{2}}\|\nabla w_{i}(t)\|_{2}^{\frac{3}{2}}$$
$$\leq -\sum_{j=1}^{3}\|\nabla w_{j}(t)\|_{2}^{\frac{3}{2}}\left[\nu\|\nabla w_{j}(t)\|_{2}^{\frac{1}{2}} - c^{2}\sum_{i=1}^{3}\|D_{i}v_{j}(t)\|_{2}\|w_{j}(t)\|_{2}^{\frac{1}{2}}\right]$$
$$\leq -\sum_{j=1}^{n}\|\nabla w_{j}(t)\|_{2}^{\frac{3}{2}}\left[\nu\lambda_{1}^{\frac{1}{4}} - c^{2}\sum_{i=1}^{n}\|D_{i}v_{j}(t)\|_{2}\right]\|w_{j}(t)\|_{2}^{\frac{1}{2}}.$$

From here is easy follows, that if  $\nu \lambda_1^{\frac{1}{4}} \geq c^2 \sum_{i=1}^{3} \|D_i v_j(t)\|_2$  then the considered problem  $(1.1^1)$ -(1.3) has only unique solution (and a solution is stable). Thus is proved

**Theorem 4.** Let  $\Omega \in \mathbb{R}^3$  be a bounded domain of Lipschitz class and  $(u_0, f) \in$  $(H(\Omega))^3 \times L^2(0,T;V^*(\Omega))$  then as is well-known weak solution u(t,x) of problem  $(1.1^1)$ -(1.3) exists and  $u \in \mathcal{V}(Q^T)$ . Then if  $\int_{\Omega} |F(t,x)| \, dx \ge 0$ , or  $\int_{\Omega} |F(t,x)| \, dx < 0$ 

and  $\nu \lambda_{1}^{\frac{1}{4}} \geq c^{2} \sum_{i=1}^{3} \left\| D_{i} u_{j}(t) \right\|_{2}$  are fulfilled then weak solution u(t, x) is unique.

## 4. Restriction of problem $(1.1^1) - (1.3)$

From Lemma 3 it follows that for the investigation of the posed question it is enough to investigate this problem on the suitable cross-sections of the domain  $Q^T \equiv (0, T) \times \Omega$ . We introduce the following concept.

**Definition 3.** A bounded domain  $\Omega \subset \mathbb{R}^3$  with the boundary  $\partial\Omega$  is class  $Lip_{loc}$  iff  $\partial\Omega$  is a locally Lipschitz hypersurface. This means that in a neighbourhood of any point  $x \in \partial\Omega$ ,  $\partial\Omega$  admits a representation as a hypersurface  $y_3 = \psi(y_1, y_2)$  where  $\psi$  is a Lipschitz function, and  $(y_1, y_2, y_3)$  are rectangular coordinates in  $\mathbb{R}^3$  in a basis that may be different from the canonical basis  $(e_1, e_1, e_3)$ .

According to [28] one can draw the conclusion: It is useful for the sequel of this section to note that a set  $\Omega$  satisfying (1.4) is "locally star-shaped". This means that each point  $x_j \in \partial \Omega$ , has an open neighbourhood  $U_j$  such that  $U'_j = \Omega \cap U_j$  is star-shaped with respect to one of its points. According to  $\Omega$  is a locally Lipschitz we may, moreover, suppose that the boundary  $U'_j$ ,  $j \in J$  is Lipschitz, or  $\partial \Omega \in Lip_{loc}$ . If  $\partial \Omega$  is bounded, it can be covered by a finite family of such sets  $U'_j$ ,  $j \in J$ . Consequently for every cross-section  $\Omega_L \equiv \Omega \cap L \neq \emptyset$  of  $\Omega$  with arbitrary hyperplain L exists, at least, one coordinate subspace  $((x_j, x_k))$  which possesses a domain  $P_{x_i}\Omega_L$  (or union of domains) whit the Lipschitz class boundary since  $\partial \Omega_L \equiv \partial \Omega \cap L \neq \emptyset$  and isomorphically defining of  $\Omega_L$  with the affine representation, in addition  $\partial \Omega_L \iff \partial P_{x_i}\Omega_L$ .

Thus, with use of the representation  $P_{x_i}L$  of the hyperplane L we get that  $\Omega_L$  can be written in the form  $P_{x_i}\Omega_L$ , therefore an integral on  $\Omega_L$  also will defined by the respective representation, i. e. as the integral on  $P_{x_i}\Omega_L$ .

It should be noted that  $\Omega_L$  can consist of many parts then  $P_{x_i}\Omega_L$  will be such as  $\Omega_L$ . Consequently in this case  $\Omega_L$  will be as the union of domains and the following relation

$$\Omega_L = \bigcup_{r=1}^m \Omega_L^r \iff P_{x_i} \Omega_L = \bigcup_{r=1}^m P_{x_i} \Omega_L^r, \quad \infty > m \ge 1,$$

will holds by virtue of the definition 3. Therefore, each of  $P_{x_i}\Omega_L^r$  will be the domain and one can investigate these separately, because  $\Omega_L^r \subset \Omega$  and  $\partial \Omega_L^r \subset \partial \Omega$  takes place.

So, we will define subdomains of  $Q^T \equiv (0,T) \times \Omega$  as follows  $Q_L^T \equiv (0,T) \times (\Omega \cap L)$ , where L is arbitrary fixed hyperplane of the dimension two and  $\Omega \cap L \neq \emptyset$ . Therefore, we will study the problem on the subdomain defined by use of the cross-section of  $\Omega$  by arbitrary fixed hyperplane dimension two L, i.e. by the  $co \dim_{R^3} L = 1$  ( $\Omega \cap L$ , namely on  $Q_L^T \equiv (0,T) \times (\Omega \cap L)$ ).

Consequently, we will investigate uniqueness of the problem  $(1.1^1) - (1.3)$  on the "cross-section"  $Q^T$  defined by the cross-section of  $\Omega$ , where  $\Omega \subset R^3$ . This cross-section we understand in the following sense: Let L be a hyperplane in  $R^3$ , i.e. with  $co \dim_{R^3} L = 1$ , that is equivalent to  $R^2$ . We denote by  $\Omega_L$  the cross-section of the form  $\Omega_L \equiv \Omega \cap L \neq \emptyset$ ,  $mes_{R^2}(\Omega_L) > 0$ , in the particular case  $L \equiv (x_1, x_2, 0)$ . In other words, if L is the hyperplane in  $R^3$  then we can determine it as  $L \equiv \{x \in R^3 \mid a_1x_1 + a_2x_2 + a_3x_3 = b\}$ , where coefficients  $a_i, b \in R^1$  (i = 1, 2, 3) are the arbitrary fixed constants. Whence follows, that  $a_3x_3 = b - a_1x_1 - a_2x_2$  or  $x_3 = \frac{1}{a_3}(b - a_1x_1 - a_2x_2)$  if we assume  $a_i \neq 0$  (i = 1, 2, 3), or if we take substitutions:  $\frac{b}{a_3} \Longrightarrow b, \frac{a_1}{a_3} \Longrightarrow a_1$  and  $\frac{a_2}{a_3} \Longrightarrow a_2$  we derive  $x_3 \equiv \psi_3(x_1, x_2) = b - a_1x_1 - a_2x_2$  in the new coefficients.

Thus, we have

(4.1) 
$$D_3 \equiv \frac{\partial x_1}{\partial x_3} D_1 + \frac{\partial x_2}{\partial x_3} D_2 = -a_1^{-1} D_1 - a_2^{-1} D_2 \quad \&$$

(4.2) 
$$D_3^2 = a_1^{-2}D_1^2 + a_2^{-2}D_2^2 + 2a_1^{-1}a_2^{-1}D_1D_2, \quad D_i = \frac{\partial}{\partial x_i}, i = 1, 2, 3.$$

For the application of our approach we need to assume that functions  $u_0$  and f posseses some smoothness.

Now we take account the following conditions of Theorem 2  $u_0 \in \mathcal{H}^{1/2}(\Omega)$ ,  $f \in L^2(0,T;\mathcal{H}^{1/2}(\Omega))$  hold, consequently their restrictions on  $[0,T) \times \Omega_L$  are defined.

Let L be arbitrary hyperplane such that  $\Omega_L \neq \emptyset$  and  $u \in \mathcal{V}(Q^T)$  be a solution of the problem  $(1.1^1) - (1.3)$ . It is need to note the restriction of problem  $(1.1^1)$ - (1.3) to  $[0,T) \times \Omega_L$  mean the restriction on  $[0,T) \times \Omega_L$  of a solution u that is defined on  $[0,T) \times \Omega$ . As function u belong to  $\mathcal{V}(Q^T)$  therefore the restriction u on  $[0,T) \times \Omega_L$  is well defined. Then by making the restriction we obtain the following problem on  $[0,T) \times \Omega_L$ , by virtue of the above conditions of the main theorem, here T > 0 some number,

$$\frac{\partial u}{\partial t} - \nu \Delta u + \sum_{j=1}^{3} u_j D_j u = \frac{\partial u_L}{\partial t} - \nu \left( D_1^2 + D_2^2 + D_3^2 \right) u_L + u_{L1} D_1 u_L + u_{L2} D_2 u_L + u_{L3} D_3 u_L = \frac{\partial u_L}{\partial t} - \nu \left[ D_1^2 + D_2^2 + a_1^{-2} D_1^2 + a_2^{-2} D_2^2 + 2a_1^{-1} a_2^{-1} D_1 D_2 \right] u_L + u_{L1} D_1 u_L + u_{L2} D_2 u_L - u_{L3} a_1^{-1} D_1 u_L - u_{L3} a_2^{-1} D_2 u_L = \frac{\partial u_L}{\partial t} - \nu \left[ \left( 1 + a_1^{-2} \right) D_1^2 + \left( 1 + a_2^{-2} \right) D_2^2 \right] u_L - u_{L3} a_2^{-1} D_2 u_L = \frac{\partial u_L}{\partial t} - \nu \left[ \left( 1 + a_1^{-2} \right) D_1^2 + \left( 1 + a_2^{-2} \right) D_2^2 \right] u_L - u_{L3} u_L + u_{L3} D_1 u_L - u_{L3} u_L + u_{L3} D_2 u_L = \frac{\partial u_L}{\partial t} - \nu \left[ \left( 1 + a_1^{-2} \right) D_1^2 + \left( 1 + a_2^{-2} \right) D_2^2 \right] u_L - u_{L3} u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L - u_{L3} u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L - u_{L3} u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L - u_{L3} u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L - u_{L3} u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L - u_{L3} u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L + u_{L3} D_1 u_L - u_{L3} u_L + u_{L3} D_1 u_L + u_{L3} u_L + u_{L3}$$

(4.3)  $2\nu a_1^{-1}a_2^{-1}D_1D_2u_L + (u_{L1} - a_1^{-1}u_{L3})D_1u_L + (u_{L2} - a_2^{-1}u_{L3})D_2u_L = f_L$ on  $(0,T) \times \Omega_L$ , by virtue of (4.1) and (4.2). We get

(4.4) div 
$$u_L = D_1 \left( u_L - a_1^{-1} u_{L3} \right) + D_2 \left( u_L - a_2^{-1} u_{L3} \right) = 0, \quad x \in \Omega_L, \ t > 0$$

(4.5) 
$$u_L(0,x) = u_{L0}(x), \quad (t,x) \in [0,T] \times \Omega_L; \quad u_L \mid_{(0,T) \times \partial \Omega_L} = 0$$

by using of same way.

Consequently, we restricted the problem  $(1.1^1)$  - (1.3) to problem (4.3) - (4.5) the study of which give we possibility to define properties of solutions u of problem  $(1.1^1)$  - (1.3) on each cross-section  $[0, T) \times \Omega_L \equiv Q_L^T$ .

In the beginning it is necessary to investigate the existence of the solution of problem (4.3) - (4.5) and determine the space where the existing solutions are contained. Consequently, for ending the proof of the uniqueness theorem for main problem it is enough to prove the existence theorem and the uniqueness theorem for the derived problem (4.3) - (4.5), in this case. So now we will investigate of problem (4.3) - (4.5).

We would like to note: Let  $L \subset R^3$  is the hyperplane for which  $\Omega \cap L \neq \emptyset$  then there is, at least, one 2-dimensional subspace in the given coordinat system that one can determine as  $(x_i, x_j)$ , consequently,  $P_{x_k}L = R^2$ , (i, j, k = 1, 2, 3) i.e.

$$L \equiv \{ x \in R^3 \mid x = (x_i, x_j, \psi_L(x_i, x_j)), (x_i, x_j) \in R^2 \}$$

and

$$\Omega \cap L \equiv \{x \in \Omega \mid x = (x_i, x_j, \psi_L(x_i, x_j)), (x_i, x_j) \in P_{x_k}(\Omega \cap L)\}$$

hold, where  $\psi_L$  is the affine function such as the mentioned above function and it is the bijection.

Thereby, in this case by applying of the mentioned restriction to functions

 $u(t, x_1, x_2, x_3), f(t, x_1, x_2, x_3), u_0(x_1, x_2, x_3)$ 

we obtain the following representations

$$u(t, x_i, x_j, \psi_L(x_i, x_j)) \equiv v(t, x_i, x_j), \ f(t, x_i, x_j, \psi_L(x_i, x_j) \equiv \phi(x_i, x_j)$$

and

$$u_0(x_i, x_j, \psi_L(x_i, x_j)) \equiv v_0(x_i, x_j) \quad \text{on } (0, T) \times P_{x_k} \Omega_L,$$

respectively.

So, each of the functions obtained by the previous transformation depends only on the independent variables:  $t, x_i$  and  $x_j$ .

4.1. On Dirichlet to Neumann map. As is known ([35], [36], [39], [37], [38] etc.) the Dirichlet to Neumann map is single-value maping if the homogeneous Dirichlet problem for elliptic equation has only trivial solution, i. e. zero not is eigenvalue of this problem. Consequently, we need to show that the homogeneous Dirichlet problem for elliptic equation appropriate to considered problem satisfies of this property. So, we will show that the obtained here problem satisfies of the corresponding condition of results of such type from mentioned articles.

**Proposition 3.** The homogeneous Dirichlet. problem for elliptic part of problem (3.3) - (3.5) has only trivial solution.

*Proof.* If consider the elliptic part of problem (3.3) - (3.5) then we get the problem

$$-\Delta u_L + Bu_L \equiv -\nu \left[ \left( 1 + a_1^{-2} \right) D_1^2 + \left( 1 + a_2^{-2} \right) D_2^2 + 2a_1^{-1}a_2^{-1}D_1D_2 \right] u_{Li} +$$

 $\left(u_{L1} - a_1^{-1} u_{L3}\right) D_1 u_L + \left(u_{L2} - a_2^{-1} u_{L3}\right) D_2 u_L = 0, \ x \in \Omega_L, \quad u_L \mid_{\partial \Omega_L} = 0,$ where  $\Omega_L = \Omega \cap L.$ 

We assume this problem has a nontrivial solution and we will show that it is unpossible. Let  $u_L \in V(\Omega_L)$  be nontrivial solution of this problem then we get the following equation

$$0 = \langle -\Delta u_L + B u_L, u_L \rangle_{P_{x_3}\Omega_L}$$

hence

$$= -\nu \sum_{i=1}^{3} \left\langle \left[ \left( D_{1}^{2} + D_{2}^{2} \right) + \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right)^{2} \right] u_{Li}, u_{Li} \right\rangle_{P_{x_{3}}\Omega_{L}} + \\ \sum_{i=1}^{3} \int_{P_{x_{3}}\Omega_{L}} \left[ u_{L1}D_{1}u_{Li}u_{Li} + u_{L2}D_{2}u_{Li}u_{Li} + \\ u_{L3} \left( -a_{1}^{-1}D_{1} - a_{2}^{-1}D_{2} \right) u_{Li}u_{Li} \right] dx_{1}dx_{2} = \\ \nu \sum_{i=1}^{3} \int_{P_{x_{3}}\Omega_{L}} \left\{ \left( D_{1}u_{Li} \right)^{2} + \left( D_{2}u_{Li} \right)^{2} + \left[ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right]^{2} \right\} dx_{1}dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right]^{2} \right\} dx_{1}dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right]^{2} \left\{ dx_{1}dx_{2} + \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right\}^{2} \right\} dx_{1}dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right\}^{2} dx_{1}dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right]^{2} \left\{ dx_{1}dx_{2} + \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right\}^{2} dx_{1}dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right\}^{2} dx_{1}dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} \right]^{2} dx_{1}dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{1} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{1} dx_{2} + \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{1} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{1} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li} dx_{2} + \\ \left( a_{1}^{-1}D_{1} + a_{2}^{-1}D_{2} \right) u_{Li$$

$$\frac{1}{2}\sum_{i=1}^{3}\int_{P_{x_{3}}\Omega_{L}} \left[u_{L1}D_{1}\left(u_{Li}\right)^{2} + u_{L2}D_{2}\left(u_{Li}\right)^{2} + u_{L3}\left(-a_{1}^{-1}D_{1} - a_{2}^{-1}D_{2}\right)\left(u_{Li}\right)^{2}\right] dx_{1}dx_{2} \geq \nu\sum_{i=1}^{3}\int_{P_{x_{3}}\Omega_{L}} \left[|D_{1}u_{Li}|^{2} + |D_{2}u_{Li}|^{2}\right] dx_{1}dx_{2} + \left[\frac{1}{2}\sum_{i=1}^{3}\int_{P_{x_{3}}\Omega_{L}} \left[D_{1}u_{L1} + D_{2}u_{L2} + \left(-a_{1}^{-1}D_{1} - a_{2}^{-1}D_{2}\right)u_{L3}\right]|u_{Li}|^{2} dx_{1}dx_{2} = 0$$
by (4.4)
$$\frac{3}{2} = \int_{P_{x_{3}}\Omega_{L}} \left[D_{1}u_{L1} + D_{2}u_{L2} + \left(-a_{1}^{-1}D_{1} - a_{2}^{-1}D_{2}\right)u_{L3}\right]|u_{Li}|^{2} dx_{1}dx_{2} = 0$$

$$\nu \sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ |D_1 u_{Li}|^2 + |D_2 u_{Li}|^2 \right] dx_1 dx_2 - \frac{1}{2} \sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} |u_{Li}|^2 \operatorname{div} u_L dx_1 dx_2 = \nu \sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ |D_1 u_{Li}|^2 + |D_2 u_{Li}|^2 \right] dx_1 dx_2 > 0.$$

Thus the obtained contradiction shows that function  $u_L$  need be zero, i.e.  $u_L = 0$  holds.

Consequently, the Dirichlet to Neumann map is single-value operator.  $\Box$ 

It is well-known that operator  $-\Delta : H_0^1(\Omega_L) \longrightarrow H^{-1}(\Omega_L)$  generates of the  $C_0$  semigroup on  $H(\Omega_L)$  and since the inclusion  $H_0^1(\Omega_L) \subset H^{-1}(\Omega_L)$  is compact therefore  $(-\Delta)^{-1}$  is the compact operator in  $H^{-1}(\Omega_L)$ . Moreover,  $-\Delta : H^{1/2}(\partial\Omega_L) \longrightarrow H^{-1/2}(\partial\Omega_L)$  and the operator  $B : H^{1/2}(\partial\Omega_L) \longrightarrow H^{-1/2}(\partial\Omega_L)$  also possess appropriate properties of such types.

### 5. EXISTENCE OF SOLUTION OF PROBLEM (3.3) - (3.5)

So, assume conditons of Theorem 2 fulfilled, i. e.

$$u_0 \in \mathcal{H}^{1/2}(\Omega), \quad f \in L^2\left(0, T; \mathcal{H}^{1/2}(\Omega)\right),$$

then restrictions of these functions on  $\Omega_L$ ,  $Q_L^T$ , respectively, are correctly defined and belong in  $H(\Omega_L)$ ,  $L^2(0,T;V^*(\Omega_L))$ , respectively. Consequently, it is enough to study the restricted problem under conditions  $u_{0L} \in H(\Omega_L)$  and  $f_L \in L^2(0,T;V^*(\Omega_L))$ , as independent problem.

To carry out the known argument started by Leray ([1], see, also [7], [28]) we can determine the following spaces

$$V(\Omega_L) = \left\{ v \mid v \in \left( W_0^{1,2}(\Omega_L) \right)^3 \equiv \left( H_0^1(\Omega_L) \right)^3, \quad \operatorname{div} v = 0 \right\},$$

where div is regarded in the sense (4.4) and

$$V\left(Q_{L}^{T}\right) \equiv L^{2}\left(0,T;V\left(\Omega_{L}\right)\right) \cap L^{\infty}\left(0,T;\left(H\left(\Omega_{L}\right)\right)^{3}\right).$$

More exactly we will adduce definitions of these spaces such way as in Definition 1, i. e.  $V(\Omega_L)$  is the closure in  $(H_0^1(\Omega_L))^3$  of

$$\left\{\varphi \mid \varphi \in \left(C_0^{\infty}\left(\Omega_L\right)\right)^3, \operatorname{div} \varphi = 0\right\}$$

the dual  $V(\Omega_L)$  is determined as  $V^*(\Omega_L)$  and  $(H(\Omega_L))^3$  is the closure in  $(L^2(\Omega_L))^3$  of

$$\left\{\varphi \mid \varphi \in \left(C_0^{\infty}\left(\Omega_L\right)\right)^3, \operatorname{div} \varphi = 0\right\}.$$

Here  $\Omega \subset \mathbb{R}^3$  is bounded domain of  $Lip_{loc}$  and  $\Omega_L \subset \mathbb{R}^2$  is subdomain defined in Section 4 therefore,  $\Omega_L$  is Lipschitz,  $Q_L^T \equiv (0,T) \times \Omega_L$ .

Consequently, a solution of this problem will be understood as follows: Let  $f_L \in L^2(0,T; V^*(\Omega_L))$  and  $u_{0L} \in (H(\Omega_L))^3$ .

So, we can call the solution of this problem: a function  $u_L \in \mathcal{V}(Q_L^T)$  is called a solution of the problem (4.3) - (4.5) if  $u_L(t, x')$  satisfy the equation and initial condition

(5.1) 
$$\frac{d}{dt} \langle u_L, v \rangle_{\Omega_L} - \langle \nu \Delta u_L, v \rangle_{\Omega_L} + \left\langle \sum_{j=1}^3 u_{Lj} D_j u_L, v \right\rangle_{\Omega_L} = \langle f_L, v \rangle_{\Omega_L},$$
$$\langle u_L(t), v \rangle|_{t=0} = \langle u_{0L}, v \rangle,$$

for any  $v \in V(\Omega_L)$  a. e. on (0,T) in the sense of H, here  $\langle \circ, \circ \rangle_{\Omega_L}$  is the dual form for the pair of spaces  $(V(\Omega_L), V^*(\Omega_L))$  and  $\Omega_L$  is Lipschitz, where  $x' \equiv (x_1, x_2)$ and

$$\mathcal{V}\left(Q_{L}^{T}\right) \equiv \left\{w \mid w \in V\left(Q_{L}^{T}\right), \ w' \in L^{2}\left(0, T; V^{*}\left(\Omega_{L}\right)\right)\right\}.$$

We will lead of the proof of this problem in five-steps as independent problem.

5.1. A priori estamations. For this we assume in (5.1)  $u_L$  instead of v then we get

(5.2) 
$$\frac{d}{dt} \langle u_L, u_L \rangle_{\Omega_L} - \langle \nu \Delta u_L, u_L \rangle_{\Omega_L} + \left\langle \sum_{j=1}^3 u_{Lj} D_j u_L, u_L \right\rangle_{\Omega_L} = \langle f_L, u_L \rangle_{\Omega_L}.$$

Thence, by making the known calculations and taking into account of the condition on  $\Omega_L$  and of calculations (4.1) and next (4.4) that carried out in previous Section, we obtain

$$\frac{1}{2} \frac{d}{dt} \|u_L\|^2_{(H(\Omega_L))^3}(t) + \nu \left(1 + a_1^{-2}\right) \|D_1 u_L\|^2_{(H(\Omega_L))^3}(t) +$$

$$(5.3)$$

$$\nu \left(1 + a_2^{-2}\right) \|D_2 u_L\|^2_{(H(\Omega_L))^3}(t) + 2\nu a_1^{-1} a_2^{-1} \langle D_1 u_L, D_2 u_L \rangle_{\Omega_L}(t) = \langle f_L, u_L \rangle_{\Omega_L},$$

where  $\langle g,h\rangle_{\Omega_L} = \sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} g_i h_i dx_1 dx_2$  for any  $g,h \in (H(\Omega_L))^3$ , or  $g \in (H^1(\Omega_L))^3$ 

and  $h \in (H^{-1}(\Omega_L))^3$ , respectively. We will show the correctness of (5.3), and to this end we shall prove the correctness of each term of this sum, separately.

So, using (4.3) we get

$$-\nu \left\langle \Delta u_{L}\left(t\right), u_{L}\left(t\right) \right\rangle_{\Omega_{L}} =$$

$$-\nu \sum_{i=1}^{3} \left\langle \left[ \left( 1 + a_{1}^{-2} \right) D_{1}^{2} + \left( 1 + a_{2}^{-2} \right) D_{2}^{2} + 2a_{1}^{-1}a_{2}^{-1}D_{1}D_{2} \right] u_{Li}, u_{Li} \right\rangle_{P_{x_{3}}\Omega_{L}} = \nu \sum_{i=1}^{3} \int_{P_{x_{3}}\Omega_{L}} \left[ \left( 1 + a_{1}^{-2} \right) \left( D_{1}u_{Li} \right)^{2} + \left( 1 + a_{2}^{-2} \right) \left( D_{2}u_{Li} \right)^{2} + 2a_{1}^{-1}a_{2}^{-$$

$$2a_1^{-1}a_2^{-1}D_1u_{Li}D_2u_{Li} dx_1 dx_2 \ge$$

thus is obtained the sum reducible in (5.3); if we estimate of the last adding in the above mentioned sum then we get

(5.4) 
$$\nu \left[ \left\| D_1 u_L \right\|_{(H(\Omega_L))^3}^2(t) + \left\| D_2 u_L \right\|_{(H(\Omega_L))^3}^2(t) \right].$$

Now consider the trilinear form from (5.2)

$$\left\langle \sum_{j=1}^{3} u_{Lj} D_j u_L, u_L \right\rangle_{\Omega_L} =$$

due to (4.3) we get

$$\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} [u_{L1}D_1u_{Li}u_{Li} + u_{L2}D_2u_{Li}u_{Li} + u_{L3}(-a_1^{-1}D_1 - a_2^{-1}D_2)u_{Li}u_{Li}] dx_1 dx_2 = \frac{1}{2} \sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}D_1(u_{Li})^2 + u_{L2}D_2(u_{Li})^2 + u_{L3}(-a_1^{-1}D_1 - a_2^{-1}D_2)(u_{Li})^2 \right] dx_1 dx_2 = -\frac{1}{2} \sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ D_1u_{L1} + D_2u_{L2} + \left(-a_1^{-1}D_1 - a_2^{-1}D_2\right)u_{L3} \right] (u_{Li})^2 dx_1 dx_2 = 0$$

hence by (4.4)

(5.5) 
$$-\frac{1}{2}\sum_{i=1}^{3}\int_{P_{x_3}\Omega_L} (u_{Li})^2 \operatorname{div} u_L dx_1 dx_2 = 0.$$

Consequently, the correctness of equation (5.3) follows from (5.4)-(5.5), that give we the following inequation

$$\frac{1}{2}\frac{d}{dt}\left\|u_{L}\right\|_{(H(\Omega_{L}))^{3}}^{2}(t)+$$

(5.6) 
$$\nu \sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ (D_1 u_{Li})^2 + (D_2 u_{Li})^2 \right] dx_1 dx_2 \le \int_{P_{x_3}\Omega_L} |(f_L \cdot u_L)| \, dx_1 dx_2$$

Namely, from here we obtain the following a priori estimations

(5.7) 
$$||u_L||_{(H(\Omega_L))^3}(t) \le C(f_L, u_{L0}, mes\Omega),$$

(5.8) 
$$\|D_1 u_L\|_{(H(\Omega_L))^3} + \|D_2 u_L\|_{(H(\Omega_L))^3} \le C(f_L, u_{L0}, mes\Omega),$$

where  $C(f_L, u_{L0}, mes\Omega) > 0$  is the constant that is independent of  $u_L$ . Consequently, any possible solution of this problem belong to a bounded subset of the space  $V(Q_L^T)$ .

So, if we will obtain the estimation for  $u'_L$  as well then we will have of the necessary a priori estimations, which are sufficient for the proof of the existence theorem.<sup>2</sup>

5.2. Boundedness of the trilinear form. Now we must study the trilinear form of (5.1) that one can also call as  $b_L(u_L, u_L, v)$ .

**Proposition 4.** Let  $u_L \in V(Q_L^T)$ ,  $v \in V(\Omega_L)$  and B is the operator defined by

$$\langle B(u_L), v \rangle_{\Omega_L} = b_L(u_L, u_L, v) = \left\langle \sum_{j=1}^3 u_{Lj} D_j u_L, v \right\rangle_{\Omega_L}$$

then  $B(u_L)$  belong to  $L^2(0,T; V^*(\Omega_L))$ .

*Proof.* At first we will show boundedness of the operator B from  $V(\Omega_L)$  to  $V^*(\Omega_L)$  for a. e.  $t \in (0, T)$ . We have

$$\langle B(u_L), v \rangle_{\Omega_L} = \left\langle \sum_{j=1}^3 u_{Lj} D_j u_L, v \right\rangle_{\Omega_L} =$$

due of (4.4) and of the definition 3

$$\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}D_1u_{Li}v_i + u_{L2}D_2u_{Li}v_i + u_{L3}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) u_{Li}v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}\left( -a_1^{-1}D_1 - a_2^{-1}D_2 \right) v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_2v_i + u_{L3}u_{Li}u_{Li} \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L2}u_{Li}D_1v_i + u_{L3}u_{Li}D_1v_i + u_{L3}u_{Li}u_{Li}D_1v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L3}u_{Li}D_1v_i + u_{L3}u_{Li}D_1v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L3}u_{Li}D_1v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} \left[ u_{L1}u_{Li}D_1v_i + u_{L3}u_{Li}D_1v_i \right] dx_1 dx_2 = -\sum_{i=1}^{3} \int_{P_$$

(5.9) 
$$-\sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} u_{Li} \left[ \left( u_{L1} - a_1^{-1} u_{L3} \right) D_1 v_i + \left( u_{L2} - a_2^{-1} u_{L3} \right) D_2 v_i \right] dx_1 dx_2.$$

Hence follows

$$|\langle B(u_L), v \rangle| \le \sum_{i=1}^{3} \int_{P_{x_3}\Omega_L} |u_L|^2 (|D_1v_i| + |D_2v_i|) dx_1 dx_2 \le$$

(5.10) 
$$c \|u_L\|_{L^4(\Omega_L)}^2 \|v\|_{V(\Omega_L)} \Longrightarrow \|B(u_L)\|_{V^*} \le c \|u_L\|_V^2$$

due of  $V(\Omega_L) \subset L^4(\Omega_L)$ . This also shows that operator  $B: V(\Omega_L) \longrightarrow V^*(\Omega_L)$  is continuous for a. e. t > 0.

<sup>&</sup>lt;sup>2</sup>It should be noted that if the representation of  $\Omega_L$  by coordinate system  $(x_1, x_2)$  not is best for the definition of the appropriate integral, then we will select other coordinate system: either  $(x_1, x_3)$  or  $(x_2, x_3)$  instead of  $(x_1, x_2)$  that is best for our goal, that must exist by virtue of the definition of  $\Omega$ .

Finally, we obtain needed result using above mentioned inequality and the wellknown inequality (see, [2], [7], [28]), which is correct for the space with two dimension

$$\int_{0}^{T} \|B(u_{L}(t))\|_{V^{*}}^{2} dt \leq c \int_{0}^{T} \|u_{L}(t)\|_{L^{4}}^{4} dt \leq c_{1} \int_{0}^{T} \|u_{L}(t)\|_{L^{2}}^{2} \|u_{L}\|_{V}^{2} dt \leq c_{1} \|u_{L}\|_{L^{\infty}(0,T;H)}^{2} \int_{0}^{T} \|u_{L}\|_{V}^{2} dt \Longrightarrow$$

(5.11) 
$$\|B(u_L)\|_{L^2(0,T;V^*)} \le c_1 \|u_L\|_{L^\infty(0,T;H)} \|u_L\|_{L^2(0,T;V)}.$$

What was to be proved.

5.3. Boundedness of u'. Sketch of the proof of the inclusion: u' belong to bounded subset of  $L^2(0,T; V^*(\Omega_L))$ . It is possible to draw the following conclusion based on receiving of a priori estimates, on proposition 4 and on reflexivity of all used spaces: If we were used of the Faedo-Galerkin's method for investigation we could obtain estimations for the approximate solutions the same as 5.7, 5.8 and 5.11. Since  $V(\Omega_L)$  is a separable there exists a sequence of linearly independent elements  $\{w_i\}_{i=1}^{\infty} \subset V(\Omega_L)$ , which is total in  $V(\Omega_L)$ . For each m we define an approximate solution  $u_{Lm}$  of (4.3) or (5.1) as follows:

(5.12) 
$$u_{Lm} = \sum_{i=1}^{m} u_{Lm}^{i}(t) w_{i}, \quad m = 1, 2, \dots$$

where  $u_{Lm}^{i}(t)$ ,  $i = \overline{1, m}$  be unknown functions that will be determined as solutions of following system of the differential equations that is received according to equation (5.1)

$$\left\langle \frac{d}{dt} u_{Lm}, w_j \right\rangle_{\Omega_L} = \left\langle \nu \Delta u_{Lm}, w_j \right\rangle_{\Omega_L} + b_L \left( u_{Lm}, u_{Lm}, w_j \right) + \dots$$

(5.13) 
$$\langle f_L, w_j \rangle_{\Omega_L}, \quad t \in (0,T], \quad j = \overline{1,m},$$

$$u_{Lm}\left(0\right) = u_{0Lm}$$

where  $\{u_{0Lm}\}_{m=1}^{\infty} \subset H(\Omega_L)$  is some sequence such that  $u_{0Lm} \longrightarrow u_{0L}$  in  $H(\Omega_L)$ as  $m \longrightarrow \infty$ . (Since  $V(\Omega_L)$  is everywhere dense in  $H(\Omega_L)$  we can determine  $u_{0Lm}$ by using the total system  $\{w_i\}_{i=1}^{\infty}$ ).

With use (5.12) in (5.13) we have

$$\sum_{j=1}^{m} \langle w_j, w_i \rangle_{\Omega_L} \frac{d}{dt} u_{Lm}^j(t) - \nu \sum_{j=1}^{m} \langle \Delta w_j, w_i \rangle_{\Omega_L} u_{Lm}^j(t) + \sum_{j,k=1}^{m} b_L(w_j, w_k, w_i) u_{Lm}^j(t) u_{Lm}^k(t) = \langle f_L(t), w_i \rangle_{\Omega_L}, \ i = \overline{1, m}$$

and taking into account the nonsingularity of matrix  $\langle w_i, w_j \rangle_{\Omega_L}$ ,  $i, j = \overline{1, m}$  we get to system of differential equations for  $u_{Lm}^i(t)$ , i = 1, ..., m

(5.14) 
$$\frac{du_{Lm}^{i}(t)}{dt} = \sum_{j=1}^{m} c_{i,j} \langle f_{L}(t), w_{j} \rangle_{\Omega_{L}} - \nu \sum_{j=1}^{m} d_{i,j} u_{Lm}^{j}(t) + \sum_{j,k=1}^{m} h_{ijk} u_{Lm}^{j}(t) u_{Lm}^{k}(t) ,$$

$$u_{Lm}^{i}(0) = u_{0Lm}^{i}, \quad i = 1, ..., m, \quad m = 1, 2, ...$$

here  $u_{0Lm}^i$  is  $i^{th}$  component of  $u_{0L}$  of the representation  $u_{0L} = \sum_{k=1}^{\infty} u_{0Lm}^k w_k$ .

The Cauchy problem for the nonlinear differential system (5.14) has solution, which is defined on the whole interval (0, T] by virtue of uniformity of estimations received in subsections 5.1 and 5.2. Consequently, the approximate solution  $u_{Lm}$ belong to a bounded subset of  $W^{1,2}(0,T;V^*(\Omega_L))$  for every m = 1, 2, ... since the right side of (5.14) belong to the bounded subset of  $L^2(0,T;V^*(\Omega_L))$  as were proved in subsections 5.1 and 5.2, by virtue of the lemma that is adduced below.

**Lemma 4.** ([28]) Let X be a given Banach space with dual  $X^*$  and let u and g be two functions belonging to  $L^1(a, b; X)$ . Then, the following three conditions are equivalent

(i) u is a. e. equal to a primitive function of g,

$$u(t) = \xi + \int_{a}^{t} g(s) ds, \quad \xi \in X, \quad a.e. \ t \in [a, b]$$

(ii) For each test function  $\varphi \in D((a, b))$ ,

$$\int_{a}^{b} u(t) \varphi'(t) dt = -\int_{a}^{b} g(t) \varphi(t) dt, \quad \varphi' = \frac{d\varphi}{dt}$$

(iii) For each  $\eta \in X^*$ ,

$$\frac{d}{dt}\left\langle u,\eta\right\rangle = \left\langle g,\eta\right\rangle$$

in the scalar distribution sense, on (a, b). If (i) - (iii) are satisfied u, in particular, is a. e. equal to a continuous function from [a, b] into X.

It not is difficult to see that if passing to limit at  $m \to \infty$  in equation (5.13) (maybe by a subsequence  $\{u_{Lm_l}\}_{l=1}^{\infty}$  as is known such subsequence exists) and to take  $\forall v \in V(\Omega_L)$  instead of  $w_k$  then we get

$$\left\langle \frac{d}{dt} u_L, v \right\rangle_{\Omega_L} = \left\langle f_L + \nu \Delta u_L - \chi, v \right\rangle_{\Omega_L},$$

as  $\{w_i\}_{i=1}^{\infty}$  is total in  $V(\Omega_L)$ , where  $\chi$  belong to  $L^2(0,T; V^*(\Omega_L))$  and is determined

$$\lim_{l \to \infty} \left\langle B\left(u_{Lm_l}\right), v \right\rangle_{\Omega_L} = \left\langle \chi, v \right\rangle_{\Omega_L}.$$

So, according to above a priori estimations and Proposition 4 we obtain that the right side belong to  $L^{2}(0,T)$  then the left side also belongs to  $L^{2}(0,T)$ , i. e.

$$\frac{du_{L}}{dt} \in L^{2}\left(0, T; V^{*}\left(\Omega_{L}\right)\right).$$

Consequently, the function  $u_L$  belong to a bounded subset of the space  $\mathcal{V}(Q_L^T)$ , where

(5.15) 
$$\mathcal{V}\left(Q_{L}^{T}\right) \equiv V\left(Q_{L}^{T}\right) \cap W^{1,2}\left(0,T;V^{*}\left(\Omega_{L}\right)\right),$$

by virtue of the above mentioned lemma and abstract form of Riesz-Fischer theorem.

Thus for the proof that  $u_l$  is the solution of equation (4.3) or (5.1) remains to prove that  $\chi = B(u_L)$  or  $\langle \chi, v \rangle_{\Omega_L} = b_L(u_L, u_L, v)$  for  $\forall v \in V(\Omega_L)$ .

## 5.4. Weakly compactness of operator B.

**Proposition 5.** Operator  $B : \mathcal{V}(Q_L^T) \longrightarrow L^2(0,T; V^*(\Omega_L))$  is weakly compact operator, *i. e.* any weakly convergent sequence  $\{u_L^m\}_1^\infty \subset \mathcal{V}(Q_L^T)$  possesses a subsequence  $\{u_L^{m_k}\}_1^\infty \subset \{u_L^m\}_1^\infty$  such that  $\{B(u_L^{m_k})\}_1^\infty$  weakly converge in  $L^2(0,T; V^*(\Omega_L))$ .

*Proof.* Let a sequence  $\{u_L^m\}_1^\infty \subset \mathcal{V}(Q_L^T)$  be weakly converge to  $u_L^0$  in  $\mathcal{V}(Q_L^T)$ . Then there exists a subsequence  $\{u_L^{m_k}\}_1^\infty \subset \{u_L^m\}_1^\infty$  such that  $u_L^{m_k} \longrightarrow u_L^0$  in  $L^2(0,T;H)$ according to known embedding theorems, i. e. since the inclusion

$$L^{2}(0,T;V(\Omega_{L})) \cap W^{1,2}(0,T;V^{*}(\Omega_{L})) \subset L^{2}(0,T;H)$$

is compact (see, e. g. [7], [28]). Indeed, for us it is enough to show that the operator generated by  $\sum_{j=1}^{3} u_{Lj} D_j u_L$  is weakly compact from  $\mathcal{V}\left(Q_L^T\right)$  to  $L^2\left(0,T; V^*\left(\Omega_L\right)\right)$ . From a priori estimations and Proposition 4 follows that operator  $B: \mathcal{V}\left(Q_L^T\right) \longrightarrow$  $L^2\left(0,T; V^*\left(\Omega_L\right)\right)$  is bounded, consequently  $B\left(\{u_L^{m_k}\}_1^\infty\right)$  belongs to bounded subset of the space  $L^2\left(0,T; V^*\left(\Omega_L\right)\right)$ . This lead the weak convergence

(5.16) 
$$B(u_L^{m_k}) \rightharpoonup \chi \quad \text{in } L^2(0,T;V^*(\Omega_L))$$

according by reflexivity of this space (at least, there exists such subsequence that this occurs).

Introduce the vector space

$$\mathcal{C}^{1}\left(\overline{Q}_{L}\right) \equiv \left\{ v \mid v_{i} \in C^{1}\left(\left[0, T\right]; C_{0}^{1}\left(\overline{\Omega_{L}}\right)\right), \ i = 1, 2, 3 \right\}$$

and consider the trilinear form

$$\int_{0}^{T} \left\langle B\left(u_{L}^{m}\right), v \right\rangle_{\Omega_{L}} dt = \int_{0}^{T} b\left(u_{L}^{m}, u_{L}^{m}, v\right) dt = \int_{0}^{T} \left\langle \sum_{j=1}^{3} u_{Lj}^{m} D_{j} u_{L}^{m}, v \right\rangle_{\Omega_{L}} dt =$$

where  $v \in \mathcal{C}^1(\overline{Q}_L)$ , then according to (5.9) we get

$$-\sum_{i=1}^{3} \int_{0}^{T} \int_{P_{x_{3}}\Omega_{L}} \left[ \left( u_{Li}^{m} u_{L1}^{m} - a_{1}^{-1} u_{Li}^{m} u_{L3}^{m} \right) D_{1} v_{i} + \left( u_{Li}^{m} u_{L2}^{m} - a_{2}^{-1} u_{Li}^{m} u_{L3}^{m} \right) D_{2} v_{i} \right] dx_{1} dx_{2} dt_{2} dt_{3} dx_{1} dx_{2} dt_{3} dx_{3} dx_{3} dx_{4} dx_{4} dx_{4} dx_{5} dx_$$

If in this sum separately we take arbitrary therm then it is not difficult to see that the following convergence is true

$$\int_{0}^{T} \int_{P_{x_3}\Omega_L} u_{Li}^m u_{L1}^m D_1 v_i dx_1 dx_2 dt \longrightarrow \int_{0}^{T} \int_{P_{x_3}\Omega_L} u_{Li} u_{L1} D_1 v_i dx_1 dx_2 dt$$

because  $u_{Li}^{m_k} \longrightarrow u_{Li}$  in  $L^2(0,T;H)$  and  $u_{Li}^{m_k} \rightharpoonup u_{Li}$  in  $L^2(0,T;H)$  (at least) since  $u_L^m$  belong, at least, to a bounded subset of  $\mathcal{V}(Q_L^T)$  and (5.16) holds for each therm. Thus we obtain

$$\chi = B\left(u_L\right) \Longrightarrow B\left(u_L^{m_k}\right) \rightharpoonup B\left(u_L\right) \quad \text{ in } L^2\left(0, T; V^*\left(\Omega_L\right)\right)$$

by using the density of  $\mathcal{C}^1(\overline{Q}_L)$  in  $\mathcal{V}(Q_L^T)$ .

Consequently, we proved the existence of the function  $u_L \in \mathcal{V}(Q_L^T)$  that satisfies equation (5.1) by applying to this problem of the Faedo-Galerkin method by virtue of the above mentioned results.

5.5. **Realisation of the initial condition.** The proof of the realisation of initial condition we can lead according to same way as in [28] (see, also [2], [7]), we will act in just the same way.

Let  $\phi$  be a continuously differentiable function on [0, T] with  $\phi(T) = 0$ . We multiply (5.13) by  $\phi(t)$ , and then integrate the first term by parts. This leads to the equation

$$-\int_{0}^{T} \left\langle u_{Lm}, \frac{d}{dt} \phi(t) w_{j} \right\rangle_{\Omega_{L}} dt = \int_{0}^{T} \left\langle \nu \Delta u_{Lm}, \phi(t) w_{j} \right\rangle_{\Omega_{L}} dt + \int_{0}^{T} b \left( u_{Lm}, u_{Lm}, \phi(t) w_{j} \right) dt + \int_{0}^{T} \left\langle f_{L}, \phi(t) w_{j} \right\rangle_{\Omega_{L}} dt + \left\langle u_{0Lm}, \phi(0) w_{j} \right\rangle_{\Omega_{L}} dt$$

We can pass to the limit with respect to subsequence  $\{u_{Lm_l}\}_{l=1}^{\infty}$  of the sequence  $\{u_{Lm}\}_{m=1}^{\infty}$  in all of terms by virtue of results which are proved in above subsections. Then we find the equation

$$(5.17) \qquad \int_{0}^{T} \left\langle u_{L}, \frac{d}{dt} \phi(t) w_{j} \right\rangle_{\Omega_{L}} dt = \int_{0}^{T} \left\langle \nu \Delta u_{L}, \phi(t) w_{j} \right\rangle_{\Omega_{L}} dt + \int_{0}^{T} b\left( u_{L}, u_{L}, \phi(t) w_{j} \right) dt + \int_{0}^{T} \left\langle f_{L}, \phi(t) w_{j} \right\rangle_{\Omega_{L}} dt + \left\langle u_{0L}, \phi(0) w_{j} \right\rangle_{\Omega_{L}} dt$$

that holds for each  $w_j$ , j = 1, 2, ... Consequently, this equation holds for any finite linear combination of the  $w_j$  and moreover because of a continuity (5.17) remains true for any  $v \in V(\Omega_L)$ .

Whence, one can draw conclusion that function  $u_L$  satisfies equation (5.1) in the distribution sense.

Now if we multiply (5.1) by  $\phi(t)$ , and integrate with respect to t after integrating the first term by parts, we get

$$-\int_{0}^{T} \left\langle u_{L}, v \frac{d}{dt} \phi(t) \right\rangle_{\Omega_{L}} dt - \int_{0}^{T} \left\langle \nu \Delta u_{L}, \phi(t) v \right\rangle_{\Omega_{L}} dt + \int_{0}^{T} \left\langle \sum_{j=1}^{3} u_{Lj} D_{j} u_{L}, \phi(t) v \right\rangle_{\Omega_{L}} dt = \int_{0}^{T} \left\langle f_{L}, \phi(t) v \right\rangle_{\Omega_{L}} dt + \left\langle u_{L}\left(0\right), \phi(0) v \right\rangle_{\Omega_{L}} dt$$

If we will compare this with (5.17) after replacing  $w_j$  with any  $v \in V(\Omega_L)$  then we obtain

$$\phi(0) \left\langle u_L(0) - u_{0L}, v \right\rangle_{\Omega_L} = 0.$$

Hence, we get the realisation of the initial condition by virtue of arbitrariness of  $v \in V(\Omega_L)$  and  $\phi$ , as one can choose  $\phi(0) \neq 0$ .

Consequently, the following result is proved.

**Theorem 5.** Under above mentioned conditions for any

$$u_{0L} \in (H(\Omega_L))^3$$
,  $f_L \in L^2(0,T; V^*(\Omega_L))$ 

problem (4.3) - (4.5) has weak solution  $u_L(t, x)$  that belongs to  $\mathcal{V}(Q_L^T)$ .

**Remark 1.** From the obtained a priori estimations and Propositions 4 and 5 follows of the fulfilment of all conditions of the general theorem of the compactness method (see, e. g. [31], [32], and for complementary informations see, [34], [33]). We would like to note also that the general theorems also were proved by using of the Faedo-Galerkin method and  $\varepsilon$ -regularization.

Here we would like to note we could prove the existence of problem (4.3) - (4.5) by other way with using of the following general existence theorem (see, [31], [32]) if in the adduced below theorem the pn-space to replace onto  $V(\Omega_L)$ .

Let X and Y be Banach spaces with duals  $X^*$  and  $Y^*$  respectively, Y be a reflexive Banach space,  $\mathcal{M}_0 \subseteq X$  be a weakly complete "reflexive" pn-space (see, Appendix A [32] or [31]),  $X_0 \subseteq \mathcal{M}_0 \cap Y$  be a separable vector topological space such that  $\overline{X_0}^{\mathcal{M}_0} \equiv \mathcal{M}_0, \overline{X_0}^Y \equiv Y$ . Consider the following problem:

(5.18) 
$$\frac{dx}{dt} + f(t, x(t)) = y(t), \quad y \in L^{p_1}(0, T; Y); \quad x(0) = 0$$

Let the following conditions be fulfilled:

i)  $f: \mathbf{P}_{0^{1,p_0,p_1}}(0,T;\mathcal{M}_0,Y) \to L^{p_1}(0,T;Y)$  is a weakly compact operator, where

$$\mathbf{P}_{0}^{1,p_{0},p_{1}}(0,T;\mathcal{M}_{0},Y) \equiv L^{p_{0}}(0,T;\mathcal{M}_{0}) \cap W^{1,p_{1}}(0,T;Y) \cap \{x(t) \mid x(0) = 0\},\$$

 $1 < \max\{p_1, p_1\} \le p_0 < \infty, \ p_1' = \frac{p_1}{p_1 - 1};$ 

(ii) there is a linear continuous operator  $L: W^{s,p_2}(0,T;X_0) \to W^{s,p_2}(0,T;Y^*)$ ,  $s \ge 0, p_2 \ge 1$  such that L commutes with  $\frac{d}{dt}$  and the conjugate operator  $L^*$  has  $ker(L^*) = \{0\}$ ;

(iii) there exist a continuous function  $\varphi : R^1_+ \cup \{0\} \longrightarrow R^1$  and numbers  $\tau_0 \ge 0$ and  $\tau_1 > 0$  such that  $\varphi(r)$  is nondecreasing for  $\tau \ge \tau_0$ ,  $\varphi(\tau_1) > 0$  and operators f

and L satisfy the following inequality for any  $x \in L^{p_0}(0,T;X_0)$ 

$$\int_{0}^{T} \langle f(t, x(t)), Lx(t) \rangle dt \ge \varphi\left( [x]_{L^{p_0}(\mathcal{M}_0)} \right) [x]_{L^{p_0}(\mathcal{M}_0)};$$

(iv) there exist a linear bounded operator  $L_0 : X_0 \to Y$  and constants  $C_0 > 0$ ,  $C_1, C_2 \ge 0, \nu > 1$  such that the inequalities

$$\int_{0}^{T} \langle \xi(t), L\xi(t) \rangle dt \geq C_{0} \|L_{0}\xi\|_{L^{p_{1}}(0,T;Y)}^{\nu} - C_{2}, \\
\int_{0}^{t} \langle \frac{dx}{d\tau}, Lx(\tau) \rangle d\tau \geq C_{1} \|L_{0}x\|_{Y}^{\nu}(t) - C_{2}, \quad a.e. \ t \in (0,T]$$

hold for any  $x \in W^{1,p_0}(0,T;X_0)$  and  $\xi \in L^{p_0}(0,T;X_0)$ .

**Theorem 6.** Assume that conditions (i) - (iv) are fulfilled. Then the Cauchy problem (5.18) is solvable in  $\mathbf{P}_{0,p_0,p_1}(0,T;\mathcal{M}_0,Y)$  in the following sense

$$\int_{0}^{T} \left\langle \frac{dx}{dt} + f(t, x(t)), y^{*}(t) \right\rangle dt = \int_{0}^{T} \left\langle y(t), y^{*}(t) \right\rangle dt, \quad \forall y^{*} \in L^{p_{1}'}(0, T; Y^{*}),$$

for any  $y \in G \subseteq L^{p_1}(0,T;Y)$ , where  $G \equiv \underset{r \geq \tau_1}{\cup} G_r$ :

$$G_{r} \equiv \left\{ y \in L^{p_{1}}(0,T;Y) \left| \int_{0}^{T} |\langle y(t), Lx(t) \rangle| \right| dt \leq \int_{0}^{T} \langle f(t,x(t)), Lx(t) \rangle dt - c, \\ \forall x \in L^{p_{0}}(0,T;X_{0}), \ [x]_{L^{p_{0}}(0,T;\mathcal{M}_{0})} = r \right\}, \ C_{2} < c < \infty.$$

The next proposition follows immediately from the theorem 6.

**Corollary 1.** Under assumptions of Theorem 6 the problem (5.18) is solvable in  $\mathbf{P}_{0^{1,p_0,p_1}}(0,T;\mathcal{M}_0,Y)$  for any  $y \in L^{p_1}(0,T;Y)$  satisfying the condition: there is r > 0 such that the inequality

$$\|y\|_{L^{p_1}(0,T;Y)} \le \varphi\left([x]_{L^{p_0}(0,T;\mathcal{M}_0)}\right)$$

holds for any  $x \in L^{p_0}(0,T;X_0)$  with  $[x]_{L^{p_0}(\mathcal{M}_0)} \geq r$ . Furthermore, if  $\varphi(\tau) \nearrow \infty$ as  $\tau \nearrow \infty$  then the problem (5.18) is solvable in  $\mathbf{P}_{0^{1,p_0,p_1}}(0,T;\mathcal{M}_0,Y)$  for any  $y \in L^{p_1}(0,T;Y)$  satisfying the inequality

$$\sup\left\{\frac{1}{[x]_{L^{p_{0}}\left(0,T;\mathcal{M}_{0}\right)}}\int_{0}^{T}\langle y\left(t\right),Lx\left(t\right)\rangle \ dt \ \mid x\in L^{p_{0}}\left(0,T;X_{0}\right)\right\}<\infty.$$

### KAMAL N. SOLTANOV

### 6. Uniqueness of Solution of Problem (3.3) - (3.5)

For the study of the uniqueness of the solution as usually: we will assume that the posed problem have two different solutions  $u = (u_1, u_2, u_3)$ ,  $v = (v_1, v_2, v_3)$ and we will investigate its difference: w = u - v. (Here for brevity we won't specify indexes for functions, which shows that we investigate problem (4.3) - (4.5) on  $Q_L^T$ .) Then for w we obtain the following problem

$$\begin{aligned} \frac{\partial w}{\partial t} &- \nu \left[ \left( 1 + a_1^{-2} \right) D_1^2 + \left( 1 + a_2^{-2} \right) D_2^2 \right] w - 2\nu a_1^{-1} a_2^{-1} D_1 D_2 w + \\ & \left( u_1 - a_1^{-1} u_3 \right) D_1 u - \left( v_1 - a_1^{-1} v_3 \right) D_1 v + \left( u_2 - a_2^{-1} u_3 \right) D_2 u - \\ & \left( v_2 - a_2^{-1} v_3 \right) D_2 v = 0, \\ & \text{div} \, w = D_1 \left[ \left( u - a_1^{-1} u_3 \right) - \left( v - a_1^{-1} v_3 \right) \right] + D_2 \left[ \left( u - a_2^{-1} u_3 \right) - \\ & \left( v - a_2^{-1} v_3 \right) \right] = D_1 w + D_2 w - \left( a_1^{-1} D_1 + a_2^{-1} D_2 \right) w_3 = 0, \end{aligned}$$
(6.1)

(6.2) 
$$w(0,x) = 0, \quad x \in \Omega \cap L; \quad w \mid_{(0,T) \times \partial \Omega_L} = 0.$$

Hence, we derive

$$\frac{1}{2}\frac{d}{dt} \|w\|_{2}^{2} + \nu \left[ \left(1 + a_{1}^{-2}\right) \|D_{1}w\|_{2}^{2} + \left(1 + a_{2}^{-2}\right) \|D_{2}w\|_{2}^{2} \right] + 2\nu a_{1}^{-1}a_{2}^{-1} \langle D_{1}w, D_{2}w \rangle_{\Omega_{L}} + \left\langle \left(u_{1} - a_{1}^{-1}u_{3}\right) D_{1}u - \left(v_{1} - a_{1}^{-1}v_{3}\right) D_{1}v, w \right\rangle_{\Omega_{L}} + \left\langle \left(u_{2} - a_{2}^{-1}u_{3}\right) D_{2}u - \left(v_{2} - a_{2}^{-1}v_{3}\right) D_{2}v, w \right\rangle_{\Omega_{L}} = 0$$

or

$$\frac{1}{2}\frac{d}{dt} \|w\|_{2}^{2} + \nu \left(\|D_{1}w\|_{2}^{2} + \|D_{2}w\|_{2}^{2}\right) + \nu \left[a_{1}^{-2} \|D_{1}w\|_{2}^{2} + a_{2}^{-2} \|D_{2}w\|_{2}^{2} + 2a_{1}^{-1}a_{2}^{-1} \langle D_{1}w, D_{2}w \rangle_{\Omega_{L}}\right] + \langle u_{1}D_{1}u - v_{1}D_{1}v, w \rangle_{\Omega_{L}} + \langle u_{2}D_{2}u - v_{2}D_{2}v, w \rangle_{\Omega_{L}} - (6.3) \qquad a_{1}^{-1} \langle u_{3}D_{1}u - v_{3}D_{1}v, w \rangle_{\Omega_{L}} - a_{2}^{-1} \langle u_{3}D_{2}u - v_{3}D_{2}v, w \rangle_{\Omega_{L}} = 0.$$

If we consider the last 4 added elements of left part (6.3), separately, and if we simplify these by calculations then we get

$$\begin{split} \langle w_{1}D_{1}u,w\rangle_{\Omega_{L}} + \langle v_{1}D_{1}w,w\rangle_{\Omega_{L}} + \langle w_{2}D_{2}u,w\rangle_{\Omega_{L}} + \langle v_{2}D_{2}w,w\rangle_{\Omega_{L}} - a_{1}^{-1} \langle w_{3}D_{1}u,w\rangle_{\Omega_{L}} - a_{1}^{-1} \langle v_{3}D_{1}w,w\rangle_{\Omega_{L}} - a_{2}^{-1} \langle w_{3}D_{2}u,w\rangle_{\Omega_{L}} - a_{2}^{-1} \langle v_{3}D_{2}w,w\rangle_{\Omega_{L}} = \\ \langle w_{1}D_{1}u,w\rangle_{\Omega_{L}} + \frac{1}{2} \langle v_{1},D_{1}w^{2}\rangle_{\Omega_{L}} + \langle w_{2}D_{2}u,w\rangle_{\Omega_{L}} + \frac{1}{2} \langle v_{2},D_{2}w^{2}\rangle_{\Omega_{L}} - \\ a_{1}^{-1} \langle w_{3}D_{1}u,w\rangle_{\Omega_{L}} - \frac{1}{2}a_{1}^{-1} \langle v_{3},D_{1}w^{2}\rangle_{\Omega_{L}} - a_{2}^{-1} \langle w_{3}D_{2}u,w\rangle_{\Omega_{L}} - \\ \frac{1}{2}a_{2}^{-1} \langle v_{3},D_{2}w^{2}\rangle = \frac{1}{2} \langle v_{1}-a_{1}^{-1}v_{3},D_{1}w^{2}\rangle_{\Omega_{L}} + \frac{1}{2} \langle v_{2}-a_{2}^{-1}v_{3},D_{2}w^{2}\rangle_{\Omega_{L}} + \\ \langle (w_{1}-a_{1}^{-1}w_{3})w,D_{1}u\rangle_{\Omega_{L}} + \langle (w_{2}-a_{2}^{-1}w_{3})w,D_{2}u\rangle_{\Omega_{L}} = \\ \langle (w_{1}-a_{1}^{-1}w_{3})w,D_{1}u\rangle_{\Omega_{L}} + \langle (w_{2}-a_{2}^{-1}w_{3})w,D_{2}u\rangle_{\Omega_{L}} . \end{split}$$

In the last equality we use the equation  $\operatorname{div} v = 0$  (see, (4.4)) and the condition (6.2).

If we take into account this equality in equation (6.3) then we get the equation

$$\frac{1}{2}\frac{d}{dt}\|w\|_{2}^{2} + \nu\left(\|D_{1}w\|_{2}^{2} + \|D_{2}w\|_{2}^{2}\right) + \nu\left[a_{1}^{-2}\|D_{1}w\|_{2}^{2} + \frac{1}{2}\frac{d}{dt}\|w\|_{2}^{2}\right]$$

ON UNIQUENESS OF SOLUTIONS OF NAVIER-STOKES EQUATIONS

$$a_{2}^{-2} \|D_{2}w\|_{2}^{2} + 2a_{1}^{-1}a_{2}^{-1} \langle D_{1}w, D_{2}w \rangle_{\Omega_{L}} \Big] + \left\langle \left(w_{1} - a_{1}^{-1}w_{3}\right)w, D_{1}u \right\rangle_{\Omega_{L}} + \left((6.4)\right) \left\langle \left(w_{2} - a_{2}^{-1}w_{3}\right)w, D_{2}u \right\rangle_{\Omega_{L}} = 0, \quad (t, x) \in (0, T) \times \Omega_{L}$$

Consequently, we derive the Cauchy problem for the equation (6.4) with the initial condition

$$\|w\|_2(0) = 0.$$

Hence giving rise to the differential inequality we get the following Cauchy problem for the differential inequality

(6.6) 
$$\frac{1}{2} \frac{d}{dt} \|w\|_{2}^{2} + \nu \left(\|D_{1}w\|_{2}^{2} + \|D_{2}w\|_{2}^{2}\right) \leq \left|\left\langle \left(w_{1} - a_{1}^{-1}w_{3}\right)w, D_{1}u\right\rangle_{\Omega_{L}}\right| + \left|\left\langle \left(w_{2} - a_{2}^{-1}w_{3}\right)w, D_{2}u\right\rangle_{\Omega_{L}}\right|$$

with the initial condition (6.5).

We have the following estimate for the right side of (6.6)

$$\left| \left\langle \left( w_1 - a_1^{-1} w_3 \right) w, D_1 u \right\rangle_{\Omega_L} \right| + \left| \left\langle \left( w_2 - a_2^{-1} w_3 \right) w, D_2 u \right\rangle_{\Omega_L} \right| \le \\ \left( \left\| w_1 - a_1^{-1} w_3 \right\|_4 + \left\| w_2 - a_2^{-1} w_3 \right\|_4 \right) \left\| w \right\|_4 \left\| \nabla u \right\|_2 \le$$

whence with the use of Gagliardo-Nirenberg-Sobolev inequality ([30]) we have

$$\left(1 + \max\left\{\left|a_{1}^{-1}\right|, \left|a_{2}^{-1}\right|\right\}\right) \|w\|_{4}^{2} \|\nabla u\|_{2} \leq c \|w\|_{2} \|\nabla w\|_{2} \|\nabla u\|_{2}.$$

It need to note that

$$(w_1 - a_1^{-1}w_3)w, (w_2 - a_2^{-1}w_3)w \in L^2(0,T; V^*(\Omega_L)),$$

by virtue of (5.15).

Now taking this into account in (6.6) one can arrive the following Cauchy problem for differential inequality

$$\begin{split} & \frac{1}{2} \frac{d}{dt} \left\| w \right\|_{2}^{2}(t) + \nu \left\| \nabla w \right\|_{2}^{2}(t) \leq c \left\| w \right\|_{2}(t) \left\| \nabla w \right\|_{2}(t) \left\| \nabla u \right\|_{2}(t) \leq c \left\| v \right\|_{2}(t) \left\| \nabla w \right\|_{2}^{2}(t) \left\| v \right\|_{2}^{2}(t) + \nu \left\| \nabla w \right\|_{2}^{2}(t), \quad \left\| w \right\|_{2}(0) = 0 \end{split}$$

since  $w \in L^{\infty}(0,T;(H(\Omega_L))^3)$ , and consequently,  $||w||_2 ||\nabla w||_2 \in L^2(0,T)$  by the virtue of the above existence theorem  $w \in \mathcal{V}(Q_L^T)$ , here  $C(c,\nu) > 0$  is constant.

Thus we obtain the problem

$$\frac{d}{dt} \|w\|_{2}^{2}(t) \leq 2C(c,\nu) \|\nabla u\|_{2}^{2}(t) \|w\|_{2}^{2}(t), \quad \|w\|_{2}(0) = 0,$$

if we denote  $\left\|w\right\|_{2}^{2}(t) \equiv y(t)$  then

$$\frac{d}{dt}y(t) \le 2C(c,\nu) \|\nabla u\|_{2}^{2}(t) y(t), \quad y(0) = 0.$$

Consequently, we obtain  $||w||_{2}^{2}(t) \equiv y(t) = 0$ , i.e. the following result is proved:

**Theorem 7.** Under above mentioned conditions for any

$$(f, u_0) \in L^2(0, T; V^*(\Omega_L)) \times (H(\Omega_L))^3$$

problem (4.3) - (4.5) has a unique weak solution u(t, x) that is contained in  $\mathcal{V}(Q_L^T)$ .

#### KAMAL N. SOLTANOV

#### 7. Proof of Theorem 2

*Proof.* (of Theorem 2). As is known ([1], [23], [9], [7]), problem  $(1.1^1)$  - (1.3) is solvable and possesses weak solution that is contained in the space  $\mathcal{V}(Q^T)$ . So, assume problem  $(1.1^1)$  - (1.3) has, at least, two different solutions under conditions of Theorem 2.

It is clear that if the problem have more than one solution then there is, at least, some subdomain of  $Q^T \equiv (0,T) \times \Omega$ , on which this problem has, at least, two solutions such, that each from the other are different. Consequently, starting from the above Lemma 3 we need to investigate the existence and uniqueness of the posed problem on arbitrary fixed subdomain on which it is possible that our problem can possess more than one solution, more exactly in the case when the subdomain is generated by an arbitrary fixed hyperplane by the virtue of Lemma 3. It is clear that, for us it is enough to prove that no such subdomain generated by a hyperplane on which more than single solutions of problem  $(1.1^1)$  - (1.3) exists, again by virtue of Lemma 3. In other words, for us it remains to use the above results (i.e. Theorems 5 and 7) in order to end the proof.

From the proved theorems above we obtain that there does not exist a subdomain, defined in the previous section, on which problem  $(1.1^1)$  - (1.3) reduced on this subdomain might possesses more than one weak solution. Consequently, taking Lemma 3 into account we obtain that the problem  $(1.1^1)$  - (1.3) (i.e.) under conditions of Theorem 2 possesses only one weak solution.  $\square$ 

So, under conditions of Theorem 2 the uniqueness of the weak solution u(x,t)(of velocity vector) of the problem obtained from the mixed problem for the incompressible Navier-Stokes equation by using approach of the Hopf-Leray in three dimension case is proved as noted in Notation 1.

Hence one can make the following conclusion

8. CONCLUSION

Let us

$$f \in L^{2}(0,T; V^{*}(\Omega)), \ u_{0} \in (H(\Omega))^{3}$$

It is well-known that the following inclusions are dense

$$L^{2}\left(0,T;H^{1/2}\left(\Omega\right)\right) \subset L^{2}\left(Q^{T}\right); \ \mathcal{H}^{1/2}\left(\Omega\right) \subset \left(H\left(\Omega\right)\right)^{3} \quad \&$$
$$L^{2}\left(0,T;\mathcal{H}^{1/2}\left(\Omega\right)\right) \subset L^{2}\left(0,T;\left(H^{-1}\left(\Omega\right)\right)^{3}\right)$$

consequently, there exist sequences

$$\left\{u_{0m}\right\}_{m=1}^{\infty} \subset \mathcal{H}^{1/2}\left(\Omega\right); \left\{f_{m}\right\}_{m=1}^{\infty} \subset L^{2}\left(0, T; \mathcal{H}^{1/2}\left(\Omega\right)\right)$$

such that  $u_{0m} \longrightarrow u_0$  in  $(H(\Omega))^3$  and  $||u_{0m}||_{(H(\Omega))^3} \leq ||u_0||_{(H(\Omega))^3}, f_m \longrightarrow f$  in  $L^{2}\left(0,T;\left(H^{-1}\left(\Omega\right)\right)^{3}\right) \text{ and } \|f_{m}\|_{L^{2}\left(0,T;\left(H^{-1}\left(\Omega\right)\right)^{3}\right)} \leq \|f\|_{L^{2}\left(0,T;\left(H^{-1}\left(\Omega\right)\right)^{3}\right)}.$ Consequently, for any  $\varepsilon > 0$  there exist  $m\left(\varepsilon\right) \geq 1$  such that under  $m \geq m\left(\varepsilon\right)$  for

the corresponding elements  $u_{0m}$ ,  $f_m$  of the above sequences

$$\|u_0 - u_{0m}\|_{(H(\Omega))^3} < \varepsilon; \ \|f - f_m\|_{L^2(0,T;(H^{-1}(\Omega))^3)} < \varepsilon$$

hold, and also the claim of Theorem 2 is valid for problem  $(1.1^1)$  - (1.3) with these datums.

One can note that the space that is everywhere dense subset of the necessary space possesses the minimal smoothness with respect to the needed space and also is sufficient for the application of our approach. So, we establish:

**Theorem 8.** Let  $\Omega$  be a Lipschitz open bounded set in  $\mathbb{R}^3$ . Then the existing by Theorem 1 weak solution  $u \in \mathcal{V}(Q^T)$  of the system  $(1.1^1) - (1.3)$  is unique, if the given functions f and  $u_0$  satisfy of conditions  $f \in L^2(0,T; \mathcal{H}^{1/2}(\Omega))$  and  $u_0 \in \mathcal{H}^{1/2}(\Omega)$ , where a solution be understood in the sense of Definition 2, as is well-known, spaces  $L^2(0,T; \mathcal{H}^{1/2}(\Omega))$  and  $\mathcal{H}^{1/2}(\Omega)$  are everywhere dense in spaces  $L^2(0,T; V^*(\Omega))$  and  $(H(\Omega))^3$ , respectively.

Acknowledgement 1. The author express the gratitude to russian mathematicians of Moscow for their giving useful suggestions and comments, which have helped to correct and sufficiently much to improve this paper.

### References

- J. Leray, Sur le Mouvement d'un Liquide Visquex Emplissent l'Espace, Act aMat h. J. 63 (1934), 193–248.
- [2] O. Ladyzhenskaya, The Mathematical Theory of Viscous Incompressible Flows (2nd edition), Gordon and Breach, (1969).
- [3] Majda, Andrew J.; Bertozzi, Andrea L. Vorticity and incompressible flow, Cambridge Texts in Applied Mathematics, 27, Cambridge University Press, Cambridge, (2002).
- [4] P. Constantin, Some open problems and research directions in the mathematical study of fluid dynamics, in Mathematics Unlimited–2001 and Beyond, (2001), Springer, Berlin, 353–360.
- [5] Charles L. Fefferman, Existence and smoothness of the Navier-Stokes Equation, Millennium Problems, Clay Math. Inst. (2006)
- [6] Galdi G.P., An introduction to the mathematical theory of the Navier-Stokes equations. Steady-state problems. Second edition. Springer Monographs in Mathematics. Springer, New York, 2011. xiv+1018 pp.
- [7] J.-L. Lions, Quelques methodes de resolution des problemes aux limites non lineaires, (1969), Dunod, Gauthier-Villars, Paris
- [8] J.-L. Lions, G. Prodi, Un theoreme d'existence et unicite dans les equations de Navier-Stokes en dimension 2, C. R. Acad. Sc. Paris, 248, (1959), 3519-3521.
- [9] O. Ladyzhenskaya, Ob odnoznachnoj razreshimosti v tselom trekhmernoj zadachi Cauchy dkja uravnenija Navier-Stokes pri nalichii osevoj simmetrii, Zap. nauch. sem. LOMI, 7, (1968), 155-177
- [10] V. Scheffer, Turbulence and Hausdorff dimension, in Turbulence and the Navier–Stokes Equations, Lecture Notes in Math. No. 565, Springer Verlag, 1976, pp. 94–112.
- [11] A. Shnirelman, On the nonuniqueness of weak solutions of the Euler equation, Comm. Pure & Appl. Math. 50 (1997), 1260–1286.
- [12] V. Scheffer, An inviscid flow with compact support in spacetime, J. Geom. Analysis 3, No. 4 (1993), 343–401.
- [13] A. V. Fursikov, Control problems and theorems concerning the unique solvability of a mixed boundary value problem for the three-dimensional Navier-Stokes and Euler equations, Math. USSR Sbornik, 43, 2 (1982), 251-273
- [14] L. Caffarelli, R. Kohn, L. Nirenberg, Partial regularity of suitable weak solutions of the Navier–Stokes equations, Comm. Pure & Appl. Math. 35 (1982), 771–831.
- [15] F.-H. Lin, A new proof of the Caffarelli-Kohn-Nirenberg theorem, Comm. Pure. & Appl. Math. 51 (1998), 241–257.
- [16] P. Constantin, I. Kukavica, V. Vicol, On the inviscid limit of the Navier-Stokes equations, Proc. AMS, 143, 7 (2015), 3075-3090
- [17] C. Foias, O. Manley, R. Rosa, R. Temam, Turbulence and Navier-Stokes equations, Cambridge University Press, (2001).

#### KAMAL N. SOLTANOV

- [18] C. Foias, R. Rosa and R.Temam, A note on statistical solutions of the three-dimensional Navier-Stokes equations: the time-dependent case, C. R. Acad. Sc. Paris, Ser. I, 348, (2010), 235-240.
- [19] C. Foias, R. Rosa, R. Temam, A note on statistical solutions of the three-dimensional Navier-Stokes equations: the stationary case, C. R. Acad. Sci. Paris, Ser. I, 348, (2010), 347-353
- [20] C. Foias, R. M. S. Rosa, R. M. Temam, Convergence of Time Averages of Weak Solutions of the Three-Dimensional Navier-Stokes Equations, J. Statistical Phys., 160, 3, (2015), 519-531
- [21] F. R. Gantmacher, The theory of matrices, v.I, Chelsea Publ. Comp., N.-Y. (1959)
- [22] N. Glatt-Holtz, V. Sverak, V. Vicol, On inviscid limits for the stochastic Navier-Stokes equations and related models, Archive for rational mech. and analysis, 217, 2, (2015), 619-649
- [23] E. Hopf, On nonlinear partial differential equations, Lecture Series of the Symposium on Partial Differential Equations, Berkeley, 1955, Ed. The Univ. of Kansas (1957), I-29.
- [24] X. Huang, Y. Wang, Global strong solution of 3D inhomogeneous Navier-Stokes equations with density-dependent viscosity, J. Diff. Eq., 259, 4, (2015), 1606-1627
- [25] Ch. Perrin, E. Zatorska, Free/Congested two-phase model from weak solutions to multidimensional compressible Navier-Stokes equations, Comm. in PDE, 40, 8, (2015), 1558-1589
- [26] K. N. Soltanov, Perturbation of the mapping and solvability theorems in the Banach space. Nonlinear Analysis: T.M.&A, 72, 1, (2010)
- [27] K. N. Soltanov, Some applications of the nonlinear analysis to the differential equations, Baku, ELM, (in Russian) (2002)
- [28] R. Temam, Navier-Stokes Equations Theory and Numerical Analysis, North-Holland Pub. Comp., in English, Third revised edition, (1984).
- [29] J.-L. Lions, E.Magenes, Problemes aux limites non homogenes et applications. Vol. 1, Dunod, Paris, (1968); English transl., Springer-Verlag, (1972).
- [30] O. V. Besov, V.P. Il'in, S. M. Nikol'skii, Integral'nye predstavleniya funktsii i teoremy vlozheniya (Integral Representations of Functions and Embedding Theorems), 2nd ed., rev. and compl., Moscow: Nauka, (1996).
- [31] K. N. Soltanov, On some modification of the Navier-Stokes equations. Nonlinear Analysis, T. M. & A., 52, 3, (2003), 769-793.
- [32] K. N. Soltanov, M. Ahmadov, Solvability of Equation of Prandtl-von Mises type, Theorems of Embedding. Transactions NAS of Azerbaijan, Ser. Phys.-Tech. and Math. Sci., 37, 1, (2017).
- [33] K. N. Soltanov, Some imbedding theorems and nonlinear differential equations. Trans. Ac. Sci. Azerb., ser. Math. & Phys.-Techn, 19, 5, (1999), 125 - 146.
- [34] K. N. Soltanov and J. Sprekels, Nonlinear equations in nonreflexive Banach spaces and strongly nonlinear equations, Adv. Math. Sci. Appl., 9, 2, (1999), 939-972.
- [35] A. I. Nachman Reconstructions from boundary measurements, Annals of Mathematics, 128 (1988), 531-576.
- [36] J. Behrndt, A.F.M. ter Elst, Dirichlet-to-Neumann maps on bounded Lipschitz domains, J. Diff. Eq., 259, (2015), 5903-5926
- [37] R. Haller-Dintelmann, J. Rehberg, Maximal parabolic regularity for divergence operators including mixed boundary conditions, J. Funct. Anal. 247, (2009), 1354–1396
- [38] M. Bellassoued, M. Choulli, Stability estimate for an inverse problem for the magnetic Schrödinger equation from the Dirichlet-to-Neumann map, J. Funct. Anal., 258, (2010), 161– 195
- [39] R. Denk, J. Prüss, R. Zacher, Maximal Lp-regularity of parabolic problems with boundary dynamics of relaxation type, J. Funct. Anal., 255 (2008), 3149–3187

INSTITUTE OF MATHEMATICS AND MECHANICS NATIONAL ACADEMY OF SCIENCES OF AZERBAIJAN, AZERBAIJAN;

E-mail address: sultan\_kamal@hotmail.com