

Negacyclic codes over the local ring $\mathbb{Z}_4[v]/\langle v^2 + 2v \rangle$ of oddly even length and their Gray images

Yuan Cao, Yonglin Cao*

*School of Mathematics and Statistics, Shandong University of Technology, Zibo,
Shandong 255091, China*

Abstract

Let $R = \mathbb{Z}_4[v]/\langle v^2 + 2v \rangle = \mathbb{Z}_4 + v\mathbb{Z}_4$ ($v^2 = 2v$) and n be an odd positive integer. Then R is a local non-principal ideal ring of 16 elements and there is a \mathbb{Z}_4 -linear Gray map from R onto \mathbb{Z}_4^2 which preserves Lee distance and orthogonality. First, a canonical form decomposition and the structure for any negacyclic code over R of length $2n$ are presented. From this decomposition, a complete classification of all these codes is obtained. Then the cardinality and the dual code for each of these codes are given, and self-dual negacyclic codes over R of length $2n$ are presented. Moreover, all $23 \cdot (4^p + 5 \cdot 2^p + 9)^{\frac{2^p-2}{p}}$ negacyclic codes over R of length $2M_p$ and all $3 \cdot (4^p + 5 \cdot 2^p + 9)^{\frac{2^{p-1}-1}{p}}$ self-dual codes among them are presented precisely, where $M_p = 2^p - 1$ is a Mersenne prime. Finally, 36 new and good self-dual 2-quasi-twisted linear codes over \mathbb{Z}_4 with basic parameters $(28, 2^{28}, d_L = 8, d_E = 12)$ and of type $2^{14}4^7$ and basic parameters $(28, 2^{28}, d_L = 6, d_E = 12)$ and of type $2^{16}4^6$ which are Gray images of self-dual negacyclic codes over R of length 14 are listed.

Keywords: Negacyclic code; Dual code; Self-dual code; Local ring; Finite chain ring

Mathematics Subject Classification (2000) 94B05, 94B15, 11T71

*corresponding author.

E-mail addresses: ylcao@sdut.edu.cn (Yonglin Cao), yuancao@sdut.edu.cn (Yuan Cao).

1. Introduction

The catalyst for the study of codes over rings was the discovery of the connection between the Kerdock and Preparata codes, which are non-linear binary codes, and linear codes over \mathbb{Z}_4 (see [3] and [4]). Soon after this discovery, codes over many different rings were studied. This led to many new discoveries and concretized the study of codes over rings as an important part of the coding theory discipline. Since \mathbb{Z}_4 is a chain ring, it was natural to expand the theory to focus on alphabets that are finite commutative chain rings and other special rings (See [1], [2], [5], [7–12], [14], [16–22], for examples).

In 1999, Wood in [23] showed that for certain reasons finite Frobenius rings are the most general class of rings that should be used for alphabets of codes. Then self-dual codes over commutative Frobenius rings were investigated by Dougherty et al. [13]. Especially, in 2014, codes over an extension ring of \mathbb{Z}_4 were studied in [24] and [25], here the ring was described as $\mathbb{Z}_4[u]/\langle u^2 \rangle = \mathbb{Z}_4 + u\mathbb{Z}_4$ ($u^2 = 0$) which is a local non-principal ring.

In this paper, all rings are associative and commutative. Let A be an arbitrary finite ring with identity $1 \neq 0$, A^\times the multiplicative group of units of A and $a \in A$. We denote by $\langle a \rangle_A$, or $\langle a \rangle$ for simplicity, the ideal of A generated by a , i.e. $\langle a \rangle_A = aA$. For any ideal I of A , we will identify the element $a + I$ of the residue class ring A/I with $a \pmod{I}$ in this paper.

For any positive integer N , let $A^N = \{(a_0, a_1, \dots, a_{N-1}) \mid a_i \in A, i = 0, 1, \dots, N-1\}$ which is an A -module with componentwise addition and scalar multiplication by elements of A . Then an A -submodule \mathcal{C} of A^N is called a *linear code* of length N over A . For any vectors $a = (a_0, a_1, \dots, a_{N-1})$, $b = (b_0, b_1, \dots, b_{N-1}) \in A^N$. The usual *Euclidian inner product* of a and b is defined by $[a, b] = \sum_{j=0}^{N-1} a_j b_j \in A$. Let \mathcal{C} be a linear code over A of length N . The *dual code* of \mathcal{C} is defined by $\mathcal{C}^\perp = \{a \in A^N \mid [a, b] = 0, \forall b \in \mathcal{C}\}$, and \mathcal{C} is said to be *self-dual* if $\mathcal{C} = \mathcal{C}^\perp$.

A linear code \mathcal{C} over A of length N is said to be *negacyclic* if

$$(-a_{N-1}, a_0, a_1, \dots, a_{N-2}) \in \mathcal{C}, \forall (a_0, a_1, \dots, a_{N-1}) \in \mathcal{C}.$$

We will use the natural connection of negacyclic codes to polynomial rings, where $c = (c_0, c_1, c_2, \dots, c_{N-1}) \in A^N$ is viewed as $c(x) = \sum_{j=0}^{N-1} c_j x^j$ and the negacyclic code \mathcal{C} is an ideal in the polynomial residue ring $A[x]/\langle x^N + 1 \rangle$.

In this paper, let n be an odd positive integer and denote

$$R = \mathbb{Z}_4[v]/\langle v^2 + 2v \rangle = \{a + bv \mid a, b \in \mathbb{Z}_4\} = \mathbb{Z}_4 + v\mathbb{Z}_4 \quad (v^2 = 2v)$$

in which the operations are defined by:

$$\alpha + \beta = (a + b) + v(c + d) \text{ and } \alpha\beta = ac + (ad + bc + 2bd)v.$$

for any $\alpha = a + bv, \beta = c + dv \in \mathbb{Z}_4 + v\mathbb{Z}_4$ with $a, b, c, d \in \mathbb{Z}_4$. Then R is a local Frobenius non-principal ideal ring of 16 elements.

Linear codes over R were studied in [15]. In the paper, a duality preserving Gray map was given and used to present MacWilliams identities and self-dual codes. Some extremal Type II \mathbb{Z}_4 -codes were provided as images of codes over this ring. \mathbb{Z}_4 -codes that are images of linear codes over R were characterised and some well-known families of \mathbb{Z}_4 -codes were proved to be linear over R . As in [15] Section 3, we define a map $\varrho : R \rightarrow \mathbb{Z}_4^2$ by

$$\varrho(\alpha) = (a + b, b), \forall \alpha = a + bv \in R \text{ where } a, b \in \mathbb{Z}_4$$

and let $\theta : R^N \rightarrow \mathbb{Z}_4^{2N}$ be such that $\theta(\alpha_1, \dots, \alpha_N) = (\varrho(\alpha_1), \dots, \varrho(\alpha_N))$, for all $\alpha_1, \dots, \alpha_N \in R$. Let w_L denote the Lee weight on \mathbb{Z}_4 defined by:

$$w_L(0) = 0, w_L(1) = w_L(3) = 1 \text{ and } w_L(2) = 2.$$

We extend w_L on the ring R in a natural way that

$$w_L(a + bv) = w_L(a + b) + w_L(b), \forall a, b \in \mathbb{Z}_4.$$

With this distance and Gray map definition, the following conclusions have been verified by Martínez-Moro et al. [15].

Lemma 1.1 ([15] Theorem 3.1) *Let \mathcal{C} be a linear code over R of length N and minimum Lee distance d . Then $\theta(\mathcal{C})$ is a linear code over \mathbb{Z}_4 of length $2N$, $|\theta(\mathcal{C})| = |\mathcal{C}|$ and is of minimum Lee distance d .*

Lemma 1.2 ([15] Proposition 3.3) *Let \mathcal{C} be a linear code over R of length N . Then $\theta(\mathcal{C}^\perp) = \theta(\mathcal{C})^\perp$. In particular, if \mathcal{C} is self-dual, then $\theta(\mathcal{C})$ is an self-dual code over \mathbb{Z}_4 of length $2N$ and has the same Lee weight distribution.*

Moreover, we have the following properties for Negacyclic codes R .

Proposition 1.3 *Let \mathcal{C} be a negacyclic code R of length N . Then $\theta(\mathcal{C})$ is a 2-quasi-twisted code over \mathbb{Z}_4 of length $2N$.*

Proof. Let $\underline{\alpha} = (\alpha_0, \alpha_1, \dots, \alpha_{N-1}) \in \mathcal{C}$, where $\alpha_i = a_i + b_i v$, $a_i, b_i \in \mathbb{Z}_4$, for all $i = 0, 1, \dots, N-1$. Then $\theta(\underline{\alpha}) = (a_0 + b_0, b_0, a_1 + b_1, b_1, \dots, a_{N-1} + b_{N-1}, b_{N-1})$

$b_{N-1}, b_{N-1}) \in \theta(\mathcal{C})$. As \mathcal{C} is negacyclic, we have $(-\alpha_{N-1}, \alpha_0, \alpha_1, \dots, \alpha_{N-2}) \in \mathcal{C}$. This implies $((-1)(a_{N-1}+b_{N-1}), -b_{N-1}, a_0+b_0, b_0, a_1+b_1, b_1, \dots, a_{N-2}+b_{N-2}, b_{N-2}) \in \theta(\mathcal{C})$. So $\theta(\mathcal{C})$ is a 2-quasi-twisted \mathbb{Z}_4 -code of length $2N$. \square

Since n is odd, the map $\varphi : R[x]/\langle x^n - 1 \rangle \rightarrow R[x]/\langle x^n + 1 \rangle$ defined by

$$\varphi(a(x)) = a(-x) = \sum_{j=0}^{n-1} a_j(-x)^j \quad (\forall a(x) = \sum_{j=0}^{n-1} a_j x^j \in R[x]/\langle x^n - 1 \rangle)$$

is an isomorphism of rings preserving Lee distance. Hence \mathcal{C} is a negacyclic code over R of n if and only if there is a unique cyclic code \mathcal{D} over R of length n such that $\varphi(\mathcal{D}) = \mathcal{C}$. Moreover, \mathcal{C} and \mathcal{D} has the same Lee weight distribution. A complete classification for cyclic codes over R of odd length and self-dual codes among them had been studied in [5]. In the paper, some good self-dual codes over \mathbb{Z}_4 of length 30 and extremal binary self-dual codes with parameters $[60, 30, 12]$ were obtained from self-dual cyclic codes over R of length 15. In this paper, we study negacyclic codes over R of length $2n$.

The present paper is organized as follows. In Section 2, we sketch the basic theory of finite rings and linear codes over finite rings needed in this paper. In Section 3, we decompose the ring $\mathbb{Z}_4[x]/\langle x^{2n} + 1 \rangle$ into a direct product of finite chain rings of length 4. In Section 4, we give a canonical form decomposition for any negacyclic code over R of length $2n$ and present all distinct codes by their generator sets. Using this decomposition, we give the number of codewords for each of these codes and an enumeration for all these codes. In Section 5, we present the dual code and its self-duality for each negacyclic code over R of length $2n$. In Section 6, we focus our attention on negacyclic code over R of length $2M_p$, where p is a prime and $M_p = 2^p - 1$ is a Mersenne prime. Especially, we present explicitly all 293687 negacyclic code over R of length 14 and 339 self-dual codes among them. Finally, we obtain 36 new and good self-dual 2-quasi-twisted codes over \mathbb{Z}_4 of length 28.

2. Preliminaries

In this section, we sketch the basic theory of finite chain rings and linear codes over finite chain rings needed in this paper.

Lemma 2.1 ([10] Proposition 2.1) *Let \mathcal{K} be a finite ring with identity. Then the following conditions are equivalent:*

- (i) \mathcal{K} is a local ring and the maximal ideal M of \mathcal{K} is principal, i.e. $M = \langle \pi \rangle$ for some $\pi \in \mathcal{K}$;
- (ii) \mathcal{K} is a local principal ideal ring;
- (iii) \mathcal{K} is a chain ring with all ideals given by: $\langle \pi^i \rangle = \pi^i \mathcal{K}$, $0 \leq i \leq s$, where s is the nilpotency of π .

Lemma 2.2 ([10] Proposition 2.2) *Let \mathcal{K} be a finite chain ring, with maximal ideal $M = \langle \pi \rangle$, and let s be the nilpotency of π . Then*

- (i) *For some prime p and positive integer m , $|\mathcal{K}/\langle \pi \rangle| = q$ where $q = p^m$, $|\mathcal{K}| = q^s$, and the characteristic of $\mathcal{K}/\langle \pi \rangle$ and \mathcal{K} are powers of p ;*
- (ii) *For $i = 0, 1, \dots, s$, $|\langle \pi^i \rangle| = q^{s-i}$.*

Lemma 2.3 ([16] Lemma 2.4) *Using the notations in Lemma 2.2, let $\mathcal{T} \subseteq \mathcal{K}$ be a system of representatives for the equivalence classes of \mathcal{K} under congruence modulo π . (Equivalently, we can define \mathcal{T} to be a maximal subset of \mathcal{K} with the property that $t_1 - t_2 \notin \langle \pi \rangle$ for all $t_1, t_2 \in \mathcal{T}$, $t_1 \neq t_2$.) Then*

- (i) *Every element a of \mathcal{K} has a unique π -expansion: $a = \sum_{j=0}^{s-1} t_j \pi^j$, $t_0, t_1, \dots, t_{s-1} \in \mathcal{T}$.*
- (ii) *$|\mathcal{K}/\langle \pi \rangle| = |\mathcal{T}|$ and $|\langle \pi^i \rangle| = |\mathcal{T}|^{s-i}$ for $0 \leq i \leq s$.*

From now on, let \mathcal{K} be an arbitrary finite chain ring with $1 \neq 0$, π be a fixed generator of the maximal ideal of \mathcal{K} with nilpotency index 4, and $F = \mathcal{K}/\langle \pi \rangle$. In this case, \mathcal{K} is called a *finite chain ring of length 4*. Using the notations of Lemma 2.3, every element $a \in \mathcal{K}$ has a unique π -adic expansion:

$$t_0 + \pi t_1 + \pi^2 t_2 + \pi^3 t_3, \quad t_0, t_1, t_2, t_3 \in \mathcal{T}.$$

Hence $|\mathcal{K}| = |F|^4$. If $a \neq 0$, the π -degree of a is defined as the least index $j \in \{0, 1, 2, 3\}$ for which $t_j \neq 0$ and written for $\|a\|_\pi = j$. If $a = 0$ we write $\|a\|_\pi = 4$. It is clear that $a \in \mathcal{K}^\times$ if and only if $t_0 \neq 0$, i.e. $\|a\|_\pi = 0$. Hence $|\mathcal{K}^\times| = (|F| - 1)|F|^3$. Moreover, we have $\mathcal{K}/\langle \pi^0 \rangle = \{0\}$ and $\mathcal{K}/\langle \pi^l \rangle = \{\sum_{i=0}^{l-1} \pi^i a_i \mid a_0, \dots, a_{l-1} \in \mathcal{T}\}$ with $|\mathcal{K}/\langle \pi^l \rangle| = |F|^l$, $1 \leq l \leq 3$.

Let L be a positive integer and $\mathcal{K}^L = \{(\alpha_1, \dots, \alpha_L) \mid \alpha_1, \dots, \alpha_L \in \mathcal{K}\}$ that is a free \mathcal{K} -module under componentwise addition and scalar multiplication with elements from \mathcal{K} . Then \mathcal{K} -submodules of \mathcal{K}^L are linear codes over \mathcal{K} of length L . Let C be a linear code over \mathcal{K} of length L . By [16] Definition 3.1, a matrix G is called a *generator matrix* for C if the rows of G span C and none of them can be written as a \mathcal{K} -linear combination of the other rows of

G . Furthermore, a generator matrix G is said to be *in standard form* if there is a suitable permutation matrix U of size $L \times L$ such that

$$G = \begin{pmatrix} \pi^0 I_{k_0} & M_{0,1} & M_{0,2} & M_{0,3} & M_{0,4} \\ 0 & \pi I_{k_1} & \pi M_{1,2} & \pi M_{1,3} & \pi M_{1,4} \\ 0 & 0 & \pi^2 I_{k_2} & \pi^2 M_{2,3} & \pi^2 M_{2,4} \\ 0 & 0 & 0 & \pi^3 I_{k_3} & \pi^3 M_{3,4} \end{pmatrix} U \quad (1)$$

where the columns are grouped into blocks of sizes k_0, k_1, k_2, k_3, k with $k_i \geq 0$ and $k = L - (k_0 + k_1 + k_2 + k_3)$. Of course, if $k_i = 0$, the matrices $\pi^i I_{k_i}$ and $\pi^i M_{i,j}$ ($i < j$) are suppressed in G . From [16] Proposition 3.2 and Theorem 3.5, we deduce the following.

Lemma 2.4 *Let C be a nonzero linear code of length L over \mathcal{K} . Then C has a generator matrix in standard form as in (1). In this case, the number of codewords in C is equal to $|C| = |F|^{4k_0+3k_1+2k_2+k_3} = |\mathcal{T}|^{4k_0+3k_1+2k_2+k_3}$.*

All distinct nontrivial linear codes of length 2 over \mathcal{K} has been listed (cf. Cao [6] Lemma 2.2 and Example 2.5). In particular, we have

Theorem 2.5 *Using the notations above, let $\omega \in \mathcal{K}^\times$. Then every linear code C over \mathcal{K} of length 2 satisfying the following condition*

$$(0, a + \omega\pi^2 b) \in C, \quad \forall (a, b) \in C \quad (2)$$

has one and only one of the following matrices G as its generator matrix in standard form:

- (I) $G = (\pi^2(a + b\pi), 1)$, where $a, b \in \mathcal{T}$.
- (II) $G = (0, \pi^3)$; $G = (\pi^3 b, \pi^2)$ where $b \in \mathcal{T}$; $G = (\pi^3 a, \pi)$ where $a \in \mathcal{T}$.
- (III) $G = \pi^k I_2$ where I_2 is the identity matrix of order 2, $0 \leq k \leq 4$.
- (IV) $G = \begin{pmatrix} 0 & 1 \\ \pi & 0 \end{pmatrix}$; $G = \begin{pmatrix} \pi^{t-1}z & 1 \\ \pi^t & 0 \end{pmatrix}$ where $z \in \mathcal{T}$ and $t = 2, 3$.
- (V) $G = \begin{pmatrix} \pi^2 z & \pi \\ \pi^3 & 0 \end{pmatrix}$ where $z \in \mathcal{T}$, and $G = \begin{pmatrix} 0 & \pi^{t-1} \\ \pi^t & 0 \end{pmatrix}$ where $t = 2, 3$.

Therefore, the number of linear codes over \mathcal{K} of length 2 satisfying Condition (2) is equal to $|\mathcal{T}|^2 + 5|\mathcal{T}| + 9$.

Proof. See Appendix. □

3. A direct sum decomposition of the ring $\mathbb{Z}_4[x]/\langle x^{2n} + 1 \rangle$

From now on, let n be an odd positive integer. In this section, we decompose the ring $\mathbb{Z}_4[x]/\langle x^{2n} + 1 \rangle$ into a direct product of finite chain rings of length 4. This direct sum decomposition will be needed in the following sections.

It is known that any element a of \mathbb{Z}_4 is unique expressed as $a = a_0 + 2a_1$ where $a_0, a_1 \in \mathbb{F}_2 = \{0, 1\}$ in which we regard \mathbb{F}_2 as a subset of \mathbb{Z}_4 . Denote $\bar{a} = a_0 \in \mathbb{F}_2$. Then $\bar{\cdot} : a \mapsto \bar{a}$ ($\forall a \in \mathbb{Z}_4$) is a ring isomorphism from \mathbb{Z}_4 onto \mathbb{F}_2 , and $\bar{\cdot}$ can be extended to a ring isomorphism from $\mathbb{Z}_4[x]$ onto $\mathbb{F}_2[x]$ by: $\bar{f}(x) = \overline{f(x)} = \sum_{i=0}^m \bar{b}_i x^i$ for all $f(x) = \sum_{i=0}^m b_i x^i \in \mathbb{Z}_4[x]$.

A monic polynomial $f(x) \in \mathbb{Z}_4[x]$ is said to be *basic irreducible* if $\bar{f}(x)$ is an irreducible polynomial in $\mathbb{F}_2[x]$. Then we have the following conclusions for monic basic irreducible polynomials in $\mathbb{Z}_4[x]$.

Lemma 3.1 *Let $f(x)$ be a monic basic irreducible polynomial in $\mathbb{Z}_4[x]$ of degree m and denote $\Gamma = \mathbb{Z}_4[x]/\langle f(x) \rangle$. Then*

(i) ([22] Theorem 6.1) Γ is a Galois ring of characteristic 4 and cardinality 4^m and $\Gamma = \mathbb{Z}_4[\zeta]$, where $\zeta = x + \langle f(x) \rangle \in \Gamma$ satisfying $\zeta^{2^m-1} = 1$ in Γ .

Denote $\bar{\Gamma} = \mathbb{F}_2[x]/\langle \bar{f}(x) \rangle$ and $\bar{\zeta} = x + \langle \bar{f}(x) \rangle \in \bar{\Gamma}$. Then $\bar{\Gamma} = \mathbb{F}_2[\bar{\zeta}]$ which is a finite field of cardinality 2^m , $\bar{f}(x) = \prod_{k=0}^{m-1} (x - \bar{\zeta}^{2^k})$ and that $\bar{\cdot}$ can be extended to a ring isomorphism from Γ onto $\bar{\Gamma}$ by $\xi \mapsto \bar{\xi} = \sum_{j=0}^{m-1} \bar{a}_j \bar{\zeta}^j$, for all $\xi = \sum_{j=0}^{m-1} a_j \zeta^j \in \Gamma$ where $a_0, a_1, \dots, a_{m-1} \in \mathbb{Z}_4$.

(ii) ([22] Proposition 6.14 or [7] Lemma 2.3(ii)) $f(x) = \prod_{k=0}^{m-1} (x - \zeta^{2^k})$.

Theorem 3.2 *Let $f(x)$ be a monic basic irreducible polynomial in $\mathbb{Z}_4[x]$ of degree m , denote $\mathcal{K}_f = \mathbb{Z}_4[x]/\langle f(-x^2) \rangle = \{\sum_{j=0}^{2m-1} a_j x^j \mid a_0, a_1, \dots, a_{2m-1} \in \mathbb{Z}_4\}$ in which the arithmetic is done modulo $f(-x^2)$, and set $\mathcal{T}_f = \{\sum_{j=0}^{m-1} b_j x^j \mid b_0, b_1, \dots, b_{m-1} \in \mathbb{F}_2\} \subseteq \mathcal{K}_f$ in which we regard \mathbb{F}_2 as a subset of \mathbb{Z}_4 . Then*

(i) *There is an invertible element $g(x) \in \mathcal{K}_f$ such that $f(x)^2 = 2g(x)$. Hence $\langle f(x)^2 \rangle = \langle 2 \rangle$ as ideals of \mathcal{K}_f .*

(ii) *\mathcal{K}_f is a finite chain ring with maximal ideal $\langle f(x) \rangle$ generated by $f(x)$, the nilpotency index of $f(x)$ is equal to 4 and $\mathcal{K}_f/\langle f(x) \rangle$ is a finite field of cardinality 2^m .*

(iii) *Every element α of \mathcal{K}_f has a unique $f(x)$ -adic expansion given by: $\alpha = \sum_{j=0}^3 b_j(x) f(x)^j$, $b_0(x), b_1(x), b_2(x), b_3(x) \in \mathcal{T}_f$.*

Moreover, we have $\mathcal{T}_f = \mathcal{K}_f/\langle f(x) \rangle$ as sets and $|\mathcal{T}_f| = 2^m$.

Proof. (i) By Lemma 3.1(ii), we have

$$f(-x^2) = \prod_{k=0}^{m-1} (-x^2 - \zeta^{2^k}) = (-1)^m \prod_{k=0}^{m-1} (x^2 + \zeta^{2^k}).$$

By Lemma 3.1(i), we know that $\zeta^{2^m} = \zeta = \zeta^{2^0}$. From this, by Lemma 3.1(ii) and $(x - \zeta^{2^k})^2 = (x^2 + \zeta^{2^{k+1}}) - 2\zeta^{2^k}x$ we deduce that

$$\begin{aligned} f(x)^2 &= \prod_{k=0}^{m-1} (x - \zeta^{2^k})^2 = \prod_{k=0}^{m-1} (x^2 + \zeta^{2^{k+1}}) - 2g(x) \\ &= (-1)^m f(-x^2) - 2g(x), \end{aligned}$$

where $g(x) = x \sum_{k=0}^{m-1} \zeta^{2^k} \prod_{0 \leq j \neq k \leq m-1} (x^2 + \zeta^{2^{j+1}}) \in \mathbb{Z}_4[x]$, since $g(x) = \frac{1}{2}((-1)^m f(-x^2) - f(x)^2) \in \mathbb{Z}[x]$, as a polynomial in $\mathbb{F}_2[x]$ we have that

$$\bar{g}(x) = x \sum_{k=0}^{m-1} \bar{\zeta}^{2^k} \prod_{0 \leq j \neq k \leq m-1} (x^2 + \bar{\zeta}^{2^{j+1}}) = x \sum_{k=0}^{m-1} \bar{\zeta}^{2^k} \prod_{0 \leq j \neq k \leq m-1} (x - \bar{\zeta}^{2^j})^2.$$

This implies $\bar{g}(\bar{\zeta}^{2^k}) = \bar{\zeta}^{2^{k+1}} \prod_{0 \leq j \neq k \leq m-1} (\bar{\zeta}^{2^k} - \bar{\zeta}^{2^j})^2 \neq 0$ for all $k = 0, 1, 2, \dots, m-1$, since $\bar{\zeta}, \bar{\zeta}^2, \dots, \bar{\zeta}^{2^{m-1}}$ are distinct root of $\bar{f}(x)$ in the finite field $\bar{\Gamma}$ by Lemma 3.1(i). From this and by $\overline{f(-x^2)} = \bar{f}(x^2) = (\bar{f}(x))^2 = \sum_{k=0}^{m-1} (x - \bar{\zeta}^{2^k})^2$, we deduce that $\gcd(\overline{f(-x^2)}, \bar{g}(x)) = 1$ in $\mathbb{F}_2[x]$. Therefore, $f(-x^2)$ and $g(x)$ are coprime in $\mathbb{Z}_4[x]$. Hence there exist $a(x), b(x) \in \mathbb{Z}_4[x]$ such that $a(x)g(x) + b(x)f(-x^2) = 1$. This implies that $g(x)$ is an invertible element of the residue class ring $\mathcal{K}_f = \mathbb{Z}_4[x]/\langle f(-x^2) \rangle$ and $g(x)^{-1} = a(x) \pmod{f(-x^2)}$. Then by $f(x)^2 = (-1)^d f(-x^2) - 2g(x)$ in $\mathbb{Z}_4[x]$ it follows that $f(x)^2 = 2g(x)$ in \mathcal{K}_f . Hence $\langle 2 \rangle = \langle f(x)^2 \rangle$ as ideals of \mathcal{R}_f .

(ii) Let $M = \langle 2, f(x) \rangle$ be the ideal of \mathcal{K}_f generated by 2 and $f(x)$. Then

$$\mathcal{K}_f/M = (\mathbb{F}_2[x]/\langle \overline{f(-x^2)} \rangle)/\langle \bar{f}(x) \rangle = (\mathbb{F}_2[x]/\langle \bar{f}(x)^2 \rangle)/\langle \bar{f}(x) \rangle = \mathbb{F}_2[x]/\langle \bar{f}(x) \rangle$$

up to natural ring isomorphisms, where $\mathbb{F}_2[x]/\langle \bar{f}(x) \rangle$ is a finite field of 2^m elements by Lemma 3.1(i). Hence M is a maximal ideal of \mathcal{K}_f .

By $f(x)^4 = (2g(x))^2 = 0$, we see that both 2 and $f(x)$ are nilpotent elements of \mathcal{K}_f . From this one can verify easily that every element of $\mathcal{K}_f \setminus M$ is invertible. This implies that \mathcal{R} is a local ring with M as its unique maximal

ideal. Furthermore, by $\langle 2 \rangle = \langle f(x)^2 \rangle$ in (i) we conclude that $M = \langle f(x) \rangle$. Hence $\mathcal{K}_f / \langle f(x) \rangle \cong \mathbb{F}_2[x] / \langle \bar{f}(x) \rangle$.

As stated above, by Lemma 2.1 we see that \mathcal{K}_f is a finite chain ring. Let s be the nilpotency index of $f(x)$. By Lemma 2.2(i) it follows that $|\mathcal{K}_f| = |\mathbb{F}_2[x] / \langle \bar{f}(x) \rangle|^s = 2^{sm}$. On the other hand, by $\deg(f(-x^2)) = 2m$ it follows that $|\mathcal{K}_f| = |\mathbb{Z}_4[x] / \langle f(-x^2) \rangle| = 4^{2m} = 2^{4m}$. Therefore, $s = 4$.

(iii) By $\mathcal{K}_f / \langle f(x) \rangle \cong \mathbb{F}_2[x] / \langle \bar{f}(x) \rangle = \{\sum_{j=0}^{m-1} b_j x^j \mid b_0, b_1, \dots, b_{m-1} \in \mathbb{F}_2\}$, we see that $\mathbb{F}_2[x] / \langle \bar{f}(x) \rangle = \mathcal{T}_f \subseteq \mathcal{K}_f$ as sets. Hence \mathcal{T}_f is a system of representatives for the equivalence classes of \mathcal{K}_f under congruence modulo $f(x)$. Then the conclusion follows from Lemma 2.3(i) immediately. \square

In the rest of this paper, let

$$x^n - 1 = f_1(x)f_2(x) \dots f_r(x), \quad (3)$$

where $f_1(x), f_2(x), \dots, f_r(x)$ are pairwise coprime monic basic irreducible polynomials in $\mathbb{Z}_4[x]$. We assume $\deg(f_i(x)) = m_i$ and denote

$$\mathcal{K}_i = \mathbb{Z}_4[x] / \langle f_i(-x^2) \rangle, \quad \mathcal{T}_i = \left\{ \sum_{j=0}^{m_i-1} b_j x^j \mid b_0, b_1, \dots, b_{m_i-1} \in \mathbb{F}_2 \right\} \subseteq \mathcal{K}_i,$$

for each integer i , $1 \leq i \leq r$. Then by Theorem 3.2, we know that

- There is an invertible element $g_i(x) \in \mathcal{K}_i$ such that

$$f_i(x)^2 = 2g_i(x) \text{ in } \mathcal{K}_i \quad (4)$$

where $g_i(x) = \frac{1}{2}((-1)^{m_i} f_i(-x^2) - f_i(x)^2)$ as a polynomial in $\mathbb{Z}[x] \pmod{f_i(-x^2), \text{ mod } 2}$. Hence $\langle f_i(x)^2 \rangle = \langle 2 \rangle$ as ideals of \mathcal{K}_i .

- \mathcal{K}_i is a finite chain ring with maximal ideal $\langle f_i(x) \rangle$, the nilpotency index of $f_i(x)$ is equal to 4 and $\mathcal{K}_i / \langle f_i(x) \rangle$ is a finite field of cardinality 2^{m_i} .

- Every element α of \mathcal{K}_i has a unique $f_i(x)$ -adic expansion: $\alpha = b_0(x) + b_1(x)f_i(x) + b_2(x)f_i(x)^2 + b_3(x)f_i(x)^3$, where $b_j(x) \in \mathcal{T}_i$ for all $j = 0, 1, 2, 3$. Moreover, we have $\mathcal{T}_i = \mathcal{K}_i / \langle f_i(x) \rangle$ as sets, $|\mathcal{T}_i| = 2^{m_i}$ and $|\mathcal{K}_i| = 4^{2m_i}$.

For each $1 \leq i \leq r$, denote $F_i(x) = \frac{x^n - 1}{f_i(x)} \in \mathbb{Z}_4[x]$. Since $F_i(x)$ and $f_i(x)$ are coprime, there are polynomials $a_i(x), b_i(x) \in \mathbb{Z}_4[x]$ such that

$$a_i(x)F_i(x) + b_i(x)f_i(x) = 1. \quad (5)$$

Substituting $-x^2$ for x in (3) and (5), we obtain

$$-(x^{2n} + 1) = f_1(-x^2)f_2(-x^2) \dots f_r(-x^2)$$

and $a_i(-x^2)F_i(-x^2) + b_i(-x^2)f_i(-x^2) = 1$ in the ring $\mathbb{Z}_4[x]$ respectively. In the rest of this paper, we set

$$\varepsilon_i(x) \equiv a_i(-x^2)F_i(-x^2) = 1 - b_i(-x^2)f_i(-x^2) \pmod{x^{2n} + 1}. \quad (6)$$

Then from classical ring theory, we deduce the following conclusions.

Theorem 3.3 *Denote $\mathcal{A} = \mathbb{Z}_4[x]/\langle x^{2n} + 1 \rangle$. We have the following:*

(i) $\varepsilon_1(x) + \dots + \varepsilon_r(x) = 1$, $\varepsilon_i(x)^2 = \varepsilon_i(x)$ and $\varepsilon_i(x)\varepsilon_j(x) = 0$ in the ring \mathcal{A} , for all $1 \leq i \neq j \leq r$.

(ii) $\mathcal{A} = \mathcal{A}_1 \oplus \dots \oplus \mathcal{A}_r$, where $\mathcal{A}_i = \varepsilon_i(x)\mathcal{A}$ and its multiplicative identity is $\varepsilon_i(x)$. Moreover, this decomposition is a direct sum of rings in that $\mathcal{A}_i\mathcal{A}_j = \{0\}$ for all integers i and j , $1 \leq i \neq j \leq r$.

(iii) For each $1 \leq i \leq r$, define a mapping $\phi_i : a(x) \mapsto \varepsilon_i(x)a(x)$ ($\forall a(x) \in \mathcal{K}_i = \mathbb{Z}_4[x]/\langle f_i(-x^2) \rangle$). Then ϕ_i is a ring isomorphism from \mathcal{K}_i onto \mathcal{A}_i . Hence $|\mathcal{A}_i| = 16^{m_i}$.

(iv) Define $\phi : (a_1(x), \dots, a_r(x)) \mapsto \phi_1(a_1(x)) + \dots + \phi_r(a_r(x))$, i.e.

$$\phi(a_1(x), \dots, a_r(x)) = \sum_{i=1}^r \varepsilon_i(x)a_i(x) \pmod{x^{2n} + 1},$$

for all $a_i(x) \in \mathcal{K}_i$, $i = 1, \dots, r$. Then ϕ is a ring isomorphism from $\mathcal{K}_1 \times \dots \times \mathcal{K}_r$ onto \mathcal{A} .

4. Structure of negacyclic codes over R of length $2n$

In this section, we list all distinct negacyclic codes of length $2n$ over the ring $R = \mathbb{Z}_4 + v\mathbb{Z}_4$ ($v^2 = 2v$), i.e. all distinct ideals of the ring $R[x]/\langle x^{2n} + 1 \rangle$. Using the notation of Theorem 3.3, we denote

$$\mathcal{A} = \mathbb{Z}_4[x]/\langle x^{2n} + 1 \rangle = \left\{ \sum_{j=0}^{2n-1} a_j x^j \mid a_j \in \mathbb{Z}_4, j = 0, 1, \dots, 2n-1 \right\}$$

in which the arithmetic is done modulo $x^{2n} + 1$, and set

$$\mathcal{A} + v\mathcal{A} = \mathcal{A}[v]/\langle v^2 + 2v \rangle = \{ \alpha + \beta v \mid \alpha, \beta \in \mathcal{A} \} \quad (v^2 = 2v)$$

in which the operations are defined by: for any $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathcal{A}$,

$$\begin{aligned}
(\alpha_1 + \beta_1 v) + (\alpha_2 + \beta_2 v) &= (\alpha_1 + \alpha_2) + v(\beta_1 + \beta_2), \\
(\alpha_1 + \beta_1 v)(\alpha_2 + \beta_2 v) &= \alpha_1 \alpha_2 + v(\alpha_1 \beta_2 + \beta_1 \alpha_2 + 2\beta_1 \beta_2).
\end{aligned}$$

Then $\mathcal{A} + v\mathcal{A}$ is a finite commutative ring containing \mathcal{A} as its subring.

Let $\alpha, \beta \in \mathcal{A}$. Then α and β can be uniquely expressed as $\alpha = \sum_{i=0}^{2n-1} a_i x^i$ and $\beta = \sum_{i=0}^{2n-1} b_i x^i$ respectively, where $a_i, b_i \in \mathbb{Z}_4$ for all $i = 0, 1, \dots, 2n-1$. Now, we define a map $\Xi : \mathcal{A} + v\mathcal{A} \rightarrow R[x]/\langle x^n - 1 \rangle$ by

$$\Xi : \alpha + \beta v \mapsto \sum_{i=0}^{2n-1} \xi_i x^i, \text{ where } \xi_i = a_i + b_i v \in R, \ i = 0, 1, \dots, 2n-1.$$

Then one can easily verify the following conclusion.

Lemma 4.1 *The map Ξ defined above is an isomorphism of rings from $\mathcal{A} + v\mathcal{A}$ onto $R[x]/\langle x^{2n} + 1 \rangle$.*

In the following, we will identify $\mathcal{A} + v\mathcal{A}$ with $R[x]/\langle x^{2n} - 1 \rangle$ under the ring isomorphism Ξ . Therefore, in order to determine all negacyclic codes over R of length $2n$, we only need to determine all ideals of the ring $\mathcal{A} + v\mathcal{A}$. To do this, we need to investigate the structure of the ring $\mathcal{A} + v\mathcal{A}$. In the rest of the paper, for each integer $1 \leq i \leq r$ we denote

$$\mathcal{K}_i + v\mathcal{K}_i = \mathcal{K}_i[v]/\langle v^2 + 2v \rangle = \{\alpha + \beta v \mid \alpha, \beta \in \mathcal{K}_i\}$$

in which the operations are defined by: $\xi_1 + \xi_2 = (\alpha_1 + \alpha_2) + v(\beta_1 + \beta_2)$ and $\xi_1 \xi_2 = \alpha_1 \alpha_2 + v(\alpha_1 \beta_2 + \beta_1 \alpha_2 + 2\beta_1 \beta_2)$, for any $\xi_1 = \alpha_1 + \beta_1 v$ and $\xi_2 = \alpha_2 + \beta_2 v$ with $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathcal{K}_i$.

We give the structure for any negacyclic code over R of length $2n$.

Theorem 4.2 *Using the notations above, we have the following conclusions.*

(i) Define $\Phi(\xi_1, \dots, \xi_r) = \sum_{i=1}^r \varepsilon_i(x) \xi_i \pmod{x^{2n} + 1}$ ($\forall \xi_i \in \mathcal{K}_i + v\mathcal{K}_i, i = 1, 2, \dots, r$). Then Φ is an isomorphism of rings from $(\mathcal{K}_1 + v\mathcal{K}_1) \times (\mathcal{K}_2 + v\mathcal{K}_2) \times \dots \times (\mathcal{K}_r + v\mathcal{K}_r)$ onto $\mathcal{A} + v\mathcal{A}$.

(ii) \mathcal{C} is a negacyclic code over R of length $2n$ if and only if for each integer $i, 1 \leq i \leq r$, there is a unique ideal C_i of the ring $\mathcal{K}_i + v\mathcal{K}_i$ such that

$$\mathcal{C} = \varepsilon_1(x)C_1 \oplus \varepsilon_2(x)C_2 \oplus \dots \oplus \varepsilon_r(x)C_r \pmod{x^{2n} + 1}$$

where $\varepsilon_i(x)C_i = \{\varepsilon_i(x)\alpha + v\varepsilon_i(x)\beta \mid \alpha + \beta v \in C_i, \alpha, \beta \in \mathcal{K}_i\} \subseteq \mathcal{A} + v\mathcal{A}$ for all $i = 1, 2, \dots, r$. Hence the number of codewords in \mathcal{C} is $\prod_{i=1}^r |C_i|$.

Proof (i) Let $\xi = (\xi_1, \xi_2, \dots, \xi_r) \in (\mathcal{K}_1 + v\mathcal{K}_1) \times (\mathcal{K}_1 + v\mathcal{K}_1) \times \dots \times (\mathcal{K}_r + v\mathcal{K}_r)$ where $\xi_i = \alpha_i + v\beta_i$ and $\alpha_i, \beta_i \in \mathcal{K}_i$ for all $1 \leq i \leq r$. By the definition ϕ defined in Theorem 3.3(iv), we have

$$\begin{aligned}\Phi(\xi) &= \sum_{i=1}^r \varepsilon_i(x) \xi_i = \sum_{i=1}^r \varepsilon_i(x) (\alpha_i + v\beta_i) = \sum_{i=1}^r \varepsilon_i(x) \alpha_i + v \sum_{i=1}^r \varepsilon_i(x) \beta_i \\ &= \phi(\alpha_1, \alpha_2, \dots, \alpha_r) + v\phi(\beta_1, \beta_2, \dots, \beta_r).\end{aligned}$$

By Theorem 3.3, we know that ϕ is a ring isomorphism from $\mathcal{K}_1 \times \mathcal{K}_2 \times \dots \times \mathcal{K}_r$ onto \mathcal{A} . Then for any $\eta = (\eta_1, \eta_2, \dots, \eta_r)$, where $\eta_i = \gamma_i + v\delta_i$ with $\gamma_i, \delta_i \in \mathcal{K}_i$ for all i , by $\Phi(\eta) = \phi(\gamma_1, \gamma_2, \dots, \gamma_r) + v\phi(\delta_1, \delta_2, \dots, \delta_r)$, $v^2 = 2v$ and direct calculations one can easily verify that $\Phi(\xi + \eta) = \Phi(\xi) + \Phi(\eta)$ and

$$\begin{aligned}\Phi(\xi\eta) &= \Phi(\xi_1\eta_1, \dots, \xi_r\eta_r) \\ &= \Phi(\alpha_1\gamma_1 + v(\alpha_1\delta_1 + \beta_1\gamma_1 + 2\beta_1\delta_1), \dots, \alpha_r\gamma_r + v(\alpha_r\delta_r + \beta_r\gamma_r + 2\beta_r\delta_r)) \\ &= \phi(\alpha_1\gamma_1, \dots, \alpha_r\gamma_r) + v\phi(\alpha_1\delta_1 + \beta_1\gamma_1 + 2\beta_1\delta_1, \dots, \alpha_r\delta_r + \beta_r\gamma_r + 2\beta_r\delta_r) \\ &= \phi(\alpha_1, \dots, \alpha_r)\phi(\gamma_1, \dots, \gamma_r) + v(\phi(\alpha_1, \dots, \alpha_r)\phi(\delta_1, \dots, \delta_r) \\ &\quad + \phi(\beta_1, \dots, \beta_r)\phi(\gamma_1, \dots, \gamma_r) + 2\phi(\beta_1, \dots, \beta_r)\phi(\delta_1, \dots, \delta_r)) \\ &= (\phi(\alpha_1, \dots, \alpha_r) + v\phi(\beta_1, \dots, \beta_r))(\phi(\gamma_1, \dots, \gamma_r) + v\phi(\delta_1, \dots, \delta_r)) \\ &= \Phi(\xi) \cdot \Phi(\eta).\end{aligned}$$

Hence Φ is a ring isomorphism from $(\mathcal{K}_1 + v\mathcal{K}_1) \times \dots \times (\mathcal{K}_r + v\mathcal{K}_r)$ onto $\mathcal{A} + v\mathcal{A}$.

(ii) From the properties of ring isomorphisms and direct product rings, by (i) we conclude that \mathcal{C} is a negacyclic code over R of length $2n$, i.e. \mathcal{C} is an ideal of $\mathcal{A} + v\mathcal{A}$, if and only if for each integer i , $1 \leq i \leq r$, there is a unique ideal C_i of the ring $\mathcal{K}_i + v\mathcal{K}_i$ such that

$$\begin{aligned}\mathcal{C} &= \Phi(C_1 \times C_2 \times \dots \times C_r) = \{\Phi(\xi_1, \xi_2, \dots, \xi_r) \mid \xi_i \in C_i, 1 \leq i \leq r\} \\ &= \left\{ \sum_{i=1}^r \varepsilon_i(x) \xi_i \mid \xi_i \in C_i, 1 \leq i \leq r \right\} = \sum_{i=1}^r \varepsilon_i(x) \{\xi_i \mid \xi_i \in C_i\}.\end{aligned}$$

Hence $\mathcal{C} = \bigoplus_{i=1}^r \varepsilon_i(x) C_i$ and $|\mathcal{C}| = |C_1 \times C_2 \times \dots \times C_r| = \prod_{i=1}^r |C_i|$. \square

Using the notations of Theorem 3.3, $\mathcal{C} = \bigoplus_{i=1}^r \varepsilon_i(x) C_i$ is called the *canonical form decomposition* of the negacyclic code \mathcal{C} over R of length $2n$.

Obviously, $\mathcal{K}_i + v\mathcal{K}_i$ is a free \mathcal{K}_i -module with basis $\{1, v\}$. Let $\mathcal{K}_i^2 = \{(\alpha, \beta) \mid \alpha, \beta \in \mathcal{K}_i\}$. Then \mathcal{K}_i^2 is a free \mathcal{K}_j -module of rank 2 with the componentwise addition and scalar multiplication. Now, define

$$\sigma : \mathcal{K}_i^2 \rightarrow \mathcal{K}_i + v\mathcal{K}_i \text{ via } (\alpha, \beta) \mapsto \alpha + \beta v \ (\forall \alpha, \beta \in \mathcal{K}_i).$$

Then σ is an isomorphism of \mathcal{K}_i -modules from \mathcal{K}_i^2 onto $\mathcal{K}_i + v\mathcal{K}_i$. Moreover, we have the following key conclusion.

Lemma 4.3 (cf. [5] Lemma 3.4) *Let $1 \leq i \leq r$. Then C_i is an ideal of the ring $\mathcal{K}_i + v\mathcal{K}_i$ if and only if there is a unique \mathcal{K}_i -submodule S_i of \mathcal{K}_i^2 satisfying*

$$(0, \alpha + 2\beta) \in S_i, \ \forall (\alpha, \beta) \in S_i \quad (7)$$

such that $\sigma(S_i) = C_i$.

For any ideal C_i of $\mathcal{K}_i + v\mathcal{K}_i$, its *annihilating ideal* is defined by $\text{Ann}(C_i) = \{\beta \in \mathcal{K}_i + v\mathcal{K}_i \mid \alpha\beta = 0, \ \forall \alpha \in C_i\}$. Now, we list all distinct ideals and their annihilating ideals of the ring $\mathcal{K}_i + v\mathcal{K}_i$ by the following theorem.

Theorem 4.4 *Let $0 \leq i \leq r$. Then all distinct ideals C_i and their annihilating ideals $\text{Ann}(C_i)$ of the ring $\mathcal{K}_i + v\mathcal{K}_i$ are given by the following table.*

| N | C_i | $ C_i $ | $\text{Ann}(C_i)$ |
|------------|--------------------------------------------|------------|------------------------------------------------|
| 2^{2m_i} | $\langle 2(a(x) + b(x)f_i(x)) + v \rangle$ | 4^{2m_i} | $\langle 2(1 + a(x) + b(x)f_i(x)) + v \rangle$ |
| 1 | $\langle 2vf_i(x) \rangle$ | 2^{m_i} | $\langle f_i(x), v \rangle$ |
| 2^{m_i} | $\langle 2(f_i(x)b(x) + v) \rangle$ | 2^{2m_i} | $\langle f_i(x)b(x) + v, 2 \rangle$ |
| 2^{m_i} | $\langle f_i(x)(2a(x) + v) \rangle$ | 2^{3m_j} | $\langle 2(1 + a(x)) + v, 2f_i(x) \rangle$ |
| 1 | $\langle 1 \rangle$ | 4^{4m_i} | $\langle 0 \rangle$ |
| 1 | $\langle f_i(x) \rangle$ | 4^{3m_i} | $\langle 2f_i(x) \rangle$ |
| 1 | $\langle 2 \rangle$ | 4^{2m_i} | $\langle 2 \rangle$ |
| 1 | $\langle 2f_i(x) \rangle$ | 4^{m_i} | $\langle f_i(x) \rangle$ |
| 1 | $\langle 0 \rangle$ | 1 | $\langle 1 \rangle$ |
| 1 | $\langle f_i(x), v \rangle$ | 2^{7m_i} | $\langle 2vf_i(x) \rangle$ |
| 2^{m_i} | $\langle f_i(x)b(x) + v, 2 \rangle$ | 2^{6m_j} | $\langle 2(f_i(x)b(x) + v) \rangle$ |
| 2^{m_i} | $\langle 2a(x) + v, 2f_i(x) \rangle$ | 2^{5m_i} | $\langle f_i(x)(2(1 + a(x)) + v) \rangle$ |
| 1 | $\langle 2, vf_i(x) \rangle$ | 2^{5m_i} | $\langle 2f_i(x), 2v \rangle$ |
| 1 | $\langle 2f_i(x), 2v \rangle$ | 2^{3m_i} | $\langle 2, vf_i(x) \rangle$ |
| 2^{m_i} | $\langle 2b(x) + vf_i(x), 2f_i(x) \rangle$ | 4^{2m_i} | $\langle 2b(x) + vf_i(x), 2f_i(x) \rangle$ |

where $a(x), b(x) \in \mathcal{T}_i$ and N is the number of ideals in the same row. Then the number of ideals in $\mathcal{K}_i + v\mathcal{K}_i$ is equal to $2^{2m_i} + 5 \cdot 2^{m_i} + 9$.

Proof Let C_i be an ideal of $\mathcal{K}_i + v\mathcal{K}_j$. By Lemma 4.3 there is a unique \mathcal{K}_i -submodule S of \mathcal{K}_i^2 satisfying condition (7) such that $C_i = \sigma(S)$.

By Equation (4) there is an invertible element $g_i(x) \in \mathcal{K}_j$ such that $f_i(x)^2 = 2g_i(x)$. Let $\omega_i(x) = g_i(x)^{-1} \in \mathcal{K}_j$. Then we have $2 = \omega_i(x)f_i(x)^2$. From this we deduce that S satisfying condition (7) if and only if S satisfies the following condition:

$$(0, \alpha + \omega_i(x)f_i(x)^2\beta) \in S_i, \quad \forall(\alpha, \beta) \in S.$$

Recall that \mathcal{K}_i -submodules S of \mathcal{K}_i^2 are called linear codes over \mathcal{K}_i of length 2. Since \mathcal{K}_i is a finite chain ring with maximal ideal $\langle f_i(x) \rangle$, the nilpotency index of $f_i(x)$ is equal to 4 and $\mathcal{T}_i = \mathcal{K}_i / \langle f_i(x) \rangle$ as sets with cardinality 2^{m_i} , by Theorem 2.5 we conclude that S has one of the following matrix G as its generator matrix in standard form:

(I) $G = (f_i(x)^2(a(x) + b(x)f_i(x)), 1)$, where $a(x), b(x) \in \mathcal{T}_i$. Since $g_i(x) \in \mathcal{K}_i^\times$, we have $2g_i(x) = 2\bar{g}_i(x)$ where $0 \neq \bar{g}_i(x) \equiv g_i(x) \pmod{2, \text{ mod } \bar{f}_i(x)}$. Hence $(a(x), b(x)) \mapsto (\bar{g}_i(x)a(x), \bar{g}_i(x)b(x)) \pmod{\bar{f}_i(x)}$ is a permutation on the set $\mathcal{T}_i \times \mathcal{T}_i$. Then by $f_i(x)^2 = 2g_i(x)$ we have

$$\begin{aligned} C_i &= \langle \sigma(f_i(x)^2(a(x) + b(x)f_i(x)), 1) \rangle = \langle f_i(x)^2(a(x) + b(x)f_i(x)) + v \rangle \\ &= \langle 2g_i(x)(a(x) + b(x)f_i(x)) + v \rangle = \langle 2(a'(x) + b'(x)f_i(x)) + v \rangle, \end{aligned}$$

where $a'(x) = \bar{g}_i(x)a(x), b'(x) = \bar{g}_i(x)b(x) \in \mathcal{T}_i$. Moreover, by Lemma 2.4 it follows that $|C_i| = |S| = |\mathcal{T}_i|^{4 \cdot 1} = (2^{m_i})^4 = 4^{2m_i}$. Obviously, the number of ideals is equal to $|\mathcal{T}_i \times \mathcal{T}_i| = |\mathcal{T}_i|^2 = 2^{2m_i}$ in this case.

(II) We have one of the following three subcases:

(II-1) $G = (0, f_i(x)^3)$. Then $C_i = \langle \sigma(0, f_i(x)^3) \rangle = \langle v f_i(x)^3 \rangle = \langle 2v f_i(x) \rangle$ by Equation (4). Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^1 = 2^{m_i}$.

(II-2) $G = (f_i(x)^3 b(x), f_i(x)^2)$ where $b(x) \in \mathcal{T}_i$. In this case, we have $C_i = \langle \sigma(f_i(x)^3 b(x), f_i(x)^2) \rangle = \langle f_i(x)^3 b(x) + v f_i(x)^2 \rangle = \langle 2(f_i(x)b(x) + v) \rangle$ by Equation (4). Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{2 \cdot 1} = 2^{2m_i}$.

(II-3) $G = (f_i(x)^3 a(x), f_i(x))$ where $a(x) \in \mathcal{T}_i$. By $g_i(x) \in \mathcal{K}_i^\times$, an argument similar to (I) shows that the map $a(x) \mapsto \bar{g}_i(x)a(x) \pmod{\bar{f}_i(x)}$ is a permutation on the set \mathcal{T}_i . Then by Equation (4) we have

$$\begin{aligned} C_i &= \langle \sigma(f_i(x)^3 a(x), f_i(x)^2) \rangle = \langle f_i(x)^3 a(x) + v f_i(x) \rangle \\ &= \langle 2f_i(x)g_i(x)a(x) + v f_i(x) \rangle = \langle 2f_i(x)a'(x) + v f_i(x) \rangle, \end{aligned}$$

where $a'(x) = \bar{g}_i(x)a(x) \in \mathcal{T}_i \pmod{\bar{f}_i(x)}$. Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{3 \cdot 1} = 2^{3m_i}$.

(III) $G = f_i(x)^k I_2$ where $0 \leq k \leq 4$. In this case, we have $C_i = \langle \sigma(f_i(x)^k, 0), \sigma(0, f_i(x)^k) \rangle = \langle f_i(x)^k, v f_i(x)^k \rangle = \langle f_i(x)^k \rangle$. Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{(4-k) \cdot 2} = 2^{2(4-k)m_i} = 4^{(4-k)m_i}$. Precisely, by Equation (4) it follows that $\langle f_i(x)^2 \rangle = \langle 2 \rangle$ and $\langle f_i(x)^3 \rangle = \langle 2f_i(x) \rangle$.

(IV) We have one of the following three subcases:

$$(IV-1) \ G = \begin{pmatrix} 0 & 1 \\ f_i(x) & 0 \end{pmatrix}. \text{ Then } C_i = \langle \sigma(0, 1), \sigma(f_i(x), 0) \rangle = \langle f_i(x), v \rangle.$$

Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{4 \cdot 1 + 3 \cdot 1} = 2^{7m_i}$.

(IV-2) $G = \begin{pmatrix} f_i(x)b(x) & 1 \\ f_i(x)^2 & 0 \end{pmatrix}$ where $b(x) \in \mathcal{T}_i$. In this case, we have $C_i = \langle \sigma(f_i(x)b(x), 1), \sigma(f_i(x)^2, 0) \rangle = \langle f_i(x)b(x) + v, f_i(x)^2 \rangle = \langle f_i(x)b(x) + v, 2 \rangle$ by Equation (4). Then by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{4 \cdot 1 + 2 \cdot 1} = 2^{6m_i}$.

$$(IV-3) \ G = \begin{pmatrix} f_i(x)^2 a(x) & 1 \\ f_i(x)^3 & 0 \end{pmatrix} \text{ where } a(x) \in \mathcal{T}_i. \text{ By Equation (4) we have}$$

$$\begin{aligned} C_i &= \langle \sigma(f_i(x)^2 a(x), 1), \sigma(f_i(x)^3, 0) \rangle = \langle f_i(x)^2 a(x) + v, f_i(x)^3 \rangle \\ &= \langle 2g_i(x)a(x) + v, 2f_i(x) \rangle = \langle 2a'(x) + v, 2f_i(x) \rangle, \end{aligned}$$

where $a'(x) = \bar{g}_i(x)a(x) \in \mathcal{T}_i \pmod{\bar{f}_i(x)}$. Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{4 \cdot 1 + 1 \cdot 1} = 2^{5m_i}$.

(V) We have one of the following three subcases:

(V-1) $G = \begin{pmatrix} 0 & f_i(x) \\ f_i(x)^2 & 0 \end{pmatrix}$. In this case, we have $C_i = \langle \sigma(0, f_i(x)), \sigma(f_i(x)^2, 0) \rangle = \langle v f_i(x), f_i(x)^2 \rangle = \langle 2, v f_i(x) \rangle$ by Equation (4). Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{3 \cdot 1 + 2 \cdot 1} = 2^{5m_i}$.

(V-2) $G = \begin{pmatrix} 0 & f_i(x)^2 \\ f_i(x)^3 & 0 \end{pmatrix}$. In this case, we have $C_i = \langle \sigma(0, f_i(x)^2), \sigma(f_i(x)^3, 0) \rangle = \langle v f_i(x)^2, f_i(x)^3 \rangle = \langle 2f_i(x), 2v \rangle$ by Equation (4). Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{2 \cdot 1 + 1 \cdot 1} = 2^{3m_i}$.

(V-3) $G = \begin{pmatrix} f_i(x)^2 b(x) & f_i(x) \\ f_i(x)^3 & 0 \end{pmatrix}$ where $b(x) \in \mathcal{T}_i$. Then by Equation (4) we have

$$\begin{aligned} C_i &= \langle \sigma(f_i(x)^2 b(x), f_i(x)), \sigma(f_i(x)^3, 0) \rangle = \langle f_i(x)^2 b(x) + v f_i(x), f_i(x)^3 \rangle \\ &= \langle 2g_i(x)b(x) + v f_i(x), 2f_i(x) \rangle = \langle 2b'(x) + v f_i(x), 2f_i(x) \rangle, \end{aligned}$$

where $b'(x) = \bar{g}_i(x)b(x) \in \mathcal{T}_i \pmod{\bar{f}_i(x)}$. Moreover, by Lemma 2.4 we have $|C_i| = |S| = |\mathcal{T}_i|^{3 \cdot 1 + 1 \cdot 1} = 2^{4m_i}$. By Theorem 2.5, we see that the number of ideals in $\mathcal{K}_i + v\mathcal{K}_i$ is equal to $|\mathcal{T}_i|^2 + 5|\mathcal{T}_i| + 9 = 2^{2m_i} + 5 \cdot 2^{m_i} + 9$.

Since $\mathcal{K}_i + v\mathcal{K}_i$ is a local ring with maximal ideal $\langle f_i(x), v \rangle$, one can deduce the conclusions for annihilating ideals from $v^2 = 2v$, $\omega_i(x)f_i(x)^2 = 2$ and a direct computation. \square

Finally, from Theorems 4.2 and 4.4 we deduce the following corollary.

Corollary 4.5 *Every negacyclic code over R of length $2n$ can be constructed by the following two steps:*

(i) *For each $i = 1, \dots, r$, choose an ideal C_i of $\mathcal{K}_i + v\mathcal{K}_i$ listed in Theorem 4.4.*

(ii) *Set $\mathcal{C} = \bigoplus_{i=1}^r \varepsilon_i(x)C_i \pmod{x^{2n} + 1}$.*

The number of codewords in \mathcal{C} is equal to $|\mathcal{C}| = \prod_{i=1}^r |C_i|$.

Moreover, the number of negacyclic codes over R of length $2n$ is equal to: $\prod_{i=1}^r (2^{2m_i} + 5 \cdot 2^{m_i} + 9)$.

5. Dual codes of negacyclic codes over R of length $2n$

In this section, we give the dual code of each negacyclic code over $R = \mathbb{Z}_4 + v\mathbb{Z}_4$ of length $2n$ and investigate the self-duality of these codes.

Let $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_{2n-1}), \beta = (\beta_0, \beta_1, \dots, \beta_{2n-1}) \in R^{2n}$, where $\alpha_j, \beta_j \in R$ for all $j = 0, 1, \dots, 2n-1$. As usual, we will identify the vector α with $\alpha(x) = \sum_{j=0}^{2n-1} \alpha_j x^j \in R[x]/\langle x^{2n} + 1 \rangle$ in this paper. Then we define

$$\mu(\alpha(x)) = \alpha(x^{-1}) = \alpha_0 - \sum_{j=1}^{2n-1} \alpha_j x^{2n-j} \in R[x]/\langle x^{2n} + 1 \rangle.$$

It is clear that μ is a ring automorphism of $R[x]/\langle x^{2n} + 1 \rangle$ satisfying $\mu^{-1} = \mu$. Now, by a direct calculation we get the following lemma.

Lemma 5.1 *Let $\alpha, \beta \in R^{2n}$. Then $[\alpha, \beta] = \sum_{j=0}^{2n-1} \alpha_j \beta_j = 0$ if $\alpha(x)\mu(\beta(x)) = 0$ in $R[x]/\langle x^{2n} + 1 \rangle$.*

Using the notations of Section 4, we have $R[x]/\langle x^{2n} + 1 \rangle = \mathcal{A} + v\mathcal{A}$ where $\mathcal{A} = \mathbb{Z}_4[x]/\langle x^{2n} + 1 \rangle$ and $v^2 = 2v$. It is obvious that the restriction of μ to \mathcal{A} is a ring automorphism of \mathcal{A} . We still denote this automorphism by μ ,

i.e. $\mu(a(x)) = a(x^{-1})$ for any $a(x) \in \mathcal{A}$. Let $1 \leq i \leq r$. By Equation (6) in Section 3, we have

$$\mu(\varepsilon_i(x)) = a_i(-x^{-2})F_i(-x^{-2}) = 1 - b_i(-x^{-2})f_i(-x^{-2}) \text{ in } \mathcal{A} \quad (8)$$

For any polynomial $f(x) = \sum_{j=0}^m c_j x^j \in \mathbb{Z}_4[x]$ of degree $m \geq 1$, recall that the *reciprocal polynomial* of $f(x)$ is defined as $\tilde{f}(x) = x^m f(\frac{1}{x}) = \sum_{j=0}^m c_j x^{m-j}$, and $f(x)$ is said to be *self-reciprocal* if $\tilde{f}(x) = f(x)$ or $-f(x)$. Then by Equation (3) in Section 3, we have

$$x^n - 1 = -(\widetilde{x^n - 1}) = -\tilde{f}_1(x)\tilde{f}_2(x)\dots\tilde{f}_r(x).$$

Since $f_1(x), f_2(x), \dots, f_r(x)$ are pairwise coprime monic basic irreducible polynomials in $\mathbb{Z}_4[x]$, $\tilde{f}_1(x), \tilde{f}_2(x), \dots, \tilde{f}_r(x)$ are pairwise coprime basic irreducible polynomials in $\mathbb{Z}_4[x]$ as well. Hence for each integer i , $1 \leq i \leq r$, there is a unique integer i' , $1 \leq i' \leq r$, such that $\tilde{f}_i(x) = \delta_i f_{i'}(x)$ where $\delta_i \in \{1, -1\}$. We assume that $\deg(f_i(x)) = d_i$ and $f_i(x) = \sum_{j=0}^{m_i} c_j x^j$ where $c_j \in \mathbb{Z}_4$. Then

$$\begin{aligned} x^{2m_i} f_i(-x^{-2}) &= (x^2)^{m_i} \sum_{j=0}^{m_i} (-1)^j c_j (x^2)^{-j} = (-1)^{m_i} \sum_{j=0}^{m_i} c_j (-x^2)^{m_i-j} \\ &= (-1)^{m_i} \tilde{f}_i(-x^2) = (-1)^{m_i} \delta_i f_{i'}(-x^2). \end{aligned}$$

From this, by Equation (6) and $x^{2n} = -1$ in the ring \mathcal{A} , we deduce that

$$\begin{aligned} \mu(\varepsilon_i(x)) &= 1 + x^{2n-2(\deg(b_i(x))+m_i)} (x^{2\deg(b_i(x))} b_i(-x^{-2})) (x^{2m_i} f_i(-x^{-2})) \\ &= 1 + (-1)^{\deg(b_i(x))+m_i} x^{2n-2(\deg(b_i(x))+m_i)} \tilde{b}_i(-x^2) \tilde{f}_i(-x^2) \\ &= 1 - h_i(x) f_{i'}(-x^2) \end{aligned}$$

where $h_i(x) = (-1)^{\deg(b_i(x))+m_i+1} \delta_i x^{2n-2(\deg(b_i(x))+m_i)} \tilde{b}_i(-x^2) \in \mathcal{A}$. Similarly, by (6) it follows that $\mu(\varepsilon_i(x)) = g_i(x) F_{i'}(-x^2)$ for some $g_i(x) \in \mathcal{A}$. Then from these and by (6), we deduce that $\mu(\varepsilon_i(x)) = \varepsilon_{i'}(x)$.

As stated above, we see that for each $1 \leq i \leq r$ there is a unique integer i' , $1 \leq i' \leq r$, such that $\mu(\varepsilon_i(x)) = \varepsilon_{i'}(x)$. We still use μ to denote this map $i \mapsto i'$. Then $\mu(\varepsilon_i(x)) = \varepsilon_{\mu(i)}(x)$.

Whether μ denotes the automorphism of \mathcal{A} or this map on the set $\{1, 2, \dots, r\}$ can be determined by the context. The next lemma shows the compatibility of the two uses of μ .

Lemma 5.2 *Using the notations above, we have the following conclusions.*

- (i) μ is a permutation on $\{1, \dots, r\}$ satisfying $\mu^{-1} = \mu$.
- (ii) After a rearrangement of $f_1(x), \dots, f_r(x)$ there are integers λ and ϵ such that $\mu(i) = i$ for all $i = 1, \dots, \lambda$ and $\mu(\lambda + j) = \lambda + \epsilon + j$ for all $j = 1, \dots, \epsilon$, where $\lambda \geq 1, \epsilon \geq 0$ and $\lambda + 2\epsilon = r$.
- (iii) For each integer i , $1 \leq i \leq r$, there is a unique element δ_i of $\{1, -1\}$ such that $\tilde{f}_i(x) = \delta_i f_{\mu(i)}(x)$.
- (iv) For any integer i , $1 \leq i \leq r$, $\mu(\varepsilon_i(x)) = \varepsilon_{\mu(i)}(x)$ and $\mu(\mathcal{A}_i) = \mathcal{A}_{\mu(i)}$ in the ring \mathcal{A} .
- (v) For any $c(x) \in \mathcal{K}_i = \mathbb{Z}_4[x]/\langle f_i(-x^2) \rangle$, define

$$\mu_i(c(x)) = c(x^{-1}) = c(-x^{2n-1}) \pmod{f_{\mu(i)}(-x^2)}.$$

Then $\mu_i = \phi_{\mu(i)}^{-1} \mu \phi_i$ which is a ring isomorphism from \mathcal{K}_i onto $\mathcal{K}_{\mu(i)}$.

- (vi) The ring isomorphism $\mu_i : \mathcal{K}_i \rightarrow \mathcal{K}_{\mu(i)}$ induces a ring isomorphism from $\mathcal{K}_i + v\mathcal{K}_i$ onto $\mathcal{K}_{\mu(i)} + v\mathcal{K}_{\mu(i)}$ ($v^2 = 2v$) defined by

$$\alpha + v\beta \mapsto \mu_i(\alpha) + v\mu_i(\beta), \quad \forall \alpha, \beta \in \mathcal{K}_i.$$

We still denote this isomorphism by μ_i . Then $\mu_i^{-1} = \mu_{\mu(i)}$.

Proof. (i)–(iii) follow from the definition of the map μ .

(iv) By $\mu(\varepsilon_i(x)) = \varepsilon_{\mu(i)}(x)$ and $\mathcal{A}_i = \varepsilon_i(x)\mathcal{A}$, it follows that $\mu(\mathcal{A}_i) = \mu(\varepsilon_i(x))\mu(\mathcal{A}) = \varepsilon_{\mu(i)}(x)\mathcal{A} = \mathcal{A}_{\mu(i)}$.

(v) Since $x^{2n} = -1$ in the ring \mathcal{A} , i.e. $x^{2n} \equiv -1 \pmod{x^{2n} + 1}$, and $f_i(-x^2)$ is a factor of $x^{2n} + 1$, it follows that $x^{2n} \equiv -1 \pmod{f_i(-x^2)}$. This implies $x^{-1} = -x^{2n-1}$ in the ring \mathcal{K}_i for all i , $1 \leq i \leq r$. Now, let $c(x) \in \mathcal{K}_i$. By Theorem 3.3(iii) and $\varepsilon_{\mu(i)}(x) = \mu(\varepsilon_i(x)) = 1 - h_i(x)f_{i'}(-x^2) = 1 - h_i(x)f_{\mu(i)}(-x^2)$, we have

$$\begin{aligned} (\phi_{\mu(i)}^{-1} \mu \phi_i)(c(x)) &= (\phi_{\mu(i)}^{-1} \mu)(\varepsilon_i(x)c(x)) = \phi_{\mu(i)}^{-1}(\mu(\varepsilon_i(x))c(x^{-1})) \\ &= (1 - h_i(x)f_{\mu(i)}(-x^2))c(x^{-1}) \pmod{f_{\mu(i)}(-x^2)} \\ &\equiv c(x^{-1}) \pmod{f_{\mu(i)}(-x^2)}. \end{aligned}$$

This implies $\mu_i(c(x)) = (\phi_{\mu(i)}^{-1} \mu \phi_i)(c(x))$ for all $c(x) \in \mathcal{K}_i$. Hence $\mu_i = \phi_{\mu(i)}^{-1} \mu \phi_i$. Since μ is a ring automorphism of \mathcal{A} , by Theorem 3.3(iii) we conclude that μ_i is a ring isomorphism from \mathcal{K}_i onto $\mathcal{K}_{\mu(i)}$.

(vi) Obviously, μ_i can be extended to a ring isomorphism from $\mathcal{K}_i[v]$ onto $\mathcal{K}_{\mu(i)}[v]$ in the natural way that $\mu_i : \sum_j \alpha_j v^j \mapsto \sum_j \mu_i(\alpha_j) v^j$, $\forall \alpha_j \in \mathcal{K}_i$. Therefore, $\alpha + v\beta \mapsto \mu_i(\alpha) + v\mu_i(\beta)$ ($\forall \alpha, \beta \in \mathcal{K}_i$) is a ring isomorphism from $\mathcal{K}_i[v]/\langle v^2 + 2v \rangle = \mathcal{K}_i + v\mathcal{K}_i$ onto $\mathcal{K}_{\mu(i)}[v]/\langle v^2 + 2v \rangle = \mathcal{K}_{\mu(i)} + v\mathcal{K}_{\mu(i)}$. \square

By Theorem 4.2(i), each element $\xi \in \mathcal{A} + v\mathcal{A}$ can be uniquely expressed as: $\xi = \sum_{i=1}^r \phi_i(\xi_i) = \sum_{i=1}^r \varepsilon_i(x) \xi_i$, where $\xi_i \in \mathcal{K}_i + v\mathcal{K}_i$ for all $i = 1, \dots, r$.

Lemma 5.3 *Let $\xi = \sum_{i=1}^r \varepsilon_i(x) \xi_i$, $\eta = \sum_{i=1}^r \varepsilon_i(x) \eta_i \in \mathcal{A} + v\mathcal{A}$, where $\xi_i, \eta_i \in \mathcal{K}_i + v\mathcal{K}_i$. Then $\xi \cdot \mu(\eta) = \sum_{i=1}^r \varepsilon_i(x) (\xi_i \cdot \mu_i^{-1}(\eta_{\mu(i)}))$.*

Proof. By Lemma 5.2(v) we have $\mu_i^{-1}(\eta_{\mu(i)}) \in \mu_i^{-1}(\mathcal{K}_{\mu(i)} + v\mathcal{K}_{\mu(i)}) = \mathcal{K}_i + v\mathcal{K}_i$. Hence $\xi_i \cdot \mu_i^{-1}(\eta_{\mu(i)}) \in \mathcal{K}_i + v\mathcal{K}_i$ for all i . If $j \neq \mu(i)$, then $i \neq \mu(j)$, which implies $\varepsilon_i(x) \varepsilon_{\mu(j)}(x) = 0$ by Theorem 3.3(i). Therefore,

$$\begin{aligned} \xi \cdot \mu(\eta) &= \sum_{i,j=1}^r \varepsilon_i(x) \xi_i \cdot \mu(\varepsilon_j(x) \eta_j) = \sum_{i,j=1}^r \varepsilon_i(x) \xi_i \cdot \mu(\varepsilon_j(x)) \mu_j(\eta_j) \\ &= \sum_{i,j=1}^r \varepsilon_i(x) \xi_i \cdot \varepsilon_{\mu(j)}(x) \mu_j(\eta_j) = \sum_{i=1}^r \varepsilon_i(x) \xi_i \cdot \varepsilon_i(x) \mu_{\mu(i)}(\eta_{\mu(i)}). \end{aligned}$$

This implies $\xi \cdot \mu(\eta) = \sum_{i=1}^r \varepsilon_i(x) (\xi_i \cdot \mu_i^{-1}(\eta_{\mu(i)}))$ by $\mu_{\mu(i)} = \mu_i^{-1}$. \square

Lemma 5.4 *Let $1 \leq i \leq r$. Then $\mu_i(f_i(x)) = -\delta_i x^{2n-m_i} f_{\mu(i)}(x)$.*

Proof. Since $f_i(-x^2) | (x^{2n} + 1)$ in $\mathbb{Z}_4[x]$, we have $x^{2n} = -1$ in \mathcal{K}_i . By the definition of μ_i and $\tilde{f}_i(x) = \delta_i f_{\mu(i)}(x)$ in Lemma 5.2 (v) and (iii), we have that $\mu_i(f_i(x)) = f_i(x^{-1}) = -x^{2n-m_i} (x^{m_i} f_i(x^{-1})) = -x^{2n-m_i} \tilde{f}_i(x) = -\delta_i x^{2n-m_i} f_{\mu(i)}(x)$. \square

Now, we can give the dual code of each negacyclic code over the ring $R = \mathbb{Z}_4 + v\mathbb{Z}_4$ of length $2n$ by the following theorem, where $h(x^{-1}) = h(-x^{2n-1}) \pmod{f_{\mu(i)}(-x^2)}$ for any $h(x) \in \mathcal{K}_i$.

Theorem 5.5 *Let \mathcal{C} be a negacyclic code over R of length $2n$ with $\mathcal{C} = \bigoplus_{i=1}^r \varepsilon_i(x) C_i$, where C_i is an ideal of $\mathcal{K}_i + v\mathcal{K}_i$. Then the dual code \mathcal{C}^\perp is also a negacyclic code over R of length $2n$. Precisely, we have*

$$\mathcal{C}^\perp = \bigoplus_{j=1}^r \varepsilon_j(x) D_j,$$

where D_j is an ideal of $\mathcal{K}_j + v\mathcal{K}_j$ determined by the following table:

| $C_i \pmod{f_i(-x^2)}$ | $D_{\mu(i)} \pmod{f_{\mu(i)}(-x^2)}$ |
|--------------------------------------------|--------------------------------------------------------------------------------------|
| $\langle 2(a(x) + b(x)f_i(x)) + v \rangle$ | $\langle 2(1 + a(x^{-1}) - \delta_i x^{2n-m_i} b(x^{-1}) f_{\mu(i)}(x)) + v \rangle$ |
| $\langle 2vf_i(x) \rangle$ | $\langle f_{\mu(i)}(x), v \rangle$ |
| $\langle 2(f_i(x)b(x) + v) \rangle$ | $\langle -f_{\mu(i)}(x)\delta_i x^{2n-m_i} b(x^{-1}) + v, 2 \rangle$ |
| $\langle f_i(x)(2a(x) + v) \rangle$ | $\langle 2(1 + a(x^{-1})) + v, 2f_{\mu(i)}(x) \rangle$ |
| $\langle 1 \rangle$ | $\langle 0 \rangle$ |
| $\langle f_i(x) \rangle$ | $\langle 2f_{\mu(i)}(x) \rangle$ |
| $\langle 2 \rangle$ | $\langle 2 \rangle$ |
| $\langle 2f_i(x) \rangle$ | $\langle f_{\mu(i)}(x) \rangle$ |
| $\langle 0 \rangle$ | $\langle 1 \rangle$ |
| $\langle f_i(x), v \rangle$ | $\langle 2vf_{\mu(i)}(x) \rangle$ |
| $\langle f_i(x)b(x) + v, 2 \rangle$ | $\langle 2(-f_{\mu(i)}(x)\delta_i x^{2n-m_i} b(x^{-1}) + v) \rangle$ |
| $\langle 2a(x) + v, 2f_i(x) \rangle$ | $\langle f_{\mu(i)}(x)(2(1 + a(x^{-1})) + v) \rangle$ |
| $\langle 2, vf_i(x) \rangle$ | $\langle 2f_{\mu(i)}(x), 2v \rangle$ |
| $\langle 2f_i(x), 2v \rangle$ | $\langle 2, vf_{\mu(i)}(x) \rangle$ |
| $\langle 2b(x) + vf_i(x), 2f_i(x) \rangle$ | $\langle 2b(x^{-1}) - vf_{\mu(i)}(x)\delta_i x^{2n-m_i}, 2f_{\mu(i)}(x) \rangle$ |

where $a(x), b(x) \in \mathcal{T}_i$.

Proof. Let $1 \leq i \leq r$. By Theorem 4.4 we see that $C_i \cdot \text{Ann}(C_i) = \{0\}$ and $|C_i| |\text{Ann}(C_i)| = 4^{4m_i}$. Since μ_i is a ring isomorphism from $\mathcal{K}_i + v\mathcal{K}_i$ onto $\mathcal{K}_{\mu(i)} + v\mathcal{K}_{\mu(i)}$, $\mu_i(\text{Ann}(C_i))$ is an ideal of $\mathcal{K}_{\mu(i)} + v\mathcal{K}_{\mu(i)}$. Denote $D_{\mu(i)} = \mu_i(\text{Ann}(C_i))$ and set $\mathcal{D} = \sum_{i=1}^r \varepsilon_{\mu(i)}(x) D_{\mu(i)}$. Then $\mathcal{D} = \bigoplus_{j=1}^r \varepsilon_j(x) D_j$ by Lemma 5.2(i) and Theorem 3.3(i). From this and by Theorem 4.4(ii), we deduce that \mathcal{D} is a negacyclic code over R of length $2n$.

As $\mu_i^{-1}(D_{\mu(i)}) = \text{Ann}(C_i)$ for all i , by Lemma 5.3 it follows that

$$\mathcal{C} \cdot \mu(\mathcal{D}) = \sum_{i=1}^r \varepsilon_i(x) (C_i \cdot \mu_i^{-1}(D_{\mu(i)})) = \sum_{i=1}^r \varepsilon_i(x) (C_i \cdot \text{Ann}(C_i)) = \{0\}.$$

This implies $\mathcal{D} \subseteq \mathcal{C}^\perp$ by Lemma 5.1. On the other hand, we have

$$\begin{aligned} |\mathcal{C}| |\mathcal{D}| &= \left(\prod_{i=1}^r |C_i| \right) \left(\prod_{i=1}^r |D_{\mu(i)}| \right) = \prod_{i=1}^r |C_i| |\text{Ann}(C_i)| = 4^{4 \sum_{i=1}^r m_i} = 4^{4n} \\ &= |(\mathbb{Z}_4 + v\mathbb{Z}_4)[x] / \langle x^{2n} + 1 \rangle| = |(\mathbb{Z}_4 + v\mathbb{Z}_4)^{2n}| \end{aligned}$$

by Theorem 4.2(ii). Since $\mathbb{Z}_4 + v\mathbb{Z}_4$ is a Frobenius ring, from the theory of linear codes over Frobenius rings (see [13]) we deduce that $\mathcal{C}^\perp = \mathcal{D}$.

Finally, for any integer i , $1 \leq i \leq r$, the expression for each ideal

$$D_{\mu(i)} = \mu_i(\text{Ann}(C_i)) = \{\mu_i(h(x)) \mid h(x) \in \text{Ann}(C_i)\} \pmod{f_{\mu(i)}(-x^2)}$$

follows from Theorem 4.4 and Lemma 5.4 immediately. \square

As the end of this section, we list all distinct self-dual negacyclic codes over R by Theorems 4.2 and 5.5.

Theorem 5.6 *Using the notations in Theorem 5.5 and Lemma 5.2(ii), denote*

$$\begin{aligned} \mathcal{W}_i^{(1)} = & \{(a(x), b(x)) \mid a(x) + a(x^{-1}) + 1 \equiv b(x) + x^{2n-m_i}b(x^{-1}) \equiv 0 \\ & \pmod{2, \pmod{\bar{f}_i(x)}}, a(x), b(x) \in \mathbb{F}_2[x]/\langle \bar{f}_i(x) \rangle\} \end{aligned}$$

and

$$\mathcal{W}_i^{(2)} = \{b(x) \in \mathbb{F}_2[x]/\langle \bar{f}_i(x) \rangle \mid b(x) + x^{m_i}b(x^{-1}) \equiv 0 \pmod{2, \pmod{\bar{f}_i(x)}}\}$$

for any $1 \leq i \leq \lambda$. Let \mathcal{C} be a negacyclic code over R of length $2n$ with $\mathcal{C} = \bigoplus_{i=1}^r \varepsilon_i(x)C_i$, where C_i is an ideal of $\mathcal{K}_i + v\mathcal{K}_i$. Then \mathcal{C} is self-dual if and only if C_i satisfies the following conditions:

(i) If $i = \lambda + j$ where $1 \leq j \leq \epsilon$, $(C_i, C_{i+\epsilon})$ is given by the following table:

| $C_i \pmod{f_i(-x^2)}$ | $C_{i+\epsilon} \pmod{f_{i+\epsilon}(-x^2)}$ |
|--------------------------------------------|------------------------------------------------------------------------------------------|
| $\langle 2(a(x) + b(x)f_i(x)) + v \rangle$ | $\langle 2(1 + a(x^{-1}) - \delta_i x^{2n-m_i}b(x^{-1})f_{i+\epsilon}(x)) + v \rangle$ |
| $\langle 2vf_i(x) \rangle$ | $\langle f_{i+\epsilon}(x), v \rangle$ |
| $\langle 2(f_i(x)b(x) + v) \rangle$ | $\langle -f_{i+\epsilon}(x)\delta_i x^{2n-m_i}b(x^{-1}) + v, 2 \rangle$ |
| $\langle f_i(x)(2a(x) + v) \rangle$ | $\langle 2(1 + a(x^{-1})) + v, 2f_{i+\epsilon}(x) \rangle$ |
| $\langle 1 \rangle$ | $\langle 0 \rangle$ |
| $\langle f_i(x) \rangle$ | $\langle 2f_{i+\epsilon}(x) \rangle$ |
| $\langle 2 \rangle$ | $\langle 2 \rangle$ |
| $\langle 2f_i(x) \rangle$ | $\langle f_{i+\epsilon}(x) \rangle$ |
| $\langle 0 \rangle$ | $\langle 1 \rangle$ |
| $\langle f_i(x), v \rangle$ | $\langle 2vf_{i+\epsilon}(x) \rangle$ |
| $\langle f_i(x)b(x) + v, 2 \rangle$ | $\langle 2(-f_{i+\epsilon}(x)\delta_i x^{2n-m_i}b(x^{-1}) + v) \rangle$ |
| $\langle 2a(x) + v, 2f_i(x) \rangle$ | $\langle f_{i+\epsilon}(x)(2(1 + a(x^{-1})) + v) \rangle$ |
| $\langle 2, vf_i(x) \rangle$ | $\langle 2f_{i+\epsilon}(x), 2v \rangle$ |
| $\langle 2f_i(x), 2v \rangle$ | $\langle 2, vf_{i+\epsilon}(x) \rangle$ |
| $\langle 2b(x) + vf_i(x), 2f_i(x) \rangle$ | $\langle 2b(x^{-1}) - vf_{i+\epsilon}(x)\delta_i x^{2n-m_i}, 2f_{i+\epsilon}(x) \rangle$ |

where $a(x), b(x) \in \mathcal{T}_i$.

(ii) If $1 \leq i \leq \lambda$, C_i is given by one of the following three cases:

(ii-1) $C_i = \langle 2 \rangle$.

(ii-2) $C_i = \langle 2(a(x) + b(x)f_i(x)) + v \rangle$, where $(a(x), b(x)) \in \mathcal{W}_i^{(1)}$.

(ii-3) $C_i = \langle 2b(x) + vf_i(x), 2f_i(x) \rangle$, where $b(x) \in \mathcal{W}_i^{(2)}$.

The number of self-dual negacyclic codes over R of length $2n$ is equal to

$$\prod_{i=1}^{\lambda} (1 + |\mathcal{W}_i^{(1)}| + |\mathcal{W}_i^{(2)}|) \prod_{j=1}^{\epsilon} (2^{2m_{\lambda+j}} + 5 \cdot 2^{m_{\lambda+j}} + 9).$$

Proof. By Lemma 5.2 (i) and (ii), we have that $\mu(i) = i$ for all $1 \leq i \leq \lambda$, $\mu(i) = i + \epsilon$ and $\mu(i + \epsilon) = i$ for all $\lambda + 1 \leq i \leq \lambda + \epsilon$. From this by Theorem 5.5 and its proof, we deduce that

$$\begin{aligned} \mathcal{C}^\perp &= \left(\bigoplus_{i=1}^{\lambda} \varepsilon_i(x) D_i \right) \oplus \left(\bigoplus_{i=\lambda+1}^{\lambda+\epsilon} (\varepsilon_{\mu(i)}(x) D_{\mu(i)} \oplus \varepsilon_{\mu(i+\epsilon)}(x) D_{\mu(i+\epsilon)}) \right) \\ &= \left(\bigoplus_{i=1}^{\lambda} \varepsilon_i(x) D_i \right) \oplus \left(\bigoplus_{i=\lambda+1}^{\lambda+\epsilon} (\varepsilon_{i+\epsilon}(x) D_{i+\epsilon} \oplus \varepsilon_i(x) D_i) \right) \end{aligned}$$

Then by Theorem 4.2, we see that $\mathcal{C} = \mathcal{C}^\perp$ if and only if $C_i = D_i$ for all $i = 1, \dots, r$. Now, we have one of the following two cases.

(i) Let $i = \lambda + j$ where $1 \leq j \leq \epsilon$. By $\mu(i) = i + \epsilon$, we have $C_{i+\epsilon} = D_{i+\epsilon} = D_{\mu(i)}$. Then the conclusions follow from Theorem 5.5 immediately.

(ii) $1 \leq i \leq \lambda$. In this case, $\mu(i) = i$. Then by $|C_i| = |D_i|$ and Theorem 5.5, we have one of the following three subcases:

(ii-1) $C_i = D_i = \langle 2 \rangle$.

(ii-2) $C_i = \langle 2(a(x) + b(x)f_i(x)) + v \rangle$ and $D_i = \langle v + 2(1 + a(x^{-1}) - \delta_i x^{2n-m_i} b(x^{-1}) f_i(x)) \rangle$ as ideals of $\mathcal{K}_i + v\mathcal{K}_i$. Then by Theorem 4.4, Theorem 2.5 and Theorem 3.2(iii), we see that $C_i = D_i$ if and only if $a(x) = 1 + a(x^{-1})$ and $b(x) = -\delta_i x^{2n-m_i} b(x^{-1})$ as elements of the finite field $\mathbb{F}_2[x]/\langle \bar{f}_i(x) \rangle$. As $-\delta_i = 1$ in \mathbb{F}_2 , these conditions are equivalent to $(a(x), b(x)) \in \mathcal{W}_i^{(1)}$.

(ii-3) $C_i = \langle 2b(x) + vf_i(x), 2f_i(x) \rangle$ and $D_i = \langle 2\hat{b}(x) - vf_i(x)\delta_i x^{2n-m_i}, 2f_i(x) \rangle$. In this case, $C_i = D_i$ if and only if $b(x) = \frac{1}{-\delta_i x^{2n-m_i}} b(x^{-1})$ in $\mathbb{F}_2[x]/\langle \bar{f}_i(x) \rangle$. Obviously, the latter is equivalent to $b(x) \in \mathcal{W}_i^{(2)}$. \square

6. Negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length $2M_p$

Recall that $2^p - 1$ is called a *Mersenne prime*, denoted by

$$M_p = 2^p - 1,$$

if p is a prime and $2^p - 1$ is a prime as well. For example, when $p = 2, 3, 5, 7, 13, 17, 19, 31$, M_p is a prime. In this section, we present negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ ($v^2 = 2v$) of length $2M_p$.

Theorem 6.1 *Let $n = M_p$ where $p \geq 3$. Then the number of all negacyclic codes over $\mathbb{Z}_4 + u\mathbb{Z}_4$ of length $2n$ and the number of self-dual negacyclic codes over $\mathbb{Z}_4 + u\mathbb{Z}_4$ of length $2n$ are equal to*

$$23 \cdot (4^p + 5 \cdot 2^p + 9)^{2 \cdot \frac{2^{p-1}-1}{p}} \text{ and } 3 \cdot (4^p + 5 \cdot 2^p + 9)^{\frac{2^{p-1}-1}{p}}, \text{ respectively.}$$

Proof. Since $n = M_p$ and M_p is a Mersenne prime, for any integer s , $1 \leq s \leq n-1$, the 2-cyclotomic coset modulo n containing s is $J_s^{(2)} = \{s, 2s, \dots, 2^{p-1}s\}$ with $|J_s^{(2)}| = p$. Therefore, the number of 2-cyclotomic cosets modulo n is equal to $1 + \frac{n-1}{p} = 1 + \frac{2^p-1-1}{p} = 1 + 2 \cdot \frac{2^{p-1}-1}{p}$.

Suppose that $2^j s \equiv -s \pmod{n}$ for some $1 \leq j \leq p-1$. Then $(2^p - 1) | s(2^j + 1)$. Since $\gcd(s, 2^p - 1) = 1$, we have $(2^p - 1) | (2^j + 1)$. This implies $2^p - 1 \leq 2^j + 1$. On the other hand, by $p \geq 3$ and $1 \leq j < p$, we have $(2^p - 1) - (2^j + 1) = 2^j(2^{p-j} - 1) - 2 > 0$. This implies $(2^p - 1) > (2^j + 1)$, and we get a contradiction. Therefore, $J_{-s}^{(2)} \neq J_s^{(2)}$ for all $1 \leq s \leq n-1$.

Using the notations of Lemma 5.2(ii), we have $r = 1 + 2 \cdot \frac{2^{p-1}-1}{p}$, $\lambda = 1$, $\epsilon = \frac{2^{p-1}-1}{p}$, $m_1 = |J_0^{(2)}| = 1$ and $m_i = |J_1^{(2)}| = p$ for all $i = 2, 3, \dots, r$.

By Corollary 4.5, the number of all negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length $2n$ is $(2^2 + 5 \cdot 2 + 9) \prod_{i=2}^r (2^{2p} + 5 \cdot 2^p + 9) = 23 \cdot (4^p + 5 \cdot 2^p + 9)^{2 \cdot \frac{2^{p-1}-1}{p}}$.

It is obvious that $f_1(x) = x - 1$ and $\tilde{f}_1(x) = 1 - x = \delta_1 f_1(x)$ with $\delta_1 = -1$. Hence $\mathcal{K}_1 = \mathbb{Z}_4[x] / \langle f_1(-x^2) \rangle = \mathbb{Z}_4[x] / \langle x^2 + 1 \rangle$. Using the notations in Section 3, by $m_1 = 1$ we have $\mathcal{T}_1 = \{0, 1\}$ and $\mathbb{F}_2[x] / \langle \tilde{f}_1(x) \rangle = \mathbb{F}_2$ where $\tilde{f}_1(x) = x - 1 \in \mathbb{F}_2[x]$. Then we have

$$\mathcal{W}_1^{(1)} = \{(a, b) \mid a + a + 1 = b + b = 0, a, b \in \mathbb{F}_2\} = \emptyset \text{ with } |\mathcal{W}_1^{(1)}| = 0;$$

$$\mathcal{W}_1^{(2)} = \{b \in \mathbb{F}_2 \mid b + b = 0\} = \mathbb{F}_2 \text{ with } |\mathcal{W}_1^{(2)}| = 2.$$

Therefore, using the notations in the proof of Theorem 4.6(i) we see that there are 3 ideals of $\mathcal{K}_1 + v\mathcal{K}_1$ satisfying $C_1 = D_1$:

$$\langle 2 \rangle, \langle v(x-1), 2(x-1) \rangle, \langle 2 + v(x-1), 2(x-1) \rangle.$$

For any $i = 1 + j$, where $1 \leq j \leq \epsilon = \frac{2^{p-1}-1}{p}$, By Theorem 4.4 and $m_i = p$ we know that the number of pairs of $(C_i, C_{i+\epsilon})$ in Theorem 5.6(ii) is equal to $2^{2p} + 5 \cdot 2^p + 9 = 4^p + 5 \cdot 2^p + 9$.

As stated above, by Theorem 5.6 we conclude that the number of self-dual negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length $2(2^p - 1)$, where both p and $2^p - 1$ are prime integers, is equal to $3 \cdot (4^p + 5 \cdot 2^p + 9)^{\frac{2^{p-1}-1}{p}}$. \square

For example, when $p = 5$, we have $M_5 = 31$. Hence the number of self-dual negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length 62 is equal to

$$3 \cdot (4^5 + 5 \cdot 2^5 + 9)^{\frac{2^{5-1}-1}{5}} = 3 \cdot 1193^3 = 5093808171 \approx 5 \times 10^9.$$

All these codes can be listed by Theorem 5.6. From these codes, by Proposition 1.3 we obtain 5093808171 self-dual 2-quasi-twisted codes over \mathbb{Z}_4 of length 124, and 5093808171 self-dual 4-quasi-twisted binary codes of length 248 by the Gray map from \mathbb{Z}_4 onto \mathbb{F}_2^2 : $0 \mapsto 00, 1 \mapsto 01, 2 \mapsto 11, 3 \mapsto 10$.

When $p = 7$, we have $M_7 = 127$. Hence the number of self-dual negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length 254 is equal to

$$3 \cdot (4^7 + 5 \cdot 2^7 + 9)^{\frac{2^{7-1}-1}{7}} = 3 \cdot 17033^9 \approx 3.62 \times 10^{38}.$$

All these codes can be listed by Theorem 5.6.

Now, let $p = 3$. Then $n = M_3 = 2^3 - 1 = 7$. By Theorem 6.1 we know that the number of all negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length 14 is

$$23 \cdot (4^3 + 5 \cdot 2^3 + 9)^2 = 23 \cdot 113^2 = 293687,$$

and the number of self-dual negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length 14 is equal to $3 \cdot (4^3 + 5 \cdot 2^3 + 9) = 3 \cdot 113 = 339$.

Precisely, we have that $x^7 - 1 = f_1(x)f_2(x)f_3(x)$, where $f_1(x) = x - 1$, $f_2(x) = x^3 + 2x^2 + x + 3$ and $f_3(x) = x^3 + 3x^2 + 2x + 3$ are pairwise coprime basic irreducible polynomials in $\mathbb{Z}_4[x]$. Hence $r = 3$, $m_1 = 1$ and $m_2 = m_3 = 3$. Obviously, $\tilde{f}_1(x) = \delta_1 f_1(x)$ and $\tilde{f}_2(x) = \delta_2 f_3(x)$ where $\delta_1 = \delta_2 = -1$, which implies that $\mu(1) = 1$ and $\mu(2) = 3$.

For each integer i , $1 \leq i \leq 3$, we denote $F_i(x) = \frac{x^7-1}{f_i(x)}$, and find polynomials $a_i(x), b_i(x) \in \mathbb{Z}_4[x]$ satisfying $a_i(x)F_i(x) + b_i(x)f_i(x) = 1$. Then we set $\varepsilon_i(x) \equiv a_i(-x^2)F_i(-x^2) \pmod{x^{14} + 1}$. Precisely, we have

$$\begin{aligned} \varepsilon_1(x) &= 3 + x^2 + 3x^4 + x^6 + 3x^8 + x^{10} + 3x^{12}; \\ \varepsilon_2(x) &= 1 + x^2 + 3x^4 + 2x^6 + 3x^8 + 2x^{10} + 2x^{12}; \end{aligned}$$

$$\varepsilon_3(x) = 1 + 2x^2 + 2x^4 + x^6 + 2x^8 + x^{10} + 3x^{12}.$$

Using the notations in Section 3, we have $\mathcal{K}_i = \mathbb{Z}_4[x]/\langle f_i(-x^2) \rangle$ for $i = 1, 2, 3$, and $\mathcal{T}_2 = \{c_0 + c_1x + c_2x^2 \mid c_0, c_1, c_2 \in \{0, 1\}\}$. As $x^{14} = -1$, we have $x^{-1} = -x^{13}$ in \mathcal{K}_2 . Hence for any $a(x) = a_0 + a_1x + a_2x^2 \in \mathcal{T}_2$, it follows that

$$a(x^{-1}) = a_0 + 2(a_1 + a_2) + 3a_2x^2 + 3a_1x^3 + 3a_2x^4 + 3a_1x^5 \pmod{f_3(-x^2)}$$

and $\bar{a}(x^{-1}) = a_0 + a_2x^2 + a_1x^3 + a_2x^4 + a_1x^5$. By Theorem 5.6, all distinct self-dual negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length 14 are given by

$$\mathcal{C} = \varepsilon_1(x)C_1 \oplus \varepsilon_2(x)C_2 \oplus \varepsilon_3(x)C_3 \pmod{x^{14} + 1},$$

where C_i is an ideal of the ring $\mathcal{K}_i + v\mathcal{K}_i$ ($\mathcal{K}_i = \mathbb{Z}_4[x]/\langle f_i(-x^2) \rangle$) satisfying one of the following conditions:

- C_1 is one of the following 3 ideals: $\langle 2 \rangle$, $\langle v(x-1), 2(x-1) \rangle$, $\langle 2 + v(x-1), 2(x-1) \rangle$.
- (C_2, C_3) is given by one of the following 113 cases, where $a(x), b(x) \in \mathcal{T}_2$:

| $C_2 \pmod{f_2(-x^2)}$ | $C_3 \pmod{f_3(-x^2)}$ |
|--------------------------------------------|----------------------------------------------------------------|
| $\langle 2(a(x) + b(x)f_2(x)) + v \rangle$ | $\langle 2(1 + a(x^{-1}) + x^{11}b(x^{-1})f_3(x)) + v \rangle$ |
| $\langle 2vf_2(x) \rangle$ | $\langle f_3(x), v \rangle$ |
| $\langle 2(f_2(x)b(x) + v) \rangle$ | $\langle f_3(x)x^{11}b(x^{-1}) + v, 2 \rangle$ |
| $\langle f_2(x)(2a(x) + v) \rangle$ | $\langle 2(1 + a(x^{-1})) + v, 2f_3(x) \rangle$ |
| $\langle 1 \rangle$ | $\langle 0 \rangle$ |
| $\langle f_2(x) \rangle$ | $\langle 2f_3(x) \rangle$ |
| $\langle 2 \rangle$ | $\langle 2 \rangle$ |
| $\langle 2f_2(x) \rangle$ | $\langle f_3(x) \rangle$ |
| $\langle 0 \rangle$ | $\langle 1 \rangle$ |
| $\langle f_2(x), v \rangle$ | $\langle 2vf_3(x) \rangle$ |
| $\langle f_2(x)b(x) + v, 2 \rangle$ | $\langle 2(f_3(x)x^{11}b(x^{-1}) + v) \rangle$ |
| $\langle 2a(x) + v, 2f_2(x) \rangle$ | $\langle f_3(x)(2(1 + a(x^{-1})) + v) \rangle$ |
| $\langle 2, vf_2(x) \rangle$ | $\langle 2f_3(x), 2v \rangle$ |
| $\langle 2f_2(x), 2v \rangle$ | $\langle 2, vf_3(x) \rangle$ |
| $\langle 2b(x) + vf_2(x), 2f_2(x) \rangle$ | $\langle 2b(x^{-1}) + vf_3(x)x^{11}, 2f_3(x) \rangle$ |

Among the 339 self-dual negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of length 14, we obtain 36 codes \mathcal{C} whose Gray image $\theta(\mathcal{C})$ are new good self-dual 2-quasi-twisted code over \mathbb{Z}_4 of length 28 (these codes does not exist in [21] too).

The new codes we obtained have new type, better minimum Lee weight, more algebraic structures and properties.

◇ When $C_1 = \langle v(x-1), 2(x-1) \rangle$ or $C_1 = \langle 2 + v(x-1), 2(x-1) \rangle$, and (C_2, C_3) is given by one of the following 12 cases:

| $C_2 \pmod{f_2(-x^2)}$ | $C_3 \pmod{f_3(-x^2)}$ |
|----------------------------------------------|------------------------------------------------------------------|
| $\langle 2(x^2 f_2(x) + v) \rangle$ | $\langle f_3(x)x^{11}(x^4 + x^2) + v, 2 \rangle$ |
| $\langle 2(x f_2(x) + v) \rangle$ | $\langle f_3(x)x^{11}(x^5 + x^3) + v, 2 \rangle$ |
| $\langle 2((x^2 + x)f_2(x) + v) \rangle$ | $\langle f_3(x)x^{11}(x^5 + x^4 + x^3 + x^2) + v, 2 \rangle$ |
| $\langle 2((x^2 + 1)f_2(x) + v) \rangle$ | $\langle f_3(x)x^{11}(x^4 + x^2 + 1) + v, 2 \rangle$ |
| $\langle 2((x + 1)f_2(x) + v) \rangle$ | $\langle f_3(x)x^{11}(x^5 + x^3 + 1) + v, 2 \rangle$ |
| $\langle 2((x^2 + x + 1)f_2(x) + v) \rangle$ | $\langle f_3(x)x^{11}(x^5 + x^4 + x^3 + x^2 + 1) + v, 2 \rangle$ |
| $\langle x^2 f_2(x) + v, 2 \rangle$ | $\langle 2(f_3(x)x^{11}(x^4 + x^2 + 1) + v) \rangle$ |
| $\langle x f_2(x) + v, 2 \rangle$ | $\langle 2(f_3(x)x^{11}(x^5 + x^3) + v) \rangle$ |
| $\langle (x^2 + x)f_2(x) + v, 2 \rangle$ | $\langle 2(f_3(x)x^{11}(x^5 + x^4 + x^3 + x^2) + v) \rangle$ |
| $\langle (x^2 + 1)f_2(x) + v, 2 \rangle$ | $\langle 2(f_3(x)x^{11}(x^4 + x^2 + 1) + v) \rangle$ |
| $\langle (x + 1)f_2(x) + v, 2 \rangle$ | $\langle 2(f_3(x)x^{11}(x^5 + x^3 + 1) + v) \rangle$ |
| $\langle (x^2 + x + 1)f_2(x) + v, 2 \rangle$ | $\langle 2(f_3(x)x^{11}(x^5 + x^4 + x^3 + x^2 + 1) + v) \rangle$ |

we obtain 24 new good self-dual \mathbb{Z}_4 codes $\theta(\mathcal{C})$ with basic parameters $(28, 2^{28}, d_L = 8, d_E = 12)$ and of type $2^{14}4^7$, where d_L is the minimum Lee weight and d_E is the minimum Euclidean weight of \mathcal{C} .

◇ When $C_1 = \langle 2 \rangle$ and (C_2, C_3) is given by one of the above 12 cases, we obtain 12 new good self-dual \mathbb{Z}_4 codes $\theta(\mathcal{C})$ with basic parameters $(28, 2^{28}, d_L = 6, d_E = 12)$ and of type $2^{16}4^6$.

Remark The existing self-dual codes over \mathbb{Z}_4 are listed in the database [26] and the maximal code length is 19 in the table. In [27], the existing linear codes over \mathbb{Z}_4 of length 24 have basic parameters $(24, 2^{28}, d_L = 6)$ and $(24, 2^{28}, d_L = 5)$ and both of type 4^{14} .

7. Conclusions and further research

We have developed a theory for negacyclic codes over the ring $\mathbb{Z}_4 + v\mathbb{Z}_4$ ($v^2 = 2v$) of oddly even length, including the enumeration and construction of these codes, the dual code and self-duality for each of these codes. These codes enjoy a rich algebraic structure (which makes the search process much simpler) compared to arbitrary linear codes. Our further interest is to consider negacyclic codes over $\mathbb{Z}_4 + v\mathbb{Z}_4$ of arbitrary even length. On the other

hand, obtaining some bounds for minimum distance such as BCH-like of a negacyclic code over the ring $\mathbb{Z}_4 + v\mathbb{Z}_4$ by just looking at the canonical form decomposition would be rather interesting.

Acknowledgments Part of this work was done when Yonglin Cao was visiting Chern Institute of Mathematics, Nankai University, Tianjin, China. Yonglin Cao would like to thank the institution for the kind hospitality. This research is supported in part by the National Natural Science Foundation of China (Grant Nos. 11671235, 11471255).

Appendix: Proof of Theorem 2.5

Using the notations of Section 2, by [6] Example 2.5 we know that the number of linear codes over the finite chain ring \mathcal{K} of length 2 is equal to $\sum_{i=0}^4 (2i+1)|\mathcal{T}|^{4-i}$. Moreover, every linear code C over \mathcal{K} of length 2 has one and only one of the following matrices G as their generator matrices:

- (i) $G = (1, a)$, $a \in \mathcal{K}$. (ii) $G = (\pi^k, \pi^k a)$, $a \in \mathcal{K}/\langle \pi^{4-k} \rangle$, $1 \leq k \leq 3$.
- (iii) $G = (\pi b, 1)$, $b \in \mathcal{K}/\langle \pi^3 \rangle$.
- (iv) $G = (\pi^{k+1}b, \pi^k)$, $b \in \mathcal{K}/\langle \pi^{4-k-1} \rangle$, $1 \leq k \leq 3$.
- (v) $G = \pi^k I_2$, $0 \leq k \leq 4$. (vi) $G = \begin{pmatrix} 1 & c \\ 0 & \pi^t \end{pmatrix}$, $c \in \mathcal{K}/\langle \pi^t \rangle$, $1 \leq t \leq 3$.
- (vii) $G = \begin{pmatrix} \pi^k & \pi^k c \\ 0 & \pi^{k+t} \end{pmatrix}$, $c \in \mathcal{K}/\langle \pi^t \rangle$, $1 \leq t \leq 4-k-1$, $1 \leq k \leq 2$.
- (viii) $G = \begin{pmatrix} c & 1 \\ \pi^t & 0 \end{pmatrix}$, where $c \in \pi(\mathcal{K}/\langle \pi^t \rangle)$ and $1 \leq t \leq 3$.
- (ix) $G = \begin{pmatrix} \pi^k c & \pi^k \\ \pi^{k+t} & 0 \end{pmatrix}$, $c \in \pi(\mathcal{K}/\langle \pi^t \rangle)$, $1 \leq t \leq 4-k-1$, $1 \leq k \leq 2$.

Therefore, we only need to consider the nine cases listed above:

(i) Suppose that C satisfies Condition (2). By $(1, a) \in C$, we have $(0, 1 + \omega\pi^2 a) \in C$. Since G is the generator matrix of C , there exists $b \in \mathcal{K}$ such that $(0, 1 + \omega\pi^2 a) = b(1, a) = (b, ba)$, i.e. $0 = b$ and $1 + \omega\pi^2 a = ba$. This implies $1 + \omega\pi^2 a = 0$. So we get a contradiction, since $1 + \omega\pi^2 a$ is an invertible element of \mathcal{K} . Hence C does not satisfy Condition (2) in this case.

(ii) Suppose that C satisfies Condition (2). By $(\pi^k, \pi^k a) \in C$, we have $(0, \pi^k + \omega\pi^{k+2}a) = (0, \pi^k + \omega\pi^2 \cdot \pi^k a) \in C$. Then there exists $b \in \mathcal{K}$ such that $(0, \pi^k + \omega\pi^{k+2}a) = b(\pi^k, \pi^k a) = (\pi^k b, \pi^k ba)$. This implies $0 = \pi^k b$ and $\pi^k(1 + \omega\pi^2 a) = \pi^k ba$. Hence $\pi^k(1 + \omega\pi^2 a) = 0$. Since $1 + \omega\pi^2 a$ is an invertible

element of \mathcal{K} , we deduce $\pi^k = 0$, which contradict that $k \leq 3$.

(iii) In this case, C satisfies Condition (2) if and only if there exists $a \in \mathcal{K}$ such that $(0, \pi b + \omega\pi^2) = (0, \pi b + \omega\pi^2 \cdot 1) = a(\pi b, 1) = (\pi ab, a)$, i.e. $0 = \pi ab$ and $\pi(b + \omega\pi) = \pi b + \omega\pi^2 = a$. These conditions are equivalent to that b satisfies $\pi^2(b + \omega\pi)b = 0$, i.e. $(b + \omega\pi)b \in \pi^2\mathcal{K}$. The latter condition is equivalent to $\|(b + \omega\pi)b\|_\pi \geq 2$. As $b \in \mathcal{K}/\langle\pi^3\rangle$, b has a unique π -expansion: $b = r_0 + r_1\pi + r_2\pi^2$ with $r_0, r_1, r_2 \in \mathcal{T}$. This implies $b + \omega\pi = r_0 + (\omega + r_1)\pi + r_2\pi^2$. Suppose that $r_0 \neq 0$. Then $\|b\|_\pi = \|b + \omega\pi\|_\pi = 0$. This implies $\|(b + \omega\pi)b\|_\pi = 0$ and we get a contradiction. Now, let $r_0 = 0$. Then by $(b + \omega\pi)b = \pi^2(\omega + r_1 + r_2\pi)(r_1 + r_2\pi)$ it follow that $\|(b + \omega\pi)b\|_\pi \geq 2$. Hence $b = \pi(r_1 + r_2\pi)$ for all $r_1, r_2 \in \mathcal{T}$ in this case.

(iv) In this case, C satisfies Condition (2) if and only if there exists $a \in \mathcal{K}$ such that $(0, \pi^{k+1}(b + \omega\pi)) = (0, \pi^{k+1}b + \omega\pi^2 \cdot \pi^k) = a(\pi^{k+1}b, \pi^k) = (\pi^{k+1}ab, \pi^k a)$, i.e. $0 = \pi^{k+1}ab = \pi b \cdot \pi^k a$ and $\pi^{k+1}(b + \omega\pi) = \pi^k a$. These conditions are equivalent to that b satisfies $\pi^{k+2}(b + \omega\pi)b = 0$, i.e. $\|(b + \omega\pi)b\|_\pi \geq 4 - k - 2$ where $1 \leq k \leq 3$. Then we have one of the following three cases:

(iv-1) Let $k = 3$. Then $\|(b + \omega\pi)b\|_\pi \geq 4 - k - 2 = -1$ for any $b \in \mathcal{K}/\langle\pi^{4-3-1}\rangle = \mathcal{K}/\langle 1 \rangle = \{0\}$. Hence $G = (0, \pi^3)$.

(iv-2) Let $k = 2$. $\|(b + \omega\pi)b\|_\pi \geq 4 - k - 2 = 0$ for any $b \in \mathcal{K}/\langle\pi^{4-2-1}\rangle = \mathcal{K}/\langle\pi\rangle = \mathcal{T}$. In this case, we have $G = (\pi^3 b, \pi^2)$.

(iv-3) Let $k = 1$. $\|(b + \omega\pi)b\|_\pi \geq 4 - k - 2 = 1$ if and only if $b \in \pi(\mathcal{K}/\langle\pi^{4-1-1}\rangle) = \pi(\mathcal{K}/\langle\pi^2\rangle)$. In this case, we have $G = (\pi^2 b, \pi) = (\pi^3 a, \pi)$ for all $b = \pi a$ with $a \in \mathcal{T}$.

(v) In this case, C satisfies Condition (2) for all $0 \leq k \leq 4$.

(vi) Suppose that C satisfies Condition (2). Then there exist $a, b \in \mathcal{K}$ such that $(0, 1 + \omega\pi^2 c) = a(1, c) + b(0, \pi^t) = (a, ac + \pi^t b)$, i.e. $0 = a$ and $1 + \omega\pi^2 c = ac + \pi^t b$. This implies $1 = \pi(\pi^{t-1}b - \omega\pi c)$ and we get a contradiction. Hence C does not satisfy Condition (2) in this case.

(vii) Suppose that C satisfies Condition (2). Then there exist $a, b \in \mathcal{K}$ such that $(0, \pi^k + \omega\pi^2 \cdot \pi^k c) = a(\pi^k, \pi^k c) + b(0, \pi^{k+t}) = (\pi^k a, \pi^k ac + \pi^{k+t} b)$, i.e. $0 = \pi^k a$ and $\pi^k + \omega\pi^{k+2} c = \pi^k a \cdot c + \pi^{k+t} b$. This implies $\pi^k = \pi^{k+1}(\pi^{t-1}b - c\omega\pi)$, and we get a contradiction as $1 \leq k \leq 3$.

(viii) It is clear that $(0, \pi^t) = \pi^t(c, 1) - c(\pi^t, 0) \in C$. Hence C satisfies Condition (2) if and only if there exist $a, b \in \mathcal{K}$ such that $(0, c + \omega\pi^2 \cdot 1) = a(c, 1) + b(\pi^t, 0) = (ac + \pi^t b, a)$, i.e. $0 = ac + \pi^t b$ and $c + \omega\pi^2 = a$, which are equivalent to that $(c + \omega\pi^2)c = -\pi^t b \in \pi^t\mathcal{K}$ for some $b \in \mathcal{K}$, i.e. $\|(c + \omega\pi^2)c\|_\pi \geq t$, where $c \in \pi(\mathcal{K}/\langle\pi^t\rangle)$ and $1 \leq t \leq 3$. Then we have one of

the following three subcases:

(viii-1) When $t = 1$, then $c \in \pi(\mathcal{K}/\langle\pi\rangle) = \{0\}$, i.e. $c = 0$. In this case, $\|(c + \omega\pi^2)c\|_\pi = \|0\|_\pi = 4 > t$. Hence $G = \begin{pmatrix} 0 & 1 \\ \pi & 0 \end{pmatrix}$.

(viii-2) When $t = 2$, then $c \in \pi(\mathcal{K}/\langle\pi^2\rangle) = \pi\mathcal{T}$, i.e. $c = \pi z$ where $z \in \mathcal{T}$. In this case, $\|(c + \omega\pi^2)c\|_\pi = \|(\pi h + \omega\pi^2) \cdot \pi h\|_\pi \geq 2 = t$. Hence $G = \begin{pmatrix} \pi z & 1 \\ \pi^2 & 0 \end{pmatrix}$.

(viii-3) When $t = 3$, then $c \in \pi(\mathcal{K}/\langle\pi^3\rangle)$, i.e. $c = r_1\pi + r_2\pi^2$ where $r_1, r_2 \in \mathcal{T}$. Suppose $r_1 \neq 0$. Then $\|c\|_\pi = \|c + \omega\pi^2\|_\pi = 1$. This implies $\|(c + \omega\pi^2)c\|_\pi = 2 < 3 = t$. Now, let $r_1 = 0$. Then it is obvious that $\|(c + \omega\pi^2)c\|_\pi = 4 > t$ for all $c = \pi^2 z$ where $z = r_2 \in \mathcal{T}$. Hence $G = \begin{pmatrix} \pi^2 z & 1 \\ \pi^3 & 0 \end{pmatrix}$.

(xi) It is clear that $(0, \pi^{k+t}) = \pi^t(\pi^k c, \pi^k) - c(\pi^{k+t}, 0) \in C$. Hence C satisfies Condition (2) if and only if there exist $a, b \in \mathcal{K}$ such that $(0, \pi^k c + \omega\pi^2 \cdot \pi^k) = a(\pi^k c, \pi^k) + b(\pi^{k+t}, 0) = (\pi^k a c + \pi^{k+t} b, \pi^k a)$, i.e. $0 = \pi^k a c + \pi^{k+t} b$ and $\pi^k(c + \omega\pi^2) = \pi^k a$, which are equivalent to that $\pi^k(c + \omega\pi^2)c = -\pi^{k+t}b \in \pi^{k+t}\mathcal{K}$, i.e. $\|(c + \omega\pi^2)c\|_\pi \geq t$, where $c \in \pi(\mathcal{K}/\langle\pi^t\rangle)$, $1 \leq t \leq 3 - k$ and $1 \leq k \leq 2$. Then we have one of the following two subcases:

(xi-1) Let $k = 1$. Then $1 \leq t \leq 2$. If $t = 1$, then $c \in \pi(\mathcal{K}/\langle\pi\rangle) = \{0\}$, i.e. $c = 0$. Obviously, $\|(c + \omega\pi^2)c\|_\pi = 4 > t$. Hence $G = \begin{pmatrix} 0 & \pi \\ \pi^2 & 0 \end{pmatrix}$.

If $t = 2$, then $c \in \pi(\mathcal{K}/\langle\pi^2\rangle) = \pi\mathcal{T}$, i.e. $c = \pi z$ where $z \in \mathcal{T}$. Obviously, $\|(c + \omega\pi^2)c\|_\pi = 2 \geq t$. Hence $G = \begin{pmatrix} \pi^2 z & \pi \\ \pi^3 & 0 \end{pmatrix}$.

(xi-2) Let $k = 2$. Then $t = 1$ and $c = 0$. Hence $G = \begin{pmatrix} 0 & \pi^2 \\ \pi^3 & 0 \end{pmatrix}$.

References

- [1] Abualrub T., Siap I.: Cyclic codes over the ring $\mathbb{Z}_2 + u\mathbb{Z}_2$ and $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2$, Des. Codes Cryptogr. **42** (2007), 273–287.
- [2] Bonnecaze, A., Udaya, P: Cyclic codes and self-dual codes over $\mathbb{F}_2 + u\mathbb{F}_2$, IEEE Trans. Inform. Theory **45** (1999), 1250–1255.
- [3] Calderbank, A.R., Hammons Jr., A.R., Kumar, P.V., Sloane, N.J.A. and Solé, P.: A linear construction for certain Kerdock and Preparata

- codes, Bulletin of the American Mathematical Society **29**(2) (1993), 218–222.
- [4] Calderbank, A.R., Hammons Jr., A.R., Kumar, P.V., Sloane, N.J.A. and Solé, P.: The \mathbb{Z}_4 -linearity of Kerdock, Preparata, Goethals, and related codes, IEEE Trans. Inform. Theory **40** (1994), 301–319.
 - [5] Cao, Y., Cao, Y.: Complete classification for simple root cyclic codes over local rings $\mathbb{Z}_{p^s}[v]/\langle v^2 - pv \rangle$, <https://www.researchgate.net/publication/320620031>.
 - [6] Cao, Y., Gao, Y.: Repeated root cyclic \mathbb{F}_q -linear codes over \mathbb{F}_{q^l} , Finite Fields Appl. **31** (2015), 202–227.
 - [7] Cao, Y.: On constacyclic codes over finite chain rings, Finite Fields Appl. **24** (2013), 124–135.
 - [8] Cao, Y., Cao, Y., Li, Q.: The concatenated structure of cyclic codes over \mathbb{Z}_{p^2} , J. Appl. Math. Comput. **52** (2016), 363–385.
 - [9] Cao, Y., Cao, Y., Li, Q.: Concatenated structure of cyclic codes over \mathbb{Z}_4 of length $4n$, Appl. Algebra in Engrg. Comm. Comput. **10** (2016), 279–302.
 - [10] Dinh, H. Q., López-Permouth, S. R.: Cyclic and negacyclic codes over finite chain rings, IEEE Trans. Inform. Theory **50** (2004), 1728–1744.
 - [11] Dinh, H. Q.: Negacyclic codes of length 2^s over Galois rings, IEEE Trans. Inform. Theory **51** (2005), 4252–4262.
 - [12] Dinh H. Q.: Complete distance of all negacyclic codes of length 2^s over \mathbb{Z}_{2^a} , IEEE Trans. Inf. Theory **53** (2007), 147–161.
 - [13] Dougherty S. T., Kim J-L, Kulosman H., Liu H.: Self-dual codes over commutative Frobenius rings, Finite Fields Appl. **16**, 14–26 (2010).
 - [14] Kanwar, P. and López-Permouth, S. R.: Cyclic codes over the integers modulo p^m , Finite Fields Appl. **4**:3 (1997), 334–352.
 - [15] Martínez-Moro, E., Szabo, S., Yildiz, B.: Linear codes over $\frac{\mathbb{Z}_4[x]}{\langle x^2+2x \rangle}$, Int. J. Information and Coding Theory, Vol. 3, No. 1 (2015), 78–96.

- [16] Norton, G., Sălăgean-Mandache, A.: On the structure of linear and cyclic codes over finite chain rings, *Appl. Algebra in Engrg. Comm. Comput.* **10** (2000), 489–506.
- [17] Shi, M., Zhang, Y.: Quasi-twisted codes with constacyclic constituent codes, *Finite Fields Appl.* **39** (2016), 159–178.
- [18] Shi, M., Guan, Y., Solé, P.: Two new families of two-weight codes, *IEEE Trans. Inform. Theory* **63**(10) (2017), 6240–6246.
- [19] Shi, M., Xu, L., Yang, G.: A note on one weight and two weight projective \mathbb{Z}_4 -codes, *IEEE Trans. Inform. Theory* **63**(1) (2017), 177–182.
- [20] Shi, M., Solé, P., Wu, B.: Cyclic codes and the weight enumerators over $\mathbb{F}_2 + v\mathbb{F}_2 + v^2\mathbb{F}_2$, *Applied and Computational Mathematics* **12**(2) (2013), 247–255.
- [21] Shi, M., Qian, L., Sok, L., Aydin, N., Solé, P.: On constacyclic codes over $\mathbb{Z}_4[u]/\langle u^2 - 1 \rangle$ and their Gray images, *Finite Fields Appl.* **45**:3 (2017), 86–95.
- [22] Wan Z.-X.: *Quaternary Codes*. World Scientific Pub Co Inc. 1997.
- [23] Wood, J. A.: Duality for modules over finite rings and applications to coding theory, *American Journal of Mathematics*, Vol. 121, No. 3 (1999), 555–575.
- [24] Yildiz B., Karadeniz S.: Linear codes over $\mathbb{Z}_4 + u\mathbb{Z}_4$: MacWilliams identities, projections, and formally self-dual codes, *Finite Fields Appl.* **27**, 24–40 (2014).
- [25] Yildiz, B. and Aydin, N.: Cyclic codes over $\mathbb{Z}_4 + u\mathbb{Z}_4$ and \mathbb{Z}_4 images, *International Journal of Information and Coding Theory*, Vol. 2, No. 4 (2014), 226–237.
- [26] Self-dual codes over integers modulo 4 (accessed on 04 Dec. 2017), <http://www.math.is.tohoku.ac.jp/~Emunemasa/research/codes/sd4.htm>.
- [27] The \mathbb{Z}_4 database (accessed on 04 Dec. 2017), <http://www.asamov.com/Z4Codes/CODES/ShowCODESTablePage.aspx>.