Do thermodynamically stable rigid solids exist?

Parswa Nath,¹ Saswati Ganguly,² Jürgen Horbach,² Peter Sollich,³ Smarajit Karmakar,¹ and Surajit Sengupta¹

¹TIFR Centre for Interdisciplinary Sciences, 36/P Gopanpally, Hyderabad 500107, India

Heinrich Heine-Universität Düsseldorf, Universitätsstraße 1, 40225 Düsseldorf, Germany

³King's College London, Department of Mathematics, Strand, London WC2R 2LS, U.K.

(Dated: March 9, 2024)

Customarily, crystalline solids are defined to be *rigid* since they resist changes of shape determined by their boundaries. However, rigid solids cannot exist in the thermodynamic limit where boundaries become irrelevant. Particles in the solid may rearrange to adjust to shape changes eliminating stress without destroying crystalline order. Rigidity is therefore valid only in the metastable state that emerges because these particle rearrangements in response to a deformation, or strain, are associated with slow collective processes. Here, we show that a thermodynamic collective variable may be used to quantify particle rearrangements that occur as a solid is deformed at zero strain rate. Advanced Monte Carlo simulation techniques are then employed to obtain the equilibrium free energy as a function of this variable. Our results lead to a new view on rigidity: While at zero strain a rigid crystal coexists with one that responds to infinitesimal strain by rearranging particles and expelling stress, at finite strain the rigid crystal is metastable, associated with a free energy barrier that decreases with increasing strain. The rigid phase becomes thermodynamically stable by switching on an external field, which penalises particle rearrangements. This produces a line of first-order phase transitions in the field - strain plane that intersects the origin. Failure of a solid once strained beyond its elastic limit is associated with kinetic decay processes of the metastable rigid crystal deformed with a finite strain rate. These processes can be understood in quantitative detail using our computed phase diagram as reference.

I. INTRODUCTION

The ability to resist changes of shape, or rigidity, has been explained as a consequence of the spontaneous breaking of continuous translational symmetry in crystalline solids [1, 2]. Quite surprisingly, this result is at the same time, paradoxical. It may be shown quite rigorously [3] that any homogeneous bulk deformation created within a solid to conform to changes of shape of the boundary may always be accommodated instead by surface distortions involving particle rearrangements [4, 5]. This automatically suggests that any internal stress generated *in equilibrium* within a macroscopically large solid in response to a change of shape must necessarily vanish [4]. Given enough time, a solid always flows to release this stress under any external mechanical load, however small [5]. The emergence of rigid solids is therefore associated with inherently long-lived *metastable* states [4, 6]. This fundamental result has, quite understandably, wide ranging connotations in the science and technology of materials, especially with regard to properties like high temperature creep, fatigue, fracture and plastic flow [7, 8].

While the immediate paradox is resolved, we still need to address the question of how a rigid crystal, when deformed, releases internal stress and transforms to a flowing state. A fundamental understanding of this process should also reveal under what conditions thermodynamically stable rigid crystals may exist. Here, a comparison to fluids in the limit of zero strain rate is very instructive. While fluids subjected to small stresses exhibit Newtonian flow with a constant viscosity [1], no such regime exists for stressed solids whose viscosity diverges with vanishing stress [5]. Does this singular behaviour of the viscosity imply an underlying phase transition? Moreover, distinct from the fluid state, flow in a crystal is triggered by the formation of slip planes [4] causing rearrangements of particle neighbourhoods. Rigid solids composed of distinguishable particles are thus also associated with the breaking of *discrete* permutation symmetry.

In this work, we show, to the best of our knowledge for the first time, that a phase transition indeed occurs at zero strain rate. A static, *equilibrium*, first-order phase transition describes the transformation of one crystal to another with identical crystal structure but with differently arranged local neighbourhoods. Stress relaxation occurs as a consequence of these rearrangements. As expected for a first-order transition, the transformation *kinetics* of the metastable rigid solid to the stable unstressed solid at finite and sufficiently small strain rates may be described by a *nucleation* process. A parameterfree prediction of the strain rate dependent, mean, limiting deformation beyond which this nucleation occurs and a rigid crystal first begins to flow, is one of the verifiable outcomes of our work.

Essential for these findings is the identification of a thermodynamic variable X, the order parameter of the transition, and its conjugated field h_X , which we define shortly. We show that thermodynamically stable rigid solids exist for finite, negative h_X . In addition, we also obtain a line of first order transitions from a rigid solid to a solid state with zero stress. In the thermodynamic limit, this phase boundary extrapolates to $h_X \to 0^-$ giving rise to the aforementioned, experimentally observable

²Institut für Theoretische Physik II: Weiche Materie,

transition associated with stress relaxation. We thus follow here a procedure analogous to many other condensed matter systems (for a classic example see [9]) wherein deeper insight is obtained, leading to quantitative predictions, by first introducing a field h_X and then letting $h_X \to 0$ after taking the thermodynamic limit.

Consider, therefore, a (single phase) crystalline solid completely enclosed by a deformable boundary. The solid is composed of macroscopic, classical particles, e.g. a colloidal crystal [10]. Our main conclusions are summarised in Fig. 1 where we plot a schematic phase diagram of the crystalline solid under changes of boundary shape, parametrised by a pure shear or uniaxial strain ε , and h_X . There are two distinct ways in which the solid may respond to ε , either resisting it by producing internal stress (the rigid "normal" \mathcal{N} phase) or deforming plastically to conform to the shape of the boundary, expelling stress from the bulk (the "Meissner" \mathcal{M} phase [11, 12]). Since the \mathcal{M} phase deforms by slipping over an integral number of lattice spacings, crystallinity is preserved. The resulting \mathcal{M} solid is *structurally identical* to the undeformed \mathcal{N} crystal, save for the presence of surface steps. Each slip line, however, leaves in its wake a set of particles whose neighbourhoods have been *rearranged*. The field, h_X , assigns a bulk free energy cost for these rearrangements in a manner we describe below, and explicitly breaks the discrete, permutation symmetry, causing a first order transition and $\mathcal{N}-\mathcal{M}$ phase coexistence. The phases co-exist across a phase boundary which extrapolates to $\varepsilon = 0$ as $h_X \to 0^-$. The \mathcal{N} phase is metastable for all ε on the $h_X = 0$ line and eventually decays by a nucleation process [5] with an ε -dependent rate. For fixed observation time, this decay process manifests itself as a sudden drop of stress at some $\varepsilon = \varepsilon^*$, where plasticity initiates [7]. This dynamical transition point extends into a smooth transition line in the $h_X - \varepsilon$ plane that intersects the $h_X = 0$ axis at the observable value, ε^* . We show that thermodynamic parameters obtained from our equilibrium study may then be used to predict time-dependent, dynamic properties of this transition.

The field couples to a collective coordinate X = $N^{-1}\sum_{i=1}^{N} \chi_i$ in the Hamiltonian $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_X = \mathcal{H}_0 - Nh_X X$, with \mathcal{H}_0 representing terms in the Hamiltonian that do not depend explicitly on X. For particle i, the local positive definite quantity χ_i with dimensions of length squared is the least squares error [13] made by fitting a local affine deformation to relative displacements within a coarse graining volume surrounding particle i. The deformation is measured from a set of fixed reference coordinates. In a series of papers [14–19] (see also Appendix A for details) some of us have worked out in detail the statistical thermodynamics of χ_i , which quantifies the "non-affine" component of the particle displacements, analytically and numerically for a number of two dimensional (2d) crystals at finite temperature. At any T > 0, X behaves as a regular thermodynamic variable with a well-defined mean and variance $\sim N^{-1}$. Apart from this thermal contribution, X also tracks local non-



FIG. 1. Schematic phase diagram in the h_X - pure shear, ε , plane showing regions of stability of the two (initially square) crystalline phases \mathcal{N} (orange region) and \mathcal{M} (blue region). Insets show ε acting on a square (dashed line) box (1) and depict the \mathcal{N} (2) and \mathcal{M} (3) phases schematically; black dashed lines with end dots denote slip planes. The equilibrium phase boundary is shown as a blue solid line and the locus of the $\mathcal{N} \to \mathcal{M}$ dynamical transition as a blue dashed line. The intersection of the dashed line with the $h_X = 0$ axis is the conventional, strain rate dependent, yield point (grey filled circle).

affine rearrangements of particles such as those resulting from the creation of defects [15, 19]. The ensemble average $\langle X \rangle$ can be tailored using h_X consistent with standard fluctuation response relations [15] and h_X can also modify the probability of defects. Note that since X is defined in terms of relative displacements, the term proportional to h_X in \mathcal{H} does not explicitly break translational invariance [20].

Positive values of h_X help create non-affine rearrangements away from the reference configuration. Specific rearrangements, such as a slip by a lattice spacing, map the crystal onto itself and do not change lattice symmetry but still contribute to the energy \mathcal{H} for non-zero h_X . Since X has an upper bound $\sim L^2 \propto N^{2/d}$ where L is a typical linear size, \mathcal{H} and the corresponding free energy is unbounded below ($\sim -N^{1+2/d}$) in the thermodynamic limit. Therefore, there is no well-defined global free energy minimum for $h_X > 0$ although multiple local minima may exist as long-lived metastable states. At $h_X = 0$, of course, all states differing only in their value of X are degenerate.

On the other hand, a negative h_X suppresses rearrangements and makes the reference configuration the thermodynamically stable phase at $\varepsilon = 0$. As ε increases at constant h_X , there is a possibility of an equilibrium first order transition which may be understood from the following T = 0 argument. In the \mathcal{N} phase, ε is the elastic strain [1] and the bulk energy density is $\Delta E = \frac{1}{2}\sigma\epsilon = \frac{1}{2}G\epsilon^2$, where σ is the elastic stress and G is an elastic modulus. In the \mathcal{M} crystal non-affineness proportional to $|\varepsilon, X = \ell^2 |\varepsilon|$ with some length-scale ℓ , is produced instead by slipping of lattice planes and $\Delta E = -\rho h_X \ell^2 |\varepsilon|$, except for a surface contribution arising from steps that are formed as a consequence of the slips. Here, ρ is the number density of the solid.



FIG. 2. Plots of the dimensionless free energy, $-\log P(X)$, at T = 0.8 and density $\rho = 1.1547$ (i.e. lattice parameter a = 1.0) as a function of h_X at fixed $\varepsilon = 0.04$ (a) and as a function of ε at fixed $h_X = -3.0$ (b.) as obtained from sequential umbrella sampling of 2d LJ crystals for N = 1024. The minima at small (large) X values represent the $\mathcal{N}(\mathcal{M})$ phase. A first order $\mathcal{N} \to \mathcal{M}$ transition occurs as $-h_X$ or ε is increased. The numbers on the graphs correspond to values of either h_X (a) or ε (b). Note the high barriers ($\sim 50k_BT$) between the phases at coexistence.

Note that X = 0 in the \mathcal{N} solid [14] while $\sigma = 0$ in the \mathcal{M} solid at coexistence. Equating, we get the \mathcal{N} — \mathcal{M} coexistence boundary as $-h_X = G|\varepsilon|/2\rho\ell^2$ (Fig. 1). The "thermodynamic stress" $\varsigma = \partial_{\varepsilon}\Delta E \neq 0$ for both \mathcal{N} and \mathcal{M} phases.

How do entropic contributions alter these arguments? Consider the case $\varepsilon = 0$ and $h_X < 0$. Permuting particles over a distance R should give a configurational entropy gain scaling as $\sim N \log R$. The energy penalty for this, given that $X \sim R^2$, will be $\sim |h_X|NR^2$. Minimizing the free energy $-TN \log(R/a) + N|h_X|R^2$ gives $R \sim (T/|h_X|)^{1/2}$. Once R falls below some fixed small value (say, the lattice constant a) this argument breaks down, implying that for $|h_X| > \text{const} \times T$ entropic effects may be neglected and the T = 0 considerations hold. For smaller $|h_X|$, however, rearrangements are possible up to a cutoff distance $\sim (T/|h_X|)^{1/2}$ that diverges as $h_X \to 0^-$. Once rearrangements are thermodynamically favoured, kinetic considerations become important. In particular, if rearrangements happen only by diffusion, then the associated *timescales* in solids are very large [7] so that spontaneous transitions between different free energy minima, corresponding to distinct rearrangements, will become effectively unobservable within realistic times. We return to the question of dynamics later.

Up to now, our discussion has been quite general and works for any crystal in any dimension. We now specialise to the case of the crystalline 2d LJ solid to study the $\mathcal{N}-\mathcal{M}$ transition at T > 0 in detail. Accurate numerical results may be obtained for this case within reasonable computational times. Further, our results have experimental consequences. These may be relatively easily verified for 2d colloidal crystals [10], for which a LJ interaction is a plausible model.

II. THE EQUILIBRIUM FIRST ORDER PHASE TRANSITION

We use a shifted and truncated LJ potential (see Appendix B) and standard LJ units for length, energy and time [21]. For our results of the equilibrium structures and transitions at T > 0 we employed the sequential umbrella sampling (SUS) technique (see Appendix C) coupled to Monte Carlo [22, 23] in the constant number N, area $A = L_x \times L_y$, ε and temperature T ensemble. Advanced sampling techniques such as SUS are necessary to overcome the large barriers between the \mathcal{N} and \mathcal{M} phases, enabling the equilibrium transition to be observed [17]. We show results for T = 0.8 and density $\rho = 1.1547$, corresponding to the choice a = 1.0 for the lattice parameter. Other T and ρ far from the 2d LJ melting line give similar results. Finally, our results for finite N are extrapolated to draw conclusions on the equilibrium transition in the thermodynamic limit.

The main output of the SUS calculations is an accurate estimate of P(X), the equilibrium probability distribution of X as a function of h_X and ε . In Fig. 2a and b we plot $-\log P(X)$, the free energy in units of $k_B T$ where k_B is the Boltzmann constant. To obtain these results we use the efficient histogram reweighting method [24, 25] starting from a few chosen h_X and ε . The two minima at $X_{\mathcal{N}}$ and $X_{\mathcal{M}}$ correspond to the two competing phases, with a first order transition from \mathcal{N} to \mathcal{M} occurring as a function of either h_X or ε . The barrier between the phases at coexistence is high and hence the phase transition is impossible to observe using standard simulation techniques. The large value of $X_{\mathcal{M}}$ results from a finite density of percolating slip bands with large local χ . Configurations for $X_{\mathcal{N}} < X < X_{\mathcal{M}}$ at coexistence show mixed phases similar to other systems with first order transitions [26].

In Fig. 3a we show a mixed phase configuration at coexistence where a portion of the solid slips locally, decreasing stress. This is apparent from the map of local χ values, which are largest at the slip band. The slip band is composed of a "proto" dislocation dipole with a large overlap between the individual defect cores [7] lying on one of the close packed atomic lines of the triangular lattice. Scanning over configurations for X between $X_{\mathcal{N}}$ and $X_{\mathcal{M}}$ reveals a slip band of increasing linear size until it percolates the whole solid, wraps about the periodic boundaries a few times, commensurate with the aspect ratio of the box and finally annihilates with itself at $X \approx X_{\mathcal{M}}$. In a periodically repeated scheme, therefore, this configuration corresponds to a finite slip band density $\sim \mathcal{O}(1)$. The proportion of the two phases follow a lever rule typical of first order transitions [26]. The strongly first order nature of the transition is obvious from the equations of state X vs h_X at fixed ε obtained by plotting the expectation value of X computed from the P(X) in Fig. 3b. The phase transition for each ε is shown by a horizontal tieline, and labelled by ε on the right axis. The end points of the tielines thus also give



FIG. 3. **a.** Plot of the local $log(\chi)$ values (colour map) superimposed on the particle displacements (grey arrows) for a N = 1024 2d LJ solid at $\varepsilon = 0.05$ and $h_X^{\text{coex}} = -5.07$. The data was averaged over 500 configurations corresponding to X = 0.0358 where a mixed phase configuration is observed. Note the presence of a slip band (red dashed line) with large χ . Inset shows δu_{\parallel} , the relative displacement of particles along the direction of the slip band, consistent with that expected from a pair of dislocations with opposite signs and large overlapping cores. **b.** Equations of state $-h_X$ (left axis) vs X. These are shown for several fixed ε but overlap to a large extent, except where jumps in X produce horizontal tielines (labelled by the corresponding ε on the right). The large jump between the coexisting phases indicates a strongly first order transition. c. Plot of the local stress σ (colour map) superimposed on the particle positions (grey circles) for the same set of configurations as in (a) showing a prominent stress interface (red dashed lines). **d.** Plot of the stress $\bar{\sigma}(x)$ averaged in the vertical y direction for three values of ε along the phase boundary. The interface between the \mathcal{N} phase (high stress) and the \mathcal{M} phase (low stress) is clearly visible.

the X- ε phase diagram. While the discontinuity in X at the transition appears to decrease with increasing ε , it cannot vanish since a slip band always creates local χ .

The nature of the \mathcal{M} phase for a finite sized box is complex. While a deformation ε needs a linear density of slip bands $\sim |\varepsilon|/a$ for complete stress relaxation, the number of slip bands actually observed depends on the size as well as the shape of the box. For example, if the required linear density of slip bands is below L^{-1} then no bands can be accommodated and stress cannot be relaxed, i.e. the ${\mathcal M}$ phase cannot exist even if it is the stable phase in the thermodynamic limit. In general, there may be multiple minima in X associated with distinct fami*lies* of parallel slip bands at crystallographically allowed angles. At large ε , these further transitions involving additional slipping occur at values of X higher than those shown here. Phases with a larger density of slip lines have lower σ , down to $\sigma = 0$. The exact sequence of these higher order transitions depends on the details of

the simulation box. We do not pursue this here as the deformation *dynamics* and departure from rigidity will be determined by the kinetics of the first transition.

In Fig. 3c we return to the mixed phase configuration shown in **a** and study the local internal stress σ conjugate to ε superimposed on the particle positions. A prominent *interface* between the two coexisting phases is clearly seen. The \mathcal{M} phase eliminates stress from its bulk by particle rearrangements, i.e. slip, as expected, while stress is retained in the bulk of the \mathcal{N} phase. As the amount of the second phase grows, total stress is proportionately reduced.

Capillary fluctuations of this interface [26] around the mean position are also seen in Fig. 3c. Averaging the local stress $\bar{\sigma}(x)$ in the vertical direction and plotting it as a function of the horizontal coordinate x reveals an interface where these capillary fluctuations are averaged out. This is shown in Fig. 3d. We have plotted $\bar{\sigma}(x)$ for a few values of ε on the phase boundary. While the jump in σ decreases with ε , the interface remains, nevertheless, sharp.

We must emphasise here that this is an *equilibrium* interface between two *co-existing* phases with different values of σ but both with bulk crystalline order. Such stable interfaces do not form at $h_X = 0$ for $\varepsilon \neq 0$ in the thermodynamic limit and have, therefore, never been described before.

A. Finite size scaling

We carry out a finite size scaling [22, 26] analysis at $\varepsilon = 0.05$ to establish that the transition between the two phases is indeed first order. At a first order transition, corrections to order parameters and to the transition point scale as $\sim L^{-d} \sim 1/N$ [26]. This is apparent from Fig. 4a i-iii where the coexisting h_X^{coex} , the values of $\langle X \rangle$ and stress σ for the two phases show the expected scaling behaviour. Finite size corrections to the properties of the \mathcal{M} phase are observed to be quite substantial due to the commensurability issues discussed before. To show that the free energy cost of creating two parallel interfaces (lines) scales as $L^{d-1} = \sqrt{N}$, we plot $-\ln P(X)$ obtained at coexistence for different N using scaled coordinates (Fig. 4 **a** iv). The region corresponding to mixed $\mathcal{N} - \mathcal{M}$ configurations collapses onto a single horizontal line as expected. Finally the T > 0 equilibrium phase boundary in h_X and ε is shown in Fig. 4b. for different N. The phase boundary is quite linear showing that our rather simplistic T = 0 calculation gives a qualitatively correct result. The offset in ε at $h_X \to 0^-$ (~ ε produced by a single slip band) is expected to vanish in the thermodynamic limit. For a thermodynamically large solid, we obtain phase coexistence only for $h_X \leq 0$ with the phase boundary intersecting the origin i.e. $h_X = \varepsilon = 0$.



FIG. 4. **a.** Finite size scaling at $\varepsilon = .05$ of the value of h_X^{coex} (i), the coexisting X_N, X_M at the two minima in $-\log P(X)$ (ii), the coexisting stress values across the $\mathcal{N} - \mathcal{M}$ interface (iii) and collapse of the scaled surface free energy in $-\log P(X)$ (iv). **b.** The $h_X - \varepsilon$ phase boundary showing that as $N \to \infty$, the boundary tends to intersect the origin (schematic: orange and blue shaded regions and blue line as in Fig. 1).

1

This completes our description of the *equilibrium* first order, $\mathcal{N} - \mathcal{M}$ phase transition. The transition is *reversible* with ε , a thermodynamic variable, being applied quasi-statically, i.e. $\dot{\varepsilon} = 0$. We study below the implications of the equilibrium transition on the dynamics of deformation ($\dot{\varepsilon} \neq 0$).

III. NUCLEATION DYNAMICS AND PLASTIC DEFORMATION

In the limit of $h_X \to 0^-$, a macroscopically large rigid (\mathcal{N}) solid is metastable for all $\varepsilon > 0$. This is illustrated in Fig. 5a for a N = 1024 solid where we plot $-\log P(X)$ as a function of ε extrapolated to $h_X = 0$. As ε increases, therefore, the \mathcal{N} solid form (and grow) within the body of the \mathcal{N} phase. At the end of this process, the equilibrium, stress free, \mathcal{M} crystal thus formed is *identical* in all respects to the unstrained \mathcal{N} crystal and with its other infinitely many copies differing only by their values of X. We show below how the free energies calculated using SUS may be used to study the dynamics of this nucleation process in quantitative detail.

A. Nucleation barriers

Following standard classical nucleation theory [1, 5, 27, 28] (CNT) we write the dimensionless excess free energy of a configuration containing a circular droplet of the

 \mathcal{M} phase of size R surrounded by the stressed \mathcal{N} crystal as $\Delta \mathcal{F} = -\pi R^2 \Delta_b + 2\pi R \gamma$. The first term represents the bulk free energy gain, which assuming that the \mathcal{M} solid is stress free, gives $\Delta_b = \frac{1}{2}\sigma\varepsilon/(k_BT)$. The second term involves the equilibrium interfacial free energy γ between \mathcal{N} and \mathcal{M} phases. Since equilibrium interfaces exist only at coexistence [26, 27], we need to obtain γ from $-\log P(X)$ along the phase boundary. The height of the horizontal region in $-\log P(X)$ (see Fig 4 **a** iv) relative to the depth of the minima is given by $2\gamma L_{\mu}$ + subdominant (~ $\mathcal{O}(\log L)$ etc.) contributions. Factoring out the length of the pair of parallel interfaces then gives the *finite size scaled* value for γ , which is shown in the inset of Fig. 5b as a function of ε for various N. Since the subdominant contributions are small and unobservable, we use a linear fit through the data to obtain $\gamma(\varepsilon)$ along the phase boundary. Extrapolation to $\varepsilon = 0$ gives $\gamma = 1.8 \pm 0.1$ as the surface free energy of the infinite solid, which enters the CNT calculation for nucleation of the \mathcal{M} phase at $h_X = 0$. The size of the critical nucleus is $R_c = \gamma / \Delta_b$ and the nucleation barrier is $\Delta \mathcal{F} = \pi \gamma^2 / \Delta_b$. The latter is plotted as a function of ε in Fig. 5b as a solid curve. The free energy barrier has been approximately modelled earlier using specific, correlated defect structures such as arrays of dislocation loops [5, 29, 30], at $h_X = 0$. The full dynamical problem of many interacting dislocations in crystals is complex and remains unsolved [7, 30], necessitating many simplifying assumptions. In our description γ is obtained without assuming any specific dislocation structure.

We may now obtain the mean first passage nucleation time as $\tau_{FP} = \tau_0 \exp(\Delta \mathcal{F})$, in the limit of large $\Delta \mathcal{F}$ where τ_0 is a relevant time scale [1, 28]. Formally, τ_0 is the time taken for nucleation when the barrier vanishes, but this interpretation is problematic because in that limit the nucleation picture itself fails. We show later how

 $^{^1}$ The phase diagram in the h_X and ς plane can also be drawn using our SUS data but does not contain substantial new information.



FIG. 5. **a.** Equilibrium free energy from SUS for a N = 1024 solid for various ε at $h_X = 0$ for comparison. **b.** The nucleation barrier $\Delta \mathcal{F}$ (solid purple line) as a function of ε , the cyan lines mark the uncertainties of our results. Inset shows the interfacial free energy γ for various N obtained from SUS (symbols have same meaning as in Fig. 4b) plotted against ε together with the linear fit $\gamma = -12.420\varepsilon + 1.806$.

this τ_0 may be extracted from molecular dynamics (MD) data. Since the \mathcal{N} phase is always metastable, τ_{FP} is finite for all $\varepsilon > 0$ and diverges as $\varepsilon \to 0$. Thus, if one waits long enough, a transition from $\mathcal{N} \to \mathcal{M}$ is inevitable at any $\varepsilon \neq 0$. We compare the CNT estimate to measured nucleation times in MD simulations below.

B. Molecular dynamics

We perform MD simulations [21, 31] (see also Appendix D) for $128 \times 128 = 16384$ LJ particles in the NA(shape)T ensemble at the same density as the SUS calculations for 0.2 < T < 1.2. The direct way of comparing SUS and MD is to compute transition times by holding the solid at various strain values. This protocol has technical issues because applying a finite strain suddenly to a solid causes transient shock waves that make extraction of meaningful data impossible. In the molecular dynamics simulations, therefore, the strain is ramped up from zero in steps of $\Delta \varepsilon$, waiting for a time t_W to obtain an average strain rate $\dot{\varepsilon} = \Delta \varepsilon / t_W$. We look for a drop in stress to mark the beginning of plasticity at the yield point ε^* .

C. X as a reaction coordinate

To proceed any further, we must first establish that the plastic event at ε^* does indeed represent an $\mathcal{N} \to \mathcal{M}$ transition described by X. In other words we need to show that X is also the relevant *reaction coordinate* for deformation.

The $\sigma(\varepsilon)$ curves obtained for various h_X , including $h_X = 0$, at a fixed value of $\dot{\varepsilon}$ are shown in Fig. 6a. These show a large stress drop at $\varepsilon = \varepsilon^*$ while $\langle X \rangle$ increases at the same value of strain indicating that particle re-



FIG. 6. **a.** Plots of σ (axis on left) and $\log\langle X \rangle$ (axis on right) as a function of ε from MD simulations at $\dot{\varepsilon} = 3.33 \times 10^{-5}$. Note that the jumps in σ and $\langle X \rangle$ at the dynamical transition coincide. The dashed line shows the limiting value $X_L = 0.0165$ before the transition. **b.** Plot of the $\mathcal{N} - \mathcal{M}$ dynamic transition in the (ϵ, h_X) plane. The open symbols mark the dynamical transition $\epsilon^*(h_X)$ predicted from LRT calculations assuming $\langle X \rangle \to X_L$. The black curve is a parabola fitted through the points. The red filled circles are transition points obtained from MD simulations. **c.** Scaling of X_L with temperature showing that X_L/T is a constant.

arrangements occur. The angular brackets here denote a time average as well as an average over several initial conditions. The jump in $\langle X \rangle$ with a simultaneous drop in stress indicates that these particle rearrangements, at the same time, relieve stress. We find that $\langle X \rangle$ always attains the same value $\langle X \rangle = X_L$ with $X_N < X_L < X_M$ just before the transition regardless of h_X . As the solid is strained, $\langle X \rangle$ increases; when it reaches X_L , enough thermal energy is available to the solid in order to cross the $\mathcal{N} \to \mathcal{M}$ barrier and the \mathcal{M} phase begins to nucleate.

The \mathcal{N} phase decays if either h_X or ε is increased. If X is the relevant coordinate for this transition, then this single quantity should describe yielding regardless of h_X and ε . In other words, $\langle X(h_X, \varepsilon) \rangle = X_L$ should trace out a unique curve in $h_X - \varepsilon$ space, beyond which the \mathcal{N} crystal decays. Since $\langle X \rangle$ grows linearly with h_X to leading order and quadratically with ε in the \mathcal{N} phase [14], this curve is a parabola. In the \mathcal{N} phase, we use linear response theory [15] to obtain the h_X values where $\langle X \rangle \to X_L$ for any given ε . To do this we record fluctuations of X and spatial correlations of χ at $h_X = 0$ for a series of ε values starting from zero. Linear response then gives $\langle X(h_X, \varepsilon) \rangle =$ $\langle X(0,\varepsilon)\rangle + h_X \langle [\Delta \chi(0,\varepsilon)]^2 \rangle \Sigma_{\mathbf{R}} C_{\chi(0,\varepsilon)}(\mathbf{R},0) = X_L$, where $C_{\chi(0,\varepsilon)}(\mathbf{R},0)$ is a two point correlation function. The predicted values for the location of the dynamical $\mathcal{N} \to \mathcal{M}$ transition, which follow the expected parabolic relation, are given as open symbols in Fig 6b. Note that we now predict the location of a *plastic* event based on *equilibrium* thermal fluctuations of a thermodynamic variable in configurations corresponding to small ε (and $h_X = 0$) where dislocations or other defects may not even



FIG. 7. **a.** Stress - strain curves from MD at T = 0.8 at $h_X = 0$ for four $\dot{\varepsilon}$ varying over three decades. **b.** The calculated value for the yield strain ε^* as a function of the strain rate $\dot{\varepsilon}$ in LJ units. The lines have the same meaning as in Fig. 5**b**. The points are from our MD results. Inset shows a plot of $\log_{10} \tau_0^{-1} = \log_{10} (\dot{\varepsilon}/\tilde{\varepsilon})$ as a function of $\dot{\varepsilon}$ obtained from our MD. We have used a value of $\log_{10} \tau_0^{-1} = 2.0$ as obtained from the plateau value in the inset to compare our MD data with the SUS predictions.

be present. The filled symbols in Fig. 6b show $\epsilon^*(h_X)$ obtained from MD. The extrapolation of the calculated curve to $h_X = 0$ agrees extremely well with the measured value. At large negative h_X the linear response prediction ceases to be valid. Given that in the \mathcal{N} -phase thermal fluctuations are primarily responsible for making X non-zero [14], X_L should scale with temperature. Fig. 6c shows that on varying only T, this expectation is justified and $X_L(T)/T$ is a constant. We expect this behaviour to be generic and easily verifiable in experiments on colloidal solids [10] constituting a stringent test for our theory.

Since ε is ramped up from zero at a fixed rate, $\dot{\varepsilon}$, the dynamical transition i.e. the values of ε^* and X_L , depend on $\dot{\varepsilon}$. We show now that this rate dependence may be predicted using parameters extracted from the equilibrium phase transition.

D. Self consistent classical nucleation theory

We are now in a position to compare predictions of CNT with MD results. In Fig 7a, we plot $\sigma(\varepsilon)$ curves at $h_X = 0$ for four deformation protocols where $\dot{\varepsilon}$ varies over three decades. The yield point ε^* is a function of $\dot{\varepsilon}$ and appears to vanish as $\dot{\varepsilon} \to 0$. This result is consistent with the \mathcal{N} phase being metastable for all $\varepsilon > 0$.

To obtain predictions for ϵ^* using the barriers obtained from our SUS calculations, however, we need a small modification of the theory. For the protocol followed in MD, the barrier is not constant, but varies as the strain is ramped up in time. If the variation is smooth and slow, we may use the self consistent formula $\tau_{FP} = \tau_0 \exp[\Delta \mathcal{F}(\dot{\epsilon} \tau_{FP})]$ [32]. Noting that $\dot{\epsilon} \tau_{FP} = \varepsilon^*$, we get a self consistency equation for ε^* which needs to be solved numerically,

$$\Delta \mathcal{F}(\varepsilon^*) = \log \varepsilon^* - \log(\tilde{\varepsilon}),$$

where $\tilde{\dot{\varepsilon}} = \tau_0 \dot{\varepsilon}$. This approximation should be valid in the small $\dot{\varepsilon}$ regime. Since small $\dot{\varepsilon}$ gives small ε^* , and therefore large barriers, the regimes of validity of the self consistent approximation and CNT itself coincide.

In principle τ_0 could be taken as a fitting parameter which links energies to time scales. But fitting τ_0 to MD data is uncertain because the range of validity of CNT is not known a priori. It has also been interpreted as the time taken for a dislocation dipole (in 2d) or loop (3d) to form [5], although obtaining an estimate for τ_0 using this interpretation requires additional assumptions. The nucleation rate of dislocations is a technologically important quantity and has been measured using experiments and computer simulations [33–36]. For a real 3d solid, the pre-factor can be written as $\tau_0^{-1} = \Gamma f_c^+$ where Γ is the Zeldovich factor and f_c^+ the "attachment" rate [28]. These have been estimated for Cu single crystals and give $\tau_0 = 3.43 \times 10^{-14}$ s [36]. This number is also consistent with τ_0^{-1} compared with typical Debye frequencies $\omega_D \sim 10^{13}$ Hz [1]. In LJ units, where the unit of time is about a pico-second (10⁻¹²), we obtain $\tau_0 = 0.0343$. Below we show how τ_0 can be alternatively obtained by appealing to the internal consistency of CNT without using dislocation nucleation times as input. Remarkably, we also determine, at the same time, the range of validity of CNT for our MD data.

From our self consistent CNT theory we obtain $\dot{\varepsilon}$ for each ε^* obtained from MD. We now use the fact that τ_0 should be independent of $\dot{\varepsilon}$ if CNT is valid and obtain $\dot{\varepsilon}/\dot{\varepsilon}$ for each ε^* . This is shown in the inset of Fig. 7b. Note that for small $\dot{\varepsilon}$ we obtain a plateau in the values of $\tau_0 = 0.01$ thus calculated. Deviations from the plateau value begin from $\dot{\varepsilon} \approx 10^{-5}$ or $\varepsilon^* \approx 0.11$. Comparing with Fig.5b we observe that this corresponds to a barrier height of about 10 - 20 k_BT, which is completely consistent with expectations. The data from MD is compared with the results of the self consistent theory, using τ_0 obtained from the plateau value in the limit $\dot{\varepsilon} \to 0$, in Fig. 7. Our SUS and MD data are in excellent agreement for the smallest $\dot{\varepsilon}$ values showing that the decay of the metastable \mathcal{N} solid sets the time scale for microscopic processes responsible for stress relaxation. Our $\varepsilon^*(\dot{\varepsilon})$ curve therefore is a prediction for the yield point at strain rates that are relevant for slow deformation of solids under experimental conditions [7]. Such processes are impossible to probe in standard MD simulations and it is remarkable that SUS allows us access to these regimes. Although we have presented results for a single temperature T = 0.8, our predictions follow from the identification of X as the reaction coordinate, with X_L as its limiting value in the \mathcal{N} phase. Since X_L scales simply with T, we expect our $\varepsilon^*(\dot{\varepsilon})$ to do the same as long as the temperature (and density) is not close to melting. Note that there are no adjustable parameters in our calculation.

IV. DISCUSSION AND CONCLUSIONS

We began our investigation by asking whether rigid solids can ever exist as a stable thermodynamic phase. We introduced a new collective variable X that keeps track of non-affine particle rearrangements by comparing positions of distinguishable particles with a set of reference coordinates. By turning on a field h_X conjugate to X one can bias particle rearrangements and explicitly break permutation symmetry. We show that the breaking of this discrete symmetry leads to a first order transition and phase coexistence between a rigid solid and one where particles rearrange to eliminate stress. The first order transition is quite conventional in all respects and both bulk and surface properties of the coexisting phases scale in the expected manner. By measuring these properties in the presence of this fictitious field and subsequently taking both the thermodynamic limit and the limit $h_X \rightarrow 0$ we were able to obtain quantities that predict the dynamics of deformation of the solid in the absence of field without any fitting parameters.

Irreversibility of plastic deformation is easy to understand. At large strain rates the product state is not in equilibrium and the process is irreversible. However, deforming a solid even with $\dot{\varepsilon} \to 0$ causes irreversibility if $h_X = 0$, due to the nature of the free energy landscape. Consider the reverse transformation from $\mathcal{M} \to \mathcal{N}$. Straining the \mathcal{M} solid in the reverse direction now increases its free energy and it decays to a stable unstressed state. However all rearranged versions of the original \mathcal{N} crystal have the same free energy and are equivalent candidate products. The solid at the end of the process is likely to reach one of these states instead of the original crystal, with overwhelmingly large probability, producing irreversible particle rearrangements. The large degeneracy of phases at $h_X = 0$ thus makes the reverse transformation non-unique and deformation irreversible. This is, of course, not true if $h_X < 0$ where equilibrium transformations between \mathcal{N} and \mathcal{M} are always reversible.

So far, h_X has been introduced as a device for understanding the relation between non-affine particle displacements and deformation in a solid with contact being made with experiments only at $h_X = 0$. For solids where individual particles can be distinguished and tracked, one should be able to realise h_X in the laboratory and check our predictions in the full $h_X - \varepsilon$ plane. Indeed, it has already been discussed in detail [15, 17, 20] how this may be accomplished in the future for colloidal particles in 2d using dynamic laser traps. Briefly, the set of reference coordinates is read in and a laser tweezer is used to exert additional forces, $\mathcal{F}_{\chi}(\mathbf{r}_i) = -\partial \mathcal{H}_X / \partial \mathbf{r}_i$ to each particle i which bias displacement fluctuations. Since the additional forces depend on *instantaneous* particle positions, they need to be updated continuously through real time particle tracking. This is possible because timescales of colloidal diffusion are large [10]. We believe that this procedure will be achievable in the near future using current

video microscopic and spatial light modulation technology [37]. Colloidal \mathcal{N} crystals stabilised under an artificially produced h_X field should show new and interesting properties, such as high failure strengths with small elastic constants or vice versa, resistance to creep, very small defect concentrations etc. These properties, coupled to the fact that they are reversible and may be switched on or off or precisely tuned, may have some technological use.

At large times, after the first plastic event at ε^* , a crystal under a constant deformation rate $\dot{\varepsilon}$ reaches a non-equilibrium steady state. Flow of a crystal with a vanishing strain rate $\dot{\varepsilon} \to 0$ may be understood as a succession of, perpetually occurring, $\mathcal{N} \to \mathcal{M}$ nucleation events, which cause deformation while attempting to reset stress to zero [5, 8]. In this regime, $\dot{\varepsilon} = \sigma/(G\tau_{FP})$, which may be interpreted as flow with a viscosity that diverges as $\sigma = G\varepsilon \to 0$. At large strain rates, $\Delta \mathcal{F} \to 0$ so that nucleation ceases to be the relevant dynamical process. The \mathcal{N} solid decays to a truly non-equilibrium steady state unrelated to \mathcal{M} . Further, in this regime one obtains critical like behaviour and scale-free avalanche driven deformation [38–42]. Our work suggests that such behaviour need not be universal but a consequence of large strain rates used in those studies.

Our work is easily generalised to other realistic crystalline solids in 3d, for example Cu, Au or Al single crystals for which a wealth of experimental data already exists [34–36]. We can obtain γ by turning on h_X and subsequently set $h_X \rightarrow 0$ after taking the thermodynamic limit to predict $\varepsilon^*(\dot{\varepsilon})$ for these solids without adjustable parameters. The introduction of h_X , of course, does not depend on any particular model interaction used and can be implemented within any simulation scheme. It also does not change any of the elastic properties of these solids. Work along these lines is in progress and will be published elsewhere.

In a departure from *all* known literature on plasticity of crystalline solids [5, 7, 8, 30, 34–36, 38–42], no property of dislocations such as core energies or dislocation - dislocation interactions [7] etc. enter our discussion.² This fact imparts a greater range of applicability to our work than those explicitly based on the language of dislocations. We have used almost the same language to describe pleating of two dimensional sheets modelled as a network of permanently bonded vertices [17, 43]. In such systems dislocations cannot form at all, although non-trivial fluctuations in the form of pleats can still be described using non-affine displacements. Similar phase coexistence between a stressed network and one where stress is relaxed by pleating is observed.

 $^{^2}$ The equilibrium ${\cal N}$ and ${\cal M}$ phases are both defect free. For very large systems and at elevated temperatures a small defect concentration is expected on entropic considerations. Kinetic jamming effects may also contribute to increase the defect concentration in the ${\cal M}$ phase relative to ${\cal N}.$

A collective variable, similar to X used by us, was initially defined to characterize local particle rearrangements in glasses under deformation [13, 44, 45]. So our method, with perhaps a few modifications and/or generalisations, could be also applied to amorphous solids. The response of an amorphous solid to a deformation in the zero strain rate limit ($\dot{\varepsilon} \rightarrow 0$) is, however, expected to be very different from that of crystals. In this case, two scenarios are possible: (i) For $\dot{\varepsilon} = 0$, the system is not a glass any more and behaves like a Newtonian fluid. This implies that no broken symmetry is involved, such as the breaking of permutation symmetry, associated with the flow of a crystal. So there is no underlying first-order transition as in the crystal. (ii) The system is in an ideal glass state. As a consequence, there is a nonzero yield stress [46] $\sigma_{\rm y}$, i.e. as a response to any deformation with a given strain rate, the ideal glass state transforms eventually to a flowing state with a finite stress $\sigma \geq \sigma_y$ (at $\dot{\varepsilon} = 0, \ \sigma = \sigma_{\rm v}$). The nature of this transformation is an open issue (see, e.g., Ref. [47]). Also the onset of flow in a glass is different in nature from that of a crystal. While in a crystal flow occurs via the formation of slip bands, in a glass flow is initiated by a percolating cluster of mobile regions [48], associated with a transition in the directed percolation universality class. Furthermore, amorphous solids driven far from equilibrium under external shear stresses exhibit complex deformation behaviour [46–57]. A deeper understanding of all these issues is a challenge that we wish to pursue using our methods in the future.

Appendix A: The projection formalism and the non-affine field

Choose a reference configuration with N particles where particle i (i = 1, ..., N) has position \mathbf{R}_i . Displacing particle i to \mathbf{r}_i the instantaneous position of the particle, produces $\mathbf{u}_i = \mathbf{r}_i - \mathbf{R}_i$. Within neighbourhood Ω around *i*, the relative displacements $\Delta_j = \mathbf{u}_j - \mathbf{u}_i$ with respect to particle $j \neq i \in \Omega$. To obtain the "best fit" [13] local affine deformation D one minimises $\sum_{i} [\mathbf{\Delta}_{i} - \mathsf{D}(\mathbf{R}_{i} - \mathbf{R}_{i})]^{2}$ so that $\chi(\mathbf{R}_{i}) > 0$ is the minimum value of this quantity. This procedure also amounts to taking a projection [14] of Δ_i onto a subspace defined by the projection operator P so that, $\chi(\mathbf{R}_i) = \mathbf{\Delta}^{\mathrm{T}} \mathsf{P} \mathbf{\Delta}$ where we use Δ , the column vector constructed out of Δ_i . In $\mathsf{P} = \mathsf{I} - \mathsf{R}(\mathsf{R}^{\mathrm{T}}\mathsf{R})^{-1}\mathsf{R}^{\mathrm{T}}$, the $Nd \times d^2$ elements of $\mathsf{R}_{j\alpha,\gamma\gamma'} = \delta_{\alpha\gamma} R_{j\gamma'}$ centering Ω at the origin. The global non-affine parameter, $X = N^{-1} \sum_{i}^{N} \chi(\mathbf{R}_{i})$ couples to h_{X} in the Hamiltonian $\mathcal{H} = \mathcal{H}_{0} - Nh_{X}X$, with \mathcal{H}_{0} as the Hamiltonian of any solid. Note that $\mathbf{u}_i \to \mathbf{u}_i + \mathbf{c}$, where \mathbf{c} is an arbitrary translation, remains a symmetry of \mathcal{H} . The statistics of $\chi(h_X)$ and $X(h_X)$ may be computed using standard methods of statistical mechanics [14, 15].

Appendix B: The Lennard-Jones Model

The shifted and truncated LJ model is defined by $\mathcal{H}_0 = \sum_{i=1}^{N} \frac{\mathbf{p}_i^2}{2m} + \sum_{i=1}^{N-1} \sum_{j>i} v_{\text{LJ}}(r_{ij})$ with \mathbf{p}_i the momentum and m = 1.0 the mass of a

with \mathbf{p}_i the momentum and m = 1.0 the mass of a particle. The interaction potential for a pair of particles, separated by a distance r, is $v_{\rm LJ} = 4\phi[(r_0/r)^{12} - (r_0/r_c)^{12} - (r_0/r_c)^6]$ for $r \leq r_c = 2.5r_0$ and $v_{\rm LJ} = 0$ for $r > r_c$. Energy and length scales of the LJ model are set by $\phi = 1$ and $r_0 = 1$, respectively. The unit of time is given by, $\tau = \sqrt{mr_0^2/\phi}$.

Appendix C: Sequential Umbrella Sampling

SUS-MC [21-23] in the NAT ensemble is implemented in a manner identical to that used in Ref. [17] using a periodically repeated rectangular box of dimensions $L_x \times L_y$. The range of X is divided into small windows and sampled successively starting at X = 0. We keep track of how often each value of X within the nth window is realised and the resulting histograms H(n) thus obtained are used to compute the probability P(X). The SUS-MC runs were done for systems with N = 2500, 1600, 1024, 900 and 576 LJ particles at T = 0.8 and the density $\rho = 1.1547$ (a = 1.0). The entire range of X (which varies depending on N) is divided into 800 - 1000 sampling windows with $\approx 1 \times 10^8$ MC moves attempted in each window. In each MC move, maximal displacements of 0.2 a - 0.4 a along the x and y directions are allowed. The SUS-MC computations were done for various h_X and ε . The simulation box is rescaled setting $L_x \to L_x(1+\varepsilon)$ and $L_y \to L_y(1-\varepsilon)$ which keeps the area constant up o linear order. The local stress is obtained from the SUS configurations in the usual way from averaging the virial, taking care of the three body terms implicit in the terms involving h_X [17, 21].

Appendix D: Molecular Dynamics

The MD simulations for the LJ were done for $128 \times$ 128 = 16384 particles within a periodically repeated box identical to that used in the SUS-MC at the same density $\rho = 1.1547$ and several T using a velocity Verlet algorithm [21]. In most simulations, the MD time step $\delta t = 0.001$ in LJ time units; only near yielding, and for $h_X \ge 0.5$, a smaller time step of $\delta t = 0.0001$ is used. In the LJ case, the system is coupled to a Berendsen thermostat [21, 31]. The solid is first equilibrated in the absence of the h_X for t = 500, followed by equilibration runs at different values of h_X , in each case for over t = 1000. Pure shear is applied by rescaling the box in steps of $\Delta \varepsilon = 0.001$ with a waiting time $3 < t_W < 3000$ at each step. The mean strain rate $\dot{\varepsilon} = \Delta \varepsilon / t_W$. For each value of h_X and t_W , 6-8 independent runs were performed.

We thank M. Barma, M. Rao and S. Ramaswamy for discussions. Funding from the FP7-PEOPLE-2013-IRSES grant no: 612707, DIONICOS is acknowledged.

- Chaikin P and Lubensky T (1995) Principles of Condensed Matter Physics (Cambridge Press, Cambridge).
- [2] Szamel G and Ernst MH (1993) Slow modes in crystals: A method to study elastic constants Phys. Rev. B48:112.
- [3] Ruelle D (1969) Statistical Mechanics, (Benjamin, New York).
- [4] Penrose O (2002) Statistical mechanics of nonlinear elasticity Markov Processes Relat. Fields 8:351.
- [5] Sausset F, Biroli G and Kurchan J (2010) Do Solids Flow? J. Stat. Phys. 140:718.
- [6] Shaw S and Harrowell P (2016) Rigidity in Condensed Matter and Its Origin in Configurational Constraint Phys. Rev. Lett.116:137801.
- [7] Phillips R (2004) Crystals, defects and microstructures: Modeling across scales (Cambridge Press, Cambridge).
- [8] Barnes HA (1999) The yield stress a review or 'παντα ρει' - everything flows? J. Non-Newtonian Fluid Mech., 81(1-2): 133-178.
- [9] Griffiths R B (1970) Thermodynamics near the two-fluid critical mixing point in He³-He⁴ Phys. Rev. Lett.24:715.
- [10] Ivlev A, Löwen H, Morfill G and Royall P (2012) Complex Plasmas and Colloidal Dispersions, (World Scientific, Singapore).
- [11] Tinkham M (2004) Introduction to Superconductivity, 2^{nd} Ed. (Dover Publications, New York).
- [12] Toledano P (2007) Theory of the amorphous solid state: Non-directional elastic vortices and a superhard crystal state *Europhys. Lett.* 78:46003.
- [13] Falk ML and Langer JS (1998) Dynamics of viscoplastic deformation in amorphous solids Phys. Rev. E57:7192.
- [14] Ganguly S, Sengupta S, Sollich P, and Rao M (2013) Nonaffine displacements in crystalline solids in the harmonic limit Phys. Rev. E87:042801.
- [15] Ganguly S, Sengupta S, and Sollich P (2015) Statistics of non-affine defect precursors: tailoring defect densities in colloidal crystals using external fields *Soft Matter* 11:4517.
- [16] Mitra A, Ganguly S, Sengupta S, and Sollich P (2015) Non-affine fluctuations and the statistics of defect precursors in the planar honeycomb lattice JSTAT P06025.
- [17] Ganguly S, Nath P, Horbach J, Sollich P, Karmakar S and Sengupta S (2017) Equilibrium and dynamic pleating of a crystalline bonded network J. Chem. Phys. 146:124501.
- [18] Ganguly S and Sengupta S (2017) Excess vibrational modes of a crystal in an external non-affine field J. Chem. Sci. 129:891.
- [19] Ganguly S, Mohanty PS, Schurtenberger P, Sengupta S, and Yethiraj A (2017) Contrasting the dynamics of elastic and non-elastic deformations across an experimental colloidal Martensitic transition *Soft Matter* 13:4689.
- [20] Popli P, Ganguly S and Sengupta S (2018) Translationally invariant colloidal crystal templates Soft Matter 14:104.

- [21] Frenkel D and Smit B (2002) Understanding Molecular Simulations (Academic Press, San Diego).
- [22] Binder K and Heermann D (2010) Monte Carlo Simulation in Statistical Physics: An Introduction, 5th Ed. (Springer, Berlin).
- [23] Virnau P and Müller M (2004) Calculation of free energy through successive umbrella sampling J. Chem. Phys. 120:10925.
- [24] Borgs C and Kotecký R (1990) A rigorous theory of finitesize scaling at first-order phase transitions J. Stat. Phys. 61:79.
- [25] Ferrenberg AM and Swendsen RH (1988) New Monte Carlo technique for studying phase transitions Phys. Rev. Lett.61:2635.
- [26] Binder K (1987) Theory of First-Order Phase Transitions Rep. Prog. Phys. 50:783.
- [27] Statt A, Virnau P and Binder K (2015) Finite-size Effects on liquid-solid phase coexistence and the estimation of crystal nucleation barriers Phys. Rev. Lett.114:026101.
- [28] Becker R, Döring W (1935) The kinetic treatment of nuclear formation in supersatu- rated vapors. Ann Phys (Weinheim, Ger) 24:719?752.
- [29] Bruinsma R, Halperin BI, Zippelius A (1982) Motion of defects and stress relaxation in two-dimensional crystals Phys. Rev. B25:579.
- [30] Bulatov V, Abraham FF, Kubin L, Devincre B, Yip S (1998) Connecting atomistic and mesoscale simulations of crystal plasticity. *Nature*, 391:669.
- [31] Allen MP and Tildesley DJ (1987) Computer Simulation of Liquids (Oxford University Press, Oxford).
- [32] J. Shillcock and U. Seifert (1998) Escape from a metastable well under a time-ramped force *Phys. Rev. E*, 57, 7301.
- [33] Schall P, Cohen I, Weitz DA and Spaepen F(2006) Visualizing dislocation nucleation by indenting colloidal crystals. *Nature* 440:319.
- [34] J. Li, K. J. Van Vilet, T. Zhu, S. Yip, and S. Suresh, Nature (London) 418, 307 (2002).
- [35] J. Li, T. Zhu, S. Yip, K. J. Van Vliet, and S. Suresh, Mat. Sc. and Eng. A365, 25 (2004).
- [36] S. Ryu, K. Kang and W. Cai (2011) Entropic effect on the rate of dislocation nucleation *Proc. Natl. Acad. Sc.* 108, 5174.
- [37] Spalding GC, Courtial J and Leonardo RD (2008) in D. L. Andrews Ed., *Structured Light and its Applications* (Elsevier, Oxford).
- [38] Miguel M-C, Vespignani A, Zapperi S, Weiss J, Grasso J-R (2001) Intermittent dislocation flow in viscoplastic deformation. *Nature*, 410:667.
- [39] Zaiser M (2006), Scale invariance in plastic flow of crystalline solids, Advances in Physics, 55:185.
- [40] Dimiduk DM et al. (2006) Scale-Free Intermittent Flow in Crystal Plasticity. *Science* 312:1188.

- [41] Papanikolaou S (2012) Quasi-periodic events in crystal plasticity and the self-organized avalanche oscillator. Nature, 490:517.
- [42] Tsekenis G, Goldenfeld N, and Dahmen KA (2011) Determination of the universality class of crystal plasticity, Phys. Rev. Lett., 106:105501.
- [43] S. Ganguly, D. Das, Horbach J, Sollich P, Karmakar S and Sengupta S, manuscript in prepraration.
- [44] Falk ML and Langer JS (2010) Deformation and Failure of Amorphous, Solid like Materials. Annu. Rev. Condens. Matter Phys. 2:353.
- [45] Binder K and Kob W (2011) Glassy Materials and Disordered Solids: An Introduction to Their Statistical Mechanics (World Scientific, Singapore).
- [46] Chaudhuri P and Horbach J (2013) Onset of flow in a confined colloidal glass under an imposed shear stress. *Phys. Rev. E* 88:040301(R)
- [47] Barrat J L and Lemaître A (2011), Chapter 8 in Dynamical Heterogeneities in Glasses, Colloids, and Granular Materials, edited by Berthier L, Biroli G, Bouchaud J P, Cipelletti L, and van Saarloos W, (Oxford University Press, Oxford)
- [48] Shrivastav GP, Chaudhuri P, and Horbach J (2016) Yielding of glass under shear: A directed percolation transition precedes shear-band formation Phys. Rev. E94:042605.
- [49] Lin J et al. (2014) Scaling description of the yielding transition in soft amorphous solids at zero temperature.

- [50] Chikkadi V et al. (2011) Long-Range Strain Correlations in Sheared Colloidal Glasses. Phys. Rev. Lett.107:198303.
- [51] Zaccone A, Schall P and Terentjev EM (2014) Microscopic origin of nonlinear nonaffine deformation in bulk metallic glasses. Phys. Rev. B, 90:140203(R).
- [52] Nagamanasa KH et al. (2014) Experimental signatures of a nonequilibrium phase transition governing the yielding of a soft glass. Phys. Rev. E, 89:062308.
- [53] Denisov DV et al. (2015) Sharp symmetry-change marks the mechanical failure transition of glasses. *Sci. Rep.* 5:14359.
- [54] Parisi G, et al. (2017) Shear bands as manifestation of a criticality in yielding amorphous solids. Proc. Nat. Acad. Sci. 114:5577?5582
- [55] Chaudhuri P and Horbach J (2014) Poiseuille flow of soft glasses in narrow channels: From quiescence to steady state. *Phys. Rev. E* 90:040301(R)
- [56] Shrivastav G P, Chaudhuri P, and Horbach J (2016) Heterogeneous dynamics during yielding of glasses: Effect of aging. J. Rheol. 60:835
- [57] Sentjabrskaja T, Chaudhuri P, Hermes M, Poon W C K, Horbach J, Egelhaaf S U, and Laurati M (2015) Creep and flow of glasses: strain response linked to the spatial distribution of dynamical heterogeneities. *Sci. Rep.* 5:11884