

A summation formula for a ${}_3F_2(1)$ hypergeometric series

R. B. PARIS

*Division of Computing and Mathematics,
Abertay University, Dundee DD1 1HG, UK*

Abstract

A summation formula is derived for the hypergeometric series of unit argument ${}_3F_2(1, 1, c; d, n+2; 1)$, where $n = 0, 1, 2, \dots$ and $\Re(d - c + n) > 0$.

Mathematics Subject Classification: 33C15, 33C20, 33C50

Keywords: Clausen's series; Hypergeometric series of unit argument; summation formula

1. Introduction

In this note we give two proofs of the following summation theorem for a particular type of ${}_3F_2(1)$ hypergeometric series (also known as Clausen's series):

Theorem 1. *Let $n = 0, 1, 2, \dots$ and c, d be complex constants satisfying $\Re(d - c + n) > 0$. Then*

$${}_3F_2\left(\begin{matrix} 1, 1, c \\ d, n+2 \end{matrix}; 1\right) = \frac{(n+1)\Gamma(d)}{(1-c)_{n+1}} \left\{ \frac{(d-c)_n}{\Gamma(d-1)} [\psi(d-c+n) - \psi(d-1)] \right. \\ \left. - \sum_{k=1}^n \binom{n}{k} \binom{n+1-c}{k} \frac{D_k(n)}{\Gamma(d-n-1+k)} \right\}, \quad (1.1)$$

where

$$D_k(n) := k! \{ \psi(n+1) - \psi(n+1-k) \}, \quad (1.2)$$

$(a)_k = \Gamma(a+k)/\Gamma(a)$ is Pochhammer's symbol and $\psi(a) = \Gamma'(a)/\Gamma(a)$ is the psi function.

The case $c = d$ (with $n \geq 1$) can be excluded from consideration since the resulting series contracts to a Gauss hypergeometric series which can be summed by the well-known Gauss summation theorem.

2. Proof 1

In Miller and Paris [1, (1.7)] the following summation theorem was established:

$${}_3F_2\left(\begin{matrix} a, c, m \\ d, m+p \end{matrix}; 1\right) = (p)_m \sum_{k=0}^{p-1} (-)^k (m)_k \binom{p-1}{k} \frac{(1-d)_{k+m}}{(1-a)_{k+m}(1-c)_{k+m}} \\ + (m)_p \frac{\Gamma(d)\Gamma(d-a-c)}{\Gamma(d-a)\Gamma(d-c)} \sum_{k=0}^{m-1} (-)^k (p)_k \binom{m-1}{k} \frac{(d-a-c)_{k+p}}{(1-a)_{k+p}(1-c)_{k+p}}$$

for positive integers m and p . In this formula we set $m = 1$ and apply a limiting process to deal with the case $a \rightarrow 1$. We therefore obtain

$${}_3F_2\left(\begin{matrix} a, c, 1 \\ d, p+1 \end{matrix}; 1\right) = \frac{p}{1-a} \sum_{k=0}^{p-1} (-)^k k! \binom{p-1}{k} \frac{(1-d)_{k+1}}{(1-c)_{k+1}(2-a)_k} \\ + \frac{p!}{1-a} \frac{\Gamma(d)\Gamma(d-a-c+p)}{\Gamma(d-a)\Gamma(d-c)(1-c)_p} \frac{\Gamma(2-a)}{\Gamma(1-a+p)}$$

upon use of the identity $(1-a)_{k+1} = (1-a)(2-a)_k$.

Now let $a = 1 - \epsilon$, where $\epsilon \rightarrow 0$. Then

$${}_3F_2\left(\begin{matrix} 1-\epsilon, c, 1 \\ d, p+1 \end{matrix}; 1\right) = \frac{p}{\epsilon} \sum_{k=0}^{p-1} (-)^k k! \binom{p-1}{k} \frac{(1-d)_{k+1}\Gamma(1+\epsilon)}{(1-c)_{k+1}\Gamma(k+1+\epsilon)} \\ + \frac{p!}{\epsilon} \frac{\Gamma(d)\Gamma(d-c+p-1+\epsilon)}{\Gamma(d-1+\epsilon)\Gamma(d-c)(1-c)_p} \frac{\Gamma(1+\epsilon)}{\Gamma(p+\epsilon)} \\ = \frac{p}{\epsilon} \sum_{k=0}^{p-1} (-)^k \binom{p-1}{k} \frac{(1-d)_{k+1}}{(1-c)_{k+1}} \left\{ 1 - \epsilon(\gamma + \psi(k+1)) + O(\epsilon^2) \right\} \\ + \frac{p}{\epsilon} \frac{\Gamma(d)\Gamma(d-c+p-1)}{\Gamma(d-1)\Gamma(d-c)(1-c)_p} \left\{ 1 - \epsilon(\gamma + \psi(p) - \psi(d-c+p-1) + \psi(d-1)) + O(\epsilon^2) \right\}$$

upon use of the expansion $\Gamma(z+\epsilon) = \Gamma(z)\{1 + \epsilon\psi(z) + O(\epsilon^2)\}$ and $\psi(1) = -\gamma$, where $\gamma = 0.55721\dots$ is the Euler-Mascheroni constant.

By means of the evaluation

$$\sum_{k=0}^{p-1} (-)^k \binom{p-1}{k} \frac{(1-d)_{k+1}}{(1-c)_{k+1}} = -\frac{\Gamma(d)\Gamma(d-c+p-1)}{\Gamma(d-1)\Gamma(d-c)(1-c)_p},$$

we then find that

$${}_3F_2\left(\begin{matrix} 1, 1, c \\ d, p+1 \end{matrix}; 1\right) = \frac{p\Gamma(d)\Gamma(d-c+p-1)}{\Gamma(d-1)\Gamma(d-c)(1-c)_p} \{\psi(d-c+p-1) - \psi(d-1)\} \\ + p \sum_{k=0}^{p-1} (-)^k \binom{p-1}{k} \frac{(1-d)_{k+1}}{(1-c)_{k+1}} \{\psi(p) - \psi(k+1)\}.$$

Now let $p \rightarrow n+1$ and replace the summation index n in the finite sum by $n-k$, followed by a little rearrangement, to obtain

$${}_3F_2\left(\begin{matrix} 1, 1, c \\ d, n+2 \end{matrix}; 1\right) = \frac{(n+1)\Gamma(d)}{(1-c)_{n+1}} \left\{ \frac{(d-c)_n}{\Gamma(d-1)} \{\psi(d-c+n) - \psi(d-1)\} \right. \\ \left. - \sum_{k=1}^n \binom{n}{k} \binom{n+1-c}{k} \frac{k! \{\psi(n+1) - \psi(n+1-k)\}}{\Gamma(d-n-1+k)} \right\} \quad (2.1)$$

for $n = 0, 1, 2, \dots$. This is the result stated in Theorem 1. \square

3. Proof 2

In a recent paper [2], the author examined convergent series expansions for the sum

$$S(a, b) = \Lambda \sum_{m=1}^{\infty} \frac{J_{\mu}(am)J_{\nu}(bm)}{m^{\alpha}}, \quad \Lambda = \frac{2^{\mu+\nu}}{a^{\mu}b^{\nu}},$$

where $J_{\nu}(x)$ is the Bessel function of the first kind and the factor Λ is added for convenience. It is supposed that $\mu, \nu \geq 0$, $a, b > 0$ and $\alpha > 0$ for absolute convergence. This was carried out using a Mellin transform approach and evaluation of residues.

In the particular case $\vartheta := \alpha - \mu - \nu = 2n+1$, $n = 0, 1, 2, \dots$ terms in $\log a$ appear in the expansions resulting from a double pole. When $a = b$, it is found that [2, Theorem 2]

$$S(a, a) = \sum_{\substack{m=0 \\ m \neq n}}^{\infty} A_m \left(\frac{1}{2}a\right)^{2m} + \frac{(-)^n \left(\frac{1}{2}a\right)^{2n} \Gamma(\alpha) \Upsilon_n(a)}{\Gamma(n+1+\mu)\Gamma(n+1+\nu)\Gamma(n+1+\mu+\nu)n!} \quad (3.1)$$

for $0 < a \leq \pi$, where

$$A_m = \frac{(-)^m}{m!} \frac{\Gamma(1+\mu+\nu+2m) \zeta(\vartheta-2m)}{\Gamma(1+\mu+m)\Gamma(1+\nu+m)\Gamma(1+\mu+\nu+m)},$$

and

$$\Upsilon_n(a) = \gamma - \log\left(\frac{1}{2}a\right) - \psi(\alpha) + \frac{1}{2}\psi(n+1) + \frac{1}{2}\psi(n+1+\mu) \\ + \frac{1}{2}\psi(n+1+\nu) + \frac{1}{2}\psi(n+1+\mu+\nu),$$

with $\zeta(s)$ being the Riemann zeta function.

When $a \neq b$ (with $a > b$), it is found that [2, Theorem 4]

$$S(a, b) = \frac{1}{\Gamma(1+\nu)} \sum_{\substack{m=0 \\ m \neq n}}^{\infty} B_m \left(\frac{1}{2}a\right)^{2m} + \frac{(-)^n \left(\frac{1}{2}a\right)^{2n}}{\Gamma(1+\nu)\Gamma(n+1+\mu)n!} \left\{ \hat{\Upsilon}_n(a) F_n(\mu, \chi) - \frac{1}{2} \Delta_n(\chi) \right\} \quad (3.2)$$

for $0 < a + b \leq 2\pi$, where $\chi := b^2/a^2$. The coefficients B_m are

$$B_m = \frac{(-)^m \zeta(\vartheta - 2m)}{m! \Gamma(1 + \mu + m)} F_m(\mu, \chi), \quad F_m(\mu, \chi) = {}_2F_1\left(\begin{matrix} -m, -m - \mu \\ 1 + \nu \end{matrix}; \chi\right),$$

where ${}_2F_1$ is the Gauss hypergeometric function. The other quantities appearing in (3.2) are

$$\hat{\Upsilon}_n(a) = \gamma - \log \frac{1}{2}a + \frac{1}{2}\psi(n+1+\mu) + \frac{1}{2}\psi(n+1)$$

and

$$\begin{aligned} \Delta_n(\chi) = \sum_{k=1}^n \binom{n}{k} \binom{n+\mu}{k} \frac{\mathcal{D}_k(n, \mu) \chi^k}{(1+\nu)_k} \\ + \frac{(\mu)_{n+1} \chi^{n+1}}{(1+\nu)_{n+1} (n+1)} {}_3F_2\left(\begin{matrix} 1, 1, 1 - \mu \\ n + \nu + 2, n + 2 \end{matrix}; \chi\right), \end{aligned} \quad (3.3)$$

where

$$\mathcal{D}_k(n, \mu) = D_k(n) + k! \{\psi(n+1+\mu) - \psi(n+1+\mu-k)\}$$

with $D_k(n)$ defined in (1.2).

If we now let $\chi = 1$ and use the Gauss summation formula to show that

$$F_m(\mu, 1) = \frac{\Gamma(1+\nu)\Gamma(1+\mu+\nu+2m)}{\Gamma(1+\nu+m)\Gamma(1+\mu+\nu+m)},$$

then it is easily seen that $A_m = B_m/\Gamma(1+\nu)$ ($m \neq n$). Hence, from (3.1) and (3.2),

$$\frac{\Gamma(\alpha)\Upsilon_n(a)}{\Gamma(1+\nu_n)\Gamma(1+\mu+\nu+n)} = \frac{1}{\Gamma(1+\nu)} \left\{ \hat{\Upsilon}_n(a) F_n(\mu, 1) - \frac{1}{2} \Delta_n(1) \right\}$$

from which we obtain

$$\Delta_n(1) = \frac{\Gamma(\alpha)\Gamma(1+\nu)}{\Gamma(1+\nu+n)\Gamma(1+\mu+\nu+n)} \left\{ 2\psi(\alpha) - \psi(1+\nu+n) - \psi(1+\mu+\nu+n) \right\}.$$

From (3.3) we then find that

$$\begin{aligned} & {}_3F_2\left(\begin{matrix} 1, 1, 1 - \mu \\ n + \nu + 2, n + 2 \end{matrix}; 1\right) \\ &= \frac{(n+1)(1+\nu)_{n+1}}{(\mu)_{n+1}} \left\{ \frac{(1+\mu+\nu+n)_n}{(1+\nu)_n} \left\{ 2\psi(\alpha) - \psi(1+\nu+n) - \psi(1+\mu+\nu+n) \right\} \right. \\ & \quad \left. - \sum_{k=1}^n \binom{n}{k} \binom{n+\mu}{k} \frac{\mathcal{D}_k(n, \mu)}{(1+\nu)_k} \right\}. \end{aligned} \quad (3.4)$$

If we now make the substitutions $c = 1 - \mu$ and $d = n + \nu + 2$ in (3.4), so that $1 + \mu + \nu + n = d - c$, $1 + \nu = d - n - 1$ and $\alpha = d - c + n$, we obtain after a little algebra

$${}_3F_2\left(\begin{matrix} 1, 1, c \\ d, n + 2 \end{matrix}; 1\right) = \frac{(n+1)\Gamma(d)}{(1-c)_{n+1}} \left\{ \frac{(d-c)_n}{\Gamma(d-1)} \{ 2\psi(d-c+n) - \psi(d-1) - \psi(d-c) \} \right.$$

$$- \sum_{k=1}^n \binom{n}{k} \binom{n+1-c}{k} \frac{\mathcal{D}_k(n, 1-c)}{\Gamma(d-n-1+k)} \Big\}.$$

Use of the identity

$$\sum_{k=1}^n \binom{n}{k} \binom{n+1-c}{k} \frac{k! \{\psi(n+2-c) - \psi(n+2-c-k)\}}{\Gamma(d-n-1+k)} = \frac{(d-c)_n}{\Gamma(d-1)} \{\psi(d-c+n) - \psi(d-c)\}$$

to remove some of the ψ functions, we then obtain the result stated in (1.1).

The result in (3.4) was obtained on the assumption that $\mu, \nu \geq 0$. But by the principle of analytic continuation the result in (1.1) will hold in the domain of the parameters c and d where both sides of (1.1) are analytic. This concludes the proof. \square

When $n = 0$ and $n = 1$ we have the summations

$${}_3F_2 \left(\begin{matrix} 1, 1, c \\ d, n+2 \end{matrix}; 1 \right) = \begin{cases} \frac{(d-1)}{1-c} \{\psi(d-c) - \psi(d-1)\} & (n=0) \\ \frac{2(d-1)(d-c)}{(1-c)_2} \{\psi(d-c+1) - \psi(d-1)\} \\ \quad + \frac{2(d-1)}{c-1} & (n=1), \end{cases}$$

where we have used the property $\psi(z+1) = \psi(z) + 1/z$. These summations agree with those presented in [3, p. 452]. In the case when $c = 1$ ($n = 0$) and $c = 1, 2$ ($n = 1$), the summations are given by their limiting forms. Numerical checks with the aid of *Mathematica* have confirmed the validity of (1.1).

References

- [1] A.R. Miller and R.B. Paris, Clausen's series ${}_3F_2(1)$ with integral parameter differences and transformations of the hypergeometric function ${}_2F_2(x)$. *Integral Transforms and Special Functions* **23** (2012) 21–33.
- [2] R.B. Paris, The evaluation of infinite sums of products of two Bessel functions. arXiv:1803.02757 (2018).
- [3] A.P. Prudnikov, Yu.A. Brychkov and O.I. Marichev, *Integrals and Series: More Special Functions* Vol. 3, Gordon and Breach, New York, 1990.