

Multiparticle quantum interference in Bogoliubov bosonic transformations

Michael G. Jabbour^{1,2,3,*} and Nicolas J. Cerf^{1,†}

¹*Quantum Information and Communication, École polytechnique de Bruxelles,
CP 165/59, Université libre de Bruxelles, 1050 Brussels, Belgium*

²*Department of Applied Mathematics and Theoretical Physics,
University of Cambridge, Cambridge CB3 0WA, United Kingdom*

³*Department of Physics, Technical University of Denmark, 2800 Kongens Lyngby, Denmark*

We explore the multiparticle transition probabilities in Gaussian unitaries effected by a two-mode Bogoliubov bosonic transformation on the mode annihilation and creation operators. We show that the transition probabilities can be characterized by remarkably simple, yet unsuspected recurrence equations involving a linear combination of probabilities. The recurrence exhibits an interferometric suppression term – a *negative* probability – which generalizes the seminal Hong-Ou-Mandel effect to more than two indistinguishable photons impinging on a beam splitter of rational transmittance. Unexpectedly, interferences thus originate in this description from the cancellation of probabilities instead of amplitudes. Our framework, which builds on the generating function of the non-Gaussian matrix elements of Gaussian unitaries in Fock basis, is illustrated here for the most common passive and active linear coupling between two optical modes driven by a beam splitter or a parametric amplifier. Hence, it also allows us to predict unsuspected multiphoton interference effects in an optical amplifier of rational gain. In particular, we confirm the newly found two-photon interferometric suppression effect in an amplifier of gain 2 originating from timelike indistinguishability [Proc. Natl. Acad. Sci. **117**, 33107 (2020)]. Overall, going beyond standard two-mode optical components, we expect our method will prove valuable for describing general quantum circuits involving Bogoliubov bosonic transformations.

I. INTRODUCTION

Quantum interference is a cornerstone of quantum physics. While it challenges our understanding of the universe as for instance witnessed in Young’s celebrated double slit-experiment, it has various applications such as quantum computing [1], quantum cryptography [2], or superconducting quantum interference devices [3]. Quantum interference is notably a key to implementing future quantum technologies with photonic integrated devices, which has resulted in a vigorous research effort on harnessing multimode multiphoton interferences over the last decade, see e.g. [4, 5]. This is also significant in connection with the boson sampling paradigm [6], which builds on the computational hardness of simulating the coherent propagation of many identical bosons through a multimode linear interferometer and holds the promise of substantiating the advantage of quantum computers [7–10]. More generally, this has led to a revived interest for quantum interferometry going beyond the celebrated Hong-Ou-Mandel (HOM) effect [11], e.g., the generalized bunching effect in linear networks [12], the signatures of nonclassicality in a multimode interferometer [13], the observation of intrinsically 3-photon interference [14, 15], or the suppression laws in a 8-mode optical Fourier interferometer [16].

Formally, quantum interferences originate from adding up the amplitudes of (often a large number of) possible paths. Since amplitudes are complex, taking the square

modulus of the resulting sum typically gives rise to constructive or destructive interferences. The HOM effect is a paradigmatic example of two-photon quantum interference: the probability of detecting two photons in coincidence at the output of a 50:50 beam splitter (one in each mode) vanishes when one photon impinges on each of the two input modes. The sum of the amplitudes of the two possible paths (both photons being either reflected or transmitted) vanishes, giving rise to destructive interference. In a nutshell, when only two paths of amplitudes α_1 and α_2 interfere, the resulting probability is $p = |\alpha_1 + \alpha_2|^2 = p_1 + p_2 + 2\sqrt{p_1 p_2} \cos \theta$, where $p_1 = |\alpha_1|^2$, $p_2 = |\alpha_2|^2$, and θ is the relative phase.

In this paper, we explore multiparticle quantum interferences that emerge in Bogoliubov bosonic transformations. Bogoliubov transformations are ubiquitous in physics, appearing in various fields such as superconductivity, superfluidity, nuclear physics and quantum field theory. They are also essential in understanding phenomena such as Hawking radiation [17, 18] and the Unruh effect [19, 20]. While our methods and results could be applied in various situations involving bosonic systems, we choose to illustrate them here by focusing on the quantum optics framework. Specifically, we investigate the optical Gaussian unitaries effected by Bogoliubov transformations in phase space, which closely model a great amount of operations performed in quantum optics experiments [21]. We start by examining the generic case of i and k photons impinging on the two input modes of a beam splitter, one of the simplest yet most essential operations in any optical setting. The probability $B_n^{(i,k)}$ of any output pattern is known to be expressible as a multiple summation involving four binomial coefficients

* mgija@dtu.dk

† ncerf@ulb.ac.be

[see Eq. (9)]. This complicated expression, owing to the many interfering paths, can of course be evaluated but cannot easily be exploited analytically. Here, we derive an unexpectedly simple formula (Theorem 1) which governs these probabilities. Counterintuitively, it involves a simple linear combination of probabilities (with no usual $\sqrt{p_1 p_2}$ terms) and the discrepancy with respect to the corresponding classical formula for distinguishable photons appears as a *negative* probability [22].

Our technique relies on calculating the generating function of the matrix elements of Gaussian unitaries in Fock basis, which can be expressed in a simple closed form with the Gaussian toolbox although it encapsulates complex non-Gaussian features such as the multiphoton transition probabilities $B_n^{(i,k)}$. It allows us to extend the HOM effect to many photons by predicting a simple negative contribution to the transition probability. More generally, our framework is suited to Gaussian unitaries describing the passive but also the active linear coupling between two bosonic modes. Hence, we predict a similar interference suppression term – a negative probability – in an optical amplifier, such as a nonlinear crystal pumped in the nondegenerate parametric amplification regime or a four-wave mixer (see Theorem 2). This corroborates and extends the recent finding of a two-boson interference effect in a gain-2 amplifier originating from timelike indistinguishability (bosons from the past and future cannot be distinguished) [23]. Active optical components are essential in continuous-variable quantum information processing [24, 25] as they give access to invaluable resources and protocols, such as universal computing with Gaussian cluster states [26–28], optical multimode entanglement [29], Gaussian quantum steering [30], or Gaussian quantum cloning [31].

As a last result, we provide a further generalization of the HOM effect and two-boson active interference effect [23] by predicting a full interferometric suppression for any rational value of the transmittance (or gain) of a passive (or active) transformation provided specific photon numbers are chosen. Furthermore, we also briefly show that the asymptotic behavior of interferences with large photon numbers can easily be accessed based on generating functions. This illustrates the potential of our framework for describing multiparticle interferences in quantum circuits involving bosonic Bogoliubov transformations in phase space.

II. MODEL AND DERIVATIONS

A. Bosonic Gaussian unitaries

Bosonic modes are common carriers of continuous-variable quantum information [24, 25]. A bosonic mode (e.g., a quantized mode of the electromagnetic field) is modelled by a quantum harmonic oscillator in an infinite-dimensional Fock space. It is associated with the usual pair of bosonic mode operators $\hat{a}$ and $\hat{a}^\dagger$, which must

satisfy the commutation relation $[\hat{a}, \hat{a}^\dagger] = \mathbb{I}$. In this context, Bogoliubov transformations [32] (i.e., linear canonical transformations in $\hat{a}$ and $\hat{a}^\dagger$) are of particular interest as they correspond to Gaussian unitaries (i.e., quadratic Hamiltonians in $\hat{a}$ and $\hat{a}^\dagger$). They are especially valuable in the framework of quantum optics, where they conserve Gaussian-shaped Wigner functions in phase space and, most importantly, model ubiquitous transformations in experimental conditions and form the core of Gaussian quantum information [21]. They can be divided into passive and active transformations as effected by linear optical interferometry or parametric amplification, respectively. In this work, we illustrate our method for the most fundamental passive and active two-mode Gaussian unitaries, namely the beam splitter (BS) and two-mode squeezer (TMS). The BS unitary U_η^{BS} effects an energy-conserving linear coupling between two modes and acts in the Heisenberg picture as

$$\begin{aligned} U_\eta^{\text{BS}\dagger} \hat{a} U_\eta^{\text{BS}} &= \sqrt{\eta} \hat{a} + \sqrt{1-\eta} \hat{b}, \\ U_\eta^{\text{BS}\dagger} \hat{b} U_\eta^{\text{BS}} &= -\sqrt{1-\eta} \hat{a} + \sqrt{\eta} \hat{b}, \end{aligned} \quad (1)$$

where $\hat{a}$ and $\hat{b}$ are the mode operators, while η is the transmittance. Similarly, the TMS unitary U_λ^{TMS} models the generation of pairs of entangled photons by parametric amplification due to the pumping of a nonlinear crystal, and acts on mode operators as

$$\begin{aligned} U_\lambda^{\text{TMS}\dagger} \hat{a} U_\lambda^{\text{TMS}} &= \cosh(r) \hat{a} + \sinh(r) \hat{b}^\dagger, \\ U_\lambda^{\text{TMS}\dagger} \hat{b} U_\lambda^{\text{TMS}} &= \sinh(r) \hat{a}^\dagger + \cosh(r) \hat{b}. \end{aligned} \quad (2)$$

with $\lambda := \tanh^2(r)$ for a parametric gain $g := \cosh^2(r)$. The transformations characterized by Eqs. (1) and (2) happen to be useful in various contexts involving the evolution of bosonic systems. For instance, they can be exploited in black hole theory, where they describe the interaction of a Gaussian bosonic state with an already formed Schwarzschild black hole [33].

B. Generating functions

The generating function (GF) of a sequence $\{c_n\}_{n \in \mathbb{N}_0}$ is defined as

$$g(z) := \mathcal{T}_n[c_n](z) = \sum_{n=0}^{\infty} c_n z^n, \quad z \in \mathbb{C}. \quad (3)$$

It is a powerful tool as $g(z)$ encapsulates the entire sequence via $c_n = n!^{-1}(\partial^n g / \partial z^n)_{z=0}$. Here, we exploit the properties of GFs in quantum optics when applied to the squared modulus of the matrix elements of Gaussian unitaries in Fock basis. Unlike the matrix elements in a coherent (Gaussian) basis, these happen to be quite difficult to handle because Fock states are non Gaussian, so it is helpful to characterize them via their GFs. Consider the 4-dimensional sequence of transition probabilities $|\langle n, m | U | i, k \rangle|^2$ for some unitary U , where $|i\rangle$, $|k\rangle$,

$|n\rangle$, and $|m\rangle$ denote Fock states ($i, k, n, m \in \mathbb{N}_0$). Its 4-variate GF can be written as (see [34])

$$f(\mathbf{v}) = \frac{\text{Tr}[(\tau_z \otimes \tau_w)U(\tau_x \otimes \tau_y)U^\dagger]}{(1-x)(1-y)(1-z)(1-w)}, \quad (4)$$

where we chose $\mathbf{v} := (x, y, z, w)$ such that $(x, y) \in [0, 1]^2$ and $(w, z) \in [0, 1]^2$, with the conventions shown in Fig. 1. Thus, $f(\mathbf{v})$ is proportional to the overlap between two Gaussian states, one of which being the product of two thermal states of the form $\tau_x := (1-x) \sum_{n=0}^{\infty} x^n |n\rangle \langle n|$, while the other is the product of two thermal states processed through the unitary U . This makes $f(\mathbf{v})$ very easy to compute when U is Gaussian, regardless of the complexity of $|\langle n, m | U | i, k \rangle|^2$ itself, by exploiting the Gaussian formalism in phase space. Recalling that the overlap between two zero-mean Gaussian states ρ_1 and ρ_2 with covariance matrices V_1 and V_2 is given by $\text{Tr}[\rho_1 \rho_2] = 1/\sqrt{\det[(V_1 + V_2)/2]}$ [35], the GF of $|\langle n, m | U_\eta^{\text{BS}} | i, k \rangle|^2$ can be expressed using standard tools of quantum optics as [34]

$$f_\eta^{\text{BS}}(\mathbf{v}) = \frac{1}{1 - \eta(xz + yw) - \bar{\eta}(xw + yz) + xyzw}, \quad (5)$$

where $\bar{\eta} := 1 - \eta$, while the GF of $|\langle n, m | U_\lambda^{\text{TMS}} | i, k \rangle|^2$ can be written as [34]

$$f_\lambda^{\text{TMS}}(\mathbf{v}) = \frac{\bar{\lambda}}{1 - \lambda(xy + zw) - \bar{\lambda}(xz + yw) + xyzw}, \quad (6)$$

where $\bar{\lambda} := 1 - \lambda$. As a consistency check, we note that

$$\begin{aligned} f_\eta^{\text{BS}}(\mathbf{0}) &= |\langle 0, 0 | U_\eta^{\text{BS}} | 0, 0 \rangle|^2 = 1, \\ f_\lambda^{\text{TMS}}(\mathbf{0}) &= |\langle 0, 0 | U_\lambda^{\text{TMS}} | 0, 0 \rangle|^2 = \bar{\lambda}, \end{aligned} \quad (7)$$

while normalization $\sum_{n,m=0}^{\infty} |\langle n, m | U | i, k \rangle|^2 = 1, \forall i, k$, translates into

$$f^{\text{BS/TMS}}(x, y, 1, 1) = (1-x)^{-1}(1-y)^{-1}. \quad (8)$$

Interestingly, energy conservation in U_η^{BS} manifests itself through $f_\eta^{\text{BS}}(x, y, z, w) = f_\eta^{\text{BS}}(tx, ty, z/t, w/t), \forall t$, while the conservation of the photon number difference in U_λ^{TMS} is reflected by $f_\lambda^{\text{TMS}}(x, y, z, w) = f_\lambda^{\text{TMS}}(tx, y/t, z/t, tw), \forall t$, see [34].

C. Multiphoton transition probabilities

We use Eq. (5) to derive a surprisingly simple recurrence equation for the multiphoton transition probabilities in a BS, denoted as $B_n^{(i,k)} := |\langle n, i+k-n | U_\eta^{\text{BS}} | i, k \rangle|^2$, with $i, k, n \in \mathbb{N}_0$. Incidentally, note that a direct calculation yields [34]

$$B_n^{(i,k)} = \eta^k \bar{\eta}^i \sum_{m,j=\max(0,n-k)}^{\min(i,n)} (-1)^{m+j} \gamma_{n,m,j}^{(i,k)} \left(\frac{\eta}{\bar{\eta}}\right)^{m+j-n}, \quad (9)$$




FIG. 1. Conventions in the definition of $f(x, y, z, w)$, which is the generating function of the transition probability for sending the Fock states $|i\rangle$ and $|k\rangle$ respectively in modes $\hat{a}$ and $\hat{b}$ and, after processing through the unitary U , measuring the Fock states $|n\rangle$ and $|m\rangle$ respectively in modes $\hat{a}$ and $\hat{b}$.

where

$$\gamma_{n,m,j}^{(i,k)} := \binom{i}{m} \binom{k}{n-m} \binom{n}{j} \binom{i+k-n}{i-j}, \quad (10)$$

which is quite cumbersome to manipulate. Nevertheless, the following theorem provides an alternative.

Theorem 1. *If $i=k=n=0$, then $B_n^{(i,k)} = 1$, else,*

$$\begin{aligned} B_n^{(i,k)} &= \eta B_{n-1}^{(i-1,k)} + \eta B_n^{(i,k-1)} \\ &\quad + \bar{\eta} B_n^{(i-1,k)} + \bar{\eta} B_{n-1}^{(i,k-1)} - B_{n-1}^{(i-1,k-1)}. \end{aligned} \quad (11)$$

The definition of $B_n^{(i,k)}$ is extended here to all integers i, k, n , setting it to zero when either of them is negative.

Proof. We set $\mathbf{u} := (x, y, z)$, $\mathbf{j} := (i, k, n)$ and denote by $g_\eta^{\text{BS}}(\mathbf{u})$ the 3-variate GF of $B_n^{(i,k)}$ with the conventions of Fig. 1. Since $g_\eta^{\text{BS}}(\mathbf{u}) = f_\eta^{\text{BS}}(x, y, z, 1)$, Eq. (5) implies

$$[1 - \eta(xz + y) - \bar{\eta}(x + yz) + xyz] g_\eta^{\text{BS}}(\mathbf{u}) = 1. \quad (12)$$

Using the shifting property of the GFs and the notation of Eq. (3), it can easily be shown that multiplying the GF by $\mathbf{u}_l$ for $l = 1, 2, 3$ corresponds to decreasing the index $\mathbf{j}_l$ of $B_n^{(i,k)}$ by one unit, so that for instance

$$\mathcal{T}_j [B_{n-1}^{(i-1,k)}] (\mathbf{u}) = xz \mathcal{T}_j [B_n^{(i,k)}] (\mathbf{u}) = xz g_\eta^{\text{BS}}(\mathbf{u}). \quad (13)$$

In addition, we know that the 3-variate GF of the product $\delta_{i,0} \delta_{k,0} \delta_{n,0}$ of three Kronecker deltas is 1. Using this, we see that Eq. (12) is equivalent to the relation

$$\begin{aligned} B_n^{(i,k)} &- \eta [B_{n-1}^{(i-1,k)} + B_n^{(i,k-1)}] \\ &- \bar{\eta} [B_n^{(i-1,k)} + B_{n-1}^{(i,k-1)}] + B_{n-1}^{(i-1,k-1)} = \delta_{i,0} \delta_{k,0} \delta_{n,0}, \end{aligned}$$

which proves the theorem. $\square$

This recurrence can be nicely interpreted in the context of the HOM effect. As illustrated in Fig. 2, the first four terms of the right-hand side of Eq. (11) corroborate the classical intuition one may have about $B_n^{(i,k)}$: one should add the probabilities corresponding to the different scenarios in which the n th photon has not reached




FIG. 2. Classical components of the recurrence formula (11) for the transition probability $B_n^{(i,k)}$ in a BS.

the BS yet, multiplied by the right probability (η or $\bar{\eta}$) depending on which path it takes. For example, $B_{n-1}^{(i-1,k)}$ must be multiplied by η since the extra photon must be injected on the input mode a and exit on the output mode a . Crucially, as a consequence of bosonic statistics, a fifth term appears in Eq. (11) with a minus sign that accounts for quantum interference and may be viewed as an interference suppression term. In the special case where $i = k = 1$ and $\eta = 1/2$, we recover the standard HOM effect,

$$\begin{aligned} B_1^{(1,1)} &= \frac{1}{2} B_0^{(0,1)} + \frac{1}{2} B_1^{(1,0)} + \frac{1}{2} B_1^{(0,1)} + \frac{1}{2} B_0^{(1,0)} - B_0^{(0,0)} \\ &= 4 \times \frac{1}{2} \times \frac{1}{2} - 1 = 0. \end{aligned} \quad (14)$$

Let us stress that this is a very unconventional proof of the HOM effect as Eq. (14) does not involve a linear combination of amplitudes but of probabilities. The first two terms account for both photons being transmitted while the third and fourth terms correspond to both of them being reflected. The fifth (negative) term has no classical counterpart. Note that if $k = 0$ and $i \geq 0$, the interference term disappears in Eq. (11) and one gets the recurrence $B_n^{(i,0)} = \eta B_{n-1}^{(i-1,0)} + \bar{\eta} B_n^{(i-1,0)}$, which had been derived in the context of majorization theory applied to bosonic transformations [36].

D. Distinguishable photons

It is instructive to give Eq. (11) further interpretation by juxtaposing it with its classical counterpart for distinguishable photons, which may for instance happen if the incident photons occupy different temporal modes. The classical probability of detecting n photons on output mode a when i and k photons impinge on input modes a and b is given by the convolution $p_{n|i,k} = \sum_{n'=0}^n p_{n'|i}^A p_{n-n'|k}^B$, where $p_{n|i}^A$ (or $p_{n|k}^B$) is the probability of getting n photons if i (or k) distinguishable photons enter mode a (or b), which itself follows a binomial distribution of parameter η (or $\bar{\eta}$), see [34] for details. Hence, the 3-variate GF of $p_{n|i,k}$ is given by $g_\eta^{\text{cl}}(\mathbf{u}) = g_\eta^A(x, z) g_\eta^B(y, z)$, where $g_\eta^A(x, z)$ and $g_\eta^B(y, z)$ are the 2-variate GFs of $p_{n|i}^A$ and $p_{n|k}^B$. For instance, it is easy to show that $g_\eta^A(x, z) = 1/(1 - \eta x z - \bar{\eta} x)$, so that $g_\eta^{\text{cl}}(\mathbf{u})$ satisfies the relation

$$(1 - \eta x z - \bar{\eta} x) g_\eta^{\text{cl}}(\mathbf{u}) = 1(x) g_\eta^B(y, z). \quad (15)$$

where $1(x) \equiv 1$ is a constant function of x . Using again the shifting property of GFs, Eq. (15) implies the classical recurrence relation

$$p_{n|i,k} = \eta p_{n-1|i-1,k} + \bar{\eta} p_{n|i-1,k}, \quad i > 0, \quad (16)$$

where we have used the fact that $1(x) g_\eta^B(y, z)$ is the GF of $\delta_{i,0} p_{n|k}^B$ and can be ignored for $i > 0$. Interchanging $p_{n|i}^A$ and $p_{n|k}^B$, a similar reasoning yields

$$p_{n|i,k} = \eta p_{n|i,k-1} + \bar{\eta} p_{n-1|i,k-1}, \quad k > 0. \quad (17)$$

We notice here that Eq. (16) coincides with the first and third terms in Eq. (11), while Eq. (17) coincides with the second and fourth terms. If either $i = 0$ or $k = 0$ (i.e., no photon in one of the two input modes), then Eq. (11) reduces to the classical recurrence (for instance, Eq. (16) covers the case $k = 0$). As advertised, the fifth (negative) term in Eq. (11) thus captures quantum interference (it appears as soon as $i, k > 0$) since it is absent from the classical formulas (16) and (17). Note also that removing this negative quantum term in Eq. (11) would then lead to twice the classical probability.

E. Active Gaussian transformations

An even more appealing application of our framework is to explore multiphoton interferences in an active transformation, such as a TMS. As proven in [23], a TMS may be viewed as a BS undergoing “partial time reversal”, namely $\langle n, m | U_\lambda^{\text{TMS}} | i, k \rangle = \sqrt{1 - \lambda} \langle n, k | U_{1-\lambda}^{\text{BS}} | i, m \rangle$. Indeed, indices k and m are interchanged, which may be interpreted as reverting the arrow of time of mode b [37]. Similarly, interchanging variables y and w , we see that the GFs are connected by $f_\lambda^{\text{TMS}}(x, y, z, w) = (1 - \lambda) f_{1-\lambda}^{\text{BS}}(x, w, z, y)$, which is consistent with Eqs. (5) and (6). This allows us to write a recurrence for the transition probabilities $A_n^{(i,k)} := |\langle n, k - i + n | U_\lambda^{\text{TMS}} | i, k \rangle|^2$ in a TMS (the definition of $A_n^{(i,k)}$ is extended to all integers i, k, n , setting it to zero when either of them is negative).

Theorem 2. *If $i = k = n = 0$, then $A_n^{(i,k)} = \bar{\lambda}$, else,*

$$\begin{aligned} A_n^{(i,k)} &= \lambda A_n^{(i-1,k-1)} + \lambda A_{n-1}^{(i,k)} \\ &\quad + \bar{\lambda} A_{n-1}^{(i-1,k)} + \bar{\lambda} A_n^{(i,k-1)} - A_{n-1}^{(i-1,k-1)}. \end{aligned} \quad (18)$$

Proof. The relation can be easily proven by making use of Theorem 1, exploiting the fact that $A_n^{(i,k)} = \bar{\lambda} B_n^{(i,k-i+n)}$ with $\eta = \bar{\lambda}$ (or $\eta = 1/g$), see [34]. $\square$




FIG. 3. Classical components of the recurrence formula (18) for the transition probabilities $A_n^{(i,k)}$ in a TMS.

Equation (18) is quite intriguing at first sight, as it is unclear how interferences take place in an active medium. However, as illustrated in Fig. 3, we may build an interpretation of Eq. (18) by considering all possible classical scenarios. The first term corresponds to the stimulated annihilation of an extra input photon pair, while the second term corresponds to the stimulated emission of an extra output photon pair (both occurring with probability $\propto \lambda$). The third and fourth terms correspond to an extra photon crossing the nonlinear medium without stimulating pair emission nor absorption (both with probability $\propto \lambda$). Most importantly, the fifth (negative) term is again responsible for an unsuspected quantum interference effect, which has no classical counterpart. In the special case where $i = k = 1$ and $\lambda = 1/2$, we predict a complete extinction of the output state $|1\rangle|1\rangle$, which confirms a newly discovered two-photon interference effect in an amplifier of gain 2 [23] originating from *timelike* indistinguishability between the input and output photon pairs (exactly like the HOM effect can be viewed as a consequence of *spacelike* indistinguishability between two photons entering a BS of transmittance $1/2$). Here again, we find a surprising explanation of this effect based on the cancellation of probabilities (not amplitudes), namely

$$\begin{aligned} A_1^{(1,1)} &= \frac{1}{2} A_1^{(0,0)} + \frac{1}{2} A_0^{(1,1)} + \frac{1}{2} A_0^{(0,1)} + \frac{1}{2} A_1^{(1,0)} - A_0^{(0,0)} \\ &= 4 \times \frac{1}{2} \times \frac{1}{4} - \frac{1}{2} = 0. \end{aligned} \quad (19)$$

The first two terms account for events consisting of the stimulated annihilation of the input photon pair accompanied with the stimulated emission of a distinct out-

put pair, while the third and fourth terms correspond to events where both photons cross the TMS. The fifth term is intrinsically quantum. Note that for $k = 0$, Eq. (18) reduces to the recurrence $A_n^{(i,0)} = \lambda A_{n-1}^{(i,0)} + \bar{\lambda} A_{n-1}^{(i-1,0)}$ implying a majorization relation in a bosonic amplifier channel that was proven in [38].

F. Rational transmittance and gain

Coming back to passive BS transformations, it is easy to predict the existence of a HOM-like suppression effect for any rational value of the transmittance $\eta < 1$ provided some specific numbers of impinging photons are considered, namely

$$B_1^{(i,k)} = 0 \quad \text{if } \eta = k/(i+k), \quad (20)$$

as illustrated in Fig. 4. This can be understood as the result of amplitude cancellation between two scenarios, taking as a reference the situation where $i-1$ photons on mode a are reflected and $k-1$ photons on mode b are transmitted. The single photon observed on the output mode a may come from input mode a or b . Either the i th photon on mode a is transmitted (there are i possible choices) and all k photons on mode b are transmitted, which yields an amplitude $\propto i\eta$, or the k th photon on mode b is reflected (there are k possible choices) and all i photons on mode a are reflected, which yields an amplitude $\propto k\bar{\eta}$. Hence, we have $B_1^{(i,k)} \propto (i\eta - k\bar{\eta})^2$, which




FIG. 4. Probability $B_1^{(i,k)}$ of observing a single photon on mode a at the output of a BS for $k = 5$ and for three different values $i = 3, 6, 9$ as a function of the transmittance η . We observe a HOM-like suppression effect for the corresponding rational values of $\eta = 5/8, 5/11, 5/14$.




FIG. 5. Probability $A_1^{(i,i+k-1)}$ of observing a single photon on mode a at the output of a TMS for $k = 5$ and for three different values $i = 3, 6, 9$ as a function of the parameter λ . Note that k denotes here the number of output photons on mode b . We observe a suppression effect akin to the HOM effect for the corresponding rational values of $\lambda = 3/8, 6/11, 9/14$, or equivalently for values of the gain $g = 8/5, 11/5, 14/5$.

is consistent with Eq. (20). However, we provide a distinct interpretation in terms of probability cancellation as implied by Eq. (11), see [34]. For a quantum optical amplifier, we observe a similar effect for any rational value of the gain $g > 1$, namely

$$A_1^{(i,i+k-1)} = 0 \quad \text{if } \lambda = i/(i+k), \quad (21)$$

corresponding to $g = 1 + i/k$ (see Fig. 5). This heretofore unknown interference effect can again be viewed as a consequence of probability cancellation in Eq. (18), see [34].

III. CONCLUSION AND OUTLOOK

Gaussian bosonic unitaries are readily described as affine transformations in phase space. Yet, addressing their action on Fock states typically leads to cumbersome calculations, which makes multiphotonic interferences in common Gaussian optical components hard to grasp. As a consequence, it is often an intractable task to prove fundamental entropy inequalities for Gaussian bosonic channels, while these are of major importance in optical quantum communication (see, e.g., the entropy photon-number inequality [39–41]). Here, we have shown that the generating function of the matrix elements of a BS or TMS in Fock space can be expressed in a closed form, which, as a central consequence, yields simple recurrence equations for the multiphoton transition probabilities. In spite of the many interfering paths, Theorems 1 and 2 then provide a simple, intuitively appealing picture of multiphoton interference in passive and active bosonic circuits. It is amazing that such a simple account of quantum interferences in terms of probabilities (instead of amplitudes) in so well-studied optical components had yet remained unnoticed.

We have then predicted several multiphoton generalizations of the HOM effect in a BS of rational transmittance and have exploited the correspondence between a BS and TMS under partial time reversal [23] in order to reveal the existence of similar interferometric suppression effects in a quantum optical amplifier of rational gain. Interestingly, these predicted effects seem to escape the general framework for quantum suppression laws that has been derived in refs. [42, 43].

Let us stress that the generating function of transition probabilities can also be useful in studying other properties of Gaussian unitaries, for example their asymptotic behavior. Using Tauberian theorems, which state that if $g(z) \sim 1/(1-z)$ for $z \rightarrow 1$, then $\sum_{l=0}^n c_l \sim n$ for $n \rightarrow \infty$, it is indeed possible to approximate $B_n^{(i,i)}$ when $i \rightarrow \infty$ [34]. For $\eta = 1/2$, this exactly coincides with the asymptotic analysis of a BS with a large photon number in both input ports [44]. Note also that the generating function has recently been exploited in order to connect boson sampling with Fock-state inputs to boson sampling with thermal-state inputs [45], which is reminiscent of Eq. (4). Moreover, the technique developed here yields a powerful tool for characterizing certain non-Gaussian bosonic channels (those that are Gaussian-dilatable), for example photon-added or photon-subtracted channels [46–48] as well as the linear coupling of a signal mode together with a passive environment [49].

Overall, beyond the results for a BS and TMS highlighted in this paper, we expect that our framework can be amenable to address any Bogoliubov transformation acting on an arbitrary number of modes. The special case of a multimode linear interferometer has already been considered in [50, 51]. Although it does not seem to have implications for the complexity of simulating bosonic interferences, it provides a neat description of multimode multiphoton interference involving negative probabilities. Furthermore, we may anticipate other applications of this framework going beyond photonic systems. The same approach should indeed prove valuable for nonphotonic bosonic systems as well, since the transformations described by Eqs. (1) and (2) are not restricted to optical components but have quite a broad range of applications. In short, we have at hand a distinct approach to quantum multiparticle interferences in (passive and active) Bogoliubov transformations acting on any bosonic quantum systems.

M.G.J. acknowledges support from the Wiener-Anspach Foundation and from the Carlsberg Foundation. This work was supported by the F.R.S.-FNRS under project no. PDR T.0224.18 and by the EC under project ShoQC within the ERA-NET Cofund in Quantum Technologies (QuantERA) program.

-
- [1] T. D. Ladd, F. Jelezko, R. Laflamme, Y. Nakamura, C. Monroe, and J. L. O’Brien, Quantum computers, *Nature* **464**, 45 (2010).
 - [2] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, Quantum cryptography, *Rev. Mod. Phys.* **74**, 145 (2002).
 - [3] D. Vasyukov, Y. Anahory, L. Embon, D. Halbertal, J. Cuppens, L. Neeman, A. Finkler, Y. Segev, Y. Myasoedov, M. L. Rappaport, M. E. Huber, and E. Zeldov, A scanning superconducting quantum interference device with single electron spin sensitivity, *Nature Nanotechnol-*

- ogy* **8**, 639 (2013).
- [4] A. Peruzzo, A. Laing, A. Politi, T. Rudolph, and J. L. O’Brien, Multimode quantum interference of photons in multiport integrated devices, *Nat. Commun.* **2**, 224 (2011).
- [5] B. A. Bell, G. S. Thekkadath, R. Ge, X. Cai, and I. A. Walmsley, Testing multi-photon interference on a silicon chip, *Optics Express* **27**, 35646 (2019).
- [6] S. Aaronson and A. Arkhipov, The computational complexity of linear optics, in *Proceedings of the Forty-*

- Third Annual ACM Symposium on Theory of Computing*, STOC '11 (Association for Computing Machinery, New York, NY, USA, 2011) p. 333–342.
- [7] M. Tillmann, B. Dakić, R. Heilmann, S. Nolte, A. Szameit, and P. Walther, Experimental boson sampling, *Nat. Photonics* **7**, 540 (2013).
 - [8] A. Crespi, R. Osellame, R. Ramponi, D. J. Brod, E. F. Galvão, N. Spagnolo, C. Vitelli, E. Maiorino, P. Mataloni, and F. Sciarrino, Integrated multimode interferometers with arbitrary designs for photonic boson sampling, *Nat. Photonics* **7**, 545 (2013).
 - [9] M. A. Broome, A. Fedrizzi, S. Rahimi-Keshari, J. Dove, S. Aaronson, T. C. Ralph, and A. G. White, Photonic boson sampling in a tunable circuit, *Science* **339**, 794 (2013).
 - [10] J. Carolan, J. D. A. Meinecke, P. J. Shadbolt, N. J. Russell, N. Ismail, K. Wörhoff, T. Rudolph, M. G. Thompson, J. L. O'Brien, J. C. F. Matthews, and A. Laing, On the experimental verification of quantum complexity in linear optics, *Nat. Photonics* **8**, 621 (2014).
 - [11] C. K. Hong, Z. Y. Ou, and L. Mandel, Measurement of subpicosecond time intervals between two photons by interference, *Phys. Rev. Lett.* **59**, 2044 (1987).
 - [12] V. S. Shchesnovich, Universality of generalized bunching and efficient assessment of boson sampling, *Phys. Rev. Lett.* **116**, 123601 (2016).
 - [13] L. Rigovacca, C. Di Franco, B. J. Metcalf, I. A. Walmsley, and M. S. Kim, Nonclassicality criteria in multiport interferometry, *Phys. Rev. Lett.* **117**, 213602 (2016).
 - [14] S. Agne, T. Kauten, J. Jin, E. Meyer-Scott, J. Z. Salvail, D. R. Hamel, K. J. Resch, G. Weihs, and T. Jennewein, Observation of genuine three-photon interference, *Phys. Rev. Lett.* **118**, 153602 (2017).
 - [15] A. J. Menssen, A. E. Jones, B. J. Metcalf, M. C. Tichy, S. Barz, W. S. Kolthammer, and I. A. Walmsley, Distinguishability and many-particle interference, *Phys. Rev. Lett.* **118**, 153603 (2017).
 - [16] A. Crespi, R. Osellame, R. Ramponi, M. Bentivegna, F. Flamini, N. Spagnolo, N. Viggianiello, L. Innocenti, P. Mataloni, and F. Sciarrino, Suppression law of quantum states in a 3d photonic fast Fourier transform chip, *Nat. Commun.* **7**, 10469 (2016).
 - [17] S. W. Hawking, Particle creation by black holes, *Commun. Math. Phys.* **43**, 199 (1975).
 - [18] J. Traschen, An Introduction to Black Hole Evaporation, in *Mathematical Methods in Physics*, edited by A. A. Bytsenko and F. L. Williams (World Scientific, 2000) p. 180.
 - [19] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 1982).
 - [20] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas, The unruh effect and its applications, *Rev. Mod. Phys.* **80**, 787 (2008).
 - [21] C. Weedbrook, S. Pirandola, R. García-Patrón, N. J. Cerf, T. C. Ralph, J. H. Shapiro, and S. Lloyd, Gaussian quantum information, *Rev. Mod. Phys.* **84**, 621 (2012).
 - [22] R. P. Feynman, Negative probability, in *Quantum Implications: Essays in Honour of David Bohm* (Routledge & Kegan Paul Ltd, London & New York, 1987) pp. 235–248.
 - [23] N. J. Cerf and M. G. Jabbour, Two-boson quantum interference in time, *Proc. Natl. Acad. Sci.* **117**, 33107 (2020).
 - [24] S. L. Braunstein and P. van Loock, Quantum information with continuous variables, *Rev. Mod. Phys.* **77**, 513 (2005).
 - [25] N. J. Cerf, G. Leuchs, and E. S. Polzik, *Quantum Information with Continuous Variables of Atoms and Light* (Imperial College Press, 2007).
 - [26] P. van Loock, C. Weedbrook, and M. Gu, Building Gaussian cluster states by linear optics, *Phys. Rev. A* **76**, 032321 (2007).
 - [27] M. Gu, C. Weedbrook, N. C. Menicucci, T. C. Ralph, and P. van Loock, Quantum computing with continuous-variable clusters, *Phys. Rev. A* **79**, 062318 (2009).
 - [28] M. V. Larsen, X. Guo, C. R. Breum, J. S. Neergaard-Nielsen, and U. L. Andersen, Deterministic generation of a two-dimensional cluster state, *Science* **366**, 369 (2019).
 - [29] S. Gerke, J. Sperling, W. Vogel, Y. Cai, J. Roslund, N. Treps, and C. Fabre, Full multipartite entanglement of frequency-comb Gaussian states, *Phys. Rev. Lett.* **114**, 050501 (2015).
 - [30] I. Kogias, A. R. Lee, S. Ragy, and G. Adesso, Quantification of Gaussian quantum steering, *Phys. Rev. Lett.* **114**, 060403 (2015).
 - [31] N. J. Cerf, A. Ipe, and X. Rottenberg, Cloning of continuous quantum variables, *Phys. Rev. Lett.* **85**, 1754 (2000).
 - [32] N. N. Bogoljubov, On a new method in the theory of superconductivity, *Il Nuovo Cimento* (1955-1965) **7**, 794 (1958).
 - [33] K. Brádler and C. Adami, Black holes as bosonic Gaussian channels, *Phys. Rev. D* **92**, 025030 (2015).
 - [34] See Supplemental Material for details on the calculation of the multiphoton transition probabilities and associated generating function for a beam splitter (and similarly for a two-mode squeezer), the classical benchmark of distinguishable photons, and the asymptotic calculation of the transition probabilities based on the generating function.
 - [35] H. Scutaru, Fidelity for displaced squeezed thermal states and the oscillator semigroup, *J. Phys. A: Math. Theor.* **31**, 3659 (1998).
 - [36] C. N. Gagatsos, O. Oreshkov, and N. J. Cerf, Majorization relations and entanglement generation in a beam splitter, *Phys. Rev. A* **87**, 042307 (2013).
 - [37] N. J. Cerf, The optical beam splitter under partial time reversal (2012), 9th Central European Workshop on Quantum Optics (CEWQO 2012), Sinaia, Romania.
 - [38] R. García-Patrón, C. Navarrete-Benlloch, S. Lloyd, J. H. Shapiro, and N. J. Cerf, Majorization theory approach to the Gaussian channel minimum entropy conjecture, *Phys. Rev. Lett.* **108**, 110505 (2012).
 - [39] S. Guha, B. Erkmen, and J. Shapiro, The entropy photon-number inequality and its consequences, in *2008 Information Theory and Applications Workshop - Conference Proceedings, ITA* (2008) pp. 128–130.
 - [40] R. König and G. Smith, The entropy power inequality for quantum systems, *IEEE Trans. Inf. Theory* **60**, 1536 (2014).
 - [41] G. De Palma, A. Mari, and V. Giovannetti, A generalization of the entropy power inequality to bosonic quantum systems, *Nat. Photonics* **8**, 958 (2014).
 - [42] C. Dittel, G. Dufour, M. Walschaers, G. Weihs, A. Buchleitner, and R. Keil, Totally destructive many-particle interference, *Phys. Rev. Lett.* **120**, 240404 (2018).
 - [43] C. Dittel, G. Dufour, M. Walschaers, G. Weihs, A. Buchleitner, and R. Keil, Totally destructive interference for permutation-symmetric many-particle states, *Phys. Rev. A* **97**, 062116 (2018).
 - [44] H. Nakazato, S. Pascasio, M. Stobińska, and K. Yuasa,

- Photon distribution at the output of a beam splitter for imbalanced input states, *Phys. Rev. A* **93**, 023845 (2016).
- [45] Y. Kim, K.-H. Hong, Y.-H. Kim, and J. Huh, Connection between BosonSampling with quantum and classical input states, *Optics Express* **28**, 6929 (2020).
- [46] C. Navarrete-Benlloch, R. García-Patrón, J. H. Shapiro, and N. J. Cerf, Enhancing quantum entanglement by photon addition and subtraction, *Phys. Rev. A* **86**, 012328 (2012).
- [47] K. K. Sabapathy and A. Winter, Non-Gaussian operations on bosonic modes of light: Photon-added Gaussian channels, *Phys. Rev. A* **95**, 062309 (2017).
- [48] S. M. Barnett, G. Ferenczi, C. R. Gilson, and F. C. Speirits, Statistics of photon-subtracted and photon-added states, *Phys. Rev. A* **98**, 013809 (2018).
- [49] M. G. Jabbour and N. J. Cerf, Fock majorization in bosonic quantum channels with a passive environment, *J. Phys. A: Math. Theor.* **52**, 105302 (2019).
- [50] M. G. Jabbour, Bosonic systems in quantum information theory: Gaussian-dilatable channels, passive states, and beyond, (Ph. D. thesis, Université libre de Bruxelles, 2018).
- [51] M. G. Jabbour and N. J. Cerf, Revisiting quantum interferences in multimode Gaussian circuits, (in preparation).
- [52] M. S. Kim, W. Son, V. Bužek, and P. L. Knight, Entanglement by a beam splitter: Nonclassicality as a prerequisite for entanglement, *Phys. Rev. A* **65**, 032323 (2002).
- [53] J. S. Ivan, K. K. Sabapathy, and R. Simon, Operator-sum representation for bosonic Gaussian channels, *Phys. Rev. A* **84**, 042311 (2011).
- [54] P. Flajolet and R. Sedgewick, *Analytic Combinatorics* (Cambridge University Press, UK, 2009).
- [55] G. H. Hardy and J. E. Littlewood, Tauberian theorems concerning power series and Dirichlet's series whose coefficients are positive, *Proceedings of the London Mathematical Society* **s2-13**, 174 (1914).
- [56] J. Karamata, Über die Hardy-Littlewoodschen umkehrungen des abelschen stetigkeitssatzes, *Mathematische Zeitschrift* **32**, 319 (1930).

SUPPLEMENTAL MATERIAL

The present material provides supplementary information to the main text. It includes calculations which are standard in both quantum optics and the theory of generating functions, which we chose not to cover in the main document. Section 1 provides the calculation of the explicit expression of the multiphoton transition probabilities in a beam splitter, which is used as a benchmark to exhibit the interest of our developed framework. Section 2 covers the calculation of the generating functions of these probabilities as well as of the corresponding probabilities for a two-mode squeezer (whose explicit form we chose not to include as its complexity makes it of little interest). The recurrence equations on the multiphoton transition probabilities that can be deduced from these generating functions (see main text) are then discussed in the case of a beam splitter with a rational transmittance or two-mode squeezer with a rational gain. The calculation of the classical transition probabilities for distinguishable photons impinging on a beam splitter, which serves as a reference in the analysis of our main results, is summarized in Section 3. Finally, Section 4 explores the asymptotic behavior of the transition probabilities, illustrating the power of generating functions.

1. Multiphoton transition probabilities in a beam splitter

Consider a beam splitter (BS) of transmittance $\eta \in [0, 1]$ characterized by the unitary U_η^{BS} of the form

$$U_\eta^{\text{BS}} = \exp \left[\theta \left(\hat{a}^\dagger \hat{b} - \hat{a} \hat{b}^\dagger \right) \right], \quad \eta = \cos^2 \theta. \quad (22)$$

An expression for the transition amplitudes $b_n^{(i,k)} := \langle n, m | U_\eta^{\text{BS}} | i, k \rangle$ (for $i, k, n \in \mathbb{N}_0$ and $m = i + k - n$) can be computed by first deriving an expression for the following state:

$$|\psi^{(i,k)}\rangle = U_\eta^{\text{BS}} |i, k\rangle = \frac{1}{\sqrt{k!}} U_\eta^{\text{BS}} (\hat{b}^\dagger)^k U_\eta^{\text{BS}\dagger} U_\eta^{\text{BS}} |i, 0\rangle = \frac{1}{\sqrt{k!}} \left(U_\eta^{\text{BS}} \hat{b}^\dagger U_\eta^{\text{BS}\dagger} \right)^k |\psi^{(i,0)}\rangle. \quad (23)$$

Exploiting the action of the BS in phase space, we get

$$|\psi^{(i,k)}\rangle = \frac{1}{\sqrt{k!}} \left(\sqrt{1-\eta} \hat{a}^\dagger + \sqrt{\eta} \hat{b}^\dagger \right)^k |\psi^{(i,0)}\rangle = \frac{1}{\sqrt{k!}} \sum_{m=0}^k \binom{k}{m} \left(\sqrt{1-\eta} \hat{a}^\dagger \right)^m \left(\sqrt{\eta} \hat{b}^\dagger \right)^{k-m} |\psi^{(i,0)}\rangle. \quad (24)$$

Similarly, we have

$$|\psi^{(i,0)}\rangle = \frac{1}{\sqrt{i!}} \left(\sqrt{\eta} \hat{a}^\dagger - \sqrt{1-\eta} \hat{b}^\dagger \right)^i |0, 0\rangle = \sum_{n=0}^i (-1)^{i-n} \sqrt{\binom{i}{n} \eta^n (1-\eta)^{i-n}} |n, i-n\rangle. \quad (25)$$

Combining Eqs. (24) and (25), we obtain

$$|\psi^{(i,k)}\rangle = \sum_{n,m=0}^{i,k} (-1)^{i-n} \sqrt{\Gamma_{n,m}^{(i,k)} \eta^{n+k-m} (1-\eta)^{i-n+m}} |n+m, i-n+k-m\rangle, \quad (26)$$

where we defined

$$\Gamma_{n,m}^{(i,k)} = \binom{i}{n} \binom{k}{m} \binom{n+m}{n} \binom{i-n+k-m}{i-n}. \quad (27)$$

Form this, we obtain

$$b_n^{(i,k)} = \sum_{m=\max(0,n-k)}^{\min(i,n)} (-1)^{i-m} \sqrt{\Gamma_{m,n-m}^{(i,k)} \eta^{2m+k-n} (1-\eta)^{i-2m+n}}, \quad (28)$$

where

$$\Gamma_{n,m}^{(i,k)} := \binom{i}{n} \binom{k}{m} \binom{n+m}{n} \binom{i-n+k-m}{i-n}. \quad (29)$$

Thus, the transition probabilities $B_n^{(i,k)} := |b_n^{(i,k)}|^2$ in the BS U_η^{BS} are given by

$$B_n^{(i,k)} = \sum_{m,j=\max(0,n-k)}^{\min(i,n)} (-1)^{m+j} \sqrt{\Gamma_{m,n-m}^{(i,k)} \Gamma_{j,n-j}^{(i,k)}} \eta^{k-n+m+j} (1-\eta)^{i+n-m-j}. \quad (30)$$

Now, it can easily be shown that

$$\Gamma_{m,n-m}^{(i,k)} \Gamma_{j,n-j}^{(i,k)} = \gamma_{n,m,j}^{(i,k)} \gamma_{n,j,m}^{(i,k)}, \quad (31)$$

where we defined

$$\gamma_{n,m,j}^{(i,k)} := \binom{i}{m} \binom{k}{n-m} \binom{n}{j} \binom{i+k-n}{i-j}. \quad (32)$$

Since $\gamma_{n,m,j}^{(i,k)} = \gamma_{n,j,m}^{(i,k)}$, we end up with $\Gamma_{m,n-m}^{(i,k)} \Gamma_{j,n-j}^{(i,k)} = (\gamma_{n,m,j}^{(i,k)})^2$, so that

$$B_n^{(i,k)} = \sum_{m,j=\max(0,n-k)}^{\min(i,n)} (-1)^{m+j} \gamma_{n,m,j}^{(i,k)} \eta^{k-n+m+j} (1-\eta)^{i+n-m-j}. \quad (33)$$

This expression can of course be evaluated, but it is rather cumbersome and hence not very useful for the analytical investigation of multiphoton interferometry in a beam splitter. For instance, it is written as a double summation over terms with alternating signs, so that the positivity of the transition probability is not obvious from the expression. Similar derivations have for instance been given in [52],[53] and [47].

2. Generating functions of the multiphoton transition probabilities

a. Case of a beam splitter

The generating function (GF) of the transition probability $|\langle n, m | U_\eta^{\text{BS}} | i, k \rangle|^2$ in a BS is given by a function $f_\eta^{\text{BS}} : [0, 1]^2 \times [0, 1]^2 \rightarrow [0, \infty)$ defined as

$$f_\eta^{\text{BS}}(x, y, z, w) := \sum_{i,k,n,m} |\langle n, m | U_\eta^{\text{BS}} | i, k \rangle|^2 x^i y^k z^n w^m, \quad (34)$$

where, when we omit limits in summations, it means that the summation is carried out over all natural numbers in $\mathbb{N}_0$ (including 0). The trick is to realize that it can be rewritten as

$$\begin{aligned} f_\eta^{\text{BS}}(x, y, z, w) &= \text{Tr} \left[U_\eta^{\text{BS}} \left(\sum_i x^i |i\rangle \langle i| \otimes \sum_k y^k |k\rangle \langle k| \right) U_\eta^{\text{BS}\dagger} \left(\sum_n z^n |n\rangle \langle n| \otimes \sum_m w^m |m\rangle \langle m| \right) \right] \\ &= \frac{1}{(1-x)(1-y)(1-z)(1-w)} \text{Tr} [U_\eta^{\text{BS}}(\tau_x \otimes \tau_y) U_\eta^{\text{BS}\dagger}(\tau_z \otimes \tau_w)] \end{aligned}$$

where τ_t is a Gaussian thermal state of parameter t [21], i.e.,

$$\tau_t := (1-t) \sum_m t^m |m\rangle \langle m|. \quad (35)$$

Now, the object $\rho_1 := U_\eta^{\text{BS}}(\tau_x \otimes \tau_y) U_\eta^{\text{BS}\dagger}$ actually represents the effect of a beam-splitter unitary on the tensor product of two Gaussian thermal states, making it a two-mode Gaussian state. The object $\rho_2 := \tau_z \otimes \tau_w$ is obviously a two-mode Gaussian state as well. This means that $f_\eta^{\text{BS}}(x, y, z, w)$ is proportional to the overlap $\text{Tr}[\rho_1 \rho_2]$ between the two Gaussian states ρ_1 and ρ_2 ,

$$f_\eta^{\text{BS}}(x, y, z, w) = \frac{1}{(1-x)(1-y)(1-z)(1-w)} \text{Tr}[\rho_1 \rho_2]. \quad (36)$$

The above quantity can therefore be computed easily using standard tools of Gaussian quantum optics, *i.e.*, the symplectic formalism applied to the phase-space representation of bosonic quantum systems [21]. Since the first moments of each of the two Gaussian states ρ_1 and ρ_2 is zero, their overlap can be computed using the formula [35]

$$\text{Tr}[\rho_1 \rho_2] = \left(\det \left[\frac{V_1 + V_2}{2} \right] \right)^{-\frac{1}{2}} = \frac{4}{\sqrt{\det[V_1 + V_2]}}, \quad (37)$$

where V_1 and V_2 are the respective covariance matrices of ρ_1 and ρ_2 . Some easy matrix algebra involving covariance matrices and the symplectic matrix of the BS in phase space finally yields

$$f_\eta^{\text{BS}}(x, y, z, w) = \frac{1}{1 - \eta x z - (1 - \eta) x w - \eta y w - (1 - \eta) y z + x y z w}. \quad (38)$$

The conservation of energy in the BS can be easily verified using the GF given by the above equation. Define the function $\tilde{f}_\eta^{\text{BS}} : [0, 1]^2 \times [0, 1]^3 \rightarrow [0, \infty)$ as

$$\tilde{f}_\eta^{\text{BS}}(x, y, z, w, t) := \sum_{i, k, n, m} |\langle n, m | U_\eta^{\text{BS}} | i, k \rangle|^2 x^i y^k z^n w^m t^{i+k-n-m} = f_\eta^{\text{BS}}(xt, yt, \frac{z}{t}, \frac{w}{t}). \quad (39)$$

From Eq. (38), we have

$$\tilde{f}_\eta^{\text{BS}}(x, y, z, w, t) = f_\eta^{\text{BS}}(x, y, z, w), \quad \forall t. \quad (40)$$

This actually means that $\tilde{f}_\eta^{\text{BS}}$ as defined in Eq. (39) does not depend on variable t , so that the only non-zero elements in the sums of the right-hand side of Eq. (39) verify $i + k - n - m = 0$. Consequently,

$$\langle n, m | U_\eta^{\text{BS}} | i, k \rangle = 0 \quad \text{if} \quad i + k \neq n + m. \quad (41)$$

b. Symmetric inputs to the beam splitter

We now consider the case in which the same Fock states impinge on both inputs of the BS and compute the GF of the corresponding transition probabilities, which will be useful when investigating the asymptotic behavior of $B_n^{(i, i)}$ (see Section 4). The sequence $B_n^{(i, k)}$ depends on 3 indices only, index m in $|\langle n, m | U_\eta^{\text{BS}} | i, k \rangle|^2$ being redundant as a consequence of energy conservation in a BS. The GF of $B_n^{(i, k)}$ is then simply given by

$$\begin{aligned} \mathcal{T}_{i, k, n} [B_n^{(i, k)}](x, y, z) &:= \sum_{i, k, n} B_n^{(i, k)} x^i y^k z^n \\ &= \sum_{i, k, n} \left(\sum_m |\langle n, m | U_\eta^{\text{BS}} | i, k \rangle|^2 \right) x^i y^k z^n \\ &= f_\eta^{\text{BS}}(x, y, z, 1). \end{aligned}$$

In order to derive the GF of the diagonal elements $B_n^{(i,i)}$, we force the relation $k = i$ in the GF of $B_n^{(i,k)}$ by only considering the elements which satisfy it. Using the notation $c_n = [z^n]g(z)$ to mean that we select the coefficient of the z^n term in $g(z) := \sum_{n=0}^{\infty} c_n z^n$, we write

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \right] (x, z) = [y^0] \sum_{i,k,n} B_n^{(i,k)} x^i y^{k-i} z^n = [y^0] f_{\eta}^{\text{BS}} \left(\frac{x}{y}, y, z, 1 \right). \quad (42)$$

By Cauchy's integral formula for any function $g(z)$, one has

$$g(a) = \frac{1}{2\pi i} \oint \frac{g(z)}{z-a} dz. \quad (43)$$

Applying this to our case, we get that, for some circle γ_x around $y = 0$,

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \right] (x, z) = \frac{1}{2\pi i} \int_{\gamma_x} dy \frac{f_{\eta}^{\text{BS}}(x/y, y, z, 1)}{y}. \quad (44)$$

Now, using the Residue Theorem, the above equation amounts to

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \right] (x, z) = \sum_l \text{Res} \left[\frac{f_{\eta}^{\text{BS}}(x/y, y, z, 1)}{y}; y = s_l(x, z, \eta) \right], \quad (45)$$

where the s_l represent the singularities of $f_{\eta}^{\text{BS}}(x/y, y, z, 1)/y$ satisfying $\lim_{x \rightarrow 0} s_l(x, z, \eta) = 0$. Some standard calculations yield

$$s_{1,2}(x, z, \eta) = \frac{1 + xz \pm \sqrt{(1+xz)^2 - 4(\eta + (1-\eta)z)(x(1-\eta) + \eta xz)}}{2(\eta + z(1-\eta))}, \quad (46)$$

with the subscript 1(2) corresponding to $+$ ($-$) in the $\pm$ in the above equation. If we take their limits for x approaching zero, we obtain

$$\lim_{x \rightarrow 0} s_1(x, z, \eta) = 0 \quad \text{and} \quad \lim_{x \rightarrow 0} s_2(x, z, \eta) = \frac{1}{\eta + z(1-\eta)}. \quad (47)$$

The residue of the function we are interested in reduces to

$$\text{Res} \left[\frac{f_{\eta}^{\text{BS}}(x/y, y, z, 1)}{y}; y = s_1(x, z, \eta) \right] = \frac{1}{\sqrt{(1+xz)^2 - 4(\eta + (1-\eta)z)(x(1-\eta) + \eta xz)}}, \quad (48)$$

so that

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \right] (x, z) = \frac{1}{\sqrt{(1+xz)^2 - 4(\eta + (1-\eta)z)(x(1-\eta) + \eta xz)}}. \quad (49)$$

If we particularize this to a balanced BS ($\eta = 1/2$), we obtain the simple expression

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \right]_{\eta=1/2} (x, z) = \frac{1}{\sqrt{(1-x)(1-z^2x)}}, \quad (50)$$

which is the GF in i, n of the diagonal sequence $B_n^{(i,i)}$ for $\eta = 1/2$. This will be useful for analyzing the asymptotic behavior of the transition probabilities in Section 4.

c. Case of a two-mode squeezer

The GF of the transition probability $|\langle n, m | U_{\lambda}^{\text{TMS}} | i, k \rangle|^2$ in a two-mode squeezer (TMS) is given by a function $f_{\lambda}^{\text{TMS}} : [0, 1)^2 \times [0, 1]^2 \rightarrow [0, \infty)$ defined as

$$f_{\lambda}^{\text{TMS}}(x, y, z, w) := \sum_{i,k,n,m} |\langle n, m | U_{\lambda}^{\text{TMS}} | i, k \rangle|^2 x^i y^k z^n w^m, \quad (51)$$

with $\lambda = \tanh^2(r)$. One could compute it from scratch similarly as in the previous section. A much more elegant option is to use a fundamental relation that links the BS and the TMS, which was proven in [23]. There, it was shown that the TMS may be viewed as a BS undergoing “partial time reversal” [37], $|\langle n, m | U_\lambda^{\text{TMS}} | i, k \rangle|^2 = (1-\lambda) |\langle n, k | U_{1-\lambda}^{\text{BS}} | i, m \rangle|^2$. As explained in the main text, it implies that the GFs are connected by the relation

$$f_\lambda^{\text{TMS}}(x, y, z, w) = (1 - \lambda) f_{1-\lambda}^{\text{BS}}(x, w, z, y), \quad (52)$$

which then yields

$$f_\lambda^{\text{TMS}}(x, y, z, w) = \frac{1 - \lambda}{1 - \lambda(xy + zw) - (1 - \lambda)(xz + yw) + xyzw}. \quad (53)$$

d. Rational value of the transmittance or gain

We now discuss the interferometric suppression effect that exists for a BS of rational transmittance $\eta < 1$ or for a TMS of rational gain $g > 1$. It can be checked that

$$B_1^{(i,k)} = 0 \quad \text{if } \eta = \frac{k}{i+k}, \quad \text{where } i, k \in \mathbb{N}_0, i+k > 0, \quad (54)$$

which gives rise to a full interferometric suppression for any rational $\eta < 1$, extending the HOM effect for $i = k = 1$ and $\eta = 1/2$. Indeed, using the mode transformation characterizing the BS (see main text), a closed expression for $B_1^{(i,k)}$ can easily be written, which entails the sum of the amplitudes where either a single photon from mode a is transmitted (all k photons on mode b being transmitted) or a single photon from mode b is reflected (all i photons on mode a being reflected) weighted with the appropriate combinatorial factors, namely

$$B_1^{(i,k)} = \frac{i+k-1}{i k} (i\eta - k\bar{\eta})^2 B_0^{(i-1,k-1)}, \quad (55)$$

where we have used the trivial expression

$$B_0^{(i,k)} = \binom{i+k}{i} \bar{\eta}^i \eta^k. \quad (56)$$

Note that given the symmetry between the two modes as well as the input-output symmetry, we have three associated interferometric suppressions (having in common a single photon in either one of the input or output mode):

$$\begin{aligned} B_{i+k-1}^{(i,k)} &= 0 & \text{if } \eta = \frac{i}{i+k}, i+k > 0, \\ B_n^{(1,k+n-1)} &= 0 & \text{if } \eta = \frac{k}{k+n}, k+n > 0, \\ B_n^{(i+n-1,1)} &= 0 & \text{if } \eta = \frac{n}{n+i}, i+n > 0. \end{aligned} \quad (57)$$

It is instructive to examine the interference effect (54) by using the recurrence equation for a BS derived in the main text, yielding

$$B_1^{(i,k)} = \eta B_0^{(i-1,k)} + \eta B_1^{(i,k-1)} + \bar{\eta} B_1^{(i-1,k)} + \bar{\eta} B_0^{(i,k-1)} - B_0^{(i-1,k-1)}. \quad (58)$$

Note first that applying the recurrence equation to $B_0^{(i,k)}$ instead of $B_1^{(i,k)}$ gives

$$B_0^{(i,k)} = \eta B_0^{(i,k-1)} + \bar{\eta} B_0^{(i-1,k)}, \quad (59)$$

which, using Eq. (56), simply reduces to Pascal’s formula for binomial coefficients

$$\binom{i+k}{i} = \binom{i+k-1}{i} + \binom{i+k-1}{i-1}. \quad (60)$$

Thus, as expected, the recurrence equation (59) is classical and can be recovered using simple combinatorial analysis. By comparison, Eq. (58) gives a more interesting recurrence. Using the above closed expression for $B_0^{(i,k)}$, it can be rewritten as

$$B_1^{(i,k)} = \eta B_1^{(i,k-1)} + \bar{\eta} B_1^{(i-1,k)} - \kappa B_0^{(i-1,k-1)}, \quad \text{with } \kappa = 1 - (i+k-1) \left(\frac{\eta^2}{k} + \frac{\bar{\eta}^2}{i} \right). \quad (61)$$

The closed expression (55) is of course a solution of Eq. (61), but we see that the recurrence here is more interesting as it exhibits again a negative probability term. If we choose $\eta = k/(i+k)$ (with $i+k > 0$), then $\kappa = 1/(i+k)$, so the interferometric suppression effect $B_1^{(i,k)} = 0$ translates Eq. (61) into

$$k B_1^{(i,k-1)} + i B_1^{(i-1,k)} - B_0^{(i-1,k-1)} = 0. \quad (62)$$

The first and second terms can be interpreted classically : starting from the case where the output mode a already contains a single photon, the first term accounts for the k th photon in mode b being transmitted (so it does not give an extra photon on output mode a) while the second term accounts for the i th photon in mode a being reflected (so it does not give an extra photon on output mode a). Remarkably, the third (negative) term has no classical meaning and results in the full cancellation of probability $B_1^{(i,k)}$.

The same analysis can be applied to a quantum optical amplifier. As shown in the main text, we have

$$A_n^{(1,k)} = 0 \quad \text{if } \lambda = \frac{n}{n+k} \text{ or } g = 1 + \frac{n}{k}, \quad \text{where } n, k \in \mathbb{N}_0, k > 0, \quad (63)$$

which gives rise to full a interferometric suppression for any rational $\lambda < 1$ (or rational gain $g = 1/(1-\lambda) > 1$). When $k = n = 1$ and $\lambda = 1/2$, we confirm the existence of an interferometric suppression effect in a parametric amplifier of gain $g = 2$ [23]. Similarly as for a BS, we also have three associated interferometric suppressions in a TMS, namely

$$\begin{aligned} A_{i+n-1}^{(i,1)} &= 0 & \text{if } \lambda = \frac{n}{i+n} \text{ or } g = 1 + \frac{n}{i}, i > 0, \\ A_1^{(i,i+k-1)} &= 0 & \text{if } \lambda = \frac{i}{i+k} \text{ or } g = 1 + \frac{k}{i}, i > 0, \\ A_n^{(k+n-1,k)} &= 0 & \text{if } \lambda = \frac{k}{k+n} \text{ or } g = 1 + \frac{k}{n}, n > 0. \end{aligned} \quad (64)$$

Let us examine the interference Eq. (63) by using the recurrence equation for a TMS,

$$A_n^{(1,k)} = \lambda A_n^{(0,k-1)} + \lambda A_{n-1}^{(1,k)} + \bar{\lambda} A_{n-1}^{(0,k)} + \bar{\lambda} A_n^{(1,k-1)} - A_{n-1}^{(0,k-1)}. \quad (65)$$

Using the closed expression

$$A_n^{(0,k)} = (1-\lambda) B_0^{(n,k)} = \binom{n+k}{k} \lambda^n \bar{\lambda}^{k+1}, \quad (66)$$

we may reexpress it as

$$A_n^{(1,k)} = \lambda A_{n-1}^{(1,k)} + \bar{\lambda} A_n^{(1,k-1)} - \kappa' A_{n-1}^{(0,k-1)}, \quad \text{with } \kappa' = 1 - (n+k-1) \left(\frac{\lambda^2}{n} + \frac{\bar{\lambda}^2}{k} \right). \quad (67)$$

Similarly as for a BS, if we choose $\lambda = n/(n+k)$ (or $g = 1 + n/k$, with $k > 0$), the interferometric suppression $A_n^{(1,k)} = 0$ implies

$$n A_{n-1}^{(1,k)} + k A_n^{(1,k-1)} - A_{n-1}^{(0,k-1)} = 0 \quad (68)$$

Here, taking as a reference the situation where the input mode a contains a single photon, the first term accounts for the stimulated emission of a photon pair at the output (with probability $\propto \lambda$) while the second term accounts for the single photon in mode b being transmitted (with probability $\propto \bar{\lambda}$). Again, the third (negative) term has no classical interpretation and is responsible for the cancellation $A_n^{(1,k)} = 0$.

3. Multiphoton transition probabilities in a beam splitter with distinguishable photons

Consider a situation in which the photons impinging on the two input modes a and b of the BS are distinguishable. The incident photons may for instance have different polarizations. We now count the photons exiting the BS in mode a' , without making a distinction between different photons. The fact that they are distinguishable will however affect the distributions of photons in the output modes. The probability $\mathbb{P}(n \text{ in } a' | i \text{ in } a)$ that we detect n photons in output mode a' when we sent i photons in mode a is given by a simple binomial of parameter η , *i.e.*,

$$p_a(n|i) := \mathbb{P}(n \text{ in } a' | i \text{ in } a) = \binom{i}{n} \eta^n (1-\eta)^{i-n}. \quad (69)$$

Similarly, the probability $\mathbb{P}(n \text{ in } a' | k \text{ in } b)$ that we detect n photons in mode a' when we sent k photons in mode b is given by

$$p_b(n|k) := \mathbb{P}(n \text{ in } a' | k \text{ in } b) = \binom{k}{n} (1-\eta)^n \eta^{k-n}. \quad (70)$$

Using this, the probability $p(n|i, k) := \mathbb{P}(n \text{ in } a' | i \text{ in } a \text{ and } k \text{ in } b)$ that we detect n photons in mode a' when we sent i photons in mode a and k photons in mode b can be calculated using a convolution

$$p(n|i, k) = \sum_{n'=0}^n p_a(n'|i) p_b(n-n'|k). \quad (71)$$

The 3-variate GF of the sequence $p(n|i, k)$ is given by a function $g_\eta^{\text{cl}} : [0, 1]^2 \times [0, 1] \rightarrow [0, \infty)$ defined as

$$g_\eta^{\text{cl}}(x, y, z) := \mathcal{T}_{i,k,n} [p(n|i, k)](x, y, z) = \sum_{i,k,n} p(n|i, k) x^i y^k z^n. \quad (72)$$

Since the sequence $p(n|i, k)$ is given by a convolution over index n of the sequences $p_a(n|i)$ and $p_b(n|k)$, the GF $g_\eta^{\text{cl}}(x, y, z)$ is simply given by the product of their two respective GFs $g_\eta^a : [0, 1] \times [0, 1]$ and $g_\eta^b : [0, 1] \times [0, 1]$, which can simply be computed as

$$g_\eta^a(x, z) = \frac{1}{1 - \eta x z - (1 - \eta)x}, \quad g_\eta^b(y, z) = \frac{1}{1 - \eta y - (1 - \eta)yz}. \quad (73)$$

This means that g_η^{cl} satisfies the relation

$$[1 - \eta x z - (1 - \eta)x] g_\eta^{\text{cl}}(x, y, z) = g_\eta^b(y, z) = \sum_{i=0}^{\infty} g_\eta^b(y, z) \delta_{i0} x^i, \quad (74)$$

where $\delta_{..}$ denotes a Kronecker delta. Using the shifting property of the GFs, the counterpart of Eq. (74) for sequences is

$$p(n|i, k) - \eta p(n-1|i-1, k) - \bar{\eta} p(n|i-1, k) = \delta_{i0} p_a(n|k), \quad (75)$$

where $\bar{\eta} = 1 - \eta$ and $p(n|i, k) = 0$ if either of the indices n, i, k is negative. A similar reasoning yields

$$p(n|i, k) - \bar{\eta} p(n-1|i, k-1) - \eta p(n|i, k-1) = \delta_{k0} p_b(n|i). \quad (76)$$

Obviously, by summing the two relations one can always write the weaker relation

$$p(n|i, k) = \nu [\eta p(n-1|i-1, k) + \bar{\eta} p(n|i-1, k) + \delta_{i0} p_b(n|k)] \\ + (1 - \nu) [\bar{\eta} p(n-1|i, k-1) + \eta p(n|i, k-1) + \delta_{k0} p_a(n|i)] \quad (77)$$

for any $\nu \in [0, 1]$, which amounts to

$$p(n|i, k) = \begin{cases} 1, & \text{if } i = 0 \text{ and } k = 0, \\ \nu p_b(n|k) + (1 - \nu) [\bar{\eta} p(n-1|i, k-1) + \eta p(n|i, k-1)], & \text{if } i = 0 \text{ and } k \neq 0, \\ \nu [\eta p(n-1|i-1, k) + \bar{\eta} p(n|i-1, k)] + (1 - \nu) p_a(n|i), & \text{if } i \neq 0 \text{ and } k = 0, \\ \nu [\eta p(n-1|i-1, k) + \bar{\eta} p(n|i-1, k)] \\ \quad + (1 - \nu) [\bar{\eta} p(n-1|i, k-1) + \eta p(n|i, k-1)], & \text{else.} \end{cases} \quad (78)$$

The last relation can be written more simply as

$$p(n|i, k) = \begin{cases} 1, & \text{if } i = 0 \text{ and } k = 0, \\ p_b(n|k), & \text{if } i = 0 \text{ and } k \neq 0, \\ p_a(n|i), & \text{if } i \neq 0 \text{ and } k = 0, \\ \nu [\eta p(n-1|i-1, k) + \bar{\eta} p(n|i-1, k)] \\ + (1-\nu) [\bar{\eta} p(n-1|i, k-1) + \eta p(n|i, k-1)], & \text{else.} \end{cases} \quad (79)$$

4. Asymptotics of the transition probabilities

The asymptotic behavior of a sequence $\{c_n\}$ for a growing index can be studied by analyzing the asymptotic behavior of the corresponding GF $g(z)$ around its singularities. This is encompassed in the Tauberian theorems [54], the most famous of which being due to Hardy, Littlewood [55] and Karamata [56].

The HLK Tauberian theorem. *Let $g(z)$ be a power series with radius of convergence equal to 1, satisfying*

$$g(z) \sim \frac{1}{(1-z)^\alpha} \Lambda\left(\frac{1}{1-z}\right), \quad z \rightarrow 1, \quad (80)$$

for some $\alpha \geq 0$ with Λ a slowly varying function. Assume that the coefficients $c_n = [z^n]g(z)$ are all non-negative. Then

$$\sum_{k=0}^n c_k \sim \frac{n^\alpha}{\Gamma(\alpha+1)} \Lambda(n), \quad n \rightarrow \infty. \quad (81)$$

A function Λ is said to be slowly varying at infinity if and only if, for any $\beta > 0$, one has

$$\frac{\Lambda(\beta x)}{\Lambda(x)} \rightarrow 1 \quad \text{as } x \rightarrow +\infty. \quad (82)$$

Our aim is now to use Tauberian theorems in order to study the asymptotic behavior of $B_n^{(i,i)}$ for $\eta = 1/2$. The HLK Tauberian theorem can be generalized, and in case of multiple singularities, each one can be analyzed separately, and the different contributions can be combined in the end [54]. In our case, this must be done in two steps, since our sequence has two indices i and n . We begin by analyzing the behavior of

$$[z^n] \mathcal{T}_{i,n} \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x, z) = \mathcal{T}_i \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x), \quad (83)$$

the GF in i , by studying the behavior of

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x, z), \quad (84)$$

the GF in i and n . We then investigate the resulting

$$\mathcal{T}_i \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x) \quad (85)$$

in order to conclude about

$$B_n^{(i,i)} \Big|_{\eta=1/2}. \quad (86)$$

Behavior of $\mathcal{T}_i \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x)$. The function given in Eq. (50) has two singularities $z_1(x) := 1/\sqrt{x}$ and $z_2(x) := -1/\sqrt{x}$. First,

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x, z) \sim \frac{1}{\sqrt{2(1-x)(1-\sqrt{x}z)}}, \quad \text{when } z \rightarrow z_1(x). \quad (87)$$

Define the sequence $\beta_{i,n}^{(1)}$ such that

$$\sum_{i,n} \beta_{i,n}^{(1)} x^i z^n = \frac{1}{\sqrt{2(1-x)(1-\sqrt{xz})}}. \quad (88)$$

In other words,

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x, z) \sim \sum_{i,n} \beta_{i,n}^{(1)} x^i z^n, \quad \text{when } z \rightarrow z_1(x). \quad (89)$$

Equation (88) is the same as (x is positive)

$$\sum_{i,n} \beta_{i,n}^{(1)} x^i \left(\frac{z}{\sqrt{x}} \right)^n = \frac{1}{\sqrt{2(1-x)(1-z)}}, \quad (90)$$

or,

$$\sum_{i,n} \beta_{i,n}^{(1)} x^{i-\frac{n}{2}} z^n = \frac{1}{\sqrt{2(1-x)(1-z)}}. \quad (91)$$

Now, for n increasing, according to the Tauberian theorems,

$$[z^n] \frac{1}{\sqrt{2(1-x)(1-z)}} \sim \frac{1}{\sqrt{2(1-x)\pi n}}, \quad (92)$$

so that

$$[z^n] \sum_{i,n} \beta_{i,n}^{(1)} x^{i-\frac{n}{2}} z^n \sim \frac{1}{\sqrt{2(1-x)\pi n}}, \quad (93)$$

$$[z^n] \sum_{i,n} \beta_{i,n}^{(1)} x^i z^n \sim \frac{x^{\frac{n}{2}}}{\sqrt{2(1-x)\pi n}}. \quad (94)$$

Using Definition (88), we end up with

$$[z^n] \frac{1}{\sqrt{2(1-x)(1-\sqrt{xz})}} \sim \frac{x^{\frac{n}{2}}}{\sqrt{2(1-x)\pi n}}. \quad (95)$$

Secondly,

$$\mathcal{T}_{i,n} \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x, z) \sim \frac{1}{\sqrt{2(1-x)(1+\sqrt{xz})}}, \quad \text{when } z \rightarrow z_2(x). \quad (96)$$

We can do the same analysis, and obtain

$$[z^n] \frac{1}{\sqrt{2(1-x)(1+\sqrt{xz})}} \sim \frac{(-1)^n x^{\frac{n}{2}}}{\sqrt{2(1-x)\pi n}}. \quad (97)$$

As we explained earlier, in the case of two singularities (having the same absolute value), the two asymptotic contributions can be added up [54], so that

$$[z^n] \mathcal{T}_{i,n} \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x, z) \sim \frac{x^{\frac{n}{2}}}{\sqrt{2(1-x)\pi n}} + \frac{(-1)^n x^{\frac{n}{2}}}{\sqrt{2(1-x)\pi n}}, \quad (98)$$

or,

$$\mathcal{T}_i \left[B_n^{(i,i)} \Big|_{\eta=1/2} \right] (x) \sim \frac{1 + (-1)^n}{\sqrt{2\pi n}} \frac{x^{\frac{n}{2}}}{\sqrt{1-x}}. \quad (99)$$

The zero contribution for odd n is consistent with the fact that the total input photon number $2i$ is even.

Behavior of $B_n^{(i,i)}|_{\eta=1/2}$. The function on the right-hand side of Eq. (99) has only one singularity, $x_0 := 1$. Since the dominant factor is $1/\sqrt{1-x}$ (compared to $x^{\frac{n}{2}}$) when $x \rightarrow x_0$, we can focus on it. We have [54]

$$[x^i] \frac{1}{\sqrt{1-x}} \sim \frac{1}{\sqrt{\pi i}}, \quad (100)$$

meaning that

$$[x^i] \left(x^{-\frac{n}{2}} \mathcal{T}_i \left[B_n^{(i,i)}|_{\eta=1/2} \right] (x) \right) \sim \frac{1 + (-1)^n}{\sqrt{2\pi n}} \frac{1}{\sqrt{\pi i}}. \quad (101)$$

Now,

$$\begin{aligned} x^{-\frac{n}{2}} \mathcal{T}_i \left[B_n^{(i,i)}|_{\eta=1/2} \right] (x) &= x^{-\frac{n}{2}} \sum_{i=0}^{\infty} B_n^{(i,i)}|_{\eta=1/2} x^i \\ &= \sum_{i=0}^{\infty} B_n^{(i,i)}|_{\eta=1/2} x^{i-\frac{n}{2}} \\ &= \sum_{j=-n/2}^{\infty} B_n^{(j+\frac{n}{2}, j+\frac{n}{2})}|_{\eta=1/2} x^j, \end{aligned}$$

and $B_n^{(i,i)} = 0$ if $n > 2i$, so that

$$x^{-\frac{n}{2}} \mathcal{T}_i \left[B_n^{(i,i)}|_{\eta=1/2} \right] (x) = \sum_{j=0}^{\infty} B_n^{(j+\frac{n}{2}, j+\frac{n}{2})}|_{\eta=1/2} x^j, \quad (102)$$

and

$$[x^i] \left(x^{-\frac{n}{2}} \mathcal{T}_i \left[B_n^{(i,i)}|_{\eta=1/2} \right] (x) \right) = B_n^{(i+\frac{n}{2}, i+\frac{n}{2})}|_{\eta=1/2}. \quad (103)$$

As a consequence of Equation (101),

$$B_n^{(i+\frac{n}{2}, i+\frac{n}{2})}|_{\eta=1/2} \sim \frac{1 + (-1)^n}{\sqrt{2\pi n}} \frac{1}{\sqrt{\pi i}}, \quad (104)$$

or,

$$B_n^{(i,i)}|_{\eta=1/2} \sim \frac{1 + (-1)^n}{\sqrt{2\pi n}} \frac{1}{\sqrt{\pi \left(i - \frac{n}{2}\right)}}. \quad (105)$$

After some simplification, we obtain

$$B_n^{(i,i)}|_{\eta=1/2} \sim \frac{1 + (-1)^n}{\pi \sqrt{n(2i - n)}}. \quad (106)$$

which exactly coincides with the result of the analysis performed in [44]. The output terms around $n \sim i$ are maximally suppressed, which is reminiscent of the HOM effect. Interestingly, we can again exploit partial time reversal and extend this analysis to a TMS with $\lambda = 1/2$, giving

$$A_k^{(i,k)} \sim \frac{1 + (-1)^i}{2\pi \sqrt{i(2k - i)}}, \quad k, i \rightarrow \infty. \quad (107)$$