UNIQUENESS OF CONVEX ANCIENT SOLUTIONS TO MEAN CURVATURE FLOW IN HIGHER DIMENSIONS

SIMON BRENDLE AND KYEONGSU CHOI

ABSTRACT. In this paper, we consider noncompact ancient solutions to the mean curvature flow in \mathbb{R}^{n+1} $(n \geq 3)$ which are strictly convex, uniformly two-convex, and noncollapsed. We prove that such an ancient solution is a rotationally symmetric translating soliton.

1. INTRODUCTION

Throughout this paper, we fix an integer $n \geq 3$. Our goal in this paper is to classify all noncompact ancient solutions to mean curvature flow in \mathbb{R}^{n+1} which are convex, uniformly two-convex, and noncollapsed in the sense of Sheng and Wang [12]:

Theorem 1.1. Let M_t , $t \in (-\infty, 0]$, be a noncompact ancient solution of mean curvature flow in \mathbb{R}^{n+1} which is strictly convex, uniformly two-convex, and noncollapsed. Then M_t is a rotationally symmetric translating soliton.

If we evolve a closed, embedded, two-convex hypersurface by mean curvature flow, then it is well known that any blow-up limit is an ancient solution which is weakly convex, uniformly two-convex, and noncollapsed (see [7], Theorem 1.10, or [14],[15]). If we combine this result with Theorem 1.1, we obtain the following result:

Corollary 1.2. Consider an arbitrary closed, embedded, two-convex hypersurface in \mathbb{R}^{n+1} , and evolve it by mean curvature flow. At the first singular time, the only possible blow-up limits are shrinking round spheres; shrinking round cylinders; and the unique rotationally symmetric translating soliton.

In a recent paper [2], we obtained a classification of noncompact ancient solutions in \mathbb{R}^3 which are convex and noncollapsed. The proof of Theorem 1.1 draws on similar techniques. In Section 2, we derive asymptotic estimates for the solution in the cylindrical region. These estimates tell us that, for -t large, the rescaled surface $(-t)^{-\frac{1}{2}} M_t \cap B_{5n}(0)$ is $O((-t)^{-\frac{1}{2}})$ -close to a cylinder of radius $\sqrt{2(n-1)}$. In Section 3, we combine this estimate with a barrier argument in the spirit of [1] to conclude that $\liminf_{t\to-\infty} H_{\max}(t) >$ 0, where $H_{\max}(t)$ denotes the supremum of the mean curvature of M_t .

The first author was supported by the National Science Foundation under grant DMS-1649174 and by the Simons Foundation. The second author was supported by the National Science Foundation under grant DMS-1811267.

In Section 4, we establish a higher-dimensional version of the Neck Improvement Theorem in [2]. This step requires significant modifications in the higher-dimensional setting. In order to formulate the Neck Improvement Theorem, we need a notion of ε -symmetry in higher dimensions, which generalizes the one introduced in [2]. We say that a point (\bar{x}, \bar{t}) in spacetime is ε -symmetric if there exists a collection of normalized rotation vector fields $\mathcal{K} = \{K_{\alpha} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ (all having a common axis of rotation) such that $|K_{\alpha}| H \leq 10n$ at (\bar{x}, \bar{t}) and $|\langle K_{\alpha}, \nu \rangle| H \leq \varepsilon$ in the parabolic neighborhood $\hat{\mathcal{P}}(\bar{x}, \bar{t}, 10, 100)$. The main difference between the two-dimensional case and the higher-dimensional case is that, instead of a single rotation vector field in ambient space, we need to consider a collection of normalized rotation vector fields which share a common axis. The statement of the Neck Improvement Theorem can be summarized as follows: if (\bar{x}, \bar{t}) lies on a neck and every point in a sufficiently large parabolic neighborhood of (\bar{x}, \bar{t}) is ε -symmetric, then the point (\bar{x}, \bar{t}) itself is $\frac{\varepsilon}{2}$ -symmetric.

In Section 5, we iterate the Neck Improvement Theorem to conclude that any ancient solution which satisfies the assumption of Theorem 1.1 is rotationally symmetric. Finally, in Section 6, we classify ancient solutions with rotational symmetry, thereby completing the proof of Theorem 1.1.

2. Asymptotic analysis as $t \to -\infty$

Suppose that M_t , $t \in (-\infty, 0]$, is a noncompact ancient solution of mean curvature flow in \mathbb{R}^{n+1} which is strictly convex, uniformly two-convex, and noncollapsed. We consider the rescaled flow $\bar{M}_{\tau} = e^{\frac{\tau}{2}} M_{-e^{-\tau}}$. The hypersurfaces \bar{M}_{τ} move with velocity $-(H - \frac{1}{2} \langle x, \nu \rangle)\nu$.

Proposition 2.1 (cf. Hashofer-Kleiner [7], Theorem 1.11). Consider a sequence $\tau_j \to -\infty$. After passing to a subsequence, the rescaled hypersurfaces \bar{M}_{τ_j} converge in C_{loc}^{∞} to a cylinder of radius $\sqrt{2(n-1)}$ with axis passing through the origin.

Proof. This is proved in [7] on p. 533. We briefly sketch the argument for the convenience of the reader. Let $t_j := -e^{-\tau_j}$. A standard barrier argument implies dist $(0, M_{t_j}) \leq C(n) (-t_j)^{\frac{1}{2}}$ for j sufficiently large. By the speed limit lemma (cf. Lemma 3.4 in [7]), there exist a time $\bar{t}_j \in [t_j, \frac{t_j}{2}]$ and a point $\bar{p}_j \in M_{\bar{t}_j}$ such that $|\bar{p}_j| \leq C(n) (-t_j)^{\frac{1}{2}}$ and $H(\bar{p}_j, \bar{t}_j) \leq C(n) (-t_j)^{-\frac{1}{2}}$. Using the global curvature estimate in [7] we conclude that, after passing to a subsequence, the rescaled hypersurfaces $\bar{M}_{\tau_j} = (-t_j)^{-\frac{1}{2}} M_{t_j}$ converge in C_{loc}^{∞} to a smooth limit. By Huisken's monotonicity formula [9], the limit must be a self-similar shrinker. Results of Colding and Minicozzi [3] then imply that the limit is either a sphere of radius $\sqrt{2n}$ centered at the origin or a cylinder of radius $\sqrt{2(n-1)}$ with axis passing through the origin. Since the ancient solution M_t is noncompact, the backward limit cannot be a sphere. This completes the proof of Proposition 2.1.

In the following, we denote by $\Sigma = \{x \in \mathbb{R}^{n+1} : x_1^2 + \ldots + x_n^2 = 2(n-1)\}$ the cylinder of radius $\sqrt{2(n-1)}$ around the x_{n+1} -axis.

Proposition 2.2. For each τ , we have

$$\int_{\bar{M}_{\tau}} e^{-\frac{|x|^2}{4}} \le \int_{\Sigma} e^{-\frac{|x|^2}{4}}.$$

Proof. Let us consider an arbitrary sequence $t_j \to -\infty$. The local version of Huisken's monotonicity formula [9] implies that, for each j and each r > 0, the function

$$t \mapsto (-t)^{-\frac{n}{2}} \int_{M_t} e^{-\frac{|x|^2}{(-4t)}} \left(1 - \frac{|x|^2 + 2nt}{(-r^2 t_j)}\right)_+^3$$

is monotone decreasing for $t \in [t_j, 0)$ (see [4], pp. 64–65). This gives

$$(-t)^{-\frac{n}{2}} \int_{M_t} e^{-\frac{|x|^2}{(-4t)}} \left(1 - \frac{|x|^2}{(-r^2 t_j)}\right)_+^3$$

$$\leq (-t_j)^{-\frac{n}{2}} \int_{M_{t_j}} e^{-\frac{|x|^2}{(-4t_j)}} \left(1 + \frac{2n}{r^2} - \frac{|x|^2}{(-r^2 t_j)}\right)_+^3$$

for each $t \in [t_j, 0)$ and each r > 0. We now send $j \to \infty$, keeping t and r fixed. By the monotone convergence theorem,

$$\lim_{j \to \infty} (-t)^{-\frac{n}{2}} \int_{M_t} e^{-\frac{|x|^2}{(-4t)}} \left(1 - \frac{|x|^2}{(-r^2 t_j)}\right)_+^3 = (-t)^{-\frac{n}{2}} \int_{M_t} e^{-\frac{|x|^2}{(-4t)}}$$

Moreover,

$$\lim_{j \to \infty} (-t_j)^{-\frac{n}{2}} \int_{M_{t_j}} e^{-\frac{|x|^2}{(-4t_j)}} \left(1 + \frac{2n}{r^2} - \frac{|x|^2}{(-r^2t_j)} \right)_+^3 = \int_{\Sigma} e^{-\frac{|x|^2}{4}} \left(1 + \frac{2n}{r^2} - \frac{|x|^2}{r^2} \right)_+^3$$

by Proposition 2.1. Thus, we conclude that

$$(-t)^{-\frac{n}{2}} \int_{M_t} e^{-\frac{|x|^2}{(-4t)}} \le \int_{\Sigma} e^{-\frac{|x|^2}{4}} \left(1 + \frac{2n}{r^2} - \frac{|x|^2}{r^2}\right)_+^3$$

for each t and each r > 0. Sending $r \to \infty$, we obtain

$$(-t)^{-\frac{n}{2}} \int_{M_t} e^{-\frac{|x|^2}{(-4t)}} \le \int_{\Sigma} e^{-\frac{|x|^2}{4}}.$$

This completes the proof of Proposition 2.2.

Lemma 2.3. We can find a smooth function $\rho(\tau)$ and a function $Q(\tau)$ taking values in SO(n+1) with the following properties:

- $\lim_{\tau \to -\infty} \rho(\tau) = \infty$. $-\rho(\tau) \le \rho'(\tau) \le 0$.

• In the ball $B_{2\rho(\tau)}(0)$, the rotated surface $Q(\tau)M_{\tau}$ can be written as a graph over the standard cylinder Σ , and the C^4 -norm of the height function is less than $\rho(\tau)^{-8}$.

Proof. We can find a smooth function $Q(\tau)$ taking values in SO(n+1) such that the rotated surfaces $Q(\tau)\bar{M}_{\tau}$ converge to the standard cylinder Σ in C_{loc}^{∞} . Hence, we can find a function $\bar{\rho}(\tau)$ with the following properties:

• $\lim_{\tau \to -\infty} \bar{\rho}(\tau) = \infty$.

4

• In the ball $B_{2\bar{\rho}(\tau)}(0)$, the rotated surface $Q(\tau)\bar{M}_{\tau}$ can be written as a graph over the standard cylinder Σ , and the C^4 -norm of the height function is less than $\bar{\rho}(\tau)^{-8}$.

We define a function $\rho(\tau)$ by $\rho(\tau) := \inf_{\tau'} \max\{e^{\tau'-\tau}, 1\} \bar{\rho}(\tau')$. Clearly, $\rho(\tau) \leq \bar{\rho}(\tau)$ for each τ , and $\rho(\tau) \to \infty$ as $\tau \to -\infty$. Moreover, the function $\tau \mapsto \rho(\tau)$ is monotone decreasing, and the function $\tau \mapsto e^{\tau} \rho(\tau)$ is monotone increasing. This gives $-\rho(\tau) \leq \rho'(\tau) \leq 0$. Finally, a standard convolution argument allows us to replace the function $\rho(\tau)$ by a smooth function. This completes the proof of Lemma 2.3.

As in [2], it is necessary to fine tune the choice of the rotation matrix. Let $\varphi \ge 0$ be a smooth cutoff function such that $\varphi = 1$ on $\left[-\frac{1}{2}, \frac{1}{2}\right]$ and $\varphi = 0$ outside $\left[-\frac{2}{3}, \frac{2}{3}\right]$.

Proposition 2.4. Let $\rho(\tau)$ be chosen as in Lemma 2.3. We can find a function $S(\tau)$ taking values in SO(n+1) with the following properties:

• In the ball $B_{2\rho(\tau)}(0)$, the rotated surface $\tilde{M}_{\tau} := S(\tau) \bar{M}_{\tau}$ can be written as a graph over the standard cylinder Σ of some function $u(\cdot, \tau)$; that is,

 $\{x+u(x,\tau)\nu_{\Sigma}(x): x\in \Sigma\cap B_{2\rho(\tau)}(0)\}\subset \tilde{M}_{\tau},$

where ν_{Σ} denotes the unit normal to Σ and $||u(\cdot, \tau)||_{C^4(\Sigma \cap B_{2\rho(\tau)}(0))} \leq \rho(\tau)^{-4}$.

• The function $u(\cdot, \tau)$ satisfies the orthogonality relations

$$\int_{\Sigma \cap B_{\rho(\tau)}(0)} e^{-\frac{|x|^2}{4}} \langle Ax, \nu_{\Sigma} \rangle u(x,\tau) \varphi\left(\frac{x_{n+1}}{\rho(\tau)}\right) = 0$$

for every matrix $A \in so(n+1)$.

• The matrix $A(\tau) := S'(\tau)S(\tau)^{-1} \in so(n+1)$ satisfies $A(\tau)_{ij} = 0$ for $i, j \in \{1, ..., n\}$.

Proof. Given any unit vector $\omega \in S^n$, we consider the rotated cylinder $\Sigma_{\omega} := \{x \in \mathbb{R}^{n+1} : |x - \langle x, \omega \rangle \omega| = \sqrt{2(n-1)}\}$. Note that $\Sigma_{\omega} = \Sigma$ if $\omega = (0, \ldots, 0, 1)$. For each $\omega \in S^n$, we denote by $\pi_{\omega} : \mathbb{R}^{n+1} \to \Sigma_{\omega}$ the radial projection to Σ_{ω} . In other words,

$$\pi_{\omega}(y) = \langle y, \omega \rangle \omega + \sqrt{2(n-1)} \frac{y - \langle y, \omega \rangle \omega}{|y - \langle y, \omega \rangle \omega|}$$

For each $\omega \in S^n$, we define a vector $I_{\tau}(\omega) \in \mathbb{R}^{n+1}$ by

$$I_{\tau}(\omega) := \int_{\bar{M}_{\tau}} e^{-\frac{2(n-1)+\langle y,\omega\rangle^2}{4}} \left(|y-\langle y,\omega\rangle\omega| - \sqrt{2(n-1)} \right) \varphi\left(\frac{\langle y,\omega\rangle}{\rho(\tau)}\right)$$
$$\cdot \det D(\pi_{\omega}|_{\bar{M}_{\tau}})(y) \cdot \langle y,\omega\rangle \frac{y-\langle y,\omega\rangle\omega}{|y-\langle y,\omega\rangle\omega|}.$$

Clearly, $I_{\tau}(\omega)$ is orthogonal to ω .

Let $Q(\tau)$ be the rotation matrix defined in Lemma 2.3, and define $\zeta(\tau) \in S^n$ by $Q(\tau)\zeta(\tau) = (0, \ldots, 0, 1)$. Using Lemma 2.3, we obtain

 $|I_{\tau}(\zeta(\tau))| \le C \,\rho(\tau)^{-8}.$

Moreover, an analysis of the linearization of the function $I_{\tau}(\omega)$ near the point $\zeta(\tau)$ gives

$$|I_{\tau}(\omega) - I_{\tau}(\tilde{\omega}) + c(n) (\omega - \tilde{\omega})| \le o(1) |\omega - \tilde{\omega}|$$

if $|\omega - \zeta(\tau)| \leq \rho(\tau)^{-6}$ and $|\tilde{\omega} - \zeta(\tau)| \leq \rho(\tau)^{-6}$. Here, c(n) is a non-zero constant that depends only on n. We now apply a standard fixed point theorem to the map which sends $\omega \in S^n$ to $\frac{I_{\tau}(\omega) + c(n)\omega}{|I_{\tau}(\omega) + c(n)\omega|} \in S^n$. Hence, if $-\tau$ is sufficiently large, then we can find a vector $\tilde{\zeta}(\tau) \in S^n$ such that $|\zeta(\tau) - \tilde{\zeta}(\tau)| \leq C \rho(\tau)^{-8}$ and

$$\frac{I_{\tau}(\tilde{\zeta}(\tau)) + c(n)\,\tilde{\zeta}(\tau)}{|I_{\tau}(\tilde{\zeta}(\tau)) + c(n)\,\tilde{\zeta}(\tau)|} = \tilde{\zeta}(\tau).$$

Since $I_{\tau}(\tilde{\zeta}(\tau))$ is orthogonal to $\tilde{\zeta}(\tau)$, we conclude that $I_{\tau}(\tilde{\zeta}(\tau)) = 0$. Moreover, $\tilde{\zeta}(\tau)$ depends smoothly on τ .

In the next step, we choose a smooth function $S(\tau)$ taking values in SO(n+1) such that $S(\tau) \,\tilde{\zeta}(\tau) = (0, \ldots, 0, 1)$. Moreover, we can choose $S(\tau)$ in such a way that the matrix $A(\tau) := S'(\tau)S(\tau)^{-1} \in so(n+1)$ satisfies $A(\tau)_{ij} = 0$ for $i, j \in \{1, \ldots, n\}$. (Otherwise, we replace $S(\tau)$ by $R(\tau)S(\tau)$, where $R(\tau) \in SO(n) \subset SO(n+1)$ is a rotation which fixes the x_{n+1} -axis.)

Let $\tilde{M}_{\tau} := S(\tau) \, \bar{M}_{\tau}$. Using Lemma 2.3 and the estimate $|\zeta(\tau) - \tilde{\zeta}(\tau)| \leq C \, \rho(\tau)^{-8}$, we conclude that the surface \tilde{M}_{τ} can be written as a graph over the standard cylinder Σ of some function $u(\cdot, \tau)$, where $||u(\cdot, \tau)||_{C^4(\Sigma \cap B_{2\rho(\tau)}(0))} \leq \rho(\tau)^{-4}$. Finally, the identity $I_{\tau}(\tilde{\zeta}(\tau)) = 0$ gives

$$\int_{\tilde{M}_{\tau}} e^{-\frac{2(n-1)+\langle y,\omega\rangle^2}{4}} \left(|y-\langle y,\omega\rangle\omega| - \sqrt{2(n-1)} \right) \varphi\left(\frac{\langle y,\omega\rangle}{\rho(\tau)}\right)$$
$$\cdot \det D(\pi_{\omega}|_{\tilde{M}_{\tau}})(y) \cdot \langle y,\omega\rangle \frac{y-\langle y,\omega\rangle\omega}{|y-\langle y,\omega\rangle\omega|} = 0$$

if $\omega = (0, \ldots, 0, 1)$ is the vertical unit vector. This finally implies

$$\int_{\Sigma} e^{-\frac{2(n-1)+x_{n+1}^2}{4}} u(x) \varphi\left(\frac{x_{n+1}}{\rho(\tau)}\right) x_{n+1} x_i = 0$$

for $i \in \{1, ..., n\}$. From this, the desired orthogonality relations follow. This completes the proof of Proposition 2.4.

We next recall the shrinker foliation constructed in [1]. By [1], the union of all the leaves in this foliation contains a truncated cone of the form $\{x \in \mathbb{R}^{n+1} : |x_{n+1}| \geq z_0, x_1^2 + \ldots + x_n^2 \leq b_0^2 x_{n+1}^2\}$ for some large constant z_0 and some small constant $b_0 > 0$. We denote by ν_{fol} the unit normal vector field to this foliation. Moreover, we denote by Δ_{τ} the region in between the cylinder Σ and the surface \tilde{M}_{τ} .

Proposition 2.5. There exists a constant L_0 such that for all $L \in [L_0, \rho(\tau)]$

$$\int_{\tilde{M}_{\tau} \cap \{|x_{n+1}| \ge L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_{n+1}| \ge L\}} e^{-\frac{|x|^2}{4}} \ge - \int_{\Delta_{\tau} \cap \{|x_{n+1}| = L\}} e^{-\frac{|x|^2}{4}} |\langle \omega, \nu_{\text{fol}} \rangle|.$$

Here, $\omega = (0, \ldots, 0, 1)$ denotes the vertical unit vector in \mathbb{R}^{n+1} .

Proof. This follows from the identity $\operatorname{div}(e^{-\frac{|x|^2}{4}}\nu_{\text{fol}}) = 0$ together with the divergence theorem. See [2], Proposition 2.2, for details.

Proposition 2.6. There exists a constant L_0 such that

$$\int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} |\nabla^{\Sigma} u(x,\tau)|^2 \le C \int_{\Sigma \cap \{|x_{n+1}| \le \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u(x,\tau)^2$$

and

$$\int_{\Sigma \cap \{\frac{L}{2} \le |x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} u(x,\tau)^2 \le CL^{-2} \int_{\Sigma \cap \{|x_{n+1}| \le \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u(x,\tau)^2$$

for all $L \in [L_0, \rho(\tau)]$.

Proof. Let $\omega = (0, \ldots, 0, 1)$ denote the vertical unit vector in \mathbb{R}^{n+1} . Lemma 4.11 in [1] implies that $|\langle \omega, \nu_{\text{fol}} \rangle| \leq CL^{-1} |x_1^2 + \ldots + x_n^2 - 2(n-1)|$ for each point $x \in \Delta_{\tau} \cap \{|x_{n+1}| = L\}$. This gives

$$\int_{\Delta_{\tau} \cap \{|x_{N+1}| = L\}} e^{-\frac{|x|^2}{4}} |\langle \omega, \nu_{\text{fol}} \rangle|$$

$$\leq CL^{-1} \int_{\Delta_{\tau} \cap \{|x_{n+1}| = L\}} e^{-\frac{|x|^2}{4}} |x_1^2 + \ldots + x_n^2 - 2(n-1)|$$

$$\leq CL^{-1} \int_{\Sigma \cap \{|x_{n+1}| = L\}} e^{-\frac{|x|^2}{4}} u^2.$$

Combining this estimate with Proposition 2.5 yields

$$\int_{\tilde{M}_{\tau} \cap \{|x_{n+1}| \ge L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_{n+1}| \ge L\}} e^{-\frac{|x|^2}{4}} \ge -CL^{-1} \int_{\Sigma \cap \{|x_{n+1}| = L\}} e^{-\frac{|x|^2}{4}} u^2.$$

We next observe that

$$\begin{split} \int_{\tilde{M}_{\tau} \cap \{|x_{n+1}| \leq L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_{n+1}| \leq L\}} e^{-\frac{|x|^2}{4}} \\ &= \int_{-L}^{L} \left(\int_{S^{n-1}} e^{-\frac{z^2}{4}} \left[e^{-\frac{(\sqrt{2(n-1)}+u)^2}{4}} \left(\sqrt{2(n-1)}+u\right)^{n-2} \right. \\ &\left. \cdot \sqrt{\left(\sqrt{2(n-1)}+u\right)^2 \left(1+\left(\frac{\partial u}{\partial z}\right)^2\right) + |\nabla^{S^{n-1}}u|^2} \right. \\ &\left. - e^{-\frac{n-1}{2}} \sqrt{2(n-1)}^{n-1} \right] \right) dz. \end{split}$$

By assumption, the height function u satisfies $|u| + |\frac{\partial u}{\partial z}| + |\nabla^{S^{n-1}}u| \le o(1)$ for $|x_{n+1}| \le L$. From this, we deduce that

$$\begin{split} &\int_{\tilde{M}_{\tau} \cap \{|x_{n+1}| \leq L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_{n+1}| \leq L\}} e^{-\frac{|x|^2}{4}} \\ &\geq \int_{-L}^{L} \left(\int_{S^{n-1}} e^{-\frac{z^2}{4}} \left[e^{-\frac{(\sqrt{2(n-1)}+u)^2}{4}} \left(\sqrt{2(n-1)} + u\right)^{n-1} \right. \\ &\left. - e^{-\frac{n-1}{2}} \sqrt{2(n-1)}^{n-1} + \frac{1}{C} \left| \nabla^{\Sigma} u \right|^2 \right] \right) dz \\ &\geq \int_{-L}^{L} \left(\int_{S^{n-1}} e^{-\frac{z^2}{4}} \left[-Cu^2 + \frac{1}{C} \left| \nabla^{\Sigma} u \right|^2 \right] \right) dz \end{split}$$

where C is a large constant that depends only on n. Putting these facts together, we conclude that

$$\int_{\tilde{M}_{\tau}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma} e^{-\frac{|x|^2}{4}} \ge \int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} \left[-Cu^2 + \frac{1}{C} |\nabla^{\Sigma} u|^2 \right] \\ - CL^{-1} \int_{\Sigma \cap \{|x_{n+1}| = L\}} e^{-\frac{|x|^2}{4}} u^2.$$

Combining this estimate with Proposition 2.2, we deduce that

$$\int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} |\nabla^{\Sigma} u|^2$$

$$\leq C \int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} u^2 + CL^{-1} \int_{\Sigma \cap \{|x_{n+1}| = L\}} e^{-\frac{|x|^2}{4}} u^2.$$

On the other hand, using the divergence theorem, we obtain

$$L \int_{\Sigma \cap \{|x_{n+1}| = L\}} e^{-\frac{|x|^2}{4}} u^2 = \int_{\Sigma \cap \{|x_{n+1}| \le L\}} \operatorname{div}_{\Sigma} (e^{-\frac{|x|^2}{4}} u^2 x^{\tan})$$

$$= \int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} \left(u^2 - \frac{1}{2} x_{n+1}^2 u^2 + 2u \langle x^{\tan}, \nabla^{\Sigma} u \rangle \right)$$

$$\leq \int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} \left(u^2 - \frac{1}{4} x_{n+1}^2 u^2 + 4 |\nabla^{\Sigma} u|^2 \right),$$

and consequently

$$L^{2} \int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^{2}}{4}} u^{2} + L \int_{\Sigma \cap \{|x_{n+1}| = L\}} e^{-\frac{|x|^{2}}{4}} u^{2}$$

$$\leq C \int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^{2}}{4}} |\nabla^{\Sigma} u|^{2} + CL^{2} \int_{\Sigma \cap \{|x_{n+1}| \le \frac{L}{2}\}} e^{-\frac{|x|^{2}}{4}} u^{2}.$$

To summarize, we have shown that

$$\int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} |\nabla^{\Sigma} u|^2$$

$$\leq CL^{-2} \int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} |\nabla^{\Sigma} u|^2 + C \int_{\Sigma \cap \{|x_{n+1}| \le \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u^2.$$

If L is sufficiently large, we can absorb the first term on the right hand side into the left hand side. This gives

$$\int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} |\nabla^{\Sigma} u|^2 \le C \int_{\Sigma \cap \{|x_{n+1}| \le \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u^2.$$

This proves the first statement. Using the inequality

$$0 \le \int_{\Sigma \cap \{|x_{n+1}| \le L\}} e^{-\frac{|x|^2}{4}} \left(u^2 - \frac{1}{4} x_{n+1}^2 u^2 + 4 |\nabla^{\Sigma} u|^2 \right),$$

the second statement follows. This completes the proof of Proposition 2.6.

Let us denote by \mathcal{H} the space of all functions f on Σ such that

$$||f||_{\mathcal{H}}^2 = \int_{\Sigma} e^{-\frac{|x|^2}{4}} f^2 < \infty$$

We define an operator \mathcal{L} on the cylinder Σ by

$$\mathcal{L}f = \Delta_{\Sigma}f - \frac{1}{2} \langle x^{\tan}, \nabla^{\Sigma}f \rangle + f.$$

This operator can be rewritten as

$$\mathcal{L}f = \frac{\partial^2}{\partial z^2}f + \frac{1}{2(n-1)}\Delta_{S^{n-1}}f - \frac{1}{2}z\frac{\partial}{\partial z}f + f.$$

Let Y_m be a basis of eigenfunctions of $\Delta_{S^{n-1}}$, and let λ_m denote the corresponding eigenvalues. We assume that the eigenfunctions Y_m are normalized so that $\int_{S^{n-1}} Y_m(\theta)^2 d\theta = \frac{1}{n} |S^{n-1}|$. Note that $\lambda_0 = 0, \lambda_1 = \ldots = \lambda_n = n-1$, and $\lambda_{n+1} = 2n$. Moreover, $Y_0(\theta) = \frac{1}{\sqrt{n}}, Y_1(\theta) = \theta_1, \ldots, Y_n(\theta) = \theta_n$ for $\theta \in S^{n-1}$, where $\theta_1, \ldots, \theta_n$ denote the Cartesian coordinates of θ .

The eigenfunctions of \mathcal{L} are of the form $H_l(\frac{z}{2}) Y_m$, where H_l denotes the Hermite polynomial of degree l. The corresponding eigenvalues are given by $1 - \frac{l}{2} - \frac{\lambda_m}{2(n-1)}$. Thus, there are n+2 eigenfunctions that correspond to positive eigenvalues of \mathcal{L} , and these are given by $1, x_{n+1}, x_1, \ldots, x_n$, up to scaling. The span of these eigenfunctions will be denoted by \mathcal{H}_+ . Moreover, there are n+1 eigenfunctions of \mathcal{L} with eigenvalue 0, and these are given by

 $x_{n+1}^2 - 2, x_1 x_{n+1}, \ldots, x_n x_{n+1}$, up to scaling. The span of these eigenfunctions will be denoted by \mathcal{H}_0 . The span of all remaining eigenfunctions will be denoted by \mathcal{H}_- . With this understood, we have

$$\langle \mathcal{L}f, f \rangle_{\mathcal{H}} \geq \frac{1}{2} \|f\|_{\mathcal{H}}^{2} \qquad \text{for } f \in \mathcal{H}_{+},$$

$$\langle \mathcal{L}f, f \rangle_{\mathcal{H}} = 0 \qquad \text{for } f \in \mathcal{H}_{0},$$

$$\langle \mathcal{L}f, f \rangle_{\mathcal{H}} \leq -\frac{1}{n-1} \|f\|_{\mathcal{H}}^{2} \qquad \text{for } f \in \mathcal{H}_{-}.$$

As in Lemma 2.4 in [2], we can show that the function $u(x, \tau)$ satisfies

$$\frac{\partial}{\partial \tau} u = \mathcal{L}u + E + \langle A(\tau)x, \nu_{\Sigma} \rangle,$$

where E is an error term satisfying $|E| \leq O(\rho(\tau)^{-1}) (|u| + |\nabla^{\Sigma} u| + |A(\tau)|)$. We next define $\hat{u}(x,\tau) = u(x,\tau) \varphi(\frac{x_{n+1}}{\rho(\tau)})$. The function $\hat{u}(x,\tau)$ satisfies

$$\frac{\partial}{\partial \tau} \hat{u} = \mathcal{L}\hat{u} + \hat{E} + \langle A(\tau)x, \nu_{\Sigma} \rangle \varphi\Big(\frac{x_{n+1}}{\rho(\tau)}\Big),$$

where \hat{E} is an error term satisfying $\|\hat{E}\|_{\mathcal{H}} \leq O(\rho(\tau)^{-1}) (\|\hat{u}\|_{\mathcal{H}} + |A(\tau)|)$ (cf. [2], Lemma 2.5). Moreover, the orthogonality relations in Proposition 2.4 imply that the function $\hat{u}(x,\tau)$ is orthogonal (with respect to the inner product on \mathcal{H}) to the function $\langle Ax, \nu_{\Sigma} \rangle$ for every τ and every matrix $A \in so(n+1)$.

Lemma 2.7. We have $|A(\tau)| \leq O(\rho(\tau)^{-1}) \|\hat{u}\|_{\mathcal{H}}$ and $\left\|\frac{\partial}{\partial \tau}\hat{u} - \mathcal{L}\hat{u}\right\|_{\mathcal{H}} \leq O(\rho(\tau)^{-1}) \|\hat{u}\|_{\mathcal{H}}.$

Proof. Analogous to [2], Lemma 2.6.

We now define

$$U_{+}(\tau) := \|P_{+}\hat{u}(\cdot,\tau)\|_{\mathcal{H}}^{2},$$

$$U_{0}(\tau) := \|P_{0}\hat{u}(\cdot,\tau)\|_{\mathcal{H}}^{2},$$

$$U_{-}(\tau) := \|P_{-}\hat{u}(\cdot,\tau)\|_{\mathcal{H}}^{2},$$

where P_+, P_0, P_- denote the orthogonal projections to $\mathcal{H}_+, \mathcal{H}_0, \mathcal{H}_-$, respectively. Using Lemma 2.7, we obtain

$$\begin{aligned} \frac{d}{d\tau} U_{+}(\tau) &\geq U_{+}(\tau) - O(\rho(\tau)^{-1}) \left(U_{+}(\tau) + U_{0}(\tau) + U_{-}(\tau) \right), \\ \left| \frac{d}{d\tau} U_{0}(\tau) \right| &\leq O(\rho(\tau)^{-1}) \left(U_{+}(\tau) + U_{0}(\tau) + U_{-}(\tau) \right), \\ \frac{d}{d\tau} U_{-}(\tau) &\leq -\frac{2}{n-1} U_{-}(\tau) + O(\rho(\tau)^{-1}) \left(U_{+}(\tau) + U_{0}(\tau) + U_{-}(\tau) \right). \end{aligned}$$

Clearly, $U_{+}(\tau) + U_{0}(\tau) + U_{-}(\tau) = \| \hat{u} \|_{\mathcal{H}}^{2} \to 0 \text{ as } \tau \to -\infty. \end{aligned}$

Lemma 2.8. We have $U_0(\tau) + U_-(\tau) \le o(1)U_+(\tau)$.

Proof. The ODE lemma of Merle and Zaag (cf. Lemma 5.4 in [1] or Lemma A.1 in [11]) implies that either $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau)$ or $U_+(\tau) + U_-(\tau) \leq o(1)U_0(\tau)$.

We now explain how to rule out the second case. If $U_{+}(\tau) + U_{-}(\tau) \leq o(1)U_{0}(\tau)$, then $\frac{\hat{u}(\cdot,\tau)}{\|\hat{u}(\cdot,\tau)\|_{\mathcal{H}}}$ converges with respect to $\|\cdot\|_{\mathcal{H}}$ to the subspace $\mathcal{H}_{0} = \operatorname{span}\{x_{n+1}^{2} - 2, x_{1}x_{n+1}, \ldots, x_{n}x_{n+1}\}$. The orthogonality relations in Proposition 2.4 imply that $\hat{u}(\cdot,\tau)$ is orthogonal to $\langle Ax, \nu_{\Sigma} \rangle$ for each $A \in so(n+1)$. In other words, $\hat{u}(\cdot,\tau)$ is orthogonal to $x_{1}x_{n+1}, \ldots, x_{n}x_{n+1}$. Therefore, $\frac{\hat{u}(\cdot,\tau)}{\|\hat{u}(\cdot,\tau)\|_{\mathcal{H}}}$ converges (with respect to $\|\cdot\|_{\mathcal{H}}$) to a non-zero multiple of $x_{n+1}^{2} - 2$.

Let $\tilde{\Omega}_{\tau}$ denote the region enclosed by \tilde{M}_{τ} , and let $\mathcal{A}(z,\tau)$ denote the area of the intersection $\tilde{\Omega}_{\tau} \cap \{x_{n+1} = z\}$. By the Brunn-Minkowski inequality, the function $z \mapsto \mathcal{A}(z,\tau)^{\frac{1}{n}}$ is concave. Since \tilde{M}_{τ} is noncompact, it follows that the function $z \mapsto \mathcal{A}(z,\tau)^{\frac{1}{n}}$ is monotone for each τ .

Note that $\mathcal{A}(z,\tau) = \frac{1}{n} \int_{S^{n-1}} (\sqrt{2(n-1)} + u(\cdot,\tau))^n$ for $|z| \leq \rho(\tau)$. Consequently, the function $z \mapsto \int_{S^{n-1}} [(\sqrt{2(n-1)} + u(\cdot,\tau))^n - \sqrt{2(n-1)}^n] d\theta$ is monotone. In particular, we either have

$$\begin{split} &\int_{-3}^{-1} \left(\int_{S^{n-1}} \left[(\sqrt{2(n-1)} + u(\cdot,\tau))^n - \sqrt{2(n-1)}^n \right] \right) dz \\ &\leq \int_{-1}^1 \left(\int_{S^{n-1}} \left[(\sqrt{2(n-1)} + u(\cdot,\tau))^n - \sqrt{2(n-1)}^n \right] \right) dz \\ &\leq \int_{1}^3 \left(\int_{S^{n-1}} \left[(\sqrt{2(n-1)} + u(\cdot,\tau))^n - \sqrt{2(n-1)}^n \right] \right) dz \end{split}$$

or

$$\begin{split} &\int_{-3}^{-1} \left(\int_{S^{n-1}} \left[(\sqrt{2(n-1)} + u(\cdot, \tau))^n - \sqrt{2(n-1)}^n \right] \right) dz \\ &\geq \int_{-1}^1 \left(\int_{S^{n-1}} \left[(\sqrt{2(n-1)} + u(\cdot, \tau))^n - \sqrt{2(n-1)}^n \right] \right) dz \\ &\geq \int_{1}^3 \left(\int_{S^{n-1}} \left[(\sqrt{2(n-1)} + u(\cdot, \tau))^n - \sqrt{2(n-1)}^n \right] \right) dz. \end{split}$$

On the other hand, we know that $\sup |u(\cdot,\tau)| \to 0$, and $\frac{\hat{u}(\cdot,\tau)}{\|\hat{u}(\cdot,\tau)\|_{\mathcal{H}}}$ converges (with respect to $\|\cdot\|_{\mathcal{H}}$) to a non-zero multiple of $x_{n+1}^2 - 2$. Consequently, the function

$$\frac{(\sqrt{2(n-1)} + \hat{u}(\cdot,\tau))^n - \sqrt{2(n-1)}^n}{\|\hat{u}(\cdot,\tau)\|_{\mathcal{H}}}$$

converges (with respect to $\|\cdot\|_{\mathcal{H}}$) to a non-zero multiple of $x_{n+1}^2 - 2$. This is a contradiction. The proof of Lemma 2.8 is now complete.

Lemma 2.9. For each $\varepsilon > 0$, we have $||u(\cdot, \tau)||_{C^4(S^{n-1} \times [-10n, 10n])} \le o(e^{\frac{(1-\varepsilon)\tau}{2}})$ and $|A(\tau)| \le o(e^{\frac{(1-\varepsilon)\tau}{2}})$.

Proof. Lemma 2.8 gives $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau)$. Substituting this back into the ODE for $U_+(\tau)$ gives

$$\frac{d}{d\tau}U_{+}(\tau) \ge U_{+}(\tau) - o(1) U_{+}(\tau).$$

Consequently, for every $\varepsilon > 0$, we have $U_{+}(\tau) \leq o(e^{(1-\varepsilon)\tau})$. Using the estimate $U_{0}(\tau) + U_{-}(\tau) \leq o(1)U_{+}(\tau)$, we obtain

$$\|\hat{u}\|_{\mathcal{H}}^2 = U_+(\tau) + U_0(\tau) + U_-(\tau) \le o(e^{(1-\varepsilon)\tau}).$$

This implies $|A(\tau)| \leq o(1) \|\hat{u}\|_{\mathcal{H}} \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$. Moreover, standard interpolation inequalities imply $\|u(\cdot,\tau)\|_{C^4(S^{n-1}\times[-10n,10n])} \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$. This completes the proof of Lemma 2.9.

It follows from the estimate $|A(\tau)| \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$ that the limit $\lim_{\tau \to -\infty} S(\tau)$ exists. Without loss of generality, we may assume that $\lim_{\tau \to -\infty} S(\tau) = id$. Then $|S(\tau) - id| \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$.

Lemma 2.10. We have

$$\sup_{\bar{M}_{\tau} \cap \{|x_{n+1}| \le e^{-\frac{\tau}{10}}\}} |x_1^2 + \ldots + x_n^2 - 2(n-1)| \le e^{\frac{\tau}{10}}$$

if $-\tau$ is sufficiently large.

Proof. Lemma 2.9 implies

$$\sup_{x \in \bar{M}_{\tau} \cap B_{5n}(0)} |x_1^2 + \ldots + x_n^2 - 2(n-1)| \le o(e^{\frac{(1-\varepsilon)\tau}{2}}).$$

In view of the convexity of \overline{M}_{τ} , it follows that

$$\sup_{\bar{M}_{\tau} \cap \{|x_{n+1}| \le e^{-\frac{\tau}{10}}\}} (x_1^2 + \ldots + x_n^2) \le 2(n-1) + e^{\frac{\tau}{10}}$$

if $-\tau$ is sufficiently large. Let

$$\Sigma_a = \{ x \in \mathbb{R}^{n+1} : x_1^2 + \ldots + x_n^2 = u_a(-x_{n+1})^2, \ -a \le x_{n+1} \le 0 \}$$

be the self-similar shrinker constructed in [1]. By Lemma 4.4 in [1], $u_a(2) \leq \sqrt{2(n-1)} - a^{-2}$. Since \bar{M}_{τ} converges to Σ in $C_{\rm loc}^{\infty}$, the surface $\bar{M}_{\tau} \cap \{x_{n+1} \leq -2\}$ encloses the surface $\Sigma_a \cap \{x_{n+1} \leq -2\}$ if $-\tau$ is sufficiently large (depending on *a*). On the other hand, the estimate $\inf_{x \in \bar{M}_{\tau} \cap B_{5n}(0)}(x_1^2 + \ldots + x_n^2) \geq 2(n-1) - o(e^{\frac{(1-\varepsilon)\tau}{2}})$ guarantees that the boundary $\bar{M}_{\tau} \cap \{x_{n+1} = -2\}$ encloses the boundary $\Sigma_a \cap \{x_{n+1} = -2\}$ provided that $-\tau$ is sufficiently large and

 $a \leq e^{-\frac{(1-\varepsilon)\tau}{4}}$. By the maximum principle, the surface $\bar{M}_{\tau} \cap \{x_{n+1} \leq -2\}$ encloses $\Sigma_a \cap \{x_{n+1} \leq -2\}$ whenever $-\tau$ is sufficiently large and $a \leq e^{-\frac{(1-\varepsilon)\tau}{4}}$. By Theorem 8.2 in [1], $u_a(y) \geq \sqrt{2(n-1)(1-a^{-2}y^2)}$. This gives

$$\inf_{\bar{M}_{\tau} \cap \{-e^{-\frac{\tau}{10}} \le x_{n+1} \le -2\}} (x_1^2 + \ldots + x_n^2) \ge 2(n-1) - e^{\frac{\tau}{10}}$$

if $-\tau$ is sufficiently large. An analogous argument gives

$$\inf_{\bar{M}_{\tau} \cap \{2 \le x_{n+1} \le e^{-\frac{\tau}{10}}\}} (x_1^2 + \ldots + x_n^2) \ge 2(n-1) - e^{\frac{\tau}{10}}$$

if $-\tau$ is sufficiently large. Putting these facts together, we conclude that

$$\inf_{\bar{M}_{\tau} \cap \{|x_{n+1}| \le e^{-\frac{\tau}{10}}\}} (x_1^2 + \ldots + x_n^2) \ge 2(n-1) - e^{\frac{\tau}{10}}$$

if $-\tau$ is sufficiently large. This completes the proof of Lemma 2.10.

Lemma 2.11. Let $\varepsilon_0 > 0$ be given. If $-\tau$ is sufficiently large (depending on ε_0), then every point in $\overline{M}_{\tau} \cap \{|x_{n+1}| \leq \frac{1}{2}e^{-\frac{\tau}{10}}\}$ lies at the center of an ε_0 -neck. Moreover, the radius of that neck is uniformly bounded from above and below.

Proof. By Lemma 2.10, the surface $\overline{M}_{\tau} \cap \{|x_{n+1}| \leq e^{-\frac{\tau}{10}}\}$ lies inside the cylinder $\{x_1^2 + \ldots + x_n^2 = 2(n-1) + e^{\frac{\tau}{10}}\}$, and outside the cylinder $\{x_1^2 + \ldots + x_n^2 = 2(n-1) - e^{\frac{\tau}{10}}\}$. In other words, the surface $M_t \cap \{|x_{n+1}| \leq (-t)^{\frac{1}{2} + \frac{1}{10}}\}$ lies inside the cylinder $\{x_1^2 + \ldots + x_n^2 = -2(n-1)t + (-t)^{1-\frac{1}{10}}\}$, and outside the cylinder $\{x_1^2 + \ldots + x_n^2 = -2(n-1)t - (-t)^{1-\frac{1}{10}}\}$. Since M_t bounds a convex domain, it follows that the surface $M_t \cap \{|x_{n+1}| \leq \frac{4}{5}(-t)^{\frac{1}{2} + \frac{1}{10}}\}$ can be written as a graph over the cylinder. Classical estimates of Ecker and Huisken [5] then imply that $H \leq C(-t)^{-\frac{1}{2}}$ on $M_t \cap \{|x_{n+1}| \leq \frac{3}{5}(-t)^{\frac{1}{2} + \frac{1}{10}}\}$. On the other hand, using the noncollapsing condition, we obtain $H \geq \frac{1}{C}(-t)^{-\frac{1}{2}}$ on $M_t \cap \{|x_{n+1}| \leq \frac{3}{5}(-t)^{\frac{1}{2} + \frac{1}{10}}\}$. Consequently, the mean curvature of the rescaled hypersurface $\overline{M}_{\tau} \cap \{|x_{n+1}| \leq \frac{3}{5}e^{-\frac{\tau}{10}}\}$ is uniformly bounded from above and below. Finally, since $\overline{M}_{\tau} \cap \{|x_{n+1}| \leq e^{-\frac{\tau}{10}}\}$ is c^0-close to a cylinder, it follows that every point on $\overline{M}_{\tau} \cap \{|x_{n+1}| \leq \frac{1}{2}e^{-\frac{\tau}{10}}\}$ lies on an ε_0 -neck if $-\tau$ is sufficiently large (depending on ε_0).

Proposition 2.12. We have

$$\sup_{x \in \bar{M}_{\tau} \cap B_{5n}(0)} |x_1^2 + \ldots + x_n^2 - 2(n-1)| \le O(e^{\frac{\tau}{2}}).$$

Proof. We repeat the argument above, this time with $\rho(\tau) = e^{-\frac{\tau}{1000}}$. It follows from Lemma 2.10 that, in the ball $B_{2e^{-\frac{\tau}{1000}}}(0)$, the surface \bar{M}_{τ} can be written as a graph over the cylinder, and the height function has

 C^4 -norm less than $e^{\frac{\tau}{100}}$. Arguing as above, we can construct a new function $S(\tau)$ taking values in SO(n+1) with the following properties:

• In the ball $B_{2e^{-\frac{\tau}{1000}}}(0)$, the rotated surface $\tilde{M}_{\tau} = S(\tau)\bar{M}_{\tau}$ can be written as a graph over the cylinder of some function $u(\cdot, \tau)$; that is,

$$\{x+u(x,\tau)\nu_{\Sigma}(x): x\in\Sigma\cap B_{2e^{-\frac{\tau}{1000}}}(0)\}\subset \tilde{M}_{\tau},$$

where $||u(\cdot, \tau)||_{C^4(\Sigma \cap B_{2e^{-\frac{\tau}{1000}}(0))}} \leq O(e^{\frac{\tau}{200}}).$ • The function $u(\cdot, \tau)$ satisfies the orthogonality relations

$$\int_{\Sigma \cap B_{e^{-\frac{\tau}{1000}}}(0)} e^{-\frac{|x|^2}{4}} \langle Ax, \nu_{\Sigma} \rangle u(x,\tau) \varphi(e^{\frac{\tau}{1000}} x_{n+1}) = 0$$

for all $A \in so(n+1)$.

Moreover, $\lim_{\tau \to -\infty} S(\tau) = id$. The function $\hat{u}(x,\tau) = u(x,\tau) \varphi(e^{\frac{\tau}{1000}} x_{n+1})$ satisfies

$$\left\|\frac{\partial}{\partial \tau}\hat{u} - \mathcal{L}\hat{u}\right\|_{\mathcal{H}} \le O(e^{\frac{\tau}{1000}}) \,\|\hat{u}\|_{\mathcal{H}}.$$

Hence, if we define

$$U_{+}(\tau) := \|P_{+}\hat{u}(\cdot,\tau)\|_{\mathcal{H}}^{2},$$

$$U_{0}(\tau) := \|P_{0}\hat{u}(\cdot,\tau)\|_{\mathcal{H}}^{2},$$

$$U_{-}(\tau) := \|P_{-}\hat{u}(\cdot,\tau)\|_{\mathcal{H}}^{2},$$

then

$$\begin{aligned} \frac{d}{d\tau} U_{+}(\tau) &\geq U_{+}(\tau) - O(e^{\frac{\tau}{1000}}) \left(U_{+}(\tau) + U_{0}(\tau) + U_{-}(\tau) \right), \\ \left| \frac{d}{d\tau} U_{0}(\tau) \right| &\leq O(e^{\frac{\tau}{1000}}) \left(U_{+}(\tau) + U_{0}(\tau) + U_{-}(\tau) \right), \\ \frac{d}{d\tau} U_{-}(\tau) &\leq -U_{-}(\tau) + O(e^{\frac{\tau}{1000}}) \left(U_{+}(\tau) + U_{0}(\tau) + U_{-}(\tau) \right). \end{aligned}$$

The ODE lemma of Merle and Zaag (cf. Lemma 5.4 in [1]) implies that either $U_0(\tau) + U_-(\tau) \le o(1)U_+(\tau)$ or $U_+(\tau) + U_-(\tau) \le o(1)U_0(\tau)$. As above, the latter case can be ruled out. Therefore, $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau)$. This gives

$$\frac{d}{d\tau}U_{+}(\tau) \ge U_{+}(\tau) - O(e^{\frac{\tau}{1000}}) U_{+}(\tau),$$

hence $U_{+}(\tau) \leq O(e^{\tau})$. Thus, $U_{0}(\tau) + U_{-}(\tau) \leq o(1) U_{+}(\tau) \leq O(e^{\tau})$. Consequently, $\|\hat{u}\|_{\mathcal{H}} \leq O(e^{\frac{\tau}{2}})$. Arguing as in Lemma 2.7, we obtain $|A(\tau)| \leq$ $O(e^{\frac{\tau}{2}})$. Since $\lim_{\tau \to -\infty} S(\tau) = \mathrm{id}$, we conclude that $|S(\tau) - \mathrm{id}| \leq O(e^{\frac{\tau}{2}})$. Finally, u satisfies an equation of the form $\frac{\partial}{\partial \tau} u = \tilde{\mathcal{L}} u + \langle A(\tau) x, \nu_{\Sigma} \rangle$, where $\tilde{\mathcal{L}}$ is an elliptic operator of second order whose coefficients depend on $u, \nabla u$, $\nabla^2 u$, and $A(\tau)$. As $\tau \to -\infty$, the coefficients of $\tilde{\mathcal{L}}$ converge smoothly to the corresponding coefficients of \mathcal{L} . Hence, standard interior estimates for parabolic equations imply that $||u(\cdot, \tau)||_{C^4(S^{n-1}\times[-10n,10n])} \leq O(e^{\frac{\tau}{2}})$. Combining this estimate with the estimate $|S(\tau) - \mathrm{id}| \leq O(e^{\frac{\tau}{2}})$, we conclude that

$$\sup_{x \in \bar{M}_{\tau} \cap B_{5n}(0)} |x_1^2 + \ldots + x_n^2 - 2(n-1)| \le O(e^{\frac{\tau}{2}}).$$

3. Lower bound for $H_{\max}(t)$ as $t \to -\infty$

Let $M_t, t \in (-\infty, 0]$, be a noncompact ancient solution of mean curvature flow in \mathbb{R}^{n+1} which is strictly convex, uniformly two-convex, and noncollapsed. For each t, we define $H_{\max}(t)$ to be the supremum of the mean curvature of M_t .

Proposition 3.1. For each t, $H_{\max}(t) < \infty$.

Proof. Let us fix a time t and a small number $\varepsilon > 0$. It follows from Proposition 3.1 in [8] that every point in M_t which lies outside some large compact set must lie at the center of an ε -neck. Hence, if $H_{\max}(t) = \infty$, then the surface M_t contains a sequence of ε -necks with radii converging to 0, but this cannot happen in a convex hypersurface.

Proposition 3.2. The function $H_{\max}(t)$ is continuous and monotone increasing in t.

Proof. We first show that $H_{\max}(t)$ is continuous in t. It follows from work of Haslhofer and Kleiner [7],[8] that $|\frac{\partial}{\partial t}H| \leq CH^3$ for some uniform constant C. Let us fix a time \bar{t} and a positive number $\varepsilon > 0$. By definition of $H_{\max}(\bar{t})$, there exists a point on $M_{\bar{t}}$ where the mean curvature lies in the interval $\left[\left(1-\frac{\varepsilon}{2}\right)H_{\max}(\bar{t}),H_{\max}(\bar{t})\right]$. If t is sufficiently close to \bar{t} , then the estimate $\left|\frac{\partial}{\partial t}H\right| \leq CH^3$ implies that there exists a point on M_t where the mean curvature lies in the interval $[(1-\varepsilon) H_{\max}(\bar{t}), (1+\varepsilon) H_{\max}(\bar{t})]$. In particular, $H_{\max}(t) \ge (1-\varepsilon) H_{\max}(\bar{t})$ if t is sufficiently close to \bar{t} . Suppose next that there is a point on M_t where the mean curvature is equal to $(1+\varepsilon) H_{\max}(t)$. If t is sufficiently close to \bar{t} , then the estimate $|\frac{\partial}{\partial t}H| \leq CH^3$ implies that there exists a point on $M_{\bar{t}}$ where the mean curvature lies in the interval $\left[\left(1+\frac{\varepsilon}{2}\right)H_{\max}(t),\left(1+2\varepsilon\right)H_{\max}(t)\right]$, which contradicts the definition of $H_{\max}(\bar{t})$. Hence, if t is sufficiently close to \bar{t} , there is no point on M_t where the mean curvature is equal to $(1 + \varepsilon) H_{\max}(\bar{t})$. On the other hand, since M_t bounds a convex domain which is not a slab, we know that M_t is connected (cf. [13], Theorem V). Consequently, $H_{\max}(t) \leq (1 + \varepsilon) H_{\max}(\bar{t})$ if t is sufficiently close to \bar{t} .

Thus, $H_{\max}(t)$ is a continuous function of t. In particular, $H_{\max}(t)$ is uniformly bounded from above on every compact time interval. Consequently, $H_{\max}(t)$ is monotone increasing in t by Hamilton's Harnack inequality [6]. This completes the proof of Proposition 3.2.

Proposition 3.3. We have $\liminf_{t\to -\infty} H_{\max}(t) > 0$.

Proof. Proposition 2.12 implies that

$$\sup_{\substack{x \in (-t)^{-\frac{1}{2}} (M_t \cap B_{5n(-t)^{\frac{1}{2}}}(0))}} |x_1^2 + \ldots + x_n^2 - 2(n-1)| \le O((-t)^{-\frac{1}{2}}).$$

Since M_t has exactly one end, we can assume without loss of generality that $M_t \cap \{x_{n+1} \ge 0\}$ is noncompact and $M_t \cap \{x_{n+1} \le 0\}$ is compact. There exists a large constant K with the following property: if -t is sufficiently large, then the cross-section

$$(-t)^{-\frac{1}{2}} (M_t \cap \{x_{n+1} = -2(-t)^{\frac{1}{2}}\})$$

lies outside the sphere

$$\{x_1^2 + \ldots + x_n^2 = (\sqrt{2(n-1)} - K(-t)^{-\frac{1}{2}})^2, x_{n+1} = -2\}.$$

We now recall the self-similar shrinkers constructed in [1]. For a > 0 large, there exists a self-similar shrinker

$$\Sigma_a = \{ x \in \mathbb{R}^{n+1} : x_1^2 + \ldots + x_n^2 = u_a(-x_{n+1})^2, \ -a \le x_{n+1} \le 0 \}$$

satisfying $H = \frac{1}{2} \langle x, \nu \rangle$. Consequently, the hypersurfaces

$$\Sigma_{a,t} := (-t)^{\frac{1}{2}} \Sigma_a + (0, \dots, 0, Ka^2)$$

= $\{x \in \mathbb{R}^{n+1} : x_1^2 + \dots + x_n^2 = (-t) u_a ((-x_{n+1} + Ka^2)(-t)^{-\frac{1}{2}})^2, Ka^2 - a(-t)^{\frac{1}{2}} \le x_{n+1} \le Ka^2\}$

evolve by mean curvature flow.

As in [2], we can use the hypersurfaces $\Sigma_{a,t} \cap \{x_{n+1} \leq -2(-t)^{\frac{1}{2}}\}$ as barriers. As $t \to -\infty$, the rescaled surfaces $(-t)^{-\frac{1}{2}} M_t$ converge in C_{loc}^{∞} to the cylinder $\{x_1^2 + \ldots + x_n^2 = 2(n-1)\}$. Furthermore, the rescaled surfaces $(-t)^{-\frac{1}{2}} (\Sigma_{a,t} \cap \{x_{n+1} \leq -2(-t)^{\frac{1}{2}}\})$ converge to $\Sigma_a \cap \{x_{n+1} \leq -2\}$ as $t \to -\infty$. It follows from Lemma 4.4 in [1] that the set $\Sigma_a \cap \{x_{n+1} \leq -2\}$ is a compact subset of $\{x_1^2 + \ldots + x_n^2 < 2(n-1)\}$. Consequently, $\Sigma_{a,t} \cap \{x_{n+1} \leq -2(-t)^{\frac{1}{2}}\}$ lies inside $M_t \cap \{x_{n+1} \leq -2(-t)^{\frac{1}{2}}\}$ if -t is sufficiently large (depending on a).

By our choice of K, the cross-section

$$(-t)^{-\frac{1}{2}} (M_t \cap \{x_{n+1} = -2(-t)^{\frac{1}{2}}\})$$

lies outside the sphere

$$\{x_1^2 + \ldots + x_n^2 = (\sqrt{2(n-1)} - K(-t)^{-\frac{1}{2}})^2, x_{n+1} = -2\}.$$

Moreover, the cross-section

$$(-t)^{-\frac{1}{2}} (\Sigma_{a,t} \cap \{x_{n+1} = -2(-t)^{\frac{1}{2}}\})$$

is a sphere

$$\{x_1^2 + \ldots + x_n^2 = u_a(2 + Ka^2(-t)^{-\frac{1}{2}})^2, x_{n+1} = -2\}.$$

Using Lemma 4.4 in [1], we obtain $u_a(2) \leq \sqrt{2(n-1)}$ and $u_a(2) - u_a(1) \leq -a^{-2}$ if a is sufficiently large. Since the function u_a is concave, we obtain

$$u_a(2 + Ka^2(-t)^{-\frac{1}{2}}) \le u_a(2) + Ka^2(-t)^{-\frac{1}{2}}(u_a(2) - u_a(1))$$
$$\le \sqrt{2(n-1)} - K(-t)^{-\frac{1}{2}}$$

for $-t \ge 4K^2a^2$. Consequently, the cross-section $\Sigma_{a,t} \cap \{x_{n+1} = -2(-t)^{\frac{1}{2}}\}$ lies inside the cross-section $M_t \cap \{x_{n+1} = -2(-t)^{\frac{1}{2}}\}$ whenever $-t \ge 4K^2a^2$ and a is sufficiently large. By the maximum principle, the hypersurface $\Sigma_{a,t} \cap \{x_{n+1} \le -2(-t)^{\frac{1}{2}}\}$ lies inside the hypersurface $M_t \cap \{x_{n+1} \le -2(-t)^{\frac{1}{2}}\}$ whenever $-t \ge 4K^2a^2$ and a is sufficiently large. For $-t = 4K^2a^2$, the tip of $\Sigma_{a,t}$ has distance $a(-t)^{\frac{1}{2}} - Ka^2 = Ka^2 = -\frac{t}{4K}$ from the origin. Consequently, the intersection $M_t \cap \{x_1 = \ldots = x_n = 0, x_{n+1} \le \frac{t}{4K}\}$ is non-empty if -t is sufficiently large. In particular, $\limsup_{t\to -\infty} H_{\max}(t) > 0$. Since $H_{\max}(t)$ is monotone increasing in t, it follows that $\liminf_{t\to -\infty} H_{\max}(t) > 0$.

4. The NECK IMPROVEMENT THEOREM

In this section, we show that a neck becomes more symmetric under the evolution.

Definition 4.1. Let $\mathcal{K} = \{K_{\alpha} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ be a collection of vector fields in \mathbb{R}^{n+1} . We say that \mathcal{K} is a normalized set of rotation vector fields if there exists an orthonormal basis $\{J_{\alpha} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ of $so(n) \subset so(n+1)$, a matrix $S \in O(n+1)$ and a point $q \in \mathbb{R}^{n+1}$ such that

$$K_{\alpha}(x) = SJ_{\alpha}S^{-1}(x-q).$$

Note that we require that the vector fields K_{α} all have a common axis of rotation, but we do not require that the axis of rotation passes through the origin.

Lemma 4.2. For each n, we can find a large constant C and a small constant $\varepsilon_0 > 0$ with the following property. Let M be a hypersurface in \mathbb{R}^{n+1} with boundary. Assume that, after suitable rescaling, M is ε_0 -close (in the C⁴-norm) to a cylinder $S^{n-1} \times [-5,5]$ of radius 1. Suppose that $\mathcal{K}^{(1)} = \{K_{\alpha}^{(1)} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ and $\mathcal{K}^{(2)} = \{K_{\alpha}^{(2)} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ are two normalized sets of rotation vector fields with the following properties:

• $\max_{\alpha} |K_{\alpha}^{(1)}| H \leq 10n \text{ and } \max_{\alpha} |K_{\alpha}^{(2)}| H \leq 10n \text{ at the point } \bar{x}.$

•
$$\max_{\alpha} |\langle K_{\alpha}^{(1)}, \nu \rangle| H \leq \varepsilon \text{ and } \max_{\alpha} |\langle K_{\alpha}^{(2)}, \nu \rangle| H \leq \varepsilon \text{ on } M.$$

Then

$$\inf_{\omega \in O(\frac{n(n-1)}{2})} \sup_{B_{100nH(\bar{x})^{-1}}(\bar{x})} \max_{\alpha} \left| K_{\alpha}^{(1)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta} K_{\beta}^{(2)} \right| H(\bar{x}) \le C\varepsilon.$$

Proof. The proof is by contradiction. If the assertion is false, then we can find a sequence of hypersurfaces $M^{(l)}$, a sequence of points $\bar{x}_l \in M^{(l)}$, a sequence $\mathcal{K}^{(1,l)} = \{K^{(1,l)}_{\alpha} : 1 \le \alpha \le \frac{n(n-1)}{2}\}$, a sequence $\mathcal{K}^{(2,l)} = \{K^{(2,l)}_{\alpha} : 1 \le \alpha \le \frac{n(n-1)}{2}\}$ $1 \le \alpha \le \frac{n(n-1)}{2}$, and a sequence of positive numbers $\varepsilon_l \to 0$ such that the following statements hold:

- (i) The hypersurfaces M_l converge in C^4 to the standard cylinder $S^{n-1} \times$ [-5, 5] of radius 1 around the x_{n+1} -axis.
- (ii) $\max_{\alpha} |K_{\alpha}^{(1,l)}| \leq 100$ and $\max_{\alpha} |K_{\alpha}^{(2,l)}| \leq 100$ at the point \bar{x}_l . (iii) $\max_{\alpha} |\langle K_{\alpha}^{(1,l)}, \nu \rangle| \leq \varepsilon_l$ and $\max_{\alpha} |\langle K_{\alpha}^{(2,l)}, \nu \rangle| \leq \varepsilon_l$ on $M^{(l)}$.
- (iv) We have

$$\varepsilon_l^{-1} \inf_{\omega \in O(\frac{n(n-1)}{2})} \sup_{B_{1000}(\bar{x}_l)} \max_{\alpha} \left| K_{\alpha}^{(1,l)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta} K_{\beta}^{(2,l)} \right| \to \infty$$

as $l \to \infty$.

Let us write

$$K_{\alpha}^{(1,l)}(x) = S^{(1,l)} J_{\alpha}^{(1,l)} (S^{(1,l)})^{-1} (x - q^{(1,l)})$$

and

$$K_{\alpha}^{(2,l)}(x) = S^{(2,l)} J_{\alpha}^{(2,l)} (S^{(2,l)})^{-1} (x - q^{(2,l)}).$$

Here, $\{J_{\alpha}^{(1,l)}: 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ and $\{J_{\alpha}^{(2,l)}: 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ are orthonormal bases of $so(n) \subset so(n+1), S^{(1,l)}, S^{(2,l)} \in O(n+1), \text{ and } q^{(1,l)}, q^{(2,l)} \in \mathbb{R}^{n+1}.$

The condition (ii) implies that the axis of rotation of $\mathcal{K}^{(1,l)}$ has bounded distance from the origin. It is easy to see that the axis of rotation of $\mathcal{K}^{(1,l)}$ converges to the x_{n+1} -axis as $l \to \infty$. Thus, $S^{(1,l)}(0,\ldots,0,1) \to$ $\pm(0,\ldots,0,1)$ as $l \to \infty$. Without loss of generality, we may assume that $S^{(1,l)} \to \text{id as } l \to \infty$. Similarly, we may assume that $S^{(2,l)} \to \text{id as } l \to \infty$. Moreover, we can arrange that $q^{(1,l)}$ and $q^{(2,l)}$ are orthogonal to the x_{n+1} axis. (To achieve this, we add a multiple of $S^{(1,l)}(0,\ldots,0,1)$ to $q^{(1,l)}$, and a multiple of $S^{(2,l)}(0,\ldots,0,1)$ to $q^{(2,l)}$. This does not change the vector fields $K_{\alpha}^{(1,l)}$ and $K_{\alpha}^{(2,l)}$.) This gives $q^{(1,l)} \to 0$ and $q^{(2,l)} \to 0$ as $l \to \infty$.

Let us write

$$(S^{(1,l)})^{-1}S^{(2,l)} = \exp(\sigma^{(l)}) U^{(l)},$$

where $U^{(l)} \in O(n) \subset O(n+1)$ (in other words, $U^{(l)}$ is an isometry of \mathbb{R}^{n+1} which fixes the x_{n+1} -axis) and $\sigma^{(l)} \in so(n+1)$ is an anti-symmetric matrix satisfying $\sigma_{ij}^{(l)} = 0$ for $i, j \in \{1, \ldots, n\}$. Finally, we can find a matrix $\omega^{(l)} \in O(\frac{n(n-1)}{2})$ such that

$$\sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta}^{(l)} J_{\beta}^{(2,l)} = (U^{(l)})^{-1} J_{\alpha}^{(1,l)} U^{(l)}.$$

This gives

$$\sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta}^{(l)} K_{\beta}^{(2,l)}(x) = S^{(2,l)} (U^{(l)})^{-1} J_{\alpha}^{(1,l)} U^{(l)} (S^{(2,l)})^{-1} (x - q^{(2,l)})$$
$$= S^{(1,l)} \exp(\sigma^{(l)}) J_{\alpha}^{(1,l)} \exp(-\sigma^{(l)}) (S^{(1,l)})^{-1} (x - q^{(2,l)})$$

Let $\delta_l := |q^{(1,l)} - q^{(2,l)}| + |\sigma^{(l)}|$. Clearly,

$$\sup_{B_{1000}(\bar{x}_l)} \max_{\alpha} \left| K_{\alpha}^{(1,l)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta}^{(l)} K_{\beta}^{(2,l)} \right| \le C(n) \,\delta_l.$$

Hence, property (iv) implies $\varepsilon_l^{-1} \delta_l \to \infty$ as $l \to \infty$. Let

$$V_{\alpha}^{(l)} := \delta_l^{-1} \left(K_{\alpha}^{(1,l)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta}^{(l)} K_{\beta}^{(2,l)} \right).$$

As $l \to \infty$, the vector fields $V_{\alpha}^{(l)}$ converge to a limiting vector field V_{α} of the form

$$V_{\alpha}(x) = -[\sigma, J_{\alpha}]x - J_{\alpha}\zeta,$$

where $J_{\alpha} := \lim_{l \to \infty} J_{\alpha}^{(1,l)}$, $\zeta := \lim_{l \to \infty} \delta_l^{-1} (q^{(1,l)} - q^{(2,l)})$, and $\sigma := \lim_{l \to \infty} \delta_l^{-1} \sigma^{(l)}$. Note that $\{J_{\alpha} : 1 \le \alpha \le \frac{n(n-1)}{2}\}$ is an orthonormal basis of $so(n) \subset so(n+1)$, $\zeta \in \mathbb{R}^{n+1}$ is orthogonal to the x_{n+1} -axis, and $\sigma \in so(n+1)$ is an antisymmetric matrix satisfying $\sigma_{ij} = 0$ for $i, j \in \{1, \ldots, n\}$. Moreover, by definition of δ_l , we have $|\zeta| + |\sigma| = \lim_{l \to \infty} \delta_l^{-1} (|q^{(1,l)} - q^{(2,l)}| + |\sigma^{(l)}|) = 1$.

Using property (iii), we obtain

$$\sup_{M^{(l)}} \max_{\alpha} |\langle V_{\alpha}^{(l)}, \nu \rangle| \leq C(n) \, \delta_l^{-1} \sup_{M^{(l)}} \max_{\alpha} (|\langle K_{\alpha}^{(1,l)}, \nu \rangle| + |\langle K_{\alpha}^{(2,l)}, \nu \rangle|) \leq C(n) \, \delta_l^{-1} \, \varepsilon_l$$

$$\to 0$$

Consequently, the limiting vector fields V_{α} are tangential to the cylinder $S^{n-1} \times [-5,5]$. From this, we deduce that $\zeta = 0$ and $\sigma = 0$. This is a contradiction. This completes the proof of Lemma 4.2.

Following [10], pp. 189–190, we denote by $\mathcal{P}(\bar{x}, \bar{t}, r, \tau)$ the set of all points (x, t) in space-time such that $x \in B_{g(\bar{t})}(\bar{x}, r)$ and $t \in [\bar{t} - \tau, \bar{t}]$. With this understood, we define $\hat{\mathcal{P}}(\bar{x}, \bar{t}, L, \theta) = \mathcal{P}(\bar{x}, \bar{t}, (n-1)L H(\bar{x}, \bar{t})^{-1}, (n-1)^2 \theta H(\bar{x}, \bar{t})^{-2})$. We say that (\bar{x}, \bar{t}) lies on an ε -neck if the parabolic neighborhood $\hat{\mathcal{P}}(\bar{x}, \bar{t}, 100, 100)$ is, after rescaling, ε -close (in the C^{10} -norm), to a family of shrinking cylinders.

Definition 4.3. Let M_t be a solution of mean curvature flow. We say that a point (\bar{x}, \bar{t}) is ε -symmetric if there exists a normalized set of rotation

18

vector fields $\mathcal{K} = \{K_{\alpha} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ such that $\max_{\alpha} |K_{\alpha}| H \leq 10n$ at the point (\bar{x}, \bar{t}) and $\max_{\alpha} |\langle K_{\alpha}, \nu \rangle| H \leq \varepsilon$ in the parabolic neighborhood $\hat{\mathcal{P}}(\bar{x}, \bar{t}, 10, 100)$.

Note that the condition that $\max_{\alpha} |K_{\alpha}| H \leq 10n$ at the point (\bar{x}, \bar{t}) ensures that the distance of the point \bar{x} from the axis of rotation of $\mathcal{K} = \{K_{\alpha} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ is at most $C(n) H(\bar{x}, \bar{t})^{-1}$.

Theorem 4.4 (Neck Improvement Theorem). Given n, we can find a large constant L and a small constant ε_1 with the following property. Suppose that M_t is a solution of mean curvature flow. Moreover, suppose that (\bar{x}, \bar{t}) is a point in space-time with the property that every point in $\hat{\mathcal{P}}(\bar{x}, \bar{t}, L, L^2)$ lies at the center of an ε_1 -neck and is ε -symmetric, where $\varepsilon \leq \varepsilon_1$. Then (\bar{x}, \bar{t}) is ε_2^2 -symmetric.

Proof. Without loss of generality, we assume $\bar{t} = -1$ and $H(\bar{x}, -1) = \sqrt{\frac{n-1}{2}}$. We will assume throughout that L is sufficiently large depending on n, and ε_1 is sufficiently small depending on L. Note that, in the parabolic neighborhood $\hat{\mathcal{P}}(\bar{x}, \bar{t}, L, L^2)$, the solution can be approximated by a family of shrinking cylinders $S^{n-1}(\sqrt{-2(n-1)t}) \times \mathbb{R}$, up to errors which are bounded by $C(L)\varepsilon_1$ in the C^{10} -norm.

Step 1: Given any point $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$, we can find a normalized set of rotation vector fields $\mathcal{K}^{(x_0,t_0)} = \{K^{(x_0,t_0)}_{\alpha} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ such that $\max_{\alpha} |\langle K^{(x_0,t_0)}_{\alpha}, \nu \rangle| H \leq \varepsilon$ on the parabolic neighborhood $\hat{\mathcal{P}}(x_0, t_0, 10, 100)$. Note that the axis of rotation depends on (x_0, t_0) . By a repeated application of Lemma 4.2, we obtain

$$\inf_{\omega \in O(\frac{n(n-1)}{2})} \sup_{B_{10nL}(0)} \max_{\alpha} \left| K_{\alpha}^{(\bar{x},-1)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta} K_{\beta}^{(x_0,t_0)} \right| \le C(L)\varepsilon$$

for each point $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$. Without loss of generality, we may assume

$$\sup_{B_{10nL}(0)} \max_{\alpha} |K_{\alpha}^{(\bar{x},-1)} - K_{\alpha}^{(x_0,t_0)}| \le C(L)\varepsilon$$

for each point $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$. This implies

$$\max_{\alpha} |\langle K_{\alpha}^{(\bar{x},-1)},\nu\rangle| \le C(L)\varepsilon$$

at each point $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$.

For abbreviation, we put $\overline{\mathcal{K}} := \mathcal{K}^{(\overline{x},-1)}$. Without loss of generality, we may assume that the axis of rotation of $\overline{\mathcal{K}} = \{\overline{K}_{\alpha} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ is the x_{n+1} axis; that is, $\overline{K}_{\alpha}(x) = J_{\alpha}x$ for some orthonormal basis $\{J_{\alpha} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ of $so(n) \subset so(n+1)$. Finally, we may assume that the point \overline{x} lies in the hyperplane $\{x_{n+1} = 0\}$. Step 2: It will be convenient to write M_t as a graph over the x_{n+1} -axis, so that

$$\left\{ (r(\theta, z, t) \, \theta, z) : \theta \in S^{n-1}, \, z \in \left[-\frac{3L}{4}, \frac{3L}{4} \right] \right\} \subset M_t.$$

Note that the function $r(\theta, z, t) - (-2(n-1)t)^{\frac{1}{2}}$ is bounded by $C(L)\varepsilon_1$ in the C^{10} -norm. A straightforward computation gives

$$\nu = \frac{1}{\sqrt{1 + r^{-2} |\nabla^{S^{n-1}} r|^2 + (\frac{\partial r}{\partial z})^2}} \left(\theta - r^{-1} \nabla^{S^{n-1}} r, -\frac{\partial r}{\partial z}\right),$$

hence

$$\langle \bar{K}_{\alpha}, \nu \rangle = -\frac{\langle J_{\alpha}\theta, \nabla^{S^{n-1}}r \rangle}{\sqrt{1 + r^{-2} |\nabla^{S^{n-1}}r|^2 + (\frac{\partial r}{\partial z})^2}}$$

where $\nabla^{S^{n-1}}r$ represents the gradient of the function r with respect to the angular variables.

We have shown in Step 1 that $\max_{\alpha} |\langle \bar{K}_{\alpha}, \nu \rangle| \leq C(L)\varepsilon$ on the parabolic neighborhood $\hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$. Moreover, our assumptions imply that $|\nabla^{S^{n-1}}r| + |\frac{\partial r}{\partial z}| \leq C(L)\varepsilon_1$. This gives $|\nabla^{S^{n-1}}r| \leq C(L)\varepsilon$ and

$$|\langle \bar{K}_{\alpha}, \nu \rangle + \langle J_{\alpha}\theta, \nabla^{S^{n-1}}r \rangle| \leq C(L)\varepsilon_{1}\varepsilon.$$

Moreover, the identity $\operatorname{div}_{S^{n-1}}(J_{\alpha}\theta) = 0$ gives $\operatorname{div}_{S^{n-1}}(r(\theta, z, t) J_{\alpha}\theta) = \langle J_{\alpha}\theta, \nabla^{S^{n-1}}r \rangle$, hence

$$|\langle \bar{K}_{\alpha}, \nu \rangle + \operatorname{div}_{S^{n-1}}(r(\theta, z, t) J_{\alpha}\theta)| \le C(L)\varepsilon_{1}\varepsilon.$$

Step 3: Let us fix an index $\alpha \in \{1, \ldots, \frac{n(n-1)}{2}\}$. For each point $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$, the vector field $K_{\alpha}^{(x_0, t_0)}$ satisfies

$$|\langle K_{\alpha}^{(x_0,t_0)},\nu\rangle| \le C\varepsilon \,(-t_0)^{\frac{1}{2}}$$

on the parabolic neighborhood $\hat{\mathcal{P}}(x_0, t_0, 10, 100)$. On the other hand, we have shown in Step 1 that $\sup_{B_{10nL}(0)} \max_{\alpha} |\bar{K}_{\alpha} - K_{\alpha}^{(x_0, t_0)}| \leq C(L)\varepsilon$. Hence, there exist real numbers $a_1, \ldots, a_n, b_1, \ldots, b_n$ (depending on (x_0, t_0)) such that

$$|a_1| + \ldots + |a_n| \le C(L)\varepsilon,$$

$$|b_1| + \ldots + |b_n| \le C(L)\varepsilon,$$

and

$$|\langle \bar{K}_{\alpha} - K_{\alpha}^{(x_0, t_0)}, \nu \rangle - (a_1\theta_1 + \ldots + a_n\theta_n) - (b_1\theta_1 + \ldots + b_n\theta_n)z| \le C(L)\varepsilon_1\varepsilon_2$$

on the parabolic neighborhood $\mathcal{P}(x_0, t_0, 10, 100)$. Consequently, the function $u = \langle \bar{K}_{\alpha}, \nu \rangle$ satisfies

$$|u(\theta, z, t) - (a_1\theta_1 + \ldots + a_n\theta_n) - (b_1\theta_1 + \ldots + b_n\theta_n)z| \le C\varepsilon (-t_0)^{\frac{1}{2}} + C(L)\varepsilon_1\varepsilon_0$$
on the parabolic neighborhood $\hat{\mathcal{P}}(x_0, t_0, 10, 100).$

20

Step 4: We again fix an index $\alpha \in \{1, \ldots, \frac{n(n-1)}{2}\}$. The function u = $\langle \bar{K}_{\alpha}, \nu \rangle$ satisfies the evolution equation

$$\frac{\partial}{\partial t}u = \Delta_{M_t}u + |A|^2u.$$

We have shown in Step 1 that $|u| \leq C(L)\varepsilon$ on the parabolic neighborhood $\hat{P}(\bar{x}, -1, L, L^2)$. Using standard interior estimates for parabolic equations, we obtain $|\nabla u| + |\nabla^2 u| \leq C(L)\varepsilon$ on the parabolic neighborhood $\hat{\mathcal{P}}(\bar{x}, -1, \frac{L}{2}, \frac{L^2}{4})$. This implies

$$\left|\frac{\partial}{\partial t}u - \frac{\partial^2}{\partial z^2}u - \frac{1}{(-2(n-1)t)}\Delta_{S^{n-1}}u - \frac{1}{(-2t)}u\right| \le C(L)\varepsilon_1\varepsilon$$

for $z \in \left[-\frac{L}{4}, \frac{L}{4}\right]$ and $t \in \left[-\frac{L^2}{16}, -1\right]$. We denote by \tilde{u} the solution of the linear equation

$$\frac{\partial}{\partial t}\tilde{u} = \frac{\partial^2}{\partial z^2}\tilde{u} + \frac{1}{(-2(n-1)t)}\Delta_{S^{n-1}}\tilde{u} + \frac{1}{(-2t)}\tilde{u}$$

in the parabolic cylinder $\{z \in [-\frac{L}{4}, \frac{L}{4}], t \in [-\frac{L^2}{16}, -1]\}$ such that $\tilde{u} = u$ on the parabolic boundary $\{|z| = \frac{L}{4}\} \cup \{t = -\frac{L^2}{16}\}$. The maximum principle gives

 $|u - \tilde{u}| < C(L)\varepsilon_1\varepsilon$

in the parabolic cylinder $\{z \in [-\frac{L}{4}, \frac{L}{4}], t \in [-\frac{L^2}{16}, -1]\}$. Step 5: In order to analyze the PDE for \tilde{u} , we perform separation of variables. As above, let Y_m be a basis of eigenfunctions of $\Delta_{S^{n-1}}$, and let λ_m denote the corresponding eigenvalues. We assume that the eigenfunctions Y_m are normalized so that $\frac{n}{|S^{n-1}|} \int_{S^{n-1}} Y_m(\theta)^2 d\theta = 1$. Clearly, $\sup_{\theta \in S^{n-1}} |Y_m(\theta)| \leq C \|Y_m\|_{H^{n-1}(S^{n-1})} \leq C\lambda_m^{n-1} \text{ for } m \geq 1. \text{ Moreover,}$ $\lambda_m \sim m^{\frac{2}{n-1}}$ as $m \to \infty$. We recall that $\lambda_0 = 0, \ \lambda_1 = \ldots = \lambda_n = n-1$, and $\lambda_{n+1} = 2n$. Moreover, $Y_0(\theta) = \frac{1}{\sqrt{n}}, \ Y_1(\theta) = \theta_1, \ldots, Y_n(\theta) = \theta_n$, where $\theta_1, \ldots, \theta_n$ denote the Cartesian coordinates of $\theta \in S^{n-1}$. Let us write

$$\tilde{u}(\theta, z, t) = \sum_{m=0}^{\infty} v_m(z, t) Y_m(\theta),$$

where

$$v_m(z,t) = \frac{n}{|S^{n-1}|} \int_{S^{n-1}} \tilde{u}(\theta, z, t) Y_m(\theta) d\theta.$$

Then

$$\frac{\partial}{\partial t}v_m(z,t) = \frac{\partial^2}{\partial z^2}v_m(z,t) + \frac{n-1-\lambda_m}{(-2(n-1)t)}v_m(z,t).$$

Hence, the rescaled function $\hat{v}_m(z,t) = (-t)^{\frac{1}{2} - \frac{\lambda_m}{2(n-1)}} v_m(z,t)$ satisfies

$$\frac{\partial}{\partial t}\hat{v}_m(z,t) = \frac{\partial^2}{\partial z^2}\hat{v}_m(z,t).$$

We first consider the case when $m \ge n+1$, so that $\lambda_m \ge 2n$. Using the results in Step 3 and Step 4, we obtain

$$|v_m(z,t)| \le (C\varepsilon + C(L)\varepsilon_1\varepsilon)(-t)^{\frac{1}{2}},$$

and consequently

$$|\hat{v}_m(z,t)| \le (C\varepsilon + C(L)\varepsilon_1\varepsilon) (-t)^{1-\frac{\lambda_m}{2(n-1)}}$$

in the parabolic cylinder $\{z \in [-\frac{L}{4}, \frac{L}{4}], t \in [-\frac{L^2}{16}, -1]\}$. Using the solution formula for the one-dimensional heat equation with Dirichlet boundary condition on the rectangle $[-\frac{L}{4}, \frac{L}{4}] \times [-\frac{L^2}{16}, -1]$, we obtain

$$\begin{aligned} |\hat{v}_m(z,t)| &\leq C \, L^{-1} \int_{-\frac{L}{4}}^{\frac{L}{4}} |\hat{v}_m(z, -\frac{L^2}{16})| \, dz \\ &+ C \, L \int_{-\frac{L^2}{16}}^{t} e^{-\frac{L^2}{100(t-s)}} \, (t-s)^{-\frac{3}{2}} \left(|\hat{v}_m(\frac{L}{4}, s)| + |\hat{v}_m(-\frac{L}{4}, s)| \right) \, ds, \end{aligned}$$

hence

$$\begin{split} |\hat{v}_m(z,t)| &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L}{4}\right)^{2-\frac{\lambda_m}{n-1}} \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L \int_{-\frac{L^2}{16}}^{t} e^{-\frac{L^2}{100(t-s)}} (t-s)^{-\frac{3}{2}} (-s)^{1-\frac{\lambda_m}{2(n-1)}} ds \\ &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L}{4}\right)^{2-\frac{\lambda_m}{n-1}} \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-\frac{1}{n-1}} \int_{-\frac{L^2}{16}}^{t} e^{-\frac{L^2}{200(t-s)}} (-s)^{\frac{1-\lambda_m}{2(n-1)}} ds \\ &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L}{4}\right)^{2-\frac{\lambda_m}{n-1}} \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-\frac{1}{n-1}} \int_{-\frac{L^2}{16}}^{(1+\frac{1}{\sqrt{\lambda_m}})t} e^{-\frac{L^2}{200(t-s)}} (-s)^{\frac{1-\lambda_m}{2(n-1)}} ds \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-\frac{1}{n-1}} \int_{(1+\frac{1}{\sqrt{\lambda_m}})t}^{t} e^{-\frac{L^2}{200(t-s)}} (-s)^{\frac{1-\lambda_m}{2(n-1)}} ds \\ &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-\frac{1}{n-1}} \int_{(1+\frac{1}{\sqrt{\lambda_m}})t}^{t} e^{-\frac{L^2}{200(t-s)}} (-s)^{\frac{1-\lambda_m}{2(n-1)}} ds \\ &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-\frac{1}{n-1}} \left(1 + \frac{1}{\sqrt{\lambda_m}}\right)^{\frac{2n-1-\lambda_m}{2(n-1)}} (-t)^{\frac{2n-1-\lambda_m}{2(n-1)}} \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-\frac{1}{n-1}} e^{-\frac{L^2\sqrt{\lambda_m}}{200(t-t)}} (-t)^{\frac{2n-1-\lambda_m}{2(n-1)}} \end{split}$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$. This implies

$$\begin{aligned} |v_m(z,t)| &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L^2}{16(-t)}\right)^{1-\frac{\lambda_m}{2(n-1)}} \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-\frac{1}{n-1}} \left(1 + \frac{1}{\sqrt{\lambda_m}}\right)^{\frac{2n-1-\lambda_m}{2(n-1)}} \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-\frac{1}{n-1}} e^{-\frac{L^2\sqrt{\lambda_m}}{200(-t)}} \end{aligned}$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$. We now sum over all $m \ge n+1$. Using the estimate $\sup_{\theta \in S^{n-1}} |Y_m(\theta)| \le C \lambda_m^{n-1}$, we obtain

$$\left|\sum_{m=n+1}^{\infty} v_m(z,t) Y_m(\theta)\right| \le C \sum_{m=n+1}^{\infty} \lambda_m^{n-1} |v_m(z,t)| \le C L^{-\frac{1}{n-1}} \varepsilon + C(L) \varepsilon_1 \varepsilon_2$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$.

We next consider the case when $1 \leq m \leq n$, so that $\lambda_m = n - 1$ and $Y_m(\theta) = \theta_m$. In this case, the function $v_m(z,t)$ satisfies

$$\frac{\partial}{\partial t}v_m(z,t) = \frac{\partial^2}{\partial z^2}v_m(z,t)$$

The results in Step 3 and Step 4 imply that, given any point $(z_0, t_0) \in [-\frac{L}{4}, \frac{L}{4}] \times [-\frac{L^2}{16}, -1]$, we can find real numbers $a_1, \ldots, a_n, b_1, \ldots, b_n$ (depending on (z_0, t_0)) such that

$$|a_1| + \ldots + |a_n| \le C(L)\varepsilon,$$

 $|b_1| + \ldots + |b_n| \le C(L)\varepsilon,$

and

$$|v_m(z,t) - (a_m + b_m z)| \le C\varepsilon(-t_0)^{\frac{1}{2}} + C(L)\varepsilon_1\varepsilon_1$$

for $z \in [z_0 - (-t_0)^{\frac{1}{2}}, z_0 + (-t_0)^{\frac{1}{2}}]$ and $t \in [2t_0, t_0]$. Using interior estimates for the linear heat equation, we obtain

$$\left|\frac{\partial^2 v_m}{\partial z^2}(z,t)\right| \le \left(C\varepsilon + C(L)\varepsilon_1\varepsilon\right)(-t)^{-\frac{1}{2}}$$

in the parabolic cylinder $\{z \in [-\frac{L}{4}, \frac{L}{4}], t \in [-\frac{L^2}{16}, -1]\}$. Using the solution formula for the one-dimensional heat equation with Dirichlet boundary

condition on the rectangle $\left[-\frac{L}{4}, \frac{L}{4}\right] \times \left[-\frac{L^2}{16}, -1\right]$, we obtain

$$\begin{aligned} \frac{\partial^2 v_m}{\partial z^2}(z,t) \Big| &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L}{4}\right)^{-1} \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L \int_{-\frac{L^2}{16}}^t e^{-\frac{L^2}{100(t-s)}} (t-s)^{-\frac{3}{2}} (-s)^{-\frac{1}{2}} ds \\ &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L}{4}\right)^{-1} \\ &+ (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-2} \int_{-\frac{L^2}{16}}^t (-s)^{-\frac{1}{2}} ds \\ &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-1} \end{aligned}$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$. Consequently, there exist real numbers $A_1, \ldots, A_n, B_1, \ldots, B_n$ such that

$$|v_m(z,t) - (A_m + B_m z)| \le CL^{-1}\varepsilon + C(L)\varepsilon_1\varepsilon$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$.

Finally, we consider the case m = 0. By the results in Step 2, the function $u = \langle \bar{K}_{\alpha}, \nu \rangle$ satisfies

$$|u(\theta, z, t) + \operatorname{div}_{S^{n-1}}(r(\theta, z, t) J_{\alpha}\theta)| \le C(L)\varepsilon_1\varepsilon_2$$

Integrating over $\theta \in S^{n-1}$ gives

$$\left| \int_{S^{n-1}} u(\theta, z, t) \, d\theta \right| \le C(L) \varepsilon_1 \varepsilon_2$$

hence

$$|v_0(z,t)| \le C(L)\varepsilon_1\varepsilon$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$. Putting everything together, we conclude that

 $|\tilde{u}(\theta, z, t) - (A_1\theta_1 + \ldots + A_n\theta_n) - (B_1\theta_1 + \ldots + B_n\theta_n)z| \le CL^{-\frac{1}{n-1}}\varepsilon + C(L)\varepsilon_1\varepsilon$ for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1].$

Step 6: By combining the results in Step 4 and Step 5, we can draw the following conclusion. For each $\alpha \in \{1, \ldots, \frac{n(n-1)}{2}\}$ we can find real numbers $A_{\alpha,1}, \ldots, A_{\alpha,n}, B_{\alpha,1}, \ldots, B_{\alpha,n}$ such that

$$|A_{\alpha,1}| + \ldots + |A_{\alpha,n}| \le C(L)\varepsilon,$$

$$|B_{\alpha,1}| + \ldots + |B_{\alpha,n}| \le C(L)\varepsilon,$$

and

$$|u_{\alpha}(\theta, z, t) - (A_{\alpha,1}\theta_{1} + \ldots + A_{\alpha,n}\theta_{n}) - (B_{\alpha,1}\theta_{1} + \ldots + B_{\alpha,n}\theta_{n})z|$$

$$\leq CL^{-\frac{1}{n-1}}\varepsilon + C(L)\varepsilon_{1}\varepsilon$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$, where $u_{\alpha}(\theta, z, t) := \langle \bar{K}_{\alpha}, \nu \rangle$.

Step 7: For each $i \in \{1, \ldots, n\}$, we define

$$E_{i} = \frac{n}{|S^{n-1}|} \int_{S^{n-1}} r(\theta, 0, -1) \,\theta_{i} \, d\theta$$

and

$$F_i = \frac{n}{2|S^{n-1}|} \int_{S^{n-1}} [r(\theta, 1, -1) - r(\theta, -1, -1)] \theta_i \, d\theta.$$

We have shown in Step 2 that $|\nabla^{S^{n-1}}r| \leq C(L)\varepsilon$. This implies $|E_i|, |F_i| \leq C(L)\varepsilon$ for $i \in \{1, \ldots, n\}$. By the results in Step 2, the function $u_{\alpha} = \langle \bar{K}_{\alpha}, \nu \rangle$ satisfies

$$u_{\alpha}(\theta, z, t) + \operatorname{div}_{S^{n-1}}(r(\theta, z, t) J_{\alpha}\theta) \le C(L)\varepsilon_1\varepsilon_2$$

A direct calculation gives

$$\operatorname{div}_{S^{n-1}}(r(\theta, z, t) J_{\alpha}\theta) \theta_i = \operatorname{div}_{S^{n-1}}(r(\theta, z, t) \theta_i J_{\alpha}\theta) - r(\theta, z, t) \sum_{j=1}^n J_{\alpha, ij} \theta_j,$$

where $J_{\alpha,ij}$ denote the components of the anti-symmetric matrix J_{α} . Putting these facts together, we obtain

$$\left| u_{\alpha}(\theta, z, t) \,\theta_{i} + \operatorname{div}_{S^{n-1}}(r(\theta, z, t) \,\theta_{i} \,J_{\alpha}\theta) - r(\theta, z, t) \sum_{j=1}^{n} J_{\alpha, ij} \,\theta_{j} \right| \leq C(L)\varepsilon_{1}\varepsilon_{1}$$

for all $i \in \{1, \ldots, n\}$. Integrating over $\theta \in S^{n-1}$ gives

$$\max_{\alpha,i} \left| \frac{n}{|S^{n-1}|} \int_{S^{n-1}} u_{\alpha}(\theta, 0, -1) \,\theta_i \, d\theta - \sum_{j=1}^n J_{\alpha,ij} E_j \right| \le C(L) \varepsilon_1 \varepsilon$$

and

$$\max_{\alpha,i} \left| \frac{n}{2 \left| S^{n-1} \right|} \int_{S^{n-1}} \left[u_{\alpha}(\theta, 1, -1) - u_{\alpha}(\theta, -1, -1) \right] \theta_i \, d\theta - \sum_{j=1}^n J_{\alpha,ij} F_j \right| \le C(L) \varepsilon_1 \varepsilon_2$$

On the other hand, using the estimate for $u_{\alpha}(\theta, z, t) - (A_{\alpha,1}\theta_1 + \ldots + A_{\alpha,n}\theta_n) - (B_{\alpha,1}\theta_1 + \ldots + B_{\alpha,n}\theta_n)z$ in Step 6, we obtain

$$\max_{\alpha,i} \left| \frac{n}{|S^{n-1}|} \int_{S^{n-1}} u_{\alpha}(\theta, 0, -1) \theta_i \, d\theta - A_{\alpha,i} \right| \le CL^{-\frac{1}{n-1}} \varepsilon + C(L) \varepsilon_1 \varepsilon$$

and

$$\begin{split} & \max_{\alpha,i} \left| \frac{n}{2 \left| S^{n-1} \right|} \int_{S^{n-1}} \left[u_{\alpha}(\theta, 1, -1) - u_{\alpha}(\theta, -1, -1) \right] \theta_i \, d\theta - B_{\alpha,i} \\ & \leq C L^{-\frac{1}{n-1}} \varepsilon + C(L) \varepsilon_1 \varepsilon. \end{split}$$

Putting these facts together, we obtain

$$\max_{\alpha,i} \left| A_{\alpha,i} - \sum_{j=1}^{n} J_{\alpha,ij} E_j \right| \le C L^{-\frac{1}{n-1}} \varepsilon + C(L) \varepsilon_1 \varepsilon$$

and

$$\max_{\alpha,i} \left| B_{\alpha,i} - \sum_{j=1}^{n} J_{\alpha,ij} F_j \right| \le C L^{-\frac{1}{n-1}} \varepsilon + C(L) \varepsilon_1 \varepsilon.$$

Substituting this back into the estimate for $u_{\alpha}(\theta, z, t) - (A_{\alpha,1}\theta_1 + \ldots +$ $A_{\alpha,n}\theta_n$) – $(B_{\alpha,1}\theta_1 + \ldots + B_{\alpha,n}\theta_n)z$ in Step 6, we finally conclude

$$\max_{\alpha} \left| \langle \bar{K}_{\alpha}, \nu \rangle - \sum_{i,j=1}^{n} J_{\alpha,ij} (E_j \theta_i + F_j \theta_i z) \right| \le C L^{-\frac{1}{n-1}} \varepsilon + C(L) \varepsilon_1 \varepsilon$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$. Step 8: Let us define a vector $q \in \mathbb{R}^{n+1}$ by $q_i = E_i$ for $i \in \{1, \ldots, n\}$ and $q_{n+1} = 0$. Moreover, let us define an anti-symmetric matrix $\sigma \in so(n+1)$ by $\sigma_{ij} = 0$ for $i, j \in \{1, \ldots, n\}$ and $\sigma_{i,n+1} = F_i$ for $i \in \{1, \ldots, n\}$. Clearly, $|q| \leq C(L)\varepsilon$ and $|\sigma| \leq C(L)\varepsilon$. We define a normalized set of rotation vector fields $\tilde{\mathcal{K}} = \{\tilde{K}_{\alpha} : 1 \le \alpha \le \frac{n(n-1)}{2}\}$ by

$$\tilde{K}_{\alpha}(x) := SJ_{\alpha}S^{-1}(x-q),$$

where $S := \exp(\sigma) \in SO(n+1)$. Then

$$\max_{\alpha} \left| \langle \tilde{K}_{\alpha} - \bar{K}_{\alpha}, \nu \rangle + \sum_{i,j=1}^{n} J_{\alpha,ij} (E_j \theta_i + F_j \theta_i z) \right| \le C(L) \varepsilon_1 \varepsilon$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$. Combining this estimate with the estimate in Step 7, we conclude that

$$\max_{\alpha} |\langle \tilde{K}_{\alpha}, \nu \rangle| \le CL^{-\frac{1}{n-1}} \varepsilon + C(L) \varepsilon_1 \varepsilon$$

for $z \in [-20n, 20n]$ and $t \in [-400n^2, -1]$. Consequently, the point $(\bar{x}, -1)$ is $(CL^{-\frac{1}{n-1}}\varepsilon + C(L)\varepsilon_1\varepsilon)$ -symmetric. In particular, if we choose L sufficiently large and ε_1 sufficiently small (depending on L), then $(\bar{x}, -1)$ is $\frac{\varepsilon}{2}$ -symmetric. This completes the proof of the Neck Improvement Theorem.

5. Proof of rotational symmetry

Let $M_t, t \in (-\infty, 0]$, be a noncompact ancient solution of mean curvature flow in \mathbb{R}^{n+1} which is strictly convex, uniformly two-convex, and noncollapsed. For each t, the hypersurface M_t bounds a convex domain which we denote by Ω_t . As in [2], if -t is sufficiently large, there exists a unique point $p_t \in M_t$ where the mean curvature attains its maximum. Moreover, this is a non-degenerate maximum in the sense that the Hessian of the mean curvature at p_t is negative definite, and the hypersurface M_t looks like the bowl soliton near p_t .

Let ε_1 and L be the constants in the Neck Improvement Theorem. Recall that $H_{\text{max}}(t)$ is uniformly bounded from below. By Proposition 3.1 in [8], we can find a large constant Λ such that the following holds. If x is a point on M_t such that $|x - p_t| \ge \Lambda$, then x lies at the center of an ε_1 -neck and furthermore $H(x,t) |x - p_t| \ge 10^6 n^2 L.$

26

Lemma 5.1. There exists a time T < 0 with the property that $\frac{d}{dt}|x-p_t| < 0$ whenever $t \leq T$, $x \in M_t \cup \Omega_t$, and $|x-p_t| \geq \Lambda$.

Proof. If -t is sufficiently large, then M_t looks like the bowl soliton near the point p_t . Hence, if -t is sufficiently large, then the vector $\frac{d}{dt}p_t$ is almost parallel to $-\nu(p_t, t)$. Moreover, using the convexity of Ω_t , we obtain $-\langle x - p_t, \nu(p_t, t) \rangle \ge c |x - p_t|$ whenever $t \le T$, $x \in M_t \cup \Omega_t$, and $|x - p_t| \ge \Lambda$. Here, c is a small positive constant. Putting these facts together, we conclude that $\langle x - p_t, \frac{d}{dt}p_t \rangle > 0$ whenever $t \le T$, $x \in M_t \cup \Omega_t$, and $|x - p_t| \ge \Lambda$. This gives $\frac{1}{2}\frac{d}{dt}|x - p_t|^2 = -\langle x - p_t, \frac{d}{dt}p_t \rangle < 0$ whenever $t \le T$, $x \in M_t \cup \Omega_t$, and $|x - p_t| \ge \Lambda$.

Lemma 5.2. Let T be defined as in Lemma 5.1. Suppose that $\overline{t} \leq T$, and \overline{x} is a point on $M_{\overline{t}}$ satisfying $|\overline{x} - p_{\overline{t}}| \geq \Lambda$. Then $|\overline{x} - p_t| \geq |\overline{x} - p_{\overline{t}}|$ for all $t \leq \overline{t}$.

Proof. We argue by contradiction. Suppose that the assertion is false, and let $\tilde{t} := \sup\{t \leq \bar{t} : |\bar{x} - p_t| < |\bar{x} - p_{\bar{t}}|\}$. By Lemma 5.1, we have $\frac{d}{dt}|\bar{x} - p_t||_{t=\bar{t}} < 0$. Consequently, $\tilde{t} < \bar{t}$. Moreover, for each $t \in [\tilde{t}, \bar{t}]$, we have $\bar{x} \in M_t \cup \Omega_t$ and $|\bar{x} - p_t| \geq |\bar{x} - p_{\bar{t}}| \geq \Lambda$. Using Lemma 5.1, we obtain $\frac{d}{dt}|\bar{x} - p_t| < 0$ for all $t \in [\tilde{t}, \bar{t}]$. Consequently, $|\bar{x} - p_{\bar{t}}| > |\bar{x} - p_{\bar{t}}|$. This contradicts the definition of \tilde{t} . This completes the proof of Lemma 5.2.

Proposition 5.3. Let T be defined as in Lemma 5.1. If $t \leq T$, $x \in M_t$ and $|x - p_t| \geq 2^{\frac{j}{400}} \Lambda$, then (x, t) is $2^{-j} \varepsilon_1$ -symmetric.

Proof. The proof is by induction on j. It follows from our choice of Λ that the assertion is true for j = 0. Suppose now that $j \ge 1$ and the assertion holds for j - 1. We claim that the assertion holds for j. Suppose this is false. Then there exists a time $\bar{t} \le T$ and a point $\bar{x} \in M_{\bar{t}}$ such that $|\bar{x}-p_{\bar{t}}| \ge 2^{\frac{j}{400}} \Lambda$ and (\bar{x}, \bar{t}) is not $2^{-j} \varepsilon_1$ -symmetric. By the Neck Improvement Theorem, there exists a point $(x, t) \in \hat{\mathcal{P}}(\bar{x}, \bar{t}, L, L^2)$ such that either (x, t) is not $2^{-j+1}\varepsilon_1$ -symmetric or (x, t) does not lie at the center of an ε_1 -neck. In view of the induction hypothesis, we conclude that $|x - p_t| \le 2^{\frac{j-1}{400}} \Lambda$. Since $t \le \bar{t} \le T$, Lemma 5.2 gives $|\bar{x} - p_{\bar{t}}| \le |\bar{x} - p_t|$. Putting these facts together, we obtain

$$\begin{split} |\bar{x} - p_{\bar{t}}| &\leq |\bar{x} - p_t| \\ &\leq |x - p_t| + |x - \bar{x}| \\ &\leq 2^{\frac{j-1}{400}} \Lambda + 10 \, n^2 L \, H(\bar{x}, \bar{t})^{-1} \\ &\leq 2^{-\frac{1}{400}} \, |\bar{x} - p_{\bar{t}}| + \frac{1}{1000} \, |\bar{x} - p_{\bar{t}}| \\ &< |\bar{x} - p_{\bar{t}}|. \end{split}$$

This is a contradiction.

Theorem 5.4. Let T be defined as in Lemma 5.1. Then the hypersurface M_t is rotationally symmetric for each $t \leq T$.

Proof. The argument is similar to [2]. We fix a time $\overline{t} \leq T$. For each j, let $\Omega^{(j)}$ be the set of all points (x,t) in space-time satisfying $t \leq \overline{t}$ and $|x - p_t| \leq 2^{\frac{j}{400}} \Lambda$. If j is sufficiently large, then $H(x,t) \geq n \cdot 2^{-\frac{j}{400}}$ for each point $(x,t) \in \Omega^{(j)}$. Proposition 5.3 guarantees that every point $(x,t) \in \partial \Omega^{(j)}$ is $2^{-j}\varepsilon_1$ -symmetric. Consequently, given any point $(x,t) \in \partial \Omega^{(j)}$, we can find a normalized set of rotation vector fields $\mathcal{K}^{(x,t)} = \{K_{\alpha}^{(x,t)} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ such that $\max_{\alpha} |\langle K_{\alpha}^{(x,t)}, \nu \rangle| H \leq 2^{-j}\varepsilon_1$ on $\hat{\mathcal{P}}(x,t,10,100)$. Using Lemma 4.2, we are able to control how the axis of rotation of $\mathcal{K}^{(x,t)}$ varies as we vary the point (x,t). More precisely, if (x_1,t_1) and (x_2,t_2) are points in $\partial \Omega^{(j)}$ satisfying $(x_2,t_2) \in \hat{\mathcal{P}}(x_1,t_1,\frac{1}{10n},\frac{1}{100n^2})$, then

$$\inf_{\substack{\omega \in O(\frac{n(n-1)}{2})}} \sup_{B_{10nH(x_2,t_2)}-1} \max_{\alpha} \left| K_{\alpha}^{(x_1,t_1)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta} K_{\beta}^{(x_2,t_2)} \right| \\
\leq C \, 2^{-j} \, H(x_2,t_2)^{-1}.$$

Now, if (x_2, t_2) is a point in $\partial \Omega^{(j)}$, then the mean curvature $H(x_2, t_2)$ satisfies the estimate $\frac{1}{C} 2^{-\frac{j}{400}} \leq H(x_2, t_2) \leq C$. Hence, if (x_1, t_1) and (x_2, t_2) are points in $\partial \Omega^{(j)}$ satisfying $(x_2, t_2) \in \hat{\mathcal{P}}(x_1, t_1, \frac{1}{10n}, \frac{1}{100n^2})$, then

$$\inf_{\substack{\omega \in O(\frac{n(n-1)}{2})}} \sup_{B_{1/C}(x_2)} \max_{\alpha} \left| K_{\alpha}^{(x_1,t_1)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta} K_{\beta}^{(x_2,t_2)} \right| \\
\leq C 2^{-j + \frac{j}{400}},$$

and consequently

$$\inf_{\substack{\omega \in O(\frac{n(n-1)}{2})}} \sup_{B_{2^{j/20}}(x_2)} \max_{\alpha} \left| K_{\alpha}^{(x_1,t_1)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta} K_{\beta}^{(x_2,t_2)} \right| \\
\leq C \, 2^{-j+\frac{j}{10}}.$$

By a repeated application of this estimate, we can show that, for each j, there exists a normalized set of rotation vector fields $\mathcal{K}^{(j)} = \{K_{\alpha}^{(j)} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ with the following property: if (x,t) is a point in $\partial\Omega^{(j)}$ satisfying $\bar{t} - 2\frac{j}{100} \leq t \leq \bar{t}$, then

$$\inf_{\omega \in O(\frac{n(n-1)}{2})} \max_{\alpha} \left| K_{\alpha}^{(j)} - \sum_{\beta=1}^{\frac{n(n-1)}{2}} \omega_{\alpha\beta} K_{\beta}^{(x,t)} \right| \le C \, 2^{-\frac{j}{2}}$$

at the point (x, t).

For each point $(x,t) \in \partial \Omega^{(j)}$, we have $\max_{\alpha} |\langle K_{\alpha}^{(x,t)}, \nu \rangle| \leq 2^{-j}$ at the point (x,t). Consequently, $\max_{\alpha} |\langle K_{\alpha}^{(j)}, \nu \rangle| \leq C 2^{-\frac{j}{2}}$ for all points $(x,t) \in \partial \Omega^{(j)}$ satisfying $\overline{t} - 2^{\frac{j}{100}} \leq t \leq \overline{t}$. Finally, we note that, for each $t \in [\overline{t} - 2^{\frac{j}{100}}, \overline{t}]$, the point p_t has distance at most $C 2^{\frac{j}{100}}$ from the point $p_{\overline{t}}$. This implies $\max_{\alpha} |\langle K_{\alpha}^{(j)}, \nu \rangle| \leq C 2^{\frac{j}{100}}$ for all points $(x,t) \in \Omega^{(j)}$ with $t \in [\overline{t} - 2^{\frac{j}{100}}, \overline{t}]$.

For each $\alpha \in \{1, \dots, \frac{n(n-1)}{2}\}$, we define a function $f_{\alpha}^{(j)} : \Omega^{(j)} \to \mathbb{R}$ by

$$f_{\alpha}^{(j)} := \exp(-2^{-\frac{j}{200}}(\bar{t}-t)) \frac{\langle K_{\alpha}^{(j)}, \nu \rangle}{H - 2^{-\frac{j}{400}}}$$

Since $\max_{\alpha} |\langle K_{\alpha}^{(j)}, \nu \rangle| \leq C 2^{-\frac{j}{2}}$ for all points $(x, t) \in \partial \Omega^{(j)}$ satisfying $\bar{t} - 2^{\frac{j}{100}} \leq t \leq \bar{t}$, we conclude that

$$\max_{\alpha} |f_{\alpha}^{(j)}(x,t)| \le C \, 2^{-\frac{j}{4}}$$

for all points $(x,t) \in \partial \Omega^{(j)}$ satisfying $\bar{t} - 2^{\frac{j}{100}} \leq t \leq \bar{t}$. The same estimate holds for all points $(x,t) \in \Omega^{(j)}$ with $t = \bar{t} - 2^{\frac{j}{100}}$. On the other hand, the function $f_{\alpha}^{(j)}$ satisfies the evolution equation

$$\frac{\partial}{\partial t} f_{\alpha}^{(j)} = \Delta f_{\alpha}^{(j)} + \frac{2}{H - 2^{-\frac{j}{400}}} \left\langle \nabla H, \nabla f_{\alpha}^{(j)} \right\rangle - 2^{-\frac{j}{400}} \left(\frac{|A|^2}{H - 2^{-\frac{j}{400}}} - 2^{-\frac{j}{400}} \right) f_{\alpha}^{(j)}.$$

On the set $\Omega^{(j)}$, we have

$$\frac{|A|^2}{H - 2^{-\frac{j}{400}}} - 2^{-\frac{j}{400}} \ge \frac{1}{n} \frac{H^2}{H - 2^{-\frac{j}{400}}} - 2^{-\frac{j}{400}} \ge \frac{1}{n} H - 2^{-\frac{j}{400}} \ge 0.$$

By the maximum principle, we obtain

$$\sup_{\substack{(x,t)\in\Omega^{(j)}, \bar{t}-2^{j} \\ 100 \le t \le \bar{t} \\ \le \max\left\{\sup_{\substack{(x,t)\in\partial\Omega^{(j)}, \bar{t}-2^{j} \\ (x,t)\in\partial\Omega^{(j)}, \bar{t}-2^{j} \\ 100 \le t \le \bar{t} \\ \le C 2^{-\frac{j}{4}}} |f_{\alpha}^{(j)}(x,t)|, \sup_{\substack{(x,t)\in\Omega^{(j)}, t=\bar{t}-2^{j} \\ 100 \\$$

for each $\alpha \in \{1, \ldots, \frac{n(n-1)}{2}\}$. Thus, we conclude that $\max_{\alpha} |\langle K_{\alpha}^{(j)}, \nu \rangle| \leq C 2^{-\frac{i}{4}}$ for all points $(x,t) \in \Omega^{(j)}$ with $t = \bar{t}$. In particular, the distance of the point $p_{\bar{t}}$ from the axis of rotation of $\mathcal{K}^{(j)} = \{K_{\alpha}^{(j)} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ is bounded from above by a uniform constant which is independent of j. Sending $j \to \infty$, the sequence $\mathcal{K}^{(j)} = \{K_{\alpha}^{(j)} : 1 \leq \alpha \leq \frac{n(n-1)}{2}\}$ converges to a normalized set of rotation vector fields which are tangential along $M_{\bar{t}}$. This completes the proof of Theorem 5.4.

Once we know that M_t is rotationally symmetric for -t sufficiently large, it follows from standard arguments that M_t is rotationally symmetric for all $t \in (-\infty, 0]$ (see [2], Proposition 5.5).

6. UNIQUENESS OF ANCIENT SOLUTIONS WITH ROTATIONAL SYMMETRY

Let M_t be an ancient solution to the mean curvature flow in \mathbb{R}^{n+1} which is strictly convex, uniformly two-convex, and noncollapsed. We may assume that M_t is symmetric with respect to the x_{n+1} -axis. Let us write M_t as a graph of a rotationally symmetric function f on \mathbb{R}^n . The function f satisfies the equation

$$f_t = \frac{f_{rr}}{1 + f_r^2} + \frac{n - 1}{r} f_r$$

Note that f(r, t) may not be defined for all r.

Conversely, we may write the radius r as a function of (z, t), so that

$$f(r(z,t),t) = z.$$

Then r(z,t) satisfies the equation

$$r_t = \frac{r_{zz}}{1 + r_z^2} - \frac{n - 1}{r}.$$

Since M_t is convex, we have

$$r_z > 0, \qquad r_t < 0, \qquad r_{zz} < 0.$$

Without loss of generality, we assume that the tip of M_0 is at the origin. In other words, f(0,0) = 0 and r(0,0) = 0.

As in [2], let $q_t = (0, \ldots, 0, f(0, t))$ denote the tip of M_t , and let $H_{\text{tip}}(t)$ denote the mean curvature of M_t at the tip q_t . By the Harnack inequality [6], the function $t \mapsto H_{\text{tip}}(t)$ is monotone increasing. Hence, the limit $\mathcal{H} := \lim_{t \to -\infty} H_{\text{tip}}(t)$ exists. Using results in Section 3, we obtain $|q_t| \ge c(-t)$ for -t sufficiently large. This gives $\mathcal{H} > 0$.

We first prove that $f_t(r,t)$ is monotone increasing in t.

Proposition 6.1. We have $f_{tt}(r,t) \ge 0$ everywhere.

Proof. This is a consequence of Hamilton's Harnack inequality for mean curvature flow [6]. See [2] for details.

We next show that $f_t(r,t)$ is bounded from below.

Proposition 6.2. We have $f_t(r,t) \ge \mathcal{H}$ at each point in space-time. Moreover, for each $r_0 > 0$,

$$\lim_{t \to -\infty} \sup_{r \le r_0} f_t(r, t) = \mathcal{H}.$$

Proof. Consider an arbitrary sequence of times $t_j \to -\infty$, and define $M_t^{(j)} := M_{t+t_j} - q_{t_j}$. Since the flows $M_t^{(j)}$ have uniformly bounded curvature, the sequence $M_t^{(j)}$ converges in C_{loc}^{∞} to a smooth eternal solution, which is

rotationally symmetric. At each point in time, the mean curvature at the tip of the limit solution equals \mathcal{H} . Consequently, we are in the equality case in the Harnack inequality. By [6], the limit solution must be a self-similar translator which is moving with speed \mathcal{H} . This gives

$$\lim_{j \to \infty} \sup_{r \le r_0} |f_t(r, t_j) - \mathcal{H}| = 0$$

for every $r_0 > 0$. Since $f_{tt}(r,t) \ge 0$ by Proposition 6.1, it follows that $f_t(r,t) \ge \mathcal{H}$ for all r and t.

In the next step, we show that $f_t(r, t)$ is monotone increasing in r.

Proposition 6.3. We have $f_{tr}(r,t) \ge 0$ everywhere.

Proof. Consider a time t_0 and a radius r_0 such that $f(r_0, t_0)$ is defined. Moreover, let t_j be a sequence of times such that $t_j \to -\infty$. For j large, we denote by Q_j the parabolic cylinder $Q_j = \{x_1^2 + \ldots + x_n^2 \leq r_0^2, t \in [t_j, t_0]\}$. It follows from the evolution equations for H and $\langle \omega, \nu \rangle$ that the maximum $\sup_{Q_j} H \langle \omega, \nu \rangle^{-1}$ must be attained on the parabolic boundary of Q_j . This gives

$$\sup_{\substack{x_1^2+\ldots+x_n^2 \le r_0^2, t=t_0}} H \langle \omega, \nu \rangle^{-1}$$

$$\leq \max \Big\{ \sup_{\substack{x_1^2+\ldots+x_n^2=r_0^2, t_j \le t \le t_0}} H \langle \omega, \nu \rangle^{-1}, \sup_{\substack{x_1^2+\ldots+x_n^2 \le r_0^2, t=t_j}} H \langle \omega, \nu \rangle^{-1} \Big\}.$$

Since $f_t(r,t) = H \langle \omega, \nu \rangle^{-1}$, it follows that

$$\sup_{r \le r_0} f_t(r, t_0) \le \max \left\{ \sup_{t_j \le t \le t_0} f_t(r_0, t), \sup_{r \le r_0} f_t(r, t_j) \right\}$$
$$= \max \left\{ f_t(r_0, t_0), \sup_{r \le r_0} f_t(r, t_j) \right\}.$$

Note that in the last step we have used Proposition 6.1. Finally, we pass to the limit as $j \to \infty$. Using Proposition 6.2, we obtain $\sup_{r \le r_0} f_t(r, t_0) \le \max\{f_t(r_0, t_0), \mathcal{H}\} = f_t(r_0, t_0)$. This completes the proof of Proposition 6.3.

By assumption, M_t is strictly convex, uniformly two-convex, and noncollapsed. Moreover, $H_{tip}(t)$ is bounded from below by \mathcal{H} . Hence, there exists a small constant $\varepsilon_0 \in (0, \frac{1}{20})$ and a decreasing function $\Lambda : (0, \varepsilon_0] \to \mathbb{R}$ such that given any $\varepsilon \in (0, \varepsilon_0]$, if $|\bar{x} - q_t| \ge \Lambda(\varepsilon)$, then (\bar{x}, \bar{t}) is a center of ε -neck (cf. [8], Proposition 3.1). We recall three estimates from [2]. These results were stated for n = 2 in [2], but the arguments carry over directly to higher dimensions.

Lemma 6.4. On every ε_0 -neck, $rr_z = \frac{r}{f_r} \leq (1+2\varepsilon_0)(n-1)\mathcal{H}^{-1}$.

Proof. This follows from the inequality $f_t \geq \mathcal{H}$ established in Proposition 6.2. See [2], Lemma 6.4, for details.

Lemma 6.5. There exists a constant $C_0 \ge 1$ such that $r^m \left| \frac{\partial^m}{\partial z^m} r \right| \le C_0$ holds for m = 1, 2, 3 at the center of ε_0 -necks with $r \ge 1$.

Proof. Analogous to Lemma 6.5 in [2].

Proposition 6.6. There exist constants C_1 and C_2 with the following property. If $r \ge C_1$, then $0 \le -r_{zz}(z,t) \le C_2 r(z,t)^{-\frac{5}{2}}$.

Proof. Analogous to Proposition 6.6 in [2].

For each z < 0, we define a real number $\mathcal{T}(z)$ by

$$r(z,t) > 0$$
 for $t < \mathcal{T}(z)$, $\lim_{t \to \mathcal{T}(z)} r(z,t) = 0.$

In other words, if $t = \mathcal{T}(z)$, then f(0,t) = z, and the tip of M_t is located at $(0, \ldots, 0, z)$. The following result allows us to estimate r(z, t) in terms of $\mathcal{T}(z) - t$.

Corollary 6.7. We have

 $2(n-1) [\mathcal{T}(z) - t] \le r(z,t)^2 \le 2(n-1) [\mathcal{T}(z) - t] + 8C_2 [\mathcal{T}(z) - t]^{\frac{1}{4}} + C_1^2$ if z < 0 and r(z,t) is sufficiently large.

Proof. Let us fix a point (\bar{z}, \bar{t}) . The inequality $(r^2+2(n-1)t)_t = \frac{2rr_{zz}}{1+r_z^2} < 0$ implies

$$r(\bar{z},\bar{t})^2 \ge 2(n-1)\left[\mathcal{T}(\bar{z})-\bar{t}\right].$$

Moreover, if $r \ge C_1$, then $(r^2+2(n-1)t)_t = \frac{2rr_{zz}}{1+r_z^2} \ge -2C_2r^{-\frac{3}{2}}$ by Proposition 6.6. Let $\tilde{t} \le \mathcal{T}(\bar{z})$ be chosen so that $r(\bar{z}, \tilde{t}) = C_1$. Then

$$\begin{aligned} r(\bar{z},\bar{t})^2 &= C_1^2 + 2(n-1)(\tilde{t}-\bar{t}) - \int_{\bar{t}}^{\bar{t}} (r(\bar{z},t)^2 + 2(n-1)t)_t \, dt \\ &\leq C_1^2 + 2(n-1)(\tilde{t}-\bar{t}) + 2C_2 \int_{\bar{t}}^{\bar{t}} r(\bar{z},t)^{-\frac{3}{2}} \, dt \\ &\leq C_1^2 + 2(n-1)(\tilde{t}-\bar{t}) + 2C_2 \int_{\bar{t}}^{\bar{t}} [\mathcal{T}(\bar{z})-t]^{-\frac{3}{4}} \, dt \\ &\leq C_1^2 + 2(n-1)(\tilde{t}-\bar{t}) - 8C_2 [\mathcal{T}(\bar{z})-\bar{t}]^{\frac{1}{4}} + 8C_2 [\mathcal{T}(\bar{z})-\bar{t}]^{\frac{1}{4}} \\ &\leq C_1^2 + 2(n-1) [\mathcal{T}(\bar{z})-\bar{t}] + 8C_2 [\mathcal{T}(\bar{z})-\bar{t}]^{\frac{1}{4}}. \end{aligned}$$

This proves the assertion.

Lemma 6.8. Let δ be an arbitrary positive real number. Then

$$r(0,t)r_z(0,t) \ge (n-1)(\mathcal{H}^{-1}-\delta)$$

provided that -t is sufficiently large.

Proof. We may assume that $R = r(0,t) \ge C_1$. Then every point (x,t) with $x_{n+1} = 0$ lies at the center of an ε_0 -neck. Consequently, $|r(z,t) - R| \le \varepsilon_0 R$ for $|z| \le 2R$. Moreover, we have $rr_z \le (1+2\varepsilon_0) (n-1) \mathcal{H}^{-1}$ by Lemma 6.4, and $|(rr_z)_z| = |rr_{zz} + r_z^2| \le C_3 R^{-\frac{3}{2}}$ by Proposition 6.6. Hence, if $R^{\frac{1}{2}} \ge 4C_3\delta^{-1}$, then we have

$$|r(z,t)r_z(z,t) - r(0,t)r_z(0,t)| \le 2C_3 R^{-\frac{1}{2}} \le \frac{\delta}{2}$$

for all $z \in [-2R, 2R]$.

Using Corollary 6.7, we obtain

$$r(-R,t)^2 \ge 2(n-1) \left[\mathcal{T}(-R) - t\right],$$

 $r(-2R,t)^2 \ge 2(n-1) \left[\mathcal{T}(-2R) - t\right]$

and

$$r(-2R,t)^{2} \leq 2(n-1) \left[\mathcal{T}(-2R) - t \right] + 8C_{2} \left[\mathcal{T}(-2R) - t \right]^{\frac{1}{4}} + C_{1}^{2}$$
$$\leq 2(n-1) \left[\mathcal{T}(-2R) - t \right] + 8C_{2}r(-2R,t)^{\frac{1}{2}} + C_{1}^{2}$$
$$\leq 2(n-1) \left[\mathcal{T}(-2R) - t \right] + 8C_{2}R^{\frac{1}{2}} + C_{1}^{2},$$

where in the last step we have used the inequality $r(-2R, t) \leq r(0, t) = R$. From this, we deduce that

$$r(-R,t)^2 - r(-2R,t)^2 \ge 2(n-1)\left[\mathcal{T}(-R) - \mathcal{T}(-2R)\right] - 8C_2R^{\frac{1}{2}} - C_1^2.$$

On the other hand, using the identity $f_t(0,t) = H_{tip}(t)$, we obtain $\frac{d}{dz}\mathcal{T}(z) = H_{tip}(\mathcal{T}(z))^{-1}$. Hence, if R is sufficiently large, then $\frac{d}{dz}\mathcal{T}(z) \geq \mathcal{H}^{-1} - \frac{\delta}{4}$ for $z \in [-2R, -R]$. This gives

$$\mathcal{T}(-R) - \mathcal{T}(-2R) \ge \left(\mathcal{H}^{-1} - \frac{\delta}{4}\right) R$$

if R is sufficiently large. Putting these facts together, we obtain

$$r(-R,t)^2 - r(-2R,t)^2 \ge 2(n-1)\left(\mathcal{H}^{-1} - \frac{\delta}{2}\right)R$$

hence

$$\sup_{z \in [-2R, -R]} r(z, t) r_z(z, t) \ge (n-1) \left(\mathcal{H}^{-1} - \frac{\delta}{2} \right)$$

if -t is sufficiently large. Thus,

$$r(0,t)r_z(0,t) \ge (n-1)(\mathcal{H}^{-1}-\delta)$$

if -t is sufficiently large.

Proposition 6.9. Given $\delta > 0$, there exists a time $\overline{t} \in (-\infty, 0]$ (depending on δ) such that

$$r(z,t)r_z(z,t) \ge (n-1)\left(\mathcal{H}^{-1} - 2\delta\right),$$

holds for all $z \ge 0$ and $t \le \overline{t}$.

Proof. Let $\psi(z,t)$ denote the solution of the Dirichlet problem for the one-dimensional heat equation on the half line with initial condition $\lim_{t\to 0} \psi(z,t) = 1$ (see [2], Proposition 6.9). By Lemma 6.8, we can find a time \bar{t} so that $r(0,t)r_z(0,t) \ge (n-1)(\mathcal{H}^{-1}-\delta)$ for $t \le \bar{t}$. Moreover, we can arrange that $r(z,t) \ge C_1 + C_2$ for $z \ge 0$ and $t \le \bar{t}$. Given any $s < \bar{t}$, we define a function $\psi^{\delta,s}(z,t)$ by

$$\psi^{\delta,s}(z,t) = (n-1) \left(\mathcal{H}^{-1} - 2\delta - \mathcal{H}^{-1} \psi(2z,t-s) \right)$$

for $t \in (s, \overline{t}]$. We will show that $rr_z > \psi^{\delta, s}$ for all $z \ge 0$ and all $t \in (s, \overline{t}]$.

It is straightforward to verify that $r(0,t)r_z(0,t) \ge (n-1)(\mathcal{H}^{-1}-\delta) > \lim \sup_{z\to 0} \psi^{\delta,s}(z,t)$ for each $t \in (s,\bar{t}]$; $\liminf_{z\to\infty} r(z,t)r_z(z,t) \ge 0 > \lim \sup_{z\to\infty} \psi^{\delta,s}(z,t)$ for each $t \in (s,\bar{t}]$; and $r(z,s)r_z(z,s) \ge 0 > \limsup_{t\to s} \psi^{\delta,s}(z,t)$ for each z > 0.

On the other hand, for z>0 and $t\in(s,\bar{t}],$ we have , we have $1+rr_{zz}\geq0,$ hence

$$(rr_z)_t = \frac{(rr_z)_{zz}}{1+r_z^2} - \frac{2r_z r_{zz}(1+r_z^2+rr_{zz})}{(1+r_z^2)^2} > \frac{(rr_z)_{zz}}{1+r_z^2}.$$

Moreover,

$$(\psi^{\delta,s})_t = \frac{1}{4} \, (\psi^{\delta,s})_{zz} \le \frac{(\psi^{\delta,s})_{zz}}{1+r_z^2}$$

Using the maximum principle, we conclude that $rr_z > \psi^{\delta,s}$ for all $z \ge 0$ and all $t \in (s, \bar{t}]$. Sending $s \to -\infty$ gives the desired result.

Corollary 6.10. We can find a time $T \in (-\infty, 0]$ such that $r(z,t)^2 \ge (n-1)\mathcal{H}^{-1}z$ for all $z \ge 0$ and $t \le T$. In particular, if $t \le T$, then the function f(r,t) is defined for all $r \in [0,\infty)$.

Proof. By Proposition 6.9, we can find a time $T \in (-\infty, 0]$ such that $r(z,t)r_z(z,t) \ge \frac{1}{2}(n-1)\mathcal{H}^{-1}$ for all $z \ge 0$ and all $t \le T$. If we integrate over z, the assertion follows.

Proposition 6.11. For each $t \leq T$, we have $\lim_{z\to\infty} r(z,t)r_z(z,t) = (n-1)\mathcal{H}^{-1}$.

Proof. It follows from Lemma 6.4 that $\limsup_{z\to\infty} r(z,t)r_z(z,t) \leq (n-1)\mathcal{H}^{-1}$. Hence, it suffices to prove that $\liminf_{z\to\infty} r(z,t)r_z(z,t) \geq (n-1)\mathcal{H}^{-1}$ for each $t \leq T$. By Proposition 6.9, we know that $\liminf_{z\to\infty} r(z,t)r_z(z,t) \geq (n-1)(\mathcal{H}^{-1}-2\delta)$ if -t is sufficiently large. On the other hand, Lemma 6.5 gives

$$|(rr_z)_t| = |r_t r_z + rr_{zt}| = \left|\frac{r_z r_{zz}}{1 + r_z^2} + \frac{rr_{zzz}}{1 + r_z^2} - \frac{2rr_z r_{zz}^2}{(1 + r_z^2)^2}\right| \le \frac{4C_0^3}{r^2}$$

for $r \ge C_1$. Using Corollary 6.10, we conclude that the quantity $\liminf_{z\to\infty} r(z,t)r_z(z,t)$ has the same value for each $t \le T$. Putting these facts together, we conclude

0

that $\liminf_{z\to\infty} r(z,t)r_z(z,t) \ge (n-1)\mathcal{H}^{-1}$ for each $t \le T$.

Theorem 6.12. For each $t \leq T$, the solution M_t is a rotationally symmetric translating solution.

Proof. Since $rr_z = \frac{r}{f_r}$, Proposition 6.11 implies

$$\lim_{r \to \infty} \frac{f_r(r,t)}{r} = \frac{1}{n-1} \mathcal{H}$$

for each $t \leq T$. Moreover, Lemma 6.5 gives $\limsup_{z\to\infty} r^2 (-r_{zz}) < \infty$, hence $\limsup_{r\to\infty} \frac{r^2 f_{rr}}{f_r^2} < \infty$. Consequently,

$$\lim_{r \to \infty} \frac{f_{rr}(r,t)}{1 + f_r(r,t)^2} = 0.$$

Using the evolution equation for f(r, t), we obtain

$$\lim_{r \to \infty} f_t(r, t) = \lim_{r \to \infty} \frac{(n-1) f_r(r, t)}{r} = \mathcal{H}$$

for each $t \leq T$. Using Proposition 6.3, we conclude that $f_t(r,t) \leq \mathcal{H}$ for all $r \geq 0$ and all $t \leq T$. Therefore, Proposition 6.2 gives $f_t(r,t) = \mathcal{H}$ for all $r \geq 0$ and all $t \leq T$. Consequently, M_t is a translating soliton for each $t \leq T$.

Once we know that M_t is a translating soliton for -t sufficiently large, it follows from standard arguments that M_t is a translating soliton for all $t \in (-\infty, 0]$ (see [2], Proposition 6.14). This completes the proof of Theorem 1.1.

References

- S. Angenent, P. Daskalopoulos, and N. Šešum, Unique asymptotics of ancient convex mean curvature flow solutions, J. Diff. Geom. 111, 381–455 (2019)
- [2] S. Brendle and K. Choi, Uniqueness of convex ancient solutions to mean curvature flow in R³, Invent. Math. 217, 35–76 (2019)
- [3] T.H Colding and W.P. Minicozzi, II, Generic mean curvature flow I: generic singularities, Ann. of Math. 175, 755–833 (2012)
- [4] K. Ecker, Regularity Theory for Mean Curvature Flow, Birkhäuser, Boston, 2004
- K. Ecker and G. Huisken, Interior estimates for hypersurfaces moving by mean curvature, Invent. Math. 105, 547–569 (1991)
- [6] R. Hamilton, Harnack estimate for the mean curvature flow, J. Diff. Geom. 41, 215– 226 (1995)
- [7] R. Hashofer and B. Kleiner, Mean curvature flow of mean convex hypersurfaces, Comm. Pure Appl. Math. 70, 511–546 (2017)
- [8] R. Hashofer and B. Kleiner, Mean curvature flow with surgery, Duke Math. J. 166, 1591–1626 (2017)
- [9] G. Huisken, Asymptotic behavior for singularities of mean curvature flow, J. Diff. Geom. 31, 285–299 (1990)
- [10] G. Huisken and C. Sinestrari, Mean curvature flow with surgeries of two-convex hypersurfaces, Invent. Math. 175, 137–221 (2009)

- [11] F. Merle and H. Zaag, Optimal estimates for blowup rate and behavior for nonlinear heat equations, Comm. Pure Appl. Math. 51, 139–196 (1998)
- [12] W. Sheng and X.J. Wang, Singularity profile in the mean curvature flow, Methods Appl. Anal. 16, 139–155 (2009)
- [13] J.J. Stoker, Unbounded convex point sets, Amer. J. Math. 62, 165–179 (1940)
- B. White, The size of the singular set in mean curvature flow of mean convex sets, J. Amer. Math. Soc. 13, 665–695 (2000)
- B. White, The nature of singularities in mean curvature flow of mean convex sets, J. Amer. Math. Soc. 16, 123–138 (2003)

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK, NY 10027, USA.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVENUE, CAMBRIDGE MA 02138, USA.