## ON NORMALIZING DISCRETE MORSE FUNCTIONS

NICOLÁS A. CAPITELLI

Departamento de Ciencias Básicas, Universidad Nacional de Luján Buenos Aires, Argentina.

ABSTRACT. We present a way to *normalize* a combinatorial Morse function into an integer-valued canonical representative of the set of discrete Morse functions inducing a given gradient field.

**Introduction.** In his seminal paper [3], Forman showed how to assign a combinatorial Morse function to a given (admissible) discrete vector field W over a simplicial complex K. The value of this function on a given simplex  $\sigma$  is linked to the length of gradient paths of W starting in  $\sigma$ . In this note, we use a simplified version of Forman's construction to produce a Morse function  $h: K \to \mathbb{Z}_{\geq 0}$  taking the minimum possible value for each simplex. We call one such function *normalized*. Normalized Morse functions are natural representatives of all functions inducing the same gradient vector field.

Throughout, K will represent a finite simplicial complex and we shall write  $\sigma \prec \tau$  if  $\sigma$  is an immediate face of  $\tau$  in K (i.e.  $\dim(\sigma) = \dim(\tau) - 1$ ). A discrete vector field over K is a map  $W: K \to K \cup \{0\}$  satisfying

(W1) if  $W(\sigma) \neq 0$  then  $\sigma \prec W(\sigma)$ , (W2) if  $W(\sigma) = W(\sigma') \neq 0$  then  $\sigma = \sigma'$ , and (W3)  $W^2 = 0$ .

The simplices in  $W^{-1}(0) \setminus W(K)$  are called *critical*. A combinatorial Morse function over K is a map  $f: K \to \mathbb{R}$  satisfying for every  $\sigma \in K$ 

- (M1)  $|\{\eta \prec \sigma \mid f(\eta) \ge f(\sigma)\}| \le 1$  and
- $(M2) |\{\tau \succ \sigma \,|\, f(\tau) \leq f(\sigma)\}| \leq 1.$

Here |X| denotes the cardinality of the set X. The gradient vector field of a combinatorial Morse function f is the discrete vector field  $V_f$  over K defined by

$$V_f(\sigma) = \begin{cases} \tau & \text{if } \tau \succ \sigma \text{ and } f(\sigma) \ge f(\tau) \\ 0 & \text{otherwise.} \end{cases}$$

A simplex  $\sigma$  is critical for f if both cardinalities in (M1) and (M2) are zero (equivalently,  $\sigma$  is critical for  $V_f$ ). Two Morse functions f and g are said to be equivalent if they induce the same gradient vector field (or equivalently,  $f(\sigma) < f(\tau)$  if and only if  $g(\sigma) < g(\tau)$ for every  $\sigma \prec \tau$ ; see e.g. [1]). Given a discrete vector field W over K, a path of index k (relative to W) is a sequence of k-simplices  $\sigma_0, \ldots, \sigma_r \in W^{-1}(K)$  such that  $\sigma_i \neq \sigma_{i+1}$ and  $\sigma_{i+1} \prec W(\sigma_i)$  for all  $0 \le i \le r-1$ . The path is closed if  $\sigma_0 = \sigma_r$  and non-stationary if r > 0. By [3, Theorem 9.3], W is the gradient vector field of a combinatorial Morse function if and only if W has no non-stationary closed paths (admissible).

<sup>2010</sup> Mathematics Subject Classification. 05E45, 52B05.

Key words and phrases. Discrete Morse theory, combinatorial Morse functions, discrete vector fields. This work was partially supported by a Postdoc grant from CONICET. E-mail address: ncapitelli@unlu.edu.ar.

The function underlying a gradient field. For an admissible vector field W over K define a relation  $\triangleleft$  on K by taking the transitive closure of the following *atomic relations*:

- (1)  $\sigma \lessdot \tau$  if  $\sigma \prec \tau$ , and
- (2)  $\sigma > \tau$  if also  $\tau = W(\sigma)$ .


We shall write  $\sigma \sim \tau$  whenever  $\sigma \leq \tau$  and  $\sigma \geq \tau$ . Define the *height* of a simplex  $\sigma$  in  $(K, \leq)$  as

$$h(\sigma) := \max\{n \in \mathbb{Z}_{\geq 0} \mid \exists \sigma = \sigma_n \geqslant \sigma_{n-1} \geqslant \cdots \geqslant \sigma_0, \sigma_i \not\approx \sigma_{i+1}\}.$$

We call  $h: K \to \mathbb{Z}_{\geq 0}$  the height function associated to W.

EXAMPLE 1. The dimension function  $\dim(-): K \to \mathbb{Z}_{\geq 0}$  is the height function of the null vector field.

EXAMPLE 2. For the boundary of the 2-simplex on the vertices  $\{a, b, c\}$ , Figure 1 shows the height function of the vector field W defined by W(a) = ab, W(b) = bc and W = 0 otherwise.



**Figure 1.** Computing the height function h for a vector field W.

THEOREM 3. With the notations as above, h is combinatorial Morse function over K with gradient vector field W.

*Proof.* First we show that if  $\sigma \prec \tau$  are such that  $\sigma \sim \tau$  then  $W(\sigma) = \tau$ . Indeed, let

$$\sigma = \eta_0 \geqslant \eta_1 \geqslant \cdots \geqslant \eta_r = \tau$$

be a chain of atomic relations. Since  $\dim(\sigma) = \dim(\tau) - 1$  and  $\eta_{i+1}$  is either an immediate face or coface of  $\eta_i$  then r must be odd and

$$|\{0 \le i \le r - 1 \mid \eta_{i+1} \prec \eta_i\}| = |\{0 \le i \le r - 1 \mid \eta_i \prec \eta_{i+1}\}| + 1.$$

Assume  $r \geq 3$ . Note that we cannot have  $\eta_i \prec \eta_{i+1} \prec \eta_{i+2}$  for it would contradict that  $W^2 = 0$ . So we must have  $\eta_i \prec \eta_{i+1} \succ \eta_{i+2}$  (with  $W(\eta_i) = \eta_{i+1}$ ) for all odd *i* and  $\eta_i \succ \eta_{i+1} \prec \eta_{i+2}$  (with  $W(\eta_{i+1}) = \eta_{i+2}$ ) for all even *i*. This gives rise to a non-stationary closed path of index dim $(\sigma)$ , contrary to the hypotheses. We conclude that r = 1 and thus  $W(\sigma) = \tau$ .

Suppose now that  $\tau \succ \sigma$  satisfies  $h(\tau) \leq h(\sigma)$ . If  $\tau \not\approx \sigma$  then adding  $\tau$  to a chain that realizes the height of  $\sigma$  gives us  $h(\tau) \geq h(\sigma) + 1$ , a contradiction. So  $\tau \not\sim \sigma$  and  $\tau = W(\sigma)$ is the only immediate co-face of  $\sigma$  satisfying  $h(\tau) \leq h(\sigma)$ . On the other hand, if  $\eta \prec \sigma$ is such that  $h(\eta) \geq h(\sigma)$  then  $W(\eta) = \sigma$  and  $\eta$  is the only immediate face of  $\sigma$  satisfying this by (W2). This proves that h fulfills (M1) and (M2).

Finally, it is straightforward to see that  $V_h = W$ .

As mentioned earlier, the function h is nothing but a simplified version of Forman's construction of the discrete Morse function produced in the proof of [3, Theorem 9.3]. Forman uses a more complex (inductive) argument in order to get some desirable extra properties for this function. Theorem 3 provides a simpler proof to the fact that W is a gradient field.

 $\mathbf{2}$ 

REMARK 4. It is straightforward to check that if  $\sigma \sim \eta \sim \varphi$  is an equivalence of *atomic* relations  $(\sigma, \varphi \neq \eta)$  then necessarily  $\sigma = \varphi$ .

The normalization of a Morse function. Let  $f : K \to \mathbb{R}$  be a discrete Morse function and let  $h_f : K \to \mathbb{Z}_{\geq 0}$  be the height function associated to  $V_f$ . We call  $h_f$  the normalization of f. This function is a natural representative of the equivalence class of f in the following way.

PROPOSITION 5. For every  $\sigma \in K$ ,  $h_f(\sigma) \leq g(\sigma)$  whenever  $g: K \to \mathbb{Z}_{\geq 0}$  is a combinatorial Morse function equivalent to f.

Proof. Suppose otherwise and let  $\sigma \in \{\eta \in K : g(\eta) < h_f(\eta)\}$  for which  $g(\sigma)$  is minimum (then  $h_f(\sigma) = r \ge 1$ ). Let C be a chain  $\sigma = \sigma_r \gg \sigma_{r-1} \gg \cdots \gg \sigma_0$  realizing  $h_f(\sigma)$  and note that  $h_f(\sigma_{r-1}) = h_f(\sigma) - 1$ . By Remark 4 and the maximality of C, any reduced chain of atomic relations realizing  $\sigma \gg \sigma_{r-1}$  (i.e. not containing repeated simplices) must be of the form  $\sigma \sim \eta > \eta' \sim \sigma_{r-1}$ , with  $\eta' \approx \eta$ . Hence  $\eta' \prec \eta$  and  $f(\eta') < f(\eta)$ .

By the equivalence  $h_f \sim f \sim g$  we have  $g(\eta') < g(\eta)$ . We claim that  $g(\eta) \leq g(\sigma)$ . By the equivalence  $f \sim g$  this is straightforward if  $\eta \succ \sigma$ , so we must discard the case  $\eta \prec \sigma$  (with  $f(\sigma) \leq f(\eta)$ ). There are two sub-cases for this case (which arise from the possibilities for  $\sigma_{r-1} \sim \eta'$ ):

- If  $\sigma_{r-1} \prec \eta'$  with  $f(\sigma_{r-1}) \ge f(\eta')$  then  $\sigma_{r-1} \prec \eta' \prec \eta \prec \sigma$ . Let  $\rho$  be the only other simplex aside  $\eta'$  that fulfills  $\sigma_{r-1} \prec \rho \prec \eta$ . Then the chain of atomic relations  $\sigma_{r-1} < \rho < \eta \sim \sigma$  contradicts the maximality of  $h_f(\sigma)$ .
- If  $\sigma_{r-1} \succ \eta'$  with  $f(\sigma_{r-1}) \leq f(\eta')$  then  $\sigma_{r-1} \succ \eta' \prec \eta \prec \sigma$ . Whether  $\sigma_{r-1}$  is or is not an immediate face of  $\sigma$ , we reach the contradictions to the maximality of  $h_f$  shown in Figure 2.



Figure 2. On the left. In the case  $\sigma_{r-1} \prec \sigma$ , the chain  $\sigma_{r-1} \prec \rho \prec \phi \prec \sigma$  contradicts the maximality of  $h_f(\sigma)$ . On the right. In the case that  $\sigma_{r-1}$  is not an immediate face of  $\sigma$ , the chain  $\sigma_{r-1} \sim \eta' \prec \rho \prec \sigma$  contradicts the maximality of  $h_f(\sigma)$ .

This proves the claim. Finally, by the minimality condition of  $\sigma$  we have

$$h_f(\sigma_{r-1}) = h_f(\eta') \le g(\eta') < g(\sigma) < h_f(\sigma)$$

from where  $h_f(\sigma) - h_f(\sigma_{r-1}) \ge 2$ , a contradiction.

COROLLARY 6. If f and g are equivalent Morse functions then  $h_f = h_q$ .

The condition in the statement of Proposition 5 is trivially seen to be an alternative definition for  $h_f$ . Thus, the normalization of f is the non-negative integer-valued Morse function equivalent to f taking the smallest possible values over all simplices. This justifies the term *normalized*.

We collect some basic properties of  $h_f$  in the following

LEMMA 7. With the notations as above, the function  $h_f$  satisfies: (1)  $h_f(\sigma) \ge \dim(\sigma)$  for all  $\sigma \in K$ . 

## N. A. CAPITELLI

(2)  $h_f(\sigma) = 0$  if and only if  $\sigma$  is a critical vertex. (3) If  $\tau = V_f(\sigma)$  then  $h_f(\sigma) = h_f(\tau)$ . (4) h takes all the integer values between 0 and  $\max_{\sigma \in K} \{h_f(\sigma)\}$ .

*Proof.* (1) and (2) follow from the fact that any simplex of positive dimension has at least two immediate faces, one of which must be a critical face. (3) is a direct consequence of the definition of the height function and (4) follows from Proposition 5.  $\Box$ 

**Computing**  $h_f$ . Finally, we present two algorithms for computing normalized functions: one from acyclic matchings over the Hasse diagram of the complex and one from the values of a given Morse function.

Recall that the Hasse diagram  $\mathcal{H}_K$  of a complex K is the digraph whose vertices are the simplices of K and whose directed edges  $\tau \to \sigma$  arise for every  $\tau \succ \sigma$ . The pairing of simplices defined by a gradient vector field V determines a matching  $\mathcal{M}_V$  in  $\mathcal{H}_K$ . If the arrows in this matching are reversed in  $\mathcal{H}_K$ , the resulting digraph  $\mathcal{H}_M$  is acyclic. On the other hand, an *acyclic matching*  $\mathcal{M}$  over  $\mathcal{H}_K$  determines a discrete gradient field  $V_{\mathcal{M}}$  on K where  $V_{\mathcal{M}}(\sigma) = \tau$  if and only if the edge  $\overline{\sigma\tau} \in \mathcal{M}$  (see e.g. [2]).

Algorithm 1 Normalization from an acyclic matching (SAGE)

In the following code, the argument D is the *dictionary of covering vertices* of the face poset: a vertex is assigned the list of vertices covering it. A typical entry is e.g. "2:[3,7,13]". Entries of maximal vertices/simplices appear e.g. as "30:[]". On the other hand, the argument M is the list of paired vertices given by the acyclic matching over D. A typical entry in M is e.g. "[6,27]", which is also a list.

```
def Normalization(D,M):
for i in range(len(M)):
    D[M[i][0]].remove(M[i][1])
    D[M[i][1]].append(M[i][0])
G=DiGraph(D)
V=len(G.vertices())
for j in range(V):
    n=0
    L=G.all_simple_paths(ending_vertices=[j])
    if L!=[]:
        values=set()
        for 1 in L:
            k=0
            l.reverse()
            for i in range(len(l)-1):
                if [1[i],1[i+1]] not in M:
                    k=k+1
            values.add(k)
        n=max(values)
        print 'Simplex', j, 'takes the value', n
```

The height function associated to an acyclic matching can be characterized in terms of the number of edges in simple paths not belonging to the matching.

4

PROPOSITION 8. Let V be a gradient field over K,  $\mathcal{M}$  the associated matching and let  $P_{\sigma}$  stand for the set of (directed) simple paths in  $\mathcal{H}_{\mathcal{M}}$  starting in  $\sigma$ . Then,

$$h(\sigma) = \max_{p \in P_{\sigma}} |\{edges \ in \ p \ not \ in \ \mathcal{M}_V\}|.$$

*Proof.* A simple path in  $\mathcal{H}_{\mathcal{M}}$  starting in  $\sigma$  is a sequence  $\sigma = \sigma_n \to \sigma_{n-1} \to \cdots \to \sigma_0$  where either

- (1)  $\sigma_{i+1} \succ \sigma_i$  and  $V(\sigma_i) \neq V(\sigma_{i+1})$  or
- (2)  $\sigma_{i+1} \prec \sigma_i$  and  $V(\sigma_{i+1}) = \sigma_i$ .

Hence, one such path gives rise to a sequence  $\sigma_n > \sigma_{n-1} > \cdots > \sigma_0$  where  $\sigma_i \sim \sigma_{i+1}$  only for the *i*'s corresponding to simplices in the case (2). By removing these arrows we get a chain with no equivalencies. Since these arrows are the "inverted arrows" in  $\mathcal{M}$ , the result follows.

Algorithm 1 is a SAGE implementation of the algorithm implicit in Proposition 8.

For the second algorithm we rely on the alternative characterization of  $h_f$  given in Proposition 5. Note that one can easily scale and perturb a Morse function f to get another one with the same gradient and such that  $\min_{\sigma \in K} \{f(\sigma)\} = 0$  and  $f(K) \subset \mathbb{Z}_{\geq 0}$ . Algorithm 2 computes the normalization of one such function.

Algorithm 2 Normalization from a Morse function f

**Require:**  $f: K \to \mathbb{Z}_{>0}, \min\{f(\sigma)\} = 0.$ **Ensure:**  $h_f$ 1: **procedure** NORMALIZE(f)Order  $K = \{\sigma_1, \dots, \sigma_r\}$  such that  $\begin{cases} f(\sigma_i) < f(\sigma_j) & i < j \\ \dim(\sigma_i) \ge \dim(\sigma_j) & f(\sigma_i) = f(\sigma_j) \end{cases}$ 2:  $f_0 \leftarrow f$ 3: for  $0 \le s < r$  do 4:  $\begin{aligned} & \int_{s}^{0} \leftarrow f_{s} \\ & \mathbf{for} \ 0 < k \leq r \mathbf{do} \\ & \mathbf{for} \ 0 < k \leq r \mathbf{do} \end{aligned}$ 5:6:  $f_{s}^{k}(\sigma) = f_{s}^{k-1}(\sigma) \text{ if } \sigma \neq \sigma_{k}$   $f_{s}^{k}(\sigma_{k}) = \begin{cases} \max\{f_{s}^{k-1}(\eta) \mid \eta \prec \sigma_{k}\} + 1 & \sigma_{k} \text{ is critical for } f_{s}^{k-1} \\ f_{s}^{k-1}(\tau) & f_{s}^{k-1}(\sigma_{k}) \geq f_{s}^{k-1}(\tau) \\ \max\{f_{s}^{k-1}(\eta') \mid \eta' \prec \sigma_{k}, \ \eta' \neq \eta\} + 1 & f_{s}^{k-1}(\eta) \geq f_{s}^{k-1}(\sigma_{k}) \end{cases}$ 7: 8: end for 9:  $f_{s+1} \leftarrow f_s^r$ 10: end for 11:  $h_f \leftarrow f_r$ 12:13: end procedure

**PROPOSITION 9.** Algorithm 2 computes the normalization of f.

Proof. First we show by induction that  $f_s^k$  is a Morse function with gradient  $V_f$  for all s, k. Since the loop corresponding to a given s and k only alters the value of  $\sigma_k$ , it suffices to show that the order relations between the values of  $\sigma_k$  and its immediate faces and cofaces are preserved in the transition  $f_s^{k-1} \to f_s^k$  (note that this comprises the transition between subindexes s as well). For simplicity, let us write  $\sigma_k = \sigma$ ,  $f_s^k = f_k$  and let  $\mathcal{O}$  stand for the order the algorithm introduces on the simplices of K. We analyze the possible cases of  $\sigma$ : (a)  $\sigma$  is critical for  $f_{k-1}$ , (b)  $f_{k-1}(\sigma) \geq f_{k-1}(\tau)$  for some  $\tau \succ \sigma$  and (c)  $f_{k-1}(\eta) \geq f_{k-1}(\sigma)$  for some  $\eta \prec \sigma$ .

In the case (a) a direct computation shows that for  $\nu \prec \sigma \prec \rho$  we have

## N. A. CAPITELLI

- $f_k(\sigma) = \max\{f_{k-1}(\eta) \mid \eta \prec \sigma\} + 1 \le f_{k-1}(\sigma) < f_{k-1}(\rho) = f_k(\rho) \text{ and }$
- $f_k(\nu) = f_{k-1}(\nu) < \max\{f_{k-1}(\eta) \mid \eta \prec \sigma\} + 1 = f_k(\sigma).$

Suppose we are in the conditions of case (b). On one hand,  $f_k(\sigma) = f_{k-1}(\tau) = f_k(\tau)$ . On the other hand, since  $f_{k-1}$  is Morse, if  $\sigma \prec \rho \neq \tau$  then

$$f_k(\sigma) = f_{k-1}(\tau) \le f_{k-1}(\sigma) < f_{k-1}(\rho) = f_k(\rho)$$

Finally, for  $\nu \prec \sigma$  let  $\sigma' \prec \tau$  be such that  $\nu = \sigma \cap \sigma'$ . Since  $f_{k-1}$  is Morse then  $f_{k-1}(\sigma') < f_{k-1}(\tau)$ . In particular  $\sigma'$  appears earlier than  $\sigma$  in  $\mathcal{O}$ ; so either  $f_{k-1}(\nu) < f_{k-1}(\sigma')$  or  $f_{k-1}(\nu) = f_{k-1}(\sigma')$  (if  $f(\nu) \ge f(\sigma')$ ). In any case,

$$f_k(\nu) = f_{k-1}(\nu) < f_{k-1}(\tau) = f_k(\sigma).$$

Let us now deal with case (c). If  $\rho' \succ \sigma$  then

$$f_k(\sigma) = \max\{f_{k-1}(\eta') \mid \eta' \prec \sigma, \, \eta' \neq \eta\} + 1 < f_{k-1}(\sigma) < f_{k-1}(\rho') = f_k(\rho').$$

On the other hand, if  $\sigma \succ \nu \neq \eta$  then

$$f_k(\sigma) = \max\{f_{k-1}(\eta') \mid \eta' \prec \sigma, \ \eta' \neq \eta\} + 1 > f_{k-1}(\nu) = f_k(\nu).$$

Finally, using that  $f_{k-1}$  is Morse,

$$f_k(\sigma) = \max\{f_{k-1}(\eta') \mid \eta' \prec \sigma, \, \eta' \neq \eta\} + 1 \le f_{k-1}(\sigma) \le f_{k-1}(\eta) = f_k(\eta).$$

This proves that  $f_r$  is a Morse function with the same gradient field as f. In order to show that  $f_r = h_f$  we prove the minimality of  $f_r$  according to Proposition 5. Assume there exists a Morse function  $g: K \to \mathbb{Z}_{\geq 0}$  with  $V_g = V_{f_r}$  contradicting this minimality and let  $\sigma = \sigma_k$  be the earliest simplex in the order  $\mathcal{O}$  such that  $g(\sigma) < f_r(\sigma)$ . If  $\sigma$  is in case (a) then all the immediate faces of  $\sigma$  appear earlier in  $\mathcal{O}$  so

$$g(\sigma) < f_r(\sigma) = f_k(\sigma) = \max\{f_{k-1}(\eta) \mid \eta \prec \sigma\} + 1$$
  
= 
$$\max\{f_r(\eta) \mid \eta \prec \sigma\} + 1$$
  
$$\leq \max\{g(\eta) \mid \eta \prec \sigma\} + 1.$$

Thus if  $\nu \prec \sigma$  realizes this last maximum then  $g(\sigma) \leq g(\nu)$ , contrary to the hypothesis of (a). If  $\sigma$  is in the conditions of (b) then  $\tau$  appears earlier than  $\sigma$  in  $\mathcal{O}$  and we reach the following contradiction:

$$f_r(\tau) \le g(\tau) \le g(\sigma) < f_r(\sigma) = f_k(\sigma) = f_{k-1}(\tau) = f_r(\tau).$$

Finally, if we are in case (c), let  $\nu \prec \sigma$  be such that  $f_k(\nu) = \max\{f_{k-1}(\eta') \mid \eta' \prec \sigma, \eta' \neq \eta\}$ . Since  $\nu$  appears earlier than  $\sigma$  in  $\mathcal{O}$  then

$$g(\nu) < g(\sigma) < f_r(\sigma) = f_k(\sigma_k) = f_{k-1}(\nu) + 1 = f_r(\nu) + 1$$

which contradicts the choice of  $\sigma$  as a minimum.

## References

- R. Ayala, L. M. Fernández and J. A. Vilches. *Characterizing equivalent discrete Morse functions*. Bull. Braz. Math. Soc 40 (2009), 225-235.
- [2] M. K. Chari. On discrete Morse functions and combinatorial decompositions. Discrete Math. 217 (2000), No. 1-3, 101-113.
- [3] R. Forman. Morse theory for cell complexes. Adv. Math. 134 (1998), No. 1, 90-145.

6