Esseen–Rozovskii type estimates for the rate of convergence in the Lindeberg theorem^{*}

Ruslan Gabdullin¹, Vladimir Makarenko¹, and Irina Shevtsova^{1,2}

Abstract

We present structural improvements of Esseen's (1969) and Rozovskii's (1974) estimates for the rate of convergence in the Lindeberg theorem and also compute the appearing absolute constants. We introduce the asymptotically exact constants in the constructed inequalities and obtain upper bounds for them. We analyze the values of Esseen's, Rozovskii's, and Lyapunov's fractions, compare them pairwise and provide some extremal distributions. As an auxiliary statement, we prove a sharp inequality for the quadratic tails of an arbitrary distribution (with finite second order moment) and its convolutional symmetrization.

1 Introduction

Let X_1, X_2, \ldots, X_n be independent random variables (r.v.'s) on a certain probability space $(\Omega, \mathcal{A}, \mathsf{P})$ with distribution functions (d.f.'s) $F_k(x) := \mathsf{P}(X_k < x), k = 1, \ldots, n$, and such that

$$\mathsf{E}X_k = 0, \quad \sigma_k^2 \coloneqq \mathsf{E}X_k^2 < \infty, \quad B_n^2 \coloneqq \sum_{k=1}^n \sigma_k^2 > 0. \tag{1}$$

Denote

$$S_n = \sum_{k=1}^n X_k, \quad \overline{F}_n(x) = \mathsf{P}(S_n < xB_n), \quad \Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt,$$
$$\Delta_n = \Delta_n(F_1, \dots, F_n) \coloneqq \sup_{x \in \mathbb{R}} \left| \overline{F}_n(x) - \Phi(x) \right|, \quad n \in \mathbb{N},$$

$$\sigma_k^2(z) = \mathsf{E}X_k^2 \mathbf{1}(|X_k| \ge z), \quad L_n(z) \coloneqq \frac{1}{B_n^2} \sum_{k=1}^n \sigma_k^2(zB_n) = \frac{1}{B_n^2} \sum_{k=1}^n \mathsf{E}X_k^2 \mathbf{1}(|X_k| \ge zB_n), \quad z \ge 0,$$

so that $\sup_{z\geq 0} \sigma_k^2(z) = \sigma_k^2(0) = \sigma_k^2$ for all k = 1, ..., n and $L_n(0) = 1$. In 1922 Lindeberg [20] proved that $\Delta_n \to 0$, if the Lindeberg fraction L_n satisfies the condition

$$\lim_{n \to \infty} L_n(z) = 0 \quad \text{for every } z > 0. \tag{L}$$

In 1935 Feller [9] completed Lindeberg's theorem by proving the necessity of condition (L) for the CLT to hold, if the random summands are asymptotically negligible in the sense that

$$\lim_{n \to \infty} B_n^{-2} \max_{1 \le k \le n} \sigma_k^2 = 0.$$
 (F)

Condition (F) is called the *Feller condition*.

^{*}Research supported by the Russian Foundation for Basic Research (projects 15-07-02984-a and 16-31-60110-mol_a_dk) and by the grant of the President of Russia No. MD-2116.2017.1.

¹Faculty of Computational Mathematics and Cybernetics, Lomonosov Moscow State University, Moscow, Russia

²Institute of Informatics Problems of Federal Research Center "Computer Science and Control," Russian Academy of Sciences, Moscow, Russia; e-mail: ishevtsova@cs.msu.ru

Lindeberg's theorem yields the celebrated Lyapunov theorem [19], according to which $\Delta_n \to 0$, if the Lyapunov fraction defined below tends to zero:

$$L_{2+\delta,n} \coloneqq \frac{1}{B_n^{2+\delta}} \sum_{k=1}^n \mathsf{E} |X_k|^{2+\delta} \to 0, \quad n \to \infty, \text{ for certain } \delta > 0.$$

In the case of independent and identically distributed (i.i.d.) random summands the Lyapunov fraction

$$L_{2+\delta,n} = \frac{\mathsf{E}|X_1|^{2+\delta}}{\sigma_1^{2+\delta} n^{\delta/2}}$$

is of order $O(n^{-\delta/2})$ as $n \to \infty$ for every fixed distribution F_1 of the random summands.

The numerical demonstration to the Lyapunov theorem is given by the inequality

$$\Delta_n \le C_{BE}(\delta) \cdot L_{2+\delta,n}, \quad 0 < \delta \le 1, \tag{2}$$

 $C_{BE}(\delta)$ being some absolute constants for every $\delta \in (0, 1]$, which was proved by Lyapunov himself in the same paper [19] for $0 < \delta < 1$, and, 40 years later, independently by Berry [3] in the i.i.d. case and Esseen [6] in the general situation for $\delta = 1$. Inequality (2) in the i.i.d. case with $\delta = 1$ takes the form

$$\Delta_n \le \frac{C_{BE}(1)}{\sqrt{n}} \cdot \frac{\mathsf{E}|X_1|^3}{\sigma_1^3} \tag{3}$$

and establishes the exact order $O(n^{-1/2})$ of the rate of convergence in the CLT for distributions with finite thirdorder moments. The history of evaluation of the constants $C_{BE}(\delta)$, especially, for $\delta = 1$, as well as the discussion of the structural improvements of the Berry-Esseen inequality is worthy of a particular narration for which we refer to the papers [15, 18, 33, 34]. In these papers some structural improvements of inequality (2) are also presented.

The numerical demonstration to the Lindeberg theorem was given in 1966 by Osipov's inequality [22] in the form

$$\Delta_n \le C(\Lambda_n(\varepsilon) + L_n(\varepsilon)) \quad \text{for every } \varepsilon > 0, \tag{4}$$

where

$$\Lambda_n(\varepsilon) \coloneqq \frac{1}{B_n^3} \sum_{k=1}^n \mathsf{E} |X_k|^3 \mathbf{1} (|X_k| < \varepsilon B_n), \quad \varepsilon > 0,$$

and C is an absolute constant. The quantity $\Lambda_n(\varepsilon) + L_n(\varepsilon)$ is called the Osipov fraction with the parameter $\varepsilon > 0$. Since $\Lambda_n(\varepsilon) \le \varepsilon B_n^{-2} \sum_{k=1}^n \mathsf{E} X_k^2 \mathbf{1}(|X_k| \le \varepsilon B_n) \le \varepsilon$, inequality (4) yields the bound

$$\Delta_n \leq C(\varepsilon + L_n(\varepsilon)) \quad \text{for every } \varepsilon > 0,$$

and hence, the Lindeberg theorem. Thus, in the case of asymptotically negligible random summands satisfying condition (F), by the Feller theorem, the left-hand and right-hand sides of (4) either tend or do not tend to zero simultaneously. In other words, inequality (4) relates the *criteria* of convergence with the *rate* of convergence in the classical CLT and hence, according to Zolotarev's terminology [36], it can be called a *natural* convergence rate estimate.

It is easy to see that inequality (4) with some $\varepsilon > 0$ yields the Berry–Esseen inequality (2) for every $\delta \in (0, 1]$, since

$$\Lambda_n(\varepsilon) + L_n(\varepsilon) = \sum_{k=1}^n \left(\frac{\mathsf{E}|X_k|^3 \mathbf{1}(|X_k| < \varepsilon B_n)}{B_n^3} + \frac{\mathsf{E}X_k^2 \mathbf{1}(|X_k| \ge \varepsilon B_n)}{B_n^2} \right) \le \sum_{k=1}^n \frac{\mathsf{E}|X_k|^{2+\delta}}{B_n^{2+\delta}} = L_{2+\delta,n}.$$

However, inequality (4) with $\varepsilon = 1$

$$\Delta_n \le C(\Lambda_n(1) + L_n(1)) = \frac{C}{B_n^3} \sum_{k=1}^n \left(\mathsf{E}|X_k|^3 \mathbf{1}(|X_k| < B_n) + B_n \mathsf{E}X_k^2 \mathbf{1}(|X_k| \ge B_n) \right) = \frac{C}{B_n^2} \sum_{k=1}^n \mathsf{E}X_k^2 \min\left\{ 1, \frac{|X_k|}{B_n} \right\} = C \int_0^1 L_n(z) dz, \quad (5)$$

is of great interest, because $\varepsilon = 1$ minimizes the right-hand side of (4). Indeed, to find the minimizer one can observe, following the outline of the reasoning in [25], that for every $\varepsilon > 0$

$$\Lambda_n(\varepsilon) + L_n(\varepsilon) = \sum_{k=1}^n \left[\mathsf{E} \Big| \frac{X_k}{B_n} \Big|^3 \mathbf{1} \Big(\Big| \frac{X_k}{B_n} \Big| < \varepsilon \Big) + \mathsf{E} \Big(\frac{X_k}{B_n} \Big)^2 \mathbf{1} \Big(\Big| \frac{X_k}{B_n} \Big| \ge \varepsilon \Big) \right]$$

and for every r.v. $X \coloneqq X_k/B_n$, k = 1..., n, with A being arbitrary Borel subset of \mathbb{R} , we have

$$\mathsf{E}|X|^{3}\mathbf{1}(|X|<1) + \mathsf{E}X^{2}\mathbf{1}(|X|\geq 1) = \mathsf{E}X^{2}\min\{1,|X|\}(\mathbf{1}(X\in A) + \mathbf{1}(X\notin A)) \leq \mathsf{E}X^{2}\max\{1,|X|\}(\mathbf{1}(X\in A) + \mathbf{1}(X\notin A)) \leq \mathsf{E}X^{2}\min\{1,|X|\}(\mathbf{1}(X\in A) + \mathbf{1}(X\notin A)) \leq \mathsf{E}X^{2}\max\{1,|X|\}(\mathbf{1}(X\in A) + \mathbf{1}(X\otimes A)) \leq \mathsf{E}X^{2}\max\{1,|X|\}(\mathbf{1}(X\otimes A) + \mathbf{1}(X\otimes A)) \leq \mathsf{E}X^{2}\max\{1,|X|\}(\mathbf{1}(X\otimes A) + \mathbf{1}(X\otimes A)) \leq \mathsf{E}X^{2}\max\{1,|X|\}(\mathbf{1}(X\otimes A) + \mathbf{1}(X\otimes A)) \leq \mathsf{E}X^{2}\max\{1,|X$$

 $\leq \mathsf{E}|X|^3 \mathbf{1}(X \in A) + \mathsf{E}X^2 \mathbf{1}(X \notin A)$

(here the optimality of the set A = [-1, 1] was first noted, probably, by Loh [21] whose hardly available thesis we cite by [2]; however, in [25], this fact was proved independently). Now the claim follows by taking $A \coloneqq [-\varepsilon, \varepsilon]$.

Inequality (5) has an interesting history. First of all we note that it is a trivial corollary to earlier and formally more general results of Katz [12] and Petrov [26], but this connection remained unnoticed for a long time, even by Petrov himself (for example, see [27, Ch. 5, § 3, Theorems 5 and 8]). Furthermore, inequality (5) was re-proved formally in a more general, but in fact, in an equivalent form, by Feller [10] with C = 6 and by Paditz [23,24] in the i.i.d. case with C = 4.77. Two years later Paditz [25] announced a sharper bound $C \leq 3.51$ without a complete proof. The works of Feller and Paditz did not refer to the above mentioned work of Osipov. Inequality (4) was also re-proved by Barbour and Hall [2] who cited Feller's result, but applied the new Tikhomirov–Stein method and obtained a rougher bound $C \leq 18$. Chen and Shao [5] who cited only Feller's work re-proved (4) with C = 4.1. Finally, Korolev and his disciples successively improved the upper bounds for C to 2.011 in [16, 17] and to 1.87 in [14].

Thus, inequalities (4) and (5) attracted much attention. However, as far back as in 1969 Esseen [8] proved the bound

$$\Delta_n \le \frac{C_1}{B_n^3} \sum_{k=1}^n \sup_{z>0} \left\{ \left| \mathsf{E} X_k^3 \mathbf{1}(|X_k| < z) \right| + z \mathsf{E} X_k^2 \mathbf{1}(|X_k| \ge z) \right\} = C_1 \cdot \widehat{L}^3_{\mathsf{E},n}(\infty), \tag{6}$$

where

$$\widehat{L}^3_{\scriptscriptstyle \mathrm{E},n}(\varepsilon) \coloneqq \frac{1}{B_n^3} \sum_{k=1}^n \sup_{0 < z < \varepsilon B_n} \left\{ |\mu_k(z)| + z \sigma_k^2(z) \right\}, \quad \mu_k(z) \coloneqq \mathsf{E} X_k^3 \mathbf{1}(|X_k| < z), \quad \varepsilon > 0, \ z \ge 0,$$

and, as its corollary, by the use of the traditional truncation techniques, deduced that

$$\Delta_n \le C_2 \cdot \widehat{L}^3_{\mathrm{E},n}(1),\tag{7}$$

with C_1 and C_2 being some absolute constants. Obviously, $\hat{L}^3_{\text{E},n}(1) \leq \hat{L}^3_{\text{E},n}(\infty)$. Moreover, due to the left-continuity of the functions $\mu_k(z)$, $\sigma_k^2(z)$, $k = 1, \ldots, n$, for z > 0, the least upper bound in the definition of $\hat{L}^3_{\text{E},n}(\varepsilon)$ can be replaced by the one over the set $z \in (0, \varepsilon B_n]$.

Comparing inequalities (6) and (7) with (5) we observe, first, that the fractions $\hat{L}^3_{\text{E},n}(\infty)$ and $\hat{L}^3_{\text{E},n}(1)$ here depend not on the *absolute*, but on the *algebraic* third-order truncated moments, which may vanish, for example, in the case of symmetrically distributed random summands. Second, Esseen's fraction $\hat{L}^3_{\text{E},n}(1)$ never exceeds the Osipov fraction $\Lambda_n(\varepsilon) + L_n(\varepsilon)$ with $\varepsilon = 1$ and hence with arbitrary $\varepsilon > 0$. Indeed, for every random summand $X \coloneqq X_k, k = 1, \ldots, n$, we have

$$\left|\mathsf{E} X^{3} \mathbf{1}(|X| < z)\right| + z \mathsf{E} X^{2} \mathbf{1}(|X| \ge z) \le \mathsf{E} |X|^{3} \mathbf{1}(|X| < z) + z \mathsf{E} X^{2} \mathbf{1}(|X| \ge z) \eqqcolon h(z),$$

with h(z) being monotonically increasing with respect to $z \ge 0$, due to

$$h(u) - h(v) = \mathsf{E}\left(|X|^3 - vX^2\right) \mathbf{1}(v \le |X| < u) + (u - v)\mathsf{E}X^2 \mathbf{1}(|X| \ge u) \ge 0, \quad u \ge v \ge 0,$$

and, hence, writing $h_k(z)$ for the analogous function of the distribution of X_k , k = 1, ..., n, we get

$$\widehat{L}^{3}_{\text{E},n}(1) \leq \frac{1}{B_{n}^{3}} \sum_{k=1}^{n} \sup_{0 < z \leq B_{n}} h_{k}(z) = \frac{1}{B_{n}^{3}} \sum_{k=1}^{n} h_{k}(B_{n}) = \Lambda_{n}(1) + L_{n}(1) = \inf_{\varepsilon > 0} \left\{ \Lambda_{n}(\varepsilon) + L_{n}(\varepsilon) \right\}.$$

This fact implies, in turn, that $C \leq C_2$, in particular, inequality (7) is also a natural convergence rate estimate in the Lindeberg theorem.

Third, Esseen's inequality (7) not only relates the criteria of convergence with the rate of convergence as Osipov's inequality does, but also provides a numerical demonstration to the *criteria of the rate of convergence* to be of the traditional order $\mathcal{O}(n^{-1/2})$, $n \to \infty$, provided by the Berry–Esseen inequality for distributions with finite third-order moments in the i.i.d. case. In fact, the condition on the finiteness of the third-order moments can be relaxed. Namely, in 1966 Ibragimov [11] proved that in the i.i.d. case $\Delta_n = \mathcal{O}(n^{-1/2})$ as $n \to \infty$ if and only if

$$\mu_1(z) = \mathcal{O}(1), \quad z\sigma_1^2(z) = \mathcal{O}(1), \quad z \to \infty.$$

It is easy to see that Ibragimov's condition is weaker than the condition $\mathsf{E}|X_1|^3 < \infty$, and Esseen's inequality (7) in the i.i.d. case

$$\Delta_n \le \frac{C_2}{\sigma_1^3 \sqrt{n}} \sup_{0 < z \le \sigma_1 \sqrt{n}} \left\{ |\mu_1(z)| + z \sigma_1^2(z) \right\}$$

trivially yields the "if" part of Ibragimov's criteria.

In 1974 Rozovskii [30] generalized Ibragimov's theorem and proved another estimate involving algebraic truncated third-order moments

$$M_n(z) \coloneqq \frac{1}{B_n^3} \sum_{k=1}^n \mu_k(zB_n) = \frac{1}{B_n^3} \sum_{k=1}^n \mathsf{E} X_k^3 \mathbf{1}(|X_k| < zB_n), \quad z > 0,$$

(8)

in the form

where C_3 is an absolute constant and

$$L_{\mathbf{R},n}^{3} \coloneqq |M_{n}(1)| + \sup_{0 < z \le 1} zL_{n}(z) = \frac{1}{B_{n}^{3}} \left(\left| \sum_{k=1}^{n} \mu_{k}(B_{n}) \right| + \sup_{0 < z \le B_{n}} z\sum_{k=1}^{n} \sigma_{k}^{2}(z) \right).$$

 $\Delta_n \le C_3 \cdot L^3_{\mathrm{B},n},$

At first glance, Rozovskii's inequality (8) is more favorable than Esseen's inequalities (6) and (7). Indeed, the right-hand side of (8) is always finite, while the right-hand side of (6) may be infinite. In (8) the first term may vanish not only in the symmetric, but also in the non-symmetric case, for example, for even n if the appearing truncated third-order moments $\mu_k(B_n)$, $k = 1, \ldots, n$, have the same absolute value, but alternating signs. One more advantage of inequality (8) over (6) and (7) is that the values of $\mu_k(z)$ are used in (8) only in one and the same point $z = B_n$ for all $k = 1, \ldots, n$, while the right-hand sides of (6) and (7) require the information on $\mu_k(z)$ in every point of the interval $z \in (0, B_n]$. However, a deeper analysis (see Theorem 6 (ii) below) shows that Rozovskii's fraction $L^3_{\mathbf{R},n}$ may take greater values than each of Esseen's fractions $\hat{L}^3_{\mathbf{E},n}(1), \hat{L}^3_{\mathbf{E},n}(\infty)$ and even greater than Lyapunov's fraction $L_{3,n}$, while Esseen's fractions always satisfy

$$\widehat{L}^{3}_{\mathrm{E},n}(\infty) \leq L_{3,n}, \qquad \widehat{L}^{3}_{\mathrm{E},n}(1) \leq \Lambda_{n}(1) + L_{n}(1) \leq L_{2+\delta,n}, \quad \delta \in (0,1].$$
(9)

Thus the choice between inequalities (7) and (8) depends not only on the concrete values of the fractions $\hat{L}_{\text{E},n}^3$ and $L_{\text{R},n}^3$, but also on the values of the appearing absolute constants C_2 and C_3 . However, the values of these constants, as well as of C_1 , remain unknown.

In the present paper, we compute upper bounds for the constants C_1, C_2 , and C_3 . Moreover, we prove new natural convergence rate estimates in the Lindeberg–Feller theorem generalizing Esseen's (6), (7) and Rozovskii's (8) inequalities, and provide an explicit analytical representation with an algorithm of evaluation of the appearing constants. Namely, we introduce a *truncation* parameter $\varepsilon > 0$ and a *balancing* parameter $\gamma > 0$ and denote

$$\begin{split} L^3_{\scriptscriptstyle \mathrm{E},n}(\varepsilon,\gamma) &\coloneqq \sup_{0 < z \le \varepsilon} \left\{ \gamma \left| M_n(z) \right| + zL_n(z) \right\} = \frac{1}{B_n^3} \sup_{0 < z \le \varepsilon B_n} \left\{ \gamma \left| \sum_{k=1}^n \mu_k(z) \right| + z \sum_{k=1}^n \sigma_k^2(z) \right\}, \\ L^3_{\scriptscriptstyle \mathrm{R},n}(\varepsilon,\gamma) &\coloneqq \gamma \left| M_n(\varepsilon) \right| + \sup_{0 < z \le \varepsilon} zL_n(z) = \frac{1}{B_n^3} \left\{ \gamma \left| \sum_{k=1}^n \mu_k(\varepsilon B_n) \right| + \sup_{0 < z \le \varepsilon B_n} z \sum_{k=1}^n \sigma_k^2(z) \right\}, \end{split}$$

where the least upper bounds with respect to $0 < z \leq \ldots$ can be replaced by those over the open sets $0 < z < \ldots$, due to the left-continuity of $\mu_k(\cdot)$ and $\sigma_k^2(\cdot)$. It is easy to see that

$$L^{3}_{\mathrm{E},n}(\varepsilon,1) = \sup_{0 < z \le \varepsilon} \{ |M_{n}(z)| + zL_{n}(z) \} \le \sup_{\substack{0 < z \le \varepsilon \\ A}} \{ \Lambda_{n}(z) + zL_{n}(z) \} = \Lambda_{n}(\varepsilon) + \varepsilon L_{n}(\varepsilon),$$

$$\sup_{\gamma \in (0,1]} L^3_{\mathbb{E},n}(\varepsilon,\gamma) \leq \frac{1}{B^3_n} \sum_{k=1}^n \sup_{0 < z \leq \varepsilon B_n} \left[\mathsf{E}|X_k|^3 \mathbf{1}(|X_k| < z) + z \mathsf{E} X^2_k \mathbf{1}(|X_k| \geq z) \right]$$
$$\leq \frac{1}{B^3_n} \sum_{k=1}^n \sup_{0 < z \leq \varepsilon B_n} z^{1-\delta} \mathsf{E}|X_k|^{2+\delta} = \varepsilon^{1-\delta} L_{2+\delta,n} \quad \text{for every} \quad \varepsilon > 0, \ \delta \in (0,1], \quad (10)$$

moreover, $L^3_{\text{R},n}(1,1) = L^3_{\text{R},n}, L^3_{\text{E},n}(1,1) \leq \hat{L}^3_{\text{E},n}(1)$ with equality sign, for example, in the i.i.d. case, and $L^3_{\text{E},n}(\varepsilon,\gamma) = L^3_{\text{R},n}(\varepsilon,\gamma)$ in the symmetric case.

Theorem 1. Under the above assumptions, for every $\gamma > 0$

$$\Delta_n \le C_{\scriptscriptstyle \rm E}(\varepsilon,\gamma) \cdot L^3_{\scriptscriptstyle \rm E,n}(\varepsilon,\gamma), \quad \varepsilon \in (0,\infty], \tag{11}$$

$$\Delta_n \le C_{\mathsf{R}}(\varepsilon, \gamma) \cdot L^3_{\mathsf{R},n}(\varepsilon, \gamma), \quad \varepsilon \in (0, \infty), \tag{12}$$

where $C_{\rm E}(\varepsilon,\gamma)$ and $C_{\rm R}(\varepsilon,\gamma)$ depend only on the arguments in the brackets, take finite values for every $\gamma > 0$ and ε specified above and can be computed for every ε,γ under consideration by an algorithm provided below in the proof. Both $C_{\rm E}(\varepsilon,\gamma)$ and $C_{\rm R}(\varepsilon,\gamma)$ are monotonically decreasing with respect to $\gamma > 0$, $C_{\rm E}(\varepsilon,\gamma)$ is also monotonically decreasing with respect to $\varepsilon > 0$. In particular,

$$\begin{split} C_{\rm E}(\infty,1) &\leq \max\left\{C_{\rm E}(\infty,0.97), \ C_{\rm E}(4.35,1)\right\} \leq 2.66, \quad C_{\rm E}(1,1) \leq C_{\rm E}(1,0.72) \leq 2.73, \\ C_{\rm E}(\infty,\infty) &\leq \max\left\{C_{\rm E}(\infty,1.43), \ C_{\rm E}(4,1.62), \ C_{\rm E}(2.74,3), \ C_{\rm E}(2.56,\infty)\right\} \leq 2.65, \\ C_{\rm R}(1,1) &\leq \sup_{\gamma \geq \gamma_*} C_{\rm R}(1,\gamma) \leq 2.73, \quad \inf_{\varepsilon > 0, \gamma > 0} C_{\rm R}(\varepsilon,\gamma) \leq \inf_{\varepsilon > 0} \sup_{\gamma \geq \gamma_*} C_{\rm R}(\varepsilon,\gamma) \leq C_{\rm R}(2.12,\gamma_*) \leq 2.66, \\ \sup_{\gamma \geq 0.4} \max\left\{C_{\rm R}(2.63,\gamma), \ C_{\rm R}(1.76,\gamma)\right\} \leq 2.70, \quad \sup_{\gamma \geq 0.2} \max\left\{C_{\rm R}(5.39,\gamma), \ C_{\rm R}(1.21,\gamma)\right\} \leq 2.87, \end{split}$$

where

$$\gamma_* \coloneqq \frac{1}{\sqrt{6\varkappa}} = 0.5599\dots, \quad \varkappa \coloneqq x^{-2}\sqrt{(\cos x - 1 + x^2/2)^2 + (\sin x - x)^2} \Big|_{x=x_0} = 0.531551\dots,$$

and $x_0 = 5.487414...$ is the unique root of the equation $8(\cos x - 1) + 8x \sin x - 4x^2 \cos x - x^3 \sin x = 0$ on the interval $(\pi, 2\pi)$.

Remark 1. Within the method used: (i) $C_{\rm R}(1,\gamma)$ does not depend on γ for $\gamma \geq \gamma_*$; (ii) further increase of $\gamma \geq 0.73$ does not reduce the constructed upper bound 2.73 for $C_{\rm E}(1,\gamma)$ by more than 0.01; (iii) the same concerns the presented upper bounds for $C_{\rm E}(\infty, 1)$ and $C_{\rm E}(\infty, \infty)$.

Remark 2. Within the method used, $C_{\text{E}}(1,1) = C_{\text{R}}(1,1)$, due to Remark 9.

The plot of the level curve $\gamma = \gamma(\varepsilon)$ delivering the constant value 2.65 to $C_{\text{E}}(\varepsilon, \gamma)$ is given on Fig. 2 (right). The plot of the function $C_{\text{R}}(\varepsilon, \gamma_*)$ constructed in the proof is given on Fig. 3 (right).

Theorem 1 trivially yields

Corollary 1. The constants C_1 , C_2 , and C_3 in inequalities (6), (7), and (8) satisfy

$$C_1 \le C_{\rm E}(\infty, 1) \le 2.66, \quad C_2 \le C_{\rm E}(1, 1) \le 2.73, \quad C_3 \le C_{\rm R}(1, 1) \le 2.73.$$

The upper bounds for the constants C_1, C_2, C_3 presented in Corollary 1 are slightly greater than the best known upper bound 1.87, obtained in [14], for the absolute constant C (which is no greater than C_2) in Osipov's inequality (5). In Theorem 6 (iv) of Section 5 we provide various examples of symmetric and non-symmetric distributions of the random summands X_1, \ldots, X_n for which the right-hand sides of both inequalities (11) and (12) with $\varepsilon = \gamma = 1$, $C_{\rm E}(1,1) = C_{\rm R}(1,1) = 2.73$ are strictly less than the right-hand side of the Osipov inequality (5) with C = 1.87.

Similarly to Kolmogorov [13], where the classical Berry–Esseen inequality was discussed, we also introduce here the so-called *asymptotically exact constants* in (11), (12)

$$C_{\rm E}^*(\varepsilon,\gamma) \coloneqq \limsup_{\ell \to 0} \sup_{n,F_1,\dots,F_n} \left\{ \frac{\Delta_n(F_1,\dots,F_n)}{\ell} \colon L^3_{{\rm E},n}(\varepsilon,\gamma) = \ell \right\}, \quad \varepsilon, \ \gamma > 0, \tag{13}$$

$$C_{\mathbf{R}}^{*}(\varepsilon,\gamma) \coloneqq \limsup_{\ell \to 0} \sup_{n,F_{1},\dots,F_{n}} \left\{ \frac{\Delta_{n}(F_{1},\dots,F_{n})}{\ell} \colon L_{\mathbf{R},n}^{3}(\varepsilon,\gamma) = \ell \right\}, \quad \varepsilon, \ \gamma > 0,$$
(14)

and present their upper bounds for every $\varepsilon > 0$ and $\gamma > 0$.

Theorem 2. For every $\varepsilon > 0$ and $\gamma > 0$ we have

$$C_{\rm E}^*(\varepsilon,\gamma) \le \frac{4}{\sqrt{2\pi}} + \frac{1}{\pi} \left[\frac{\varkappa}{\varepsilon} \Upsilon\left(1, \frac{t_{\gamma}^2}{2\varepsilon^2}\right) + \frac{\varepsilon}{12} \Upsilon\left(2, \frac{t_{\gamma}^2}{2\varepsilon^2}\right) + \frac{\sqrt{2(6\varkappa\gamma^2 + 1)}}{6\gamma} \Gamma\left(\frac{3}{2}, \frac{t_{\gamma}^2}{2\varepsilon^2}\right) \right] =: \widehat{C_{\rm E}^*}(\varepsilon,\gamma), \tag{15}$$

$$C_{\mathbf{R}}^{*}(\varepsilon,\gamma) \leq \frac{4}{\sqrt{2\pi}} + \frac{1}{\pi} \Big[\frac{\varkappa}{\varepsilon} \Upsilon \Big(1, \frac{t_{1,\gamma}^{2}}{2\varepsilon^{2}} \Big) + \frac{\varepsilon}{12} \Upsilon \Big(2, \frac{t_{1,\gamma}^{2}}{2\varepsilon^{2}} \Big) + \frac{\varepsilon}{6} \Gamma \Big(2, \frac{t_{2,\gamma}^{2}}{2\varepsilon^{2}} \Big) + \frac{\sqrt{2}}{6\gamma} \Big(\frac{\sqrt{\pi}}{2} - \Upsilon \Big(\frac{3}{2}, \frac{t_{1,\gamma}^{2}}{2\varepsilon^{2}} \Big) - \Gamma \Big(\frac{3}{2}, \frac{t_{2,\gamma}^{2}}{2\varepsilon^{2}} \Big) \Big] =: \widehat{C}_{\mathbf{R}}^{*}(\varepsilon,\gamma) \quad (16)$$

where $\Gamma(r,x) \coloneqq \int_x^\infty t^{r-1} e^{-t} dt$, $\Upsilon(r,x) \coloneqq \int_0^x t^{r-1} e^{-t} dt = \Gamma(r) - \Gamma(r,x)$, r, x > 0, are the upper and the lower incomplete gamma functions,

$$t_{\gamma} \coloneqq \frac{2}{\gamma} \big(\sqrt{(\gamma/\gamma_*)^2 + 1} - 1 \big), \quad t_{2,\gamma} = 2 \max \big\{ \gamma^{-1}, \gamma_*^{-1} \big\}, \quad t_{1,\gamma} \coloneqq t_{2,\gamma} \cdot \big(1 - \sqrt{(1 - (\gamma/\gamma_*)^2)_+} \big),$$

 $\varkappa = 0.5315...$ and $\gamma_* = 0.5599...$ are defined in Theorem 1. Moreover, the function $\widehat{C}^*_{\scriptscriptstyle E}(\varepsilon, \gamma)$ is monotonically decreasing with respect to $\varepsilon > 0$ and $\gamma > 0$, the function $\widehat{C}^*_{\scriptscriptstyle R}(\varepsilon, \gamma)$ is monotonically decreasing with respect $\gamma > 0$ being constant for $\gamma \ge \gamma_*$ for every fixed $\varepsilon > 0$. In particular,

$$C_{\rm E}^*(\infty,\gamma) \le \frac{1}{\sqrt{2\pi}} \left(4 + \frac{1}{6}\sqrt{\gamma_*^{-2} + \gamma^{-2}} \right) = \begin{cases} \frac{1}{\sqrt{2\pi}} \left(4 + \frac{\sqrt{2}}{6}\gamma_*^{-1} \right) = 1.7636\dots, & \gamma = \gamma_*, \\ \frac{1}{\sqrt{2\pi}} \left(4 + \frac{1}{6}\sqrt{\gamma_*^{-2} + 1} \right) = 1.7318\dots, & \gamma = 1, \\ \frac{1}{\sqrt{2\pi}} \left(4 + \frac{1}{6\gamma_*} \right) = 1.7145\dots, & \gamma \to \infty, \end{cases}$$
(17)

$$\sup_{\gamma \ge \gamma_*} C^*_{\mathsf{R}}(\varepsilon,\gamma) \le \widehat{C}^*_{\mathsf{R}}(\varepsilon,\gamma_*) = \frac{4}{\sqrt{2\pi}} + \frac{1}{\pi} \left[\frac{\varkappa}{\varepsilon} \Upsilon \left(1, \frac{2}{(\varepsilon\gamma_*)^2} \right) + \frac{\varepsilon}{12} \left(1 + \Gamma \left(2, \frac{2}{(\varepsilon\gamma_*)^2} \right) \right) \right], \quad \varepsilon > 0,$$
(18)
$$\inf_{\varepsilon > 0} \inf_{\gamma > 0} C^*_{\mathsf{R}}(\varepsilon,\gamma) \le \inf_{\varepsilon > 0} \sup_{\gamma \ge \gamma_*} C^*_{\mathsf{R}}(\varepsilon,\gamma) \le \widehat{C}^*_{\mathsf{R}}(1.89,\gamma_*) \le 1.75.$$

The values of the functions $\widehat{C}_{E}^{*}(\varepsilon,\gamma)$ and $\widehat{C}_{R}^{*}(\varepsilon,\gamma)$ for some $\varepsilon > 0$ and $\gamma > 0$ are given in the third columns of tables 2 and 3, respectively. The plot of the level curve $\gamma = \gamma(\varepsilon)$ with the constant value of $\widehat{C}_{E}^{*}(\varepsilon,\gamma) = 1.72$, closest to the minimal one $\widehat{C}_{E}^{*}(\infty,\infty)$ up to 10^{-2} , is given on Fig. 2 (left). The plot of the function $\widehat{C}_{R}^{*}(\varepsilon,\gamma_{*})$ is given on Fig. 3 (left, solid line). The graphs of the functions t_{γ} , $t_{1,\gamma}$, $t_{2,\gamma}$ are given on Fig. 1.

Remark 3. Within the method used, the functions $C_{\text{E}}(\varepsilon, \gamma)$ and $C_{\text{R}}(\varepsilon, \gamma)$ constructed here in the proof of Theorem 1 are bounded below by the functions $\widehat{C}_{\text{E}}^*(\varepsilon, \gamma)$ and $\widehat{C}_{\text{R}}^*(\varepsilon, \gamma)$, respectively.

Remark 4. Since

$$\lim_{\varepsilon \to 0} L_{\mathrm{E},n}(\varepsilon,\gamma) = \lim_{\varepsilon \to 0} L_{\mathrm{R},n}(\varepsilon,\gamma) = 0 \quad \text{for every} \ \gamma > 0,$$

the functions $C_{\text{E}}(\varepsilon,\gamma)$, $C_{\text{R}}(\varepsilon,\gamma)$, $C_{\text{E}}^{*}(\varepsilon,\gamma)$, and $C_{\text{R}}^{*}(\varepsilon,\gamma)$ must be unbounded as $\varepsilon \to 0$.

Remark 5. Though the first terms $\mu_k(\cdot)$, k = 1, 2, ..., n, in the definitions of $L_{\text{E},n}(\varepsilon, \gamma)$, $L_{\text{R},n}(\varepsilon, \gamma)$ vanish for symmetric distributions of random summands, one cannot, in general, get rid of the third truncated moments $\mu_k(\cdot)$ in (11), (12), that is, the "constants" $C_{\text{E}}(\varepsilon, 0+)$, $C_{\text{R}}(\varepsilon, 0+)$, $C_{\text{E}}^*(\varepsilon, 0+)$, and $C_{\text{R}}^*(\varepsilon, 0+)$ are no more bounded. This fact follows from the observation that each of the above constants is bounded below by the so-called asymptotically best constant (we follow here the terminology introduced in [32])

$$\sup_{F_1=\ldots=F_n: B_n>0} \limsup_{n\to\infty} \frac{\Delta_n(F_1,\ldots,F_n)}{\sup_{0< z\leq \varepsilon} zL_n(z)}$$

for which, in Theorem 7 of Section 5, an infinite lower bound is constructed for every $\varepsilon > 0$.

The proofs of Theorems 1 and 2 are based on the method of characteristic functions (ch.f.'s) realized in the Prawitz smoothing inequality [28] (see Lemma 8 below) and new estimates for ch.f.'s presented in Section 3. As an auxiliary result used in bounding the absolute value of a ch.f., we prove a sharp inequality

$$\mathsf{E}(X - X')^2 \mathbf{1}(|X - X'| \ge z) \le 4\mathsf{E}X^2 \mathbf{1}(|X| \ge z/2), \quad z \ge 0,$$

for arbitrary centered i.i.d. r.v.'s X and X' with finite second moment, where the constant factor 4 on the R.H.S. cannot be made less (see Theorem 3 in Section 2). Similar inequality was proved and used in the preceding works with the constant factor 40 by Esseen [8], 12 and 8 by Rozovskii [30,31], respectively.

The proof of the main Theorem 1 is given in Section 4 and consists of two steps. First, we consider the case of small values of the fractions $L_{E,n} \leq L$ and $L_{R,n} \leq L$ with L being small enough including the limit $L \to 0$, where the upper bounds for the corresponding "constants" (in fact, depending on L) are obtained in the analytical form (see Subsection 4.1) yielding, as a particular case, Theorem 2. Then we consider the remaining case, where the upper bounds for the appearing "constants" also depend on the concrete value of the fraction $L \coloneqq L_{E,n}$ or $L \coloneqq L_{R,n}$ and have a more complicated form assuming numerical evaluation by the use of a computer (see Subsection 4.2).

In Section 5 we compare the fractions $L^3_{E,n}(\varepsilon,\gamma)$, $L^3_{R,n}(\varepsilon,\gamma)$, $L_{3,n}$ and also the right-hand sides of Osipov's inequality (5) with our new inequalities (11), (12).

All the figures and tables are presented in the concluding Appendix.

2 An inequality for quadratic tails

Theorem 3. Let X and X' be i.i.d. with the d.f. F and EX = 0. Then with

$$\begin{split} \sigma^2(z) &\coloneqq \mathsf{E} X^2 \mathbf{1}(|X| \ge z) = \int_{|x| \ge z} x^2 dF(x), \\ \sigma_s^2(z) &\coloneqq \mathsf{E} (X - X')^2 \mathbf{1}(|X - X'| \ge z) = \int_{|x-y| \ge z} (x - y)^2 dF(x) dF(y), \quad z \ge 0, \end{split}$$

we have

$$\sigma_s^2(z) \le 2\sigma^2(\alpha z) + 2\sigma^2((1-\alpha)z), \quad z \ge 0, \quad \alpha \in [0,1].$$

In particular, with $\alpha = 1/2$ we have

$$\sigma_s^2(z) \le 4\sigma^2(z/2), \quad z \ge 0, \tag{19}$$

where the constant factor 4 on the right-hand side is the best possible in the sense of

$$\sup_{X: \mathsf{E}X=0} \frac{\sigma_s^2(z)}{\sigma^2(z/2)} = 4 \quad for \; every \quad z > 0, \tag{20}$$

where the least upper bound is taken over all distributions of the r.v. X with $\mathsf{E}X = 0$ and delivered by the sequence of two-point distributions of the form $\mathsf{P}(X = qz) = p$, $\mathsf{P}(X = -pz) = q \coloneqq 1 - p$ with $p \to \frac{1}{2} + .$

Remark 6. In the original work of Esseen [8] inequality (19) was proved with the constant factor 40 on the right-hand side instead of 4, and in the works of Rozovskii [30, 31] this factor was successively lowered to 12 and 8.

Proof. By $F(x,y) = F(x) \cdot F(y)$ denote the d.f. of the random vector (X, X'). Then for every $\alpha \in [0, 1]$ with $\beta := 1 - \alpha$ we have

$$\sigma_s^2(z) = \int_{|x-y| \ge z} (x-y)^2 dF(x,y) \le \int_{|x| \ge \alpha z} + \int_{|y| \ge \beta z} (x-y)^2 dF(x,y) = \int_{|x| \ge \alpha z} + \int_{|y| \ge \beta z} (x^2 - 2xy + y^2) dF(x,y) = \sigma^2(\alpha z) + \sigma^2(\beta z) + \sigma^2 \int_{|x| \ge \alpha z} dF(x) + \sigma^2 \int_{|y| \ge \beta z} dF(y),$$

since $\mathsf{E}X = \mathsf{E}X' = 0$. Observing that

=

$$\sigma^2 \int_{|x| \ge \alpha z} dF(x) = \int_{|x| \ge \alpha z} dF(x) \int_{\mathbb{R}} y^2 dF(y) = \int_{|y| < \alpha z \le |x|} y^2 dF(x) dF(y) + \sigma^2(\alpha z) \int_{|x| \ge \alpha z} dF(x) \le \frac{1}{7} \int_{|x| \ge \alpha z} dF(x) dF(y) dF$$

$$\leq \int_{|y|<\alpha z} dF(y) \int_{|x|\geq \alpha z} x^2 dF(x) + \sigma^2(\alpha z) \int_{|x|\geq \alpha z} dF(x) = \sigma^2(\alpha z),$$

we finally obtain

$$\sigma_s^2(z) \le 2\sigma^2(\alpha z) + 2\sigma^2(\beta z), \quad z \ge 0.$$

To prove (20), fix arbitrary z > 0 and consider a centred two-point distribution

$$X = \begin{cases} a, & \frac{b}{a+b} \\ -b, & \frac{a}{a+b} \end{cases}$$

where the choice of b > a > 0 will depend on z. Then, with X' being an independent copy of X, we have

$$X - X' = \begin{cases} a + b, & \frac{ab}{(a+b)^2}, \\ 0, & \frac{a^2 + b^2}{(a+b)^2}, \\ -a - b, & \frac{ab}{(a+b)^2}, \end{cases}, \quad \sigma_s^2(z) = \begin{cases} 2ab, & 0 \le z \le a + b, \\ 0, & z > a + b, \end{cases}, \quad \sigma^2\left(\frac{z}{2}\right) = \begin{cases} ab, & 0 \le \frac{z}{2} \le a \\ \frac{ab^2}{a+b}, & a < \frac{z}{2} \le b \\ 0, & z/2 > b. \end{cases}$$

Now let a and b satisfy a + b = z. Then $\sigma_s^2(z) = 2ab$ and $\sigma^2(z/2) = ab^2/(a+b) = ab^2/z$, and hence for every z > 0

$$\sup_{0 < a < b: \ a+b=z} \frac{\sigma_s^2(z)}{\sigma^2\left(\frac{z}{2}\right)} = \sup_{z/2 < b < z} \frac{2z}{b} = \lim_{b \to z/2+} \frac{2z}{b} = 4.$$

3 Estimates for characteristic functions

In what follows we shall omit the arguments ε, γ of the fractions $L_{E,n}(\varepsilon, \gamma)$ and $L_{R,n}(\varepsilon, \gamma)$ assuming $\gamma \in (0, \infty)$ and $\varepsilon \in (0, \infty]$ for $L_{E,n}(\varepsilon, \gamma)$ or $\varepsilon \in (0, \infty)$ for $L_{R,n}(\varepsilon, \gamma)$ being fixed and even simply using the notation L for each of the fractions $L_{E,n}, L_{R,n}$. Furthermore, we shall also assume that the random summands X_1, \ldots, X_n are normalized so that

$$B_n^2 \coloneqq \sum_{k=1}^n \sigma_k^2 = 1.$$

For $k = 1, \ldots, n$ denote

$$f_{k}(t) \coloneqq \mathsf{E}e^{itX_{k}}, \quad \overline{f}_{n}(t) = \mathsf{E}\exp\left\{\frac{itS_{n}}{\sqrt{\mathsf{D}S_{n}}}\right\} = \prod_{k=1}^{n} f_{k}\left(t\right), \quad t \in \mathbb{R},$$

the ch.f.'s of X_k and of the (already normalized) sum S_n . Recall that $e^{-t^2/2}$, $t \in \mathbb{R}$, is the ch.f. of the standard normal distribution on \mathbb{R} .

3.1 Estimation of $|\overline{f}_n(t)|$

This section is aimed at bounding the absolute value $|\overline{f}_n(t)|$ of the ch.f. of the normalized sum above. We provide two kinds of such estimates. The first estimate which is the most exact, but has a rather complicated form, is used below in the numerical method described in Subsection 4.2 for the case of $L_{\text{R},n}$, $L_{\text{E},n}$ separated from the origin. The second estimate has a simpler form convenient for analytical work and is used below in Subsection 4.1 for the case of small $L_{\text{R},n}$, $L_{\text{E},n}$.

We also note here that the form of the estimates for $|\overline{f}_n(t)|$ presented below is the same for the both fractions $L_{\text{E},n}^3$ and $L_{\text{R},n}^3$ which is explained by the independence of the constructed bounds of the function $M_n(\cdot)$.

Lemma 1 (see [29]). For every $\theta \in [0, 2\pi]$ and $x \in \mathbb{R}$ we have

$$\cos x \le 1 - a(\theta)x^2 + b(\theta)x^4,\tag{21}$$

with equality if and only if $x \in \{-\theta, 0, \theta\}$, where

$$a(\theta) = 2\frac{1 - \cos\theta}{\theta^2} - \frac{\sin\theta}{2\theta}, \quad b(\theta) = \frac{1 - \cos\theta}{\theta^4} - \frac{\sin\theta}{2\theta^3} \quad \text{for } \theta \in (0, 2\pi)$$

and $a(0) \coloneqq 1/2$, $b(0) \coloneqq 1/24$ are defined by continuity.

Lemma 2. (i) The functions $a(\theta)$ and $b(\theta)$ on the open interval $\theta \in (0, 2\pi)$ are both strictly monotonically decreasing and strictly positive varying within the range

$$\frac{1}{2} = a(0) > a(\theta) > a(2\pi) = 0, \quad \frac{1}{24} = b(0) > b(\theta) > b(2\pi) = 0.$$

(ii) The function $a(\theta)/b(\theta)$ is strictly monotonically increasing for $\theta \in [0, 2\pi)$, in particular,

$$\sup_{0 < \theta < 2\pi} \frac{a(\theta)}{b(\theta)} = \lim_{\theta \to 2\pi^-} \frac{a(\theta)}{b(\theta)} = 4\pi^2$$

Proof. For $\theta \in (0, 2\pi)$ denote $g(\theta) \coloneqq a(\theta)/b(\theta)$. Then, omitting the argument θ of the functions $a(\theta)$ and $b(\theta)$ for short, we have

$$a' = -4\frac{1-\cos\theta}{\theta^3} + 5\frac{\sin\theta}{2\theta^2} - \frac{\cos\theta}{\theta}, \quad b' = -4\frac{1-\cos\theta}{\theta^5} + 5\frac{\sin\theta}{2\theta^4} - \frac{\cos\theta}{\theta^3} = \frac{a'(\theta)}{\theta^2},$$
$$g'(\theta) = \frac{a'b-b'a}{b^2} = -\frac{b'(a-\theta^2b)}{b^2} = -\frac{b'}{b^2} \cdot \frac{1-\cos\theta}{\theta^2},$$
(22)

hence, $\operatorname{sgn} g'(\theta) = -\operatorname{sgn} b'(\theta)$ for all $\theta \in (0, 2\pi)$.

Observe that

$$b_1(\theta) \coloneqq 2\theta^5 b'(\theta) = -8(1 - \cos\theta) + 5\theta \sin\theta - \theta^2 \cos\theta, \quad b_1(0+) = 0, \quad b_1(2\pi -) = -4\pi^2, \tag{23}$$

$$b_1'(\theta) = (\theta^2 - 3)\sin\theta + 3\theta\cos\theta, \quad b_1'(0+) = 0, \quad b_1'(2\pi -) = 6\pi,$$
(24)

$$b_2(\theta) \coloneqq \frac{b_1''(\theta)}{\theta^2} = \cos \theta - \frac{\sin \theta}{\theta}, \quad b_2(0+) = 0, \quad b_2(\pi) = -1, \quad b_2(2\pi) = 1.$$
(25)

It is easy to see that there exists a unique $\theta_0 \in (\pi, 2\pi)$ such that

$$\begin{aligned} b_2(\theta) &< 0, \qquad 0 < \theta < \theta_0, \\ b_2(\theta) &> 0, \qquad \theta_0 < \theta < 2\pi, \end{aligned}$$

With the account of $\operatorname{sgn} b_2 = \operatorname{sgn} b_1''(\theta)$ and $b_1'(0+) = 0$, $b_1'(2\pi-) > 0$, we conclude that b_1' has a unique sign change in a point $\theta_1 \in (\theta_0, 2\pi) \subset (\pi, 2\pi)$ and

$$b_1'(\theta) < 0, \qquad 0 < \theta < \theta_1, \\ b_1'(\theta) > 0, \qquad \theta_1 < \theta < 2\pi$$

Hence, b_1 has the unique point of minimum θ_1 on $(0, 2\pi)$ and for all $\theta \in (0, 2\pi)$

$$b_1(\theta) < \max\{b_1(0+), b_1(2\pi-)\} = 0,$$

i. e., $b'(\theta) = \frac{b_1(\theta)}{2\theta^5} < 0$ for $\theta \in (0, 2\pi)$. Since also $\operatorname{sgn} a' = \operatorname{sgn} b' < 0$ on $(0, 2\pi)$, we immediately obtain part (i) of the lemma. Finally, recalling that $\operatorname{sgn} g' = -\operatorname{sgn} b' > 0$ on $(0, 2\pi)$ we get part (ii) of the lemma.

Theorem 4. For $\varepsilon > 0$, $\tau \ge 0$, $u \ge 0$, and $\theta \in (0, 2\pi]$ let

$$\begin{split} k(\tau, u, \theta) &\coloneqq a(\theta) - 4\tau u^{-1} \left(a(\theta) + b(\theta) u^2 \right) \text{ for } u > 0, \quad k(0, 0, \cdot) \coloneqq a(\theta), \quad k(\tau, 0, \cdot) \coloneqq -\infty \text{ for } \tau > 0, \\ k(\tau, u) &\coloneqq \sup \left\{ k(\tau, v, \theta) \colon 0 \le v \le u \land \sqrt{\frac{a(\theta)}{b(\theta)}}, \ \theta \in (0, 2\pi) \right\} = \sup_{0 < \theta < 2\pi} k\left(\tau, u \land \sqrt{\frac{a(\theta)}{b(\theta)}}, \theta \right), \\ k\left(\tau\right) &= \sup_{\theta \in (0, 2\pi], u > 0} k(\tau, u, \theta) = \sup_{0 < \theta < 2\pi} k\left(\tau, \sqrt{\frac{a(\theta)}{b(\theta)}}, \theta \right) = \sup_{0 < \theta \le 2\pi} \left\{ a(\theta) - 8\tau \sqrt{a(\theta)b(\theta)} \right\}, \end{split}$$

where the functions $a(\theta)$, $b(\theta)$ are defined in the formulation of Lemma 1. Then for each of the fractions $L \in \{L_{\text{R},n}, L_{\text{E},n}\}$ we have

(i) for every $t \in \mathbb{R}$

$$\left|\overline{f}_{n}(t)\right| \leq \exp\left\{-k\left(L^{3}|t|, 2\varepsilon|t|\right)t^{2}\right\};$$
(26)

(ii) for every $L_0 \ge L$, $\tau \ge \underline{\tau}_1(L_0, \varepsilon) \coloneqq \pi L_0^3/\varepsilon$ and $|t| \le \tau/L^3$

$$\left|\overline{f}_{n}(t)\right| \leq \exp\left\{-k(\tau)t^{2}\right\},\tag{27}$$

with $k(\tau) > 0$ if and only if $\tau < \overline{\tau}_1 := \frac{\pi}{4} = 0.7853...$ and the interval $[\underline{\tau}_1(L_0, \varepsilon), \overline{\tau}_1)$ being nonempty if and only if $L_0 < (\varepsilon/4)^{1/3}$.

(iii) Statements (i) and (ii) remain true with $L^3 = \sup_{0 < z \le \varepsilon} zL_n(z)$.

Remark 7. The right-hand side of (26) is monotonically increasing with respect to L > 0, monotonically decreasing with respect to $\varepsilon > 0$ and does not depend on γ with L being fixed. The right-hand side of (27) does not depend neither on L, nor on ε and γ .

Proof. Due to the symmetry, one can assume that $t \ge 0$. Moreover, inequalities (26) and (27) hold trivially true with t = 0, so that, in what follows, we shall assume that t > 0.

Consider the absolute value of the ch.f. of every single random summand. Namely, let X and X' be i.i.d. centred r.v.'s with the d.f. F, the ch.f. f, and $\sigma^2 := \mathsf{E}X^2 < \infty$. Denote $F^s(x) := \mathsf{P}(X - X' < x), x \in \mathbb{R}$, the symmetrization of F,

$$\sigma^2(z) \coloneqq \int_{|x| \ge z} x^2 dF(x), \quad \text{and} \quad \sigma^2_s(z) \coloneqq \int_{|x| \ge z} x^2 dF^s(x), \quad z \ge 0.$$

Then for every t under consideration and $\theta \in (0, 2\pi]$ we have

$$|f(t)|^{2} = \mathsf{E}\cos t \left(X - X' \right) = 1 - 2a(\theta)\sigma^{2}t^{2} + J,$$

with $a(\theta)$ defined in Lemma 1 and

$$J = J(t,\theta) \coloneqq \int_{\mathbb{R}} \left(\cos tx - 1 + a(\theta)t^2x^2 \right) dF^s(x)$$

Separating the range of integration in J into two domains: $|x| \ge z$ and |x| < z with arbitrary $z \ge 0$, we use the inequality $\cos x - 1 \le 0$, $x \in \mathbb{R}$, to bound the integrand in the first integral and Lemma 1 to bound the second and obtain

$$J \le at^2 \int_{|x| \ge z} x^2 dF^s(x) + bt^4 \int_{|x| < z} x^4 dF^s(x) = \left(a - bt^2 z^2\right) t^2 \sigma_s^2(z) + 2bt^4 \int_0^z x \sigma_s^2(x) dx,$$

where, for short, we omitted the argument θ of the functions $a(\theta)$ and $b(\theta)$ defined in Lemma 1 and used the representation

$$\int_{|x|
(28)$$

with $F := F^s$, which is valid for arbitrary d.f. F with finite second moment. Now assuming that $a \ge bt^2 z^2$ we may apply Theorem 3 to bound $\sigma_s^2(\cdot)$ in the majorant for J above and obtain

$$J \le 4t^2 \left(a - bt^2 z^2 \right) \sigma^2(z/2) + 8bt^4 \int_0^z x \sigma^2 \left(\frac{x}{2} \right) dx, \quad z \ge 0.$$

Now repeating the same procedure for every random summand $X \coloneqq X_k, k = 1, \ldots, n$, we construct an estimate for the absolute value of the corresponding ch.f. $|f_k(t)|$ in the form

$$|f_k(t)|^2 = 1 - 2a\sigma_k^2 t^2 + J_k,$$

where

$$J_k = J_k(t,\theta) \le 4t^2 \left(a - bt^2 z^2\right) \sigma_k^2(z/2) + 8bt^4 \int_0^z x \sigma_k^2\left(\frac{x}{2}\right) dx, \quad z \ge 0,$$

with

$$\sum_{k=1}^{n} J_{k} \leq 4t^{2} \left(a - bt^{2}z^{2}\right) L_{n}\left(\frac{z}{2}\right) + 16bt^{4} \int_{0}^{z} \frac{x}{2} L_{n}\left(\frac{x}{2}\right) dx \leq \\ \leq \left(8t^{2}z^{-1} \left(a - bt^{2}z^{2}\right) + 16bt^{4}z\right) \sup_{0 < x \leq z/2} x L_{n}\left(x\right) \leq 8t^{2}z^{-1} \left(a + bt^{2}z^{2}\right) \sup_{0 < x \leq \varepsilon} x L_{n}\left(x\right)$$

for all $\varepsilon \ge z/2 \ge 0$, provided that $a \ge bt^2 z^2$. Thus, using the inequality $1 + x \le e^x$, $x \in \mathbb{R}$, and observing that

$$\sup_{0 < x \le \varepsilon} x L_n(x) \le \min\{L^3_{\mathrm{E},n}, L^3_{\mathrm{R},n}\} \le L^3$$

(compare with part (iii) of the theorem), we obtain

$$\left|\overline{f}_{n}(t)\right| = \prod_{k=1}^{n} |f_{k}(t)| \leq \exp\left\{-at^{2} + \frac{1}{2}\sum_{k=1}^{n} J_{k}\right\} \leq \exp\left\{-at^{2} + 4L^{3}t^{2}z^{-1}(a+bt^{2}z^{2})\right\} = \exp\left\{-t^{2}\left(a+4L^{3}z^{-1}(a+bt^{2}z^{2})\right)\right\} =:\exp\left\{-t^{2} \cdot k(L^{3}t,tz,\theta)\right\}$$
(29)

for all $t \in (0,\infty)$, $0 < z \leq \frac{1}{t} \sqrt{\frac{a(\theta)}{b(\theta)}} \wedge 2\varepsilon \rightleftharpoons z_*(t,\theta,\varepsilon)$, and $\theta \in (0,2\pi]$. Note that the function

$$k(\tau, u, \theta) = a - 4\tau u^{-1} \left(a + bu^2\right)$$

attains its global maximum with respect to $u \in (0, \infty)$ for every fixed $\tau > 0$ and $\theta \in (0, 2\pi)$ at the point $u = \sqrt{\frac{a(\theta)}{b(\theta)}} =: u_*(\theta)$ with

$$\max_{u>0} k(\tau, u, \theta) = k\left(\tau, u_*(\theta), \theta\right) = a(\theta) - 8\tau \sqrt{a(\theta)b(\theta)}, \quad \tau > 0, \quad \theta \in (0, 2\pi),$$

being monotonically increasing for $0 < u < u_*(\theta)$ and monotonically decreasing for $u > u_*(\theta)$ (in fact, $k(\tau, u, \theta)$ is concave with respect u > 0 for every fixed $\tau > 0$ and $\theta \in (0, 2\pi)$). Now choosing $z = z_*$ to minimize the right-hand side of (29) with respect to $z \in (0, z_*]$ (which is equivalent to the choice of $u = u_*(\theta) \land 2\varepsilon t$) and optimizing then with respect to $\theta \in (0, 2\pi]$ we arrive at (26) with the right-hand side being a monotonically increasing function of L > 0 as the least upper bound to a family of monotonically decreasing functions.

Let us prove that (26) (or (29)) yields (27) under the assumptions stated in the formulation of part (ii) of the theorem. For this purpose, observe that for every $\theta \in (0, 2\pi)$

$$-k(L^{3}t, tz_{*}, \theta) + a = \begin{cases} 2aL^{3}\varepsilon^{-1} + 8bL^{3}\varepsilon t^{2} \leq 2aL^{3}\varepsilon^{-1} + 4L^{3}t\sqrt{ab}, & 2\varepsilon t \leq \sqrt{a/b}, \\ 8L^{3}t\sqrt{ab}, & 2\varepsilon t > \sqrt{a/b}, \end{cases}$$

so that for all $t \in \mathbb{R}$ we have

$$-k(L^{3}t, tz_{*}, \theta) + a \leq \max\left\{2aL^{3}\varepsilon^{-1} + 4L^{3}t\sqrt{ab}, 8L^{3}t\sqrt{ab}\right\}$$

Now let τ be an arbitrary positive number. Then for $0 < t \le \tau/L^3$ we trivially obtain

$$-k(L^{3}t, tz_{*}, \theta) + a \leq \max\left\{2aL^{3}\varepsilon^{-1} + 4\tau\sqrt{ab}, 8\tau\sqrt{ab}\right\} = 8\tau\sqrt{ab}, \quad \text{if} \ \tau \geq \frac{L^{3}}{2\varepsilon}\sqrt{\frac{a}{b}}$$

and if for $L_0 \ge L$ the uniform condition

$$\tau \geq \frac{L_0^3}{2\varepsilon} \sup_{0 < \theta < 2\pi} \sqrt{\frac{a(\theta)}{b(\theta)}} = \frac{L_0^3}{2\varepsilon} \lim_{\theta \to 2\pi-} \sqrt{\frac{a(\theta)}{b(\theta)}} = \frac{\pi L_0^3}{\varepsilon} \eqqcolon \underline{\tau}_1(L_0, \varepsilon)$$

holds, where the equality sign is due to Lemma 2, then also

$$-k(L^{3}t, 2\varepsilon t) \coloneqq -\sup_{0<\theta<2\pi} k(L^{3}t, tz_{*}(t, \theta, \varepsilon), \theta) \leq -\sup_{0<\theta<2\pi} \left\{ a(\theta) - 8\tau \sqrt{a(\theta)b(\theta)} \right\} \eqqcolon -k(\tau)$$

for every $t \leq \tau/L^3$ and $L \geq L_0$. Note that for fixed ε the quantity $\underline{\tau}_1(L_0, \varepsilon)$ can be made arbitrarily small by the choice of $L_0 = L_0(\varepsilon) > 0$ small enough.

Finally, observe that the function $h(\tau, \theta) \coloneqq a(\theta) - 8\tau \sqrt{a(\theta)b(\theta)}, \tau > 0, \theta \in (0, 2\pi)$ is strictly positive if and only if $\tau < \frac{1}{8}\sqrt{\frac{a(\theta)}{b(\theta)}} \eqqcolon \tau_1(\theta)$, and hence, $k(\tau) \coloneqq \sup_{0 < \theta < 2\pi} h(\tau, \theta) > 0$ if and only if there exists $\theta \in (0, 2\pi)$ such that $\tau < \tau_1(\theta)$. The latest condition is equivalent to $\tau < \sup_{0 < \theta < 2\pi} \tau_1(\theta) = \pi/4 \eqqcolon \overline{\tau}_1$. The proof is completed by the remark that the set of admissible values of $\tau \in [\underline{\tau}_1(L_0, \varepsilon), \overline{\tau}_1)$ is not empty if and only if

$$\underline{\tau}_1(L_0,\varepsilon) \coloneqq \pi L_0^3/\varepsilon < \pi/4 \eqqcolon \overline{\tau}_1,$$

i. e. $L_0 < (\varepsilon/4)^{1/3}$.

3.2

Estimation of $|\overline{f}_n(t) - e^{-t^2/2}|$.

The present section is aimed at bounding the absolute value of the difference $|\overline{f}_n(t) - e^{-t^2/2}|$ between the ch.f. of the normalized sum and that of the standard normal law. Similarly to the preceding section, we construct two estimates, the first being the most exact and used in Subsection 4.2 and the second being the most convenient for analytical work in Subsection 4.1.

However, unlike in the previous section, here significant distinctions arise in the dependence of which of the fractions $L_{E,n}^3$ or $L_{R,n}^3$ is used. We pay special attention to the appearing distinctions and explain their reasons.

Before passing to the main theorem of the present subsection we proof four auxiliary statements.

Lemma 3. For every $x \in \mathbb{R}$ we have

$$\left|e^{ix} - 1 - ix - (ix)^2/2\right| \le \varkappa \cdot x^2,$$

where $\varkappa := x^{-2} \sqrt{(\cos x - 1 + x^2/2)^2 + (\sin x - x)^2} \Big|_{x=x_0} = 0.531551...$ and $x_0 = 5.487414...$ is the unique root of the equation

 $8(\cos x - 1) + 8x\sin x - 4x^2\cos x - x^3\sin x = 0$

on the interval $(0, 2\pi]$ lying in $(\pi, 2\pi)$.

Proof. Due to the symmetry and triviality of the stated inequality for x = 0 we may assume that x > 0. Observe that the inequality of interest is equivalent to the relation

$$\max_{x>0} g(x) = g(x_0) = 0.2825... = \varkappa^2, \quad \text{where} \quad g(x) := \frac{\left(\cos x - 1 + x^2/2\right)^2 + \left(\sin x - x\right)^2}{x^4}, \quad x > 0.$$

We have

$$h(x) := x^5 g'(x) = 8(\cos x - 1) + 8x \sin x - 4x^2 \cos x - x^3 \sin x, \quad h'(x) = x^2 (\sin x - x \cos x).$$

Note that the function h'(x) has a unique root $x_1 \in (\pi, 3\pi/2)$ on $(0, 2\pi]$, which is the point of maximum of h(x). Since, h(0) = 0 and h is strictly increasing on $(0, x_1)$, we conclude that $h(x_1) > 0$. Moreover, $h(2\pi) = -16\pi^2 < 0$, hence, h has a unique sign change on $(0, 2\pi]$ and this sign change occurs in the point $x_0 \in (x_1, 2\pi) \subset (\pi, 2\pi)$ from + to -. Thus, the function g has a unique stationary point $x_0 \in (\pi, 2\pi)$ on $(0, 2\pi]$, which is the point of maximum. So, it remains to prove that $g(x) \leq g(x_0)$ for $x > 2\pi$.

If $x = 2\pi + y \in (2\pi, 9\pi/4]$, then, with the account of the monotone increase of $\sin y$ and monotone decrease of $\cos y$ for $y \in (0, \pi/4]$, we have

$$\frac{h'(x)}{x^2} = \sin y - (y + 2\pi)\cos y \le \sin \frac{\pi}{4} - 2\pi \cos \frac{\pi}{4} = \frac{\sqrt{2}}{2}(1 - 2\pi) < 0.$$

Hence, h(x) is strictly monotonically decreasing for $x \in (2\pi, 9\pi/4]$ with

$$\sup_{x \in (2\pi, 9\pi/4]} h(x) = h(2\pi +) = -16\pi^2 < 0,$$

so that $g'(x) \coloneqq x^{-5}h(x) < 0$ for $x \in (2\pi, 9\pi/4]$ and

$$\sup_{x \in (2\pi, 9\pi/4]} g(x) = g(2\pi +) = \frac{1}{4} \left(1 + \pi^{-2} \right) = 0.2753 \dots < 0.2825 \dots = \varkappa^2.$$

Finally, for $x > 9\pi/4$ we use the trivial bound

$$g(x) = \frac{2(1-\cos x) + x^4/4 + x^2\cos x - 2x\sin x}{x^4} < \left(\frac{1}{4} + \frac{1}{x^2} + \frac{2}{x^3}\right) \bigg|_{x=9\pi/4} < 0.2757 < \varkappa^2,$$

which completes the proof of the lemma.

Lemma 4. For $\varepsilon > 0$ denote

$$\alpha(\varepsilon) := \inf_{0 < x < \varepsilon \land 1} \left(x - x^3 \right)^{-2/3} = \begin{cases} \left(\varepsilon - \varepsilon^3 \right)^{-2/3}, & \text{if } \varepsilon \le 3^{-1/2}, \\ 3 \cdot 2^{-2/3} = 1.8898 \dots, & \text{otherwise.} \end{cases}$$

Then for every $\varepsilon > 0$ we have

$$\max_{1 \le k \le n} \sigma_k^2 \le \alpha(\varepsilon) \sup_{0 < x \le \varepsilon} \left(x L_n(x) \right)^{2/3}.$$

Proof. From the definition of $L_n(z) \coloneqq \sum_{k=1}^n \sigma_k^2(z)$ it follows that

$$\sup_{0 < x \le \varepsilon} x L_n(x) \ge z \sigma_k^2(z)$$

for every $z \in (0, \varepsilon]$ and k = 1, ..., n. On the other hand, by the definition of $\sigma_k^2 := \mathsf{E} X_k^2$, we have

$$\sigma_k^2 = \mathsf{E} X_k^2 \mathbf{1}(|X_k| < z) + \mathsf{E} X_k^2 \mathbf{1}(|X_k| \ge z) \le z^2 + \sigma_k^2(z), \quad z > 0, \quad k = 1, \dots, n$$

whence it follows that

$$\sup_{0 < x \le \varepsilon} x L_n(x) \ge z \left(\sigma_k^2 - z^2\right) \quad \text{for } 0 < z \le \varepsilon.$$

Now the choice of $z \coloneqq \sigma_k / \sqrt{3} \wedge \varepsilon \sigma_k \leq \varepsilon$ as a maximizer to the right-hand side of the latest inequality completes the proof of the lemma.

Lemma 5. For all $t \in \mathbb{R}$ and $0 < z \le \varepsilon \le \infty$ we have

$$\sum_{k=1}^{n} |f_k(t) - 1|^2 \le t^4 \left[\frac{z}{2} \sup_{0 < x \le \varepsilon} x L_n(x) + \frac{1}{4} \sum_{k=1}^{n} \sigma_k^2 \sigma_k^2(z) \right] \le t^4 \left[\frac{z}{2} \sup_{0 < x \le \varepsilon} x L_n(x) + \frac{\alpha(\varepsilon)}{4z} \sup_{0 < x \le \varepsilon} (x L_n(x))^{5/3} \right]$$

Proof. For every random summand X with the d.f. F, the ch.f. f and $\mathsf{E}X = 0$, $\sigma^2(z) \coloneqq \mathsf{E}X^2 \mathbf{1}(|X| \ge z)$, $z \ge 0$, $\sigma^2 \coloneqq \sigma^2(0) = \mathsf{E} X^2$ we have

$$|f(t) - 1|^{2} \leq \left(\frac{t^{2}\sigma^{2}}{2}\right)^{2} = \frac{t^{4}}{4} \left[\int_{|x| < z} x^{2} dF(x) + \sigma^{2}(z) \right]^{2} =$$
$$= \frac{t^{4}}{4} \left[\left(\int_{|x| < z} x^{2} dF(x) \right)^{2} + 2\sigma^{2}(z) \int_{|x| < z} x^{2} dF(x) + (\sigma^{2}(z))^{2} \right], \quad z \geq 0$$

Using the Jensen inequality for the first term, both the representation $\int_{|x| < z} x^2 dF(x) = \sigma^2 - \sigma^2(z)$ and the bound $\int_{|x| < z} x^2 dF(x) \leq z^2$ for the second term in square brackets in the first step below, and then equality (28) we obtain

$$|f(t) - 1|^2 \le \frac{t^4}{4} \left[\int_{|x| < z} x^4 dF(x) + z^2 \sigma^2(z) + \sigma^2 \sigma^2(z) \right] = \frac{t^4}{4} \left[2 \int_0^z x \sigma^2(x) dx + \sigma^2 \sigma^2(z) \right].$$

Summing up the constructed bounds for every random summand $X := X_k$ over all k = 1, ..., n, we get

$$\sum_{k=1}^{n} |f_k(t) - 1|^2 \le \frac{t^4}{4} \left[2 \int_0^z x L_n(x) dx + \sum_{k=1}^n \sigma_k^2 \sigma_k^2(z) \right] \le t^4 \left[\frac{z}{2} \sup_{0 < x \le z} x L_n(x) + \frac{1}{4} \sum_{k=1}^n \sigma_k^2 \sigma_k^2(z) \right].$$
13

Now the observation that the least upper bound with respect to $x \in (0, z]$ is a non-decreasing function of z yields the first claim of the lemma. The second claim follows from Lemma 4 yielding the chain of inequalities

$$\sum_{k=1}^{n} \sigma_k^2 \sigma_k^2(z) \le \alpha(\varepsilon) \sup_{0 < x \le \varepsilon} \left(x L_n(x) \right)^{2/3} L_n(z) \le \frac{\alpha(\varepsilon)}{z} \sup_{0 < x \le \varepsilon} \left(x L_n(x) \right)^{5/3}, \quad z \in (0, \varepsilon].$$

Lemma 6. For all $t \in \mathbb{R}$, $\varepsilon > 0$, and $\gamma > 0$ we have

$$\sum_{k=1}^{n} \left(f_k(t) - 1 + \sigma_k^2 t^2 / 2 \right) \bigg| \le p_{\mathsf{E}}(t,\varepsilon,\gamma) \cdot L^3_{\mathsf{E},n}(\varepsilon,\gamma), \tag{30}$$

$$\left|\sum_{k=1}^{n} \left(f_k(t) - 1 + \sigma_k^2 t^2 / 2 \right) \right| \le p_{\mathsf{R}}(t,\varepsilon,\gamma) \cdot L^3_{\mathsf{R},n}(\varepsilon,\gamma), \tag{31}$$

where

$$p_{\mathsf{E}}(t,\varepsilon,\gamma) \coloneqq t^2 \min_{0 < z \le \varepsilon} \left[\frac{zt^2}{12} + \max\left\{ \frac{|t|}{6\gamma}, \frac{\varkappa}{z} - \frac{zt^2}{24} \right\} \right] = \begin{cases} \frac{\varkappa t^2}{\varepsilon} + \frac{\varepsilon t^4}{24}, & \varepsilon |t| \le t_\gamma, \\ \frac{\sqrt{6\varkappa\gamma^2 + 1}}{6\gamma} |t|^3, & \varepsilon |t| > t_\gamma, \end{cases}$$
(32)

$$p_{\mathbf{R}}(t,\varepsilon,\gamma) \coloneqq t^{2} \max\left\{\frac{|t|}{6\gamma}, \frac{\varkappa}{\varepsilon} + \frac{\varepsilon t^{2}}{24}, \frac{\varepsilon t^{2}}{12}\right\} = \begin{cases} \frac{\varkappa t^{2}}{\varepsilon} + \frac{\varepsilon t^{4}}{24}, \quad \varepsilon|t| \leq t_{\infty}, \qquad \gamma \geq \gamma_{*}, \\ \frac{\varepsilon t^{4}}{12}, \qquad \varepsilon|t| > t_{\infty}, \qquad \gamma \geq \gamma_{*}, \\ \left\{\frac{\frac{\varkappa t^{2}}{\varepsilon} + \frac{\varepsilon t^{4}}{24}, \quad \varepsilon|t| \leq t_{1,\gamma}, \qquad = p_{\mathbf{R}}(t,\varepsilon,\gamma \wedge \gamma_{*}), \quad (33) \\ \frac{\frac{|t|^{3}}{6\gamma}, \qquad t_{1,\gamma} < \varepsilon|t| \leq t_{2,\gamma}, \qquad \gamma < \gamma_{*}, \\ \frac{\varepsilon t^{4}}{12}, \qquad \varepsilon|t| > t_{2,\gamma}, \end{cases}$$

$$t_{\gamma} \coloneqq \frac{2}{\gamma} \left(\sqrt{(\gamma/\gamma_{*})^{2} + 1} - 1\right), \quad t_{\infty} \coloneqq \lim_{\gamma \to \infty} t_{\gamma} = 2/\gamma_{*} = 3.5717 \dots, \\ t_{2,\gamma} \coloneqq 2 \max\left\{\gamma^{-1}, \gamma_{*}^{-1}\right\}, \quad t_{1,\gamma} \coloneqq t_{2,\gamma} \left(1 - \sqrt{(1 - (\gamma/\gamma_{*})^{2})_{+}}\right), \end{cases}$$

 $\gamma_* = 1/\sqrt{6\varkappa} = 0.5599\ldots$ and \varkappa is defined in Lemma 3. Moreover, the functions t_{γ} and $t_{1,\gamma}$ are monotonically increasing with respect to γ , and $t_{\gamma} \leq t_{1,\gamma} \leq t_{\infty}$ for all $\gamma > 0$, the functions $p_{\rm E}(t,\varepsilon,\gamma)$, $p_{\rm R}(t,\varepsilon,\gamma)$ are monotonically decreasing with respect to $\gamma > 0$ with $p_{\rm R}(t,\varepsilon,\gamma)$ do not depending on γ for $\gamma \geq \gamma_*$, and $p_{\rm E}(t,\varepsilon,\gamma)$ is also monotonically decreasing with respect to ε .

The values of the function $t_{\gamma} = 2\gamma^{-1} (\sqrt{(\gamma/\gamma_*)^2 + 1} - 1)$ for some γ are given in Table 1. The plots of the functions t_{γ} and $t_{1,\gamma}$ are given on Fig. 1 in the Appendix.

γ	0 +	0.25	0.41	γ_*	0.73	1	1.25	1.5	∞
t_{γ}	0	0.7611	1.1678	1.4794	1.7617	2.0935	2.3137	2.4791	3.5717

Table 1. Values of the function $t_{\gamma} = 2\gamma^{-1} \left(\sqrt{(\gamma/\gamma_*)^2 + 1} - 1 \right)$ for some γ (rounded down).

Remark 8. For $\gamma \geq \gamma_*$ we have $t_{1,\gamma} = t_{2,\gamma} = 2/\gamma_* = t_{\infty}$.

Remark 9. The functions $p_{\rm E}(t,\varepsilon,\gamma)$ and $p_{\rm R}(t,\varepsilon,\gamma)$ coincide on the interval $|t| \leq t_{\gamma}/\varepsilon$ for every $\gamma > 0$.

Proof. Since $\mathsf{E}X_k = 0$ for all $k = 1, \ldots, n$, for every $t \in \mathbb{R}$ and z > 0 we have

$$\begin{split} I &\coloneqq \left| \sum_{k=1}^{n} \left(f_k(t) - 1 + \sigma_k^2 t^2 / 2 \right) \right| = \left| \sum_{k=1}^{n} \int_{\mathbb{R}} \left(e^{itx} - 1 - itx - \frac{(itx)^2}{2} \right) dF_k(x) \right| \le \\ &\le \sum_{k=1}^{n} \int_{|x| < z} \left| e^{itx} - 1 - itx - \frac{(itx)^2}{2} - \frac{(itx)^3}{6} \right| dF_k(x) + \\ &+ \left| \sum_{k=1}^{n} \int_{|x| < z} \frac{(itx)^3}{6} dF_k(x) \right| + \sum_{k=1}^{n} \int_{|x| \ge z} \left| e^{itx} - 1 - itx - \frac{(itx)^2}{2} \right| dF_k(x). \end{split}$$

Using the inequalities $|e^{iy} - 1 - iy - \frac{1}{2}(iy)^2 - \frac{1}{6}(iy)^3| \le \frac{1}{24}y^4$ and $|e^{iy} - 1 - iy - \frac{1}{2}(iy)^2| \le \varkappa y^2$ valid for all $y \in \mathbb{R}$ and with the account of (28) we obtain

$$I \leq \frac{t^4}{24} \sum_{k=1}^n \int_{|x| < z} x^4 dF_k(x) + \frac{|t|^3}{6} |M_n(z)| + \varkappa t^2 \sum_{k=1}^n \sigma_k^2(z) =$$

$$= \frac{t^4}{12} \int_0^z x L_n(x) dx - \frac{z^2 t^4}{24} L_n(z) + \frac{|t|^3}{6} |M_n(z)| + \varkappa t^2 L_n(z) \leq$$

$$\leq \frac{z t^4}{12} \sup_{0 < x \leq z} x L_n(x) + \left(\frac{\varkappa t^2}{z} - \frac{z t^4}{24}\right) z L_n(z) + \frac{|t|^3}{6} |M_n(z)|, \quad z > 0, \ t \in \mathbb{R}.$$
(34)

To prove (30), observe that (34) yields

$$I \leq \frac{zt^4}{12} \sup_{0 < x \leq z} xL_n(x) + \max\left\{\frac{\varkappa t^2}{z} - \frac{zt^4}{24}, \frac{|t|^3}{6\gamma}\right\} (zL_n(z) + \gamma |M_n(z)|)$$
$$\leq t^2 \left(\frac{zt^2}{12} + \max\left\{\frac{\varkappa}{z} - \frac{zt^2}{24}, \frac{|t|}{6\gamma}\right\}\right) \sup_{0 < x \leq z} \{\gamma |M_n(x)| + xL_n(x)\}$$

for all $t \in \mathbb{R}$ and z > 0. Now choosing $z = \frac{2}{\gamma|t|} \left(\sqrt{6\varkappa\gamma^2 + 1} - 1 \right) \wedge \varepsilon$ to minimize the R.H.S. of the expression in the large brackets with respect to $z \in (0, \varepsilon]$ and observing that

$$\sup_{0 < x \le z} \left\{ \gamma \left| M_n(x) \right| + x L_n(x) \right\} = L_{\mathbf{E},n}(z,\gamma) \le L_{\mathbf{E},n}(\varepsilon,\gamma)$$

for every $0 < z \leq \varepsilon$ and $\gamma > 0$, we arrive at (30).

To prove (31), observe that (34) yields

$$I \leq \left(\frac{zt^4}{12} + \left(\frac{\varkappa t^2}{z} - \frac{zt^4}{24}\right)_+\right) \sup_{0 < x \leq z} xL_n(x) + \frac{|t|^3}{6} |M_n(z)| \leq \\ \leq \max\left\{\frac{zt^4}{12} + \left(\frac{\varkappa t^2}{z} - \frac{zt^4}{24}\right)_+, \frac{|t|^3}{6\gamma}\right\} \left(\sup_{0 < x \leq z} xL_n(x) + \gamma |M_n(z)|\right) = p_{\mathsf{R}}(t, z, \gamma)L^3_{\mathsf{R}, n}(z, \gamma)$$

for every z > 0 and $t \in \mathbb{R}$. The choice of $z = \varepsilon$ completes the proof of inequality (31). The stated properties of the functions t_{γ} , \tilde{t}_{γ} , $p_{\rm E}$, and $p_{\rm R}$, as well as representations for $p_{\rm E}$ and $p_{\rm R}$ in the right-hand sides of (32) and (33) are proved by use of elementary analysis.

Now we are ready to prove the main result of the present subsection.

Theorem 5. For $\varepsilon > 0$, $\gamma > 0$, L > 0, $\tau > 0$ denote

$$\alpha(\varepsilon) := \inf_{0 < x < \varepsilon \land 1} (x - x^3)^{-2/3} = \begin{cases} (\varepsilon - \varepsilon^3)^{-2/3}, & \text{if } \varepsilon \le 1/\sqrt{3} = 0.5773 \dots, \\ 3 \cdot 2^{-2/3} = 1.8898 \dots, & \text{otherwise}, \end{cases}$$

$$\overline{\tau}_0(\varepsilon) \coloneqq \sqrt{\frac{2}{\alpha(\varepsilon)}}, \qquad \overline{L}_0(\varepsilon) \coloneqq \varepsilon \sqrt{\frac{2}{\alpha(\varepsilon)}} = \varepsilon \overline{\tau}_0(\varepsilon),$$

$$\alpha_1(\varepsilon, L) = \min_{0 < u \le \varepsilon L^{-1}} \left(\frac{u}{2} + \frac{\alpha(\varepsilon)}{4u}\right) = \begin{cases} \sqrt{\alpha(\varepsilon)/2}, & L \le \overline{L}_0(\varepsilon), \\ \frac{\varepsilon}{2L} + \frac{\alpha(\varepsilon)}{4\varepsilon}L, & L > \overline{L}_0(\varepsilon), \end{cases}$$

$$= \sup_{x > 0} x^{-2} \sqrt{(\cos x - 1 + x^2/2)^2 + (\sin x - x)^2} = 0.5315 \dots, \quad \gamma_* = 1/\sqrt{6\varkappa} = 0.5599 \dots$$

For $\varepsilon > 0$, $\gamma > 0$, L > 0, $0 < \tau < \overline{\tau}_0(\varepsilon)$ also introduce the functions

$$\begin{split} B(\tau,\varepsilon,L) &= -\frac{4\alpha_1(\varepsilon,L)}{\alpha^2(\varepsilon)\tau^4} \Big[\ln\left(1 - \frac{1}{2}\alpha(\varepsilon)\tau^2\right) + \frac{1}{2}\alpha(\varepsilon)\tau^2 \Big], \quad B(\tau,\varepsilon) = \frac{\sqrt{\alpha(\varepsilon)/2}}{2 - \alpha(\varepsilon)\tau^2}, \\ A(\tau,\varepsilon) &= \exp\left\{\tau^4 B(\tau,\varepsilon)\right\}, \quad A_{\rm E}(\tau,\varepsilon,\gamma,L) = A(\tau,\varepsilon)\exp\left\{\frac{\tau^3}{6\gamma} + \sqrt{\frac{\varkappa}{3}}\tau^3 + \frac{\varkappa}{\varepsilon}L\tau^2\right\}, \\ A_{\rm R}(\tau,\varepsilon,\gamma,L) &= A(\tau,\varepsilon)\exp\left\{\frac{\tau^3}{6\gamma}\mathbf{1}(\gamma<\gamma_*) + \frac{\varkappa}{\varepsilon}L\tau^2\right\}, \end{split}$$

(i) For every $L \in \{L_{E,n}(\varepsilon,\gamma), L_{R,n}(\varepsilon,\gamma)\}, \varepsilon > 0, \gamma > 0, and L|t| < \overline{\tau}_0(\varepsilon)$ we have

$$r_n(t) \coloneqq \left|\overline{f}_n(t) - e^{-t^2/2}\right| \le \left(\exp\left\{L^3 p(t,\varepsilon,\gamma) + L^4 t^4 B(L|t|,\varepsilon,L)\right\} - 1\right) e^{-t^2/2},\tag{35}$$

where $p = p_{\rm E}$ for $L = L_{{\rm E},n}(\varepsilon,\gamma)$ and $p = p_{\rm R}$ for $L = L_{{\rm R},n}(\varepsilon,\gamma)$ with $p_{\rm E}$, $p_{\rm R}$ defined in Lemma 6. In particular, for $L = L_{{\rm E},n}(\varepsilon,\gamma)$ with $\varepsilon = \infty$ we have

$$r_n(t) \le \left(\exp\left\{ \frac{\sqrt{6\varkappa\gamma^2 + 1}}{6\gamma} L^3 - 4\sqrt{2} \cdot 3^{-3/2} \left[\ln\left(1 - 3 \cdot 2^{-5/3}\tau^2 \right) + 3 \cdot 2^{-5/3}\tau^2 \right] \tau^{-4} t^4 L^4 \right\} - 1 \right) e^{-t^2/2} \left[\ln\left(1 - 3 \cdot 2^{-5/3}\tau^2 \right) + 3 \cdot 2^{-5/3}\tau^2 \right] \tau^{-4} t^4 L^4 \right] = 1 e^{-t^2/2} \left[\ln\left(1 - 3 \cdot 2^{-5/3}\tau^2 \right) + 3 \cdot 2^{-5/3}\tau^2 \right] \tau^{-4} t^4 L^4 \right]$$

for $L|t| < 2^{5/6}/\sqrt{3} = 1.0287...$ Moreover, the right-hand side of (35) is monotonically increasing with respect to L > 0 and monotonically decreasing with respect to $\gamma > 0$ and, for $L = L_{E,n}$, also with respect to $\varepsilon > 0$. The right-hand side of (35) with $L = L_{R,n}(\varepsilon, \gamma)$ does not depend on γ for $\gamma \ge \gamma_*$. (ii) For $L = L_{E,n}(\varepsilon, \gamma) \le \overline{L}_0(\varepsilon)$ and $L|t| < \tau \le \overline{\tau}_0(\varepsilon)$, we have

$$r_n(t) \le A_{\rm E}(\tau,\varepsilon,\gamma,L) \left(p_{\rm E}(t,\varepsilon,\gamma) + B(\tau,\varepsilon)Lt^4 \right) \cdot L^3 e^{-t^2/2},\tag{36}$$

in particular, with $\varepsilon = \infty$,

 \mathcal{X}

$$r_n(t) \le \exp\left\{\frac{\tau^3}{6\gamma} + \sqrt{\frac{\varkappa}{6}}\tau^3 + \tau^4 B(\tau)\right\} \left(\frac{\sqrt{6\varkappa\gamma^2 + 1}}{6\gamma} + B(\tau)L|t|\right) \cdot L^3|t|^3 e^{-t^2/2}, \quad t \in \mathbb{R},$$

where $B(\tau) := B(\tau, \infty) = (\sqrt{3} \cdot 2^{-1/6}) / (2^{5/3} - 3\tau^2)$. Moreover, the right-hand side of (36) is monotonically increasing with respect to L > 0 and monotonically decreasing with respect to $\varepsilon > 0$ and $\gamma > 0$.

For $L = L_{R,n}(\varepsilon, \gamma) \leq \overline{L}_0(\varepsilon)$ and $L|t| < \tau \leq \overline{\tau}_0(\varepsilon)$ with $B \coloneqq B(\tau, \varepsilon)$, we have

$$r_n(t) \le A_{\mathrm{R}}(\tau,\varepsilon,\gamma,L) \left(p_{\mathrm{R}}(t,\varepsilon,\gamma) + B(\tau,\varepsilon)Lt^4 \right) \cdot L^3 e^{-h(\tau,\varepsilon,L)t^2/2},\tag{37}$$

where $h(\tau, \varepsilon, L) \coloneqq 1 - \varepsilon \tau^2 L/6 > 0$ for all $\tau \in (0, \overline{\tau}_0(\varepsilon))$ if and only if $L \leq 6/(\varepsilon \overline{\tau}_0^2(\varepsilon))$. Moreover, the right-hand side of (37) is monotonically increasing with respect to L > 0 and monotonically decreasing with respect to $\gamma > 0$.

Remark 10. The right-hand sides of (35) with $L = L_{R,n}(\varepsilon, \gamma)$ and (37) are unbounded as $\varepsilon \to \infty$.

Proof. Fix arbitrary $\varepsilon, \gamma > 0$ and let $L = L_{E,n}(\varepsilon, \gamma)$ or $L = L_{R,n}(\varepsilon, \gamma)$. With $u_k \coloneqq f_k(t) - 1$ Taylor's formula and Lemma 4 yield

$$|u_k| = |f_k(t) - 1| \le \frac{t^2 \sigma_k^2}{2} \le \frac{\alpha(\varepsilon)}{2} L^2 t^2$$
(38)

for every $k = 1, \ldots, n$ and $t \in \mathbb{R}$, whence it follows that $|u_k| < 1$ for $L|t| < \sqrt{2/\alpha(\varepsilon)} =: \overline{\tau}_0(\varepsilon)$ and that $\overline{f}_n(t) = \prod_{k=1}^n (1+u_k)$ does not vanish for $L|t| < \overline{\tau}_0(\varepsilon)$. Hence, the logarithm $\ln \overline{f}_n(t)$ is well-defined for $L|t| < \overline{\tau}_0(\varepsilon)$. Considering the main branch of the logarithm and using the inequality $|e^z - 1| \le e^{|z|} - 1$, $z \in \mathbb{C}$, we get

$$r_n(t) = \left| \exp\left\{ \frac{t^2}{2} + \ln \overline{f}_n(t) \right\} - 1 \right| e^{-t^2/2} \le \left(e^{\delta_n(t)} - 1 \right) e^{-t^2/2} \le \delta_n(t) e^{\delta_n(t)} e^{-t^2/2}, \quad L|t| < \overline{\tau}_0(\varepsilon), \tag{39}$$

where

$$\begin{split} \delta_n(t) &\coloneqq \left| \frac{t^2}{2} + \ln \overline{f}_n(t) \right| = \left| \sum_{k=1}^n \left(\ln \left(1 + u_k \right) + \frac{\sigma_k^2 t^2}{2} \right) \right| \le \sum_{k=1}^n \left| \ln \left(1 + u_k \right) - u_k \right| + \left| \sum_{k=1}^n \left(u_k + \frac{\sigma_k^2 t^2}{2} \right) \right| \le \sum_{k=1}^n \sum_{j=2}^\infty \frac{|u_k|^j}{j} + \left| \sum_{k=1}^n \left(f_k(t) - 1 + \frac{\sigma_k^2 t^2}{2} \right) \right| =: I_2 + I_1. \end{split}$$

By Lemma 6 we have

 $I_1 \le p(t,\varepsilon,\gamma) \cdot L^3 \quad \text{for} \quad t \in \mathbb{R}, \ \varepsilon > 0, \ \gamma > 0,$

where $p = p_{\text{E}}$ if $L = L_{\text{E},n}(\varepsilon, \gamma)$ and $p = p_{\text{R}}$ if $L = L_{\text{R},n}(\varepsilon, \gamma)$.

To bound I_2 above, observe that inequality (38) yields

$$I_{2} \leq \sum_{k=1}^{n} |u_{k}|^{2} \sum_{j=2}^{\infty} \frac{1}{j} \left(\frac{\alpha(\varepsilon)}{2} L^{2} t^{2} \right)^{j-2} = -\frac{4}{\alpha^{2}(\varepsilon) L^{4} t^{4}} \left[\ln \left(1 - \frac{\alpha(\varepsilon)}{2} L^{2} t^{2} \right) + \frac{\alpha(\varepsilon)}{2} L^{2} t^{2} \right] \sum_{k=1}^{n} |u_{k}|^{2}$$

for $L|t| < \overline{\tau}_0(\varepsilon)$, while Lemma 5, with the account of the inequality $\sup_{0 < x \leq \varepsilon} xL_n(x) \leq L^3$, implies that

$$\sum_{k=1}^{n} |u_k|^2 = \sum_{k=1}^{n} |f_k(t) - 1|^2 \le t^4 \left[\frac{z}{2} L^3 + \frac{\alpha(\varepsilon)}{4z} L^5 \right] =: t^4 L^4 \cdot \widetilde{\alpha}_1 \left(\varepsilon, z L^{-1} \right), \quad t \in \mathbb{R}, \ z \in [0, \varepsilon],$$

where

$$\widetilde{\alpha}_1(\varepsilon, u) = \frac{u}{2} + \frac{\alpha(\varepsilon)}{4u}, \quad \varepsilon > 0, \ u > 0.$$

Choosing $u = zL^{-1} = \sqrt{\alpha(\varepsilon)/2} \wedge \varepsilon L^{-1} \coloneqq u_*(L,\varepsilon)$ to minimize $\widetilde{\alpha}_1(\varepsilon, u)$ with respect to $u \in (0, \varepsilon L^{-1}]$ we obtain $\sum_{k=1}^n |u_k|^2 \leq \alpha_1(\varepsilon, L) \cdot L^4 t^4$ with $\alpha_1(\varepsilon, L) \coloneqq \min_{0 < u \leq \varepsilon L^{-1}} \widetilde{\alpha}_1(\varepsilon, u) = \alpha_1(\varepsilon, u_*(L,\varepsilon))$ as in the formulation of the lemma. Thus, for $L|t| < \overline{\tau}_0(\varepsilon)$ we have

$$I_2 \le -\frac{4\alpha_1(\varepsilon, L)}{\alpha^2(\varepsilon)} \Big[\ln\left(1 - \frac{1}{2}\alpha(\varepsilon)L^2t^2\right) + \frac{1}{2}\alpha(\varepsilon)L^2t^2 \Big] \eqqcolon B(L|t|, \varepsilon, L) \cdot L^4t^4,$$

and

$$\delta_n(t) \le I_1 + I_2 \le p(t,\varepsilon,\gamma) \cdot L^3 + B(L|t|,\varepsilon,L) \cdot L^4 t^4, \tag{40}$$

whence, with the account of the penultimate inequality in (39), we obtain (35). The observation that $p(t, \varepsilon, \gamma)$ is monotonically decreasing with respect to γ , the functions $p_{\rm E}(t, \varepsilon, \gamma)$, $\alpha(\varepsilon)$, $\alpha_1(\varepsilon, L)$ are monotonically decreasing with respect to ε and

$$B(L|t|,\varepsilon,L) = \alpha_1(\varepsilon,L) \sum_{j=0}^{\infty} \frac{(\alpha(\varepsilon)L^2t^2/2)^j}{j+2}$$

is monotonically increasing with respect to L > 0 and monotonically decreasing with respect to $\varepsilon > 0$ yields the stated properties of the right-hand side of (35).

Now let us assume that L is small and prove slightly rougher bounds than (35) that are more convenient for analytical work. By virtue of the inequality $-(\ln(1-x) + x) = \sum_{j=2}^{\infty} \frac{x^j}{j} \le \frac{1}{2} \cdot \frac{x^2}{1-x}, \ 0 \le x < 1$, applied to $x \coloneqq \alpha(\varepsilon) L^2 t^2/2$ we have

$$B(L|t|,\varepsilon,L) \le \frac{\alpha_1(\varepsilon,L)}{2-\alpha(\varepsilon)L^2t^2} \le \frac{\alpha_1(\varepsilon,L)}{2-\alpha(\varepsilon)\tau^2} \quad \text{for} \quad L|t| < \tau \le \sqrt{2/\alpha(\varepsilon)} \eqqcolon \overline{\tau}_0(\varepsilon),$$
17

and if, in addition, $L \leq \varepsilon \sqrt{2/\alpha(\varepsilon)} =: \overline{L}_0(\varepsilon)$, then $\alpha_1(\varepsilon, L) = \sqrt{\alpha(\varepsilon)/2}$ and we also get

$$B(L|t|,\varepsilon,L) \le \frac{\sqrt{\alpha(\varepsilon)/2}}{2-\alpha(\varepsilon)\tau^2} \eqqcolon B(\tau,\varepsilon),$$

whence, with the account of (39), we obtain the factors $(p(t, \varepsilon, \gamma) + B(\tau, \varepsilon)Lt^4) \cdot L^3 e^{-t^2/2}$ in (36) and (37), which are monotonically decreasing with respect to $\gamma > 0$ and $\varepsilon > 0$ and monotonically increasing with respect to L > 0. Finally, to bound $e^{\delta_n(t)}$ in (39) we observe that for all $L|t| \leq \tau$

$$p_{\mathsf{E}}(t,\varepsilon,\gamma) = t^2 \min_{0 < z \le \varepsilon} \left[\frac{zt^2}{24} + \max\left\{ \frac{|t|}{6\gamma} + \frac{zt^2}{24}, \frac{\varkappa}{z} \right\} \right] \le L^{-2} \tau^2 \min_{0 < z \le \varepsilon} \left[\frac{zt^2}{24} + \max\left\{ \frac{\tau}{6\gamma L} + \frac{zt^2}{24}, \frac{\varkappa}{z} \right\} \right].$$

Majorizing now the maximum of two non-negative numbers by their sum we obtain

$$\begin{split} L^{3}p_{\mathrm{E}}(t,\varepsilon,\gamma) &\leq \frac{\tau^{3}}{6\gamma} + L\tau^{2} \min_{0 < z \leq \varepsilon} \left\{ \frac{\varkappa}{z} + \frac{zt^{2}}{12} \right\} = \frac{\tau^{3}}{6\gamma} + L\tau^{2} \times \begin{cases} \frac{\varepsilon t^{2}}{12} + \frac{\varkappa}{\varepsilon} \leq \sqrt{\frac{\varkappa}{12}} |t| + \frac{\varkappa}{\varepsilon}, & \varepsilon |t| \leq \sqrt{12\varkappa}, \\ \sqrt{\frac{\varkappa}{3}} |t|, & \varepsilon |t| > \sqrt{12\varkappa}, \end{cases} \leq \\ &\leq \frac{\tau^{3}}{6\gamma} + L\tau^{2} \left(\sqrt{\frac{\varkappa}{3}} |t| + \frac{\varkappa}{\varepsilon} \right) \leq \frac{\tau^{3}}{6\gamma} + \sqrt{\frac{\varkappa}{3}} \tau^{3} + \frac{\varkappa}{\varepsilon} L\tau^{2}, \end{split}$$

with the latest expression being monotonically decreasing with respect to $\gamma > 0$ and $\varepsilon > 0$ and monotonically increasing with respect to L > 0. Thus the desired estimate takes the form

$$e^{\delta_n(t)} \le \exp\left\{\frac{\tau^3}{6\gamma} + \sqrt{\frac{\varkappa}{3}}\tau^3 + \frac{\varkappa}{\varepsilon}L\tau^2 + B(\tau,\varepsilon)\tau^4\right\} \eqqcolon A_{\scriptscriptstyle \rm E}(\tau,\varepsilon,\gamma,L) \quad \text{for} \ L|t| \le \tau,$$

whence, with the account of (39) and the above constructed estimate for $\delta_n(t)$, we obtain (36). Similarly, for $L = L_{\text{R},n}$, using the explicit representation for p_{R} in the right-hand side of (33), we have

$$L^{3}p_{\mathrm{R}}(t,\varepsilon,\gamma) \leq L^{3}\left(\frac{|t|^{3}}{6\gamma}\mathbf{1}(\gamma<\gamma_{*}) + \frac{\varkappa t^{2}}{\varepsilon} + \frac{\varepsilon t^{4}}{12}\right) \leq \frac{\tau^{3}}{6\gamma}\mathbf{1}(\gamma<\gamma_{*}) + \frac{\varkappa}{\varepsilon}L\tau^{2} + \frac{\varepsilon L\tau^{2}}{12}t^{2},$$
$$e^{\delta_{n}(t)} \leq \exp\left\{\frac{\tau^{3}}{6\gamma}\mathbf{1}(\gamma<\gamma_{*}) + \frac{\varkappa}{\varepsilon}L\tau^{2} + \frac{\varepsilon L\tau^{2}}{12}t^{2} + B(\tau,\varepsilon)\tau^{4}\right\} \eqqcolon A_{\mathrm{R}}(\tau,\varepsilon,\gamma,L)e^{\varepsilon L\tau^{2}t^{2}/12} \quad \text{for} \quad L|t| \leq \tau,$$

with the right-hand side monotonically increasing with respect to L > 0 and monotonically decreasing with respect to $\gamma > 0$. Hence, with the account of (39), we obtain (37).

4 Proofs of the main results

The proof of Theorem 1 will be given simultaneously for both fractions $L = L_{E,n}(\varepsilon, \gamma)$ and $L = L_{R,n}(\varepsilon, \gamma)$ with $\varepsilon > 0$ and $\gamma > 0$ being fixed. In what follows by $C = C(\varepsilon, \gamma)$ we mean $C_{E}(\varepsilon, \gamma)$ or $C_{R}(\varepsilon, \gamma)$, sometimes omitting the arguments ε, γ . Following the outline of Zolotarev's reasoning employed in [35] we construct an upper bound for $C(\varepsilon, \gamma)$ in the form $\sup_{L>0} C(\varepsilon, \gamma, L)$, where $C(\varepsilon, \gamma, L)$ is such a function of L that the inequality

$$\Delta_n \le C(\varepsilon, \gamma, L) \cdot L^3,\tag{41}$$

holds for all $n \in \mathbb{N}$ and all distributions of independent centered r.v.'s X_1, \ldots, X_n with fixed value of the fraction of interest $L_{E,n}(\varepsilon, \gamma) = L$ or $L_{R,n}(\varepsilon, \gamma) = L$ for every L > 0. Due to the boundedness of the Kolmogorov distance $\Delta_n \leq 1$ for arbitrary d.f.'s, one can exclude from consideration large values of L. Moreover, the region of values of L to be considered can be restricted even more by use of the following sharpened upper bound for Δ_n for standardized distributions.

Lemma 7 (see [1,4]). For arbitrary r.v. X with $0 < DX < \infty$ we have

$$\sup_{x \in \mathbb{R}} \left| \mathsf{P}\left(\frac{X - \mathsf{E}X}{\sqrt{\mathsf{D}X}} < x\right) - \Phi(x) \right| \le \sup_{x > 0} \left| \frac{1}{1 + x^2} - \Phi(-x) \right| = 0.5409 \dots$$

Lemma 7 implies that, in order to prove the inequality $\Delta_n \leq C \cdot L^3$ with the constant $C \geq C_{\min} \approx 2$, say, it suffices to consider $L < (0.541/C_{\min})^{1/3} < 0.65 \approx L_1$. As this is so, we separate then the interval $L \in (0, L_1]$ into the two regions:

(i) $L \in (0, L_0]$ with some $L_0 = L_0(\varepsilon)$ small enough, where mostly analytical techniques is used;

(ii) $L \in (L_0, L_1]$, where numerical computations with the help of a computer are needed;

and construct an upper bound for C as the maximum of the two corresponding bounds and the absolute constant C_{\min} :

$$C(\varepsilon,\gamma) \le \max\left\{C_{\min}, \max_{0 < L \le L_1} C(\varepsilon,\gamma,L)\right\} = \max\left\{C_{\min}, \max_{0 < L \le L_0} C(\varepsilon,\gamma,L), \max_{L_0 < L \le L_1} C(\varepsilon,\gamma,L)\right\},$$

where, in fact, the third term turns to be extremal. However, upper bounds for the asymptotically exact constants $C_{\rm E}^*(\varepsilon,\gamma), C_{\rm R}^*(\varepsilon,\gamma)$ are obtained as limiting values of $C(\varepsilon,\gamma,L)$ as $L \to 0$.

To bound Δ_n above on each of the intervals $(0, L_0]$ and $(L_0, L_1]$, we use the method of characteristic functions realized by the Prawitz smoothing inequality.

Lemma 8 (see [28]). For arbitrary d.f. F with the ch.f. f and for all $0 < T_0 \leq T_1$ we have

$$\begin{split} \sup_{x \in \mathbb{R}} |F(x) - \Phi(x)| &\leq \frac{2}{T_1} \int_0^{T_0} \left| K\left(\frac{t}{T_1}\right) \right| \cdot \left| f(t) - e^{-t^2/2} \right| dt + \frac{2}{T_1} \int_{T_0}^{T_1} \left| K\left(\frac{t}{T_1}\right) \right| \cdot |f(t)| \, dt + \\ &+ \frac{2}{T_1} \int_0^{T_0} \left| K\left(\frac{t}{T_1}\right) - \frac{iT_1}{2\pi t} \right| e^{-t^2/2} dt + \frac{1}{\pi} \int_{T_0}^{\infty} e^{-t^2/2} \frac{dt}{t}, \end{split}$$

where

$$K(t) = \frac{1}{2} \left(1 - |t| \right) + \frac{i}{2} \left[\left(1 - |t| \right) \cot \pi t + \frac{\operatorname{sgn} t}{\pi} \right], \quad t \in (-1, 1) \setminus \{0\}.$$

and $K(\pm 1) \coloneqq 0$ defined by continuity. Moreover, the function K(t) for all $t \in [-1,1] \setminus \{0\}$ satisfies

$$|K(t)| \le \frac{1.0253}{2\pi |t|}, \quad \left|K(t) - \frac{i}{2\pi t}\right| \le \frac{1}{2} \left(1 - |t| + \frac{\pi^2 t^2}{18}\right) \le \frac{1}{2}.$$
(42)

By Lemma 8 we have

$$\Delta_n \le I_1 + I_2 + I_3 + I_4, \tag{43}$$

where

$$I_{1} = \frac{2}{T_{1}} \int_{0}^{T_{0}} \left| K\left(\frac{t}{T_{1}}\right) \right| \cdot \left| \overline{f}_{n}(t) - e^{-t^{2}/2} \right| dt, \quad I_{2} = \frac{2}{T_{1}} \int_{T_{0}}^{T_{1}} \left| K\left(\frac{t}{T_{1}}\right) \right| \cdot \left| \overline{f}_{n}(t) \right| dt,$$
$$I_{3} = \frac{2}{T_{1}} \int_{0}^{T_{0}} \left| K\left(\frac{t}{T_{1}}\right) - \frac{iT_{1}}{2\pi t} \right| e^{-t^{2}/2} dt, \quad I_{4} = \frac{1}{\pi} \int_{T_{0}}^{\infty} e^{-t^{2}/2} \frac{dt}{t}, \quad 0 < T_{0} \leq T_{1}.$$

In what follows we use the notation

$$T_0(\tau_0, L) \coloneqq \frac{\tau_0}{L}, \quad T_1(\tau_1, L) \coloneqq \frac{\tau_1}{L^3}, \quad \text{where} \quad \tau_0 \in (0, \overline{\tau}_0(\varepsilon)), \quad \tau_1 \in (0, \overline{\tau}_1)$$

are free parameters to be chosen below with $\overline{\tau}_1 \coloneqq \pi/4$, $\overline{\tau}_0(\varepsilon) \coloneqq \sqrt{2/\alpha(\varepsilon)}$ defined in Theorems 4, 5, respectively.

4.1 The case (i) $L \in (0, L_0]$ and the proof of Theorem 2

The purpose of the present subsection is to bound $C(\varepsilon, \gamma, L)$ above for $L \in (0, L_0]$ by an increasing function of L with the best possible (within the method used) limiting value as $L \to 0$.

Let $L_0 > 0$ satisfy the conditions of Theorem 4 (ii) and Theorem 5 (ii):

$$L_0 < (\varepsilon/4)^{1/3}$$
 and $L_0 \le \overline{L}_0(\varepsilon) \coloneqq \varepsilon \sqrt{2/\alpha(\varepsilon)} = \varepsilon \overline{\tau}_0(\varepsilon),$

so that for $\varepsilon \ge 0.0557$ arbitrary $L_0 \le 0.03$ surely fits. For $L = L_{\text{R},n}$ we additionally assume that $L_0 \le 6/(\varepsilon \overline{\tau}_0^2(\varepsilon)) = 3\alpha(\varepsilon)/\varepsilon$, which provides the positiveness of the exponent in the right-hand side of (37). Since $\alpha(\varepsilon) \ge 3 \cdot 2^{-2/3}$, this assumption is surely satisfied if

$$\varepsilon L_0 \le 9 \cdot 2^{-2/3} = 5.6696 \dots$$

in particular, for $\varepsilon \leq 188$ we may consider arbitrary $L_0 \leq 0.03$.

Let us describe the process of estimation of each term in (43) assuming that $T_0 \coloneqq T_0(\tau_0, L), T_1 \coloneqq T_1(\tau_1, L)$ with τ_1 also satisfying the condition of Theorem 4 (ii): $\tau_1 \ge \pi L_0^3/\varepsilon \rightleftharpoons \underline{\tau}_1(L_0, \varepsilon)$. Adding and substracting $(iT_1)/(2\pi t)$ from $K(\frac{t}{T_1})$ under the modulus sign in the integrand in I_1 and applying

Adding and substracting $(iT_1)/(2\pi t)$ from $K\left(\frac{t}{T_1}\right)$ under the modulus sign in the integrand in I_1 and applying then the inequality $\left|K\left(\frac{t}{T_1}\right) - \frac{iT_1}{2\pi t}\right| \leq \frac{1}{2}, |t| \leq T_1$, from Lemma 8, we obtain

$$I_{1} \leq \frac{1}{\pi} \int_{0}^{T_{0}} \frac{r_{n}(t)}{t} dt + \frac{2}{T_{1}} \int_{0}^{T_{0}} \left| K\left(\frac{t}{T_{1}}\right) - \frac{iT_{1}}{2\pi t} \right| r_{n}(t) dt \leq I_{11} + I_{12},$$

where

$$r_n(t) \coloneqq \left| \overline{f}_n(t) - e^{-t^2/2} \right|, \quad I_{11} \coloneqq \frac{1}{\pi} \int_0^{T_0} \frac{r_n(t)}{t} \, dt, \quad I_{12} \coloneqq \frac{1}{T_1} \int_0^{T_0} r_n(t) \, dt.$$

Further we use inequalities (36) and (37) from Theorem 5 to estimate the integrands $r_n(t)$ in I_{11} and I_{12} and enlarge then the region of integration from $(0, T_0)$ to $(0, \infty)$. With the definitions of the upper and the lower incomplete gamma-functions yielding

$$\int_{x}^{\infty} t^{s} e^{-kt^{2}} dt = \frac{k^{-\frac{s+1}{2}}}{2} \Gamma\left(\frac{s+1}{2}, kx^{2}\right), \quad \int_{0}^{x} t^{s} e^{-kt^{2}} dt = \frac{k^{-\frac{s+1}{2}}}{2} \Upsilon\left(\frac{s+1}{2}, kx^{2}\right), \quad s, k, x > 0, \tag{44}$$

• for the Esseen-type fraction $L = L_{E,n}(\varepsilon, \gamma)$ with arbitrary $\varepsilon, \gamma > 0$ we have

$$\begin{split} \frac{I_{11}}{L^3} &\leq \frac{A_{\rm E}}{\pi} \bigg[\int_0^{t_\gamma/\varepsilon} \left(\frac{\varkappa t}{\varepsilon} + \frac{\varepsilon t^3}{24}\right) e^{-t^2/2} dt + \frac{\sqrt{6\varkappa\gamma^2 + 1}}{6\gamma} \int_{t_\gamma/\varepsilon}^{\infty} t^2 e^{-t^2/2} dt + BL \int_0^{\infty} t^3 e^{-t^2/2} dt \bigg] = \\ &= \frac{A_{\rm E}}{\pi} \bigg[\frac{\varkappa}{\varepsilon} \Upsilon \Big(1, \frac{t_\gamma^2}{2\varepsilon^2} \Big) + \frac{\varepsilon}{12} \Upsilon \Big(2, \frac{t_\gamma^2}{2\varepsilon^2} \Big) + \frac{\sqrt{2(6\varkappa\gamma^2 + 1)}}{6\gamma} \Gamma \Big(\frac{3}{2}, \frac{t_\gamma^2}{2\varepsilon^2} \Big) + 2BL \bigg], \end{split}$$

$$\begin{split} \frac{I_{12}}{L^6} &\leq \frac{A_{\rm E}}{\tau_1} \bigg[\int_0^{t_{\gamma}/\varepsilon} \left(\frac{\varkappa t^2}{\varepsilon} + \frac{\varepsilon t^4}{24} \right) e^{-t^2/2} dt + \frac{\sqrt{6\varkappa\gamma^2 + 1}}{6\gamma} \int_{t_{\gamma}/\varepsilon}^{\infty} t^3 e^{-t^2/2} dt + BL \int_0^{\infty} t^4 e^{-t^2/2} dt \bigg] = \\ &= \frac{A_{\rm E}}{\tau_1} \bigg[\frac{\sqrt{2}\varkappa}{\varepsilon} \,\Upsilon \Big(\frac{3}{2}, \frac{t_{\gamma}^2}{2\varepsilon^2} \Big) + \frac{\sqrt{2}\varepsilon}{6} \Upsilon \Big(\frac{5}{2}, \frac{t_{\gamma}^2}{2\varepsilon^2} \Big) + \frac{\sqrt{6\varkappa\gamma^2 + 1}}{3\gamma} \Gamma \Big(2, \frac{t_{\gamma}^2}{2\varepsilon^2} \Big) + \frac{3}{2} \sqrt{2\pi} BL \bigg], \end{split}$$

where $A_{\rm E} = A_{\rm E}(\tau_0, \varepsilon, \gamma, L), B = B(\tau_0, \varepsilon);$

• in particular, for the Esseen-type fraction $L = L_{E,n}(\infty, \gamma)$ with $\varepsilon = \infty$ we obtain

$$\frac{I_{11}}{L^3} \le \frac{A_{\rm E}}{\pi} \left[\frac{\sqrt{2\pi (6\varkappa\gamma^2 + 1)}}{12\gamma} + 2BL \right], \quad \frac{I_{12}}{L^6} \le \frac{A_{\rm E}}{\tau_1} \left[\frac{\sqrt{6\varkappa\gamma^2 + 1}}{3\gamma} + \frac{3}{2}\sqrt{2\pi}BL \right];$$

• for the Rozovskii-type fraction $L = L_{R,n}(\varepsilon, \gamma)$ with arbitrary $\varepsilon, \gamma > 0$ we have

$$\begin{split} \frac{I_{11}}{L^3} &\leq \frac{A_{\mathrm{R}}}{\pi} \bigg[\int_0^{t_{1,\gamma}/\varepsilon} \left(\frac{\varkappa t}{\varepsilon} + \frac{\varepsilon t^3}{24} \right) e^{-ht^2/2} dt + \frac{1}{6\gamma} \int_{t_{1,\gamma}/\varepsilon}^{t_{2,\gamma}/\varepsilon} t^2 e^{-ht^2/2} dt + \frac{\varepsilon}{12} \int_{t_{2,\gamma}/\varepsilon}^{\infty} t^3 e^{-ht^2/2} dt + \\ &\quad + BL \int_0^{\infty} t^3 e^{-ht^2/2} dt \bigg] = \frac{A_{\mathrm{R}}}{\pi} \bigg[\frac{\varkappa}{\varepsilon h} \Upsilon \bigg(1, \frac{ht_{1,\gamma}^2}{2\varepsilon^2} \bigg) + \frac{\varepsilon}{12h^2} \Upsilon \bigg(2, \frac{ht_{1,\gamma}^2}{2\varepsilon^2} \bigg) + \\ &\quad + \frac{\sqrt{2}}{6\gamma h^{3/2}} \bigg(\frac{\sqrt{\pi}}{2} - \Upsilon \bigg(\frac{3}{2}, \frac{ht_{1,\gamma}^2}{2\varepsilon^2} \bigg) - \Gamma \bigg(\frac{3}{2}, \frac{ht_{2,\gamma}^2}{2\varepsilon^2} \bigg) \bigg) + \frac{\varepsilon}{6h^2} \Gamma \bigg(2, \frac{ht_{2,\gamma}^2}{2\varepsilon^2} \bigg) + \frac{2BL}{h^2} \bigg], \end{split}$$

$$\begin{split} \frac{I_{12}}{L^6} &\leq \frac{A_{\rm R}}{\tau_1} \bigg[\int_0^{t_{1,\gamma}/\varepsilon} \left(\frac{\varkappa t^2}{\varepsilon} + \frac{\varepsilon t^4}{24} \right) e^{-ht^2/2} dt + \frac{1}{6\gamma} \int_{t_{1,\gamma}/\varepsilon}^{t_{2,\gamma}/\varepsilon} t^3 e^{-ht^2/2} dt + \frac{\varepsilon}{12} \int_{t_{2,\gamma}/\varepsilon}^{\infty} t^4 e^{-ht^2/2} dt + \\ &+ BL \int_0^{\infty} t^4 e^{-ht^2/2} dt \bigg] = \frac{A_{\rm R}}{\tau_1} \bigg[\frac{\sqrt{2}\varkappa}{\varepsilon h^{3/2}} \Upsilon \bigg(\frac{3}{2}, \frac{ht_{1,\gamma}^2}{2\varepsilon^2} \bigg) + \frac{\sqrt{2}\varepsilon}{12h^{5/2}} \Upsilon \bigg(\frac{5}{2}, \frac{ht_{1,\gamma}^2}{2\varepsilon^2} \bigg) + \\ &+ \frac{1}{3\gamma h^2} \bigg(1 - \Upsilon \bigg(2, \frac{ht_{1,\gamma}^2}{2\varepsilon^2} \bigg) - \Gamma \bigg(2, \frac{ht_{2,\gamma}^2}{2\varepsilon^2} \bigg) \bigg) + \frac{\sqrt{2}\varepsilon}{6h^{5/2}} \Gamma \bigg(\frac{5}{2}, \frac{ht_{2,\gamma}^2}{2\varepsilon^2} \bigg) + \frac{3\sqrt{2\pi}BL}{2h^{5/2}} \bigg], \end{split}$$

where $A_{\rm R} = A_{\rm R}(\tau_0, \varepsilon, \gamma, L)$, $B = B(\tau_0, \varepsilon)$, $h = h(\tau_0, \varepsilon, L)$ provided that $\varepsilon L_0 < 9 \cdot 2^{-2/3} = 5.6696 \dots$; • in particular, for the Rozovskii-type fraction $L = L_{\rm R,n}(\varepsilon, \gamma)$ with $\gamma \ge \gamma_*$, taking into account that $t_{1,\gamma} = t_{2,\gamma} = t_{\infty} = 2/\gamma_*$, we obtain

$$\frac{I_{11}}{L^3} \leq \frac{A_{\rm R}}{\pi} \bigg[\frac{\varkappa}{\varepsilon h} \Upsilon \bigg(1, \frac{2h}{(\varepsilon\gamma_*)^2} \bigg) + \frac{\varepsilon}{12h^2} \bigg(1 + \Gamma \bigg(2, \frac{2h}{(\varepsilon\gamma_*)^2} \bigg) \bigg) + \frac{2BL}{h^2} \bigg],$$
$$\frac{I_{12}}{L^6} \leq \frac{A_{\rm R}}{\tau_1} \bigg[\frac{\sqrt{2}\varkappa}{\varepsilon h^{3/2}} \Upsilon \bigg(\frac{3}{2}, \frac{2h}{(\varepsilon\gamma_*)^2} \bigg) + \frac{\sqrt{2}\varepsilon}{12h^{5/2}} \bigg(\frac{3}{4} \sqrt{\pi} + \Gamma \bigg(\frac{5}{2}, \frac{2h}{(\varepsilon\gamma_*)^2} \bigg) \bigg) + \frac{3\sqrt{2\pi}BL}{2h^{5/2}} \bigg].$$

Note that the constructed upper bounds for I_{11}/L^3 and I_{12}/L^6 are monotonically increasing with respect to L, monotonically decreasing with respect to $\gamma > 0$ and, for $L = L_{E,n}$, also with respect to $\varepsilon > 0$ as integrals of the functions possessing the stated properties. Moreover, they do not depend on γ for $L = L_{R,n}(\varepsilon, \gamma)$ with $\gamma \ge \gamma_*$, so that the further increase of γ has no effect on the value of the resulting constant $C(\varepsilon, \gamma, L)$, hence, the value $\gamma = \gamma_*$ is an optimal one.

To estimate I_2 , we use the first inequality in (42) from Lemma 8 and bound (27) from Theorem 4 to get

$$\frac{I_2}{L^3} \le \frac{1.0253}{\pi L^3} \int_{T_0}^{T_1} \frac{\left|\overline{f}_n(t)\right|}{t} dt \le \frac{1.0253}{\pi L^3 T_0^3} \int_{T_0}^{\infty} t^2 e^{-k(\tau_1)t^2} dt = \frac{1.0253}{2\pi \tau_0^3 \left(k(\tau_1)\right)^{3/2}} \Gamma\left(\frac{3}{2}, \frac{k(\tau_1)\tau_0^2}{L^2}\right),$$

with $k(\tau_1)$ defined in the formulation of Theorem 4 (recall that $k(\tau_1) > 0$ iff $0 < \tau_1 < \overline{\tau}_1$). We also observe that the constructed upper bound for I_2/L^3 holds true for both fractions $L = L_{\text{E},n}(\varepsilon, \gamma)$ and $L = L_{\text{R},n}(\varepsilon, \gamma)$, is independent of ε and γ , and is monotonically increasing with respect to L, moreover, $I_2 = \mathcal{O}(L^{\nu})$ as $L \to 0$ for every $\nu > 0$.

Using the inequality $\left|K\left(\frac{t}{T_1}\right) - \frac{iT_1}{2\pi t}\right| \leq \frac{1}{2}$ from Lemma 8 once again to bound I_3 and the condition $1 \leq t^3/T_0^3$ defining the region of integration in I_4 , we estimate the sum of the integrals I_3 and I_4 as

$$\frac{I_3 + I_4}{L^3} \le \frac{1}{\tau_1} \int_0^\infty e^{-t^2/2} dt + \frac{1}{\pi L^3 T_0^3} \int_{T_0}^\infty t^2 e^{-t^2/2} dt = \frac{1}{\tau_1} \sqrt{\frac{\pi}{2}} + \frac{\sqrt{2}}{\pi \tau_0^3} \Gamma\left(\frac{3}{2}, \frac{\tau_0^2}{2L^2}\right)$$

with the latest expression being monotonically increasing with respect to L and independent of ε and γ .

Summing up the obtained bounds for the integrals $I_{11}, I_{12}, I_2, I_3, I_4$, we obtain an upper bound for Δ_n in the form

$$\Delta_n \le L^3 \cdot C_0(\varepsilon, \gamma, L, \tau_0, \tau_1) \tag{45}$$

with the function $C_0(\varepsilon, \gamma, L, \tau_0, \tau_1)$ being monotonically increasing with respect to L, monotonically decreasing with respect to $\gamma > 0$ and, for $L = L_{E,n}(\varepsilon, \gamma)$, also with respect to $\varepsilon > 0$ and satisfying

$$\begin{split} C_{0}(\varepsilon,\gamma,\tau_{0},\tau_{1}) &\coloneqq \lim_{L \to 0} C_{0}(\varepsilon,\gamma,L,\tau_{0},\tau_{1}) = \frac{1}{\tau_{1}}\sqrt{\frac{\pi}{2}} + \\ &+ \begin{cases} \frac{A_{\text{E}}(\tau_{0},\varepsilon,\gamma,0+)}{\pi} \Big[\frac{\varkappa}{\varepsilon} \Upsilon\left(1,\frac{t_{\gamma}^{2}}{2\varepsilon^{2}}\right) + \frac{\varepsilon}{12} \Upsilon\left(2,\frac{t_{\gamma}^{2}}{2\varepsilon^{2}}\right) + \frac{\sqrt{2(6\varkappa\gamma^{2}+1)}}{6\gamma} \Gamma\left(\frac{3}{2},\frac{t_{\gamma}^{2}}{2\varepsilon^{2}}\right) \Big], \quad L = L_{\text{E},n}(\varepsilon,\gamma), \\ \frac{A_{\text{R}}(\tau_{0},\varepsilon,\gamma,0+)}{\pi} \Big[\frac{\varkappa}{\varepsilon\hbar} \Upsilon\left(1,\frac{ht_{1,\gamma}^{2}}{2\varepsilon^{2}}\right) + \frac{\varepsilon}{12h^{2}} \Upsilon\left(2,\frac{ht_{1,\gamma}^{2}}{2\varepsilon^{2}}\right) + \frac{\varepsilon}{6\hbar^{2}} \Gamma\left(2,\frac{ht_{2,\gamma}^{2}}{2\varepsilon^{2}}\right) + \\ &+ \frac{\sqrt{2}}{6\gamma\hbar^{3/2}} \Big(\frac{\sqrt{\pi}}{2} - \Upsilon\left(\frac{3}{2},\frac{ht_{1,\gamma}^{2}}{2\varepsilon^{2}}\right) - \Gamma\left(\frac{3}{2},\frac{ht_{2,\gamma}^{2}}{2\varepsilon^{2}}\right) \Big], \qquad L = L_{\text{R},n}(\varepsilon,\gamma), \end{split}$$

where $h = h(\tau_0, \varepsilon, L) \coloneqq 1 - \varepsilon \tau_0^2 L/6$.

Inequality (45) yields the following upper bound for the function $C(\varepsilon, \gamma, L)$ from (41):

$$C(\varepsilon,\gamma,L) \le C_0(\varepsilon,\gamma,L) \coloneqq \inf \left\{ C_0(\varepsilon,\gamma,L,\tau_0,\tau_1) \colon \tau_0 \in \left(0,\overline{\tau}_0(\varepsilon)\right), \ \tau_1 \in \left(\underline{\tau}_1(L_0,\varepsilon),\overline{\tau}_1\right), \ \tau_1 \ge L^2 \tau_0 \right\}$$
(46)

for every $\varepsilon > 0$, $\gamma > 0$, and $L \leq L_0$ with

$$L_0 \leq \overline{L}_0(\varepsilon) \wedge (\varepsilon/4)^{1/3}$$
, also $L_0 \leq 9 \cdot 2^{-2/3} \varepsilon^{-1}$ for $L = L_{\mathbb{R},n}$, (47)

and $C_0(\varepsilon, \gamma, L)$ being a monotonically increasing function of L as the greatest lower bound to the increasing function $C_0(\varepsilon, \gamma, L, \tau_0, \tau_1)$, where the greatest lower bound is taken over a decreasing system of sets. Hence,

$$\max_{0 < L \le L_0} C(\varepsilon, \gamma, L) \le \max_{0 < L \le L_0} C_0(\varepsilon, \gamma, L) = C_0(\varepsilon, \gamma, L_0).$$

Moreover, $C_0(\varepsilon, \gamma, L)$ is monotonically decreasing with respect to $\gamma > 0$ and, for $L = L_{E,n}(\varepsilon, \gamma)$, also with respect to $\varepsilon > 0$.

Inequality (45) yields upper bounds for the asymptotically exact constants $C^* \in \{C_{\rm E}^*, C_{\rm R}^*\}$ defined in (13), (14). Namely, with the observation that conditions (47) are trivially satisfied for every $\varepsilon > 0$ if $L \to 0$, we have

$$C^*(\varepsilon,\gamma) \le \lim_{L \to 0} C(\varepsilon,\gamma,L) \le C_0(\varepsilon,\gamma,0+) \le \inf \left\{ C_0(\varepsilon,\gamma,\tau_0,\tau_1) \colon \tau_0 \in \left(0,\overline{\tau}_0(\varepsilon)\right), \ \tau_1 \in \left(0,\overline{\tau}_1\right) \right\} = C^*(\varepsilon,\gamma)$$

 $= C_0(\varepsilon, \gamma, 0+, \overline{\tau}_1 -)$ for every $\varepsilon, \gamma > 0$.

Hence, recalling that $\overline{\tau}_1 = \pi/4$, $A_{\text{E}}(0+,\varepsilon,\gamma,0+) = A_{\text{R}}(0+,\varepsilon,\gamma,0+) = A(0+,\varepsilon) = 1$, $h(\tau_0,\varepsilon,0+) = 1$, for every $\varepsilon > 0$ and $\gamma > 0$ we obtain:

• in the Esseen-type inequality

$$C_{\rm E}^*(\varepsilon,\gamma) \le C_0\big(\varepsilon,\gamma,0+,\overline{\tau}_1-\big) = \frac{4}{\sqrt{2\pi}} + \frac{1}{\pi} \Big[\frac{\varkappa}{\varepsilon} \Upsilon\big(1,\frac{t_\gamma^2}{2\varepsilon^2}\big) + \frac{\varepsilon}{12} \Upsilon\big(2,\frac{t_\gamma^2}{2\varepsilon^2}\big) + \frac{\sqrt{2(6\varkappa\gamma^2+1)}}{6\gamma} \Gamma\big(\frac{3}{2},\frac{t_\gamma^2}{2\varepsilon^2}\big) \Big] = \widehat{C_{\rm E}^*}(\varepsilon,\gamma),$$

• in the Rozovskii-type inequality

$$\begin{split} C^*_{\scriptscriptstyle \rm E}(\varepsilon,\gamma) &\leq C_0\big(\varepsilon,\gamma,0+,\overline{\tau}_1-\big) = \frac{4}{\sqrt{2\pi}} + \frac{1}{\pi}\Big[\frac{\varkappa}{\varepsilon}\Upsilon\Big(1,\frac{t_{1,\gamma}^2}{2\varepsilon^2}\Big) + \frac{\varepsilon}{12}\Upsilon\Big(2,\frac{t_{1,\gamma}^2}{2\varepsilon^2}\Big) + \frac{\varepsilon}{6}\Gamma\Big(2,\frac{t_{2,\gamma}^2}{2\varepsilon^2}\Big) + \\ &+ \frac{\sqrt{2}}{6\gamma}\Big(\frac{\sqrt{\pi}}{2} - \Upsilon\Big(\frac{3}{2},\frac{t_{1,\gamma}^2}{2\varepsilon^2}\Big) - \Gamma\Big(\frac{3}{2},\frac{t_{2,\gamma}^2}{2\varepsilon^2}\Big)\Big)\Big] = \widehat{C^*_{\scriptscriptstyle \rm R}}(\varepsilon,\gamma), \end{split}$$

with $\widehat{C}_{E}^{*}(\varepsilon,\gamma), \widehat{C}_{R}^{*}(\varepsilon,\gamma)$ defined in the formulation of Theorem 2 (see (15), (16)). Moreover, the function $\widehat{C}_{E}^{*}(\varepsilon,\gamma)$ decreases with respect to $\varepsilon > 0$ and $\gamma > 0$ as an integral and then a limit of a function with the similar properties and is unbounded as $\varepsilon \to 0+$ or $\gamma \to 0+$. Similarly, the function $\widehat{C}^*_{\mathbb{R}}(\varepsilon,\gamma)$ decreases with respect to $\gamma > 0$ being constant for $\gamma \geq \gamma_*$ for every fixed $\varepsilon > 0$ and infinitely grows as $\varepsilon \to 0+$, $\varepsilon \to \infty$, or $\gamma \to 0+$.

The case (ii) $L \in [L_0, L_1]$ 4.2

Let $0 < T_0 \leq T_0(\overline{\tau}_0(\varepsilon), L)$, $T_0 \leq T_1 \leq T_1(\overline{\tau}_1, L)$, and $0 < L_0 \leq L \leq L_1 < \infty$. Though the function K(t) has a singularity of order $\mathcal{O}(|t|^{-1})$ as $t \to 0$, the integrands in I_1 and I_3 have no singularities due to the presence of the factor $r_n(t) = \mathcal{O}(t^2), t \to 0$, in the integrand in I_1 and to the boundedness of the function $|K(\frac{t}{T_1}) - \frac{iT_1}{2\pi t}|$ by (42) in I_3 . Using estimates (35) and (26) from Theorems 5 and 4 to bound integrands in I_1 and I_2 , we obtain, by (43),

an upper bound for Δ_n in the form

$$\Delta_n \le D(\varepsilon, \gamma, L, T_0, T_1),$$

which is uniform in the class of all distributions of random summands with fixed value of the fraction under consideration $L \in [L_0, L_1]$, where $T_0 \leq T_1$ are free parameters. Moreover, the function $D(\varepsilon, \gamma, L, T_0, T_1)$ here is monotonically increasing with respect to L, monotonically decreasing with respect to $\gamma > 0$ and, for $L = L_{\text{E},n}$, also with respect to $\varepsilon > 0$. Hence, we may construct an upper bound for $C(\varepsilon, \gamma, L)$ on the interval $L_0 \leq L \leq L_1$ in the form

$$C(\varepsilon,\gamma,L) \le C_1(\varepsilon,\gamma,L) := \inf\left\{\frac{D(\varepsilon,\gamma,L,T_0,T_1)}{L^3} : 0 < T_0 < \frac{\overline{\tau}_0(\varepsilon)}{L}, \ T_0 < T_1 < \frac{\overline{\tau}_1}{L^3}\right\},\tag{48}$$

where $\overline{\tau}_1 \coloneqq \pi/4$ and $\overline{\tau}_0(\varepsilon) \coloneqq \sqrt{2/\alpha(\varepsilon)}$ are defined in Theorems 4 and 5, respectively. The maximum value $\max_{L_0 \leq L \leq L_1} C_1(\varepsilon, \gamma, L)$ can be estimated similarly to [35] by computation of $C_1(\varepsilon, \gamma, L)$ in a finite number of points using the inequality

$$\max_{L' \leq L \leq L''} C_1(\varepsilon, \gamma, L) \leq C_1(\varepsilon, \gamma, L'') \cdot \left(L''/L'\right)^3,$$

which is valid due to the monotone growth of the function $C_1(\varepsilon, \gamma, L) \cdot L^3$ as the greatest lower bound to the monotonically increasing function $D(\varepsilon, \gamma, L, T_0, T_1)$ over a decreasing system of sets. Furthermore, since the function $D(\varepsilon, \gamma, L, T_0, T_1)$ is monotonically decreasing with respect to γ and, for $L = L_{E,n}$, with respect to ε , so is $\max_{L_0 \le L \le L_1} C_1(\varepsilon, \gamma, L).$

4.3 Numerical results

Summarizing what was said above, as the constants $C(\varepsilon, \gamma) \in \{C_{\rm E}(\varepsilon, \gamma), C_{\rm R}(\varepsilon, \gamma)\}$ in inequalities (11) and (12) we can take

$$C(\varepsilon,\gamma) \coloneqq \max\left\{C_{\min}, \sup_{0 < L \le L_1} C(\varepsilon,\gamma,L)\right\} \text{ with } C(\varepsilon,\gamma,L) \coloneqq \begin{cases} C_0(\varepsilon,\gamma,L), & 0 < L \le L_0(\varepsilon), \\ C_1(\varepsilon,\gamma,L), & L_0(\varepsilon) < L \le L_1, \end{cases}$$

 $C_{\min} \coloneqq 2, L_1 \coloneqq 0.65 > (0.541/C_{\min})^{1/3}, L_0(\varepsilon) \coloneqq (\varepsilon/4)^{1/3} \wedge \varepsilon \overline{\tau}_0(\varepsilon) \text{ if } L = L_{\varepsilon,n} \text{ and } L_0(\varepsilon) \coloneqq (\varepsilon/4)^{1/3} \wedge \varepsilon \overline{\tau}_0(\varepsilon) \wedge 6/(\varepsilon \overline{\tau}_0^2(\varepsilon)) \text{ if } L = L_{\varepsilon,n}, \text{ for every } \varepsilon, \gamma > 0.$ Since $C_0(\varepsilon, \gamma, L)$ and $C_1(\varepsilon, \gamma, L)$ are both monotonically decreasing with respect to $\gamma > 0$ and, for $L = L_{\varepsilon,n}$, also with respect to ε , so is $C(\varepsilon, \gamma)$.

The concrete numerical values of $C_0(\varepsilon, \gamma, L)$ and $C_1(\varepsilon, \gamma, L)$ are processed with the help of a computer. Our computations were carried out in Python 3.6 using the library Scipy 1.0.0. The values of $\max_{0 \le L \le L_0} C_0(\varepsilon, \gamma, L) = C_0(\varepsilon, \gamma, L_0)$ for some ε and γ with $L_0 = 0.001$ and $L_0 = 0.03$ are given for the Esseen-type fraction $L = L_{\text{E},n}(\varepsilon, \gamma)$ in table 2 in the fourth and seventh columns, respectively, accompanied by the optimal values of the parameters τ_0 and τ_1 in (46). The values of $\max_{0.03 \le L \le L_1} C_1(\varepsilon, \gamma, L) = C_1(\varepsilon, \gamma, L_*)$ for some ε and γ are given in table 4 for the Esseen fraction $L = L_{\text{E},n}(\varepsilon, \gamma)$ accompanied by the optimal values of the parameters T_0 and T_1 in (48) specified in the form $\tau_0 \coloneqq T_0 L$ and $\tau_1 \coloneqq T_1 L^3$, in the extremal point $L = L_*$ which is also given in the fourth column of the same table. Columns 7–10 contain the normalized contributions of the integrals I_k/L_*^3 , k = 1, 2, 3, 4, into the extreme value $C_1(\varepsilon, \gamma, L_*)$, so that $C_1(\varepsilon, \gamma, L_*) = (I_1 + I_2 + I_3 + I_4)/L_*^3$. Tables 3 and 5 contain similar results for the Rozovskii-type fraction $L = L_{\text{R},n}(\varepsilon, \gamma)$.

For example, for the minimum value $C_{\rm E}(\infty,\infty)$ of the constant $C_{\rm E}(\varepsilon,\gamma)$ in the Esseen-type inequality (11) we have

$$\sup_{0 < L \le 0.03} C(\infty, \infty, L) \le C_0(\infty, \infty, 0.03) \le 2.28, \quad \sup_{0.03 < L \le L_1} C(\infty, \infty, L) \le C_1(\infty, \infty, 0.4833...) \le 2.65$$

(the plot of the function $C_1(\infty, \infty, L)$ for $L := L_{\mathbb{E},n}(\infty, \infty) \in [0.03, L_1]$ is given on Fig. 4 (left)), hence,

$$\inf_{\varepsilon > 0, \gamma > 0} C_{\scriptscriptstyle\rm E}(\varepsilon, \gamma) = C_{\scriptscriptstyle\rm E}(\infty, \infty) \le \max\{C_{\rm min}, \, 2.28, \, 2.65\} = 2.65.$$

However, the same upper bound $C_{\text{E}}(\varepsilon, \gamma) \leq 2.65$ can be reached, due to the rounding gap, already for finite values of ε and $\gamma = \gamma(\varepsilon)$ plotted on Fig. 2 (right) and also at the points with one infinite component, for example,

 $(\varepsilon, \gamma) \in \{(2.56, \infty), (2.74, 3), (4, 1.62), (\infty, 1.43)\},\$

for which, due to the monotonicity and according to tables 2, 4, we have

$$\max \left\{ C(2.56, \infty, L), C(2.74, 3, L), C(4, 1.62, L), C(\infty, 1.43, L) \right\} = \int C_0(0.6, 0.3, 0.03) \le 2.64, \qquad L \le 0.03 \quad (\text{see table } 2),$$

$$- \left\{ \max \left\{ C_1(2.56, \infty, 0.4833 \dots), \dots \right\} \le 2.65, \quad 0.03 < L \le L_1 \quad \text{(see table 4)}. \right\}$$

Similar level curve $\{(\varepsilon, \gamma) : \widehat{C}^*_{\scriptscriptstyle E}(\varepsilon, \gamma) = 1.72\}$ of the upper bound $\widehat{C}^*_{\scriptscriptstyle E}(\varepsilon, \gamma) \coloneqq C_0(\varepsilon, \gamma, 0+)$ for the asymptotically exact constant $C^*_{\scriptscriptstyle E}(\varepsilon, \gamma)$ is plotted on Fig. 2 (left).

Since the constructed upper bound for $C_{\text{R}}(\varepsilon, \gamma)$ is monotonically decreasing with respect to γ and is constant for $\gamma \geq \gamma_*$, its global minimum with respect to γ is attained at $\gamma = \gamma_* = 0.5599...$. Plot of the function $\max_{0.03 \leq L \leq L_1} C_1(\varepsilon, \gamma_*, L)$ with respect to ε is given on Fig. 3 (right, solid line), and numerical computations show that its minimum is attained around the point $\varepsilon = 2.12$, for which, according to tables 5 and 3, we have

$$C_0(2.12, \gamma_*, 0.03) \le 2.29, \quad \max_{0.03 \le L \le L_1} C_1(2.12, \gamma_*, L) = C_1(2.12, \gamma_*, 0.4827...) \le 2.66$$

(the plot of the function $C_1(2.12, \gamma_*, L)$ for $L \coloneqq L_{\mathbb{R},n}(2.12, \gamma_*) \in [0.03, L_1]$ is given on Fig. 4 (right)). Hence,

$$\inf_{\varepsilon > 0, \gamma > 0} C_{\mathbf{R}}(\varepsilon, \gamma) \le C_{\mathbf{R}}(2.12, \gamma_*) \le \max\{C_{\min}, 2.29, 2.66\} = 2.66.$$

We also provide the least value of ε (up to the second decimal digit), for which, within the numerical method used, the same upper bound 2.66 holds,

$$\inf \left\{ \varepsilon > 0 \colon C_{\mathsf{R}}(\varepsilon, \gamma_*) \le 2.66 \right\} \approx 1.99.$$

The interest to exactly the least value of ε is stipulated by that the second term $(\sup_{0 \le z \le \varepsilon} \{\ldots\})$ in the definition of $L_{\mathbb{R},n}(\varepsilon,\gamma)$ is monotonically increasing with respect $\varepsilon > 0$.

Similarly, the constructed upper bound $\widehat{C}^*_{\mathbf{R}}(\varepsilon,\gamma)$ (defined in (18)) for the asymptotically exact constant $C^*_{\mathbf{R}}(\varepsilon,\gamma)$ (defined in (14)) attains its minimum value at the point $\gamma = \gamma_*, \varepsilon \approx 1.89$ with

$$\inf_{\varepsilon>0,\gamma>0}\widehat{C_{\mathbf{R}}^*}(\varepsilon,\gamma)\leq \widehat{C_{\mathbf{R}}^*}(1.89,\gamma_*)\leq 1.75;$$

the plot of $\widehat{C}_{\mathbf{R}}^*(\varepsilon,\gamma_*)$ is given on Fig. 3 (left, solid line), while the least value of ε , for which the inequality $\widehat{C}_{\mathbf{R}}^*(\varepsilon,\gamma_*) \leq 1.75$ still holds, is around 1.52. Furthermore, as Fig. 3 demonstrates, with the decrease of γ , the graphs of $\widehat{C}_{\mathbf{R}}^*(\cdot,\gamma)$ and $C_{\mathbf{R}}(\cdot,\gamma)$ become more and more flat around the points of minimum, so that the intervals, where the values of the functions $\widehat{C}_{\mathbf{R}}^*(\cdot,\gamma)$ and $C_{\mathbf{R}}(\cdot,\gamma)$ and $C_{\mathbf{R}}(\cdot,\gamma)$ and $C_{\mathbf{R}}(\cdot,\gamma)$ differ slightly from their minimal values, $\min_{\varepsilon>0} \widehat{C}_{\mathbf{R}}^*(\varepsilon,\gamma)$ and $\min_{\varepsilon>0} C_{\mathbf{R}}(\varepsilon,\gamma)$, enlarges. For example, for $\gamma = 0.4$ (see Fig. 3 (left, dashdot line) and table 3) the point of minimum of $\widehat{C}_{\mathbf{R}}^*(\cdot,\gamma)$ is located around $\varepsilon \approx 1.99$ with $\widehat{C}_{\mathbf{R}}^*(1.99, 0.4) \leq 1.78$, while the inequality $\widehat{C}_{\mathbf{R}}^*(\varepsilon, 0.4) \leq 1.78$ still holds for $\varepsilon = 1.41$; similarly, for $\gamma = 0.2$

$$\inf_{\varepsilon>0}\widehat{C}^*_{\scriptscriptstyle \rm R}(\varepsilon,0.2) \le \widehat{C}^*_{\scriptscriptstyle \rm R}(2.77,0.2) \le 1.93, \quad \inf\{\varepsilon>0\colon \widehat{C}^*_{\scriptscriptstyle \rm R}(\varepsilon,0.2) \le 1.93\} \approx 1.89.5$$

see Fig. 3 (left, dotted line) and table 3. For the "absolute" constant $C_{\rm R}(\varepsilon, \gamma)$ with $\gamma \in \{0.2, 0.4\}$ we have

$$\begin{split} \inf_{\varepsilon>0,\,\gamma\geq0.4} C_{\rm R}(\varepsilon,\gamma) &\leq C_{\rm R}(2.63,0.4) \leq \max\{C_{\rm min},\,C_0(2.63,0.4,0.03),C_1(2.63,0.4,0.4822\ldots)\} \leq 2.70,\\ \inf_{\varepsilon>0,\,\gamma\geq0.2} C_{\rm R}(\varepsilon,0.4) \leq 2.70\} &\approx 1.76, \quad C_{\rm R}(1.76,0.4) \leq \max\{2,\,2.37,\,2.70\} = 2.70;\\ \inf_{\varepsilon>0,\,\gamma\geq0.2} C_{\rm R}(\varepsilon,\gamma) \leq C_{\rm R}(5.39,0.2) \leq \max\{2,\,2.68,\,2.87\} = 2.87,\\ \inf_{\varepsilon>0,\,\gamma\geq0.2} C_{\rm R}(\varepsilon,0.2) \leq 2.87\} &\approx 1.21, \quad C_{\rm R}(1.21,0.2) \leq \max\{2,\,2.64,\,2.87\} = 2.87; \end{split}$$

see Fig. 3 (right, dashdot and dotted lines) and tables 3 and 5.

Another particular case is concerned with the historical values $\gamma = 1$ and $\varepsilon \in \{1, \infty\}$, for which, according to tables 2, 3, 4, and 5, we have

$$C_{\mathsf{R}}(1,1) \le C_{\mathsf{R}}(1,\gamma_*) \le 2 \lor C_0(1,\gamma_*,0.03) \lor C_1(1,\gamma_*,0.4834\ldots) \le 2 \lor 2.35 \lor 2.73 = 2.73,$$
$$C_{\mathsf{E}}(1,1) \le C_{\mathsf{E}}(1,0.72) \le 2.73 \ge C_{\mathsf{E}}(1,\infty),$$

 $C_{\rm E}(\infty, 1) \le 2.66$, which also follows from $C_{\rm E}(\infty, 0.97) \le 2.66$ or $C_{\rm E}(4.35, 1) \le 2.66$.

It is interesting to note that, as it follows from Tables 4 and 5, the largest contribution into extreme values of $C(\varepsilon, \gamma) = C_1(\varepsilon, \gamma, L_*)$ in both inequalities (11), (12) for all the presented values of ε and γ is provided by the integral I_3 which depends on the constructed estimates for characteristic functions through the maximal length of the interval, where the absolute value of a characteristic function can be estimated by a majorant strictly less than 1. Hence, to get further improvements of the constructed upper bounds for $C(\varepsilon, \gamma)$, one should improve, in first turn, upper bounds for absolute values of characteristic functions presented in Theorem 4.

5 The comparison of Osipov's, Lyapunov's, and modified Esseen's and Rozovskii's fractions. Lower bound for $\gamma \to 0$.

In the present section we compare the fractions $L^3_{E,n}(\varepsilon, 1)$, $L^3_{R,n}(\varepsilon, 1)$ with $L_{3,n}$ and $\Lambda_n(1) + L_n(1)$, and demonstrate that our new inequalities (11), (12) with $C_E(1,1) = 2.73 = C_R(1,1)$ may be sharper than Osipov's inequality (5) with the best known constant C = 1.87 [14]. In what follows we emphasize the dependence of the above fractions on the distributions of random summands X_1, \ldots, X_n with the d.f.'s F_1, \ldots, F_n by writing $L^3_{E,n}(\varepsilon, \gamma, F_1, \ldots, F_n)$, $L^3_{R,n}(\varepsilon, \gamma, F_1, \ldots, F_n)$, and $L_{3,n}(F_1, \ldots, F_n)$ using the three-argument notation $L^3_{E,n}(\varepsilon, \gamma, F)$, $L_{3,n}(F)$, $n \in \mathbb{N}$, in the i.i.d. case, that is, for $F_1 = \ldots = F_n = F$. Let \mathcal{F} denote the set of all d.f.'s on \mathbb{R} with zero mean and finite second-order moment and $F_p \in \mathcal{F}$ be the d.f. of the two-point distribution prescribing the masses $p \in [\frac{1}{2}, 1)$ and q = 1 - p to the points $\sqrt{q/p}$ and $-\sqrt{p/q}$. It easy to see that F_p has zero mean and unit variance.

Theorem 6. (i) For all $n \in \mathbb{N}$, $\varepsilon > 0$, and $F_1, \ldots, F_n \in \mathcal{F}$ such that $B_n > 0$ we have

$$L^3_{\mathrm{E},n}(\varepsilon,1) \le L_{3,n},$$

where the equality takes place for every $n \in \mathbb{N}$ and $\varepsilon > 0$ such that $n\varepsilon^2 \ge 1$. As for the extremal, one can take a common d.f. $F_1 = \ldots = F_n = F_p$ with arbitrary $p \in \left[\frac{1}{2}, 1\right)$ satisfying $p/(1-p) \le n\varepsilon^2$. (ii) For all $n \in \mathbb{N}$, $\varepsilon > 0$, and $p \in \left(\frac{1}{2}, \frac{\sqrt{5}-1}{2}\right)$ such that $n\varepsilon^2 > (1-p)/p$ we have

$$L^3_{\mathbf{R},n}(\varepsilon,1,F_p) > L_{3,n}(F_p).$$

in particular, $L^3_{\text{R},n}(\varepsilon, 1, F_p) > L^3_{\text{E},n}(\infty, 1, F_p)$, $L^3_{\text{R},n}(\varepsilon, 1, F_p) > \widehat{L}^3_{\text{E},n}(\infty)$. (iii) There exists a d.f. $F \in \mathcal{F}$ such that for all $n \in \mathbb{N}$ and $\varepsilon > 0$ satisfying the condition $n\varepsilon^2 \ge \frac{49}{25} = 1.96$ we have

$$L^3_{\mathrm{r},n}(\varepsilon,1,F) < L^3_{\mathrm{e},n}(\varepsilon,1,F)$$

in particular, $\sup_{\varepsilon>0} L^3_{R,n}(\varepsilon,1,F) < L_{3,n}(F)$, where one can consider F to be a discrete tree-point d.f.

(iv) There exists a common distribution of random summands with the d.f. from \mathcal{F} such that Esseen–Rozovskiitype inequalities (11), (12) with $C_{\rm E}(1,1) = C_{\rm R}(1,1) = 2.73$ are sharper than Osipov's inequality (5) with C = 1.87, namely,

 $2.73 \cdot \max \{L_{\mathbb{F},n}^3(1,1), L_{\mathbb{R},n}^3(1,1)\} < 1.87(\Lambda_n(1) + L_n(1)) \text{ for every } n \geq 9.$

Proof. The inequality in (i) follows from (10) with $\delta = 1$. To prove that equality also occurs, consider the sequence of i.i.d. r.v.'s X_1, \ldots, X_n with the d.f. $F_p \in \mathcal{F}$ for $p \in [\frac{1}{2}, 1)$:

$$X_1 = \begin{cases} \sqrt{\frac{q}{p}}, & p, \\ -\sqrt{\frac{p}{q}}, & q \coloneqq 1 - p. \end{cases}$$

We have $B_n = \sqrt{n}$,

$$\sqrt{n}L_{3,n} = \int_{\mathbb{R}} |x|^3 dF_p(x) = \frac{p^2 + q^2}{\sqrt{pq}} \quad \text{for all } n \in \mathbb{N},$$

$$\mu(z) \coloneqq \int_{|x| < z} x^3 dF_p(x) = \begin{cases} 0, & 0 < z \le \sqrt{\frac{q}{p}}, \\ \frac{q^2}{\sqrt{pq}}, & \sqrt{\frac{q}{p}} < z \le \sqrt{\frac{p}{q}}, \\ \frac{p-q}{\sqrt{pq}}, & z > \sqrt{\frac{p}{q}}, \end{cases} \quad \sigma^2(z) \coloneqq \int_{|x| \ge z} x^2 dF_p(x) = \begin{cases} 1, & 0 < z \le \sqrt{\frac{q}{p}}, \\ p, & \sqrt{\frac{q}{p}} < z \le \sqrt{\frac{p}{q}}, \\ 0, & z > \sqrt{\frac{p}{q}}. \end{cases}$$

With the account of $\mu(z) \ge 0$ for all z > 0 and of the left-continuity of the functions $\mu(z)$ and $\sigma^2(z)$ for z > 0, for all $n \in \mathbb{N}$ and $p \in \left[\frac{1}{2}, 1\right)$ we obtain for every $\varepsilon > 0$

$$\begin{split} \sqrt{n}L^3_{\mathbf{E},n}(\varepsilon,1,F_p) &= \sup_{0 < z \le \varepsilon \sqrt{n}} \left\{ \mu(z) + z\sigma^2(z) \right\} = \\ &= \begin{cases} \varepsilon \sqrt{n}, & n\varepsilon^2 \le \frac{q}{p}, \\ \max\left\{\sqrt{\frac{q}{p}}, \frac{q^2}{\sqrt{pq}} + p\varepsilon \sqrt{n}\right\} = \frac{q^2}{\sqrt{pq}} + p\varepsilon \sqrt{n}, & \frac{q}{p} < n\varepsilon^2 \le \frac{p}{q}, \\ \max\left\{\sqrt{\frac{q}{p}}, \frac{q^2}{\sqrt{pq}} + p\sqrt{\frac{p}{q}}, \frac{p-q}{\sqrt{pq}}\right\} = \frac{q^2+p^2}{\sqrt{pq}}, & n\varepsilon^2 > \frac{p}{q}, \end{cases} \end{split}$$

$$\begin{split} \sqrt{n}L^3_{\mathbf{R},n}(\varepsilon,1,F_p) &= \mu\left(\varepsilon\sqrt{n}\right) + \sup_{0 < z \le \varepsilon\sqrt{n}} z\sigma^2(z) = \\ &= \begin{cases} \varepsilon\sqrt{n}, & n\varepsilon^2 \le \frac{q}{p}, \\ \frac{q^2}{\sqrt{pq}} + \max\left\{\sqrt{\frac{q}{p}}, p\varepsilon\sqrt{n}\right\} = \frac{q^2 + \max\left\{q, \varepsilon p\sqrt{npq}\right\}}{\sqrt{pq}}, & \frac{q}{p} < n\varepsilon^2 \le \frac{q}{q}, \\ \frac{p-q}{\sqrt{pq}} + \max\left\{\sqrt{\frac{q}{p}}, p\sqrt{\frac{p}{q}}\right\} = \frac{p-q + \max\left\{q, p^2\right\}}{\sqrt{pq}} = \begin{cases} \frac{p}{\sqrt{pq}}, & \frac{1}{2} \le p \le \frac{\sqrt{5}-1}{2}, \\ \frac{p-q+p^2}{\sqrt{pq}}, & \frac{\sqrt{5}-1}{2} < p < 1, \end{cases} & n\varepsilon^2 > \frac{p}{q}. \end{split}$$

Now it is easy to see that for all $p \in \left[\frac{1}{2}, 1\right)$ and $n\varepsilon^2 \ge p/q$,

$$\sqrt{n}L^3_{E,n}(\varepsilon, 1, F_p) = \frac{q^2}{\sqrt{pq}} + p\sqrt{\frac{p}{q}} = \frac{q^2 + p^2}{\sqrt{pq}} = \sqrt{n}L_{3,n}(F_p),$$

where, for $n\varepsilon^2 = 1$, the unique value $p = \frac{1}{2}$ is admissible. Thus, (i) is proved. Now let $\frac{1}{2} . If <math>n\varepsilon^2 > p/q$, then

$$\sqrt{n}L_{\mathbf{R},n}^{3}(\varepsilon,1,F_{p}) = \frac{p}{\sqrt{pq}} > \frac{p+q(q-p)}{\sqrt{pq}} = \frac{p^{2}+q^{2}}{\sqrt{pq}} = \sqrt{n}L_{3,n}(F_{p}),$$

while for $q/p < n\varepsilon^2 \leq p/q$ we have $q > p^2 \geq p\varepsilon \sqrt{npq}$ and, hence,

$$\sqrt{n}L^3_{{}_{\mathrm{R},n}}(\varepsilon,1,F_p) = \frac{q^2+q}{\sqrt{pq}} > \frac{q^2+p^2}{\sqrt{pq}} = \sqrt{n}L_{3,n}(F_p),$$

so that $L^3_{\text{\tiny R},n}(\varepsilon,1,F_p) > L_{3,n}(F_p) \ge \sup_{\varepsilon'>0} L^3_{\text{\tiny E},n}(\varepsilon',1,F_p)$ for all $n \in \mathbb{N}$, $\varepsilon > 0$, and $p \in \left(\frac{1}{2},\frac{\sqrt{5}-1}{2}\right)$ such that $n\varepsilon^2 > (1-p)/p$, which proves (ii).

To prove (iii), consider the common three-point distribution of the random summands X_1, \ldots, X_n concentrated in the points

$$x_1 = \frac{4}{5}, \quad x_2 = -1, \quad x_3 = \frac{7}{5},$$

with masses $p, q, r \ge 0$ such that

$$\begin{cases} p+q+r=1,\\ \mathsf{E}X_1=px_1+qx_2+rx_3=0,\\ \mathsf{E}X_1^2=px_1^2+qx_2^2+rx_3^2=1, \end{cases}$$

that is,

$$p = \frac{10}{27} = 0.3703..., \quad q = \frac{53}{108} = 0.4907..., \quad r = \frac{5}{36} = 0.1388...$$

Now for $n\varepsilon^2 \ge (|x_1| \lor |x_2| \lor |x_3|)^2 = x_3^2 = \frac{49}{25} = 1.96$ with $B_n = \sqrt{n}$ we have

$$\begin{split} \sqrt{n} L^3_{\text{E},n}(\varepsilon,1) &= \max\left\{ \left| x_1 \right|, \left| x_1^3 p \right| + (1 - x_1^2 p) \left| x_2 \right|, \left| x_1^3 p + x_2^3 q \right| + x_3^2 r \cdot \left| x_3 \right|, \left| x_1^3 p + x_2^3 q + x_3^3 r \right| \right\} = \\ &= \frac{643}{675} = 0.9525 \dots, \end{split}$$

 $\sqrt{n}L_{\mathbf{R},n}^{3}(\varepsilon,1) = \left|x_{1}^{3}p + x_{2}^{3}q + x_{3}^{3}r\right| + \max\left\{\left|x_{1}\right|, \left(1 - x_{1}^{2}p\right)\left|x_{2}\right|, x_{3}^{2}r \cdot \left|x_{3}\right|\right\} = \frac{22}{25} = 0.88 < \sqrt{n}L_{\mathbf{E},n}^{3}(\varepsilon,1),$

which proves (iii) for $n\varepsilon^2 \ge \frac{49}{25}$.

To prove (iv), let us consider the common symmetric four-point distribution of the random summands X_1, \ldots, X_n of the form

$$X_1 = \begin{cases} \pm x_1, & p/2 \ge 0, \\ \pm x_2, & q/2 \ge 0, \end{cases} \quad p+q = 1,$$

and such that $\mathsf{E}X_1 = 0$, $\mathsf{E}X_1^2 = 1$. Setting $x_1 = 0.9$ and $x_2 = 3$, we find $p = \frac{800}{819} = 0.9768\ldots$ and $q = \frac{19}{819} = 0.0231\ldots$, so that for $n\varepsilon^2 \ge x_1^2 \lor x_2^2 = x_2^2 = 9$ with $B_n = \sqrt{n}$ we have

$$\sqrt{n}L^{3}_{E,n}(\varepsilon,1) = \sqrt{n}L^{3}_{R,n}(\varepsilon,1) = \max\left\{x_{1}, x_{2}\left(1-px_{1}^{2}\right)\right\} = 0.9,$$

while for $n \ge x_1^2 \lor x_2^2 = 9$ we obtain

$$\sqrt{n}(\Lambda_n(1) + L_n(1)) = x_1^3 p + x_2^3 q = \frac{87}{65} = 1.3384\dots$$

In particular, for $\varepsilon = 1$ and $n \ge 9$

$$2.73 \cdot \sqrt{n} \max\{L_{\text{E},n}^3(1,1), L_{\text{R},n}^3(1,1)\} = 2.457 < 2.5029 \dots = 1.87 \cdot \sqrt{n}(\Lambda_n(1) + L_n(1)),$$

which proves (iv). We can also propose another "non-symmetric" example for even n. Let X_1, \ldots, X_n be independent r.v.'s with the distributions $X_k \stackrel{d}{=} X$ if k is odd and $X_k \stackrel{d}{=} -X$ if k is even, where X is a three-point r.v. with

$$X = \begin{cases} x_1 = 1/2, & p = \frac{4}{9}, \\ x_2 = -1, & q = \frac{4}{9}, \\ x_3 = 2, & r = \frac{1}{9}, \end{cases} \quad p + q + r = 1, \text{ so that } \mathsf{E}X = 0, \quad \mathsf{E}X^2 = 1.$$

Taking into account that $\sum_{k=1}^{n} \mathsf{E} X_k^3 \mathbf{1} (|X_k| < z) = 0$ for even n and all z > 0, for even $n \ge \max\{x_1^2, x_2^2, x_3^2\} = 4$ we have

$$\sqrt{n}L_{\text{E},n}^{3}(1,1) = \sqrt{n}L_{\text{R},n}^{3}(1,1) = \sup_{0 < z \le \sqrt{n}} \left\{ z\mathsf{E}X^{2}\mathbf{1} \left(|X| \ge z \right) \right\} = \max \left\{ x_{1}, \left(1 - px_{1}^{2}\right) |x_{2}|, rx_{3}^{3} \right\} = \frac{8}{9},$$

$$\sqrt{n}(\Lambda_{n}(1) + L_{n}(1)) = px_{1}^{3} + q |x_{2}|^{3} + rx_{3}^{3} = \frac{25}{18} = 1.3888...,$$

$$2.73 \cdot \sqrt{n} \max \left\{ L_{\text{E},n}^{3}(1,1), L_{\text{R},n}^{3}(1,1) \right\} = 2.4266... < 2.5972... = 1.87 \cdot \sqrt{n}(\Lambda_{n}(1) + L_{n}(1)).$$

Theorem 7. For the asymptotically best constant we have

$$\sup_{F_1 = \ldots = F_n \in \mathcal{F}: \ B_n > 0} \quad \limsup_{n \to \infty} \frac{\Delta_n(F_1, \ldots, F_n)}{\sup_{0 < z \le \varepsilon} z L_n(z)} = \infty \quad for \ every \quad \varepsilon > 0.$$

Proof. Using his asymptotic expansion, Esseen deduced [7] that in the i.i.d. case with $\mathsf{E}X_1 = 0$, $\mathsf{E}X_1^2 = \sigma^2 > 0$, $\mathsf{E}|X_1|^3 < \infty$ the limit below exists and

$$\lim_{n \to \infty} \sqrt{n} \Delta_n = \frac{|\alpha_3| + 3h\sigma^2}{6\sqrt{2\pi}\sigma^3},$$

where $\alpha_3 := \mathsf{E} X_1^3$, h > 0 is the span in case of a lattice distribution of X_1 , and h := 0 otherwise. Now let us consider an absolute continuous distribution of X_1 whose d.f. F_{θ} is defined by the density

$$p_{\theta}(x) = \begin{cases} ax^{-4-\theta}, & x > 1\\ 0, & |x| \le 1, \\ b|x|^{-5}, & x < -1, \end{cases} \text{ with } a = a(\theta) \coloneqq \frac{4(2+\theta)(3+\theta)}{17+7\theta}, \ b = b(\theta) \coloneqq \frac{12(3+\theta)}{17+7\theta}, \ \theta \in (0,1).$$

Then h = 0, $EX_1 = 0$,

$$\begin{split} \sigma^2 &= \sigma^2(\theta) = \frac{a}{1+\theta} + \frac{b}{2} = \frac{2(3+\theta)(7+5\theta)}{(1+\theta)(17+7\theta)}, \quad \alpha_3 = \alpha_3(\theta) = \frac{a}{\theta} - b = \frac{4(3+\theta)(2-8\theta-3\theta^2)}{\theta(17+7\theta)}, \\ & \mathsf{E}X_1^2 \mathbf{1}(|X_1| \ge z) = \begin{cases} \sigma^2, & z \in (0,1], \\ \frac{a}{1+\theta} z^{-1-\theta} + \frac{b}{2} z^{-2}, & z > 1, \end{cases} \\ & \lim_{n \to \infty} \sqrt{n} \sup_{0 < z \le \varepsilon} z L_n(z) = \sigma^{-3} \lim_{n \to \infty} \sup_{0 < z \le \varepsilon \sigma \sqrt{n}} z \mathsf{E}X_1^2 \mathbf{1}(|X_1| \ge z) = \sigma^{-3} \sup_{z > 0} z \mathsf{E}X_1^2 \mathbf{1}(|X_1| \ge z) = \sigma^{-1}, \end{split}$$

so that

$$\sup_{F_1 = \ldots = F_n \in \mathcal{F} \colon B_n > 0} \limsup_{n \to \infty} \frac{\Delta_n(F_1, \ldots, F_n)}{\sup_{0 < z \le \varepsilon} z L_n(z)} \ge \sup_{\theta \in (0,1]} \frac{\lim_{n \to \infty} \sqrt{n} \Delta_n(F_\theta, \ldots, F_\theta)}{\lim_{n \to \infty} \sqrt{n} \sup_{0 < z \le \varepsilon} z L_n(z)} = \frac{1}{6\sqrt{2\pi}} \lim_{\theta \to 0} \frac{|\alpha_3(\theta)|}{\sigma^2(\theta)} = \infty.$$

6 Appendix: Figures and Tables



Figure 1. Graphs of the functions $t_{\gamma} \coloneqq \frac{2}{\gamma} \left(\sqrt{(\gamma/\gamma_*)^2 + 1} - 1 \right)$ (solid line) and $t_{1,\gamma} \coloneqq \frac{2}{\gamma} \left(1 - \sqrt{(1 - (\gamma/\gamma_*)^2)_+} \right)$ (dashdot line) for $\gamma > 0$; dashed line represents the limiting value $t_{\infty} \coloneqq \lim_{\gamma \to \infty} t_{\gamma} = t_{1,\gamma_*} = 2/\gamma_* = 3.5717 \dots$



Figure 2. Level curves $\gamma = \gamma(\varepsilon)$ for the upper bounds to the asymptotically exact "constant" $C_{\text{E}}^*(\varepsilon, \gamma)$ defined in (13) and to the absolute $C_{\text{E}}(\varepsilon, \gamma)$ "constant" in the Esseen-type inequality (11). Left: $\{(\varepsilon, \gamma): \widehat{C_{\text{E}}^*}(\varepsilon, \gamma) = 1.72\}$ with $\widehat{C_{\text{E}}^*}(\varepsilon, \gamma) \coloneqq C_0(\varepsilon, \gamma, 0+)$ defined in (15). Right: $\{(\varepsilon, \gamma): \max_{L_0 \leq L \leq L_1} C_1(\varepsilon, \gamma, L) = 2.65\}$ with $C_1(\varepsilon, \gamma, L)$ defined in (48).



Figure 3. Left: Graphs of the function $\widehat{C}_{\mathbb{R}}^{*}(\varepsilon,\gamma)$ (see (18)) which bounds from above the asymptotically exact constant $C_{\mathbb{R}}^{*}(\varepsilon,\gamma)$ (see (14)), with respect to ε , for $\gamma \geq \gamma_{*} = 0.5599...$ (solid line), $\gamma = 0.4$ (dashdot line), $\gamma = 0.3$ (dashed line), and $\gamma = 0.2$ (dotted line).

Right: Graphs of the upper bounds $\max_{L_0 \leq L \leq L_1} C_1(\varepsilon, \gamma, L)$ (see (48)) for the "constants" $C_{\mathbb{R}}(\varepsilon, \gamma)$ in the Rozovskii-type inequality, with respect to ε , for $\gamma \geq \gamma_* = 0.5599...$ (solid line), $\gamma = 0.4$ (dashdot line), $\gamma = 0.3$ (dashed line), and $\gamma = 0.2$ (dotted line).



Figure 4. Demonstration to the evaluation of the absolute constants $C_{\text{E}}(\infty, \infty)$ and $C_{\text{R}}(2.12, \gamma_*)$: Plots of the functions $C_1(\infty, \infty, L), L \in [L_0, L_1]$, defined in (48) with $L \coloneqq L_{\text{E},n}(\infty, \infty)$ (left) and $L \coloneqq L_{\text{R},n}(2.12, \gamma_*)$ (right).

c	\sim	$C_0(\varepsilon, \gamma, 0+)$	L = 0.001	Opt	imal	L = 0.03	Opt	imal
E	·γ	$C_0(\varepsilon, \gamma, 0+)$	$C_0(\varepsilon, \gamma, L)$	$ au_0$	$ au_1$	$C_0(\varepsilon, \gamma, L)$	$ au_0$	$ au_1$
0.6	0.3	1.92245	1.98369	0.33196	0.76348	2.63565	0.61269	0.58888
1.21	0.2	1.95457	2.02040	0.30850	0.76192	2.70866	0.58750	0.58212
2.06	0.2	1.94999	2.01563	0.30911	0.76197	2.70121	0.58906	0.58256
∞	0.2	1.94879	2.01437	0.30943	0.76199	2.69812	0.58998	0.58279
1.48	0.4	1.80997	1.86401	0.37487	0.76589	2.45637	0.65732	0.59975
∞	0.4	1.80005	1.85350	0.37849	0.76607	2.44001	0.66177	0.60081
1.89	γ_*	1.77136	1.82167	0.39960	0.76705	2.38689	0.68022	0.60488
2.03	γ_*	1.76995	1.82017	0.40115	0.76711	2.38467	0.68083	0.60502
∞	γ_*	1.76370	1.81351	0.40416	0.76725	2.37413	0.68386	0.60563
1	γ_*	1.80596	1.85831	0.38771	0.76651	2.43955	0.66679	0.60193
1	0.67	1.79961	1.85099	0.39328	0.76673	2.42541	0.67230	0.60312
1	∞	1.79149	1.83892	0.42035	0.76791	2.38889	0.69303	0.60759
2.24	1	1.73996	1.78661	0.43002	0.76828	2.32719	0.70218	0.60952
∞	1	1.73186	1.77796	0.44157	0.76870	2.31385	0.70646	0.61040
3.07	∞	1.71998	1.76313	0.45717	0.76925	2.28233	0.72259	0.61358
3.2	5	1.71997	1.76354	0.45694	0.76926	2.28502	0.72045	0.61316
3.28	4	1.71999	1.76368	0.45267	0.76914	2.28573	0.71991	0.61306
4	2.4	1.71998	1.76401	0.45217	0.76907	2.28788	0.71820	0.61272
5	2.06	1.71997	1.76413	0.45034	0.76902	2.28870	0.71753	0.61259
5.37	2	1.72000	1.76420	0.45155	0.76907	2.28892	0.71737	0.61256
∞	1.83	1.71995	1.76423	0.45135	0.76907	2.28913	0.71712	0.61251
∞	∞	1.71451	1.75725	0.46485	0.76952	2.27337	0.72554	0.61416

Table 2. Demonstration to the evaluation of the upper bound $C_0(\varepsilon, \gamma, L_0)$ defined in (46) for the constant $C_{\rm E}(\varepsilon, \gamma)$ in the Esseen-type inequality (11) for small values of $L \coloneqq L_{{\rm E},n}(\varepsilon, \gamma) \leq L_0$ and some ε, γ : Values of $C_0(\varepsilon, \gamma, L_0)$, rounded up, for $L_0 = 0.001$ (the fourth column) and $L_0 = 0.03$ (the seventh column) accompanied by the corresponding optimal values of the parameters τ_0, τ_1 in (46). The third column provides values of the function $\widehat{C}^*_{\rm E}(\varepsilon, \gamma) \coloneqq C_0(\varepsilon, \gamma, 0+)$ defined in (15) which bounds from above the asymptotically exact constant $C^*_{\rm E}(\varepsilon, \gamma)$ defined in (13). Recall that $\gamma_* = 0.5599...$

c	\sim	$C_0(\varepsilon \propto 0)$	L = 0.001	Opt	imal	L = 0.03	Optimal	
C	Ŷ	$C_0(\varepsilon,\gamma,0)$	$C_0(\varepsilon, \gamma, L)$	$ au_0$	$ au_1$	$C_0(\varepsilon, \gamma, L)$	$ au_0$	$ au_1$
1.21	0.2	1.93474	1.99463	0.33997	0.76401	2.63576	0.62018	0.59077
1.89	0.2	1.92998	1.98967	0.33862	0.76391	2.62929	0.62130	0.59105
2.77	0.2	1.92890	1.98855	0.33849	0.76389	2.62857	0.62119	0.59103
5.39	0.2	1.95832	2.01917	0.33339	0.76358	2.67331	0.61213	0.58874
1.41	0.4	1.77974	1.82650	0.42470	0.76809	2.37242	0.69715	0.60847
1.76	0.4	1.77249	1.81886	0.43083	0.76830	2.36208	0.69946	0.60894
1.99	0.4	1.77128	1.81759	0.43528	0.76848	2.36048	0.69982	0.60902
2.63	0.4	1.77841	1.82511	0.42595	0.76814	2.37148	0.69707	0.60846
0.5	γ_*	1.94743	1.99139	0.43510	0.76847	2.54844	0.68154	0.60513
1	γ_*	1.79154	1.83112	0.48527	0.77017	2.34604	0.72358	0.61378
1.52	γ_*	1.74995	1.78796	0.49835	0.77051	2.29031	0.73728	0.61641
1.89	γ_*	1.74383	1.78159	0.50808	0.77078	2.28222	0.73934	0.61679
1.99	γ_*	1.74412	1.78189	0.50748	0.77083	2.28271	0.73918	0.61676
2.12	γ_*	1.74542	1.78324	0.50216	0.77063	2.28462	0.73863	0.61666
3	γ_*	1.77092	1.80977	0.48990	0.77029	2.32025	0.72921	0.61487
5	γ_*	1.86500	1.90688	0.45666	0.76924	2.44632	0.70070	0.60920

Table 3. Demonstration to the evaluation of the upper bound $C_0(\varepsilon, \gamma, L_0)$ defined in (46) for the constant $C_{\mathbb{R}}(\varepsilon, \gamma)$ in the Rozovskii-type inequality (12) for small values of $L \coloneqq L_{\mathbb{R},n}(\varepsilon, \gamma) \leq L_0$ and some ε, γ : Values of $C_0(\varepsilon, \gamma, L_0)$, rounded up, for $L_0 = 0.001$ (the fourth column) and $L_0 = 0.03$ (the seventh column) accompanied by the corresponding optimal values of the parameters τ_0, τ_1 in (46). The third column provides values of the function $\widehat{C}^*_{\mathbb{R}}(\varepsilon, \gamma) \coloneqq C_0(\varepsilon, \gamma, 0+)$ defined in (18) which bounds from above the asymptotically exact constant $C^*_{\mathbb{R}}(\varepsilon, \gamma)$ defined in (14). Recall that $\gamma_* = 0.5599...$

c	5	$C_1 \leq$	L	Optima	$1 T_0, T_1$	Contributions of I_k/L_*^3				
C	7	01 2	L_*	T_0L_*	$T_{1}L_{*}^{3}$	k = 1	k = 2	k = 3	k = 4	
1.21	0.2	2.89038	0.47850	0.87755	0.69402	0.44928	0.81565	1.51412	0.11135	
1.24	0.2	2.88998	0.47856	0.87744	0.69400	0.44866	0.81589	1.51400	0.11144	
∞	0.2	2.88457	0.47857	0.87749	0.69398	0.44331	0.81581	1.51405	0.11142	
1.76	0.4	2.73593	0.48170	0.92023	0.69306	0.35252	0.76761	1.52685	0.08897	
5.94	0.4	2.73000	0.48173	0.92020	0.69311	0.34652	0.76781	1.52668	0.08900	
∞	0.4	2.72984	0.48173	0.92020	0.69311	0.34635	0.76781	1.52669	0.08900	
1	γ_*	2.73662	0.48267	0.92963	0.69278	0.36492	0.75795	1.52899	0.08477	
1.87	γ_*	2.69989	0.48245	0.92958	0.69285	0.32825	0.75774	1.52921	0.08471	
∞	γ_*	2.69190	0.48244	0.92960	0.69285	0.32028	0.75770	1.52924	0.08470	
1	0.72	2.72979	0.48304	0.93393	0.69261	0.36342	0.75335	1.53015	0.08289	
1	∞	2.72857	0.48346	0.93595	0.69239	0.36458	0.75136	1.53053	0.08211	
4.35	1	2.66000	0.48305	0.93735	0.69266	0.29795	0.74958	1.53117	0.08131	
∞	1	2.65879	0.48305	0.93734	0.69266	0.29673	0.74959	1.53117	0.08132	
∞	0.97	2.65985	0.48303	0.93710	0.69266	0.29749	0.74984	1.53111	0.08142	
2.56	∞	2.64999	0.48339	0.94137	0.69256	0.29282	0.74546	1.53211	0.07962	
2.62	5	2.64992	0.48338	0.94121	0.69256	0.29257	0.74561	1.53207	0.07968	
2.65	4	2.64996	0.48342	0.94113	0.69253	0.29249	0.74571	1.53205	0.07974	
2.74	3	2.64995	0.48335	0.94092	0.69257	0.29224	0.74592	1.53200	0.07980	
3.13	2	2.64997	0.48329	0.94031	0.69259	0.29153	0.74652	1.53188	0.08005	
4	1.62	2.64999	0.48325	0.93979	0.69259	0.29091	0.74705	1.53176	0.08027	
5.37	1.5	2.64993	0.48323	0.93954	0.69260	0.29055	0.74732	1.53170	0.08038	
∞	1.43	2.64991	0.48321	0.93935	0.69260	0.29030	0.74750	1.53166	0.08046	
∞	∞	2.64082	0.48338	0.94138	0.69255	0.28367	0.74541	1.53214	0.07961	

Table 4. Demonstration to the evaluation of the upper bound $\max_{L_0 \leq L \leq L_1} C_1(\varepsilon, \gamma, L)$ (see (48)) for the constant $C_{\rm E}(\varepsilon, \gamma)$ in the Esseen-type inequality (11) for moderate values of $L := L_{\rm E,n}(\varepsilon, \gamma) \in [L_0, L_1] = [0.03, 0.65]$ and some ε, γ : Extreme values of $C_1(\varepsilon, \gamma, L) \leq C_1(\varepsilon, \gamma, L_*)$ on the interval $L \in [L_0, L_1]$ (column 3); maximizer L_* , rounded down (column 4); optimal values of the parameters T_0 and T_1 in (48) multiplied by L and L^3 in the extremal point $L = L_*$, rounded down (columns 5, 6); values of the normalized integrals I_k/L_*^3 , k = 1, 2, 3, 4, rounded down (columns 7–10), so that $C_1(\varepsilon, \gamma, L_*) = (I_1 + I_2 + I_3 + I_4)/L_*^3$. Recall that $\gamma_* = 0.5599\ldots$

c	γ	$C_1 \leq$	L_*	Optimal T_0, T_1		Contributions of I_k/L_*^3			
C				T_0L_*	$T_{1}L_{*}^{3}$	k = 1	k = 2	k = 3	k = 4
1.21	0.2	2.86991	0.47901	0.88360	0.69388	0.43734	0.80872	1.51596	0.10790
5.39	0.2	2.86343	0.47902	0.88371	0.69384	0.43101	0.80851	1.51608	0.10784
1.76	0.4	2.69985	0.48249	0.93002	0.69288	0.32875	0.75737	1.52923	0.08452
2.63	0.4	2.69323	0.48229	0.92948	0.69296	0.32154	0.75779	1.52922	0.08470
0.5	γ_*	3.03953	0.50011	0.92200	0.66074	0.52576	0.85947	1.55924	0.09507
1	γ_*	2.72857	0.48346	0.93595	0.69239	0.36458	0.75136	1.53053	0.08211
1.99	γ_*	2.65991	0.48300	0.93912	0.69274	0.30011	0.74764	1.53169	0.08048
2.12	γ_*	2.65925	0.48273	0.93728	0.69293	0.29727	0.74960	1.53117	0.08122
3	γ_*	2.67687	0.48125	0.92365	0.69352	0.29824	0.76363	1.52790	0.08710
5	γ_*	2.75611	0.47832	0.89074	0.69456	0.33467	0.79931	1.51873	0.10341

Table 5. Demonstration to the evaluation of the upper bound $\max_{L_0 \leq L \leq L_1} C_1(\varepsilon, \gamma, L)$ (see (48)) for the constant $C_{\mathbb{R}}(\varepsilon, \gamma)$ in the Rozovskii-type inequality (12) for moderate values of $L \coloneqq L_{\mathbb{R},n}(\varepsilon, \gamma) \in [L_0, L_1] = [0.03, 0.65]$ and some ε, γ : Extreme values of $C_1(\varepsilon, \gamma, L) \leq C_1(\varepsilon, \gamma, L_*)$ on the interval $L \in [L_0, L_1]$ (column 3); maximizer L_* , rounded down (column 4); optimal values of the parameters T_0 and T_1 in (48) multiplied by L and L^3 in the extremal point $L = L_*$, rounded down (columns 5, 6); values of the normalized integrals I_k/L_*^3 , k = 1, 2, 3, 4, rounded down (columns 7–10), so that $C_1(\varepsilon, \gamma, L_*) = (I_1 + I_2 + I_3 + I_4)/L_*^3$. Recall that $\gamma_* = 0.5599 \dots$

References

- [1] R. P. Agnew, "Estimates for global central limit theorems", Ann. Math. Stat., 28, 26–42 (1957).
- [2] A. D. Barbour and P. Hall, "Stein's method and the Berry-Esseen theorem", Australian Journal of Statistics, 26, 8–15 (1984).
- [3] A. C. Berry, "The accuracy of the Gaussian approximation to the sum of independent variates", Trans. Amer. Math. Soc., 49, 122–136 (1941).
- [4] R. N. Bhattacharya and R. Ranga Rao, Normal Approximation and Asymptotic Expansions, Wiley, New York (1976).
- [5] L. H. Y. Chen and Q. M. Shao, "A non-uniform Berry-Esseen bound via Stein's method", Probab. Theory Relat. Fields, 120, 236-254 (2001).
- [6] C.-G. Esseen, "On the Liapounoff limit of error in the theory of probability", Ark. Mat. Astron. Fys., A28, No. 9, 1–19 (1942).
- [7] C.-G. Esseen, "Fourier analysis of distribution functions. A mathematical study of the Laplace–Gaussian law", Acta Math., 77, No. 1, 1–125 (1945).
- [8] C.-G. Esseen, "On the remainder term in the central limit theorem", Arkiv för Matematik, 8, No. 1, 7–15 (1969).
- [9] W. Feller, "Über den zentralen Genzwertsatz der Wahrscheinlichkeitsrechnung", Math. Z., 40, 521–559 (1935).
- [10] W. Feller, "On the Berry–Esseen theorem", Z. Wahrsch. Verw. Geb., 10, 261–268 (1968).
- [11] I. A. Ibragimov, "On the accuracy of the approximation of distribution functions of sums of independent variables by the normal distribution", *Theory Probab. Appl.*, **11**, No. 4, 559–579 (1966).
- [12] M. L. Katz, "Note on the Berry–Esseen theorem", Ann. Math. Statist., 34, 1107–1108 (1963).
- [13] A. N. Kolmogorov, "Some recent works in the field of limit theorems of probability theory", Bulletin of Moscow University [in Russian], 10, No. 7, 29–38 (1953).
- [14] V. Korolev and A. Dorofeyeva, "Bounds of the accuracy of the normal approximation to the distributions of random sums under relaxed moment conditions", *Lith. Math. J.*, 57, No. 1, 38–58 (2017).
- [15] V. Korolev and I. Shevtsova, "An improvement of the Berry-Esseen inequality with applications to Poisson and mixed Poisson random sums", Scand. Actuar. J., 2012, No. 2, 81–105 (2012). Available online since 04 June 2010.
- [16] V. Yu. Korolev and S. V. Popov, "An improvement of convergence rate estimates in the central limit theorem under absence of moments higher than the second", *Theory Probab. Appl.*, 56, No. 4, 682–691 (2012).
- [17] V. Yu. Korolev and S. V. Popov, "Improvement of convergence rate estimates in the central limit theorem under weakened moment conditions", *Dokl. Math.*, 86, No. 1, 506–511 (2012).
- [18] V. Yu. Korolev and I. G. Shevtsova, "On the upper bound for the absolute constant in the Berry-Esseen inequality", *Theory Probab. Appl.*, 54, No. 4, 638–658 (2010).
- [19] A. Liapunoff, "Nouvelle forme du théorème sur la limite de probabilité", Mém. Acad. Sci. St-Pétersbourg, 12, No. 5, 1–24 (1901).
- [20] J. W. Lindeberg, "Eine neue Herleitung des Exponentialgesetzes in der Wahrscheinlichkeitsrechnung", Mathematische Zeitschrift, 15, No. 1, 211–225 (1922).
- [21] W. Y. Loh, On the normal approximation for sums of mixing random variables, Master thesis, Department of Mathematics, University of Singapore (1975).
- [22] L. V. Osipov, "Refinement of Lindeberg's theorem", Theory Probab. Appl., 10, No.2, 299–302 (1966).

- [23] L. Paditz, "Bemerkungen zu einer Fehlerabschätzung im zentralen Grenzwertsatz", Wiss. Z. Hochschule für Verkehrswesen "Friedrich List". Dresden., 27, No. 4, 829–837 (1980).
- [24] L. Paditz, "On error-estimates in the central limit theorem for generalized linear discounting", Math. Operationsforsch. u. Statist., Ser. Statistics, 15, No. 4, 601–610 (1984).
- [25] L. Paditz, "Über eine globale Fehlerabschätzung im zentralen Grenzwertsatz", Wiss. Z. Hochschule für Verkehrswesen "Friedrich List". Dresden., 33, No. 2, 399–404 (1986).
- [26] V. V. Petrov, "An estimate of the deviation of the distribution function of a sum of independent random variables from the normal law", Soviet Math. Dokl., 6, No. 5, 242–244 (1965).
- [27] V. V. Petrov, Sums of independent random variables, Springer-Verlag, Berlin-Heidelberg (1975).
- [28] H. Prawitz, "Limits for a distribution, if the characteristic function is given in a finite domain", Skand. Aktuarietidskr., 55, 138–154 (1972).
- [29] H. Prawitz, "Noch einige Ungleichungen f
 ür charakteristische Funktionen", Scand. Actuar. J., No. 1, 49–73 (1991).
- [30] L. V. Rozovskii, "On the rate of convergence in the Lindeberg–Feller theorem", Bulletin of Leningrad University [in Russian], No. 1, 70–75 (1974).
- [31] L. V. Rozovskii, "An estimate of the speed of convergence in the multidmensional central limit theorem without moment hypotheses", *Math. Notes*, **23**, No. 4, 343–351 (1978).
- [32] I. G. Shevtsova, "On the asymptotically exact constants in the Berry-Esseen-Katz inequality", Theory Probab. Appl., 55, No. 2, 225–252 (2011).
- [33] I. G. Shevtsova, "On the absolute constant in the Berry–Esseen inequality and its structural and non-uniform improvements", *Informatics and its Applications* [in Russian], 7, No. 1, 124–125 (2013).
- [34] I. G. Shevtsova, "On the absolute constants in the Berry–Esseen-type inequalities", Dokl. Math., 89, No.3, 378–381 (2014).
- [35] V. M. Zolotarev, "An absolute estimate of the remainder term in the central limit theorem", Theory Probab. Appl., 11, No. 1, 95–105 (1966).
- [36] V. M. Zolotarev, Modern Theory of Summation of Random Variables, VSP, Utrecht, The Netherlands (1997).