

Tree lengths for general Λ -coalescents and the asymptotic site frequency spectrum around the Bolthausen-Sznitman coalescent

Christina S. Diehl* and Götz Kersting*

March 13, 2022

Abstract

We study tree lengths in Λ -coalescents without a dust component from a sample of n individuals. For the total length of all branches and the total length of all external branches we present laws of large numbers in full generality. The other results treat regularly varying coalescents with exponent 1, which cover the Bolthausen-Sznitman coalescent. The theorems contain laws of large numbers for the total length of all internal branches and of internal branches of order a (i.e. branches carrying a individuals out of the sample). These results transform immediately to sampling formulas in the infinite sites model. In particular, we obtain the asymptotic site frequency spectrum of the Bolthausen-Sznitman coalescent. The proofs rely on a new technique to obtain laws of large numbers for certain functionals of decreasing Markov chains.

MSC 2010 subject classification. 60J75 (primary), 60F05, 60J27, 92D25

Keywords and phrases. Λ -coalescent, Bolthausen-Sznitman coalescent, law of large numbers, tree length, infinite sites model, sampling formula, site frequency spectrum, decreasing Markov chain

1 Introduction and main results

Λ -coalescents are established models for family trees of a sample of individuals from some large population. Its most prominent representative, the Kingman coalescent [18], is widely used in population genetics. More recently the Bolthausen-Sznitman coalescent [6] gained attention for models including selection. The class of Beta-coalescents with parameter $1 < \alpha < 2$ has been applied to marine populations [5]. In this paper we investigate branch lengths and sampling formulas in the infinite sites model for the whole class of Λ -coalescents without a dust component, which cover all these special cases.

Λ -coalescents have been introduced by Pitman [20] and Sagitov [21] as Markov processes whose states are partitions of \mathbb{N} and whose evolution may be imagined as a random tree. In this paper we identify a Λ -coalescent with an induced sequence of n -coalescents, $n \in \mathbb{N}$, by restricting partitions to the subsets $\{1, \dots, n\}$ of \mathbb{N} . These n -coalescents are considered to be Markovian models for the family trees of a sample of n individuals. If such a tree contains

*Institut für Mathematik, Goethe Universität, Frankfurt am Main, Germany
diehl@math.uni-frankfurt.de, kersting@math.uni-frankfurt.de

Work partially supported by the DFG Priority Programme SPP 1590 “Probabilistic Structures in Evolution”

$b \in \{2, \dots, n\}$ lineages at the moment $t \geq 0$ backwards in time (representing the ancestors living at that moment), then it is assumed that out of them $k \in \{2, \dots, b\}$ specified lines merge at rate

$$\lambda_{b,k} := \int_{[0,1]} p^k (1-p)^{b-k} \frac{\Lambda(dp)}{p^2}$$

to one line. Here Λ denotes any finite, non-vanishing measure on $[0, 1]$. The resulting tree consists of n leaves, τ_n merging events, and a root at time $\tilde{\tau}_n > 0$ representing the MCRA (most recent common ancestor). Its branches have lengths which specify lifetimes. There are external branches ending in leaves on the one hand and internal branches ending in mergers on the other. For the detailed partition valued picture we refer to the survey [2].

In the sequel we work with the Markovian counting process $N_n = (N_n(t))_{t \geq 0}$, where $N_n(t)$ denotes the number of lineages at time $t \geq 0$. Thus $N_n(0) = n$ and $N_n(\tilde{\tau}_n) = 1$. For convenience we set $N_n(t) := 1$ for $t > \tilde{\tau}_n$. Then for $b = 2, \dots, n$ the numbers

$$\lambda(b) = \sum_{k=2}^b \binom{b}{k} \lambda_{b,k}$$

give the jump rates of the process N_n and

$$\mu(b) = \sum_{k=2}^b (k-1) \binom{b}{k} \lambda_{b,k}$$

its rate of decrease, since a merger of k lineages results in a downward jump of the block counting process of size $k-1$. These two sequences can be naturally extended to positive continuous functions $\lambda, \mu : [2, \infty) \rightarrow \mathbb{R}$, see the formulas (4) and (5) below.

In this paper we focus on the class of Λ -coalescents without a dust component. To put it briefly, in this case the rate at which, within the n -coalescents a single lineage merges with some others, diverges as the sample size n tends to infinity. They are characterized by the condition (Pitman [20])

$$\int_{[0,1]} \frac{\Lambda(dp)}{p} = \infty .$$

In particular, they cover Λ -coalescents coming down from infinity. These are the coalescents with the property that the absorption times $\tilde{\tau}_n$ are bounded in probability uniformly in n . They are distinguished by the criterion (Schweinsberg [22])

$$\int_2^\infty \frac{dx}{\mu(x)} < \infty .$$

We shall analyse the lengths of the whole n -coalescents as well as of different parts. They play an important role in the infinite sites model introduced by Kimura [17]. In this model each mutation effects a different site on the DNA. The mutations are distributed along the branches of the coalescent depending on their appearance in the past. Mathematically they build a homogeneous Poisson point process with rate $\theta > 0$. Their total number $S(n)$ counts the segregating sites in the sample of size n underlying the coalescent and are thus closely tied to its total length ℓ_n . Mutations which are located on an external branch appear only once in the sample, these are the singleton polymorphisms (see Wakeley [25], page 103). Accordingly their number is linked to the total length $\tilde{\ell}_n$ of all external branches. Likewise the number of mutations visible repeatedly in the sample corresponds to the total internal length $\check{\ell}_n$.

Notation. For two sequences A_n and B_n of positive random variables we write $A_n \stackrel{1}{\sim} B_n$ and $A_n \stackrel{P}{\sim} B_n$ if the sequence A_n/B_n converges to 1 in the L_1 -sense or in probability, respectively. The notation $A_n = O_P(B_n)$ means that the sequence A_n/B_n is tight and $A_n = o_P(B_n)$ that A_n/B_n converges to 0 in probability.

Now we come to the main results of this paper. As a first notion let

$$\ell_n := \int_0^{\tilde{\tau}_n} N_n(t) dt, \quad n \geq 1,$$

be the total length of the coalescent tree.

Theorem 1. *Assume that the Λ -coalescent has no dust component. Then as $n \rightarrow \infty$*

$$\ell_n \stackrel{1}{\sim} \int_2^n \frac{x}{\mu(x)} dx. \quad (1)$$

This is an intuitive approximation: if the counting process N_n takes the value x , then there are currently x lines and $1/\mu(x)$, the reciprocal of the rate of decrease, indicates how long on the average N_n will stay close to x . Observe that the right-hand integral diverges for $n \rightarrow \infty$: because of Lemma 1 (i) below the function $\mu(x)/(x(x-1))$ is decreasing, thus $\int_2^n \frac{x}{\mu(x)} dx \geq \frac{2}{\mu(2)} \log(n-1)$. This lower bound is attained by the Kingman coalescent.

Berestycki et al [3] conjectured Theorem 1 and proved it for Λ -coalescents coming down from infinity in the framework of convergence in probability by using instead of $\mu(x)$ the asymptotically equivalent quantity

$$\psi(x) := \int_{[0,1]} (e^{-xp} - 1 + xp) \frac{\Lambda(dp)}{p^2}.$$

For the Kingman coalescent the result was earlier obtained by Watterson [26] and for the Bolthausen-Sznitman coalescent by Drmota et al [10].

In the context of the infinite sites model Theorem 1 may be restated directly in terms of the number $S(n)$ of segregating sites as

$$S(n) \stackrel{1}{\sim} \theta \int_2^n \frac{x}{\mu(x)} dx,$$

due to the assumption that mutations appear according to a homogeneous Poisson point process with rate θ .

Next we define the total external length

$$\widehat{\ell}_n := \int_0^{\tilde{\tau}_n} \widehat{N}_n(t) dt,$$

where $\widehat{N}_n(t)$ denotes the number of external branches extant at time t . In contrast to the previous theorem the following result contains a statement only of convergence in probability (but see Theorem 4 below showing that, indeed, L_1 -convergence holds under the stronger conditions of that theorem).

Theorem 2. *Suppose that the Λ -coalescent has no dust component. Then as $n \rightarrow \infty$*

$$\widehat{\ell}_n \stackrel{P}{\sim} \frac{n^2}{\mu(n)}. \quad (2)$$

An intuitive explanation goes as follows: μ_n/n is the rate of decrease per individual at time 0. Therefore it is plausible that each external branch length is of order $n/\mu(n)$, and the external branch length in total results approximately in $n^2/\mu(n)$. The sequence $\mu(n)/n^2$, $n \geq 1$, has the limit $\Lambda(\{0\})/2$, see formula (8) below. Thus the total external lengths diverge in probability if and only if $\Lambda(\{0\}) = 0$. We point out that $x/\mu(x)$ is a decreasing function (see Lemma 1 (i) below), therefore the integral appearing in (1) exceeds the corresponding term in (2).

The proof rests on a close relation between the functions λ and μ which seems not to be observed until now. To describe it let us introduce another function $\kappa : [2, \infty) \rightarrow \mathbb{R}$ given by

$$\kappa(x) := \frac{\mu(x)}{x},$$

which could be named the rate of decrease per capita. Then we have the approximation

$$\lambda(x) \sim x^2 \kappa'(x)$$

as $x \rightarrow \infty$ (see Lemma 1 (ii) below).

In the special case of Beta-coalescents Theorem 2 follows from [7] and [9]. For Λ -coalescents with a dust component the picture is rather different. Then ℓ_n/n as well as $\widehat{\ell}_n/n$ converge in distribution to one and the same non-degenerate limit law, see [19]. Fluctuation results on the total length or the total external length of Λ -coalescents with no dust component are known only in special cases [7, 10, 13, 14, 24].

Theorem 2 again allows a reformulation in terms of the infinite site model. Letting $M_1(n)$ be the number of singletons, we obtain for $\Lambda(\{0\}) = 0$

$$M_1(n) \stackrel{P}{\sim} \theta \frac{n^2}{\mu(n)},$$

whereas for $\Lambda(\{0\}) > 0$ it follows that $M_1(n)$ is asymptotically Poisson with parameter $2\theta/\Lambda(\{0\})$.

Remark. From our proofs we will gain further insight in the structure of the coalescents. Let for $0 < c < 1$

$$\tilde{\rho}_n := \inf\{t \geq 0 : N_n(t) \leq cn\}$$


be the first moment, when the number of lineages falls below cn , and let

$$\ell_n^* := \int_0^{\tilde{\rho}_n} N_n(t) dt, \quad \widehat{\ell}_n^* := \int_0^{\tilde{\rho}_n} \widehat{N}_n(t) dt$$

be the respective lengths up to this moment. Then from (23) and (32) below we have

$$\ell_n^* \stackrel{1}{\sim} \int_{cn}^n \frac{x}{\mu(x)} dx \quad \text{and} \quad \widehat{\ell}_n^* \stackrel{P}{\sim} (1-c) \frac{n^2}{\mu(n)}.$$

The picture gives an illustration.



ℓ_n^* and $\widehat{\ell}_n^*$ are the areas of the total grey region and of the lighter part, respectively. \square

It is tempting to expect an analogous result for the total internal length

$$\check{\ell}_n := \ell_n - \widehat{\ell}_n .$$

This is certainly true in cases where the total external length $\widehat{\ell}_n$ does not exceed the total internal length $\check{\ell}_n$, then we have

$$\check{\ell}_n \stackrel{P}{\sim} \int_2^n \frac{x}{\mu(x)} dx - \frac{n^2}{\mu(n)} \sim \int_2^n \left(\frac{x}{\mu(x)} - \frac{n}{\mu(n)} \right) dx .$$

To provide some examples let us introduce the following class of coalescents.

Definition. We call the Λ -coalescent regularly varying with exponent $0 \leq \alpha \leq 2$ if $\Lambda(\{0\}) = 0$ and if as $y \rightarrow 0$

$$\int_{(y,1]} \frac{\Lambda(dp)}{p^2} \sim y^{-\alpha} L\left(\frac{1}{y}\right)$$

with a function L slowly varying at infinity.

This generalises the notion of Berestycki et al [3] of a strong regularly varying coalescent.

Examples. (i) The Kingman case: If $\Lambda(\{0\}) > 0$, then $\mu(x)/x^2 \sim \Lambda(\{0\})/2$, thus

$$\ell_n \stackrel{1}{\sim} \frac{2 \log n}{\Lambda(\{0\})} \quad \text{and} \quad \widehat{\ell}_n \stackrel{P}{\sim} \frac{2}{\Lambda(\{0\})}$$

as $n \rightarrow \infty$. Here the internal total length completely dominates the external ones, and we have

$$\check{\ell}_n \stackrel{1}{\sim} \frac{2 \log n}{\Lambda(\{0\})} .$$

(ii) Regularly varying coalescents with exponent $1 < \alpha < 2$ come down from infinity as can be easily checked by the above criterion. They fulfil (see Lemma 2 (ii) below)

$$\mu(x) \sim \frac{\Gamma(2-\alpha)}{\alpha-1} x^\alpha L(x)$$

as $x \rightarrow \infty$. Hence (having in mind how to integrate regularly varying functions, see Thm. 1 (b), Section VIII.9 in Feller [11])

$$\ell_n \stackrel{1}{\sim} \frac{\alpha-1}{(2-\alpha)\Gamma(2-\alpha)} \frac{n^{2-\alpha}}{L(n)} \quad \text{and} \quad \widehat{\ell}_n \stackrel{P}{\sim} \frac{\alpha-1}{\Gamma(2-\alpha)} \frac{n^{2-\alpha}}{L(n)}$$

and consequently

$$\check{\ell}_n \stackrel{P}{\sim} \frac{(\alpha - 1)^2}{(2 - \alpha)\Gamma(2 - \alpha)} \frac{n^{2-\alpha}}{L(n)} \stackrel{P}{\sim} \frac{\alpha - 1}{2 - \alpha} \widehat{\ell}_n .$$

Here the external and the internal length are of the same order. As an application, one may use the quantity $2 - \widehat{\ell}_n/\ell_n$ as an estimator for the parameter α . In the special case of Beta($2 - \alpha, \alpha$)-coalescents this was already discussed in more detail in Example 9 of [16]. \square

In some other cases $\check{\ell}_n$ is of smaller order than $\widehat{\ell}_n$. Then a large part of internal length will be located close to the coalescent tree's root, where extremely large mergers may take over (which is definitely the case for coalescents not coming down from infinity). Still the internal length may obey the law of large numbers suggested above. The following theorem presents a situation of special interest. Define for a slowly varying function L and for $x \geq 1$

$$L^*(x) := \int_1^x \frac{L(y)}{y} dy .$$

Theorem 3. *Assume that the Λ -coalescent has no dust component and is regularly varying with exponent $\alpha = 1$. Then*

$$\check{\ell}_n \stackrel{1}{\sim} \int_2^n \left(\frac{x}{\mu(x)} - \frac{n}{\mu(n)} \right) dx$$

and

$$\int_2^n \left(\frac{x}{\mu(x)} - \frac{n}{\mu(n)} \right) dx \sim \frac{nL(n)}{L^*(n)^2}$$

as $n \rightarrow \infty$.

Below we prove (Lemma 2 (ii)) that the function L^* is slowly varying at infinity, too, and that $L(x) = o(L^*(x))$ as $x \rightarrow \infty$. In comparison with Theorem 2 we see that $\check{\ell}_n = o_P(\widehat{\ell}_n)$ for regularly varying coalescents with exponent 1. In particular, for the Bolthausen-Sznitman coalescent Theorems 2 and 3 yield

$$\widehat{\ell}_n \stackrel{P}{\sim} \frac{n}{\log n} \quad \text{and} \quad \check{\ell}_n \stackrel{1}{\sim} \frac{n}{\log^2 n} .$$

These approximations were already obtained in Dhersin and Möhle [9] and Kersting, Pardo and Siri-Jégousse [15], respectively.

Our last object of investigation concerns lengths of higher order. A branch of order $a \geq 2$ is by definition an internal branch carrying a subtree with a leaves out of the original sample. In this context we consider external branches as branches of order 1. Denote the number of branches of order $a \geq 1$ present at time $t \geq 0$ as $\widehat{N}_{n,a}(t)$, notably $\widehat{N}_{n,1}(t) = \widehat{N}_n(t)$. Then the total length of all these branches is given by

$$\widehat{\ell}_{n,a} = \int_0^{\bar{\tau}_n} \widehat{N}_{n,a}(t) dt , \quad a \geq 1 .$$

Theorem 4. *Suppose that the Λ -coalescent has no dust component and is regularly varying with exponent $\alpha = 1$. Then*

$$\widehat{\ell}_{n,1} \stackrel{1}{\sim} \frac{n}{L^*(n)},$$

whereas for $a \geq 2$

$$\widehat{\ell}_{n,a} \stackrel{1}{\sim} \frac{1}{(a-1)a} \frac{nL(n)}{L^*(n)^2}$$

as $n \rightarrow \infty$.

These formulas have interesting applications to the site frequency spectrum. It consists of the counts $M_a(n)$, $a \geq 1$, specifying the numbers of mutations located on branches of order a , which can be distinguished in the various DNAs of the sample. The theorem yields

$$M_1(n) \stackrel{1}{\sim} \theta \frac{n}{L^*(n)} \quad \text{and} \quad M_a(n) \stackrel{1}{\sim} \frac{\theta}{(a-1)a} \frac{nL(n)}{L^*(n)^2}, \quad a \geq 2.$$

A corresponding result in the so-called infinite allele model was obtained by Basdevant and Goldschmidt [1]. They deal with the allele frequency spectrum and consider the special case of the Bolthausen-Sznitman coalescent. Analogue results for Beta-coalescents coming down from infinity were presented by Berestycki, Berestycki and Schweinsberg [4] and more generally for strongly regular varying coalescents with exponent $1 < \alpha < 2$ by Berestycki, Berestycki and Limic [3]. We conjecture that these results can be further extended to regular varying coalescents. Computational procedures for the general site frequency spectrum are due to Spence et al [23].

Theorem 4 illustrates that mutations show mainly up on external branches, whereas the tree structure of the coalescent becomes visible only at branches of higher order. This reflects that for regularly varying coalescents with exponent $\alpha = 1$ mergers occur preferentially at a late time and close to the MRCA. Our proof will show that with probability asymptotically equal to 1 any mutation seen in exactly $a \geq 2$ individuals stems from an internal branch of order a arose from one single merger. This is similar to findings of Basdevant and Goldschmidt.

Closing this introduction, we briefly discuss our methods of proof, which differ from other approaches in the literature. They rest upon L_2 -considerations and elementary martingale estimates, and they may well find further applications as indicated in two examples below. These methods deal with the time-discrete Markov chain $n = X_0 > X_1 > \dots > X_{\tau_n} = 1$, embedded in the Markov process N_n (or more generally with decreasing Markov chains). Let

$$\Delta_i := X_{i-1} - X_i, \quad i \geq 1,$$

denote its downwards jump size resulting from the i -th merger. We shall present different laws of large numbers for expressions of the form $\sum_{i=0}^{\rho_n-1} f(X_i)$, with some function $f : [2, \infty) \rightarrow \mathbb{R}$ and with stopping times ρ_n of the form

$$\rho_n := \min\{i \geq 0 : X_i \leq r_n\},$$

where r_n , $n \geq 1$, is some sequence of positive numbers.

Our approach specifies the following intuition:

$$\sum_{i=0}^{\rho_n-1} f(X_i) \approx \sum_{i=0}^{\rho_n-1} f(X_i) \frac{\Delta_{i+1}}{\nu(X_i)} \approx \int_{r_n}^n f(x) \frac{dx}{\nu(x)} \quad (3)$$

with

$$\nu(x) := \frac{\mu(x)}{\lambda(x)}, \quad x \geq 2,$$

thus

$$\nu(b) = \sum_{k=2}^b (k-1) \binom{b}{k} \frac{\lambda_{b,k}}{\lambda(b)} = \sum_{k=2}^b (k-1) \mathbf{P}(\Delta_{i+1} = k-1 \mid X_i = b)$$

or

$$\nu(b) = \mathbf{E}[\Delta_{i+1} \mid X_i = b], \quad b \geq 2.$$

The rationale behind this intuition is that the differences of both sums in (3) are small, because they stem from a martingale, whereas the second sum may be considered as a Riemann approximation of the right-hand integral. This latter approximation requires that the jump sizes Δ_{i+1} are small compared to the values X_i of the Markov chain and is only ensured if the time

$$\tilde{\rho}_n := \inf\{t \geq 0 : N_n(t) = X_{\rho_n}\} = \inf\{t \geq 0 : N_n(t) \leq r_n\}$$

of entrance into the interval $[1, r_n]$ converges to 0 in probability. (Observe that these random times generalise the notion used in the above remark.) Thus we strive towards small time approximations.

The quadratic variation of the just mentioned martingale will be estimated along the following lines: Since $\Delta_{i+1} = o_P(X_i)$ in the range of the small time approximation, we have

$$\sum_{i=0}^{\rho_n-1} f(X_i)^2 \frac{\Delta_{i+1}^2}{\nu(X_i)^2} = o_P\left(\sum_{i=0}^{\rho_n-1} f(X_i)^2 \frac{X_i \Delta_{i+1}}{\nu(X_i)^2}\right) = o_P\left(\max_{r_n \leq x \leq n} \frac{xf(x)}{\nu(x)} \sum_{i=0}^{\rho_n-1} f(X_i) \frac{\Delta_{i+1}}{\nu(X_i)}\right).$$

Under suitable conditions not only the right-hand sum but also the maximum is of order $\int_{r_n}^n f(x) \frac{dx}{\nu(x)}$ resulting in

$$\sum_{i=0}^{\rho_n-1} f(X_i)^2 \frac{\Delta_{i+1}^2}{\nu(X_i)^2} = o_P\left(\left(\int_{r_n}^n f(x) \frac{dx}{\nu(x)}\right)^2\right).$$

According to this pattern we may control the martingale's quadratic variation and its fluctuations.

The paper is organized as follows: Section 2 deals in detail with the rate functions. Section 3 contains our general laws of large numbers. Finally, our theorems are proved in Sections 4 to 7.

2 Properties of the rate functions

We extend the above sequences $\lambda(b)$ and $\mu(b)$, $b \geq 2$, of rates to positive continuous functions $\lambda, \mu : [2, \infty) \rightarrow \mathbb{R}$ via the definitions

$$\lambda(x) := \int_{[0,1]} (1 - (1-p)^x - xp(1-p)^{x-1}) \frac{\Lambda(dp)}{p^2}, \quad (4)$$

$$\mu(x) := \int_{[0,1]} (xp - 1 + (1-p)^x) \frac{\Lambda(dp)}{p^2}, \quad (5)$$

with $x \geq 2$. Note that $\mu(x) \geq \lambda(x)$ for all $x \geq 2$, since the integrands fulfil the corresponding inequality. Recall our notion

$$\kappa(x) := \frac{\mu(x)}{x}, \quad x \geq 2.$$

Lemma 1. (i) The functions $\lambda(x)$ and $\kappa(x)$, $x \geq 2$, are increasing in x , and the functions $\lambda(x)/(x(x-1))$ and $\kappa(x)/(x-1)$, $x \geq 2$, are decreasing.

(ii) For any $0 < \chi < 1$ as $x \rightarrow \infty$

$$\lambda(x) = x^2 \kappa'(x) (1 + O(x^{-\chi})) .$$

(iii) The Λ -coalescent has no dust component if and only if $\kappa(x) \rightarrow \infty$ as $x \rightarrow \infty$ and then $\kappa(x) = o(\lambda(x))$.

Proof. (i) The function $(1-p)^x + xp(1-p)^{x-1} = (1-p)^{x-1}(1+(x-1)p)$, $x \geq 2$, respectively its logarithm, has a negative derivative. From (4) we thus obtain monotonicity of $\lambda(x)$. Moreover, a partial integration yields

$$\frac{\lambda(x)}{x(x-1)} = \frac{\Lambda(\{0\})}{2} + \int_0^1 y(1-y)^{x-2} \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy \quad (6)$$

implying that $\lambda(x)/(x(x-1))$ is decreasing for $x \geq 2$.

Similarly, from (5) and a partial integration we have

$$\kappa(x) = \frac{\mu(x)}{x} = \frac{x-1}{2} \Lambda(\{0\}) + \int_0^1 (1-(1-y)^{x-1}) \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy , \quad (7)$$

which is increasing in x , and by another partial integration

$$\frac{\mu(x)}{x(x-1)} = \frac{\Lambda(\{0\})}{2} + \int_0^1 (1-z)^{x-2} \int_z^1 \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy dz , \quad (8)$$

a decreasing function in x .

(ii) From (7) we have

$$\kappa'(x) = \frac{\Lambda(\{0\})}{2} + \int_0^1 (1-y)^{x-1} \log \frac{1}{1-y} \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy . \quad (9)$$

From concavity $(1-y) \log \frac{1}{1-y} \leq y$ for $y \in [0, 1]$, also $(1-y) \log \frac{1}{1-y} \sim y + O(y^2)$ for $y \rightarrow 0$. Thus, combining (6) and (9), for $0 < \chi < 1$ and a suitable $c \geq 1$

$$\begin{aligned} 0 &\leq \frac{\lambda(x)}{x(x-1)} - \kappa'(x) \\ &\leq c \int_0^{x^{-\chi}} y^2 (1-y)^{x-2} \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy + \int_{x^{-\chi}}^1 y(1-y)^{x-2} \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy \\ &\leq cx^{-\chi} \int_0^{x^{-\chi}} y(1-y)^{x-2} \int_y^1 \frac{\Lambda(dp)}{p^2} dy + (1-x^{-\chi})^{x-2} \int_{x^{-\chi}}^1 y \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy \\ &\leq cx^{-\chi} \int_0^1 y(1-y)^{x-2} \int_y^1 \frac{\Lambda(dp)}{p^2} dy + e^{-x^{-\chi}(x-2)} \int_0^1 y \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy \\ &\leq cx^{-\chi} \frac{\lambda(x)}{x(x-1)} + e^{2-x^{1-\chi}} \Lambda([0, 1]) . \end{aligned}$$

Since λ is an increasing function, this implies

$$\kappa'(x) = \frac{\lambda(x)}{x^2} + O\left(\frac{\lambda(x)}{x^{2+\chi}}\right),$$

which is equivalent to our claim.

(iii) From (5) and the definition of κ it follows by monotone convergence that

$$\kappa(x) \rightarrow \int_{[0,1]} \frac{\Lambda(dp)}{p}.$$

This formula implies our first claim. As to the second one we have for $2 \leq x_0 \leq x$

$$\kappa(x) = \kappa(x_0) + \int_{x_0}^x \kappa'(y) dy.$$

From part (ii) and since λ is increasing, if x_0 is sufficiently large

$$\kappa(x) \leq \kappa(x_0) + 2 \int_{x_0}^x \frac{\lambda(y)}{y^2} dy \leq \kappa(x_0) + 2\lambda(x) \int_{x_0}^x \frac{dy}{y^2} \leq \kappa(x_0) + 2\frac{\lambda(x)}{x_0}.$$

Thus, from $\kappa(x) \rightarrow \infty$ it follows that $\lambda(x) \rightarrow \infty$ as $x \rightarrow \infty$, and we obtain for any $x_0 \geq 2$

$$\limsup_{x \rightarrow \infty} \frac{\kappa(x)}{\lambda(x)} \leq \frac{2}{x_0}.$$

This implies our second claim. □

First consequences of the previous results are contained in the next lemma.

Lemma 2. *Let the Λ -coalescent be regularly varying with exponent α .*

(i) *If $0 \leq \alpha < 2$, then*

$$\lambda(x) \sim \Gamma(2 - \alpha)x^\alpha L(x)$$

as $x \rightarrow \infty$, and for $k \geq 2$

$$\binom{b}{k} \frac{\lambda_{b,k}}{\lambda(b)} \rightarrow \frac{\alpha}{\Gamma(2 - \alpha)} \frac{\Gamma(k - \alpha)}{k!}$$

as $b \rightarrow \infty$.

(ii) *The Λ -coalescent has no dust component if and only if $\int_1^\infty x^{\alpha-2} L(x) dx = \infty$. Then $\alpha \geq 1$ and we have*

$$\kappa(x) \sim \begin{cases} \frac{\Gamma(2-\alpha)}{\alpha-1} x^{\alpha-1} L(x) & \text{for } 1 < \alpha < 2 \\ L^*(x) & \text{for } \alpha = 1 \end{cases}$$

as $x \rightarrow \infty$, with a function L^* given by

$$L^*(x) := \int_1^x \frac{L(y)}{y} dy, \quad x \geq 1.$$

L^* is slowly varying at infinity and satisfies $L(x) = o(L^*(x))$ as $x \rightarrow \infty$.

For $1 < \alpha < 2$ the convergence result on $\lambda_{b,k}$ has already been obtained by Delmas et al [8]. In this case they also have asymptotic estimates on $\lambda(x)$ and $\mu(x)$ under the assumption that the slowly varying function L is constant.

Proof. (i) Let $1 \leq k \leq b$ be natural numbers. Starting from the identity

$$1 - (1-p)^b - bp(1-p)^{b-1} - \dots - \binom{b}{k} p^k (1-p)^{b-k} = \binom{b}{k} (b-k) \int_0^p y^k (1-y)^{b-k-1} dy ,$$

we obtain

$$\begin{aligned} \lambda(b) - \sum_{j=2}^k \binom{b}{j} \lambda_{b,j} &= \binom{b}{k} (b-k) \int_{[0,1]} \int_0^p y^k (1-y)^{b-k-1} dy \frac{\Lambda(dp)}{p^2} \\ &= \binom{b}{k} (b-k) \int_0^1 y^k (1-y)^{b-k-1} \int_{(y,1]} \frac{\Lambda(dp)}{p^2} dy . \end{aligned}$$

Taking account of the definition of regular varying coalescents, it is no loss of generality to specify the function L in such a way that $\int_{(y,1]} p^{-2} \Lambda(dp) = y^{-\alpha} L(1/y)$ for $0 < y \leq 1$. It follows that

$$\begin{aligned} \lambda(b) - \sum_{j=2}^k \binom{b}{j} \lambda_{b,j} &= \binom{b}{k} (b-k) \int_0^1 y^{k-\alpha} (1-y)^{b-k-1} L\left(\frac{1}{y}\right) dy \\ &= \binom{b}{k} (b-k) b^{\alpha-k-1} L(b) \int_0^b z^{k-\alpha} \left(1 - \frac{z}{b}\right)^{b-k-1} \frac{L(b/z)}{L(b)} dz , \end{aligned}$$

where for $k = 1$ we set the value of the left-hand sum equal to 0.

Since the function L is slowly varying, the right-hand integrand converges pointwise to the limit $z^{k-\alpha} e^{-z}$. Also by the fundamental representation theorem of slowly varying functions (see Feller [11] Section VIII.9, Corollary) we have $L(b) \sim c \exp(\int_0^b \eta(z) z^{-1} dz)$ with a constant $c > 0$ and a function $\eta(z) = o(z)$ as $z \rightarrow \infty$. This implies that for any $\varepsilon > 0$ we have $L(b/z) \leq z^\varepsilon L(b)$ for $z \geq 1$ and $L(b/z) \leq z^{-\varepsilon} L(b)$ for $z \leq 1$ if only b is sufficiently large. Hence, for large b we may dominate the above integrand by the function $z^{k-\alpha} \max(z^\varepsilon, z^{-\varepsilon}) e^{-z/2}$. By the assumption $\alpha < 2$ it is integrable for $k \geq 1$ if we choose ε small enough. Thus by dominated convergence

$$\lambda(b) - \sum_{j=2}^k \binom{b}{j} \lambda_{b,j} \sim \frac{b^\alpha L(b)}{k!} \int_0^\infty z^{k-\alpha} e^{-z} dz = \frac{b^\alpha L(b) \Gamma(k - \alpha + 1)}{k!}$$

as $b \rightarrow \infty$, or in other terms

$$\lambda(b) \sim \Gamma(2 - \alpha) b^\alpha L(b) \quad \text{and} \quad \binom{b}{k} \lambda_{b,k} \sim \frac{\alpha \Gamma(k - \alpha)}{k!} b^\alpha L(b)$$

for $k \geq 2$. This implies our claim.

(ii) From part (i) and Lemma 1 (ii) we obtain

$$\kappa'(x) \sim \Gamma(2 - \alpha) x^{\alpha-2} L(x)$$

as $x \rightarrow \infty$. In view of Lemma 1 (iii) we are in the dustless case if and only if $\kappa(x) \rightarrow \infty$, that is if and only if the integral $\int_1^\infty x^{\alpha-2} L(x) dx$ is divergent. This implies $\alpha \geq 1$ and the claimed asymptotic formulas for κ .

Finally, since L is slowly varying at infinity, we have for $c > 1$

$$L^*(cx) - L^*(x) = \int_x^{cx} \frac{L(y)}{y} dy \sim L(x) \int_x^{cx} \frac{1}{y} dy = L(x) \log c \quad (10)$$

as $x \rightarrow \infty$. Because L^* is increasing, this implies for any $d > 1$ and large x

$$0 \leq \frac{L^*(cx)}{L^*(x)} - 1 \leq \frac{L^*(cx) - L^*(x)}{L^*(x) - L^*(x/d)} \sim \frac{\log c}{\log d}$$

as $x \rightarrow \infty$. Since d may be chosen arbitrarily large, it follows that L^* is slowly varying. Consequently, choosing $c = e$ in (10),

$$\frac{L(x)}{L^*(x)} \sim \frac{L^*(ex) - L^*(x)}{L^*(x)} = o(1) .$$

This finishes our proof. □

Examples. We consider regularly varying coalescents with exponent $\alpha = 1$.

(i) Let $L(x) = (\log x)^\chi$ with exponent $\chi > -1$. Then

$$L^*(x) = \int_0^{\log x} y^\chi dy = \frac{1}{\chi + 1} (\log x)^{\chi+1} = \frac{1}{\chi + 1} L(x) \log x .$$

For $-1 < \chi \leq 0$ these coalescents neither have a dust component nor come down from infinity, and for $\chi > 0$ they come down from infinity. The case $\chi = 0$ covers the Bolthausen-Sznitman coalescent.

(ii) Let $L(x) = e^{(\log x)^\chi}$ with $0 < \chi < 1$. Then

$$L^*(x) = \int_0^{\log x} e^{y^\chi} dy \sim \frac{1}{\chi} e^{(\log x)^\chi} (\log x)^{1-\chi} = \frac{1}{\chi} L(x) (\log x)^{1-\chi} .$$

These coalescents come down from infinity. □

3 Some general laws of large numbers

The laws of large numbers of this section apply not only to branch lengths of Λ -coalescents. As explained in the introduction, they concern the discrete-time Markov chains $X = (X_i)_{i \in \mathbb{N}_0}$ embedded in the lineage counting processes. For notational ease we do not account here for the dependence of the chains X on n . Thus $n = X_0 > X_1 > \dots > X_{\tau_n-1} > X_{\tau_n} = 1$ denote the states which the process N_n is successively visiting, and $\tau_n := \min\{i \geq 0 : X_i = 1\}$ is the respective number of merging events. For convenience we set $X_i = 1$ for all natural numbers $i > \tau_n$. Also let for $i \geq 1$ be W_i the waiting times of the process N_n in the states X_i and

$$\Delta_i := X_{i-1} - X_i .$$

Recall that

$$\mathbf{E}[\Delta_{i+1} \mid X_i = b] = \nu(b) := \frac{\mu(b)}{\lambda(b)} .$$

For sequences $r_n, s_n \geq 2$, $n \geq 1$, of positive numbers we set

$$\rho_n := \min\{i \geq 0 : X_i \leq r_n\} , \quad \sigma_n := \min\{i \geq 0 : X_i \leq s_n\} , \quad \tilde{\rho}_n := \inf\{t \geq 0 : N_n(t) \leq r_n\} .$$

Proposition 1. *Assume that the coalescent has no dust component. Let $f : [2, \infty) \rightarrow \mathbb{R}$ be a non-negative function with the property that $x^\beta f(x)$ is increasing and $x^{-\beta} f(x)$ is decreasing in x for some $\beta > 0$. Let $2 \leq r_n \leq s_n \leq n$, $n \geq 1$, be two sequences of numbers fulfilling*

$$r_n \leq \gamma s_n \tag{11}$$

for all $n \geq 1$ and some $\gamma < 1$. Also assume that

$$\tilde{\rho}_n \xrightarrow{P} 0 \tag{12}$$

as $n \rightarrow \infty$. Then, as $n \rightarrow \infty$

$$\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) \stackrel{1}{\sim} \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} .$$

Also, as $n \rightarrow \infty$

$$\mathbf{E} \left[\sum_{i=\rho_n}^{\tau_n-1} f(X_i) \right] = O \left(\int_2^{r_n} f(x) \frac{dx}{\nu(x)} \right) .$$

Due to the assumption (12) this result addresses the coalescent's evolution in only a short initial time interval. This takes double effect: first, as seen from the next lemma, the chain is kept away from the occurrence of huge jumps Δ_i being of the same order as the chain's values. Second, the chain X is prevented from taking too small values where larger fluctuations may become obstructive. Note that for any Λ -coalescent the passage times $\inf\{t \geq 0 : N_n(t) \leq r\}$ below some number $r > 1$ are bounded away from 0 uniformly in $n \in \mathbb{N}$. Therefore the assumption (12) enforces that

$$r_n \rightarrow \infty \tag{13}$$

as $n \rightarrow \infty$. Actually, both requirements are equivalent if the coalescent comes down from infinity, otherwise the assumption (12) is the more incisive one.

Example: total number of mergers. If we choose $f \equiv 1$ and $s_n = n$, then $\sum_{i=0}^{\rho_n-1} f(X_i)$ is equal to the number of mergers up to time $\tilde{\rho}_n$. For coalescents coming down from infinity we may consider any divergent sequence $r_n \leq \gamma n$. In particular, since $\int_2^\infty \frac{dx}{\nu(x)} = \infty$, we may choose the r_n in such a way that $r_n = o\left(\int_{r_n}^n \frac{dx}{\nu(x)}\right)$. This implies that the number of mergers after the moment $\tilde{\rho}_n$ are of negligible order and that for the total number τ_n of mergers we have

$$\tau_n \stackrel{1}{\sim} \int_2^n \frac{dx}{\nu(x)} .$$

Away from coalescents coming down from infinity the scope of this formula is unclear. For the Bolthausen-Sznitman coalescent it is valid (see [12]). \square

The proof of Proposition 1 is prepared by the next lemma.

Lemma 3. (i) *If the coalescent has no dust component and if $\int_{r_n}^n \mu(x)^{-1} dx \rightarrow 0$, then as $n \rightarrow \infty$*

$$\mathbf{E}[\tilde{\rho}_n] \rightarrow 0 .$$

(ii) *If $\tilde{\rho}_n \xrightarrow{P} 0$, then for any $\eta > 0$ as $n \rightarrow \infty$*

$$\mathbf{P}(\Delta_{i+1} > \eta X_i \text{ for some } i < \rho_n) \rightarrow 0 .$$

Proof. (i) Given X the waiting times W_i are exponential with expectation $1/\lambda(X_i)$. Therefore

$$\mathbf{E}[\tilde{\rho}_n] = \mathbf{E}\left[\sum_{i=0}^{\rho_n-1} W_i\right] = \sum_{i=0}^{n-1} \mathbf{E}\left[\frac{1}{\lambda(X_i)}; X_i > r_n\right].$$

Also $\mathbf{E}[\Delta_{i+1} | X_i] = \nu(X_i)$ a.s., thus by the Markov property

$$\mathbf{E}[\tilde{\rho}_n] = \sum_{i=0}^{n-1} \mathbf{E}\left[\frac{\Delta_{i+1}}{\lambda(X_i)\nu(X_i)}; X_i > r_n\right] = \mathbf{E}\left[\sum_{i=0}^{\rho_n-1} \frac{\Delta_{i+1}}{\mu(X_i)}\right].$$

From Lemma 1 (i) we have that $\mu(x) = x\kappa(x)$ is increasing. Also $\Delta_{i+1} \leq X_i$, hence

$$\mathbf{E}[\tilde{\rho}_n] \leq \mathbf{E}\left[\sum_{i=0}^{\rho_n-2} \int_{X_{i+1}}^{X_i} \frac{dx}{\mu(x)} + \frac{X_{\rho_n-1}}{\mu(X_{\rho_n-1})}\right] = \mathbf{E}\left[\int_{X_{\rho_n-1}}^n \frac{dx}{\mu(x)} + \frac{1}{\kappa(X_{\rho_n-1})}\right],$$

and since κ is an increasing function, we end up with the estimate

$$\mathbf{E}[\tilde{\rho}_n] \leq \int_{r_n}^n \frac{dx}{\mu(x)} + \frac{1}{\kappa(r_n)}.$$

Our assumptions imply $r_n \rightarrow \infty$. Therefore, since there is no dust component, we have $\kappa(r_n) \rightarrow \infty$ by Lemma 1 (iii). This entails our claim.

(ii) For $b \geq 2/\eta$ we have

$$\begin{aligned} \mathbf{P}(\Delta_{i+1} > \eta X_i | X_i = b) &= \sum_{k > \eta b} \frac{1}{\lambda(b)} \int_{[0,1]} \binom{b}{k} p^k (1-p)^{b-k} \frac{\Lambda(dp)}{p^2} \\ &\leq \frac{1}{\lambda(b)} \int_{[0,1]} \sum_{k > \eta b} \frac{2}{\eta^2} \binom{b-2}{k-2} p^{k-2} (1-p)^{(b-2)-(k-2)} \Lambda(dp) \\ &\leq \frac{c}{\lambda(b)}, \end{aligned}$$

with $c := 2\Lambda([0,1])/\eta^2$.

Let $\chi_0 := 0$ and $\chi_i := W_0 + \dots + W_{i-1}$, $i \geq 1$, which is the moment of the i -th jump. Then

$$\begin{aligned} \mathbf{P}(\Delta_{i+1} > \eta X_i \text{ for some } i < \rho_n, \tilde{\rho}_n \leq 1) &\leq \mathbf{P}(\Delta_{i+1} > \eta X_i, X_i > r_n, \chi_i \leq 1 \text{ for some } i < n) \\ &\leq \sum_{i=0}^{n-1} \mathbf{E}[\mathbf{P}(\Delta_{i+1} > \eta X_i | X_i); X_i > r_n, \chi_i \leq 1] \\ &\leq \sum_{i=0}^{n-1} \mathbf{E}\left[\frac{c}{\lambda(X_i)}; X_i > r_n, \chi_i \leq 1\right]. \end{aligned}$$

Also, because $\lambda(x)$ is increasing, for $X_i \geq 2$

$$\mathbf{E}[W_i I_{\{W_i \leq 1\}} | X_i] = \int_0^1 t \lambda(X_i) e^{-\lambda(X_i)t} dt = \frac{1}{\lambda(X_i)} \int_0^{\lambda(X_i)} u e^{-u} du \geq \frac{d}{\lambda(X_i)}$$

with $d := \int_0^{\lambda(2)} ue^{-u} du > 0$. This allows for the estimate

$$\begin{aligned}
& \mathbf{P}(\Delta_{i+1} > \eta X_i \text{ for some } i < \rho_n, \tilde{\rho}_n \leq 1) \\
& \leq \frac{c}{d} \sum_{i=0}^{n-1} \mathbf{E} \left[W_i; X_i > r_n, W_i \leq 1, W_0 + \dots + W_{i-1} \leq 1 \right] \\
& \leq \frac{c}{d} \mathbf{E} \left[\sum_{i=0}^{\rho_n-1} W_i I_{\{W_0 + \dots + W_i \leq 2\}} \right] \\
& \leq \frac{c}{d} \mathbf{E} \left[2 \wedge \sum_{i=0}^{\rho_n-1} W_i \right] \\
& = \frac{c}{d} \mathbf{E} [2 \wedge \tilde{\rho}_n],
\end{aligned}$$

and consequently

$$\mathbf{P}(\Delta_{i+1} > \eta X_i \text{ for some } i < \rho_n) \leq \frac{c}{d} \mathbf{E} [\tilde{\rho}_n \wedge 2] + \mathbf{P}(\tilde{\rho}_n > 1).$$

Thus by assumption and dominated convergence our claim follows. \square

Proof of Proposition 1. (i) We start with some preliminary estimates. Since $x^{-\beta}f(x)$ is a decreasing function, we have for $0 < \chi < 1$, $0 \leq y \leq \chi z$ and $y \leq z - 2$

$$yf(z) \leq z^\beta \int_{z-y}^z x^{-\beta} f(x) dx \leq z^\beta (z-y)^{-\beta} \int_{z-y}^z f(x) dx \leq (1-\chi)^{-\beta} \int_{z-y}^z f(x) dx.$$

Similarly, using that $x^\beta f(x)$ is increasing, we obtain a lower bound. Altogether, for $0 < \chi < 1$, $0 \leq y \leq \chi z$ and $y \leq z - 2$

$$(1-\chi)^\beta \int_{z-y}^z f(x) dx \leq yf(z) \leq (1-\chi)^{-\beta} \int_{z-y}^z f(x) dx. \tag{14}$$

Moreover, for any $\varepsilon > 0$ there is an $\eta > 0$ such that

$$(1-\varepsilon) \int_{(1-\eta)r_n}^{s_n} f(x) dx \leq \int_{r_n}^{s_n} f(x) dx \leq (1+\varepsilon) \int_{r_n}^{(1-\eta)s_n} f(x) dx \tag{15}$$

for all n . We prove the left-hand inequality. Let a be the affine function mapping the interval $[(1-\eta)r_n, r_n]$ onto $[(1-\eta)r_n, r_n/\gamma]$. Substituting $y = a(x)$, this implies

$$dy = \frac{\eta + \gamma^{-1} - 1}{\eta} dx.$$

Moreover, since $\gamma < 1$, $a(x) \geq x$ for $x \geq (1-\eta)r_n$. Therefore by monotonicity of x^β and $x^\beta f(x)$

we have with γ as in (11)

$$\begin{aligned}
\int_{(1-\eta)r_n}^{r_n} f(x) dx &\leq ((1-\eta)r_n)^{-\beta} \int_{(1-\eta)r_n}^{r_n} f(x)x^\beta dx \\
&\leq ((1-\eta)r_n)^{-\beta} \int_{(1-\eta)r_n}^{r_n} f(a(x))a(x)^\beta dx \\
&= ((1-\eta)r_n)^{-\beta} \frac{\eta}{\eta + \gamma^{-1} - 1} \int_{(1-\eta)r_n}^{r_n/\gamma} f(y)y^\beta dy \\
&\leq (\gamma(1-\eta))^{-\beta} \frac{\eta}{\eta + \gamma^{-1} - 1} \int_{(1-\eta)r_n}^{r_n/\gamma} f(y) dy .
\end{aligned}$$

From condition (11) it follows that

$$\int_{(1-\eta)r_n}^{r_n} f(x) dx \leq (\gamma(1-\eta))^{-\beta} \frac{\eta}{\eta + \gamma^{-1} - 1} \int_{(1-\eta)r_n}^{s_n} f(x) dx \leq \varepsilon \int_{(1-\eta)r_n}^{s_n} f(x) dx$$

for sufficiently small $\eta > 0$. This implies the left-hand inequality of (15). The other one follows similarly, now using the monotonicity of $x^{-\beta}f(x)$.

We note that $(x-1)/x^2$ is a decreasing function for $x \geq 2$. Therefore, in view of Lemma 1 (i) the functions $\lambda(x)$ and $\mu(x)$ are increasing and $\lambda(x)/x^3$ and $\mu(x)/x^3$ are decreasing for $x \geq 2$. Thus the function $f(x)/\nu(x) = f(x)\lambda(x)/\mu(x)$ fulfils the same assumptions as $f(x)$, with β replaced by $\beta + 3$. Accordingly, we shall use the preceding estimates with $f(x)$ replaced by $f(x)/\nu(x)$ (and β replaced by $\beta + 3$).

(ii) Next we develop Riemann approximations of certain random sums. For $0 < \eta < 1$ and $b \geq 2$ let

$$\mu_\eta(b) := \sum_{2 \leq k \leq \eta b} (k-1) \binom{b}{k} \int_{[0,1]} p^k (1-p)^{b-k} \frac{\Lambda(dp)}{p^2} .$$

In view of the well-known formula for the second factorial moment of binomials,

$$\begin{aligned}
\mu(b) - \mu_\eta(b) &= \sum_{\eta b < k \leq b} (k-1) \binom{b}{k} \int_{[0,1]} p^k (1-p)^{b-k} \frac{\Lambda(dp)}{p^2} \\
&\leq \frac{1}{\eta b} \int_{[0,1]} \sum_{k=0}^b k(k-1) \binom{b}{k} p^k (1-p)^{b-k} \frac{\Lambda(dp)}{p^2} \\
&= \frac{1}{\eta b} \int_{[0,1]} b(b-1)p^2 \frac{\Lambda(dp)}{p^2} = \frac{(b-1)\Lambda([0,1])}{\eta} .
\end{aligned}$$

From Lemma 1 (iii) we have $\mu(b)/b \rightarrow \infty$ in the dustless case, hence we obtain for any $\eta > 0$

$$\frac{\mu_\eta(b)}{\mu(b)} \rightarrow 1 \tag{16}$$

as $b \rightarrow \infty$.

Also let

$$\nu_\eta(b) := \frac{\mu_\eta(b)}{\lambda(b)} = \mathbf{E}[\Delta_{i+1} I_{\{\Delta_{i+1} \leq \eta b\}} \mid X_i = b] .$$

Note that $\nu_\eta(b) > 0$ for $b \geq 2/\eta$. Therefore, for $0 < \eta < 1$ and natural numbers n satisfying $r_n \geq 2/\eta$ we may define the random variables

$$R_n = R_{n,\eta} := \sum_{i=\sigma_n}^{\rho_n-1} f(X_i) \frac{\Delta_{i+1} I_{\{\Delta_{i+1} \leq \eta X_i\}}}{\nu_\eta(X_i)} .$$

Given η these random variables are in view of (13) well-defined up to finitely many n . We shall use them below as an intermediate approximation to $\sum_{i=\sigma_n}^{\rho_n-1} f(X_i)$. We estimate R_n from above and below. From (14) with $z = X_i$, $y = \Delta_{i+1}$ and $\chi = \eta$ and with $f(x)/\nu(x)$ replacing $f(x)$ we have on the event that $\Delta_{i+1} \leq \eta X_i$

$$\frac{f(X_i)}{\nu(X_i)} \Delta_{i+1} \leq (1-\eta)^{-\beta-3} \int_{X_{i+1}}^{X_i} f(x) \frac{dx}{\nu(x)} ,$$

consequently

$$R_n \leq (1-\eta)^{-\beta-3} \sum_{i=\sigma_n}^{\rho_n-1} \frac{\nu(X_i)}{\nu_\eta(X_i)} I_{\{\Delta_{i+1} \leq \eta X_i\}} \int_{X_{i+1}}^{X_i} f(x) \frac{dx}{\nu(x)}$$

and by definition of ρ_n

$$R_n \leq (1-\eta)^{-\beta-3} \sup_{b \geq r_n} \frac{\mu(b)}{\mu_\eta(b)} \int_{r_n(1-\eta)}^{s_n} f(x) \frac{dx}{\nu(x)} . \quad (17)$$

Therefore, in view of (15) and (16) and since $r_n \rightarrow \infty$, there is for given $\varepsilon > 0$ an $\eta > 0$ fulfilling

$$R_n \leq (1+\varepsilon) \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} \quad (18)$$

for all n sufficiently large. Similarly, for given $\varepsilon > 0$ there is an $\eta > 0$ satisfying for large n the inequality

$$R_n \geq (1-\varepsilon) \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} \quad \text{on the event } \{\Delta_{i+1} \leq \eta X_i \text{ for all } i < \rho_n\} . \quad (19)$$

(iii) Now observe that the random variables $M_0 := 0$ and

$$M_k := \sum_{i=0}^{k \wedge \tau_n - 1} \left(f(X_i) \frac{\Delta_{i+1} I_{\{\Delta_{i+1} \leq \eta X_i\}}}{\nu_\eta(X_i)} - f(X_i) \right) , \quad k \geq 1 ,$$

build a martingale $M = (M_k)_{k \geq 0}$. The optional sampling theorem yields

$$\begin{aligned} \mathbf{E} \left[\left(R_n - \sum_{i=\sigma_n}^{\rho_n-1} f(X_i) \right)^2 \right] &= \mathbf{E} [(M_{\rho_n} - M_{\sigma_n})^2] \\ &\leq \mathbf{E} \left[\sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)^2 \Delta_{i+1}^2}{\nu_\eta(X_i)^2} I_{\{\Delta_{i+1} \leq \eta X_i\}} \right] \\ &\leq \eta \mathbf{E} \left[\sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)^2 X_i \Delta_{i+1}}{\nu_\eta(X_i)^2} I_{\{\Delta_{i+1} \leq \eta X_i\}} \right] . \end{aligned}$$

Letting x_n be the point where the function $xf(x)/\nu_\eta(x)$ takes its maximal value within the interval $[r_n, s_n]$, it follows

$$\mathbf{E}\left[\left(R_n - \sum_{i=\sigma_n}^{\rho_n-1} f(X_i)\right)^2\right] \leq \eta \frac{x_n f(x_n)}{\nu_\eta(x_n)} \mathbf{E}\left[\sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)\Delta_{i+1}}{\nu_\eta(X_i)} I_{\{\Delta_{i+1} \leq \eta X_i\}}\right] = \eta \frac{x_n f(x_n)}{\nu_\eta(x_n)} \mathbf{E}[R_n] .$$

By the assumption (11) it follows that there are numbers ξ_n , $n \geq 1$, such that

$$r_n \leq \gamma \xi_n \leq x_n \leq \xi_n \leq s_n .$$

Using monotonicity of $x^{\beta+3}f(x)/\nu(x)$ and (14) with $z = \xi_n$, $y = (1 - \gamma)\xi_n$, $\chi = 1 - \gamma$, it follows that

$$(1 - \gamma) \frac{x_n f(x_n)}{\nu(x_n)} \leq \left(\frac{\xi_n}{x_n}\right)^{\beta+2} (1 - \gamma) \frac{\xi_n f(\xi_n)}{\nu(\xi_n)} \leq \gamma^{-2\beta-5} \int_{\gamma \xi_n}^{\xi_n} f(x) \frac{dx}{\nu(x)} \leq \gamma^{-2\beta-5} \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} ,$$

hence

$$\mathbf{E}\left[\left(R_n - \sum_{i=\sigma_n}^{\rho_n-1} f(X_i)\right)^2\right] \leq \eta \frac{\gamma^{-2\beta-5}}{1 - \gamma} \sup_{x \geq 2/\eta} \frac{\nu(x)}{\nu_\eta(x)} \mathbf{E}[R_n] \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} .$$

Finally, the formulas (16) and (18) yield that for any $\varepsilon > 0$ there is an $\eta > 0$ fulfilling

$$\mathbf{E}\left[\left(R_n - \sum_{i=\sigma_n}^{\rho_n-1} f(X_i)\right)^2\right] \leq \varepsilon \left(\int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)}\right)^2 . \quad (20)$$

(iv) Putting pieces together, we obtain for given $\varepsilon > 0$ and $\eta > 0$

$$\begin{aligned} & \mathbf{P}\left(\left|\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) - \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)}\right| \geq \varepsilon \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)}\right) \\ & \leq \mathbf{P}\left(\left|R_n - \sum_{i=\sigma_n}^{\rho_n-1} f(X_i)\right| \geq \frac{\varepsilon}{2} \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)}\right) \\ & \quad + \mathbf{P}\left(\left|R_n - \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)}\right| \geq \frac{\varepsilon}{2} \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} , \Delta_{i+1} \leq \eta X_i \text{ for all } i < \rho_n\right) \\ & \quad + \mathbf{P}(\Delta_{i+1} > \eta X_i \text{ for some } i < \rho_n) . \end{aligned}$$

From (20) for suitably chosen $\eta > 0$ the first right-hand term becomes smaller than ε and from (18) and (19) the second one vanishes for large n . Consulting also Lemma 3 (ii), we arrive at

$$\mathbf{P}\left(\left|\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) - \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)}\right| \geq \varepsilon \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)}\right) \leq \varepsilon$$

for n large enough. This means that

$$\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) \stackrel{P}{\sim} \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} .$$

To show L_1 -convergence, it is by a convergence criterion due to F. Riesz sufficient to have

$$\mathbf{E} \left[\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) \right] \sim \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)}$$

as $n \rightarrow \infty$. From convergence in probability and Fatou's lemma we have

$$\liminf_{n \rightarrow \infty} \mathbf{E} \left[\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) \right] / \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} \geq 1 .$$

On the other hand, (18) yields for given $\varepsilon > 0$, suitable $\eta > 0$ and large n

$$\mathbf{E} \left[\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) \right] = \mathbf{E}[R_n] \leq (1 + \varepsilon) \int_{r_n}^{s_n} f(x) \frac{dx}{\nu(x)} .$$

This gives our first claim.

(v) For the second claim we again use the random variables R_n , now for some $\eta > 0$ (say $\eta = 1/2$) with $r_n = r := 2/\eta$. Recall that for this choice the R_n are well-defined for all n . From inequality (17) we obtain the estimate

$$R_n \leq (1 - \eta)^{-\beta-3} \sup_{b \geq 2/\eta} \frac{\mu(b)}{\mu_\eta(b)} \int_2^{s_n} f(x) \frac{dx}{\nu(x)} .$$

It follows

$$\mathbf{E} \left[\sum_{i=\sigma_n}^{\tau_n-1} f(X_i) \right] \leq \sum_{b < 2/\eta} f(b) + \mathbf{E} \left[\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) \right] = \sum_{b < 2/\eta} f(b) + \mathbf{E}[R_n] .$$

Putting both estimates together and then replacing σ_n by ρ_n (which is just a change in notation), we arrive at our second claim. \square

The next proposition presents a version of Proposition 1 in continuous time. As before, let $\tilde{\rho}_n$ and $\tilde{\sigma}_n$ denote the times, when the process N_n falls below r_n and s_n , respectively, while $\tilde{\tau}_n$ is the absorption time of N_n .

Proposition 2. *Under the assumptions of Proposition 1, as $n \rightarrow \infty$*

$$\int_{\tilde{\sigma}_n}^{\tilde{\rho}_n} f(N_n(t)) dt \stackrel{\sim}{\sim} \int_{r_n}^{s_n} f(x) \frac{dx}{\mu(x)} .$$

Also, as $n \rightarrow \infty$

$$\mathbf{E} \left[\int_{\tilde{\rho}_n}^{\tilde{\tau}_n} f(N_n(t)) dt \right] = O \left(\int_2^{r_n} f(x) \frac{dx}{\mu(x)} \right) .$$

Example. For $f(x) \equiv 1$ we obtain under the assumptions of Proposition 2

$$\tilde{\rho}_n - \tilde{\sigma}_n \stackrel{1}{\sim} \int_{r_n}^{s_n} \frac{dx}{\mu(x)},$$

in particular,

$$\tilde{\rho}_n \stackrel{1}{\sim} \int_{r_n}^n \frac{dx}{\mu(x)}$$

as $n \rightarrow \infty$. □

Proof of Proposition 2. We have

$$\int_{\tilde{\sigma}_n}^{\tilde{\rho}_n} f(N_n(t)) dt = \sum_{i=\sigma_n}^{\rho_n-1} f(X_i) W_i$$

and

$$\mathbf{E} \left[\sum_{i=\sigma_n}^{\rho_n-1} f(X_i) W_i \mid X \right] = \sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)}{\lambda(X_i)}.$$

Thus for any $\eta > 0$ by the Markov property

$$\mathbf{E} \left[\left(\int_{\tilde{\sigma}_n}^{\tilde{\rho}_n} f(N_n(t)) dt - \sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)}{\lambda(X_i)} \right)^2 \right] = \mathbf{E} \left[\sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)^2}{\lambda(X_i)^2} \right] = \mathbf{E}[R_n],$$

where we now set

$$R_n := \sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)^2}{\lambda(X_i)^2} \frac{\Delta_{i+1} I_{\{\Delta_{i+1} \leq \eta X_i\}}}{\nu_\eta(X_i)}.$$

Using (18) with $\varepsilon = 1$, it follows

$$\mathbf{E} \left[\left(\int_{\tilde{\sigma}_n}^{\tilde{\rho}_n} f(N_n(t)) dt - \sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)}{\lambda(X_i)} \right)^2 \right] \leq 2 \int_{r_n}^{s_n} \frac{f(x)^2}{\lambda(x)^2 \nu(x)} dx = 2 \int_{r_n}^{s_n} \frac{f(x)^2}{\lambda(x) \mu(x)} dx.$$

Furthermore, since we are in the dustless case, due to Lemma 1 (iii) we have

$$\mathbf{E} \left[\left(\int_{\tilde{\sigma}_n}^{\tilde{\rho}_n} f(N_n(t)) dt - \sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)}{\lambda(X_i)} \right)^2 \right] \leq o \left(\int_{r_n}^{s_n} \frac{x f(x)^2}{\mu(x)^2} dx \right).$$

Letting x_n be the point, where $x f(x) / \mu(x)$ takes its maximum in the interval $[r_n, s_n]$, we obtain in the same manner as in the preceding proof

$$\mathbf{E} \left[\left(\int_{\tilde{\sigma}_n}^{\tilde{\rho}_n} f(N_n(t)) dt - \sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)}{\lambda(X_i)} \right)^2 \right] = o \left(\frac{x_n f(x_n)}{\mu(x_n)} \int_{r_n}^{s_n} \frac{f(x)}{\mu(x)} dx \right) = o \left(\left(\int_{r_n}^{s_n} \frac{f(x)}{\mu(x)} dx \right)^2 \right).$$

On the other hand, Proposition 1 implies

$$\sum_{i=\sigma_n}^{\rho_n-1} \frac{f(X_i)}{\lambda(X_i)} \stackrel{1}{\sim} \int_{r_n}^{s_n} \frac{f(x)}{\lambda(x) \nu(x)} dx = \int_{r_n}^{s_n} \frac{f(x)}{\mu(x)} dx.$$

These last two formulas imply our first statement. Moreover

$$\mathbf{E} \left[\int_{\tilde{\rho}_n}^{\tilde{\tau}_n} f(N_n(t)) dt \right] = \mathbf{E} \left[\sum_{i=\rho_n}^{\tau_n-1} \frac{f(X_i)}{\lambda(X_i)} \right],$$

therefore our second claim follows from the second statement of Proposition 1. \square

Now we turn to the special case that $f(x) = 1/x$, $x \geq 2$, where Proposition 1 can be considerably extended. Here we content ourselves with the case $s_n = n$. Observe that the two next statements do not imply each other.

Proposition 3. *Assume that the Λ -coalescent has no dust component. Let $2 \leq r_n \leq n$, $n \geq 1$, be a sequence of numbers fulfilling*

$$r_n \leq \gamma n$$

for all n sufficiently large and some $\gamma < 1$. Also assume that

$$\tilde{\rho}_n \xrightarrow{P} 0$$

as $n \rightarrow \infty$. Then

$$\sum_{i=0}^{\rho_n-1} \frac{1}{X_i} \underset{1}{\sim} \log \frac{\kappa(n)}{\kappa(r_n)}$$

and

$$\max_{1 \leq j \leq \rho_n} \left| \sum_{i=0}^{j-1} \frac{1}{X_i} - \log \frac{\kappa(n)}{\kappa(X_j)} \right| = o_P(1).$$

Proof. (i) The first statement is a special case of Proposition 1: due to Lemma 1 (ii) we have $1/(x\nu(x)) = \lambda(x)/(x\mu(x)) \sim \kappa'(x)/\kappa(x)$ as $x \rightarrow \infty$. Therefore

$$\int_{r_n}^n \frac{dx}{x\nu(x)} \sim \int_{r_n}^n \frac{\kappa'(x) dx}{\kappa(x)} = \log \frac{\kappa(n)}{\kappa(r_n)}.$$

(ii) For the proof of the second statement we proceed along similar lines as in the proof of Proposition 1. Using the second factorial moment of a binomial distribution, we have as a first estimate

$$\begin{aligned} \mathbf{E} \left[\frac{\Delta_{i+1}^2}{X_i^2} \mid X_i = b \right] &= \frac{1}{b^2 \lambda(b)} \sum_{k=2}^b (k-1)^2 \binom{b}{k} \int_{[0,1]} p^k (1-p)^{b-k} \frac{\Lambda(dp)}{p^2} \\ &\leq \frac{1}{b^2 \lambda(b)} \int_{[0,1]} \sum_{k=0}^b k(k-1) \binom{b}{k} p^k (1-p)^{b-k} \frac{\Lambda(dp)}{p^2} \\ &= \frac{1}{b^2 \lambda(b)} \int_{[0,1]} p^2 b(b-1) \frac{\Lambda(dp)}{p^2} \\ &\leq \frac{\Lambda([0,1])}{\lambda(b)}. \end{aligned}$$

This bound yields for large n

$$\begin{aligned}
\mathbf{E} \left[\sum_{i=0}^{\rho_n-1} \frac{\Delta_{i+1}^2}{X_i^2 \nu(X_i)} \right] &\leq \sum_{i=0}^{n-1} \mathbf{E} \left[\frac{1}{\mu(X_i)} ; X_i > r_n \right] \Lambda([0, 1]) \\
&= \sum_{i=0}^{n-1} \mathbf{E} \left[\frac{\Delta_{i+1}}{\mu(X_i) \nu_{1/2}(X_i)} I_{\{\Delta_{i+1} \leq X_i/2\}} ; X_i > r_n \right] \Lambda([0, 1]) \\
&\leq c \mathbf{E} \left[\sum_{i=0}^{\rho_n-1} \frac{\Delta_{i+1}}{\mu(X_i) \nu(X_i)} I_{\{\Delta_{i+1} \leq X_i/2\}} \right]
\end{aligned}$$

with $c := \Lambda([0, 1]) \sup_{b \geq 4} \nu(b)/\nu_{1/2}(b) < \infty$. In view of Lemma 1 (i) the function

$$x \mapsto (x-1)\mu(x)\nu(x)/x = \kappa(x)^2/(\lambda(x)/x(x-1))$$

is increasing, which entails $\Delta_{i+1}/(\mu(X_i)\nu(X_i)) \leq \int_{X_{i+1}}^{X_i} x((x-1)\mu(x)\nu(x))^{-1} dx$. By means of Lemma 1 (ii) $\mu(x)\nu(x) \sim \kappa(x)^2/\kappa'(x)$, therefore

$$\mathbf{E} \left[\sum_{i=0}^{\rho_n-1} \frac{\Delta_{i+1}^2}{X_i^2 \nu(X_i)} \right] \leq c \int_{r_n/2}^n \frac{x dx}{(x-1)\mu(x)\nu(x)} \sim c \int_{r_n/2}^n \frac{\kappa'(x)}{\kappa^2(x)} dx \leq \frac{c}{\kappa(r_n/2)},$$

and consequently, since $\kappa(x) \rightarrow \infty$ in the dustless case,

$$\mathbf{E} \left[\sum_{i=0}^{\rho_n-1} \frac{\Delta_{i+1}^2}{X_i^2 \nu(X_i)} \right] = o(1).$$

(iii) Now we consider the martingale $M = (M_k)_{k \geq 0}$ given by $M_0 = 0$ a.s. and

$$M_k := \sum_{i=0}^{k \wedge \tau_n - 1} \left(\frac{\Delta_{i+1}}{(X_i - 1)\nu(X_i)} - \frac{1}{X_i - 1} \right), \quad k \geq 1.$$

By means of the optional sampling theorem, since ρ_n is a stopping time, we have, because $\nu(b) \geq 1$ for all $b \geq 2$,

$$\mathbf{E} [M_{\rho_n}^2] \leq \mathbf{E} \left[\sum_{i=0}^{\rho_n-1} \frac{\Delta_{i+1}^2}{X_i^2 \nu(X_i)^2} \right] = o(1).$$

Thus by means of Doob's maximal inequality

$$\max_{1 \leq j \leq \rho_n} \left| \sum_{i=0}^{j-1} \frac{\Delta_{i+1}}{(X_i - 1)\nu(X_i)} - \sum_{i=0}^{j-1} \frac{1}{X_i - 1} \right| = o_P(1).$$

Also for $j \leq \rho_n$

$$0 \leq \sum_{i=0}^{j-1} \frac{1}{X_i - 1} - \sum_{i=0}^{j-1} \frac{1}{X_i} \leq \sum_{m=X_{\rho_n-1}}^{\infty} \left(\frac{1}{m-1} - \frac{1}{m} \right) = \frac{1}{X_{\rho_n-1} - 1} \leq \frac{1}{r_n - 1},$$

and consequently

$$\max_{1 \leq j \leq \rho_n} \left| \sum_{i=0}^{j-1} \frac{\Delta_{i+1}}{(X_i - 1)\nu(X_i)} - \sum_{i=0}^{j-1} \frac{1}{X_i} \right| = o_P(1). \quad (21)$$

(iv) Note that in view of Lemma 1 (i) the function $(x-1)\nu(x) = \kappa(x)/(\lambda(x)/x(x-1))$ is increasing and $\nu(x)/(x-1)^2 = (x/(x-1)) \cdot (\kappa(x)/(x-1))/\lambda(x)$ is decreasing. For $X_i \geq 2$ this yields

$$\begin{aligned} 0 &\leq \frac{1}{(X_{i+1}-1)\nu(X_{i+1})} - \frac{1}{(X_i-1)\nu(X_i)} \\ &= \frac{(X_{i+1}-1)^2}{\nu(X_{i+1})} \frac{1}{(X_{i+1}-1)^3} - \frac{(X_i-1)^2}{\nu(X_i)} \frac{1}{(X_i-1)^3} \\ &\leq \frac{(X_i-1)^2}{\nu(X_i)} \left(\frac{1}{(X_{i+1}-1)^3} - \frac{1}{(X_i-1)^3} \right) \\ &= \frac{((X_i-1)^2 + (X_i-1)(X_{i+1}-1) + (X_{i+1}-1)^2)\Delta_{i+1}}{\nu(X_i)(X_i-1)(X_{i+1}-1)^3} \\ &\leq \frac{3(X_i-1)}{\nu(X_i)(X_{i+1}-1)^3} \Delta_{i+1} \\ &\leq \frac{24X_i}{\nu(X_i)X_{i+1}^3} \Delta_{i+1}. \end{aligned}$$

It follows

$$\begin{aligned} &\mathbf{E} \left[\max_{1 \leq j \leq \rho_n} \left| \sum_{i=0}^{j-1} \frac{\Delta_{i+1}}{(X_i-1)\nu(X_i)} - \sum_{i=0}^{j-1} \frac{\Delta_{i+1}}{(X_{i+1}-1)\nu(X_{i+1})} \right|; \Delta_{i+1} \leq X_i/2 \text{ for all } i < \rho_n \right] \\ &\leq \mathbf{E} \left[\sum_{i=0}^{\rho_n-1} \left| \frac{1}{(X_i-1)\nu(X_i)} - \frac{1}{(X_{i+1}-1)\nu(X_{i+1})} \right| \Delta_{i+1}; \Delta_{i+1} \leq X_i/2 \text{ for all } i < \rho_n \right] \\ &\leq \mathbf{E} \left[\sum_{i=0}^{\rho_n-1} \frac{8 \cdot 24}{X_i^2 \nu(X_i)} \Delta_{i+1}^2 \right] = o(1). \end{aligned}$$

Also, since $(x-1)\nu(x)$ is increasing

$$\frac{1}{(X_i-1)\nu(X_i)} \Delta_{i+1} \leq \int_{X_{i+1}}^{X_i} \frac{dx}{(x-1)\nu(x)} \leq \frac{1}{(X_{i+1}-1)\nu(X_{i+1})} \Delta_{i+1},$$

and we obtain

$$\max_{1 \leq j \leq \rho_n} \left| \sum_{i=0}^{j-1} \frac{1}{(X_i-1)\nu(X_i)} \Delta_{i+1} - \int_{X_j}^n \frac{dx}{(x-1)\nu(x)} \right| = o_P(1) \quad (22)$$

on the event that $\Delta_{i+1} \leq X_i/2$ for all $i < \rho_n$. Lemma 3 (ii) says that the complementary event has an asymptotically vanishing probability.

(v) Finally, by Lemma 1 (ii) with $r_n \leq y \leq n$ for $\chi < 1$

$$\begin{aligned} \int_y^n \frac{dx}{(x-1)\nu(x)} &= \int_y^n \frac{\lambda(x)/x^2}{\mu(x)/x} dx + \int_y^n \frac{dx}{x(x-1)\nu(x)} \\ &= \int_y^n \frac{\kappa'(x)}{\kappa(x)} dx + O\left(\int_{r_n}^n \frac{\kappa'(x) dx}{\kappa(x)x^\chi}\right) + O(r_n^{-1}), \end{aligned}$$

and recalling $r_n \rightarrow \infty$

$$\int_{r_n}^n \frac{\kappa'(x) dx}{\kappa(x)x^\chi} \sim \int_{r_n}^n \frac{dx}{x\nu(x)x^\chi} \leq \int_{r_n}^n \frac{dx}{x^{1+\chi}} = o(1).$$

Altogether we obtain

$$\max_{r_n \leq y \leq n} \left| \int_y^n \frac{dx}{(x-1)\nu(x)} - \log \frac{\kappa(n)}{\kappa(y)} \right| = o(1).$$

Combining this formula with (21) and (22) and recalling the definition of ρ_n , we arrive at

$$\max_{1 \leq j \leq \rho_n} \left| \sum_{i=0}^{j-1} \frac{1}{X_i} - \log \frac{\kappa(n)}{\kappa(X_j)} \right| = o_P(1).$$

This is our claim. □

4 Proof of Theorem 1

Under the assumptions of Proposition 2 we have

$$\ell_n^* := \int_0^{\tilde{\rho}_n} N_n(t) dt \stackrel{1}{\sim} \int_{r_n}^n \frac{x}{\mu(x)} dx. \quad (23)$$

In particular, this formula holds for $r_n := cn$ with $0 < c < 1$ as anticipated in the introduction's remark. Here the assumptions of Proposition 2 are satisfied because of Lemma 3 and

$$\int_{cn}^n \frac{dx}{\mu(x)} \leq (1-c) \frac{n}{\mu(cn)} = o(1),$$

which in turn is valid in view of Lemma 1 (iii).

In order to fill the gap up to ℓ_n , we construct a distinguished sequence of real numbers. We construct the numbers $2 \leq r_n \leq n$, $n \geq 1$, satisfying

$$\int_{r_n}^n \frac{dx}{\mu(x)} \rightarrow 0 \quad \text{and} \quad \int_2^{r_n} \frac{x}{\mu(x)} dx = o\left(\int_{r_n}^n \frac{x}{\mu(x)} dx\right) \quad (24)$$

as $n \rightarrow \infty$. From Lemma 3 we get that $\tilde{\rho}_n = o_P(1)$. Also, since by Lemma 1 (i) $x/\mu(x)$ is decreasing

$$\int_2^{r_n} \frac{x}{\mu(x)} dx \geq \frac{r_n(r_n-2)}{\mu(r_n)} \quad \text{and} \quad \int_{r_n}^n \frac{x}{\mu(x)} dx \leq \frac{r_n(n-r_n)}{\mu(r_n)}.$$

Therefore the second statement in (24) entails $r_n - 2 = o(n - r_n)$ as $n \rightarrow \infty$ and consequently $r_n = o(n)$. Hence the sequence r_n , $n \geq 1$, fulfils all requirements of Proposition 2.

For the construction of the numbers r_n note that from Lemma 1 (i) we have $x/\mu(x) \geq 2/(\mu(2)(x-1))$ for $x \geq 2$ and consequently

$$\int_2^\infty \frac{x}{\mu(x)} dx = \infty. \quad (25)$$

We distinguish two cases. If $\int_2^\infty \frac{dx}{\mu(x)} < \infty$, then the required sequence is easily obtained, because the first condition of (24) is fulfilled for any divergent sequence $r_n \leq n$ and the second one by reason of (25), if only r_n is diverging slowly enough.

Thus let us assume $\int_2^\infty \frac{dx}{\mu(x)} = \infty$ and let $r_{n,m} \geq 2$ for given $m \in \mathbb{N}$ be the solution of the equation

$$\int_{r_{n,m}}^n \frac{dx}{\mu(x)} = \frac{1}{m},$$

which exists for $n \geq 3$ and $m \geq 1/\int_2^3 \frac{dx}{\mu(x)}$. Since $\int_2^\infty \frac{dx}{\mu(x)} = \infty$, we have $r_{n,m} \rightarrow \infty$ as $n \rightarrow \infty$. It follows

$$\int_{r_{n,m}}^n \frac{x}{\mu(x)} dx \geq r_{n,m}/m \quad \text{and} \quad \int_2^{r_{n,m}} \frac{x}{\mu(x)} dx = o(r_{n,m})$$

as $n \rightarrow \infty$ because of $x = o(\mu(x))$ from Lemma 1 (iii). Therefore there are natural numbers $n_1 < n_2 < \dots$ such that

$$\int_2^{r_{n,m}} \frac{x}{\mu(x)} dx \leq \frac{1}{m} \int_{r_{n,m}}^n \frac{x}{\mu(x)} dx$$

for all $n \geq n_m$. Now letting $r_n := r_{n,m}$ for $n = n_m, \dots, n_{m+1} - 1$, we obtain

$$\int_{r_n}^n \frac{dx}{\mu(x)} \leq \frac{1}{m} \quad \text{and} \quad \int_2^{r_n} \frac{x}{\mu(x)} dx \leq \frac{1}{m} \int_{r_n}^n \frac{x}{\mu(x)} dx$$

for all $n \geq n_m$. This implies (24).

Applying now Proposition 2 with $f(x) = x$, we obtain from (24)

$$\int_0^{\tilde{\rho}_n} N_n(t) dt \stackrel{1}{\sim} \int_{r_n}^n \frac{x}{\mu(x)} dx \sim \int_2^n \frac{x}{\mu(x)} dx$$

and

$$\mathbf{E} \left[\int_{\tilde{\rho}_n}^{\tilde{\tau}_n} N_n(t) dt \right] = O \left(\int_2^{r_n} \frac{x}{\mu(x)} dx \right) = o \left(\int_2^n \frac{x}{\mu(x)} dx \right).$$

This implies our claim. \square

5 Proof of Theorem 2

Again, let r_n , $n \geq 1$, be any sequence fulfilling the assumptions of Proposition 1. We investigate the lengths

$$\widehat{\ell}_n^* := \int_0^{\tilde{\rho}_n} \widehat{N}_n(t) dt = \sum_{i=0}^{\rho_n-1} W_i Y_i, \quad (26)$$

which are the total lengths of the external branches up to the ρ_n -th merger. Here Y_i , $i \geq 0$, denotes the number of external branches extant after the first i merging events and W_i as above

the waiting time at the state X_i . In the proof we approximate $\widehat{\ell}_n^*$ by its conditional expectation given the block-counting process N_n , which in turn can be handled by means of Proposition 2 and 3. We shall employ the representation

$$Y_i = \sum_{k=1}^n I_{\{\zeta_k \geq i\}} \quad (27)$$

where ζ_k denotes the number of coalescent events before the k -th external branch (out of n) merges with some other branches within the coalescent.

Lemma 4. *We have for $i, j \geq 0, k, l = 1, \dots, n$*

$$\mathbf{P}(\zeta_k \geq i \mid N_n) = \frac{X_i - 1}{n - 1} \prod_{m=0}^{i-1} \left(1 - \frac{1}{X_m}\right) \quad \text{a.s.}$$

and for $k \neq l$

$$\mathbf{P}(\zeta_k \geq i, \zeta_l \geq j \mid N_n) \leq \mathbf{P}(\zeta_k \geq i \mid N_n) \mathbf{P}(\zeta_l \geq j \mid N_n) \quad \text{a.s.}$$

Proof. Let A be a subset of $\{1, \dots, n\}$ with $a \geq 1$ elements, and let ζ_A be the number of mergers before one of the branches ending in A gets involved into a merging event. Given Δ_1 the first merger consists of a uniformly random choice of $\Delta_1 + 1$ members out of $X_0 = n$ elements. Therefore we have

$$\mathbf{P}(\zeta_A \geq 1 \mid N_n) = \frac{\binom{X_0 - a}{\Delta_1 + 1}}{\binom{X_0}{\Delta_1 + 1}} = \frac{(X_0 - a) \cdots (X_1 - a)}{X_0 \cdots X_1} = \frac{(X_1 - 1) \cdots (X_1 - a)}{X_0 \cdots (X_0 - a + 1)} \quad \text{a.s.}$$

or

$$\mathbf{P}(\zeta_A \geq 1 \mid N_n) = \frac{(X_1 - 1) \cdots (X_1 - a)}{(X_0 - 1) \cdots (X_0 - a)} \left(1 - \frac{a}{X_0}\right) \quad \text{a.s.}$$

Because of the Markov property we may iterate this formula yielding

$$\mathbf{P}(\zeta_A \geq i \mid N_n) = \frac{(X_i - 1) \cdots (X_i - a)}{(X_0 - 1) \cdots (X_0 - a)} \prod_{m=0}^{i-1} \left(1 - \frac{a}{X_m}\right) \quad \text{a.s.} \quad (28)$$

In particular, with $A = \{k\}$ and $a = 1$ our first claim follows.

Similar for $k \neq l$ and $i \leq j$ with $\zeta'_{\{l\}} := \zeta_{\{l\}} - \zeta_{\{k,l\}}$ and $N'_{X_i}(t) := N_n(t + W_0 + \cdots + W_{i-1})$, $t \geq 0$, by means of the Markov property

$$\begin{aligned} \mathbf{P}(\zeta_k \geq i, \zeta_l \geq j \mid N_n) &= \mathbf{P}(\zeta_{\{k,l\}} \geq i \mid N_n) \mathbf{P}(\zeta'_{\{l\}} \geq j - i \mid N'_{X_i}) \\ &= \frac{(X_i - 1)(X_i - 2)}{(X_0 - 1)(X_0 - 2)} \prod_{m=0}^{i-1} \left(1 - \frac{2}{X_m}\right) \times \frac{X_j - 1}{X_i - 1} \prod_{m=i}^{j-1} \left(1 - \frac{1}{X_m}\right) \quad \text{a.s.} \end{aligned}$$

Since $X_i \leq X_0$ and $(1 - 2/X_m) \leq (1 - 1/X_m)^2$, this implies

$$\begin{aligned} \mathbf{P}(\zeta_k \geq i, \zeta_l \geq j \mid N_n) &\leq \frac{(X_i - 1)^2}{(X_0 - 1)^2} \prod_{m=0}^{i-1} \left(1 - \frac{1}{X_m}\right)^2 \times \frac{X_j - 1}{X_i - 1} \prod_{m=i}^{j-1} \left(1 - \frac{1}{X_m}\right) \\ &= \mathbf{P}(\zeta_k \geq i \mid N_n) \mathbf{P}(\zeta_l \geq j \mid N_n) \quad \text{a.s.}, \end{aligned}$$

which is our second claim. \square

Proof of Theorem 2. (i) First, we consider $\mathbf{E}[\widehat{\ell}_n^* | N_n]$. Due to (27) and Lemma 4 we have

$$\begin{aligned}\mathbf{E}[\widehat{\ell}_n^* | N_n] &= \sum_{i=0}^{\rho_n-1} \sum_{k=1}^n W_i \mathbf{P}(\zeta_k \geq i | N_n) \\ &= \frac{n}{n-1} \sum_{i=0}^{\rho_n-1} W_i (X_i - 1) \prod_{m=0}^{i-1} \left(1 - \frac{1}{X_m}\right) \quad \text{a.s.}\end{aligned}\quad (29)$$

Since $\sum_{m=0}^{i-1} X_m^{-2} \leq \sum_{a=X_{i-1}}^{\infty} a^{-2} \leq (X_{i-1} - 1)^{-1} \leq (r_n - 1)^{-1}$ for $i \leq \rho_n$ and in view of Proposition 3,

$$\prod_{m=0}^{i-1} \left(1 - \frac{1}{X_m}\right) = \exp\left(-\sum_{m=0}^{i-1} \frac{1}{X_m} + O(r_n^{-1})\right) = \frac{\kappa(X_i)}{\kappa(n)} \exp(o_P(1)),$$

where the $o_P(1)$ may be taken uniformly in $i < \rho_n$ in the sense of Proposition 3. Thus we obtain

$$\mathbf{E}[\widehat{\ell}_n^* | N_n] \stackrel{P}{\sim} \frac{1}{\kappa(n)} \sum_{i=0}^{\rho_n-1} W_i (X_i - 1) \kappa(X_i) = \frac{1}{\kappa(n)} \int_0^{\tilde{\rho}_n} f(N_n(t)) dt$$

with $f(x) := (x-1)\kappa(x)$. This function satisfies the assumption of Proposition 2. Because of $f(x) \sim \mu(x)$, $r_n \rightarrow \infty$ we obtain

$$\mathbf{E}[\widehat{\ell}_n^* | N_n] \stackrel{P}{\sim} \frac{1}{\kappa(n)} \int_{r_n}^n (x-1)\kappa(x) \frac{dx}{\mu(x)} \sim \frac{n-r_n}{\kappa(n)} = \frac{n(n-r_n)}{\mu(n)}. \quad (30)$$

(ii) Next, we have

$$\begin{aligned}\mathbf{E}[(\widehat{\ell}_n^* - \mathbf{E}[\widehat{\ell}_n^* | N_n])^2 | N_n] &= \mathbf{E}\left[\left(\sum_{i=0}^{\rho_n-1} \sum_{k=1}^n \left(W_i I_{\{\zeta_k \geq i\}} - W_i \mathbf{P}(\zeta_k \geq i | N_n)\right)\right)^2 | N_n\right] \\ &= \sum_{i,j=0}^{\rho_n-1} \sum_{k,l=1}^n W_i W_j (\mathbf{P}(\zeta_k \geq i, \zeta_l \geq j | N_n) - \mathbf{P}(\zeta_k \geq i | N_n) \mathbf{P}(\zeta_l \geq j | N_n)) \quad \text{a.s.}\end{aligned}$$

Applying Lemma 4, it follows

$$\begin{aligned}\mathbf{E}[(\widehat{\ell}_n^* - \mathbf{E}[\widehat{\ell}_n^* | N_n])^2 | N_n] &\leq \sum_{i,j=0}^{\rho_n-1} \sum_{k=1}^n W_i W_j \mathbf{P}(\zeta_k \geq i \vee j | N_n) \\ &\leq \sum_{i,j=0}^{\rho_n-1} W_i W_j \sum_{k=1}^n \mathbf{P}(\zeta_k \geq i | N_n) \\ &= \mathbf{E}[\widehat{\ell}_n^* | N_n] \sum_{j=0}^{\rho_n-1} W_j \\ &= \tilde{\rho}_n \mathbf{E}[\widehat{\ell}_n^* | N_n] \quad \text{a.s.}\end{aligned}$$

Since by assumption $\tilde{\rho}_n = o_P(1)$ and $r_n \leq \gamma n$, (30) yields

$$\mathbf{E}[(\widehat{\ell}_n^* - \mathbf{E}[\widehat{\ell}_n^* | N_n])^2 | N_n] = o_P\left(\frac{n(n-r_n)}{\mu(n)}\right) = o_P\left(\frac{n^2}{\mu(n)}\right).$$

Because of Lemma 1 (i) $n(n-1)/\mu(n)$ is increasing, which implies

$$\widehat{\ell}_n^* - \mathbf{E}[\widehat{\ell}_n^* | N_n] = o_P\left(\frac{n}{\sqrt{\mu(n)}}\right) = o_P\left(\frac{n^2}{\mu(n)}\right) \quad (31)$$

and because of (30)

$$\widehat{\ell}_n^* \stackrel{P}{\sim} \frac{n(n-r_n)}{\mu(n)}. \quad (32)$$

In particular, as discussed in the proof of Theorem 1 and addressed in the introduction, this approximation is valid for the sequence $r_n = cn$ with $0 < c < 1$.

(iii) Finally, let us switch to the numbers r_n , $n \geq 1$, constructed in the proof of Theorem 1 and fulfilling (24) as well as $r_n = o(n)$. As above

$$\begin{aligned} \mathbf{E}[\widehat{\ell}_n - \widehat{\ell}_n^* | N_n] &= \frac{n}{n-1} \sum_{i=\rho_n}^{\tau_n-1} W_i (X_i - 1) \prod_{m=0}^{i-1} \left(1 - \frac{1}{X_m}\right) \\ &\leq 2 \exp\left(-\sum_{m=0}^{\rho_n-1} \frac{1}{X_m}\right) \sum_{i=\rho_n}^{\tau_n-1} W_i X_i \quad \text{a.s.} \end{aligned}$$

From the Markov property and Theorem 1, applied to the coalescent with initial value X_{ρ_n} , we obtain

$$\sum_{i=\rho_n}^{\tau_n-1} W_i X_i \stackrel{P}{\sim} \int_2^{X_{\rho_n}} \frac{x}{\mu(x)} dx \leq \int_2^{r_n} \frac{x}{\mu(x)} dx.$$

By (24) and by monotonicity of $x/\mu(x)$ it follows

$$\sum_{i=\rho_n}^{\tau_n-1} W_i X_i = o_P\left(\int_{r_n}^n \frac{x}{\mu(x)} dx\right) = o_P\left(\frac{r_n}{\mu(r_n)} n\right).$$

Moreover, from Proposition 2

$$\exp\left(-\sum_{m=0}^{\rho_n-1} \frac{1}{X_m}\right) \stackrel{P}{\sim} \frac{\mu(r_n)}{r_n} \frac{n}{\mu(n)},$$

and we arrive at

$$\mathbf{E}[\widehat{\ell}_n - \widehat{\ell}_n^* | N_n] = o_P\left(\frac{n^2}{\mu(n)}\right).$$

Hence

$$\widehat{\ell}_n - \widehat{\ell}_n^* = o_P\left(\frac{n^2}{\mu(n)}\right).$$

This estimate in combination with (32) and $r_n = o(n)$ proves our theorem. \square

6 Proof of Theorem 3

(i) As to the second claim,

$$\int_2^n \left(\frac{x}{\mu(x)} - \frac{n}{\mu(n)} \right) dx = \int_2^n \int_x^n \frac{\kappa'(y)}{\kappa(y)^2} dy dx = \int_2^n (y-2) \frac{\kappa'(y)}{\kappa(y)^2} dy ,$$

and by Lemma 1 (ii) and Lemma 2

$$\int_2^n \left(\frac{x}{\mu(x)} - \frac{n}{\mu(n)} \right) dx \sim \int_2^n y \frac{\lambda(y)}{\mu(y)^2} dy \sim \int_2^n \frac{L(y)}{L^*(y)^2} dy \sim \frac{nL(n)}{L^*(n)^2} .$$

(ii) Turning to the first claim, we now strive for a lower bound for $\check{\ell}_n$. We resort to the definitions (23) and (26) and set

$$\check{\ell}_n^* := \ell_n^* - \widehat{\ell}_n^* .$$

Again, we first investigate its conditional expectation, given N_n . Note that $\mathbf{E}[\ell_n^* | N_n] = \ell_n^*$ a.s., therefore in view of (31)

$$\check{\ell}_n^* - \mathbf{E}[\check{\ell}_n^* | N_n] = \mathbf{E}[\widehat{\ell}_n^* | N_n] - \widehat{\ell}_n^* = o_P\left(\frac{n}{\sqrt{\mu(n)}}\right) .$$

Lemma 1 (i) implies $\mu(n) \geq n\mu(2)/2$ for $n \geq 2$, hence

$$\check{\ell}_n^* - \mathbf{E}[\check{\ell}_n^* | N_n] = o_P(n^{1/2}) . \quad (33)$$

We like to estimate $\mathbf{E}[\check{\ell}_n^* | N_n]$ from below. For this purpose we specify our choice of the numbers r_n . We fix $h \in \mathbb{N}$, we define stopping times $0 = \rho_{n,h} \leq \rho_{n,h-1} \leq \dots \leq \rho_{n,1}$ and the corresponding times $\tilde{\rho}_{n,g}$ as

$$\rho_{n,g} := \min\{i \geq 0 : X_i \leq \frac{g}{h}n\} , \quad \tilde{\rho}_{n,g} := \inf\{t \geq 0 : N_n(t) \leq \frac{g}{h}n\} , \quad g = 1, \dots, h , . \quad (34)$$

As we already argued in the proof of Theorem 1, we may apply Propositions 2 and 3 to these stopping times. We proceed now in the same manner as in (29). Using $X_i \leq n$ respectively $X_i \geq n(X_i - 1)/(n - 1)$, we obtain

$$\begin{aligned} \mathbf{E}\left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i(X_i - Y_i) | N_n\right] &= \sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i \left(X_i - \frac{n}{n-1}(X_i - 1) \prod_{m=0}^{i-1} \left(1 - \frac{1}{X_m}\right) \right) \\ &\geq \sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i X_i \left(1 - \prod_{m=0}^{i-1} \left(1 - \frac{1}{X_m}\right)\right) \\ &\geq \left(1 - \prod_{m=0}^{\rho_{n,g}-1} \left(1 - \frac{1}{X_m}\right)\right) \sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i X_i \\ &\geq \left(1 - \exp\left(-\sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m}\right)\right) \int_{\tilde{\rho}_{n,g}}^{\tilde{\rho}_{n,g-1}} N_n(t) dt \quad \text{a.s.} \end{aligned}$$

Proposition 2 together with Lemma 2 (ii) implies

$$\int_{\tilde{\rho}_{n,g}}^{\tilde{\rho}_{n,g-1}} N_n(t) dt \stackrel{1}{\sim} \int_{(g-1)n/h}^{gn/h} \frac{x}{\mu(x)} dx \sim \frac{n}{hL^*(n)} ,$$

and Proposition 3 yields

$$\sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m} \stackrel{1}{\sim} \log \frac{\kappa(n)}{\kappa(gn/h)} \sim \log \frac{L^*(n)}{L^*(gn/h)}.$$

Because L^* is slowly varying and because of (10), we have

$$\log \frac{L^*(n)}{L^*(gn/h)} \sim \frac{L^*(n) - L^*(gn/h)}{L^*(gn/h)} \sim \frac{L(n)}{L^*(n)} \log \frac{h}{g},$$

hence because of Lemma 2 (ii)

$$\sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m} \stackrel{1}{\sim} \frac{L(n)}{L^*(n)} \log \frac{h}{g} = o_P(1) \quad (35)$$

and

$$1 - \exp\left(-\sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m}\right) \stackrel{P}{\sim} \frac{L(n)}{L^*(n)} \log \frac{h}{g}.$$

Altogether

$$\mathbf{E}\left[\sum_{i=\rho_{n,g}}^{\rho_{n,g}-1} W_i(X_i - Y_i) \mid N_n\right] \geq (1 + o_P(1)) \frac{nL(n)}{L^*(n)^2} \frac{1}{h} \log \frac{h}{g}$$

and consequently

$$\begin{aligned} \mathbf{E}[\check{\ell}_n^* \mid N_n] &= \mathbf{E}\left[\sum_{i=0}^{\rho_{n,1}-1} W_i(X_i - Y_i) \mid N_n\right] \\ &\geq (1 + o_P(1)) \frac{nL(n)}{L^*(n)^2} \sum_{g=2}^h \frac{1}{h} \log \frac{h}{g} \\ &\geq (1 + o_P(1)) \frac{nL(n)}{L^*(n)^2} \int_{2/h}^1 \log \frac{1}{z} dz \\ &= (1 + o_P(1)) \frac{nL(n)}{L^*(n)^2} \left(1 - \frac{2}{h} - \frac{2}{h} \log \frac{2}{h}\right). \end{aligned}$$

In view of (33) this estimate transfers to $\check{\ell}_n^*$. We have $\check{\ell}_n \geq \check{\ell}_n^*$, thus letting $h \rightarrow \infty$, we obtain

$$\check{\ell}_n \geq (1 + o_P(1)) \frac{nL(n)}{L^*(n)^2} \quad (36)$$

as $n \rightarrow \infty$.

(iii) Coming to an upper bound, we have in view of $\prod_{m=0}^{i-1} (1 - 1/X_m) \geq 1 - \sum_{m=0}^{i-1} 1/X_m$

$$\begin{aligned} \mathbf{E}[\check{\ell}_n] &= \mathbf{E}[\mathbf{E}[\check{\ell}_n \mid N_n]] \\ &= \mathbf{E}\left[\sum_{i=0}^{\tau_n-1} W_i\left(X_i - \frac{n}{n-1}(X_i - 1) \prod_{m=0}^{i-1} \left(1 - \frac{1}{X_m}\right)\right)\right] \\ &\leq \mathbf{E}\left[\sum_{i=0}^{\tau_n-1} W_i\left(1 + \frac{n}{n-1} X_i \sum_{m=0}^{i-1} \frac{1}{X_m}\right)\right]. \end{aligned} \quad (37)$$

From Proposition 2 and since $x = o(\mu(x))$ (see Lemma 1 (iii)),

$$\mathbf{E}\left[\sum_{i=0}^{\tau_n-1} W_i\right] = \mathbf{E}[\tilde{\tau}_n] = O\left(\int_2^n \frac{dx}{\mu(x)}\right) = o\left(\int_2^n \frac{dx}{x}\right) = o(\log n). \quad (38)$$

Furthermore, by means of the Markov property

$$\begin{aligned} \mathbf{E}\left[\sum_{i=0}^{\tau_n-1} W_i X_i \sum_{m=0}^{i-1} \frac{1}{X_m}\right] &= \sum_{i=0}^{n-1} \mathbf{E}\left[\frac{X_i}{\lambda(X_i)} \sum_{m=0}^{i-1} \frac{1}{X_m}; X_i > 2\right] \\ &= \sum_{i=0}^{n-1} \mathbf{E}\left[\frac{X_i \Delta_{i+1}}{\mu(X_i)} \sum_{m=0}^{i-1} \frac{1}{X_m}; X_i > 2\right] \\ &= \mathbf{E}\left[\sum_{m=0}^{\tau_n-1} \frac{1}{X_m} \sum_{i=m+1}^{\tau_n-1} \frac{X_i \Delta_{i+1}}{\mu(X_i)}\right]. \end{aligned}$$

Since $x/\mu(x)$ is decreasing, we have $\sum_{i=m+1}^{\tau_n-1} X_i \Delta_{i+1}/\mu(X_i) \leq \int_1^{X_{m+1}} x/\mu(x) dx$, where we let $x/\mu(x) := 2/\mu(2)$ for $1 \leq x \leq 2$. Thus

$$\mathbf{E}\left[\sum_{i=0}^{\tau_n-1} W_i X_i \sum_{m=0}^{i-1} \frac{1}{X_m}\right] \leq \mathbf{E}\left[\sum_{m=0}^{\tau_n-1} \frac{1}{X_m} \int_1^{X_{m+1}} \frac{x}{\mu(x)} dx\right].$$

Invoking the Markov property once again, we obtain

$$\mathbf{E}\left[\sum_{i=0}^{\tau_n-1} W_i X_i \sum_{m=0}^{i-1} \frac{1}{X_m}\right] \leq \mathbf{E}\left[\sum_{m=0}^{\tau_n-1} \frac{\Delta_{m+1}}{(X_m - 1)\nu(X_m)} \int_1^{X_{m+1}} \frac{x}{\mu(x)} dx\right],$$

and taking now into account that the functions $(x-1)\nu(x)$ and $\int_1^x z/\mu(z) dz$ are increasing, we end up with

$$\mathbf{E}\left[\sum_{i=0}^{\tau_n-1} W_i X_i \sum_{m=0}^{i-1} \frac{1}{X_m}\right] \leq \int_1^n \frac{1}{(x-1)\nu(x)} \int_1^x \frac{z}{\mu(z)} dz dx, \quad (39)$$

where we let $(x-1)\nu(x) := \nu(2)$ for $1 \leq x \leq 2$. Using Lemma 2 (ii) it follows

$$\begin{aligned} \int_1^n \frac{1}{(x-1)\nu(x)} \int_1^x \frac{z}{\mu(z)} dz dx &\sim \int_2^n \frac{L(x)}{xL^*(x)} \int_2^x \frac{1}{L^*(z)} dz dx \\ &\sim \int_2^n \frac{L(x)}{xL^*(x)} \frac{x}{L^*(x)} dx \\ &\sim \frac{nL(n)}{L^*(n)^2}. \end{aligned}$$

Together with (37), (38) and (39) we obtain all in all

$$\mathbf{E}[\check{\ell}_n] \leq (1 + o(1)) \frac{nL(n)}{L^*(n)^2}.$$

Combining this result with (36), we get

$$\check{\ell}_n \stackrel{P}{\sim} \frac{nL(n)}{L^*(n)^2},$$

and invoking once again the convergence criterion of F. Riesz, also L_1 -convergence follows. This finishes the proof of Theorem 3. \square

7 Proof of Theorem 4

(i) Concerning the first claim, we have from Theorem 2 and Lemma 2 (ii)

$$\widehat{\ell}_{n,1} = \widehat{\ell}_n \stackrel{P}{\sim} \frac{n^2}{\mu(n)} \sim \frac{n}{L^*(n)},$$

where L^* is slowly varying. To obtain also L_1 -convergence, we use Theorem 1 saying that

$$\ell_n \stackrel{1}{\sim} \int_2^n \frac{x}{\mu(x)} dx \sim \int_2^n \frac{dx}{L^*(x)} \sim \frac{n}{L^*(n)}.$$

In particular the random variables $L^*(n)\ell_n/n$, $n \geq 1$, make a uniformly integrable sequence. Since $\widehat{\ell}_{n,1} \leq \ell_n$, this holds true also for the random variables $L^*(n)\widehat{\ell}_{n,1}/n$, $n \geq 1$. Therefore the above convergence in probability converts to L_1 -convergence.

(ii) Now we turn to the case $a \geq 2$. Preliminary we study lengths of the form

$$\widehat{\ell}_{n,a}^* = \int_{\widehat{\sigma}_n}^{\widehat{\rho}_n} \widehat{N}_{n,a}(t) dt = \sum_{i=\sigma_n}^{\rho_n-1} W_i Y_{i,a},$$

where $Y_{i,a}$ denotes the number of internal branches of order a present in the coalescent after i merging events. We are going to bound these numbers from below. Let A denote a subset of $\{1, \dots, n\}$ with a elements. For $1 \leq k \leq i$ let $E_{i,k,A}$ be the event that the external branches ending in A are not involved in the first $k-1$ mergers, next coalesce with the k -th merger to one lineage without any other branch participating, and then remain untouched by merging events till the i -th merger. These are disjoint events, which all contribute to $Y_{i,a}$, therefore

$$Y_{i,a} \geq Y'_{i,a} := \sum_{k=1}^i \sum_A I_{E_{i,k,A}},$$

where the second sum is taken over all $A \subset \{1, \dots, n\}$ with a elements.

Lemma 5. *Let both $A, A' \subset \{1, \dots, n\}$ have $a \geq 1$ elements. Then for $1 \leq k \leq i$*

$$\mathbf{P}(E_{i,k,A} | N_n) = \frac{a!}{(X_0 - 1) \cdots (X_0 - a)} \frac{X_i}{X_{k-1}} \prod_{m=0}^{k-2} \left(1 - \frac{a}{X_m}\right) \prod_{m=k}^i \left(1 - \frac{1}{X_m}\right) I_{\{\Delta_k = a-1\}} \text{ a.s.}$$

and for $A \neq A'$ or $k \neq l$ and for $1 \leq j \leq \ell$

$$\mathbf{P}(E_{i,k,A} \cap E_{j,l,A'} | N_n) \leq (1 + O(X_{k-1}^{-1}) + O(X_{l-1}^{-1})) \mathbf{P}(E_{i,k,A} | N_n) \mathbf{P}(E_{j,l,A'} | N_n) \text{ a.s.}$$

Proof. We proceed similar as in the proof of Lemma 4. From (28) and the Markov property

$$\begin{aligned} \mathbf{P}(E_{i,k,A} | N_n) &= \frac{(X_{k-1} - 1) \cdots (X_{k-1} - a)}{(X_0 - 1) \cdots (X_0 - a)} \prod_{m=0}^{k-2} \left(1 - \frac{a}{X_m}\right) \\ &\quad \times \binom{X_{k-1}}{a}^{-1} I_{\{\Delta_k = a-1\}} \times \frac{X_i - 1}{X_k - 1} \prod_{m=k}^{i-1} \left(1 - \frac{1}{X_m}\right) \text{ a.s.} \end{aligned}$$

Note that $X_k - 1 = X_{k-1} - a$ on the event $\{\Delta_k = a - 1\}$. Thus all factors containing X_{k-1} cancel up to the term X_{k-1} in the denominator. Replacing also $X_i - 1$ by $X_i(1 - 1/X_i)$, the first statement follows.

For the second claim note that in the case $A \neq A'$ and $A \cap A' \neq \emptyset$ or in the case $A = A'$ and $k \neq l$ the events $E_{i,k,A}$ and $E_{j,l,A'}$ are disjoint, then our claim is obvious. Therefore we may assume that $A \cap A' = \emptyset$. Let us consider the case $k < l < i \leq j$. Then from (28) and the Markov property

$$\begin{aligned} \mathbf{P}(E_{i,k,A} \cap E_{j,l,A'} \mid N_n) &= \frac{(X_{k-1} - 1) \cdots (X_{k-1} - 2a)}{(X_0 - 1) \cdots (X_0 - 2a)} \prod_{m=0}^{k-2} \left(1 - \frac{2a}{X_m}\right) \\ &\quad \times \binom{X_{k-1}}{a}^{-1} I_{\{\Delta_k = a-1\}} \times \frac{(X_{l-1} - 1) \cdots (X_{l-1} - a - 1)}{(X_k - 1) \cdots (X_k - a - 1)} \prod_{m=k}^{l-2} \left(1 - \frac{a+1}{X_m}\right) \\ &\quad \times \binom{X_{l-1}}{a}^{-1} I_{\{\Delta_l = a-1\}} \times \frac{(X_i - 1)(X_i - 2)}{(X_l - 1)(X_l - 2)} \prod_{m=l}^{i-1} \left(1 - \frac{2}{X_m}\right) \\ &\quad \times \frac{X_j - 1}{X_i - 1} \prod_{m=i}^{j-1} \left(1 - \frac{1}{X_m}\right) \text{ a.s.} \end{aligned}$$

Here the product $(X_k - 1) \cdots (X_k - a - 1) = (X_{k-1} - a) \cdots (X_{k-1} - 2a)$ cancels out on the event $\{\Delta_k = a - 1\}$, and again only the factor X_{k-1} remains. Similarly $X_l - 2$ and $X_{l-1} - a - 1$ cancel. By increasing some other terms, we get

$$\begin{aligned} \mathbf{P}(E_{i,k,A} \cap E_{j,l,A'} \mid N_n) &\leq \frac{a!a!}{(X_0 - 1) \cdots (X_0 - 2a)} \prod_{m=0}^{k-2} \left(1 - \frac{a}{X_m}\right)^2 \\ &\quad \times I_{\{\Delta_k = a-1\}} \frac{1}{X_{k-1}} \prod_{m=k}^{l-2} \left(1 - \frac{a}{X_m}\right) \left(1 - \frac{1}{X_m}\right) \\ &\quad \times I_{\{\Delta_l = a-1\}} \frac{1}{X_{l-1}} \prod_{m=l}^{i-1} \left(1 - \frac{1}{X_m}\right) \left(1 - \frac{1}{X_m}\right) \\ &\quad \times (X_i - 1)(X_j - 1) \prod_{m=i}^{j-1} \left(1 - \frac{1}{X_m}\right) \text{ a.s.} \end{aligned}$$

Replacing also $X_i - 1$ and $X_j - 1$ as above and comparing with our first formula, we obtain

$$\begin{aligned} &\mathbf{P}(E_{i,k,A} \cap E_{j,l,A'} \mid N_n) \\ &\leq \frac{(X_0 - 1) \cdots (X_0 - a)}{(X_0 - a - 1) \cdots (X_0 - 2a)} \left(1 - \frac{a}{X_{k-1}}\right)^{-1} \mathbf{P}(E_{i,k,A} \mid N_n) \mathbf{P}(E_{j,l,A'} \mid N_n) \text{ a.s.} \end{aligned}$$

This implies our second claim. Other cases like $k < i < l < j$ are treated similarly. \square

(iii) Coming back to the proof of the theorem's second claim, we first consider, similarly as above, conditional expectations given N_n . Recall from (34) the definition of $\rho_{n,g}$. We have from Lemma 5 for $g \geq 2$

$$\begin{aligned} \mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] &= \sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i \sum_{k=1}^i \sum_A \mathbf{P}(E_{i,k,A} \mid N_n) \\ &= \frac{X_0}{X_0 - a} \sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i \sum_{k=1}^i \frac{X_i}{X_{k-1}} \prod_{m=0}^{k-2} \left(1 - \frac{a}{X_m}\right) \prod_{m=k}^i \left(1 - \frac{1}{X_m}\right) I_{\{\Delta_k=a-1\}} \text{ a.s.} \end{aligned}$$

The products may be estimated from above by 1 and, by means of Proposition 3 and the bound $(1-z) \geq \exp(-cz)$ for $z \leq 1/2$ and a suitable $c > 0$, from below uniformly in i, k by

$$\prod_{m=0}^{\rho_{n,1}-1} \left(1 - \frac{a}{X_m}\right) \geq \exp \left(-ca \sum_{m=0}^{\rho_{n,1}-1} \frac{1}{X_m} \right) \stackrel{P}{\sim} \left(\frac{\kappa(n/h)}{\kappa(n)} \right)^{ca} \sim \left(\frac{L^*(n/h)}{L^*(n)} \right)^{ca} \sim 1.$$

Consequently we may replace the products by 1 and obtain

$$\mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] \stackrel{P}{\sim} \sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i X_i \sum_{k=1}^i \frac{1}{X_{k-1}} I_{\{\Delta_k=a-1\}} \quad (40)$$

as $n \rightarrow \infty$.

From Lemma 2 (i) and (35),

$$\sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m} \mathbf{P}(\Delta_{m+1} = a-1) \stackrel{P}{\sim} \frac{1}{(a-1)a} \sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m} \stackrel{P}{\sim} \frac{1}{(a-1)a} \frac{L(n)}{L^*(n)} \log \frac{h}{g}.$$

By the Markov property

$$\begin{aligned} &\mathbf{E} \left[\left(\sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m} (I_{\{\Delta_{m+1}=a-1\}} - \mathbf{P}(\Delta_{m+1} = a-1)) \right)^2 \right] \\ &= \mathbf{E} \left[\sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m^2} (I_{\{\Delta_{m+1}=a-1\}} - \mathbf{P}(\Delta_{m+1} = a-1))^2 \right] \\ &\leq \mathbf{E} \left[\frac{1}{X_{\rho_{n,g-1}-1}} \right] \leq \frac{1}{(gn/h) - 1}. \end{aligned}$$

Therefore

$$\sum_{m=0}^{\rho_{n,g}-1} \frac{1}{X_m} I_{\{\Delta_{m+1}=a-1\}} \stackrel{P}{\sim} \frac{1}{a(a-1)} \frac{L(n)}{L^*(n)} \log \frac{h}{g},$$

also once more by Proposition 2 and Lemma 2 (ii)

$$\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i X_i = \int_{\tilde{\rho}_{n,g}}^{\tilde{\rho}_{n,g-1}} N_n(t) dt \stackrel{1}{\sim} \int_{(g-1)n/h}^{gn/h} \frac{x}{\mu(x)} dx \sim \frac{n}{hL^*(n)}.$$

These formulas yield together with (40) the following lower and upper bounds for $g \geq 2$

$$\mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] \geq (1 + o_P(1)) \frac{1}{a(a-1)} \frac{nL(n)}{L^*(n)^2} \frac{1}{h} \log \frac{h}{g} \quad (41)$$

and

$$\mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] \leq (1 + o_P(1)) \frac{1}{a(a-1)} \frac{nL(n)}{L^*(n)^2} \frac{1}{h} \log \frac{h}{g-1} . \quad (42)$$

(iv) Now we estimate the difference between $\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a}$ and its conditional expectation given N_n . We have

$$\begin{aligned} & \mathbf{E} \left[\left(\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} - \mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] \right)^2 \mid N_n \right] \\ &= \mathbf{E} \left[\left(\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i \sum_{k=1}^i \sum_A (I_{E_{i,k,A}} - \mathbf{P}(E_{i,k,A} \mid N_n)) \right)^2 \mid N_n \right] \\ &= \sum_{i,j=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i W_j \sum_{k=1}^i \sum_{l=1}^j \sum_{A,A'} (\mathbf{P}(E_{i,k,A} \cap E_{j,l,A'} \mid N_n) - \mathbf{P}(E_{i,k,A} \mid N_n) \mathbf{P}(E_{j,l,A'} \mid N_n)) \text{ a.s.} \end{aligned}$$

For $A = A'$, $k = l$ and $i \leq j$ we use the estimate

$$\mathbf{P}(E_{i,k,A} \cap E_{j,k,A} \mid N_n) \leq \mathbf{P}(E_{i,k,A} \mid N_n) \text{ a.s.}$$

Taking also account of Lemma 5, we obtain with some $c > 0$

$$\begin{aligned} & \mathbf{E} \left[\left(\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} - \mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] \right)^2 \mid N_n \right] \\ & \leq 2 \sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} \sum_{j=i}^{\rho_{n,g-1}-1} \sum_{k=1}^i \sum_A W_i W_j \mathbf{P}(E_{i,k,A} \mid N_n) \\ & \quad + c \sum_{i,j=\rho_{n,g}}^{\rho_{n,g-1}-1} \sum_{k=1}^i \sum_{l=1}^j \sum_{A,A'} W_i W_j \left(\frac{1}{X_k} + \frac{1}{X_l} \right) \mathbf{P}(E_{i,k,A} \mid N_n) \mathbf{P}(E_{j,l,A'} \mid N_n) \\ & \leq 2 \sum_{j=0}^{\rho_{n,g-1}-1} W_j \sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} \sum_{k=1}^i \sum_A W_i \mathbf{P}(E_{i,k,A} \mid N_n) \\ & \quad + \frac{2c}{X_{\rho_{n,g-1}-1}} \left(\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} \sum_{k=1}^i \sum_A W_i \mathbf{P}(E_{i,k,A} \mid N_n) \right)^2 \\ & = 2\tilde{\rho}_{n,g-1} \mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] + \frac{2ch}{n} \left(\mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] \right)^2 \text{ a.s.} \end{aligned}$$

Using (42), $L_n = o(L^*(n))$ and the fact that $L^*(n)$ is increasing, this implies for $g \geq 2$

$$\mathbf{E} \left[\left(\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} - \mathbf{E} \left[\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \mid N_n \right] \right)^2 \mid N_n \right] = o(n)$$

and thus from (41) for $g \geq 2$

$$\sum_{i=\rho_{n,g}}^{\rho_{n,g-1}-1} W_i Y'_{i,a} \geq (1 + o_P(1)) \frac{1}{a(a-1)} \frac{nL(n)}{L^*(n)^2} \frac{1}{h} \log \frac{h}{g}.$$

(v) The last formula implies for $a \geq 2$ that

$$\begin{aligned} \widehat{\ell}_{n,a} &= \sum_{i=0}^{\tau_n-1} W_i Y_{i,a} \geq \sum_{i=\rho_{n,h-1}}^{\rho_{n,1}-1} W_i Y'_{i,a} \\ &\geq (1 + o_P(1)) \frac{1}{a(a-1)} \frac{nL(n)}{L^*(n)^2} \sum_{g=2}^{h-1} \frac{1}{h} \log \frac{h}{g} \\ &\geq (1 + o_P(1)) \frac{1}{a(a-1)} \frac{nL(n)}{L^*(n)^2} \int_{2/h}^1 \log \frac{1}{z} dz. \end{aligned}$$

Letting $h \rightarrow \infty$, we obtain the lower estimate

$$\widehat{\ell}_{n,a} \geq (1 + o_P(1)) \frac{1}{a(a-1)} \frac{nL(n)}{L^*(n)^2}.$$

For an upper estimate note that $\check{\ell}_n = \sum_{a \geq 2} \widehat{\ell}_{n,a}$. This formula and Theorem 3 imply for any natural number r

$$\begin{aligned} \widehat{\ell}_{n,a} &\leq \check{\ell}_n - \sum_{2 \leq b \leq r, b \neq a} \widehat{\ell}_{b,n} \\ &\leq (1 + o_P(1)) \frac{nL(n)}{L^*(n)^2} \left(1 - \sum_{2 \leq b \leq r, b \neq a} \frac{1}{(b-1)b} \right) \\ &\stackrel{P}{\sim} \frac{nL(n)}{L^*(n)^2} \left(\frac{1}{r} + \frac{1}{(a-1)a} \right). \end{aligned}$$

Letting $r \rightarrow \infty$ yields the upper estimates and we obtain altogether

$$\widehat{\ell}_{n,a} \stackrel{P}{\sim} \frac{1}{a(a-1)} \frac{nL(n)}{L^*(n)^2}.$$

In order to achieve also L_1 -convergence we deduce from Theorem 3 that the random variables $\check{\ell}_n L^*(n)^2 / (nL(n))$ form a uniformly integrable sequence. Since $\widehat{\ell}_{n,a} \leq \check{\ell}_n$, the same holds true for the random variables $\widehat{\ell}_{n,a} L^*(n)^2 / (nL(n))$. Thus also L_1 -convergence follows. This finishes the proof of Theorem 4. \square

References

- [1] BASDEVANT, A.-L. and GOLDSCHMIDT, C. (2008). Asymptotics of the Allele Frequency Spectrum Associated with the Bolthausen-Sznitman Coalescent. *Electron. J. Probab.* **13**, 486–512.
- [2] BERESTYCKI, N. (2009). Recent progress in coalescent theory. *Ensaïos Matemáticos* **16**, 1–193.
- [3] BERESTYCKI, J., BERESTYCKI, N. and LIMIC, V. (2014). Asymptotic sampling formulae for Λ -coalescents. *Ann. Henri Poincaré* **50**, 715–731.
- [4] BERESTYCKI, J., BERESTYCKI, N. and SCHWEINSBERG, J. (2007). Beta-coalescents and continuous stable random trees. *Ann. Probab.* **35**, 1835–1887.
- [5] BIRKNER, M., BLATH, J. and ELDON, B. (2013). Statistical properties of the site-frequency spectrum associated with Λ -coalescents. *Genetics* **195**, 1037–1053.
- [6] BOLTHAUSEN, E. and SZNITMAN, A.-S. (1998). On Ruelle’s probability cascades and an abstract cavity method. *Commun. Math. Phys.* **197**, 247–276.
- [7] DAHMER, I., KERSTING, G. and WAKOLBINGER, A. (2014). The total external branch length of Beta-coalescents. *Combinatorics Prob. Comput.* **23**, 1010–1027.
- [8] DELMAS, J.-F., DHERSIN, J.-S. and SIRI-JÉGOUSSE, A. (2008). Asymptotic results on the length of coalescent trees. *Ann. Appl. Probab.* **18**, 997–1025.
- [9] DHERSIN, J.-S. and MÖHLE, M. (2013). On the external branches of coalescents with multiple collisions. *Electron. J. Prob.* **18**, 1–11.
- [10] DRMOTA, M., IKSANOV, A., MÖHLE, M. and RÖSLER, U. (2007). Asymptotic results concerning the total branch length of the Bolthausen-Sznitman coalescent. *Stoch. Process. Appl.* **117**, 1404–1421.
- [11] FELLER, W. (1971). *An Introduction to Probability Theory and Its Applications*, Vol. 2. John Wiley & Sons, New York.
- [12] GOLDSCHMIDT, C. and MARTIN, J. (2005). Random Recursive Trees and the Bolthausen-Sznitman Coalescent. *Electron. J. Probab.* **10**, 718–745.
- [13] JANSON, S. and KERSTING, G. (2011). On the total external length of the Kingman coalescent. *Electron. J. Prob.* **16**, 2203–2218.
- [14] KERSTING, G. (2012). The asymptotic distribution of the length of Beta-coalescent trees. *Ann. Appl. Probab.* **22**, 2086–2107.
- [15] KERSTING, G., PARDO, J. C. and SIRI-JÉGOUSSE, A. (2014). Total internal and external lengths of the Bolthausen-Sznitman coalescent. *J. Appl. Probab.* **51A**, 73–86.
- [16] KERSTING, G., SCHWEINSBERG, J. and WAKOLBINGER, A. (2014). The evolving beta coalescent. *Electron. J. Prob.* **19**, 1–27.

- [17] KIMURA, M. (1969). The Number of Heterozygous Nucleotide Sites Maintained in a Finite Population Due to Steady Flux of Mutations. *Genetics* **61**, 893–903.
- [18] KINGMAN, J. F. C. (1982). On the genealogies of large populations. *J. Appl. Probab.* **19A**, 27–43.
- [19] MÖHLE, M. (2010). Asymptotic results for coalescent processes without proper frequencies and applications to the two-parameter Poisson-Dirichlet coalescent. *Stoch. Process. Appl.* **120**, 2159–2173.
- [20] PITMAN, J. (1999). Coalescents with multiple collisions. *Ann. Probab.* **27**, 1870–1902.
- [21] SAGITOV, S. (1999). The general coalescent with asynchronous mergers of ancestral lines. *J. Appl. Probab.* **36**, 1116–1125.
- [22] SCHWEINSBERG, J. (2000). A necessary and sufficient condition for the Λ -coalescent to come down from infinity. *Electron. Comm. Probab.* **5**, 1–11.
- [23] SPENCE, J. P., KAMM, J. A. and SONG, Y. S. (2016). The Site Frequency Spectrum for General Coalescents. *Genetics* **202**, 1549–1561.
- [24] TAVARÉ, S. (2004). Ancestral inference in population genetics, in: Lectures on Probability and Statistics 1–188, in: *Lecture Notes in Mathematics*, Vol. 1837, Springer, Berlin.
- [25] WAKELEY, J. (2009). *Coalescent Theory: An Introduction*. Roberts and Company Publishers, Greenwood Village, USA.
- [26] WATTERSON, G. A. (1975). On the number of segregating sites in genetical models without recombination. *Theor. Pop. Biol.* **7**, 256–276.