

Aldous' Spectral Gap Conjecture for Normal Sets

Ori Parzanchevski and Doron Puder

February 26, 2022

Abstract

Let S_n denote the symmetric group on n elements, and $\Sigma \subseteq S_n$ a symmetric subset of permutations. Aldous' spectral gap conjecture, proved by Caputo, Liggett and Richthammer [CLR10], states that if Σ is a set of transpositions, then the second eigenvalue of the Cayley graph $\text{Cay}(S_n, \Sigma)$ is identical to the second eigenvalue of the Schreier graph on n vertices depicting the action of S_n on $\{1, \dots, n\}$. Inspired by this seminal result, we study similar questions for other types of sets in S_n . Specifically, we consider normal sets: sets that are invariant under conjugation. Relying on character bounds due to Larsen and Shalev [LS08], we show that for large enough n , if $\Sigma \subset S_n$ is a full conjugacy class, then the second eigenvalue of $\text{Cay}(S_n, \Sigma)$ is roughly identical to the second eigenvalue of the Schreier graph depicting the action of S_n on ordered 4-tuples of elements from $\{1, \dots, n\}$. We further show that this type of result does not hold when Σ is an arbitrary normal set, but a slightly weaker one does hold. We state a conjecture in the same spirit regarding an arbitrary symmetric set $\Sigma \subset S_n$, which yields surprisingly strong consequences.

Contents

1	Introduction	1
2	A Single Conjugacy Class	5
3	Arbitrary Normal Sets	11
4	A Conjecture for Arbitrary Symmetric Sets and Its Consequences	13

1 Introduction

Consider a finite group G and a symmetric subset $\Sigma \subseteq G$, namely, $g \in \Sigma \Rightarrow g^{-1} \in \Sigma$. The $|G| \times |G|$ adjacency matrix A of the Cayley graph $\text{Cay}(G, \Sigma)$ is symmetric and equals

$$\sum_{g \in \Sigma} \rho_{\text{reg}}(g),$$

where reg is the right regular representation of G , namely, $\rho_{\text{reg}}(g)$ is the permutation matrix depicting multiplication from the right by g . Recall that the regular representation of G decomposes as a direct sum of all complex irreducible representations of G (irreps for short), each appearing with multiplicity identical to its dimension¹. An appropriate change of basis thus turns A into a block-diagonal matrix, with $\dim(\rho)$ blocks of size $\dim(\rho) \times \dim(\rho)$ for every irrep ρ of G . The value of each of these $\dim(\rho)$ blocks is $\sum_{g \in \Sigma} \rho(g)$. This shows that the multiset of eigenvalues of A can be partitioned into sub-multisets, each of which is associated

¹Throughout this paper we use some standard facts from the theory of group representations and, more specifically, from the theory of representations of the symmetric groups S_n . Good references are [FH91] for the general theory, [Ful97] for representations of S_n , and [Dia88] for applications to probability.

with some $\rho \in \widehat{G}$, where \widehat{G} is the set of (isomorphism types of) irreps of G . For example, the largest eigenvalue of A is $|\Sigma|$: this is the eigenvalue corresponding to the constant eigenfunction, and it is associated with the trivial representation of G .

The current work focuses on the symmetric group S_n . We consider the eigenvalues associated with the trivial and the sign representations to be *trivial*, and denote by $\lambda(S_n, \Sigma)$ the largest non-trivial eigenvalue of a symmetric set $\Sigma \subseteq S_n$. If we think of Σ as the element $\sum_{\sigma \in \Sigma} \sigma$ of $\mathbb{R}[S_n]$, we can thus write

$$\lambda(S_n, \Sigma) = \max_{\rho \in \widehat{S_n} \setminus \{\text{triv}, \text{sgn}\}} \lambda_1(\rho(\Sigma));$$

here $\lambda_1(\rho(\Sigma))$ marks the largest eigenvalue of the matrix $\rho(\Sigma) = \sum_{\sigma \in \Sigma} \rho(\sigma)$, which has only real eigenvalues as Σ is symmetric. Alternatively, $\lambda(S_n, \Sigma)$ is simply the largest eigenvalue of A whose eigenvector is orthogonal both to the constant functions, and to the indicator function of $A_n \leq S_n$.

Many properties of a regular graph are related to the value of its second largest eigenvalue. Primarily, the spectral gap $\lambda_1 - \lambda_2$ is a good measure for the extent to which the graph is “expanding” (see, e.g., the surveys [HLW06, Lub12]). Around 1992, David Aldous conjectured the following: whenever $\Sigma \subset S_n$ is a set of transpositions, the largest non-trivial eigenvalue $\lambda(S_n, \Sigma)$ is equal to the largest eigenvalue associated with the *standard* irrep,

$$\text{std} = \{(x_1, \dots, x_n) \in \mathbb{C}^n \mid x_1 + \dots + x_n = 0\},$$

which corresponds to the Young diagram $(n-1, 1)$.² This conjecture was proved in 2009 by Caputo, Liggett and Richthammer. In fact, they proved a stronger version applying to weighted Cayley graphs as well:

Theorem 1.1 (Aldous’ spectral gap conjecture, [CLR10]). *Let $\Sigma \in \mathbb{R}[S_n]$ be supported on transpositions, with non-negative coefficients. Then the second eigenvalue of the weighted Cayley graph $\text{Cay}(S_n, \Sigma)$ is equal to the largest eigenvalue of the standard representation. Namely,*

$$\lambda_2(\text{Cay}(S_n, \Sigma)) = \lambda_1(\text{std}(\Sigma)).$$

This theorem can be equivalently stated in terms of Schreier graphs. Given $\Sigma \subset S_n$, denote $[n] \stackrel{\text{def}}{=} \{1, \dots, n\}$ and by $\text{Sch}(S_n \curvearrowright [n], \Sigma)$ the Schreier graph depicting the action of S_n on $[n]$ with respect to the subset Σ . This is the $|\Sigma|$ -regular graph with n vertices labeled $1, \dots, n$, and with an edge (i, j) for every $\sigma \in \Sigma$ with $\sigma(i) = j$ (there may be loops and multiple edges). The n eigenvalues of (the adjacency matrix of) $\text{Sch}(S_n \curvearrowright [n], \Sigma)$ are a sub-multiset of the $n!$ eigenvalues of $\text{Cay}(S_n, \Sigma)$: this can be seen directly by lifting every eigenfunction on the Schreier graph to an eigenfunction with the same eigenvalue in the Cayley graph. Both the Schreier graph and the Cayley graph have trivial eigenvalue $\lambda_1 = |\Sigma|$. It follows that if Σ is symmetric then

$$\lambda_2(\text{Sch}(S_n \curvearrowright [n], \Sigma)) \leq \lambda_2(\text{Cay}(S_n, \Sigma)). \quad (1.1)$$

Aldous’ conjecture states that whenever Σ is a set of transpositions, there is equality in (1.1). This is equivalent to (the non-weighted version of) Theorem 1.1, since the representation $\mathbb{C}^{[n]}$ decomposes into the constant functions in $\mathbb{C}^{[n]}$, which form the trivial representation, and *std*.

Aldous’ conjecture cannot be naïvely extended to arbitrary symmetric sets $\Sigma \subset S_n$, as illustrated by the following three examples:

Example 1.2. 1. If Σ generates a proper transitive subgroup $H \leq S_n$, $H \neq A_n$, such as $\langle (12 \dots n) \rangle$, then $\text{Cay}(S_n, \Sigma)$ has at least three connected components and so $\lambda(S_n, \Sigma) = |\Sigma|$, whereas the Schreier graph $\text{Sch}(S_n \curvearrowright [n], \Sigma)$ is connected and so $\lambda_1(\text{std}(\Sigma)) = \lambda_2(\text{Sch}(S_n \curvearrowright [n], \Sigma)) \leq |\Sigma|$.

²Occasionally, several different irreps give rise to an eigenvalue which is equal to $\lambda(S_n, \Sigma)$. Aldous’ conjecture says that when Σ is a set of transpositions, the standard irrep is always one of these irreps.

2. For $\Sigma = \{\text{id}, (1\ 2), (1\ 2\ \dots\ n)^{\pm 1}\}$, it follows from [DSC93] that $\lambda_1(\text{std}(\Sigma)) < \max(\lambda_1(\rho_{(n-2,2)}(\Sigma)), \lambda_1(\rho_{(n-2,1,1)}(\Sigma)))$, where ρ_λ is the irrep with Young diagram λ ; we elaborate in Example 4.2.
3. Full conjugacy classes also occasionally have $\lambda(S_n, \Sigma)$ not associated with std. For example, denoting by σ^{S_n} the conjugacy class of σ , if n is odd and $\sigma = (1\ 2)(3\ 4)\dots(n-2\ n-1)$, then $\lambda_1(\text{std}(\sigma^{S_n})) = 0$ while $\lambda(S_n, \sigma^{S_n}) = |\sigma^{S_n}|/n$. We explain this computation below.

But what if instead of considering the Schreier graph depicting the action of S_n on $[n]$, we consider the Schreier graph depicting the action on 2-tuples, 4-tuples or k -tuples of distinct elements from $[n]$, for some fixed k ? We denote this Schreier graph, which is a $|\Sigma|$ -regular graph on $n(n-1)\dots(n-k+1)$ vertices, by $\text{Sch}(S_n \curvearrowright [n]_k, \Sigma)$.

A more sophisticated variation of Aldous' conjecture is then suggested by the classification of multiply-transitive finite groups. This classification, which follows from the classification of finite simple groups, gives a full description of (faithful) k -transitive actions of finite groups, for every $k \geq 2$. In particular, a finite 4-transitive group is either S_n ($n \geq 4$), A_n ($n \geq 6$), or one of the four Mathieu groups M_{11} , M_{12} , M_{23} and M_{24} (where M_n is a subgroup of S_n and is 4-transitive in its action on $[n]$, for $n = 11, 12, 23, 24$) [Cam99, Theorem 4.11]. It follows that for $n \geq 25$, if $\Sigma \subset S_n$ does not generate A_n or S_n , then $\langle \Sigma \rangle$ is not 4-transitive, and so the Schreier graph associated with the action of S_n on 4-tuples of distinct elements in $[n]$ is not connected. This shows that the mere existence of a (positive) spectral gap is captured by $\text{Sch}(S_n \curvearrowright [n]_4, \Sigma)$, and this resolves the issue illustrated in Example 1.2(1). Is it possible that $\text{Sch}(S_n \curvearrowright [n]_4, \Sigma)$ can capture not only the existence of a spectral gap but also its exact value?

As elaborated in Example 4.2 below, replacing $\text{Sch}(S_n \curvearrowright [n], \Sigma)$ with $\text{Sch}(S_n \curvearrowright [n]_2, \Sigma)$ resolves the issue with Example 1.2(2) as well. As for a full conjugacy class, it turns out that occasionally one needs to consider also the “signed” action of S_n on k -tuples. We define it more generally, for weighted subsets, keeping the generality of Theorem 1.1:

Definition 1.3. For an action of S_n on a set X and an element $\Sigma = \sum_{\sigma \in S_n} \alpha_\sigma \sigma \in \mathbb{R}[S_n]$, let $\rho_X(\sigma)$ be the permutation matrix depicting the action of σ on X , and denote

$$\rho_X(\Sigma) = \sum_{\sigma \in S_n} \alpha_\sigma \rho_X(\sigma), \quad \rho_X^{\text{sgn}}(\Sigma) = (\rho_X \otimes \text{sgn})(\Sigma) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) \alpha_\sigma \rho_X(\sigma).$$

We denote by $\lambda(k, \Sigma)$ (resp. $\lambda^{\text{sgn}}(k, \Sigma)$) the largest eigenvalue of $\rho_{[n]_k}(\Sigma)$ (resp. $\rho_{[n]_k}^{\text{sgn}}(\Sigma)$) corresponding to an eigenvector orthogonal to the constant functions. $\lambda(k, \Sigma)$
 $\lambda^{\text{sgn}}(k, \Sigma)$

We call Σ *symmetric* if $\alpha_\sigma = \alpha_{\sigma^{-1}}$ for every $\sigma \in S_n$ and *non-negative* if $\alpha_\sigma \geq 0$ for every $\sigma \in S_n$. It follows from the decomposition of the regular representation that for every symmetric non-negative Σ and $n > k$

$$\max(\lambda(k, \Sigma), \lambda^{\text{sgn}}(k, \Sigma)) \leq \lambda(S_n, \Sigma),$$

similarly to (1.1). The discussion above leads to the following potential generalization of Theorem 1.1, which was raised during discussions between Gady Kozma and the second author.

Question 1.4. *Is there a fixed $k \geq 1$ such that for every large enough n and every symmetric non-negative $\Sigma \in \mathbb{R}[S_n]$, we have*

$$\max(\lambda(k, \Sigma), \lambda^{\text{sgn}}(k, \Sigma)) = \lambda(S_n, \Sigma)?$$

This question has an equivalent representation-theoretic formulation. When k is fixed, the decomposition of the (signed) action of S_n on k -tuples into irreps, contains a fixed number of natural families of irreps. For example, for $n \geq 8$, the action on 4-tuples decomposes precisely to the 12 irreps associated with Young diagrams with at most four blocks outside the first row. These families are formally defined as follows:

Definition 1.5. A series of irreps $\{\rho_n \in \hat{S}_n\}_{n \geq n_0}$ is called a *family of irreps* if one of the following conditions holds:

1. Either the structure of the associated Young diagram outside the first row is constant, namely, for every $n \geq n_0$, ρ_{n+1} is obtained from ρ_n by adding a block to the first row, or
2. The structure of the associated Young diagram outside the first column is constant, namely, for every $n \geq n_0$, ρ_{n+1} is obtained from ρ_n by adding a block to the first column.

The representation $\mathbb{C}^{[n]_k}$ of S_n decomposes into all irreps with at most k blocks outside the first row (with some multiplicities). Similarly, the *signed* action on k -tuples, $\text{sgn} \otimes \mathbb{C}^{[n]_k}$, decomposes into all irreps with at most k blocks outside the first column. Thus, the following question is equivalent to Question 1.4.

Question 1.6. *Is there a finite set of families of irreps $\rho^{(1)}, \dots, \rho^{(m)}$ as in Definition 1.5 such that for every large enough n and every symmetric non-negative $\Sigma \in \mathbb{R}[S_n]$, we have that*

$$\lambda(S_n, \Sigma) = \max_{i=1}^m \lambda_1(\rho_n^{(i)}(\Sigma))?$$

This work studies a special case of this question, where Σ is *normal*, in the sense that the coefficients α_σ are constant on every conjugacy class³. In this case, the eigenvalues of the different irreps can be computed directly from character values (see Lemma 2.1 below). Our first result gives a positive answer to Question 1.6 (and to Question 1.4) when Σ is a single conjugacy class. For every $n \geq 8$, consider the following set of eight irreps:

$$\text{eight}_n = \left\{ \begin{array}{ccccc} (n-1, 1) & (n-2, 2) & (n-3, 3) & (n-3, 2, 1) & (n-4, 4) \\ (n-1, 1)^t & (n-2, 2)^t & (n-2, 1, 1)^t & & \end{array} \right\} \subset \widehat{S}_n;$$

Here ρ^t denotes the representation $\text{sgn} \otimes \rho$, whose Young diagram is given by transposing that of ρ , e.g. $(n-1, 1)^t = (2, 1, 1, \dots, 1)$.

Theorem 1.7. *There exists $N_0 \in \mathbb{N}$ such that for every $n \geq N_0$, if $\Sigma \in S_n$ is a full, single conjugacy class, then $\lambda(S_n, \Sigma)$ is attained by one of the eight irreps in eight_n :*

$$\lambda(S_n, \Sigma) = \max_{\rho \in \text{eight}_n} \lambda_1(\rho(\Sigma)).$$

In the language of Definition 1.3 and Question 1.4, this implies that for n large enough, if $\Sigma \in S_n$ is a conjugacy class, then $\lambda(S_n, \Sigma) = \max(\lambda(4, \Sigma), \lambda^{\text{sgn}}(2, \Sigma))$. The proof of Theorem 1.7 relies heavily on asymptotically sharp character bounds due to Larsen and Shalev [LS08] – see Section 2. The statement of Theorem 1.7 does not hold for $n = 16$: for $\Sigma = [(1\ 2\ 3\ 4\ 5)(6\ 7\ 8\ 9\ 10)(11\ 12\ 13\ 14\ 15)]^{S_{16}}$, the largest non-trivial eigenvalue $\lambda(S_{16}, \Sigma)$ is associated with the irreps $(11, 5)$ and its transpose. However, simulations suggest that this is the largest counter-example:

Conjecture 1.8. *Theorem 1.7 holds with $N_0 = 17$.*

When $\Sigma \in \mathbb{R}[S_n]$ is a general non-negative normal element, not only are the eight irreps from Theorem 1.7 insufficient, but no finite set of families of irreps as in Definition 1.5 suffices to capture $\lambda(S_n, \Sigma)$, and so the answer to Questions 1.4 and 1.6 turns out to be negative:

Theorem 1.9. *For every $k \geq 1$ and every large enough n , there is a non-negative normal element $\Sigma \in \mathbb{R}[S_n]$ such that*

$$\max(\lambda(k, \Sigma), \lambda^{\text{sgn}}(k, \Sigma)) \leq \lambda(\text{Cay}(S_n, \Sigma)).$$

³Note that a normal element $\Sigma \in S_n$ is symmetric, as σ^{-1} is conjugate to σ for every $\sigma \in S_n$.

Stated differently, no family of irreps as in Definition 1.5 suffices to capture $\lambda(S_n, \Sigma)$ for general non-negative normal Σ . However, our analysis for the case of a single conjugacy class does readily show the following.

Theorem 1.10. *Let $\Sigma = \sum \alpha_\sigma \sigma \in \mathbb{R}[S_n]$ be non-negative and normal, and denote $|\Sigma| \stackrel{\text{def}}{=} \rho_{\text{triv}}(\Sigma) = \sum \alpha_\sigma$. Then the spectral gap $|\Sigma| - \lambda(S_n, \Sigma)$ is bounded by the spectral gap of the Schreier graph $\text{Sch}(S_n \curvearrowright [n], \Sigma)$ multiplied by a decaying multiplicative factor:*

$$|\Sigma| - \lambda(S_n, \Sigma) \geq [|\Sigma| - \lambda(1, \Sigma)] \cdot [1 - o_n(1)].$$

All the evidence we have so far supports the following generalization of Aldous' spectral gap conjecture (Theorem 1.1), which was raised, as was Question 1.4 above, during discussions between Gady Kozma and the second author:

Conjecture 1.11 (Kozma-Puder). *There is a number $k \geq 4$ and a universal constant $0 < c < 1$, such that for large enough n and for every symmetric non-negative $\Sigma \in \mathbb{R}[S_n]$,*

$$|\Sigma| - \lambda(S_n, \Sigma) \geq c \cdot [|\Sigma| - \max(\lambda(k, \Sigma), \lambda^{\text{sgn}}(k, \Sigma))].$$

This conjecture, if true, would have far-reaching consequences. It would yield that random pairs of permutations in S_n give rise to a uniform family of expanders, which is a long standing open question (e.g., [Lub12, Problem 2.28]). It would also yield a conjecture of Babai [BS92, Conjecture 1.7] that for every generating set Σ of A_n , the diameter of $\text{Cay}(A_n, \Sigma)$ is bounded by some n^c where c is a universal constant. See Section 4 for more details.

Remark 1.12. There have been a few attempts to find phenomena as the ones described here in families of groups other than the symmetric groups. Recently, Cesi found an analog of Aldous' conjecture in signed symmetric groups [Ces20]. Greenhut found a small set of irreps of the groups $\text{SL}_n(\mathbb{F}_q)$ which detect the mere existence of a spectral gap for all n and prime powers q [Gre20].

The paper is organized as follows. In Section 2 we consider sets consisting of a single conjugacy class and prove Theorem 1.7. Section 3 deals with arbitrary normal sets and contains the proofs of Theorems 1.9 and 1.10. In Section 4 we further discuss Conjecture 1.11 and its consequences.

2 A Single Conjugacy Class

We start with the following standard lemma⁴, which explains why all eigenvalues of $\text{Cay}(G, \Sigma)$ can be read off from the character table of G when $\Sigma \in \mathbb{R}[G]$ is normal. The important quantity here is the normalized character of $\rho \in \widehat{S_n}$ which we denote by $\tilde{\chi}_\rho$:

$$\tilde{\chi}_\rho(\sigma) \stackrel{\text{def}}{=} \frac{\chi_\rho(\sigma)}{\chi_\rho(1)} \quad (\sigma \in S_n),$$

where $\chi_\rho: \sigma \mapsto \text{trace}(\rho(\sigma))$ is the character of ρ . The character table of S_n consists only of integers (cf. [Ful97]), and $|\chi_\rho(\sigma)| \leq \chi_\rho(1)$, so that $\tilde{\chi}_\rho(\sigma) \in \mathbb{Q} \cap [-1, 1]$ for every $\rho \in \widehat{S_n}$, $\sigma \in S_n$.

Lemma 2.1. *Let G be a finite group, $\Sigma \in \mathbb{R}[G]$ a normal element, and $\rho \in \widehat{G}$. Denoting by α_C the coefficient in Σ of each σ in the conjugacy class C , and by $\tilde{\chi}_\rho(C)$ the value of $\tilde{\chi}_\rho$ on C , the matrix $\rho(\Sigma)$ equals the scalar*

$$\sum_{C \in \text{Conj}(G)} \alpha_C |C| \tilde{\chi}_\rho(C). \quad (2.1)$$

⁴This lemma was popularized by Diaconis, for example in his book [Dia88].

Proof. Since Σ is in the center of $\mathbb{C}[G]$, $\rho(\Sigma)$ is an endomorphism of an irreducible representation, hence a scalar by Schur's Lemma, and the trace of $\rho(\Sigma)$ is $\sum_C \alpha_C |C| \chi_\rho(C)$. \square

This section studies the case of a single conjugacy class $\Sigma = \mathbf{1}_C$, which means that

$$\lambda(S_n, \Sigma) = |C| \cdot \max_{\rho \in \widehat{S_n} \setminus \{\text{triv}, \text{sgn}\}} \tilde{\chi}_\rho(C). \quad (2.2)$$

In this case Theorem 1.7 states that for every $n \geq N_0$ and $\sigma \in S_n$ we have

$$\max_{\rho \in \widehat{S_n} \setminus \{\text{triv}, \text{sgn}\}} \tilde{\chi}_\rho(\sigma) = \max_{\rho \in \text{right}_n} \tilde{\chi}_\rho(\sigma). \quad (2.3)$$

For a given $\sigma \in S_n$, if the maximum in the left hand side of (2.3) is obtained by some $\rho \in \widehat{S_n} \setminus \{\text{triv}, \text{sgn}\}$, we say that “ ρ rules for σ in S_n ”.

ρ rules

Our main tool in analyzing the normalized characters $\tilde{\chi}_\rho(\sigma)$ is the following asymptotically sharp character bounds established by Larsen and Shalev. We use the notation $c_\ell(\sigma)$ for the number of ℓ -cycles in the permutation $\sigma \in S_n$. For example, $c_1(\sigma)$ is the number of fixed points.

$c_\ell(\sigma)$

Theorem 2.2 ([LS08, Theorem 1.3]). *Let $\sigma \in S_n$ and let $f = \max(c_1(\sigma), 1)$. For every irrep $\rho \in \widehat{S_n}$, its character χ_ρ satisfies*

$$\tilde{\chi}_\rho(\sigma) \leq |\chi_\rho(1)|^{-\frac{\log(n/f)}{2 \log n} + \varepsilon_n}, \quad (2.4)$$

where ε_n is a real number tending to 0 as $n \rightarrow \infty$.

Our strategy in proving Theorem 1.7 is as follows: using Theorem 2.2, we show that for large enough n , if $\sigma \in S_n$ has **exactly two** fixed points, then the standard representation std rules, namely, the maximal normalized character is $\tilde{\chi}_{\text{std}}(\sigma) = \frac{1}{n-1}$. Using a simple induction argument we then show that the same is true for every large enough n when $c_1(\sigma) \geq 2$. Finally, we use Theorem 2.2 again to deal with the case $c_1(\sigma) \in \{0, 1\}$. Indeed, Theorem 1.7 follows immediately from the following two propositions, which we prove in the following two subsections.

Proposition 2.3. *There is some $N_1 \in \mathbb{N}$ such that for every $n \geq N_1$ and every $\sigma \in S_n$ with $c_1(\sigma) \geq 2$, the standard irrep $\text{std} = (n-1, 1)$ rules.*

Proposition 2.4. *There is some $N_2 \in \mathbb{N}$ such that for every $n \geq N_2$ and every $\sigma \in S_n$ with $c_1(\sigma) \leq 1$, one of the irreps in right_n rules.*

2.1 The case $c_1(\sigma) \geq 2$

The following lemma goes back to Frobenius (cf. [Mac95, §I.7.14]). We give its proof for completeness.

Lemma 2.5. *Let $\{\rho_n \in \widehat{S_n}\}_{n \geq n_0}$ be a family of irreps as in Definition 1.5 with constant structure outside the first row, so that the first row of ρ_n has exactly $n-k$ blocks. Then there is a polynomial $p \in \mathbb{Q}[c_1, \dots, c_k]$ so that for $n \geq 2k$ and $\sigma \in S_n$,*

$$\begin{aligned} \chi_{\rho_n}(\sigma) &= p(c_1(\sigma), \dots, c_k(\sigma)), \text{ and} \\ \chi_{\rho_n^t}(\sigma) &= \text{sgn}(\sigma) \cdot p(c_1(\sigma), \dots, c_k(\sigma)). \end{aligned}$$

In particular, $\dim \rho_n = \chi_{\rho_n}(1)$ is given by a polynomial in n , equal to $p(n, 0, \dots, 0)$.

The fact that $\chi_{\rho_n}(1) = \dim \rho_n$ is given by a polynomial in n is also evident from the hook length formula (e.g., [FH91, Formula 4.12]). The polynomials associated with every family of irreps with at most four blocks outside the first row are listed in Table 1.

ρ	$\dim(\rho) = \chi_\rho(1)$	$\chi_\rho(\sigma)$ when $c_i(\sigma) = c_i$
(n)	1	1
$(n-1, 1)$	$n-1$	$c_1 - 1$
$(n-2, 2)$	$\frac{n(n-3)}{2}$	$\frac{c_1(c_1-3)}{2} + c_2$
$(n-2, 1, 1)$	$\frac{(n-1)(n-2)}{2}$	$\frac{(c_1-1)(c_1-2)}{2} - c_2$
$(n-3, 3)$	$\frac{n(n-1)(n-5)}{6}$	$\frac{c_1(c_1-1)(c_1-5)}{6} + (c_1-1)c_2 + c_3$
$(n-3, 2, 1)$	$\frac{n(n-2)(n-4)}{3}$	$\frac{c_1(c_1-2)(c_1-4)}{3} - c_3$
$(n-3, 1, 1, 1)$	$\frac{(n-1)(n-2)(n-3)}{6}$	$\frac{(c_1-1)(c_1-2)(c_1-3)}{6} - (c_1-1)c_2 + c_3$
$(n-4, 4)$	$\frac{n(n-1)(n-2)(n-7)}{24}$	$\frac{c_1(c_1-1)(c_1-2)(c_1-7)}{24} + \frac{(c_1^2-3c_1-1)c_2}{2} + \frac{c_2^2}{2} + (c_1-1)c_3 + c_4$
$(n-4, 3, 1)$	$\frac{n(n-1)(n-3)(n-6)}{8}$	$\frac{c_1(c_1-1)(c_1-3)(c_1-6)}{8} + \frac{(c_1^2-3c_1+3)c_2}{2} - \frac{c_2^2}{2} - c_4$
$(n-4, 2, 2)$	$\frac{n(n-1)(n-4)(n-5)}{12}$	$\frac{c_1(c_1-1)(c_1-4)(c_1-5)}{12} + (c_2-2)c_2 - (c_1-1)c_3$
$(n-4, 2, 1, 1)$	$\frac{n(n-2)(n-3)(n-5)}{8}$	$\frac{c_1(c_1-2)(c_1-3)(c_1-5)}{8} - \frac{(c_1^2-3c_1-1)c_2}{2} - \frac{c_2^2}{2} + c_4$
$(n-4, 1, 1, 1, 1)$	$\frac{(n-1)(n-2)(n-3)(n-4)}{24}$	$\frac{(c_1-1)(c_1-2)(c_1-3)(c_1-4)}{24} - \frac{(c_1^2-3c_1+3)c_2}{2} + \frac{c_2^2}{2} + (c_1-1)c_3 - c_4$

Table 1: Dimensions and characters of irreps with at most four blocks outside the first row.

Proof. For a partition $\lambda = (\lambda^{(1)}, \dots, \lambda^{(\ell)}) \vdash n$, let M^λ denote the reducible representation associated with λ : this is the permutation representation describing the action of S_n on partitions of $\{1, \dots, n\}$ with sizes of blocks given by λ (see [Ful97, §7.2]). It is not hard to see that the statement of the lemma holds for families $\{M^{\lambda_n}\}_{n \geq n_0}$ when λ_{n+1} is obtained from λ_n by increasing $\lambda^{(1)}$ by one. For example, if $\lambda_n = (n-3, 2, 1)$, then $\chi_{M^{\lambda_n}}(\sigma)$, which equals the number of fixed points of σ in this action of S_n , is $c_1 \cdot \binom{c_1-1}{2} + c_1 c_2$.

For every $\lambda \vdash n$ as above, the character χ_{ρ^λ} of the irrep ρ^λ corresponding to λ is given by a linear combination with integer coefficients of the representations $\{M^\mu \mid \mu \triangleleft \lambda\}$, where “ \triangleleft ” marks the dominance relation (see [Ful97, §7.2]). Moreover, when $\lambda^{(1)} = n - k \geq \frac{n}{2}$, this linear combination is independent of n , namely, the coefficient of every $\mu \triangleleft \lambda$ depends only on the structure of μ outside the first row. For example,

$$\chi_{\rho^{(n-2,1,1)}} = \chi_{M^{(n-2,1,1)}} - \chi_{M^{(n-2,2)}} - \chi_{M^{(n-1,1)}} + \chi_{M^{(n)}}.$$

The statement of the lemma follows. \square

Lemma 2.6. *Let $n \geq 13$, and let $\rho \in \widehat{S}_n$ be an irrep whose Young diagram has at least three blocks outside the first row and at least three blocks outside the first column. Then $\dim \rho \geq n^{2.05}$.*

Proof. If ρ has exactly three blocks outside the first row then ρ is one of $(n-3, 3)$, $(n-3, 2, 1)$ or $(n-3, 1, 1, 1)$, in which case its dimension is $\frac{n(n-1)(n-5)}{6}$, $\frac{n(n-2)(n-4)}{3}$ or $\frac{(n-1)(n-2)(n-3)}{6}$, respectively. In each of these cases $\dim \rho \geq n^{2.05}$ for $n \geq 13$. The transpose case where ρ has exactly three blocks outside the first column is identical.

For the case ρ has at least four blocks outside the first row/column we use induction on n . It is easy to check directly (on a computer) that the statement is true for $n = 13, 14$: all 93 irreps of S_{13} and 127 irreps of S_{14} satisfying the assumption of the lemma have dimension $\geq n^{2.05}$. For $n \geq 15$, we assume the statement holds for $n-1$ and for $n-2$ and that $\rho \in \widehat{S}_n$ has at least four blocks outside the first row and at least four blocks outside the first column. By the branching rule, $\dim \rho = \sum_{\rho' = \rho - \square} \dim \rho'$, the sum being over all $\rho' \in \widehat{S}_{n-1}$ obtained from ρ by removing one block. If the Young diagram corresponding to ρ is not a rectangle, there are at least two such ρ' , each with at least three blocks outside the first row and outside the first column, and we are done as $2(n-1)^{2.05} \geq n^{2.05}$ for $n \geq 15$. Finally, if ρ is a rectangle, it has at

least $\frac{n}{2} > 4$ blocks outside the first row and outside the first column, and there are exactly two ways to remove two blocks from ρ . The branching rule now gives $\dim \rho = \dim \rho_1 + \dim \rho_2$ with $\rho_1, \rho_2 \in \widehat{S_{n-2}}$ satisfying the assumption in the lemma. We are done as $2(n-2)^{2.05} \geq n^{2.05}$ for $n \geq 15$. \square

Lemma 2.7. *There is some $N_3 \in \mathbb{N}$ such that std rules for every $n \geq N_3$ and $\sigma \in S_n$ with $c_1(\sigma) = 2$. Namely, $\tilde{\chi}_\rho(\sigma) \leq \tilde{\chi}_{\text{std}}(\sigma) = \frac{1}{n-1}$ for every $\rho \in \widehat{S_n} \setminus \{\text{triv}, \text{sgn}\}$.*

Proof. Fix $N_3 \geq 13$ so that for every $n \geq N_3$, the term ε_n in (2.4) satisfies

$$2.05 \cdot \left(-\frac{1}{2} + \frac{\log 2}{2 \log n} + \varepsilon_n \right) \leq -1.$$

Then, if $n \geq N_3$ and $\rho \in \widehat{S_n}$ satisfies the assumptions in Lemma 2.6, it follows from Theorem 2.2 that

$$\tilde{\chi}_\rho(\sigma) \leq (n^{2.05})^{-\frac{1}{2} + \frac{\log 2}{2 \log n} + \varepsilon_n} \leq \frac{1}{n} \leq \frac{1}{n-1}.$$

Finally, if ρ is one of the five remaining irreps $(n-1, 1)^t$, $(n-2, 2)$, $(n-2, 2)^t$, $(n-2, 1, 1)$ and $(n-2, 1, 1)^t$, we use the explicit expressions in Table 1 to show that $\tilde{\chi}_\rho(\sigma) \leq \frac{1}{n-1}$. For example, if $\rho = (n-2, 1, 1)^t$ and σ is odd, we get

$$\tilde{\chi}_\rho(\sigma) = \frac{2c_2(\sigma)}{(n-1)(n-2)} \leq \frac{2 \cdot \frac{n-2}{2}}{(n-1)(n-2)} = \frac{1}{n-1},$$

so at worst there is a tie between $(n-1, 1)$ and $(n-2, 1, 1)^t$ when $c_1(\sigma) = 2$. \square

The following easy but crucial lemma says that the normalized character of a permutation with at least one fixed point (which we can take without loss of generality to be n) is bounded from both sides by normalized characters with one block omitted.

Lemma 2.8. *Let $\sigma \in S_n$ be a permutation satisfying $\sigma(n) = n$. Then σ can be considered also as an element of S_{n-1} , and for every $\rho \in \widehat{S_n}$, the normalized character $\tilde{\chi}_\rho(\sigma)$ is a weighted average of the normalized characters $\{\tilde{\chi}_{\rho'}(\sigma)\}_{\rho'=\rho-\square \in \widehat{S_{n-1}}}$.*

Here ρ' runs over all Young diagrams obtained from ρ by deletion of a single block.

Proof. By the branching rule $\chi_\rho(1) = \sum_{\rho'=\rho-\square} \chi_{\rho'}(1)$, and, similarly $\chi_\rho(\sigma) = \sum_{\rho'=\rho-\square} \chi_{\rho'}(\sigma)$. For any real numbers x_1, \dots, x_k and positive real numbers y_1, \dots, y_k , $\frac{x_1+\dots+x_k}{y_1+\dots+y_k}$ is a convex combination of $\frac{x_1}{y_1}, \dots, \frac{x_k}{y_k}$. \square

Corollary 2.9. *For $\sigma \in S_n$ with $\sigma(n) = n$, if std rules for σ in S_{n-1} , it also rules for σ in S_n .*

Proof. Denote the standard irrep in S_{n-1} by std' . Then

$$\tilde{\chi}_{\text{std}}(\sigma) = \frac{c_1(\sigma) - 1}{n-1} \geq \frac{c_1(\sigma) - 2}{n-2} = \tilde{\chi}_{\text{std}'}(\sigma).$$

On the other hand, if $\rho \in \widehat{S_n} \setminus \{\text{triv}, \text{sgn}, \text{std}, \text{std}^t\}$, then removing a block from ρ does not yield neither the trivial nor the sign irreps of S_{n-1} , and so by Lemma 2.8,

$$\tilde{\chi}_\rho(\sigma) \leq \tilde{\chi}_{\text{std}'}(\sigma).$$

Finally, regarding std^t , if σ is even then $\tilde{\chi}_{\text{std}^t}(\sigma) = \tilde{\chi}_{\text{std}}(\sigma)$, and if σ is odd then as std rules for σ in S_{n-1} , we have $c_1(\sigma) \geq 2$ and $\tilde{\chi}_{\text{std}^t}(\sigma) = -\tilde{\chi}_{\text{std}}(\sigma) < \tilde{\chi}_{\text{std}}(\sigma)$. \square

Recall that the *support* of $\sigma \in S_n$ is $\text{supp}(\sigma) = \{i \in \{1, \dots, n\} \mid \sigma(i) \neq i\}$.

supp

Corollary 2.10. *Let N_3 be as in Lemma 2.7. Then std rules for every $n \geq N_3$ and $\sigma \in S_n$ with $c_1(\sigma) \geq 2$ and $|\text{supp}(\sigma)| \geq N_3 - 2$.*

Proof. Let $k = |\text{supp}(\sigma)| + 2 \geq N_3$. By omitting $n - k$ fixed points from σ , we may think of σ as representing a conjugacy class in S_k with exactly two fixed points. By Lemma 2.7, std rules for σ in S_k . By applying Corollary 2.9 $n - k$ times, we deduce that std also rules for σ in S_n . \square

Lemma 2.11. *Let $\sigma \in S_r$ and consider it as a permutation in S_n for any $n \geq r$ by appending $n - r$ fixed points. For large enough n , std rules for σ in S_n .*

Proof. By enlarging r a bit first, we can assume without loss of generality that $r \geq 5$, so every $\rho \in \widehat{S_r} \setminus \{\text{triv}, \text{sgn}\}$ is faithful, and also that $c_1(\sigma) \geq 1$ in S_r , so if std^t rules for some $n \geq r$, so does std. Assume that some $\rho \in \widehat{S_r} \setminus \{\text{triv}, \text{sgn}\}$ rules for σ in S_r . As ρ is faithful, $\tilde{\chi}_\rho(\sigma) < 1$. As n increases, as long as std does not rule, all normalized characters of irreps in $\widehat{S_n} \setminus \{\text{triv}, \text{sgn}\}$ are bounded from above by $\tilde{\chi}_\rho(\sigma)$: indeed, by induction this is true for all such irreps in $\widehat{S_{n-1}}$, and by Lemma 2.8 this is also true for every irrep in $\widehat{S_n} \setminus \{\text{triv}, \text{sgn}, \text{std}, \text{std}^t\}$. For std and std^t this is true by the assumption that std does not rule. In contrast, the normalized character of std, which is $\frac{n - |\text{supp}(\sigma)| - 1}{n - 1}$, tends to 1 as $n \rightarrow \infty$. Therefore, for some n_0 , std rules. Corollary 2.9 then implies that std rules for σ for every $n \geq n_0$. \square

Corollary 2.12. *Let $M \in \mathbb{N}$ be a constant. There is some $N_4 = N_4(M) \in \mathbb{N}$ such that for every $n \geq N_4$ and every $\sigma \in S_n$ with $|\text{supp}(\sigma)| \leq M$, std rules for σ in S_n .*

Proof. Every such σ belongs to a conjugacy class in S_n which is obtained from blowing up (by appending fixed point) a fixed-point-free conjugacy class in S_k for some $k \leq M$. This is a finite set of starting points, which means we need to apply Lemma 2.11 finitely many times. \square

Proof of Proposition 2.3. Let N_3 be the constant from Lemma 2.7 and Corollary 2.10, and $N_4 = N_4(M)$ with $M = N_3 - 3$ be the constant from Corollary 2.12. Set $N_1 = \max(N_3, N_4)$. Then std rules for every $n \geq N_1$ and for every $\sigma \in S_n$ with $c_1(\sigma) \geq 2$. \square

2.2 The case $c_1(\sigma) \leq 1$

The proof strategy of Proposition 2.4 is the same as in the case $c_1(\sigma) = 2$, albeit significantly more tedious. The difference is that the largest normalized character is not necessarily of order $\frac{1}{n}$ as in the $c_1(\sigma) = 2$ case, but can be of order as low as $\frac{1}{n^3}$. For example, this is the case when σ is even, $c_1(\sigma) = c_3(\sigma) = 1$ and $c_2(\sigma) = 0$ (and see Table 2).

Lemma 2.13. *Let $\{\rho_n\}_{n \geq n_0}$ be a family of irreps $\rho_n \in \widehat{S_n}$ as in Definition 1.5 with k blocks outside the first row or outside the first column. Every monomial $c_1^{\alpha_1} c_2^{\alpha_2} \cdots c_k^{\alpha_k}$ in the associated polynomial p from Lemma 2.5 satisfies $\sum_{i=1}^k i \cdot \alpha_i \leq k$. In addition, the polynomial $p(c_1, 0, \dots, 0)$ giving the dimension of ρ_n is of degree exactly k .*

Proof. The first statement holds because it holds for the polynomials depicting the characters of the reducible representations M^λ (see the proof of Lemma 2.5), and the character of ρ_n is equal to a linear combination of M^λ 's with at most k blocks outside the first row (namely, $\lambda^{(1)} \geq n - k$). The second statement is immediate from the hook length formula. \square

Lemma 2.14. *For every large enough n and every $\sigma \in S_n$*

$$f(\sigma) \stackrel{\text{def}}{=} \max_{\rho \in \text{eight}_n} \tilde{\chi}_\rho(\sigma) \geq \frac{3}{n(n-2)(n-4)}.$$

Proof. We deal with six different cases in the following table (the values of normalized characters can be read from Table 1):

Assumptions on σ	lower bound on $f(\sigma)$
$c_1 \geq 2$	$f(\sigma) \geq \tilde{\chi}_{\text{std}}(\sigma) \geq \frac{1}{n-1}$
$c_1 = 1, c_3 \geq 1$	$f(\sigma) \geq \tilde{\chi}_{(n-3,3)}(\sigma) = \frac{6c_3}{n(n-1)(n-5)} \geq \frac{6}{n(n-1)(n-5)}$
$c_1 = 1, c_3 = 0$	$f(\sigma) \geq \tilde{\chi}_{(n-3,2,1)}(\sigma) = \frac{3}{n(n-2)(n-4)}$
$c_1 = 0, c_2 \geq 1$	$f(\sigma) \geq \tilde{\chi}_{(n-2,2)}(\sigma) = \frac{2c_2(\sigma)}{n(n-3)} \geq \frac{2}{n(n-3)}$
$c_1 = 0, c_2 = 0, \sigma$ is even	$f(\sigma) \geq \tilde{\chi}_{(n-2,1,1)^t}(\sigma) = \frac{2}{(n-1)(n-2)}$
$c_1 = 0, c_2 = 0, \sigma$ is odd	$f(\sigma) \geq \tilde{\chi}_{\text{std}^t}(\sigma) = \frac{1}{(n-1)}$

□

Lemma 2.15. *Let $\{\rho_n\}_{n \geq n_0}$ be a family of irreps $\rho_n \in \widehat{S}_n$ as in Definition 1.5 with $k \geq 5$ blocks outside the first row or outside the first column. Then for large enough n and every $\sigma \in S_n$ with $c_1(\sigma) \leq 1$,*

$$\tilde{\chi}_{\rho_n}(\sigma) \leq f(\sigma) \stackrel{\text{def}}{=} \max_{\rho \in \text{eigh}_n} \tilde{\chi}_{\rho}(\sigma).$$

Proof. If $c_2(\sigma) \geq 2$ then

$$f(\sigma) \geq \tilde{\chi}_{(n-2,2)}(\sigma) \geq \frac{2(c_2(\sigma) - 1)}{n(n-3)} \geq \frac{2}{n(n-3)}.$$

In contrast, by Lemma 2.13, as $c_2(\sigma), \dots, c_k(\sigma) \leq n$, we have $\chi_{\rho_n}(\sigma) \leq O(n^{\lfloor k/2 \rfloor})$ and $\chi_{\rho_n}(1)$ is a polynomial in n of degree k . We deduce that the normalized character satisfies $\tilde{\chi}_{\rho_n}(\sigma) \leq O(n^{\lfloor k/2 \rfloor - k}) = O(n^{-3})$ as $k \geq 5$.

If $c_2(\sigma) \leq 1$ then by Lemma 2.14 $f(\sigma) \geq \frac{3}{n(n-2)(n-4)}$ whereas, by Lemma 2.13, $\chi_{\rho_n}(\sigma) \leq O(n^{\lfloor k/3 \rfloor})$ and so $\tilde{\chi}_{\rho_n}(\sigma) \leq O(n^{\lfloor k/3 \rfloor - k}) = O(n^{-4})$ as $k \geq 5$. □

Of course, Lemma 2.15 suffices to deal with every family of irreps separately, but not with all irreps uniformly. For this, we need to use Larsen-Shalev's Theorem 2.2. First, as above, we need a uniform lower bound on the dimension of almost all irreps:

Lemma 2.16. *Let $n \geq 39$, and let $\rho \in \widehat{S}_n$ be an irrep represented by a Young diagram with at least 14 blocks outside the first row and at least 14 blocks outside the first column. Then $\dim \rho \geq n^{6.05}$.*

Proof. We verified the statement numerically for $n = 39, \dots, 48$. For the general case, assume that ρ has exactly 14 blocks outside the first row (the transpose case is identical). Consider the hook lengths at the first row of the Young diagram associated with ρ . The hook length at the 15th block is $n - 28$, since the second row is of length ≤ 14 . The hook length at the 16th block is $n - 29$ and so on. For $1 \leq i \leq 14$, the first $(i - 1)$ columns contain at least $2(i - 1)$ blocks, so the hook length of the i th block is at most $n - 2(i - 1)$. The product of hook lengths of all blocks outside the first row is at most $14!$ (by the hook length formula for Young diagrams with 14 blocks). Thus, by the hook length formula, we obtain that

$$\begin{aligned} \dim \rho &= \frac{n!}{\text{product of hook lengths}} \\ &\geq \frac{n!}{n(n-2)(n-4) \cdots (n-26) \cdot (n-28)! \cdot 14!} = \frac{(n-1)(n-3)(n-5) \cdots (n-27)}{14!}, \end{aligned}$$

which is greater than $n^{6.05}$ for $n \geq 47$. For $n \geq 49$ and $\rho \in \widehat{S}_n$ with at least 15 blocks outside the first row or outside the first column, we proceed by induction exactly as in the proof of Lemma 2.6. □

Proof of Proposition 2.4. As in the proof of Lemma 2.7, we deduce from Theorem 2.2 and Lemma 2.16 that for large enough n , for all $\sigma \in S_n$ with $c_1(\sigma) \leq 1$ and for all $\rho \in \widehat{S}_n$ with at least 14 blocks outside the first row or outside the first column, we have

$$\tilde{\chi}_\rho(\sigma) \leq \frac{1}{n^3},$$

which, by Lemma 2.14, is less than the maximal normalized character of σ among the irreps in \mathbf{eight}_n . For the finite number of families of irreps with $5 \leq k \leq 13$, where k is the number of blocks outside the first row or column, we use Lemma 2.15. Finally, the normalized characters of all irreps with $k \leq 4$ appear in Table 1. Comparing them for large n yields that for every $\sigma \in S_n$ with $c_1(\sigma) \leq 1$, some $\rho \in \mathbf{eight}_n$ rules. We omit the technical details of this comparison, but see Tables 2 and 3 for the ruling irrep in each case. \square

ρ	ρ rules for even σ when:
$(n-1, 1)$ $(n-1, 1)^t$	$c_1 \geq 2$
$(n-2, 2)$ $(n-2, 2)^t$	$c_1 = 1, c_2 \geq 2$ $c_1 = 0, c_2 \geq 1$
$(n-2, 1, 1)$ $(n-2, 1, 1)^t$	$c_1 = 0, c_2 = 0, c_3 \leq \frac{n-4}{3}$
$(n-3, 3)$ $(n-3, 3)^t$	$c_1 = 1, c_2 = 1, c_3 \geq 1$ $c_1 = 1, c_2 = 0, c_3 \geq 2$ $c_1 = 1, c_2 = 0, c_3 = 1, c_4 \leq \frac{n-5}{4}$ $c_1 = 0, c_2 = 0, c_3 = \frac{n}{3}$
$(n-3, 2, 1)$ $(n-3, 2, 1)^t$	$c_1 = 1, c_2 = 1, c_3 = 0, c_4 \leq \frac{n+3}{8}$ $c_1 = 1, c_2 = 0, c_3 = 0, c_4 \leq \frac{n-5}{8}$
$(n-4, 4)$ $(n-4, 4)^t$	$c_1 = 1, c_2 = 1, c_3 = 0, c_4 \geq \frac{n+4}{8}$ $c_1 = 1, c_2 = 0, c_3 = 1, c_4 = \frac{n-4}{4}$ $c_1 = 1, c_2 = 0, c_3 = 0, c_4 \geq \frac{n-4}{8}$

Table 2: For large enough n , this table shows which $\rho \in \mathbf{eight}_n$ rules for every **even** $\sigma \in S_n$. Note that at least one of every pair of irreps in the left column belongs to \mathbf{eight}_n .

ρ	ρ rules for odd σ when:
$(n-1, 1)$ $(n-1, 1)^t$	$c_1 \geq 2$ $c_1 = 0, c_2 \leq \frac{n-3}{2}$
$(n-2, 2)$ $(n-2, 2)^t$	$c_1 = 0, c_2 = \frac{n}{2}$ $c_1 = 1, c_2 = 0$
$(n-2, 1, 1)^t$	$c_1 = 2, c_2 = \frac{n-2}{2}$ $c_1 = 1, c_2 \geq 2$ $c_1 = 1, c_2 = 1, c_3 \leq \frac{n-4}{3}$
$(n-3, 3)$	$c_1 = 1, c_2 = 1, c_3 = \frac{n-3}{3}$

Table 3: For large enough n , this table shows which $\rho \in \mathbf{eight}_n$ rules for every **odd** $\sigma \in S_n$.

3 Arbitrary Normal Sets

In this section we prove the negative result, Theorem 1.9, stating that no finite set of families of irreps is enough to capture $\lambda(S_n, \Sigma)$ for an arbitrary symmetric non-negative element $\Sigma \in \mathbb{R}[S_n]$,

nor even for a normal non-negative element $\Sigma \in \mathbb{R}[S_n]$. We also prove the positive result, Theorem 1.10, which says that for every normal non-negative $\Sigma \in \mathbb{R}[S_n]$, std alone captures the spectral gap $|\Sigma| - \lambda(S_n, \Sigma)$ up to a decaying multiplicative factor.

3.1 A negative result

Before proving Theorem 1.9, we remark that the irreducible characters of S_n constitute a linear basis for the space of class functions on S_n , hence there are normal elements $\Sigma \in \mathbb{R}[S_n]$ giving any prescribed values for the eigenvalues of every irreps as in (2.1). One could still hope that as we restrict to *non-negative* normal elements, the answer to Question 1.6 would still be affirmative in the case of normal sets. However, this is not the case, as we now prove.

Proof of Theorem 1.9. Let $\rho^{(1)}, \dots, \rho^{(m)}$ be arbitrary families of non-trivial irreps of S_n as in Definition 1.5. Assume each of these families has at most k blocks outside the first row/column. In particular, for $n \geq 2k$, the evaluation of the characters $\rho^{(1)}, \dots, \rho^{(m)}$ on $\sigma \in S_n$ depends only on the numbers $c_1(\sigma), \dots, c_k(\sigma)$ of short cycles – see Lemma 2.5. Moreover, as the characters are given by polynomials in c_1, \dots, c_k , their expected values for some $\Sigma \in \mathbb{R}[S_n]$ depend only on the *joint distribution* of $\{(c_1(\sigma), \dots, c_k(\sigma))\}_{\sigma \in \Sigma}$, and even more particularly on the distribution in Σ of the monomials $c_1^{\alpha_1} \dots c_k^{\alpha_k}$ with $\sum i \cdot \alpha_i \leq k$ (see Lemma 2.13).

Now consider the uniform distribution on S_{2k} . By orthogonality of irreducible characters, for every $\text{triv} \neq \rho \in \widehat{S_{2k}}$, the expected value of $\chi_\rho(\sigma)$ is zero. Now, for every $n \geq 2k$, if $\Sigma \in \mathbb{R}[S_n]$ is a normal non-negative element with the same joint distribution of c_1, \dots, c_k as the uniform distribution in S_{2k} , we get that the eigenvalues of $\text{Cay}(S_n, \Sigma)$ associated with $\rho_n^{(1)}, \dots, \rho_n^{(m)}$ all vanish. For large enough n , one can construct such Σ with large values of, say, c_{k+1} , and supported on odd conjugacy classes. This would assure that there is some irrep ρ with $k+1$ blocks outside the first row or outside the first column with large positive average value of the character χ_ρ . Hence $\rho^{(1)}, \dots, \rho^{(m)}$ do not rule for such Σ . \square

Example 3.1. Let us illustrate the proof for the families $\rho^{(1)} = (n-1, 1)$, $\rho^{(2)} = (n-2, 2)$ and $\rho^{(3)} = (n-2, 1, 1)^t$ and $n = 100$. So we can take $k = 2$ and consider the joint distribution of (c_1, c_2) in S_4 :

c_1	c_2	probability
4	0	1/24
2	1	1/4
1	0	1/3
0	2	1/8
0	0	1/4

Consider the following permutations in S_{100} :

$$\begin{aligned}
\sigma_1 &= (1 \ 2 \ 3) (4 \ 5 \ 6) \dots (88 \ 89 \ 90) (91 \dots 96) \\
\sigma_2 &= (1 \ 2 \ 3) (4 \ 5 \ 6) \dots (94 \ 95 \ 96) (97 \ 98) \\
\sigma_3 &= (1 \ 2 \ 3) (4 \ 5 \ 6) \dots (91 \ 92 \ 93) (94 \dots 99) \\
\sigma_4 &= (1 \ 2 \ 3) (4 \ 5 \ 6) \dots (88 \ 89 \ 90) (91 \dots 96) (97 \ 98) (99 \ 100) \\
\sigma_5 &= (1 \ 2 \ 3) (4 \ 5 \ 6) \dots (94 \ 95 \ 96) (97 \dots 100)
\end{aligned}$$

and notice that they all have many 3-cycles and are all odd. Define $\Sigma \in \mathbb{R}[S_{100}]$ by

$$\Sigma = \sum_{i=1}^5 \alpha_i \cdot \sigma_i^{S_{100}}$$

so that $\alpha_i \cdot |\sigma_i^{S_{100}}|$ is equal to the probability in the i th line in the table. For example, $\alpha_1 \cdot |\sigma_1^{S_{100}}| = \frac{1}{24}$. This choice of Σ assures that for $i = 1, \dots, m$, $\rho^{(i)}(\Sigma)$ is the zero matrix, but there is an

irrep with three blocks outside the first row or column with $\rho(\Sigma)$ being a positive scalar matrix (in our case, this is true for the irreps $(97, 3)$, $(97, 2, 1)^t$ and $(97, 1, 1, 1)$).

Remark 3.2. In fact, it seems that the proof of Theorem 1.9 can work also with the joint distribution of (c_1, \dots, c_k) induced from the uniform distribution on S_k , rather than on S_{2k} . For example, when $k = 2$, the joint distribution in S_2 of (c_1, c_2) is $(2, 0)$ and $(0, 1)$ each with probability $\frac{1}{2}$. For any normal $\Sigma \in \mathbb{R}[S_n]$ with such joint distribution on (c_1, c_2) , the matrix $\rho(\Sigma)$ is zero for $\rho \neq \text{triv}, \text{sgn}$ with at most 2 blocks outside the first row or the first column.

Remark 3.3. Theorem 1.9 is true also if one restricts attention to normal sets in S_n , namely, to normal elements $\Sigma \in \mathbb{R}[S_n]$ with 0–1 coefficients. The point is that there is a lot of flexibility in construction Σ in the proof, so as $n \rightarrow \infty$ one can construct normal sets with joint distribution of (c_1, \dots, c_k) tending to the one in S_{2k} (or in S_k), while keeping the average value of c_{k+1} at least, say, $\frac{n}{2(k+1)}$, and all conjugacy classes odd. This assures that some irrep with $k+1$ blocks outside the first row/column eventually beats every irrep with at most k blocks outside the first row/column.

3.2 A positive result

We now prove Theorem 1.10, based on the following lemma:

Lemma 3.4. *Let N_0 be the constant from Theorem 1.7. For every $n \geq N_0$ there is a constant $\delta_n > 0$ tending to zero as $n \rightarrow \infty$, so that for every $\sigma \in S_n$ and $\rho \in \widehat{S_n} \setminus \{\text{triv}, \text{sgn}\}$ we have*

$$1 - \tilde{\chi}_\rho(\sigma) \geq [1 - \tilde{\chi}_{\text{std}}(\sigma)] \cdot (1 - \delta_n). \quad (3.1)$$

Proof. By Proposition 2.3 (and assuming $N_0 \geq N_1$), if $c_1(\sigma) \geq 2$ then std rules and (3.1) holds even with $\delta_n = 0$. If $c_1(\sigma) \leq 1$ then one of $\rho \in \text{eight}_n$ rules, so it is enough to check (3.1) for each of the seven irreps in $\text{eight}_n \setminus \{\text{std}\}$. And, indeed, (3.1) holds in this cases. The worst case is when σ has $c_1(\sigma) = 0$ and $c_2(\sigma) = \frac{n}{2}$, where $(n-2, 2)$ rules, and $1 - \tilde{\chi}_{(n-2,2)}(\sigma) = 1 - \frac{1}{n-3}$, whereas $1 - \tilde{\chi}_{\text{std}}(\sigma) = 1 + \frac{1}{n-1}$. Hence one can take $\delta_n = \frac{2(n-2)}{n(n-3)}$. \square

Proof of Theorem 1.10. By Lemma 2.1, for any non-negative normal element $\Sigma \in \mathbb{R}[S_n]$, the eigenvalues associated with $\rho \in \widehat{S_n}$ are all equal to $\sum_C \alpha_C |C| \tilde{\chi}_\rho(C)$. The gap between this eigenvalue and $|\Sigma|$ is

$$|\Sigma| - \sum_C \alpha_C |C| \tilde{\chi}_\rho(C) = \sum_C \alpha_C |C| (1 - \tilde{\chi}_\rho(C)).$$

By Lemma 3.4, if $n \geq N_0$, this is at least

$$\sum_C \alpha_C |C| [1 - \tilde{\chi}_{\text{std}}(\sigma)] \cdot (1 - \delta_n) = [|\Sigma| - \lambda_1(\text{std}, \Sigma)] \cdot (1 - \delta_n). \quad \square$$

4 A Conjecture for Arbitrary Symmetric Sets and Its Consequences

This section gives more justification, motivation and background to Conjecture 1.11.

Background and Examples

Recall that Conjecture 1.11 says that there is some $k \geq 4$ such that the (signed and unsigned) actions of S_n on k -tuples determine the spectral gap $|\Sigma| - \lambda(S_n, \Sigma)$ up to a universal multiplicative factor, for every large enough n and every symmetric non-negative element $\Sigma \in \mathbb{R}[S_n]$. In terms of Definition 1.5, this means that there is a finite number of families of irreps which are nearly dominant spectrally, in the same sense. In fact, even the following stronger version of Conjecture 1.11 is conceivable:

Question 4.1. *Is it true that for every symmetric non-negative $\Sigma \in \mathbb{R}[S_n]$, the spectral gap of the Cayley graph $\text{Cay}(S_n, \Sigma)$ is equal, up to a decaying multiplicative factor, to the spectral gap given by the action on 4-tuples and the signed action on 4-tuples, namely, that*

$$|\Sigma| - \lambda(S_n, \Sigma) \geq [|\Sigma| - \max(\lambda(4, \Sigma), \lambda^{\text{sgn}}(4, \Sigma))] \cdot [1 - o_n(1)]?$$

Perhaps even the non-signed action suffices, namely, is it true that

$$|\Sigma| - \lambda(S_n, \Sigma) \geq [|\Sigma| - \lambda(4, \Sigma)] \cdot [1 - o_n(1)]?$$

Note that this question differs from Conjecture 1.11 both by giving a specific set of irreps, and by suggesting that the multiplicative factor tends to 1 as $n \rightarrow \infty$. The evidence we gathered so far supports Conjecture 1.11 and even its stronger version – Question 4.1. The conjecture is certainly true, and with std alone, for normal elements (Theorem 1.10) and for elements supported on transpositions (Theorem 1.1). The example of non-generating sets, where the spectral gap is zero (see Section 1), shows that std alone is not sufficient. The latter is also demonstrated by the following example relating to the generating set consisting of an n -cycle and a sole transposition:

Example 4.2. Let

$$\Sigma = \frac{1}{4} \left[\text{id} + (1\ 2) + (1\ 2 \dots n) + (1\ 2 \dots n)^{-1} \right] \in \mathbb{R}[S_n].$$

In this case, one of $(n-2, 2)$ or $(n-2, 1, 1)$ rules, at least up to a multiplicative constant factor on the spectral gap, whereas $1 - \lambda_1(\text{std}(\Sigma))$ is of a different order. More concretely, it is known [DSC93, Example 1, Page 2139] that

$$1 - \lambda(S_n, \Sigma) \geq \frac{1}{18n^3},$$

and that $\frac{1}{n^3}$ is the right order of the spectral gap. The proof of the upper bound for the spectral gap in [DSC93] can be adapted as follows: Consider $\text{Sch}(S_n \curvearrowright [n]_2, \Sigma)$, and let f be a function on its vertices given by $f((x, y)) = (x - y \bmod n) - \frac{n}{2}$. As f is orthogonal to the constant functions, its Rayleigh quotient $\frac{\langle (I-A)f, f \rangle}{\langle f, f \rangle}$, which roughly equals $\frac{6}{n^3}$, gives a lower bound on the spectral gap of this Schreier graph, where A is the adjacency operator of the graph. The adjacency operator of this graph decomposes into the irreps $(n-2, 1, 1)$, $(n-2, 2)$, $\text{std} = (n-1, 1)$ and $\text{triv} = (n)$, so the second eigenvalue comes from one of $(n-2, 1, 1)$, $(n-2, 2)$ or $(n-1, 1)$. However, $(n-1, 1)$ is not possible because $1 - \lambda_1(\text{std}(\Sigma))$ is of order $\frac{1}{n^2}$: this is the spectral gap of the connected 4-regular graph $\text{Sch}(S_n \curvearrowright [n], \Sigma)$, and it follows from the discrete Cheeger inequality [HLW06, Thm. 4.11] that $1 - \lambda_1(\text{std}(\Sigma)) \geq \frac{1}{32n^2}$. (It can be shown, in fact, that $n^2 \cdot (1 - \lambda_1(\text{std}(\Sigma))) \xrightarrow{n \rightarrow \infty} 1$.) Simulations for small values of n suggest that $(n-2, 1, 1)$ rules for this Σ , but we do not know whether this is true for all n .

Let us also mention another attempt to generalize Theorem 1.1 (Aldous' conjecture) which is attributed to Caputo in [Ces16, Page 301] and is named there “the α -shuffles conjecture”. For $A \subseteq [n] = \{1, \dots, n\}$, let $J_A = \sum_{\sigma \in S_n \text{ s.t. } \text{supp}(\sigma) \subseteq A} \sigma \in \mathbb{R}[S_n]$. The conjecture states that for every linear combination with non-negative coefficients of the J_A 's

$$\Sigma = \sum_{A \subseteq [n]} \alpha_A \cdot J_A, \quad \alpha_A \geq 0$$

the standard representation rules, namely, $\lambda(S_n, \Sigma) = \lambda_1(\text{std}(\Sigma))$. Some special cases of this conjecture were proven in work in progress by Gil Alon, Gady Kozma and the second author.

Consequence 1: Random pairs of permutations expand

It is well known that if $g, h \in S_n$ are chosen uniformly and independently at random then they generate A_n or S_n with probability tending to 1 as $n \rightarrow \infty$ [Dix69]. So with high probability $\lambda(S_n, \frac{1}{4}(g + g^{-1} + h + h^{-1})) < 1$. But do random pairs of S_n also generate a family of expander Cayley graphs? Namely,

Question 4.3. *Is there some $\varepsilon > 0$ so that for uniformly random $g, h \in S_n$*

$$\text{Prob} \left[1 - \lambda \left(S_n, \frac{1}{4}(g + g^{-1} + h + h^{-1}) \right) \geq \varepsilon \right] \xrightarrow{n \rightarrow \infty} 1? \quad (4.1)$$

This question is asked for all finite non abelian simple groups in [Lub12, Problem 2.28] (with uniform ε for all), and is proven in [BGGT15] for finite simple group of Lie type and bounded Lie rank. However, the case of A_n remains wide open. The best known bound is given in [HSZ15, Theorem 1.2], where (4.1) is proven with ε replaced with $\frac{c_1}{n^3(\log n)^{c_2}}$ for some absolute constants c_1, c_2 . In fact, it was a major challenge to show that A_n (or S_n) can even be turned into an expanding family – this was done in [Kas07], and it is still not known whether two generators suffice for this goal (see also [BL18, Problem 3.2(a)]).

Conjecture 1.11, if true, would yield a positive answer to Question 4.3. This is based on the fact, proven in [FJR⁺98], that for every fixed k and r , the Schreier graphs depicting the action of S_n on k -tuples of different elements in $[n]$ with respect to r random permutations form an expander family with high probability, namely, there is some $\varepsilon_k > 0$ so that⁵

$$\text{Prob} \left[1 - \lambda \left(k, \frac{1}{4}(g + g^{-1} + h + h^{-1}) \right) \geq \varepsilon_k \right] \xrightarrow{n \rightarrow \infty} 1.$$

A small adaptation of the argument in [FJR⁺98] allows one to prove an analogous statement for the *signed* action on ℓ -tuples: introducing a random sign to the action on ℓ -tuples only adds more randomness and thus can only increase the expected spectral gap. This explains how Conjecture 1.11, if true, yields a positive answer to Question 4.3.

Remark 4.4. A weaker version of Conjecture 1.11 states that in order to approximate the spectral gap $|\Sigma| - \lambda(S_n, \Sigma)$, it is enough to consider, for every n , a set $\mathcal{L}_n \subseteq \widehat{S_n}$ of irreps with at most ℓ_n blocks outside the first row/column, where ℓ_n grows slowly with n (say, $\ell_n \sim c \cdot \log n$). This may still be enough to yield a positive answer to Question 4.3. We remark that the analog result for the mere existence of a spectral gap, namely, that S_n has no $3 \log n$ -transitive subgroups but itself and A_n , was proved by Jordan in 1895 [Jor95]. Unlike the fact mentioned in Section 1 that there are no 4-transitive subgroups of S_n other than A_n and S_n (for $n \geq 25$), this weaker result of Jordan has elementary proofs, which, in particular, do not depend on the classification of finite simple groups – see [BS87] and the references therein.

⁵In fact, there is evidence to that for every fixed k , the Schreier graph $\text{Sch}(S_n \curvearrowright [n]_k, \Sigma)$ with Σ symmetric and random of fixed size $r \geq 3$, is nearly Ramanujan with high probability, namely, that for every $\varepsilon > 0$, $\lambda(k, \Sigma) < 2\sqrt{r-1} + \varepsilon$ with probability tending to 1 as $n \rightarrow \infty$. This is true for $k = 1$ [Fri08] and for $k = 2$ [BC19], and it seems plausible that the latter proof may be extended to every fixed value of k .

Consequence 2: A bound on the diameter of Cayley graphs of A_n

In [BS92, Conjecture 1.7], Babai conjectured that for every finite simple group G and every generating set $\Sigma \subset G$,

$$\text{diam}(\text{Cay}(G, \Sigma)) \leq (\log |G|)^{O(1)},$$

where the implied constant is absolute. The special case of $G = A_n$ is referred to as a “folklore” conjecture. In this case, the conjecture translates to that $\text{diam}(\text{Cay}(A_n, \Sigma)) \leq n^{O(1)}$. The best upper bound to date is quasi-polynomial and is due to Helfgott and Seress [HS14].

Consider $\text{Sch}(A_n \curvearrowright [n]_k, \Sigma)$, the Schreier graph of A_n on k -tuples with respect to $\Sigma \subset A_n$. This graph has $n(n-1) \cdots (n-k+1) \leq n^k$ vertices, and if Σ is generating, it is connected. By [ACTT18], the spectral gap of the simple random walk on this graph is bounded from below by $(1 + o_n(1)) \frac{54}{n^{3k}}$ (slightly weaker bounds were known before – see the references in [ACTT18]). If conjecture 1.11 holds, this gives a lower bound of the same order on the spectral gap of the simple random walk on the Cayley graph $\text{Cay}(A_n, \Sigma)$. Finally, a lower bound of this kind on the spectral gap yields a polynomial upper bound on the diameter of the Cayley graph: for instance, [SC04, Equation (6.6)] says that if $(1 - \lambda_2)$ is the spectral gap of the simple random walk on a Cayley graph of a finite group G , then the diameter of this graph is at most

$$\frac{3 \log |G|}{\sqrt{1 - \lambda_2}}.$$

Acknowledgments We are grateful to Gady Kozma for highly valuable discussions. We also thank Gil Alon and Avi Wigderson for beneficial comments. D.P. learned about Aldous’ conjecture and its proof by Caputo et al. in a mini-course by Kozma as part of a summer school in Budapest in 2014. We thank the organizers of the school, Miklós Abért, Ágnes Backhausz, László Lóvász, Balázs Szegedy and Bálint Virág, for this wonderful event. Computer simulations mentioned in this work were carried out using Sage. The research was supported by the Israel Science Foundation, ISF grant 1031/17 of O.P. and ISF grant 1071/16 of D.P.

References

- [ACTT18] Sinan G Aksoy, Fan Chung, Michael Tait, and Josh Tobin. The maximum relaxation time of a random walk. *Advances in Applied Mathematics*, 101:1–14, 2018. 16
- [BC19] Charles Bordenave and Benoît Collins. Eigenvalues of random lifts and polynomials of random permutation matrices. *Annals of Mathematics*, 190(3):811–875, 2019. 15
- [BGGT15] Emmanuel Breuillard, Ben J. Green, Robert M. Guralnick, and Terence Tao. Expansion in finite simple groups of Lie type. *Journal of the European Mathematical Society*, 17(6):1367–1434, 2015. 15
- [BL18] Emmanuel Breuillard and Alexander Lubotzky. Expansion in simple groups. arXiv:1807.03879, 2018. 15
- [BS87] László Babai and Ákos Seress. On the degree of transitivity of permutation groups: a short proof. *Journal of Combinatorial Theory, Series A*, 45(2):310–315, 1987. 15
- [BS92] László Babai and Ákos Seress. On the diameter of permutation groups. *European journal of combinatorics*, 13(4):231–243, 1992. 5, 16
- [Cam99] Peter J. Cameron. *Permutation groups*, volume 45 of *Student Texts*. Cambridge University Press, 1999. 3

- [Ces16] Filippo Cesi. A few remarks on the octopus inequality and Aldous’ spectral gap conjecture. *Communications in Algebra*, 44(1):279–302, 2016. 15
- [Ces20] Filippo Cesi. On the spectral gap of some Cayley graphs on the Weyl group $W(B_n)$. *Linear Algebra and its Applications*, 586:274–295, 2020. 5
- [CLR10] Pietro Caputo, Thomas Liggett, and Thomas Richthammer. Proof of Aldous’ spectral gap conjecture. *Journal of the American Mathematical Society*, 23(3):831–851, 2010. 1, 2
- [Dia88] Persi Diaconis. Group representations in probability and statistics. *Lecture Notes-Monograph Series*, 11:i–192, 1988. 1, 5
- [Dix69] John D. Dixon. The probability of generating the symmetric group. *Mathematische Zeitschrift*, 110(3):199–205, 1969. 15
- [DSC93] Persi Diaconis and Laurent Saloff-Coste. Comparison techniques for random walk on finite groups. *The Annals of Probability*, pages 2131–2156, 1993. 3, 14
- [FH91] William Fulton and Joe Harris. *Representation theory: a first course*. Springer, 1991. 1, 6
- [FJR⁺98] Joel Friedman, Antoine Joux, Yuval Roichman, Jacques Stern, and Jean-Pierre Tillich. The action of a few permutations on r -tuples is quickly transitive. *Random Structures and Algorithms*, 12(4):335–350, 1998. 15
- [Fri08] J. Friedman. *A proof of Alon’s second eigenvalue conjecture and related problems*, volume 195 of *Memoirs of the AMS*. AMS, september 2008. 15
- [Ful97] William Fulton. Young tableaux, with applications to representation theory and geometry. *London Mathematical Society Student Texts*, 35, 1997. 1, 5, 7
- [Gre20] Ziv Greenhut. A generation criterion for subsets of special linear groups over finite fields. arXiv preprint arXiv:2002.06461, 2020. 5
- [HLW06] Shlomo Hoory, Nathan Linial, and Avi Wigderson. Expander graphs and their applications. *Bulletin of the American Mathematical Society*, 43(4):439–561, 2006. 2, 14
- [HS14] Harald A. Helfgott and Akos Seress. On the diameter of permutation groups. *Annals of mathematics*, 179(2):611–658, 2014. 16
- [HSZ15] Harald A. Helfgott, Ákos Seress, and Andrzej Zuk. Random generators of the symmetric group: diameter, mixing time and spectral gap. *Journal of Algebra*, 421:349–368, 2015. 15
- [Jor95] Camille Jordan. Nouvelles recherches sur la limite de transitivité des groupes qui ne contiennent pas le groupe alterné. *J. de Math. Pures et Appliquées*, 1:35–60, 1895. 15
- [Kas07] Martin Kassabov. Symmetric groups and expander graphs. *Inventiones mathematicae*, 170(2):327–354, 2007. 15
- [LS08] Michael Larsen and Aner Shalev. Characters of symmetric groups: sharp bounds and applications. *Inventiones mathematicae*, 174(3):645, 2008. 1, 4, 6

- [Lub12] Alexander Lubotzky. Expander graphs in pure and applied mathematics. *Bulletin of the American Mathematical Society*, 49(1):113–162, 2012. 2, 5, 15
- [Mac95] I.G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford U., 1995. 6
- [SC04] Laurent Saloff-Coste. Random walks on finite groups. In *Probability on discrete structures*. Springer, 2004. 16

ORI PARZANCHEVSKI, EINSTEIN SCHOOL OF MATHEMATICS, THE HEBREW UNIVERSITY
 parzan@math.huji.ac.il

DORON PUDER, SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY
 doronpuder@gmail.com