

The freezing Rènyi quantum discord

Xiao-Yu Li ^{1,*}, Qin-Sheng Zhu ^{2,†}, Ming-Zheng Zhu ², Hao Wu ¹, Shao-Yi Wu ¹, and Ming-Chuan Zhu ¹

¹ *School of information and software engineering,*

University of Electronic Science and Technology of China, Chengdu, 610054, P.R.China and

² *School of Physical, University of Electronic Science and Technology of China, Chengdu, 610054, P.R.China*

(Dated: March 9, 2024)

As a universal quantum character of quantum correlation, the freezing phenomenon is researched by geometry and quantum discord methods, respectively. In this paper, the properties of Rènyi discord is studied for two independent Dimer System coupled to two correlated Fermi-spin environments under the non-Markovian condition. We further demonstrate that the freezing behaviors still exist for Rènyi discord and study the effects of different parameters on this behaviors.

PACS numbers: 03.65.Yz, 03.65.Ud, 02.30.Yy, 03.67.-a.

Key words: Rènyi discord; the freezing phenomenon; quantum correlation.

I. INTRODUCTION

As an important part of the quantum theory, the quantum correlation has aroused extensive attention in lots of physical fields, such as quantum information[1-3], condensed matter physics [4-5] and gravitation wave [6] due to some unimaginable properties in a composite quantum system which can not be reproduced by a classical system. In the past twenty years, entanglement was considered as the quantum correlation and gradually understood. But, the quantum discord concept has been put forward by Ollivier and Zurek [7] and Henderson and Vedral [8] with the deep understanding of quantum correlation. It was clearly demonstrated that entanglement represents only a portion of the quantum correlations and can entirely cover the latter only for a global pure state [9]. Later, many efforts have been devoted to quantify quantum correlation from the view of geometry [9-15] and entropy [7-8,16-19].

Since the systematic correlation contains two parts: the classical correlations and quantum correlation, Maziero et al. [20] found the frozen behavior of the classical correlations for phase-flip, bit-flip, and bit-phase flip channels. As for the possible similar behaviors for the quantum correlation, Mazzola, Piilo, and Maniscalco [21] displayed the similar behavior of the quantum correlations under the nondissipative-independent-Markovian reservoirs for special choices of the initial state. In the same year, Lang and Caves [22] provided a complete geometry picture of the freezing discord phenomenon for Bell-diagonal states. Later, some effort has been devoted to discuss the condition for the frozen-discord with some Non-Markovian processes and initial states[9,14-15,23] (Bell-diagonal states, X states and SCI atates). In conclusion, the freezing discord shows a robust feature of a family of two-qubit models subject to nondissipative decoherence. Although different measures of discords lead to some different conditions for the freezing phenomenon,

this phenomenon of quantum correlation reflects a deeper physical interpretation, such as some relationship with quantum phase transition [24].

Recently, the Rènyi entropy

$$S_\alpha(\rho) = \frac{1}{1-\alpha} \log \text{Tr}[\rho^\alpha] \quad (1)$$

arouses much attention because it is easier to implement in the experiment than the von Neumann entropy which needs the tool of tomography. Here the parameter $\alpha \in (0, 1) \cup (1, \infty)$ and the logarithm is in base 2. Notably, the Rènyi entropy will reduce to the von Neumann entropy when $\alpha \rightarrow 1$. As a natural extension of quantum discord, the Rènyi entropy discord (RED)[25] is also put forward. Therefore, it is valuable to study the properties of RED and the condition for the freezing phenomenon of RED in quantum information field.

II. THE QUANTUM CORRELATION OF DIMER SYSTEM

A. The definition of Rènyi discord

At first, Ollivier and Zurek [7] gave the concept of quantum discord (QD)

$$D(\rho_{AB}) = \min_{\Pi_k^A} \sum_k p_k S(\rho_k^B) + S(\rho_B) - S(\rho_{AB}) \quad (2)$$

to quantify the quantum correlation, where the von Neumann entropy $S(X) = -\text{tr}(\rho_X \log_2 \rho_X)$ is for the density operator ρ_X of system X , $\rho_{A(B)} = \text{Tr}_{B(A)}(\rho_{AB})$ is the reduced density matrix by tracing out the degree of the system $B(A)$, $p_k = \text{Tr}((\Pi_k^A)^\dagger \rho_{AB} \Pi_k^A)$ and $\rho_k^B = \text{Tr}_A((\Pi_k^A)^\dagger \rho_{AB} (\Pi_k^A)) / p_k$.

Later, an equivalent description is introduced in ref [25-26]. The main idea of this equivalent description is to apply an isometry extension of the measurement map $U_{A \rightarrow EX}$ from A to a composite system EX . This method reveals that any channel from A to A' can be used to describe the composite system EX when we discard the

* xiaoyu33521@163.com

† zhuqinsheng@gmail.com

freedom of E . Finally, the quantum discord is rewritten as:

$$D(\rho_{AB}) = \inf_{\Pi_k^A} I(E; B | X)_{\tau_{XEB}} \quad (3)$$

where the optimization is with respect to all possible POVMs Π_k^A of system A with the classical output X. E is an environment for the measurement map and

$$\begin{aligned} \tau_{XEB} &= U_{A \rightarrow EX} \rho_{AB} U_{A \rightarrow EX}^\dagger \\ U_{A \rightarrow EX} |\psi_A\rangle &= \sum_k |k\rangle_X \otimes (\sqrt{\Pi_k^A} |\psi_A\rangle \otimes |k\rangle_E) \end{aligned} \quad (4)$$

The conditional mutual information $I(E; B | X)_{\tau_{XEB}}$ satisfy:

$$\begin{aligned} I(E; B | X)_{\tau_{XBE}} &= S(EX)_{\tau_{XEB}} + S(BX)_{\tau_{XEB}} \\ &\quad - S(X)_{\tau_{XEB}} - S(EBX)_{\tau_{XEB}} \end{aligned} \quad (5)$$

As an extension of quantum discord, the Rènyi quantum discord of ρ_{AB} is defined for $\alpha \in (0, 1) \cup (1, 2]$ as [25]

$$D_\alpha(\rho_{AB}) = \inf_{\Pi_k^A} I_\alpha(E; B | X)_{\tau_{XEB}} \quad (6)$$

where the Rènyi conditional mutual information $I_\alpha(E; B | X)_{\tau_{XEB}}$ satisfy:

$$\begin{aligned} I_\alpha(E; B | X)_{\tau_{XBE}} &= \frac{\alpha}{\alpha - 1} \log \text{Tr} \{ (\rho_X^{\frac{\alpha-1}{2}} \text{Tr}_E \{ \rho_{EX}^{\frac{1-\alpha}{2}} \rho_{EBX}^\alpha \\ &\quad \rho_{EX}^{\frac{1-\alpha}{2}} \} \rho_X^{\frac{\alpha-1}{2}})^\frac{1}{\alpha} \} \end{aligned} \quad (7)$$

The properties of the Rènyi quantum discord are shown in Table 2 of Ref.[25].

B. The Hamiltonian of the open system

We consider two independent dimer systems which are coupled to two correlated Fermi-spin environments, respectively, as shown in Fig.1. The Hamiltonian of the total system has the following form [27]:

$$H = H_d + \sum_{i=1,2} H_{B_i} + \sum_{i,j=1,2} H_{d_i B_j} + q S_1^z S_2^z \quad (8)$$

where $H_d = H_{d_1} + H_{d_2}$ and H_{B_i} describe two independent dimer system and Fermi-spin environments, respectively. $H_{d_i B_j}$ and $S_1^z S_2^z$ denote the interaction between dimer system and that between environments and the environments, respectively. The various parts of the Hamiltonian can be written as following forms:

$$\begin{aligned} H_{d_1} &= \varepsilon_1 |1\rangle\langle 1| + \varepsilon_2 |2\rangle\langle 2| + J_1 (|1\rangle\langle 2| + |2\rangle\langle 1|) \\ H_{d_2} &= \varepsilon_3 |3\rangle\langle 3| + \varepsilon_4 |4\rangle\langle 4| + J_2 (|3\rangle\langle 4| + |4\rangle\langle 3|) \\ H_{d_1 B_1} &= \gamma_1 |1\rangle\langle 1| S_1^z; H_{d_1 B_2} = \gamma_2 |2\rangle\langle 2| S_2^z \\ H_{d_2 B_1} &= \gamma_3 |3\rangle\langle 3| S_1^z; H_{d_2 B_2} = \gamma_4 |4\rangle\langle 4| S_2^z \\ H_{B_i} &= \alpha_i S_i^z \end{aligned}$$




FIG. 1. (Color online) The two dimer systems interacting with two interaction spin-environments (S_1 and S_2). The red and blue balls denote the spin particles of environments (B_1, B_2), respectively. The $|1\rangle$ and $|2\rangle$ ($|3\rangle$ and $|4\rangle$) denote the energy level states of dimer system H_{d_1} (H_{d_2}). q denotes the interaction strength between the spin-environments S_1 and S_2 . The arrows between the energy level states and spin particles of environments denote the interaction between dimer systems and environments.

Here, each environment B_i consists of N_i particles ($i = 1, 2$) with spin $\frac{1}{2}$; ε_a and $|a\rangle$ ($a = 1, 2, 3, 4$) are the energy levels and the energy states of the dimer system, J_1 and J_2 are the amplitudes of transition. The collective spin operators are defined as $S_i^z = \sum_{k=1}^{N_i} \frac{\sigma_z^{k,i}}{2}$, where $\sigma_z^{k,i}$ are the Pauli matrices and α_i is the frequency of $\sigma_z^{k,i}$. So $q S_1^z S_2^z$ describes an Ising-type correlation between the environments with strength q . The cases $q = 0$ and $q \neq 0$, describe independent and correlated spin bath, respectively.

C. The dynamics evolution of the dimer system

The formal solution of the von Neumann equation ($\hbar = 1$)

$$\frac{d}{dt} \rho(t) = \mathcal{L} \rho(t) = -i[H, \rho(t)] \quad (9)$$

can be solved as

$$\rho(t) = e^{\mathcal{L}t} \rho(0) \quad (10)$$

where $\rho(t)$ denotes the density matrix of the total system.

The dynamics of the reduced density matrix $\rho_d(t)$ is obtained by the partial trace method which discards the freedom of the environments. That is

$$\rho_d(t) = \text{Tr}_B(e^{\mathcal{L}t} \rho(0)) \quad (11)$$

Here, the states $|j, m\rangle$ denote the orthogonal bases in the environment Hilbert space H_B which satisfy [28]:

$$\begin{aligned} S^2 |j, m\rangle &= j(j+1) |j, m\rangle; \\ S^z |j, m\rangle &= m |j, m\rangle; S^2 = (S^x)^2 + (S^y)^2 + (S^z)^2 \\ j &= 0, \dots, \frac{N}{2}; m = j, \dots, -j \end{aligned}$$

For the initial state $\rho(0) = \rho_d(0) \otimes \rho_B(0)$ condition, the reduced density matrices $\rho_d(t)$ of the dimer system is

$$\rho_d(t) = \frac{1}{Z} \sum_{j_1=0}^{N_1/2} \sum_{m_1=-j_1}^{j_1} \sum_{j_2=0}^{N_2/2} \sum_{m_2=-j_2}^{j_2} \frac{\nu(N_1, j_1) \nu(N_2, j_2)}{e^{\beta q m_1 m_2} e^{\beta \alpha_1 m_1} e^{\beta \alpha_2 m_2}} \times A^\dagger U^\dagger \rho'_d(0) U A \quad (12)$$

where $\nu(N_i, j_i)$ denotes the degeneracy of the spin bath[28-29]. $\rho'_d(0)$ is the matrix form of the density operator $\rho_d(0)$ under the basis states of dimer system Hilbert space $A^\dagger = (|3\rangle\langle 1| |3\rangle\langle 2| |4\rangle\langle 1| |4\rangle\langle 2|)$. The symbol U in Eq.(12) denotes the 4×4 matrix and equal to MBQ (here, M , B and Q are also 4×4 matrices [27]).

In order to obtain Eq.(12), the environment is given as the canonical distribution

$$\rho_B(0) = \frac{1}{Z} e^{\beta q S_1^z S_2^z} \prod_{i=1}^2 e^{-\beta \alpha_i S_i^z}$$

with $\beta = \frac{1}{K_B T}$ (T is temperature and K_B is Boltzmann constant). The partition function Z is

$$Z = \sum_{j_1=0}^{N_1/2} \sum_{m_1=-j_1}^{j_1} \sum_{j_2=0}^{N_2/2} \sum_{m_2=-j_2}^{j_2} \frac{\nu(N_1, j_1) \nu(N_2, j_2)}{e^{\beta q m_1 m_2} e^{\beta \alpha_1 m_1} e^{\beta \alpha_2 m_2}}$$

D. The properties of Rènyi discord

In this section, the changing behaviors of the quantum correlation are discussed for the two-qubit X [9] and special canonical initial (SCI)[23] under different parameters, respectively. The two-qubit X state is widely used in condensed matter systems and quantum dynamics[9-10,19,30-31]. Under the basis vectors $|00\rangle$, $|01\rangle$, $|10\rangle$ and $|11\rangle$ (here 0(1) denotes the spin up (down) state), the density matrix of a two-qubit X state can be written as

$$\rho_d(0) = \begin{bmatrix} a & 0 & 0 & \delta \\ 0 & b & \beta & 0 \\ 0 & \beta^* & c & 0 \\ \delta^* & 0 & 0 & d \end{bmatrix} \quad (13)$$

satisfying $a, b, c, d \geq 0, a + b + c + d = 1, \|\delta\|^2 \leq ad$ and $\|\beta\|^2 \leq bc$.

Unlike X states, the class of canonical initial (CI)states [23] have the density matrix

$$\rho_d(0) = \frac{1}{4} \begin{bmatrix} 1 + C_{33} & C_{01} & C_{10} & C_{11} - C_{22} \\ C_{01}^* & 1 - C_{33} & C_{11} + C_{22} & C_{10} \\ C_{10}^* & C_{11} + C_{22} & 1 - C_{33} & C_{01} \\ C_{11} - C_{22} & C_{10}^* & C_{01}^* & 1 + C_{33} \end{bmatrix} \quad (14)$$

and the SCI states satisfy:

$$\begin{cases} C_{22}/C_{33} = -C_{11} \\ C_{10}/C_{01} = C_{11} \\ (C_{33})^2 + (C_{01})^2 \leq 1 \end{cases} \quad (15)$$




FIG. 2. (Color online) The properties of Rènyi discord as a function of time t for X and SCI initial states. The parameters are $\alpha_1 = 250ps^{-1}$, $\alpha_2 = 200ps^{-1}$, $\Delta_1 = 20ps^{-1}$, $\Delta_2 = 10ps^{-1}$, $\Delta_3 = 22ps^{-1}$, $\Delta_4 = 12ps^{-1}$, $q = 30ps^{-1}$, $\beta = 1/77$, $N_1 = 20$, $N_2 = 22$, $\gamma_1 = 1ps^{-1}$, $\gamma_2 = 1.1ps^{-1}$, $\gamma_3 = 0.9ps^{-1}$, $\gamma_4 = 1.2ps^{-1}$, $J_1 = 10ps^{-1}$, $J_2 = 12ps^{-1}$ and $\alpha = 0.9$.

In view of the freezing phenomenon for X (SCI) states [9-10,14,19,23,30-31] by geometry and von Neumann entropy discords, X and SCI initial states are chosen here.

In Fig.2, the changing behaviors of quantum correlation are shown for X (Fig.2(a)) and SCI(Fig.2(b)) initial states. With the time evolution, the quantum correlation displays the non-Markov behaviors, especially the peak at about 3 second for some initial states. It indicates the feedback of quantum information (quantum correlation) with the non-Markov process. There are the freezing phenomena for some initial states, with the purple and red solid lines denoting the freezing phenomena for two initial states, respectively. It also hints that the freezing phenomenon of quantum correlation is a universal quantum character and has a deep physical meaning. In the




FIG. 3. (Color online) The changes of freezing Rényi discord based on X (Fig.3(a)) and SCI (Fig.3(b)) initial states (blue solid line of Fig.(1)) with time t for different environment coupling parameters q . The other parameters are same in Figure 2.

inset box, it shows the partially enlarged drawing of the max freezing quantum correlation. Simultaneously, the blue solid line shows the spring of the quantum correlation at $t = 0$. It means that we can generate the quantum correlation by the environments of this quantum system.

Although the general results are obtained, the intrinsic parameters may play an important role in the changing behaviors of the quantum correlation [14-15,17,32], especially the freezing behavior. In Fig.3, the environment coupling parameter q can strongly affect the occurrence of freezing phenomenon, and the quantum correlation appears the quasi-periodic oscillations for X (Fig.3(a)) and SCI (Fig.3(b)) initial states. However, the oscillation behaviors are depressed with $q \geq 50$ for X and SCI initial states. The freezing phenomena of quantum correlation




FIG. 4. (Color online) The changes of freezing Rényi discord based on X (Fig.4(a)) and SCI (Fig.4(b)) initial states (blue solid line of Fig.(1)) with time t for different environment temperatures T . The other parameters are same in Figure 2.

also spring up with increasing q for X state. But, the freezing phenomena of SCI state first appears with increasing q , and then disappears with $q \geq 70$. Particularly, the freezing phenomena always exist throughout the time when q excesses 90. Furthermore, the value of quantum correlation increases with increasing q for X state. From the perspective of the non-Markovian dynamical process, the larger q means the more information flowing from the system into the environment than that from the environment into the system. Therefore, a reasonable value of q is important for the maintenance of quantum correlation.

Except for the parameter q , the temperature T is also important for the quantum correlation. According to our previous works [24,25], a higher temperature may depress the activity of quantum correlation. How does temperature affect the frozen platform? The effect of temper-




FIG. 5. (Color online) The changes of freezing Rényi discord based on X (Fig.4(a)) and SCI (Fig.4(b)) initial states (blue solid line of Fig.(1)) with time t for different parameter α . The other parameters are the same as Figure 2.

ature on the frozen platform is shown in Fig.4. With increasing temperature T , the frozen platform collapses and then reappears at $T \geq 150$. Simultaneously, the platform height is reduced.

Fig. 5 display the effect of different parameters α which is an important parameter of Rényi entropy for Rényi discord. With the increase of α , the monotonicity of Rényi discord is well displayed [25]. For X and SCI initial states, the freezing platform appears in the range of $\alpha \in [0.7, 1.4]$ and $\alpha \in [0.1, 1.6]$, respectively. When the freezing platform appears, there is a particular scope of parameter alpha, which depends on the initial states. Finally, compared the black line ($\alpha = 1$) with others, the quantum discord only shows part of the nature of quantum correlation which quantifies by one of entropy discord, while the others correspond to different entropy discord. This otherness maybe supply the help to discuss the difference between quantum discord and geometric discord, especially, the occurrence of freezing phenomenon has different conditions.

III. CONCLUSION

In this paper, the changing properties of Rényi discord are shown for two independent Dimer System coupled to two correlated Fermi-spin environments. Three main results are presented: 1) the freezing behaviors still exist for Rényi discord which is not just a mathematical coincidence. 2) the freezing platform depends on the parameter α under the same conditions due to the divergence and nonlinearity properties of Rényi entropy. 3) for larger parameter q and lower temperature T , the collapse of the freezing quantum correlation is depressed. These results would supply help to the measurement of quantum correlation and the research of quantum information.

-
- [1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*, (Cambridge University Press, Cambridge, England, 2000).
 - [2] J. Sh. Xu, X. Y. Xu, Ch. F. Li, Ch. J. Zhang, X. B. Zou and G. C. Guo, Nature Commun. **1**, (2010)7.
 - [3] J.S. Xu, K. Sun, C.F. Li, X.Y. Xu, G.C. Guo, E. Andersson, R. Lo Franco, G. Compagno, Nature Commun. **4**, (2013)2851.
 - [4] Y. Yao, H.W. Li, C.M. Zhang, Z.Q. Yin, W. Chen, G.C. Guo, Z.F. Han, Phys. Rev. A **86** (2012) 042102.
 - [5] K. Modi, A. Brodutch, H. Cable, T. Paterek, V. Vedral, Rev. Modern Phys. **84**, (2012)1655.
 - [6] Dmitriy Podolskiy and Robert Lanza, Annalen der physik, **528**,(2016)663.
 - [7] W.H. Zurek, Ann. Phys. (Leipzig) **9**, (2000)855; H. Ollivier, W.H. Zurek, Phys. Rev. Lett. **88**, (2001)017901.
 - [8] L. Henderson, V. Vedral, J. Phys. A **34**, (2001)6899.
 - [9] M. Cianciaruso, T.R. Bromley, W. Roga, R. Lo Franco, G. Adesso, Sci. Rep. **5**, (2015)10177.
 - [10] B. Dakic, V. Vedral, C. Brukner, Phys. Rev. Lett. **105** (2010)190502; S. M. Giampaolo, A. Streltsov, W. Roga, D. Bruß and F. Illuminati, Phys. Rev. A **87**, 012313 (2013).
 - [11] W. K. Wootters, Phys. Rev. D **23**, (1980)357.
 - [12] D. Girolami, A. M. Souza, V. Giovannetti, T. Tufarelli, J. G. Filgueiras, R. S. Sarthour, D. O. Soares-Pinto, I. S. Oliveira, G. Adesso, Phys. Rev. Lett. **112**, 210401 (2014); M. N. Bera, arXiv: 1405.5357.
 - [13] Benjamin Aaronson, Rosario Lo Franco, Giuseppe Compagno and Gerardo Adesso, New Journal of Physics **15**, 093022(2013); D. Girolami, T. Tufarelli, and G. Adesso, Phys. Rev. Lett. **110**, 240402 (2013); L. Chang and S.

- Luo, Phys. Rev. A **87**, 062303 (2013).
- [14] Chang-Chun Ding, Qin-Sheng Zhu, Shao-Yi Wu, and Wei Lai, Annalen der physik, **529**,(2017)1700014.
 - [15] Qinsheng Zhu, Changchun Ding, Shaoyi Wu, Wei Lai, Physica A **458** (2016)67.
 - [16] L. Henderson and V. Vedral, J. Phys. A: Math. Gen. **34** (2001)6899.
 - [17] Q. S. Zhu, C. C. Ding, S. Y. Wu, and W. Lai, Eur. Phys. J. D **69** (2015)231.
 - [18] S. Luo, Phys. Rev. A **77** 022301 (2008); T. Yu, J.H. Eberly, Quantum Inf. Comput. **7** 459 (2007); Q. Chen, C. Zhang, S. Yu,X.X. Yi, C.H. Oh, Phys. Rev.A **84** 042313 (2011); X.-M. Lu, J.Ma, Z. Xi, X.Wang, Phys. Rev. A **83** 012327(2011); M. Ali, A.R.P. Rau, G. Alber, Phys. Rev. A **81** 042105(2010); M. Ali, A.R.P. Rau, G. Alber, Phys. Rev. A **82** 069902(E)(2010).
 - [19] ROSARIO LO FRANCO, BRUNO BELLOMO,SABRINA MANISCALCO and GIUSEPPE COMPAGNO, International Journal of Modern Physics B, **27**, (2013)1245053.
 - [20] Maziero, J., L. C. Céleri, R. M. Serra, and V. Vedral, Phys. Rev. A **80** 009)044102.
 - [21] Mazzola, L., J. Piilo, and S. Maniscalco, Phys. Rev. Lett. **104** (2010)200401.
 - [22] Lang, M. D., and C. M. Caves, Phys. Rev. Lett. textbf105 (2010)150501.
 - [23] T. Chanda, A. K. Pal, A. Biswas, A. Sen, and U. Sen, Phys. Rev. A **91** (2015)062119.
 - [24] Y. Yao, H.W. Li, C.M. Zhang, Z.Q. Yin, W. Chen, G.C. Guo, Z.F. Han, Phys. Rev. A **86** (2012)042102.
 - [25] Mario Berta, Kaushik P. Seshadreesan, and Mark M. Wilde, Journal of Mathematical Physics **56** (2015)022205; Phys. Rev. A **91** (2015)022333.
 - [26] M. Piani,Phys. Rev. A **86** (2012)034101.
 - [27] Qin-Sheng Zhu, Chuan-Ji Fu, and Wei Lai, Z. Naturforsch. **68a** (2013)272.
 - [28] H.-P. Breuer, D. Burgarth, and F. Petruccione, Phys. Rev. B **70** (2004)45323.
 - [29] J. Wesenberg and K. Molmer, Phys. Rev. A **65** (2002)62304; Y. Hamdouni, M. Fannes, and F. Petruccione, Phys. Rev. B **73** (2006)245323.
 - [30] C. Benedetti, Matteo G.A. Paris, S. Maniscalco, Phys. Rev. A **89** (2014)012114; P. Haikka, S. Maniscalco, <https://arxiv.org/abs/1403.2156>; M. Rossi, C. Benedetti, Matteo G.A. Paris, Int. J. Quantum Inf. **12** (2014)1560003.
 - [31] Y. Huang, Phys. Rev. A **88**, (2013)014302 ; M. Namkung, J. Chang, J. Shin, and Y. Kwon, arXiv: 1404.6329 [quant-ph] (2014); S. Luo, Phys. Rev. A **77** (2008)022301.
 - [32] Q. Sh. Zhu, W. Lai, and D. L. Wu,Z. Nat.forsch. A **67a** (2012)559.