

# Portfolio Optimization under Fast Mean-reverting and Rough Fractional Stochastic Environment

Jean-Pierre Fouque\*

Ruimeng Hu†

April 7, 2024

## Abstract

Fractional stochastic volatility models have been widely used to capture the non-Markovian structure revealed from financial time series of realized volatility. On the other hand, empirical studies have identified scales in stock price volatility: both fast-time scale on the order of days and slow-scale on the order of months. So, it is natural to study the portfolio optimization problem under the effects of dependence behavior which we will model by fractional Brownian motions with Hurst index  $H$ , and in the fast or slow regimes characterized by small parameters  $\epsilon$  or  $\delta$ . For the slowly varying volatility with  $H \in (0, 1)$ , it was shown that the first order correction to the problem value contains two terms of order  $\delta^H$ , one random component and one deterministic function of state processes, while for the fast varying case with  $H > \frac{1}{2}$ , the same form holds at order  $\epsilon^{1-H}$ . This paper is dedicated to the remaining case of a fast-varying rough environment ( $H < \frac{1}{2}$ ) which exhibits a different behavior. We show that, in the expansion, only one deterministic term of order  $\sqrt{\epsilon}$  appears in the first order correction.

**Keywords:** Optimal portfolio, rough stochastic volatility, fractional Ornstein–Uhlenbeck process, martingale distortion, asymptotic optimality.

## 1 Introduction

Portfolio optimization in continuous time was originally studied by Merton [1969, 1971]. In this celebrated work, explicit solutions are provided on how to allocate wealth between risky and risk-less assets, and how to consume wealth so that expected utility is maximized. The model there for the underlying assets is the Black–Scholes–Merton model, and the utility functions are of specific type, for instance, of Constant Relative Risk Aversion (CRRA) type.

In this paper, we study the Merton problem with power utility under a non-Markovian stochastic environment that is fast mean-reverting. Such proposed modeling is supported by recent studies. It is shown that stochastic volatility has rapidly decaying correlations at the origin. A common approach for modeling short-range dependence is by using fractional Brownian motion (fBm) processes with  $H < 1/2$ , see Gatheral et al. [2014]. On the other hand, empirical study in Fouque et al. [2003] shows that, in order to fit well the implied volatility, it is appropriate to consider at least short-time scales volatility. To combine these two features, we let the underlying asset follow a Geometric Brownian motion-like model, and we let its return and volatility be driven by a fast-varying factor  $Y_t^{\epsilon, H}$  characterized as the solution of

$$dY_t^{\epsilon, H} = -\frac{a}{\epsilon} Y_t^{\epsilon, H} dt + \frac{1}{\epsilon^H} dW_t^{(H)}.$$

For a full justification/discussion about this model, we refer to Garnier and Sølna [2017b], where  $Y_t^{\epsilon, H}$  is firstly introduced. Here  $\epsilon \ll 1$  is a small parameter, which makes  $Y_t^{\epsilon, H}$  fast-varying and its mean-reversion time scale proportional to  $\epsilon$ , and,  $W_t^{(H)}$  denotes a fBm with Hurst index  $H \in (0, \frac{1}{2})$ . The solution to the above stochastic differential equation (SDE) is actually a fractional Ornstein–Uhlenbeck process. Some properties regarding  $Y_t^{\epsilon, H}$  are discussed in Section 2.2 as a preparation for the asymptotic derivations in Section 3. For further references on fBm and fOU processes, we refer to Mandelbrot and Van Ness [1968], Cheridito et al. [2003], Coutin [2007], Biagini et al. [2008], Kaarakka and Salminen [2011].

\*Department of Statistics & Applied Probability, University of California, Santa Barbara, CA 93106-3110, [fouque@pstat.ucsb.edu](mailto:fouque@pstat.ucsb.edu). Work supported by NSF grant DMS-1409434 and DMS-1814091.

†Department of Statistics, Columbia University, New York, NY 10027-4690, [rh2937@columbia.edu](mailto:rh2937@columbia.edu). The work was mainly done when RH was a graduate student at the University of California, Santa Barbara.

**Motivation and related literature.** Main features and reasons to consider such a problem setup are the following.

Firstly, the fOU process  $Y_t^{\epsilon, H}$  is Gaussian and it admits a convenient moving-average representation with respect to a well-studied kernel (cf. (2.10)). This maintains analytic tractability and simplifies the derivation of needed estimates. However, the modeling of  $Y_t^{\epsilon, H}$  is not limited to fOU processes, and more general kernels can be considered. For example, an alternative model is analyzed in [Garnier and Sølna, 2017b, Appendix B] for the linear pricing problem.

Secondly, we choose to focus on power utilities and one factor model, which means the return and volatility of the underlying asset are driven by only one process. The reason is mainly due to tractability. In this case, a convenient martingale distortion transformation (MDT) is available to express the problem value and the optimal strategy, even under the non-Markovian modeling of  $Y_t^{\epsilon, H}$ . However, we remark that partial results can still be obtained under general utilities where MDT is not available, following a similar argument as in our previous work Fouque and Hu [2017b, 2018]. The details of this generalization will not be included here.

Thirdly, the fOU process  $Y_t^{\epsilon, H}$  is rough ( $H < \frac{1}{2}$ ) and fast mean-reverting ( $\epsilon$  small). This is the missing case in our previous work Fouque and Hu [2017b, 2018], where the asset allocation problem is studied under a slowly varying fractional stochastic environment (fSE) with  $H \in (0, 1)$ , corresponding to  $\epsilon := 1/\delta$  large, and under a fast varying fSE ( $\epsilon$  small) with  $H > \frac{1}{2}$ . Therefore, this paper completes the full picture of the analysis of the portfolio optimization problem in single-factored fractional stochastic environments.

Fourthly, although it is natural to consider multiscale factor models for risky assets, with a fast factor and a slow factor as in Fouque et al. [2015] in a Markovian framework, the analysis requires more technical details, as the MDT is not available. This will be presented in another paper in preparation (Hu [2018b]).

We now summarize the related existing literature in the following table.

Table 1: Problems, models and expansion results. The acronyms used are: SV = Stochastic Volatility, SE = Stochastic Environment, AO = Asymptotic Optimality.

Paper	Problem	Model	Form of Solution*
Fouque et al. [2011] Garnier and Sølna [2017a] Garnier and Sølna [2016] Garnier and Sølna [2017b]	<b>Linear</b> Option Pricing	SV + Multiscale fSV + Slow ( $H \in (0, 1)$ ) fSV + Fast ( $H > \frac{1}{2}$ ) fSV + Fast ( $H < \frac{1}{2}$ )	$P^{(0)} + \sqrt{\epsilon}P^{(1,0)} + \sqrt{\delta}P^{(0,1)}$ $P^{(0)} + \phi^\delta + \delta^H P^{(1)}$ $P^{(0)} + \phi^\epsilon + \epsilon^{1-H} P^{(1)}$ $P^{(0)} + \sqrt{\epsilon}P^{(1)}$
Fouque et al. [2015] Fouque and Hu [2017a] Hu [2018a] Fouque and Hu [2017b] Fouque and Hu [2018] This paper	<b>Nonlinear</b> Portfolio Optimization	SE + Multiscale SE + Slow SE + Fast fSE + Slow ( $H \in (0, 1)$ ) fSE + Fast ( $H > \frac{1}{2}$ ) fSE + Fast ( $H < \frac{1}{2}$ )	$v^{(0)} + \sqrt{\epsilon}v^{(1,0)} + \sqrt{\delta}v^{(0,1)\dagger}$ AO of a zeroth order strategy AO of a zeroth order strategy $v^{(0)} + \phi^\delta + \delta^H v^{(1)}$ $v^{(0)} + \phi^\epsilon + \epsilon^{1-H} v^{(1)}$ $v^{(0)} + \sqrt{\epsilon}v^{(1)}$

\* We denote by  $h^{(0)}$  the leading order term in the expansion, and by  $h^{(0,1)}$ ,  $h^{(1,0)}$ ,  $h^{(1)}$  the first order corrections, where  $h = P$  for option pricing, and  $h = v$  for problem value of the optimization problem. The notation  $\phi^\delta$  (resp.  $\phi^\epsilon$ ) means a random component of order  $\delta^H$  (resp.  $\epsilon^{1-H}$ ).

† The expansion is heuristic except for the case of power utility and one factor.

**Main results.** In this paper, we focus on one-factor models and we study the effect of a fast time-scale on the optimal allocation problem under power utility. When the problem is Markovian, a PDE approach is preferred, since after a distortion transformation, firstly discovered in Zariphopoulou [1999], the PDE becomes linear where perturbation techniques usually work well. However, in the current setting, the fast factor is driven by a rough fBm and possesses short-range dependence. Nevertheless, the nice MDT is available and gives a representation of the value process as well as the optimal strategy. This was proved by Tehranchi [2004] via a conditional Hölder inequality, and by Frei and Schweizer [2008] via a BSDE approach in the case of exponential utility. Recently, it has been restated in Fouque and Hu [2017b] under the setup (2.1) with a short proof based on a verification argument.

Starting by applying MDT, we obtain the representation of the problem value and the optimal portfolio. We then expand them using the “ergodic property” of  $Y_t^{\epsilon, H}$ , and we deduce approximation results for both quantities. Unlike in the long-range dependent case  $H > \frac{1}{2}$  in Fouque and Hu [2018], here, the first order correction to the value process appear at  $\sqrt{\epsilon}$  and contains only one term, which is a explicit function of the state processes. And, surprisingly, there is no correction term at order  $\sqrt{\epsilon}$  for the optimal strategy. However, we are still able to show that the leading order strategy  $\pi_t^{(0)}$  itself generates the value process up to its first order  $\sqrt{\epsilon}$  correction, which is obtained by the “epsilon-martingale decomposition” method. This approach was firstly introduced in Fouque et al. [2000, 2001] and frequently used in problems with small parameters, especially in non-Markovian settings as in Garnier and Sølna [2017a, 2016, 2017b].

We remark that our expansion is only valid when  $H \in (0, \frac{1}{2})$  where  $Y_t^{\epsilon, H}$  does not appear in the leading order nor in the correction of the value process. Moreover, up to order  $\sqrt{\epsilon}$ , the expansion does not really depend on  $H$ , except through constants  $\overline{D}$  and  $\overline{\lambda}$ , see (3.1)–(3.2). We also observe that the limit  $H \uparrow \frac{1}{2}$  does not commute with the limit  $\epsilon \rightarrow 0$  (see Section 3.5), which makes impossible to recover the results in the Markovian case provided in Fouque et al. [2015].

**Organization of the paper.** In Section 2, we revisit the martingale distortion transformation under general stochastic volatility models. This is derived in the Markovian case in Zariphopoulou [1999], and in non-Markovian settings in Tehranchi [2004], Frei and Schweizer [2008], Fouque and Hu [2017b]. Then the fast mean-reverting rough fractional stochastic environment (RFSE) is precisely described, that is,  $Y_t^{\epsilon, H}$  follows a  $\epsilon$ -scaled fractional Ornstein–Uhlenbeck process. In Sections 3.1 and 3.2, we derive the asymptotic results under this modeling for the value process and optimal portfolio respectively. The asymptotic optimality of the leading order strategy  $\pi^{(0)}$  is proved within all admissible strategies in Section 3.3. Finally, we compare the results with the Markovian case  $H = \frac{1}{2}$ , and we comment on the influence of rough fractional model. We make conclusive remarks in Section 4.

## 2 Problem setup and preliminaries

Denote by  $S_t$  the risky asset price at time  $t$  with both returns and volatility driven by a stochastic factor  $Y_t$ :

$$dS_t = S_t [\mu(Y_t) dt + \sigma(Y_t) dW_t]. \quad (2.1)$$

Here  $Y_t$  is a general stochastic process adapted to the filtration  $\mathcal{G}_t \equiv \sigma(W_s^Y, s \leq t)$  of a Brownian motion  $W^Y$ . The two Brownian motions  $W$  and  $W^Y$  are imperfectly correlated:

$$d\langle W_t, W_t^Y \rangle = \rho dt, \quad |\rho| < 1. \quad (2.2)$$

We define  $(\mathcal{F}_t)$  as the natural filtration generated by the two Brownian motions  $(W_t, W_t^Y)$ , and we shall use  $Y_t$  to model the one factor stochastic environment. Later, in Section 3,  $Y_t$  will be replaced by  $Y_t^{\epsilon, H}$ , which will be a fast mean-reverting fOU process.

To formulate the Merton problem under such a stochastic environment, we introduce the following notations. Denote by  $\pi_t \in \mathcal{F}_t$  the amount of money invested in the risky asset  $S_t$  at time  $t$ , and by  $X_t^\pi$  the corresponding wealth process. The rest,  $X_t^\pi - \pi_t$  is put into the bank account earning a constant interest rate  $r$ . Then, under self-financing, and, without loss of generality, assuming a zero interest rate, the dynamics of  $X_t^\pi$  is given by:

$$dX_t^\pi = \pi_t \mu(Y_t) dt + \pi_t \sigma(Y_t) dW_t.$$

In the Merton problem, the agent aims at finding the optimal allocation  $\pi$  so as to optimize the expected utility of her terminal wealth  $X_T^\pi$ . Mathematically, it consists in identifying the value process  $V_t$  defined by

$$V_t \equiv \operatorname{ess\,sup}_{\pi \in \mathcal{A}_t} \mathbb{E}[U(X_T^\pi) | \mathcal{F}_t], \quad (2.3)$$

and the corresponding optimal strategy  $\pi^*$ . Here  $U(\cdot)$  is a utility function describing the agent's preference. Throughout this paper, we shall work with power utilities, namely

$$U(x) = \frac{x^{1-\gamma}}{1-\gamma}, \quad \gamma > 0, \quad \gamma \neq 1.$$

The set  $\mathcal{A}_t$  is the collections of admissible strategies:

$$\mathcal{A}_t \equiv \{\pi \text{ is } (\mathcal{F}_t)\text{-adapted} : X_s^\pi \text{ stays nonnegative } \forall s \geq t, \text{ given } \mathcal{F}_t\},$$

with the following integrability condition:

$$\sup_{t \in [0, T]} \mathbb{E} \left[ (X_t^\pi)^{2p(1-\gamma)} \right] < +\infty, \text{ for some } p > 1, \quad \text{and} \quad \mathbb{E} \left[ \int_0^T (X_t^\pi)^{-2\gamma} \pi_t^2 \sigma^2(Y_t) dt \right] < \infty.$$

In addition, Assumption 2.1 in Fouque and Hu [2018] is enforced throughout the paper.

In Fouque and Hu [2017b], the representation of value process (2.3) and optimal strategy  $\pi^*$  are given via a *martingale distortion transformation*. The rest of this section is a preparation for the main results presented in the next section. The martingale distortion transformation is reviewed, and the fast mean-reverting RFSE is introduced.

## 2.1 Martingale distortion transformation

The martingale distortion transformation was derived in Tehranchi [2004] with a slightly different utility function, and, recently, stated in Fouque and Hu [2017b] under the same setup as in this paper. Since the results in next section heavily rely on this transformation, for reader's convenience, we re-state it here.

Let  $\tilde{\mathbb{P}}$  be an equivalent probability measure determined by the Radon–Nikodym derivative

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} = \exp \left\{ - \int_0^T a_s dW_s^Y - \frac{1}{2} \int_0^T a_s^2 ds \right\}, \quad (2.4)$$

with  $a_t = -\rho \left( \frac{1-\gamma}{\gamma} \right) \lambda(Y_t)$ , and Sharpe-ratio  $\lambda(\cdot) = \mu(\cdot)/\sigma(\cdot)$ . Then, define the  $\tilde{\mathbb{P}}$ -martingale

$$M_t \equiv \tilde{\mathbb{E}} \left[ e^{\frac{1-\gamma}{2q\gamma} \int_0^T \lambda^2(Y_s) ds} \middle| \mathcal{G}_t \right], \quad (2.5)$$

where the parameter  $q$  is given in terms of the utility's relative risk-aversion  $\gamma$  and the correlation coefficient  $\rho$  by

$$q = \frac{\gamma}{\gamma + (1-\gamma)\rho^2}. \quad (2.6)$$

The martingale  $(M_t)_{t \in [0, T]}$  admits the representation

$$dM_t = M_t \xi_t d\tilde{W}_t^Y. \quad (2.7)$$

**Proposition 2.1** (Martingale Distortion Transformation). *Under model assumptions, the value process  $V_t$  defined in (2.3) is expressed by*

$$V_t = \frac{X_t^{1-\gamma}}{1-\gamma} \left[ \tilde{\mathbb{E}} \left( e^{\frac{1-\gamma}{2q\gamma} \int_t^T \lambda^2(Y_s) ds} \middle| \mathcal{G}_t \right) \right]^q.$$

The expectation  $\tilde{\mathbb{E}}[\cdot]$  is computed with respect to  $\tilde{\mathbb{P}}$  introduced in (2.4), and the parameter  $q$  is given in (2.6).

The optimal strategy  $\pi^*$  is

$$\pi_t^* = \left[ \frac{\lambda(Y_t)}{\gamma\sigma(Y_t)} + \frac{\rho q \xi_t}{\gamma\sigma(Y_t)} \right] X_t, \quad (2.8)$$

where  $\xi_t$  is given by the Martingale Representation Theorem in (2.7).

*Proof.* See [Fouque and Hu, 2017b, Proposition 2.2] for a detailed proof. Discussions about special cases (uncorrelated  $\rho = 0$ , degenerate Sharpe-ratio  $\lambda(y) = \lambda_0$ , etc.) and generalization to multi-asset case can also be found therein.  $\square$

**Remark 2.2.** *The model assumptions consist of the existence of a unique strong solution to (2.1), the coincidence between the filtration generated by  $Y_t$  and  $\mathcal{G}_t$ , the regularity condition on  $\lambda(\cdot)$  and integrability condition on  $\xi_t$ :*

$$\mathbb{E}[e^{c \int_0^T \xi_t^2 dt}] < \infty, \text{ for some constant } c. \quad (2.9)$$

*They are actually the same as in our previous paper Fouque and Hu [2018]. So, for the sake of space, we omit the detailed description here, and refer to Assumption 2.1 and Remark 2.2 therein for further discussion.*

## 2.2 Modeling of the fast mean-reverting RFSE

To accommodate the fast mean-reverting property in the stochastic environment, we introduce the small parameter  $\epsilon$  in the modeling of  $Y_t$ , and we switch to the notation  $Y_t^{\epsilon, H}$  to emphasize this dependence. The other superscript  $H$  comes from the Hurst index of the fractional Brownian motion that drives  $Y_t^{\epsilon, H}$ , see (2.11). Then, in the rest of this paper,  $Y_t^{\epsilon, H}$  plays the role of  $Y_t$  in equation (2.1).

Following Garnier and Sølna [2017b], we define our fast factor  $Y_t^{\epsilon, H}$  by

$$Y_t^{\epsilon, H} = \int_{-\infty}^t \mathcal{K}^\epsilon(t-s) dW_s^Y, \quad \mathcal{K}^\epsilon(t) = \frac{1}{\sqrt{\epsilon}} \mathcal{K}\left(\frac{t}{\epsilon}\right), \quad (2.10)$$

where  $\epsilon \ll 1$  is a small parameter,  $\mathcal{K}(t)$  is a non-negative kernel taking the form

$$\mathcal{K}(t) = \frac{1}{\Gamma(H + \frac{1}{2})} \left[ t^{H-\frac{1}{2}} - a \int_0^t (t-s)^{H-\frac{1}{2}} e^{-as} ds \right],$$

with  $H \in (0, \frac{1}{2})$ ,  $(W_t^Y)_{t \in \mathbb{R}^+}$  is the standard Brownian motion (Bm) that is correlated with  $W_t$  as given in (2.2), and  $(W_t^Y)_{t \in \mathbb{R}^-} := (B_{-t})_{t \in \mathbb{R}^-}$  is another Bm independent of  $(W_t^Y)_{t \in \mathbb{R}^+}$  and  $(W_t)$ . The case  $H = \frac{1}{2}$  recovers the usual Markovian OU process. Such  $Y_t^{\epsilon, H}$  is actually the unique (in distribution) stationary solution to the rescaled fractional Ornstein–Uhlenbeck (fOU) SDE:

$$dY_t^{\epsilon, H} = -\frac{a}{\epsilon} Y_t^{\epsilon, H} dt + \frac{1}{\epsilon^H} dW_t^{(H)}, \quad (2.11)$$

where  $W_t^{(H)}$  is the fractional Brownian motion (fBm) with Hurst index  $H$ . Properties regarding this process (with or without scaling) has been widely studied, for instance, see Cheridito et al. [2003], Garnier and Sølna [2017b]. For the sake of simplicity, here, we only mention results that are related to our derivations in the sequel.

The process  $Y_t^{\epsilon, H}$  is Gaussian with mean zero and (co)variance structure:

$$\begin{aligned} \mathbb{E} \left[ \left( Y_t^{\epsilon, H} \right)^2 \right] &\equiv \sigma_{ou}^2 = \frac{1}{2} a^{-2H} \sin(\pi H)^{-1}, \quad \mathbb{E} \left[ Y_t^{\epsilon, H} Y_{t+s}^{\epsilon, H} \right] = \sigma_{ou}^2 \mathcal{C}_Y \left( \frac{s}{\epsilon} \right), \\ \mathcal{C}_Y(s) &\equiv \frac{2 \sin(\pi H)}{\pi} \int_0^\infty \cos(asx) \frac{x^{1-2H}}{1+x^2} dx. \end{aligned} \quad (2.12)$$

The form of covariance shows that the natural scale of  $Y_t^{\epsilon, H}$  is  $\epsilon$  as desired. The variance of  $Y_t^{\epsilon, H}$  stays invariant, which validates the way  $\mathcal{K}^\epsilon(t)$  is rescaled in (2.10). The kernel  $\mathcal{K}(t)$  is in  $L^2$  with  $\int_0^\infty \mathcal{K}^2(u) du = \sigma_{ou}^2$ , and  $\mathcal{K}(t) \in L^1$ , when  $H < \frac{1}{2}$ .

Denote by  $\overline{\lambda}^2$  the average with respect to the invariant distribution of fOU process  $\mathcal{N}(0, \sigma_{ou}^2)$ :

$$\overline{\lambda}^2 \equiv \int_{\mathbb{R}} \lambda^2(z) \frac{1}{\sqrt{2\pi\sigma_{ou}^2}} e^{-\frac{z^2}{2\sigma_{ou}^2}} dz$$

Then, the following differences between time averages and spacial averages are of importance in the derivations:

$$I_t^\epsilon \equiv \int_0^t \left( \lambda^2(Y_s^{\epsilon,H}) - \bar{\lambda}^2 \right) ds, \quad (2.13)$$

$$\phi_t^\epsilon \equiv \mathbb{E} \left[ \int_t^T \lambda^2(Y_s^{\epsilon,H}) - \bar{\lambda}^2 ds \middle| \mathcal{G}_t \right]. \quad (2.14)$$

By the ergodicity of  $Y_t^{\epsilon,H}$ , these differences are small and of order  $\epsilon^{1-H}$ , see Lemma A.1. More properties and estimates regarding  $Y_t^{\epsilon,H}$  are also stated therein.

Furthermore, our fast factor  $Y_t^{\epsilon,H}$  satisfies the model assumption, and we state it as follows.

**Lemma 2.3.** *Under model assumptions, the fast mean-reverting stationary fractional Ornstein–Uhlenbeck process  $Y_t^{\epsilon,H}$  defined in (2.10) satisfies the integrability assumption (2.9).*

*Proof.* This can be easily checked using the argument in [Fouque and Hu, 2017b, Lemma 3.1] and the fact that  $A^\epsilon(T) \equiv \int_0^T \mathcal{K}^\epsilon(s) ds$  is of order  $\sqrt{\epsilon}$ . Therefore, we omit the details here.  $\square$

### 3 Merton problem under fast-varying RFSE

In this section, we study the Merton problem (2.3) with power utility, when the stochastic environment is modeled by  $Y_t^{\epsilon,H}$  with  $H$  restricted to  $H < \frac{1}{2}$ . By the nature of  $Y_t^{\epsilon,H}$ , this nonlinear problem is non-Markovian. This immediately rules out the Hamilton-Jacobi-Bellman PDE approach, which is usually an efficient tool to analyze and find approximations to control problems with small parameters involved. Nevertheless, Proposition 2.1 is available and we will start with applying it to our problem and then find expansions based on it.

To be specific, we will give approximations of both the value process, denoted by  $V_t^\epsilon$  and the corresponding optimal strategy  $\pi^*$ . This is done by applying Proposition 2.1 with  $Y_t = Y_t^{\epsilon,H}$ , then by expanding the expressions (3.1) based on the properties mentioned in Section 2.2. We also show that the “leading order” strategy alone can produce the given approximation of  $V_t^\epsilon$ . Finally, we compare the results with the Markovian case, and we comment on the effects of taking into account the short-range dependence.

#### 3.1 First order approximation to the value process

Let  $S_t$  be the price of a risky asset in the fast mean-reverting RFSE:

$$dS_t = S_t \left[ \mu(Y_t^{\epsilon,H}) dt + \sigma(Y_t^{\epsilon,H}) dW_t \right],$$

where  $Y_t^{\epsilon,H}$  is the  $\epsilon$ -scaled stationary fOU process with  $H < \frac{1}{2}$ , defined in (2.10). Accordingly, the wealth process  $X_t^\pi$  becomes

$$dX_t^\pi = \pi_t \mu(Y_t^{\epsilon,H}) dt + \pi_t \sigma(Y_t^{\epsilon,H}) dW_t,$$

and the value process is denoted by  $V_t^\epsilon$ :

$$V_t^\epsilon \equiv \operatorname{ess\,sup}_{\pi \in \mathcal{A}_t^\epsilon} \mathbb{E}[U(X_T^\pi) | \mathcal{F}_t].$$

We add the superscript  $\epsilon$  to the problem value  $V_t^\epsilon$  and to the admissible set  $\mathcal{A}_t^\epsilon$  to emphasize the dependence on  $\epsilon$  brought by  $Y_t^{\epsilon,H}$ .

Directly applying Proposition 2.1 with  $Y_t = Y_t^{\epsilon,H}$  gives the following expression for  $V_t^\epsilon$ :

$$V_t^\epsilon = \frac{X_t^{1-\gamma}}{1-\gamma} \left[ \widetilde{\mathbb{E}} \left( e^{\frac{1-\gamma}{2q\gamma} \int_t^T \lambda^2(Y_s^{\epsilon,H}) ds} \middle| \mathcal{G}_t \right) \right]^q. \quad (3.1)$$

**Theorem 3.1.** *In the regime of  $\epsilon$  small, under model assumptions, for fixed  $t \in [0, T]$ ,  $V_t^\epsilon$  takes the form*

$$V_t^\epsilon = Q_t^\epsilon(t, X_t) + o(\sqrt{\epsilon}), \quad (3.2)$$

where

$$Q^\epsilon(t, x) = v^{(0)}(t, x) + \sqrt{\epsilon} \rho \overline{D} v^{(1)}(t, x), \quad (3.3)$$

with  $v^{(0)}$  and  $v^{(1)}$  defined as

$$v^{(0)}(t, x) \equiv \frac{x^{1-\gamma}}{1-\gamma} e^{\frac{1-\gamma}{2\gamma} \overline{\lambda}^2 (T-t)}, \quad \text{and} \quad v^{(1)}(t, x) \equiv \frac{1-\gamma}{\gamma^2} (T-t) x^{1-\gamma} e^{\frac{1-\gamma}{2\gamma} \overline{\lambda}^2 (T-t)},$$

and the coefficient  $\overline{D}$  defined by

$$\overline{D} \equiv \int_0^\infty \left[ \iint_{\mathbb{R}^2} \lambda(\sigma_{ou} z) (\lambda \lambda') (\sigma_{ou} z') p_{C_Y(s)}(z, z') dz dz' \right] \mathcal{K}(s) ds.$$

The function  $p_C(z, z')$  is the pdf of the bivariate normal distribution with mean zero and covariance matrix  $\begin{bmatrix} 1 & C \\ C & 1 \end{bmatrix}$ , and  $C_Y(s)$  is given in (2.12). As usual, the notation  $o(\sqrt{\epsilon})$  denotes an  $\mathcal{F}_t$ -adapted random variable whose order is higher than  $\sqrt{\epsilon}$  in the  $L^p$  sense, for any  $1 \leq p < 2(1-H)$ .

*Proof.* Given the representation (3.1) for  $V^\epsilon$ , the expansion result (3.2)-(3.3) can be obtained by firstly expanding

$$\Psi_t^\epsilon \equiv \widetilde{\mathbb{E}} \left[ e^{\frac{1-\gamma}{2q\gamma} \int_t^T (\lambda^2(Y_s^{\epsilon,H}) - \overline{\lambda}^2) ds} \middle| \mathcal{G}_t \right], \quad (3.4)$$

and then applying Taylor formula to the function  $x^q$ .

Using the fact that  $I_t^\epsilon$  is “small” and Taylor expansion of  $e^x$  in  $x$ , one deduces

$$\begin{aligned} \Psi_t^\epsilon &= \widetilde{\mathbb{E}} \left[ 1 + \frac{1-\gamma}{2q\gamma} \int_t^T (\lambda^2(Y_s^{\epsilon,H}) - \overline{\lambda}^2) ds + R_{[t,T]} \middle| \mathcal{G}_t \right] \\ &= 1 + \frac{1-\gamma}{q\gamma} \widetilde{\mathbb{E}} \left[ \frac{1}{2} \int_t^T (\lambda^2(Y_s^{\epsilon,H}) - \overline{\lambda}^2) ds \middle| \mathcal{G}_t \right] + \widetilde{\mathbb{E}} [R_{[t,T]} | \mathcal{G}_t], \end{aligned} \quad (3.5)$$

where  $R_{[t,T]} = e^\chi \left[ \frac{1-\gamma}{2q\gamma} \int_t^T (\lambda^2(Y_s^{\epsilon,H}) - \overline{\lambda}^2) ds \right]^2$  with  $\chi$  being the bounded Lagrange remainder. Observing that  $\|\widetilde{\mathbb{E}}[R_{[t,T]} | \mathcal{G}_t]\|_p \sim \|R_{[t,T]}\|_p$  by the conditional Hölder inequality and the boundedness of  $\lambda$ , and that  $\int_0^T (\lambda^2(Y_s^{\epsilon,H}) - \overline{\lambda}^2) ds \sim o(\sqrt{\epsilon})$  in  $L^2$ , we claim that  $\widetilde{\mathbb{E}} [R_{[t,T]} | \mathcal{G}_t]$  is of order higher than  $\sqrt{\epsilon}$  in  $L^p$  for  $1 \leq p < 2(1-H)$ .

Define the  $\widetilde{\mathbb{P}}$ -martingale  $\widehat{\psi}_t^\epsilon$  with its martingale representation by

$$\widehat{\psi}_t^\epsilon \equiv \widetilde{\mathbb{E}} \left[ \int_0^T G(Y_s^{\epsilon,H}) ds \middle| \mathcal{G}_t \right], \quad G(y) = \frac{1}{2} (\lambda^2(y) - \overline{\lambda}^2), \quad d\widehat{\psi}_t^\epsilon = \widehat{\vartheta}_t^\epsilon d\widetilde{W}_t^Y.$$

It remains to find the expansion of  $\widehat{\psi}_t^\epsilon$  up to order  $\sqrt{\epsilon}$ . To condense the notation in the following derivation, we also define

$$\psi_t^\epsilon \equiv \mathbb{E} \left[ \frac{1}{2} \int_0^T (\lambda^2(Y_s^{\epsilon,H}) - \overline{\lambda}^2) ds \middle| \mathcal{G}_t \right], \quad (3.6)$$

$$\vartheta_t^\epsilon \equiv \int_t^T \mathbb{E} [G'(Y_s^{\epsilon,H}) | \mathcal{G}_t] \mathcal{K}^\epsilon(s-t) ds, \quad (3.7)$$

$$\widetilde{\psi}_t^\epsilon \equiv \widetilde{\mathbb{E}} \left[ \frac{1}{2} \int_0^T (\lambda^2(\widetilde{Y}_s^{\epsilon,H}) - \overline{\lambda}^2) ds \middle| \mathcal{G}_t \right], \quad (3.8)$$

$$\widetilde{\vartheta}_t^\epsilon \equiv \int_t^T \widetilde{\mathbb{E}} [G'(\widetilde{Y}_s^{\epsilon,H}) | \mathcal{G}_t] \mathcal{K}^\epsilon(s-t) ds, \quad (3.9)$$



where  $\psi_t^\epsilon$  is a  $\mathbb{P}$ -martingale satisfying  $d\psi_t^\epsilon = \vartheta_t^\epsilon dW_t^Y$ , see Lemma A.1(i) for details. By a similar argument, we have the  $\tilde{\mathbb{P}}$ -martingale  $\tilde{\psi}_t^\epsilon$  admitting the representation  $d\tilde{\psi}_t^\epsilon = \tilde{\vartheta}_t^\epsilon d\tilde{W}_t^Y$ . The difference between  $\vartheta_t^\epsilon$  and  $\tilde{\vartheta}_t^\epsilon$  is of order  $\epsilon$  and this is discussed in Lemma A.2(ii).

With all above preparations, we deduce:

$$\begin{aligned}
\hat{\psi}_t^\epsilon &= \tilde{\mathbb{E}} \left[ \int_0^T G(Y_s^{\epsilon,H}) ds \middle| \mathcal{G}_0 \right] + \int_0^t \hat{\vartheta}_s^\epsilon d\tilde{W}_s^Y \\
&\quad \text{(Taylor expanding } G(y) \text{ at } y = \tilde{Y}_s^{\epsilon,H}) \\
&= \tilde{\mathbb{E}} \left[ \int_0^T G(\tilde{Y}_s^{\epsilon,H}) ds \middle| \mathcal{G}_0 \right] + \tilde{\mathbb{E}} \left[ \int_0^T G'(\tilde{Y}_s^{\epsilon,H})(Y_s^{\epsilon,H} - \tilde{Y}_s^{\epsilon,H}) ds \middle| \mathcal{G}_0 \right] \\
&\quad + \tilde{\mathbb{E}} \left[ \int_0^T G''(\chi_s)(Y_s^{\epsilon,H} - \tilde{Y}_s^{\epsilon,H})^2 ds \middle| \mathcal{G}_0 \right] + \int_0^t \vartheta_s^\epsilon d\tilde{W}_s^Y + \int_0^t (\hat{\vartheta}_s^\epsilon - \vartheta_s^\epsilon) d\tilde{W}_s^Y \\
&\quad (Y_s^{\epsilon,H} - \tilde{Y}_s^{\epsilon,H} \sim O(\sqrt{\epsilon}), \hat{\vartheta}_s^\epsilon - \vartheta_s^\epsilon \sim O(\epsilon) \text{ and the relation between } W^Y \text{ and } \tilde{W}^Y) \\
&= \tilde{\mathbb{E}} \left[ \int_0^T G(\tilde{Y}_s^{\epsilon,H}) ds \middle| \mathcal{G}_0 \right] + \tilde{\mathbb{E}} \left[ \int_0^T G'(\tilde{Y}_s^{\epsilon,H}) \int_0^s \mathcal{K}^\epsilon(s-u) \rho \left( \frac{1-\gamma}{\gamma} \right) \lambda(Y_u^{\epsilon,H}) du ds \middle| \mathcal{G}_0 \right] \\
&\quad + \int_0^t \vartheta_s^\epsilon dW_s^Y - \int_0^t \vartheta_s^\epsilon \rho \left( \frac{1-\gamma}{\gamma} \right) \lambda(Y_s^{\epsilon,H}) ds + \mathcal{O}(\epsilon) \\
&\quad (\tilde{Y}_s^{\epsilon,H} | \mathcal{G}_0 \stackrel{\mathcal{D}}{=} Y_s^{\epsilon,H} | \mathcal{G}_0 \text{ and the definition of } \kappa_t^\epsilon) \\
&= \mathbb{E} \left[ \int_0^T G(Y_s^{\epsilon,H}) ds \middle| \mathcal{G}_0 \right] + \int_0^t \vartheta_s^\epsilon dW_s^Y + \rho \left( \frac{1-\gamma}{\gamma} \right) \tilde{\mathbb{E}} \left[ \int_0^T \int_u^T \tilde{\mathbb{E}}[G'(\tilde{Y}_s^{\epsilon,H}) | \mathcal{G}_u] \mathcal{K}^\epsilon(s-u) ds \lambda(Y_u^{\epsilon,H}) du \middle| \mathcal{G}_0 \right] \\
&\quad - \rho \left( \frac{1-\gamma}{\gamma} \right) (\sqrt{\epsilon} \overline{D}t + \kappa_t^\epsilon) + \mathcal{O}(\epsilon) \\
&\quad \text{(expression of } \psi_t^\epsilon \text{ and } \tilde{\psi}_t^\epsilon, \text{ and } \|\kappa_t^\epsilon\|_2 \sim o(\sqrt{\epsilon})) \\
&= \psi_t^\epsilon + \rho \left( \frac{1-\gamma}{\gamma} \right) \tilde{\mathbb{E}} \left[ \int_0^T \tilde{\vartheta}_u^\epsilon \lambda(Y_u^{\epsilon,H}) du \middle| \mathcal{G}_0 \right] - \rho \left( \frac{1-\gamma}{\gamma} \right) \sqrt{\epsilon} \overline{D}t + o(\sqrt{\epsilon}) \\
&\quad (\vartheta_t^\epsilon - \tilde{\vartheta}_t^\epsilon \sim \mathcal{O}(\epsilon)) \\
&= \psi_t^\epsilon + \rho \left( \frac{1-\gamma}{\gamma} \right) \tilde{\mathbb{E}} \left[ \int_0^T \vartheta_u^\epsilon \lambda(Y_u^{\epsilon,H}) du \middle| \mathcal{G}_0 \right] - \rho \left( \frac{1-\gamma}{\gamma} \right) \sqrt{\epsilon} \overline{D}t + o(\sqrt{\epsilon}) \\
&= \psi_t^\epsilon + \rho \left( \frac{1-\gamma}{\gamma} \right) \sqrt{\epsilon} \overline{D}(T-t) + \rho \left( \frac{1-\gamma}{\gamma} \right) \tilde{\mathbb{E}} \left[ \int_0^T \vartheta_u^\epsilon \lambda(Y_u^{\epsilon,H}) - \sqrt{\epsilon} \overline{D} du \middle| \mathcal{G}_0 \right] + o(\sqrt{\epsilon}) \\
&\quad (\|\kappa_t^\epsilon\|_2 \sim o(\sqrt{\epsilon})) \\
&= \psi_t^\epsilon + \rho \left( \frac{1-\gamma}{\gamma} \right) \sqrt{\epsilon} \overline{D}(T-t) + o(\sqrt{\epsilon}).
\end{aligned}$$

All reasonings are mentioned in the parentheses from line to line and detailed statements can be found in Lemmas A.1–A.3. Subtracting  $\int_0^t G(Y_u^{\epsilon,H}) du$  from both sides of the above expansion, together with (3.5), (2.14) and (3.6), brings

$$\begin{aligned}
\Psi_t^\epsilon &= 1 + \frac{1-\gamma}{q\gamma} \left( \phi_t^\epsilon + \sqrt{\epsilon} \rho \left( \frac{1-\gamma}{\gamma} \right) \overline{D}(T-t) \right) + o(\sqrt{\epsilon}) \\
&= 1 + \frac{1-\gamma}{q\gamma} \sqrt{\epsilon} \rho \left( \frac{1-\gamma}{\gamma} \right) \overline{D}(T-t) + o(\sqrt{\epsilon}).
\end{aligned} \tag{3.10}$$



The last step follows from  $\phi_t^\epsilon \sim \mathcal{O}(\epsilon^{1-H})$  (see Lemma A.1(ii)). Now, Taylor expanding  $x^q$  produces the desired result

$$\begin{aligned} V_t^\epsilon &= \frac{X_t^{1-\gamma}}{1-\gamma} e^{\frac{1-\gamma}{2\gamma} \bar{\lambda}^2 (T-t)} (\Psi_t^\epsilon)^q \\ &= \frac{X_t^{1-\gamma}}{1-\gamma} e^{\frac{1-\gamma}{2\gamma} \bar{\lambda}^2 (T-t)} \left\{ 1 + \sqrt{\epsilon} \rho \left( \frac{1-\gamma}{\gamma} \right)^2 \bar{D}(T-t) \right\} + o(\sqrt{\epsilon}), \end{aligned}$$

where  $o(\sqrt{\epsilon})$  is in  $L^p$  sense, for any  $1 \leq p < 2(1-H)$ .  $\square$

Note that, unlike in the long-range dependent  $H > 1/2$  case studied in Fouque and Hu [2018] where the first order corrections consist of two terms (one random component  $\phi_t^\epsilon$  and one deterministic function in  $(t, X_t)$ ) at order  $\epsilon^{1-H}$ , here, the correction appears at order  $\sqrt{\epsilon}$  and contains only a deterministic function of  $(t, X_t)$ . In other words, except for the constant  $\bar{D}$  and  $\bar{\lambda}$ , neither the fast factor  $Y_t^{\epsilon, H}$  nor the Hurst index  $H$  is visible in the leading order term nor in the first order correction.

### 3.2 First order expansion of the optimal strategy

The optimal portfolio  $\pi^*$  is also of interest, if not the most important quantity. Under the RFSE described by  $Y_t^{\epsilon, H}$ , using the results in Proposition 2.1, the optimal strategy (2.8) takes the form

$$\pi_t^* = \left[ \frac{\lambda(Y_t^{\epsilon, H})}{\gamma \sigma(Y_t^{\epsilon, H})} + \frac{\rho q \xi_t}{\gamma \sigma(Y_t^{\epsilon, H})} \right] X_t. \quad (3.11)$$

The term  $\xi_t$  is defined by the martingale representation of  $\tilde{\mathbb{P}}$ -martingale  $M_t$ , and is in general not known explicitly. This brings extra difficulty when one wants to implement the optimal strategy  $\pi^*$  to attain the problem value. However, at least in the regime of  $\epsilon$  small, we can give the following approximation result for  $\xi$  and  $\pi^*$ .

**Theorem 3.2.** *Under model assumptions, we have the following approximation of the optimal strategy  $\pi_t^*$ :*

$$\left( \mathbb{E} \int_0^T |\pi_t^* - \pi_t^{(0)}|^p dt \right)^{1/p} \sim o(\sqrt{\epsilon}), \quad \forall 1 \leq p < 2(1-H),$$

where  $\pi_t^{(0)}$  is the leading order strategy:

$$\pi_t^{(0)} := \frac{\lambda(Y_t^{\epsilon, H})}{\gamma \sigma(Y_t^{\epsilon, H})} X_t. \quad (3.12)$$

*Proof.* This is done by obtaining the expansion of  $\xi_t$  from its definition (2.7). We rewrite  $M_t$  in terms of  $\Psi_t^\epsilon$  by comparing (2.5) to (3.4),

$$M_t = \Psi_t^\epsilon e^{\frac{1-\gamma}{2q\gamma} \int_0^t \lambda^2(Y_s^{\epsilon, H}) ds} e^{\frac{1-\gamma}{2q\gamma} \bar{\lambda}^2 (T-t)},$$

and then, we use the approximation (3.10) of  $\Psi_t^\epsilon$ . It has been shown that

$$\Psi_t^\epsilon = a_\epsilon(t) + R_\epsilon(t),$$

where  $a_\epsilon(t) = 1 + \frac{1-\gamma}{q\gamma} \sqrt{\epsilon} \rho \left( \frac{1-\gamma}{\gamma} \right) \bar{D}(T-t)$  is of finite variation, and  $R_\epsilon(t)$  is of order  $o(\sqrt{\epsilon})$  in  $L^p$ , for  $1 \leq p < 2(1-H)$ . This ensures that  $\langle R_\epsilon(\cdot), W(\cdot) \rangle_t \sim o(\sqrt{\epsilon})$  in  $L^p$  by the following reasoning. Using  $L^2$  representation theorem, for each  $t \in [0, T]$ , there exists an adapted process  $\beta(\cdot)$ , such that  $\mathbb{E} \int_0^t \beta^2(u) du < \infty$  and  $R_\epsilon(t)$  admits the representation

$$R_\epsilon(t) = \mathbb{E}[R_\epsilon(t)] + \int_0^t \beta(u) dW_u.$$

Since  $|\mathbb{E}[R_\epsilon(t)]| \leq (\mathbb{E}|R_\epsilon(t)|^p)^{1/p} \sim o(\sqrt{\epsilon})$ , we deduce

$$\mathbb{E} \left| \int_0^t \beta(u) dW_u \right|^p = \mathbb{E} |R_\epsilon(t) - \mathbb{E}[R_\epsilon(t)]|^p \leq C(\mathbb{E}|R_\epsilon(t)|^p + \mathbb{E}^p |R_\epsilon(t)|) \sim o(\sqrt{\epsilon}^p),$$

and

$$\mathbb{E} |\langle R_\epsilon, W \rangle_t|^p \leq \mathbb{E} \int_0^t |\beta(u)|^p du \leq C \mathbb{E} \left| \int_0^t \beta(u) dW_u \right|^p \leq o(\sqrt{\epsilon}^p).$$

Now applying Itô's formula to the above expression of  $M_t$ , none of the three terms will contribute to the diffusion part at order  $\sqrt{\epsilon}$ , meaning that  $\int_0^t M_u \xi_u du \sim o(\sqrt{\epsilon})$  in  $L^p$ , for every  $t \in [0, T]$ . By model assumptions,  $M_t$  is bounded, therefore

$$\int_0^t \xi_u du \sim o(\sqrt{\epsilon}) \text{ in } L^p, \quad \forall t \in [0, T].$$

Denote by  $\|\xi\|_p := \left( \mathbb{E} \int_0^T |\xi_t|^p dt \right)^{1/p}$  the  $L^p(\Omega \times [0, T])$  norm, together with the estimates of  $\xi_t$  in Lemma 2.3, that is,  $|\xi| \leq C\sqrt{\epsilon}$  for any  $t \in [0, T]$ , we have the desired approximation (3.12):

$$\begin{aligned} \left( \mathbb{E} \int_0^T |\pi_t^* - \pi_t^{(0)}|^p dt \right)^{1/p} &= \|\pi^* - \pi^{(0)}\|_p \leq C \|\xi X\|_p = C \|\xi\|_{pr} \|X\|_{pq} \\ &\leq C' \left( \|\xi\|_p (\sqrt{\epsilon})^{r-1} \right)^{1/r}, \text{ where } \frac{1}{r} + \frac{1}{q} = 1 \text{ and } r, q > 1 \\ &\sim o(\sqrt{\epsilon}). \end{aligned}$$

□

In the next subsection, we show the asymptotic optimality property of  $\pi^{(0)}$  defined in (3.12). That is, by only implementing  $\pi^{(0)}$ , the agent is able to obtain the first order approximation (3.3) of the optimal value (3.1).

### 3.3 Asymptotic optimality of $\pi_t^{(0)}$

Let  $X_t^{\pi^{(0)}}$  be the wealth process associated to  $\pi_t^{(0)} = \frac{\lambda(Y_t^{\epsilon, H})}{\gamma \sigma(Y_t^{\epsilon, H})} X_t$ :

$$\begin{aligned} dX_t^{\pi^{(0)}} &= \mu(Y_t^{\epsilon, H}) \pi_t^{(0)} dt + \sigma(Y_t^{\epsilon, H}) \pi_t^{(0)} dW_t \\ &= \frac{\lambda^2(Y_t^{\epsilon, H})}{\gamma} X_t^{\pi^{(0)}} dt + \frac{\lambda(Y_t^{\epsilon, H})}{\gamma} X_t^{\pi^{(0)}} dW_t. \end{aligned} \tag{3.13}$$

By the boundedness of  $\lambda(\cdot)$ ,  $X_t^{\pi^{(0)}}$  has  $p^{th}$ -moment for any  $p$ , and this ensures the admissibility of  $\pi^{(0)}$ .

To systematically simplify the notation in the derivation in Proposition 3.3, we introduce the risk-tolerance function  $R(t, x)$  and the differential operator  $D_k$ , as in Fouque et al. [2015], by:

$$R(t, x) \equiv -\frac{v_x^{(0)}(t, x)}{v_{xx}^{(0)}(t, x)} = \frac{x}{\gamma}, \quad \text{and} \quad D_k \equiv R(t, x)^k \partial_x^k, \quad \forall k \in \mathbb{N}^+. \tag{3.14}$$

Then, the wealth process  $X_t^{\pi^{(0)}}$  can be written as

$$dX_t^{\pi^{(0)}} = \lambda^2(Y_t^{\epsilon, H}) R(t, X_t^{\pi^{(0)}}) dt + \lambda(Y_t^{\epsilon, H}) R(t, X_t^{\pi^{(0)}}) dW_t.$$

We also introduce the nonlinear operator  $\mathcal{L}_{t,x}(\lambda)$  by

$$\mathcal{L}_{t,x}(\lambda) \equiv \partial_t + \frac{1}{2} \lambda^2 D_2 + \lambda^2 D_1,$$

and one can check by straightforward calculation that  $v^{(0)}$  satisfies

$$\mathcal{L}_{t,x}(\bar{\lambda})v^{(0)}(t,x) = 0. \quad (3.15)$$

Denote by  $V_t^{\pi^{(0)},\epsilon}$  the corresponding value process

$$V_t^{\pi^{(0)},\epsilon} := \mathbb{E} \left[ U \left( X_T^{\pi^{(0)}} \right) \middle| \mathcal{F}_t \right].$$

In what follows, we aim to find the approximation of  $V_t^{\pi^{(0)},\epsilon}$  in the regime of  $\epsilon$  small. This will be obtained via “epsilon-martingale decomposition”, firstly introduced in Fouque et al. [2000] to solve the linear pricing problem, and later developed in Fouque et al. [2001], Garnier and Sølna [2017a, 2016, 2017b], Fouque and Hu [2017b, 2018]. Roughly speaking, we need to construct an explicit function  $Q_t$  for  $V_t^{\pi^{(0)},\epsilon}$  in the form of a martingale plus something small (non-martingale part), which has the same terminal condition as  $V_t^{\pi^{(0)},\epsilon}$ . Then, this ansatz is indeed the approximation to  $V_t^{\pi^{(0)},\epsilon}$  up to order of the non-martingale part. Detailed explanation can be found in the above references.

**Proposition 3.3.** *Under model assumptions, for fixed  $t \in [0, T)$  and the observed value  $X_t$ ,  $V_t^{\pi^{(0)},\epsilon}$  is approximated by*

$$V_t^{\pi^{(0)},\epsilon} = Q_t^\epsilon(X_t) + o(\sqrt{\epsilon}),$$

where  $Q_t^\epsilon$  is given in (3.3).

The above Proposition combined with Theorem 3.1 immediately gives:

$\pi_t^{(0)}$  is asymptotically optimal within all admissible strategy  $\mathcal{A}_t^\epsilon$  up to order  $\sqrt{\epsilon}$ .

This is because  $V_t^{\pi^{(0)},\epsilon} - V_t^\epsilon$  is of order  $o(\sqrt{\epsilon})$ , which indicates that  $\pi_t^{(0)}$  already generates the leading order term plus the correction of order  $\sqrt{\epsilon}$  given by (3.3).

*Proof of Proposition 3.3.* Based on the epsilon-martingale decomposition approach, it is enough to find a decomposition  $M_t^\epsilon + R_t^\epsilon$  for  $Q_t^\epsilon$ , with  $M_t^\epsilon$  being a true martingale, and  $R_t^\epsilon$  being of order  $o(\sqrt{\epsilon})$ . We present how  $M_t^\epsilon$  and  $R_t^\epsilon$  are determined, with actual proofs delayed to Appendix A. Note that terms of order  $\sqrt{\epsilon}$  are included in  $M_t^\epsilon$  so that  $R_t^\epsilon$  is pushed to a higher order.

Applying Itô formula to  $v^{(0)}(t, X_t^{\pi^{(0)}})$  brings

$$\begin{aligned} dv^{(0)}(t, X_t^{\pi^{(0)}}) &= \mathcal{L}_{t,x}(\lambda(Y_t^{\epsilon,H}))v^{(0)}(t, X_t^{\pi^{(0)}}) dt + \sigma(Y_t^{\epsilon,H})\pi^{(0)}(t, X_t^{\pi^{(0)}})v_x^{(0)}(t, X_t^{\pi^{(0)}}) dW_t \\ &= \frac{1}{2} \left( \lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2 \right) D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) dt + dM_t^{(1)}, \end{aligned} \quad (3.16)$$

where  $M_t^{(1)}$  is the martingale given by

$$dM_t^{(1)} = \sigma(Y_t^{\epsilon,H})\pi^{(0)}(t, X_t^{\pi^{(0)}})v_x^{(0)}(t, X_t^{\pi^{(0)}}) dW_t, \quad (3.17)$$

and the relations (3.15) and  $D_1 v^{(0)}(t, x) = -D_2 v^{(0)}(t, x)$  have been used.

Recall  $\phi_t^\epsilon$  and  $\psi_t^\epsilon$  defined in (2.14) and (3.6) respectively, then, we have  $d\psi_t^\epsilon - d\phi_t^\epsilon = \frac{1}{2} \left( \lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2 \right) dt$ , and the first term in (3.16) becomes

$$\frac{1}{2} \left( \lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2 \right) D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) dt = D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) (d\psi_t^\epsilon - d\phi_t^\epsilon).$$

To further simplify  $D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) d\phi_t^\epsilon$ , which corresponds to finding the corrector to  $v^{(0)}(t, X_t^{\pi^{(0)}})$  at order  $\sqrt{\epsilon}$ , we compute the total differential of  $D_1 v^{(0)}(t, X_t^{\pi^{(0)}})\phi_t^\epsilon$  (the arguments of  $v^{(0)}(t, X_t^{\pi^{(0)}})$  will be omitted

systematically in the following):

$$\begin{aligned}
d \left( D_1 v^{(0)} \phi_t^\epsilon \right) &= D_1 v^{(0)} d\phi_t^\epsilon + \phi_t^\epsilon \mathcal{L}_{t,x}(\lambda(Y_t^{\epsilon,H})) D_1 v^{(0)} dt + \phi_t^\epsilon \sigma(Y_t^{\epsilon,H}) \pi^{(0)}(t, X_t^{\pi^{(0)}}) \partial_x D_1 v^{(0)} dW_t \\
&\quad + \sigma(Y_t^{\epsilon,H}) \pi^{(0)}(t, X_t^{\pi^{(0)}}) \partial_x D_1 v^{(0)} d \langle W, \phi^\epsilon \rangle_t \\
&= D_1 v^{(0)} d\phi_t^\epsilon + \phi_t^\epsilon \left[ \frac{1}{2} (\lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2) (D_2 + 2D_1) D_1 v^{(0)} \right] dt \\
&\quad + \phi_t^\epsilon \lambda(Y_t^{\epsilon,H}) D_1^2 v^{(0)} dW_t + \rho \lambda(Y_t^{\epsilon,H}) D_1^2 v^{(0)} d \langle W^Y, \psi^\epsilon \rangle_t
\end{aligned}$$

In the derivation, we have used the definition of  $D_1$  and  $R(t, x)$  (cf. (3.14)), and

$$\mathcal{L}_{t,x}(\bar{\lambda}) D_1 v^{(0)} = D_1 \mathcal{L}_{t,x}(\bar{\lambda}) v^{(0)} = 0, \text{ and } d \langle W, \phi^\epsilon \rangle_t = \rho d \langle W^Y, \psi^\epsilon \rangle_t.$$

The results in Lemma A.1(i):  $d \langle W^Y, \psi^\epsilon \rangle_t = \vartheta_t^\epsilon dt$ , together with the above derivation produce

$$\begin{aligned}
d \left( D_1 v^{(0)} \phi_t^\epsilon \right) &= -\frac{1}{2} \left( \lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2 \right) D_1 v^{(0)} dt + \phi_t^\epsilon \left[ \frac{1}{2} (\lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2) (D_2 + 2D_1) D_1 v^{(0)} \right] dt \\
&\quad + \rho \lambda(Y_t^{\epsilon,H}) D_1^2 v^{(0)} \vartheta_t^\epsilon dt + dM_t^{(2)},
\end{aligned} \tag{3.18}$$

and

$$dM_t^{(2)} = D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) d\psi_t^\epsilon + \phi_t^\epsilon \lambda(Y_t^{\epsilon,H}) D_1^2 v^{(0)}(t, X_t^{\pi^{(0)}}) dW_t. \tag{3.19}$$

From Lemma A.1(ii), we see that the second term in (3.18) is of order  $\epsilon^{1-H}$ , and thus it remains to analyze the third term  $\rho \lambda(Y_t^{\epsilon,H}) D_1^2 v^{(0)} \vartheta_t^\epsilon dt$ . To this end, we rewrite  $v^{(1)}$  in terms of  $D_1$  and  $v^{(0)}$  by  $v^{(1)} = D_1^2 v^{(0)}(T-t)$ , and observe that it satisfies  $\mathcal{L}_{t,x}(\bar{\lambda}) v^{(1)} = -D_1^2 v^{(0)}$ . Apply Itô's formula again to  $v^{(1)}$  gives,

$$\begin{aligned}
dv^{(1)}(t, X_t^{\pi^{(0)}}) &= \mathcal{L}_{t,x}(\lambda(Y_t^{\epsilon,H})) v^{(1)}(t, X_t^{\pi^{(0)}}) dt + \sigma(Y_t^{\epsilon,H}) \pi^{(0)}(t, X_t^{\pi^{(0)}}) v_x^{(1)}(t, X_t^{\pi^{(0)}}) dW_t \\
&= \frac{1}{2} (\lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2) (D_2 + 2D_1) v^{(1)}(t, X_t^{\pi^{(0)}}) dt - D_1^2 v^{(0)}(t, X_t^{\pi^{(0)}}) dt + dM_t^{(3)},
\end{aligned} \tag{3.20}$$

where  $M_t^{(3)}$  is the martingale defined by

$$dM_t^{(3)} = \sigma(Y_t^{\epsilon,H}) \pi^{(0)}(t, X_t^{\pi^{(0)}}) v_x^{(1)}(t, X_t^{\pi^{(0)}}) dW_t. \tag{3.21}$$

Defining the quantity  $\tilde{Q}_t^\epsilon$  by

$$\tilde{Q}_t^\epsilon(x) = v^{(0)}(t, x) + D_1 v^{(0)}(t, x) \phi_t^\epsilon + \sqrt{\epsilon} \rho \bar{D} v^{(1)}(t, x),$$

and combining equation (3.16), (3.18) and (3.20) yields

$$\begin{aligned}
d\tilde{Q}_t^\epsilon(X_t^{\pi^{(0)}}) &= d \left( v^{(0)}(t, X_t^{\pi^{(0)}}) + D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) \phi_t^\epsilon + \sqrt{\epsilon} \rho \bar{D} v^{(1)}(t, X_t^{\pi^{(0)}}) \right) \\
&= \phi_t^\epsilon \left[ \frac{1}{2} (\lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2) (D_2 + 2D_1) D_1 v^{(0)} \right] dt + \rho \left( \lambda(Y_t^{\epsilon,H}) \vartheta_t^\epsilon - \sqrt{\epsilon} \bar{D} \right) D_1^2 v^{(0)} dt \\
&\quad + \frac{1}{2} \sqrt{\epsilon} \rho \bar{D} (\lambda^2(Y_t^{\epsilon,H}) - \bar{\lambda}^2) (D_2 + 2D_1) v^{(1)}(t, X_t^{\pi^{(0)}}) dt \\
&\quad + dM_t^{(1)} + dM_t^{(2)} + \sqrt{\epsilon} \rho \bar{D} dM_t^{(3)}.
\end{aligned}$$

Denote by  $R_{t,T}^{(j)}$ ,  $j = 1, 2, 3$  the first three terms in the above expression

$$R_{t,T}^{(1)} := \int_t^T \phi_s^\epsilon \left[ \frac{1}{2} (\lambda^2(Y_s^{\epsilon,H}) - \bar{\lambda}^2) (D_2 + 2D_1) D_1 v^{(0)}(s, X_s^{\pi^{(0)}}) \right] ds, \tag{3.22}$$

$$R_{t,T}^{(2)} := \int_t^T \rho \left( \lambda(Y_s^{\epsilon,H}) \vartheta_s^\epsilon - \sqrt{\epsilon} \bar{D} \right) D_1^2 v^{(0)}(s, X_s^{\pi^{(0)}}) ds,$$

$$R_{t,T}^{(3)} := \int_t^T \frac{1}{2} \sqrt{\epsilon} \rho \bar{D} (\lambda^2(Y_s^{\epsilon,H}) - \bar{\lambda}^2) (D_2 + 2D_1) v^{(1)}(s, X_s^{\pi^{(0)}}) ds. \tag{3.23}$$

It is proved in Lemma A.5 that they are  $o(\sqrt{\epsilon})$  terms in  $L^1$ :

$$\lim_{\epsilon \rightarrow 0} \epsilon^{-1/2} \mathbb{E} \left| R_{t,T}^{(j)} \right| = 0, \quad \forall j = 1, 2, 3.$$

Lemma A.4 also shows that  $M_t^{(j)}$ ,  $j = 1, 2, 3$  are indeed true  $\mathbb{P}$ -martingales.

Therefore, define the martingale  $M_t^\epsilon$  and the non-martingale part  $R_t^\epsilon$  respectively by

$$\begin{aligned} M_t^\epsilon &:= \int_0^t dM_s^{(1)} + dM_s^{(2)} + \sqrt{\epsilon} \rho \overline{D} dM_s^{(3)}, \\ R_T^\epsilon - R_t^\epsilon &:= R_{t,T}^{(1)} + R_{t,T}^{(2)} + R_{t,T}^{(3)}, \end{aligned}$$

and observe that  $\tilde{Q}_T^\epsilon(x) = v^{(0)}(T, x) = U(x)$  (since  $\phi_T^\epsilon = v^{(1)}(T, x) = 0$  by definition), and  $D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) \phi_t^\epsilon$  is of order  $o(\sqrt{\epsilon})$  in  $L^1$  (by Lemma A.1(ii) and integrability of  $D_1 v^{(0)}$ ), we obtain the desired result

$$\begin{aligned} V_t^{\pi^{(0)}, \epsilon} &= \mathbb{E} \left[ \tilde{Q}_T^\epsilon(X_T^{\pi^{(0)}}) | \mathcal{F}_t \right] = \tilde{Q}_t^\epsilon(X_t^{\pi^{(0)}}) + \mathbb{E}[M_T^\epsilon - M_t^\epsilon | \mathcal{F}_t] + \mathbb{E}[R_T^\epsilon - R_t^\epsilon | \mathcal{F}_t] \\ &= Q_t^\epsilon(X_t^{\pi^{(0)}}) + D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) \phi_t^\epsilon + \mathbb{E}[R_{t,T}^{(1)} + R_{t,T}^{(2)} + R_{t,T}^{(3)} + R_{t,T}^{(4)} | \mathcal{F}_t] \\ &= Q_t^\epsilon(X_t^{\pi^{(0)}}) + o(\sqrt{\epsilon}). \end{aligned}$$

□

**Remark 3.4.** *Expansion results of  $V^{\pi^{(0)}, \epsilon}$  can be extended to the case with general utility functions, as in [Fouque and Hu, 2017b, 2018, Section 4]. This is accomplished using the properties of the risk-tolerance function  $R(t, x)$  studied in Fouque and Hu [2017a].*

### 3.4 Numerical Implementation

Numerical implementing  $\pi^{(0)}$  needs to track  $Y_t^{\epsilon, H}$ . By the high oscillation nature of  $Y_t^{\epsilon, H}$ , this usually requires high frequency data and to deal with micro-structure issues. This is not practical for long-period investment and agents usually prefer not to tackle this issue. Instead, they would look into strategies which do not depend on the factor  $Y_t^{\epsilon, H}$ . As shown in Fouque and Hu [2018], such a strategy takes the form of classical Merton optimal strategies under Black-Scholes setting with drift  $\bar{\mu}$  and volatility  $\bar{\sigma} := \sqrt{\langle \sigma^2 \rangle}$ :

$$\bar{\pi}_t^{(0)} = \frac{\bar{\mu}}{\gamma \bar{\sigma}^2} X_t.$$

To measure the utility loss of using  $\bar{\pi}_t^{(0)}$ , we define the associated problem value  $V_t^{\bar{\pi}^{(0)}, \epsilon}$

$$V_t^{\bar{\pi}^{(0)}, \epsilon} = \mathbb{E}[U(X_T^{\bar{\pi}^{(0)}}) | \mathcal{F}_t]. \quad (3.24)$$

Using the ergodic property of  $Y_t^{\epsilon, H}$ :

$$\int_t^T (\mu(Y_s^{\epsilon, H}) - \bar{\mu}) ds \sim o(1), \quad \text{and} \quad \int_t^T (\sigma^2(Y_s^{\epsilon, H}) - \bar{\sigma}^2) ds \sim o(1),$$

one can deduces the optimal leading order term

$$\frac{X_t^{1-\gamma}}{1-\gamma} e^{\frac{1-\gamma}{2\gamma} \frac{\bar{\mu}^2}{\bar{\sigma}^2} (T-t)}.$$

This can be interpreted as the optimal value with Sharpe ratio  $\bar{\mu}/\bar{\sigma}$ . Then, the principal term of utility loss of using  $\bar{\pi}_t^{(0)}$  is quantified by comparing the above term with the leading order term of  $V_t^\epsilon$  given in (3.2)-(3.3):

$$\frac{X_t^{1-\gamma}}{1-\gamma} e^{\frac{1-\gamma}{2\gamma} \bar{\lambda}^2 (T-t)},$$

and is measured by the Cauchy-Schwarz gap

$$\bar{\lambda}^2 = \left\langle \frac{\mu^2}{\sigma^2} \right\rangle \geq \frac{\langle \mu^2 \rangle}{\langle \sigma^2 \rangle} \geq \frac{\bar{\mu}^2}{\bar{\sigma}^2},$$

as in the Markovian setup in Fouque et al. [2015], and in the long-memory fractional setup in Fouque and Hu [2018].

Next, we illustrate numerically the asymptotic optimality property of  $\pi_t^{(0)}$  and the sub-optimality of  $\bar{\pi}_t^{(0)}$ . To this end, we numerically evaluate  $V_t^\epsilon$ ,  $V_t^{\pi^{(0)}, \epsilon}$ , and  $V_t^{\bar{\pi}^{(0)}, \epsilon}$  at time  $t = 0$ , and compare their differences. Applying a change of measure to equation (3.1), we have

$$V_0^\epsilon = \frac{X_0^{1-\gamma}}{1-\gamma} \left[ \mathbb{E} \left( e^{\left(\frac{1-\gamma}{2\gamma}\right) \int_0^T \lambda^2(Y_s^{\epsilon, H}) ds + \rho \left(\frac{1-\gamma}{\gamma}\right) \int_0^T \lambda(Y_s^{\epsilon, H}) dW_s^Y} \middle| \mathcal{G}_0 \right) \right]^q.$$

Solving the SDE for  $X_t^{\pi^{(0)}}$  and plugging the solution into the definition of  $V_t^{\pi^{(0)}, \epsilon}$  bring

$$V_0^{\pi^{(0)}, \epsilon} = \frac{X_0^{1-\gamma}}{1-\gamma} \mathbb{E} \left( e^{\left(\frac{-2\gamma^2+3\gamma-1}{2\gamma^2}\right) \int_0^T \lambda^2(Y_s^{\epsilon, H}) ds + \left(\frac{1-\gamma}{\gamma}\right) \int_0^T \lambda(Y_s^{\epsilon, H}) dW_s} \middle| \mathcal{F}_0 \right).$$

Similarly, the value process defined in (3.24) following the  $Y$ -independent strategy  $\bar{\pi}_t^{(0)}$  is given by

$$V_0^{\bar{\pi}^{(0)}, \epsilon} = \frac{X_0^{1-\gamma}}{1-\gamma} \mathbb{E} \left( e^{\left(\frac{1-\gamma}{\gamma}\right) \frac{\bar{\pi}}{\bar{\sigma}^2} \int_0^T \mu(Y_s^{\epsilon, H}) ds - \left(\frac{1-\gamma}{2\gamma^2}\right) \frac{\bar{\pi}^2}{\bar{\sigma}^4} \int_0^T \sigma^2(Y_s^{\epsilon, H}) ds + \left(\frac{1-\gamma}{\gamma}\right) \frac{\bar{\pi}}{\bar{\sigma}^2} \int_0^T \sigma(Y_s^{\epsilon, H}) dW_s} \middle| \mathcal{F}_0 \right).$$

A full study of the quality of the approximation to optimal strategy involves precise simulations of the fractional OU process  $Y_t^{\epsilon, H}$  and Brownian motion  $W_t$  jointly, and mesh-size determination in  $t$  so that the numerical error is negligible comparing to the loss in value function given by the approximated strategy. This is beyond the scope of this paper, and we therefore compute the values for only a few “omegas” for a purpose of illustration.

The model parameters are chosen as:

$$T = 1, \quad H = 0.1, \quad a = 1, \quad \gamma = 0.4, \quad \rho = -0.5, \quad \mu(y) = \frac{0.1 \times \lambda(y)}{0.1 + \lambda(y)}, \quad \lambda^2(y) = \frac{1}{2} \int_{-\infty}^{y/\sigma_{ou}} p(z/2) dz,$$

where  $p(z)$  denotes the standard normal density. Our choice  $\lambda$  satisfies the standing assumptions in this paper. Also, both  $\mu(y)$  and  $\sigma^2(y) = \mu^2(y)/\lambda^2(y)$  are integrable with respect to the stationary distribution of  $Y^{\epsilon, H}$ , and  $\bar{\mu} = 0.087$  and  $\bar{\sigma}^2 = 0.0176$ .

Notice that  $\mathcal{F}_0$  and  $\mathcal{G}_0$  are not trivial  $\sigma$ -algebra. We first generate a “historical” path  $W_t^Y$  between  $-M$  and 0, and then evaluate each conditional expectation by the average of 500,000 paths. By the short-range dependence of  $Y_t^{\epsilon, H}$ , we  $M = (T/\Delta t)^{0.5}$  (cf. Bardet et al. [2003]), and by its roughness, we choose a fine mesh-size  $\Delta t = 10^{-4}$ . Then, the fast-varying factor  $(Y_t^{\epsilon, H})_{t \in [0, T]}$  (2.10) is generated using the Euler scheme. We mention again that, the numerical results in Table 2 are only for illustration purpose for the reason aforementioned.

Table 2: The value processes  $V_0^\epsilon$  vs.  $V_0^{\pi^{(0)},\epsilon}$  vs.  $V_0^{\bar{\pi}^{(0)},\epsilon}$  for the power utility case.

		#1	#2	#3
$\epsilon = 1$	$V_0^\epsilon$	1.4645	1.4296	1.4075
	$V_0^\epsilon - V_0^{\pi^{(0)},\epsilon}$	0.0021	0.0021	0.0021
	$V_0^\epsilon - V_0^{\bar{\pi}^{(0)},\epsilon}$	0.0538	0.0495	0.0509
$\epsilon = 0.5$	$V_0^\epsilon$	1.4530	1.4328	1.4191
	$V_0^\epsilon - V_0^{\pi^{(0)},\epsilon}$	0.0017	0.0018	0.0017
	$V_0^\epsilon - V_0^{\bar{\pi}^{(0)},\epsilon}$	0.0524	0.0475	0.0473
$\epsilon = 0.1$	$V_0^\epsilon$	1.4442	1.4464	1.4465
	$V_0^\epsilon - V_0^{\pi^{(0)},\epsilon}$	0.0010	0.0010	0.0010
	$V_0^\epsilon - V_0^{\bar{\pi}^{(0)},\epsilon}$	0.0423	0.0405	0.0400
$\epsilon = 0.05$	$V_0^\epsilon$	1.4456	1.4503	1.4522
	$V_0^\epsilon - V_0^{\pi^{(0)},\epsilon}$	0.0006	0.0007	0.0007
	$V_0^\epsilon - V_0^{\bar{\pi}^{(0)},\epsilon}$	0.0369	0.0366	0.0365
$\epsilon = 0.01$	$V_0^\epsilon$	1.4507	1.4541	1.4563
	$V_0^\epsilon - V_0^{\pi^{(0)},\epsilon}$	0.0002	0.0002	0.0003
	$V_0^\epsilon - V_0^{\bar{\pi}^{(0)},\epsilon}$	0.0224	0.0249	0.0254

As expected, utility losses for both approximated strategies tend to decrease as  $\epsilon$  goes to zero. The zeroth order strategy  $\pi^{(0)}$  performs well, even for not so small values of  $\epsilon$ . The relative utility loss is below 0.2%. Again, as expected, the “lazy” strategy  $\bar{\pi}_t^{(0)}$  produces a larger utility loss, and thus underperforms  $\pi^{(0)}$ , but we still consider its performance good after observing that the  $(V_0^\epsilon - V_0^{\bar{\pi}^{(0)},\epsilon})/V_0^\epsilon$  is below 4%.

### 3.5 Comparison with the Markovian case

In the Markovian case, corresponding to  $H = \frac{1}{2}$  in the modeling of  $Y_t^{\epsilon,H}$  (2.10), approximations to the value function and the optimal portfolio have been rigorously derived in Fouque et al. [2015], and are of the form:

$$\begin{aligned}
 V^\epsilon(t, X_t) &= \frac{X_t^{1-\gamma}}{1-\gamma} e^{\frac{1-\gamma}{2\gamma}\bar{\lambda}^2(T-t)} \left[ 1 - \sqrt{\epsilon}\rho \left( \frac{1-\gamma}{\gamma} \right)^2 \frac{\langle \lambda\theta' \rangle}{2}(T-t) \right] + \mathcal{O}(\epsilon), \\
 \pi^*(t, X_t, Y_t^{\epsilon,H}) &= \left[ \frac{\lambda(Y_t^{\epsilon,H})}{\gamma\sigma(Y_t^{\epsilon,H})} + \sqrt{\epsilon} \frac{\rho(1-\gamma)}{\gamma^2\sigma(Y_t^{\epsilon,H})} \frac{\theta'(Y_t^{\epsilon,H})}{2} \right] X_t + \mathcal{O}(\epsilon),
 \end{aligned} \tag{3.25}$$

where  $\theta(y)$  solves the Poisson equation  $\frac{1}{2}\theta''(y) - ay\theta'(y) = \lambda^2(y) - \bar{\lambda}^2$ . These can be viewed as the limits  $\lim_{\epsilon \rightarrow 0} \lim_{H \uparrow \frac{1}{2}}$  of our current setup.

However, the two limits apparently do not commute for the optimal control, since there is no correction term at order  $\sqrt{\epsilon}$  in (3.11). This is also the case for the problem value  $V^\epsilon$ , even the first order correction in (3.3) turns out to be of order  $\sqrt{\epsilon}$ . Formally, letting  $H \uparrow \frac{1}{2}$  in equation (3.3), we obtain

$$V^\epsilon(t, X_t) = \frac{X_t^{1-\gamma}}{1-\gamma} e^{\frac{1-\gamma}{2\gamma}\bar{\lambda}^2(T-t)} \left[ 1 + \sqrt{\epsilon}\rho \left( \frac{1-\gamma}{\gamma} \right)^2 \bar{D}'(T-t) \right] + o(\sqrt{\epsilon})$$

where  $\bar{D}'$  is the limit of  $\bar{D}$  as  $H$  approaches  $\frac{1}{2}$ ,

$$\bar{D}' = \lim_{H \uparrow \frac{1}{2}} \bar{D} = \int_0^\infty \left[ \iint_{\mathbb{R}^2} \lambda(\sigma_{ou}z)(\lambda\lambda')(\sigma_{ou}z') \tilde{p}_{CY(s)}(z, z') \right] e^{-as} ds$$



and  $\tilde{p}_{C_Y(s)}(z, z')$  is the bivariate normal density with mean zero and variance matrix  $\begin{bmatrix} 1 & e^{-as} \\ e^{-as} & 1 \end{bmatrix}$ . This is not the same constant  $\langle \lambda \theta' \rangle$  as in equation (3.25).

Although the expansions of the two cases ( $H = \frac{1}{2}$  vs.  $H \in (0, \frac{1}{2})$ ) surprisingly share the same form (a leading order term plus the first order correction at order  $\sqrt{\epsilon}$ ), the coefficients are not identical. This is because our derivations in Theorem 3.1 and 3.2 are only valid for  $H \in (0, \frac{1}{2})$ , and the singular perturbation is “singular” at  $H = \frac{1}{2}$ . Consequently, the order of limits  $H \uparrow \frac{1}{2}$  and  $\epsilon \rightarrow 0$  is not interchangeable, and this leads to different expansion results. We also remark that, unlike in the case  $H > \frac{1}{2}$  where the Hurst index  $H$  influences the order of first correction ( $\epsilon^{1-H}$ ), here it is  $\sqrt{\epsilon}$  no matter what value  $H$  takes in  $(0, \frac{1}{2})$ .

## 4 Conclusion

In this paper, we treated the portfolio optimization problem in a one-factor stochastic environment when the investor’s utility is of power type. To accommodate recent empirical studies, we model this factor using a fast mean-reverting process driven by a fractional Brownian motion with Hurst index  $H < \frac{1}{2}$ . Thus, its paths are rougher than the standard Bm and it has short-range dependence. Under this setup, the value process can be represented explicitly thanks to the martingale distortion transformation (MDT), which enables us to perform an asymptotic expansion and obtain an approximation of the form: leading order term plus a correction term at order  $\sqrt{\epsilon}$ . Surprisingly, the order of the correction is not associated to the Hurst index  $H$ , and the fast factor  $Y_t^{\epsilon, H}$  appears in neither terms, which is a different behavior than in the cases studied in our previous work Fouque and Hu [2017b, 2018]. The approximation of the optimal strategy is also analyzed, and it turns out that there is no correction at the order  $\sqrt{\epsilon}$ . Nevertheless, we are still able to show that the leading order strategy, derived in Section 3.2, is able to reproduce the problem value up to order  $\sqrt{\epsilon}$ , and therefore, it is asymptotically optimal within all admissible strategies. We remark that this is only proved in the case of power utility, as in general, the MDT is not available either with general utility or with multi-factor models, as well as the expansion of the full problem value. However, one can work within a smaller class of admissible strategies and easily extend the “epsilon-martingale decomposition” argument in Section 3.3 to obtain a weaker optimality of  $\pi^{(0)}$ . For the general utility case, this argument is very similar to our previous work [Fouque and Hu, 2017b, 2018, Section 4], and we did not include it here. The multi-factor case involves more techniques and will be presented in another paper in preparation Hu [2018a].

## A Technical Lemmas

In this section, we present several lemmas used in Section 3. Note that the constants  $K, K'$  in all lemmas do not depend on  $\epsilon$  and may vary from line to line, and we denote the function  $G(y)$  as

$$G(y) = \frac{1}{2}(\lambda^2(y) - \bar{\lambda}^2),$$

and  $\|X\|_p := (\mathbb{E}X^p)^{1/p}$  as the  $L^p$ -norm of  $X$ .

**Lemma A.1.**

(i) The martingale  $\psi_t^\epsilon$  defined in (3.6):

$$\psi_t^\epsilon = \mathbb{E} \left[ \int_0^T G(Y_s^{\epsilon, H}) ds \middle| \mathcal{G}_t \right],$$

satisfies

$$d\psi_t^\epsilon = \vartheta_t^\epsilon dW_t^Y, \quad \vartheta_t^\epsilon := \int_t^T \mathbb{E} [G'(Y_s^{\epsilon, H}) | \mathcal{G}_t] \mathcal{K}^\epsilon(s - t) ds.$$

Moreover, for all  $t \in [0, T]$  we have

$$\mathbb{E}[\lambda(Y_t^{\epsilon, H})\vartheta_t^\epsilon] = \sqrt{\epsilon} \bar{D} + \tilde{D}_t^\epsilon,$$

where  $\overline{D}$  is the deterministic constant and  $\tilde{D}_t^\epsilon$  is of higher order than  $\sqrt{\epsilon}$ :

$$\sup_{\epsilon \in [0,1]} \sup_{t \in [0,T]} \epsilon^{-1/2} \left| \tilde{D}_t^\epsilon \right| < \infty, \quad \text{and} \quad \forall t \in [0, T], \lim_{\epsilon \rightarrow 0} \epsilon^{-1/2} \left| \tilde{D}_t^\epsilon \right| = 0.$$

Define the random process  $\kappa_t^\epsilon$  by

$$\kappa_t^\epsilon = \int_0^t (\vartheta_s^\epsilon \lambda(Y_s^{\epsilon,H}) - \sqrt{\epsilon} \overline{D}) \, ds. \quad (\text{A.1})$$

It is of higher order than  $\sqrt{\epsilon}$  in  $L^2$  sense uniformly in  $t \in [0, T]$ :

$$\lim_{\epsilon \rightarrow 0} \epsilon^{-1/2} \sup_{t \in [0,T]} \|\kappa_t^\epsilon\|_2 = 0.$$

(ii) The random component  $\phi_t^\epsilon$  defined in (2.14) has the form

$$\phi_t^\epsilon = \mathbb{E} \left[ \int_t^T G(Y_s^{\epsilon,H}) \, ds \middle| \mathcal{G}_t \right].$$

It is a random variable with mean zero and standard deviation of order  $\epsilon^{1-H}$  uniformly in  $t \in [0, T]$ :

$$\sup_{\epsilon \in [0,1]} \sup_{t \in [0,T]} \epsilon^{1-H} \|\phi_t^\epsilon\|_2 < \infty.$$

(iii) The random process  $I_t^\epsilon$  defined in (2.13)

$$I_t^\epsilon = \int_0^t \left( \lambda^2(Y_s^{\epsilon,H}) - \overline{\lambda}^2 \right) \, ds,$$

satisfies

$$\lim_{\epsilon \rightarrow 0} \sup_{t \in [0,T]} \mathbb{E}[(I_t^\epsilon)^2] \leq K \epsilon^{1-H}.$$

*Proof.* All results are slightly different versions or straightforward generalizations of lemmas in [Garnier and Sølna, 2017b, Appendix A], thus we omit the details here.  $\square$

**Lemma A.2.**

(i) Denote by  $\tilde{Y}_t^{\epsilon,H}$  the  $\tilde{\mathbb{P}}$ -stationary fractional Ornstein–Uhlenbeck process, whose moving average representation is of the form

$$\tilde{Y}_t^{\epsilon,H} := \int_{-\infty}^t \mathcal{K}^\epsilon(t-s) \, d\tilde{W}_s^Y.$$

Then,  $\sup_{t \in [0,T]} \left| \tilde{Y}_t^{\epsilon,H} - Y_t^{\epsilon,H} \right| \leq K \sqrt{\epsilon}$ .

(ii) Recall the stochastic process  $\vartheta_t^\epsilon$  defined in (3.7), and  $\tilde{\vartheta}_t^\epsilon$  defined in (3.9):

$$\vartheta_t^\epsilon := \int_t^T \mathbb{E}[G'(Y_s^{\epsilon,H}) | \mathcal{G}_t] \mathcal{K}^\epsilon(s-t) \, ds, \quad \tilde{\vartheta}_t^\epsilon := \int_t^T \tilde{\mathbb{E}}[G'(\tilde{Y}_s^{\epsilon,H}) | \mathcal{G}_t] \mathcal{K}^\epsilon(s-t) \, ds.$$

Then,  $\sup_{t \in [0,T]} \left| \tilde{\vartheta}_t^\epsilon - \vartheta_t^\epsilon \right| \leq K \epsilon$ .

*Proof.* Both are proved by using the arguments in [Fouque and Hu, 2018, Lemma A.3] and the fact  $\mathcal{K}(t) \in L^1$ , and we omit the details for simplicity.  $\square$

**Lemma A.3.** Recall the  $\widetilde{\mathbb{P}}$ -martingale  $\widehat{\psi}_t^\epsilon$  defined in (3.8)

$$\widehat{\psi}_t^\epsilon = \widetilde{\mathbb{E}} \left[ \int_0^T G(Y_s^{\epsilon,H}) ds \middle| \mathcal{G}_t \right].$$

Denote its martingale representation by

$$d\widehat{\psi}_t^\epsilon = \widehat{\vartheta}_t^\epsilon d\widetilde{W}_t^Y.$$

Then, the process  $\widehat{\vartheta}_t^\epsilon$  satisfies  $\sup_{t \in [0, T]} |\vartheta_t^\epsilon - \widehat{\vartheta}_t^\epsilon| \leq K\epsilon$ .

*Proof.* We first claim that, for all  $t \in [0, T]$

$$\int_t^T |\widetilde{\mathcal{D}}_t Y_s^{\epsilon,H}| ds \leq K\sqrt{\epsilon}, \quad (\text{A.2})$$

where  $\widetilde{\mathcal{D}}$  denotes the Malliavin derivative with respect to  $\widetilde{W}$ . This is obtained by applying the derivation in [Fouque and Hu, 2017b, Lemma 3.1] and the fact that  $\int_0^T \mathcal{K}^\epsilon(s) ds$  is bounded by  $K\sqrt{\epsilon}$ .

Then,  $\widehat{\vartheta}_t^\epsilon$  is computed as:

$$\begin{aligned} \widehat{\vartheta}_t^\epsilon &= \widetilde{\mathbb{E}} \left[ \int_0^T G'(Y_s^{\epsilon,H}) \widetilde{\mathcal{D}}_t Y_s^{\epsilon,H} ds \middle| \mathcal{G}_t \right] \\ &= \int_t^T \widetilde{\mathbb{E}} [G'(Y_s^{\epsilon,H}) | \mathcal{G}_t] \mathcal{K}^\epsilon(s-t) ds + \widetilde{\mathbb{E}} \left[ \int_t^T G'(Y_s^{\epsilon,H}) \int_t^s \mathcal{K}^\epsilon(s-u) \rho \left( \frac{1-\gamma}{\gamma} \right) \lambda'(Y_u^{\epsilon,H}) \widetilde{\mathcal{D}}_t Y_u^{\epsilon,H} du ds \middle| \mathcal{G}_t \right] \\ &= \widetilde{\vartheta}_t^\epsilon + R^\epsilon = \vartheta_t^\epsilon + (\widetilde{\vartheta}_t^\epsilon - \vartheta_t^\epsilon) + R^\epsilon, \end{aligned}$$

with  $R^\epsilon$  defined as:

$$\begin{aligned} R^\epsilon &:= \int_t^T \widetilde{\mathbb{E}} \left[ G''(\chi_s)(Y_s^{\epsilon,H} - \widetilde{Y}_s^{\epsilon,H}) | \mathcal{G}_t \right] \mathcal{K}^\epsilon(s-t) ds \\ &\quad + \widetilde{\mathbb{E}} \left[ \int_t^T G'(Y_s^{\epsilon,H}) \int_t^s \mathcal{K}^\epsilon(s-u) \rho \left( \frac{1-\gamma}{\gamma} \right) \lambda'(Y_u^{\epsilon,H}) \widetilde{\mathcal{D}}_t Y_u^{\epsilon,H} du ds \middle| \mathcal{G}_t \right], \end{aligned}$$

and  $\chi_s$  is the remainder of Taylor expansion. Given Lemma A.2(ii), it remains to show that  $R^\epsilon$  is bounded by  $K\epsilon$ . The first term in  $R^\epsilon$  is guaranteed by Lemma A.2(i) and  $\mathcal{K}(t) \in L^1$ , while the second term is by the boundedness of  $\lambda$  and its derivatives,  $\mathcal{K}(t) \in L^1$  and (A.2).  $\square$

**Lemma A.4.** The processes  $M_t^{(j)}$ ,  $j = 1, 2, 3$  defined in (3.17), (3.19) and (3.21) are true  $\mathbb{P}$ -martingales.

*Proof.* By the Burkholder–Davis–Gundy inequality, it suffices to show  $\mathbb{E} [\langle M^{(j)} \rangle_T^{1/2}] < \infty$ , for  $j = 1, 2, 3$ .

For the case  $j = 1$ , we compute

$$d \langle M^{(1)} \rangle_t = \lambda^2(Y_t^{\epsilon,H}) \left( D_1 v^{(0)}(t, X_t^{\pi^{(0)}}) \right)^2 dt \leq K \left( X_t^{\pi^{(0)}} \right)^{2-2\gamma} dt,$$

by the boundedness of  $\lambda$ . Then since  $X_t^{\pi^{(0)}}$  has  $p^{th}$ -moment for any  $p$  uniformly in  $t \in [0, T]$ , we obtain

$$\mathbb{E} \left[ \langle M^{(1)} \rangle_T^{1/2} \right] \leq \left[ \mathbb{E} \int_0^T K (X_t^{\pi^{(0)}})^{2-2\gamma} dt \right]^{1/2} \leq K \sup_{t \in [0, T]} \left( \mathbb{E} [(X_t^{\pi^{(0)}})^{2-2\gamma}] \right)^{1/2} < \infty.$$

The martingality of  $M_t^{(3)}$  is obtained in the same way; while for  $M_t^{(2)}$ , additional properties such as the boundedness (uniform in  $\delta$ ) of  $\vartheta_t^\epsilon$  and  $\phi_t^\epsilon$  are used.  $\square$

**Lemma A.5.** The random variable  $R_{t,T}^{(j)}$ ,  $j = 1, 2, 3$  defined in (3.22)-(3.23)

$$\begin{aligned} R_{t,T}^{(1)} &:= \int_t^T \phi_s^\epsilon \left[ \frac{1}{2} (\lambda^2(Y_s^{\epsilon,H}) - \bar{\lambda}^2)(D_2 + 2D_1) D_1 v^{(0)}(s, X_s^{\pi^{(0)}}) \right] ds, \\ R_{t,T}^{(2)} &:= \int_t^T \rho(\lambda(Y_s^{\epsilon,H}) \vartheta_s^\epsilon - \sqrt{\epsilon} \bar{D}) D_1^2 v^{(0)}(s, X_s^{\pi^{(0)}}) ds, \\ R_{t,T}^{(3)} &:= \int_t^T \frac{1}{2} \sqrt{\epsilon} \rho \bar{D} (\lambda^2(Y_s^{\epsilon,H}) - \bar{\lambda}^2)(D_2 + 2D_1) v^{(1)}(s, X_s^{\pi^{(0)}}) ds, \end{aligned}$$

are of order  $o(\sqrt{\epsilon})$ :

$$\lim_{\epsilon \rightarrow 0} \epsilon^{-1/2} \mathbb{E} \left| R_{t,T}^{(j)} \right| = 0, \quad \forall j = 1, 2, 3. \quad (\text{A.3})$$

*Proof.* The proofs here are similar to the ones in [Garnier and Sølna, 2017b, Proposition 4.1].

For the case  $j = 1$ , using the definition of  $v^{(0)}$  and  $D_k$ , the property  $\phi_t^\epsilon \sim \mathcal{O}(\epsilon^{1-H})$  in  $L^2$  (cf. Lemma A.1(ii)), and the boundedness of  $\lambda$ , one deduces

$$\mathbb{E} \left| R_{t,T}^{(1)} \right| \leq K \mathbb{E} \left| \int_t^T \phi_s^\epsilon \left( X_s^{\pi^{(0)}} \right)^{1-\gamma} ds \right| \leq K \sup_{t \in [0, T]} \|\phi_t^\epsilon\|_2 \left( \mathbb{E} \int_t^T \left( X_s^{\pi^{(0)}} \right)^{2-2\gamma} ds \right)^{1/2}.$$

Since  $X_t^{\pi^{(0)}}$  has  $p^{th}$  moment for any  $p$ , the above expectation is of order  $\epsilon^{1-H}$ , and thus (A.3) is satisfied for  $j = 1$ .

To prove (A.3) with  $j = 2$ , we denote  $t_k = t + (T-t)k/N$ ,  $Z_s^{(2)} = D_1^2 v^{(0)}(s, X_s^{\pi^{(0)}}) = K(X_s^{\pi^{(0)}})^{1-\gamma}$  and recall  $\kappa_t^\epsilon$  defined in (A.1), thus  $R_{t,T}^{(2)}$  can be written as

$$\begin{aligned} R_{t,T}^{(2)} &= \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} Z_s^{(2)} \frac{d\kappa_s^\epsilon}{ds} ds = \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} Z_{t_k}^{(2)} \frac{d\kappa_s^\epsilon}{ds} ds + \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} (Z_s^{(2)} - Z_{t_k}^{(2)}) \frac{d\kappa_s^\epsilon}{ds} ds \\ &= \sum_{k=0}^{N-1} Z_{t_k}^{(2)} (\kappa_{t_{k+1}}^\epsilon - \kappa_{t_k}^\epsilon) + \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} (Z_s^{(2)} - Z_{t_k}^{(2)}) \frac{d\kappa_s^\epsilon}{ds} ds \\ &:= R_{t,T}^{(2,a)} + R_{t,T}^{(2,b)}. \end{aligned}$$

We then claim two properties for  $Z_t^{(2)}$ : (a) it has finite second moment uniformly in  $\epsilon$  and  $s \in [0, T]$ , i.e.

$$\sup_{\epsilon \in [0, 1]} \sup_{t \in [0, T]} \mathbb{E}[(Z_t^{(2)})^2] < \infty;$$

and (b) its increments are Lipschitz in  $L^2$ :

$$\mathbb{E}[(Z_u^{(2)} - Z_v^{(2)})^2] \leq K |u - v|. \quad (\text{A.4})$$

These are ensured by the boundedness of  $\lambda(\cdot)$ , and the formulation (3.13) of  $X_t^{\pi^{(0)}}$  in the case of power utility. In general (utility), the two properties regarding  $Z_t^{(2)}$  are also satisfied under proper assumptions, see [Fouque and Hu, 2018, Lemma A.6] for further details.

Now we proceed to the analysis of  $R_{t,T}^{(2,a)}$  and  $R_{t,T}^{(2,b)}$ :

$$\mathbb{E} \left| R_{t,T}^{(2,a)} \right| \leq \sqrt{2} \sum_{k=0}^{N-1} \left\| Z_{t_k}^{(2)} \right\|_2 [\mathbb{E}(\kappa_{t_k}^\epsilon)^2 + \mathbb{E}(\kappa_{t_{k+1}}^\epsilon)^2]^{1/2} \leq 2N \sup_{s \in [t, T]} \left\| Z_s^{(2)} \right\|_2 \sup_{s \in [t, T]} \|\kappa_s^\epsilon\|_2$$

and is of order  $o(\sqrt{\epsilon})$  for any fixed  $N$  by Lemma A.1(i). For the second term, using (A.4) and the fact that  $\frac{d\kappa_s^\epsilon}{ds}$  is of (or higher than) order  $\mathcal{O}(\sqrt{\epsilon})$ , one computes

$$\mathbb{E} \left| R_{t,T}^{(2,b)} \right| \leq K \sqrt{\epsilon} \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} \left\| Z_s^{(2)} - Z_{t_k}^{(2)} \right\|_2 ds \leq K \sqrt{\epsilon} \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} (s - t_k)^{1/2} ds = K \sqrt{\epsilon} \frac{1}{\sqrt{N}},$$

and claims

$$\lim_{\epsilon \rightarrow 0} \epsilon^{-1/2} \mathbb{E} \left| R_{t,T}^{(1,b)} \right| \leq \frac{K}{\sqrt{N}}$$

holds for any  $N$ . Thus, we have the desired result, by letting  $N \rightarrow \infty$ .

Proof of (A.3) for  $j = 3$  is given by repeating the same argument as in the previous case to  $Z_s^{(3)} = (D_2 + 2D_1)v^{(1)}(s, X_s^{\pi^{(0)}})$  which also satisfies the two properties since  $Z_s^{(3)} = K \left( X_s^{\pi^{(0)}} \right)^{1-\gamma}$ , and to  $\frac{1}{2}\sqrt{\epsilon}\rho\overline{D}(\lambda^2(Y_s^{\epsilon,H}) - \overline{\lambda}^2)$  which is clearly of order  $\sqrt{\epsilon}$ . □

## References

- J.-M. Bardet, G. Lang, G. Oppenheim, A. Philippe, and M. S. Taqqu. Generators of long-range dependent processes: a survey. *Theory and applications of long-range dependence*, pages 579–623, 2003.
- F. Biagini, Y. Hu, B. Øksendal, and T. Zhang. *Stochastic calculus for fractional Brownian motion and applications*. Springer Science & Business Media, 2008.
- P. Cheridito, H. Kawaguchi, and M. Maejima. Fractional ornstein-uhlenbeck processes. *Electronic Journal of Probability*, 8(3):1–14, 2003.
- L. Coutin. An introduction to (stochastic) calculus with respect to fractional brownian motion. In *Séminaire de Probabilités XL*, pages 3–65. Springer, 2007.
- J.-P. Fouque and R. Hu. Asymptotic optimal strategy for portfolio optimization in a slowly varying stochastic environment. *SIAM Journal on Control and Optimization*, 5(3), 2017a.
- J.-P. Fouque and R. Hu. Optimal portfolio under fractional stochastic environment. *arXiv preprint arXiv:1703.06969*, 2017b.
- J.-P. Fouque and R. Hu. Optimal portfolio under fast mean-reverting fractional stochastic environment. *SIAM Journal on Financial Mathematics*, 2018. to appear.
- J.-P. Fouque, G. Papanicolaou, and R. Sircar. *Derivatives in financial markets with stochastic volatility*. Cambridge University Press, 2000.
- J.-P. Fouque, G. Papanicolaou, and R. Sircar. Stochastic volatility and epsilon-martingale decomposition. In *Trends in Mathematics, Birkhauser Proceedings of the Workshop on Mathematical Finance*, pages 152–161. Springer, 2001.
- J.-P. Fouque, G. Papanicolaou, R. Sircar, and K. Sølna. Short time-scale in s&p500 volatility. *Journal of Computational Finance*, 6(4):1–24, 2003.
- J.-P. Fouque, G. Papanicolaou, R. Sircar, and K. Sølna. *Multiscale Stochastic Volatility for Equity, Interest-Rate and Credit Derivatives*. Cambridge University Press, 2011.
- J.-P. Fouque, R. Sircar, and T. Zariphopoulou. Portfolio optimization & stochastic volatility asymptotics. *Mathematical Finance*, 2015.
- C. Frei and M. Schweizer. Exponential utility indifference valuation in two brownian settings with stochastic correlation. *Advances in Applied Probability*, 40(2):401–423, 2008.
- J. Garnier and K. Sølna. Option pricing under fast-varying long-memory stochastic volatility. *arXiv preprint arXiv:1604.00105*, 2016.
- J. Garnier and K. Sølna. Correction to black-scholes formula due to fractional stochastic volatility. *SIAM Journal on Financial Mathematics*, 8(1), 2017a.

- J. Garnier and K. Sølna. Option pricing under fast-varying and rough stochastic volatility. *arXiv preprint arXiv:1707.00610*, 2017b.
- J. Gatheral, T. Jaisson, and M. Rosenbaum. Volatility is rough. *arXiv preprint arXiv:1410.3394*, 2014.
- R. Hu. Asymptotic optimal portfolio in fast mean-reverting stochastic environments. *arXiv preprint arXiv:1803.07720*, 2018a.
- R. Hu. Asymptotic methods for portfolio optimization problem in multiscale stochastic environments, 2018b. In preparation.
- T. Kaarakka and P. Salminen. On fractional ornstein-uhlenbeck process. *Communications on Stochastic Analysis*, 5(1):121–133, 2011.
- B. B. Mandelbrot and J. W. Van Ness. Fractional brownian motions, fractional noises and applications. *SIAM review*, 10(4):422–437, 1968.
- R. C. Merton. Lifetime portfolio selection under uncertainty: The continuous-time case. *Review of Economics and statistics*, 51:247–257, 1969.
- R. C. Merton. Optimum consumption and portfolio rules in a continuous-time model. *Journal of economic theory*, 3(4):373–413, 1971.
- M. Tehranchi. Explicit solutions of some utility maximization problems in incomplete markets. *Stochastic Processes and their Applications*, 114(1):109–125, 2004.
- T. Zariphopoulou. Optimal investment and consumption models with non-linear stock dynamics. *Mathematical Methods of Operations Research*, 50(2):271–296, 1999.