

Maximum likelihood estimation for Gaussian processes under inequality constraints

François Bachoc¹, Agnès Lagnoux¹, and Andrés F. López-Lopera²

¹Institut de Mathématiques de Toulouse, France.

²Mines Saint-Étienne, UMR CNRS 6158, LIMOS, F-42023 Saint-Étienne, France.

Abstract

We consider covariance parameter estimation for a Gaussian process under inequality constraints (boundedness, monotonicity or convexity) in fixed-domain asymptotics. We first show that the (unconstrained) maximum likelihood estimator has the same asymptotic distribution, unconditionally and conditionally, to the fact that the Gaussian process satisfies the inequality constraints. Then, we study the recently suggested constrained maximum likelihood estimator. We show that it has the same asymptotic distribution as the (unconstrained) maximum likelihood estimator. In addition, we show in simulations that the constrained maximum likelihood estimator is generally more accurate on finite samples.

1 Introduction

Kriging (Rasmussen and Williams, 2006; Stein, 1999) consists in inferring the values of a Gaussian random field given observations at a finite set of points. It has become a popular method for a large range of applications, such as geostatistics (Matheron, 1970), numerical code approximation (Bachoc et al., 2016; Sacks et al., 1989; Santner et al., 2003) and calibration (Bachoc et al., 2014; Paulo et al., 2012), global optimization (Jones et al., 1998), and machine learning (Rasmussen and Williams, 2006).

When considering a Gaussian process, one has to deal with the estimation of its covariance function. Usually, it is assumed that the covariance function belongs to a given parametric family (see (Abrahamsen, 1997) for a review of classical families). In this case, the estimation boils down to estimating the corresponding covariance parameters. The main estimation techniques that have been considered are based on maximum likelihood (Stein, 1999), cross-validation (Bachoc, 2013, 2014; Zhang and Wang, 2010) and variation estimators (Anderes, 2010; Istas and Lang, 1997).

In this paper, we address maximum likelihood estimation of covariance parameters under fixed-domain asymptotics (Stein, 1999). The fixed domain asymptotic setting corresponds to observation points for the Gaussian process that become dense in a fixed bounded domain. Under fixed-domain asymptotics, two types of covariance parameters can be distinguished: microergodic and non-microergodic parameters (Ibragimov and Rozanov, 1978; Stein, 1999). It is said that a covariance parameter is microergodic if, when it takes two different values, the two corresponding Gaussian measures are orthogonal (Ibragimov and Rozanov, 1978; Stein, 1999). Then it is said to be non-microergodic if, even for two different values, the

corresponding Gaussian measures are equivalent. Although non-microergodic parameters cannot be estimated consistently, they have an asymptotically negligible impact on prediction (Stein, 1988, 1990a,b; Zhang, 2004). On the contrary, it is at least possible to consistently estimate microergodic covariance parameters, and misspecifying them can have a strong negative impact on predictions.

It is still challenging to obtain results on maximum likelihood estimation of microergodic parameters that would hold for very general classes of covariance functions. Nevertheless, significant contributions have been made for specific types of covariance functions. In particular, when considering the isotropic Matérn family of covariance functions, for input space dimension $d = 1, 2, 3$, a reparameterized quantity obtained from the variance and correlation length parameters is microergodic (Zhang, 2004). It has been shown in (Kaufman and Shaby, 2013), from previous results in (Du et al., 2009) and (Wang and Loh, 2011), that the maximum likelihood estimator of this microergodic parameter is consistent and asymptotically Gaussian distributed. Anterior results on the exponential covariance function have been also obtained in (Ying, 1991, 1993).

In this paper, we shall consider the situation where the trajectories of the Gaussian process are known to satisfy either boundedness, monotonicity or convexity constraints. Indeed, Gaussian processes with inequality constraints provide suitable regression models in application fields such as computer networking (monotonicity) (Golchi et al., 2015), social system analysis (monotonicity) (Riihimäki and Vehtari, 2010) and econometrics (monotonicity or positivity) (Cousin et al., 2016). Furthermore, it has been shown that taking the constraints into account may considerably improve the predictions and the predictive intervals for the Gaussian process (Da Veiga and Marrel, 2012; Golchi et al., 2015; Riihimäki and Vehtari, 2010).

Recently, a constrained maximum likelihood estimator (cMLE) for the covariance parameters has been suggested in (López-Lopera et al., 2017). Contrary, to the (unconstrained) maximum likelihood estimator (MLE) discussed above, the cMLE explicitly takes into account the additional information brought by the inequality constraints. In (López-Lopera et al., 2017), it is shown, essentially, that the consistency of the MLE implies the consistency of the cMLE under boundedness, monotonicity or convexity constraints.

The aim of this paper is to study the asymptotic conditional distributions of the MLE and the cMLE, given that the Gaussian process satisfies the constraints. We consider the estimation of a single variance parameter and the estimation of the microergodic parameter in the isotropic Matérn family of covariance functions. In both cases, we show that the asymptotic conditional distributions of the MLE and the cMLE are identical to the unconditional asymptotic distribution of the MLE. Hence, it turns out that the impact of the constraints on covariance parameter estimation is asymptotically negligible. To the best of our knowledge, this paper is the first work on the asymptotic distribution of covariance parameter estimators for constrained Gaussian processes. The proofs involve tools from asymptotic spatial statistics, extrema of Gaussian processes and reproducing kernel Hilbert spaces. In simulations, we confirm that for large sample sizes, the MLE and the cMLE have very similar empirical distributions, that are close to the asymptotic Gaussian distribution. For small or moderate sample sizes, we observe that the cMLE is generally more accurate than the MLE, so that taking the constraints into account is beneficial.

The rest of the manuscript is organized as follows. In Section 2, we introduce in details the constraints and the MLE and the cMLE. In Section 3 we provide the asymptotic results for the estimation of the variance parameter. In Section 4, we provide the asymptotic results for the isotropic Matérn family of covariance functions. In Section 5, we report the simulation outcomes. Concluding remarks are given in Section 6. Intermediate results for the proofs are given in Appendix A. The proof of Theorems 3.2, 3.3, 4.2, 4.3 and 4.4 for boundedness, monotonicity and convexity constraints are given in the Appendix B, C and D (respectively).

2 Gaussian processes under inequality constraints

2.1 Framework and purpose of the paper

We consider a parametric set of covariance functions $\{k_\theta; \theta \in \Theta\}$ defined on $[0, 1]^d$, where Θ is compact in \mathbb{R}^p . We also assume that, for each $\theta \in \Theta$, there exists a Gaussian process with continuous realizations having mean function zero and covariance function k_θ . We refer to, e.g., (Adler, 1990) for mild smoothness conditions on k_θ ensuring this. We consider an application

$$Y : (\Omega, \mathcal{A}) \rightarrow (\mathcal{C}([0, 1]^d), \mathcal{B}),$$

where (Ω, \mathcal{A}) is a measurable space, $\mathcal{C}([0, 1]^d)$ is the set of continuous functions from $[0, 1]^d$ to \mathbb{R} , and \mathcal{B} is the Borel Sigma algebra on $\mathcal{C}([0, 1]^d)$ corresponding to the L^∞ norm. For each $\theta \in \Theta$, let \mathbb{P}_θ be a probability measure on Ω for which

$$Y : (\Omega, \mathcal{A}, \mathbb{P}_\theta) \rightarrow (\mathcal{C}([0, 1]^d), \mathcal{B})$$

has the distribution of a Gaussian process with mean function zero and covariance function k_θ .

Now consider a triangular array $(x_i^{(n)})_{n \in \mathbb{N}, i=1, \dots, n}$ of observation points in $[0, 1]^d$, where we write for concision $(x_1, \dots, x_n) = (x_1^{(n)}, \dots, x_n^{(n)})$. We assume that $(x_i^{(n)})$ is dense, that is $\sup_{x \in [0, 1]^d} \inf_{i=1, \dots, n} |x - x_i^{(n)}| \rightarrow 0$ as $n \rightarrow \infty$. Let y be the Gaussian vector defined by $y_i = Y(x_i)$ for $i = 1, \dots, n$. For $\theta \in \Theta$, let $R_\theta = [k_\theta(x_i, x_j)]_{1 \leq i, j \leq n}$ and

$$\mathcal{L}_n(\theta) = -\frac{n}{2} \log(2\pi) - \frac{1}{2} \log(|R_\theta|) - \frac{1}{2} y^\top R_\theta^{-1} y \quad (2.1)$$

be the log likelihood function. Here, $|R_\theta|$ stands for $\det(R_\theta)$. Maximizing $\mathcal{L}_n(\theta)$ with respect to θ yields the widely studied and applied MLE (Santner et al., 2003; Stein, 1999; Ying, 1993; Zhang, 2004).

In this paper, we assume that the information $\{Y \in \mathcal{E}_\kappa\}$ is available where \mathcal{E}_κ is a convex set of functions defined by inequality constraints. We will consider

$$\begin{aligned} \mathcal{E}_0 &= \{f \in \mathcal{C}([0, 1]^d, \mathbb{R}) \mid \text{s.t. } \ell \leq f(x) \leq u, \forall x \in [0, 1]^d\}, \\ \mathcal{E}_1 &= \{f \in \mathcal{C}^1([0, 1]^d, \mathbb{R}) \mid \text{s.t. } \partial f(x)/\partial x_i \geq 0, \forall x \in [0, 1]^d, i \in \{1, \dots, d\}\}, \\ \mathcal{E}_2 &= \{f \in \mathcal{C}^2([0, 1]^d, \mathbb{R}) \mid \text{s.t. } f \text{ is convex}\}, \end{aligned}$$

which correspond to boundedness, monotonicity and convexity constraints. For \mathcal{E}_0 , the bounds $-\infty \leq \ell < u \leq +\infty$ are fixed and known.

First, we will study the conditional asymptotic distribution of the (unconstrained) MLE obtained by maximizing (2.1), given $\{Y \in \mathcal{E}_\kappa\}$. Nevertheless, a drawback of this MLE is that it does not exploit the information $\{Y \in \mathcal{E}_\kappa\}$. Then we study the cMLE introduced in (López-Lopera et al., 2017). This estimator is obtained by maximizing the logarithm of the probability density function of y , conditionally to $\{Y \in \mathcal{E}_\kappa\}$, with respect to the probability measure \mathbb{P}_θ on Ω . This logarithm of conditional density is given by

$$\begin{aligned}\mathcal{L}_{n,c}(\theta) &= \mathcal{L}_n(\theta) - \log \mathbb{P}_\theta(Y \in \mathcal{E}_\kappa) + \log \mathbb{P}_\theta(Y \in \mathcal{E}_\kappa | y) \\ &= \mathcal{L}_n(\theta) + A_n(\theta) + B_n(\theta),\end{aligned}\tag{2.2}$$

say, where $\mathbb{P}_\theta(\cdot)$ and $\mathbb{P}_\theta(\cdot|\cdot)$ are defined in Section 2.2. In (López-Lopera et al., 2017), the cMLE is studied and compared to the MLE. The authors show that the cMLE is consistent when the MLE is. In this paper, we aim at providing more quantitative results regarding the asymptotic distribution of the MLE and the cMLE, conditionally to $\{Y \in \mathcal{E}_\kappa\}$.

2.2 Notation

In the paper, $0 < c < +\infty$ stands for a generic constant that may differ from one line to another. It is convenient to have short expressions for terms that converge in probability to zero. Following (van der Vaart, 1998), the notation $o_{\mathbb{P}}(1)$ (respectively $O_{\mathbb{P}}(1)$) stands for a sequence of random variables (r.v.'s) that converges to zero in probability (resp. is bounded in probability) as $n \rightarrow \infty$. More generally, for a sequence of r.v.'s R_n ,

$$\begin{aligned}X_n &= o_{\mathbb{P}}(R_n) \quad \text{means} \quad X_n = Y_n R_n \quad \text{with} \quad Y_n \xrightarrow{\mathbb{P}} 0, \\ X_n &= O_{\mathbb{P}}(R_n) \quad \text{means} \quad X_n = Y_n R_n \quad \text{with} \quad Y_n = O_{\mathbb{P}}(1).\end{aligned}$$

For deterministic sequences X_n and R_n , the stochastic notation reduce to the usual o and O . For a sequence of random vectors or variables $(X_n)_{n \in \mathbb{N}}$ on \mathbb{R}^l , that are functions of Y , and for a probability distribution μ on \mathbb{R}^l , we write

$$X_n \xrightarrow[n \rightarrow \infty]{\mathcal{L}|Y \in \mathcal{E}_\kappa} \mu$$

when, for any bounded continuous function $g : \mathbb{R}^l \rightarrow \mathbb{R}$, we have

$$\mathbb{E}[g(X_n) | Y \in \mathcal{E}_\kappa] \xrightarrow[n \rightarrow \infty]{} \int_{\mathbb{R}^l} g(x) \mu(dx).$$

We also write $X_n = o_{\mathbb{P}|Y \in \mathcal{E}_\kappa}(1)$ when for all $\varepsilon > 0$ we have $\mathbb{P}(|X_n| \geq \varepsilon | Y \in \mathcal{E}_\kappa) \rightarrow 0$ as $n \rightarrow \infty$.

For any two functions $f(Y)$ and $g(Y)$, let $\mathbb{E}_\theta[f(Y)]$ (respectively $\mathbb{E}_\theta[f(Y)|g(Y)]$) be the expectation (resp. the conditional expectation) with respect to the measure \mathbb{P}_θ on Ω . We define similarly $\mathbb{P}_\theta(A(Y))$ and $\mathbb{P}_\theta(A(Y)|g(Y))$ when $A(Y)$ is an event with respect to Y . Let $\theta_0 \in \Theta$ be fixed. We consider θ_0 as the true unknown covariance parameter and we let $\mathbb{E}[\cdot]$, $\mathbb{E}[\cdot|\cdot]$, $\mathbb{P}(\cdot)$, and $\mathbb{P}(\cdot|\cdot)$ be shorthands for $\mathbb{E}_{\theta_0}[\cdot]$, $\mathbb{E}_{\theta_0}[\cdot|\cdot]$, $\mathbb{P}_{\theta_0}(\cdot)$, and $\mathbb{P}_{\theta_0}(\cdot|\cdot)$. When a quantity is said to converge, say, in probability or almost surely, it is also implicit that we consider the measure \mathbb{P}_{θ_0} on Ω .

For $a > 0$, let $f_a : (0, \infty) \rightarrow \mathbb{R}$ be defined by $f_a(t) = -\log(t) - a/t$. We will repeatedly use the fact that f_a has a unique global maximum at a and $f_a''(t) = 1/t^2 - 2a/t^3$.

Finally, let $\xi_* = \inf_{x \in [0,1]^d} \xi(x)$, $\xi^* = \sup_{x \in [0,1]^d} \xi(x)$, and $\xi^{**} = \sup_{x \in [0,1]^d} |\xi(x)|$ for any stochastic process $\xi : [0, 1]^d \rightarrow \mathbb{R}$.

2.3 Conditions on the observation points

In some cases, we will need to assume that as $n \rightarrow \infty$, the triangular array of observation points contains finer and finer tensorized grids.

Condition-Grid. There exist d sequences $(v_i^{(j)})_{i \in \mathbb{N}}$ for $j = 1, \dots, d$ that are dense in $[0, 1]$ and so that for all $N \in \mathbb{N}$, there exists $n_0 \in \mathbb{N}$ such that for $n \geq n_0$ we have $\{(v_{i_1}^{(1)}, \dots, v_{i_d}^{(d)}), 1 \leq i_1, \dots, i_d \leq N\} \subset (x_i)_{i=1, \dots, n}$.

In our opinion, Condition-Grid is reasonable and natural. Its purpose is to guarantee that the partial derivatives of Y are consistently estimable from y everywhere on $[0, 1]^d$ (see, for instance, the proof of Theorems 3.2 and 3.3 for $\kappa = 1$ in the supplementary material). We believe that, for the results for which Condition-Grid is assumed, one could replace it by a milder condition and prove similar results. Then the proofs would be based on essentially the same ideas as the current ones, but could be more cumbersome.

In some other cases, we only need to assume that the observation points constitute a sequence.

Condition-Sequence. For all $n \in \mathbb{N}$ and $i \leq n$, we have $x_i^{(n)} = x_i^{(i)}$.

Condition-Sequence implies that sequences of conditional expectations with respect to the observations are martingales. This condition is necessary in some of the proofs (for instance, that of Theorem 3.3) where convergence results for martingales are used.

3 Variance parameter estimation

3.1 Model and assumptions

In this section, we focus on the estimation of a single variance parameter when the correlation function is known. Hence, we let $p = 1$, $\theta = \sigma^2$, and for $x, x' \in [0, 1]^d$,

$$k_{\sigma^2}(x, x') = \sigma^2 k_1(x, x'), \quad (3.1)$$

where k_1 is a fixed known correlation function. We make the following assumption.

Condition-Var. Let κ be fixed in $\{0, 1, 2\}$.

- k_1 is stationary, that is $k_1(x, x') = k_1(x - x')$ for x and x' in $[0, 1]^d$.

- If $\kappa = 0$, k_1 is α -Hölder, which means that there exist nonnegative constants C and α such that

$$|k_1(t) - k_1(t')| \leq C \|t - t'\|^\alpha$$

for all t and t' in $[0, 1]^d$, where $\|\cdot\|$ is the Euclidean norm. Furthermore, the Fourier transform \widehat{k}_1 of k_1 satisfies, for some fixed $P < \infty$,

$$\widehat{k}_1(\omega) \|\omega\|^P \xrightarrow{\omega \rightarrow \infty} \infty. \quad (3.2)$$

- If $\kappa = 1$, the Gaussian process Y is differentiable in quadratic mean. For $i = 1, \dots, d$, let $k_{1,i}$ be the covariance function of $\partial Y / \partial x_i$. Then $k_{1,i}$ is α -Hölder for a fixed $\alpha > 0$. Also, (3.2) holds with \widehat{k}_1 replaced by the Fourier transform $\widehat{k}_{1,i}$ of $k_{1,i}$ for $i = 1, \dots, d$.
- If $\kappa = 2$, the Gaussian process Y is twice differentiable in quadratic mean. For $i, j = 1, \dots, d$, let $k_{2,i,j}$ be the covariance function of $\partial^2 Y / (\partial x_i \partial x_j)$. Then $k_{2,i,j}$ is α -Hölder for a fixed $\alpha > 0$. Also, (3.2) holds with \widehat{k}_1 replaced by the Fourier transform $\widehat{k}_{1,i,j}$ of $k_{1,i,j}$ for $i, j = 1, \dots, d$.

These assumptions make the conditioning by $\{Y \in \mathcal{E}_\kappa\}$ valid for $\kappa = 0, 1, 2$ as established in the following lemma.

Lemma 3.1. *Assume that Condition-Var holds. Then for all $\kappa \in \{0, 1, 2\}$ and for any compact K in $(0, +\infty)$, we have*

$$\inf_{\sigma^2 \in K} \mathbb{P}_{\sigma^2}(Y \in \mathcal{E}_\kappa) > 0.$$

Proof of Lemma 3.1. It suffices to follow the same lines as in the proof of Lemma A.6 in (López-Lopera et al., 2017). \square

3.2 Asymptotic conditional distribution of the maximum likelihood estimator

The log-likelihood function in (2.1) for σ^2 can be written as

$$\mathcal{L}_n(\sigma^2) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2} \ln(|R_1|) - \frac{1}{2\sigma^2} y^\top R_1^{-1} y, \quad (3.3)$$

where $R_1 = [k_1(x_i, x_j)]_{1 \leq i, j \leq n}$. Then the standard MLE is given by

$$\bar{\sigma}_n^2 \in \operatorname{argmax}_{\sigma^2 > 0} \mathcal{L}_n(\sigma^2). \quad (3.4)$$

Now we show that, for $\kappa = 0, 1, 2$, $\sqrt{n}(\bar{\sigma}_n^2 - \sigma_0^2)$ is asymptotically Gaussian distributed conditionally to $\{Y \in \mathcal{E}_\kappa\}$.

Theorem 3.2. *For $\kappa = 1, 2$, we assume that Condition-Grid holds. Under Condition-Var, the MLE $\bar{\sigma}_n^2$ of σ_0^2 defined by (3.4) conditioned on $\{Y \in \mathcal{E}_\kappa\}$ is asymptotically Gaussian distributed. More precisely, for $\kappa = 0, 1, 2$,*

$$\sqrt{n} (\bar{\sigma}_n^2 - \sigma_0^2) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}|Y \in \mathcal{E}_\kappa} \mathcal{N}(0, 2\sigma_0^4).$$

It is well known that $\sqrt{n} (\bar{\sigma}_n^2 - \sigma_0^2)$ converges (unconditionally) to the $\mathcal{N}(0, 2\sigma_0^4)$ distribution. Hence, conditioning by $\{Y \in \mathcal{E}_\kappa\}$ has no impact on the asymptotic distribution of the MLE.

3.3 Asymptotic conditional distribution of the constrained maximum likelihood estimator

Here, we assume that the compact set Θ is $[\sigma_l^2, \sigma_u^2]$ with $0 < \sigma_l^2 < \sigma_0^2 < \sigma_u^2 < +\infty$ and we consider the cMLE $\hat{\sigma}_{n,c}^2$ of σ_0^2 derived by maximizing on the compact set Θ the constrained log-likelihood in (2.2):

$$\hat{\sigma}_{n,c}^2 \in \operatorname{argmax}_{\sigma^2 \in \Theta} \mathcal{L}_{n,c}(\sigma^2). \quad (3.5)$$

Now we show that the conditional asymptotic distribution of the cMLE is the same as that of the MLE.

Theorem 3.3. *For $\kappa = 1, 2$, we assume that Condition-Grid holds. Under Condition-Var and Condition-Sequence, the cMLE $\hat{\sigma}_{n,c}^2$ of σ_0^2 defined in (3.5) is asymptotically Gaussian distributed. More precisely, for $\kappa = 0, 1, 2$,*

$$\sqrt{n} (\hat{\sigma}_{n,c}^2 - \sigma_0^2) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}|Y \in \mathcal{E}_\kappa} \mathcal{N}(0, 2\sigma_0^4).$$

4 Microergodic parameter estimation for the isotropic Matérn model

4.1 Model and assumptions

In this section, we let $d = 1, 2$ or 3 and we consider the isotropic Matérn family of covariance functions on $[0, 1]^d$. We refer to, e.g., (Stein, 1999) for more details. Here $k_\theta = k_{\theta,\nu}$ is given by, for $x, x' \in [0, 1]^d$,

$$k_{\theta,\nu}(x, x') = \sigma^2 K_\nu \left(\frac{\|x - x'\|}{\rho} \right) = \frac{\sigma^2}{\Gamma(\nu) 2^{\nu-1}} \left(\frac{\|x - x'\|}{\rho} \right)^\nu \kappa_\nu \left(\frac{\|x - x'\|}{\rho} \right).$$

The parameter $\sigma^2 > 0$ is the variance of the process, $\rho > 0$ is the correlation length parameter that controls how fast the covariance function decays with the distance, and $\nu > 0$ is the regularity parameter of the process. The function κ_ν is the modified Bessel function of the second kind of order ν (see (Abramowitz and Stegun, 1964)). We assume in the sequel that the smoothness parameter ν is known. Then $\theta = (\sigma^2, \rho)$ and $p = 2$.

Condition- ν . For $\kappa = 0$ (respectively $\kappa = 1$ and $\kappa = 2$), we assume that $\nu > 0$ (resp. $\nu > 1$ and $\nu > 2$).

We remark that Condition- ν naturally implies Condition-Var so that the conditioning by $\{Y \in \mathcal{E}_\kappa\}$ is valid for any $\kappa = 0, 1, 2$ as established in the next lemma. We refer to (Stein, 1999) for a reference on the impact of ν on the smoothness of the Matérn covariance function and on its Fourier transform.

Lemma 4.1. *Assume that Condition- ν holds. Then for all $\kappa \in \{0, 1, 2\}$ and for any compact K of $(0, \infty)^2$, we have*

$$\inf_{(\sigma^2, \rho) \in K} \mathbb{P}_{\sigma^2, \rho}(Y \in \mathcal{E}_\kappa) > 0.$$

Proof of Lemma 4.1. This lemma is a special case of Lemma A.6 in (López-Lopera et al., 2017). \square

4.2 Asymptotic conditional distribution of the maximum likelihood estimator

The log-likelihood function in (2.1) for σ^2 and ρ under the Matérn model with fixed parameter ν can be written as

$$\mathcal{L}_n(\sigma^2, \rho) = -\frac{n}{2} \log 2\pi - \frac{1}{2} n \ln(\sigma^2) - \frac{1}{2} \ln(|R_{\rho, \nu}|) - \frac{1}{2\sigma^2} y^\top R_{\rho, \nu}^{-1} y, \quad (4.1)$$

where $R_{\rho, \nu} = [K_\nu(\|x_i - x_j\|/\rho)]_{1 \leq i, j \leq n}$. Let $\Theta = [\sigma_l^2, \sigma_u^2] \times [\rho_l, \rho_u]$ with fixed $0 < \sigma_l^2 < \sigma_u^2 < \infty$ and fixed $0 < \rho_l < \rho_u < \infty$. Assume moreover that the true parameters are such that $\sigma_l^2/(\rho_l^{2\nu}) < \sigma_0^2/(\rho_0^{2\nu}) < \sigma_u^2/(\rho_u^{2\nu})$. Then the MLE is given by

$$(\hat{\sigma}_n^2, \hat{\rho}_n) \in \operatorname{argmax}_{(\sigma^2, \rho) \in \Theta} \mathcal{L}_n(\sigma^2, \rho). \quad (4.2)$$

It has been shown in (Zhang, 2004) that the parameters σ_0^2 and ρ_0 can not be estimated consistently but that the microergodic parameter $\sigma_0^2/\rho_0^{2\nu}$ can. Furthermore, it is shown in (Kaufman and Shaby, 2013) that $\sqrt{n}(\hat{\sigma}_n^2/\hat{\rho}_n^{2\nu} - \sigma_0^2/\rho_0^{2\nu})$ converges to a $\mathcal{N}(0, 2(\sigma_0^2/\rho_0^{2\nu})^2)$ distribution. In the next theorem, we show that this asymptotic normality also holds conditionally to $\{Y \in \mathcal{E}_\kappa\}$.

Theorem 4.2. *For $\kappa = 1, 2$, we assume that Condition-Grid holds. Under Condition- ν , the estimator $\hat{\sigma}_n^2/\hat{\rho}_n^{2\nu}$ of the microergodic parameter $\sigma_0^2/\rho_0^{2\nu}$ defined by (4.2) and conditioned on $\{Y \in \mathcal{E}_\kappa\}$ is asymptotically Gaussian distributed. More precisely, for $\kappa = 0, 1, 2$,*

$$\sqrt{n} \left(\frac{\hat{\sigma}_n^2}{\hat{\rho}_n^{2\nu}} - \frac{\sigma_0^2}{\rho_0^{2\nu}} \right) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}|Y \in \mathcal{E}_\kappa} \mathcal{N} \left(0, 2 \left(\frac{\sigma_0^2}{\rho_0^{2\nu}} \right)^2 \right).$$

4.3 Asymptotic conditional distribution of the constrained maximum likelihood estimator

We turn to the constrained log-likelihood and its maximizer. We consider two types of estimation settings obtained by maximizing the constrained log-likelihood (2.2) under the Matérn model. In the first setting, $\rho = \rho_1$ is fixed and (2.2) is maximized over σ^2 (in the case $\rho_1 = \rho_0$ this setting is already covered by Theorem 3.3). In the second setting, (2.2) is maximized over both σ^2 and ρ . Under the two settings, we show that the cMLE has the same asymptotic distribution as the MLE, conditionally to $\{Y \in \mathcal{E}_\kappa\}$.

Theorem 4.3 (Fixed correlation length parameter ρ_1). *For $\kappa = 1, 2$, we assume that Condition-Grid holds. Assume that Condition- ν and Condition-Sequence hold. Let for $\rho \in [\rho_l, \rho_u]$,*

$$\hat{\sigma}_{n,c}^2(\rho) \in \operatorname{argmax}_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathcal{L}_{n,c}(\sigma^2, \rho). \quad (4.3)$$

Let $\rho_1 \in [\rho_l, \rho_u]$ be fixed. Then $\hat{\sigma}_{n,c}^2(\rho_1)$ is asymptotically Gaussian distributed. More precisely, for $\kappa = 0, 1, 2$,

$$\sqrt{n} \left(\frac{\hat{\sigma}_{n,c}^2(\rho_1)}{\rho_1^{2\nu}} - \frac{\sigma_0^2}{\rho_0^{2\nu}} \right) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}|Y \in \mathcal{E}_\kappa} \mathcal{N} \left(0, 2 \left(\frac{\sigma_0^2}{\rho_0^{2\nu}} \right)^2 \right).$$

Theorem 4.4 (Estimated correlation length parameter). *Assume that Condition- ν holds. Let $\hat{\sigma}_{n,c}^2(\rho)$ be defined as in (4.3) and let $(\hat{\sigma}_{n,c}^2, \hat{\rho}_{n,c})$ be defined by*

$$(\hat{\sigma}_{n,c}^2, \hat{\rho}_{n,c}) \in \operatorname{argmax}_{(\sigma^2, \rho) \in \Theta} \mathcal{L}_{n,c}(\sigma^2, \rho).$$

Notice that $\hat{\sigma}_{n,c}^2 = \hat{\sigma}_{n,c}^2(\hat{\rho}_{n,c})$.

(i) *For $\kappa = 0$, assume that one of the following two conditions hold.*

- a) *We have $\nu > 1$, $d = 1$ and $\max_{x \in [0,1]} \min_{i=1, \dots, n} |x - x_i| = o(1/\sqrt{n})$.*
- b) *We have $\nu > 2$ and there exists a sequence $(a_n)_{n \in \mathbb{N}}$ with $a_n = o(1/n^{1/4})$ as $n \rightarrow \infty$, so that, for all $x \in [0, 1]^d$, there exists $d + 1$ points v_1, \dots, v_{d+1} with $\{v_1, \dots, v_{d+1}\} \subset \{x_1, \dots, x_n\}$, so that x belongs to the convex hull of v_1, \dots, v_{d+1} and $\max_{j=1, \dots, d+1} \|x - v_j\| \leq a_n$.*

(ii) *For $\kappa = 1, 2$, assume that one of the following two conditions hold.*

- a) *We have $\nu > \kappa + 1$, $d = 1$ and $\max_{x \in [0,1]} \min_{i=1, \dots, n} |x - x_i| = o(1/\sqrt{n})$.*
- b) *We have $\nu > \kappa + 2$ and the observation points $\{x_1, \dots, x_n\}$ are so that, for all $n \geq 2^d$, with $N = \lfloor n^{1/d} \rfloor$,*

$$\{x_1, \dots, x_n\} \supset \left\{ \left(\frac{i_1}{N-1}, \dots, \frac{i_d}{N-1} \right), 0 \leq i_1, \dots, i_d \leq N-1 \right\}.$$

Then $\widehat{\sigma}_{n,c}^2/\widehat{\rho}_{n,c}^{2\nu}$ is asymptotically Gaussian distributed. More precisely, for $\kappa = 0, 1, 2$,

$$\sqrt{n} \left(\frac{\widehat{\sigma}_{n,c}^2}{\widehat{\rho}_{n,c}^{2\nu}} - \frac{\sigma_0^2}{\rho_0^{2\nu}} \right) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}|Y \in \mathcal{E}_\kappa} \mathcal{N} \left(0, 2 \left(\frac{\sigma_0^2}{\rho_0^{2\nu}} \right)^2 \right).$$

In Theorem 4.4, we assume that ν is larger than in Condition- ν , and we assume that the observation points have specific quantitative space filling properties. The condition i) b) also implies that a portion of the observation points are located in the corners and borders of $[0, 1]^d$. Furthermore, the condition ii) b) implies that the majority of the observation points are located on regular grids. We believe that these two last conditions could be replaced by milder ones, at the cost of proofs similar to but more cumbersome than the present ones.

We make stronger assumptions in Theorem 4.4 than in Theorem 4.3 because the former is more challenging than the latter. Indeed, since $\rho = \rho_1$ is fixed in Theorem 4.3, we can use the equivalence of two fixed Gaussian measures in order to obtain asymptotic properties of the conditional mean function of Y under $k_{1,\rho_1,\nu}$ (see the developments following (B.12) in the proofs). This is not possible anymore when considering the conditional mean function of Y under $k_{1,\widehat{\rho}_{n,c},\nu}$, where $\widehat{\rho}_{n,c}$ is random. Hence, we use other proof techniques, based on reproducing kernel Hilbert spaces, for studying this conditional mean function, for which the above additional conditions are needed. We refer for instance to the developments following (B.17) in the supplementary material for more details.

5 Numerical results

In this section, we illustrate numerically the conditional asymptotic normality of the MLE and the cMLE of the microergodic parameter for the Matérn 5/2 covariance function. The numerical experiments were implemented using the R package “LineqGPR” (López-Lopera, 2017).

5.1 Experimental settings

We let $d = 1$ in the rest of the section. Since the event $Y \in \mathcal{E}_\kappa$ can not be simulated exactly in practice, we consider the piecewise affine interpolation Y_m of Y at $t_1, \dots, t_m \in [0, 1]$, with $m > n$ (López-Lopera et al., 2017; Maatouk and Bay, 2017). Then, the event $Y \in \mathcal{E}_\kappa$ is approximated by the event $Y_m \in \mathcal{E}'_\kappa$, where \mathcal{E}'_0 (respectively $\mathcal{E}'_1, \mathcal{E}'_2$) is the set of continuous bounded between ℓ and u (resp. increasing, convex) functions. We can simulate efficiently Y_m conditionally to $\{Y_m \in \mathcal{E}'_\kappa\}$ by using Markov Chain Monte Carlo procedures (see, for instance, (Pakman and Paninski, 2014)).

In Section 5, we consider the Matérn 5/2 covariance function defined by

$$k_{\theta,5/2}(x, x') = \sigma^2 \left(1 + \frac{\sqrt{5}|x - x'|}{\rho} + \frac{5}{3} \frac{(x - x')^2}{\rho^2} \right) \exp \left\{ - \frac{\sqrt{5}|x - x'|}{\rho} \right\},$$

for $x, x' \in [0, 1]$ and with $\theta = (\sigma^2, \rho)$. Remark that $k_{\theta,5/2}$ is obtained by the parametrization of (López-Lopera, 2017; Roustant et al., 2012) rather than that of Section 4.1. For an easy reading, we keep the same notation.

5.2 Numerical results when ρ_0 is known

We let $m = 300$ and x_1, \dots, x_n be equispaced in $[0, 1]$ in the rest of Section 5. For $\kappa = 0, 1$, we generate $N = 1,000$ trajectories of Y_m given $\{Y_m \in \mathcal{E}'_\kappa\}$. For each of these trajectories, we compute the MLE $\bar{\sigma}_{m,n}^2(\rho_0)$ with $\bar{\sigma}_{m,n}^2(\rho) = y^\top R_\rho^{-1} y / n = \arg\max_{\sigma^2 \in (0, \infty)} \mathcal{L}_n(\sigma^2, \rho)$, where $R_\rho = [k_{1,\rho,5/2}(x_i, x_j)]_{i,j=1,\dots,d}$ for $\rho \in (0, \infty)$. Then we evaluate the cMLE as follows. We let $m_{m,n,\rho,y}$ be the conditional mean function of Y_m given y under covariance function $k_{1,\rho,5/2}$. We simulate $n_t = 1,000$ trajectories Z_1, \dots, Z_{n_t} of a Gaussian process with zero mean function and covariance function $k_{m,n,1,\rho_0,5/2}$, where $k_{m,n,1,\rho,5/2}$ is the covariance function of Y_m given y under covariance function $k_{1,\rho,5/2}$. Then we let $B_n(\sigma^2, \rho_0)$ be approximated by $\log((1/n_t) \sum_{i=1}^{n_t} \mathbb{1}_{m_{m,n,\rho_0,y} + \sigma Z_i \in \mathcal{E}'_\kappa})$. The term $A_n(\sigma^2, \rho_0)$ can be easily approximated as it does not depend on the trajectory of Y_m under consideration. We maximize the resulting approximation of $\mathcal{L}_{n,c}(\sigma^2, \rho_0)$ on 1,000 equispaced values of σ^2 between $(1 - 4\sqrt{2/n})\sigma_0^2$ and $(1 + 4\sqrt{2/n})\sigma_0^2$, yielding the approximated cMLE estimator $\hat{\sigma}_{m,n,c}^2(\rho_0)$.

In Figure 1, we report the results for $\kappa = 0$ (boundedness constraints) with $(\sigma_0^2, \rho_0) = (2, 0.2)$ and $n = 20, 50, 80$. We show the probability density functions obtained from the samples $\{n^{1/2}(\bar{\sigma}_{m,n}^2(\rho_0)_i - \sigma_0^2)\}_{i=1,\dots,N}$ and $\{n^{1/2}(\hat{\sigma}_{m,n,c}^2(\rho_0)_i - \sigma_0^2)\}_{i=1,\dots,N}$ obtained as discussed above. We also plot the probability density function of the limit $\mathcal{N}(0, 2\sigma_0^4)$ distribution. We observe that for a small number of observations, e.g. $n = 20$, the distribution of the cMLE is closer to the limit distribution than that of the MLE in terms of median value. We also observe that, as n increases, both distributions become more similar to the limit one. Nevertheless, the cMLE exhibits faster convergence.

In Figure 2, we report the same quantities for $\kappa = 1$ (monotonicity constraints) and for $(\sigma_0^2, \rho_0) = (0.5^2, 1)$. In this case, we observe that the distributions of both the MLE and the cMLE are close to the limit one even for small values of n ($n = 5, 20$).

5.3 Numerical results when ρ_0 is unknown

We let $\kappa = 0$, $(\sigma_0^2, \rho_0) = (2, 0.2)$ and $n = 20, 50, 80$. We proceed similarly as in the case where ρ_0 is known. To compute the MLE $(\bar{\sigma}_{m,n}^2(\hat{\rho}_{m,n}), \hat{\rho}_{m,n})$ of (σ_0^2, ρ_0) , we maximize $\mathcal{L}_n(\bar{\sigma}_{m,n}^2(\rho), \rho)$ over a finite grid of values for ρ . To compute the cMLE $(\hat{\sigma}_{m,n,c}^2(\hat{\rho}_{m,n,c}), \hat{\rho}_{m,n,c})$ of (σ_0^2, ρ_0) , we evaluate $\log((1/n_t) \sum_{i=1}^{n_t} \mathbb{1}_{m_{m,n,\rho,y} + \sigma Z_{\rho,i} \in \mathcal{E}'_\kappa})$ over 100^2 pairs $(\sigma_{i,j}^2, \rho_i)_{i,j=1,\dots,100}$. Here $Z_{\rho,i}$ is generated as in Section 5.2 but with ρ_0 replaced by ρ , for $i = 1, \dots, 100$. Then $\rho_1, \dots, \rho_{100}$ are equispaced in $[0.1, 0.3]$ and for $i = 1, \dots, 100$, $\sigma_{i,1}^2, \dots, \sigma_{i,100}^2$ are equispaced in

$$\rho_i^5 \left[\frac{\sigma_0^2}{\rho_0^5} - \frac{4\sqrt{2}}{\sqrt{n}} \frac{\sigma_0^2}{\rho_0^5}, \frac{\sigma_0^2}{\rho_0^5} + \frac{4\sqrt{2}}{\sqrt{n}} \frac{\sigma_0^2}{\rho_0^5} \right].$$

Hence, the estimator of the microergodic parameter σ_0^2/ρ_0^5 is restricted to be at distance less than 4 times the asymptotic standard deviation of the microergodic parameter.

In Figure 3, we show the probability density functions obtained from the samples $\{n^{1/2}(\bar{\sigma}_{m,n}^2(\hat{\rho}_{m,n})_i / \hat{\rho}_{m,n,i}^5 - \sigma_0^2/\rho_0^5)\}_{i=1,\dots,N}$ and $\{n^{1/2}(\hat{\sigma}_{m,n,c}^2(\hat{\rho}_{m,n,c})_i / \hat{\rho}_{m,n,c,i}^5 - \sigma_0^2/\rho_0^5)\}_{i=1,\dots,N}$,




Figure 1: Asymptotic conditional distribution of the variance parameter estimators under boundedness constraints. Here $(\sigma_0^2, \rho_0) = (2, 0.2)$. Each panel shows: the limit conditional distribution $\mathcal{N}(0, 2\sigma_0^4)$ (red, solid lines), and the distributions of the MLE (green, dashed lines) and of the cMLE (blue, dotted lines). The vertical lines represent the median values of the distributions. Each subcaption shows the number of observations n used for the estimations.

with $N = 1,000$. Similarly to Section 5.2, we observe that the distribution of the cMLE tends to be closer to the limit one, than that of the MLE. Moreover, the convergence with the cMLE is faster than with the MLE in terms of median value.

6 Concluding remarks

We have shown that the MLE and the cMLE are asymptotically Gaussian distributed, conditionally to the fact that the Gaussian process satisfies either boundedness, monotonicity or convexity constraints. Their asymptotic distributions are identical to the unconditional asymptotic distribution of the MLE. In simulations, we confirm that the MLE and the cMLE have very similar performances when the number n of observation points becomes large enough. We also observe that the cMLE is more accurate for small or moderate values of n .




Figure 2: Asymptotic conditional distribution of the variance parameter estimators under monotonicity constraints. Here $(\sigma_0^2, \rho_0) = (0.5^2, 1)$. Panel description is the same as in Figure 1.




Figure 3: Asymptotic conditional distribution of the microergodic parameter estimators for the isotropic $\nu = 5/2$ Matérn model under boundedness constraints. Here $(\sigma_0^2, \rho_0) = (2, 0.2)$. Panel description is the same as in Figure 1, with $\mathcal{N}(0, 2\sigma_0^4)$ replaced by $\mathcal{N}(0, 2(\sigma_0^2/\rho_0^5)^2)$.

Hence, since the computation of the cMLE is more challenging than that of the MLE, we recommend to use the MLE for large data sets and the cMLE for smaller ones. In the proofs of the asymptotic behavior of the cMLE, one of the main steps is to show that $\mathbb{P}_\theta(Y \in \mathcal{E}_\kappa|y)$ converges to one as n goes to infinity. Hence, in practice, one may evaluate $\mathbb{P}_\theta(Y \in \mathcal{E}_\kappa|y)$, for some values of θ , in order to gauge whether this conditional probability is not too close to 1 so that it is worth using the cMLE despite the additional computational cost.

Our theoretical results could be extended in different ways. First, we remark that the techniques we have used to show that $\mathbb{P}_\theta(Y \in \mathcal{E}_\kappa)$ and $\mathbb{P}_\theta(Y \in \mathcal{E}_\kappa|y)$ are asymptotically negligible (see (A.6) and (A.7)) can be used for more general families of covariance functions. Hence, other results on the (unconditional) asymptotic distribution of the MLE could be extended to the case of constrained Gaussian processes in future work. These types of results exist for instance for the product exponential covariance function (Ying, 1993) and the generalized Wendland covariance function (Bevilacqua et al., 2016).

Also, in practice, computing the cMLE requires a discretization of the constraints, for instance using a piecewise affine interpolation as in Section 5, or a finite set of constrained points (Da Veiga and Marrel, 2012). Thus it would be interesting to extend our results by taking this discretization into account.

A Intermediate results

Lemma A.1. *Let $(X_n)_{n \in \mathbb{N}}$ be a sequence of r.v.'s and $(m_{k,n})_{n,k \in \mathbb{N}, k \leq n}$ and $(M_{k,n})_{n,k \in \mathbb{N}, k \leq n}$ be two triangular arrays of r.v.'s. We consider a random vector $(m, M)^\top$ such that $m \leq m_{k,n} \leq M_{k,n} \leq M$ for all $k \leq n$. We assume that $\mathbb{P}(m = \ell) = \mathbb{P}(M = u) = 0$ and $\mathbb{P}(\ell \leq m \leq M \leq u) > 0$ for some fixed ℓ and $u \in \mathbb{R}$. Moreover, we consider a sequence $(k_n)_{n \in \mathbb{N}}$ so that, $k_n \leq n$, $k_n \rightarrow_{n \rightarrow \infty} \infty$ and*

$$(m_{k_n,n}, M_{k_n,n})^\top \xrightarrow[n \rightarrow +\infty]{a.s.} (m, M)^\top. \quad (\text{A.1})$$

Then for any $a \in \mathbb{R}$,

$$\lim_{n \rightarrow +\infty} \left| \mathbb{P}(X_n \leq a | \ell \leq m_{k_n,n} \leq M_{k_n,n} \leq u) - \mathbb{P}(X_n \leq a | \ell \leq m \leq M \leq u) \right| = 0. \quad (\text{A.2})$$

Proof of Lemma A.1. For the sake of simplicity, we denote by $E_{k,n}$ (respectively E) the event $\{\ell \leq m_{k,n} \leq M_{k,n} \leq u\}$ (resp. $\{\ell \leq m \leq M \leq u\}$). Then

$$\begin{aligned} |\mathbb{P}(X_n \leq a | E_{k_n,n}) - \mathbb{P}(X_n \leq a | E)| &\leq \frac{|\mathbb{P}(X_n \leq a, E_{k_n,n}) - \mathbb{P}(X_n \leq a, E)|}{\mathbb{P}(E_{k_n,n})} \\ &\quad + \left| \frac{1}{\mathbb{P}(E_{k_n,n})} - \frac{1}{\mathbb{P}(E)} \right| \mathbb{P}(X_n \leq a, E). \end{aligned} \quad (\text{A.3})$$

(i) By (A.1), $\mathbb{P}(E_{k_n,n})$ goes to $\mathbb{P}(E) = \mathbb{P}(\ell \leq m \leq M \leq u) > 0$ as n goes to $+\infty$. Thus $1/\mathbb{P}(E_{k_n,n})$ is well-defined for large values of n and bounded as $n \rightarrow \infty$. Moreover, by trivial arguments of set theory, one gets

$$|\mathbb{P}(X_n \leq a, E_{k_n,n}) - \mathbb{P}(X_n \leq a, E)| \leq \mathbb{P}(E_{k_n,n} \Delta E) = \mathbb{P}(E_{k_n,n} \setminus E)$$

since $\mathbb{P}(E \setminus E_{k_n,n}) = 0$. Now let $\varepsilon > 0$. One has

$$\begin{aligned}\mathbb{P}(E_{k_n,n} \setminus E) &= \mathbb{P}(\ell \leq m_{k_n,n} \leq M_{k_n,n} \leq u, (m, M) \notin [\ell, u]^2) \\ &\leq \mathbb{P}(\ell \leq m_{k_n,n} \leq M_{k_n,n} \leq u, m < \ell) + \mathbb{P}(\ell \leq m_{k_n,n} \leq M_{k_n,n} \leq u, M > u) \\ &\leq \mathbb{P}(\ell \leq m_{k_n,n}, m < \ell) + \mathbb{P}(M_{k_n,n} \leq u, M > u).\end{aligned}$$

One may decompose $\mathbb{P}(\ell \leq m_{k_n,n}, m < \ell)$ into

$$\begin{aligned}\mathbb{P}(\ell + \varepsilon \leq m_{k_n,n}, m < \ell) &+ \mathbb{P}(\ell \leq m_{k_n,n} \leq \ell + \varepsilon, m < \ell) \\ &\leq \mathbb{P}(|m_{k_n,n} - m| > \varepsilon) + \mathbb{P}(\ell \leq m_{k_n,n} \leq \ell + \varepsilon).\end{aligned}$$

The first term in the right hand-side goes to 0 as n goes to infinity. By Portemanteau's lemma and (A.1),

$$\limsup_{n \rightarrow +\infty} \mathbb{P}(\ell \leq m_{k_n,n} \leq \ell + \varepsilon) \leq \mathbb{P}(\ell \leq m \leq \ell + \varepsilon) \xrightarrow{\varepsilon \rightarrow 0} 0.$$

We handle similarly the term $\mathbb{P}(M_{k_n,n} \leq u, M > u)$. Hence, in the r.h.s. of (A.3), the first term goes to 0 as $n \rightarrow \infty$.

(ii) Now we turn to the control of the second term in (A.3). Upper bounding $\mathbb{P}(X_n \leq a, E)$ by 1, it remains to control $\left| \frac{1}{\mathbb{P}(E_{k_n,n})} - \frac{1}{\mathbb{P}(E)} \right|$ which is immediate by the convergence in distribution of $(m_{k_n,n}, M_{k_n,n})^\top$ as n goes to infinity (implied by the a.s. convergence) and the fact that $\mathbb{P}(E) > 0$ and $\mathbb{P}(m = \ell) = \mathbb{P}(M = u) = 0$. The proof is now complete. \square

Lemma A.2. *Consider three sequences of random functions $f_n, g_n, h_n : [x_{inf}, x_{sup}] \rightarrow \mathbb{R}$, with $0 < x_{inf} < x_{sup} < \infty$ fixed. Consider that for all $x \in [x_{inf}, x_{sup}]$, $f_n(x)$, $g_n(x)$, and $h_n(x)$ are functions of Y and x only. Let*

$$\hat{x}_n \in \operatorname{argmax}_{x \in [x_{inf}, x_{sup}]} f_n(x).$$

Assume the following properties.

(i) *There exists $A > 0$, $B > 0$ and $\delta > 0$ such that*

$$\sup_{\substack{|x - \hat{x}_n| \leq \delta \\ x \in [x_{inf}, x_{sup}]}} f_n(x) - f_n(\hat{x}_n) \leq -An(x - \hat{x}_n)^2 \quad (\text{A.4})$$

and

$$\sup_{\substack{|x - \hat{x}_n| > \delta \\ x \in [x_{inf}, x_{sup}]}} f_n(x) - f_n(\hat{x}_n) \leq -Bn, \quad (\text{A.5})$$

with probability going to 1 as $n \rightarrow \infty$.

(ii) *There exists $C > 0$ such that for all $x_1, x_2 \in [x_{inf}, x_{sup}]$*

$$|g_n(x_1) - g_n(x_2)| \leq C|x_1 - x_2|, \quad (\text{A.6})$$

with probability going to 1 as $n \rightarrow \infty$.

(iii) One has, for $\kappa = 0, 1, 2$,

$$\sup_{x_1, x_2 \in [x_{inf}, x_{sup}]} |h_n(x_1) - h_n(x_2)| = o_{\mathbb{P}|Y \in \mathcal{E}_\kappa}(1). \quad (\text{A.7})$$

Then, with

$$\widehat{\widehat{x}}_n \in \operatorname{argmax}_{x \in [x_{inf}, x_{sup}]} \{f_n(x) + g_n(x) + h_n(x)\}$$

we have

$$\sqrt{n}|\widehat{\widehat{x}}_n - \widehat{x}_n| = o_{\mathbb{P}|Y \in \mathcal{E}_\kappa}(1). \quad (\text{A.8})$$

Proof of Lemma A.2. Let $\varepsilon > 0$. First, we have, with probability (conditionally to $\{Y \in \mathcal{E}_\kappa\}$) going to 1 as $n \rightarrow \infty$, from (A.4), (A.6) and (A.7)

$$\begin{aligned} & \sup_{\substack{|x - \widehat{x}_n| \geq \varepsilon/\sqrt{n} \\ |x - \widehat{x}_n| \leq 1/n^{1/4}}} (f_n(x) + g_n(x) + h_n(x) - f_n(\widehat{x}_n) - g_n(\widehat{x}_n) - h_n(\widehat{x}_n)) \\ & \leq -An \left(\frac{\varepsilon}{\sqrt{n}} \right)^2 + \frac{C}{n^{1/4}} + o_{\mathbb{P}|Y \in \mathcal{E}_\kappa}(1) = -A\varepsilon^2 + o_{\mathbb{P}|Y \in \mathcal{E}_\kappa}(1). \end{aligned}$$

Second, from (A.4), (A.6) and (A.7), we have, with probability (conditionally to $\{Y \in \mathcal{E}_\kappa\}$) going to 1 as $n \rightarrow \infty$,

$$\begin{aligned} & \sup_{\substack{|x - \widehat{x}_n| \geq 1/n^{1/4} \\ |x - \widehat{x}_n| \leq \delta}} (f_n(x) + g_n(x) + h_n(x) - f_n(\widehat{x}_n) - g_n(\widehat{x}_n) - h_n(\widehat{x}_n)) \\ & \leq -An \left(\frac{1}{n^{1/4}} \right)^2 + C\delta + o_{\mathbb{P}|Y \in \mathcal{E}_\kappa}(1) \xrightarrow{n \rightarrow \infty} -\infty. \end{aligned}$$

Third, from (A.5), (A.6) and (A.7), we have, with probability (conditionally to $\{Y \in \mathcal{E}_\kappa\}$) going to 1 as $n \rightarrow \infty$,

$$\begin{aligned} & \sup_{|x - \widehat{x}_n| \geq \delta} (f_n(x) + g_n(x) + h_n(x) - f_n(\widehat{x}_n) - g_n(\widehat{x}_n) - h_n(\widehat{x}_n)) \\ & \leq -Bn + C(x_{sup} - x_{inf}) + o_{\mathbb{P}|Y \in \mathcal{E}_\kappa}(1) \xrightarrow{n \rightarrow \infty} -\infty. \end{aligned}$$

Finally, for all $\varepsilon > 0$ there exists $c > 0$ so that, with probability (conditionally to $\{Y \in \mathcal{E}_\kappa\}$) going to 1 as $n \rightarrow \infty$,

$$\sup_{|x - \widehat{x}_n| \geq \varepsilon/\sqrt{n}} (f_n(x) + g_n(x) + h_n(x) - f_n(\widehat{x}_n) - g_n(\widehat{x}_n) - h_n(\widehat{x}_n)) \leq -c.$$

Hence, we have, by definition of $\widehat{\widehat{x}}_n$

$$\sqrt{n}|\widehat{\widehat{x}}_n - \widehat{x}_n| = o_{\mathbb{P}|Y \in \mathcal{E}_\kappa}(1).$$

□

Lemma A.3. *Let $\{k_\theta; \theta \in \Theta\}$ be the set of covariance functions in Section 2 where Θ is compact. Assume that k_θ satisfies Condition-Var in the case $\kappa = 0$, where C and α can be chosen independently of θ . Let $Z_{n,\theta}$ be a Gaussian process with mean function zero and covariance function $(x_1, x_2) \mapsto \text{Cov}_\theta(Y(x_1), Y(x_2)|y)$. Then, we have*

$$\sup_{\theta \in \Theta} \mathbb{E} \left[\sup_{x \in [0,1]^d} |Z_{n,\theta}(x)| \right] \xrightarrow{n \rightarrow \infty} 0.$$

Proof of Lemma A.3. This result is proved as an intermediate result in the proof of Lemma A.3 in (López-Lopera et al., 2017). There, the result was for fixed θ , but it can be made uniform over $\theta \in \Theta$ with no additional difficulties. \square

B Proofs - Boundedness

We let $\kappa = 0$ throughout Section B.

B.1 Estimation of the variance parameter

Proof of Theorem 3.2 under boundedness constraints.

1) Let $m_{k,n} = \min_{i=1,\dots,k} y_i$, $M_{k,n} = \max_{i=1,\dots,k} y_i$, and $(m, M)^\top = (Y_*, Y^*)^\top$, where Y_* and Y^* have been defined in Section 2.2. We clearly have $m \leq m_{k,n} \leq M_{k,n} \leq M$. Since $(x_i)_{i \in \mathbb{N}}$ is dense, for any sequence $(k_n)_{n \in \mathbb{N}}$ so that $k_n \rightarrow \infty$ as $n \rightarrow \infty$ and $k_n \leq n$, we have $(m_{k_n,n}, M_{k_n,n})^\top \rightarrow (m, M)$ a.s. as $n \rightarrow \infty$ (up to reindexing x_1, \dots, x_n).

2) Let $k \in \mathbb{N}$ be fixed. We have

$$\sqrt{n} (\bar{\sigma}_n^2 - \sigma_0^2) = \frac{1}{\sqrt{n}} \left(y^\top R_1^{-1} y - n\sigma_0^2 \right).$$

Writing the Gaussian probability density function of y as the product of the conditional probability density functions of y_i given y_1, \dots, y_{i-1} leads to

$$\frac{1}{\sqrt{n}} \left(y^\top R_1^{-1} y - n\sigma_0^2 \right) = \frac{\sigma_0^2}{\sqrt{n}} \sum_{i=1}^n \left(\frac{(y_i - \mathbb{E}[y_i|y_1, \dots, y_{i-1}])^2}{\text{Var}(y_i|y_1, \dots, y_{i-1})} - 1 \right).$$

The terms in the sum above are independent. Indeed,

$$\text{Cov}(y_l, y_i - \mathbb{E}[y_i|y_1, \dots, y_{i-1}]) = 0, \quad \text{for any } l \leq i-1$$

and the Gaussianity then leads to independence. Therefore,

$$\begin{aligned} \frac{1}{\sqrt{n}} \left(y^\top R_1^{-1} y - n\sigma_0^2 \right) &= \frac{\sigma_0^2}{\sqrt{n}} \sum_{i=1}^k \left(\frac{(y_i - \mathbb{E}[y_i|y_1, \dots, y_{i-1}])^2}{\text{Var}(y_i|y_1, \dots, y_{i-1})} - 1 \right) \\ &\quad + \frac{\sigma_0^2}{\sqrt{n}} \sum_{i=k+1}^n \left(\frac{(y_i - \mathbb{E}[y_i|y_1, \dots, y_{i-1}])^2}{\text{Var}(y_i|y_1, \dots, y_{i-1})} - 1 \right). \end{aligned}$$

The first term is $o_{\mathbb{P}}(1)$ being the sum of k r.v.'s (whose variances are all equal to 2) divided by the square root of n . Because $\mathbb{P}_{\sigma^2}(\ell \leq \min_{i=1,\dots,k} y_i \leq \max_{i=1,\dots,k} y_i \leq u) > 0$, the first term is also $o_{\mathbb{P}}(1)$ conditionally to $\left\{ \ell \leq \min_{i=1,\dots,k} y_i \leq \max_{i=1,\dots,k} y_i \leq u \right\}$. The second term is equal to σ_0^2/\sqrt{n} times the sum of $n - k$ independent variables with zero mean and variance 2 and is also independent of y_1, \dots, y_k . Hence, from the central limit theorem and Slutsky's lemma (van der Vaart, 1998, Lemma 2.8), we obtain that

$$\frac{1}{\sqrt{n}} \left(y^\top R_1^{-1} y - n\sigma_0^2 \right) \xrightarrow[n \rightarrow \infty]{\mathcal{L}|y \in \mathcal{E}_{0,k}} \mathcal{N}(0, 2\sigma_0^4),$$

where $\mathcal{E}_{0,k} := \left\{ y : \ell \leq \min_{i=1,\dots,k} y_i \leq \max_{i=1,\dots,k} y_i \leq u \right\}$ and $\frac{\mathcal{L}|y \in \mathcal{E}_{0,k}}{n \rightarrow \infty}$ is defined similarly as $\frac{\mathcal{L}|Y \in \mathcal{E}_0}{n \rightarrow \infty}$.

3) Hence, for $x \in \mathbb{R}$, there exists a sequence $k_n \xrightarrow[n \rightarrow \infty]{} \infty$ satisfying $k_n = o(n)$ as $n \rightarrow \infty$ so that:

$$\mathbb{P} \left(\sqrt{n} (\bar{\sigma}_n^2 - \sigma_0^2) \leq x \mid \ell \leq \min_{i=1,\dots,k_n} y_i \leq \max_{i=1,\dots,k_n} y_i \leq u \right) \xrightarrow[n \rightarrow \infty]{} \mathbb{P}(V \leq x)$$

with $V \sim \mathcal{N}(0, 2\sigma_0^4)$. Therefore, from Lemma A.1,

$$\mathbb{P} \left(\sqrt{n} (\bar{\sigma}_n^2 - \sigma_0^2) \leq x \mid \ell \leq Y(x) \leq u, \forall x \in [0, 1]^d \right) \xrightarrow[n \rightarrow \infty]{} \mathbb{P}(V \leq x).$$

This concludes the proof. \square

Proof of Theorem 3.3 under boundedness constraints. We apply Lemma A.2 to the sequences of functions f_n , g_n and h_n defined by $f_n(\sigma^2) = \mathcal{L}_n(\sigma^2)$, $g_n(x) = A_n(\sigma^2)$, and $h_n(\sigma^2) = B_n(\sigma^2)$. Here we recall that for $\sigma^2 \in \Theta$,

$$A_n(\sigma^2) = -\log \mathbb{P}_{\sigma^2}(Y \in \mathcal{E}_0) \text{ and } B_n(\sigma^2) = \log \mathbb{P}_{\sigma^2}(Y \in \mathcal{E}_0 | y).$$

In order to apply Lemma A.2, we need to check that the conditions (A.4) to (A.7) hold.

1) By (3.3), one has

$$\mathcal{L}_n(\sigma^2) = -\frac{n}{2} \log 2\pi - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2} \ln(|R_1|) - \frac{1}{2\sigma^2} y^\top R_1^{-1} y.$$

Now $y^\top R_1^{-1} y$ is the square of the norm of a Gaussian vector with variance-covariance matrix $\sigma_0^2 I_n$, where I_n stands for the identity matrix of dimension n . Thus one way write $y^\top R_1^{-1} y$ as the sum of the squares of n independent and identically distributed r.v.'s ε_i , where ε_i is Gaussian distributed with mean 0 and variance σ_0^2 . We prove that (A.4) is satisfied. One may rewrite $\mathcal{L}_n(\sigma^2)$ as

$$\mathcal{L}_n(\sigma^2) = -\frac{n}{2} \log(2\pi) - \frac{1}{2} \log(|R_1|) + \frac{n}{2} f_{\sigma_0^2 + o_{\mathbb{P}}(1)}(\sigma^2), \quad (\text{B.1})$$

where the $o_{\mathbb{P}}(1)$ above does not depend on σ^2 and f_a has been introduced in Section 2.2. By a Taylor expansion and the definition of $\bar{\sigma}_n^2$, we have, with probability going to 1 as $n \rightarrow \infty$,

$$\begin{aligned}\mathcal{L}_n(\sigma^2) - \mathcal{L}_n(\bar{\sigma}_n^2) &= (\sigma^2 - \bar{\sigma}_n^2)\mathcal{L}'_n(\bar{\sigma}_n^2) + \frac{1}{2}(\sigma^2 - \bar{\sigma}_n^2)^2\mathcal{L}''_n(\bar{\sigma}_n^2) \\ &= \frac{n}{4}f''_{\sigma_0^2+o_{\mathbb{P}}(1)}(\bar{\sigma}_n^2)(\sigma^2 - \bar{\sigma}_n^2)^2 \\ &= \frac{n}{4}\left(\frac{1}{\bar{\sigma}_n^4} - 2\frac{\sigma_0^2 + o_{\mathbb{P}}(1)}{\bar{\sigma}_n^6}\right)(\sigma^2 - \bar{\sigma}_n^2)^2,\end{aligned}$$

with $\bar{\sigma}_n^2$ in the interval with endpoints σ^2 and $\bar{\sigma}_n^2$. Hence, non-random constants $A > 0$ and $\delta > 0$ exist for which (A.4) is satisfied.

2) Second, let us prove that (A.5) holds with the previous $\delta > 0$ and for some $B > 0$. From (B.1), $2\mathcal{L}_n/n + \log(2\pi) + (1/n)\log(|R_1|)$ converges uniformly on $[\sigma_l^2, \sigma_u^2]$ as n goes to infinity to $f_{\sigma_0^2}$. The function $f_{\sigma_0^2}$ attains its unique maximum at σ_0^2 , which implies the result since $\bar{\sigma}_n^2$ converges to σ_0^2 in probability. Hence (A.5) holds.

3) Now we consider (A.6). Let us introduce the Gaussian process Y_r with mean function zero and covariance function k_1 . Let $\sigma_1^2 \leq \sigma_2^2$. Then, one has:

$$\begin{aligned}\left| \exp\{-A_n(\sigma_1^2)\} - \exp\{-A_n(\sigma_2^2)\} \right| &= \left| \mathbb{P}(\sigma_1 Y_r \in \mathcal{E}_0) - \mathbb{P}(\sigma_2 Y_r \in \mathcal{E}_0) \right| \\ &\leq \mathbb{P}\left(\frac{u}{\sigma_2} \leq Y_r(x) \leq \frac{u}{\sigma_1}, \forall x \in [0, 1]^d\right) \\ &\quad + \mathbb{P}\left(\frac{\ell}{\sigma_2} \leq Y_r(x) \leq \frac{\ell}{\sigma_1}, \forall x \in [0, 1]^d\right) \\ &\leq c \left| \frac{1}{\sigma_1} - \frac{1}{\sigma_2} \right| \\ &\leq c|\sigma_2^2 - \sigma_1^2|\end{aligned}$$

by Tsirelson theorem in (Azaïs and Wschebor, 2009). Then, from Lemma 3.1, (A.6) holds.

4) We turn to

$$B_n(\sigma^2) = \log \mathbb{P}_{\sigma^2}(Y \in \mathcal{E}_0|y) = \log \mathbb{P}_{\sigma^2}(\ell \leq Y(x) \leq u, \forall x \in [0, 1]^d|y).$$

Let $m_{n,y}$ and $\sigma^2 k_n$ be the conditional mean and covariance functions of Y given y , under the probability measure \mathbb{P}_{σ^2} . Using Borell-TIS inequality (Adler and Taylor, 2007), with Z_{n,σ^2} a Gaussian process with mean function zero and covariance function $\sigma^2 k_n$, we obtain

$$\begin{aligned}\mathbb{P}_{\sigma^2}(Y^* > u|y) &\leq \mathbb{P}_{\sigma^2}\left(Z_{n,\sigma^2}^* > u - \sup_{x \in [0,1]^d} m_{n,y}(x)|y\right) \\ &\leq \mathbb{P}_{\sigma^2}\left(Z_{n,\sigma^2}^{**} > u - \sup_{x \in [0,1]^d} m_{n,y}(x)|y\right) \\ &\leq \exp \left\{ - \frac{\left(\left(u - \sup_{x \in [0,1]^d} m_{n,y}(x) - \mathbb{E}[Z_{n,\sigma^2}^{**}] \right)_+ \right)^2}{2 \sup_{x \in [0,1]^d} \mathbb{E}[Z_{n,\sigma^2}(x)^2]} \right\}.\end{aligned}\tag{B.2}$$

But by Lemma A.3, $\sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathbb{E}[Z_{n,\sigma^2}^{**}] \rightarrow 0$ as $n \rightarrow +\infty$. Additionally, one can simply show that $\sup_{x \in [0,1]^d} \mathbb{E}[Z_{n,\sigma^2}(x)^2] = \sup_{x \in [0,1]^d} \sigma^2 k_n(x, x)$ goes to zero uniformly in $\sigma^2 \in [\sigma_l^2, \sigma_u^2]$ as $n \rightarrow \infty$. By (Bect et al., 2016, Proposition 2.8) and because the sequence of observation points is dense,

$$\sup_{x \in [0,1]^d} |m_{n,y}(x) - Y(x)| \xrightarrow[n \rightarrow +\infty]{\text{a.s.}} 0$$

from which we deduce that on $\{Y^* < u - \delta\}$, a.s.

$$\limsup_{n \rightarrow +\infty} \left(u - \sup_{x \in [0,1]^d} m_{n,y}(x) \right) \geq \delta.$$

Consequently, (B.2) leads to

$$\mathbb{1}_{\{Y^* < u - \delta\}} \sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathbb{P}_{\sigma^2}(Y^* > u | y) \xrightarrow[n \rightarrow +\infty]{\text{a.s.}} 0. \quad (\text{B.3})$$

Similarly, taking $-Y$ instead of Y , one may prove easily that

$$\mathbb{1}_{\{Y_* > l + \delta\}} \sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathbb{P}_{\sigma^2}(Y_* < l | y) \xrightarrow[n \rightarrow +\infty]{\text{a.s.}} 0. \quad (\text{B.4})$$

Then, we deduce that

$$\mathbb{1}_{\{\ell + \delta < Y(x) < u - \delta, \forall x \in [0,1]^d\}} \sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathbb{P}_{\sigma^2}(Y^* > u \text{ or } Y_* < \ell | y) \xrightarrow[n \rightarrow +\infty]{\text{a.s.}} 0. \quad (\text{B.5})$$

Now let $\varepsilon > 0$, $\varepsilon' = 2|\log(1 - \varepsilon)|$ and $\mathcal{E}_{0,\delta} := \{f \in \mathcal{C}([0,1]^d, \mathbb{R}) \text{ s.t. } \ell + \delta \leq f(x) \leq u - \delta, \forall x \in [0,1]^d\}$. We have:

$$\begin{aligned} & \mathbb{P}\left(\sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathbb{P}_{\sigma^2}(Y^* > u \text{ or } Y_* < \ell | y) \geq \varepsilon, Y \in \mathcal{E}_{0,\delta}\right) \xrightarrow[n \rightarrow +\infty]{} 0 \\ &= \mathbb{P}\left(\inf_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} B_n(\sigma^2) \leq -\varepsilon'/2, Y \in \mathcal{E}_{0,\delta}\right) \\ &= \mathbb{P}\left(\sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} |B_n(\sigma^2)| \geq \varepsilon'/2, Y \in \mathcal{E}_{0,\delta}\right) \\ &\geq \mathbb{P}\left(\sup_{\sigma_1^2, \sigma_2^2 \in [\sigma_l^2, \sigma_u^2]} |B_n(\sigma_1^2) - B_n(\sigma_2^2)| \geq \varepsilon', Y \in \mathcal{E}_{0,\delta}\right) \end{aligned}$$

by the triangular inequality and (B.5). Therefore,

$$\begin{aligned} & \mathbb{P}\left(\sup_{\sigma_1^2, \sigma_2^2 \in [\sigma_l^2, \sigma_u^2]} |B_n(\sigma_1^2) - B_n(\sigma_2^2)| \geq \varepsilon', Y \in \mathcal{E}_0\right) \\ &= \mathbb{P}\left(\sup_{\sigma_1^2, \sigma_2^2 \in [\sigma_l^2, \sigma_u^2]} |B_n(\sigma_1^2) - B_n(\sigma_2^2)| \geq \varepsilon', Y \in \mathcal{E}_{0,\delta}\right) \end{aligned} \quad (\text{B.6})$$

$$+ \mathbb{P}\left(\sup_{\sigma_1^2, \sigma_2^2 \in [\sigma_l^2, \sigma_u^2]} |B_n(\sigma_1^2) - B_n(\sigma_2^2)| \geq \varepsilon', Y \in \mathcal{E}_0 \setminus \mathcal{E}_{0,\delta}\right). \quad (\text{B.7})$$

As already shown, the term (B.6) converges to 0 as $n \rightarrow +\infty$ for any fixed $\delta > 0$. For (B.7), we have

$$\sup_{\substack{t_1, t_2 \in \mathbb{R} \\ t_1 \neq t_2}} \frac{|\mathbb{P}_{\sigma_0^2}(Y^* \leq t_1) - \mathbb{P}_{\sigma_0^2}(Y^* \leq t_2)|}{|t_1 - t_2|} < +\infty.$$

This follows from Tsirelson theorem in (Azaïs and Wschebor, 2009). Hence for all $\varepsilon > 0$, there exists $\delta^* > 0$ such that,

$$\mathbb{P}_{\sigma_0^2}(Y^* \in [u - \delta^*, u]) \leq \varepsilon. \quad (\text{B.8})$$

Similarly, for all $\varepsilon > 0$, there exists $\delta_* > 0$ such that,

$$\mathbb{P}_{\sigma_0^2}(Y_* \in [\ell + \delta_*, \ell]) \leq \varepsilon. \quad (\text{B.9})$$

Taking $\delta = \min(\delta_*, \delta^*)$, we conclude the proof of (A.7).

5) Finally, we remark that with probability going to one as $n \rightarrow \infty$, $\bar{\sigma}_n^2 = \operatorname{argmax}_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathcal{L}_n(\sigma^2)$.

Hence, one may apply Lemma A.2 to obtain

$$\sqrt{n}|\hat{\sigma}_{n,c}^2 - \bar{\sigma}_n^2| = o_{\mathbb{P}|Y \in \mathcal{E}_0}(1).$$

By Theorem 3.2 and Slutsky's lemma, we conclude the proof. \square

B.2 Isotropic Matérn process

Before proving Theorems 4.2, 4.3 and 4.4, we establish an intermediate result useful in the sequel.

Lemma B.1. *For $\rho \in [\rho_l, \rho_u]$, let*

$$\bar{\sigma}_n^2(\rho) \in \operatorname{argmax}_{\sigma^2 \in (0, \infty)} \mathcal{L}_n(\sigma^2, \rho)$$

and

$$\hat{\sigma}_n^2(\rho) \in \operatorname{argmax}_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathcal{L}_n(\sigma^2, \rho).$$

Then, we have

$$\sup_{\rho_1, \rho_2 \in [\rho_l, \rho_u]} \left| \frac{\bar{\sigma}_n^2(\rho_1)}{\rho_1^{2\nu}} - \frac{\bar{\sigma}_n^2(\rho_2)}{\rho_2^{2\nu}} \right| = o_{\mathbb{P}}(1/\sqrt{n}) \quad (\text{B.10})$$

and

$$\sup_{\rho_1, \rho_2 \in [\rho_l, \rho_u]} \left| \frac{\hat{\sigma}_n^2(\rho_1)}{\rho_1^{2\nu}} - \frac{\hat{\sigma}_n^2(\rho_2)}{\rho_2^{2\nu}} \right| = o_{\mathbb{P}}(1/\sqrt{n}). \quad (\text{B.11})$$

Proof of Lemma B.1. We have $\bar{\sigma}_n^2(\rho_1)/(\rho_1^{2\nu}) = (1/n)(1/(\rho_1^{2\nu}))y^\top R_{\rho_1, \nu}^{-1}y$ so that from Lemma 1 in (Kaufman and Shaby, 2013), we get $\frac{\bar{\sigma}_n^2(\rho_1)}{\rho_1^{2\nu}} \geq \frac{\bar{\sigma}_n^2(\rho_2)}{\rho_2^{2\nu}}$ for $\rho_1 \leq \rho_2$. Thus,

$$\sup_{\rho_1, \rho_2 \in [\rho_l, \rho_u]} \left| \frac{\bar{\sigma}_n^2(\rho_1)}{\rho_1^{2\nu}} - \frac{\bar{\sigma}_n^2(\rho_2)}{\rho_2^{2\nu}} \right| = \left| \frac{\bar{\sigma}_n^2(\rho_l)}{\rho_l^{2\nu}} - \frac{\bar{\sigma}_n^2(\rho_u)}{\rho_u^{2\nu}} \right|.$$

Then, it is shown in the proof of Theorem 5 in (Du et al., 2009) that

$$\left| \frac{\bar{\sigma}_n^2(\rho_l)}{\rho_l^{2\nu}} - \frac{\bar{\sigma}_n^2(\rho_0)}{\rho_0^{2\nu}} \right| = o_{\mathbb{P}}(1/\sqrt{n})$$

and similarly for ρ_u . Hence, (B.10) follows. Also, let $p_{\sigma_l^2, \sigma_u^2}$ be the function from $(0, \infty)$ to $[\sigma_l^2, \sigma_u^2]$ defined by $p_{\sigma_l^2, \sigma_u^2}(t) = \min(\max(t, \sigma_l^2), \sigma_u^2)$. Then, since $\sigma^2 \mapsto \mathcal{L}_n(\sigma^2, \rho)$ is first increasing and then decreasing, we have $\hat{\sigma}_n^2(\rho) = p_{\sigma_l^2, \sigma_u^2}(\bar{\sigma}_n^2(\rho))$. Notice that $p_{\sigma_l^2, \sigma_u^2}$ is continuous and bounded by σ_u^2 . Hence, (B.11) follows. \square

Proof of Theorem 4.2 under boundedness constraints. Because $\sigma_l^2/(\rho_l^{2\nu}) < \sigma_0^2/(\rho_0^{2\nu}) < \sigma_u^2/(\rho_u^{2\nu})$, we have $\hat{\sigma}_n^2 = \hat{\sigma}_n^2(\hat{\rho}_n)$ with the notation of Lemma B.1. Also, with probability going to 1 as $n \rightarrow \infty$, $\bar{\sigma}_n^2(\hat{\rho}_n) = \bar{\sigma}_n^2(\hat{\rho}_n)$ with the notation of Lemma B.1. From Lemma B.1 and with the notation therein, we have

$$\begin{aligned} \sqrt{n} \left(\frac{\hat{\sigma}_n^2(\hat{\rho}_n)}{\hat{\rho}_n^{2\nu}} - \frac{\sigma_0^2}{\rho_0^{2\nu}} \right) &= \sqrt{n} \left(\frac{\hat{\sigma}_n^2(\rho_0)}{\rho_0^{2\nu}} - \frac{\sigma_0^2}{\rho_0^{2\nu}} \right) + \sqrt{n} \left(\frac{\hat{\sigma}_n^2(\hat{\rho}_n)}{\hat{\rho}_n^{2\nu}} - \frac{\hat{\sigma}_n^2(\rho_0)}{\rho_0^{2\nu}} \right) \\ &= \frac{1}{\sqrt{n}} \frac{1}{\rho_0^{2\nu}} \left(y^\top R_{\rho_0, \nu}^{-1}y - n\sigma_0^2 \right) + o_{\mathbb{P}}(1), \end{aligned}$$

where we have used that, with probability going to 1 as $n \rightarrow \infty$, $\hat{\sigma}_n^2(\rho_0) = \bar{\sigma}_n^2(\rho_0)$. Then we conclude by applying Theorem 3.2 when $\kappa = 0$ and with $K_0 = K_\nu(\cdot/\rho_0)$. \square

Proof of Theorem 4.3 under boundedness constraints. We apply Lemma A.2 to the sequences of functions f_n , g_n and h_n defined by $f_n(\sigma^2) = \mathcal{L}_n(\sigma^2, \rho_1)$, $g_n(x) = A_n(\sigma^2, \rho_1)$, and $h_n(\sigma^2) = B_n(\sigma^2, \rho_1)$.

1) We have, with σ_1^2 so that $\sigma_1^2/\rho_1^{2\nu} = \sigma_0^2/\rho_0^{2\nu}$,

$$\begin{aligned} \mathcal{L}_n(\sigma^2, \rho_1) &= -\frac{n}{2} \log(2\pi) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2} \ln(|R_{\rho_1, \nu}|) - \frac{1}{2\sigma^2} y^\top R_{\rho_1, \nu}^{-1}y \\ &= -\frac{n}{2} \log(2\pi) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2} \ln(|R_{\rho_1, \nu}|) - \frac{1}{2\sigma^2} \left[\frac{\rho_1^{2\nu}}{\rho_0^{2\nu}} y^\top R_{\rho_0, \nu}^{-1}y + o_{\mathbb{P}}(\sqrt{n}) \right] \end{aligned}$$

from (B.10) in Lemma B.1, observing that $\bar{\sigma}_n^2(\rho_1) = (1/n)y^\top R_{\rho_1, \nu}^{-1}y$. Thus

$$\mathcal{L}_n(\sigma^2, \rho_1) = -\frac{n}{2} \log(2\pi) - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2} \ln(|R_{\rho_1, \nu}|) - \frac{\sigma_1^2}{2\sigma^2} \left[\frac{1}{\sigma_0^2} y^\top R_{\rho_0, \nu}^{-1}y + o_{\mathbb{P}}(\sqrt{n}) \right],$$

where $(1/\sigma_0^2)y^\top R_{\rho_0, \nu}^{-1}y$ is a sum of the squares of independent standard Gaussian variables. Hence, we show (A.4) and (A.5) exactly as for the proof of Theorem 3.3 when $\kappa = 0$.

2) Assumption (A.6) is satisfied since it has been established in the proof of Theorem 3.3 when $\kappa = 0$ (for any $\rho_0 \in (0, \infty)$) and does not involve y .

3) We turn to $B_n(\sigma^2, \rho_1) = \log \mathbb{P}_{(\sigma^2, \rho_1)}(Y \in \mathcal{E}_0 | y)$. Similarly to the proof of Theorem 3.3 when $\kappa = 0$, we have that, for any $\delta > 0$:

$$\mathbb{P}_{(\sigma_1^2, \rho_1)} \left(\mathbb{1}_{\{\ell + \delta \leq Y(x) \leq u - \delta, \forall x \in [0, 1]^d\}} \sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} |B_n(\sigma^2, \rho_1)| \xrightarrow[n \rightarrow \infty]{} 0 \right) = 1. \quad (\text{B.12})$$

Now, for $(\sigma^2, \rho) \in (0, \infty)^2$, we recall that $\mathbb{P}_{(\sigma^2, \rho)}$ is the measure on Ω for which $Y : (\Omega, \mathcal{A}, \mathbb{P}_{\sigma^2, \rho}) \rightarrow (\mathcal{C}([0, 1]^d), \mathcal{B})$ has the distribution of a Gaussian process on $[0, 1]^d$ with mean function zero and covariance function $k_{\sigma^2, \rho, \nu}$. By Theorem 2 in (Zhang, 2004), the measures $\mathbb{P}_{(\sigma_0^2, \rho_0)}$ and $\mathbb{P}_{(\sigma_1^2, \rho_1)}$ are equivalent as soon as $\sigma_0^2 / \rho_0^{2\nu} = \sigma_1^2 / \rho_1^{2\nu}$ meaning that for any set $A \in \mathcal{A}$,

$$\mathbb{P}_{(\sigma_0^2, \rho_0)}(A) = 1 \Leftrightarrow \mathbb{P}_{(\sigma_1^2, \rho_1)}(A) = 1.$$

Then, one gets

$$\mathbb{P}_{(\sigma_0^2, \rho_0)} \left(\mathbb{1}_{\{\ell + \delta \leq Y(x) \leq u - \delta, \forall x \in [0, 1]^d\}} \sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} |B_n(\sigma^2, \rho_1)| \xrightarrow[n \rightarrow \infty]{} 0 \right) = 1$$

which can also be written as

$$\mathbb{1}_{\{\ell + \delta \leq Y(x) \leq u - \delta, \forall x \in [0, 1]^d\}} \sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} |B_n(\sigma^2, \rho_1)| \xrightarrow[n \rightarrow \infty]{\text{a.s.}} 0.$$

This implies (A.7), as for the proof of Theorem 3.3 when $\kappa = 0$.

4) From a special case of Theorem 4.2 when $\kappa = 0$ and with $\rho_l = \rho_u = \rho_1$, we have

$$\sqrt{n} (\hat{\sigma}_n^2(\rho_1) - \sigma_1^2) \xrightarrow[n \rightarrow +\infty]{\mathcal{L} | Y \in \mathcal{E}_0} \mathcal{N}(0, 2\sigma_1^4).$$

Therefore, by Lemma A.2 and Slutsky's lemma, we conclude the proof of Theorem 3.3 when $\kappa = 0$. \square

Proof of Theorem 4.4 under boundedness constraints. Let $\kappa = 0$ in this proof. We apply Lemma A.2 to the sequences of functions f_n , g_n and h_n defined by $f_n(x) = \mathcal{L}_n(x \hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c})$, $g_n(x) = A_n(x \hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c})$, and $h_n(x) = B_n(x \hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c})$.

1) We naturally have

$$\frac{\hat{\sigma}_{n,c}^2}{\hat{\rho}_{n,c}^{2\nu}} \in \operatorname{argmax}_{x \in [\sigma_l^2 / \hat{\rho}_{n,c}^{2\nu}, \sigma_u^2 / \hat{\rho}_{n,c}^{2\nu}]} \mathcal{L}_{n,c}(x \hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c}^{2\nu}).$$

Also, we have

$$\begin{aligned} \mathcal{L}_n(x \hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c}) &= -\frac{n}{2} \log(2\pi) - \frac{n}{2} \ln(x \hat{\rho}_{n,c}^{2\nu}) - \frac{1}{2} \ln(|R_{\hat{\rho}_{n,c}, \nu}|) - \frac{1}{2x \hat{\rho}_{n,c}^{2\nu}} y^\top R_{\hat{\rho}_{n,c}, \nu}^{-1} y \\ &= -\frac{n}{2} \log(2\pi) - \frac{n}{2} \ln(\hat{\rho}_{n,c}^{2\nu}) - \frac{1}{2} \ln(|R_{\hat{\rho}_{n,c}, \nu}|) - \frac{n}{2} \ln(x) - \frac{1}{2x} \left[\frac{y^\top R_{\rho_0, \nu}^{-1} y}{\rho_0^{2\nu}} + o_{\mathbb{P}}(\sqrt{n}) \right] \end{aligned}$$

from (B.10) in Lemma B.1, observing that $\bar{\sigma}_n^2(\rho) = (1/n)y^\top R_{\rho,\nu}^{-1}y$. Thus

$$\mathcal{L}_n(x\hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c}) = -\frac{n}{2}\log(2\pi) - \frac{n}{2}\ln(\hat{\rho}_{n,c}^{2\nu}) - \frac{1}{2}\ln(|R_{\hat{\rho}_{n,c},\nu}|) + \frac{n}{2}f_{(\sigma_0^2/\rho_0^{2\nu})+V_n+o_{\mathbb{P}}(1/\sqrt{n})}(x), \quad (\text{B.13})$$

where $\sqrt{n}V_n$ converges in distribution to a centered Gaussian distribution with variance $2(\sigma_0^2/\rho_0^{2\nu})^2$. Here V_n and the above $o_{\mathbb{P}}(1/\sqrt{n})$ do not depend on x . Hence, we show (A.4) and (A.5) exactly as for the proof of Theorem 3.3 when $\kappa = 0$.

2) One can also see that $\delta > 0$ can be chosen so that $\sigma_l^2/\rho_l^{2\nu} + \delta \leq \sigma_0^2/\rho_0^{2\nu} \leq \sigma_u^2/\rho_u^{2\nu} - \delta$. Furthermore, let $x_{inf} = \sigma_l^2/\rho_l^{2\nu}$ and $x_{sup} = \sigma_u^2/\rho_u^{2\nu}$. Then, from (B.13), one can show that with

$$\frac{\tilde{\sigma}_n^2(\hat{\rho}_{n,c})}{\hat{\rho}_{n,c}^{2\nu}} \in \underset{x \in [x_{inf}, x_{sup}]}{\operatorname{argmax}} \mathcal{L}_n(x\hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c}) \quad (\text{B.14})$$

we have:

$$\frac{\tilde{\sigma}_n^2(\hat{\rho}_{n,c})}{\hat{\rho}_{n,c}^{2\nu}} = \frac{\hat{\sigma}_n^2(\hat{\rho}_{n,c})}{\hat{\rho}_{n,c}^{2\nu}}, \quad (\text{B.15})$$

with probability going to 1 as $n \rightarrow \infty$. It is convenient to introduce $\tilde{\sigma}_n^2(\hat{\rho}_{n,c})$ because this yields a non-random optimization domain in (B.14). Hence, from Theorem 4.2 when $\kappa = 0$ and from Lemma B.1,

$$\sqrt{n} \left(\frac{\tilde{\sigma}_n^2(\hat{\rho}_{n,c})}{\hat{\rho}_{n,c}^{2\nu}} - \frac{\sigma_0^2}{\rho_0^{2\nu}} \right) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}|Y \in \mathcal{E}_0} \mathcal{N} \left(0, 2 \left(\frac{\sigma_0^2}{\rho_0^{2\nu}} \right)^2 \right).$$

Let also

$$\frac{\tilde{\sigma}_{n,c}^2(\hat{\rho}_{n,c})}{\hat{\rho}_{n,c}^{2\nu}} \in \underset{x \in [x_{inf}, x_{sup}]}{\operatorname{argmax}} \mathcal{L}_{n,c}(x\hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c}).$$

Then, if we show (A.6) and (A.7), we can show similarly as for (B.15) that:

$$\frac{\tilde{\sigma}_{n,c}^2(\hat{\rho}_{n,c})}{\hat{\rho}_{n,c}^{2\nu}} = \frac{\hat{\sigma}_{n,c}^2}{\hat{\rho}_{n,c}^{2\nu}},$$

with probability going to 1 as $n \rightarrow \infty$. Hence, from Lemmas A.2 and B.1 and Slutsky's lemma, we can obtain, if (A.6) and (A.7) hold,

$$\sqrt{n} \left(\frac{\hat{\sigma}_{n,c}^2}{\hat{\rho}_{n,c}^{2\nu}} - \frac{\sigma_0^2}{\rho_0^{2\nu}} \right) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}|Y \in \mathcal{E}_0} \mathcal{N} \left(0, 2 \left(\frac{\sigma_0^2}{\rho_0^{2\nu}} \right)^2 \right).$$

Therefore, in order to conclude the proof, it is sufficient to prove (A.6) and (A.7).

3) We turn to (A.6). Let σY_ρ be a Gaussian process with mean function zero and covariance function $k_{\sigma^2, \rho, \nu}$. Then we have, from Lemma 4.1,

$$\begin{aligned} |g_n(x_1) - g_n(x_2)| &\leq c \left| \mathbb{P}(x_1 \hat{\rho}_{n,c}^{2\nu} Y_{\hat{\rho}_{n,c}} \in \mathcal{E}_0) - \mathbb{P}(x_2 \hat{\rho}_{n,c}^{2\nu} Y_{\hat{\rho}_{n,c}} \in \mathcal{E}_0) \right| \\ &\leq c |x_1 - x_2| \sup_{\substack{\rho \in [\rho_l, \rho_u] \\ t_1, t_2 \in [l/(x_{sup} \rho_u^{2\nu}), u/(x_{inf} \rho_l^{2\nu})] \\ t_1 \neq t_2}} \frac{|\mathbb{P}(Y_\rho^* \leq t_1) - \mathbb{P}(Y_\rho^* \leq t_2)|}{|t_1 - t_2|}. \end{aligned}$$

We introduce the following notation:

$$F_\rho(t) = \mathbb{P}\left(\sup_{x \in [0,1]^d} Y_\rho(x) \leq t\right)$$

and assume that

$$\sup_{\substack{\rho \in [\rho_l, \rho_u] \\ t \in [l/(x_{sup} \rho_u^{2\nu}), u/(x_{inf} \rho_l^{2\nu})]}} \sup_{\varepsilon > 0} \frac{F_\rho(t + \varepsilon) - F_\rho(t)}{\varepsilon} = +\infty.$$

Therefore, there exists a sequence $(\rho_k, t_k, \varepsilon_k)_{k \in \mathbb{N}}$ such that

$$\frac{F_{\rho_k}(t_k + \varepsilon_k) - F_{\rho_k}(t_k)}{\varepsilon_k} \xrightarrow{k \rightarrow +\infty} +\infty. \quad (\text{B.16})$$

We extract from $(\rho_k, t_k, \varepsilon_k)_{k \in \mathbb{N}}$ a subsequence (still denoted $(\rho_k, t_k, \varepsilon_k)_{k \in \mathbb{N}}$) such that $(\rho_k)_k$ is convergent and we denote by $\bar{\rho}$ its limit. Let Φ be the cumulative distribution function of a standard Gaussian random variable. Then by the mean value theorem,

$$\frac{\Phi^{-1} \circ F_{\rho_k}(t_k + \varepsilon_k) - \Phi^{-1} \circ F_{\rho_k}(t_k)}{\varepsilon_k} \geq \frac{F_{\rho_k}(t_k + \varepsilon_k) - F_{\rho_k}(t_k)}{\varepsilon_k} \inf_{p \in [0,1]} (\Phi^{-1})'(p) \xrightarrow{k \rightarrow +\infty} +\infty$$

noticing that $\inf_{p \in [0,1]} (\Phi^{-1})'(p) > 0$ and using (B.16).

But, using the concavity of $\Phi^{-1} \circ F_\rho$ (see Theorem 10 in (Lifshits, 1995, Section 11)), one gets

$$\begin{aligned} \frac{\Phi^{-1} \circ F_{\rho_k}(t_k + \varepsilon_k) - \Phi^{-1} \circ F_{\rho_k}(t_k)}{\varepsilon_k} &\leq \Phi^{-1} \circ F_{\rho_k} \left(\frac{l}{x_{sup} \rho_u^{2\nu}} \right) - \Phi^{-1} \circ F_{\rho_k} \left(\frac{l}{x_{sup} \rho_u^{2\nu}} - 1 \right) \\ &\xrightarrow{k \rightarrow \infty} \Phi^{-1} \circ F_{\bar{\rho}} \left(\frac{l}{x_{sup} \rho_u^{2\nu}} \right) - \Phi^{-1} \circ F_{\bar{\rho}} \left(\frac{l}{x_{sup} \rho_u^{2\nu}} - 1 \right). \end{aligned}$$

The convergence comes from the continuity of the function $\rho \mapsto F_\rho(t)$ for a fixed t (see the proof of lemma A.6 in (López-Lopera et al., 2017)). From Lemma 4.1, the above limit is finite, which is contradictory with (B.16). Hence, (A.6) is proved.

4) Finally, we turn to (A.7). We let $m_{n,\rho,y}$ and $\sigma^2 k_{n,\rho}$ be the mean and covariance functions of Y given y under covariance function $k_{\sigma^2, \rho, \nu}$. Our first aim is to show that, for any $\varepsilon > 0$, with probability going to 1 as $n \rightarrow \infty$,

$$\sup_{\rho \in [\rho_l, \rho_u]} \sup_{x \in [0,1]^d} (m_{n,\rho,y}(x) - Y^*) \leq \varepsilon \quad (\text{B.17})$$

and

$$\sup_{\rho \in [\rho_l, \rho_u]} \sup_{x \in [0,1]^d} (Y_* - m_{n,\rho,y}(x)) \leq \varepsilon. \quad (\text{B.18})$$

Now we use tools from the theory of reproducing kernel Hilbert spaces (RKHSs) and refer to, e.g., (Wendland, 2004) for the definitions and properties of RKHSs used in the rest of the proof. For $\rho \in [\rho_l, \rho_u]$, the function $m_{n,\rho,y}$ belongs to the RKHS of the covariance function $k_{1,\rho,\nu}$. Its RKHS norm $\|m_{n,\rho,y}\|_{k_{1,\rho,\nu}}$ can be simply shown to satisfy

$$\|m_{n,\rho,y}\|_{k_{1,\rho,\nu}}^2 = y^\top R_{\rho,\nu}^{-1} y.$$

Hence, from Lemma B.1, observing that $\bar{\sigma}_n^2(\rho) = (1/n)y^\top R_{\rho,\nu}^{-1} y$, we have, with probability going to 1 as $n \rightarrow \infty$,

$$\sup_{\rho \in [\rho_l, \rho_u]} \|m_{n,\rho,y}\|_{k_{1,\rho,\nu}} \leq c\sqrt{n}. \quad (\text{B.19})$$

Consider the case a). Since $\nu > 1$, the covariance function $k_{1,\rho,\nu}$ is twice continuously differentiable on $[0,1]$. Hence, we have from Theorem 1 in (Zhou, 2008),

$$\sup_{\rho \in [\rho_l, \rho_u]} \sup_{x \in [0,1]} |m'_{n,\rho,y}(x)| \leq c \sup_{\rho \in [\rho_l, \rho_u]} \|m_{n,\rho,y}\|_{k_{1,\rho,\nu}} \leq c\sqrt{n}$$

with probability going to 1 as $n \rightarrow \infty$. Hence, since for $i = 1, \dots, n$ $m_{n,\rho,y}(x_i) = y_i \leq Y^*$, and from the assumption $\max_{x \in [0,1]} \min_{i=1,\dots,n} |x - x_i| = o(1/\sqrt{n})$, it follows that (B.17) holds. Similarly, one can show that, with probability going to 1 as $n \rightarrow \infty$, (B.18) holds.

Consider the case b). Since $\nu > 2$, the covariance function $k_{1,\rho,\nu}$ is four times continuously differentiable on $[0,1]^d$. Hence, we have also from Theorem 1 in (Zhou, 2008),

$$\sup_{\rho \in [\rho_l, \rho_u]} \max_{i,j=1,\dots,d} \sup_{x \in [0,1]^d} \left| \frac{\partial^2 m_{n,\rho,y}}{\partial x_i \partial x_j}(x) \right| \leq c \sup_{\rho \in [\rho_l, \rho_u]} \|m_{n,\rho,y}\|_{k_{1,\rho,\nu}} \leq c\sqrt{n}, \quad (\text{B.20})$$

with probability going to 1 as $n \rightarrow \infty$. For $\varepsilon > 0$, consider the event

$$\left\{ \sup_{\rho \in [\rho_l, \rho_u]} \sup_{x \in [0,1]^d} (m_{n,\rho,y}(x) - Y^*) \geq \varepsilon \right\}. \quad (\text{B.21})$$

Then, there exists $\bar{\rho} \in [\rho_l, \rho_u]$ and $\bar{x} \in [0,1]^d$ for which $m_{n,\bar{\rho},y}(\bar{x}) \geq Y^* + \varepsilon$. Let $H_x m_{n,\bar{\rho},y}$ be the Hessian matrix of $m_{n,\bar{\rho},y}$ at x and $\|H_x m_{n,\bar{\rho},y}\|$ be its largest singular value.

- For $d = 1$, we consider $\{v_1, v_2\} \subset \{x_1, \dots, x_n\}$ for which $v_1 \leq \bar{x} \leq v_2$ with $|v_1 - v_2| \leq 2a_n$ (the existence is assumed in the case b)). Then, necessarily $v_1 < \bar{x} < v_2$. Because, $m_{n,\bar{\rho},y}(v_1) \leq \max_{i=1,\dots,n} y_i \leq Y^*$, it follows that there exists $w_1 \in [v_1, \bar{x}]$ for which $m'_{n,\bar{\rho},y}(w_1) \geq \varepsilon/a_n$. Similarly there exists $w_2 \in [\bar{x}, v_2]$ for which $m'_{n,\bar{\rho},y}(w_2) \leq -\varepsilon/a_n$. Hence, there exists $w_3 \in [v_1, v_2]$ for which $m''_{n,\bar{\rho},y}(w_3) \leq -\varepsilon/a_n^2$ so that $\sup_{x \in [0,1]} \|H_x m_{n,\bar{\rho},y}\| \geq c\varepsilon/a_n^2$.

- For $d = 2$, we consider $\{v_1, v_2, v_3\} \subset \{x_1, \dots, x_n\}$ for which \bar{x} belongs to the convex hull of v_1, v_2, v_3 . Then, if \bar{x} belongs to one of the three segments with end points v_1, v_2 or

v_2, v_3 or v_1, v_3 , from the previous step with $d = 1$, it follows that $\sup_{x \in [0,1]^2} \|H_x m_{n,\bar{\rho},y}\| \geq c\varepsilon/a_n^2$.

Consider now that \bar{x} does not belong to one of these segments and consider the (unique) intersection point r of the line with direction $v_1 - \bar{x}$ and of the segment with endpoints v_2 and v_3 . If $m_{n,\bar{\rho},y}(r) \leq Y^* + \varepsilon/2$, by considering the triplet (v_1, \bar{x}, r) , from the reasoning of the case $d = 1$, it follows that $\sup_{x \in [0,1]^2} \|H_x m_{n,\bar{\rho},y}\| \geq c\varepsilon/a_n^2$. If $m_{n,\bar{\rho},y}(r) \geq Y^* + \varepsilon/2$, by considering

the triplet (v_2, r, v_3) , it also follows that $\sup_{x \in [0,1]^2} \|H_x m_{n,\bar{\rho},y}\| \geq c\varepsilon/a_n^2$.

• For $d = 3$, we consider $\{v_1, v_2, v_3, v_4\} \subset \{x_1, \dots, x_n\}$ for which \bar{x} belongs to the convex hull of v_1, v_2, v_3, v_4 . Let $\text{ch}(z_1, z_2, z_3)$ be the convex hull of $z_1, z_2, z_3 \in [0,1]^d$ (a two-dimensional triangle). If \bar{x} belongs to one of the four triangles $\text{ch}(v_1, v_2, v_3)$, $\text{ch}(v_1, v_2, v_4)$, $\text{ch}(v_1, v_3, v_4)$, $\text{ch}(v_2, v_3, v_4)$, then from the previous step with $d = 2$, it follows that

$\sup_{x \in [0,1]^3} \|H_x m_{n,\bar{\rho},y}\| \geq c\varepsilon/a_n^2$. Now if \bar{x} does not belong to one of these triangles, then there

exists a plane P_l containing \bar{x} , intersecting $\text{ch}(v_1, v_2, v_3)$, $\text{ch}(v_1, v_2, v_4)$, $\text{ch}(v_1, v_3, v_4)$, and being parallel to $\text{ch}(v_2, v_3, v_4)$. Let E be the intersection of this plane P_l and of $\text{ch}(v_1, v_2, v_3) \cup \text{ch}(v_1, v_2, v_4) \cup \text{ch}(v_1, v_3, v_4)$. If there exists $x \in E$ so that $m_{n,y,\bar{\rho}}(x) \geq Y^* + \varepsilon/2$, then from the previous step with $d = 2$, it follows that $\sup_{x \in [0,1]^3} \|H_x m_{n,\bar{\rho},y}\| \geq c\varepsilon/a_n^2$. If for all $x \in E$,

$m_{n,y,\bar{\rho}}(x) \leq Y^* + \varepsilon/2$, then there exists $z_1, z_2, z_3 \in E$ so that $\bar{x} \in \text{ch}(z_1, z_2, z_3)$ and hence we obtain $\sup_{x \in [0,1]^3} \|H_x m_{n,\bar{\rho},y}\| \geq c\varepsilon/a_n^2$.

Hence eventually, in all the configurations of the case b), we have $\sup_{x \in [0,1]^d} \|H_x m_{n,\bar{\rho},y}\| \geq c\varepsilon/a_n^2$, under the event (B.21). Hence, from (B.20), (B.17) follows in the case b). Analogously, (B.18) holds in that case.

Similarly to the proof of (A.7) in the proof of Theorem 3.3 when $\kappa = 0$, one can show that $\sup_{(\sigma^2, \rho) \in \bar{\Theta}} \mathbb{E}[Z_{n,\sigma^2,\rho}^{**}] \rightarrow 0$ as $n \rightarrow +\infty$ and that $\sup_{x \in [0,1]^d} \mathbb{E}[Z_{n,\sigma^2,\rho}(x)^2] = \sigma^2 \sup_{x \in [0,1]^d} k_{n,\rho}(x, x)$ goes to zero uniformly in $(\sigma^2, \rho) \in \bar{\Theta}$ as $n \rightarrow \infty$ for any compact $\bar{\Theta} \subset (0, \infty)^2$. Here $Z_{n,\sigma^2,\rho}$ is defined as in Lemma A.3. We also have as in the proof of Theorem 3.3 when $\kappa = 0$ that

$$\mathbb{P}_{\sigma^2,\rho}(Y^* > u|y) \leq \exp \left\{ - \frac{\left(\left(u - \sup_{x \in [0,1]^d} m_{n,\rho,y}(x) - \mathbb{E}[Z_{n,\sigma^2,\rho}^{**}] \right)_+ \right)^2}{2 \sup_{x \in [0,1]^d} \mathbb{E}[Z_{n,\sigma^2,\rho}(x)^2]} \right\}.$$

Similarly, we bound the probability of the event $\{Y_* < \ell\}$ conditionally to y . Hence, we conclude (as in the proof of Theorem 3.3 when $\kappa = 0$), also from (B.8) and (B.9), that

$$\sup_{(\sigma^2, \rho) \in \bar{\Theta}} |B_n(\sigma^2, \rho)| = o_{\mathbb{P}|Y \in \mathcal{E}_0}(1).$$

Consequently, (A.7) follows and the proof is concluded. \square

C Proofs - Monotonicity

We let $\kappa = 1$ throughout Appendix C.

C.1 Estimation of the variance parameter

Proof of Theorem 3.2 under monotonicity constraints. The proof is similar to that of Theorem 3.2 when $\kappa = 0$ and is also divided into the three steps **1)**, **2)** and **3)**.

1) For $n \in \mathbb{N}$, let N_n be the greatest integer such that Condition-Grid holds. Now we define

$$m_{i,n} = \min \left\{ \frac{1}{v_{j_i}^{(i)} - v_{j_{i-1}}^{(i)}} \left(Y \left(v_{j_1}^{(1)}, \dots, v_{j_{i-1}}^{(i-1)}, v_{j_i}^{(i)}, v_{j_{i+1}}^{(i+1)}, \dots, v_{j_d}^{(d)} \right) \right. \right. \\ \left. \left. - Y \left(v_{j_1}^{(1)}, \dots, v_{j_{i-1}}^{(i-1)}, v_{j_{i-1}}^{(i)}, v_{j_{i+1}}^{(i+1)}, \dots, v_{j_d}^{(d)} \right) \right), \right. \\ \left. j_k \in \{1, \dots, N_n\}, \forall k \in \{1, \dots, d\} \setminus \{i\}, j_i \in \{2, \dots, N_n\} \right\}$$

for all $i = 1, \dots, d$. Since $N_n \rightarrow \infty$ as $n \rightarrow \infty$, $m_{i,n} \xrightarrow[n \rightarrow +\infty]{\text{a.s.}} \inf_{x \in [0,1]^d} \partial Y(x) / \partial x_i$ since Y is \mathcal{C}^1 a.s. by Condition-Var. Now we notice that $m_{i,n} = g_{i,n}(y_1, \dots, y_n)$ and we define $m_{k,i,n} = g_{i,k}(y_1, \dots, y_k)$. One can see that a slightly different version of Lemma A.1 can be shown (up to reindexing x_1, \dots, x_n) with $m_{k,n} = \min\{m_{k,1,n}, \dots, m_{k,d,n}\}$, $m = \inf_{x \in [0,1]^d} \min_{i=1, \dots, d} \partial Y(x) / \partial x_i$, and $M_{k,n} = M = u = +\infty$. After applying this different version, points **2)** and **3)** in the proof of Theorem 3.2 when $\kappa = 0$ remain unchanged. This concludes the proof. \square

Proof of Theorem 3.3 under monotonicity constraints. The proof is similar to that of Theorem 3.3 when $\kappa = 0$ and is also divided into the five steps **1)** to **5)**. We apply Lemma A.2 to the sequences of functions f_n , g_n and h_n defined by $f_n(\sigma^2) = \mathcal{L}_n(\sigma^2)$, $g_n(x) = A_n(\sigma^2)$ and $h_n(\sigma^2) = B_n(\sigma^2)$. Here we recall that for $\sigma^2 \in \Theta$,

$$A_n(\sigma^2) = -\log \mathbb{P}_{\sigma^2}(Y \in \mathcal{E}_1) \text{ and } B_n(\sigma^2) = \log \mathbb{P}_{\sigma^2}(Y \in \mathcal{E}_1 | y).$$

In order to apply Lemma A.2, we need to check that the conditions (A.4) to (A.7) hold.

1) and **2)** The proof that (A.4) and (A.5) are satisfied is identical to the proof for Theorem 3.3 when $\kappa = 0$, as (A.4) and (A.5) do not involve the event $\{Y \in \mathcal{E}_1\}$.

3) Let us introduce the Gaussian process Y_r with mean function zero and covariance function k_1 . Then we have

$$A_n(\sigma^2) = -\log \mathbb{P} \left(\forall x \in [0, 1]^d, \forall i = 1, \dots, d, \sigma \frac{\partial}{\partial x_i} Y_r \geq 0 \right).$$

Hence $A_n(\sigma^2)$ does not depend on σ^2 so that (A.6) holds.

4) We turn to

$$B_n(\sigma^2) = \log \mathbb{P}_{\sigma^2} \left(\forall x \in [0, 1]^d, \forall i = 1, \dots, d, \frac{\partial}{\partial x_i} Y \geq 0 \middle| y \right)$$

For $i = 1, \dots, d$, let $m_{n,y}^{(1,i)}$ and $\sigma^2 k_n^{(1,i)}$ be the conditional mean and covariance function of $\partial Y / \partial x_i$ given y , under the probability measure \mathbb{P}_{σ^2} . We obtain using Borell-TIS inequality (Adler and Taylor, 2007) and a union bound, with $Z_{n,\sigma^2}^{(1,i)}$ a Gaussian process with mean function zero and covariance function $\sigma^2 k_n^{(1,i)}$,

$$\begin{aligned} & \mathbb{P}_{\sigma^2} \left(\exists x \in [0, 1]^d, \exists i = 1, \dots, d, \frac{\partial}{\partial x_i} Y(x) \leq 0 \middle| y \right) \\ & \leq \sum_{i=1}^d \mathbb{P}_{\sigma^2} \left(\sup_{x \in [0, 1]^d} \left(-Z_{n,\sigma^2}^{(1,i)}(x) \right) \geq \inf_{x \in [0, 1]^d} m_{n,y}^{(1,i)}(x) \middle| y \right) \\ & \leq \sum_{i=1}^d \mathbb{P}_{\sigma^2} \left(\sup_{x \in [0, 1]^d} \left| Z_{n,\sigma^2}^{(1,i)}(x) \right| \geq \inf_{x \in [0, 1]^d} m_{n,y}^{(1,i)}(x) \middle| y \right) \\ & \leq \sum_{i=1}^d \exp \left\{ - \frac{\left(\left(\inf_{x \in [0, 1]^d} m_{n,y}^{(1,i)}(x) - \mathbb{E} \left[\sup_{x \in [0, 1]^d} \left| Z_{n,\sigma^2}^{(1,i)}(x) \right| \right] \right)_+ \right)^2}{2 \sup_{x \in [0, 1]^d} \mathbb{E}[Z_{n,\sigma^2}^{(1,i)}(x)^2]} \right\}. \end{aligned} \quad (\text{C.1})$$

One can see that Lemma A.3 can also be shown when $Z_{n,\theta}$ is replaced by $Z_{n,\theta}^{(1,i)}$ (here $\theta = \sigma^2$). Hence $\sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathbb{E} \left[\sup_{x \in [0, 1]^d} \left| Z_{n,\sigma^2}^{(1,i)}(x) \right| \right]$ goes to 0 as $n \rightarrow \infty$. Additionally, one can simply show that $\sup_{x \in [0, 1]^d} \mathbb{E}[Z_{n,\sigma^2}^{(1,i)}(x)^2] = \sup_{x \in [0, 1]^d} \sigma^2 k_n^{(1,i)}(x, x)$ goes to zero uniformly in $\sigma^2 \in [\sigma_l^2, \sigma_u^2]$ as $n \rightarrow \infty$.

One can see that the proof of (Bect et al., 2016, Proposition 2.8) can be adapted to establish that, for $i = 1, \dots, d$,

$$\sup_{x \in [0, 1]^d} \left| m_{n,y}^{(1,i)}(x) - \frac{\partial}{\partial x_i} Y(x) \right| \xrightarrow[n \rightarrow +\infty]{\text{a.s.}} 0$$

from which we deduce that on the set $\{Y \in \mathcal{E}_{1,\delta}\}$, where

$$\mathcal{E}_{1,\delta} := \left\{ f \in \mathcal{C}^1([0, 1]^d, \mathbb{R}) \quad \text{s.t.} \quad \partial f(x) / \partial x_i \geq \delta, \quad \forall x \in [0, 1]^d, i \in \{1, \dots, d\} \right\},$$

we have a.s., for $i = 1, \dots, d$,

$$\liminf_{n \rightarrow +\infty} \left(\inf_{x \in [0, 1]^d} m_{n,y}^{(1,i)}(x) \right) \geq \delta.$$

Consequently, from (C.1), on $\{Y \in \mathcal{E}_{1,\delta}\}$, we have:

$$\sup_{\sigma^2 \in [\sigma_l^2, \sigma_u^2]} \mathbb{P}_{\sigma^2} \left(\exists x \in [0, 1]^d, \exists i = 1, \dots, d, \frac{\partial}{\partial x_i} Y(x) \leq 0 \middle| y \right) \xrightarrow[n \rightarrow +\infty]{\text{a.s.}} 0. \quad (\text{C.2})$$

Similarly as in the proof of Theorem 3.3 when $\kappa = 0$, we can show, by applying Tsirelson theorem in (Azaïs and Wschebor, 2009) to the processes $\partial Y / \partial x_i$, that

$$\mathbb{P}_{\sigma_0^2} (Y \in \mathcal{E}_1 \cap \mathcal{E}_{1,\delta}^c) \xrightarrow{\delta \rightarrow 0} 0.$$

Hence we conclude the proof of (A.7) as for Theorem 3.3 when $\kappa = 0$.

5) We conclude the proof as in 5) for Theorem 3.3 when $\kappa = 0$. \square

C.2 Isotropic Matérn process

Proof of Theorem 4.2 under monotonicity constraints. The proof is the same as for Theorem 4.2 when $\kappa = 0$ and is concluded by applying Theorem 3.2 when $\kappa = 1$. \square

Proof of Theorem 4.3 under monotonicity constraints. The proof is similar to that of Theorem 4.3 when $\kappa = 0$ and is also divided into the four steps 1) to 4). We apply Lemma A.2 to the sequences of functions f_n , g_n and h_n defined by $f_n(\sigma^2) = \mathcal{L}_n(\sigma^2, \rho_1)$, $g_n(x) = A_n(\sigma^2, \rho_1)$, and $h_n(\sigma^2) = B_n(\sigma^2, \rho_1)$.

1) The proof that (A.4) and (A.5) are satisfied is identical to the proof of Theorem 4.3 when $\kappa = 0$, as (A.4) and (A.5) do not involve the event $\{Y \in \mathcal{E}_1\}$.

2) Let us introduce the Gaussian process Y_r with mean function zero and covariance function $k_{1,\rho_1,\nu}$. Then we have

$$A_n(\sigma^2, \rho_1) = -\log \mathbb{P} \left(\forall x \in [0, 1]^d, \forall i = 1, \dots, d, \sigma \frac{\partial}{\partial x_i} Y_r(x) \geq 0 \right).$$

Hence $A_n(\sigma^2, \rho_1)$ does not depend on σ^2 so that (A.6) holds.

3) We turn to $B_n(\sigma^2, \rho_1)$. We conclude to (A.7) following the same lines as in the proof of Theorem 4.3 when $\kappa = 0$ and using the equivalence of measures.

4) We conclude the proof of Theorem 4.3 when $\kappa = 1$ similarly as in the proof of Theorem 4.3 when $\kappa = 0$ using Theorem 4.2 when $\kappa = 1$. \square

Proof of Theorem 4.4 under monotonicity constraints. The proof follows the similar four steps of the proof Theorem 4.4 when $\kappa = 0$. We apply Lemma A.2 to the sequences of functions f_n , g_n and h_n defined by $f_n(x) = \mathcal{L}_n(x \hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c})$, $g_n(x) = A_n(x \hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c})$, and $h_n(x) = B_n(x \hat{\rho}_{n,c}^{2\nu}, \hat{\rho}_{n,c})$.

1) The proof that (A.4) and (A.5) are satisfied is identical to the proof of Theorem 4.4 when $\kappa = 0$, as (A.4) and (A.5) do not involve the event $\{Y \in \mathcal{E}_1\}$.

2) Similarly as in the proof of Theorem 4.4 when $\kappa = 0$, we show that Theorem 4.4 when $\kappa = 1$ holds if (A.6) and (A.7) are satisfied.

3) Similarly as in the proof of Theorem 4.3 when $\kappa = 1$, we show that (A.6) holds.

4) Finally, we turn to (A.7). First, consider the case a). Recall the notation $Y'_* = \inf_{x \in [0,1]} Y'(x)$ from Section 2.2. We proceed similarly as in the proof of Theorem 4.4 when $\kappa = 0$. Since $\mathbb{E}_\rho[Y'(x)|y] = m'_{n,\rho,y}(x)$, it is then sufficient to show that, for all $i = 1, \dots, d$, for any $\varepsilon > 0$, with probability going to 1 as $n \rightarrow \infty$,

$$\sup_{\rho \in [\rho_l, \rho_u]} \sup_{x \in [0,1]} (Y'_* - m'_{n,\rho,y}(x)) \leq \varepsilon \quad (\text{C.3})$$

in order to prove (A.7) as in the proof of Theorem 4.4 when $\kappa = 0$. Analogously, since $\nu > 2$, (B.20) holds. Consider $\bar{x}, \bar{\rho}$ so that $m'_{n,\bar{\rho},y}(\bar{x}) \leq Y'_* - \varepsilon$. There exists $i, j \in \{1, \dots, n\}$, $x_i < x_j$ such that $|x_i - \bar{x}| \leq a_n$ and $|x_j - \bar{x}| \leq a_n$. We have $m_{n,\bar{\rho},y}(x_j) - m_{n,\bar{\rho},y}(x_i) = Y(x_j) - Y(x_i) \geq (x_j - x_i)Y'_*$. Thus from the mean value theorem, there exists $w \in [0, 1]$ so that $|\bar{x} - w| \leq 2a_n$ and $|m'_{n,\bar{\rho},y}(w) - m'_{n,\bar{\rho},y}(\bar{x})| \geq \varepsilon$. Hence there exists $z \in [0, 1]$ so that $m''_{n,\bar{\rho},y}(z) \geq \varepsilon/(2a_n)$. Hence (C.3) holds from (B.20).

Second, consider the case b). We shall only address the case $d = 3$, the cases $d = 1, 2$ being treated similarly. For all $i = 1, 2, 3$, let us define $Y^{(1,i)} = \partial Y / \partial x_i$ and $Y_*^{(1,i)} = \inf_{x \in [0,1]^3} Y^{(1,i)}(x)$

following the notation of Section 2.2. Let also $m_{n,\rho,y}^{(1,i)}(x) = \mathbb{E}_\rho[Y^{(1,i)}(x)|y] = \partial m_{n,\rho,y}(x) / \partial x_i$. First, we want to show that, for all $i = 1, 2, 3$, for any $\varepsilon > 0$, with probability going to 1 as $n \rightarrow \infty$,

$$\sup_{\rho \in [\rho_l, \rho_u]} \sup_{x \in [2/n^{1/3}, 1-2/n^{1/3}]^3} (Y_*^{(1,i)} - m_{n,\rho,y}^{(1,i)}(x)) \leq \varepsilon. \quad (\text{C.4})$$

Assume that there exists $i \in \{1, 2, 3\}$ and $\bar{x}, \bar{\rho}$ for which $m_{n,\bar{\rho},y}^{(1,i)}(\bar{x}) \leq Y_*^{(1,i)} - \varepsilon$. There exists $\{i_1, \dots, i_8\} \in \{1, \dots, n\}$ so that, \bar{x} belongs to the hypercube C with vertices x_j , $j \in \{i_1, \dots, i_8\}$. We refer to Figure 4 for an illustration. This hypercube lies between two adjacent hypercubes C_l, C_r (with vertices in $\{x_1, \dots, x_n\}$ and edge lengths $1/(\lfloor n^{1/3} \rfloor - 1)$) which are obtained by translations (to the left and to the right) in the direction i . Note that C, C_l, C_r are disjoint and that the pairs C, C_l and C, C_r each have a common face which is orthogonal to the direction i . We now consider the 8 vertices v_1, \dots, v_8 of C_l and C_r which are parallel to the direction i . For any $j = 1, \dots, 8$, the endpoints of v_j can be written as x_a, x_b with $(x_a)_k = (x_b)_k$ for $k \neq i$ and with $(x_a)_i < (x_b)_i$. Then we have $m_{n,\bar{\rho},y}(x_b) - m_{n,\bar{\rho},y}(x_a) = Y(x_b) - Y(x_a) \geq Y_*^{(1,i)}((x_b)_i - (x_a)_i)$. Hence, from the mean value theorem, there exists $w_j \in v_j$ for which $m_{n,\bar{\rho},y}^{(1,i)}(w_j) \geq Y_*^{(1,i)}$. Also, it can be shown that \bar{x} belongs to $\text{ch}(w_1, \dots, w_8)$. In addition, it can also be shown that \bar{x} belongs to $\text{ch}(z_1, \dots, z_4)$, with $\{z_1, \dots, z_4\} \subset \{w_1, \dots, w_8\}$. Since $m_{n,\bar{\rho},y}^{(1,i)}(z_j) \geq Y_*^{(1,i)}$ for $j = 1, \dots, 4$ and since $m_{n,\bar{\rho},y}^{(1,i)}(\bar{x}) \leq Y_*^{(1,i)} - \varepsilon$, we show (C.4) as in the proof of Theorem 4.4 b) when $\kappa = 0$.




Figure 4: The point \bar{x} belongs to the hypercube C (in black) that lies between C_l (in green) and C_r (in blue). The mean value theorem ensures the existence of the 8 points w_j , $j = 1, \dots, 8$, and \bar{x} belongs to the convex hull $\text{ch}(w_1, \dots, w_8)$. Furthermore, \bar{x} also belongs to $\text{ch}(z_1, \dots, z_4)$, with $\{z_1, \dots, z_4\} \subset \{w_1, \dots, w_8\}$ (in red).

We have, for $i = 1, 2, 3$,

$$\begin{aligned}
& \mathbb{P}_{(\sigma^2, \rho)} \left(\inf_{x \in [0, 1]^3} Y^{(1, i)}(x) \leq 0 \mid y \right) \\
& \leq \mathbb{P}_{(\sigma^2, \rho)} \left(\inf_{x \in [2/n^{1/3}, 1-2/n^{1/3}]^3} Y^{(1, i)}(x) \leq \delta/2 \mid y \right) \\
& \quad + \mathbb{P}_{(\sigma^2, \rho)} \left(\inf_{x \in [0, 1]^3} Y^{(1, i)}(x) \leq 0, \inf_{x \in [2/n^{1/3}, 1-2/n^{1/3}]^3} Y^{(1, i)}(x) > \delta/2 \mid y \right) \\
& =: P_1 + P_2,
\end{aligned}$$

say. As in the proof of Theorem 4.4 b) when $\kappa = 0$, we can show from (C.4) that $\mathbb{1}_{Y_*^{(1, i)} \geq \delta} \sup_{(\sigma^2, \rho) \in \bar{\Theta}} P_1$ goes to zero in probability so that it is sufficient to show $\sup_{(\sigma^2, \rho) \in \bar{\Theta}} P_2$ goes to zero in probability where $\bar{\Theta}$ is any compact set of $(0, \infty)^2$. We have, with $Y^{(2, j, i)}(x) = \partial^2 Y(x) / (\partial x_j \partial x_i)$, and with $m_{n, \rho, y}^{(2, j, i)}(x) = \partial^2 m_{n, \rho, y}(x) / (\partial x_j \partial x_i) = \mathbb{E}[Y^{(2, j, i)}(x) | y]$,

$$\begin{aligned}
P_2 & \leq \mathbb{P}_{(\sigma^2, \rho)} \left(\max_{j=1, 2, 3} \sup_{x \in [0, 1]^3} |Y^{(2, j, i)}(x)| \geq cn^{1/3} \mid y \right) \\
& \leq \mathbb{P}_{(\sigma^2, \rho)} \left(\max_{j=1, 2, 3} \sup_{x \in [0, 1]^3} |m_{n, \rho, y}^{(2, j, i)}(x)| \geq (c/2)n^{1/3} \mid y \right) \\
& \quad + \mathbb{P}_{(\sigma^2, \rho)} \left(\max_{j=1, 2, 3} \sup_{x \in [0, 1]^3} |m_{n, \rho, y}^{(2, j, i)}(x) - Y^{(2, j, i)}(x)| \geq (c/2)n^{1/3} \mid y \right).
\end{aligned}$$

Using the Borell-TIS inequality as in the proof of 3.3 when $\kappa = 1$, we can show that

$$\sup_{\rho, \sigma^2 \in \bar{\Theta}} \mathbb{P}_{(\sigma^2, \rho)} \left(\max_{j=1,2,3} \sup_{x \in [0,1]^3} \left| m_{n,\rho,y}^{(2,j,i)}(x) - Y^{(2,j,i)}(x) \right| \geq (c/2)n^{1/3} \middle| y \right) = o_{\mathbb{P}}(1).$$

Now consider that there exist $\bar{\rho} \in [\rho_l, \rho_u]$, $\bar{j} \in \{1, 2, 3\}$ and $\bar{x} \in [0, 1]^3$ so that $|m_{n,\bar{\rho},y}^{(2,\bar{j},i)}(\bar{x})| \geq (c/2)n^{1/3}$. If $\bar{j} = i$, then by applications of the mean value theorem, by using that $m_{n,\rho,y}(x) = Y(x)$ for $x \in \{x_1, \dots, x_n\}$, we can show that there exists $\bar{w} \in [0, 1]^3$ so that $\|\bar{w} - \bar{x}\| \leq cn^{-1/3}$ and $|m_{n,\bar{\rho},y}^{(2,\bar{j},i)}(\bar{w})| \leq 2 \max_{i,j \in \{1,2,3\}} Y^{(2,j,i)**}$ where with the notation of Section 2.2, $Y^{(2,j,i)**} := \sup_{x \in [0,1]^3} |Y^{(2,j,i)}(x)|$. Hence there exists $\bar{r} \in [0, 1]^3$ so that $\max_{j,k \in \{1,2,3\}} |m_{n,\rho,y}^{(3,k,j,i)}(\bar{r})| \geq cn^{2/3}$ where $m_{n,\rho,y}^{(3,k,j,i)}(x) = \partial^3 / (\partial x_k \partial x_j \partial x_i) m_{n,\rho,y}(x)$.

If $\bar{j} \neq i$, we can consider $\bar{z} \in \{x_1, \dots, x_n\}$ so that $\|\bar{z} - \bar{x}\| \leq cn^{-1/3}$. We also consider 9 additional points $(\bar{z}_{k,l})_{k,l \in \{1,2,3\}} \subset \{x_1, \dots, x_n\}$ so that $\bar{z}_{k,l} = \bar{x} + l/(\lfloor n^{1/3} \rfloor - 1)v_k$ where $v_1 = e_i$ with e_i the i -th base column vector, $v_2 = e_{\bar{j}}$ and $v_3 = e_i + e_{\bar{j}}$. By applications of the mean value theorem, we can show that there exist \bar{w}_1 for which $|m_{n,\bar{\rho},y}^{(2,i,i)}(\bar{w}_1)| \leq 3Y^{(2,i,i)**}$, \bar{w}_2 for which $|m_{n,\bar{\rho},y}^{(2,\bar{j},\bar{j})}(\bar{w}_2)| \leq 3Y^{(2,\bar{j},\bar{j})**}$ and \bar{w}_3 for which $|m_{n,\bar{\rho},y}^{(2,i,i)}(\bar{w}_3) + m_{n,\bar{\rho},y}^{(2,\bar{j},\bar{j})}(\bar{w}_3) + 2m_{n,\bar{\rho},y}^{(2,\bar{j},i)}(\bar{w}_3)| \leq 12 \max_{i,j \in \{1,2\}} Y^{(2,j,i)**}$. If $|m_{n,\bar{\rho},y}^{(2,\bar{j},i)}(\bar{w}_3)| \leq (c/4)n^{1/3}$, then there exists $\bar{r} \in [0, 1]^3$ so that $\max_{j,k \in \{1,2,3\}} |m_{n,\rho,y}^{(3,k,j,i)}(\bar{r})| \geq cn^{2/3}$. If $|m_{n,\bar{\rho},y}^{(2,\bar{j},i)}(\bar{w}_3)| \geq (c/4)n^{1/3}$, then $|m_{n,\bar{\rho},y}^{(2,i,i)}(\bar{w}_3)| \geq (c/4)n^{1/3} - 6 \max_{i,j \in \{1,2\}} Y^{(2,j,i)**}$ or $|m_{n,\bar{\rho},y}^{(2,\bar{j},\bar{j})}(\bar{w}_3)| \geq (c/4)n^{1/3} - 6 \max_{i,j \in \{1,2\}} Y^{(2,j,i)**}$. In all the cases, there exists $\bar{r} \in [0, 1]^3$ so that $\max_{j,k \in \{1,2,3\}} |m_{n,\rho,y}^{(3,k,j,i)}(\bar{r})| \geq cn^{2/3}$.

We have also from Theorem 1 in (Zhou, 2008), and since $\nu > 3$,

$$\sup_{\rho \in [\rho_l, \rho_u]} \max_{i,j,k=1,2,3} \sup_{x \in [0,1]^3} \left| \frac{\partial^3}{\partial x_i \partial x_j \partial x_k} m_{n,\rho,y}(x) \right| \leq c \sup_{\rho \in [\rho_l, \rho_u]} \|m_{n,\rho,y}\|_{k_{1,\rho,\nu}} \leq c\sqrt{n} \quad (\text{C.5})$$

with probability going to 1 as $n \rightarrow \infty$ from (B.19).

Hence, we have that with probability going to 1 as $n \rightarrow \infty$,

$$\begin{aligned} & \sup_{\sigma^2, \rho} \mathbb{P}_{(\sigma^2, \rho)} \left(\max_{j=1,2,3} \sup_{x \in [0,1]^3} |m_{n,\rho,y}^{(2,j,i)}(x)| \geq (c/2)n^{1/3} \middle| y \right) \\ &= \mathbb{1}_{\{\max_{j=1,2,3} \sup_{x \in [0,1]^3} |m_{n,\rho,y}^{(2,j,i)}(x)| \geq (c/2)n^{1/3}\}} = 0. \end{aligned}$$

This proves that (A.7) holds so that the proof is complete. \square

D Proofs - Convexity

We let $\kappa = 2$ throughout Appendix D.

D.1 Estimation of the variance parameter

Proof of Theorem 3.2 under convexity constraints. The proof is similar to that of Theorem 3.2 when $\kappa = 1$, where the finite differences of order one are replaced by finite differences of order two. \square

Proof of Theorem 3.3 under convexity constraints. The proof is similar to that of Theorem 3.3 when $\kappa = 0$ introducing the Gaussian process V defined on $S_1 \times \mathbb{R}^d$ by $V(v, x) = v^\top H Y(x) v$ where $S_1 = \{v \in \mathbb{R}^d, \|v\| = 1\}$ and observing that

$$\mathcal{E}_2 = \left\{ f \in \mathcal{C}([0, 1]^d, \mathbb{R}), \forall x \in [0, 1]^d, \inf_{\substack{v \in \mathbb{R}^d \\ \|v\|=1}} v^\top H f(x) v \leq 0 \right\},$$

where $Hf(x)$ represents the Hessian matrix of f at x which means that $Hf(x)_{i,j} = \partial^2 f(x) / (\partial x_i \partial x_j)$. \square

D.2 Isotropic Matérn process

Proof of Theorem 4.2 under convexity constraints. The proof is the same as for Theorem 4.2 when $\kappa = 0$ and is concluded by applying Theorem 3.3 when $\kappa = 2$. \square

Proof of Theorem 4.3 under convexity constraints. The proof is similar to that of Theorem 4.3 when $\kappa = 0$. \square

Proof of Theorem 4.4 under convexity constraints. The proof follows the similar four steps of the proof Theorem 4.4 when $\kappa = 0$. Points **1)** to **3)** are identical. Turning to (A.7), point **4)** can be treated similarly as in the proof of Theorem 4.4 when $\kappa = 1$ but with more cumbersome notation and arguments. In order to ease the reading of the paper, we omit this technical proof. \square

Acknowledgement

This research was conducted within the frame of the Chair in Applied Mathematics OQUAIDO, gathering partners in technological research (BRGM, CEA, IFPEN, IRSN, Safran, Storengy) and academia (CNRS, Ecole Centrale de Lyon, Mines Saint-Étienne, University of Grenoble, University of Nice, University of Toulouse) around advanced methods for Computer Experiments.

References

- Abrahamsen, P. (1997). A review of Gaussian random fields and correlation functions. Technical report, Norwegian computing center.
- Abramowitz, M. and Stegun, I. A. (1964). *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Dover, New York, ninth Dover printing, tenth GPO printing edition.
- Adler, R. J. (1990). An introduction to continuity, extrema, and related topics for general Gaussian processes. IMS.
- Adler, R. J. and Taylor, J. E. (2007). *Random fields and geometry*. Springer Monographs in Mathematics. Springer, New York.
- Anderes, E. (2010). On the consistent separation of scale and variance for Gaussian random fields. *The Annals of Statistics*, 38:870–893.
- Azaïs, J.-M. and Wschebor, M. (2009). *Level sets and extrema of random processes and fields*. John Wiley & Sons, Inc., Hoboken, NJ.
- Bachoc, F. (2013). Cross validation and maximum likelihood estimations of hyper-parameters of Gaussian processes with model misspecification. *Computational Statistics and Data Analysis*, 66:55–69.
- Bachoc, F. (2014). Asymptotic analysis of the role of spatial sampling for covariance parameter estimation of Gaussian processes. *Journal of Multivariate Analysis*, 125:1–35.
- Bachoc, F., Ammar, K., and Martinez, J. (2016). Improvement of code behavior in a design of experiments by metamodeling. *Nuclear science and engineering*, 183(3):387–406.
- Bachoc, F., Bois, G., Garnier, J., and Martinez, J. (2014). Calibration and improved prediction of computer models by universal Kriging. *Nuclear Science and Engineering*, 176(1):81–97.
- Bect, J., Bachoc, F., and Ginsbourger, D. (2016). A supermartingale approach to Gaussian process based sequential design of experiments. *ArXiv e-print 1608.01118*.
- Bevilacqua, M., Faouzi, T., Furrer, R., and Porcu, E. (2016). Estimation and prediction using generalized Wendland covariance functions under fixed domain asymptotics. *Annals of Statistics*, in press.
- Cousin, A., Maatouk, H., and Rulli re, D. (2016). Kriging of financial term-structures. *European Journal of Operational Research*, 255(2):631–648.
- Da Veiga, S. and Marrel, A. (2012). Gaussian process modeling with inequality constraints. *Annales de la facult  des sciences de Toulouse Math matiques*, 21(3):529–555.
- Du, J., Zhang, H., and Mandrekar, V. (2009). Fixed-domain asymptotic properties of tapered maximum likelihood estimators. *The Annals of Statistics*, 37:3330–3361.

- Golchi, S., Bingham, D. R., Chipman, H., and Campbell, D. A. (2015). Monotone emulation of computer experiments. *SIAM/ASA Journal on Uncertainty Quantification*, 3(1):370–392.
- Ibragimov, I. and Rozanov, Y. (1978). *Gaussian Random Processes*. Springer-Verlag, New York.
- Istas, J. and Lang, G. (1997). Quadratic variations and estimation of the local Hölder index of a Gaussian process. *Annales de l’Institut Henri Poincaré*, 33:407–436.
- Jones, D., Schonlau, M., and Welch, W. (1998). Efficient global optimization of expensive black box functions. *Journal of Global Optimization*, 13:455–492.
- Kaufman, C. and Shaby, B. (2013). The role of the range parameter for estimation and prediction in geostatistics. *Biometrika*, 100:473–484.
- Lifshits, M. A. (1995). *Gaussian random functions*, volume 322 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht.
- López-Lopera, A., Bachoc, F., Durrande, N., and Roustand, O. (2017). Finite-dimensional Gaussian approximation with linear inequality constraints. *ArXiv e-print 1710.07453*.
- López-Lopera, A. F. (2017). LineqGPR: Gaussian Process Regression Models with Linear Inequality Constraints. R package version 0.0.1. This package will be freely available in June 2018.
- Maatouk, H. and Bay, X. (2017). Gaussian process emulators for computer experiments with inequality constraints. *Mathematical Geosciences*, 49(5):557–582.
- Matheron, G. (1970). *La Théorie des Variables Régionalisées et ses Applications*. Fascicule 5 in Les Cahiers du Centre de Morphologie Mathématique de Fontainebleau. Ecole Nationale Supérieure des Mines de Paris.
- Pakman, A. and Paninski, L. (2014). Exact Hamiltonian Monte Carlo for truncated multivariate Gaussians. *Journal of Computational and Graphical Statistics*, 23(2):518–542.
- Paulo, R., Garcia-Donato, G., and Palomo, J. (2012). Calibration of computer models with multivariate output. *Computational Statistics and Data Analysis*, 56:3959–3974.
- Rasmussen, C. and Williams, C. (2006). *Gaussian Processes for Machine Learning*. The MIT Press, Cambridge.
- Riihimäki, J. and Vehtari, A. (2010). Gaussian processes with monotonicity information. In *Journal of Machine Learning Research: Workshop and Conference Proceedings*, volume 9, pages 645–652.
- Roustant, O., Ginsbourger, D., Deville, Y., et al. (2012). DiceKriging, DiceOptim: Two R packages for the analysis of computer experiments by Kriging-based metamodeling and optimization. *Journal of Statistical Software*, 51(i01).
- Sacks, J., Welch, W., Mitchell, T., and Wynn, H. (1989). Design and analysis of computer experiments. *Statistical Science*, 4:409–423.

- Santner, T., Williams, B., and Notz, W. (2003). *The Design and Analysis of Computer Experiments*. Springer, New York.
- Stein, M. (1988). Asymptotically efficient prediction of a random field with a misspecified covariance function. *The Annals of Statistics*, 16:55–63.
- Stein, M. (1990a). Bounds on the efficiency of linear predictions using an incorrect covariance function. *The Annals of Statistics*, 18:1116–1138.
- Stein, M. (1990b). Uniform asymptotic optimality of linear predictions of a random field using an incorrect second-order structure. *The Annals of Statistics*, 18:850–872.
- Stein, M. (1999). *Interpolation of Spatial Data: Some Theory for Kriging*. Springer, New York.
- van der Vaart, A. W. (1998). *Asymptotic statistics*, volume 3 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge.
- Wang, D. and Loh, W.-L. (2011). On fixed-domain asymptotics and covariance tapering in Gaussian random field models. *Electronic Journal of Statistics*, 5:238–269.
- Wendland, H. (2004). *Scattered data approximation*, volume 17. Cambridge university press.
- Ying, Z. (1991). Asymptotic properties of a maximum likelihood estimator with data from a Gaussian process. *Journal of Multivariate Analysis*, 36:280–296.
- Ying, Z. (1993). Maximum likelihood estimation of parameters under a spatial sampling scheme. *The Annals of Statistics*, 21:1567–1590.
- Zhang, H. (2004). Inconsistent estimation and asymptotically equivalent interpolations in model-based geostatistics. *Journal of the American Statistical Association*, 99:250–261.
- Zhang, H. and Wang, Y. (2010). Kriging and cross validation for massive spatial data. *Environmetrics*, 21:290–304.
- Zhou, D.-X. (2008). Derivative reproducing properties for kernel methods in learning theory. *Journal of computational and Applied Mathematics*, 220(1):456–463.