

Bridges with random length: Gamma case

Mohamed Erraoui

Université Cadi Ayyad, Faculté des Sciences Semlalia,
Département de Mathématiques, B.P. 2390, Marrakech, Maroc
Email: erraoui@uca.ac.ma

Astrid Hilbert

School of Computer Science, Physics and Mathematics,
Linnaeus University, Vejdesplats 7, SE-351 95 Växjö, Sweden.
Email: astrid.hilbert@lnu.se

Mohammed Louriki

Université Cadi Ayyad, Faculté des Sciences Semlalia,
Département de Mathématiques, B.P. 2390, Marrakech, Maroc
Email: louriki.1992@gmail.com

March 9, 2024

Abstract

The aim objective of this paper is to show that certain basic properties of gamma bridges with deterministic length stay true also for gamma bridges with random length. Among them the Markov property as well as the canonical decomposition with respect to the usual augmentation of its natural filtration, which leads us to conclude that its completed natural filtration is right continuous.

Keywords: Lévy processes, Gamma processes, Gamma bridges, Markov Process, Bayes Theorem.

1 Introduction

The gamma process has proven very successful when modelling accumulation processes. Early studies by e.g. Hammersley [13], Moran [22], Gani [12], Kendall [19], Kingman [20] addressed

the modelling of water stored in and released from reservoirs and accumulation related to storage in general. Dufresne et al. [8] show how to employ the gamma process to model liabilities of insurance portfolios for continuous claims. For these risk models the fixed budget horizon of one year is assumed. The authors investigate furthermore the gamma process in the setting of ruin theory and supply ruin probabilities in form of tables. The gamma process replaces the compound processes used traditionally. In Eméry and Yor [11] and Yor [28] gamma bridges were studied and their application to stop loss reinsurance and credit risk management were pointed out in Brody et al [7]. This work introduces and focusses on random gamma bridges, which model accumulated losses of large credit portfolios in credit risk management. These studies were continued by Hoyle et al. in a series of papers, see e.g. Hoyle and Mengütürk [15]. In a further article pricing at an intermediate time is studied Hoyle et al [14]. Returning to the starting point accumulation processes for storage, we refer to recent developments in Chan et al [9] for further references. Numerical results may be found in Assmusen and Hobolth [2].

In this paper we generalize the concept of a gamma bridge to random times, at which the bridge is pinned, to study amongst others its Markov property and to give its decomposition semimartingale. There are two recent works in which bridges with random length are studied. The first by Bedini et al [4] studies related properties of the Brownian bridge with random length, the second by Erraoui and Louriki [10] studies Gaussian bridges with random length. In both works existence of an explicit expression for the bridge with random length is exploited. Applications for the random gamma bridge suggest itself for accumulation processes in financial mathematics, see the results of Jeanblanc and Le Cam in [16] and [17], if heavier tails are desired, electricity production in river power plants with implications to electricity prices, life insurance portfolios, where the instant of death is random, accumulation storage in the setting of just in time production.

The paper is organized as follows. Section 2 begins by recalling the definitions and some properties of gamma processes and gamma bridges of deterministic length, which will be used throughout the paper. In Section 3 we define the gamma bridge with random time τ which will be denoted by ζ and we consider the stopping time property of τ with respect to the right continuous and completed filtration $\mathbb{F}_+^{\zeta,c}$ generated by the process ζ . Moreover, we give the conditional distribution of τ and ζ_u given ζ_t for $u > t > 0$. Next we establish the Markov property of the process ζ with respect to its completed natural filtration. As a consequence, we derive Bayesian estimates for the distribution of the default time τ , given the past behaviour of the process ζ up to time t . After that we study the Markov property of the gamma bridge with random length, with respect to $\mathbb{F}_+^{\zeta,c}$. Finally we give its semimartingale decomposition with respect to $\mathbb{F}_+^{\zeta,c}$.

The following notation will be used throughout the paper: For a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$, \mathcal{N}_p denotes the collection of \mathbb{P} -null sets. If θ is a random variable, then \mathbb{P}_θ denotes the law of θ under \mathbb{P} . \mathcal{D} denotes the space of right continuous functions with left limits (càdlàg) from \mathbb{R}_+ to \mathbb{R}_+ , endowed with Skorohod's topology, under which the space \mathcal{D} is a Polish space. If E is a topological space, then the Borel σ -algebra over E will be denoted by $\mathcal{B}(E)$. The characteristic function of a set A is written \mathbb{I}_A . The symmetric difference of two sets A and B is denoted by $A\Delta B$. Finally for any process $Y = (Y_t, t \geq 0)$ on $(\Omega, \mathcal{F}, \mathbb{P})$, we define by:

(i) $\mathbb{F}^Y = \left(\mathcal{F}_t^Y := \sigma(Y_s, s \leq t), t \geq 0 \right)$ the natural filtration of the process Y .

(ii) $\mathbb{F}^{Y,c} = \left(\mathcal{F}_t^{Y,c} := \mathcal{F}_t^Y \vee \mathcal{N}_p, t \geq 0 \right)$ the completed natural filtration of the process Y .

- (iii) $\mathbb{F}_+^{Y,c} = \left(\mathcal{F}_{t^+}^{Y,c} := \bigcap_{s>t} \mathcal{F}_s^{Y,c} = \mathcal{F}_{t^+}^Y \vee \mathcal{N}_P, t \geq 0 \right)$ the smallest filtration containing \mathbb{F}^Y and satisfying the usual hypotheses of right-continuity and completeness.

2 Gamma and Gamma bridge processes

The purpose of this section is to recall the definition and some properties of the standard gamma process and the gamma bridge with deterministic length.

2.1 Gamma process

By a standard gamma process $(\gamma_t, t \geq 0)$ on $(\Omega, \mathcal{F}, \mathbb{P})$, we mean a subordinator without drift having the Lévy-Khintchine representation given by

$$\mathbb{E}(\exp(-\lambda \gamma_t)) = \exp \left(-t \int_0^\infty (1 - \exp(-\lambda x)) \frac{\exp(-x)}{x} dx \right) \quad (2.1)$$

$$= (1 + \lambda)^{-t}, \quad (2.2)$$

where $\nu(dx) = \frac{\exp(-x)}{x} \mathbb{I}_{(0,\infty)}(x) dx$ is the so-called Lévy measure. We note that the formula (2.2) is obtained from (2.1) using the Frullani formula.

The following properties, inferred from (2.2) by means of standard arguments (see, e.g., Sato [25], ch. 2 and 4), describe the paths of the gamma process.

Proposition 2.1. *The gamma process $(\gamma_t, t \geq 0)$ has the following properties:*

- (i) γ is a purely jump process;
- (ii) γ is not a compound Poisson process and its jumping times are countable and dense in $[0, \infty)$ a.s.;
- (iii) the map $t \mapsto \gamma_t$ is strictly increasing and not continuous anywhere a.s.;
- (iv) γ has sample paths of finite variation a.s.;
- (v) $\gamma_t, t > 0$, follows a gamma distribution with density

$$f_{\gamma_t}(x) = \frac{x^{t-1} \exp(-x)}{\Gamma(t)} \mathbb{I}_{(0,\infty)}(x), \quad (2.3)$$

where Γ is the gamma function.

The second property means that, for any $t > 0$, γ has infinite activity, that is, almost all paths have infinitely many jumps along any time interval of finite length. It is a direct consequence of $\nu(\mathbb{R}_+) = +\infty$, whereas the fourth property arises from $\int_0^1 x \nu(dx) < +\infty$.

Remark 2.2. 1. It is clear that the process gamma $(\gamma_t, t \geq 0)$ is a process with paths in \mathcal{D} .

2. The process $(\gamma_t - \gamma_{t-} := e_t, t \geq 0)$ of jumps of the gamma process $(\gamma_t, t \geq 0)$ is a Poisson point process whose intensity measure is the Lévy measure of $(\gamma_t, t \geq 0)$, see Bertoin [3]. For $r > 0$, let us denote by $(J_1^r \geq J_2^r \geq \dots)$ the sequel of the lengths of jumps of the process $(\gamma_t, t \in [0, r])$ ranked in decreasing order. It is not difficult to see that since the intensity measure of the Poisson point process $((t, e_t), t \geq 0)$ is $dt \frac{\exp(-x)}{x} \mathbb{I}_{(0, \infty)}(x) dx$, then the jump times (U_1^r, U_2^r, \dots) constitute a sequence of i.i.d r.v.'s with uniform law on $[0, r]$ which is independent from the sequence $(J_k^r, k \geq 1)$. Thus we have the following representation:

$$\gamma_t = \sum_{k \geq 1} J_k^r \mathbb{I}_{\{U_k^r \leq t\}}, \quad t \in [0, r]. \quad (2.4)$$

We note that : $\gamma_r = \sum_{k \geq 1} J_k^r$.

The next proposition gives three other useful properties of the gamma process.

Proposition 2.3. (i) For every $r > 0$, the σ -algebras, $\sigma(\frac{\gamma_u}{\gamma_r}, u \in [0, r])$ and $\sigma(\gamma_u, u \in [r, \infty))$ are independent.

(ii) For any $r > 0$, $(\gamma_t, 0 \leq t \leq r)$ satisfies the following equation

$$\gamma_t = M_t^r + \int_0^t \frac{\gamma_r - \gamma_s}{r - s} ds, \quad (2.5)$$

where $(M_t^r, t \in [0, r])$ is a $\mathcal{G}_t^{(r)}$ -martingale with $\mathcal{G}_t^{(r)} = \sigma(\gamma_s, s \in [0, t] \cup \{r\})$.

(iii) $(\gamma_t, t \geq 0)$ has the Markov property with respect to its natural filtration.

Proof. For (i) and (ii) See, [11]. (iii) $(\gamma_t, t \geq 0)$ has the Markov property since it is a Lévy process. \square

For a deeper investigation on the properties of the gamma process we refer to Kyprianou [21], Sato [25] and Yor [28].

2.2 Gamma bridge with deterministic length

A bridge is a stochastic process that is pinned to some fixed point at a fixed future time. In this section we define the gamma bridge with deterministic length and we give some important properties of this process. For fixed $r > 0$, we define the gamma bridge of length r by setting

Definition 2.4. Let $r \in (0, +\infty)$. The map $\zeta^r : \Omega \mapsto \mathcal{D}$ defined by

$$\zeta_t^r(\omega) := \frac{\gamma_{t \wedge r}(\omega)}{\gamma_r(\omega)}, \quad t \geq 0, \quad \omega \in \Omega, \quad (2.6)$$

is the bridge associated with the standard gamma process $(\gamma_t, t \geq 0)$. Then clearly $\zeta_0^r = 0$ and $\zeta_r^r = 1$. We refer to ζ^r as the standard gamma bridge of length r associated with γ . We note that ζ^r is also called the Dirichlet process with parameter r .

We note that the process ζ^r is really a function of the variables (r, t, ω) and for technical reasons, it is convenient to have certain joint measurability properties.

Lemma 2.5. *The map $(r, t, \omega) \mapsto \zeta_t^r(\omega)$ of $((0, +\infty) \times \mathbb{R}_+ \times \Omega, \mathcal{B}((0, +\infty)) \otimes \mathcal{B}(\mathbb{R}_+) \otimes \mathcal{F})$ into $(\mathbb{R}_+, \mathcal{B}(\mathbb{R}_+))$ is measurable. In particular, the t -section of $(r, t, \omega) \mapsto \zeta_t^r(\omega)$: $(r, \omega) \mapsto \zeta_t^r(\omega)$ is measurable with respect to the σ -algebra $\mathcal{B}((0, +\infty)) \otimes \mathcal{F}$, for all $t \geq 0$.*

Proof. Since the map $(r, t) \mapsto t \wedge r$ is Lipschitz continuous and $t \mapsto \gamma_t$ is càdlàg for all almost $\omega \in \Omega$, then the map $(r, t, \omega) \mapsto \zeta_t^r(\omega)$ can be obtained as the pointwise limit of sequences of measurable functions. So, it is sufficient to use standard results on the passage to the limit of sequences of measurable functions. \square

As a consequence we have the following corollary.

Corollary 2.6. *The map $(r, \omega) \mapsto \zeta_t^r(\omega)$ of $((0, +\infty) \times \Omega, \mathcal{B}((0, +\infty)) \otimes \mathcal{F})$ into $(\mathcal{D}, \mathcal{B}(\mathcal{D}))$ is measurable.*

A number of properties of the gamma bridge ζ^r sample paths can be easily deduced from the corresponding properties of the gamma sample paths. Hence, we have

Proposition 2.7. *The gamma bridge ζ_t^r , $t \geq 0$, has the following properties:*

- (i) ζ^r is a purely jump process and its jumping times are countable and dense in $[0, r]$ a.s.;
- (ii) the map $t \mapsto \zeta_t^r$ is strictly increasing and not continuous anywhere in $[0, r]$ a.s.;
- (iii) ζ^r has sample paths of finite variation in $[0, +\infty)$ a.s.;
- (iv) ζ^r has the following representation:

$$\zeta_t^r = \sum_{k \geq 1} \frac{J_k^r}{\sum_{j \geq 1} J_j^r} \mathbb{I}_{\{U_k^r \leq t\}}, \quad t \geq 0. \quad (2.7)$$

We now turn to distributional properties of the gamma bridge.

Proposition 2.8. (i) *For all $0 < t < r$, the random variable ζ_t^r has a beta distribution $\beta(t, r-t)$ i.e. its density function is given by*

$$\varphi_{\zeta_t^r}(x) = \frac{\Gamma(r)}{\Gamma(t)\Gamma(r-t)} x^{t-1} (1-x)^{r-t-1} \mathbb{I}_{(0,1)}(x). \quad (2.8)$$

- (ii) *For any $0 = t_0 < t_1 < \dots < t_n = r$, the vector $(\zeta_{t_1}^r - \zeta_{t_0}^r, \dots, \zeta_{t_n}^r - \zeta_{t_{n-1}}^r)$ is independent from γ_r , with density*

$$\frac{\Gamma(r)}{\prod_{i=1}^n \Gamma(t_i - t_{i-1})} \prod_{i=1}^n x_i^{t_i - t_{i-1} - 1}$$

with respect to the Lebesgue measure $dx_1 \dots dx_{n-1}$ (or, as well, $dx_2 \dots dx_n$) on the simplex

$$\{(x_1, \dots, x_n) : x_i \geq 0, x_1 + \dots + x_n = 1\}.$$

(iii) For all $t < u < r$ and $x \in (0, 1)$, the regular conditional law of ζ_u^r given $\zeta_t^r = x$ is given by:

$$\mathbb{P}(\zeta_u^r \in dy | \zeta_t^r = x) = \frac{\Gamma(r-t)}{\Gamma(u-t)\Gamma(r-u)} \frac{(y-x)^{u-t-1}(1-y)^{r-u-1}}{(1-x)^{r-t-1}} \mathbb{I}_{\{x < y < 1\}} dy, \quad (2.9)$$

In the same spirit as in the Proposition 2.3 we have

Proposition 2.9. (i) ζ^r is a Markov process with respect to its natural filtration.

(ii) ζ^r satisfies the following equation

$$\zeta_t^r = N_t^r + \int_0^t \frac{1 - \zeta_s^r}{r - s} ds, \quad t \in [0, r], \quad (2.10)$$

where $(N_t^r, t \in [0, r])$ is a \mathbb{F}^{ζ^r} -martingale.

Proof. (i) From Theorem 1.3 in Blumenthal and Gettoor [5] it suffices to prove that for every bounded measurable function g we have:

$$\mathbb{E}[g(\zeta_u^r) | \zeta_{t_1}^r, \dots, \zeta_{t_n}^r] = \mathbb{E}[g(\zeta_u^r) | \zeta_{t_n}^r], \quad (2.11)$$

for all $0 \leq t_1 < \dots < t_n < u \leq r$ and for all $n \geq 1$.

Using Proposition 2.3 (i) we have

$$\begin{aligned} \mathbb{E}[g(\zeta_u^r) | \zeta_{t_1}^r, \dots, \zeta_{t_n}^r] &= \mathbb{E}\left[g\left(\frac{\gamma_u}{\gamma_r}\right) \middle| \frac{\gamma_{t_1}}{\gamma_r}, \dots, \frac{\gamma_{t_n}}{\gamma_r}\right] \\ &= \mathbb{E}\left[g\left(\frac{\gamma_u}{\gamma_r}\right) \middle| \frac{\gamma_{t_1}}{\gamma_r}, \frac{\gamma_{t_2}}{\gamma_{t_3}}, \dots, \frac{\gamma_{t_{n-1}}}{\gamma_{t_n}}, \frac{\gamma_{t_n}}{\gamma_r}\right] \\ &= \mathbb{E}\left[g\left(\frac{\gamma_u}{\gamma_r}\right) \middle| \frac{\gamma_{t_n}}{\gamma_r}\right] \\ &= \mathbb{E}\left[g(\zeta_u^r) | \zeta_{t_n}^r\right]. \end{aligned}$$

Hence the formula (2.11) is proved, then ζ^r is a Markov process with respect to its natural filtration.

(ii) We have from Proposition 2.3 (ii) that

$$\gamma_t = M_t^r + \int_0^t \frac{\gamma_r - \gamma_s}{r - s} ds, \quad t \in [0, r],$$

where M^r is a martingale with respect to the filtration $\mathcal{G}_t^{(r)} = \sigma(\gamma_s, s \in [0, t] \cup \{r\})$. Then it is easy to see that

$$\zeta_t^r = N_t^r + \int_0^t \frac{1 - \zeta_s^r}{r - s} ds, \quad t \in [0, r], \quad (2.12)$$

where $N_t^r = \frac{M_t^r}{\gamma_r}, t \in [0, r]$. Firstly, notice that, $\mathcal{F}_t^{\zeta^r} \subset \mathcal{G}_t^{(r)}$ and γ_r is $\mathcal{G}_t^{(r)}$ -measurable for all $t \leq r$. Moreover, equation (2.12) yields that is the process N^r is \mathbb{F}^{ζ^r} -adapted. In view of these

considerations, as well as the fact that M_t^r is a $\mathcal{G}_t^{(r)}$ -martingale we obtain

$$\begin{aligned}\mathbb{E}\left[N_t^r|\mathcal{F}_s^{\zeta^r}\right] &= \mathbb{E}\left[\frac{M_t^r}{\gamma_r}|\mathcal{F}_s^{\zeta^r}\right] = \mathbb{E}\left[\mathbb{E}\left[\frac{M_t^r}{\gamma_r}|\mathcal{G}_s^{(r)}\right]|\mathcal{F}_s^{\zeta^r}\right] \\ &= \mathbb{E}\left[\frac{M_s^r}{\gamma_r}|\mathcal{F}_s^{\zeta^r}\right] = \mathbb{E}\left[N_s^r|\mathcal{F}_s^{\zeta^r}\right] = N_s^r,\end{aligned}$$

for $0 \leq s \leq t \leq r$. It follows that $(N_t^r, t \in [0, r])$ is a \mathbb{F}^{ζ^r} -martingale. Hence the equation (2.10) is satisfied. \square

Remark 2.10. We can rewrite (2.5) in the form

$$\gamma_{t \wedge r} = M_{t \wedge r}^r + \int_0^{t \wedge r} \frac{\gamma_r - \gamma_s}{r - s} ds, \quad t \geq 0. \quad (2.13)$$

Then we obtain

$$\zeta_t^r = \frac{\gamma_{t \wedge r}}{\gamma_r} = \frac{M_{t \wedge r}^r}{\gamma_r} + \int_0^{t \wedge r} \frac{1 - \zeta_s^r}{r - s} ds, \quad t \geq 0. \quad (2.14)$$

For every $t \geq 0$, we set $\widehat{N}_t^r = \frac{M_{t \wedge r}^r}{\gamma_r}$. We have thus

$$\zeta_t^r = \widehat{N}_t^r + \int_0^t \frac{1 - \zeta_s^r}{r - s} \mathbb{I}_{\{s < r\}} ds, \quad t \geq 0. \quad (2.15)$$

It follows from the above proposition that $(\widehat{N}_t^r, t \geq 0)$ is a \mathbb{F}^{ζ^r} -martingale stopped at r .

3 Gamma bridges with random length

In this section we define and study a process $(\zeta_t, t \geq 0)$ which generalizes the gamma bridge in the sense that the time r at which the bridge is pinned is substituted by an independent random time τ . We call it *gamma bridge with random length*. We prove that the random time τ is a stopping time with respect to the completed filtration $\mathbb{F}^{\zeta, c}$ and we give the regular conditional distribution of τ and (τ, ζ_\cdot) given ζ_\cdot . Moreover, we prove that the gamma bridge with random length ζ is an inhomogeneous Markov process with respect to its completed natural filtration $\mathbb{F}^{\zeta, c}$ as well as with respect to $\mathbb{F}_+^{\zeta, c}$. The last property allows us to deduce an interesting consequence that is the filtration $\mathbb{F}^{\zeta, c}$ satisfies the usual conditions of completeness and right-continuity. Finally we give the semimartingale decomposition of ζ with respect to $\mathbb{F}_+^{\zeta, c}$.

Now we give precise definition of the process $(\zeta_t, t \geq 0)$. Due to Corollary 2.6 we could substitute r by a random time τ in (2.6). Thus we obtain

Definition 3.1. Let $\tau : (\Omega, \mathcal{F}, \mathbb{P}) \mapsto (0, +\infty)$ be a strictly positive random time, with distribution function $F(t) := \mathbb{P}(\tau \leq t)$, $t \geq 0$. The map $\zeta : \Omega, \mathcal{F}) \rightarrow (\mathcal{D}, \mathcal{B}(\mathcal{D}))$ is defined by

$$\zeta_t(\omega) := \zeta_t^r(\omega)|_{r=\tau(\omega)}, \quad (t, \omega) \in \mathbb{R}_+ \times \Omega.$$

Then ζ takes the form

$$\zeta_t := \frac{\gamma_{t \wedge \tau}}{\gamma_\tau}, \quad t \geq 0. \quad (3.1)$$

Since ζ is obtained by composition of two maps $(r, t, \omega) \mapsto \zeta_t^r(\omega)$ and $(t, \omega) \mapsto (\tau(\omega), t, \omega)$, it's not hard to verify that the map $\zeta : \Omega, \mathcal{F}) \longrightarrow (\mathcal{D}, \mathcal{B}(\mathcal{D}))$ is measurable. The process ζ will be called *gamma bridge of random length τ* .

As mentioned above, we work under the following standing assumption:

Assumption 3.1. *The random time τ and the gamma process γ are independent.*

Using the fact that the process ζ is obtained by the substitution of r in ζ^r by the random time τ allows us to derive a lot of information about its path properties. Hence, we have

Proposition 3.2. *The gamma bridge $\zeta_t, t \geq 0$, has the following properties:*

- (i) *ζ is a purely jump process and its jumping times are countable and dense in $[0, \tau]$ a.s.;*
- (ii) *the map $t \mapsto \zeta_t$ is increasing and not continuous anywhere on $[0, \tau]$ a.s.;*
- (iii) *ζ has sample paths of finite variation a.s.*
- (iv) *ζ has the following representation:*

$$\zeta_t = \sum_{k \geq 1} \frac{J_k^\tau}{\sum_{j \geq 1} J_j^\tau} \mathbb{I}_{\{U_k^\tau \leq t\}}, \quad t \geq 0,$$

where the jump times $(U_1^\tau, U_2^\tau, \dots)$ constitute a sequence of r.v.'s identically distributed with the law given by

$$\begin{aligned} \mathbb{P}[U_k^\tau \leq t] &= \mathbb{P}[\tau \leq t] + \int_{(t, +\infty)} \mathbb{P}[U_k^r \leq t] \mathbb{P}_\tau(dr) \\ &= \mathbb{P}[\tau \leq t] + t \mathbb{E} \left[\frac{1}{\tau} \mathbb{I}_{(\tau > t)} \right], \quad t \geq 0, k \geq 1. \end{aligned}$$

3.1 Stopping time property of τ

The aim of this subsection is to prove that the random time τ is a stopping time with respect to $\mathbb{F}^{\zeta, c}$.

Proposition 3.3. *For all $t > 0$, we have $\mathbb{P}(\{\zeta_t = 1\} \triangle \{\tau \leq t\}) = 0$. Then τ is a stopping time with respect to $\mathbb{F}^{\zeta, c}$ and consequently it is a stopping time with respect to $\mathbb{F}_+^{\zeta, c}$.*

Proof. First we have from the definition of ζ that $\zeta_t = 1$ for $\tau \leq t$. Then $\{\tau \leq t\} \subseteq \{\zeta_t = 1\}$. On the other hand, using the formula of total probability we obtain

$$\begin{aligned} \mathbb{P}(\zeta_t = 1, t < \tau) &= \int_{(t, +\infty)} \mathbb{P}(\zeta_t = 1 | \tau = r) \mathbb{P}_\tau(dr) \\ &= \int_{(t, +\infty)} \mathbb{P}(\zeta_t^r = 1) \mathbb{P}_\tau(dr) \\ &= 0. \end{aligned}$$

The latter equality uses the fact that ζ_t^r is a random variable has a beta distribution for $0 < t < r$. Thus $\mathbb{P}(\{\zeta_t = 1\} \triangle \{\tau \leq t\}) = 0$. It follows that the event $\{\tau \leq t\}$ belongs to $\mathcal{F}_t^\zeta \vee \mathcal{N}_P$, for all $t \geq 0$. Hence τ is a stopping time with respect to $\mathbb{F}^{\zeta, c}$ and consequently it is also a stopping time with respect to $\mathbb{F}_+^{\zeta, c}$. \square

In order to determine the conditional law of the random time τ given ζ_t we will use the following

Proposition 3.4. *Let $t > 0$ such that $F(t) > 0$. Let $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a Borel function satisfying $\mathbb{E}[|g(\tau)|] < +\infty$. Then, \mathbb{P} -a.s., we have*

$$\mathbb{E}[g(\tau)|\zeta_t] = \int_{(0,t]} \frac{g(r)}{F(t)} \mathbb{P}_\tau(dr) \mathbb{I}_{\{\zeta_t=1\}} + \int_{(t,+\infty)} g(r) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}, \quad (3.2)$$

where the function $\phi_{\zeta_t^r}$ is defined on \mathbb{R} by:

$$\begin{aligned} \phi_{\zeta_t^r}(x) &= \frac{\varphi_{\zeta_t^r}(x)}{\int_{(t,+\infty)} \varphi_{\zeta_t^s}(x) \mathbb{P}_\tau(ds)} \\ &= \frac{(1-x)^r \frac{\Gamma(r)}{\Gamma(r-t)}}{\int_{(t,+\infty)} (1-x)^s \frac{\Gamma(s)}{\Gamma(s-t)} \mathbb{P}_\tau(ds)} \mathbb{I}_{(0,1)}(x), \quad x \in \mathbb{R}, \quad r \in (t, +\infty). \end{aligned} \quad (3.3)$$

Proof. Let us consider the measure μ defined on $\mathcal{B}(\mathbb{R})$ by

$$\mu(dx) = \delta_1(dx) + dx,$$

where $\delta_1(dx)$ and dx are the Dirac measure and the Lebesgue measure on $\mathcal{B}(\mathbb{R})$ respectively. Then for any $B \in \mathcal{B}(\mathbb{R})$ we have

$$\mathbb{P}(\zeta_t \in B | \tau = r) = \mathbb{P}(\zeta_t^r \in B) = \int_B q_t(r, x) \mu(dx),$$

where the function q_t is a nonnegative and measurable in the two variables jointly given by

$$q_t(r, x) = \mathbb{I}_{\{x=1\}} \mathbb{I}_{\{r \leq t\}} + \varphi_{\zeta_t^r}(x) \mathbb{I}_{\{0 < x < 1\}} \mathbb{I}_{\{t < r\}}.$$

It follows from Bayes formula (see [26] p. 272) that \mathbb{P} -a.s.:

$$\begin{aligned} \mathbb{E}[g(\tau)|\zeta_t] &= \frac{\int_{(0,+\infty)} g(r) q_t(r, \zeta_t) \mathbb{P}_\tau(dr)}{\int_{(0,+\infty)} q_t(r, \zeta_t) \mathbb{P}_\tau(dr)} \\ &= \frac{\int_{(0,t]} g(r) \mathbb{P}_\tau(dr) \mathbb{I}_{\{\zeta_t=1\}} + \int_{(t,+\infty)} g(r) \varphi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}}{F(t) \mathbb{I}_{\{\zeta_t=1\}} + \int_{(t,+\infty)} \varphi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}} \\ &= \int_{(0,t]} \frac{g(r)}{F(t)} \mathbb{P}_\tau(dr) \mathbb{I}_{\{\zeta_t=1\}} + \int_{(t,+\infty)} g(r) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}. \end{aligned}$$

\square

Corollary 3.5. *The conditional law of the random time τ given ζ_t is given by*

$$\mathbb{P}_{\tau|\zeta_t=x}(x, dr) = \frac{1}{F(t)} \mathbb{I}_{\{x=1\}} \mathbb{I}_{(0,t]}(r) \mathbb{P}_\tau(dr) + \phi_{\zeta_t^r}(x) \mathbb{I}_{\{0 < x < 1\}} \mathbb{I}_{(t,+\infty)}(r) \mathbb{P}_\tau(dr) \quad (3.4)$$

The previous proposition can be expanded as follows

Proposition 3.6. *Let $u > t > 0$ such that $F(t) > 0$. Let \mathbf{g} be a bounded measurable function defined on $(0, +\infty) \times \mathbb{R}$. Then, \mathbb{P} -a.s., we have*

$$\mathbb{E}[\mathbf{g}(\tau, \zeta_t) | \zeta_t] = \int_{(0,t]} \frac{\mathbf{g}(r, 1)}{F(t)} \mathbb{P}_\tau(dr) \mathbb{I}_{\{\zeta_t=1\}} + \int_{(t,+\infty)} \mathbf{g}(r, \zeta_t) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}, \quad (3.5)$$

and

$$\begin{aligned} \mathbb{E}[\mathbf{g}(\tau, \zeta_u) | \zeta_t] &= \int_{(0,t]} \frac{\mathbf{g}(r, 1)}{F(t)} \mathbb{P}_\tau(dr) \mathbb{I}_{\{\zeta_t=1\}} + \int_{(t,u]} \mathbf{g}(r, 1) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}} \\ &\quad + \int_{(u,+\infty)} \mathfrak{G}_{t,u}(r, \zeta_t) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}. \end{aligned} \quad (3.6)$$

Here the function $\mathfrak{G}_{t,u}(r, \cdot)$ is defined by

$$\begin{aligned} \mathfrak{G}_{t,u}(r, x) &:= \mathbb{E}[\mathbf{g}(r, \zeta_u^r) | \zeta_t^r = x] \\ &= \int_{\mathbb{R}} \mathbf{g}(r, y) \mathbb{P}_{\zeta_u^r | \zeta_t^r = x}(dy). \end{aligned} \quad (3.7)$$

Proof. First of all, it is easy to see that (3.5) is an immediate consequence of Proposition 3.4. Now to show (3.6) we begin with by splitting $\mathbb{E}[\mathbf{g}(\tau, \zeta_u) | \zeta_t]$ as follows

$$\mathbb{E}[\mathbf{g}(\tau, \zeta_u) | \zeta_t] = \mathbb{E}[\mathbf{g}(\tau, 1) \mathbb{I}_{\{\tau \leq t\}} | \zeta_t] + \mathbb{E}[\mathbf{g}(\tau, 1) \mathbb{I}_{\{t < \tau \leq u\}} | \zeta_t] + \mathbb{E}[\mathbf{g}(\tau, \zeta_u) \mathbb{I}_{\{u < \tau\}} | \zeta_t]$$

We obtain from Proposition 3.4 that

$$\mathbb{E}[\mathbf{g}(\tau, 1) \mathbb{I}_{\{\tau \leq t\}} | \zeta_t] = \int_{(0,t]} \frac{\mathbf{g}(r, 1)}{F(t)} \mathbb{P}_\tau(dr) \mathbb{I}_{\{\zeta_t=1\}}$$

and

$$\mathbb{E}[\mathbf{g}(\tau, 1) \mathbb{I}_{\{t < \tau \leq u\}} | \zeta_t] = \int_{(t,u]} \mathbf{g}(r, 1) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}.$$

Next we prove that

$$\mathbb{E}[\mathbf{g}(\tau, \zeta_u) \mathbb{I}_{\{u < \tau\}} | \zeta_t] = \int_{(u,+\infty)} \mathfrak{G}_{t,u}(r, \zeta_t) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}. \quad (3.8)$$

Indeed, for a bounded Borel function h we have

$$\begin{aligned}\mathbb{E}[\mathfrak{g}(\tau, \zeta_u)\mathbb{I}_{\{u < \tau\}}h(\zeta_t)] &= \int_{(u, +\infty)} E[\mathfrak{g}(r, \zeta_u^r)h(\zeta_t^r)]\mathbb{P}_\tau(dr) \\ &= \int_{(u, +\infty)} \mathbb{E}[\mathbb{E}[\mathfrak{g}(r, \zeta_u^r)h(\zeta_t^r)|\zeta_t^r]]\mathbb{P}_\tau(dr) \\ &= \int_{(u, +\infty)} \mathbb{E}[\mathbb{E}[\mathfrak{g}(r, \zeta_u^r)|\zeta_t^r]h(\zeta_t^r)]\mathbb{P}_\tau(dr).\end{aligned}$$

Using (3.7), for $t < u < r$, we get

$$\begin{aligned}\mathbb{E}[\mathfrak{g}(\tau, \zeta_u)\mathbb{I}_{\{u < \tau\}}h(\zeta_t)] &= \int_{(u, +\infty)} \mathbb{E}[\mathfrak{G}_{t,u}(r, \zeta_t^r)h(\zeta_t^r)]\mathbb{P}_\tau(dr) \\ &= \mathbb{E}[\mathfrak{G}_{t,u}(\tau, \zeta_t)\mathbb{I}_{\{u < \tau\}}h(\zeta_t)].\end{aligned}$$

It follows from (3.5), that \mathbb{P} -a.s.

$$\mathbb{E}[\mathfrak{G}_{t,u}(\tau, \zeta_t)\mathbb{I}_{\{u < \tau\}}|\zeta_t] = \int_{(u, +\infty)} \mathfrak{G}_{t,u}(r, \zeta_t) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}. \quad (3.9)$$

This induces that

$$\mathbb{E}[\mathfrak{g}(\tau, \zeta_u)\mathbb{I}_{\{u < \tau\}}h(\zeta_t)] = \mathbb{E}\left[\int_{(u, +\infty)} \mathfrak{G}_{t,u}(r, \zeta_t)\phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}} h(\zeta_t)\right].$$

Hence the formula (3.8) is proved and then the proof of the proposition is completed. \square

3.2 Markov property of ζ and Bayes estimate of τ

In this part we prove that the gamma bridge with random length ζ is an inhomogeneous Markov process with respect to its completed natural filtration $\mathbb{F}^{\zeta, c}$.

Theorem 3.7. *The process $(\zeta_t, t \geq 0)$ is an \mathbb{F}^ζ -Markov process. That is, for any $t \geq 0$, we have*

$$\mathbb{E}[f(\zeta_{t+h})|\mathcal{F}_t^\zeta] = \mathbb{E}[f(\zeta_{t+h})|\zeta_t], \mathbb{P} - a.s., \quad (3.10)$$

for all $t, h \geq 0$ and for every bounded measurable function f .

Proof. First, we would like to mention that since $\zeta_0 = 0$ almost surely then it is easy to see that

$$\mathbb{E}[f(\zeta_{t+h})|\mathcal{F}_0^\zeta] = \mathbb{E}[f(\zeta_{t+h})|\zeta_0].$$

Let us assume $t > 0$. As $\mathbb{I}_{\{\zeta_t=0\}} = \mathbb{I}_{\{\tau \leq t\}}$ \mathbb{P} -a.s we rewrite $\mathbb{E}[f(\zeta_{t+h})|\mathcal{F}_t^\zeta]$ as follows

$$\begin{aligned}\mathbb{E}[f(\zeta_{t+h})|\mathcal{F}_t^\zeta] &= \mathbb{E}[f(\zeta_{t+h})|\mathcal{F}_t^\zeta]\mathbb{I}_{\{\tau \leq t\}} + \mathbb{E}[f(\zeta_{t+h})|\mathcal{F}_t^\zeta]\mathbb{I}_{\{t < \tau\}} \\ &= f(1)\mathbb{I}_{\{\zeta_t=1\}} + \mathbb{E}[f(\zeta_{t+h})|\mathcal{F}_t^\zeta]\mathbb{I}_{\{t < \tau\}}.\end{aligned}$$

So it remains to show that

$$\mathbb{E}[f(\zeta_{t+h})\mathbb{I}_{\{t<\tau\}}|\mathcal{F}_t^\zeta] = \mathbb{E}[f(\zeta_{t+h})\mathbb{I}_{\{t<\tau\}}|\zeta_t], \quad \mathbb{P} - a.s.$$

To do this it is enough to verify that

$$\int_{A \cap \{t < \tau\}} f(\zeta_{t+h}) d\mathbb{P} = \int_{A \cap \{t < \tau\}} \mathbb{E}[f(\zeta_{t+h})|\zeta_t] d\mathbb{P}, \quad (3.11)$$

for all $A \in \mathcal{F}_t^\zeta$. We start by remarking that, for $t > 0$, \mathcal{F}_t^ζ is generated by

$$\zeta_{t_n}, \alpha_n := \frac{\zeta_{t_{n-1}}}{\zeta_{t_n}}, \alpha_{n-1} = \frac{\zeta_{t_{n-2}}}{\zeta_{t_{n-1}}}, \dots, \alpha_2 = \frac{\zeta_{t_1}}{\zeta_{t_2}}, \alpha_1 := \frac{\zeta_{t_0}}{\zeta_{t_1}},$$

$0 < t_0 < t_1 < \dots < t_n = t$ for n running through \mathbb{N} . Then by the monotone class theorem it is sufficient to prove (3.11) for sets A of the form $A = \{\zeta_t^z \in B, \alpha_1 \in B_1, \dots, \alpha_n \in B_n\}$ with $B, B_1, B_2, \dots, B_n \in \mathcal{B}(\mathbb{R})$, $n \geq 1$. Moreover, on the set $\{t < \tau\}$, we have

$$\beta_k := \frac{\gamma_{t_{k-1}}}{\gamma_{t_k}} = \alpha_k, \quad k = 1, \dots, n.$$

Using Proposition 2.3 (i), then for $t < r$ the vectors $(\beta_1, \dots, \beta_n)$ and $(\zeta_t^r, \zeta_{t+h}^r)$ are independent.

Now taking into account all the above considerations, we have

$$\begin{aligned}
\int_{A \cap \{t < \tau\}} f(\zeta_{t+h}) d\mathbb{P} &= \mathbb{E} [f(\zeta_{t+h}) \mathbb{I}_{B \times B_1 \times \dots \times B_n}(\zeta_t, \alpha_1, \dots, \alpha_n) \mathbb{I}_{\{t < \tau\}}] \\
&= \mathbb{E}[f(\zeta_{t+h}) \mathbb{I}_{B \times B_1 \times \dots \times B_n}(\zeta_t, \beta_1, \dots, \beta_n) \mathbb{I}_{\{t < \tau\}}] \\
&= \int_{(t, \infty)} \mathbb{E} [f(\zeta_{t+h}^r) \mathbb{I}_B(\zeta_t^r) \mathbb{I}_{B_1 \times \dots \times B_n}(\beta_1, \dots, \beta_n)] \mathbb{P}_\tau(dr) \\
&= \int_{(t, \infty)} \mathbb{E}[f(\zeta_{t+h}^r) \mathbb{I}_B(\zeta_t^r)] \mathbb{P}_\tau(dr) \mathbb{E}[\mathbb{I}_{B_1 \times \dots \times B_n}(\beta_1, \dots, \beta_n)] \\
&= \mathbb{E}[f(\zeta_{t+h}) \mathbb{I}_B(\zeta_t) \mathbb{I}_{\{t < \tau\}}] \mathbb{E}[\mathbb{I}_{B_1 \times \dots \times B_n}(\beta_1, \dots, \beta_n)] \\
&= \mathbb{E} [\mathbb{E}[f(\zeta_{t+h}) | \zeta_t] \mathbb{I}_B(\zeta_t) \mathbb{I}_{\{t < \tau\}}] \mathbb{E}[\mathbb{I}_{B_1 \times \dots \times B_n}(\beta_1, \dots, \beta_n)] \\
&= \mathbb{E}[\mathbb{E}[f(\zeta_{t+h}) | \zeta_t] \mathbb{I}_B(\zeta_t) \mathbb{I}_{\{t < \tau\}} \mathbb{I}_{B_1 \times \dots \times B_n}(\beta_1, \dots, \beta_n)] \\
&= \mathbb{E}[\mathbb{E}[f(\zeta_{t+h}) | \zeta_t] \mathbb{I}_B(\zeta_t) \mathbb{I}_{\{t < \tau\}} \mathbb{I}_{B_1 \times \dots \times B_n}(\alpha_1, \dots, \alpha_n)] \\
&= \mathbb{E}[\mathbb{E}[f(\zeta_{t+h}) | \zeta_t] \mathbb{I}_{B \times B_1 \times \dots \times B_n}(\zeta_t, \alpha_1, \dots, \alpha_n) \mathbb{I}_{\{t < \tau\}}] \\
&= \int_{A \cap \{t < \tau\}} \mathbb{E}[f(\zeta_{t+h}) | \zeta_t] d\mathbb{P}.
\end{aligned}$$

Hence (3.11) is proved and this ends the proof. \square

Corollary 3.8. *The Markov property can be extended to the completed filtration $\mathbb{F}^{\zeta, c}$.*

The aim of this proposition is to provide, using the Markov property, that the observation of ζ_t would be sufficient to give estimates of the time τ based on the observation of the information process ζ up to time t .

Proposition 3.9. *Let $0 < t < u$.*

(i) *For each bounded measurable function g defined on $(0, \infty)$, we have \mathbb{P} -a.s.*

$$\mathbb{E}[g(\tau) | \mathcal{F}_t^{\zeta, c}] = g(\tau \wedge t) \mathbb{I}_{\{\zeta_t = 1\}} + \int_{(t, +\infty)} g(r) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}. \quad (3.12)$$

(ii) For each bounded measurable function defined on $(0, +\infty) \times \mathbb{R}$, we have \mathbb{P} -a.s.

$$\mathbb{E}[g(\tau, \zeta_t) | \mathcal{F}_t^{\zeta, c}] = g(\tau \wedge t, 1) \mathbb{I}_{\{\zeta_t=1\}} + \int_{(t, +\infty)} g(r, \zeta_t) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}. \quad (3.13)$$

$$\begin{aligned} \mathbb{E}[g(\tau, \zeta_u) | \mathcal{F}_t^{\zeta, c}] &= g(\tau \wedge t, 1) \mathbb{I}_{\{\zeta_t=1\}} + \int_{(t, u]} g(r, 1) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}} \\ &+ \int_{(u, +\infty)} \int_{\mathbb{R}} g(r, y) \mathbb{P}_{\zeta_u^r | \zeta_t^r = x}(dy) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}. \end{aligned} \quad (3.14)$$

Proof. (i) Obviously, we have

$$\mathbb{E}[g(\tau) | \mathcal{F}_t^{\zeta, c}] = \mathbb{E}[g(\tau \wedge t) \mathbb{I}_{\{\tau \leq t\}} | \mathcal{F}_t^{\zeta, c}] + \mathbb{E}[g(\tau \vee t) \mathbb{I}_{\{t < \tau\}} | \mathcal{F}_t^{\zeta, c}].$$

Now since $g(\tau \wedge t) \mathbb{I}_{\{\tau \leq t\}}$ is $\mathcal{F}_t^{\zeta, c}$ -measurable then, \mathbb{P} -a.s, one has

$$\begin{aligned} \mathbb{E}[g(\tau \wedge t) \mathbb{I}_{\{\tau \leq t\}} | \mathcal{F}_t^{\zeta, c}] &= g(\tau \wedge t) \mathbb{I}_{\{\tau \leq t\}} \\ &= g(\tau \wedge t) \mathbb{I}_{\{\zeta_t=1\}}. \end{aligned}$$

On the other hand due to the facts that $g(\tau \vee t) \mathbb{I}_{\{t < \tau\}}$ is $\sigma(\zeta_s, t \leq s \leq +\infty) \vee \mathcal{N}_P$ -measurable and ζ is a Markov process with respect to its completed natural filtration we obtain \mathbb{P} -a.s.

$$\mathbb{E}[g(\tau \vee t) \mathbb{I}_{\{t < \tau\}} | \mathcal{F}_t^{\zeta, c}] = \mathbb{E}[g(\tau \vee t) \mathbb{I}_{\{t < \tau\}} | \zeta_t],$$

The result is deduced from (3.2).

(ii) The equation (3.13) is an immediate consequence of (i). Concerning the equation (3.14) we use the same method which we used to prove (i). \square

Remark 3.10. The process ζ cannot be an homogeneous $\mathbb{F}^{\zeta, z}$ -Markov process. Indeed, Proposition 3.9 enables us to see that, for $A \in \mathcal{B}(\mathbb{R})$ and $t < u$, we have \mathbb{P} -a.s.,

$$\begin{aligned} \mathbb{P}(\zeta_u \in A | \mathcal{F}_t^{\zeta}) &= \mathbb{I}_{\{1 \in A\}} \mathbb{I}_{\{\zeta_t=1\}} + \mathbb{I}_{\{1 \notin A\}} \int_{(t, u]} \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}} \\ &+ \int_{(u, +\infty)} \int_A \mathbb{I}_{\{\zeta_t < y < 1\}} \mathbb{P}_{\zeta_u^r | \zeta_t^r = x}(dy) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_t < 1\}}, \end{aligned}$$

which is clear that it doesn't depend only on $u - t$.

3.3 Markov property with respect to $\mathbb{F}_+^{\zeta, c}$

We have established, in the previous section, the Markov property of ζ with respect to its completed natural filtration $\mathbb{F}^{\zeta, c}$. In this section we are interested in the the Markov property of ζ with respect to $\mathbb{F}_+^{\zeta, c}$. It has an interesting consequence which is none other than the filtration $\mathbb{F}_+^{\zeta, c}$ satisfies the usual conditions of completeness and right-continuity. However, we need the following condition of on the integrability of τ .

Assumption 3.2. *There exists a sufficiently small $\alpha > 0$ such that*

$$\mathbb{E}(\tau^\alpha) < +\infty. \quad (3.15)$$

The next theorem shows the Markov property of ξ with respect to $\mathbb{F}_+^{\xi,c}$.

Theorem 3.11. *The process ζ is a Markov process with respect to $\mathbb{F}_+^{\zeta,c}$.*

Proof. It is sufficient to prove that for any $0 \leq t < u$ and any function bounded continuous g we have

$$\mathbb{E}[g(\zeta_u)|\mathcal{F}_{t+}^{\zeta,c}] = \mathbb{E}[g(\zeta_u)|\zeta_t], \quad \mathbb{P} - a.s. \quad (3.16)$$

Let $(t_n)_{n \in \mathbb{N}}$ be a decreasing sequence of strictly positive real numbers converging to t : that is $0 \leq t < \dots < t_{n+1} < t_n < \dots < t_1 < u$, $t_n \searrow t$ as $n \rightarrow +\infty$. Since g is bounded and $\mathcal{F}_{t+}^{\zeta,c} = \bigcap_n \mathcal{F}_{t_n}^{\zeta,c}$ then, \mathbb{P} -a.s., we have

$$\mathbb{E}[g(\zeta_u)|\mathcal{F}_{t+}^{\zeta,c}] = \lim_{n \rightarrow +\infty} \mathbb{E}[g(\zeta_u)|\mathcal{F}_{t_n}^{\zeta,c}]. \quad (3.17)$$

It follows from the Markov property of ζ with respect to $\mathbb{F}^{\zeta,c}$ that

$$\mathbb{E}[g(\zeta_u)|\mathcal{F}_{t+}^{\zeta,c}] = \lim_{n \rightarrow +\infty} \mathbb{E}[g(\zeta_u)|\zeta_{t_n}], \quad \mathbb{P} - a.s. \quad (3.18)$$

It remains to prove that

$$\lim_{n \rightarrow +\infty} \mathbb{E}[g(\zeta_u)|\zeta_{t_n}] = \mathbb{E}[g(\zeta_u)|\zeta_t], \quad \mathbb{P} - a.s. \quad (3.19)$$

The proof is splitted into two parts. In the first one we show the statment (3.16) for $t > 0$, while in the second part we consider the case $t = 0$.

Let $t > 0$. We begin by noticing that from Proposition 3.9, \mathbb{P} -a.s., we have

$$\begin{aligned} \mathbb{E}[g(\zeta_u)|\zeta_{t_n}] &= g(1) \left(\mathbb{I}_{\{\zeta_{t_n}=1\}} + \int_{(t_n, u]} \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_{t_n} < 1\}} \right) \\ &\quad + \int_{(u, +\infty)} K_{t_n, u}(r, \zeta_{t_n}) \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_{t_n} < 1\}} \\ &= g(1) \left(\mathbb{I}_{\{\tau \leq t_n\}} + \int_{(t_n, u]} \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_\tau(dr) \mathbb{I}_{\{t_n < \tau\}} \right) \\ &\quad + \int_{(u, +\infty)} K_{t_n, u}(r, \zeta_{t_n}) \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_\tau(dr) \mathbb{I}_{\{t_n < \tau\}}. \end{aligned}$$

Where the function $K_{t,u}(r, x)$ is defined on \mathbb{R} by for $0 < t < u < r$

$$\begin{aligned} K_{t,u}(r, x) &:= \mathbb{E}[g(\zeta_u^r)|\zeta_t^r = x] \\ &= \int_{\mathbb{R}} g(y) \mathbb{I}_{\{x < y < 1\}} \mathbb{P}_{\zeta_u^r|\zeta_t^r=x}(dy). \end{aligned} \quad (3.20)$$

Since $\lim_{n \rightarrow +\infty} \mathbb{I}_{\{t_n < \tau\}} = \mathbb{I}_{\{t < \tau\}}$ then assertion (3.19) will be established if we show, \mathbb{P} -a.s on $\{t < \tau\}$, that

$$\lim_{n \rightarrow +\infty} \int_{(t_n, u]} \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_\tau(dr) = \int_{(t, u]} \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr), \quad (3.21)$$

and

$$\lim_{n \rightarrow +\infty} \int_{(u, +\infty)} K_{t_n, u}(r, \zeta_{t_n}) \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_\tau(dr) = \int_{(u, +\infty)} K_{t, u}(r, \zeta_t) \phi_{\zeta_t^r}(\zeta_t) \mathbb{P}_\tau(dr). \quad (3.22)$$

We start by proving assertion (3.21). The integral on the left-hand side of (3.21) can be rewritten as

$$\begin{aligned} \int_{(t_n, u]} \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_\tau(dr) &= \frac{\int_{(t_n, u]} \varphi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_\tau(dr)}{\int_{(t_n, +\infty)} \varphi_{\zeta_{t_n}^s}(\zeta_{t_n}) \mathbb{P}_\tau(ds)} \mathbb{I}_{\{0 < \zeta_{t_n} < 1\}} \\ &= \frac{\int_{(t_n, u]} \left(1 - \zeta_{t_n}\right)^r \frac{\Gamma(r)}{\Gamma(r - t_n)} \mathbb{P}_\tau(dr)}{\int_{(t_n, +\infty)} \left(1 - \zeta_{t_n}\right)^s \frac{\Gamma(s)}{\Gamma(s - t_n)} \mathbb{P}_\tau(ds)} \mathbb{I}_{\{0 < \zeta_{t_n} < 1\}}. \end{aligned}$$

First let us remark that the function

$$(t, r, x) \longrightarrow (1 - x)^r \frac{\Gamma(r)}{\Gamma(r - t)}$$

defined on $\{(t, r) \in (0, +\infty)^2, t < r\} \times (0, 1)$ is continuous. Using the facts that ζ_{t_n} is decreasing to ζ_t and $\mathbb{P}[\zeta_t = 0] = 0$ then, \mathbb{P} -a.s on $\{t < \tau\}$, we have

$$\lim_{n \rightarrow +\infty} (1 - \zeta_{t_n})^r \frac{\Gamma(r)}{\Gamma(r - t_n)} \mathbb{I}_{\{t_n < r\}} \mathbb{I}_{\{0 < \zeta_{t_n} < 1\}} = (1 - \zeta_t)^r \frac{\Gamma(r)}{\Gamma(r - t)} \mathbb{I}_{\{t < r\}} \mathbb{I}_{\{0 < \zeta_t < 1\}}. \quad (3.23)$$

On the other hand since the function $x \mapsto (1 - x)^r$ is decreasing on $(0, 1)$ for all $r > 0$ and

$$0 \leq \frac{\Gamma(r)}{\Gamma(r - t)} = r^t \left[1 - \frac{t(t+1)}{2r} + O\left(\frac{1}{r^2}\right) \right], \quad (3.24)$$

for large enough r , see [1], p. 257, 6.1.46, then for any compact subset \mathcal{K} of $(0, +\infty) \times (0, 1)$ it yields

$$\sup_{(t, x) \in \mathcal{K}} (1 - x)^r \frac{\Gamma(r)}{\Gamma(r - t)} \mathbb{I}_{\{t < r\}} < +\infty.$$

Hence, \mathbb{P} -a.s on $\{t < \tau\}$, we have

$$\sup_{n \in \mathbb{N}} \left(1 - \zeta_{t_n}\right)^r \frac{\Gamma(r)}{\Gamma(r - t_n)} \mathbb{I}_{\{t_n < r\}} \mathbb{I}_{\{0 < \zeta_{t_n} < 1\}} < +\infty. \quad (3.25)$$

We conclude assertion (3.21) from the Lebesgue dominated convergence theorem.

Now let us prove (3.22). Recall that the function $K_{t_n,u}(r, \zeta_{t_n})$ is given by

$$\begin{aligned} K_{t_n,u}(r, \zeta_{t_n}) &= \int_{\mathbb{R}} g(y) \mathbb{I}_{\{x < y < 1\}} \mathbb{P}_{\zeta_u | \zeta_{t_n} = x}(dy) |_{x=\zeta_{t_n}} \\ &= \frac{\Gamma(r - t_n)}{\Gamma(u - t_n) \Gamma(r - u)} \int_{\mathbb{R}} g(y) \frac{(y - \zeta_{t_n})^{u-t-1} (1 - y)^{r-u-1}}{(1 - \zeta_{t_n})^{r-t-1}} \mathbb{I}_{\{\zeta_{t_n} < y < 1\}} dy. \end{aligned}$$

Since g is bounded we deduce that $K_{t_n,u}(r, \zeta_{t_n})$ is bounded. Moreover we obtain from the weak convergence that

$$\lim_{n \rightarrow +\infty} K_{t_n,u}(r, \zeta_{t_n}) = K_{t,u}(r, \zeta),$$

\mathbb{P} -a.s on $\{t < \tau\}$. Combining the fact that $K_{t_n,u}(r, \zeta_{t_n})$ is bounded, (3.23) and (3.25) assertion (3.22) is then derived from the Lebesgue dominated convergence theorem.

Next, we investigate the second part of the proof, that is the case $t = 0$. It will be carried out in two steps. In the first one we assume that there exists $\varepsilon > 0$ such that

$$\mathbb{P}(\tau > \varepsilon) = 1. \quad (3.26)$$

As in the first part, it is sufficient to verify that

$$\lim_{n \rightarrow +\infty} \mathbb{E}[g(\zeta_u) | \zeta_{t_n}] = \mathbb{E}[g(\zeta_u) | \zeta_0], \quad \mathbb{P} - a.s. \quad (3.27)$$

Without loss of generality we assume $t_n < \alpha \wedge \varepsilon$ for all $n \in \mathbb{N}$. It is easy to see that under condition (3.26), $\mathbb{E}[g(\zeta_u) | \zeta_{t_n}]$ takes the form

$$\mathbb{E}[g(\zeta_u) | \zeta_{t_n}] = g(1) \int_{(\varepsilon, u]} \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_{\tau}(dr) + \int_{(u, +\infty)} K_{t_n,u}(r, \zeta_{t_n}) \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_{\tau}(dr).$$

On the other hand we have

$$\mathbb{E}[g(\zeta_u) | \zeta_0] = \mathbb{E}[g(\zeta_u)] = g(1)F(u) + \int_{(u, +\infty)} \int_{\mathbb{R}} g(y) \varphi_{\zeta_t^r}(y) dy \mathbb{P}_{\tau}(dr).$$

Then in order to show (3.27) it is sufficient to prove, \mathbb{P} -a.s, the following

$$\lim_{n \rightarrow +\infty} \int_{(\varepsilon, u]} \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_{\tau}(dr) = F(u), \quad (3.28)$$

and

$$\lim_{n \rightarrow +\infty} \int_{(u, +\infty)} K_{t_n,u}(r, \zeta_{t_n}) \phi_{\zeta_{t_n}^r}(\zeta_{t_n}) \mathbb{P}_{\tau}(dr) = \int_{(u, +\infty)} \int_{\mathbb{R}} g(y) \varphi_{\zeta_t^r}(y) dy \mathbb{P}_{\tau}(dr). \quad (3.29)$$

First, for $r > \varepsilon$, we have

$$\lim_{n \rightarrow +\infty} (1 - \zeta_{t_n})^r \frac{\Gamma(r)}{\Gamma(r - t_n)} \mathbb{I}_{\{t_n < r\}} \mathbb{I}_{\{0 < \zeta_{t_n} < 1\}} = 1.$$

Since the gamma function is increasing on $[2, \infty)$ then, for $r \geq 2 + t_1$, we obtain

$$\sup_{n \in \mathbb{N}} \left(1 - \zeta_{t_n}\right)^r \frac{\Gamma(r)}{\Gamma(r - t_n)} < \frac{\Gamma(r)}{\Gamma(r - t_1)}. \quad (3.30)$$

It follows from (3.15) and (3.24) that the function $r \mapsto \frac{\Gamma(r)}{\Gamma(r - t_1)}$ is \mathbb{P}_τ -integrable on $(\varepsilon, +\infty)$. Hence (3.28) follows from a simple application of the Lebesgue dominated convergence theorem. In the same way as in the first case ($t > 0$) we obtain from the weak convergence that

$$\lim_{n \rightarrow +\infty} K_{t_n, u}(r, \zeta_{t_n}) = \frac{\Gamma(r)}{\Gamma(u)\Gamma(r - u)} \int_{\mathbb{R}} g(y) y^{u-1} (1 - y)^{r-u-1} \mathbb{I}_{\{0 < y < 1\}} dy,$$

then also (3.29) follows from a simple application of the Lebesgue dominated convergence theorem. Finally, we have to consider the general case, that is $\mathbb{P}(\tau > 0) = 1$. In order to prove the Markov property of ζ with respect to $\mathbb{F}_+^{\zeta, c}$ at $t = 0$ it is sufficient to show that $\mathcal{F}_{0+}^{\zeta, c}$ is \mathbb{P} -trivial. This amounts to prove that \mathcal{F}_{0+}^{ζ} is \mathbb{P} -trivial since $\mathcal{F}_{0+}^{\zeta, c} = \mathcal{F}_{0+}^{\zeta} \vee \mathcal{N}_P$. To do so, let $\varepsilon > 0$ be fixed and consider the stopping time $\tau_\varepsilon = \tau \vee \varepsilon$. We define the process $\zeta_t^{\tau_\varepsilon}$ by

$$\{\zeta_t^{\tau_\varepsilon}; t \geq 0\} := \{\zeta_t^r|_{r=\tau \vee \varepsilon}; t \geq 0\}.$$

The first remark is that the sets $(\tau_\varepsilon > \varepsilon) = (\tau > \varepsilon)$ are equal and therefore the following equality of processes holds

$$\zeta^{\tau_\varepsilon} \mathbb{I}_{(\tau > \varepsilon)} = \zeta \cdot \mathbb{I}_{(\tau > \varepsilon)}.$$

Then for each $A \in \mathcal{F}_{0+}^{\zeta}$ there exists $B \in \mathcal{F}_{0+}^{\zeta^{\tau_\varepsilon}}$ such that

$$A \cap (\tau > \varepsilon) = B \cap (\tau > \varepsilon).$$

As $\mathbb{P}(\tau_\varepsilon > \varepsilon/2) = 1$ then according to the previous case we have that $\mathcal{F}_{0+}^{\zeta^{\tau_\varepsilon}}$ is \mathbb{P} -trivial. That is $\mathbb{P}(B) = 0$ or 1 . Consequently we obtain

$$\mathbb{P}(A \cap (\tau > \varepsilon)) = 0 \text{ or } \mathbb{P}(A \cap (\tau > \varepsilon)) = \mathbb{P}(\tau > \varepsilon).$$

Now if $\mathbb{P}(A) > 0$, then there exists $\varepsilon > 0$ such that $\mathbb{P}(A \cap \{\tau > \varepsilon\}) > 0$. Therefore for all $0 < \varepsilon' \leq \varepsilon$ we have

$$\mathbb{P}(A \cap (\tau > \varepsilon')) = \mathbb{P}(\tau > \varepsilon').$$

Passing to the limit as ε' goes to 0 yields $\mathbb{P}(A \cap (\tau > 0)) = \mathbb{P}(\tau > 0) = 1$. It follows that $\mathbb{P}(A) = 1$, which ends the proof. \square

Corollary 3.12. *The filtration $\mathbb{F}^{\zeta, c}$ satisfies the usual conditions of rightcontinuity and completeness.*

Proof. See, e.g., [5], Ch. I, Proposition (8.12) \square

3.4 Semimartingale Decomposition of ζ

Our purpose is to derive the semimartingale property of ζ with respect to its own filtration $\mathbb{F}^{\zeta, c}$. Firstly, we obtain from the representation (2.15) that

$$\zeta_t = \widehat{N}_t + \int_0^t Z_s ds, \quad t \geq 0, \quad (3.31)$$

where the processes \widehat{N} and Z are defined as follows:

$$\widehat{N}_t(\omega) := \widehat{N}_t^r(\omega)|_{r=\tau(\omega)},$$

and

$$Z_t = \frac{1 - \zeta_t}{\tau - t} \mathbb{I}_{\{t < \tau\}},$$

for $(t, \omega) \in \mathbb{R}_+ \times \Omega$. Now let us consider the filtration

$$\mathbb{H} = \left(\mathcal{H}_t := \mathcal{F}_t^{\zeta, c} \vee \sigma(\tau), \quad t \geq 0 \right), \quad (3.32)$$

which is equal to the initial enlargement of the filtration $\mathbb{F}^{\zeta, c}$ by the σ -algebra $\sigma(\tau)$. Since the processes ζ and Z are \mathbb{H} -adapted it follows from equation (3.31) that \widehat{N} is \mathbb{H} -adapted. Moreover, τ is a stopping time with respect to \mathbb{H} . The next proposition will play a very important role in forthcoming developments, since it shows the semimartingale property of ζ with respect to \mathcal{H}_t .

Proposition 3.13. (i) *We have*

$$\mathbb{E} \left[\int_0^t |Z_s| ds \right] < +\infty, \forall t \geq 0.$$

(ii) *The process $\widehat{N} = (\widehat{N}_t, t \geq 0)$ defined by*

$$\widehat{N}_t = \zeta_t - \int_0^t Z_s ds, \quad t \geq 0, \quad (3.33)$$

is a \mathcal{H}_t -martingale stopped at τ .

Proof. (i) We first note that Z is a nonnegative process. Since, for $s \leq r$, ζ_s^r has a beta distribution $\beta(s, r - s)$ then $\mathbb{E}(\zeta_s^r) = s/r$. So, we can see, for any $t \geq 0$, that

$$\begin{aligned} \mathbb{E} \left[\int_0^t Z_s ds \right] &= \int_0^{+\infty} \int_0^{t \wedge r} \frac{1 - \mathbb{E}[\zeta_s^r]}{r - s} ds \mathbb{P}_\tau(dr) \\ &= \int_0^{+\infty} \int_0^{t \wedge r} \frac{1}{r} ds \mathbb{P}_\tau(dr) \leq 1. \end{aligned}$$

(ii) By assertion (i), the process $(Z_t, t \geq 0)$ is integrable with respect to the Lebesgue measure, hence \widehat{N} is well-defined. It is clear that the process \widehat{N} is \mathbb{H} -adapted and $\widehat{N}_t = \widehat{N}_\tau$, \mathbb{P} -a.s, on

the set $\{t \geq \tau\}$. Now since $(\widehat{N}_t^r, t \geq 0)$ is a \mathbb{F}^{ζ^r} -martingale stopped at r we obtain, for any $0 < t_1 < t_2 < \dots < t_n = t$, $n \in \mathbb{N}^*$, $h \geq 0$ and g a bounded Borel function, that

$$\begin{aligned}
\mathbb{E} \left[(\widehat{N}_{t+h} - \widehat{N}_t) g(\zeta_{t_1}, \dots, \zeta_{t_n}, \tau) \right] &= \int_{(0, +\infty)} \mathbb{E}[(\widehat{N}_{t+h}^r - \widehat{N}_t^r) g(\zeta_{t_1}^r, \dots, \zeta_{t_n}^r, r)] \mathbb{P}_\tau(dr) \\
&= \int_{(0, t)} \mathbb{E}[(\widehat{N}_{t+h}^r - \widehat{N}_t^r) g(\zeta_{t_1}^r, \dots, \zeta_{t_n}^r, r)] \mathbb{P}_\tau(dr) \\
&\quad + \int_{[t, t+h)} \mathbb{E}[(\widehat{N}_{t+h}^r - \widehat{N}_t^r) g(\zeta_{t_1}^r, \dots, \zeta_{t_n}^r, r)] \mathbb{P}_\tau(dr) \\
&\quad + \int_{[t+h, +\infty)} \mathbb{E}[(\widehat{N}_{t+h}^r - \widehat{N}_t^r) g(\zeta_{t_1}^r, \dots, \zeta_{t_n}^r, r)] \mathbb{P}_\tau(dr) \\
&= \int_{(0, t)} \mathbb{E}[(\widehat{N}_r^r - \widehat{N}_t^r) g(\zeta_{t_1}^r, \dots, \zeta_{t_n}^r, r)] \mathbb{P}_\tau(dr) \\
&\quad + \int_{[t, t+h)} \mathbb{E}[(\widehat{N}_r^r - \widehat{N}_t^r) g(\zeta_{t_1}^r, \dots, \zeta_{t_n}^r, r)] \mathbb{P}_\tau(dr) \\
&\quad + \int_{[t+h, +\infty)} \mathbb{E}[(\widehat{N}_{t+h}^r - \widehat{N}_t^r) g(\zeta_{t_1}^r, \dots, \zeta_{t_n}^r, r)] \mathbb{P}_\tau(dr) = 0.
\end{aligned}$$

The desired result follows by a standard monotone class argument. This completes the proof. \square

Therefore, it follows from Stricker's Theorem [27] that ζ is a semimartingale relative to its natural filtration $\mathbb{F}^{\zeta, c}$. A natural question is: what is the explicit form of its canonical decomposition? That is the problem we want to discuss. The method consists in applying the stochastic filtering theory.

Theorem 3.14. *The canonical decomposition of ζ in its natural filtration $\mathbb{F}^{\zeta, c}$ is given by*

$$\zeta_t = \widetilde{N}_t + \int_0^t (1 - \zeta_s) \int_{(s, +\infty)} \frac{1}{r - s} \phi_{\zeta_s^r}(\zeta_s) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_s < 1\}} ds, \quad (3.34)$$

where $(\widetilde{N}_t, t \geq 0)$ is an $\mathbb{F}^{\zeta, c}$ -martingale stopped at τ .

Proof. Let us start by recalling that τ is a stopping time with respect to $\mathbb{F}^{\zeta, c}$. A well known result of filtering theory, [6] (T1, p. 87) (or Theorem 8.1.1 and Remark 8.1.1 [18] for more general setting) tells us that the decomposition of ζ in its natural filtration $\mathbb{F}^{\zeta, c}$ is given by

$$\zeta_t = \widetilde{N}_t + \int_0^t \mathbb{E}(Z_s | \mathcal{F}_s^{\zeta, c}) ds, \quad (3.35)$$

where $(\widetilde{N}_t, t \geq 0)$ is an $\mathbb{F}^{\zeta, c}$ -martingale stopped at τ . Therefore, we have only to compute the conditional expectation of Z_s relative to $\mathcal{F}_s^{\zeta, c}$. Indeed, using (3.12) we have

$$\begin{aligned}
\mathbb{E}(Z_s | \mathcal{F}_s^{\zeta, c}) &= \mathbb{E}\left(\frac{1 - \zeta_s}{\tau - s} \mathbb{I}_{\{s < \tau\}} | \mathcal{F}_s^{\zeta, c}\right) = (1 - \zeta_s) \mathbb{E}\left(\frac{1}{\tau - s} \mathbb{I}_{\{s < \tau\}} | \mathcal{F}_s^{\zeta, c}\right) \\
&= (1 - \zeta_s) \int_{(s, +\infty)} \frac{1}{r - s} \phi_{\zeta_s^r}(\zeta_s) \mathbb{P}_\tau(dr) \mathbb{I}_{\{0 < \zeta_s < 1\}}.
\end{aligned}$$

Hence we derive the canonical decomposition (3.34) of ζ as a semimartingale in its own filtration $\mathbb{F}^{\zeta, c}$. \square

Remark 3.15. The results of the paper can be straightforwardly extended to a large class of gamma subordinator $(\gamma_t^{(\eta, \kappa)}, t \geq 0)$, $\eta, \kappa > 0$, with Lévy measure

$$\nu(dx) = \frac{\kappa}{x} \exp(-\eta x) \mathbb{I}_{(0, \infty)}(x) dx$$

and whose law at time t is the gamma distribution with density

$$f_{\gamma_t^{(\eta, \kappa)}}(x) = \frac{\eta^{\kappa t} x^{\kappa t - 1} \exp(-\eta x)}{\Gamma(\kappa t)} \mathbb{I}_{(0, \infty)}(x).$$

The Lévy-Khintchine representation is given by

$$\mathbb{E}(\exp(-\lambda \gamma_t^{(\eta, \kappa)})) = (1 + \frac{\lambda}{\eta})^{-\kappa t}.$$

On the other hand, they can be also easily extended to the gamma bridges of length r , starting at 0, with an arbitrary ending point $a > 0$

$$\zeta_t^r := a \frac{\gamma_{t \wedge r}^{(m)}}{\gamma_r^{(m)}}, \quad t \geq 0.$$

For the sake of simplicity, we have therefore considered only the case $\eta = \kappa = a = 1$ without loss of generality.

References

- [1] Abramowitz, M., Stegun, I. A. (Eds), Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, National Bureau of Standards, Applied Mathematics Series 55, 9th printing, Washington, 1970.
- [2] Asmussen, S., Hobolth, A. Markov Bridges, Bisection and Variance Reduction. In: Plaskota L., Wozniakowski H. (eds) Monte Carlo and Quasi-Monte Carlo Methods 2010. Springer Proceedings in Mathematics and Statistics, vol 23. Springer, Berlin, Heidelberg. 2012.
- [3] Bertoin, J. *Lévy Processes*. Cambridge University Press, Cambridge, 1996.
- [4] Bedini, M. L., Buckdahn, R., Engelbert, H. J. Brownian bridges on random intervals. Theory Probab. Appl. 61 no. 1, 15-39, 2017.
- [5] Blumenthal, R. M., Gettoor, R. K. *Markov Processes and Potential Theory*. Academic Press, New York-London 1968.
- [6] P. Brémaud. *Point Processes and Queues*. Springer Series in Statistics. Springer-Verlag, New York, 1981.

- [7] Brody, D. C., Hughston, L. P., and Macrina, A. Dam rain and cumulative gain. Proceedings of the Royal Society A, 464, 1801-1822, 2008.
- [8] Dufresne, F., Gerber, H. U., Shiu, E. S. Risk theory with the gamma process. ASTIN Bull. 21, 177-192. 1991.
- [9] Chan, Y., Talburt, J., Talley, T. Data engineering: Mining, information and intelligence. Norwell, MA: Springer. 2010.
- [10] Erraoui, M., Louriki, M. Bridges with random length: Gaussian-Markovian case. arXiv:1711.02467v1, 2017.
- [11] Emery, M., Yor, M. A parallel between Brownian bridges and gamma bridges. Publ. RIMS, Kyoto Univ., 40 (3), 669-688, 2004.
- [12] Gani, J. Problems in the probability theory of storage systems. J. Roy. Statist. Soc. Ser. B. 19, 181-206, 1957.
- [13] Hammersley, J. M. Storage problems. Math. Ann. 128, 475-478, 1955.
- [14] Hoyle, E., Hughston, L. P., Macrina, A. Stable- $\frac{1}{2}$ bridges and insurance. Advances in mathematics of finance, 95-120, Banach Center Publ., 104, Polish Acad. Sci. Inst. Math., Warsaw, 2015.
- [15] Hoyle, E., Mengütürk, L. A. Archimedean survival processes. Journal of Multivariate Analysis, 115, 1-15. 2013.
- [16] Jeanblanc M., Le Cam Y. Immersion Property and Credit Risk Modelling. Optimality and Risk Modern Trends in Mathematical Finance, pp. 99-132. Springer Berlin Heidelberg, 2010.
- [17] Jeanblanc M., Le Cam Y. Progressive Enlargement of Filtrations with Initial Times. Stoch. Process. Appl., v. 119, No. 8, 2523-2543. 2009.
- [18] Kallianpur, G. *Stochastic Filtering Theory*. Applications of Mathematics, 13. Springer Verlag, New York-Berlin, 1980.
- [19] Kendall, D. G. Some problems in the theory of dams, J. Roy. Statist. Soc., ser. B, vol. 19, 207-212, 1957.
- [20] Kingman, J. F. C. On continuous time models in the theory of dams. J. Austral. Math. Soc. 3, 480-487, 1963.
- [21] Kyprianou, A. E. *Introductory Lectures on Fluctuations of Lévy Processes with Applications*. Springer Verlag, Berlin, 2006.
- [22] Moran, P. A. P. A probability theory of a dam with a continuous release. Quart. J. Math. Oxford Ser. 7, Issue 1, 130-137, 1956.
- [23] Protter, P. *Stochastic Integration and Differential Equations*. 2nd edn. Springer, Berlin, 2005.

- [24] Revuz D., Yor M. *Continuous Martingales and Brownian Motion*. Springer-Verlag, Berlin, Third edition, 1999.
- [25] Sato, K. I. *Lévy Processes and Infinitely Divisible Distributions*. Cambridge University Press, Cambridge, 1999.
- [26] Shiryaev, A. N. *Probability. 1*. Springer, New York, Third edition, 2016.
- [27] Stricker, C. Quasimartingales, martingales locales, semimartingales et filtration naturelle. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 39, no. 1, 55-63, 1977.
- [28] Yor, M. Some remarkable properties of gamma processes. Advances in mathematical finance, 37-47, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, 2007.