

Infinite Mode Quantum Gaussian States

B.V. RAJARAMA BHAT¹, TIJU CHERIAN JOHN², AND R. SRINIVASAN³

^{1,2}*Indian Statistical Institute, Stat-Math Unit, 8th Mile, Mysore Road, R.V. College Post, Bengaluru-560059, India.*

bhat@isibang.ac.in, tijucherian@gmail.com

³*Chennai Mathematical Institute, H1, SIPCOT IT Park, Siruseri, Kelambakkam, Chennai-603103, India.*

vasanth@cmi.ac.in

March 9, 2024

Abstract

Quantum Gaussian states on Bosonic Fock spaces are quantum versions of Gaussian distributions. In this paper we explore infinite mode quantum Gaussian states. We extend many of the results of Parthasarathy in [Par10] and [Par13] to the infinite mode case, which includes various characterizations, convexity and symmetry properties.

Keywords: Quantum Gaussian states, Williamson's normal form, Infinite mode quantum systems.

2010 Mathematics Subject classification: Primary 81S05; Secondary 46L60.

1 Introduction

There is extensive literature on quasifree states on CCR (Canonical Commutation Relation) algebras variously known as quantum Gaussian states or squeezed states. Some early references being [AS72], [Ara72], [Hol71a], [Hol71b], and [vD71]. Comprehensive treatments and additional references can be seen in monographs [BR97], [Pet90] and [DG13].

Recently finite mode Gaussian states have been getting more attention in the context of its importance in quantum information theory. Some references are [ARL14], [FOP05], [WGC06] and [WHTH07]. K R Parthasarathy wrote an expository article on finite mode Gaussian states in [Par10] and followed it up with a series of articles systematically studying convexity, symmetry and dilation properties (see [Par13, Par15]). The aim of this article is to extend some of these results to the infinite mode case. Now the covariance operators are restricted by some trace class or Hilbert-Schmidt conditions. We take care of these technicalities by developing necessary notation and tools. The infinite mode setting has been studied earlier in different contexts by [HR96, DG13] and others.

The scheme is as follows. In Section 2, we review some terminology and known results. For notation and basics on Boson Fock spaces we follow mostly [Par12].

We need some minor refinements regarding existing results on Shale Operators and Bogoliubov transformations [Sha62, Par12, BS05, HR96]. We crucially make use of the notion of quasi-free states on CCR Algebra and their classification theory. For this we depend on [Pet90], [Hol71a], [Hol71b] and [vD71]. Finally, we need suitable extension of Williamson's normal form to infinite dimensions [BJ19]. The result we need has been quoted in Section 2.4.

Section 3, has the formal definition of quantum Gaussian states in the infinite mode case, by first introducing quantum characteristic function or quantum Fourier transform of states on Boson Fock space. Every quantum Gaussian state comes with a 'covariance matrix', which in our setup is a symmetric and invertible real linear operator. Making use of the theory of quasifree states we derive some necessary conditions on this operator. Some further analysis through explicit constructions finally leads to a complete characterization of covariance operators of quantum Gaussian states in infinite mode (Theorem 3.17).

Now we have the proper setting to extend the results of K R Parthasarathy [Par13] to infinite mode Gaussian states. We characterize extreme points of the set of Gaussian state covariance operators and express every interior point as midpoint of two extreme points (Theorem 4.4). Theorem 5.1 provides explicit description for every quantum Gaussian state, in terms of symplectic eigenvalues of the covariance operator. We see that every mixed Gaussian state can be purified to a pure Gaussian state.

A unitary operator in the Boson Fock space is called a Gaussian symmetry if it preserves Gaussianity of states under conjugation. The last result in this article (Theorem 6.5) is a complete characterization of Gaussian symmetries and shows that symplectic spectrum is a complete invariant for Gaussian states under conjugation by Gaussian symmetries.

2 Preliminaries and Notations

In this section we fix the notation used in this paper and recall some of the well-known results. Throughout we consider only separable Hilbert spaces.

§ 2.1 Symmetric Fock space

Let \mathcal{H} be a complex Hilbert space with inner product $\langle \cdot, \cdot \rangle$, which is anti-linear in the first variable. The symmetric Fock space or the Boson Fock space over \mathcal{H} is defined as

$$\Gamma_s(\mathcal{H}) := \bigoplus_{n=0}^{\infty} \mathcal{H}^{\otimes n}$$

where $\mathcal{H}^{\otimes n}$ is the n -fold symmetric tensor product of \mathcal{H} and $\mathcal{H}^{\otimes 0} := \mathbb{C}$. The n -th direct summand is called the n -particle subspace. When $n = 0$, it is called the *vacuum subspace*. The vector $\Phi := 1 \oplus 0 \oplus 0 \oplus \dots \in \Gamma_s(\mathcal{H})$ is called the *vacuum vector*. Any element in the n -particle subspace is called an n -particle vector. The dense linear manifold generated by the n -particle vectors, $n = 0, 1, 2, \dots$ is called the *finite particle space*, denoted by $\Gamma_s^0(\mathcal{H})$. For $f \in \mathcal{H}$, the exponential vector

$e(f) \in \Gamma_s(\mathcal{H})$ is defined by

$$e(f) = \oplus_{n=0}^{\infty} \frac{f^{\otimes n}}{\sqrt{n!}}, \quad (2.1)$$

where $f^{\otimes 0} := 1$. Then $\langle e(f), e(g) \rangle = \exp \langle f, g \rangle$. Further the set $E := \{e(f) | f \in \mathcal{H}\}$ of all exponential vectors is linearly independent and total in $\Gamma_s(\mathcal{H})$. Indeed if A is a dense set in \mathcal{H} , then the linear span of the set $\{e(f) | f \in A\}$ is dense in $\Gamma_s(\mathcal{H})$.

2.1 Example. We describe a well known identification of $\Gamma_s(\mathbb{C})$ here, this identification prove to be very useful for the later part of this work. One can refer Example 19.8 and Exercise 20.20 in [Par12] for more details. Consider $L^2(\mathbb{R})$, where \mathbb{R} is endowed with the Lebesgue measure. Identify $e(z) \in \Gamma_s(\mathbb{C})$ with the L^2 -function $x \mapsto (2\pi)^{-1/4} \exp\{-4^{-1}x^2 + zx - 2^{-1}z^2\}$, $x \in \mathbb{R}$. Then this identification extends as a Hilbert space isomorphism between $\Gamma_s(\mathbb{C})$ and $L^2(\mathbb{R})$. We may write $\Gamma_s(\mathbb{C}) = L^2(\mathbb{R})$ in this sense.

The Weyl operator, corresponding to an element $f \in \mathcal{H}$, is defined on the set of exponential vectors by

$$W(f)e(g) = \{\exp\left(-\frac{1}{2}\|f\|^2 - \langle f, g \rangle\right)\}e(f+g), \quad (2.2)$$

extends to a unitary operator on $\Gamma_s(\mathcal{H})$. The mapping $f \mapsto W(f)$ from \mathcal{H} into $\mathcal{B}(\Gamma_s(\mathcal{H}))$ is strongly continuous. Further,

$$W(-f) = W(f)^*; \quad W(f)W(g) = \exp(-i \operatorname{Im} \langle f, g \rangle)W(f+g) \quad \forall f, g \in \mathcal{H}, \quad (2.3)$$

where ‘Im’ denotes ‘imaginary part’. We denote by $p(f)$, the Stone generator of the strongly continuous one parameter unitary group $\{W(tf) | t \in \mathbb{R}\}$, that is $W(tf) = e^{-itp(f)}$, $t \in \mathbb{R}$, $f \in \mathcal{H}$. The exponential domain \mathcal{E} , the dense subspace spanned by exponential vectors in $\Gamma_s(\mathcal{H})$, is a core for $p(f)$ for all $f \in \mathcal{H}$. The space of all finite particle vectors, $\Gamma_s^0(\mathcal{H})$ is also a core for $p(f)$ for all f .

Let us fix an orthonormal basis $\{e_j\}$ for \mathcal{H} and let

$$p_j = 2^{-1/2}p(e_j), \quad q_j = -2^{-1/2}p(ie_j) \quad (2.4)$$

$$a_j = 2^{-1/2}(q_j + ip_j) \quad a_j^\dagger = 2^{-1/2}(q_j - ip_j) \quad (2.5)$$

for each $j \in \mathbb{N}$. Then we have the Lie brackets

$$[q_r, p_s] = i\delta_{rs}I, \quad [a_r, a_s^\dagger] = \delta_{rs} \quad (2.6)$$

$\forall r, s \in \mathbb{N}$. Further $\{a_r, r \in \mathbb{N}\}$ and $\{a_r^\dagger, r \in \mathbb{N}\}$ commute among themselves. We call p_j and q_j as the j -th *momentum* and *position* operator, a_j and a_j^\dagger as the j -th *annihilation* and *creation* operator for all $j \in \mathbb{N}$. Let $z \in \mathcal{H}$ be such that $z = \sum_{j=1}^n \alpha_j e_j$, where $\alpha_j = x_j + iy_j$, $x_j, y_j \in \mathbb{R}$, $\forall j$, then

$$W(z) = e^{-i\sqrt{2} \sum_{j=1}^n (x_j p_j - y_j q_j)}; \quad p(z) = \sqrt{2} \sum_{j=1}^n (x_j p_j - y_j q_j).$$

If T is any contraction on \mathcal{H} then we define $\Gamma(T)$ on the exponential vectors by

$$\Gamma(T)(e(f)) = e(Tf), \quad (2.7)$$

which extends to a contraction on $\Gamma_s(\mathcal{H})$. Then $\Gamma(T)$ is called as the *second quantization* of T . It is possible to extend the definition of $\Gamma(T)$ via (2.7) even if T is a contraction mapping \mathcal{H} to a different Hilbert space \mathcal{K} .

If $U : \mathcal{H} \rightarrow \mathcal{K}$ is a unitary then we take $\Gamma_s(U) := \Gamma(U)$. Then $\Gamma_s(U)$ is a unitary and we have

$$\Gamma_s(U)^{-1} = \Gamma_s(U^{-1}); \quad \Gamma_s(U)W(u)\Gamma_s(U)^{-1} = W(Uu). \quad (2.8)$$

Later we will extend the definition of $\Gamma_s(\cdot)$ for a large collection of operators called Shale operators.

If $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$, then there is a unique unitary isomorphism between $\Gamma_s(\mathcal{H})$ and $\Gamma_s(\mathcal{H}_1) \otimes \Gamma_s(\mathcal{H}_2)$ satisfying $e(f \oplus g) \mapsto e(f) \otimes e(g)$. Under this isomorphism we have $W(f \oplus g) = W(f) \otimes W(g)$.

Let $\mathcal{H} = \bigoplus_{n=1}^{\infty} \mathcal{H}_n$, where $\mathcal{H}_n, n = 1, 2, 3, \dots$ is a sequence of Hilbert spaces. Consider the infinite tensor product $\bigotimes_{n=1}^{\infty} \Gamma_s(\mathcal{H}_n)$, constructed using the stabilizing sequence $\{\Phi_n\}$, where $\Phi_n \in \Gamma_s(\mathcal{H}_n)$ is the vacuum vector for every n . Then $\Gamma_s(\mathcal{H}) = \bigotimes_{n=1}^{\infty} \Gamma_s(\mathcal{H}_n)$ under the natural isomorphism. In this identification, for $\bigoplus_{n=1}^{\infty} x_n \in \mathcal{H}$, and contractions $A_n \in \mathcal{B}(\mathcal{H}_n), n \geq 1$,

$$e(\bigoplus_{n=1}^{\infty} x_n) = \bigotimes_{n=1}^{\infty} e(x_n) \quad := \lim_{N \rightarrow \infty} \bigotimes_{j=1}^N e(x_j) \otimes \bigotimes_{n=N}^{\infty} \Phi_n \quad (2.9)$$

$$W(\bigoplus_{n=1}^{\infty} x_n) = \bigotimes_{n=1}^{\infty} W(x_n) \quad := \text{s-lim}_{N \rightarrow \infty} \bigotimes_{j=1}^N W(x_j) \otimes \bigotimes_{n=N}^{\infty} I \quad (2.10)$$

$$\Gamma(\bigoplus_{n=1}^{\infty} A_n) = \bigotimes_{n=1}^{\infty} \Gamma(A_n) \quad := \text{s-lim}_{N \rightarrow \infty} \bigotimes_{j=1}^N \Gamma(A_j) \otimes \bigotimes_{n=N}^{\infty} I \quad (2.11)$$

We refer to Section 20 of [Par12] for more details and for proofs of all the above facts.

§ 2.2 CCR Algebra in Weyl form

In this Section we list some basic facts about symplectic spaces and quasifree states of CCR algebras. For more details see [Pet90, DG13, BR97].

Let H be a real linear space. A bilinear form $\sigma : H \times H \rightarrow \mathbb{R}$ is called a *symplectic form* if $\sigma(f, g) = -\sigma(g, f)$, for every $f, g \in H$. The pair (H, σ) is called a *symplectic space*. A symplectic form σ on H is called *nondegenerate* if $\sigma(f, g) = 0, \forall g \in H$ implies $f = 0$. A symplectic space (H, σ) is called a *standard (symplectic) space* if H is a Hilbert space over \mathbb{C} with respect to some inner product $\langle \cdot, \cdot \rangle$ and $\sigma(\cdot, \cdot) = \text{Im} \langle \cdot, \cdot \rangle$. It is called *separable* if there exists in H a countable family of vectors $\{f_k\}$ such that $\sigma(f, f_k) = 0$ for all k implies $f = 0$. Note that standard symplectic spaces are nondegenerate and separable.

Let (H, σ) be a nondegenerate symplectic space. There exists a universal C^* -algebra generated by unitary elements $\{W(f) : f \in H\}$ satisfying

$$W(-f) = W(f)^*; \quad W(f)W(g) = \exp(i\sigma(f, g))W(f + g), \quad \forall f, g \in H, \quad (2.12)$$

which is unique up to isomorphism. This C^* -algebra is known as the algebra of canonical commutation relations in Weyl form and is denoted by $CCR(H, \sigma)$. In view of (2.12), the linear hull of $\{W(f) : f \in H\}$ is dense in $CCR(H, \sigma)$.

§ 2.2.1 Quasifree States on $CCR(H, \sigma)$

Let \mathcal{A} be a C^* -algebra and let ϕ be a state on it. Let $(H_\phi, \Pi_\phi, \Omega_\phi)$ be the *GNS* triple of (\mathcal{A}, ϕ) . A state ϕ on \mathcal{A} is called *primary* or a *factor state* if the von Neumann algebra $(\Pi_\phi(\mathcal{A}))''$ corresponding to the *GNS*-representation is a factor. It is called *type I* if the von Neumann algebra $(\Pi_\phi(\mathcal{A}))''$ corresponding to the *GNS*-representation is a Type I factor.

2.2 Definition. Two representations Π_1 and Π_2 of \mathcal{A} are said to be *quasiequivalent* if there exists a $*$ -isomorphism of von Neumann algebras

$$\Theta : \Pi_1(\mathcal{A})'' \mapsto \Pi_2(\mathcal{A})''$$

satisfying $\Theta(\Pi_1(x)) = \Pi_2(x)$ for all $x \in \mathcal{A}$. Two states ϕ and ψ on \mathcal{A} are said to be *quasiequivalent* if the corresponding *GNS*-representations are quasiequivalent.

Let (H, σ) be a standard symplectic space and let $\alpha : H \times H \rightarrow \mathbb{R}$ be a real inner product on H with respect to which H is complete. Assume further that there exists a bounded, invertible real linear operator A on (H, α) such that $\alpha(f, g) = \sigma(Af, g)$, for all $f, g \in H$. Then there exists a state (see [Pet90]) ϕ_A on $CCR(H, \sigma)$ such that

$$\phi_A(W(f)) = \exp\left(-\frac{1}{2}\alpha(f, f)\right) \quad \forall f \in H. \quad (2.13)$$

if and only if

$$\sigma(f, g)^2 \leq \alpha(f, f)\alpha(g, g), \quad \forall f, g \in H. \quad (2.14)$$

A state ϕ_A on $CCR(H, \sigma)$ determined in the form of (2.13) is called a *quasifree state*. It is known that any quasifree state ϕ_A is primary (see Proposition 1 and 2 of [Hol71a]). Further in this case,

$$A^\tau = -A; \quad -A^2 - I \geq 0 \quad (2.15)$$

on $(H, \alpha(\cdot, \cdot))$. By A^τ we denote the transpose with respect to the inner product α . Let us denote by H_A the real Hilbert space $(H, \alpha(\cdot, \cdot))$. One major tool of the subject is the following theorem of Holevo.

2.3 Theorem ([Hol71a]). *Two primary quasifree states ϕ_A and ϕ_B on a standard $CCR(H, \sigma)$ are quasi equivalent if and only if $A - B$ and $\sqrt{-A^2 - I} - \sqrt{-B^2 - I}$ are Hilbert-Schmidt operators on H_A .*

§ 2.3 Shale maps and Bogoliubov transformations

Let \mathcal{H} be a complex Hilbert space with inner product $\langle \cdot, \cdot \rangle$. Often we consider \mathcal{H} as real Hilbert space also with $\langle \cdot, \cdot \rangle_{\mathbb{R}} = \text{Re} \langle \cdot, \cdot \rangle$. Let $H \subset \mathcal{H}$ be a real subspace such that \mathcal{H} is the complexification $H + iH$ of H . For any real linear operator S on \mathcal{H} , define operators S_{ij} on H such that

$$S(x + iy) = S_{11}x + iS_{21}x + S_{12}y + iS_{22}y.$$

Define the operator S_0 on $H \oplus H$ by $S_0 = \begin{bmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{bmatrix}$. We identify S with S_0 as a real linear operator and often switch between them freely. We do the same identification

for any $S : \mathcal{H} \rightarrow \mathcal{K}$, for some other complex Hilbert space $\mathcal{K} = K + iK$. Note that if S is a complex linear operator then $S_{11} = S_{22}$ and $S_{12} = -S_{21}$.

Let J be the operator of multiplication by $-i$ on \mathcal{H} considered as a real linear map. Then

$$J_0 = \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix}.$$

Clearly $J^* = J^{-1} = -J$ or equivalently $J_0^\tau = J_0^{-1} = -J_0$.

A real linear bijective map $L : \mathcal{H} \rightarrow \mathcal{K}$ is said to be a *symplectic transformation* if it satisfies

- (i) L and L^{-1} are continuous (bounded) ;
- (ii) $\text{Im} \langle Lz, Lw \rangle = \text{Im} \langle z, w \rangle$ for all $z, w \in \mathcal{H}$.

Correspondingly, L_0 will also be called as symplectic transformation. If $\mathcal{K} = \mathcal{H}$ then we call it a symplectic automorphism.

2.4 Proposition (Section 22 in [Par12]). *$L : \mathcal{H} \rightarrow \mathcal{K}$ is symplectic if and only if*

$$L_0^\tau J_0 L_0 = J_0.$$

Here J_0 on left side is on $K \oplus K$ and that on the right side is on $H \oplus H$.

2.5 Example. Let $A \in \mathcal{B}(H, K)$ be any invertible operator on H then the operator $T : \mathcal{H} \mapsto \mathcal{K}$ defined by $T(u + iv) = Au + i(A^*)^{-1}v$ is a symplectic transformation of \mathcal{H} .

We will have occasions to deal with the complexification of complex Hilbert spaces, considered as real Hilbert spaces under the inner product $\text{Re}\langle \cdot, \cdot \rangle$. We saw above that $(\mathcal{H}, \text{Re}\langle \cdot, \cdot \rangle)$ has a canonical isomorphism to $H \oplus H$ as a real Hilbert space. Let $\hat{\mathcal{H}}$ denote its complexification, which is $\mathcal{H} \oplus \mathcal{H}$. If A is a real linear operator on \mathcal{H} then we denote by \hat{A} , the complexification of A , defined by $\hat{A}(z + iw) = Az + iAw$. It is easy to verify, for a real linear operator S on \mathcal{H} with $S_0 = \begin{bmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{bmatrix}$, that

$$\hat{S} = \begin{bmatrix} \hat{S}_{11} & \hat{S}_{12} \\ \hat{S}_{21} & \hat{S}_{22} \end{bmatrix} \text{ on } \mathcal{H} \oplus \mathcal{H}. \text{ In particular we have } \hat{J} = \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix} \text{ on } \mathcal{H} \oplus \mathcal{H}.$$

By the same analysis as in Proposition 22.1 in [Par12] and by applying polar decomposition, we arrive at the following Proposition.

2.6 Proposition. *Let \mathcal{H}, \mathcal{K} be complex Hilbert spaces and let $S : \mathcal{H} \rightarrow \mathcal{K}$ be symplectic. Then it admits a decomposition:*

$$S = UTV \tag{2.16}$$

where $U : \mathcal{H} \rightarrow \mathcal{K}$ and $V : \mathcal{H} \rightarrow \mathcal{H}$ are unitaries and $T : \mathcal{H} \rightarrow \mathcal{H}$ has the form

$$T(u + iv) = Au + iA^{-1}v,$$

where $A \in \mathcal{B}(H)$ is a positive and invertible operator.

Suppose \mathcal{H} is a complex Hilbert space. We look at automorphisms of the collection of Weyl operators on $\Gamma_s(\mathcal{H})$. Shale's theorem [Sha62] characterizes the set of all bounded, invertible real linear maps $L : \mathcal{H} \rightarrow \mathcal{H}$ which admit some unitary operator Z (unique up to a scalar multiple of modulus one) on the Fock space $\Gamma_s(\mathcal{H})$ such that $ZW(u)Z^* = W(Lu)$, $u \in \mathcal{H}$. A lucid presentation of this can be seen in Section 22 of [Par12]. We state the theorem here with some obvious modifications required for our work and sketch a proof of the same based on the construction given in [BS05].

Let \mathcal{H}, \mathcal{K} be two Hilbert spaces, define the set of *Shale Operators* from \mathcal{H} to \mathcal{K} , $\mathcal{S}(\mathcal{H}, \mathcal{K})$ by

$$\mathcal{S}(\mathcal{H}, \mathcal{K}) = \{L \in \mathcal{B}_{\mathbb{R}}(\mathcal{H}, \mathcal{K}) : L \text{ is symplectic and } L^T L - I \text{ is Hilbert-Schmidt.}\}$$

We denote $\mathcal{S}(\mathcal{H}) := \mathcal{S}(\mathcal{H}, \mathcal{H})$. Notice from Example 2.5 that there exists symplectic transformations T such that $T \notin \mathcal{S}(\mathcal{H}, \mathcal{K})$.

2.7 Theorem. 1. *Let $L \in \mathcal{B}_{\mathbb{R}}(\mathcal{H}, \mathcal{K})$ be a symplectic operator. Then there exists a unitary operator $\Gamma_s(L) : \Gamma_s(\mathcal{H}) \rightarrow \Gamma_s(\mathcal{K})$ such that*

$$\Gamma_s(L)W(u)\Gamma_s(L)^* = W(Lu), \forall u \in \mathcal{H}, \quad (2.17)$$

if and only if $L \in \mathcal{S}(\mathcal{H}, \mathcal{K})$. In such a case, $\Gamma_s(L)$ is determined uniquely up to a scalar of modulus unity.

2. *A unitary $\Gamma_s(L)$ satisfying (2.17) can be chosen such that it satisfies*

$$\langle \Gamma_s(L)\Phi_{\mathcal{H}}, \Phi_{\mathcal{K}} \rangle \in \mathbb{R}^+, \quad (2.18)$$

where $\Phi_{\mathcal{H}}$ and $\Phi_{\mathcal{K}}$ are vacuum vectors in $\Gamma_s(\mathcal{H})$ and $\Gamma_s(\mathcal{K})$ respectively, this choice makes $\Gamma_s(L)$ unique. In this case,

$$\Gamma_s(L^{-1}) = \Gamma_s(L)^* \quad (2.19)$$

3. *Let $\mathcal{H}_1, \mathcal{H}_2, \mathcal{H}_3$ be three Hilbert spaces and $L_1 \in \mathcal{S}(\mathcal{H}_1, \mathcal{H}_2)$, $L_2 \in \mathcal{S}(\mathcal{H}_2, \mathcal{H}_3)$. Then*

$$\Gamma_s(L_2 L_1) = \sigma(L_2, L_1) \Gamma_s(L_2) \Gamma_s(L_1), \quad (2.20)$$

where $\sigma(L_2, L_1) \in \mathbb{C}$, $|\sigma(L_2, L_1)| = 1$.

Proof. 1. Assume that $L \in \mathcal{S}(\mathcal{H}, \mathcal{K})$. We will prove the existence of $\Gamma_s(L)$ based on the construction in [BS05]. By Proposition 2.6 there exist unitaries $U : \mathcal{H} \rightarrow \mathcal{K}$, $V : \mathcal{H} \rightarrow \mathcal{H}$ such that $L = UTV$ where T is a symplectic automorphism of H such that

$$T(u + iv) = Au + iA^{-1}v$$

where $A \in \mathcal{B}(H)$ is positive and invertible. It can be seen from the proof of Proposition 2.6 that

$$L_0 = U_0 \begin{bmatrix} A & 0 \\ 0 & A^{-1} \end{bmatrix} V_0$$

for some orthogonal transformations $U_0 \in \mathcal{B}(H, K)$ and $V_0 \in \mathcal{B}(H)$. Now it can be seen that

$$L_0^T L_0 = V_0^{-1} \begin{bmatrix} A^2 & 0 \\ 0 & A^{-2} \end{bmatrix} V_0.$$

Therefore $L_0^T L_0 - I = V_0^{-1} \left(\begin{bmatrix} A^2 & 0 \\ 0 & A^{-2} \end{bmatrix} - \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix} \right) V_0$. Hence we get that $A^2 - I$ is Hilbert-Schmidt and since A is positive, Theorem 2.1 of [BS05] applies. Thus there exists $\Gamma_s(T)$ such that

$$\Gamma_s(T)W(u)\Gamma_s(T)^* = W(Tu), \forall u \in \mathcal{H}, \quad (2.21)$$

$$\langle \Gamma_s(T)\Phi_{\mathcal{H}}, \Phi_{\mathcal{K}} \rangle \in \mathbb{R}^+. \quad (2.22)$$

Define

$$\Gamma_s(L) := \Gamma_s(U)\Gamma_s(T)\Gamma_s(V), \quad (2.23)$$

where $\Gamma_s(U)$ and $\Gamma_s(V)$ are the second quantization associated with the unitary U and V . A direct computation shows that $\Gamma_s(L)$ satisfies the (2.17) (because of properties of $\Gamma_s(U), \Gamma_s(V)$ and equation 2.22) and (2.18) (because second quantizations $\Gamma_s(U_j)$ acts as identity on vacuum vector). We refer to Theorem 22.11 in [Par12] for the necessity part.

2. Equation (2.18) is automatically satisfied in our construction in (1) above because of (2.22). To see the uniqueness, let $\Gamma_s^1(L)$ and $\Gamma_s^2(L)$ satisfy (2.17) and (2.18). Therefore we get $\Gamma_s^2(L)^*\Gamma_s^1(L)W(u) = W(u)\Gamma_s^2(L)^*\Gamma_s^1(L), \forall u \in \mathcal{H}$. Therefore by irreducibility of Weyl operators (Proposition 20.9 in [Par12]), $\Gamma_s^2(L)^*\Gamma_s^1(L) = cI$ for some complex scalar of unit modulus. But now by (2.18) we get $\Gamma_s^2(L) = \Gamma_s^1(L)$.

To prove (2.19), note that $\langle \Gamma_s(L)^*\Phi_{\mathcal{K}}, \Phi_{\mathcal{H}} \rangle = \langle \Gamma_s(L)\Phi_{\mathcal{H}}, \Phi_{\mathcal{K}} \rangle \in \mathbb{R}^+$ therefore if we show that $\Gamma_s(L)^*W(u)\Gamma_s(L) = W(L^{-1}u)$ then by the uniqueness of $\Gamma_s(L^{-1})$ we get (2.19). Recall from Theorem 2.1 of [BS05] that $\Gamma_s(T^{-1}) = \Gamma_s(T)^*$ and for second quantization (2.7) unitary we have, $\Gamma_s(U_j^*) = \Gamma_s(U_j)^*$. Further by (2.23), and (2.8) we have

$$\begin{aligned} \Gamma_s(L)^*W(u)\Gamma_s(L) &= \Gamma_s(U_2)^*\Gamma_s(T)^*\Gamma_s(U_1)^*W(u)\Gamma_s(U_1)\Gamma_s(T)\Gamma_s(U_2) \\ &= W(U_2^*T^{-1}U_1^*u) \\ &= W(L^{-1}u). \end{aligned}$$

This completes the proof of 2.

3. Follows immediately from 1. □

2.8 Remarks. 1. When \mathcal{H} is finite dimensional, it is known that there exists a choice of $\Gamma_s(L)$ such that the multiplier $\sigma(L_1, L_2) = \pm 1, \forall L_1, L_2 \in Sp_{2n}(\mathbb{R})$, where $Sp_{2n}(\mathbb{R}) \subseteq M_{2n}(\mathbb{R})$ is the subgroup of all $2n \times 2n$ symplectic matrices. This is called the metaplectic representation of the symplectic group. An elementary and self-contained presentation can be found in Chapter 4 of [Fol89], Theorem 4.37 there is of particular interest in this regard. In the infinite dimensional case, [MS04] and [Tve04] are of interest.

2. The map $W(u) \mapsto W(Lu)$ is known as the *Bogoliubov transformation* of the CCR algebra, induced by L . Whenever we write $\Gamma_s(L)$, we mean the unique unitary operator satisfying (2.18). It is called the *Shale unitary* corresponding to an $L \in \mathcal{S}(\mathcal{H}, \mathcal{K})$. It is to be noted that if L is a non-unitary contraction then $\Gamma(L)$ defined by (2.7) is not a unitary and hence in such a case $\Gamma_s(L) \neq \Gamma(L)$.

§ 2.4 Williamson's normal form and symplectic spectrum

The following Theorem [BJ19] extends well-known Williamson's normal form to bounded operators on infinite dimensional real Hilbert spaces.

2.9 Theorem. (*Williamson's Normal Form*) *Let H be a real Hilbert space and let S_0 be a strictly positive invertible operator on $H \oplus H$ then there exists a real Hilbert space K , a positive invertible operator \tilde{P} on K and a symplectic transformation $M_0: H \oplus H \rightarrow K \oplus K$ such that*

$$S_0 = M_0^\tau \begin{bmatrix} \tilde{P} & 0 \\ 0 & \tilde{P} \end{bmatrix} M_0 \quad (2.24)$$

The decomposition is unique in the sense that if P_1 is any strictly positive invertible operator on a Hilbert space \tilde{H} and $\tilde{M}: H \oplus H \rightarrow \tilde{H} \oplus \tilde{H}$ is a symplectic transformation such that

$$S_0 = \tilde{M}^T \begin{bmatrix} P_1 & 0 \\ 0 & P_1 \end{bmatrix} \tilde{M}, \quad (2.25)$$

then P and P_1 are orthogonally equivalent.

For the work in this article we need a version of Williamson's Normal form which produces a diagonalization within the original Hilbert space \mathcal{H} itself. We will identify \mathcal{H} and \mathcal{K} accordingly for this purpose. We obtain this goal in Corollary 2.10. The scheme is as described below.

Since M_0 in (2.24) is an invertible operator between $H \oplus H$ and $K \oplus K$; H and K have same dimension. Let $\{h_j\}$ and $\{k_j\}$ be a basis for H and K respectively. Since $\mathcal{H} = H + iH$ and $\mathcal{K} = K + iK$, there exists a unitary, $U: \mathcal{K} \rightarrow \mathcal{H}$ such that $U(h_j) = k_j$. If we write $\tilde{\mathcal{P}}_0 = \begin{bmatrix} \tilde{P} & 0 \\ 0 & \tilde{P} \end{bmatrix}$, then $U\tilde{\mathcal{P}}U^* := \mathcal{P}$ is a complex linear, positive and invertible operator on \mathcal{H} . Recall the relationship between the operator \mathcal{P} defined on \mathcal{H} and \mathcal{P}_0 on H described at the beginning of § 2.3. Notice now, for the corresponding real linear map \mathcal{P}_0 , we have $\mathcal{P}_0 = \begin{bmatrix} P & 0 \\ 0 & P \end{bmatrix}$ for some positive invertible operator P on H . Now we have the following Corollary.

2.10 Corollary. *Let $\mathcal{H} = H + iH$ and S be a real linear positive, invertible operator on a complex Hilbert space \mathcal{H} . Then there exists a complex linear positive invertible operator \mathcal{P} (unique up to a unitary conjugation) and a symplectic automorphism L on \mathcal{H} such that*

$$S = L^\tau \mathcal{P} L. \quad (2.26)$$

Further, \mathcal{P} has the property that $\mathcal{P}_0 = \begin{bmatrix} P & 0 \\ 0 & P \end{bmatrix}$ and thus we also have

$$S_0 = L_0^\tau \begin{bmatrix} P & 0 \\ 0 & P \end{bmatrix} L_0. \quad (2.27)$$

Proof. We have $S_0 = M_0^\tau \tilde{\mathcal{P}}_0 M_0$ by Theorem 2.9. Therefore $S = M^\tau U^* U \tilde{\mathcal{P}} U^* U M$. Take $L = U M$. It should be noticed that the transpose of U considered as a real linear operator is same as the U^* . \square

2.11 Remark. Under the situation of Corollary 2.10, in view of the uniqueness of the decomposition, the spectrum of \mathcal{P} , can be defined as the *symplectic spectrum* of the positive invertible operator S .

3 Quantum Gaussian States

In this Section we present the basics of infinite mode quantum Gaussian states. Perhaps most of the results in this Section are known. Since notation as well as the set up differ in different sources (some of them consider only complex linear operators), for the convenience of the reader we present some of the steps.

Recall that a *state (or density operator)* ρ on a Hilbert space \mathcal{H} is a bounded positive operator of unit trace. Such a density operator uniquely determines a state on the C^* -algebra $\mathcal{B}(\mathcal{H})$ by $Y \mapsto \text{Tr } \rho Y$, $Y \in \mathcal{B}(\mathcal{H})$.

3.1 Definition. Let $\rho \in \mathcal{B}(\Gamma_s(\mathcal{H}))$ be a density operator. Then the complex valued function $\hat{\rho}$ on \mathcal{H} defined by

$$\hat{\rho}(z) = \text{Tr } \rho W(z), \quad z \in \mathcal{H} \quad (3.1)$$

is called the *quantum characteristic function* (or *quantum Fourier transform*) of ρ .

As the von Neumann algebra generated by Weyl unitaries is the algebra of all bounded operators it follows that the map $\rho \rightarrow \hat{\rho}$ is injective. Here is an analogue of well-known positive definiteness of characteristic functions of probability measures. If ρ is any density operator, then the kernel K_ρ on $\mathcal{H} \times \mathcal{H}$ defined by $K_\rho(z, w) = e^{i \text{Im} \langle z, w \rangle} \hat{\rho}(w - z)$ is positive definite, i.e. $\sum_{j,k=1}^n \overline{c_j} c_k K_\rho(z_j, z_k) \geq 0$, for all $(c_1, c_2, \dots, c_n) \in \mathbb{C}^n$, $n \in \mathbb{N}$. The proof is a direct computation making use of commutation relations of Weyl unitaries (see [Par10]).

If ρ is a state so is any unitary conjugation of it. By using Theorem 2.7, proof of the following proposition follows in the same way as that of Proposition 2.5 in [Par10].

3.2 Proposition. If ρ is a state on $\Gamma_s(\mathcal{K})$ and $L \in \mathcal{S}(\mathcal{H}, \mathcal{K})$ then

$$\{\Gamma_s(L)^* \rho \Gamma_s(L)\}^\wedge(\beta) = \hat{\rho}(L\beta).$$

Further, for every $\alpha \in \mathcal{H}$,

$$\{W(\alpha) \rho W(\alpha)^{-1}\}^\wedge(\beta) = \hat{\rho}(\beta) e^{2i \text{Im} \langle \alpha, \beta \rangle}.$$

3.3 Definition. Let $\rho \in \mathcal{B}(\Gamma_s(\mathcal{H}))$ be a state, ρ is said to be Gaussian if there exists $w \in \mathcal{H}$ and a symmetric, invertible $S \in \mathcal{B}_\mathbb{R}(\mathcal{H})$ such that

$$\hat{\rho}(z) = \exp \left\{ -i \text{Re} \langle w, z \rangle - \frac{1}{2} \text{Re} \langle z, Sz \rangle \right\}, \forall z \in \mathcal{H}. \quad (3.2)$$

In such a case we write $\rho = \rho_g(w, S)$.

Note that this definition determines a real linear functional $z \mapsto \text{Re} \langle w, z \rangle$ and a bounded quadratic form $z \mapsto \text{Re} \langle z, Sz \rangle$ on the real Hilbert space \mathcal{H} . Hence w and S are uniquely determined by the definition. We call w the *mean vector* and S the *covariance operator* associated with ρ . Suppose $\mathcal{H} = H + iH$, where H is a real subspace and let $w = \sqrt{2}(l - im)$, then we call l and m as *mean momentum vector* and *mean position vector* respectively.

Let $\mathbf{G}(\mathcal{H})$ denote the set of all Gaussian states on $\Gamma_s(\mathcal{H})$ and $\mathcal{K}_\mathbf{G}(\mathcal{H})$ denote the set of all Gaussian covariance operators on \mathcal{H} . We will characterize the elements of $\mathcal{K}_\mathbf{G}(\mathcal{H})$ in Theorem 3.17.

3.4 Examples. 1. For $f \in \mathcal{H}$, consider the normalized exponential vector $\psi(f) := e^{-\frac{1}{2}\|f\|^2}e(f)$. The pure state $|\psi(f)\rangle\langle\psi(f)|$ is called as the *coherent state*. The same arguments as that of Proposition 2.9 in [Par10], prove that any coherent state is a pure Gaussian state on $\Gamma_s(\mathcal{H})$ and

$$(|\psi(f)\rangle\langle\psi(f)|)^\wedge(z) = \exp\left\{-2i \operatorname{Im} \langle z, f \rangle - \frac{1}{2}\|z\|^2\right\} \quad (3.3)$$

In particular $|e(0)\rangle\langle e(0)| = \rho_g(0, I)$.

2. Let L be a symplectic automorphism on \mathcal{H} such that $L^\tau L - I$ is Hilbert-Schmidt. Define $\psi_L = \Gamma_s(L)^*|e(0)\rangle$. Then

$$\begin{aligned} (|\psi_L\rangle\langle\psi_L|)^\wedge(z) &= \operatorname{Tr}(|\psi_L\rangle\langle\psi_L| W(z)) \\ &= \langle\psi_L, W(z)\psi_L\rangle \\ &= \langle e(0), \Gamma_s(L)W(z)\Gamma_s(L)^*e(0)\rangle \\ &= \langle e(0), W(Lz)e(0)\rangle \\ &= e^{-\frac{1}{2}\langle z, L^\tau Lz \rangle}. \end{aligned}$$

Therefore, $|\psi_L\rangle\langle\psi_L| = \rho_g(0, L^\tau L)$.

3. Consider $\Gamma_s(\mathbb{C}) = L^2(\mathbb{R})$, under this isomorphism there is an orthonormal basis, $\{\psi_n\}$ of $L^2(\mathbb{R})$, in which $e(z) = \sum_{n=0}^{\infty} \frac{z^n}{\sqrt{n!}}\psi_n$. Let us notate the annihilation and creation operators as a and a^\dagger respectively. Then the number operator, $a^\dagger a$ satisfies $a^\dagger a\psi_n = n\psi_n, \forall n \in \mathbb{N}$ and

$$\operatorname{Tr} e^{-sa^\dagger a} = (1 - e^{-s})^{-1}, \quad s > 0.$$

Therefore the states

$$\rho_s = (1 - e^{-s})e^{-sa^\dagger a}, \quad s > 0 \quad (3.4)$$

are well defined. By Proposition 2.12 in [Par10] we have

$$\hat{\rho}_s(z) = \exp\left\{-\frac{1}{2}(\coth \frac{s}{2})|z|^2\right\}. \quad (3.5)$$

Therefore $\rho_s \in \mathbf{G}(\mathbb{C})$. Since the spectrum of $a^\dagger a$ is $\{0, 1, 2, \dots\}$, it is not a pure state. It may be noted that $e^{-sa^\dagger a} = \Gamma(e^{-s})$, the second quantization (2.7) of the contraction $e^{-s}I$.

All the statements in the following Proposition follow by direct computations.

3.5 Proposition. (i) Let $\alpha \in \mathcal{H}$. Then

$$W(\alpha)\rho_g(w, S)W(\alpha)^{-1} = \rho_g(w - 2i\alpha, S).$$

In particular,

$$W\left(\frac{-i}{2}w\right)\rho_g(w, S)W\left(\frac{-i}{2}w\right)^{-1} = \rho_g(0, S).$$

(ii) Let $\rho_1 = \rho_g(w, S_1)$ and $\rho_2 = \rho_g(w_2, S_2)$ be Gaussian states on $\Gamma_s(\mathcal{H}_1)$ and $\Gamma_s(\mathcal{H}_2)$ respectively. Then $\rho_1 \otimes \rho_2 = \rho_g(w_1 \oplus w_2, S_1 \oplus S_2)$.

(iii) If $\rho = \rho_g(w, S)$ on $\Gamma_s(\mathcal{K})$ and $L \in \mathcal{S}(\mathcal{H}, \mathcal{K})$ then

$$\Gamma_s(L)^* \rho \Gamma_s(L) = \rho_g(L^\tau w, L^\tau S L).$$

The above proposition asserts that the Weyl conjugation displaces the mean (hence the name displacement operator in physics literature) and the Shale unitary transforms the covariance operator.

§ 3.1 Necessary conditions on the covariance operator

We would like to characterize covariance operators of quantum Gaussian states. In the following lemma we see that if S is one such operator on \mathcal{H} then $\hat{S} - i\hat{J} \geq 0$ on $\hat{\mathcal{H}}$. It is to be noted that here \mathcal{H} is a complex Hilbert space, but we are considering it as a real Hilbert space (under the real part of the given complex inner product) and then complexifying it to get $\hat{\mathcal{H}}$.

3.6 Lemma. *Let S be a real linear symmetric and invertible operator on \mathcal{H} . Suppose the function $f : \mathcal{H} \rightarrow \mathbb{R}$ defined by $f(z) = e^{-\frac{1}{2}\text{Re}\langle z, Sz \rangle}$ is the quantum characteristic function of a density operator ρ i.e. $S \in \mathcal{K}_{\mathbf{G}}(\mathcal{H})$ then on $\hat{\mathcal{H}}$,*

$$\hat{S} - i\hat{J} \geq 0. \quad (3.6)$$

Proof. We have already noticed that on f being a quantum characteristic function, the kernel

$$K_\rho(\alpha, \beta) = e^{i\text{Im}\langle \alpha, \beta \rangle} f(\beta - \alpha), \quad \alpha, \beta \in \mathcal{H}. \quad (3.7)$$

is positive definite on \mathcal{H} . If $\alpha = x + iy, \beta = u + iv$ where $x, y, u, v \in H$ then $\text{Im}\langle \alpha, \beta \rangle = \langle \begin{pmatrix} x \\ y \end{pmatrix}, J_0 \begin{pmatrix} u \\ v \end{pmatrix} \rangle$ on $H \oplus H$ (c.f. § 2.3) we can rewrite the definition of K_ρ as

$$K_\rho(\alpha, \beta) = \exp \left\{ i \left\langle \begin{pmatrix} x \\ y \end{pmatrix}, J_0 \begin{pmatrix} u \\ v \end{pmatrix} \right\rangle - \left\langle \begin{pmatrix} u-x \\ v-y \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} u-x \\ v-y \end{pmatrix} \right\rangle \right\} \quad (3.8)$$

Now positive definiteness of K_ρ in \mathcal{H} reduces to that of L in $H \oplus H$ where

$$L((x, y), (u, v)) = \exp \left\{ i \left\langle \begin{pmatrix} x \\ y \end{pmatrix}, J_0 \begin{pmatrix} u \\ v \end{pmatrix} \right\rangle - \left\langle \begin{pmatrix} u-x \\ v-y \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} u-x \\ v-y \end{pmatrix} \right\rangle \right\} \quad (3.9)$$

This is equivalent to the positive definiteness of

$$L_t((x, y), (u, v)) = L\left(\sqrt{t}(x, y), \sqrt{t}(u, v)\right)$$

for all $t \geq 0$. But $\{L_t\}$ is a one parameter multiplicative semigroup of kernels on $H \oplus H$. By elementary properties of positive definite kernels as described in Section 1 of [PS72], positive definiteness of $L_t, t \geq 0$ is equivalent to the conditional positive definiteness of

$$N((x, y), (u, v)) = i \left\langle \begin{pmatrix} x \\ y \end{pmatrix}, J_0 \begin{pmatrix} u \\ v \end{pmatrix} \right\rangle - \left\langle \begin{pmatrix} u-x \\ v-y \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} u-x \\ v-y \end{pmatrix} \right\rangle$$

or equivalently (by the Lemma 1.7 in [PS72]), the positive definiteness of

$$\begin{aligned}
& N((x, y), (u, v)) - N((x, y), (0, 0)) - N((0, 0), (u, v)) - N((0, 0), (0, 0)) \\
&= i \langle \begin{pmatrix} x \\ y \end{pmatrix}, J_0 \begin{pmatrix} u \\ v \end{pmatrix} \rangle - \left\langle \begin{pmatrix} u-x \\ v-y \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} u-x \\ v-y \end{pmatrix} \right\rangle + \left\langle \begin{pmatrix} x \\ y \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} x \\ y \end{pmatrix} \right\rangle + \left\langle \begin{pmatrix} u \\ v \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} u \\ v \end{pmatrix} \right\rangle \\
&= i \langle \begin{pmatrix} x \\ y \end{pmatrix}, J_0 \begin{pmatrix} u \\ v \end{pmatrix} \rangle + \left\langle \begin{pmatrix} x \\ y \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} u \\ v \end{pmatrix} \right\rangle + \left\langle \begin{pmatrix} u \\ v \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} x \\ y \end{pmatrix} \right\rangle \\
&= i \langle \begin{pmatrix} x \\ y \end{pmatrix}, J_0 \begin{pmatrix} u \\ v \end{pmatrix} \rangle + \left\langle \begin{pmatrix} x \\ y \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} u \\ v \end{pmatrix} \right\rangle + \left\langle \begin{pmatrix} x \\ y \end{pmatrix}, \frac{1}{2} S_0 \begin{pmatrix} u \\ v \end{pmatrix} \right\rangle \tag{3.10}
\end{aligned}$$

$$\begin{aligned}
&= i \langle \begin{pmatrix} x \\ y \end{pmatrix}, J_0 \begin{pmatrix} u \\ v \end{pmatrix} \rangle + \langle \begin{pmatrix} x \\ y \end{pmatrix}, S_0 \begin{pmatrix} u \\ v \end{pmatrix} \rangle \\
&= i \langle \begin{pmatrix} u \\ v \end{pmatrix}, -J_0 \begin{pmatrix} x \\ y \end{pmatrix} \rangle + \langle \begin{pmatrix} u \\ v \end{pmatrix}, S_0 \begin{pmatrix} x \\ y \end{pmatrix} \rangle \tag{3.11}
\end{aligned}$$

where (3.10) follows because the real inner-product is symmetric and S_0 is symmetric, and (3.11) for a similar reason. But $H \oplus H \subset \hat{\mathcal{H}} = (H \oplus H) + i(H \oplus H)$, the positive definiteness of (3.11) lifts to the positive definiteness of

$$M(w, z) := \langle w, \{\hat{S} - i\hat{J}\} z \rangle = \left\langle \begin{pmatrix} u \\ v \end{pmatrix}, -i\hat{J}_0 \begin{pmatrix} x \\ y \end{pmatrix} \right\rangle + \left\langle \begin{pmatrix} u \\ v \end{pmatrix}, \hat{S}_0 \begin{pmatrix} x \\ y \end{pmatrix} \right\rangle \tag{3.12}$$

where M is a kernel defined (as above) in $\mathcal{H} \subset \hat{\mathcal{H}}$. Now, since $\hat{S} - i\hat{J}$ is self-adjoint, the positive definiteness of M in (3.12) is equivalent to (3.6). \square

3.7 Lemma. *Let S be a real linear, invertible operator on \mathcal{H} and $\hat{S} - i\hat{J} \geq 0$ on $\hat{\mathcal{H}}$. Then*

1. $S \geq 0$.
2. If $S = L^\tau \mathcal{P} L$ is the Williamson's normal form associated with S (as in Corollary 2.10), then $\mathcal{P} - I \geq 0$ on \mathcal{H} .
3. There exists a primary quasifree state ϕ on $CCR(\mathcal{H}, \sigma)$ such that

$$\phi(W(z)) = e^{-\frac{1}{2} \operatorname{Re} \langle z, Sz \rangle}. \tag{3.13}$$

Further, $\phi = \phi_A$, where $A = -JS$ (the notation ϕ_A is as in § 2.2.1).

Proof. 1. Note $\hat{S} - i\hat{J} \geq 0$ implies \hat{S} is symmetric, hence S is also symmetric. Let $z, w \in H \oplus H$, so that $z + iw \in \hat{\mathcal{H}}$. Then

$$\begin{aligned}
0 &\leq \langle z + iw, (\hat{S} - i\hat{J})z + iw \rangle \\
&= \operatorname{Re} \langle z, Sz \rangle + i \operatorname{Re} \langle z, Sw \rangle - i \operatorname{Re} \langle w, Sz \rangle + \operatorname{Re} \langle w, Sw \rangle \\
&\quad - i \operatorname{Re} \langle z, Jz \rangle + \operatorname{Re} \langle z, Jw \rangle - \operatorname{Re} \langle w, Jz \rangle - i \operatorname{Re} \langle w, Jw \rangle \\
&= \operatorname{Re} \langle z, Sz \rangle + \operatorname{Re} \langle w, Sw \rangle + 2 \operatorname{Re} \langle z, Jw \rangle \tag{3.14}
\end{aligned}$$

where we used the facts $\operatorname{Re} \langle z, Jz \rangle = 0$ for all z . Letting $z = w$ in the above computation, then we get $S \geq 0$.

2. Let $\mathcal{P}_0 = \begin{bmatrix} P & 0 \\ 0 & P \end{bmatrix}$. Then $\hat{\mathcal{P}} = \begin{bmatrix} \hat{P} & 0 \\ 0 & \hat{P} \end{bmatrix}$ and $\hat{J} = \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix}$ on $\hat{\mathcal{H}} = \mathcal{H} \oplus \mathcal{H}$. $\hat{S} - i\hat{J} \geq 0$ implies $\hat{L}^\tau \hat{\mathcal{P}} \hat{L} - i\hat{J} \geq 0$. By a conjugation with \hat{L}^{-1} and using the fact that L^{-1} is

symplectic we get $\begin{bmatrix} \hat{P} & 0 \\ 0 & \hat{P} \end{bmatrix} - i \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix} \geq 0$ on $\mathcal{H} \oplus \mathcal{H}$. Hence we get $\begin{bmatrix} \hat{P} & -iI \\ iI & \hat{P} \end{bmatrix} \geq 0$ on $\mathcal{H} \oplus \mathcal{H}$. But this means $\hat{P} \geq I$ on \mathcal{H} and correspondingly $\mathcal{P} \geq I$ on \mathcal{H} .

3. Since S is positive and invertible, $\alpha(z, w) := \operatorname{Re} \langle z, Sw \rangle$ defines a complete real inner product on \mathcal{H} . Therefore, by (2.14), a primary quasifree state ϕ as in (3.13) exists if and only if $\sigma(z, w)^2 \leq \alpha(z, z)\alpha(w, w)$, for all $f, g \in \mathcal{H}$. That is

$$\operatorname{Im} \langle z, w \rangle^2 \leq \operatorname{Re} \langle z, Sz \rangle \operatorname{Re} \langle w, Sw \rangle. \quad (3.15)$$

Now

$$\begin{aligned} \operatorname{Im} \langle z, w \rangle^2 &= \operatorname{Im} \langle Lz, Lw \rangle^2 \\ &\leq |\langle Lz, Lw \rangle|^2 \\ &\leq \langle Lz, Lz \rangle \langle Lw, Lw \rangle \\ &\leq \langle Lz, \mathcal{P}Lz \rangle \langle Lw, \mathcal{P}Lw \rangle \\ &= \operatorname{Re} \langle Lz, \mathcal{P}Lz \rangle \operatorname{Re} \langle Lw, \mathcal{P}Lw \rangle \\ &= \langle z, L^\tau \mathcal{P}Lz \rangle \langle w, L^\tau \mathcal{P}Lw \rangle, \end{aligned} \quad (3.16)$$

where (3.16) follows from 2). Thus we proved (3.15). Further, $\phi = \phi_A$ because $\operatorname{Re} \langle \cdot, S(\cdot) \rangle_{\mathcal{H}} = -\operatorname{Im} \langle A(\cdot), \cdot \rangle$. \square

3.8 Lemma. *Let S be a real linear symmetric and invertible operator on \mathcal{H} . Suppose the function $f : \mathcal{H} \rightarrow \mathbb{R}$ defined by $f(z) = e^{-\frac{1}{2} \operatorname{Re} \langle z, Sz \rangle}$ is the quantum characteristic function of a density operator ρ i.e. $S \in \mathcal{K}_{\mathbf{G}}(\mathcal{H})$ then on $(\mathcal{H}, \operatorname{Re} \langle \cdot, \cdot \rangle)$,*

1. $S - I$ is Hilbert-Schmidt.
2. $(\sqrt{S}J\sqrt{S})^\tau(\sqrt{S}J\sqrt{S}) - I$ is in trace class.

Proof. 1. We want to prove that $S - I$ is Hilbert-Schmidt on the real Hilbert space \mathcal{H} . Since $\hat{S} - i\hat{J} \geq 0$, by Lemma 3.7 there exists a primary quasifree state ϕ on $CCR(\mathcal{H}, \sigma)$ such that

$$\phi(W(z)) = e^{-\frac{1}{2} \operatorname{Re} \langle z, Sz \rangle}.$$

Claim : ϕ_A and $\phi_{(-J)}$ are quasi equivalent, where $A = -JS$.

Proof (of Claim). Consider the state ψ on $\mathcal{B}(\Gamma_s(\mathcal{H}))$ given by $X \mapsto \operatorname{Tr} \rho X$. The quasifree state ϕ_A is the restriction of ψ to $\mathcal{A} := CCR(\mathcal{H}, \sigma) \hookrightarrow \mathcal{B}(\Gamma_s(\mathcal{H}))$. Let $(H_\psi, \Pi_\psi, \Omega_\psi)$ be the GNS triple for $\mathcal{B}(\Gamma_s(\mathcal{H}))$ with respect to ψ . Then $(H_\psi, \Pi_\psi|_{\mathcal{A}}, \Omega_\psi)$ is the GNS triple for \mathcal{A} with respect to ϕ_A . To see this, only thing to be noticed is Ω_ψ is cyclic for $\Pi_\psi(\mathcal{A})$, which is clear since \mathcal{A} is strongly dense in $\mathcal{B}(\Gamma_s(\mathcal{H}))$. We further note that the inclusion $\mathcal{A} \subseteq \mathcal{B}(\Gamma_s(\mathcal{H}))$ is the GNS representation with respect to the vacuum state which is the quasi-free state given by ϕ_{-J} . It can be seen that the association

$$W(x) \mapsto \Pi_\psi(W(x))$$

can be extended as an isomorphism between $\mathcal{B}(\Gamma_s(\mathcal{H})) = \mathcal{A}''$ and $\Pi_\psi(\mathcal{B}(\Gamma_s(\mathcal{H})))$. Thus the claim is proved.

Since $\phi_{(-J)}$ and ϕ_A are quasi equivalent, by Theorem 2.3 we get $A + J$ is Hilbert-Schmidt on \mathcal{H}_{-J} which is same as \mathcal{H} with the real inner product $\operatorname{Re} \langle \cdot, \cdot \rangle_{\mathcal{H}}$. Since $A = -JS$, multiplying by J on $A + J$ we get $S - I$ is Hilbert-Schmidt on $(\mathcal{H}, \operatorname{Re} \langle \cdot, S(\cdot) \rangle_{\mathcal{H}})$. Now the result is an easy consequence.

2. This follows due to the same reason as that of 1) because of Theorem 2.3 itself. We get $\sqrt{-A^2 - I}$ is Hilbert-Schmidt on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$. This is same as $-A^2 - I$ is trace class on the same Hilbert space. Hence we have $-JSJS - I$ is trace class. By multiplying with \sqrt{S} on the left and $(\sqrt{S})^{-1}$ on the right we see that $-\sqrt{S}JSJ\sqrt{S} - I$ is trace class. The result follows because $J^\tau = -J$

□

It may be noted at this point that the operator $\sqrt{S}J\sqrt{S}$ in 2) of the above theorem is the skew symmetric operator B appearing in the proof of Williamson's normal form in [BJ19]. Proof of Williamson's normal form was obtained there by applying spectral theorem (as proved in [BJ19]) to B ,

$$\Gamma^T B \Gamma = \begin{bmatrix} 0 & -P \\ P & 0 \end{bmatrix}$$

where Γ is an orthogonal transformation. L was obtained by taking

$$L = \begin{bmatrix} P^{-1/2} & 0 \\ 0 & P^{-1/2} \end{bmatrix} \Gamma^\tau S^{1/2}.$$

This choice of L provides $S = L^\tau \mathcal{P} L$, where $\mathcal{P}_0 = \begin{bmatrix} P & 0 \\ 0 & P \end{bmatrix}$.

3.9 Corollary. *Assume the hypothesis of the previous Lemma 3.8. Then*

1. *If $S - I \geq 0$ then $S - I$ is trace class on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$*
2. *If S is complex linear then $S - I \geq 0$ and $S - I$ is trace class on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$.*

Proof. 1. We have $-\sqrt{S}JSJ\sqrt{S} - I$ is trace class on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$. Hence by multiplying with $(\sqrt{S})^{-1}$ on both sides $(-J)SJ - S^{-1}$ is trace class. Since $S - I \geq 0$, $(-J)SJ - I \geq 0$ and $S^{-1} \leq I$ therefore we have

$$0 \leq (-J)SJ - I \leq (-J)SJ - S^{-1}$$

and we conclude that $(-J)SJ - I$ is trace class on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$. Thus this part is proved by a conjugation with J .

2. By 3) in Lemma 3.7 we have

$$-A^2 - I \geq 0 \tag{3.17}$$

with respect to the real inner product $\text{Re} \langle \cdot, S(\cdot) \rangle$. We have $A^2 = JSJS$ but since S is complex linear it commutes with J , thus $A^2 = -S^2$ and we see that $S^2 - I \geq 0$, consequently $S \geq I$ on $(\mathcal{H}, \text{Re} \langle \cdot, S(\cdot) \rangle_{\mathcal{H}})$. But this implies $S \geq I$ on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$ since S is positive. Since S commutes with J , by 2) of Lemma 3.8 we see that $S^2 - I$ is Hilbert-Schmidt on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$. Now the claim follows because $0 \leq S - I \leq S^2 - I$. □

3.10 Remark. By 2) of Example 3.4 we have seen that for a symplectic automorphism L , $L^\tau L$ is a covariance operator whenever $L^\tau L - I$ is Hilbert-Schmidt. Now by Lemma 3.8 we get that $L^\tau L$ satisfies the conditions 1, and 2 there. This is true also for any such symplectic transformation. But since $\sqrt{L^\tau L}$ is symplectic whenever L is so, the condition 2 is just void. Also it can be proved independently that for any symplectic transformation the positivity condition (3.6) on $L^\tau L$ is true. Therefore, $L^\tau L - I$ is Hilbert-Schmidt is the only non-trivial condition here.

§ 3.2 Sufficiency

Now we proceed to prove that the conditions in Lemma 3.6 and 3.8 are sufficient to ensure that S is covariance operator of a quantum Gaussian state. This essentially involves an explicit construction of the quantum Gaussian state for positive complex linear operators and an application of Williamson normal form. We begin with some computations for this construction.

3.11 Lemma. *If $s_j > 0$ then $\sum_{j=1}^{\infty} \left(\frac{e^{-s_j}}{1-e^{-s_j}} \right) < \infty$ if and only if $\sum_{j=1}^{\infty} e^{-s_j}$ is convergent.*

Proof. Assume $\sum_{j=1}^{\infty} \left(\frac{e^{-s_j}}{1-e^{-s_j}} \right) < \infty$. Since $\frac{e^{-s_j}}{1-e^{-s_j}} > 0$ and $\frac{1}{1-e^{-s_j}} > 1$, we have $0 < \sum_{j=1}^{\infty} e^{-s_j} < \sum_{j=1}^{\infty} \left(\frac{e^{-s_j}}{1-e^{-s_j}} \right) < \infty$. Now assume that $\sum_{j=1}^{\infty} e^{-s_j} < \infty$. Then $s_j \rightarrow \infty$ and hence $\frac{1}{1-e^{-s_j}} \rightarrow 1$. This means we have $0 < \frac{1}{1-e^{-s_j}} < M, \forall j$, for some $M > 1$. Therefore, $\sum_{j=1}^{\infty} \left(\frac{e^{-s_j}}{1-e^{-s_j}} \right) < \infty$. \square

Let $\mathcal{H} = H + iH$ and $\{e_1, e_2, e_3 \dots\}$ be an orthonormal basis for H . Note that $\{e_j\}$ is also a basis for \mathcal{H} as a complex Hilbert space. Let $D = \text{Diag}(d_j)$ be a bounded diagonal operator on \mathcal{H} , with $d_j > 1$, $j = 1, 2, 3, \dots$ in the given basis. Since $d_j > 1$ there exists $s_j > 0$ such that $d_j = \coth\left(\frac{s_j}{2}\right)$ for all j . If we consider D as a real linear operator on \mathcal{H} , then $D_0 = \begin{bmatrix} D & 0 \\ 0 & D \end{bmatrix}$ on $H \oplus H$.

3.12 Lemma. *Let $D = \text{Diag}(d_j)$ be a bounded diagonal operator on \mathcal{H} , with $d_j > 1$, $j = 1, 2, 3, \dots$ with respect to a basis. Write $d_j = \coth\left(\frac{s_j}{2}\right)$ for all j . Then $D - I$ is trace class if and only if $\sum_{j=1}^{\infty} e^{-s_j}$ is convergent.*

Proof. Observe,

$$\begin{aligned}
 D - I \text{ is in trace class} &\Leftrightarrow \sum_{j=1}^{\infty} (d_j - 1) < \infty \\
 &\Leftrightarrow \sum_{j=1}^{\infty} \left(\coth\left(\frac{s_j}{2}\right) - 1 \right) < \infty \\
 &\Leftrightarrow \sum_{j=1}^{\infty} \left(\frac{1 + e^{-s_j}}{1 - e^{-s_j}} - 1 \right) \\
 &\Leftrightarrow \sum_{j=1}^{\infty} \left(\frac{e^{-s_j}}{1 - e^{-s_j}} \right) < \infty \\
 &\Leftrightarrow \sum_{j=1}^{\infty} e^{-s_j} < \infty
 \end{aligned} \tag{3.18}$$

where (3.18) follows from Lemma 3.11. \square

3.13 Proposition. *Let D be as described in Lemma 3.12 and $D - I$ is trace class. Then there exists a state ρ_D on $\Gamma_s(\mathcal{H})$ such that $\hat{\rho}_D(x) = e^{-\frac{1}{2}\langle x, Dx \rangle}$.*

Proof. Consider the diagonal operator $T = \text{Diag}(e^{-s_j})$ with respect to the same basis in which D is diagonal then the second quantization $\Gamma(T)$ is a trace class operator on the symmetric Fock space, $\Gamma_s(\mathcal{H})$ because of the following reasoning. T is positive and by Lemma 3.12 it is a trace class operator. Thus we have $s_j > 0$ and $s_j \rightarrow \infty$. Therefore we get $\sup_j(e^{-s_j}) < 1$. Now by Exercise 20.22 (iv) in [Par12], the second quantization (2.7) $\Gamma(T)$ is trace class with

$$\text{Tr } \Gamma(T) = \prod_{j=1}^{\infty} (1 - e^{-s_j})^{-1} \quad (3.19)$$

Define $\rho_D = \prod_{j=1}^{\infty} (1 - e^{-s_j}) \Gamma(T)$, then ρ is a density operator on $\Gamma_s(\mathcal{H})$. We have $\mathcal{H} = \oplus_j \mathbb{C} e_j$. Since $\Gamma(e^{-s_j}) = e^{-s_j a_j^\dagger a_j}$ on $\Gamma_s(\mathbb{C} e_j)$, $\rho_D = \prod_{j=1}^{\infty} (1 - e^{-s_j}) \Gamma(\oplus_j e^{-s_j}) = \otimes_j \rho_j$, where $\rho_j = (1 - e^{-s_j}) e^{-s_j a_j^\dagger a_j}$. Let $x = \oplus_j x_j e_j$ then

$$\begin{aligned} \hat{\rho}_D(x) &= \text{Tr } \rho W(x) \\ &= \text{Tr } \left(\prod_{j=1}^{\infty} (1 - e^{-s_j}) \Gamma(\oplus_j e^{-s_j}) W(\oplus_j x_j) \right) \\ &= \text{Tr } \left(\otimes_j (1 - e^{-s_j}) \Gamma(e^{-s_j}) W(x_j) \right) \\ &= \text{Tr } \left(\otimes_j (1 - e^{-s_j}) e^{-s_j a_j^\dagger a_j} W(x_j) \right) \\ &= \prod_{j=1}^{\infty} \text{Tr } \left((1 - e^{-s_j}) e^{-s_j a_j^\dagger a_j} W(x_j) \right) \\ &= \prod_{j=1}^{\infty} e^{-\langle x_j, \frac{1}{2} \coth\left(\frac{s_j}{2}\right) x_j \rangle} \\ &= e^{-\frac{1}{2} \langle x, D x \rangle} \end{aligned} \quad (3.20)$$

where (3.20) follows from Example 3 of Gaussian states. \square

Recall from Example 1 that the vacuum state $|e(0)\rangle\langle e(0)|$ on $\Gamma_s(\mathcal{H})$ is a Gaussian state with covariance operator I .

3.14 Proposition. *If \mathcal{P} is any complex linear operator on \mathcal{H} such that $\mathcal{P} - I$ is positive and trace class then there exists a state ρ on $\Gamma_s(\mathcal{H})$ such that the quantum characteristic function $\hat{\rho}$ associated with ρ is given by*

$$\hat{\rho}(x) = e^{-\frac{1}{2} \langle x, \mathcal{P} x \rangle}$$

for every $x \in \mathcal{H}$.

Proof. Let U be a unitary operator such that $\mathcal{P} = U^* D U$. Such a U exists by applying spectral theorem to the compact positive operator $\mathcal{P} - I$. Since $\mathcal{P} \geq I$ assume without loss of generality that $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$ is such that $D = \begin{bmatrix} D_1 & 0 \\ 0 & I \end{bmatrix}$, where we separated all the diagonal entries of D which are equal to one and not equal to one. Then D_1 satisfies the assumptions in Proposition 3.13 and ρ_{D_1} exists as a Gaussian state on $\Gamma_s(\mathcal{H}_1)$. Let ρ_0 denote the vacuum state $|e(0)\rangle\langle e(0)|$ on $\Gamma_s(\mathcal{H}_2)$, which is Gaussian by Example 1. Then $\rho_{D_1} \otimes \rho_0 = \rho_g(0, D)$. Define $\rho = \Gamma_s(U^*) \rho_{D_1} \otimes \rho_0 \Gamma_s(U)$ and the result follows from Proposition 3.2. \square

For a proof of the following Lemma see the discussion in the beginning of Section 2 in [BS05].

3.15 Lemma. *Let $C - I$ is Hilbert-Schmidt (trace class) then*

1. If $C \geq 0$ then $\sqrt{C} - I$ is Hilbert-Schmidt (trace class).
2. If C is invertible then $C^{-1} - I$ is Hilbert-Schmidt (trace class).

3.16 Lemma. *Let S be a real linear, positive and invertible operator on \mathcal{H} . Then L and \mathcal{P} as in Corollary 2.10 can be chosen such that*

1. *If $S - I$ is Hilbert-Schmidt then $L^\tau L - I$ is Hilbert Schmidt, i.e $L \in \mathcal{S}(\mathcal{H})$.*
2. *If $(\sqrt{S}J\sqrt{S})^\tau(\sqrt{S}J\sqrt{S}) - I$ is trace class then $\mathcal{P} - I$ is a trace class operator on \mathcal{H} .*

Proof. 1. It can be seen from the proof of Williamson's normal form in [BJ19] that M_0 in Theorem 2.9 can be chosen as $M = \mathcal{P}^{-1/2}\Gamma^\tau S^{1/2}$, where $\Gamma_0: K \oplus K \rightarrow H \oplus H$ is an orthogonal transformation such that the skew symmetric operator

$$B_0 := S_0^{1/2} J_0 S_0^{1/2} = \Gamma_0 \begin{bmatrix} 0 & -P \\ P & 0 \end{bmatrix} \Gamma_0^\tau. \quad (3.21)$$

and $\mathcal{P}_0 = \begin{bmatrix} P & 0 \\ 0 & P \end{bmatrix}$ (recall the notation we employed at the beginning of § 2.3 to distinguish between the two avatar's of a real linear operator defined on a complexified Hilbert space). Since $L = UM$ in the proof of Corollary 2.10,

$$L^\tau L = S^{1/2} \Gamma \mathcal{P}^{-1} \Gamma^\tau S^{1/2}. \quad (3.22)$$

But

$$\begin{bmatrix} 0 & P \\ -P & 0 \end{bmatrix} \begin{bmatrix} 0 & -P \\ P & 0 \end{bmatrix} = \begin{bmatrix} P^2 & 0 \\ 0 & P^2 \end{bmatrix}, \quad (3.23)$$

therefore if we write $\mathcal{P}_0 = \begin{bmatrix} 0 & -P \\ P & 0 \end{bmatrix}$, and recall the relationship between the operator \mathcal{P}_0 on the real Hilbert space and \mathcal{P} on the complexified Hilbert space (described at the beginning of § 2.3), we see that

$$\mathcal{P}^{-1} = (\sqrt{\mathcal{P}^\tau \mathcal{P}})^{-1} \quad (3.24)$$

Since Γ is orthogonal, by (3.21) and (3.24) we get $(\sqrt{B^\tau B})^{-1} = \Gamma \mathcal{P}^{-1} \Gamma^\tau$. Now by (3.22) we get

$$L^\tau L = S^{1/2} (\sqrt{B^\tau B})^{-1} S^{1/2}. \quad (3.25)$$

We have $S - I$ is Hilbert-Schmidt. Therefore, so is $J^\tau S J - I$. Hence $S^{1/2} J^\tau S J S^{1/2} - S$ is Hilbert-Schmidt. By adding and subtracting I and using the fact the $S - I$ is Hilbert-Schmidt we get $S^{1/2} J^\tau S J S^{1/2} - I$ is also so. In other words, we just got $B^\tau B - I$ is Hilbert-Schmidt. Now by Lemma 3.15 we get $(\sqrt{B^\tau B})^{-1} - I$ is Hilbert-Schmidt. This along with (3.25) finally allows us to conclude that $L^\tau L - I$ is Hilbert-Schmidt.

2. By keeping the notations above and using Lemma 3.15, we have $(\sqrt{B^\tau B})^{-1} - I$ is trace class and thus $S^{1/2} (\sqrt{B^\tau B})^{-1} S^{1/2} - S = L^\tau L - S$ is trace class. Since $S = L^\tau \mathcal{P} L$ we get $L^\tau (\mathcal{P} - I) L$ is trace class. Since L is invertible we see that $\mathcal{P} - I$ is trace class. \square

Here is the theorem which characterizes all covariance operators of quantum Gaussian states. It is essentially the statement that there exists a quantum Gaussian state ρ with covariance matrix S on the Boson Fock space of \mathcal{H} if and only if $\hat{\rho}|_{CCR(\mathcal{H}, \sigma)}$ is a primary quasifree state ϕ_A quasiequivalent to the vacuum state ϕ_{-J} on $CCR(\mathcal{H}, \sigma)$, where $A = -JS$.

3.17 Theorem. *Let S be a real linear, bounded, symmetric and invertible operator on \mathcal{H} . Then S is the covariance operator of a quantum Gaussian state if and only if the following holds*

1. $\hat{S} - i\hat{J} \geq 0$ on \mathcal{H} .
2. $S - I$ is Hilbert-Schmidt on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$.
3. $(\sqrt{S}J\sqrt{S})^\tau(\sqrt{S}J\sqrt{S}) - I$ is trace class on $(\mathcal{H}, \text{Re} \langle \cdot, \cdot \rangle)$.

Proof. In the last Section in Lemma 3.6 and 3.8 we have seen the necessity of conditions 1-3 for a quantum Gaussian state. Now we prove sufficiency. So assume 1-3. As $\hat{S} - i\hat{J} \geq 0$, $S \geq 0$ and by assumption S is invertible and hence it has Williamson's normal form. Thus there exists a symplectic automorphism L on \mathcal{H} such that $S = L^\tau \mathcal{P} L$ (Corollary 2.10). By Lemma 3.16, $\mathcal{P} - I$ is trace class and $L^\tau L - I$ is Hilbert-Schmidt. Now by Theorem 2.7 there exists a unitary operator $\Gamma_s(L)$ on $\Gamma_s(\mathcal{H})$ such that

$$\Gamma_s(L)W(u)\Gamma_s(L)^* = W(Lu) \quad (3.26)$$

Since $\mathcal{P} - I$ is trace class and positive, by Proposition 3.14 there exists a density operator $\rho_{\mathcal{P}}$ such that $\hat{\rho}_{\mathcal{P}}(y) = e^{-\frac{1}{2}\langle y, \mathcal{P}y \rangle}$ for every $y \in \mathcal{H}$. Define

$$\rho = \Gamma_s(L)^* \rho_{\mathcal{P}} \Gamma_s(L) \quad (3.27)$$

Claim. $\hat{\rho}(z) = e^{-\frac{1}{2}\text{Re}\langle z, Sz \rangle}$ for every $z \in \mathcal{H}$.

Proof (of Claim). By Proposition 3.2 we have

$$\begin{aligned} \hat{\rho}(z) &= \hat{\rho}_{\mathcal{P}}(Lz) \\ &= e^{-\frac{1}{2}\langle Lz, \mathcal{P}Lz \rangle} \\ &= e^{-\frac{1}{2}\text{Re}\langle Lz, \mathcal{P}Lz \rangle} \\ &= e^{-\frac{1}{2}\text{Re}\langle z, L^\tau \mathcal{P}Lz \rangle} \\ &= e^{-\frac{1}{2}\text{Re}\langle z, Sz \rangle}. \end{aligned}$$

□

It easily seen that if S is complex linear then the third condition of previous theorem is redundant. So we have the following.

3.18 Corollary. *Let S be a complex linear, self-adjoint and invertible operator on \mathcal{H} . Then S is the covariance operator of a quantum Gaussian state on $\Gamma_s(\mathcal{H})$ if and only if $\hat{S} - i\hat{J} \geq 0$ and $S - I$ is trace class.*

3.19 Corollary. *Let $S \geq I$ be real linear then S is the covariance operator of a quantum Gaussian state on $\Gamma_s(\mathcal{H})$ if and only if $S - I$ is trace class*

3.20 Remark. Notice that even in the finite mode case the condition $\hat{S} - i\hat{J} \geq 0$ does not necessarily imply that $S \geq I$. For example, one can consider the 2×2 matrix $\begin{bmatrix} 2 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}$, which is a valid covariance matrix but not greater than I .

4 Convexity Properties of Covariance Operators

The previous section described and characterized infinite mode Gaussian states. The Hilbert Schmidt conditions appearing there show that unlike finite dimensions covariance operators in infinite dimensions do not form a cone. However, the next proposition shows that they do form a convex set. This allows us to extend some beautiful symmetry properties of Gaussian states proved by Parthasarathy [Par10, Par13] in the finite mode case to this setting. In the following $\mathcal{K}_{\mathbf{G}}(\mathcal{H})$ denotes the collection of covariance operators for Gaussian states on $\Gamma_s(\mathcal{H})$ and $\mathcal{S}(\mathcal{H})$ are Shale operators on \mathcal{H} for a fixed infinite dimensional separable real Hilbert space \mathcal{H} . The following Proposition is an easy consequence of earlier discussions.

4.1 Proposition. *Consider two mean zero Gaussian states*

$$\rho_i = \rho_g(0, S_i), i = 1, 2$$

on $\Gamma_s(\mathcal{H})$. For $\theta \in \mathbb{R}$, let U_θ be the unitary operator $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ on $\mathcal{H} \oplus \mathcal{H}$. Then

$$\text{Tr}_2 (\Gamma_s(U_\theta)(\rho_1 \otimes \rho_2)\Gamma_s(U_\theta)^*) = \rho_g(0, (\cos^2 \theta)S_1 + (\sin^2 \theta)S_2)$$

where Tr_2 denotes the relative trace over the second factor of $\Gamma_s(\mathcal{H}) \otimes \Gamma_s(\mathcal{H})$. Consequently $\mathcal{K}_{\mathbf{G}}(\mathcal{H})$ is a convex set.

4.2 Lemma. *Let $P \geq I$, then there exists invertible positive operators P_1 and P_2 such that*

$$P = \frac{1}{2}(P_1 + P_2) = \frac{1}{2}(P_1^{-1} + P_2^{-1}) \quad (4.1)$$

Proof. Take $P_1 = P + \sqrt{P^2 - I}$ and $P_2 = P - \sqrt{P^2 - I}$. Then $P_1 P_2 = P_2 P_1 = I$ and (4.1) is satisfied. \square

4.3 Lemma. *Let $\mathcal{H} = H + iH$ and $\mathcal{P} \in \mathcal{B}(\mathcal{H})$ be such that $\mathcal{P} - I$ is positive and trace class, further let $\mathcal{P}_0 = \begin{bmatrix} P & 0 \\ 0 & P \end{bmatrix}$ on $H \oplus H$ (§ 2.3). Then $\mathcal{P} = \frac{1}{2}(\mathcal{P}_1 + \mathcal{P}_2)$, for some $\mathcal{P}_j \geq 0$, and $\mathcal{P}_j^{\frac{1}{2}} \in \mathcal{S}(\mathcal{H})$, $j = 1, 2$.*

Proof. Take $P_1 = P + \sqrt{P^2 - I}$ and $P_2 = P - \sqrt{P^2 - I}$, then by (4.1)

$$\mathcal{P}_0 = \frac{1}{2} \left\{ \begin{bmatrix} P_1 & 0 \\ 0 & P_1^{-1} \end{bmatrix} + \begin{bmatrix} P_2 & 0 \\ 0 & P_2^{-1} \end{bmatrix} \right\}$$

Define \mathcal{P}_j such that $\mathcal{P}_j(x + iy) = P_j x + P_j^{-1} y, \forall x, y \in H, j = 1, 2$. Then \mathcal{P}_j is symplectic and positive. To prove $\mathcal{P}_j^{\frac{1}{2}} \in \mathcal{S}(\mathcal{H})$ we should prove $\mathcal{P}_j - I$ is Hilbert-Schmidt. To this end, it is enough to show that $P_j - I$ is Hilbert-Schmidt, $j = 1, 2$. Since $P - I$ is trace class (and hence Hilbert-Schmidt) it is enough to show $\sqrt{P^2 - I}$ is Hilbert-Schmidt or equivalently $P^2 - I$ is trace class. This is true because $P^2 - I = (P - I)^2 + 2(P - I)$. \square

4.4 Theorem. *$S \in \mathcal{K}_{\mathbf{G}}(\mathcal{H})$ if and only if*

$$S = \frac{1}{2}(N^\tau N + M^\tau M) \quad (4.2)$$

for some $N, M \in \mathcal{S}(\mathcal{H})$. Further, S is an extreme point of $\mathcal{K}_{\mathbf{G}}(\mathcal{H})$ if and only if $S = N^\tau N$ for some $N \in \mathcal{S}(\mathcal{H})$.

Proof. Note that if $N \in \mathcal{S}(\mathcal{H})$ then $N^\tau N$ is a covariance operator (see 2 of Example 3.4). Now let $S \in \mathcal{K}_{\mathbf{G}}(\mathcal{H})$, let $S = L^\tau \mathcal{P} L$ be the Williamson's normal form as in Corollary 2.10. Then by Lemma 3.16 and Lemma 3.7 we have $L \in \mathcal{S}(\mathcal{H})$ and $\mathcal{P} - I$ is trace class and positive. By Corollary 2.10 we have $\mathcal{P}_0 = \begin{bmatrix} P & 0 \\ 0 & P \end{bmatrix}$. By Lemma 4.3, $\mathcal{P} = \frac{1}{2}(\mathcal{P}_1 + \mathcal{P}_2)$ with $\mathcal{P}_j \geq 0, j = 1, 2$. Therefore we have

$$S = \frac{1}{2} L^\tau (\mathcal{P}_1 + \mathcal{P}_2) L.$$

By taking $N = \mathcal{P}_1^{1/2} L$ and $M = \mathcal{P}_2^{1/2} L$ we get (4.2). An easy computation shows $N, M \in \mathcal{S}(\mathcal{H})$.

Proof of second part of the Theorem goes in an exactly similar lines to the proof of the similar statement in the finite mode case, Theorem 3 in [Par13]. We give it here for completeness. The first part also shows that for an element S of $\mathcal{K}_{\mathbf{G}}(\mathcal{H})$ to be extremal it is necessary that $S = L^\tau L$ for some $L \in \mathcal{S}(\mathcal{H})$. To prove sufficiency, suppose there exist $L \in \mathcal{S}(\mathcal{H})$ and $S_1, S_2 \in \mathcal{K}_{\mathbf{G}}(\mathcal{H})$ such that

$$L^\tau L = \frac{1}{2}(S_1 + S_2).$$

By the first part of the theorem there exist $L_j \in \mathcal{S}(\mathcal{H})$ such that

$$L^\tau L = \frac{1}{4} \sum_{j=1}^4 L_j^\tau L_j \tag{4.3}$$

where $S_1 = \frac{1}{2}(L_1^\tau L_1 + L_2^\tau L_2)$, $S_2 = \frac{1}{2}(L_3^\tau L_3 + L_4^\tau L_4)$. Left multiplication by $(L^\tau)^{-1}$ and right multiplication by L^{-1} on both sides of (4.3) gives

$$I = \frac{1}{4} \sum_{j=1}^4 M_j \tag{4.4}$$

where $M_j = (L^\tau)^{-1} L_j^\tau L_j L^{-1}$. Each $M_j \in \mathcal{S}(\mathcal{H})$ and is positive. Multiplying by J on both sides of (4.4) we get

$$J = \frac{1}{4} \sum_{j=1}^4 M_j J = \frac{1}{4} \sum_{j=1}^4 M_j J M_j M_j^{-1} = \frac{1}{4} J \sum_{j=1}^4 M_j^{-1}.$$

Thus

$$I = \frac{1}{4} \sum_{j=1}^4 M_j = \frac{1}{4} \sum_{j=1}^4 M_j^{-1} = \frac{1}{4} \sum_{j=1}^4 \frac{1}{2} (M_j + M_j^{-1}),$$

which implies

$$\sum_{j=1}^4 (M_j^{1/2} - M_j^{-1/2})^2 = 0,$$

or $M_j = I, \forall 1 \leq j \leq 4$. Thus $L_j^\tau L_j = L^\tau L, \forall j$ and $S_1 = S_2$. \square

4.5 Corollary. Let S_1, S_2 be extreme points of $\mathcal{K}_{\mathbf{G}}(\mathcal{H})$ such that $S_1 \geq S_2$. Then $S_1 = S_2$.

Proof. By Theorem 4.4, let $S_1 = L_1^\tau L_1$ and $S_2 = L_2^\tau L_2$ for some $L_1, L_2 \in \mathcal{S}(\mathcal{H})$. $L_1^\tau L_1 \geq L_2^\tau L_2$ implies that the symplectic transformation $M := L_2 L_1^{-1}$ has the property $M^\tau M \leq I$. But since $M^\tau M$ is a positive symplectic automorphism $M^\tau M = VTV^*$ for some unitary V , where $\mathcal{H} = H + iH$ and $T(x + iy) = Ax + iA^{-1}y$ for some positive invertible operator A on H . This can be seen by applying Proposition 2.6 to M . But such a $T \leq I$ if and only if $A = I$. This proves $M^\tau M = I$. But this implies $L_2^\tau L_2 = L_1^\tau L_1$ from the definition of M . \square

5 Structure of Quantum Gaussian States

If S is a Gaussian covariance matrix it satisfies the properties listed in Theorem 3.17, then by combining Lemma 3.7 and Lemma 3.16 we get a Williamson's normal form (Corollary 2.10), $S = L^\tau \mathcal{P} L$ such that $\mathcal{P} - I$ is positive and trace class. By spectral theorem, there exists a unitary U such that $\mathcal{P} = U^* D U$, where D is diagonal and positive. Since a unitary is also symplectic, whenever S is a covariance operator we may assume without loss of generality that the \mathcal{P} occurring in the Williamson's normal form is of the form $\mathcal{P} = \begin{bmatrix} D & 0 \\ 0 & I \end{bmatrix}$ on a decomposition $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$, with $D = \text{Diag}(d_1, d_2, \dots)$, $d_1 \geq d_2 \geq \dots > 1$. But now we have fixed a basis of \mathcal{H} , therefore through the identification of $\Gamma_s(\mathbb{C})$ with $L^2(\mathbb{R})$ (Lebesgue measure), $(e(z) \in \Gamma_s(\mathbb{C}))$ is identified with the L^2 -function $x \mapsto (2\pi)^{-1/4} \exp\{-4^{-1}x^2 + zx - 2^{-1}z^2\}$ (refer Example 2.1), we can assume without loss of generality that $\Gamma_s(\mathcal{H}) = \otimes_j L^2(\mathbb{R})$, with respect to the stabilising vector $e(0)$.

5.1 Theorem. *Let $\rho_g(w, S)$ be a Gaussian state in $\Gamma_s(\mathcal{H})$. Let $S = L^\tau \mathcal{P} L$ be a Williamson's normal form of S , with $L^\tau L - I$ is Hilbert-Schmidt and $\mathcal{P} = \begin{bmatrix} D & 0 \\ 0 & I \end{bmatrix}$, on a decomposition $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$, with $D = \text{Diag}(d_1, d_2, \dots)$, $d_1 \geq d_2 \geq \dots > 1$, $d_j = \coth(\frac{s_j}{2})$, $\forall j$. Then*

$$\rho_g(w, S) = W\left(\frac{-i}{2}w\right)^* \Gamma_s(L)^* [\otimes_j (1 - e^{-s_j}) e^{-s_j a_j^\dagger a_j} \otimes \rho_0] \Gamma_s(L) W\left(\frac{-i}{2}w\right). \quad (5.1)$$

where $\rho_0 = |e(0)\rangle\langle e(0)|$ is the vacuum state on $\Gamma_s(\mathcal{H}_2)$.

Proof. By Proposition 3.5, $\rho_g(w, S) = W(\frac{-i}{2}w)^{-1} \rho_g(0, S) W(\frac{-i}{2}w)$. Since $S = L^\tau \mathcal{P} L$, $\rho_g(0, S) = \Gamma_s(L)^* \rho_g(0, \mathcal{P}) \Gamma_s(L)$. Since $\mathcal{P} = D \oplus I$, $\rho_g(0, \mathcal{P}) = \rho_g(0, D) \otimes \rho_g(0, I)$. But $\rho_g(0, D) = \otimes_j (1 - e^{-s_j}) e^{-s_j a_j^\dagger a_j}$. Since both on left and right hand sides have same quantum characteristic function by proof of Proposition 3.13 and it is obvious that $\rho_g(0, I) = \rho_0$. \square

5.2 Corollary. *If $\{e_j\}$ is an orthonormal basis of \mathcal{H} , consider $\Gamma_s(\mathcal{H}) = \Gamma(\oplus_j \mathbb{C} e_j) = \otimes_j L^2(\mathbb{R})$. Then the wave function of a general pure quantum Gaussian state is of the form*

$$|\psi\rangle = W(\alpha) \Gamma_s(U) (\otimes_j |e_{\lambda_j}\rangle) \quad (5.2)$$

where $e_\lambda \in L^2(\mathbb{R})$ is defined for $\lambda > 0$ by

$$e_\lambda(x) = (2\pi)^{-1/4} \lambda^{-1/2} \exp\{-4^{-1} \lambda^{-2} x^2\}, \quad x \in \mathbb{R};$$

$\alpha \in \mathcal{H}$, U is a unitary operator on \mathcal{H} , $\Gamma_s(U)$ is the second quantization unitary operator associated with U and $\lambda_j, j \in \mathbb{N}$ are positive scalars.

Proof. The proof is essentially similar to the proof of Corollary 2 in [Par13] because of Theorem 5.1 and Proposition 2.6. \square

5.3 Theorem (Purification). *Let ρ be a mixed Gaussian state in $\Gamma_s(\mathcal{H})$. Then there exists a pure Gaussian state $|\psi\rangle$ in $\Gamma_s(\mathcal{H}) \otimes \Gamma_s(\mathcal{H})$ such that*

$$\rho = \text{Tr}_2 U |\psi\rangle\langle\psi| U^*$$

where U is a unitary and Tr_2 is the relative trace over the second factor.

Proof. Proof is same as that of Theorem 5 in [Par13]. \square

6 Symmetry group of Gaussian states

Let \mathcal{H} be a complex separable infinite dimensional Hilbert space and let $\mathbf{G}(\mathcal{H})$ denote the set of all Gaussian states on $\Gamma_s(\mathcal{H})$. In this Section, we characterize all automorphisms of $\mathcal{B}(\Gamma_s(\mathcal{H}))$ preserving the set of Gaussian states.

6.1 Definition. A unitary operator U on $\Gamma_s(\mathcal{H})$ is called a *Gaussian symmetry* if $U\rho U^* \in \mathbf{G}(\mathcal{H})$ for every $\rho \in \mathbf{G}(\mathcal{H})$.

We use \mathbb{Z}_+ to denote the set $\{0, 1, 2, 3, \dots\}$ and take $\mathbb{Z}_{+,0}^\infty := \{(k_1, k_2, \dots, k_n, 0, 0, \dots)^\tau | k_j \in \mathbb{Z}_+, j, n \in \mathbb{N}\}$. Let $\{e_j\}_{j \in \mathbb{N}}$ denote the standard orthonormal basis for $\ell^2(\mathbb{N})$, the column vector with 1 at the j^{th} position and zero elsewhere. An infinite order matrix A is said to be a permutation matrix if A corresponds to a unitary operator which maps $\{e_j\}$ to itself.

6.2 Lemma. *Let $\{s_j\}_{j \in \mathbb{N}}$ and $\{t_j\}_{j \in \mathbb{N}}$ be two sequences consisting of positive real numbers such that their positive integer linear combinations form same sets, that is,*

$$\left\{ \sum_{j=1}^n s_j k_j | k_j \in \mathbb{Z}_+ \forall j, n \in \mathbb{N} \right\} = \left\{ \sum_{j=1}^n t_j k_j | k_j \in \mathbb{Z}_+ \forall j, n \in \mathbb{N} \right\}. \quad (6.1)$$

If $\{s_j\}_{j \in \mathbb{N}}$ and $\{t_j\}_{j \in \mathbb{N}}$ are both linearly independent over the field \mathbb{Q} , then there exists a bijection $\sigma : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$ such that $s_j = t_{\sigma(j)}$ for all $j \in \mathbb{Z}_+$.

Proof. Consider $s = (s_1, s_2, s_3, \dots)^\tau$ and $t = (t_1, t_2, t_3, \dots)^\tau$ as vectors in \mathbb{R}^∞ . The condition (6.1) in particular means that each t_i is a positive integer linear combination of $\{s_j\}_{j \in \mathbb{N}}$. Hence there exists an infinite matrix A with entries from \mathbb{Z}_+ such that

$$t_i = \sum_{j=1}^{\infty} a_{ij} s_j \quad \forall i.$$

Similarly there exists another infinite matrix B with entries from \mathbb{Z}_+ such that

$$s_i = \sum_{j=1}^{\infty} b_{ij} t_j \quad \forall i.$$

Note that by construction every row of A, B have only finitely many non-zero entries. From this, it is easily verified that formal matrix multiplications AB and BA make

sense and give us matrices with entries from \mathbb{Z}_+ of same kind, namely every row has only finitely many non-zero entries. We also get

$$BA s = s, \quad AB t = t.$$

Then, by the assumption of linear independence on rationals of $\{s_j\}_{j \in \mathbb{N}}$ and $\{t_j\}_{j \in \mathbb{N}}$, we get $AB = BA = I$. In other words A and B are permutation matrices and inverses of each other. Clearly this completes the proof. \square

Now we develop some notation which would help us in our computations. Recall the action of annihilation and creation operators a, a^\dagger on $\Gamma_s(\mathbb{C})$, from Exercise 20.18(b) in [Par12]: $\Gamma_s(\mathbb{C})$ has a complete orthonormal basis $\{|k\rangle : k \in \mathbb{Z}_+\}$, with $|0\rangle$ as the vacuum vector, $a(|k\rangle) = \sqrt{k} |k-1\rangle$, $a(|0\rangle) = 0$, $a^\dagger(|k\rangle) = \sqrt{k+1} |k+1\rangle$. Further each $|k\rangle$ is an eigenvector for the number operator $a^\dagger a$ with eigenvalue k , that is, $a^\dagger a(|k\rangle) = k |k\rangle$.

Now let \mathcal{H} be an infinite dimensional Hilbert space with orthonormal basis, $\{e_j\}_{j \in \mathbb{N}}$. Then we may identify $\Gamma_s(\mathcal{H})$ with $\otimes_{j=1}^\infty \Gamma_s(\mathbb{C}e_j)$, where the sequence of vacuum vectors is chosen as the stabilizing sequence. This identification can be done with respect to any orthonormal basis of \mathcal{H} . Define

$$|\mathbf{k}\rangle = |k_1\rangle \otimes |k_2\rangle \otimes \cdots \otimes |k_N\rangle \otimes |0\rangle \otimes |0\rangle \otimes \cdots =: |k_1\rangle |k_2\rangle \cdots |k_N\rangle \quad (6.2)$$

for $\mathbf{k} = (k_1, k_2, k_3, \dots)^\tau \in \mathbb{Z}_{+,0}^\infty$. It can be seen that $\{|\mathbf{k}\rangle | \mathbf{k} \in \mathbb{Z}_{+,0}^\infty\}$ forms an orthonormal basis for $\Gamma_s(\mathcal{H})$. For $u \in \mathcal{H}$, taking $u_j = \langle e_j, u \rangle$,

$$\langle \mathbf{k} | e(u) \rangle = \prod_{j=1}^n \frac{u_j^{k_j}}{\sqrt{k_j!}} := \frac{u^{\mathbf{k}}}{\sqrt{\mathbf{k}!}},$$

where the last equality defines the multi-index notation and also we take $0^0 = 1$. Therefore we write,

$$e(u) = \sum_{\mathbf{k} \in \mathbb{Z}_{+,0}^\infty} \frac{u^{\mathbf{k}}}{\sqrt{\mathbf{k}!}} |\mathbf{k}\rangle. \quad (6.3)$$

We also have,

$$(I \otimes I \otimes \cdots \otimes I \otimes a_j^\dagger a_j \otimes I \otimes I \otimes \cdots)(|\mathbf{k}\rangle) = \begin{cases} k_j |\mathbf{k}\rangle, & \text{if } j \leq N \\ 0, & \text{otherwise} \end{cases}, \quad (6.4)$$

where $a_j^\dagger a_j$ is the number operator on $\Gamma_s(\mathbb{C}e_j)$, $j \in \mathbb{N}$.

Consider $\mathcal{H} = \oplus_{j=1}^\infty \mathbb{C}e_j$ as above. For a sequence of positive numbers $\{s_j\}_{j \in \mathbb{N}}$ such that $d_j = \coth(\frac{s_j}{2}) > 1$ and $\sum_j (d_j - 1)$ is finite, we know from Theorem 5.1

that, there exists a Gaussian state $\rho_s = \prod_{j=1}^\infty (1 - e^{-s_j}) \otimes_{j=1}^\infty e^{-s_j a_j^\dagger a_j} \in \mathcal{B}(\Gamma_s(\mathcal{H}))$. Then we have the

6.3 Lemma. *The spectrum of the Gaussian state ρ_s is,*

$$\sigma(\rho_s) = \left\{ p e^{-\sum_{j=1}^N s_j k_j} \mid k_j \in \mathbb{Z}_+, N \in \mathbb{N} \right\} \cup \{0\}, \quad (6.5)$$

where $p := \prod_{j=1}^{\infty} (1 - e^{-s_j})$. The point spectrum $\sigma_p(\rho_s) = \sigma(\rho_s) \setminus \{0\}$. Further, if $\{s_j\}_{j \in \mathbb{N}}$ is a sequence of (distinct) irrational numbers which are linearly independent over the field \mathbb{Q} then each number $pe^{-\sum_{j=1}^N s_j k_j}$ is an eigenvalue with multiplicity one.

Proof. Since ρ_s is a compact operator $\sigma(\rho_s) = \sigma_p(\rho_s) \cup \{0\}$. We will show that $\sigma_p(\rho_s) = \{pe^{-\sum_{j=1}^N s_j k_j} | k_j \in \mathbb{Z}_+, N \in \mathbb{N}\}$. We have $\otimes_{j=1}^{\infty} e^{-s_j a_j^\dagger a_j} := \text{s-lim}_{N \rightarrow \infty} \otimes_{j=1}^N e^{-s_j a_j^\dagger a_j} \otimes I \otimes I \otimes \dots$. Therefore, $\otimes_{j=1}^{\infty} e^{-s_j a_j^\dagger a_j} (e(u) \otimes e(0) \otimes e(0) \otimes \dots) = \left(\otimes_{j=1}^N e^{-s_j a_j^\dagger a_j} e(u) \right) \otimes e(0) \otimes e(0) \otimes \dots, \forall u \in \mathbb{C}^N$. Thus $\Gamma_s(\mathbb{C}^N)$ is a reducing subspace for ρ_s and $\rho_s|_{\Gamma_s(\mathbb{C}^N)} = \otimes_{j=1}^N e^{-s_j a_j^\dagger a_j}, \forall N$. Therefore,

$$\rho_s(|\mathbf{k}\rangle) = (p \otimes_{j=1}^{\infty} e^{-s_j a_j^\dagger a_j}) |\mathbf{k}\rangle = pe^{-\sum_{j=1}^{\infty} s_j k_j} |\mathbf{k}\rangle, \forall \mathbf{k} \in \mathbb{Z}_{+,0}^{\infty}. \quad (6.6)$$

Since $\{|\mathbf{k}\rangle | \mathbf{k} \in \mathbb{Z}_{+,0}^{\infty}\}$ forms a complete orthonormal basis for $\Gamma_s(\mathcal{H})$, $\{pe^{-\sum_{j=1}^N s_j k_j} | k_j \in \mathbb{Z}_+, N \in \mathbb{N}\}$ is the complete set of eigen values for ρ_s . If $\{s_j\}$ is linearly independent over \mathbb{Q} , then we see that the eigenvalues corresponding to $|\mathbf{k}_1\rangle \neq |\mathbf{k}_2\rangle$ are not same. Thus the multiplicity of each of these eigenvalues is one. \square

6.4 Lemma. Let ρ_s be as in Lemma 6.3 where $\{s_j\}_{j \in \mathbb{N}}$ is a sequence of (distinct) irrational numbers which are linearly independent over the field \mathbb{Q} . Then a unitary operator U in $\Gamma_s(\mathcal{H})$ is such that $U\rho_s U^*$ is a Gaussian state if and only if for some $\alpha \in \mathcal{H}$, $L \in \mathcal{S}(\mathcal{H})$ and a complex valued function β of modulus one on $\mathbb{Z}_{+,0}^{\infty}$

$$U = W(\alpha) \Gamma_s(L) \beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots),$$

where $\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots)$ is the unique unitary which satisfies

$$\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) |\mathbf{k}\rangle = \beta(\mathbf{k}) |\mathbf{k}\rangle, \forall \mathbf{k} \in \mathbb{Z}_{+,0}^{\infty}.$$

Proof. Since $\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots)$ commutes with ρ_s the sufficiency is immediate. To prove the necessity, suppose $U\rho_s U^*$ is Gaussian. Then by Theorem 5.1 there exists $z \in \mathcal{H}$, $M \in \mathbf{S}(\mathcal{H})$ and $\rho_t := \prod_{j=1}^{\infty} (1 - e^{-t_j}) \otimes_{j=1}^{\infty} e^{-t_j a_j^\dagger a_j} \in \mathcal{B}(\Gamma_s(\mathcal{H}))$ such that

$$U\rho_s U^* = W(z)^* \Gamma_s(M)^* \rho_t \Gamma_s(M) W(z). \quad (6.7)$$

By Lemma 6.3, ρ_s has a complete orthonormal eigenbasis with distinct eigenvalues. By (6.7) ρ_s and ρ_t are unitarily equivalent and thus their eigenvalues and multiplicities are same. Therefore by applying Lemma 6.3 to ρ_t , $\prod_{j=1}^{\infty} (1 - e^{-t_j}) = p$ (since p

is the maximum eigenvalue of ρ_s) and ρ_t has a set of distinct eigenvalues $pe^{-\sum_{j=1}^N t_j k_j}$ corresponding to the eigenvectors $|\mathbf{k}\rangle, \mathbf{k} = (k_1, k_2, \dots, k_N, 0, 0, \dots)^T \in \mathbb{Z}_{+,0}^{\infty}, N \in \mathbb{N}$.

Claim. The sequence $\{t_j\}_{j \in \mathbb{N}}$ consists of (distinct) numbers which are linearly independent over the field \mathbb{Q} .

Proof (of Claim). If $t_i = t_k$ for some $i \neq k$ then it is possible to choose distinct $\mathbf{k}, \mathbf{k}' \in \mathbb{Z}_{+,0}^{\infty}$ such that the eigenvalues of ρ_t corresponding to $|\mathbf{k}\rangle$ and $|\mathbf{k}'\rangle$ are same. This will imply that the corresponding eigenspace is at least two dimensional which

is not possible. To see the rational independence note that for any two finite subsets $I, J \subset \mathbb{N}$, $\sum_{j \in I} t_j k_j \neq \sum_{j \in J} t_j k'_j$ where $k_j, k'_j \in \mathbb{Z}_+, \forall j$. Now if

$$\sum_{j=1}^N t_j q_j = 0 \quad (6.8)$$

for a finite collection of rational numbers q_j 's, since $t_j > 0, \forall j$ then there must be negative rational numbers in the set $\{q_1, q_2, \dots, q_N\}$ (unless $q_j = 0, \forall j$). Then 6.8 can be written in the form $\sum_{j \in I} t_j k_j = \sum_{j \in J} t_j k'_j$ for two finite sets I, J , which is not possible. Thus the claim is proved.

We have $\{pe^{-\sum_{j=1}^N s_j k_j} | k_j \in \mathbb{Z}_+, N \in \mathbb{N}\} = \{pe^{-\sum_{j=1}^N t_j k_j} | k_j \in \mathbb{Z}_+, N \in \mathbb{N}\}$. Therefore $\{\sum_{j=1}^n s_j k_j | k_j \in \mathbb{Z}_+, \forall j, n \in \mathbb{N}\} = \{\sum_{j=1}^n t_j k_j | k_j \in \mathbb{Z}_+, \forall j, n \in \mathbb{N}\}$. Now by the proof of Lemma 6.2, there is a bijection $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ such that $s_j = t_{\sigma(j)}$ for all $j \in \mathbb{N}$.

By (6.7) there exists a unitary V such that

$$V \rho_s V^* = \rho_t. \quad (6.9)$$

where $V = \Gamma_s(M)W(z)U$. Let $\mathbf{k} = (k_1, k_2, \dots, k_N, 0, 0, \dots)^\tau \in \mathbb{Z}_{+,0}^\infty$ be arbitrary, by 6.9 if $|\mathbf{k}\rangle$ is an eigenvector for ρ_s then $V|\mathbf{k}\rangle$ is an eigenvector for ρ_t with the same eigenvalue. Therefore, $V|\mathbf{k}\rangle$ is an eigenvector for ρ_t with eigenvalue $pe^{-\sum_{j=1}^N s_j k_j} = pe^{-\sum_{j \in \mathbb{N}} t_{\sigma(j)} k_j}$. But defining the unitary operator A on \mathcal{H} , $A_\sigma(e_j) = e_{\sigma(j)}$, its second quantization, $\Gamma(A)\mathbf{k}$ is an eigenvector for ρ_t with same eigenvalue. Since the multiplicity for each eigenvalue is one, there exists a complex number $\beta(\mathbf{k})$ of unit modulus such that,

$$\begin{aligned} V|\mathbf{k}\rangle &= \beta(\mathbf{k}) |A\mathbf{k}\rangle \\ &= \Gamma_s(A)\beta(\mathbf{k}) |\mathbf{k}\rangle \\ &= \Gamma_s(A)\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) |\mathbf{k}\rangle \end{aligned}$$

Then by (6.9) $U = W(z)^* \Gamma_s(M)^* \Gamma_s(A) \beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots)$. Now the proof is complete due to Theorem 2.7. It should be noted that we may need to redefine β if the multiplier $\sigma(M^{-1}, A) \neq 1$ (refer Theorem 2.7). \square

The following theorem characterizes the Gaussian symmetries. Recall the definition of Shale operators from § 2.3.

6.5 Theorem. *A unitary operator $U \in \mathcal{B}(\Gamma_s(\mathcal{H}))$ is a Gaussian symmetry if and only if*

$$U = \lambda W(\alpha) \Gamma_s(L)$$

for some $\lambda \in \mathbb{C}$ with $|\lambda| = 1$, $\alpha \in \mathcal{H}$, and L is a Shale operator ($L \in \mathbf{S}(\mathcal{H})$).

Proof. The sufficiency is immediate. To prove the necessity, let us consider $\mathcal{H} = \oplus_j \mathbb{C} e_j$ with respect to some orthonormal basis $\{e_j\}$. By considering a Gaussian state ρ_s as in Lemma 6.4, we can assume without loss of generality that $U =$

$\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots)$ for some β . We will show that $U = \Gamma_s(D)$ for some unitary operator D and this will prove the theorem because of (3) of Theorem 2.7.

Let $\psi \in \Gamma_s(\mathcal{H})$ be such that $|\psi\rangle\langle\psi|$ is a pure Gaussian state. Then by assumption $|U\psi\rangle\langle U\psi|$ is also a Gaussian state and it is clearly a pure state as it is obtained from the wave function $|U\psi\rangle$. We choose the coherent state (Example 1)

$$\psi = e^{-\frac{1}{2}\|u\|^2} |e(u)\rangle = W(u) |e(0)\rangle,$$

where $u = \sum_j u_j e_j$. Now

$$|U\psi\rangle = e^{-\frac{1}{2}\|u\|^2} \beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) |e(u)\rangle \quad (6.10)$$

Recall that

$$e(u) = \sum_{\mathbf{k} \in \mathbb{Z}_{+,0}^\infty} \frac{u^{\mathbf{k}}}{\sqrt{\mathbf{k}!}} |\mathbf{k}\rangle$$

Consider any ‘finite’ vector $z = \sum_{j=1}^N z_j e_j$, for some z_1, z_2, \dots, z_N in \mathbb{C} , $N \in \mathbb{N}$. Then

$$e(z) = \sum_{\mathbf{m} \in \mathbb{Z}_+^N} \frac{z^{\mathbf{m}}}{\sqrt{\mathbf{m}!}} |\mathbf{m}\rangle, \quad (6.11)$$

where $\mathbf{m} \in \mathbb{Z}_+^N$ is considered as the vector $(m_1, m_2, \dots, m_N, 0, 0, \dots)^t \in \mathbb{Z}_{+,0}^\infty$ and $|\mathbf{m}\rangle = |m_1\rangle |m_2\rangle \cdots |m_N\rangle \in \Gamma_s(\mathcal{H})$ as in the notation of (6.2). We will evaluate the function $f(z) = \langle U\psi, e(z) \rangle$ using (6.10) and (6.15). From (6.3), (6.10), (6.11) and continuity of $\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots)$ we have

$$\begin{aligned} f(z) &= e^{-\frac{1}{2}\|u\|^2} \langle Ue(u), e(z) \rangle \\ &= e^{-\frac{1}{2}\|u\|^2} \left\langle \sum_{\mathbf{k} \in \mathbb{Z}_{+,0}^\infty} \frac{u^{\mathbf{k}}}{\sqrt{\mathbf{k}!}} U |\mathbf{k}\rangle, \sum_{\mathbf{m} \in \mathbb{Z}_+^N} \frac{z^{\mathbf{m}}}{\sqrt{\mathbf{m}!}} |\mathbf{m}\rangle \right\rangle \\ &= e^{-\frac{1}{2}\|u\|^2} \left\langle \sum_{\mathbf{k} \in \mathbb{Z}_{+,0}^\infty} \frac{u^{\mathbf{k}}}{\sqrt{\mathbf{k}!}} \beta(\mathbf{k}) |\mathbf{k}\rangle, \sum_{\mathbf{m} \in \mathbb{Z}_+^N} \frac{z^{\mathbf{m}}}{\sqrt{\mathbf{m}!}} |\mathbf{m}\rangle \right\rangle \end{aligned} \quad (6.12)$$

Thus

$$f(z) = e^{-\frac{1}{2}\|u\|^2} \sum_{\mathbf{k} \in \mathbb{Z}_+^N} \frac{(\bar{u}_1 z_1)^{k_1} (\bar{u}_2 z_2)^{k_2} \cdots (\bar{u}_N z_N)^{k_N}}{k_1! k_2! \cdots k_N!} \overline{\beta(\mathbf{k})}. \quad (6.13)$$

Since $|\beta(\mathbf{k})| = 1$, from (6.13) we see that

$$|f(z)| \leq \exp \left\{ -\frac{1}{2}\|u\|^2 + \sum_{j=1}^N |u_j| |z_j| \right\}. \quad (6.14)$$

Now we compute $f(z)$ in an alternative way. As $|U\psi\rangle\langle U\psi|$ is a pure Gaussian state, by Corollary 5.2 there exists a unitary A and an $\alpha \in \mathcal{H}$ such that

$$|U\psi\rangle = W(\alpha) \Gamma_s(A) \otimes_j |e_{\lambda_j}\rangle. \quad (6.15)$$

From the definition of $e(w)$ and e_λ in $\Gamma(\mathbb{C}) = L^2(\mathbb{R})$, we have

$$e_\lambda(x) = (2\pi)^{-1/4} \lambda^{-1/2} \exp\{-4^{-1} \lambda^{-2} x^2\}, \quad x \in \mathbb{R}, \lambda > 0$$

on $L^2(\mathbb{R})$. Further,

$$\langle e_\lambda, e(w) \rangle = \sqrt{\frac{2\lambda}{1+\lambda^2}} \exp \frac{1}{2} \left(\frac{\lambda^2 - 1}{\lambda^2 + 1} \right) w^2, \lambda > 0, w \in \mathbb{C}. \quad (6.16)$$

Using (6.15),

$$\begin{aligned} f(z) &= \langle W(\alpha) \Gamma_s(A) \otimes_j e_{\lambda_j}, e(z) \rangle \\ &= \langle \otimes_j e_{\lambda_j}, \Gamma_s(A^*) W(-\alpha) e(z) \rangle \\ &= e^{\langle \alpha, z \rangle - \frac{1}{2} \|\alpha\|^2} \langle \otimes_j e_{\lambda_j}, e(A^*(z - \alpha)) \rangle \end{aligned} \quad (6.17)$$

Since z is a finite vector and α is fixed, each coordinate of $A^*(z - \alpha)$ is a first degree polynomial in z_j 's. Therefore $e(A^*(z - \alpha)) = \otimes_j e(w_j)$ where each w_j is a first degree polynomial in z_j 's. Therefore from (6.16) and property of infinite tensor products

$$f(z) = e^{\langle \alpha, z \rangle - \frac{1}{2} \|\alpha\|^2} \lim_{n \rightarrow \infty} \prod_{j=1}^n \sqrt{\frac{2\lambda_j}{1+\lambda_j^2}} \exp \sum_{j=1}^n \frac{1}{2} \left(\frac{\lambda_j^2 - 1}{\lambda_j^2 + 1} \right) w_j^2$$

Since each w_j^2 is a second degree polynomial in z_1, z_2, \dots, z_N . This contradicts (6.14) unless $\lambda_j = 1$ for all j . Now (6.15) implies

$$\begin{aligned} |U\psi\rangle &= W(\alpha) \Gamma_s(A) |e(0)\rangle \\ &= e^{-\frac{1}{2} \|\alpha\|^2} |e(\alpha)\rangle \end{aligned}$$

Now from (6.10) we get

$$e^{-\frac{1}{2} \|u\|^2} \beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) |e(u)\rangle = e^{-\frac{1}{2} \|\alpha\|^2} |e(\alpha)\rangle \quad (6.18)$$

Thus $\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots)$ is a unitary with the following properties:

1. $\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) |\mathbf{k}\rangle = \beta(\mathbf{k}) |\mathbf{k}\rangle$ for every $\mathbf{k} \in \mathbb{Z}_{+,0}^\infty$.
2. It maps coherent vectors to coherent vectors.

We will prove that $\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) = \Gamma_s(D)$ for a diagonal unitary D . To this end we fix a $u = \sum_j u_j e_j$ in \mathcal{H} with $u_j \neq 0, \forall j$. We have $\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) |e(u)\rangle = e^{\frac{1}{2} (\|u\|^2 - \|\alpha\|^2)} |e(\alpha)\rangle$. Therefore if $\alpha = \sum_j \alpha_j e_j$ from (6.18) and (6.3) we get,

$$\sum_{\mathbf{k} \in \mathbb{Z}_{+,0}^\infty} \frac{u^{\mathbf{k}}}{\sqrt{\mathbf{k}!}} \beta(\mathbf{k}) |\mathbf{k}\rangle = e^{\frac{1}{2} (\|u\|^2 - \|\alpha\|^2)} \sum_{\mathbf{k} \in \mathbb{Z}_{+,0}^\infty} \frac{\alpha^{\mathbf{k}}}{\sqrt{\mathbf{k}!}} |\mathbf{k}\rangle$$

Therefore,

$$u^{\mathbf{k}} \beta(\mathbf{k}) = e^{\frac{1}{2} (\|u\|^2 - \|\alpha\|^2)} \alpha^{\mathbf{k}}, \forall \mathbf{k} \in \mathbb{Z}_{+,0}^\infty$$

Since $u_j \neq 0$ for all j , we see that if $\mathbf{k} = (k_1, k_2, \dots, k_m, 0, 0, \dots) \in \mathbb{Z}_{+,0}^\infty$,

$$\beta(\mathbf{k}) = e^{\frac{1}{2} (\|u\|^2 - \|\alpha\|^2)} \left(\frac{\alpha_1}{u_1} \right)^{k_1} \left(\frac{\alpha_2}{u_2} \right)^{k_2} \dots \left(\frac{\alpha_m}{u_m} \right)^{k_m}, \forall \mathbf{k} \in \mathbb{Z}_{+,0}^\infty$$

Since $|\beta(\mathbf{k})| = 1$, we get $\left|\frac{\alpha_j}{u_j}\right| = 1$ for all j . If we write $\frac{\alpha_j}{u_j} = e^{i\theta_j}$, then from (6.18) we get

$$\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) |e(u)\rangle = |e(Du)\rangle, \quad (6.19)$$

where D is the unitary $\text{Diag}(e^{i\theta_1}, e^{i\theta_2}, \dots)$, for every $u = \sum_j u_j e_j \in \mathcal{H}$ with $u_j \neq 0, \forall j$. Now it is easy to see that (6.19) holds for all $u \in \mathcal{H}$. We conclude that $\beta(a_1^\dagger a_1, a_2^\dagger a_2, \dots) = \Gamma_s(D)$. \square

6.6 Remark. Observe that the set of all Gaussian symmetries $\{\lambda W(\alpha) \Gamma_s(L) : \lambda \in \mathbb{C}, |\lambda| = 1, \alpha \in \mathcal{H}, L \in \mathcal{S}(\mathcal{H})\}$ form a group under multiplication. A typical computation in the proof of the previous statement is given below.

$$\begin{aligned} \lambda_1 W(\alpha_1) \Gamma_s(L_1) \lambda_2 W(\alpha_2) \Gamma_s(L_2) &= \lambda_1 \lambda_2 W(\alpha_1) \Gamma_s(L_1) W(\alpha_2) \Gamma_s(L_1)^{-1} \Gamma_s(L_1) \Gamma_s(L_2) \\ &= \lambda_1 \lambda_2 W(\alpha_1) W(L_1 \alpha_2) \Gamma_s(L_1 L_2) \\ &= \lambda_1 \lambda_2 e^{-i \text{Im} \langle \alpha_1, L_1 \alpha_2 \rangle} W(\alpha_1 + L_1 \alpha_2) \Gamma_s(L_1 L_2). \end{aligned}$$

The quantity in the last line is again in the form of a prototypical element in the set of Gaussian symmetries.

Combining Theorem 6.5 with Theorem 5.1, we have the following important observation.

6.7 Remark. The symplectic spectrum (Remark 2.11) of the covariance operator is a complete invariant for Gaussian states and two Gaussian states with same symplectic spectrum are conjugate to each other through a Gaussian symmetry.

Acknowledgements: We wish to thank Prof. K. R. Parthasarathy for introducing the subject to us during his 2015 lecture series at the Indian Statistical Institute, Bangalore centre. John thanks Prof. Martin Lindsay and Prof. K. B. Sinha for discussions on quasifree states and metaplectic representations respectively. Bhat thanks J C Bose Fellowship for financial support. We sincerely thank the anonymous referee for several constructive suggestions which helped us to improve the paper immensely.

References

- [Ara72] Huzihiro Araki, *On quasifree states of the canonical commutation relations. II*, Publ. Res. Inst. Math. Sci. **7** (1971/72), 121–152. MR 0313834
- [ARL14] Gerardo Adesso, Sammy Ragy, and Antony R. Lee, *Continuous variable quantum information: Gaussian states and beyond*, Open Syst. Inf. Dyn. **21** (2014), no. 1-2, 1440001, 47. MR 3190189
- [AS72] Huzihiro Araki and Masafumi Shiraishi, *On quasifree states of the canonical commutation relations. I*, Publ. Res. Inst. Math. Sci. **7** (1971/72), 105–120.
- [BJ19] B. V. Rajarama Bhat and Tiju Cherian John, *Real Normal Operators and Williamson’s Normal Form*, Acta Sci. Math. (Szeged), To appear (2019).

- [BR97] Ola Bratteli and Derek W. Robinson, *Operator algebras and quantum statistical mechanics. 2*, second ed., Texts and Monographs in Physics, Springer-Verlag, Berlin, 1997, Equilibrium states. Models in quantum statistical mechanics.
- [BS05] B. V. Rajarama Bhat and R. Srinivasan, *On product systems arising from sum systems*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **8** (2005), no. 1, 1–31. MR 2126876
- [DG13] Jan Dereziński and Christian Gérard, *Mathematics of quantization and quantum fields*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 2013. MR 3060648
- [Fol89] Gerald B. Folland, *Harmonic analysis in phase space*, *Annals of Mathematics Studies*, vol. 122, Princeton University Press, Princeton, NJ, 1989. MR 983366
- [FOP05] A. Ferraro, S. Olivares, and M. Paris, *Gaussian states in quantum information*, Napoli series on physics and astrophysics, Bibliopolis, 2005.
- [Hol71a] A. S. Holevo, *Quasi-free states on the c^* algebra of ccr*, *Theoretical and Mathematical Physics* **6** (1971), no. 1, 1–12.
- [Hol71b] ———, *Quasi-free states on the C^* -algebra of commutation relations. II*, *Teoret. Mat. Fiz.* **6** (1971), no. 2, 145–150. MR 0479144
- [HR96] Reinhard Honegger and Alfred Rieckers, *Squeezing bogoliubov transformations on the infinite mode ccr-algebra.*, *J. Math. Phys.* **37** (1996), no. 9, 4292–4309. MR 1408093
- [MS04] Taku Matsui and Yoshihito Shimada, *On quasifree representations of infinite dimensional symplectic group*, *Journal of Functional Analysis* **215** (2004), no. 1, 67 – 102.
- [Par10] K. R. Parthasarathy, *What is a Gaussian state?*, *Commun. Stoch. Anal.* **4** (2010), no. 2, 143–160. MR 2662722
- [Par12] ———, *An introduction to quantum stochastic calculus*, *Modern Birkhäuser Classics*, Birkhäuser, 2012, [2012 reprint of the 1992 original].
- [Par13] ———, *The symmetry group of Gaussian states in $l^2(\mathbb{R}^n)$* , *Prokhorov and contemporary probability theory*, *Springer Proc. Math. Stat.*, vol. 33, Springer, Heidelberg, 2013, pp. 349–369. MR 3070484
- [Par15] ———, *Symplectic dilations, gaussian states and gaussian channels*, *Indian Journal of Pure and Applied Mathematics* **46** (2015), no. 4, 419–439.
- [Pet90] Dénes Petz, *An invitation to the algebra of canonical commutation relations*, *Leuven Notes in Mathematical and Theoretical Physics. Series A: Mathematical Physics*, vol. 2, Leuven University Press, Leuven, 1990. MR 1057180

- [PS72] K. R. Parthasarathy and K. Schmidt, *Positive definite kernels, continuous tensor products, and central limit theorems of probability theory*, Lecture Notes in Mathematics, Vol. 272, Springer-Verlag, Berlin-New York, 1972. MR 0622034
- [Sha62] David Shale, *Linear symmetries of free boson fields*, Trans. Amer. Math. Soc. **103** (1962), 149–167. MR 0137504
- [Tve04] I. D. Tveritinov, *Several remarks on the representation of the infinite-dimensional symplectic group and on the construction of the metaplectic group*, Mathematical Notes **75** (2004), no. 5, 805–818.
- [vD71] A. van Daele, *Quasi-equivalence of quasi-free states on the weyl algebra*, Comm. Math. Phys. **21** (1971), 171–191. MR 0287844
- [WGC06] Michael M. Wolf, Geza Giedke, and J. Ignacio Cirac, *Extremality of gaussian quantum states*, Phys. Rev. Lett. **96** (2006), 080502.
- [WHTH07] Xiang-Bin Wang, Tohya Hiroshima, Akihisa Tomita, and Masahito Hayashi, *Quantum information with gaussian states*, Physics Reports **448** (2007), no. 1, 1 – 111.