Hidden Hamiltonian Cycle Recovery via Linear Programming

Vivek Bagaria, Jian Ding, David Tse, Yihong Wu, Jiaming Xu^{*}

March 13, 2022

Abstract

We introduce the problem of hidden Hamiltonian cycle recovery, where there is an unknown Hamiltonian cycle in an *n*-vertex complete graph that needs to be inferred from noisy edge measurements. The measurements are independent and distributed according to P_n for edges in the cycle and Q_n otherwise. This formulation is motivated by a problem in genome assembly, where the goal is to order a set of contigs (genome subsequences) according to their positions on the genome using long-range linking measurements between the contigs. Computing the maximum likelihood estimate in this model reduces to a Traveling Salesman Problem (TSP). Despite the NP-hardness of TSP, we show that a simple linear programming (LP) relaxation, namely the fractional 2-factor (F2F) LP, recovers the hidden Hamiltonian cycle with high probability as $n \to \infty$ provided that $\alpha_n - \log n \to \infty$, where $\alpha_n \triangleq -2\log \int \sqrt{dP_n dQ_n}$ is the Rényi divergence of order $\frac{1}{2}$. This condition is information-theoretically optimal in the sense that, under mild distributional assumptions, $\alpha_n \ge (1 + o(1)) \log n$ is necessary for any algorithm to succeed regardless of the computational cost.

Departing from the usual proof techniques based on dual witness construction, the analysis relies on the combinatorial characterization (in particular, the half-integrality) of the extreme points of the F2F polytope. Represented as bicolored multi-graphs, these extreme points are further decomposed into simpler "blossom-type" structures for the large deviation analysis and counting arguments. Evaluation of the algorithm on real data shows improvements over existing approaches.

^{*}V. Bagaria is with Stanford University, Stanford, USA vbagaria@stanford.edu. J. Ding is with University of Pennsylvania, Philadelphia, USA dingjian@wharton.upenn.edu. D. Tse is with Stanford University, Stanford, USA dntse@stanford.edu. Y. Wu is with Yale University, New Haven, USA yihong.wu@yale.edu. J. Xu is with Purdue University, West Lafayette, USA xu972@purdue.edu.

1 Introduction

Given an input graph, the problem of finding a subgraph satisfying certain properties has diverse applications. MAX CUT, MAX CLIQUE, TSP are a few canonical examples. Traditionally these problems have been studied in theoretical computer science from the worst-case perspective, and many such problems have been shown to be NP-hard. However, in machine learning applications, many such problems arise when an underlying ground truth subgraph needs to be recovered from the noisy measurement data represented by the entire graph. Canonical models to study such problems include planted partition models [CK01] (such as planted clique [Jer92]) in community detection and planted ranking models (such as Mallows model [Mal57]) in rank aggregation. In these models, the planted or hidden subgraph represents the ground-truth, and one is not necessarily interested in the worst-case instances but rather in only instances for which there is enough information in the data to recover the ground-truth sub-graph, i.e. when the amount of data is above the information limit. The key question is whether there exists an efficient recovery algorithm that can be successful all the way to the information limit.

In this paper, we pose and answer this question for a hidden Hamiltonian cycle recovery model:

Definition 1 (Hidden Hamiltonian cycle recovery).

Given: $n \ge 1$, and two distributions P_n and Q_n , parameterized by n.

Observation: A randomly weighted, undirected complete graph G = ([n], E) with a hidden Hamiltonian cycle C^* such that every edge has an independent weight distributed as P_n if it is on C^* and as Q_n otherwise.

Inference Problem: Recover the hidden Hamiltonian cycle C^* from the observed random graph.

Our problem is motivated from *de novo* genome assembly, the reconstruction of an organism's long sequence of A,G,C,T nucleotides from fragmented sequencing data. The first step of the standard assembly pipeline stitches together short, overlapping fragments (so-called shotgun reads) to form longer subsequences called contigs, of lengths typically tens to hundreds of thousands of nucleotides (Fig. 1). Due to coverage gaps and other issues, these individual contigs cannot be extended to the whole genome. To get a more complete picture of the genome, the contigs need to be ordered according to their positions on the genome, a process called *scaffolding*. Recent advances in sequencing assays [LAVBW⁺09, POS⁺16] aid this process by providing long range linking information between these contigs in the form of randomly sampled *Hi-C reads*. This data can be summarized by a contact map (Fig. 2), tabulating the counts of Hi-C reads linking each pair of contigs. The problem of ordering the contigs from the contact map data can be modeled by the hidden Hamiltonian cycle recovery problem, where the vertices of the graph are the contigs, the hidden Hamiltonian cycle is the true ordering of the contigs on the genome,¹ and the weights on the graph are the counts of the Hi-C reads linking the contigs. As can be seen in Fig. 2(a), there is a much larger concentration of Hi-C reads between contigs adjacent on the genome than between far-away contigs. A first order model is to choose $P_n = \text{Pois}(\lambda_n)$ and $Q_n = \text{Pois}(\mu_n)$, where λ_n is the average number of Hi-C reads between adjacent contigs and μ_n is the average number between non-adjacent contigs. The parameter $n\lambda_n + \frac{n(n-1)}{2}\mu_n$ increase with the coverage depth² of the Hi-C reads and are part of the design of the sequencing experiment.

The hidden Hamiltonian cycle can be represented as an adjacency vector $x^* \in \{0, 1\}^{\binom{n}{2}}$ such that $x_e^* = 1$ if edge e is on the Hamiltonian cycle, and $x_e^* = 0$ otherwise. Let A denote the weighted

¹Strictly speaking, this only applies to genomes which are circular. For genomes which are linear, the ordering of the contigs would correspond to a hidden Hamiltonian *path*. We show in Section 7 that our results extend to a hidden Hamiltonian path model as well.

²The coverage depth is the average number of Hi-C reads that include a given nucleotide (base pair).




Figure 1: Short reads are assembled to form contigs which are then ordered by using long-range linking Hi-C reads.




Figure 2: (a): Contact map where the rows (and columns) correspond to *ordered* contigs of human chromosome 1 [POS⁺16] and the value at entry (i, j) corresponds to the number of HiC reads between contig *i* and contig *j*. (b): Contact map of the *unordered* matrix in (a), where the contigs are randomly ordered. This is the data from which the ordering of the contigs is inferred.

adjacency matrix of G, so that A_e is distributed according to P (resp. Q) if $x_e^* = 1$ (resp. 0). The maximum likelihood (ML) estimator for the hidden Hamiltonian cycle recovery problem is equivalent to solving the traveling salesman problem (TSP) on a transformed weighted graph, where each edge weight $w_e = \log \frac{dP}{dQ}(A_e)$ is the log likelihood ratio evaluated on the weights of the observed graph:

$$\widehat{x}_{\text{ML}} = \arg \max_{x} \langle w, x \rangle$$
s.t. x is the adjacency vector of a Hamiltonian cycle in G. (1)

In the Poisson or Gaussian model where the log likelihood ratio is an affine function, we can simply take w to be A itself.

Solving TSP is NP-hard, and a natural approach is to look for a tractable relaxation. It is

well-known that TSP (1) can be cast as an integer linear program (ILP) [SWvZ13]:

$$\widehat{x}_{\text{TSP}} = \arg\max \langle w, x \rangle \tag{2}$$

s.t.
$$x(\delta(v)) = 2$$
 (3)

$$x(\delta(S)) \ge 2, \ \forall S \subset [n], \ 3 \le |S| \le n-3 \tag{4}$$

 $x_e \in \{0, 1\},$ (5)

where $\delta(S)$ denotes the set of all edges in G with exactly one endpoint in $S \subset [n]$, and $\delta(v) \triangleq \delta(\{v\})$. In particular, (3) are called *degree* constraints, enforcing each vertex to have exactly two incident edges in the graph represented by the adjacency vector x, while (4) are subtour elimination constraints, eliminating solutions whose corresponding graph is a disjoint union of subtours of length less than n. Note that there are small number of degree constraints but exponentially large number of subtour elimination constraints. If we drop the subtour elimination constraints as well as relax the integer constraints on x, we obtain the *fractional 2-factor* (F2F) LP relaxation:³

$$\widehat{x}_{F2F} = \arg \max_{x} \langle w, x \rangle$$
s.t. $x (\delta(v)) = 2$
 $x_e \in [0, 1].$
(6)

The main result of the paper is the following. We abbreviate P_n and Q_n as P and Q, respectively.

Theorem 1. Define:

$$\alpha_n \triangleq -2\log \int \sqrt{\mathrm{d}P\mathrm{d}Q} \tag{7}$$

to be the Rényi divergence of order $\frac{1}{2}$ between distributions P and Q.⁴ If

$$\alpha_n - \log n \to +\infty,\tag{9}$$

then the optimal solution of the F2F LP (6) satisfies $\min_{x^*} \mathbb{P} \{ \widehat{x}_{F2F} = x^* \} \to 1 \text{ as } n \to \infty.$

For Gaussian, Poisson, or Bernoulli weight distribution, the explicit expressions of α_n are given as follows

$$\alpha_n = \begin{cases} \mu^2/4 & \text{if } P = N(\mu, 1), \ Q = N(0, 1) \\ \left(\sqrt{\lambda} - \sqrt{\mu}\right)^2 & \text{if } P = \text{Pois}(\lambda), \ Q = \text{Pois}(\mu) \\ -2\log\left(\sqrt{pq} + \sqrt{(1-p)(1-q)}\right) & \text{if } P = \text{Bern}(p), \ Q = \text{Bern}(q). \end{cases}$$
(10)

Although the relaxation from TSP to F2F LP is quite drastic, the resulting algorithm is in fact information theoretically optimal for the hidden Hamiltonian cycle recovery problem. Specifically, under an assumption which can be easily verified for Poisson, Gaussian or Bernoulli weight distribution, we show in Section 6 that if there exists any algorithm, efficient or not, which exactly recovers x^* with high probability, then it must hold that

$$\alpha_n \ge (1 + o(1)) \log n.$$

$$D_{\rho}(P||Q) \triangleq \frac{1}{\rho - 1} \log \int (dP)^{\rho} (dQ)^{1 - \rho}.$$
(8)

It particular, for $\rho = 1/2$ it is related to the so-called Battacharyya distance B(P,Q) via $D_{\frac{1}{2}}(P||Q) = 2B(P,Q)$.

³ A 2-factor is a spanning subgraph consisting of disjoint cycles.

 $^{^4}$ The Rényi divergence of order $\rho>0$ from P to Q is defined as [Rén61]

This necessary condition together with sufficient condition (9) implies that the optimal recovery threshold is at

$$\liminf_{n \to \infty} \frac{\alpha_n}{\log n} = 1$$

achieved by the F2F LP.

Applying the results back to the geome scaff We discuss two consequences of Theorem 1. First, as a corollary of the integrality and the optimality of the F2F LP, it can be shown that the Max-Product belief propagation algorithm introduced in [BBCZ11] can be used to solve the F2F LP exactly, which, for the Gaussian or Poisson weight distribution, requires $o(n^2 \log n)$ iterations (see Section 2 for details). Second, note that we do not require the edge weights to be real-valued. Thus the formulation also encompasses the case of partial observation, by letting the weight of every edge in G takes on a special "erasure" symbol with some probability. See Appendix F for details.

In related work, a version of the hidden Hamiltonian cycle model was studied in [BFS94], where the observed graph is the superposition of a hidden Hamiltonian cycle and an Erdös-Rényi random graph with constant average degree d. Our measurement model is more general than the one in [BFS94], but more importantly, the goal in [BFS94] is not to recover the hidden Hamiltonian cycle but rather to find any Hamiltonian cycle in the observed graph, which may not coincide with the hidden one. (In fact in the regime considered there, exact recovery of the hidden cycle is information theoretically impossible.⁵) The fractional 2-factor relaxation of TSP has been wellstudied in the worst case [DFJ54, BC99, SWvZ13]. It has been shown that for metric TSP (the cost minimization formulation) where the costs are symmetric and satisfy the triangle inequality, the *integral gap* of F2F is 4/3; here the integrality gap is defined as the worst-case ratio of the cost of the optimal integral solution to the cost of the optimal relaxed solution. In contrast, our model does not make any metric assumption on the graph weights.

The rest of the paper is organized as follows. In Section 2, we describe a few other computationally efficient algorithms for the hidden Hamiltonian cycle problem and benchmark their performance against the information-theoretic limit. In Section 3, we discuss related work in more details. Sections 4 and 5 are devoted to the proof of Theorem 1, while Section 6 characterizes the information theoretic limit for the recovery problem. In Section 7 we describe the closely related hidden Hamiltonian path problem and show that it can be reduced to and from the hidden Hamiltonian cycle problem both statistically and computationally. Empirical evaluation of various algorithms on both simulated and real DNA datasets are given in Section 8.

2 Performance of Other Algorithms

It is striking to see that the simple F2F LP relaxation of the TSP achieves the optimal recovery threshold in the hidden Hamiltonian cycle model. A natural question to ask is whether there exists other efficient and perhaps even simpler estimator with provable optimality. We have considered various efficient algorithms and derived their performance guarantees. As summarized in Table 1, spectral algorithms is orderwise suboptimal; greedy methods including thresholding achieve the optimal scaling but not the sharp constant. Finally, MaxProduct Belief Propagation also achieves

⁵To see this, suppose the hidden Hamiltonian cycle is given by sequence of vertices (1, 2, ..., n, 1). If $q = \Omega(1/n)$, then with a non-vanishing probability there exist $1 \le i \le n-4$ and $i+2 \le j \le n-2$ such that (i, j) and (i+1, j+1) are edges in G. Thus we have a new Hamiltonian cycle by deleting edges (i, i+1) and (j, j+1) in the hidden one and adding edges (i, j) and (i+1, j+1), leading to the impossibility of exact recovery. See Fig. 14 for an illustration.

| Efficient Algorithms | Performance Guarantee |
|----------------------|-----------------------------------|
| F2F LP | $\alpha_n - \log n \to +\infty$ |
| MaxProduct BP | |
| Greedy Merging | $\beta_n - \log n \to +\infty$ |
| Simple Thresholding | $\alpha_n - 2\log n \to +\infty$ |
| Nearest Neighbor | $\alpha_n - 2 \log n \to +\infty$ |
| Spectral Methods | $\alpha_n \gg n^5$ (Gaussian) |

Table 1: Sufficient conditions for various efficient algorithms to achieve exact recovery.

the sharp threshold as a corollary of our result on the F2F LP. In Table 1,

$$\beta_n \triangleq -\frac{3}{2} \log \int (\mathrm{d}P)^{2/3} (\mathrm{d}Q)^{1/3}$$
 (11)

is the $\frac{1}{3}$ -Rényi divergence from Q to P; cf. (8). By Jensen's and Hölder's inequality, for any distinct P and Q, we have

$$\frac{1}{2}\alpha_n < \beta_n < \alpha_n. \tag{12}$$

For Gaussian weights with $P = \mathcal{N}(\mu, 1)$ and $Q = \mathcal{N}(0, 1)$, we have $\beta_n = \frac{1}{6}\mu^2 = \frac{2}{3}\alpha_n$. Simulation of these algorithms confirm these theoretical results. See Figure 15 in Section 8.1.

Spectral Methods Spectral algorithms are powerful methods for recovering the underlying structure in planted models based on the principal eigenvectors of the observed adjacency matrix A. Under planted models such as planted clique [AKS98] or planted partition models [McS01], spectral algorithms and their variants have been shown to achieve either the optimal recovery thresholds [Mas13, BLM15, AS15] or the best possible performance within certain relaxation hierarchies [MPW15, DM15, BHK⁺16]. The rationale behind spectral algorithms is that the principal eigenvectors of $\mathbb{E}[A]$ contains information about underlying structures and the principal eigenvectors of A are close to those of $\mathbb{E}[A]$, provided that the spectral gap (the gap between the largest few eigenvalues and the rest of them) is much larger than the spectral norm of the perturbation $||A - \mathbb{E}[A]||$. In our setting, indeed the principal eigenvectors of $\mathbb{E}[A]$ contain information about $P = \mathcal{N}(\mu, 1)$ and $Q = \mathcal{N}(0, 1)$ as an illustrating example. Then the observed matrix can be expressed as

$$A = \mu C^* + Z,$$

where with a slight abuse of notation we use A to denote the weighted adjacency matrix of Gand C^* to denote the adjacency matrix of the true Hamiltonian cycle; Z is a symmetric Gaussian matrix with zero diagonal and $Z_{ij} = Z_{ji}$ independently drawn from $\mathcal{N}(0,1)$ for i < j. Since C^* is a circulant matrix, its eigenvalues and the corresponding eigenvectors can be explicitly derived via discrete Fourier transform. It turns out that the eigenvector corresponding to the second largest eigenvalue of C^* contains perfect information about the true Hamiltonian cycle. Unfortunately, in contrast to the planted clique and planted partition models under which $\mathbb{E}[A]$ is low rank and has a large eigen-gap, here C^* is full-rank and the gap between the second and the third largest eigenvalue is on the order of $1/n^2$, which is much smaller than $||Z|| = \Theta(\sqrt{n})$. Therefore, for spectral algorithms to succeed, it requires a very high signal level $\mu^2 \gg n^5$. This agrees with the empirical performance on simulated data and is highly suboptimal as compared to the sufficient condition (9) of F2F LP: $\mu^2 - 4 \log n \to \infty$. **Greedy Methods** To recover the hidden Hamiltonian cycle, we can also resort to greedy methods. It turns out that the following simple thresholding algorithm achieves the optimal recovery threshold (9) within a factor of two: for each vertex, keep the two incident edges with the two largest weights and delete the other n-3 edges. The resulting graph has degree at most 2. It can be shown that the resulting graph coincides with C^* with high probability provided that $\alpha_n - 2 \log n \to +\infty$.

Another well-known greedy heuristic is the following nearest-neighbor algorithm. Start on an arbitrary vertex as the current vertex and find the edge with the largest weight connecting the current vertex and an unvisited vertex v, set the current vertex to v and mark v as visited. Repeat until all vertices have been visited. Let v_1, \ldots, v_n denote the sequence of visited vertices and output the Hamiltonian cycle formed by (v_1, \ldots, v_n, v_1) . It can be shown (see Appendix E) that the resulting Hamiltonian cycle coincides with C^* with high probability provided that $\alpha_n - 2 \log n \to +\infty$.

Finally, we consider a greedy merging algorithm proposed in [MBT13]: connect pairs of vertices with the largest edge weights until all vertices have degree two. The output is a 2-factor, and it can be shown that the output 2-factor coincides with C^* with high probability provided that $\beta_n - \log n \rightarrow +\infty$, strictly improving on the performance guarantee of previous two greedy algorithms.

Notice that the aforementioned greedy algorithms only exploit local information, and do not take the into account the global cycle structure. Naturally, none of them achieves the optimal threshold (9). See Appendix E for further details.

Max-Product Belief Propagation We can improve on the simple thresholding algorithm using an iterative message-passing algorithm known as max-product belief propagation. Specifically, at each time $t = 0, 1, \ldots, t_f$, each vertex *i* sends a real-valued message $m_{i\to j}(t)$ to each of its neighbors *j*. Messages are initialized by $m_{i\to j}(0) = w_e$ for all e = (i, j). For $t \ge 1$, messages transmitted by vertex *i* in iteration *t* are updated based on messages received in iteration t - 1 recursively as follows:

$$m_{i \to j}(t) = w_e - 2 \operatorname{ndmax}_{\ell \neq j} \left\{ m_{\ell \to i}(t-1) \right\},$$

where 2nd max denotes the second largest value. At the end of the final iteration t_f , for every vertex, keep the two incident edges with the two largest received message values and delete the other n-3 edges, and output the resulting graph. Note that BP with one iteration $t_f = 1$ reduces to the simple thresholding algorithm.

Belief propagation algorithm is studied in [BBCZ11] to find the *b*-factor with the maximum weight for $b \ge 1$; it is shown that if the fractional *b*-factor LP relaxation has no fractional optimum solution, then the output of BP coincides with the optimal *b*-factor when $t_f \ge \lceil 2nw^*/\epsilon \rceil$, where w^* is the weight of the optimal *b*-factor and ϵ is the difference between the weight of the optimal *b*-factor and the second largest weight of *b*-factors. Our optimality result of F2F implies that if $\alpha_n - \log n \to +\infty$, then with high probability, F2F has no fractional optimum solution and the optimal 2-factor coincides with the ground truth x^* ; Therefore, by combining our result with results of BP in [BBCZ11], we immediately conclude that the output of BP coincides with x^* with high probability after t_f iterations, provided that $\alpha_n - \log n \to +\infty$. For both the Gaussian and the Poisson model, with high probability the number of iterations t_f of the BP algorithm is in fact $o(n^2 \log n)$, nearly linear in the problem size (see Appendix G for a justification).

3 Related Work

We discuss additional related work before presenting the proof of our main results. Because of the NP-hardness of TSP, researchers have imposed structural assumptions on the costs (weights) and devised efficient approximation algorithms. One natural assumption is the metric assumption under which the costs are symmetric $(c_{ij} = c_{ji} \text{ for all } i, j \in V)$ and satisfy the triangle inequality $(c_{ik} \leq c_{ij} + c_{jk} \text{ for all } i, j, k \in V)$. Metric TSP turns out to be still NP-hard, as shown by reduction from the NP-hard Hamiltonian cycle problem [Sch03, Theorem 58.1]. The best approximation algorithm for metric TSP currently known is Christofides' algorithm which finds a Hamiltonian cycle of cost at most a factor of 3/2 times the cost of an optimal Hamiltonian cycle.

Integrality gap of LP relaxations of TSP Various relaxations of TSP has also been extensively studied under the metric assumption. To measure the tightness of LP relaxations, a commonly used figure of merit is the *integrality gap*. As is the convention in the TSP literature, the optimization is formulated as a minimization problem with nonnegative costs. In general, the integrality gap is defined as the supremum of the ratio OPT/FRAC, over all instances of the problem, where FRAC denotes the objective value of the optimal fractional solution and OPT denotes the objective value of the optimal integral solution [CT12]. Note that by definition the integrality gap is always at least one. Dropping the integer constraints (5) in ILP formulation of TSP (2) leads to a LP relaxation known as Subtour LP [DFJ54, HK70]. The integrality gap of the subtour LP is known to be between 4/3 and 3/2. The integrality gap of fractional 2-factor LP (6) is shown in [BC99, SWvZ13] to be 4/3. In contrast to the previous worst-case approximation results on metric TSP, this paper focuses on a planted instance of TSP, where we impose probabilistic assumption on the costs (weights) and the goal is to recover the hidden Hamiltonian cycle. In particular, the metric assumption is not fulfilled in our hidden Hamiltonian cycle model and hence the previous results do not apply. Our results imply that when $\alpha_n - \log n \to +\infty$, the optimal solution of F2F coincides with the optimal solution of TSP with probability tending to 1, where the probability is taken over the randomness of weights w in the hidden Hamiltonian cycle model. In other words, for "typical" instances of the hidden Hamiltonian cycle model, the optimal objective value of TSP is the same as that of F2F.

SDP relaxations of TSP Semidefinite programming (SDP) relaxations of the traveling salesman problem have also been extensively studied in the literature. A classical SDP relaxation of TSP due to [CČKV99] is obtained by imposing an extra constraint on the second largest eigenvalue of a Hamiltonian cycle in F2F LP (6). A more sophisticated SDP relaxation is derived in [ZKRW98] by viewing the TSP as a quadratic assignment problem, from which one can obtain a simpler SDP relaxation of TSP based on association schemes [DKPS08]. This SDP relaxation in [DKPS08] is shown to dominate that of [CČKV99]. Since all these SDP relaxations are tighter than the F2F LP, our results immediately imply that the optimal solutions of these SDP relaxations coincide with the true Hamiltonian cycle x^* with high probability provided $\alpha_n - \log n \to +\infty$.

Data seriation The problem of recovering a hidden Hamiltonian cycle (path) in a weighted complete graph falls into a general problem known as data seriation [Ken71] or data stringing [CCMW11]. In particular, we are given a similarity matrix Y for n objects, and are interested in seriating or stringing the data, by ordering the n objects so that similar objects i and j are near each other. Data seriation has diverse applications ranging from data visualization, DNA sequencing to functional data analysis [CCMW11] and archaeological dating [Rob51]. Most previous work on data seriation focuses on the noiseless case [Rob51, Ken71], where there is an unknown ordering of n objects so that if object j is closer than object k to object i in the ordering, then $Y_{ij} \geq Y_{ik}$, i.e., the similarity between i and j is always no less than the similarity between i and k. Such a matrix Y is called *Robinson matrix*. It is shown in [ABH98] that one can recover the underlying true ordering of objects up to a global shift by component-wisely sorting the second eigenvector of the Laplacian matrix associated with Y if Y is a Robinson matrix. The data seriation problem has also been formulated as a quadratic assignment problem and convex relaxations are derived [FJBd13, LW14] and references therein).

One interesting generalization of the hidden Hamiltonian cycle model is to extend the hidden structure from a cycle to k-regular graph for general $k \ge 2$, e.g., nearest-neighbor graphs, which can potentially better fit the genome assembly data. The underlying k-regular graph can represent the hidden geometric structure and the observed graph can be viewed as a realization of the Watts-Strogatz small-world graph [WS98] if the weight distribution is Bernoulli. Recent work [CLR17] has studied the problem of detecting and recovering the underlying k-regular graph under the small-world graph model and derived conditions for reliable detection and recovery; however, the information limit and the optimal algorithm remain open.

Finally, we mention that the Rényi divergence of order 1/2 also plays a key role in determining the exact recovery threshold for community detection under the stochastic block models [ABH16, MNS15, ZZ16, JL15, AS15].

4 Proof Techniques and a Simpler Result

The proof of the main result, Theorem 1, is quite involved. In this section, we will discuss the highlevel ideas and the difference with the conventional proof using dual certificates. As a warm-up we also prove a weaker version of the result on the 2-factor ILP. The proof of the full result is given in Section 5.

4.1 **Proof Techniques**

A standard technique for analyzing convex relaxations is the *dual certificate argument*, which amounts to constructing the dual variables so that the desired KKT conditions are satisfied for the primal variable corresponding to the ground truth. This type of argument has been widely used, for instance, for proving the optimality of SDP relaxations for community detection under stochastic block models [ABH16, HWX16a, Ban15, HWX16b, ABKK15, PW15, HWX16c]. However, for the F2F LP (6), we were only able to find explicit constructions of dual certificates that attain the optimal threshold within a factor of two. Instead, the proof of Theorem 1 is by means of a direct *primal argument*, which shows that with high probability no other vertices of the F2F polytope has a higher objective value than that of the ground truth. Nevertheless, it is still instructive to describe this dual construction before explaining the ideas of the primal proof.

To certify the optimality of x^* for F2F LP, it reduces to constructing a dual variable $u \in \mathbb{R}^n$ (corresponding to the degree constraints) such that for every edge (i, j),

$$u_i + u_j \le w_{ij}, \quad \text{if } x_{ij}^* = 1,$$
(13)

$$u_i + u_j \ge w_{ij}, \quad \text{if } x_{ij}^* = 0.$$
 (14)

A simple choice of u is

$$u_i = \frac{1}{2} \min_j \left\{ w_{ij} : x_{ij}^* = 1 \right\}.$$
 (15)

Then (13) is fulfilled automatically and (14) can be shown (see Appendix D) to hold with high probability provided that

$$\beta_n - \log n \to +\infty. \tag{16}$$

where β_n is the $\frac{1}{3}$ -Rényi divergence defined in (11). Since $\alpha_n/2 < \beta_n < \alpha_n$ by (12), this construction shows that F2F achieves the optimal recovery threshold by at most a multiplicative factor of 2. For specific distributions, this factor-of-two gap can be further improved, e.g., to $\frac{3}{2}$ for Gaussian weights, for which we have $P = N(\mu, 1)$ and Q = N(0, 1) and $\beta = \frac{1}{6}\mu^2 = \frac{2}{3}\alpha$. However, this certificate does not get us all the way to the information limit (9).

Departing from the usual dual certificate argument, our proof of the optimality of F2F relaxation relies on delicate primal analysis. In particular, we show that $\langle w, x-x^* \rangle < 0$ for any vertex (extremal point) of the F2F polytope $x \neq x^*$ with high probability. It is known that F2F polytope is not integral in the sense that some of its vertices is fractional. Fortunately, it turns out for any vertex x, its fractional entry x_e must be 1/2. Thanks to this *half-integrality* property, we can encode the difference $y \triangleq 2(x - x^*)$ as a bicolored multigraph G_y with a total weight $w(G_y) = 2\langle w, x - x^* \rangle$. Finally, we bound $w(G_y)$ via a divide-and-conquer argument by first decomposing G_y into an edge-disjoint union of graphs in a family with simpler structures and then proving that for every graph H in this family, its total weight w(H) is negative with high probability under condition (9). Our decomposition of G_y heavily exploits the fact that G_y is a balanced multigraph in the sense that every vertex has an equal number of incident red edges and blue edges, and the classical graph-theoretic result that every connected balanced multigraph has an Eulerian circuit with edges alternating in colors.

4.2 2-factor (2F) Integer Linear Programming Relaxation

The 2-factor (2F) Integer Linear Programming relaxation of the TSP is

$$\widehat{x}_{2F} = \arg \max_{x} \langle w, x \rangle$$
s.t. $x (\delta(v)) = 2$
 $x_e \in \{0, 1\}.$
(17)

The 2-factor ILP is the same as the F2F LP (6) except that the x_e 's have integrality constraints, and is therefore a tighter relaxation of the original TSP than F2F LP. As a warm-up for the optimality proof of F2F LP, we provide a much simpler proof, showing that the optimal solution of the 2F ILP coincides with the true cycle x^* with high probability, under the same condition that $\alpha_n - \log n \to +\infty$. We note that although it is not an LP, the 2F ILP can be solvable in $O(n^4)$ time using a variant of the blossom algorithm [Edm65b, Edm65a, LRT08]; however, finding an efficient and scalable implementation of the blossom algorithm can be challenging in practice [GH87].




Figure 3: The ground truth x^* is a cycle of length 6, x is a feasible solution to (17) corresponding to two disjoint triangles, and the graph G_y for $y = x - x^*$ is an alternating 4-cycle.

Let x denote the adjacency vector of a given 2-factor. To prove that x^* is the unique optimal solution to the 2F ILP, it suffices to show $\langle w, x - x^* \rangle < 0$ for the adjacency vector of any 2-factor

 $x \neq x^*$. To capture the difference between x and x^* , we define $y \in \{0, \pm 1\}^{\binom{n}{2}}$ by

$$y = x - x^*. \tag{18}$$

Define a simple graph G_y with bicolored edge whose adjacency matrix is |y| with isolated vertices removed and each edge is colored red if $y_e = -1$ and blue if $y_e = +1$ (see Fig. 3 for an example). Furthermore, for a given bicolored graph B, we define its weight as

$$w(B) \triangleq \sum_{\text{blue } e \in E(B)} w_e - \sum_{\text{red } e \in E(B)} w_e.$$

Then $w(G_y) = \langle w, x - x^* \rangle$.

A bicolored graph is *balanced* if for every vertex the number of red incident edges is equal to the number of blue incident edges. Since $x(\delta(v)) = 2$ and $x^*(\delta(v)) = 2$ for every vertex v, it follows that $y(\delta(v)) = 0$ and thus G_y is balanced. Define

$$\mathcal{B} = \{B : B \text{ is a simple, connected, and balanced bicolored graph}\}$$
$$\mathcal{B}^* = \{B \in \mathcal{B} : V(B) \subset [n], x_e^* = 1 \text{ for every red edge } e \in E(B)\},$$

where V(B) and E(B) denote the vertex set and edge set of B, respectively.

Let B_1, \ldots, B_m denote the connected components of G_y . Since each connected component of G_y is balanced, it follows that $B_i \in \mathcal{B}^*$ and

$$w(G_y) = \sum_{i=1}^m w(B_i).$$

Hence to show $w(G_y) < 0$ for all possible G_y , it reduces to proving that w(B) < 0 for all $B \in \mathcal{B}^*$.

Fix an even integer $\ell \geq 4$ and let

$$\mathcal{B}_{\ell}^* = \{ B \in \mathcal{B}^* : |E(B)| = \ell \}.$$

Fix any $B \in \mathcal{B}_{\ell}^*$. By the balancedness, B has $\ell/2$ red edges and $\ell/2$ blue edges. Hence,

$$w(B) \stackrel{d}{=} \sum_{i=1}^{\ell/2} Y_i - \sum_{i=1}^{\ell/2} X_i,$$

where X_i 's and Y_i 's are independent sequences of random variables such that X_i 's are i.i.d. copies of dP/dQ under distribution P and Y_i 's are i.i.d. copies of dP/dQ under distribution Q; the notation $\stackrel{d}{=}$ denotes equality in distribution. It follows from the Chernoff's inequality (cf. the large-deviation bound (44) in Appendix B) that

$$\mathbb{P}\left\{w(B) \ge 0\right\} = \mathbb{P}\left\{\sum_{i=1}^{\ell/2} Y_i - \sum_{i=1}^{\ell/2} X_i \ge 0\right\} \le \exp\left(-\alpha_n \ell/2\right).$$
(19)

Next we claim that there are at most $(2n)^{\ell/2}$ different graphs B in \mathcal{B}^*_{ℓ} . The proof of the claim is deferred to the end of this section. Combining the union bound with (19) gives that

$$\mathbb{P}\left\{\max_{B\in\mathcal{B}_{\ell}^{*}}w(B)\geq 0\right\}\leq |\mathcal{B}_{\ell}^{*}|\exp\left(-\alpha_{n}\ell/2\right)\leq \exp\left\{-\left(\alpha_{n}-\log(2n)\right)\ell/2\right\}$$

Taking another union bound over all integers $\ell \geq 4$, we get the desired result.

$$\mathbb{P}\left\{\max_{B\in\mathcal{B}^*} w(B) \ge 0\right\} \le \sum_{\ell=4}^{\infty} \mathbb{P}\left\{\max_{B\in\mathcal{B}^*_{\ell}} w(B) \ge 0\right\}$$
$$\le \sum_{\ell=4}^{\infty} \exp\left\{-\left(\alpha_n - \log(2n)\right)\ell/2\right\}$$
$$\le \frac{\exp\left\{-2\left(\alpha_n - \log(2n)\right)\right\}}{1 - \exp\left\{-\left(\alpha_n - \log(2n)\right)/2\right\}} \stackrel{(9)}{\to} 0.$$

We are left to show $|\mathcal{B}_{\ell}^*| \leq (2n)^{\ell/2}$. This follows from the following classical graph-theoretic result that every connected balanced multigraph G has an *alternating* Eulerian circuit, that is, the edges in the circuit alternate in colors.

Lemma 1. Every connected balanced bicolored multigraph G has an alternating Eulerian circuit.

The lemma is proved in [Kot68, Theorem 1] in a more general form (see also [Pev95, Corollary 1]). For completeness, we provide a short proof in Appendix A.

In view of Lemma 1, for every $B \in \mathcal{B}_{\ell}^*$, it must have a Eulerian circuit T given by the sequence of $(v_0, v_1, \ldots, v_{\ell-1}, v_{\ell} = v_0)$ of vertices (vertices may repeat) such that $v_i \in [n]$ and (v_i, v_{i+1}) is a red edge for even i and blue edge for odd i in B. Let \mathcal{T} denote the set of all possible such Eulerian circuits. Moreover, every Eulerian circuit $T \in \mathcal{T}$ uniquely determines a $B \in \mathcal{B}_{\ell}^*$, because the vertex set V(B) is the union of vertices v_i 's and the colored edge set E(B) is the union of colored edges (v_i, v_{i+1}) 's in T. Hence, $|\mathcal{B}_{\ell}^*| \leq |\mathcal{T}|$. To enumerate all possible $T \in \mathcal{T}$, it suffices to enumerate all the possible labelings of vertices in T. Recall that by definition, for every red edge $e = (v_i, v_{i+1})$ in T, $x_e^* = 1$. Thus the two endpoints v_i and v_{i+1} must be neighbors in the true cycle corresponding to x^* , and hence once the vertex labeling of v_i is fixed, there are at most 2 different choices for the vertex labeling of v_{i+1} . Therefore, we enumerate all possible Eulerian circuits $T \in \mathcal{T}$ by sequentially choose the vertex labeling of v_i from i = 0 to $i = \ell - 1$. Given the vertex labelings of (v_0, \ldots, v_{i-1}) , the number of choices of the vertex labeling of v_i is at most n for even i and 2 for odd i. Hence, $|\mathcal{T}| \leq (2n)^{\ell/2}$, which further implies that $|\mathcal{B}_{\ell}^*| \leq (2n)^{\ell/2}$.

5 Proof of Theorem 1

In this section, we prove that the optimal solution of the fractional 2-factor coincides with x^* with high probability, provided that $\alpha_n - \log n \to \infty$. This is the bulk of the paper.

5.1 Graph Notations

We describe several key graph-theoretic notations used in the proof. We start with multigraphs. Formally, a multigraph G is an ordered pair (V, E) with a vertex set V = V(G) and an edge *multiset* E = E(G) consisting of subsets of V(G) of size 2. Note that by definition multigraphs do not have self-loops. A multi-edge is a set of edges in E(G) with the same end points. The multiplicity of an edge is its multiplicity as an element in E(G). We call a multi-edge single and double if its edge multiplicity is 1 and 2, respectively. Note that a double edge (u, v) refers to the set of two edges connecting vertices u and v. We say a multigraph G is bicolored, if every distinct element in E(G)is colored in either red or blue and the repeated copies of an element all have the same color.

For two multigraphs G and H on the same set of vertices, we define G - H to be the multigraph induced by the edge multiset $E(G) \setminus E(H)$. The union of multigraphs G and H is the multigraph $G \cup H$ with vertex set $V(G) \cup V(H)$ and edge multiset $E(G) \cup E(H)$.⁶ By definition, the multiplicity of an element in $E(G) \cup E(H)$ is the sum of its multiplicity in E(G) and E(H). When $E(G) \cap E(H) = \emptyset$, $G \cup H$ is called an edge-disjoint union. When $V(G) \cap V(H) = \emptyset$, $G \cup H$ is called an vertex-disjoint union.

A walk in a multigraph G is a sequence (v_0, v_1, \ldots, v_m) of vertices (which may repeat) such that $(v_{i-1}, v_i) \in E(G)$ for $1 \leq i \leq m$. A trail in a multigraph G is a walk (v_0, v_1, \ldots, v_m) such that for all $1 \leq i \leq m$, the number of times that edge (v_{i-1}, v_i) appears in the walk is no more than its edge multiplicity in E(G). A trail is closed if the starting and ending vertices are the same. A circuit is a closed trail. An Eulerian trail in a multigraph G is a trail (v_0, v_1, \ldots, v_m) such that for every $e \in E(G)$, the number of times that it appears in the trail coincides with its edge multiplicity in E(G). An Eulerian circuit is a closed Eulerian trail. A path is a trail with no repeated vertex. A cycle consists a path plus an edge from its last vertex to the first.

5.2 Proof Outline

Let $x^* = (x_e^*)$ denote the adjacency vector of the hidden Hamiltonian cycle (ground truth). The feasible set of the F2F LP (6) is the F2F polytope:

$$Q \triangleq \left\{ x \in [0,1]^{\binom{n}{2}} : x(\delta(v)) = 2, \forall v \in [n] \right\}.$$
(20)

To prove that x^* is the unique optimal solution to the F2F LP with high probability, it suffices to show that $\langle w, x - x^* \rangle < 0$ holds with high probability for any vertex (extremal point) of the F2F polytope x other than x^* . It turns out that the vertices of the F2F polytope Q has the following simple characterization [Bal65, BC99, SWvZ13]. First of all, for any vertex x, its fractional entry must be a half-integer, i.e.,

$$x_e \in \{0, 1/2, 1\}, \quad \forall e.$$
 (21)

Furthermore, if we define the support graph of x as be the graph with vertex set [n] and edge set $\{e : x_e \neq 0\}$, then each connected component of the support graph of x must be one of the following two cases: it is either

- 1. a cycle of at least three vertices with $x_e = 1$ for all edges e in the cycle,
- 2. or consists of an even number of odd-sized cycles with $x_e = 1/2$ for all edges e in the cycles that are connected by paths of edges e with $x_e = 1$. In this case, if we remove the edges in the odd cycles, the resulting graph is a spanning disjoint set of paths formed by edges e with $x_e = 1$.

It turns out that, among the aforementioned characterizations of the vertices of the F2F polytope, our analysis of the LP relaxation uses only the half-integrality property (21).

⁶Here the union of multisets is defined so that the multiplicity of each elements adds up. For example, $\{a, a, b\} \cup \{a, b, c\} = \{a, a, a, b, b, c\}$.




Figure 4: (a): The support graph of a fractional vertex x of the F2F polytope with n = 6. The edges in the support graph of x^* are highlighted in red. (b): The multigraph representation of $y = 2(x - x^*)$.

To capture the difference between a given vertex x and the true solution x^* , we use the following multigraph representation: define $y \in \mathbb{R}^{\binom{n}{2}}$ by

$$y = 2(x - x^*),$$
 (22)

with $y_e = 2 (x_e - x_e^*) \in \{0, \pm 1, \pm 2\}$. Define a multi-graph G_y whose adjacency matrix is |y| with isolated vertices removed and each edge e is colored red if $y_e < 0$ or blue if $y_e > 0$. In particular, the edge multiplicity of G_y is at most 2. Comparing to (18), the extra factor of 2 in (22) is to ensure that y is still integral; as a consequence, G_y may be a multigraph with multiplicity 2 instead of a simple graph. For any given bicolored multigraph F, we define its weight as

$$w(F) \triangleq \sum_{\text{blue } e \in E(F)} w_e - \sum_{\text{red } e \in E(F)} w_e ,$$

where the summation above includes all repeated copies of e in E(F). Then $w(G_y) = \langle w, x - x^* \rangle$. Hence, to prove that x^* is the unique optimal solution to the LP program with high probability, it reduces to showing that $w(G_y) < 0$ for all possible G_y constructed from the extremal point $x \neq x^*$ with high probability.

Instead of first calculating the probability of $w(G_y) \leq 0$ and then taking a union bound on all possible G_y , our proof crucially relies on a decomposition of G_y into some suitably defined simpler graphs. In the next subsection, we will describe a family \mathcal{F}^* of graphs, and show that every possible G_y can be decomposed as a union of graphs in \mathcal{F}^* : $G_y = \bigcup_{i=1}^m F_i$ with $F_i \in \mathcal{F}^*$ for each $1 \leq i \leq m$. Since the multiplicity of e in $E(G_y)$ is equal to the sum of multiplicities of e in $E(F_i)$ over $i \in [m]$, it follows that

$$w(G_y) = \sum_{i=1}^m w(F_i).$$

Therefore, to show that $w(G_y) < 0$ for all possible G_y with high probability, it suffices to show w(F) < 0 for all graphs F in family \mathcal{F}^* .

We remark that in the analysis of the 2F ILP in Section 4.2, we have $y = x - x^*$ as opposed to $y = 2(x - x^*)$ and thus G_y is a balanced simple graph. Consequently, we can simply decompose G_y into its connected components which are connected, balanced simple graphs. In contrast, here the decomposition of G_y as a multigraph is much more sophisticated due to the existence of double edges. In particular, the weight of a double edge in F appears twice in w(F) and hence its variance is twice the total variance of two independent edge weight. For this reason, to control the deviation

of w(F) from its mean, it is essential to account for the contribution of double edges and single edges separately, which, in turn, requires us to separate the double edges from single edges in our decomposition.

5.3 Edge Decomposition

Our decomposition of G_y relies on the notion of balanced multigraph and alternating Eulerian circuit. A bicolored multi-graph is *balanced* if for every vertex the number of red incident edges is equal to the number of blue incident edges. Since $x(\delta(v)) = 2$ and $x^*(\delta(v)) = 2$ for every vertex v, it follows that $y(\delta(v)) = 0$ and thus G_y is balanced. As a result, the vertices in G_y all have even degree (in fact, either 2, 4, 6 or 8). Therefore each connected component of G_y has an Eulerian circuit. Recall that Eulerian circuit is alternating if the edges in the Eulerian circuit alternate in colors. In view of Lemma 1, each connected component of G_y has an Eulerian circuit. In the remainder, we suppress the subscript y in G_y whenever the context is clear.

Next, we describe a family \mathcal{F} of graphs and show that G is a union of graphs in this family. First, we need to introduce a few notations. For any pair of two vertices u, v in graph G, vertex identification (also known as vertex contraction) produces a graph by removing all edges between u, v and replacing u, v with a single vertex w incident to all edges formerly incident to either u or v. When u and v are adjacent, i.e., sharing two endpoints of edge e, vertex identification specializes to edge contraction of e and the resulting graph is denoted by $G \cdot e$; visually, e shrinks to a vertex. Note that edge contraction may introduce multi-edges. We define a stem as a path $(v_0, v_1, \ldots, v_{k-1})$ for some k distinct vertices such that (v_{i-1}, v_i) is a double edge for all $1 \le i \le k$ and the double edges alternate in colors. The two endpoints v_0 and v_{k-1} of the stem are identified as the tips of the stem. We say a tip of the stem is red if it is incident to the red double edge; otherwise we say it is blue. Given a stem and an even cycle C_0 consisting of only single edges of alternating colors, we define the following blossoming procedure to connect the stem with C_0 : first contract any single blue (red) edge in C_0 to a vertex v and attached to v the stem by identifying v with a blue (red) tip of the stem. The resulting graph known as *flower* has an alternating circuit and the contracted C_0 is called *blossom*. The tip of the stem not incident to the blossom is called the tip of the flower. We say a flower is red (blue) if its tip is red (blue). For example, a red flower are shown Fig. 5. Similar notions of stem, flower, blossom were introduced in [Edm65b] in the context of simple graphs.




Figure 5: A red flower consisting of a stem of 4 alternating double edges followed by a blossom of 5 single edges.

Then we introduce a family \mathcal{U} of balanced graphs. We start with an even cycle G_0 in alternating colors. At each step $t \geq 1$, construct a new balanced graph G_t from G_{t-1} as follows. Fix any cycle consisting of at least 4 edges in G_{t-1} . In this cycle, pick any edge and apply the following *flowering* procedure: contract the red (blue) edge to one vertex w and attach to w a flower by identifying w with the root of a blue (red) flower.⁷ Let \mathcal{U} denote the collection of all graphs obtained from applying the flowering procedure recursively for finitely many times. In particular, \mathcal{U} includes all

⁷Since we allow contracting an edge incident to a stem, it is possible to have a vertex with multiple stems attached.

even cycles in alternating colors. For example, the graph in Fig. 6 is in \mathcal{U} which is obtained by starting with a 10-cycle and applying the flowering procedure 4 times. By construction, any graph $H \in \mathcal{U}$ must contain even $\ell \geq 4$ number of single edges.




Figure 6: Example of graph in family \mathcal{U}




Figure 7: Tree representation of graph in Fig. 6.

Alternatively, note that each graph in the family \mathcal{U} can be viewed as cycles interconnected by stems. Thus, we can represent G_t using a tree T_t , whose nodes correspond to even cycles and links corresponds to stems. Next we describe this enumeration scheme in details. Every node in the tree represents a cycle in alternating colors, with a mark ℓ being the length of the cycle. The cycle corresponding to the root node is assumed to have a fixed ordering of edges, and the root node has an extra mark which is 1 if the color of the first edge in the corresponding cycle is blue and 0 otherwise. Every link (u, v) represents a stem consisting of only double edges of alternating colors with mark (k, i), where k is the length of the stem, and i is the index of the contracted edge of the parent vertex u. We start with tree T_0 with a single root node corresponding to $G_0 = C_0$ with the mark being the length of C_0 . At each step $t \geq 1$, we view the flowering procedure as growing to a new tree T_t as follows. For any vertex u in T_{t-1} that corresponds to an alternating cycle C, a new vertex v that corresponds to an alternating cycle C', and a stem, we connect u and v with a edge corresponding to the stem. The edge is marked with (k, i), where k is the length of the stem, and i is the index of the contracted edge in C. Then the edges in the alternating cycle C' is indexed by $1, 2, \ldots$ by starting from the contracted edge in C' and traversing C' in clockwise direction.

Finally, we need to introduce the notion of homomorphism between two bicolored multigraphs H and F. There exist multiple definitions of homomorphism between multigraphs; here we follow the convention in [Lov12, Section 5.2.1]. A node-and-edge homomorphism $H \to F$ is a pair of vertex map $\phi: V(H) \to V(F)$ and bijective⁸ edge map $\psi: E(H) \to E(F)$ such that if $e \in E(H)$

⁸Let A and B denote two multisets. Let A' and B' denote the set of distinct elements in A and B, respectively. We say $\psi: A \to B$ is bijective if $\psi: A' \to B'$ is bijective and for every element $a \in A'$, the multiplicity of a in A is the

connects *i* and *j*, then $\psi(e)$ connects $\phi(i)$ and $\phi(j)$ and has the same color as *e*. We say *H* is homomorphic to *F* if such a node-to-edge homomorphism exists. By construction, an edge *e* is incident to *u* in *H* if and only if $\psi(e)$ is incident to $\phi(u)$ in *F*. Therefore, if *H* is homomorphic to *F*, then they are either both balanced or both unbalanced. Moreover, since ψ is bijective, $H \to F$ is edge-multiplicity preserving, i.e., the multiplicity of $\psi(e)$ in E(F) is the same as the multiplicity of *e* in E(H). Hence, the number of double (single) edges in *H* and *F* are the same. Furthermore, note that for two node-and-edge homomorphisms $(\phi, \psi) : H \to F$ and $(\phi, \psi') : H \to F'$ with the same vertex map ϕ , it holds that F = F'. Hence, when the context is clear, we simply write $\phi : H \to F$ or $\phi(H) = F$ by suppressing the underlying edge map.

Let \mathcal{F} denote the collection of all graphs F such that $H \to F$ for some $H \in \mathcal{U}$. In particular, $\mathcal{F} \supseteq \mathcal{U}$, and this inclusion is strict as the example in Fig. 8 shows. The next lemma shows that \mathcal{F}




Figure 8: Example of a graph F in \mathcal{F} but not in \mathcal{U} . Here F is homomorphic to $H \in \mathcal{U}$, where the homomorphism ϕ maps both vertices 11 and 12 to 11.

includes all connected balanced simple graphs. This result serves as the base case of the induction proof of the decomposition lemma.

Lemma 2. An alternating cycle is homomorphic to any connected balanced simple graph G with equal number of edges. In particular, $G \in \mathcal{F}$.

Proof. By Lemma 1, G has an Eulerian circuit $T = (v_0, v_1, \ldots, v_{m-1}, v_m = v_0)$ of alternating colors where m is the total number of edges in G and vertices v_i 's may repeat. Let C denote any alternating cycle with m edges. We write $C = (u_0, u_1, \ldots, u_{m-1}, u_m = u_0)$ such that the edge (u_0, u_1) has the same color as (v_0, v_1) . Then we define a pair of vertex and edge map (ϕ, ψ) from C to G such that $\phi(u_i) = v_i$ and $\psi((u_i, u_{i+1})) = (v_i, v_{i+1})$ for all $0 \le i \le m-1$. Since both G and C are simple graphs, $\psi : E(C) \to E(G)$ is bijective. Hence, $(\phi, \psi) : C \to G$ is a node-and-edge homomorphism and the conclusion follows.

In contrast to connected balanced simple graphs, if a balanced graph G contains double edges then certainly no alternating cycle is homomorphic to G. What's more, it is possible that G is not homomorphic to any graph in the class \mathcal{U} , i.e., G may not belong to \mathcal{F} . See Fig. 9 for such an example. Nevertheless, the next lemma shows that if G has edge multiplicity at most 2, then it can be decomposed as a *union of elements* in \mathcal{F} .

Lemma 3 (Decomposition). Every balanced multigraph G with edge multiplicity at most 2 can be decomposed as a union of elements in \mathcal{F} .

same as the multiplicity of $\psi(a)$ in B. For example, if $A = \{a, a, b, c\}$ and $B = \{x, x, y, z\}$. Let $\psi(a) = x$, $\psi(b) = y$, and $\psi(c) = z$, then $\psi : A \to B$ is bijective.




Figure 9: Example of a graph F which is not in \mathcal{F} , but can be decomposed as $F = F_1 \cup F_2$ with $F_1, F_2 \in \mathcal{F}$.

Proof. It suffices to prove the lemma for the case where G is connected. By Lemma 1, G has an Eulerian circuit $T = (v_0, v_1, \ldots, v_{m-1}, v_m = v_0)$ of alternating colors where m is the total number of edges and vertices v_i may repeat. We proceed by induction on the number of double edges k in G. If k = 0, G is simple. By Lemma 2, $G \in \mathcal{F}$ and thus the conclusion holds.

Suppose the conclusion holds for $k \ge 0$, we aim to prove it also holds for k+1. We call a double edge (u, v) bidirectional if the Eulerian circuit T traverses it twice in two different directions, i.e., there exists an s and $t \ge s+3$ such that $(v_{s-1}, v_s) = (u, v)$ and $(v_{t-1}, v_t) = (v, u)$. Similarly, we call a double edge (u, v) unidirectional if T traverses it twice in the same direction. We divide our induction into two cases according to whether there exists a unidirectional double edge.

Case 1: There exists at least one unidirectional double edge. Let (u, v) be an arbitrary unidirectional double edge. See Fig. 9 for an illustration. By definition, in the alternating Eulerian circuit $T = (v_0, v_1, \ldots, v_{m-1}, v_0)$, there must exist an s and $t \ge s + 3$, such that $(v_{s-1}, v_s) = (v_{t-1}, v_t) = (u, v)$. Since T is alternating, it follows that $T' \triangleq (v_{s-1}, v_s, \ldots, v_{t-1})$ is also an alternating circuit. Also define T - T' to be the resulting graph by deleting the edges in the circuit T' from T, i.e.,

$$T - T' = (v_0, v_1, v_{s-1}, v_t, \dots, v_{m-1}, v_0).$$

It follows that T - T' is also a circuit in alternating colors. Hence both T' and T - T' are balanced multigraphs with edge multiplicity at most 2. Also, since the two edges connecting u and v appear separately in T' and T - T', both T' and T - T' have at most k double edges. Applying the induction hypothesis to each connected component of T' and T - T', we conclude that both T' and T - T' can be decomposed as a union of elements in \mathcal{F} . Hence, T can be decomposed as a union of elements in \mathcal{F} .

Case 2: All double edges are bidirectional. In this case, we pick an arbitrary bidirectional double edge (u, v). In the Eulerian circuit T, we find a trail S of maximal length that contains the double edge (u, v) and consists of only double edges in alternating colors (see Fig. 10). More precisely, we find the smallest s and the largest $t \ge s+1$ such that (v_{i-1}, v_i) all have edge multiplicity 2 for $s \le i \le t$, and $(v_{i-1}, v_i) = (u, v)$ for some $s \le i \le t$, and there exists $t' \ge t+3$ and $s' \ge t'$ such that

$$(v_{t'}, v_{t'+1}, \dots, v_{s'-1}) = (v_t, v_{t-1}, \dots, v_{s-1}),$$

in which the Eulerian circuit T traverses S in two different directions.

By the maximality of S, v_{t+1} and $v_{t'-1}$ must be two distinct vertices. Since the Eulerian circuit T has alternating colors, (v_t, v_{t+1}) and $(v_{t'-1}, v_{t'})$ must have the same color, and thus the circuit

$$R = (v_t, v_{t+1}, \dots, v_{t'} = v_t)$$

must have an odd length. Entirely analogously, v_{s-2} and $v_{s'}$ must be two distinct vertices, and (v_{s-2}, v_{s-1}) and $(v_{s'-1}, v_{s'})$ must be of the same color; thus the circuit

$$L = (v_0, \dots, v_{s-1}, v_{s'}, \dots, v_{m-1}, v_m = v_0)$$

must have an odd length. In particular, both L and R cannot be empty. Therefore we have $T = L \cup S \cup R$ as shown in Fig. 10. We further consider two subcases according to whether L and R share any edge or not.




Figure 10: The decomposition $T = L \cup S \cup R$ where L and R are circuits of alternating colors shown in dashed lines

Case 2.1: $E(L) \cap E(R) \neq \emptyset$. In this case, there exists a bidirectional double edge whose two simple edges appear separately in E(L) and E(R). In particular, there exist $p \in \{t+1,\ldots,t'\}$ and $q \in \{1,\ldots,s-1\} \cup \{s',\ldots,m\}$ such that $(v_{p-1},v_p) = (v_q,v_{q-1})$.

Define a circuit $T' = (v_{s-1}, \ldots, v_{p-1}, v_{q-1}, v_{q-2}, \ldots, v_{s'-1})$ by first traversing S to R until reaching $v_{p-1} = v_q$ and then traversing L in the reverse direction until reaching $v_{s'-1} = v_{s-1}$. Note that T' is an alternating circuit. It follows that both T' and T - T' are balanced multigraphs with edge multiplicity at most 2. Moreover, for each double edge in S, its two simple edges appear separately in T' and T - T'. Thus, both T and T - T' have at most k double edges. Applying the induction hypothesis to each connected component of T' and T - T', we conclude that both T' and T - T' can be decomposed as a union of elements in \mathcal{F} . Hence, T can be decomposed as a union of elements in \mathcal{F} .

Case 2.2: $E(L) \cap E(R) = \emptyset$. In this case, $T = L \cup S \cup R$ is an edge-disjoint union.

<u>Step 1</u>. First we argue that, without loss of generality, we can and will assume that (v_t, v_{t+1}) , $(v_{t'-1}, v_{t'})$, (v_{s-2}, v_{s-1}) , and $(v_{s'-1}, v_{s'})$ in Fig. 10 are all single edges. Below we only consider (v_t, v_{t+1}) . Suppose (v_t, v_{t+1}) is not a single edge. Then by assumption, it must be a bi-directional double edge. Thus there exists a $t'' \in (t, t')$ such that $(v_{t''-1}, v_{t''}) = (v_{t+1}, v_t)$. Thus $v_t = v_{t'} = v_{t''}$. Let $T' = (v_0, \ldots, v_{t''}, v_{t'+1}, \ldots, v_0)$ and $T - T' = (v_{t''}, \ldots, v_{t'})$ (cf. Fig. 11). Then T' is balanced and thus T - T' is also balanced. Moreover, T - T' has at most k double edges. Applying the induction hypothesis to each connected component of T - T', we conclude that T - T' can be decomposed as a union of elements in \mathcal{F} . Then it remains to show T' can be decomposed as a union of elements in \mathcal{F} . Then it remains to show T' can be decomposed as a union of elements in \mathcal{F} . Bedefine S by including the double edge (v_t, v_{t+1}) and redefine v_t and R accordingly. Applying the above procedure in finitely many times ensures that (v_t, v_{t+1}) becomes a single edge.

Step 2. Next we capitalize on the fact that $T = L \cup S \cup R$ is an edge-disjoint union to complete the proof of the decomposition.




Figure 11: Example of the decomposition in Step 1 in Case 2.2. The highlighted circuit is T - T', which needs to be decomposed so that the double edge (v_t, v_{t+1}) can be included in the stem S.

In R, we expand the vertex v_t to an edge $e = (u_t, v_t)$ whose color is different from the color of edge (v_t, v_{t+1}) . In particular, we add a distinct vertex u_t and edge e, and reconnect $v_{t'-1}$ from $v_{t'}$ to u_t . Let R' be the resulting circuit, i.e., $R' = (u_t, v_t, v_{t+1}, \ldots, v_{t'-1}, u_t)$. Then R' is a circuit of alternating colors. Hence R' is a balanced multigraph with edge multiplicity at most 2. Moreover, R' has at most k double edges. Applying the induction hypothesis on each connected component of R', we conclude that R' can be decomposed as a union of elements in \mathcal{F} . In this union, denote by P the element in \mathcal{F} that contains the edge e. Since u_t is only incident to v_t and $v_{t'-1}$, it follows that P must also contain the edge $(v_{t'-1}, u_t)$. Furthermore, since $(v_{t'-1}, v_{t'})$ is a single edge in R, it follows that $v_{t'-1}$ and v_t are disconnected in P and thus e is contained in a cycle of length at least 4 in P. Let $P \cdot e$ be the resulting graph by contracting e in P to vertex v_t .

Analogously, in L, we expand the vertex v_{s-1} to an edge $e' = (u_{s-1}, v_{s-1})$ whose color is different from the color of edge (v_{s-2}, v_{s-1}) . Let L' be the resulting circuit, i.e.,

$$L' = (v_0, \dots, v_{s-2}, u_{s-1}, v_{s-1}, v_{s'}, \dots, v_{m-1}, v_0).$$

Then L' is a circuit of alternating colors. By the same argument as in the case of R', we get that L' can be decomposed as a union of elements in \mathcal{F} . In this union, denote by Q the element in \mathcal{F} that contains the edge e'. Then e' is contained in a cycle of length at least 4 in Q. Let $Q \cdot e'$ be the resulting graph by contracting e' in Q to vertex v_{s-1} .

To show T is an union of elements in \mathcal{F} , it suffices to show that $M \triangleq (Q \cdot e') \cup S \cup (P \cdot e) \in \mathcal{F}$. Note that vertices may repeat in S. Nevertheless, there exists a homomorphism (ϕ_S, ψ_S) such that $S_0 \to S$ for some stem S_0 . Let w' denote an endpoint of S_0 such that $\phi_S(w') = v_{s-1}$ and w denote the other endpoint of S_0 such that $\phi_S(w) = v_t$. Moreover, since $P \in \mathcal{F}$, it follows that there exists a homomorphism (ϕ_P, ψ_P) such that $P_0 \to P$ for some $P_0 \in \mathcal{U}$. Similarly, there exists a homomorphism (ϕ_Q, ψ_Q) such that $Q_0 \to Q$ for some $Q_0 \in \mathcal{U}$. Note that e (resp. e') is a simple edge in P (resp. Q). Therefore, there exist simple edge $e_0 \in E(P_0)$ and $e'_0 \in E(Q_0)$) such that $\psi_P(e_0) = e$ and $\psi_Q(e'_0) = e'$. By the tree representation of graphs in \mathcal{U} , P_0 can be represented as a tree I with vertex i corresponding to the cycle C_i that contains the edge e_0 . Entirely analogously, Q_0 can be represented as a tree J with vertex j corresponding to the cycle C_j that contains the edge e'_0 . We form a new tree K by connecting vertex i and vertex j with an edge corresponding to stem S_0 and identifying the root vertex of I as the root vertex of K. In other words, let $Q_0 \cdot e'_0$ denote the graph obtained by contracting e'_0 in Q_0 to vertex w', and $P_0 \cdot e_0$ denote the graph obtained by contracting e_0 in P_0 to vertex w. Note that e_0 (resp. e'_0) is contained in a cycle of length at least 4 in P_0 (resp. Q_0). Let $M_0 \triangleq (Q_0 \cdot e'_0) \cup S_0 \cup (P_0 \cdot e_0)$. Then K is a tree representation of M_0 starting from a cycle represented by the root vertex of tree I and thus $M_0 \in \mathcal{U}$.

Finally, it remains to show that M_0 is homomorphic to M. Since the vertex sets $V(P_0)$, $V(Q_0)$, and $V(S_0)$ are disjoint, we can define a vertex map $\phi : V(M_0) \to V(M)$ as follows: for any vertex $v \in V(M_0),$

$$\phi(v) = \begin{cases} \phi_P(v) & \text{if } v \in V(P_0) \\ \phi_Q(v) & \text{if } v \in V(Q_0) \\ \phi_S(v) & \text{if } v \in V(S_0). \end{cases}$$

Moreover, since $M_0 = (Q_0 \cdot e'_0) \cup S_0 \cup (P_0 \cdot e_0)$ is an edge-disjoint union, we can define an edge map $\psi : E(M_0) \to E(M)$ as follows: for any edge $e \in E(M_0)$,

$$\psi(e) = \begin{cases} \psi_P(e) & \text{if } e \in E\left(P_0 \cdot e_0\right) \\ \psi_Q(e) & \text{if } e \in E\left(Q_0 \cdot e_0'\right) \\ \psi_S(e) & \text{if } e \in E(S_0). \end{cases}$$

By assumption, $E(L) \cap E(R) = \emptyset$. Therefore, $(Q \cdot e') \cup S \cup (P \cdot e)$ is also an edge-disjoint union. As a consequence, $\psi : E(M_0) \to E(M)$ is bijective, and for any $e \in E(M_0)$, the multiplicity of $\psi(e)$ is the same as the multiplicity of e. Thus $(\phi, \psi) : M_0 \to M$ is a homomorphism. Hence, $M \in \mathcal{F}$, concluding the proof.

For $k \geq 0$ and $\ell \geq 3$, we define $\mathcal{U}_{k,\ell} \subset \mathcal{U}$ as the bicolored balanced multigraphs $H \in \mathcal{U}$ with k double edges and ℓ single edges. The following lemma upper bounds the number of unlabeled graphs in $\mathcal{U}_{k,\ell}$.

Lemma 4 (Enumeration of isomorphism classes). Let $k \ge 0$ and even $\ell \ge 4$. Then the number of unlabeled graphs in $\mathcal{U}_{k,\ell}$ is at most $17^k 4^{\ell}$.

Proof. If k = 0, then by Lemma 2, all graphs in $\mathcal{U}_{0,\ell}$ are isomorphic to an alternating ℓ -cycle. Thus the number of unlabeled graphs in $\mathcal{U}_{0,\ell}$ is 1. Hence the conclusion trivially holds. Next we focus on the case where $k \geq 1$. Note that if two unlabeled multigraphs in \mathcal{U} have the same tree representation, they must be the same. Hence it suffices to upper bound the number of possible tree representations of multigraphs $H \in \mathcal{U}_{k,\ell}$.

Fix a marked tree T of m vertices that is a tree representation of a multigraph $H \in \mathcal{U}_{k,\ell}$. Since $k \geq 1$, it follows that $2 \leq m \leq k + 1$. Starting with the root node, we order the nodes and links in T via breadth first search. For $i \in [m]$, let ℓ_i denote the length of the cycle represented by node i and S_i denote the set of indices of contracted edges in the cycle. For $j \in [m-1]$, let k_j the length of the stem represented by link j. Let c denote the extra mark of the root vertex. Note that from $\{c, (\ell_i, S_i)_{i=1}^m, (k_j)_{j=1}^m\}$, we can uniquely determine the marked tree T. In particular, ℓ_i determines the mark of vertex i and k_j determines the first mark of edge j. Moreover, S_i determines the number of children of node i and the second mark on every link connecting i to its child. Therefore, to bound the number of all possible marked trees T, it suffices to bound the number of all possible $\{c, (\ell_i, S_i)_{i=1}^m, (k_j)_{i=1}^{m-1}\}$. Note that

$$\sum_{i=1}^{m} \ell_i = \ell + 2(m-1), \quad \sum_{j=1}^{m-1} k_j = k.$$

Hence, the number of possible choices of sequences (ℓ_1, \ldots, ℓ_m) is at most $\binom{\ell+2m-3}{m-1}$, and the number of possible choices of sequences (k_1, \ldots, k_{m-1}) is at most $\binom{k-1}{m-2}$. Moreover, for each *i*, there are at most 2^{ℓ_i} different choices of S_i . Hence, the number of all possible $\{c, (\ell_i, S_i)_{i=1}^m, (k_j)_{j=1}^m\}$ is at

most $2^{\ell+2m-1} \binom{\ell+2m-1}{m-1} \binom{k-1}{m-2}$. Therefore, the number of possible tree representations of multigraphs $H \in \mathcal{U}_{k,\ell}$ is at most

$$\sum_{m=2}^{k+1} 2^{\ell+2m-1} \binom{\ell+2m-3}{m-1} \binom{k-1}{m-2} \le \sum_{m=2}^{k+1} 2^{\ell+2m-1} 2^{\ell+2m-3} \binom{k-1}{m-2} = 2^{2\ell+4} \sum_{m=2}^{k+1} 2^{4(m-2)} \binom{k-1}{m-2} \le 2^{2\ell+4} 17^{k-1} \le 4^{\ell} 17^k.$$

Define⁹

 $\mathcal{F}^* = \{ F \in \mathcal{F} : V(F) \subset [n] \text{ and } |E(F)| \le 4n \text{ and for every red edge } e \in E(F), \, x_e^* = 1 \}.$

Given $H \in \mathcal{U}$, we say a homomorphism $\phi : H \to F$ is *compatible with* x^* if $\phi(H) \in \mathcal{F}^*$. Denote by Φ_H^* the set of all homomorphism $\phi : H \to F$ that is compatible with x^* . Then

$$\mathcal{F}^* = \{\phi(H) : \phi \in \Phi^*_H, H \in \mathcal{U}\}.$$

In the following, we upper bound the number of elements in Φ_H^* for a given $H \in \mathcal{U}$. We need to set up a few notations. Let H_d and H_s denote the subgraph of H induced by all the double edges and all the single edges, respectively. Then we have an edge-disjoint union $H = H_d \cup H_s$. For a vertex map $\phi \in \Phi_H^*$, let ϕ_d and ϕ_s denote ϕ restricted to $V(H_d)$ and $V(H_s)$, respectively. Note that $\phi_d(v) = \phi_s(v)$ for all $v \in V(H_d) \cap V(H_s)$. We write $\phi = (\phi_s, \phi_d)$.

Lemma 5 (Enumeration of homomorphisms). Let $k \ge 0$ and let $\ell \ge 4$ be an even integer. Fix a bicolored balanced multigraph $H \in \mathcal{U}_{k,\ell}$.

• There exists an integer $0 \le r \le \ell/2$ such that

$$\log |\Phi_{H_d}^*| \le \frac{1}{2}(k+r)\log(2n), \tag{23}$$

where

$$\Phi_{H_d}^* \triangleq \{\phi_d : \exists \phi_s, \ s.t. \ (\phi_d, \phi_s) \in \Phi_H^*\};$$

• For any fixed vertex map $\phi_d : V(H_d) \to [n]$,

$$\log |\{\phi_s : (\phi_s, \phi_d) \in \Phi_H^*\}| \le (\ell/2 - r) \log n + (\ell/2 + k) \log 2.$$
(24)

Proof. Let n_0 denote the number of stems and m_0 denote the number of distinct vertices that are tips of some stem in H. Then $n_0 \leq k$ and $m_0 \leq 2n_0$. Recall that in the tree representation of H, each link corresponds to a stem in H and each node corresponds to a cycle. Hence, there are $n_0 + 1$ nodes in the tree. Each cycle after contraction will have at least 3 single edges in H, and thus the total number of single edges $\ell \geq 3(n_0 + 1)$.

We will use the following crucial property in enumerating the vertex maps $\phi: V(H) \to [n]$ that are compatible with x^* . Given a red edge (u, v), $\phi(u)$ and $\phi(v)$ must be neighbors in the cycle corresponding to x^* . Therefore, once the label $\phi(u)$ is fixed, there are at most two different choices

⁹The constraint that $|E(F)| \leq 4n$ is due to the fact that, for any vertex x of the F2F polytope, the multigraph G_y obtained from $y = 2(x - x^*)$ has maximal degree at most 8.

for $\phi(v)$. For every vertex in H, we will assign a mark in a certain order such that the mark on a vertex is an upper bound on the number of choices of its labeling given the labelings of previous vertices. In particular, for every red edge (u, v), if one endpoint has been assigned a mark before the other endpoint, then we assign mark 2 to the other endpoint.

First, we count the vertex maps for double edges $\phi_d : V(H_d) \to [n]$ that are compatible with x^* . Fix a stem $S = (v_0, \ldots, v_s)$ of s double edges in alternating colors. We sequentially choose the vertex labelings from v_0 to v_s . We distinguish three types of stems.

- Type 1: Both tips are red. In this case, the number of different values of $\phi_d(v_i)$ given $(\phi_d(v_0), \ldots, \phi_d(v_{i-1}))$ is at most *n* for even *i* and 2 for odd *i*. See Fig. 12(a).
- Type 2: The two tips have different colors. Without loss of generality, assume (v_0, v_1) is a red double edge. Again the number of choices of $\phi_d(v_i)$ given $(\phi_d(v_0), \ldots, \phi_d(v_{i-1}))$ is at most n for even i and 2 for odd i. See Fig. 12(b).
- Type 3: Both tips are blue. In this case, the number of choices of $\phi_d(v_0)$ is at most n, and the remaining part of stem (v_1, \ldots, v_s) is of Type II. Thus, for $i \ge 1$, the number of possible maps $\phi_d(v_i)$ is given $(\phi_d(v_0), \ldots, \phi_d(v_{i-1}))$ at most n for odd i and 2 for even i. See Fig. 12(c).

In summary, the number of possible vertex labelings for a stem of length s of Type i is at most




Figure 12: Three types of stems. The number on each vertex shows the number of different choices for its labeling under ϕ_d .

Suppose in H there are n_i stems of Type i, for i = 1, 2, 3. Then the total number of different vertex maps for H_d is at most

$$n^{(k+n_1+2n_2+3n_3)/2} (k+n_1-n_3)/2$$

This bound can be further improved by taking into account the fact that some of the tips are either identical or connected by a red single edge. In the following we give a tighter upper bound by incorporating these constraints.

Let ℓ_r and ℓ_b (resp. k_r and k_b) denote the number of red and blue single (resp. double) edges in H respectively. Then we have $k_r + k_b = k$ and $k_r - k_b = n_1 - n_3$. Hence,

$$k_r = \frac{1}{2}(k+n_1-n_3), \quad k_b = \frac{1}{2}(k-n_1+n_3).$$
 (25)

Analogously, $\ell_r + \ell_b = \ell$ and $2k_r + \ell_r = 2k_b + \ell_b$ by the balancedness of H. Hence,

$$\ell_r = \frac{\ell}{2} - n_1 + n_3, \quad \ell_b = \frac{\ell}{2} + n_1 - n_3.$$
 (26)

Moreover, the number of red (resp. blue) tips is $2n_1+n_2$ (resp. n_2+2n_3). To count the total number of vertex maps for H_d , it suffices to count the labelings of tip and non-tip vertices separately. We will count the labelings sequentially: first for tip vertices and then for non-tip vertices.

Tips. Without loss of generality, we assume there is no cycle in H consisting of only red single edges. Suppose, for the sake of contradiction, there is a cycle $(u_0, \ldots, u_{\ell_1-1}, u_0)$ consisting of ℓ_1 red single edges. By the construction of \mathcal{U} , each vertex is attached to at least one flower. For any homomorphism $\phi : H \to F$, $(u_0, \ldots, u_{\ell_1-1}, u_0)$ is mapped to the ground truth cycle x^* . Therefore $\ell_1 = n$. A flower has at least 5 edges. Therefore, the total number of edges (counting with multiplicity) in E(F) is at least n + 5n = 6n, and thus ϕ is not compatible with x^* by definition. Therefore, there is no homomorphism $H \to F$ compatible with x^* .

In a given cycle of H, we define a red path to be a maximal path consisting of all red single edges. We define a vertex as a red path of length 0, if it is incident to two blue single edges in H. Then all cycles in H are segmented by blue single edges into a total of ℓ_b red paths (cf. Fig. 13 for an illustration). We distinguish two types of red paths depending on whether it is incident to a double edge or not:

- Type I: The red path is not incident to any double edge. In this case, the red path must consist of only one red single edge.
- Type II: The red path is incident to at least one double edge. In this case, at least one of the vertex in the path is a tip of some stem.

To count the vertex maps for tips of stems, it suffices to focus on red paths of Type II. Note that two red paths of Type II may be connected by a stem consisting of only one red double edge. To account for the total number of constraints induced by red paths, let n_4 denote the total number of red paths of Type II and let m_1 denotes the total number of stems consisting of one red double edge. Define a graph whose vertices are red paths of Type II, and two red paths are connected if they are connected by a one red double edge. By definition of the tree representation, such a graph must be a forest; thus the number of connected components therein is $n_4 - m_1$. Call each connected component a *red component*. The key observation is that, for each red component, once we fix the labeling for a given vertex in it, any remaining vertex has at most 2 labelings. Moreover, each red component must contain at least one vertex as the tip of some stem, and each tip is contained in some red component. Therefore, we can assign marks to tips in two steps: First, for each red component, pick an arbitrary tip and assign a mark n to it; next, we assign mark 2 to each of the remaining tips. Since there are $n_4 - m_1$ red component, there are in total $n_4 - m_1$ tips having mark n.

Non-tips. Given the labeling of tips, we count the labeling of non-tips. Note that each non-tip vertex u is incident to at least one red double edge. Consider two cases separately:

• If the red double edge is incident to a tip, since the labeling of tips have been assigned, there are at most 2 possible labelings for u. Hence we assign a mark 2 to vertex u.




Figure 13: Three cycles with attached double edges in H. There are one red path with length 2 of type II in cycle C_1 . There are three red paths in cycle C_2 : one with length 1 of type I, one with length 0 and the other of length 3 both of type II. There are two red paths in cycle C_3 : one with length 1 of type I and the other with length 1 of type II. In total, there are 2 red components formed by 4 red paths of type II: one formed by the red path in cycle C_1 and the red path with length 0 in cycle C_2 , and the other formed by the red path with length 3 in cycle C_2 and the red path with length 1 of type II in cycle C_3 .

• If the red double edge is in the interior of the stem, i.e., not incident to any tip, we assign mark n to one of its endpoint and mark 2 to the other endpoint. Recall that m_1 denotes the number of stems consisting of only one red double edge. Then there are exactly

$$k_r - 2(n_1 - m_1) - n_2 - m_1 = k_r - 2n_1 - n_2 + m_1$$

number of red double edges in the interior of the stems.

By now we have assigned marks – either n or 2 – to every vertex in H_d . Once we multiply all the marks together, we obtain an upper bound on the number of vertex maps ϕ_d for H_d . Recall that the total number of stems is $n_0 = n_1 + n_2 + n_3$ and m_0 denotes the number of distinct tip vertices in H. Then in total there are $k - n_0 + m_0$ vertices in H_d . Recall that there are $n_4 - m_1$ marks of n assigned to tips and $k_r - 2n_1 - n_2 + m_1$ marks of n assigned to non-tips. Hence, in total we have $k_r - 2n_1 - n_2 + n_4$ marks of n. The rest of marks all take values 2. In view of (25), we have

$$k_r - 2n_1 - n_2 + n_4 = \frac{1}{2} \left(k + n_1 - n_3 \right) - 2n_1 - n_2 + n_4 = \frac{1}{2} \left(k - 3n_1 - 2n_2 - n_3 + 2n_4 \right).$$

Therefore,

$$\log |\{\phi_d : \phi \in \Phi_H^*\}| \le \frac{1}{2} \left(k - 3n_1 - 2n_2 - n_3 + 2n_4\right) \log(n/2) + \left(k - n_0 + m_0\right) \log 2.$$
(27)

Next, we fix a vertex map ϕ_d for H_d and count all the possible vertex maps for H_s that are compatible with x^* . To this end, a key observation is that it suffices to count the possible labelings for the endpoints of red single edges in H_s . Indeed, for any vertex u not incident to any red single edges, by the balancedness and connectedness of H, u must be incident to at least one red double edge, which has been accounted for by ϕ_d .

For each red path of Type I, it consists of only one red single edge and thus we assign mark n to one of its endpoint and 2 to its other endpoint. For each red path of Type II, since it contains at least one tip whose vertex map has already been fixed by the vertex map for double edges, each of its other vertex has at most 2 different vertex maps. Hence, we assign mark 2 to every vertex other than tips in the red path.

Recall that there are $\ell_b - n_4$ red paths of Type I. Hence the total number of marks of value n, in view of (26), is

$$\ell_b - n_4 = \frac{\ell}{2} + n_1 - n_3 - n_4.$$

The rest of marks are 2. Recall that m_0 denotes the total number of distinct tip vertices in H, and each tip is incident to at least one single edge. Note that there are in total ℓ vertices in H_s . Thus m_0 of them are tips. Hence, there are $\ell - m_0$ vertices in H_s we need to assign marks to and thus the total number of marks of 2 in H_s is at most $\ell - m_0 - (\ell_b - n_4)$. In total, we have

$$\log |\{\phi_s : (\phi_s, \phi_d) \in \Phi_H^*\}| \le (\ell/2 + n_1 - n_3 - n_4) \log(n/2) + (\ell - m_0) \log 2$$

= $(\ell/2 - r) \log n + (\ell/2 - m_0 + r) \log 2,$ (28)

where we defined

$$r \triangleq n_3 + n_4 - n_1$$

Furthermore, recall that n_0 denotes the total number of stems. Note that $n_4 \leq m_0 \leq 2n_0$, because every red path of Type II must contain at least one tip, and each stem has at most 2 tip vertices that are distinct from the tip vertices of the other stems. Using $n_0 = n_1 + n_2 + n_3$ and $n_4 \leq 2n_0$, we get that $r \geq -3n_1 - 2n_2 - n_3 + 2n_4$. In view of (27), we have

$$\log |\{\phi_d : \phi \in \Phi_H^*\}| \le \frac{1}{2} (k+r) \log(n/2) + (k-n_0+m_0) \log 2$$

$$= \frac{1}{2} (k+r) \log n + \frac{1}{2} (k-r-2n_0+2m_0) \log 2$$

$$\le \frac{1}{2} (k+r) \log n + \frac{1}{2} (k-r+m_0) \log 2, \qquad (29)$$

where the last inequality holds because $m_0 \leq 2n_0$.

Next we show that $m_0/2 \le r \le \min\{\ell/2, m_0+n_0\}$. Indeed, by definition, we have $r \le n_3+n_4 \le n_0 + m_0$. Since $0 \le \ell_b - n_4 = \ell/2 - r$, it follows that $r \le \ell/2$. Furthermore, for any red path of Type I, it consists of precisely one red single edge, where neither of the two endpoints is a tip. Thus $\ell - m_0 \ge 2(\ell_b - n_4)$. Recall that $\ell_b - n_4 = \ell/2 + n_1 - n_3 - n_4 = \ell/2 - r$. It follows that $r \ge m_0/2$.

Finally, the desired (23) follows from (29) in view of $m_0/2 \le r$ and the desired (24) follows from (28) in view of $r \le m_0 + n_0 \le m_0 + k$.

5.4 Proof of Theorem 1

We prove that if

$$\alpha - \log n \ge 16 \log 17,\tag{30}$$

then

$$\min_{x^*} \mathbb{P}\left\{ \hat{x}_{F2F} = x^* \right\} \ge 1 - 8 \exp\left(-(\alpha - \log n)/8\right).$$
(31)

Then Theorem 1 readily follows by taking $\alpha - \log n \to +\infty$. By Lemma 3,

$$G_y = \bigcup_{i=1}^m F_i, \quad F_i \in \mathcal{F}$$

for each $1 \leq i \leq m$ and some finite m. Note that for each red edge e in G_y , $x_e^* = 1$. Therefore, $F_i \in \mathcal{F}^*$. Thus, to prove (31), it suffices to show

$$\mathbb{P}\left\{\max_{F\in\mathcal{F}^*} w(F) < 0\right\} \ge 1 - 8\exp\left(-(\alpha - \log n)/8\right).$$
(32)

Fix $k \ge 0$ and $\ell \ge 4$, define

 $\mathcal{F}_{k,\ell}^* = \{ F \in \mathcal{F}^* : E(F) \text{ consists of } k \text{ double edges and } \ell \text{ single edges } \}.$

Then

$$\mathcal{F}_{k,\ell}^* = \{\phi(H) : H \in \mathcal{U}_{k,\ell} \text{ and } \phi \in \Phi_H^*\}$$
(33)

and

$$\mathcal{F}^* = \bigcup_{k \ge 0} \bigcup_{\ell \ge 4} \mathcal{F}^*_{k,\ell}$$

In view of (33), we have

$$\max_{F \in \mathcal{F}_{k,\ell}^*} w(F) = \max_{H \in \mathcal{U}_{k,\ell}} \max_{\phi \in \Phi_H^*} w(\phi(H))$$
(34)

We first show a high-probability bound to the inner maximum for a given $H \in \mathcal{U}_{k,\ell}$. Since maximizing over ϕ is equivalent to first maximizing over ϕ_d and then maximizing over ϕ_s for a fixed ϕ_d , it follows that

$$\max_{\phi \in \Phi_H^*} w(\phi(H)) = \max_{\phi_d \in \Phi_{H_d}^*} \left(w\left(\phi_d(H_d)\right) + \max_{\phi_s:(\phi_d,\phi_s) \in \Phi_H^*} w\left(\phi_s(H_s)\right) \right)$$

Recall that X_i 's and Y_i 's are two independent sequences of random variables, where X_i 's are i.i.d. copies of dP/dQ under distribution P and Y_i 's are i.i.d. copies of dP/dQ under distribution Q. Recall that ℓ_r and ℓ_b (resp. k_r and k_b) denote the number of red and blue single (resp. double) edges in H respectively. Let $\delta = k_b - k_r = (\ell_r - \ell_b)/2$. Then $\delta \leq \min\{k, \ell/2\}$. In view of (25) and (26), for a fixed ϕ_d ,

$$w(\phi_d(H_d)) \stackrel{d}{=} 2\left(\sum_{i=1}^{(k+\delta)/2} Y_i - \sum_{i=1}^{(k-\delta)/2} X_i\right)$$

and for a fixed ϕ_s ,

$$w\left(\phi_s(H_s)\right) \stackrel{d}{=} \sum_{i=1}^{\ell/2-\delta} Y_i - \sum_{i=1}^{\ell/2+\delta} X_i$$

where $\stackrel{d}{=}$ denotes equality in distribution. Moreover, for a fixed ϕ_d , $w(\phi_d(H_d))$ is the sum of the weights on double edges, which is independent of the collection of $w(\phi_s(H_s))$ ranging over all possible ϕ_s such that $(\phi_s, \phi_d) \in \Phi_H^*$.

Recall from Lemma 5 that there exists an integer $0 \le r \le \ell/2$ such that

$$\log |\Phi_{H_d}^*| \le \frac{1}{2}(k+r)\log(2n),$$
$$\log |\{\phi_s : (\phi_s, \phi_d) \in \Phi_H^*\}| \le (\ell/2 - r)\log n + (k + \ell/2)\log 2.$$

Invoking the large deviation bound Lemma 7 in Appendix B with $s = (k - \delta)/2$, $t = \ell/2 - \delta$, $u = \delta$, and $v = r - \delta$, and noting that

$$s + u + v/2 = (k + r)/2$$

$$t - v = \ell/2 - r$$

$$s + t + u - v/2 = (k + \ell - r)/2 \ge k/2 + \ell/4,$$

and (30), we get

$$\mathbb{P}\left\{\max_{\phi\in\Phi_{H}^{*}}w(\phi(H))\geq 0\right\}\leq 5\exp\left(-(\alpha-\log n)(k/8+\ell/16)\right).$$
(35)

It follows that

$$\mathbb{P}\left\{\max_{F\in\mathcal{F}_{k,\ell}^{*}}w(F) \ge 0\right\} \stackrel{(a)}{\le} 5|\mathcal{U}_{k,\ell}|\exp\left(-(\alpha - \log n)(k/8 + \ell/16)\right)$$

$$\stackrel{(b)}{\le} 5 \times 17^{k}4^{\ell}\exp\left(-(\alpha - \log n)(k/8 + \ell/16)\right)$$

$$\stackrel{(c)}{\le} 5\exp\left(-(\alpha - \log n)(k/16 + \ell/32)\right),$$

where (a) follows from union bound; (b) follows from Lemma 4; (c) holds because $\alpha - \log n \ge 16 \log 17$ by assumption (30). Taking another union bound over $k \ge 0$ and $\ell \ge 4$, we get

$$\begin{split} \mathbb{P}\left\{\max_{F\in\mathcal{F}^*}w(F)\geq 0\right\} &= \mathbb{P}\left\{\max_{k\geq 0,\ell\geq 4}\max_{F\in\mathcal{F}^*_{k,\ell}}w(F)\geq 0\right\}\\ &\leq \sum_{k\geq 0}\sum_{\ell\geq 4}\mathbb{P}\left\{\max_{F\in\mathcal{F}^*_{k,\ell}}w(F)\geq 0\right\}\\ &\leq 5\sum_{k\geq 0}\sum_{\ell\geq 4}\exp\left(-(\alpha-\log n)(k/16+\ell/32)\right)\\ &\leq \frac{5}{1-e^{-(\alpha-\log n)/8}}\frac{e^{-(\alpha-\log n)/8}}{1-e^{-(\alpha-\log n)/32}}\\ &\leq \frac{5}{1-1/17}\frac{e^{-(\alpha-\log n)/8}}{1-1/4}\leq 8e^{-(\alpha-\log n)/8}. \end{split}$$

Therefore, we arrive at the desired (32), completing the proof of Theorem 1.

6 Information-theoretic Necessary Conditions

We first present a general necessary condition needed for any algorithm to succeed in recovering the hidden Hamiltonian cycle with high probability. Recall that X and Y are two independent random variables distributed as the log likelihood ratio $\log(dP/dQ)$ under P and Q, respectively.

Theorem 2 (Information-theoretic conditions). If there exists a sequence of estimators \widehat{C} such that $\min_{C^*} \mathbb{P}\{\widehat{C} = C^*\} \to 1 \text{ as } n \to \infty$, then

$$\sup_{\tau \in \mathbb{R}} \left\{ \log \mathbb{P} \left\{ X \le \tau \right\} + \log \mathbb{P} \left\{ Y \ge \tau \right\} \right\} + \log n \le O(1).$$
(36)

Next, we state a regularity assumption on P and Q under which it immediately follows from Theorem 2 that $\alpha_n \ge (1 + o(1)) \log n$ is necessary information-theoretically, thereby establishing the optimality of F2F LP.

Assumption 1.

$$\sup_{\tau \in \mathbb{R}} \left\{ \log \mathbb{P}\left\{ X \le \tau \right\} + \log \mathbb{P}\left\{ Y \ge \tau \right\} \right\} \ge -(1 + o(1)) \alpha_n + o(\log n).$$

Corollary 1. Suppose Assumption 1 holds. If there exists a sequence of estimators \widehat{C} such that $\min_{C^*} \mathbb{P}\{\widehat{C} = C^*\} \to 1 \text{ as } n \to \infty$, then

$$\alpha_n \ge (1+o(1))\log n. \tag{37}$$

Assumption 1 is very general and fulfilled when the weight distributions are either Poisson, Gaussian or Bernoulli as the following result shows (see Appendix C for a proof):

Lemma 6. Assumption 1 holds in the Gaussian case with $P = \mathcal{N}(\mu, 1)$ and $Q = \mathcal{N}(0, 1)$, the Poisson case with $P = \text{Pois}(\lambda)$ and $Q = \text{Pois}(\mu)$ for for $\lambda \ge \mu$ such that

$$\log(\lambda\mu) = o\left((\sqrt{\lambda} - \sqrt{\mu})^2\right) + o(\log n)$$

and the Bernoulli case with P = Bern(p) and Q = Bern(q) for $p \ge q$.

Let us now explain the intuition behind Assumption 1: Denote the log moment generating function of X and Y as

$$\psi_P(\theta) = \log \mathbb{E}[e^{\theta X}], \quad \psi_Q(\theta) = \log \mathbb{E}[e^{\theta Y}] = \psi_P(\theta - 1).$$
 (38)

Denote the Legendre transform of ψ_P and ψ_Q as

$$E_P(\tau) = \sup_{\theta \ge 0} \left\{ -\theta\tau - \psi_P(-\theta) \right\}, \quad E_Q(\tau) = \sup_{\theta \ge 0} \left\{ \theta\tau - \psi_Q(\theta) \right\}.$$
(39)

Then Chernoff's inequality gives the following large deviation bounds: for any $\tau \in \mathbb{R}$,

$$\mathbb{P}\left\{X \le \tau\right\} \le \exp\left(-E_P(\tau)\right), \quad \mathbb{P}\left\{Y \ge \tau\right\} \le \exp\left(-E_Q(\tau)\right), \tag{40}$$

Therefore,

$$\sup_{\tau \in \mathbb{R}} \left\{ \log \mathbb{P} \left\{ X \le \tau \right\} + \log \mathbb{P} \left\{ Y \ge \tau \right\} \right\} \le - \inf_{\tau \in \mathbb{R}} \left\{ E_P(\tau) + E_Q(\tau) \right\}$$

The infimum on the right-hand side is in fact equal to α_n . Indeed,

$$\begin{split} \inf_{\tau \in \mathbb{R}} E_P(\tau) + E_Q(\tau) &= \inf_{\tau \in \mathbb{R}} \sup_{\theta_1, \theta_2 \ge 0} \left\{ -\theta_1 \tau - \psi_P(-\theta_1) + \theta_2 \tau - \psi_Q(\theta_2) \right\} \\ &\geq \sup_{\theta_1, \theta_2 \ge 0} \left\{ \inf_{\tau \in \mathbb{R}} (\theta_2 - \theta_1) \tau - \psi_P(-\theta_1) - \psi_Q(\theta_2) \right\} \\ &= \sup_{\theta \ge 0} \left\{ -\psi_P(-\theta) - \psi_Q(\theta) \right\} \\ &= -\psi_P(-1/2) - \psi_Q(1/2) = -2\log \int \sqrt{\mathrm{d}P\mathrm{d}Q} = \alpha_n \end{split}$$

and the infimum over τ is in fact achieved by

$$\tau^* = \psi'_P(-1/2) = \psi'_Q(1/2) \,,$$

so that $E_P(\tau^*) + E_Q(\tau^*) = \alpha_n$. Hence,

$$\sup_{\tau \in \mathbb{R}} \left\{ \log \mathbb{P} \left\{ X \le \tau \right\} + \log \mathbb{P} \left\{ Y \ge \tau \right\} \right\} \le -E_P \left(\tau^*\right) - E_Q \left(\tau^*\right) = -\alpha_n.$$

Therefore, the point of Assumption 1 is to require that the large deviation exponents in Chernoff's inequalities (40) are asymptotically tight, so that we can reverse the Chernoff bound in the lower bound proof.

Proof of Theorem 2. To lower bound the worst-case probability of error, consider the Bayesian setting where C^* is drawn uniformly at random from all possible Hamiltonian cycles of G. Since the prior distribution of C^* is uniform, the ML estimator minimizes the error probability among all estimators. Thus, without loss of generality, we can assume the estimator \hat{C} used is \hat{C}_{ML} and the true Hamiltonian cycle C^* is given by $(1, 2, \ldots, n, 1)$. Hence, by assumption $\mathbb{P}\{\hat{C}_{ML} = C^*\} \to 1$.

Recall that the ML estimator is equivalent to finding a Hamiltonian cycle of the maximum weight. Given a Hamiltonian cycle x, define the simple graph G_x with bicolored edge whose adjacency matrix is $|x - x^*|$ and each edge is colored in red if $(x - x^*)_e = -1$ and in blue if $(x - x^*)_e = +1$. Also, each edge e has a weight $w_e(x - x^*)_e$ and hence $w(G_x) = \langle w, x - x^* \rangle$. Note that if G_x is a 4-cycle of alternating colors given by (i, i + 1, j + 1, j, i), then x corresponds to a Hamiltonian cycle constructed by deleting edges (i, i + 1), (j, j + 1) in C^* and adding edges (i, j), (i + 1, j + 1) (see Fig. 14 for an illustration). Let \mathcal{D} denote the set of all possible 4-cycles of alternating colors given by (i, i + 1, j + 1, j). Then $|\mathcal{D}| = n(n-3)/2$, because for a given i, j have (n-3) choices except i - 1, i, i + 1.




Figure 14: The cycle (1, 2, ..., i, j, j - 1, ..., i + 1, j + 1, j + 2, ..., n) and the corresponding graph G_x as a four-cycle.

Define

$$S = \sum_{D \in \mathcal{D}} \mathbf{1}_{\{w(D) \le 0\}}.$$

If S > 0, then there exists a Hamiltonian cycle $x \neq x^*$ whose weight is at least as large as the weight of C^* ; hence the likelihood function has at least two maximizers, which in turn implies the probability of exact recovery by ML estimator is at most 1/2. Therefore, $\frac{1}{2}\mathbb{P}\{S > 0\} \leq \mathbb{P}\{ML \text{ fails}\} = o(1)$. As a consequence, $\mathbb{P}\{S = 0\} \rightarrow 1$.

To explain the intuition, suppose w(D) are mutually independent for all $D \in \mathcal{D}$. Then

$$\mathbb{P}\left\{S=0\right\} = \mathbb{P}\left\{\forall D \in \mathcal{D}, w(D) > 0\right\}$$
$$= \prod_{D \in \mathcal{D}} \mathbb{P}\left\{w(D) < 0\right\}$$
$$\stackrel{(a)}{=} (1 - \mathbb{P}\left\{Y_1 + Y_2 - X_1 - X_2 \ge 0\right\})^{|\mathcal{D}|}$$
$$\le \exp\left(-|\mathcal{D}|\mathbb{P}\left\{Y_1 + Y_2 - X_1 - X_2 \ge 0\right\}\right)$$

where (a) holds because w(D) has the same distribution as $Y_1 + Y_2 - X_1 - X_2$ and the last inequality holds in view of $1 - x \le e^{-x}$. In view of $\mathbb{P}\{S = 0\} \to 1$, it follows from the last displayed equation that

$$\log |\mathcal{D}| + \log \mathbb{P} \left\{ Y_1 + Y_2 - X_1 - X_2 \ge 0 \right\} \to -\infty.$$

Furthermore, for any $\tau \in \mathbb{R}$, we have

$$\log \mathbb{P}\left\{Y_1 + Y_2 - X_1 - X_2 \ge 0\right\} \ge \log \left(\mathbb{P}\left\{Y_1 \ge \tau\right\} \mathbb{P}\left\{Y_2 \ge \tau\right\} \mathbb{P}\left\{X_1 \le \tau\right\} \mathbb{P}\left\{X_2 \le \tau\right\}\right)$$
$$= 2\log \mathbb{P}\left\{Y \ge \tau\right\} + 2\log \mathbb{P}\left\{X \le \tau\right\}.$$

Combining the last two displayed equation and recalling that $|\mathcal{D}| = n(n-3)/2$, we immediately get that

$$\log \mathbb{P}\left\{Y \ge \tau\right\} + \log \mathbb{P}\left\{X \le \tau\right\} + \log n \to -\infty,$$

Taking the supremum over $\tau \in \mathbb{R}$ of the last displayed equation yields the desired (36).

However, w(D) and w(D') are dependent if D and D' share edges. To deal with this dependency, we focus on a subset of \mathcal{D} . In particular, for any $\tau \in \mathbb{R}$, define

$$I = \{ \text{odd } i : w_{i,i+1} \le \tau \}$$

and

$$J = \{(i,j) \in I \times I : i \neq j, w_{i,j} + w_{i+1,j+1} \ge 2\tau\}$$

Then for any $(i, j) \in J$, the alternating 4-cycle given by (i, i+1, j+1, j, i) belongs to \mathcal{D} and has a non-positive weight. Hence, $|J| \leq S$ and thus $\mathbb{P}\{|J| = 0\} \geq \mathbb{P}\{S = 0\} \rightarrow 1$.

Note that $w_{i,i+1}$ has the same distribution as X. Thus for any $\tau \in \mathbb{R}$,

$$\mathbb{P}\left\{w_{i,i+1} \le \tau\right\} = \mathbb{P}\left\{X \le \tau\right\} \triangleq p$$

Also, $w_{i,i+1}$ are mutually independent for different *i*. Thus $|I| \sim \text{Binom}(\lceil n/2 \rceil, p)$. By the Chernoff's bound for binomial distribution,

$$\mathbb{P}\left\{|I| \le np/4\right\} \le \exp\left(-np/8\right).$$

Thus,

$$\begin{split} \mathbb{P}\left\{|J|=0\right\} &\leq \mathbb{P}\left\{|J|=0, |I| > np/4\right\} + \mathbb{P}\left\{|I| \le np/4\right\} \le \mathbb{P}\left\{|J|=0 \mid |I| > np/4\right\} + \exp\left(-np/8\right).\\ \text{Let } q \triangleq \mathbb{P}\left\{Y \ge \tau\right\}. \text{ Then } \mathbb{P}\left\{Y_1 + Y_2 \ge 2\tau\right\} \ge \mathbb{P}\left\{Y_1 \ge \tau\right\} \mathbb{P}\left\{Y_2 \ge \tau\right\} = q^2 \text{ and hence}\\ \mathbb{P}\left\{|J|=0 \mid |I| > np/4\right\} = \mathbb{P}\left\{\forall i < j \in I, w_{(i,j)} + w_{(i+1,j+1)} < 2\tau \mid |I| > np/4\right\}\\ &\stackrel{(a)}{\le} (1-q^2)^{\binom{np/4}{2}} \le e^{-q^2\binom{np/4}{2}}, \end{split}$$

where (a) holds because conditional on I, $w_{(i,j)} + w_{(i+1,j+1)}$ are i.i.d. copies of $Y_1 + Y_2$. Combining the last three displayed equations yields that

$$\mathbb{P}\left\{|J|=0\right\} \le e^{-q^2\binom{np/4}{2}} + e^{-np/8}.$$
(41)

Recall that $\mathbb{P}\{|J|=0\} \to 1$. It follows that

$$e^{-q^2\binom{np/4}{2}} + e^{-np/8} \ge 1 + o(1).$$
 (42)

Hence, $\log n + \log p + \log q \le O(1)$, or equivalently,

$$\log \mathbb{P}\left\{X \le \tau\right\} + \log \mathbb{P}\left\{Y \ge \tau\right\} + \log n \le O(1).$$

Taking the supremum over $\tau \in \mathbb{R}$ of the last displayed equation yields the desired (36).

7 Reduction between Hamiltonian Cycle and Hamiltonian Path

In this section, we extend our results to the hidden Hamiltonian path model. As opposed to Definition 1, G is a randomly weighted, undirected complete graph with a hidden Hamiltonian path H^* such that every edge has an independent weight drawn from P if it is on H^* and from Q otherwise.

The following theorem shows that the information-theoretic limits of hidden Hamiltonian cycle and path models coincide.

Theorem 3. Consider the ML estimator $\widehat{H}_{ML} \in \arg \min_{H \in \mathcal{H}} w(H)$, where \mathcal{H} denotes the set of all possible Hamiltonian paths in graph G and w(H) denote the total weights on the Hamiltonian path H. If

$$\alpha_n - \log n \to +\infty,$$

i.e., (9) holds, then $\min_{H^* \in \mathcal{H}} \mathbb{P}\{\widehat{H}_{\mathrm{ML}} = H^*\} \to 1 \text{ as } n \to \infty.$

Conversely, suppose Assumption 1 holds. If there exists a sequence of estimators \widehat{H} such that $\min_{H^* \in \mathcal{H}} \mathbb{P}\{\widehat{H} = H^*\} \to 1 \text{ as } n \to \infty$, then

$$\alpha_n \ge (1 + o(1)) \log n,$$

i.e., (37) holds.

Proof. We first prove the sufficiency part. We abbreviate the ML estimator \hat{H}_{ML} as \hat{H} . Let $L = |H^* \setminus \hat{H}|$. Since $|H^*| = |\hat{H}| = n - 1$, it follows that $0 \le L \le n - 1$ and $|H^* \Delta \hat{H}| = 2L$. To prove the theorem, it suffices to show that $\mathbb{P}\{L \ge 1\} = o(1)$. For any $1 \le \ell \le n - 1$,

 $\{L = \ell\} \subset \{\exists \text{ Hamiltonian path } H : |H^* \setminus H| = \ell, w(H^*) < w(H)\}.$

Note that for each fixed $H \in \mathcal{H}_n$ with $|H^* \setminus H| = \ell$,

$$w(H) - w(H^*) \stackrel{d}{=} \sum_{i=1}^{\ell} Y_i - \sum_{i=1}^{\ell} X_i,$$

where $\{X_i\}$ and $\{Y_i\}$ are independent and iid copies of the log likelihood ratio $\log(dP/dQ)$ under P and Q, respectively. Also, there are at most $\binom{n-1}{\ell}2^{\ell+1}(\ell+1)!$ different choices of Hamiltonian paths H with $|H^* \setminus H| = \ell$. To see this, note that there are at most $\binom{n-1}{\ell}$ different choices of edges in $H^* \setminus H$. For each such choice, there are at most $\ell + 1$ disjoint paths in $H^* \cap H$; thus there are at most $2^{\ell+1}$ different orientations of these paths, and at most $(\ell+1)!$ different permutations of these paths.

Applying union bound gives that

$$\mathbb{P}\{L=\ell\} \le \binom{n-1}{\ell} 2^{\ell+1} (\ell+1)! \times \mathbb{P}\left\{\sum_{i=1}^{\ell} Y_i - \sum_{i=1}^{\ell} X_i < 0\right\}$$

$$\le n^{\ell} 2^{\ell+1} (\ell+1) e^{-\ell\alpha} \le e^{-\omega(\ell)},$$

where the second inequality follows due to $\binom{n-1}{\ell}(\ell+1)! \leq n^{\ell}(\ell+1)$ and Lemma 7; the last inequality follows from the assumption (9). Therefore,

$$\mathbb{P}\{L \ge 1\} = \sum_{\ell=1}^{n-1} \mathbb{P}\{L = \ell\} \le \sum_{\ell=1}^{n-1} e^{-\omega(\ell)} = e^{-\omega(1)},$$

completing the proof for sufficiency part.

Next we show the necessity part. To lower bound the worst-case probability of error, consider the Bayesian setting where H^* is uniformly chosen among all possible Hamiltonian paths of G. Since the ML estimator minimizes the error probability among all estimators, without loss of generality, we can assume the estimator \hat{H} used is \hat{H}_{ML} and the true Hamiltonian path H^* is given by $(1, 2, \ldots, n)$. Hence, by assumption $\mathbb{P}\left\{\hat{H}_{ML} = H^*\right\} \to 1$. Note that under the hidden Hamiltonian path model, the ML estimator is equivalent to finding the maximum weighted Hamiltonian path. For $i \in \{2, \ldots, n-1\}$, define the event that

$$\mathcal{E}_i = \{ w_{1,i+1} \ge w_{i,i+1} \}.$$

Let $\mathcal{F} = \bigcup_{i=2}^{n-1} \mathcal{E}_i$, which implies the existence of vertex *i* such that the total weight of the Hamiltonian path $H^* \setminus \{(i, i+1)\} \cup \{(1, i+1)\}$ is at least as large as the total weight of H^* . Hence, on \mathcal{F} , the likelihood function has at least two maximizers, which in turn implies the probability of exact recovery by ML estimator is at most 1/2. Hence $\frac{1}{2}\mathbb{P}\{\mathcal{F}\} \leq \mathbb{P}\{ML \text{ fails}\} = o(1)$. As a consequence, $\mathbb{P}\{\mathcal{F}^c\} \to 1$. Note that

$$\mathbb{P}\left\{\mathcal{F}^{c}\right\} \stackrel{(a)}{=} \prod_{i=2}^{n-1} \mathbb{P}\left\{\mathcal{E}_{i}^{c}\right\}$$
$$\stackrel{(b)}{=} (1 - \mathbb{P}\left\{X \leq Y\right\})^{n-2}$$
$$\stackrel{(c)}{\leq} \exp\left(-(n-2)\mathbb{P}\left\{X \leq Y\right\}\right)$$

where (a) holds due the independence among \mathcal{E}_i ; (b) holds because $\mathbb{P} \{ \mathcal{E}_i \} = \mathbb{P} \{ X < Y \}$; (c) follows due to $1 + x \leq e^x$ for all $x \in \mathbb{R}$. It follows that

$$\log n + \log \mathbb{P}\left\{X \le Y\right\} = o(1).$$

Note that

$$\begin{split} \log \mathbb{P}\left\{X \leq Y\right\} &\geq \sup_{\tau \in \mathbb{R}} \left\{\log \mathbb{P}\left\{X \leq \tau\right\} + \log \mathbb{P}\left\{Y \geq \tau\right\}\right\} \\ &\geq -\left(1 + o(1)\right)\alpha + o(\log n), \end{split}$$

where the last inequality is exactly Assumption 1. Combining the last two displayed equations yields that $\alpha \ge (1 + o(1)) \log n$.

Next, we show that given a polynomial-time estimator which achieves exact recovery in one model, one can construct another polynomial-time estimator which achieves exact recovery for the other model whenever it is information-theoretically possible.

Algorithm 1 gives a procedure to construct a polynomial-time estimator of Hamiltonian path given a polynomial-time estimator of Hamiltonian cycle.

Similarly, Algorithm 2 gives a procedure to construct a polynomial-time estimator of Hamiltonian cycle given a polynomial-time estimator of Hamiltonian path.

Theorem 4. Given an estimator $\widehat{C}(G)$ with time complexity $O(T_n)$ such that $\mathbb{P}\{\widehat{C} = C^*\} \to 1$ as $n \to \infty$ under the hidden Hamiltonian cycle model. Let $\widehat{H}(G)$ denote the output of Algorithm 1 with input $\widehat{C}(G)$. Then $\widehat{H}(G)$ has time complexity $O(n^2T_n)$. Moreover, under the hidden Hamiltonian path model if $\mathbb{P}\{\widehat{H}_{ML} = H^*\} \to 1$ as $n \to \infty$, then $\mathbb{P}\{\widehat{H} = H^*\} \to 1$.

Algorithm 1 From Hamiltonian cycle estimator to Hamiltonian path estimator

- 1: Input: Graph G, an estimator $\widehat{C}(G)$ of Hamiltonian cycle in G.
- 2: **Output:** An estimator $\widehat{H}(G)$ of Hamiltonian path in G.
- 3: Generate a random variable $W \sim P$.
- 4: For every edge e = (i, j) in G, construct a weighted graph G^e by replacing the edge weight on e in G by W. If $\widehat{C}(G^e)$ is a Hamiltonian cycle containing e, let \widehat{H}^e denote the Hamiltonian path by deleting edge e from $\widehat{C}(G^e)$; otherwise, let $\widehat{H}^e = \emptyset$.
- 5: Output

$$\widehat{H} \in \arg\min_{e:\widehat{H}^e \neq \emptyset} w\left(\widehat{H}^e\right),$$

where $w\left(\widehat{H}^e\right)$ denotes the sum of log likelihood weights on edges in \widehat{H}^e .

Algorithm 2 From Hamiltonian path estimator to Hamiltonian cycle estimator

- 1: Input: Graph G, an estimator $\widehat{H}(G)$ of Hamiltonian path in G.
- 2: **Output:** An estimator $\widehat{C}(G)$ of Hamiltonian cycle in G.
- 3: Generate a random variable $W \sim Q$.
- 4: For every edge e = (i, j) in G, construct a weighted graph G^e by replacing the edge weight on e in G by W. If $\widehat{H}(G^e)$ is a Hamiltonian path connecting vertices i, j, let \widehat{C}^e denote the Hamiltonian cycle by adding edge e in $\widehat{H}(G^e)$; otherwise, let $\widehat{C}^e = \emptyset$.
- 5: Output

$$\widehat{C} \in \arg\min_{e:\widehat{C}^e \neq \emptyset} w\left(\widehat{C}^e\right)$$

where $w\left(\widehat{C}^e\right)$ denotes the sum of log likelihood weights on edges in \widehat{C}^e .

Conversely, given an estimator $\widehat{H}(G)$ with time complexity $O(T_n)$ such that $\mathbb{P}\{\widehat{H} = H^*\} \to 1$ as $n \to \infty$ under the hidden Hamiltonian path model. Let $\widehat{C}(G)$ denote the output of Algorithm 2 with input $\widehat{H}(G)$. Then $\widehat{C}(G)$ has time complexity $O(n^2T_n)$. Moreover, under the hidden Hamiltonian cycle model if $\mathbb{P}\{\widehat{C}_{ML} = C^*\} \to 1$ as $n \to \infty$, then $\mathbb{P}\{\widehat{C} = C^*\} \to 1$.

Proof. By construction in Algorithm 1, the time complexity of $\widehat{H}(G)$ is $O(n^2T_n)$. We show that $\mathbb{P}\{\widehat{H} = H^*\} \to 1$. Without loss of generality, assume that $H^* = (1, 2, ..., n)$. Then according to Algorithm 1, G^e with e = (1, n) is a realization of the hidden Hamiltonian cycle model in Definition 1 with ground truth $C^* = (1, 2, ..., n, 1)$. By assumption, $\mathbb{P}\{\widehat{C}(G^e) = C^*\} \to 1$. Hence, $\mathbb{P}\{\widehat{H}^e = H^*\} \to 1$. Define two events

$$E_1 = \left\{ \hat{H}^e = H^* \right\}, \quad E_2 = \left\{ \hat{H}_{\rm ML} = H^* \right\}.$$

By assumption $\mathbb{P}{E_2} \to 1$. Hence, $\mathbb{P}{E_1 \cap E_2} \to 1$. On the event E_2 , H^* is the unique Hamiltonian path with the maximal weight. Thus, on the event $E_1 \cap E_2$,

$$\widehat{H}^e = H^* = \arg\min_{H \in \mathcal{H}} w(H)$$

where \mathcal{H} denotes the set of all possible Hamiltonian paths. Since $\widehat{H}^e \in \mathcal{H}$ for all $\widehat{H}_e \neq \emptyset$, it follows that on the event $E_1 \cap E_2$, $\widehat{H} = \widehat{H}^e = H^*$. Thus, $\mathbb{P}\{\widehat{H} = H^*\} \to 1$.

Conversely, by construction in Algorithm 2, the time complexity of $\widehat{C}(G)$ is $O(n^2T_n)$. The proof for $\mathbb{P}\{\widehat{C}=C^*\}\to 1$ is completely analogous as before and omitted.

As an immediate corollary of Theorem 4 and Theorem 1, we show the hidden Hamiltonian path can be exactly recovered with high probability in polynomial-time, provided that $\alpha_n - \log n \to +\infty$.

Corollary 2. If the optimal solution x^* of the F2F LP (6) is an adjacency vector of a Hamiltonian cycle in G, let $\widehat{C}(G)$ denote this Hamiltonian cycle; otherwise, let $\widehat{C}(G)$ denote any arbitrary Hamiltonian cycle in G. Let $\widehat{H}(G)$ denote the output of Algorithm 1 with input $\widehat{C}(G)$. If $\alpha_n - \log n \rightarrow +\infty$, i.e., (9) holds, then $\mathbb{P}\{\widehat{H} = H^*\} \rightarrow 1$ as $n \rightarrow \infty$.

8 Empirical Results

We empirically evaluate the performance of simple thresholding, greedy merging, belief propagation (BP) and F2F (LP) on both simulated and real datasets from DNA assembly.¹⁰ Of the four, the first three algorithms output integer-valued solutions. However, the optimal solution of F2F may contain half-integral entries and we randomly round them to either 0 or 1 with equal probability. We run BP for 10000 iterations for all the experiments in this section. All the plots in this section can be reproduced from the following repository - https://github.com/bagavi/HiC_LP.

8.1 Simulation

In this subsection, we consider a completely simulated setting of hidden Hamiltonian cycle graph consisting of n = 1000 vertices, whose edge weights are independently drawn from $P = \mathcal{N}(\mu, 1)$ if the edge on the Hamiltonian cycle C^* , and $Q = \mathcal{N}(0, 1)$ otherwise.

Fig. 15 plots the error probability $\mathbb{P}\left\{\widehat{C}\neq C^*\right\}$ of the output \widehat{C} for each of the four algorithms. As predicted by our theory (cf. (10) in the Gaussian case), the error probabilities of F2F and BP

¹⁰Since nearest-neighbor algorithm is very similar to simple thresholding and has the same performance guarantee as the simple thresholding, we do not include nearest-neighbor in the empirical study.

exhibit a phase transition around $\mu^2 = 4 \log n$. Moreover, the error probabilities of greedy merging and simple thresholding drop off to f0 around $\mu^2 = 6 \log n$ and $\mu^2 = 8 \log n$, respectively, which is consistent with our theoretical performance analysis (cf. Table 1).




Figure 15: Four algorithms are evaluated under simulated setting of hidden Hamiltonian cycle with n = 1000 vertices, with $P = \mathcal{N}(\mu, 1)$ and $Q = \mathcal{N}(0, 1)$, for μ^2 between 20 and 65. The y-axis is the probability of error, i.e., the probability that the hidden Hamiltonian cycle C^* is not exactly recovered. For each value of μ^2 , the error probability is calibrated using 50 independent runs. BP algorithm is run for 10000 iterations.

Fig. 16 plots the fraction of misclassified edges $|\hat{C}\Delta C^*|/n$, of the output \hat{C} for each of the four algorithms, where Δ denotes the set difference. In other words, the fraction of misclassified edges is equal to the number of edges in C^* that are absent in the \hat{C} plus the number of edges in the \hat{C} that are absent in C^* , normalized by n. From both Figures 15 and 16, we observe that when the optimal solution of F2F is integral, the solution of BP coincides with the F2F solution, as predicted by the result in [BBCZ11]. However, when the optimal solution of F2F has half-integral entries, the solution of BP often does not coincide with the F2F solution, and in fact has larger errors. The same phenomenon is also observed with real datasets as shown later.

8.2 DNA Assembly

As described in Section 1, ordering contigs using HiC reads (or Chicago reads, a variant of HiC reads) [POS⁺16, DBO⁺17] can be modeled as recovering hidden Hamiltonian cycle where the contigs are vertices of the graph, the hidden Hamiltonian cycle is the true ordering of the contigs on the genome, and the weights on the graph are the counts of the Hi-C reads (or Chicago) linking the contigs. In this section, we empirically compare the performance of simple thresholding, greedy merging, BP and F2F on datasets from the following three organisms:

- 1. Homosapiens Sequenced from a human in Iowa, USA. [POS⁺16].
- 2. Aedes Aegypti [DBO⁺17] A breed of mosquito which spreads dengue, zika fevers, and found throughout the world.
- 3. Gasterosteus Aculeatus [PSLW16] A variety of fish which is extensively studied to understand evolution and behavioral ecology.




Figure 16: This figure uses the same experiments in Fig. 15 to plot a different measure of error – fraction of misclassified edges $|\hat{C}\Delta C^*|/n$. For each value of μ^2 , the error is averaged over 50 independent trials. BP algorithm is run for 10000 iterations.

We choose these three datasets because the DNAs of these organisms have been sequenced and assembled using other (expensive) sequencing technologies, and that gives us the ground truth– the true contig ordering. The information on data sources and the detailed procedure is provided in Appendix H. We evaluate the performance of all the four algorithms under two scenarios – equal-length contigs and variable-length contigs.

8.2.1 DNA Assembly: Equal-length contigs

For real datasets, shotgun reads are assembled to produce unordered contigs of variable lengths. However, in this subsection we consider a controlled experimental setting. We partition a chromosome into contigs of equal length 100 kbps and randomly shuffle these contigs. We choose this length because the span of HiC reads is typically between 0 and 100 kbps. As a consequence, neighboring pairs of contigs tend to have many HiC reads between them, while non-neighboring pairs tend to have relatively fewer HiC reads, as postulated under our hidden Hamiltonian cycle model. Hence, we expect our algorithms – developed for recovering the hidden Hamiltonian cycle – to perform well for finding the true ordering of contigs under this setting.

Graph Construction Firstly, the chromosome is partitioned into *equal*-length contigs of length 100 kbps, and only the first 1000 contigs are kept and shuffled randomly. Secondly, the HiC reads are aligned to these 1000 contigs. Lastly, we construct graph G = ([n], E), where every vertex i corresponds to a contig and the weight on every edge $e = (i, j) \in E$ is the number of HiC reads between the two contigs corresponding to vertices i, j. For example, the weighted adjacency matrix of G for chromosome 1 of Homosapiens is shown in Fig. 2.

Performance evaluation The performance of all four algorithms is shown in Figures 17, 18, 19 for data from Homosapiens, Gasterous Aculeatus, and Aedes Agyepti, respectively. We observe that across all chromosomes and organisms, for a fixed coverage depth of HiC reads, F2F consistently has fewer errors than the two greedy algorithms – simple thresholding and greedy merging. Moreover,

BP also has fewer errors than the two greedy algorithms in most of the experiments. However, as mentioned previously, when the optimal solution of F2F has fractional entries, the BP solution no longer matches the optimal solution of F2F and has more errors.

8.2.2 DNA Assembly: Variable-length contigs

We turn to variable-length contigs generated by shotgun assembly softwares [CHS⁺11, BMK⁺08]. In this setting, HiC reads bias towards longer contigs. In other words, there tends to be more HiC reads linking two long contigs, which is not captured by our hidden Hamiltonian cycle model. Hence, our algorithms developed for recovering the hidden Hamiltonian cycle may not perform well for finding the true ordering of contigs. Nevertheless, we find that F2F and BP still consistently have few errors than the greedy methods, which are an integral part of the state-of-the art HiC assemblers [POS⁺16, DBO⁺17, BAP⁺13, GPK⁺17]. Thus we expect these assemblers can be possibly improved by replacing their greedy algorithm component by F2F or BP algorithm.

Graph Construction Firstly, the unordered contigs are obtained as output of short-gun reads assemblers such as Meraculous [CHS⁺11] and AllPaths [BMK⁺08]. These obtained contigs have variable lengths ranging from 10 kbps to 10 mbps (million base pairs). Secondly, the HiC reads are aligned to these contigs to obtain the raw number of HiC reads between each pair of contigs. Lastly, we construct graph G = ([n], E), where every vertex *i* correspond to a contig. The weight on every edge $e = (i, j) \in E$ is equal to $w_{ij} = b_i b_j N_{i,j}$ with $N_{i,j}$ denoting the *number* of HiC reads between the two contigs corresponding to vertices i, j. The values of b_i 's are chosen such that $\sum_i w_{i,j} = 1$ for all j, in order to remove the bias due to the variable contig length. These values of b_i 's are calculated using an iterative algorithm, ICE, described in 'Online methods' section from [IFM⁺12].

Performance evaluation The performance of all four algorithms is shown in Figures 20, 21, 22 for the data from Homosapiens, Gasterous Aculeatus, and Aedes Agyepti, respectively. Similar to the results in the setting with equal-length contigs, across all chromosomes and organisms, for a fixed coverage depth of HiC reads, F2F consistently has fewer errors than the two greedy algorithms. Moreover, BP also has fewer errors than greedy methods in most of the experiments. When the optimal solution of F2F has fractional entries, the BP solution does not match the optimal solution of F2F and has more errors.




Figure 17: Homosapiens [Equal length contigs]: The x-axis is the coverage depth of Hi-C reads i.e, expected number of Hi-C reads that include a base pair (nucleotide). The y-axis is the fraction of misclassified edges. For each value of coverage depth, the corresponding misclassification error is averaged over 25 independent runs of experiment.




Figure 18: Gasterosteus Aculeatus [Equal length contigs]: The *x*-axis is the coverage depth of Hi-C reads and the *y*-axis is the fraction of misclassified edges. For each value of coverage depth, the corresponding misclassification error is averaged over 25 independent runs of experiment.




Figure 19: Aedes Agyepti [Equal length contigs]: The x-axis is the coverage depth of Hi-C reads and the y-axis is the fraction of misclassified edges. For each value of coverage depth, the corresponding misclassification error is averaged over 25 independent runs of experiment.




Figure 20: Homosapiens [Variable length contigs]: The x-axis is the coverage depth of Hi-C reads i.e, expected number of HiC-reads that include a base pair (nucleotide). The y-axis is the fraction of misclassified edges. For each value of coverage depth, the corresponding misclassification error is averaged over 25 independent runs of experiment.




Figure 21: Gasterosteus Aculeatus [Variable length contigs]: The x-axis is the coverage depth of Hi-C reads and the y-axis is the fraction of misclassified edges. For each value of coverage depth, the corresponding misclassification error is averaged over 25 independent runs of experiment.




Figure 22: Aedes Agyepti [Variable length contigs]: The x-axis is the coverage depth of Hi-C reads and the y-axis is the fraction of misclassified edges. For each value of coverage depth, the corresponding misclassification error is averaged over 25 independent runs of experiment.

Appendix A Proof of Lemma 1

By assumption, G is connected and balanced. Therefore, every vertex in G has an even degree. Hence, there exists an Eulerian circuit $T = (v_0, v_1, \ldots, v_{m-1}, v_0)$, where m is the number of edges in G. Next we show that it can be modified, if necessary, to have alternating colors.

Suppose there exists a pair of two adjacent edges (v_{i-1}, v_i) and (v_i, v_{i+1}) such that they have the same color, say blue. Then by the balancedness, vertex v_i must be visited later on the circuit, say at step t, such that $v_t = v_i$, and (v_{t-1}, v_t) and (v_t, v_{t+1}) are both red. Consider the modified tour

$$T' = (v_0, \dots, v_{i-1}, v_i, v_{t-1}, v_{t-2}, \dots, v_{i+1}, v_t, v_{t+1}, \dots, v_{m-1}, v_0)$$

formed by reversing the trail between v_{i+1} and v_{t-1} . As a result, (v_{i-1}, v_i) and (v_i, v_{t-1}) have alternating colors. Similarly for (v_{i+1}, v_t) and (v_t, v_{t+1}) . The ordering of colors on the other adjacent edges is same as that in T. Therefore, the number of pairs of adjacent edges of the same color in the modified Eulerian circuit T' decreases by 2. Hence one can apply this procedure repeatedly for a finite number of rounds to transform T to an alternating Eulerian circuit.

Appendix B Large Deviation Bounds

Recall that X_i 's and Y_i 's are two independent sequences of random variables, where X_i 's are i.i.d. copies of dP/dQ under distribution P and Y_i 's are i.i.d. copies of dP/dQ under distribution Q. Let F denote the Legendre transform of the log moment generating function of $Y_1 - X_1$, i.e.,

$$F(x) = \sup_{\theta \ge 0} \left\{ \theta x - \psi_P(-\theta) - \psi_Q(\theta) \right\}.$$
(43)

Then from the Chernoff bound we have the following large deviation inequality: for any $x \in \mathbb{R}$ and any $\ell \in \mathbb{N}$,

$$\mathbb{P}\left\{\sum_{i=1}^{\ell} (Y_i - X_i) \ge \ell x\right\} \le \exp\left(-\ell F(x)\right).$$
(44)

Moreover, it is straightforward to check that

$$F(0) = -\psi_P(-1/2) - \psi_Q(1/2) = \alpha_n.$$
(45)

and F'(0) = 1/2.

Lemma 7. Fix $s, t \in \mathbb{N}$ and $u \in \mathbb{Z}$ such that $u \ge \min\{-s, -t/2\}$ and $v \in \mathbb{Z}$ such that $-2(s+u) \le v \le t$. Let

$$A \triangleq \sum_{i=1}^{s+u} Y_i - \sum_{i=1}^{s} X_i, \quad B \triangleq \sum_{i=1}^{t} Y_i - \sum_{i=1}^{t+2u} X_i.$$

Let $\{A_i : i \in [N]\}$ be identically distributed as A and $\{B_{ij} : i \in [N], j \in [M]\}$ as B, where

$$M \le n^{t-v} 4^{s+t+u-v/2}, \quad N \le (2n)^{s+u+v/2}.$$

Assume that for each i, A_i is independent of $\{B_{ij} : j \in [M]\}$. If

$$\alpha_n - \log n \ge \log 64,\tag{46}$$

then

$$\mathbb{P}\left\{\max_{i\in[N],j\in[M]} 2A_i + B_{ij} \ge 0\right\} \le 5\exp\left(-(\alpha - \log n)(s + t + u - v/2)/4\right).$$
(47)

Proof. We abbreviate α_n as α . Define the rate function for A and B as follows: for any $x \in \mathbb{R}$,

$$F_s(x) = \sup_{\theta \ge 0} \left\{ \theta x - (s+u)\psi_Q(\theta) - s\psi_P(-\theta) \right\},$$

$$G_t(x) = \sup_{\theta \ge 0} \left\{ \theta x - t\psi_Q(\theta) - (t+2u)\psi_P(-\theta) \right\}.$$

Then Chernoff's inequality gives the large deviation bound:

$$\mathbb{P}\left\{A \ge x\right\} \le \exp(-F_s(x)), \quad \mathbb{P}\left\{B \ge x\right\} \le \exp(-G_t(x)).$$

Next we provide a lower bound on the rate function. To explain the intuition, consider the easiest situation of u = 0, then we have $F_s(x) = G_s(x) = sF(\frac{x}{s})$. In the case of large s (which can be shown to be the most adversarial scenario), we have the approximation $F(\epsilon) = F(0) + F'(0)\epsilon + o(\epsilon)$ as $\epsilon \to 0$. Recall F'(0) = 1/2. Thus we have $F_s(x) = s\alpha + x/2 + o(1)$ as $s \to \infty$, which depends linearly on both x and s. This intuition can be made precise by the following non-asymptotic bound:

$$F_s(x) \ge s\alpha + x/2 - u\psi_Q(1/2) \tag{48}$$

$$G_t(x) \ge t\alpha + x/2 - 2u\psi_P(-1/2),$$
(49)

which simply follows from substituting $\theta = 1/2$ in the definition of F_s and G_t and applying (45).

Next we bound the probability in (47). By the union bound, we have for any $\tau \in \mathbb{R}$,

$$\mathbb{P}\left\{\max_{i\in[N],j\in[M]} 2A_i + B_{ij} \ge 0\right\} \le N \cdot \mathbb{P}\left\{A \ge \tau\right\} + NM \cdot \mathbb{P}\left\{B \ge -2A, A \le \tau\right\},\tag{50}$$

where $A \perp B$. Notice that here instead of directly taking the union bound over both i and j, we first single out the event $\{\max_{i \in [N]} A_i \geq \tau\}$. The key idea is to choose τ appropriately such that event $\{\max_{i \in [N]} A_i \geq \tau\}$ has low probability. In view of the assumption $N \leq (2n)^{s+u+v/2}$ and (48), a straightforward choice of τ is such that

$$s\alpha + \tau/2 - u\psi_Q(1/2) = (s + u + v/2)\alpha.$$

Here, we inflate τ slightly by setting

$$\tau \triangleq 2\left(u\psi_Q(1/2) + (u+v/2)\alpha + \epsilon(s+t+u-v/2)\log n\right),\tag{51}$$

where

$$\epsilon = \frac{\alpha - \log n}{4\log n}.\tag{52}$$

Then for the first term in (50):

$$N\mathbb{P} \{A \ge \tau\} \le N \exp(-F_s(\tau))$$

$$\stackrel{(48)}{\le} \exp(-(s\alpha + \tau/2 - u\psi_Q(1/2)) + \log(2n)(s + u + v/2))$$

$$\stackrel{(51)}{=} \exp(-(s + u + v/2)(\alpha - \log(2n)) - \epsilon(s + t + u - v/2)\log n)$$

$$\stackrel{(46)}{\le} \exp(-\epsilon(s + t + u - v/2)\log n).$$
(53)

For the second term in (50), fix some $\delta > 0$ to be chosen later and partition $(-\infty, \tau]$ into disjoint intervals $\bigcup_{i\geq 1}(\tau_i, \tau_{i-1}]$, where $\tau_i = \tau - i\delta$. Then by union bound,

$$\mathbb{P}\left\{B \ge -2A, A \le \tau\right\} \le \sum_{i \ge 1} \mathbb{P}\left\{B \ge -2A, \tau_i \le A \le \tau_{i-1}\right\}$$
$$\le \sum_{i \ge 1} \mathbb{P}\left\{B \ge -2\tau_{i-1}\right\} \mathbb{P}\left\{A \ge \tau_i\right\}$$
$$\le \sum_{i \ge 1} \exp\left(-F_s(\tau_i) - G_t(-2\tau_{i-1})\right)$$

where, in view of (48) and (49), the combined exponent satisfies

$$\begin{aligned} F_s(\tau_i) + G_t(-2\tau_{i-1}) \\ &\geq s\alpha + \tau_i/2 - u\psi_Q(1/2) + t\alpha - \tau_{i-1} - 2u\psi_P(-1/2) \\ &= (s+t)\alpha - \tau/2 - 2u\psi_P(-1/2) - u\psi_Q(1/2) + (i-2)\delta/2 \\ \\ \stackrel{(51)}{=} (s+t)\alpha - (u\psi_Q(1/2) + (u+v/2)\alpha + \epsilon(s+t+u-v/2)\log n) - 2u\psi_P(-1/2) - u\psi_Q(1/2) + (i-2)\delta/2 \\ \stackrel{(45)}{=} (s+t)\alpha + 2u\alpha - (u+v/2)\alpha - \epsilon(s+t+u-v/2)\log n + (i-2)\delta/2 \\ &= (s+t+u-v/2)(\alpha-\epsilon) + (i-2)\delta/2 \overset{(52)}{\leq} (s+t+u-v/2)(1+3\epsilon)\log n + (i-2)\delta/2. \end{aligned}$$

Combining the last two displayed equation and noting that $NM \leq (8n)^{s+t+u-v/2}$, we have

$$\begin{split} NM\mathbb{P}\left\{B \ge -2A, A \le \tau\right\} \le \exp\left(-(3\epsilon \log n - \log 8)(s + t + u - v/2)\right) \sum_{i \ge 1} \exp\left(-(i - 2)\delta/2\right) \\ \le \frac{e^{\delta/2}}{1 - e^{-\delta/2}} \exp\left(-\epsilon(s + t + u - v/2)\log n\right), \end{split}$$

where the last inequality holds because $2\epsilon \log n = (\alpha - \log n)/2 \ge \log 8$ in view of (46). Choosing $\delta = 2 \log 2$ that minimizes the right hand side of the last displayed equation, we get

$$NM\mathbb{P}\left\{B \ge -2A, A \le \tau\right\} \le 4\exp\left(-\epsilon(s+t+u-v/2)\log n\right).$$
(54)

Finally, we arrive at (47) by combining (50), (53), and (54), and (52).

Appendix C Proof of Lemma 6

Proof. In the Gaussian case, the log likelihood ratio is given by

$$\log \frac{\mathrm{d}P}{\mathrm{d}Q}(x) = x\mu - \mu^2/2.$$

Hence,

$$\psi_P(\theta) = \frac{1}{2}\theta(1+\theta)\mu^2$$

and $\alpha_n = \mu^2/4$. It follows that $\tau^* = \psi'_P(-1/2) = 0$. Define $\Phi^c(x) \triangleq \int_x^\infty \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$ as the tail probability of the standard normal. Then

$$\log \mathbb{P} \{ X \le \tau^* \} = \log \mathbb{P} \{ \mathcal{N}(\mu, 1) \le \mu/2 \} = \log \Phi^c(\mu/2) \ge -\mu^2/8 + O(\log \mu), \\ \log \mathbb{P} \{ Y \ge \tau^* \} = \log \mathbb{P} \{ \mathcal{N}(0, 1) \ge \mu/2 \} = \log \Phi^c(\mu/2) \ge -\mu^2/8 + O(\log \mu).$$

where the last inequality holds because $\Phi^c(x) \ge \frac{x}{1+x^2}e^{-x^2/2}$. Hence Assumption 1 holds in the Gaussian case.

In the Poisson case, for integer $k \ge 0$,

$$\log \frac{\mathrm{d}P}{\mathrm{d}Q}(k) = k \log(\lambda/\mu) - (\lambda - \mu).$$

Hence,

$$\psi_P(\theta) = \lambda \left((\lambda/\mu)^{\theta} - 1 \right) - \theta(\lambda - \mu).$$

and

$$\alpha_n = \left(\sqrt{\lambda} - \sqrt{\mu}\right)^2, \quad \tau^* = \sqrt{\lambda\mu} \log(\lambda/\mu) - (\lambda - \mu).$$

Let $\eta = \sqrt{\lambda \mu}$. Then

$$\log \mathbb{P} \{ X \le \tau^* \} = \log \mathbb{P} \{ \operatorname{Pois}(\lambda) \le \eta \}$$

$$\geq \log \mathbb{P} \{ \operatorname{Pois}(\lambda) = \lfloor \eta \rfloor \}$$

$$= \lfloor \eta \rfloor \log \lambda - \lambda - \log (\lfloor \eta \rfloor!)$$

$$= \eta \log \lambda - \lambda - \eta \log(\eta/e) + O(\log(\lambda\mu)) + O(1)$$

$$= -\lambda + \eta + \frac{1}{2} \eta \log (\lambda/\mu) + o(\alpha) + o(\log n),$$

where we used the fact that $\log n! \le n \log(n/e) + (1/2) \log n + 1$ and the assumption that $\log(\lambda \mu) = o(\alpha) + o(\log n)$. Similarly,

$$\log \mathbb{P} \{ Y \ge \tau^* \} = \log \mathbb{P} \{ \operatorname{Pois}(\mu) \ge \eta \}$$

$$\geq \log \mathbb{P} \{ \operatorname{Pois}(\mu) = \lceil \eta \rceil \}$$

$$= \lceil \eta \rceil \log \mu - \mu - \log \left(\lceil \eta \rceil ! \right)$$

$$= \eta \log \mu - \mu - \eta \log(\eta/e) + +O(\log(\lambda\mu)) + O(1)$$

$$= -\mu + \eta - \frac{1}{2}\eta \log(\lambda/\mu) + o(\alpha) + o(\log n).$$

Thus Assumption 1 holds in the Poisson case.

In the Bernoulli case, let $\bar{p} = 1 = p$ and $\bar{q} = 1 - q$. Then X and Y takes the value of either $\log(p/q)$ or $\log(\bar{p}/\bar{q})$. Hence,

$$\log \mathbb{P}\left\{X \le \log(p/q)\right\} + \log \mathbb{P}\left\{Y \ge \log(p/q)\right\} = \log 1 + \log q = \log q$$
$$\log \mathbb{P}\left\{X \le \log(\bar{p}/\bar{q})\right\} + \log \mathbb{P}\left\{Y \ge \log(\bar{p}/\bar{q})\right\} = \log \bar{p} + \log 1 = \log \bar{p}.$$

Hence, to verify Assumption 1, it suffices to show

$$\max\{\log \bar{p}, \log q\} \ge -(1+o(1))\alpha + o(\log n).$$
(55)

Note that

$$-\alpha = 2\log\left(\sqrt{pq} + \sqrt{\bar{p}\bar{q}}\right) \le \max\{\log(pq), \log(\bar{p}\bar{q})\} + \log 4 \le \max\{\log q, \log \bar{p}\} + \log 4.$$

Hence, the desired (55) holds.

Appendix D Analysis for the Suboptimal Dual Certificate (15)

We first present an auxiliary lemma. Recall that

$$\beta_n = -\frac{3}{2} \log \int (\mathrm{d}P)^{2/3} (\mathrm{d}Q)^{1/3}$$

Lemma 8.

$$\mathbb{P}\left\{2Y - X_1 - X_2 \ge 0\right\} \le \exp\left(-2\beta_n\right).$$

Proof. It follows from the Chernoff bound that

$$\mathbb{P}\left\{2Y - X_1 - X_2 \ge 0\right\} \le \exp\left(-\sup_{\theta \ge 0} \left\{\psi_Q(2\theta) + 2\psi_P(-\theta)\right\}\right).$$

It turns out that the supremum is achieved at $\theta = 1/3$ so that $\psi'_Q(2/3) = \psi'_P(-1/3)$ and

$$\psi_Q(2/3) + 2\psi_P(-1/3) = -3\log \int (\mathrm{d}P)^{2/3} (\mathrm{d}Q)^{1/3} = 2\beta_n.$$

Theorem 5. If

$$\beta_n - \log n \to \infty$$

then the optimal solution of F2F (6) satisfies $\min_{x^*} \mathbb{P} \{ \widehat{x}_{F2F} = x^* \} \to 1 \text{ as } n \to \infty.$

Proof. We use the standard dual certificate argument. The Lagrangian function is given by

$$L(x, b, h, u) = -\langle w, x \rangle - \langle b, x \rangle + \langle h, x - \mathbf{1} \rangle + \sum_{i} u_{i} \left(x \left(\delta(i) \right) - 2 \right) + \sum_{i} u_{i} \left(x \left(\delta(i) \right) + \sum_{i} u_{i} \left(x \left(\delta(i) \right) - 2 \right) + \sum_{i} u_{i} \left(x \left(\delta(i) \right) + \sum_{i} u$$

where **1** denotes a $\binom{n}{2}$ -dimensional all-one vector; $b \ge 0$, $h \ge 0$, and $u \in \mathbb{R}^n$ are dual variables. To show that x^* is an optimal solution to F2F (6), it suffices to construct the dual variables to satisfy the following KKT conditions: for every edge e = (i, j),

$$-w_e - b_e + h_e + u_i + u_j = 0, (56)$$

$$h_e(x_e^* - 1) = 0, \tag{57}$$

$$b_e x_e^* = 0, \tag{58}$$

where (56) is the first-order condition, and (57) and (58) are complementary slackness conditions.

It follows from (57) that if $x_e^* = 0$, then $h_e = 0$. Similarly, it follows from (58) that if $x_e^* = 1$, then $b_e = 0$. Thus, in view of (56), for every edge e = (i, j), we must have

$$h_e = w_e - u_i - u_j$$
 if $x_e^* = 1$,
 $b_e = -w_e + u_i + u_j$ if $x_e^* = 0$,

Then to satisfy all the KKT conditions, it reduces to constructing u such that that for every e = (i, j),

$$u_i + u_j \le w_e \quad \text{if } x_e^* = 1, \tag{59}$$

$$u_i + u_j \ge w_e \quad \text{if } x_e^* = 0. \tag{60}$$

Let

$$u_i = \frac{1}{2} \min \{ w_e : e \in \delta(i), x_e^* = 1 \}.$$

Then for every e = (i, j) such that $x_e^* = 1$, $u_i \leq w_e/2$ and $u_j \leq w_e/2$, and thus $u_i + u_j \leq w_e$. Define event

$$\mathcal{E} = \bigcup_{e=(i,j):x_e^*=0} \{ w_e - u_i/2 - u_j/2 \ge 0 \}$$

We are left to show that as $n \to \infty$,

$$\mathbb{P}\left\{\mathcal{E}\right\} \to 0$$

For each e = (i, j) such that $x_e^* = 0$, by the union bound,

$$\mathbb{P}\left\{w_{e} - u_{i}/2 - u_{j}/2 \ge 0\right\} = \mathbb{P}\left\{\bigcup_{f \in \delta(i): x_{f}^{*}=1} \bigcup_{g \in \delta(j): x_{g}^{*}=1} \left\{2w_{e} - w_{f} - w_{g} \ge 0\right\}\right\}$$

$$\leq \sum_{f \in \delta(i): x_{f}^{*}=1} \sum_{g \in \delta(j): x_{g}^{*}=1} \mathbb{P}\left\{2w_{e} - w_{f} - w_{g} \ge 0\right\}$$

$$= 4\mathbb{P}\left\{2Y - X_{1} - X_{2} \ge 0\right\}$$

$$\leq 4\exp\left(-2\beta_{n}\right),$$

where the last inequality holds due to Lemma 8. Hence, by the union bound

$$\mathbb{P}\left\{\mathcal{E}\right\} \leq \sum_{e=(i,j):x_e^*=0} \mathbb{P}\left\{w_e - u_i/2 - u_j/2 \geq 0\right\}$$
$$\leq 4n^2 \exp\left(-2\beta_n\right) = o(1),$$

where the last inequality holds due to the assumption that $\beta_n - \log n \to +\infty$.

Finally, we show that x^* is the unique optimal solution to F2F on event \mathcal{E}^c . Let \tilde{x} is another optimal solution to F2F. Since KKT conditions are also necessary, it follows that \tilde{x} must satisfy $\langle \tilde{x}, b \rangle = 0$. On event \mathcal{E}^c , for every e = (i, j) such that $x_e^* = 0$, $b_e = -w_e + u_i + u_j > 0$ and hence $\tilde{x}_e = 0$. Moreover, since \tilde{x} is feasible, it follows that $\tilde{x}_e = 1$ for every e such that $x_e^* = 1$ and thus $\tilde{x} = x$.

Appendix E Greedy Methods

We first consider the simple thresholding algorithm.

| Algorithm 3 Simple Thresholding |
|--|
| 1: For every vertex keep the two edges with the largest weights and delete the rest of $n-3$ edges |
| with ties broken arbitrarily. 2. Output the resulting graph \hat{C}_{max} |

2: Output the resulting graph C_{TH} .

The following theorem provides an sufficient condition under which the simple thresholding algorithm to attain exact recovery.

Theorem 6 (Simple Thresholding). If

$$\alpha_n - 2\log n \to +\infty,\tag{61}$$

then $\min_{C^*} \mathbb{P}\left\{\widehat{C}_{\mathrm{TH}} = C^*\right\} \to 1 \text{ as } n \to \infty.$

Proof. For any vertex i, define event

$$\mathcal{H}_i = \{\min\{w_e : e \in \delta(i), x_e^* = 1\} > \max\{w_e : e \in \delta(i), x_e^* = 0\}\}.$$

Then by union bound and Chernoff's inequality (44),

$$\mathbb{P}\left\{\mathcal{H}_{i}^{c}\right\} \leq 2(n-3)\mathbb{P}\left\{X \leq Y\right\} \leq 2ne^{-\alpha}.$$

Let $\mathcal{H} = \bigcap_{i=1}^{n} \mathcal{H}_i$. Hence, applying union bound, we get that

$$\mathbb{P}\left\{\mathcal{H}^{c}\right\} \leq \sum_{i=1}^{n} \mathbb{P}\left\{\mathcal{H}^{c}_{i}\right\} \leq 2n^{2}e^{-\alpha} = o(1),$$

where the last equality holds in view of the assumption (61). The proof is complete by noting that $\mathcal{H} \subset \{\widehat{C}_{\mathrm{TH}} = C^*\}.$

Next we consider the following nearest neighbor algorithm.

Algorithm 4 Nearest Neighbor

- 1: Start on an arbitrary vertex as current vertex.
- 2: Find out the edge with the largest weight connecting current vertex and an unvisited vertex v.
- 3: Set current vertex to v and mark v as visited.
- 4: If all the vertices in domain are visited, then output Hamiltonian cycle $\widehat{C}_{NN} = (v_1, \ldots, v_n, v_1)$, where v_1, \ldots, v_n is the sequence of visited vertices; otherwise go to Step 2.

The following theorem provides an sufficient condition under which the nearest neighbor algorithm to attain exact recovery.

Theorem 7 (Nearest Neighbor). If

$$\alpha_n - 2\log n \to +\infty,\tag{62}$$

then $\min_{C^*} \mathbb{P}\left\{\widehat{C}_{NN} = C^*\right\} \to 1 \text{ as } n \to \infty.$

Proof. At step t = 1, ..., n - 2, let V_t denote the set of visited vertices and v_t denote the current vertex, and define event

$$\mathcal{H}_t = \left\{ \max\{w_{v_t,j} : j \notin V_t, \ x_{v_t,j}^* = 1\} > \max\{w_{v_t,j} : j \notin V_t, \ x_{v_t,j}^* = 0\} \right\}.$$

Define event $\mathcal{H} = \bigcap_{t=1}^{n-2} \mathcal{H}_t$. Since on event \mathcal{H}_t , v_{t+1} must be neighbor of v_t in the true Hamiltonian cycle C^* , it follows that $\{\widehat{C}_{NN} = C^*\} = \mathcal{H}$. Hence, the prove the theorem, it suffices to show $\mathbb{P}\{\mathcal{H}\} \to 1$. Using the property of conditional probability, we have

$$\mathbb{P}\left\{\mathcal{H}\right\} = \mathbb{P}\left\{\mathcal{H}_{1}\right\} \prod_{t=2}^{n-2} \mathbb{P}\left\{\mathcal{H}_{t} | \mathcal{H}_{t-1}, \dots, \mathcal{H}_{1}\right\}.$$

Note that

$$\mathbb{P}\left\{\mathcal{H}_{1}^{c}\right\} = \mathbb{P}\left\{\max\{X_{1}, X_{2}\} \leq \max_{1 \leq j \leq n-3} Y_{j}\right\}$$
$$\leq \sum_{j=1}^{n-3} \mathbb{P}\left\{\max\{X_{1}, X_{2}\} \leq Y_{j}\right\}$$
$$\leq (n-3)\mathbb{P}\left\{X \leq Y\right\} \leq (n-3)e^{-\alpha}.$$

Fix any $2 \le t \le n-2$ and condition on $(\mathcal{H}_{t-1}, \ldots, \mathcal{H}_1)$. Then there exists exactly one one unvisited neighbor of v_t in the true Hamiltonian cycle C^* . Moreover, since the edges connecting current vertex v_t and all unvisited vertices are independent of $(\mathcal{H}_{t-1}, \ldots, \mathcal{H}_1)$, it follows that

$$\mathbb{P}\left\{\mathcal{H}_{t}^{c}|\mathcal{H}_{t-1},\ldots,\mathcal{H}_{1}\right\} = \mathbb{P}\left\{X \leq \max_{1 \leq j \leq n-1-t} Y_{j}\right\} \leq (n-1-t)\mathbb{P}\left\{X \leq Y\right\} \leq (n-1-t)e^{-\alpha}$$

Combining the last three displayed equations yields that

$$\mathbb{P}\left\{\mathcal{H}\right\} = \left(1 - (n-3)e^{-\alpha}\right) \prod_{t=2}^{n-2} \left(1 - (n-1-t)e^{-\alpha}\right)$$
$$\geq 1 - (n-3)e^{-\alpha} - \sum_{t=2}^{n-2} (n-1-t)e^{-\alpha}$$
$$= 1 - (n-3)e^{-\alpha} - \frac{(n-2)(n-3)}{2}e^{-\alpha} \to 1,$$

where the first inequality holds by iteratively applying $(1-x)(1-y) \ge 1-x-y$ for $x, y \ge 0$; the last limit holds in view of assumption (62).

Finally, we consider a greedy merging algorithm, which improves over the previous two greedy algorithms.

Algorithm 5 Greedy Merging

- 1: Start with n isolated vertices.
- 2: Among all vertices with degree strictly less than 2, select two of them, say i and j, with the largest weight w_{ij} and then connect them.
- 3: If all vertices have degree 2, output the resulting graph \widehat{C}_{GM} ; otherwise go to Step 2.

Theorem 8 (Greedy Merging). If

$$\beta_n - \log n \to +\infty,$$
 (63)

then $\min_{C^*} \mathbb{P}\left\{\widehat{C}_{\mathrm{GM}} = C^*\right\} \to 1 \text{ as } n \to \infty.$

Proof. For all edge e = (i, j) such that $x_e^* = 0$, define an event

$$\mathcal{H}_e = \left\{ \min_{f \in \delta(i): x_f^* = 1} w_f > w_e \right\} \cup \left\{ \min_{g \in \delta(j): x_g^* = 1} > w_e \right\}$$

We claim that on event \mathcal{H}_e , edge e will not be added in \widehat{C} . Suppose not. Then by the definition of \mathcal{H}_e and the greedy merging algorithm, either i or j must already have degree 2 before adding edge e and hence e cannot be added. Define $\mathcal{H} = \bigcap_{e:x_e^*=0} \mathcal{H}_e$. It follows that $\{\widehat{C}_{GM} = C^*\} = \mathcal{H}$. Hence, to prove the theorem, it suffices to show $\mathbb{P}\{\mathcal{H}^c\} = o(1)$. Note that

$$\mathcal{H}_e^c = \left\{ \min_{f \in \delta(i): x_f^* = 1} w_f \le w_e \right\} \cap \left\{ \min_{g \in \delta(j): x_g^* = 1} \le w_e \right\}$$
$$= \bigcup_{f \in \delta(i): x_f^* = 1} \bigcup_{g \in \delta(j): x_g^* = 1} \left\{ w_f \le w_e, w_g \le w_e \right\}.$$

Hence, by the union bound,

$$\mathbb{P}\left\{\mathcal{H}_{e}^{c}\right\} \leq \sum_{f \in \delta(i): x_{f}^{*}=1} \sum_{g \in \delta(j): x_{g}^{*}=1} \mathbb{P}\left\{w_{f} \leq w_{e}, w_{g} \leq w_{e}\right\}$$
$$\leq 4\mathbb{P}\left\{X_{1} \leq Y, X_{2} \leq Y\right\}$$
$$\leq 4\mathbb{P}\left\{X_{1} + X_{2} \leq 2Y\right\} \leq 4e^{-2\beta},$$

where the last inequality follows from Lemma 8. Applying union bound again, we get that

$$\mathbb{P}\left\{\mathcal{H}^{c}\right\} \leq \sum_{e:x_{e}^{*}=0} \mathbb{P}\left\{\mathcal{H}_{e}^{c}\right\} \leq 4n^{2}e^{-2\beta} = o(1),$$

where the last inequality holds due to assumption (63).

Appendix F Exact Recovery under the Partial Observation Model

Consider the hidden Hamiltonian cycle recovery problem under a partial observation model, where the weight of every edge in G is erased independently with probability η_n . This is equivalent to recovering the hidden Hamiltonian cycle from G' generated under hidden Hamiltonian cycle model in Definition 1 with $P'_n = \eta_n \delta_* + (1 - \eta_n)P_n$ and $Q'_n = \eta_n \delta_* + (1 - \eta_n)Q_n$, where * is any element outside the support of P_n and Q_n denoting the erasure. In this case, the log-likelihood weight matrix is the same as before with each erasure replaced with 0. Straightforward calculation shows that

$$\alpha_n' \triangleq -2\log \int \sqrt{dP_n' dQ_n'} = -2\log\left(\eta_n + (1-\eta_n)\exp(-\alpha_n/2)\right).$$

Note that $\alpha'_n - \log n \to +\infty$ if and only if $\eta_n = o(1/\sqrt{n})$ and $\alpha_n - \log n \to \infty$. Hence, Theorem 1 implies that if $\eta_n = o(1/\sqrt{n})$ and $\alpha_n - \log n \to +\infty$, then the F2F LP recovers the hidden Hamiltonian cycle with high probability.

For the converse, suppose MLE recovers the hidden Hamiltonian cycle with high probability. Then as shown in Fig. 14, with high probability there is no 4-cycle (i, i + 1, j + 1, j, i) such that the weights on all 4 edges are *, because otherwise the likelihood function has at least two maximizers which renders MLE to fail with probability at least 1/2. The probability that all 4 edges in (i, i + 1, j + 1, j, i) have weights * is η_n^4 , and there are roughly $n^2/2$ different such 4-cycles. If the weights of all such 4-cycles were mutually independent, then it would immediately follow that $\eta_n^4 n^2 \to 0$, i.e., $\eta_n = o(1/\sqrt{n})$. However, the weights of all such 4-cycles are dependent because of the multiple appearances of (i, i + 1) edges. The dependency can be dealt with in the same way as in the proof of Theorem 2. Specifically, it directly follows from (42) that for any $\tau \in \mathbb{R}$,

$$e^{-\mathbb{P}\{Y \ge \tau\}^2 \binom{n\mathbb{P}\{X \le \tau\}/4}{2}} + e^{-n\mathbb{P}\{X \le \tau\}/8} \ge 1 + o(1).$$

Setting $\tau = 0$ and noting that $\mathbb{P}\{X \le \tau\} \ge \mathbb{P}\{X = 0\} \ge \eta_n$ and $\mathbb{P}\{Y \ge \tau\} \ge \mathbb{P}\{Y \ge 0\} \ge \eta_n$, we get that

$$e^{-\eta_n^2 \binom{n\eta_n/4}{2}} + e^{-n\eta_n/8} \ge 1 + o(1).$$

It follows that $\eta_n = o(1/\sqrt{n})$ is necessary for exact recovery.

Appendix G Time Complexity of the Max-Product BP algorithm

In this appendix we bound the number of iterations t_f of Max-Product BP algorithm mentioned in the end of Section 2 in both the Gaussian and the Poisson model under the assumption (9) that ensures the F2F LP succeeds. Recall that $t_f = \lceil 2nw^*/\epsilon \rceil$, where w^* is the weight of the optimal 2-factor and ϵ is the difference between the weight of the optimal 2-factor and the second largest weight of *b*-factors.

First we show that $\epsilon \to \infty$, which holds for any weight distribution. Following the analysis of the 2F ILP in Section 4.2, we prove a slightly stronger separation between the weight of the true Hamiltonian cycle and the weight of any other 2-factor. Let $\rho_n \triangleq \alpha_n - \log(2n)$, which tends to infinity. Then it follows from the Chernoff's inequality that

$$\mathbb{P}\left\{w(B) \ge -\rho_n\right\} = \mathbb{P}\left\{\sum_{i=1}^{\ell/2} Y_i - \sum_{i=1}^{\ell/2} X_i \ge -\rho_n\right\} \le \exp\left(-(\ell/2)F(-2\rho_n/\ell)\right)$$
$$\le \exp\left(-\alpha_n\ell/2 + \rho_n\right).$$

where F is given in (43) and the last inequality follows because by convexity of F, $F(x) \ge F(0) + F'(0)x = \alpha_n + x/2$. Then following the exactly same argument as in Section 4.2, we get that

$$\mathbb{P}\left\{\max_{B\in\mathcal{B}^*} w(B) \ge -\rho_n\right\} \le e^{\rho_n} \frac{\exp\left\{-2\left(\alpha_n - \log(2n)\right)\right\}}{1 - \exp\left\{-\left(\alpha_n - \log(2n)\right)/2\right\}} \to 0.$$

It follows that $\epsilon \geq \rho_n \to +\infty$ with high probability.

Finally, for the Gaussian and Poisson weight distribution (cf. (10)), the total weight w^* of the true Hamiltonian cycle is distributed as $N(n\mu, n)$ and $Pois(n\lambda)$, respectively. At the information-theoretic limit, we have $\mu = \Theta(\sqrt{\log n})$ and $\lambda = \Theta(\log n)$. Hence $w^* = O_P(n \log n)$ under both cases.

Appendix H Data Source and Preparation for DNA Assembly

H.1 Data Source

Table 2 contains links to all the datasets used for experiments in Section 8. Each organism requires the following three datasets:

- 1. Contigs Fasta file of unordered contigs generated from denovo assembly softwares such as Meraculous [CHS⁺11], AllPaths [BMK⁺08] etc.
- 2. HiC/Chicago Fastq file containing HiC/Chicago reads obtained from the DNA.
- 3. Chromosome Chromosomes pre-assembled using other expensive data. These chromosome comprise of the above contigs in the correct order and thus this dataset gives us the ground truth ordering of the contigs.

Goal: Order the contigs using the HiC/Chicago reads and check the accuracy of the contig ordering using the true answer obtained from pre-assembled chromosome.

| Organism | Contigs | HiC/Chicago | Chromosome |
|-------------------------------------|------------|-------------|--------------|
| HomoSapiens [POS ⁺ 16] | SRR2911057 | GL582980.1 | GRCh37 |
| Gasterosteus Aculeatus [PSLW16] | SRR4007584 | dryad.h7h32 | GCA000180675 |
| Aedes Agyepti [DBO ⁺ 17] | SRX2618715 | txid7159 | GCA000180675 |

Table 2: Datasets.

H.2 Procedure

For each chromosome of an organism, the following steps were performed to generate the contigs graph G and to evaluate the performance of F2F, BP, greedy merging and simple thresholding.

- 1. Align the HiC/Chicago reads to the contigs using HiC-Pro [SVL⁺15]. This gives us the raw number of HiC reads between each pair of contigs.
- 2. Construct the graph G using the data obtained in the previous step.
- 3. On graph G, run F2F, BP, greedy merging and simple thresholding algorithms, with outputs denoted by \tilde{C}_{F2F} , \tilde{C}_{BP} , \tilde{C}_{GM} , and \tilde{C}_{TH} , respectively.
- 4. Align the contigs on the pre-assembled chromosome to obtain the true ordering of the contigs to obtain the ground truth Hamiltonian path C^* .
- 5. Evaluate the misclassification error of the outputs of the four algorithms with respect to the ground truth path C^* .

Acknowledgment

D. Tse and J. Xu would like to thank the support of the Simons Institute, where this collaboration began. J. Xu would also like to thank Mohit Tawarmalani for inspiring discussions on fractional 2-factor LP relaxation of TSP. Y. Wu is grateful to Dan Spielman for helpful comments and to David Pollard and Dana Yang for an extremely thorough reading of the manuscript and various corrections.

V. Bagaria and D. Tse are supported by the Center for Science of Information, an NSF Science and Technology Center, under grant agreement CCF-0939370, as well as the National Human Genome Research Institute of the National Institutes of Health under award number R01HG008164. J. Ding was supported in part by the NSF Grant DMS-1757479 and an Alfred Sloan fellowship. Y. Wu was supported in part by the NSF Grant IIS-1447879, CCF-1527105, an NSF CAREER award CCF-1651588, and an Alfred Sloan fellowship. J. Xu was supported in part by a Simons-Berkeley Research Fellowship and the NSF Grant CCF-1755960.

References

- [ABH98] Jonathan E. Atkins, Erik G. Boman, and Bruce Hendrickson. A spectral algorithm for seriation and the consecutive ones problem. *SIAM Journal on Computing*, 28(1):297–310, 1998.
- [ABH16] E. Abbe, A. S. Bandeira, and G. Hall. Exact recovery in the stochastic block model. *IEEE Transactions on Information Theory*, 62(1):471–487, 2016. arXiv 1405.3267.

- [ABKK15] N. Agarwal, A. S. Bandeira, K. Koiliaris, and A. Kolla. Multisection in the stochastic block model using semidefinite programming. arXiv 1507.02323, July 2015.
- [AKS98] N. Alon, M. Krivelevich, and B. Sudakov. Finding a large hidden clique in a random graph. *Random Structures and Algorithms*, 13(3-4):457–466, 1998.
- [AS15] Emmanuel Abbe and Colin Sandon. Community detection in general stochastic block models: fundamental limits and efficient recovery algorithms. In 2015 IEEE 56th Annual Symposium on Foundations of Computer Science (FOCS), pages 670–688, 2015. arXiv 1503.00609.
- [Bal65] Michel Louis Balinski. Integer programming: methods, uses, computations. *Management Science*, 12(3):253–313, 1965.
- [Ban15] A.S. Bandeira. Random Laplacian matrices and convex relaxations. arXiv 1504.03987, April, 2015.
- [BAP⁺13] Joshua N Burton, Andrew Adey, Rupali P Patwardhan, Ruolan Qiu, Jacob O Kitzman, and Jay Shendure. Chromosome-scale scaffolding of de novo genome assemblies based on chromatin interactions. *Nature biotechnology*, 31(12):1119, 2013.
- [BBCZ11] Mohsen Bayati, Christian Borgs, Jennifer Chayes, and Riccardo Zecchina. Belief propagation for weighted b-matchings on arbitrary graphs and its relation to linear programs with integer solutions. *SIAM Journal on Discrete Mathematics*, 25(2):989– 1011, 2011.
- [BC99] Sylvia Boyd and Robert Carr. A new bound for the ratio between the 2matching problem and its linear programming relaxation. *Mathematical programming*, 86(3):499–514, 1999.
- [BFS94]Andrei Z Broder, Alan M Frieze, and Eli Shamir. Finding hidden hamiltonian cycles.
Random Structures & Algorithms, 5(3):395–410, 1994.
- [BHK⁺16] Boaz Barak, Samuel B. Hopkins, Jonathan A. Kelner, Pravesh Kothari, Ankur Moitra, and Aaron Potechin. A nearly tight sum-of-squares lower bound for the planted clique problem. In *IEEE 57th Annual Symposium on Foundations of Computer Science, FOCS*, pages 428–437, 2016.
- [BLM15] C. Bordenave, M. Lelarge, and L. Massoulié. Non-backtracking spectrum of random graphs: community detection and non-regular Ramanujan graphs. In 2015 IEEE 56th Annual Symposium on Foundations of Computer Science (FOCS), pages 1347– 1357, 2015. arXiv 1501.06087.
- [BMK⁺08] Jonathan Butler, Iain MacCallum, Michael Kleber, Ilya A Shlyakhter, Matthew K Belmonte, Eric S Lander, Chad Nusbaum, and David B Jaffe. Allpaths: de novo assembly of whole-genome shotgun microreads. Genome research, 18(5):810–820, 2008.
- [CČKV99] Dragoš Cvetković, Mirjana Čangalović, and Vera Kovačević-Vujčić. Semidefinite programming methods for the symmetric traveling salesman problem. In International Conference on Integer Programming and Combinatorial Optimization, pages 126–136. Springer, 1999.

- [CCMW11] Kun Chen, Kehui Chen, Hans-Georg Muller, and Jane-Ling Wang. Stringing highdimensional data for functional analysis. Journal of the American Statistical Association, 106(493):275–284, 2011.
- [CHS⁺11] Jarrod A Chapman, Isaac Ho, Sirisha Sunkara, Shujun Luo, Gary P Schroth, and Daniel S Rokhsar. Meraculous: de novo genome assembly with short paired-end reads. *PloS one*, 6(8):e23501, 2011.
- [CK01] A. Condon and R. M. Karp. Algorithms for graph partitioning on the planted partition model. *Random Struct. Algorithms*, 18(2):116–140, Mar. 2001.
- [CLR17] Tony Cai, Tengyuan Liang, and Alexander Rakhlin. On detection and structural reconstruction of small-world random networks. *IEEE Transactions on Network Science and Engineering*, 2017.
- [CT12] Eden Chlamtac and Madhur Tulsiani. Convex relaxations and integrality gaps. In *Handbook on semidefinite, conic and polynomial optimization*, pages 139–169. Springer, 2012.
- [DBO⁺17] Olga Dudchenko, Sanjit S Batra, Arina D Omer, Sarah K Nyquist, Marie Hoeger, Neva C Durand, Muhammad S Shamim, Ido Machol, Eric S Lander, Aviva Presser Aiden, et al. De novo assembly of the aedes aegypti genome using hi-c yields chromosome-length scaffolds. *Science*, 356(6333):92–95, 2017.
- [DFJ54] George Dantzig, Ray Fulkerson, and Selmer Johnson. Solution of a large-scale traveling-salesman problem. Journal of the operations research society of America, 2(4):393–410, 1954.
- [DKPS08] Etienne De Klerk, Dmitrii V Pasechnik, and Renata Sotirov. On semidefinite programming relaxations of the traveling salesman problem. *SIAM Journal on Optimization*, 19(4):1559–1573, 2008.
- [DM15] Y. Deshpande and A. Montanari. Improved sum-of-squares lower bounds for hidden clique and hidden submatrix problems. In *Proceedings of COLT 2015*, pages 523–562, June 2015.
- [Edm65a] Jack Edmonds. Maximum matching and a polyhedron with 0, 1-vertices. Journal of Research of the National Bureau of Standards B, 69(125-130):55–56, 1965.
- [Edm65b] Jack Edmonds. Paths, trees, and flowers. *Canadian Journal of mathematics*, 17(3):449–467, 1965.
- [FJBd13] Fajwel Fogel, Rodolphe Jenatton, Francis Bach, and Alexandre d'Aspremont. Convex relaxations for permutation problems. In Advances in Neural Information Processing Systems, pages 1016–1024, 2013.
- [GH87] Martin Grötschel and Olaf Holland. A cutting plane algorithm for minimum perfect 2-matchings. *Computing*, 39(4):327–344, Dec 1987.
- [GPK⁺17] Jay Ghurye, Mihai Pop, Sergey Koren, Derek Bickhart, and Chen-Shan Chin. Scaffolding of long read assemblies using long range contact information. *BMC genomics*, 18(1):527, 2017.

- [HK70] Michael Held and Richard M Karp. The traveling-salesman problem and minimum spanning trees. *Operations Research*, 18(6):1138–1162, 1970.
- [HWX16a] B. Hajek, Y. Wu, and J. Xu. Achieving exact cluster recovery threshold via semidefinite programming. *IEEE Transactions on Information Theory*, 62(5):2788–2797, May 2016. (arXiv 1412.6156 Nov. 2014).
- [HWX16b] B. Hajek, Y. Wu, and J. Xu. Achieving exact cluster recovery threshold via semidefinite programming: Extensions. *IEEE Trans. on Information Theory*, 62(10):5918– 5937, Oct. 2016.
- [HWX16c] Bruce Hajek, Yihong Wu, and Jiaming Xu. Semidefinite programs for exact recovery of a hidden community. In *Proceedings of Conference on Learning Theory (COLT)*, pages 1051–1095, New York, NY, Jun 2016. arXiv:1602.06410.
- [IFM⁺12] Maxim Imakaev, Geoffrey Fudenberg, Rachel Patton McCord, Natalia Naumova, Anton Goloborodko, Bryan R Lajoie, Job Dekker, and Leonid A Mirny. Iterative correction of hi-c data reveals hallmarks of chromosome organization. *Nature methods*, 9(10):999, 2012.
- [Jer92] Mark Jerrum. Large cliques elude the Metropolis process. Random Structures & Algorithms, 3(4):347–359, 1992.
- [JL15] V. Jog and P.-L. Loh. Information-theoretic bounds for exact recovery in weighted stochastic block models using the renyi divergence. arXiv 1509.06418, Sept. 2015.
- [Ken71] David G Kendall. Abundance matrices and seriation in archaeology. *Probability Theory and Related Fields*, 17(2):104–112, 1971.
- [Kot68] Anton Kotzig. Moves without forbidden transitions in a graph. *Matematický časopis*, 18(1):76–80, 1968.
- [LAVBW⁺09] Erez Lieberman-Aiden, Nynke L Van Berkum, Louise Williams, Maxim Imakaev, Tobias Ragoczy, Agnes Telling, Ido Amit, Bryan R Lajoie, Peter J Sabo, Michael O Dorschner, et al. Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *science*, 326(5950):289–293, 2009.
- [Lov12] László Lovász. Large Networks and Graph Limits, volume 60. American Mathematical Society, 2012.
- [LRT08] Adam N Letchford, Gerhard Reinelt, and Dirk Oliver Theis. Odd minimum cut sets and b-matchings revisited. *SIAM Journal on Discrete Mathematics*, 22(4):1480– 1487, 2008.
- [LW14] Cong Han Lim and Stephen Wright. Beyond the birkhoff polytope: Convex relaxations for vector permutation problems. In Advances in Neural Information Processing Systems, pages 2168–2176, 2014.
- [Mal57] Colin L Mallows. Non-null ranking models. i. Biometrika, 44(1/2):114-130, 1957.
- [Mas13] L. Massoulié. Community detection thresholds and the weak ramanujan property. arXiv:1109.3318. The conference version appeared in Proceedings of the 46th Annual ACM Symposium on Theory of Computing, 2013.

sequencing. IEEE Transactions on Information Theory, 59(10):6273–6289, 2013. [McS01]F. McSherry. Spectral partitioning of random graphs. In 42nd IEEE Symposium on Foundations of Computer Science, pages 529 – 537, Oct. 2001. [MNS15] Elchanan Mossel, Joe Neeman, and Allan Sly. Consistency thresholds for the planted bisection model. In Proceedings of the forty-seventh annual ACM symposium on Theory of computing, pages 69–75. ACM, 2015. [MPW15] R. Meka, A. Potechin, and A. Wigderson. Sum-of-squares lower bounds for planted clique. In Proceedings of the Forty-Seventh Annual ACM on Symposium on Theory of Computing, STOC '15, pages 87–96, New York, NY, USA, 2015. ACM. [Pev95] Pavel A Pevzner. Dna physical mapping and alternating eulerian cycles in colored graphs. Algorithmica, 13(1-2):77–105, 1995. $[POS^+16]$ Nicholas H Putnam, Brendan L O'Connell, Jonathan C Stites, Brandon J Rice, Marco Blanchette, Robert Calef, Christopher J Troll, Andrew Fields, Paul D Hartley, and Charles W Sugnet. Chromosome-scale shotgun assembly using an in vitro method for long-range linkage. Genome research, 26(3):342–350, 2016. [PSLW16] Catherine L Peichel, Shawn T Sullivan, Ivan Liachko, and Michael A White. Improvement of the threespine stickleback (gasterosteus aculeatus) genome using a hi-c-based proximity-guided assembly method. bioRxiv, page 068528, 2016. [PW15] W. Perry and A.S. Wein. A semidefinite program for unbalanced multisection in the stochastic block model. arXiv 1507.05605, July 2015. [Rén61] Alfréd Rényi. On measures of entropy and information. In Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics. The Regents of the University of California, 1961. [Rob51] William S Robinson. A method for chronologically ordering archaeological deposits. American antiquity, 16(4):293–301, 1951. [Sch03]Alexander Schrijver. Combinatorial optimization: polyhedra and efficiency, volume 24. Springer Science & Business Media, 2003. $[SVL^+15]$ Nicolas Servant, Nelle Varoquaux, Bryan R Lajoie, Eric Viara, Chong-Jian Chen, Jean-Philippe Vert, Edith Heard, Job Dekker, and Emmanuel Barillot. Hic-pro: an optimized and flexible pipeline for hi-c data processing. Genome biology, 16(1):259, 2015.[SWvZ13] Frans Schalekamp, David P Williamson, and Anke van Zuylen. 2-matchings, the traveling salesman problem, and the subtour LP: A proof of the Boyd-Carr conjecture. Mathematics of Operations Research, 39(2):403–417, 2013. [WS98] Duncan J Watts and Steven H Strogatz. Collective dynamics of 'small-world' networks. nature, 393(6684):440, 1998.

Abolfazl S Motahari, Guy Bresler, and David Tse. Information theory of dna shotgun

[MBT13]

- [ZKRW98] Qing Zhao, Stefan E Karisch, Franz Rendl, and Henry Wolkowicz. Semidefinite programming relaxations for the quadratic assignment problem. *Journal of Combinatorial Optimization*, 2(1):71–109, 1998.
- [ZZ16] Anderson Y Zhang and Harrison H Zhou. Minimax rates of community detection in stochastic block models. *The Annals of Statistics*, 44(5):2252–2280, 2016.