One-point boundaries of ends of clusters in percolation in \mathbb{H}^d

Jan Czajkowski

Argentinian National Research Council at the University of Buenos Aires

March 16, 2022

Abstract

Consider Bernoulli bond percolation on a graph nicely embedded in hyperbolic space \mathbb{H}^d in such a way that it admits a transitive action by isometries of \mathbb{H}^d . Let p_0 be the supremum of such percolation parameters that no point at infinity of \mathbb{H}^d lies in the boundary of the cluster of a fixed vertex with positive probability. Then for any parameter $p < p_0$, a.s. every percolation cluster has only one-point boundaries of ends.

1 Introduction

Consider a graph Γ . Fix $p \in [0, 1]$ and for each edge, mark it as "open" with probability p; do it independently for all edges of Γ . We mark the edges which are not declared open, as "closed". The *state* of an edge is the information whether it is open or closed. The (random) set ω of open edges of Γ forms, together with all the vertices of Γ , a random subgraph of Γ . We will often identify it with the set of edges ω . This is a model of percolation, which we call *Bernoulli bond percolation* on Γ with parameter p.

One of the main objects of interest in percolation theory are the connected components of the random subgraph ω , called *clusters*, and their "size". Here, "size" can mean the number of vertices, the diameter and many other properties of a cluster which measure how "large" it is. For example, one may ask if there is a cluster containing infinitely many vertices in the random subgraph, with positive probability.

Definition 1.1. We define the *critical probability* for the above percolation model on Γ as

 $p_{c}(\Gamma) := \inf\{p \in [0,1] : \text{with positive probability } \omega \text{ has some infinite cluster}\}.$

It follows from the Kolmogorov 0-1 law that if the graph Γ is connected and locally finite, then the probability $P(\omega$ has some infinite cluster) always equals 0 or 1. Since this event is an *increasing* event (see Definition 5.5), its probability is an increasing function of p. Thus, it is 0 for $p < p_c(\Gamma)$ and 1 for $p > p_c(\Gamma)$. **Definition 1.2.** A graph Γ is *transitive*, if its automorphism group acts transitively on the set of vertices of Γ . Γ is *quasi-transitive*, if there are finitely many orbits of vertices of Γ under the action of its automorphism group.

If Γ is connected, locally finite and quasi-transitive, then it is known from [NewmSchul, Theorem 1] that the number of infinite clusters in ω is a.s. constant and equal to 0, 1 or ∞ (see also [LP, thm. 7.5]). Let us focus on the question when this number is ∞ .

Definition 1.3. The *unification probability* for the above percolation model on any graph Γ is the number

 $p_{u}(\Gamma) := \inf\{p \in [0, 1] : \text{a.s. there is a unique infinite cluster in } \omega\}.$

Definition 1.4. A locally finite graph Γ is *non-amenable* if there is a constant $\Phi > 0$ such that for every non-empty finite set K of vertices of Γ , $|\mathrm{bd} K| \ge \Phi |K|$, where bd K is the set of edges of Γ with exactly one vertex in K (a kind of boundary of K). Otherwise, Γ is *amenable*.

It turns out that if a connected, transitive, locally finite graph Γ is amenable, then there is at most 1 infinite cluster a.s. for bond and site Bernoulli percolation, see [LP, thm. 7.6]. So, if $p_{\rm c}(\Gamma) < p_{\rm u}(\Gamma)$ then the graph Γ must be non-amenable. It is an interesting question whether the converse is true:

Conjecture 1.5 ([BS96]). If Γ is non-amenable and quasi-transitive, then $p_{\rm c}(\Gamma) < p_{\rm u}(\Gamma)$.

There are several classes of non-amenable graphs, for which the above conjecture has been established. Let us mention here a few of them. For an infinite regular tree it is actually folklore. It was shown for bond percolation on the Cartesian product of \mathbb{Z}^d with an infinite regular tree of sufficiently high degree in [GrimNewm]. Later, it was shown for site percolation on Cayley graphs of a wide class of Fuchsian groups in [Lal], and for site and bond percolation on transitive, non-amenable, planar graphs with one end in [BS01]. (A graph has *one end*, if after throwing out any finite set of vertices it has exactly one infinite component.) These two result concern the hyperbolic plane \mathbb{H}^2 . Similarly, the conjecture is obtained in [Cz] for many tiling graphs in \mathbb{H}^3 . There is also a rather general result in [PSN] saying that any finitely-generated non-amenable group has a Cayley graph Γ with $p_c(\Gamma) < p_u(\Gamma)$ for bond percolation.

An interested reader may consult e.g. [Grim] and [LP], which give quite wide introduction to percolation theory.

1.1 Boundaries of ends

In this paper we consider percolation clusters on graphs "naturally" embedded in \mathbb{H}^d with $d \geq 2$. We define the *boundaries of ends* of a cluster in \mathbb{H}^d as follows:

Notations 1.6. For any topological space X, by int_X and $\overline{\cdot}^X$ we mean operations of taking interior and closure, respectively, in the space X.

Definition 1.7. Let X be a completely regular Hausdorff $(T_{3\frac{1}{2}})$, locally compact topological space. Then:

- An end of a subset $C \subseteq X$ is a function e from the family of all compact subsets of X to the family of subsets of C such that:
 - for any compact $K \subseteq X$ the set e(K) is one of the component of $C \setminus K$;
 - for $K \subseteq K' \subseteq X$ both compact we have

$$e(K) \supseteq e(K').$$

Now let \hat{X} be an arbitrary compactification of X. Then

• The boundary of $C \subseteq X$ is the following:

$$\partial C = \overline{C}^{X} \setminus X.$$

• Finally the boundary of an end e of $C \subseteq X$ is

$$\partial e = \bigcap_{\substack{K \subseteq X \\ K - \text{ compact}}} \partial e(K).$$

We also put $\widehat{C} = \overline{C}^{\widehat{X}}$. Whenever we use the usual notion of boundary (taken in \mathbb{H}^d by default), we denote it by bd to distinguish it from ∂ .

We use these notions in the context of the hyperbolic space \mathbb{H}^d , where the underlying compactification is the compactification $\widehat{\mathbb{H}}^d$ of \mathbb{H}^d by its set of points at infinity¹ (see [BH, Definition II.8.1]). The role of C above will be played by percolation clusters in \mathbb{H}^d .

Thus, $\partial \mathbb{H}^d = \widehat{\mathbb{H}}^d \setminus \mathbb{H}^d$ is the set of points at infinity. If \mathbb{H}^d is considered in its Poincaré disc model², $\partial \mathbb{H}^d$ is naturally identified with the boundary sphere of the Poincaré disc.

Remark 1.8. In this paper, whenever we consider a subset of \mathbb{H}^d denoted by a symbol of the form e.g. $C_x^y(z)$, we use the notation $\widehat{C}_x^y(z)$ for its closure in $\widehat{\mathbb{H}}^d$ instead of $\widehat{C}_x^y(z)$, for aesthetic reasons.

Let us define a percolation threshold $p_{1/2}$ as the supremum of percolation parameters p such that P_p -a.s. all infinite clusters in ω have only one-point boundaries of ends. The question is if $p_c < p_{1/2} < p_u$ e.g. for some natural tiling graphs in \mathbb{H}^d for $d \geq 3$. In such case one will have an additional percolation threshold in the non-uniqueness phase. In this paper we give a sufficient condition for p-Bernoulli bond percolation to have only one-point boundaries of ends of infinite clusters, for a large class of transitive graphs embedded in \mathbb{H}^d . That sufficient condition is " $p < p_0$ ", where p_0 is a threshold defined in Definition 1.11. The key part of the proof is an adaptation of the proof of Theorem (5.4) from [Grim], which in turn is based on [Men].

In the next section we formulate the assumptions on the graph and the main theorem.

¹For \mathbb{H}^d , it is the same as its Gromov boundary—see [BH, Section III.H.3].

²It is called also Poincaré ball model.

1.2 The graph and the sufficient condition

Definition 1.9. Let for any graph Γ , $V(\Gamma)$ denote its set of vertices and $E(\Gamma)$ its set of edges. In this paper, we often think of a $\omega \subseteq E(\Gamma)$ as a sample, called *percolation configuration*. Accordingly, $2^{E(\Gamma)}$ is the sample space for modelling Bernoulli bond percolation. The accompanying σ -algebra on it consists of all Borel sets (with respect to the product topology). Taking the natural product measure P_p on $2^{E(\Gamma)}$, we treat the configuration ω as the random (set-valued) variable described above.

For any graph G embedded in arbitrary metric space, we call this embedded graph *transitive under isometries* if some group of isometries of the space acts on G by graph automorphisms transitively on its set of vertices.

A graph embedding in a topological space is *locally finite* if every compact subset of \mathbb{H}^d meets only finitely many vertices and edges of the embedded graph.

By a *simple* graph we mean a graph without loops and multiple edges.

Assumption 1.10. Throughout this paper we assume that G is a connected (simple) graph embedded in \mathbb{H}^d , such that:

- its edges are geodesic segments;
- the embedding is locally finite;
- it is transitive under isometries.

Let us also pick a vertex o (for "origin") of G and fix it once and for all.

Note that by these assumptions, V(G) is countable, G has finite degree and is a closed subset of \mathbb{H}^d .

Definition 1.11. For $v \in V(G)$, by C(v) we mean the percolation cluster of v in G. Let $\mathcal{N}(G)$ (for "null"), or \mathcal{N} for short, be defined by

$$\mathcal{N}(G) = \{ p \in [0, 1] : (\forall x \in \partial \mathbb{H}^d) (\mathcal{P}_p(x \in \partial C(o)) = 0) \}$$
(1.1)

and put

$$p_0 = p_0(G) = \sup \mathcal{N}(G). \tag{1.2}$$

Remark 1.12. In words, \mathcal{N} is the set of parameters p of Bernoulli bond percolation on G such that no point of $\partial \mathbb{H}^d$ lies in boundary of the cluster of o with positive probability. Note that \mathcal{N} is an interval (the author does not know whether it is right-open or rightclosed) because the events $\{x \in \partial C(o)\}$ for $x \in \partial \mathbb{H}^d$ are all increasing (see Definition 5.5), so $P_p(x \in \partial C(o))$ is a non-decreasing function of p (see [Grim, Theorem (2.1)]). That allows us to think of p_0 as the point of a phase transition.

We are going to make a few more remarks concerning the above definition and how p_0 may be related to the other percolation thresholds in Section 1.3.

Now, we formulate the main theorem:

Theorem 1.13. Let G satisfy the Assumption 1.10. Then, for any $0 \le p < p_0$, a.s. every cluster in p-Bernoulli bond percolation on G has only one-point boundaries of ends.

The key ingredient of the proof of this theorem is Lemma 3.9, which is a corollary of Theorem 3.6. The latter is quite interesting in its own right. They are presented (along with a proof of Lemma 3.9) in separate Section 3. The elaborate proof of Theorem 3.6, rewritten from the proof of Theorem (5.4) in [Grim], is deferred to Section 5. The proof of this theorem itself, is presented in Section 4.

Remark 1.14. In the assumptions of the above theorem, p_0 can be replaced by

$$p'_0 = \sup\{p \in [0;1] : g_p(r) \xrightarrow[r \to \infty]{} 0\}$$

with $g_p(r)$ from the Definition 3.2 because only the fact that for $p < p_0, g_p(r) \xrightarrow[r \to \infty]{r \to \infty} 0$ (implied by Claim 5.24) is used (in the proof of Lemma 5.20). Accordingly, $p'_0 \ge p_0$. Nevertheless, the author does not know if it is possible that $p'_0 > p_0$.

1.3 Remarks on the sufficient condition

In this section we give some remarks on the threshold p_0 and on the events $\{x \in \partial C(o)\}$ (used to define \mathcal{N}).

Definition 1.15. For $A, B \subseteq \mathbb{H}^d$, let $A \leftrightarrow B$ be the event that there is an open path in the percolation process (given by the context) intersecting both A and B. We say also that such path *joins* A and B. If any of the sets is of the form $\{x\}$, we write x instead of $\{x\}$ in that formula and those phrases.

Remark 1.16. For $x \in \partial \mathbb{H}^d$, the configuration property $\{x \in \partial C(o)\}$ is indeed a (measurable) random event. Even more: the set

$$A = \{ (x, \omega) \in \partial \mathbb{H}^d \times 2^{E(G)} : x \in \partial (C(o))(\omega) \}$$
(1.3)

is measurable in the product $\partial \mathbb{H}^d \times 2^{E(G)}$ (where the underlying σ -field on $\partial \mathbb{H}^d$ is the σ -field of Borel sets). To prove it, let us introduce a countable family $(H_n)_{n \in \mathbb{N}}$ of half-spaces such that the family of open discs

$$\{\operatorname{int}_{\partial \mathbb{H}^d} \partial H_n : n \in \mathbb{N}\}\$$

is a base of the topology on $\partial \mathbb{H}^d$. Then, let us rewrite the condition defining A:

$$x \in \partial C(o) \iff (\forall n) (x \in \operatorname{int}_{\partial \mathbb{H}^d} \partial H_n \Longrightarrow C(o) \cap H_n \neq \emptyset) \iff (1.4)$$

$$\Rightarrow \quad (\forall n) \big(\neg (x \in \operatorname{int}_{\partial \mathbb{H}^d} \partial H_n) \lor \tag{1.5}$$

$$\bigvee \Big(x \in \operatorname{int}_{\partial \mathbb{H}^d} \partial H_n \wedge (\exists v \in V(G) \cap H_n)(o \leftrightarrow v) \Big) \Big),$$

which is a measurable condition, as the sets

\$

$$\{(x,\omega)\in\partial\mathbb{H}^d\times 2^{E(G)}:x\in\operatorname{int}_{\partial\mathbb{H}^d}\partial H_n\}=\operatorname{int}_{\partial\mathbb{H}^d}\partial H_n\times 2^{E(G)}$$

and

$$\{(x,\omega) \in \partial \mathbb{H}^d \times 2^{E(G)} : o \leftrightarrow v \text{ in } \omega\}$$

are measurable for $n \in \mathbb{N}, v \in V(G)$.

For $x \in \partial \mathbb{H}^d$, the measurability of the event $\{x \in \partial C(o)\}$ follows the same way if we treat it as the x-section of A:

$$\{x \in \partial C(o)\} = \{\omega : (x, \omega) \in A\}.$$
(1.6)

Remark 1.17. The threshold p_0 is bounded as follows:

$$p_{\rm c} \le p_0 \le p_{\rm u}.$$

The inequality $p_c \leq p_0$ is obvious and the inequality $p_0 \leq p_u$ can be shown as follows: if p is such that P_p -a.s. there is a unique infinite cluster in G, then with some probability a > 0, o belongs to the infinite cluster and by BK-inequality (see Theorem 5.15), for any $v \in V(G)$,

$$P_p(o \leftrightarrow v) \ge a^2$$

Take $x \in \partial G$. Choose a decreasing (in the sense of set inclusion) sequence $(H_n)_n$ of half-spaces such that $\bigcap_{n=1}^{\infty} \operatorname{int}_{\partial \mathbb{H}^d} \partial H_n = \{x\}$. Since $x \in \partial G$, we have $V(G) \cap H_n \neq \emptyset$ for all n. Therefore

$$P_p(x \in \partial C(o)) = P_p\left(\bigcap_{n \in \mathbb{N}} \{(\exists v \in V(G) \cap H_n)(o \leftrightarrow v)\}\right) =$$
(1.7)

$$= \lim_{n \to \infty} \mathcal{P}_p((\exists v \in V(G) \cap H_n)(o \leftrightarrow v)) \ge a^2.$$
(1.8)

Hence, $p \notin \mathcal{N}$, so $p \geq p_0$, as desired.

The main theorem (Theorem 1.13) is interesting when $p_{\rm c} < p_0$. The author does not know what is the class of embedded graphs G (even among those arising from Coxeter reflection groups as in [Cz]) satisfying $p_{\rm c}(G) < p_0(G)$. The author suspects that $p_0 = p_{\rm u}$ for such graphs as in [Cz] in the cocompact case (see Remark 1.19; in such case most often we would have $p_0 > p_c$). On the other hand, there are examples where $p_0 < p_u$ (see Example 1.18 below). Still, the author does not know if it is possible that $p_c = p_0 < p_u$. *Example* 1.18. Let Π be an unbounded polyhedron with 6 faces in \mathbb{H}^3 whose five faces are cyclically perpendicular and the sixth one is disjoint from them. Then, the group Ggenerated by the (hyperbolic) reflections in the faces of Π is isomorphic to the free product of \mathbb{Z}_2 and the group $G_5 < \text{Isom}(\mathbb{H}^2)$ generated by the reflections in the sides of a rightangled pentagon in \mathbb{H}^2 . Let Γ and Γ_5 be the Cayley graphs of G and G_5 , respectively. Then, Γ has infinitely many ends, so from [LP, Exercise 7.12(b)] $p_u(\Gamma) = 1$. Next, if $p > p_{u}(\Gamma_{5})$, then with positive probability $\partial C(o)$ contains the whole circle $\partial (G_{5} \cdot o)$. (It is implied by Theorem 4.1 and Lemma 4.3 from [BS01].) Hence, $p_0 \leq p_u(\Gamma_5) < p_u(\Gamma)$, as $p_{\rm u}(\Gamma_5) < 1$ by [BB, Theorem 10]. Moreover, the conclusion of the main theorem (Theorem 1.13) fails for any $p > p_u(\Gamma_5)$.

Remark 1.19. This remark is hoped to explain a little the suspicion that for the Cayley graph of a cocompact Coxeter reflection group in \mathbb{H}^d , we have $p_0 = p_u$ (Remark 1.17). It is based on another suspicion: for $p < p_u$ in the same setting,

$$P_p\text{-a.s.} \left|\partial C(o)\right| = 0, \tag{1.9}$$

which is a property of the *p*-Bernoulli bond percolation quite similar to $P_p(x \in \partial C(o)) = 0$. Here $|\cdot|$ can be the Lebesgue measure on $\partial \mathbb{H}^d = \mathbb{S}^{d-1}$, or the Poisson measure on $\partial \mathbb{H}^d$ arising from the simple random walk on G starting at o. (For a definition of a simple random walk and an explanation of Poisson boundary, see [Woe], Section 1.C and Section 24., p. 260, respectively.) If one proves (1.9), then the probability vanishing in (1.1) follows for $|\cdot|$ -a.e. point $x \in \partial \mathbb{H}^d$. In addition, because the induced action of such cocompact group on $\partial \mathbb{H}^d$ has only dense orbits (see e.g. [KapBen, Proposition 4.2]), one might suspect that in such situation as above, $P_p(x \in \partial C(o)) = 0$ holds for all $x \in \partial \mathbb{H}^d$.

2 Definitions: percolation on a fragment of \mathbb{H}^d

Here we are going to introduce some notions and notations used in Theorem 3.6 and Lemma 3.9 and in the proof of the main theorem.

Notation 2.1. Let us adopt the convention saying that natural numbers include 0. We denote the set of all positive natural numbers by \mathbb{N}_+ .

Definition 2.2. For the rest of this paper, consider \mathbb{H}^d in its fixed half-space Poincaré model (being the upper half-space $\mathbb{R}^{d-1} \times (0; \infty)$) in which the point o (the distinguished vertex of G) is represented by $(0, \ldots, 0, 1)$. (It will play the role of origin of both \mathbb{H}^d and G.)

The half-space model of \mathbb{H}^d and its relation to the Poincaré ball model are explained in [BH, Chapter I.6, p. 90]. Note that the inversion of \mathbb{R}^d mapping the Poincaré ball model \mathbb{B}^d to our fixed half-space model sends one point of the sphere bd \mathbb{B}^d to infinity. In the context of the half-space model, we treat that "infinity" as an abstract point (outside \mathbb{R}^d) compactifying \mathbb{R}^d . We call it the *point at infinity* and denote it by ∞ .

Let $\widehat{\mathbb{H}}^d$ be the closure of \mathbb{H}^d in \mathbb{R}^d and $\eth \mathbb{H}^d = \widehat{\mathbb{H}}^d \setminus \mathbb{H}^d$ (so here $\widehat{\mathbb{H}}^d = \mathbb{R}^{d-1} \times [0; \infty)$ and $\eth \mathbb{H}^d = \mathbb{R}^{d-1} \times \{0\}$). We identify $\widehat{\mathbb{H}}^d$ with $\widehat{\mathbb{H}}^d \setminus \{\infty\}$ and $\eth \mathbb{H}^d$ with $\eth \mathbb{H}^d \setminus \{\infty\}$ in a natural way. Also, for any closed $A \subseteq \mathbb{H}^d$, let $\widehat{A} = \overline{A}^{\widehat{\mathbb{H}}^d}$ and $\eth A = \widehat{A} \setminus A$. (Here, for complex notation for a subset of \mathbb{H}^d (of the form e.g. $A_x^y(z)$), we use the same notational convention for $\widehat{\cdot}$ as for $\widehat{\cdot}$ —see Remark 1.8.)

Although sometimes we use the linear and Euclidean structure of \mathbb{R}^d in \mathbb{H}^d , the default geometry on \mathbb{H}^d is the hyperbolic one, unless indicated otherwise. On the other hand, by the *Euclidean metric of the disc model* we mean the metric on $\widehat{\mathbb{H}}^d$ induced by the embedding of $\widehat{\mathbb{H}}^d$ in \mathbb{R}^d (as a unit disc) arising from the Poincaré disc model of \mathbb{H}^d . Nevertheless, we are going to treat that metric as a metric on the set $\widehat{\mathbb{H}}^d \cup \{\infty\} = \widehat{\mathbb{H}}^d$, never really considering \mathbb{H}^d in the disc model.

Definition 2.3. For k > 0 and $x \in \mathbb{R}^{d-1} \times \{0\}$, by $y \mapsto k \cdot y$ and $y \mapsto y + x$ (or $k \cdot , \cdot + x$, respectively, for short) we mean always just a scaling and a translation of \mathbb{R}^d , respectively, often as isometries of \mathbb{H}^d . (Note that restricted to \mathbb{H}^d they are indeed hyperbolic isometries.)

Notations 2.4. Let $\text{Isom}(\mathbb{H}^d)$ denote the isometry group of \mathbb{H}^d .

For any $h \in (0; 1]$ and $R \in O(d)$ (the orthogonal linear group of \mathbb{R}^d) the pair (h, R) determines uniquely an isometry of \mathbb{H}^d denoted by $\Phi^{(h,R)}$ such that $\Phi^{(h,R)}(o) = (0, \ldots, 0, h)$ and $D\Phi^{(h,R)}(o) = hR$ (as an ordinary derivative of a function $\mathbb{R}^{d-1} \times (0, \infty) \to \mathbb{R}^d$).

Let $G^{(h,R)}$ denote $\Phi^{(h,R)}[G]$. Similarly, for any $\Phi \in \text{Isom}(\mathbb{H}^d)$ let $G^{\Phi} = \Phi[G]$. Further, in the same fashion, let $o^{(h,R)} = \Phi^{(h,R)}(o)$ (which is $h \cdot o$) and $o^{\Phi} = \Phi(o)$.

Definition 2.5. For any $p \in [0, 1]$, whenever we consider *p*-Bernoulli bond percolation on G^{Φ} for $\Phi \in \text{Isom}(\mathbb{H}^d)$, we just take $\Phi[\omega]$, where ω denotes the random configuration in *p*-Bernoulli bond percolation on *G*.

Remark 2.6. One can say that this is a way of coupling of the Bernoulli bond percolation processes on G^{Φ} for $\Phi \in \text{Isom}(\mathbb{H}^d)$.

Formally, the notion of "*p*-Bernoulli bond percolation on G^{Φ} " is not well-defined because for different isometries Φ_1 , Φ_2 of \mathbb{H}^d such that $G^{\Phi_1} = G^{\Phi_2}$, still the processes $\Phi_1[\omega]$ and $\Phi_2[\omega]$ are different. Thus, we are going to use the convention that the isometry Φ used to determine the process $\Phi[\omega]$ is the same as used in the notation G^{Φ} determining the underlying graph.

Notations 2.7. Let $L^h = \mathbb{R}^{d-1} \times (0; h] \subseteq \mathbb{H}^d$ and put $L = L^1$. (In other words, L^h is the complement of some open horoball in \mathbb{H}^d , which viewed in the Poincaré disc model \mathbb{B}^d is tangent to $\partial \mathbb{B}^d$ at the point corresponding to ∞ .)

Definition 2.8. Consider any closed set $A \subseteq \mathbb{H}^d$ intersecting each geodesic line only in finitely many intervals and half-lines of that line (every set from the algebra of sets generated by convex sets satisfies this condition, e.g. $A = L^h$). Then, by $G^{\Phi} \cap A$ we mean an embedded graph in A with the set of vertices consisting of $V(G^{\Phi}) \cap A$ and the points of intersection of the edges of G^{Φ} with bd A and with the edges being all the non-degenerate components of intersections of edges of G^{Φ} with A. The percolation process on $G^{\Phi} \cap A$ considered in this paper is, by default, the process $\Phi[\omega] \cap A$. The same convention as in Remark 2.6 is used for these processes.

Remark 2.9. To prove the main theorem, we use the process $\Phi^{(h,R)}[\omega] \cap L^H$ for $p \in [0;1]$ and for different H. In some sense, it is p-Bernoulli bond percolation on $G^{(h,R)} \cap L^H$: on one hand, this process is defined in terms of the independent random states of the edges of $G^{(h,R)}$, but on the other hand, some different edges of the graph $G^{(h,R)} \cap L^H$ are obtained from the same edge of $G^{(h,R)}$, so their states are stochastically dependent. Nevertheless, we are going to use some facts about Bernoulli percolation for the percolation process on $G^{(h,R)} \cap L^H$. In such situation, we consider the edges of $G^{(h,R)}$ intersecting L^H instead of their fragments obtained in the intersection with L^H .

3 Exponential decay of the cluster size distribution

We are going to treat the percolation process $\Phi^{(h,R)}[\omega] \cap L^H$ roughly as a Bernoulli percolation process on the standard lattice \mathbb{Z}^{d-1} (given graph structure by joining every pair of vertices from \mathbb{Z}^{d-1} with distance 1 by an egde). It is motivated by the fact that \mathbb{Z}^{d-1} with the graph metric is quasi-isometric to $\partial \mathbb{H}^d$ or L^H with the Euclidean metric. (Two metric spaces are *quasi-isometric* if, loosely speakig, there are mappings in both directions between them which are bi-Lipschitz up to an additive constant. For strict definition, see [BH, Definition I.8.14]; cf. also Exercise 8.16(1) there.)

In the setting of \mathbb{Z}^{d-1} , we have a theorem on exponential decay of the cluster size distribution, below the critical threshold of percolation:

Theorem 3.1 ([Grim, Theorem (5.4)]). For any $p < p_c(\mathbb{Z}^d)$ there exists $\psi(p) > 0$ such that in p-Bernoulli bond percolation on \mathbb{Z}^d

 $P_p(\text{the origin } (0, \dots, 0) \text{ is connected to the sphere of radius } n) < e^{-\psi(p)n}$ for all n,

where the spheres are considered in the graph metric on \mathbb{Z}^d .

The idea (of a bit more general theorem) comes from [Men], where a sketch of proof is given, and a detailed proof of the above statement is present in [Grim].

We adapt the idea of this theorem to the percolation process on $G^{(h,R)} \cap L$ in Theorem 3.6 and Lemma 3.9, appropriately rewriting the proof in [Grim], which is going to be the key part of the proof of main theorem. In order to consider such counterpart of the above theorem, we define a kind of tail of all the distributions of the cluster size in $G^{(h,R)} \cap L$ for $(h,R) \in (0;1] \times O(d)$ as follows:

Definition 3.2. Let π be the Euclidean orthogonal projection from \mathbb{H}^d onto $\eth \mathbb{H}^d$ and for any $x, y \in \mathbb{H}^d$,

$$d_{\mathfrak{d}}(x,y) = \|\pi(x) - \pi(y)\|_{\infty},$$

where $\|\cdot\|_{\infty}$ is the maximum (i.e. l^{∞}) norm on $\eth \mathbb{H}^d = \mathbb{R}^{d-1} \times \{0\}$. Then, for r > 0 and $x \in \mathbb{H}^d$, let

$$B_r(x) = \{ y \in \mathbb{H}^d : d_{\eth}(x, y) \le r \} \text{ and } S_r(x) = \operatorname{bd} B_r(x)$$

and for h > 0, put

$$B_r^h(x) = B_r(x) \cap L^h.$$

If x = o (or, more generally, if $\pi(x) = \pi(o)$), then we omit "(x)". At last, for $p \in [0, 1]$ and r > 0, let

$$g_p(r) = \sup_{(h,R)\in(0;1]\times O(d)} \mathcal{P}_p(o^{(h,R)} \leftrightarrow S_r \text{ in } G^{(h,R)} \cap L).$$

Remark 3.3. In the Euclidean geometry, $B_r(x)$ and $B_r^h(x)$ are just cuboids of dimensions $r \times \ldots \times r \times \infty$ (unbounded in the direction of *d*-th axis) and $r \times \ldots \times r \times h$, respectively (up to removal of the face lying in $\partial \mathbb{H}^d$).

The condition " $p < p_c(\mathbb{Z}^d)$ " in Theorem 3.1 is going to be replaced by " $p < p_0$ ", which is natural because of the remark below. Before making it, we introduce notation concerning the percolation clusters:

Notations 3.4. For $\Phi \in \text{Isom}(\mathbb{H}^d)$ and $v \in V(G^{\Phi})$ and a set $A \subseteq \mathbb{H}^d$ from the algebra generated by the convex sets, let $C^{\Phi}(v)$ and $C^{\Phi}_A(v)$ be the clusters of v in G^{Φ} and $G^{\Phi} \cap A$, respectively, in the percolation configuration. Similarly, for $(h, R) \in (0; 1] \times O(d)$ and $\Phi = \Phi^{(h,R)}$, we use notations $C^{(h,R)}(v)$ and $C^{(h,R)}_A(v)$, respectively.

If $v = \Phi(o)$, we omit "(v)" for short.

Remark 3.5. If $p \in \mathcal{N}$, then for any $\Phi \in \text{Isom}(\mathbb{H}^d)$, the cluster C^{Φ} is P_p -a.s. bounded in the Euclidean metric. The reason is as follows. Take any $p \in \mathcal{N}$ and $\Phi \in \text{Isom}(\mathbb{H}^d)$. Then, for any $x \in \partial \mathbb{H}^d$, we have $x \notin \widehat{C}(o) P_p$ -a.s. as well as $x \notin \widehat{C}^{\Phi} P_p$ -a.s. If we choose $x = \infty$ (for our half-space model of \mathbb{H}^d), then \widehat{C}^{Φ} is P_p -a.s. a compact set in $\widehat{\mathbb{H}}^d$, so C^{Φ} is bounded in the Euclidean metric.

Now, we formulate the theorem which is the counterpart of Theorem 3.1. Its proof (based on that of [Grim, Theorem (5.4)]) is deferred to Section 5.

Theorem 3.6 (exponential decay of $g_p(\cdot)$). Let a graph G embedded in \mathbb{H}^d be connected, locally finite, transitive under isometries and let its edges be geodesic segments (as in Assumption 1.10). Then, for any $p < p_0$, there exists $\psi = \psi(p) > 0$ such that for any r > 0,

$$g_p(r) \le e^{-\psi r}$$

The next lemma is a stronger version of the above one, where we take the union of all the clusters meeting some $B_{r_0}^1$ instead of the cluster of $o^{(h,R)}$ in $G^{(h,R)} \cap L$. In other words, here the role of $o^{(h,R)}$ played in Theorem 3.6 is taken over by its thickened version $B_{r_0}^1 \cap V(G^{\Phi})$ for any $\Phi \in \text{Isom}(\mathbb{H}^d)$. That leads to the following notation:

Notation 3.7. We put

$$\mathbf{o}_{\Phi} = B^1_{r_0} \cap V(G^{\Phi})$$

for $\Phi \in \text{Isom}(\mathbb{H}^d)$.

Φ

Definition 3.8. For any $C \subseteq \mathbb{H}^d$, we define its *size* by

$$r(C) = \sup_{x \in C} d_{\mathfrak{d}}(o, x).$$

Lemma 3.9. Let a graph G embedded in \mathbb{H}^d be connected, locally finite, transitive under isometries and let its edges be geodesic segments (as in Assumption 1.10). Then, for any p such that the conclusion of Theorem 3.6 holds (in particular, for $p < p_0$) and for any $r_0 > 0$, there exist $\alpha = \alpha(p, r_0), \varphi = \varphi(p, r_0) > 0$ such that for any $r \ge 0$,

$$\sup_{\Phi \in \text{Isom}(\mathbb{H}^d)} \mathsf{P}_p(r(\bigcup_{v \in \mathbf{o}_{\Phi}} C_L^{\Phi}(v)) \ge r) \le \alpha e^{-\varphi r}.$$
(3.1)

Proof. First, note that it is sufficient to prove the inequality

$$\sup_{\Phi \in \text{Isom}(\mathbb{H}^d)} \mathcal{P}_p(\mathbf{o}_\Phi \leftrightarrow S_r \text{ in } G^\Phi \cap L) \le \alpha e^{-\varphi r}$$
(3.2)

for r greater than some fixed $r_1 > 0$ in place of (3.1). Indeed, suppose there exist $\alpha, \varphi > 0$ such that (3.2) holds for all $r > r_1$. We then have:

• for any $r > r_1$ and $\varepsilon \in (0; \min(r - r_1, 1))$,

$$\sup_{\in \text{Isom}(\mathbb{H}^d)} \Pr_p(r(\bigcup_{v \in \mathbf{o}} C_L^{\Phi}(v)) \ge r) \le \sup_{\Phi \in \text{Isom}(\mathbb{H}^d)} \Pr_p(\mathbf{o}_\Phi \leftrightarrow S_{r-\varepsilon} \text{ in } G^{\Phi} \cap L) \le (3.3)$$
$$\le \alpha e^{-\varphi(r-\varepsilon)} \le (\alpha e^{\varphi}) e^{-\varphi r}, \tag{3.4}$$

• for $r \leq r_1$, the left-hand side of (3.1) is less than or equal to $1 \leq e^{\varphi r_1} e^{-\varphi r}$.

So then we will get the lemma for any $r \ge 0$ with $\max(e^{\varphi r_1}, \alpha e^{\varphi})$ put in place of α .

Now, we prove (3.2) (we pick r_1 as above later): let $r > r_0$ and $\Phi \in \text{Isom}(\mathbb{H}^d)$. The task is to pick appropriate values of α and φ independently of r and Φ .

Definition 3.10. Put $\mathbf{o} = \mathbf{o}_{\Phi}$. For $x \in \mathbb{H}^d \subseteq \mathbb{R}^d$, let h(x) denote the *d*-th coordinate of x (or: Euclidean distance from x to $\partial \mathbb{H}^d$), which we call *height* of x.

Assume for a while that $\mathbf{o} \leftrightarrow S_r$ in $G^{\Phi} \cap L$ (note that this event may have probability 0, e.g. when $\mathbf{o} = \emptyset$). Consider all open paths in $G^{\Phi} \cap L$ joining \mathbf{o} to S_r and consider all the vertices of $G\Phi$ visited by those paths, lying in B_r^1 . There is a non-zero finite number of vertices of maximal height among them because G^{Φ} is locally finite. Choose one of these vertices at random and call it $v_{\rm h}$. This \mathbb{H}^d -valued random variable is defined whenever $\mathbf{o} \leftrightarrow S_r$ in $G^{\Phi} \cap L$.)

Observation 3.11. There exists $H \ge 1$ such that a.s. if $v_{\rm h}$ is defined, then $v_{\rm h} \leftrightarrow S_{\frac{r-r_0}{2}}(v_{\rm h})$ in $G^{\Phi} \cap L^{Hh(v_{\rm h})}$.

Proof. Assume that $v_{\rm h}$ is defined and take a path P joining **o** to S_r passing through $v_{\rm h}$. Hyperbolic lengths of edges in G^{Φ} are bounded from above (by the transitivity of G^{Φ} under isometries). That implies that for any edge of G^{Φ} the ratio between the heights of any two of its points is also bounded from above by some constant $H \ge 1$ (it is going to be the H in the observation). (The reason for that are the two following basic properties of the half-space model of \mathbb{H}^d :

- The heights of points of any fixed hyperbolic ball (of finite radius) are bounded from above and from below by some positive constants.
- Any hyperbolic ball can be mapped onto any other hyperbolic ball of the same radius by a translation by vector from $\mathbb{R}^{d-1} \times \{0\}$ composed with a linear scaling of \mathbb{R}^d .

That implies that the path $P \subseteq L^{Hh(v_h)}$.

Now, because P contains some points $x \in B_{r_0}^1$ and $y \in S_r$ and, by triangle inequality, $d_{\overline{\partial}}(x, y) \geq r - r_0$, it follows that $d_{\overline{\partial}}(v_h, x)$ or $d_{\overline{\partial}}(v_h, y)$ is at least $\frac{r-r_0}{2}$ (again by triangle inequality). Hence, P intersects $S_{\frac{r-r_0}{2}}(v_h)$, which finishes the proof. \Box

Based on that observation, we estimate:

$$P_p(\mathbf{o} \leftrightarrow S_r \text{ in } G^{\Phi} \cap L) = \sum_{v \in V(G^{\Phi}) \cap B_r^1} P_p(v_h \text{ is defined and } v_h = v) \le$$
(*)

$$\leq \sum_{v \in V(G^{\Phi}) \cap B_{r}^{1}} \mathbb{P}_{p}(v \leftrightarrow S_{\frac{r-r_{0}}{2}}(v) \text{ in } G^{\Phi} \cap L^{Hh(v)}) =$$
(3.5)

$$= \sum_{v \in V(G^{\Phi}) \cap B_r^1} \Pr_p \left(\frac{1}{H} \cdot o \leftrightarrow S_{\frac{r-r_0}{2Hh(v)}} \left(\frac{1}{H} \cdot o \right) \text{ in } \frac{1}{Hh(v)} (G^{\Phi} - \pi(v)) \cap L^1 \right),$$
(3.6)

by mapping the situation via the (hyperbolic) isometry $\frac{1}{Hh(v)}(\cdot - \pi(v))$ for each v. Note that because for $v \in V(G^{\Phi}) \cap B_r^1$, $\frac{1}{H} \cdot o$ indeed is a vertex of $\frac{1}{Hh(v)}(G^{\Phi} - \pi(v))$, by the transitivity of G under isometries, we can replace the isometry $\frac{1}{Hh(v)}(\cdot - \pi(v))$ with an isometry giving the same image of G and mapping o to $\frac{1}{H} \cdot o$, hence of the form $\Phi^{(1/H,R)}$. That, combined with the assumption on p (the conclusion of Theorem 3.6), gives

$$(*) \leq \sum_{v \in V(G^{\Phi}) \cap B_{r}^{1}} g_{p}\left(\frac{r-r_{0}}{2Hh(v)}\right) \leq \sum_{v \in V(G^{\Phi}) \cap B_{r}^{1}} e^{-\psi \frac{r-r_{0}}{2Hh(v)}},$$
(3.7)

where ψ is as in Theorem 3.6.

Because $B_r^1 = [-r; r]^{d-1} \times (0; 1]$, one can cover it by $\left\lceil \frac{r}{r_0} \right\rceil^{d-1}$ translations of $B_{r_0}^1$ by vectors from $\mathbb{R}^{d-1} \times \{0\}$. So, let $\{B_{r_0}^1(x_i) : i = 1, \ldots, \left\lceil \frac{r}{r_0} \right\rceil^{d-1}\}$ be such covering. Moreover, each $B_{r_0}^1(x_i)$ can be tesselated by infinitely many isometric (in the hyperbolic sense) copies of $K = B_{r_0}^1 \setminus L^{\frac{1}{2}}$, more precisely, by: a translation of K, 2^{d-1} translations of $\frac{1}{2}K$, $(2^{d-1})^2$

translations of $\frac{1}{2^2}K$, etc., all along $\mathbb{R}^{d-1} \times \{0\}$. Let $U = \sup_{\varphi \in \text{Isom}(\mathbb{H}^d)} \#(V(G^{\Phi}) \cap \varphi[K])$ $(U < \infty$ by local finiteness of G). Then, splitting the sum from (3.7) according to those tesselations,

$$(*) \leq \sum_{i=1}^{\left\lceil \frac{r}{r_0} \right\rceil^{d-1}} \sum_{v \in V(G^{\Phi}) \cap B^1_{r_0}(x_i)} e^{-\psi \frac{r-r_0}{2Hh(v)}} \leq$$
(3.8)

$$\leq \left\lceil \frac{r}{r_0} \right\rceil^{d-1} \sum_{k=0}^{\infty} (2^{d-1})^k U \sup_{h \in \left[\frac{1}{2^{k+1}}; \frac{1}{2^k}\right]} e^{-\psi \frac{r-r_0}{2Hh}} \leq (3.9)$$

$$\leq U \left[\frac{r}{r_0} \right]^{d-1} \sum_{k=0}^{\infty} (2^{d-1})^k e^{-\frac{\psi}{H} 2^{k-1} (r-r_0)} =$$
(3.10)

$$= U \left[\frac{r}{r_0} \right]^{d-1} \sum_{k=0}^{\infty} e^{\ln 2 \cdot k(d-1) - \frac{\psi}{H} 2^{k-1}(r-r_0)}.$$
(3.11)

Now, we are going to show that the above bound is finite and tends to 0 at exponential rate with $r \to \infty$. First, we claim that there exists $k_0 \in \mathbb{N}$ such that

$$(\forall k \ge k_0)(\forall r \ge 2r_0) \left(\ln 2 \cdot k(d-1) - 2^{k-1} \frac{\psi}{H}(r-r_0) \le -kr \right).$$
(3.12)

Indeed, for sufficiently large k we have $2^{k-1}\frac{\psi}{H} - k > 0$, so for $r \ge 2r_0$

$$\left(2^{k-1}\frac{\psi}{H}-k\right)r \ge \left(2^{k-1}\frac{\psi}{H}-k\right)\cdot 2r_0 \tag{3.13}$$

and

$$2^{k-1}\frac{\psi}{H}(r-r_0) - kr \ge 2^{k-1}\frac{\psi}{H}r_0 - 2kr_0 \ge k(d-1)\ln 2$$
(3.14)

for sufficiently large k. So, let k_0 satisfy (3.12). Then, for $r \geq 2r_0$,

$$(*) \le U \left\lceil \frac{r}{r_0} \right\rceil^{d-1} \left(\sum_{k=0}^{k_0-1} (2^{d-1})^k e^{-2^{k-1} \frac{\psi}{H}(r-r_0)} + \sum_{k=k_0}^{\infty} e^{-kr} \right) \le$$
(3.15)

$$\leq U \left\lceil \frac{r}{r_0} \right\rceil^{d-1} \left(k_0 (2^{d-1})^{k_0 - 1} e^{-\frac{\psi}{2H}(r - r_0)} + e^{-k_0 r} \underbrace{\frac{1}{1 - e^{-r}}}_{\leq \frac{1}{1 - e^{-2r_0}}} \right) \leq (3.16)$$

$$\leq U \left\lceil \frac{r}{r_0} \right\rceil^{d-1} \left(D e^{-Er} \right) \tag{3.17}$$

for some constants D, E > 0. If we choose $r_1 \ge 2r_0$ such that

$$(\forall r \ge r_1) \left(\left\lceil \frac{r}{r_0} \right\rceil^{d-1} \le e^{\frac{Er}{2}} \right)$$

(which is possible), then

$$(*) \leq UDe^{\frac{-Er}{2}}$$
 for $r \geq r_1$,

which finishes the proof of the lemma.

4 Scaling—proof of the main theorem

Now we complete the proof of the main theorem:

Theorem (recalled Theorem 1.13). Let G satisfy the Assumption 1.10. Then, for any $0 \le p < p_0$, a.s. every cluster in p-Bernoulli bond percolation on G has only one-point boundaries of ends.

Proof of Theorem 1.13. Fix $p \in [0; p_0)$ and suppose towards a contradiction that with some positive probability there is some cluster with some end with a non-one-point boundary. Note that by Remark 3.5 and by the transitivity of G under isometries, for any $v \in V(G)$ a.s. C(v) is bounded in the Euclidean metric, so, a.s. all the percolation clusters in G are bounded in the Euclidean metric. Then, for some $\delta > 0$ and r > 0, there exists with probability a > 0 a cluster bounded in the Euclidean metric, with boundary of some end having Euclidean diameter greater than or equal to δ and intersecting the open disc int $\partial \mathbb{H}^d \partial B_r$. Let C and e be such cluster and its end, respectively. Let for $A \subseteq \mathbb{H}^d$, the projection diameter of A be the Euclidean diameter of $\pi(A)$. Then for h > 0

- the set $\overline{C \setminus L^h}$ is compact;
- $e(\overline{C \setminus L^h})$ is a cluster in the percolation configuration on $G \cap L^h$;
- $e(\overline{C \setminus L^h})$ has projection diameter at least δ and intersects $B_r \cap V(G)$.

All the above implies that for any $k \in \mathbb{N}$,

 $P_p(\exists \text{ a cluster in } G \cap L^{\frac{1}{2^k}} \text{ of projection diameter } \geq \delta \text{ intersecting } B_r \cap V(G)) \geq a,$

so, by scaling by 2^k in \mathbb{R}^d (which is a hyperbolic isometry), we obtain

 $P_p(\exists a \text{ cluster in } G^{2^k} \cap L \text{ of projection diameter} \geq 2^k \delta \text{ intersecting } B_{2^k r} \cap V(G^{2^k})) \geq a$

(where $G^{2^{k}}$ is the image of G under the scaling). $B_{2^{k}r} \cap L$ is a union of $(2^{k})^{d-1}$ isometric copies of $B_r \cap L$, so the left-hand side of above inequality is bounded from above by

$$(2^k)^{d-1} \sup_{\Phi \in \operatorname{Isom}(\mathbb{H}^d)} \operatorname{P}_p(\exists \text{ a cluster in } G^{\Phi} \cap L \text{ of proj. diam.} \geq 2^k \delta \text{ intersecting } B_r \cap V(G^{\Phi})) \leq 0$$

$$\leq (2^k)^{d-1} \sup_{\Phi \in \operatorname{Isom}(\mathbb{H}^d)} \operatorname{P}_p\left(r\left(\bigcup_{v \in B^1_r \cap V(G^\Phi)} C_L^\Phi(v)\right) \geq \frac{2^k \delta}{2} \right) \quad (4.1)$$

(because the size of a cluster is at least half its projection diameter), so by Lemma 3.9, for any $k \in \mathbb{N}$,

$$a \le (2^k)^{d-1} \alpha e^{-\varphi \delta 2^{k-1}}$$

where $\alpha, \varphi > 0$ are constants (as well as δ , a and r). But the right-hand side of this inequality tends to 0 with $k \to \infty$, a contradiction.

5 Proof of the exponential decay

In this section, we prove Theorem 3.6:

Theorem (recalled Theorem 3.6). Let a graph G embedded in \mathbb{H}^d be connected, locally finite, transitive under isometries and let its edges be geodesic segments (as in Assumption 1.10). Then, for any $p < p_0$, there exists $\psi = \psi(p) > 0$ such that for any r > 0,

$$g_p(r) \le e^{-\psi r}$$

Proof. As mentioned earlier, this proof is an adaptation of the proof of Theorem (5.4) in [Grim] based on the work [Men]. Its structure and most of its notation are also borrowed from [Grim], so it is quite easy to compare both the proofs. (The differences are technical and they are summarised in Remark 5.26.) The longest part of this proof is devoted to show functional inequality (5.1) and it goes roughly linearly. Then follows Lemma 5.20, whose proof, using that functional inequality, is deferred to the end of this section. Roughly speaking, that lemma provides a mild asymptotic estimate for g_p (more precisely: for \tilde{g}_p defined below), which is then sharpened to that desired in Theorem 3.6, using repeatedly inequality (5.1).

At some point, we would like to use random variables with left-continuous distribution function³ $1 - g_p$. Because $1 - g_p$ does not need to be left-continuous, we replace g_p , when needed, by its left-continuous version \tilde{g}_p defined as follows:

Definition 5.1. Put $\tilde{g}_p(r) = \lim_{\varrho \to r^-} g_p(\varrho)$ for r > 0.

As one of the cornerstones of this proof, we are going to prove the following functional inequality for $\tilde{g}_{\cdot}(\cdot)$: for any α, β s.t. $0 \leq \alpha < \beta \leq 1$ and for r > 0,

$$\tilde{g}_{\alpha}(r) \leq \tilde{g}_{\beta}(r) \exp\left(-(\beta - \alpha)\left(\frac{r}{a + \int_{0}^{r} \tilde{g}_{\beta}(m) \,\mathrm{d}m} - 1\right)\right),\tag{5.1}$$

where a is a positive constant depending only on G. Note that it implies Theorem 3.6 provided that the integral in the denominator is a bounded function of r.

We are going to approach this inequality, considering the following events depending only on a finite fragment of the percolation configuration and proving functional inequality (5.2) (see below). Cf. Remark 5.27.

Definition 5.2. Fix arbitrary $(h, R) \in (0; 1] \times O(d)$. We are going to use events $A^{\delta}(r)$ defined as follows: let $p \in [0, 1]$, r > 0 and $\delta \in (0; h]$, and define $L_{\delta} = \mathbb{R}^{d-1} \times [\delta; 1] \subseteq \mathbb{H}^d$ (not to be confused with L^{δ}). Let the event

$$A^{\delta}(r) = \{ o^{(h,R)} \leftrightarrow S_r \text{ in } G^{(h,R)} \cap L_{\delta} \}$$

and let

$$f_p^{\delta}(r) = \mathcal{P}_p(A^{\delta}(r)).$$

³By the *left-continuous distribution function* of a probability distribution (measure) μ on \mathbb{R} , we mean the function $\mathbb{R} \ni x \mapsto \mu((-\infty, x))$.

Now, we are going to show that the functions defined above satisfy a functional inequality:

$$f_{\alpha}^{\delta}(r) \le f_{\beta}^{\delta}(r) \exp\left(-(\beta - \alpha)\left(\frac{r}{a + \int_{0}^{r} \tilde{g}_{\beta}(m) \,\mathrm{d}m} - 1\right)\right)$$
(5.2)

for any $0 \le \alpha < \beta \le 1$, r > 0 and for $\delta \in (0; h)$. Having obtained this, we will pass to some limits and to supremum over (h, R), obtaining the inequality (5.1).

Note that, if there is no path joining $o^{(h,R)}$ to S_r in $G^{(h,R)} \cap L_{\delta}$ at all, then for any $p \in [0;1]$, $f_p^{\delta}(r) = 0$ and the inequality (5.2) is obvious. The same happens when $\alpha = 0$. Because in the proof of that inequality we need $f_p^{\delta}(r) > 0$ and $\alpha > 0$, now we make the following assumption (without loss of generality):

Assumption 5.3. We assume that there is a path joining $o^{(h,R)}$ to S_r in $G^{(h,R)} \cap L_{\delta}$ and that $\alpha > 0$. (Then for p > 0, $f_p^{\delta}(r) > 0$).)

The first step in proving inequality (5.2) in using Russo's formula for the events $A^{\delta}(r)$. Before we formulate it, we provide a couple of definition needed there.

Definition 5.4. Next, for a random event A in the percolation on $G^{(h,R)}$, call an edge *pivotal* for a given configuration iff changing the state of that edge (and preserving other edges' states) causes A to change its state as well (from occurring to not occurring or *vice-versa*). Then, let N(A) be the (random) number of all edges pivotal for A.

Definition 5.5. We say that an event A (being a set of configurations) is *increasing* iff for any configurations $\omega \subseteq \omega'$, if $\omega \in A$, then $\omega' \in A$.

Theorem 5.6 (Russo's formula). Consider Bernoulli bond percolation on any graph G and let A be an increasing event defined in terms of the states of only finitely many edges of G. Then

$$\frac{\mathrm{d}}{\mathrm{d}p} \mathcal{P}_p(A) = \mathcal{E}_p(N(A)).$$

This formula is proved as Theorem (2.25) in [Grim] for G being the classical lattice \mathbb{Z}^d , but the proof applies for any graph G.

Let p > 0. The events $A^{\delta}(r)$ depend on the states of only finitely many edges of $G^{(h,R)}$ (namely, those intersecting $L_{\delta} \cap B_r$), so we are able to use Russo's formula for them, obtaining

$$\frac{\mathrm{d}}{\mathrm{d}p}f_p^{\delta}(r) = \mathrm{E}_p(N(A^{\delta}(r))).$$

Now, for $e \in E(G^{(h,R)})$, the event $\{e \text{ is pivotal for } A^{\delta}(r)\}$ is independent of the state of e (which is easily seen; it is the rule for any event), so

$$P_p(A^{\delta}(r) \wedge e \text{ is pivotal for } A^{\delta}(r)) = P_p(e \text{ is open and pivotal for } A^{\delta}(r)) = (5.3)$$

$$= (\text{because } A^{\delta}(r) \text{ is increasing})$$
(5.4)

$$= p \mathcal{P}_p(e \text{ is pivotal for } A^{\delta}(r)), \qquad (5.5)$$

hence

$$\frac{\mathrm{d}}{\mathrm{d}p} f_p^{\delta}(r) = \sum_{e \in E(G^{(h,R)})} \mathcal{P}_p(e \text{ is pivotal for } A^{\delta}(r)) =$$
(5.6)

$$= \frac{1}{p} \sum_{e \in E(G^{(h,R)})} \mathcal{P}_p(A^{\delta}(r) \wedge e \text{ is pivotal for } A^{\delta}(r)) =$$
(5.7)

$$= \frac{f_p^{\delta}(r)}{p} \sum_{e \in E(G^{(h,R)})} \mathcal{P}_p(e \text{ is pivotal for } A^{\delta}(r) | A^{\delta}(r)) =$$
(5.8)

$$=\frac{f_p^{\delta}(r)}{p} \mathcal{E}_p(N(A^{\delta}(r))|A^{\delta}(r)), \qquad (5.9)$$

which can be written as

$$\frac{\mathrm{d}}{\mathrm{d}p}\ln(f_p^{\delta}(r)) = \frac{1}{p}\mathrm{E}_p(N(A^{\delta}(r))|A^{\delta}(r)).$$

For any $0 < \alpha < \beta \leq 1$, integrating over $[\alpha, \beta]$ and exponentiating the above equality gives

$$\frac{f_{\alpha}^{\delta}(r)}{f_{\beta}^{\delta}(r)} = \exp\left(-\int_{\alpha}^{\beta} \frac{1}{p} \mathcal{E}_{p}(N(A^{\delta}(r))|A^{\delta}(r)) \,\mathrm{d}p\right),$$

which implies

$$f_{\alpha}^{\delta}(r) \le f_{\beta}^{\delta}(r) \exp\left(-\int_{\alpha}^{\beta} \mathcal{E}_{p}(N(A^{\delta}(r))|A^{\delta}(r)) \,\mathrm{d}p\right).$$
(5.10)

At this point, our aim is to bound $E_p(N(A^{\delta}(r))|A^{\delta}(r))$ from below.

Definition 5.7. Let η denote the percolation configuration in $G^{(h,R)} \cap L_{\delta}$, i.e.

$$\eta = \Phi^{(h,R)}[\omega] \cap L_{\delta}.$$

Fix any r > 0 and $\delta \in (0; h]$ and assume for a while that $A^{\delta}(r)$ occurs. Let us make a picture of the cluster of $o^{(h,R)}$ in $G^{(h,R)} \cap L_{\delta}$ in the context of the pivotal edges for $A^{\delta}(r)$ (the same picture as in [Grim] and [Men]). If $e \in E(G^{(h,R)})$ is pivotal for $A^{\delta}(r)$, then if we change the percolation configuration by closing e, we cause the cluster of $C_{L_{\delta}}^{(h,R)}(o^{(h,R)})$ to be disjoint from S_r . So, in our situation, all the pivotal edges lie on any open path in $G^{(h,R)} \cap L_{\delta}$ joining $o^{(h,R)}$ to S_r and they are visited by the path in the same order and direction (regardless of the choice of the path).

Definition 5.8. Let $N = N(A^{\delta}(r))$, and let e_1, \ldots, e_N be this ordering, and denote by x_i, y_i the endvertices of e_i, x_i being the one closer to $o^{(h,R)}$ along a path as above. Also, let $y_0 = o^{(h,R)}$.

Note that because for i = 1, ..., N, there is no edge separating y_{i-1} from x_i in the open cluster in $G^{(h,R)} \cap L_{\delta}$, by Menger's theorem (see e.g. [Diest, Theorem 3.3.1, Corollary 3.3.5(ii)]), there exist two edge-disjoint open paths in that cluster joining y_{i-1} to x_i . (One can say, following the discoverer of this proof idea, that that open cluster resembles a chain of sausages.)

Definition 5.9. Now, for i = 1, ..., N, let $\rho_i = d_{\delta}(y_{i-1}, x_i)$ (this way of defining ρ_i , of which one can think as "projection length" of the *i*-th "sausage", is an adaptation of that from [Grim]).

Now we drop the assumption that $A^{\delta}(r)$ occurs. The next lemma is used to compare $(\varrho_1, \ldots, \varrho_N)$ to some renewal process with inter-renewal times of roughly the same distribution as the size of $C_L^{(h,R)}(o^{(h,R)})$.

Definition 5.10. Let a denote the maximal projection distance (in the sense of $d_{\mathfrak{d}}$) between the endpoints of a single edge $G^{(h,R)}$ crossing L.

Lemma 5.11 (cf. [Grim, Lemma (5.12)]). Let $k \in \mathbb{N}_+$ and let $r_1, \ldots, r_k \geq 0$ be such that $\sum_{i=1}^k r_i \leq r - (k-1)a$. Then for 0 ,

$$P_p(\varrho_k < r_k, \ \varrho_i = r_i \text{ for } i < k | A^{\delta}(r)) \ge (1 - g_p(r_k)) P_p(\varrho_i = r_i \text{ for } i < k | A^{\delta}(r)).$$

Remark 5.12. We use the convention that for $i \in \mathbb{N}$ such that $i > N(A^{\delta}(r))$ (i.e. e_i, ϱ_i are undefined), $\varrho_i = +\infty$ (being greater than any real number). On the other hand, whenever we mention $e_i, i \leq N(A^{\delta}(r))$.

Proof of the lemma. This proof mimics that of [Grim, Lemma (5.12)]. Let $k \ge 2$ (we defer the case of k = 1 to the end of the proof).

Definition 5.13. Let for $e \in E(G^{(h,R)} \cap L^{\delta})$, D_e be the connected component of $o^{(h,R)}$ in $\eta \setminus \{e\}$. Let B_e denote the event that the following conditions are satisfied:

- e is open;
- exactly one endvertex of e lies in D_e —call it x(e) and the other—y(e);
- D_e is disjoint from S_r ;
- there are k-1 pivotal edges for the event $\{o^{(h,R)} \leftrightarrow y(e) \text{ in } \eta\}$ (i.e. the edges each of which separates $o^{(h,R)}$ from y(e) in $D_e \cup \{e\}$)—call them $e'_1 = \{x'_1, y'_1\}, \ldots, e'_{k-1} = \{x'_{k-1}, y'_{k-1}\} = e, x'_i$ being closer to $o^{(h,R)}$ than y'_i , in the order from $o^{(h,R)}$ to y(e) (as in the Definition 5.8);
- $d_{\eth}(y'_{i-1}, x'_i) = r_i$ for i < k, where $y'_0 = o^{(h,R)}$.

Let $B = \bigcup_{e \in E(G^{(h,R)} \cap L^{\delta})} B_e$. When B_e occurs, we say that $D_e \cup \{e\}$ with y(e) marked, as a graph with distinguished vertex, is a *witness for B*.

Note that it may happen that there are more than one such witnesses (which means that B_e occurs for many different e). On the other hand, when $A^{\delta}(r)$ occurs, then B_e occurs for only one edge e, namely $e = e_{k-1}$ (in other words, $B \cap A^{\delta}(r) = \bigcup_{e \in E(G^{(h,R)} \cap L^{\delta})} (B_e \cap A^{\delta}(r))$), and there is only one witness for B. Hence,

$$P_p(A^{\delta}(r) \cap B) = \sum_{\Gamma} P_p(\Gamma \text{ a witness for } B) P_p(A^{\delta}(r) | \Gamma \text{ a witness for } B),$$

where the sum is always over all Γ being finite subgraphs of $G^{(h,R)} \cap L_{\delta}$ with distinguished vertices such that $P_p(\Gamma$ a witness for B) > 0.

For Γ a graph with distinguished vertex, let $y(\Gamma)$ denote that vertex. Under the condition that Γ is a witness for B, $A^{\delta}(r)$ is equivalent the event that $y(\Gamma)$ is joined to S_r by an open path in η which is disjoint from $V(\Gamma) \setminus \{y(\Gamma)\}$. We shortly write the latter event $\{y(\Gamma) \leftrightarrow S_r \text{ in } \eta \text{ off } \Gamma\}$. Now, the event $\{\Gamma \text{ a witness for } B\}$ depends only on the states of edges incident to vertices from $V(\Gamma) \setminus \{y(\Gamma)\}$, so it is independent of the event $\{y(\Gamma) \leftrightarrow S_r \text{ in } \eta \text{ off } \Gamma\}$. Hence,

$$P_p(A^{\delta}(r) \cap B) = \sum_{\Gamma} P_p(\Gamma \text{ a witness for } B) P_p(y(\Gamma) \leftrightarrow S_r \text{ in } \eta \text{ off } \Gamma).$$
(5.11)

A similar reasoning, performed below, gives us the estimate of $P_p(\{\varrho_k \ge r_k\} \cap A^{\delta}(r) \cap B)$. Here we use also the following fact: conditioned on the event $\{\Gamma \text{ a witness for } B\}$, the event $A^{\delta}(r) \cap \{\varrho_k \ge r_k\}$ is equivalent to each of the following:

$$(A^{\delta}(r) \wedge e_k \text{ does not exist}) \vee (A^{\delta}(r) \wedge e_k \text{ exists } \wedge \varrho_k \ge r_k) \iff$$

 $\iff (\exists \text{ two edge-disjoint paths joining } y(\Gamma) \text{ to } S_r \text{ in } \eta \text{ off } \Gamma) \lor$

 $\vee(\exists \text{ two edge-disjoint paths in } \eta \text{ off } \Gamma, \text{ joining } y(\Gamma) \text{ to } S_r \text{ and to } S_{r_k}(y(\Gamma)), \text{ resp.}) \iff \iff (\exists \text{ two edge-disjoint paths in } \eta \text{ off } \Gamma, \text{ joining } y(\Gamma) \text{ to } S_r \text{ and to } S_{r_k}(y(\Gamma)), \text{ resp.}),$

because $S_{r_k}(y(\Gamma)) \subseteq B_r$ from the assumption on $\sum_{i=1}^k r_i$. So we estimate

$$P_{p}(\{\varrho_{k} \geq r_{k}\} \cap A^{\delta}(r) \cap B) = \sum_{\Gamma} P_{p}(\Gamma \text{ a witness for } B)P_{p}(\{\varrho_{k} \geq r_{k}\} \cap A^{\delta}(r) | \Gamma \text{ a witness for } B) = \sum_{\Gamma} P_{p}(\Gamma \text{ a witness for } B)P_{p}((y(\Gamma) \leftrightarrow S_{r} \text{ in } \eta \text{ off } \Gamma) \circ (y(\Gamma) \leftrightarrow S_{r_{k}}(y(\Gamma)) \text{ in } \eta \text{ off } \Gamma)),$$

$$(5.12)$$

where the operation " \circ " is defined below:

Definition 5.14. For increasing events A and B in a percolation on any graph G, the event $A \circ B$ means that "A and B occur on disjoint sets of edges". Formally,

$$A \circ B = \{ \omega_A \cup \omega_B : \omega_A, \omega_B \subseteq E(G) \land \omega_A \in A \land \omega_B \in B \},\$$

that is, $A \circ B$ is the set of configurations containing two disjoint set of open edges (ω_A, ω_B above) which guarantee occurring of the events A and B, respectively.

Now, we are going to use the following BK inequality (proved in [Grim]):

Theorem 5.15 (BK inequality, [Grim, Theorems (2.12) and (2.15)]). For any graph G and increasing events A and B depending on the states of only finitely many edges in p-Bernoulli bond percolation on G, we have

$$P_p(A \circ B) \le P_p(A)P_p(B).$$

We use this inequality for the last term (as the events involved are increasing (see def. 5.5) and defined in terms of only the edges from $E(G^{(h,R)} \cap (L_{\delta} \cap B_r)))$, obtaining

$$\begin{split} \mathbf{P}_{p}(\{\varrho_{k} \geq r_{k}\} \cap A^{\delta}(r) \cap B) &\leq \sum_{\Gamma} \mathbf{P}_{p}(\Gamma \text{ a witness for } B) \cdot \mathbf{P}_{p}(y(\Gamma) \leftrightarrow S_{r} \text{ in } \eta \text{ off } \Gamma) \cdot \\ &\cdot \mathbf{P}_{p}(y(\Gamma) \leftrightarrow S_{r_{k}}(y(\Gamma)) \text{ in } \eta \text{ off } \Gamma) \leq \\ &\leq \left(\sum_{\Gamma} \mathbf{P}_{p}(\Gamma \text{ a witness for } B) \mathbf{P}_{p}(y(\Gamma) \leftrightarrow S_{r} \text{ in } \eta \text{ off } \Gamma)\right) g_{p}(r_{k}) = \\ &= \mathbf{P}_{p}(A^{\delta}(r) \cap B) g_{p}(r_{k}) \end{split}$$

(by (5.11)). Dividing by $P_p(A^{\delta}(r))$ (which is positive by Assumption 5.3) gives

$$P_p(\{\varrho_k \ge r_k\} \cap B | A^{\delta}(r)) \le P_p(B | A^{\delta}(r)) g_p(r_k) \quad | \quad P_p(B | A^{\delta}(r)) - \cdot$$
(5.13)

$$P_p(\{\varrho_k < r_k\} \cap B | A^{\delta}(r)) \ge P_p(B | A^{\delta}(r))(1 - g_p(r_k)).$$

$$(5.14)$$

Note that, conditioned on $A^{\delta}(r)$, B is equivalent to the event $\{\varrho_i = r_i \text{ for } i < k\}$, so the above amounts to

$$P_p(\varrho_k < r_k, \varrho_i = r_i \text{ for } i < k | A^{\delta}(r)) \ge P_p(\varrho_i = r_i \text{ for } i < k | A^{\delta}(r))(1 - g_p(r_k)), \quad (5.15)$$

which is the desired conclusion.

Now, consider the case of k = 1. In this case, similarly to (5.12) and thanks to the assumption $r_1 \leq r$,

$$P_p(\{\varrho_1 \ge r_1\} \cap A^{\delta}(r)) = P_p((o^{(h,R)} \leftrightarrow S_{r_1} \text{ in } \eta) \circ (o^{(h,R)} \leftrightarrow S_r \text{ in } \eta)) \le g_p(r_1)P_p(A^{\delta}(r)).$$
(5.16)

Further, similarly to (5.13),

$$P_p(\varrho_1 < r_1 | A^{\delta}(r)) \ge 1 - g_p(r_1), \tag{5.17}$$

which is the lemma's conclusion for k = 1.

Now, we want to do some probabilistic reasoning using random variables with the leftcontinuous distribution function $1 - \tilde{g}_p$. The function $1 - \tilde{g}_p$ is non-decreasing (because for $(h, R) \in (0; 1] \times O(d)$, $P_p(o^{(h,R)} \leftrightarrow S_r$ in $G^{(h,R)} \cap L$) is non-increasing with respect to r, so g_p and \tilde{g}_p are non-increasing as well), left-continuous, with values in [0; 1] and such that $1 - \tilde{g}_p(0) = 0$, so it is the left-continuous distribution function of a random variable with values in $[0; \infty]$.

Notations 5.16. Let M_1, M_2, \ldots be an infinite sequence of independent random variables all distributed according to $1 - \tilde{g}_p$ and all independent of the whole percolation process. Because their distribution depends on p, we will also denote them by $M_1^{(p)}, M_2^{(p)}, \ldots$ (Here, an abuse of notation is going to happen, as we are still writing P_p for the whole probability measure used also for defining the variables M_1, M_2, \ldots)

We can now state the following corollary of Lemma 5.11:

Corollary 5.17. For any r > 0, positive integer k and 0 ,

$$P_p(\varrho_1 + \dots + \varrho_k < r - (k-1)a|A^{\delta}(r)) \ge P_p(M_1 + \dots + M_k < r - (k-1)a)$$

Proof. We compose the proof of the intermediate inequalities:

$$P_{p}(\varrho_{1} + \dots + \varrho_{k} < r - (k - 1)a|A^{\delta}(r)) \geq \\ \geq P_{p}(\varrho_{1} + \dots + \varrho_{k-1} + M_{k} < r - (k - 1)a|A^{\delta}(r)) \geq \\ \dots \geq P_{p}(\varrho_{1} + M_{2} + \dots + M_{k} < r - (k - 1)a|A^{\delta}(r)) \geq \\ \geq P_{p}(M_{1} + \dots + M_{k} < r - (k - 1)a|A^{\delta}(r)) = P_{p}(M_{1} + \dots + M_{k} < r - (k - 1)a)$$

using the step:

$$P_p(\varrho_1 + \dots + \varrho_j + M_{j+1} + \dots + M_k < r - (k-1)a|A^{\delta}(r)) \ge$$
(5.18)

$$\geq P_p(\varrho_1 + \dots + \varrho_{j-1} + M_j + \dots + M_k < r - (k-1)a|A^{\delta}(r)).$$
(5.19)

for $j = k, k - 1, \dots, 2, 1$. Now we prove this step: let $j \in \{1, 2, \dots, k\}$.

Definition 5.18. Put

$$\mathcal{R}^{(h,R)} = \{ d_{\eth}(x,y) : x, y \in V(G^{(h,R)}) \}.$$

Note that it is a countable set of all possible values of ρ_i for $i = 1, \ldots, N$.

We express the considered probability as an integral, thinking of the whole probability space as Cartesian product of the space on which the percolation processes are defined and the space used for defining M_1, M_2, \ldots , and using a version of Fubini theorem for events:

$$P_p(\varrho_1 + \dots + \varrho_j + M_{j+1} + \dots + M_k < r - (k-1)a|A^{\delta}(r)) =$$

= $\int P_p(\varrho_1 + \dots + \varrho_j + S_M < r - (k-1)a|A^{\delta}(r)) d\mathcal{L}_{j+1}^k(S_M) =$

(here \mathcal{L}_{j+1}^k denotes the distribution of the random variable $M_{j+1} + \cdots + M_k$)

$$= \int \sum_{(r_1,\dots,r_{j-1})} \Pr\left(\varrho_i = r_i \text{ for } i < j \land \varrho_j < r - (k-1)a - \sum_{i=1}^{j-1} r_i - S_M \middle| A^{\delta}(r) \right) \, \mathrm{d}\mathcal{L}_{j+1}^k(S_M) \ge 0$$

(where the sum is taken over all $(r_1, ..., r_{j-1}) \in (\mathcal{R}^{(h,R)})^{j-1} : r_1 + \cdots + r_{j-1} < r - (k - 1)a - S_M$)

$$\geq \int \sum_{(r_1,\dots,r_{j-1})} \left(1 - \tilde{g}_p \left(r - (k-1)a - \sum_{i=1}^{j-1} r_i - S_M \right) \right) P_p(\varrho_i = r_i \text{ for } i < j | A^{\delta}(r)) \, \mathrm{d}\mathcal{L}_{j+1}^k(S_M) =$$

(from Lemma 5.11 and because $g_p \leq \tilde{g}_p$)

$$= \int \sum_{(r_1,\dots,r_{j-1})} P_p\left(M_j < r - (k-1)a - \sum_{i=1}^{j-1} r_i - S_M \land \varrho_i = r_i \text{ for } i < j \middle| A^{\delta}(r)\right) d\mathcal{L}_{j+1}^k(S_M) = \\ = \int P_p(\varrho_1 + \dots + \varrho_{j-1} + M_j + S_M < r - (k-1)a | A^{\delta}(r)) d\mathcal{L}_{j+1}^k(S_M) = \\ = P_p(\varrho_1 + \dots + \varrho_{j-1} + M_j + M_{j+1} + \dots + M_k < r - (k-1)a | A^{\delta}(r)).$$

That completes the proof.

Lemma 5.19 (cf. [Grim, Lemma (5.17)]). For 0 , <math>r > 0,

$$\mathbb{E}_p(N(A^{\delta}(r))|A^{\delta}(r)) \ge \frac{r}{a + \int_0^r \tilde{g}_p(m) \,\mathrm{d}m} - 1.$$

Proof. For any $k \in \mathbb{N}_+$, if $\varrho_1 + \cdots + \varrho_k < r - (k-1)a$, then e_1, \ldots, e_k exist and $N(A^{\delta}(r)) \ge k$. So, from the corollary above,

$$P_p(N(A^{\delta}(r)) \ge k | A^{\delta}(r)) \ge P_p(\sum_{i=1}^k \varrho_i < r - (k-1)a) \ge P_p(\sum_{i=1}^k M_i < r - (k-1)a).$$
(5.20)

Now, we use a calculation which relates $a + \int_0^r \tilde{g}_p(m) dm$ to the distribution of M_1 . Namely, we replace the variables M_i by

$$M_i' = a + \min(M_i, r)$$

for i = 1, 2, ... (a kind of truncated version of M_i). In this setting,

$$\sum_{i=1}^{k} M_i < r - (k-1)a \iff \sum_{i=1}^{k} \min(M_i, r) < r - (k-1)a \iff \sum_{i=1}^{k} M'_i < r + a, \quad (5.21)$$

so from (5.20),

$$E_p(N(A^{\delta}(r))|A^{\delta}(r)) = \sum_{k=1}^{\infty} P_p(N(A^{\delta}(r)) \ge k|A^{\delta}(r)) \ge$$
(5.22)

$$\geq \sum_{k=1}^{\infty} \mathcal{P}_p(\sum_{i=1}^k M'_i < r+a) = \sum_{k=1}^{\infty} \mathcal{P}_p(K \ge k+1) =$$
(5.23)

$$= E(K) - 1,$$
 (5.24)

where

 $K = \min\{k : M'_1 + \dots + M'_k \ge r + a\}.$

Let for $k \in \mathbb{N}$,

$$S_k = M_1' + \dots + M_k'.$$

By Wald's equation (see e.g. [GrimSti, p. 396]) for the random variable S_K ,

 $r + a \le \mathcal{E}(S_K) = \mathcal{E}(K)\mathcal{E}(M_1').$

In order that Wald's equation were valid for S_K , the random variable K has to satisfy $E(M'_i|K \ge i) = E(M'_i)$ for $i \in \mathbb{N}_+$. But we have

$$K \ge i \iff M'_1 + \dots + M'_{i-1} < r + a,$$

so M'_i is independent of the event $\{K \ge i\}$ for $i \in \mathbb{N}_+$, which allows us to use Wald's equation. (In fact, K is a so-called stopping time for the sequence $(M'_i)_{i=1}^{\infty}$.) Hence,

$$E_p(N(A^{\delta}(r))|A^{\delta}(r)) \ge E(K) - 1 \ge \frac{r+a}{E(M'_1)} - 1 =$$

$$= \frac{r+a}{a + \int_0^\infty P_p(\min(M_1, r) \ge m) \, \mathrm{d}m} - 1 \ge \frac{r}{a + \int_0^r \tilde{g}_p(m) \, \mathrm{d}m} - 1,$$
(5.25)
(5.26)

which finishes the proof.

Now, combining that with inequality (5.10) for $0 < \alpha < \beta \leq 1$, we have

$$f_{\alpha}^{\delta}(r) \le f_{\beta}^{\delta}(r) \exp\left(-\int_{\alpha}^{\beta} \mathcal{E}_{p}(N(A^{\delta}(r))|A^{\delta}(r)) \,\mathrm{d}p\right) \le$$
(5.27)

$$\leq f_{\beta}^{\delta}(r) \exp\left(-\int_{\alpha}^{\beta} \left(\frac{r}{a+\int_{0}^{r} \tilde{g}_{p}(m) \,\mathrm{d}m} - 1\right) \,\mathrm{d}p\right) \leq \tag{5.28}$$

$$\leq f_{\beta}^{\delta}(r) \exp\left(-(\beta - \alpha)\left(\frac{r}{a + \int_{0}^{r} \tilde{g}_{\beta}(m) \,\mathrm{d}m} - 1\right)\right)$$
(5.29)

(because $\tilde{g}_p \leq \tilde{g}_\beta$ for $p \leq \beta$), which completes the proof of inequality 5.2. (Let us now drop Assumption 5.3.)

Now, note that for any r > 0 and $p \in [0; 1]$, the event $A^{\delta}(r)$ increases as δ decreases. Thus, taking the limit with $\delta \to 0$, we have

$$\lim_{\delta \to 0^+} f_p^{\delta}(r) = \mathcal{P}_p(\bigcup_{\delta > 0} A^{\delta}(r)) = \mathcal{P}_p(o^{(h,R)} \leftrightarrow S_r \text{ in } G^{(h,R)} \cap L).$$

So for any r > 0 and $0 \le \alpha < \beta \le 1$, using this for inequality 5.2 gives

$$P_{\alpha}(o^{(h,R)} \leftrightarrow S_r \text{ in } G^{(h,R)} \cap L) \leq \\ \leq P_{\beta}(o^{(h,R)} \leftrightarrow S_r \text{ in } G^{(h,R)} \cap L) \exp\left(-(\beta - \alpha)\left(\frac{r}{a + \int_0^r \tilde{g}_{\beta}(m) \,\mathrm{d}m} - 1\right)\right).$$
(5.30)

Further, we take the supremum over $(h, R) \in (0; 1] \times O(d)$, obtaining

$$g_{\alpha}(r) \leq g_{\beta}(r) \exp\left(-(\beta - \alpha)\left(\frac{r}{a + \int_{0}^{r} \tilde{g}_{\beta}(m) \,\mathrm{d}m} - 1\right)\right).$$

At last, taking the limits with r from the left, we get the functional inequality (5.1) involving only $\tilde{g}_{\cdot}(\cdot)$:

$$\tilde{g}_{\alpha}(r) \leq \tilde{g}_{\beta}(r) \exp\left(-(\beta - \alpha)\left(\frac{r}{a + \int_{0}^{r} \tilde{g}_{\beta}(m) \,\mathrm{d}m} - 1\right)\right).$$
(5.31)

(Note that the exponent remains unchanged all the time from (5.29) till now.)

Recall that once we have

$$\int_0^\infty \tilde{g}_\beta(m) \,\mathrm{d}m = \mathcal{E}(M_1^{(\beta)}) < \infty,$$

then we obtain Theorem 3.6 for $\tilde{g}_{\alpha}(r)$, for $\alpha < \beta$. This bound is going to be established by showing the rapid decay of \tilde{g}_p , using repeatedly (5.31). The next lemma is the first step of this procedure.

Lemma 5.20 (cf. [Grim, Lemma (5.24)]). For any $p < p_0$, there exists $\delta(p)$ such that

$$\tilde{g}_p(r) \le \delta(p) \cdot \frac{1}{\sqrt{r}} \quad for \ r > 0.$$

We defer proving the above lemma to the end of this section.

Obtaining Theorem 3.6 (being proved) from Lemma 5.20 is relatively easy. First, we deduce that for r > 0 and $p < p_0$,

$$\int_0^r \tilde{g}_p(m) \, \mathrm{d}m \le 2\delta(p)\sqrt{r},$$

so if $r \ge a^2$, then

$$a + \int_0^r \tilde{g}_p(m) \,\mathrm{d}m \le (2\delta(p) + 1)\sqrt{r}.$$

Then, using (5.31), for $0 \le \alpha < \beta < p_0$, we have

$$\int_{a^2}^{\infty} \tilde{g}_{\alpha}(r) \,\mathrm{d}r \le \int_{a^2}^{\infty} \exp\left(-(\beta - \alpha)\left(\frac{r}{a + \int_0^r \tilde{g}_{\beta}(m) \,\mathrm{d}m} - 1\right)\right) \,\mathrm{d}r \le \tag{5.32}$$

$$\leq e \int_{a^2}^{\infty} \exp\left(-\underbrace{\frac{\beta - \alpha}{2\delta(\beta) + 1}}_{=C>0} \sqrt{r}\right) \mathrm{d}r =$$
(5.33)

$$= e \int_{a}^{\infty} e^{-Cx} \cdot 2x \,\mathrm{d}x,\tag{5.34}$$

 \mathbf{SO}

$$\mathcal{E}(M_1^{(\alpha)}) = \int_0^\infty \tilde{g}_\alpha(r) \,\mathrm{d}r \le a^2 + e \int_a^\infty e^{-Cx} \cdot 2x \,\mathrm{d}x < \infty,$$

as desired. Finally, we use the finiteness of $E(M_1^{(\alpha)})$ as promised: for r > 0 and $0 \le \alpha < p_0$, if we take $\alpha < \beta < p_0$, then, using (5.31) again,

$$g_{\alpha}(r) \leq \tilde{g}_{\alpha}(r) \leq \exp\left(-(\beta - \alpha)\left(\frac{r}{a + \mathcal{E}(M_{1}^{(\beta)})} - 1\right)\right) \leq e^{-\varphi(\alpha,\beta)r + \gamma(\alpha,\beta)}, \qquad (5.35)$$

for some constants $\varphi(\alpha, \beta), \gamma(\alpha, \beta) > 0$.

Now we perform a standard estimation, aiming to rule out the additive constant $\gamma(\alpha, \beta)$. For any $0 < \psi_1 < \varphi(\alpha, \beta)$, there exists $r_0 > 0$ such that for $r \ge r_0$,

$$-\varphi(\alpha,\beta)r + \gamma(\alpha,\beta) \le -\psi_1 r,$$

 \mathbf{SO}

$$g_{\alpha}(r) \le e^{-\psi_1 r}.$$

On the other hand, for any r > 0, $g_p(r)$ is no greater than the probability of opening at least on edge adjacent to o, so $g_{\alpha}(r) \leq 1 - (1 - \alpha)^{\deg(o)} < 1$, where $\deg(o)$ is the degree of o in the graph G. Hence,

$$g_{\alpha}(r) \le e^{-\psi_2(\alpha)r}$$

for $r \leq r_0$, for some sufficiently small $\psi_2(\alpha) > 0$. Taking $\psi = \min(\psi_1, \psi_2(\alpha))$ gives

$$g_{\alpha}(r) \le e^{-\psi r}$$

for any r > 0, completing the proof of Theorem 3.6.

Now we are going to prove Lemma 5.20.

Proof of Lemma 5.20. Assume without loss of generality that $\tilde{g}_p(r) > 0$ for r > 0. We are going to construct sequences $(p_i)_{i=1}^{\infty}$ and $(r_i)_{i=1}^{\infty}$ such that

$$p_0 > p_1 > p_2 > \dots > p, \quad 0 < r_1 \le r_2 \le \dots$$

and such that the sequence $(\tilde{g}_{p_i}(r_i))_{i=1}^{\infty}$ decays rapidly. The construction is by recursion: for $i \geq 1$, having constructed p_1, \ldots, p_i and r_1, \ldots, r_i , we put

$$r_{i+1} = r_i/g_i$$
 and $p_{i+1} = p_i - 3g_i(1 - \ln g_i),$ (5.36)

where $g_i = \tilde{g}_{p_i}(r_i)$. (Note that indeed, $r_{i+1} \leq r_i$ and $p_{i+1} < p_i$.) The above formula may give an incorrect value of p_{i+1} , i.e. not satisfying $p_{i+1} > p$ (this condition is needed because we want to bound values of \tilde{g}_p). In order to prevent that, we choose appropriate values of p_1, r_1 , using the following fact to bound the difference $p_1 - p_i$ by a small number independent of i.

Proposition 5.21. If we define sequence $(x_i)_{i=1}^{\infty}$ by $x_{i+1} = x_i^2$ for $i \ge 1$ (i.e. $x_i = x_1^{2^{i-1}}$) with $0 < x_1 < 1$, then

$$s(x_1) := \sum_{i=1}^{\infty} 3x_i (1 - \ln x_i)$$
(5.37)

is finite and $s(x_1) \xrightarrow[x_1 \to 0]{} 0$.

(The idea of the proof of this fact is similar to that of estimating the sum in (3.10).) To make use of it, we are going to bound g_i by x_i for any i and to make g_1 small enough. It is done thanks to the two claims below, respectively.

Claim 5.22. If $p_1, \ldots, p_i > p$ and $r_1, \ldots, r_i > 0$ are defined by (5.36) with $r_1 \ge a$, we have

$$g_{j+1} \le g_j^2$$

for j = 1, ..., i - 1.

Proof. Let $j \in \{1, \ldots, i-1\}$. From (5.31),

$$g_{j+1} \le \tilde{g}_{p_j}(r_{j+1}) \exp\left(-(p_j - p_{j+1})\left(\frac{r_{j+1}}{a + \int_0^{r_{j+1}} \tilde{g}_{p_j}(m) \,\mathrm{d}m} - 1\right)\right) \le (5.38)$$

$$\leq g_j \exp\left(1 - (p_j - p_{j+1}) \frac{r_{j+1}}{a + \int_0^{r_{j+1}} \tilde{g}_{p_j}(m) \,\mathrm{d}m}\right).$$
(5.39)

Inverse of the fraction above is estimated as follows

$$\frac{1}{r_{j+1}} \left(a + \int_0^{r_{j+1}} \tilde{g}_{p_j}(m) \, \mathrm{d}m \right) \le \frac{a}{r_{j+1}} + \frac{r_j}{r_{j+1}} + \frac{1}{r_{j+1}} \int_{r_j}^{r_{j+1}} \tilde{g}_{p_j}(m) \, \mathrm{d}m \le \tag{5.40}$$

$$\leq \frac{a}{r_{j+1}} + g_j + \frac{r_{j+1} - r_j}{r_{j+1}} \tilde{g}_{p_j}(r_j) \leq (5.41)$$

(using $r_{j+1} = r_j/g_j$ and the monotonicity of $\tilde{g}_{p_j}(\cdot)$)

$$\leq \frac{a}{r_{j+1}} + 2g_j. \tag{5.42}$$

Now, by the assumption, $r_j \ge r_1 \ge a$, so $r_{j+1} = r_j/g_j \ge a/g_j$ and

$$\frac{a}{r_{j+1}} + 2g_j \le 3g_j.$$

That gives

$$g_{j+1} \le g_j \exp\left(1 - \frac{p_j - p_{j+1}}{3g_j}\right) = g_j^2$$

by the definition of p_{j+1} .

Notation 5.23. Put

$$M^{(h,R)} = r(C^{(h,R)})$$

for $(h, R) \in (0; 1] \times O(d)$.

Note that, by Remark 3.5, for any $p \in \mathcal{N}$, P_p -a.s. $M^{(h,R)} < \infty$.

Claim 5.24. For any $p \in \mathcal{N}$,

$$\tilde{g}_p(r) \xrightarrow[r \to \infty]{} 0$$

Proof. First, note that it is sufficient to prove

$$\sup_{R \in O(d)} \mathcal{P}_p(M^{(1,R)} \ge r) \xrightarrow[r \to \infty]{} 0, \tag{5.43}$$

because

$$g_p(r) = \sup_{(h,R)\in(0;1]\times O(d)} P_p(o^{(h,R)} \leftrightarrow S_r \text{ in } G^{(h,R)} \cap L) \le$$
(5.44)

$$\leq \sup_{(h,R)\in(0;1]\times O(d)} \mathcal{P}_p(o^{(h,R)} \leftrightarrow S_{hr} \text{ in } G^{(h,R)}) = \quad (\text{because } hr \leq r) \tag{5.45}$$

$$= \sup_{R \in O(d)} \mathcal{P}_p(o \leftrightarrow S_r \text{ in } G^{(1,R)}) \le \quad \text{(by scaling the situation)}$$
(5.46)

$$\leq \sup_{R \in O(d)} \mathcal{P}_p(M^{(1,R)} \ge r),$$
(5.47)

so $g_p(r) \xrightarrow[r\to\infty]{} 0$ and, equivalently, $\tilde{g}_p(r) \xrightarrow[r\to\infty]{} 0$ will be implied. To prove (5.43), we use upper semi-continuity of the function $O(d) \ni R \mapsto \mathcal{P}_p(M^{(1,R)} \ge r)$ for any $p \in \mathcal{N}$ and r > 0. Let us fix such p and r and let $(R_n)_n$ be a sequence of elements of O(d) convergent to some R. Assume without loss of generality that the cluster $C^{(1,R)}$ is bounded in the Euclidean metric and, throughout this proof, condition on it all the events by default. We are going to show that

$$\limsup_{n \to \infty} \{ M^{(1,R_n)} \ge r \} \subseteq \{ M^{(1,R)} \ge r \}.$$
(5.48)

Definition 5.25. For any isometry Φ of \mathbb{H}^d , let $\widehat{\Phi}$ denote the unique continuous extension of Φ to $\widehat{\mathbb{H}}^d$ (which is a homeomorphism of $\widehat{\mathbb{H}}^d$ —see [BH, Corollary II.8.9]).

Put $\Phi_n = \Phi^{(1,R)} \circ (\Phi^{(1,R_n)})^{-1}$ and assume that the event $\limsup_{n\to\infty} \{M^{(1,R_n)} \ge r\}$ occurs. Then, for infinitely many values of n, all the following occur:

$$M^{(1,R_n)} \ge r \Longrightarrow \widehat{C}^{(1,R_n)} \text{ intersects } \widehat{S}_r \stackrel{\widehat{\Phi}_n(\cdot)}{\Longrightarrow} \widehat{C}^{(1,R)} = \widehat{C}^{(1,R)} \text{ intersects } \widehat{\Phi}_n(\widehat{S}_r).$$

Let for any such n, x_n be chosen from the set $\widehat{C}^{(1,R)} \cap \widehat{\Phi}_n(\widehat{S}_r)$. Because $\widehat{C}^{(1,R)}$ is compact, the sequence $(x_n)_n$ (indexed by a subset of \mathbb{N}_+) has an (infinite) subsequence $(x_{n_k})_{k=1}^{\infty}$ convergent to some point in $\widehat{C}^{(1,R)}$. On the other hand, note that $\widehat{\Phi}_n \xrightarrow[n\to\infty]{} \operatorname{Id}_{\widehat{\mathbb{H}}^d}$ uniformly in the Euclidean metric of the disc model (see Definition 2.2). Hence, the distance in that metric between $x_{n_k} \in \widehat{\Phi}_{n_k}(\widehat{S}_r)$ and \widehat{S}_r tends to 0 with $k \to \infty$, so

$$\lim_{k \to \infty} x_{n_k} \in \widehat{S}_r \cap \widehat{C}^{(1,R)} = \widehat{S}_r \cap \widehat{C}^{(1,R)},$$

which shows that $M^{(1,R)} \ge r$, as desired in 5.48. Now,

$$\limsup_{n \to \infty} \mathcal{P}_p(M^{(1,R_n)} \ge r) \le \mathcal{P}_p(\limsup_{n \to \infty} \{M^{(1,R_n)} \ge r\}) \le \quad \text{(by an easy exercise)} \quad (5.49)$$
$$\le \mathcal{P}_p(M^{(1,R)} \ge r), \tag{5.50}$$

which means exactly the upper semi-continuity of $R \mapsto P_p(M^{(1,R)} \ge r)$.

Next, note that because for $p \in \mathcal{N}$ and $R \in O(d)$, a.s. $M^{(1,R)} < \infty$, we have

$$P_p(M^{(1,R)} \ge r) \xrightarrow[r \to \infty]{} 0 \quad (decreasingly).$$

Hence, if for r > 0 and $\varepsilon > 0$ we put

$$U_{\varepsilon}(r) = \{ R \in O(d) : \mathcal{P}_p(M^{(1,R)} \ge r) < \varepsilon \},\$$

then for any fixed $\varepsilon > 0$,

$$\bigcup_{r \nearrow \infty} U_{\varepsilon}(r) = O(d).$$
(5.51)

 $U_{\varepsilon}(r)$ is always an open subset of O(d) by upper semi-continuity of $R \mapsto \mathcal{P}_p(M^{(1,R)} \geq r)$, so by the compactness of O(d), the union (5.51) is indeed finite. Moreover, because $U_{\varepsilon}(r)$ increases as r increases, it equals O(d) for some r > 0. It means that $\sup_{R \in O(d)} \mathcal{P}_p(M^{(1,R)} \geq r) \leq \varepsilon$, whence $\sup_{R \in O(d)} \mathcal{P}_p(M^{(1,R)} \geq r) \xrightarrow[r \to \infty]{} 0$, as desired. \Box

Now, taking any $p_1 \in (p, p_0)$ and $1 > x_1 > 0$ in (5.37) s.t. $s(x_1) \leq p_1 - p$ and taking $r_1 \geq a$ so large that $\tilde{g}_{p_1}(r_1) < x_1$, we obtain for $i \geq 1$, $g_i < x_i$ (by induction). Then, in the setting of (5.36),

$$p_{i+1} = p_1 - \sum_{j=1}^{i} 3g_i(1 - \ln g_i) > p_1 - \sum_{j=1}^{i} 3x_i(1 - \ln x_i) \ge$$
(5.52)

(because $x \mapsto 3x(1 - \ln x)$ is increasing for $x \in (0, 1]$)

$$\geq p_1 - s(x_1) \geq p.$$
 (5.53)

Once we know that the recursion (5.36) is well-defined, we use the constructed sequences to prove the lemma. First, note that for $k \ge 1$,

$$r_k = r_1/(g_1g_2\cdots g_{k-1}).$$

Further, the above claim implies

$$g_{k-1}^2 \le g_{k-1}g_{k-2}^2 \le \dots \le g_{k-1}g_{k-2}\dots g_2g_1^2 = \frac{r_1}{r_k}g_1 = \frac{\delta^2}{r_k},$$
(5.54)

where $\delta = \sqrt{r_1 g_1}$. Now, let $r \ge r_1$. We have $r_k \xrightarrow[k \to \infty]{} \infty$ because $\frac{r_k}{r_{k+1}} = g_k \xrightarrow[k \to \infty]{} 0$, so for some $k, r_{k-1} \le r < r_k$. Then,

$$\tilde{g}_{p}(r) \le \tilde{g}_{p_{k-1}}(r) \le \tilde{g}_{p_{k-1}}(r_{k-1}) = g_{k-1} \le \frac{\delta}{\sqrt{r_k}} < \frac{\delta}{\sqrt{r}}$$
(5.55)

(from (5.54) and the monotonicity of $\tilde{g}_p(r)$ with regard to each of p and r), which finishes the proof.

Remark 5.26. As declared in Section 5, in this remark we summarise the differences between the proof of Theorem 3.6 and the proof of Theorem (5.4) in [Grim]:

- 1. First, the skeleton structure and most of the notation of the proof here is borrowed from [Grim]. The major notation that is different here, is " $A^{\delta}(r)$ " and " S_r " (respectively A_n and $\partial S(s)$ in [Grim]).
- 2. To be strict, the proper line of the proof borrowed from [Grim] starts by considering the functions f_p instead of g_p or \tilde{g}_p , although the functional inequality (5.2) involves both functions $f_{\cdot}(\cdot)$ and $\tilde{g}_{\cdot}(\cdot)$. In fact, each of the functions $f_{\cdot}(\cdot)$, $\tilde{g}_{\cdot}(\cdot)$ and $g_{\cdot}(\cdot)$ is a counterpart of the function $g_{\cdot}(\cdot)$ from [Grim] at some stage of the proof. After proving inequality (5.2), we pass to a couple of limits with it in order to obtain inequality (5.1) involving only $\tilde{g}_{\cdot}(\cdot)$ (the step not present in [Grim]). This form is needed to perform the repeated use of inequality (5.1) at the end of the proof of Theorem 3.6.
- 3. Obviously, the geometry used here is much different from that in [Grim]. In fact, we analyse the percolation cluster in $G^{(h,R)} \cap L_{\delta}$ using the pseudometric d_{\eth} (in place of the graph metric δ in [Grim]). Consequently, the set $\mathcal{R}^{(h,R)}$ of possible values of the random variables ϱ in Lemma 5.11 is much richer than \mathbb{N} , the respective set for the graph \mathbb{Z}^d . Moreover, the functions \tilde{g}_p arise from the percolation process on the whole $G^{(h,R)} \cap L$, so the distribution of the random variables M_i is not necessarily discrete. That cause the need for using integrals instead of sums, when concerned with those random variables, especially in the proof of Corollary 5.17. All that leads also to a few other minor technical differences between the proof here and the proof in [Grim].
- 4. The author tried to clarify the use of the assumption on $\sum_{i=1}^{k} r_i$ in Lemma 5.11 and why Wald's equation can be used in the proof of Lemma 5.19, which could be found quite hidden in [Grim].

5. The proof of Lemma 5.20 itself has a little changed structure (compared to the proof of Lemma (5.24) in [Grim]) and contains a proof of the convergence $\tilde{g}_p(r) \xrightarrow[r \to \infty]{} 0$ (Claim 5.24).

Remark 5.27. In order to prove Theorem 3.6, one could try to consider the percolation processes on the whole graph $G^{(h,R)} \cap L$ (without restricting it to L_{δ}) in order to obtain functional inequality similar to (5.1), involving only one function. That approach caused many difficulties to the author, some of which have not been overcome. Restricting the situation to L_{δ} makes the event A^{δ} depend on the states of only finitely many edges. That allows e.g. to condition the event $A^{\delta}(r) \cap B$ on the family of events { Γ a witness for B}, where Γ runs over a countable set (in the proof of Lemma 5.11) or to use BK inequality and Russo's formula.

Acknowledgements

References

[BB]	E. Babson, I. Benjamini, <i>Cut sets and normed cohomology with applications to percolation</i> (English summary), Proc. Amer. Math. Soc. 127 (1999), no. 2, pp. 589–597.
[BH]	M. R. Bridson, A. Haefliger, <i>Metric Spaces of Non-Positive Curvature</i> , Springer-Verlag, Berlin, 1999.
[BK89]	R. M. Burton, M. Keane, <i>Density and uniqueness in percolation</i> , Comm. Math. Phys. 121 (1989), no. 3, pp. 501–505.
[BS96]	I. Benjamini, O. Schramm, Percolation beyond \mathbf{Z}^d , many questions and a few answers (English summary), Electron. Comm. Probab. 1 (1996), no. 8, pp. 71–82 (electronic).
[BS01]	I. Benjamini, O. Schramm, <i>Percolation in the Hyperbolic Plane</i> , J. Amer. Math. Soc. 14 (2001), no. 2, pp. 487–507.
[Cz]	J. Czajkowski, Non-uniqueness phase of percolation on reflection groups in \mathbb{H}^3 , preprint in arXiv (http://arxiv.org/abs/1303.5624).
[Diest]	R. Diestel, <i>Graph Theory</i> , 4th electronic edition (free preview), 2010 (corrected reprint 2012).
[Grim]	G. Grimmett Percolation, Springer-Verlag, New York, 1999.
[GrimNewm]	G. Grimmett, C. M. Newman, Percolation in $\infty + 1$ dimensions, Disorder in physical systems, pp. 167–190, Oxford Sci. Publ., Oxford Univ. Press, New York, 1990.
[GrimSti]	G. Grimmett, D. R. Stirzaker, <i>Probability and Random Processes</i> , 2nd Edition, Oxford Univ. Press, Oxford, 1992.

[KapBen]	I. Kapovich, N. Benakli, <i>Boundaries of hyperbolic groups</i> (English summary), Combinatorial and geometric group theory (New York, 2000/Hoboken, NJ, 2001), pp. 39–93, Contemp. Math., 296, Amer. Math. Soc., Providence, RI, 2002.
[Lal]	S. P. Lalley, <i>Percolation on Fuchsian groups</i> , Annales de l'Institut Henri Poincaré (B) Probability and Statistics, Volume 34, Issue 2, 1998, pp. 151–177.
[LP]	R. Lyons with Y. Peres, <i>Probability on Trees and Networks</i> , Cambridge University Press (in preparation, version of 5 June 2013, current version available at http://mypage.iu.edu/~rdlyons/).
[Men]	M. V. Men'shikov, Sovpadenie kriticheskikh tochek v zadachakh perkolyat- sii (Russian, Coincidence of critical points in percolation problems), Dokl. Akad. Nauk SSSR, vol. 288 (1986), no. 6, pp. 1308–1311.
[NewmSchul]	C. M. Newman, L. S. Schulman, <i>Infinite clusters in percolation models</i> , J. Statist. Phys. 26 (1981), no. 3, pp. 613–628.
[PSN]	I. Pak, T. Smirnova-Nagnibeda, On non-uniqueness of percolation on non- amenable Cayley graphs (English, French summary), C. R. Acad. Sci. Paris Ser. I Math. 330 (2000), no. 6, pp. 495-500.
[Woe]	W. Woess, Random Walks on Infinite Graphs and Groups, Cambridge University Press, Cambridge, 2000.