

EFFECTIVE FILTERING FOR MULTISCALE STOCHASTIC DYNAMICAL SYSTEMS IN HILBERT SPACES*

HUIJIE QIAO

*School of Mathematics & Jiangsu Provincial Key Laboratory of Networked Collective Intelligence,
Southeast University, Nanjing, Jiangsu 211189, China
hjiaogean@seu.edu.cn*

ABSTRACT. In the paper, effective filtering for a type of slow-fast data assimilation systems in Hilbert spaces is considered. Firstly, the system is reduced to a system on a random invariant manifold. Secondly, nonlinear filtering of the origin system can be approximated by that of the reduction system. Finally, we apply the obtained result to an example.

1. INTRODUCTION

Give a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and two separable Hilbert spaces $\mathbb{H}^1, \mathbb{H}^2$ with the inner products $\langle \cdot, \cdot \rangle_{\mathbb{H}^1}, \langle \cdot, \cdot \rangle_{\mathbb{H}^2}$ and the norms $\| \cdot \|_{\mathbb{H}^1}, \| \cdot \|_{\mathbb{H}^2}$, respectively. Consider a stochastic slow-fast system on $\mathbb{H}^1 \times \mathbb{H}^2$

$$\begin{cases} \dot{x}^\varepsilon = Ax^\varepsilon + F(x^\varepsilon, y^\varepsilon) + \sigma_1 \dot{W}_1, \\ \dot{y}^\varepsilon = \frac{1}{\varepsilon}By^\varepsilon + \frac{1}{\varepsilon}G(x^\varepsilon, y^\varepsilon) + \frac{\sigma_2}{\sqrt{\varepsilon}}\dot{W}_2, \end{cases} \quad (1)$$

where A, B are two linear operators on $\mathbb{H}^1, \mathbb{H}^2$, respectively, and the interaction functions $F : \mathbb{H}^1 \times \mathbb{H}^2 \rightarrow \mathbb{H}^1$ and $G : \mathbb{H}^1 \times \mathbb{H}^2 \rightarrow \mathbb{H}^2$ are Borel measurable. Moreover, W_1, W_2 are two-sided $\mathbb{H}^1, \mathbb{H}^2$ -valued Brownian motions with covariance operators K_1, K_2 such that $\text{tr}K_1 < \infty, \text{tr}K_2 < \infty$, respectively, and mutually independent. σ_1 and σ_2 are nonzero real noise intensities, and ε is a small positive parameter representing the ratio of two time scales. The type of systems (1) have appeared in many fields, such as engineering and science([21]). For example, the climate evolution consists of fast atmospheric and slow oceanic dynamics, and state dynamic in electric power systems include fast- and slowly-varying elements.

The research for systems (1) is various. Let us mention some references. Schmalfuß-Schneider [22] observed the invariant manifold for systems (1) in finite dimensional Hilbert spaces $\mathbb{H}^1, \mathbb{H}^2$. When $\mathbb{H}^1, \mathbb{H}^2$ are infinite dimensional, and only the fast part contains a finite dimensional noise, Fu-Liu-Duan [9] studied the invariant manifold of systems (1). Stochastic average of systems (1) is considered in [12, 17]. By comparing the slow manifold reduction with the averaging principle reduction, we point out that the conditions of the

AMS Subject Classification(2010): 60H15, 37D10, 70K70.

Keywords: Multiscale systems in Hilbert spaces, random invariant manifolds, nonlinear filtering, dimensional reduction.

*This work was supported by NSF of China (No. 11001051, 11371352) and the Jiangsu Provincial Key Laboratory of Networked Collective Intelligence under Grant No. BM2017002.

former are shorter and the reduction system of the former is easier to simulate on the computer.

Fix a separable Hilbert space \mathbb{H}^3 with the inner product $\langle \cdot, \cdot \rangle_{\mathbb{H}^3}$ and the norm $\| \cdot \|_{\mathbb{H}^3}$. The nonlinear filtering problem for the slow component x_t^ε with respect to a \mathbb{H}^3 -valued observation process $\{r_s^\varepsilon, 0 \leq s \leq t\}$ (See Subsection 5.1 in details) is to evaluate the ‘filter’ $\mathbb{E}[\phi(x_t^\varepsilon) | \mathcal{R}_t^\varepsilon]$, where ϕ is a Borel measurable function such that $\mathbb{E}|\phi(x_t^\varepsilon)| < \infty$ for $t \in [0, T]$, and $\mathcal{R}_t^\varepsilon$ is the σ -algebra generated by $\{r_s^\varepsilon, 0 \leq s \leq t\}$. In essence, the nonlinear filtering is to achieve the best estimate for the origin system state, given only noisy observations for the system. It provides an algorithm for estimating a state of a random dynamical system based on noisy measurements. Stochastic filtering is important in many practical applications, from inertial guidance of aircrafts and spacecrafts to weather and climate prediction.

When $\mathbb{H}^1, \mathbb{H}^2, \mathbb{H}^3$ are finite dimensional, the nonlinear filtering problems of multiscale systems have been widely studied. Let us recall some results. In [11], Imkeller-Namachchivaya-Perkowski-Yeong showed that the filter $\mathbb{E}[\phi(x_t^\varepsilon) | \mathcal{R}_t^\varepsilon]$ converges to the homogenized filter by double backward stochastic differential equations and asymptotic techniques. Recently, Papanicolaou-Spiliopoulos [13] also studied this convergence problem by independent version technique and then applied it to statistical inference. When jumps processes are added in the system (1), the author proved the convergence by weak convergence technique in [18]. Besides, in [19] and [25], the author and two coauthors reduced the system (1) to a system on a random invariant manifold, and showed that $\mathbb{E}[\phi(x_t^\varepsilon) | \mathcal{R}_t^\varepsilon]$ converges to the filter of the reduction system. Thus, a new method to study the nonlinear filtering problem for multiscale systems is offered.

For a general system x^ε on a Hilbert space, that is to say, there is no fast component y^ε , its nonlinear filtering problem has been studied by Sritharan [23] and Hobbs-Sritharan [10]. However, for nonlinear filtering problems of multiscale systems on Hilbert spaces, nowadays there are no related results. Moreover, the type of problems have appeared in applications.(cf. [24])

In the paper, we consider a nonlinear filtering for the system (1) in general Hilbert spaces by following up the line in [19] and [25]. Firstly, the system is reduced to a system on a random invariant manifold. Moreover, our result covers the known result in [3, 9]. Secondly, the nonlinear filtering of the origin system can be approximated by that of the reduction system. And this result generalizes the result in [19].

It is worthwhile to mention our conditions and techniques. Firstly, the linear operators A and B may be unbounded, which can contain usual differential operators. Secondly, a general method of constructing random invariant manifolds for fast-slow systems is to transform those origin systems to non-random systems by stationary solutions of auxiliary equations and then construct random invariant manifolds for the obtained systems (c.f.[22, 19, 25], etc.). However, the existence of stationary solutions usually need strong conditions. Here we directly construct a random invariant manifold in terms of the system (1) itself. Therefore, our conditions are weaker than that in [22, 19, 25]. Finally, since these stochastic evolution equations on random slow manifolds have no Markov property, some techniques, such as the Zakai equations in [14, 15, 16] and backward stochastic differential equations in [11], do not work. Therefore, we use an exponential martingale technique to treat these nonlinear filtering problems.

This paper is arranged as follows. In Section 2, we introduce basic concepts about random dynamical systems and random invariant manifolds. We present the framework of the probability space and the system in Section 3. Our method for reduced filtering is placed in Section 4. In Section 5, the nonlinear filtering problem is introduced and the approximation theorem of the filtering is proved. And then, we apply the obtained result to an example in Section 6. In Section 7, we summarize all the results in the paper. Finally, the proof of Theorem 3.3 is placed in Section 8.

The following convention will be used throughout the paper: C with or without indices will denote different positive constants (depending on the indices) whose values may change from one place to another.

2. PRELIMINARIES

In the section, we introduce some notations and basic concepts in random dynamical systems.

2.1. Notation and terminology. Let $\mathcal{B}(\mathbb{H}^1)$ be the Borel σ -algebra on \mathbb{H}^1 , and $\mathcal{B}(\mathbb{H}^1)$ be the set of all real-valued bounded Borel-measurable functions on \mathbb{H}^1 . Let $\mathcal{C}(\mathbb{H}^1)$ be the set of all real-valued continuous functions on \mathbb{H}^1 , and $\mathcal{C}_b^1(\mathbb{H}^1)$ denote the collection of all functions of $\mathcal{C}(\mathbb{H}^1)$ which are bounded and Lipschitz continuous. And set

$$\|\phi\| := \max_{x \in \mathbb{H}^1} |\phi(x)| + \max_{x_1 \neq x_2} \frac{|\phi(x_1) - \phi(x_2)|}{\|x_1 - x_2\|_{\mathbb{H}^1}}, \quad \phi \in \mathcal{C}_b^1(\mathbb{H}^1).$$

2.2. Random dynamical systems ([1]). Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, and $(\theta_t)_{t \in \mathbb{R}}$ a family of measurable transformations from Ω to Ω satisfying for $s, t \in \mathbb{R}$,

$$\theta_0 = 1_\Omega, \quad \theta_{t+s} = \theta_t \circ \theta_s. \quad (2)$$

If for each $t \in \mathbb{R}$, θ_t preserves the probability measure \mathbb{P} , i.e.,

$$\theta_t^* \mathbb{P} = \mathbb{P},$$

$(\Omega, \mathcal{F}, \mathbb{P}; (\theta_t)_{t \in \mathbb{R}})$ is called a metric dynamical system.

Definition 2.1. Let $(\mathbb{X}, \mathcal{X})$ be a measurable space. A mapping

$$\varphi : \mathbb{R} \times \Omega \times \mathbb{X} \mapsto \mathbb{X}, \quad (t, \omega, y) \mapsto \varphi(t, \omega, y)$$

is called a measurable random dynamical system (RDS), or in short, a cocycle, if these following properties hold:

- (i) *Measurability:* φ is $\mathcal{B}(\mathbb{R}) \otimes \mathcal{F} \otimes \mathcal{X} / \mathcal{X}$ -measurable,
- (ii) *Cocycle property:* $\varphi(t, \omega)$ satisfies the following conditions

$$\varphi(0, \omega) = id_{\mathbb{X}},$$

and for $\omega \in \Omega$ and all $s, t \in \mathbb{R}$

$$\varphi(t + s, \omega) = \varphi(t, \theta_s \omega) \circ \varphi(s, \omega),$$

- (iii) *Continuity:* $\varphi(t, \omega)$ is continuous for $t \in \mathbb{R}$.

2.3. Random invariant manifolds ([22]). Let φ be a random dynamical system on the normed space $(\mathbb{X}, \|\cdot\|_{\mathbb{X}})$.

A family of nonempty sets $\mathcal{M} = \{\mathcal{M}(\omega)\}_{\omega \in \Omega}$ is called a *random set* in \mathbb{X} if for $\omega \in \Omega$, $\mathcal{M}(\omega)$ is a closed set in \mathbb{X} and for every $y \in \mathbb{X}$, the mapping

$$\Omega \ni \omega \mapsto \text{dist}(y, \mathcal{M}(\omega)) := \inf_{x \in \mathcal{M}(\omega)} \|x - y\|_{\mathbb{X}}$$

is measurable. Moreover, if \mathcal{M} satisfies

$$\varphi(t, \omega, \mathcal{M}(\omega)) \subset \mathcal{M}(\theta_t \omega), \quad t \geq 0, \quad \omega \in \Omega,$$

\mathcal{M} is called (positively) invariant with respect to φ .

In the sequel, we consider a random set defined by a Lipschitz continuous function. Concretely speaking, we will define a mapping by

$$\Omega \times \mathbb{H}^1 \ni (\omega, x) \mapsto H(\omega, x) \in \mathbb{H}^2,$$

where for all $\omega \in \Omega$, $H(\omega, x)$ is globally Lipschitzian in x and for any $x \in \mathbb{H}^1$, $\omega \mapsto H(\omega, x)$ is a \mathbb{H}^2 -valued random variable. Thus, set

$$\mathcal{M}(\omega) := \{(x, H(\omega, x)) | x \in \mathbb{H}^1\},$$

and then \mathcal{M} is a random set in $\mathbb{H}^1 \times \mathbb{H}^2$ ([22, Lemma 2.1]). Moreover, the random set $\mathcal{M}(\omega)$ is called a *Lipschitz random invariant manifold* if it is (positively) invariant with respect to some random dynamical system.

3. FRAMEWORK

In the section, we present some results which will be applied in the following sections.

Let $\Omega^1 := C_0(\mathbb{R}, \mathbb{H}^1)$ be the collection of all strongly continuous functions $f : \mathbb{R} \rightarrow \mathbb{H}^1$ with $f(0) = 0$. And then it is equipped with the compact-open topology. Let \mathcal{F}^1 be its Borel σ -algebra and \mathbb{P}^1 the distribution of W_1 on Ω^1 . Set

$$\theta_t^1 \omega_1(\cdot) := \omega_1(\cdot + t) - \omega_1(t), \quad \omega_1 \in \Omega^1, \quad t \in \mathbb{R},$$

and then $(\theta_t^1)_{t \in \mathbb{R}}$ satisfy (2). Moreover, by the property of \mathbb{P}^1 we obtain that $(\Omega^1, \mathcal{F}^1, \mathbb{P}^1, \theta_t^1)$ is a metric dynamical system. Next, set $\Omega^2 := C_0(\mathbb{R}, \mathbb{H}^2)$. And then we define $\mathcal{F}^2, \mathbb{P}^2, \theta_t^2$ by the similar means to $\mathcal{F}^1, \mathbb{P}^1, \theta_t^1$. Thus, $(\Omega^2, \mathcal{F}^2, \mathbb{P}^2, \theta_t^2)$ becomes another metric dynamical system. Set

$$\Omega := \Omega^1 \times \Omega^2, \quad \mathcal{F} := \mathcal{F}^1 \times \mathcal{F}^2, \quad \mathbb{P} := \mathbb{P}^1 \times \mathbb{P}^2, \quad \theta_t := \theta_t^1 \times \theta_t^2,$$

and then $(\Omega, \mathcal{F}, \mathbb{P}, \theta_t)$ is a metric dynamical system that is used in the sequel.

Consider the slow-fast system (1) on $\mathbb{H}^1 \times \mathbb{H}^2$, i.e.

$$\begin{cases} \dot{x}^\varepsilon = Ax^\varepsilon + F(x^\varepsilon, y^\varepsilon) + \sigma_1 \dot{W}_1, \\ \dot{y}^\varepsilon = \frac{1}{\varepsilon} By^\varepsilon + \frac{1}{\varepsilon} G(x^\varepsilon, y^\varepsilon) + \frac{\sigma_2}{\sqrt{\varepsilon}} \dot{W}_2. \end{cases}$$

We make the following hypotheses:

(H₁) There exists a $\gamma_1 \geq 0$ such that

$$\|e^{At}\| \leq e^{-\gamma_1 t}, \quad t \leq 0, \tag{3}$$

where $\|e^{At}\|$ stands for the norm of the operator e^{At} , and $\{e^{At}, t \geq 0\}$ is a strongly continuous semigroup on \mathbb{H}^1 and

$$\|e^{At}\| \leq 1, \quad t \geq 0. \quad (4)$$

(H₂) There exists a $\gamma_2 > 0$ such that for any $y \in \mathbb{H}^2$,

$$\langle By, y \rangle_{\mathbb{H}^2} \leq -\gamma_2 \|y\|_{\mathbb{H}^2}^2.$$

(H₃) There exists a positive constant L such that for all $(x_1, y_1), (x_2, y_2) \in \mathbb{H}^1 \times \mathbb{H}^2$

$$\|F(x_1, y_1) - F(x_2, y_2)\|_{\mathbb{H}^1} \leq L(\|x_1 - x_2\|_{\mathbb{H}^1} + \|y_1 - y_2\|_{\mathbb{H}^2}),$$

and

$$\|G(x_1, y_1) - G(x_2, y_2)\|_{\mathbb{H}^2} \leq L(\|x_1 - x_2\|_{\mathbb{H}^1} + \|y_1 - y_2\|_{\mathbb{H}^2}),$$

and $F(0, 0) = G(0, 0) = 0$.

(H₄)

$$\gamma_2 > L.$$

Remark 3.1. By (H₂), we know that $\frac{B}{\varepsilon}$ generates a strongly continuous semigroup $\{e^{\frac{B}{\varepsilon}t}, t \geq 0\}$ on \mathbb{H}^2 and

$$\|e^{\frac{B}{\varepsilon}t}\| \leq e^{-\frac{\gamma_2}{\varepsilon}t}, \quad t \geq 0. \quad (5)$$

And (H₃) admits us to obtain that for any $x \in \mathbb{H}^1, y \in \mathbb{H}^2$

$$\|F(x, y)\|_{\mathbb{H}^1} \leq L(\|x\|_{\mathbb{H}^1} + \|y\|_{\mathbb{H}^2}), \quad \|G(x, y)\|_{\mathbb{H}^2} \leq L(\|x\|_{\mathbb{H}^1} + \|y\|_{\mathbb{H}^2}). \quad (6)$$

In the following, we give the definition of mild solutions to the system (1) and then state that the system (1) has a unique mild solution which generates a RDS. Let $\mathbb{H} := \mathbb{H}^1 \times \mathbb{H}^2$ with the norm $\|z\|_{\mathbb{H}} = \|x\|_{\mathbb{H}^1} + \|y\|_{\mathbb{H}^2}$ for $z = (x, y) \in \mathbb{H}$. Let $\mathcal{C}([a, b], \mathbb{H})$ be the collection of strongly continuous functions on $[a, b]$ with values in \mathbb{H} .

Definition 3.2. Let $s \in \mathbb{R}, T > 0$ and $z_0 = (x_0, y_0) \in \mathbb{H}$. $z^\varepsilon(t) \equiv z^\varepsilon(t, s, \omega; z_0)$ is said to be a mild solution to the system (1) on the interval $(s, s + T]$ if (i) $z^\varepsilon(s) = z_0$, (ii) $z^\varepsilon(t)$ satisfies the following integral equation

$$z^\varepsilon(t) = \begin{pmatrix} x_t^\varepsilon \\ y_t^\varepsilon \end{pmatrix} = \begin{pmatrix} e^{A(t-s)}x_0 + \int_s^t e^{A(t-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr + \int_s^t e^{A(t-r)}\sigma_1 dW_1(r) \\ e^{\frac{B}{\varepsilon}(t-s)}y_0 + \int_s^t e^{\frac{B}{\varepsilon}(t-r)}\frac{1}{\varepsilon}G(x_r^\varepsilon, y_r^\varepsilon)dr + \int_s^t e^{\frac{B}{\varepsilon}(t-r)}\frac{\sigma_2}{\sqrt{\varepsilon}}dW_2(r) \end{pmatrix}$$

for $t \in [s, s + T]$ and $\omega \in \Omega$, (iii) $z^\varepsilon(t)$ belongs to $\mathcal{C}([s, s + T], \mathbb{H})$.

Theorem 3.3. Suppose that (H₁)-(H₄) are satisfied. Let $s \in \mathbb{R}, T > 0$ and $z_0 = (x_0, y_0) \in \mathbb{H}$. Then the system (1) has a unique mild solution $z^\varepsilon(t, s, \omega; z_0)$ for $t \in [s, s + T]$ and $\omega \in \Omega$. Moreover, set $\varphi^\varepsilon(t, \omega)z_0 := z^\varepsilon(t, 0, \omega; z_0), t \geq 0$, and then $\varphi^\varepsilon(t, \omega)$ is a RDS.

The above theorem is similar to Theorem 3.3 in [3]. To the readers' convenience, we give a proof in the Appendix.

4. REDUCTION ON RANDOM INVARIANT MANIFOLDS

In the section, we prove that the origin system can be approximated by the reduction system on a random invariant manifold.

4.1. Random invariant manifolds. In the subsection, we prove that the system (1) has a random invariant manifold. Firstly, we introduce some spaces. Let

$$\begin{aligned}\mathcal{C}_{\frac{\mu}{\varepsilon},s}^{1,-} &:= \left\{ \phi \in \mathcal{C}((-\infty, s], \mathbb{H}^1) : \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\phi(t)\|_{\mathbb{H}^1} < \infty \right\}, \\ \mathcal{C}_{\frac{\mu}{\varepsilon},s}^{2,-} &:= \left\{ \phi \in \mathcal{C}((-\infty, s], \mathbb{H}^2) : \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\phi(t)\|_{\mathbb{H}^2} < \infty \right\}, \\ \mathcal{C}_{\frac{\mu}{\varepsilon}}^1 &:= \left\{ \phi \in \mathcal{C}(\mathbb{R}, \mathbb{H}^1) : \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon}t} \|\phi(t)\|_{\mathbb{H}^1} < \infty \right\}, \\ \mathcal{C}_{\frac{\mu}{\varepsilon}}^2 &:= \left\{ \phi \in \mathcal{C}(\mathbb{R}, \mathbb{H}^2) : \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon}t} \|\phi(t)\|_{\mathbb{H}^2} < \infty \right\},\end{aligned}$$

where μ is a positive constant and satisfies $\gamma_2 - L > \mu > 0$. Let $\mathcal{C}_{\frac{\mu}{\varepsilon},s}^{-} := \mathcal{C}_{\frac{\mu}{\varepsilon},s}^{1,-} \times \mathcal{C}_{\frac{\mu}{\varepsilon},s}^{2,-}$ with the norm $\|z\|_{\mathcal{C}_{\frac{\mu}{\varepsilon},s}^{-}} = \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|z(t)\|_{\mathbb{H}}$ for $z \in \mathcal{C}_{\frac{\mu}{\varepsilon},s}^{-}$, and $\mathcal{C}_{\frac{\mu}{\varepsilon}} := \mathcal{C}_{\frac{\mu}{\varepsilon}}^1 \times \mathcal{C}_{\frac{\mu}{\varepsilon}}^2$ with the norm $\|z\|_{\mathcal{C}_{\frac{\mu}{\varepsilon}}} = \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon}t} \|z(t)\|_{\mathbb{H}}$ for $z \in \mathcal{C}_{\frac{\mu}{\varepsilon}}$.

Lemma 4.1. *Suppose that (\mathbf{H}_1) – (\mathbf{H}_4) are satisfied. Let $s \in \mathbb{R}$ and $x_0 \in \mathbb{H}^1$. Then there exists a $\varepsilon_0 > 0$ such that for $0 < \varepsilon \leq \varepsilon_0$, the following integral equation has a unique solution $\bar{z}_t^\varepsilon = (\bar{x}_t^\varepsilon, \bar{y}_t^\varepsilon)$ for $t \leq s$:*

$$\bar{z}_t^\varepsilon = \begin{pmatrix} \bar{x}_t^\varepsilon \\ \bar{y}_t^\varepsilon \end{pmatrix} = \begin{pmatrix} e^{A(t-s)}x_0 - \int_t^s e^{A(t-r)}F(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon)dr - \int_t^s e^{A(t-r)}\sigma_1 dW_1(r) \\ \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)}\frac{1}{\varepsilon}G(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon)dr + \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)}\frac{\sigma_2}{\sqrt{\varepsilon}}dW_2(r) \end{pmatrix}, t \leq s, \quad (7)$$

$$\bar{x}_s^\varepsilon = x_0.$$

Proof. Set for $\bar{z}^\varepsilon = (\bar{x}^\varepsilon, \bar{y}^\varepsilon) \in \mathcal{C}_{\frac{\mu}{\varepsilon},s}^{-}$

$$\mathcal{K}(\bar{z}^\varepsilon)(t) := \begin{pmatrix} \mathcal{K}_1(\bar{z}^\varepsilon)(t) \\ \mathcal{K}_2(\bar{z}^\varepsilon)(t) \end{pmatrix} := \begin{pmatrix} e^{A(t-s)}x_0 - \int_t^s e^{A(t-r)}F(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon)dr - \int_t^s e^{A(t-r)}\sigma_1 dW_1(r) \\ \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)}\frac{1}{\varepsilon}G(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon)dr + \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)}\frac{\sigma_2}{\sqrt{\varepsilon}}dW_2(r) \end{pmatrix}, \quad t \leq s, \quad (8)$$

and then \mathcal{K} is well defined on $\mathcal{C}_{\frac{\mu}{\varepsilon},s}^{-}$. Indeed, we calculate that for $\bar{z}^\varepsilon = (\bar{x}^\varepsilon, \bar{y}^\varepsilon) \in \mathcal{C}_{\frac{\mu}{\varepsilon},s}^{-}$,

$$\begin{aligned}\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|e^{A(t-s)}x_0\|_{\mathbb{H}^1} &\leq \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} e^{-\gamma_1(t-s)} \|x_0\|_{\mathbb{H}^1} \leq \|x_0\|_{\mathbb{H}^1}, \\ \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \left\| \int_t^s e^{A(t-r)}F(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon)dr \right\|_{\mathbb{H}^1} &\leq \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \int_t^s e^{-\gamma_1(t-r)} L(\|\bar{x}_r^\varepsilon\|_{\mathbb{H}^1} + \|\bar{y}_r^\varepsilon\|_{\mathbb{H}^2})dr \\ &\leq L \left(\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\bar{z}_t^\varepsilon\|_{\mathbb{H}} \right) \sup_{t \leq s} \int_t^s e^{(\frac{\mu}{\varepsilon} - \gamma_1)(t-r)}dr \\ &= \frac{\varepsilon L}{\mu - \varepsilon \gamma_1} \left(\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\bar{z}_t^\varepsilon\|_{\mathbb{H}} \right), \quad a.s., \quad (9)\end{aligned}$$

and

$$\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \left\| \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)}\frac{1}{\varepsilon}G(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon)dr \right\|_{\mathbb{H}^2} \leq \frac{1}{\varepsilon} \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \int_{-\infty}^t e^{-\frac{\gamma_2}{\varepsilon}(t-r)} L(\|\bar{x}_r^\varepsilon\|_{\mathbb{H}^1} + \|\bar{y}_r^\varepsilon\|_{\mathbb{H}^2})dr$$

$$\begin{aligned}
&\leq \frac{L}{\varepsilon} \left(\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\bar{z}_t^\varepsilon\|_{\mathbb{H}} \right) \int_{-\infty}^t e^{(\frac{\mu}{\varepsilon} - \frac{\gamma_2}{\varepsilon})(t-r)} dr \\
&= \frac{L}{\gamma_2 - \mu} \left(\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\bar{z}_t^\varepsilon\|_{\mathbb{H}} \right), \quad a.s.. \tag{10}
\end{aligned}$$

By [5, Proposition 3.1], it holds that

$$\begin{aligned}
\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \left\| \int_t^s e^{A(t-r)} \sigma_1 dW_1(r) \right\|_{\mathbb{H}^1} &< \infty, \quad a.s., \\
\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \left\| \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)} \frac{\sigma_2}{\sqrt{\varepsilon}} dW_2(r) \right\|_{\mathbb{H}^2} &< \infty, \quad a.s..
\end{aligned}$$

In the following, we show that \mathcal{K} is contractive. For $\bar{z}^{\varepsilon,1}, \bar{z}^{\varepsilon,2} \in \mathcal{C}_{\frac{\mu}{\varepsilon},s}^-$, by the same deduction to (9) (10), one can obtain that

$$\begin{aligned}
\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\mathcal{K}_1(\bar{z}^{\varepsilon,1})(t) - \mathcal{K}_1(\bar{z}^{\varepsilon,2})(t)\|_{\mathbb{H}^1} &\leq \frac{\varepsilon L}{\mu - \varepsilon \gamma_1} \left(\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\bar{z}_t^{\varepsilon,1} - \bar{z}_t^{\varepsilon,2}\|_{\mathbb{H}} \right), \\
\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\mathcal{K}_2(\bar{z}^{\varepsilon,1})(t) - \mathcal{K}_2(\bar{z}^{\varepsilon,2})(t)\|_{\mathbb{H}^2} &\leq \frac{L}{\gamma_2 - \mu} \left(\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\bar{z}_t^{\varepsilon,1} - \bar{z}_t^{\varepsilon,2}\|_{\mathbb{H}} \right).
\end{aligned}$$

Thus, we get that

$$\begin{aligned}
\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\mathcal{K}(\bar{z}^{\varepsilon,1})(t) - \mathcal{K}(\bar{z}^{\varepsilon,2})(t)\|_{\mathbb{H}} &\leq \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\mathcal{K}_1(\bar{z}^{\varepsilon,1})(t) - \mathcal{K}_1(\bar{z}^{\varepsilon,2})(t)\|_{\mathbb{H}^1} \\
&\quad + \sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\mathcal{K}_2(\bar{z}^{\varepsilon,1})(t) - \mathcal{K}_2(\bar{z}^{\varepsilon,2})(t)\|_{\mathbb{H}^2} \\
&\leq \left(\frac{\varepsilon L}{\mu - \varepsilon \gamma_1} + \frac{L}{\gamma_2 - \mu} \right) \left(\sup_{t \leq s} e^{\frac{\mu}{\varepsilon}(t-s)} \|\bar{z}_t^{\varepsilon,1} - \bar{z}_t^{\varepsilon,2}\|_{\mathbb{H}} \right).
\end{aligned}$$

Therefore, in order to prove that \mathcal{K} is contractive, one only needs to show that

$$\frac{\varepsilon L}{\mu - \varepsilon \gamma_1} + \frac{L}{\gamma_2 - \mu} < 1.$$

Note that $\gamma_2 - L > \mu > 0$, and then

$$\frac{L}{\gamma_2 - \mu} < 1.$$

Thus, there exists a $\varepsilon_0 > 0$ such that for any $0 < \varepsilon \leq \varepsilon_0$,

$$\frac{\varepsilon L}{\mu - \varepsilon \gamma_1} + \frac{L}{\gamma_2 - \mu} < 1,$$

and furthermore \mathcal{K} is contractive. So, Eq.(7) has a unique solution. The proof is completed. \square

Next, we study a property of the solution to Eq.(7). First of all, for $u \in (-\infty, s)$, Eq.(7) can be rewritten as

$$\begin{pmatrix} \bar{x}_t^\varepsilon \\ \bar{y}_t^\varepsilon \end{pmatrix} = \begin{pmatrix} e^{A(t-u)} \bar{x}_u^\varepsilon + \int_u^t e^{A(t-r)} F(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon) dr + \int_u^t e^{A(t-r)} \sigma_1 dW_1(r) \\ e^{\frac{B}{\varepsilon}(t-u)} \bar{y}_u^\varepsilon + \int_u^t e^{\frac{B}{\varepsilon}(t-r)} \frac{1}{\varepsilon} G(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon) dr + \int_u^t e^{\frac{B}{\varepsilon}(t-r)} \frac{\sigma_2}{\sqrt{\varepsilon}} dW_2(r) \end{pmatrix}, u \leq t \leq s.$$

That is, the dynamic of Eq.(7) is the same as that of the system (1). Thus, by Theorem 3.3 it holds that for $r \leq s$

$$\bar{x}_r^\varepsilon(\theta_u \omega) = \bar{x}_{r+u}^\varepsilon(\omega), \quad \bar{y}_r^\varepsilon(\theta_u \omega) = \bar{y}_{r+u}^\varepsilon(\omega), \quad u \leq s - r. \quad (11)$$

Remark 4.2. Set

$$H^{\varepsilon,s}(\omega, x_0) := \bar{y}_s^\varepsilon = \int_{-\infty}^s e^{\frac{B}{\varepsilon}(s-r)} \frac{1}{\varepsilon} G(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon) dr + \int_{-\infty}^s e^{\frac{B}{\varepsilon}(s-r)} \frac{\sigma_2}{\sqrt{\varepsilon}} dW_2(r),$$

and then for $x_0^1, x_0^2 \in \mathbb{H}^1$, it holds that

$$\|H^{\varepsilon,s}(\omega, x_0^1) - H^{\varepsilon,s}(\omega, x_0^2)\|_{\mathbb{H}^2} \leq \frac{L e^{\frac{\mu}{\varepsilon}s}}{(\gamma_2 - \mu) \left[1 - \left(\frac{\varepsilon L}{\mu - \varepsilon \gamma_1} + \frac{L}{\gamma_2 - \mu} \right) \right]} \|x_0^1 - x_0^2\|_{\mathbb{H}^1}, \quad (12)$$

where we use (9) (10). Besides, it follows from (11) that for $t \leq s$,

$$\begin{aligned} H^{\varepsilon,s}(\theta_{t-s} \omega, x_0) &= \int_{-\infty}^s e^{\frac{B}{\varepsilon}(s-r)} \frac{1}{\varepsilon} G(\bar{x}_r^\varepsilon(\theta_{t-s} \omega), \bar{y}_r^\varepsilon(\theta_{t-s} \omega)) dr \\ &\quad + \int_{-\infty}^s e^{\frac{B}{\varepsilon}(s-r)} \frac{\sigma_2}{\sqrt{\varepsilon}} dW_2(r)(\theta_{t-s} \omega) \\ &= \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)} \frac{1}{\varepsilon} G(\bar{x}_r^\varepsilon, \bar{y}_r^\varepsilon) dr + \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)} \frac{\sigma_2}{\sqrt{\varepsilon}} dW_2(r) \\ &= \bar{y}_t^\varepsilon. \end{aligned} \quad (13)$$

Theorem 4.3. (A random invariant manifold)

Assume that $(\mathbf{H}_1) - (\mathbf{H}_4)$ are satisfied. Let $z_0 = (x_0, y_0) \in \mathbb{H}$. Then for $0 < \varepsilon \leq \varepsilon_0$, φ^ε has a random invariant manifold

$$\mathcal{M}^\varepsilon(\omega) = \{ (x, H^{\varepsilon,0}(\omega, x)), x \in \mathbb{H}^1 \},$$

where for $\omega \in \Omega$, the Lipschitz constant of $H^{\varepsilon,0}(\omega, x)$ is bounded by

$$\frac{L}{(\gamma_2 - \mu) \left[1 - \left(\frac{\varepsilon L}{\mu - \varepsilon \gamma_1} + \frac{L}{\gamma_2 - \mu} \right) \right]}.$$

Moreover, \mathcal{M}^ε is exponentially attracting in the following sense: for any $z_0 = (x_0, y_0)$, there exists a $\tilde{z}_0 = (\tilde{x}_0, \tilde{y}_0) \in \mathcal{M}^\varepsilon(\omega)$ such that

$$\|\varphi^\varepsilon(t, \omega) z_0 - \varphi^\varepsilon(t, \omega) \tilde{z}_0\|_{\mathbb{H}} \leq \frac{e^{-\frac{\mu}{\varepsilon}t}}{1 - M} \left((2 + 2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right), \quad t \geq 0, \quad (14)$$

where

$$M := \frac{\varepsilon L}{\mu - \varepsilon \gamma_1} + \frac{L}{\gamma_2 - \mu},$$

$$R(\omega) := \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \left\| \int_t^0 e^{A(t-r)} \sigma_1 dW_1(r) \right\|_{\mathbb{H}^1} + \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \left\| \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)} \frac{\sigma_2}{\sqrt{\varepsilon}} dW_2(r) \right\|_{\mathbb{H}^2}.$$

Proof. First of all, set

$$\mathcal{M}^\varepsilon(\omega) := \{ (x, H^{\varepsilon,0}(\omega, x)), x \in \mathbb{H}^1 \},$$

and then by (12), it holds that $\mathcal{M}^\varepsilon(\omega)$ is a random set. And based on (13) one can justify that $\mathcal{M}^\varepsilon(\omega)$ is invariant with respect to φ^ε . Therefore, $\mathcal{M}^\varepsilon(\omega)$ is a Lipschitz random invariant manifold with respect to φ^ε .

Next, we prove (14). For this, set

$$z_t^\varepsilon = \begin{pmatrix} x_t^\varepsilon \\ y_t^\varepsilon \end{pmatrix} := \begin{cases} \begin{pmatrix} x_0, (I - |t|B)^{-1}y_0 \end{pmatrix}, & t \leq 0, \\ \varphi^\varepsilon(t, \omega)(x_0, y_0), & t > 0, \end{cases}$$

and

$$Z_0(t) := \begin{cases} -z_t^\varepsilon + \mathcal{K}(z^\varepsilon)(t), & t \leq 0, \\ -(e^{At}x_0, e^{\frac{B}{\varepsilon}t}y_0) + (e^{At}x_0, e^{\frac{B}{\varepsilon}t}\mathcal{K}_2(z^\varepsilon)(0)), & t > 0, \end{cases}$$

where \mathcal{K} is defined in (8). And then we consider the following integral equation

$$\begin{pmatrix} X_t^\varepsilon \\ Y_t^\varepsilon \end{pmatrix} = Z_0(t) + \begin{pmatrix} -\int_t^\infty e^{A(t-r)} \left[F(x_r^\varepsilon + X_r^\varepsilon, y_r^\varepsilon + Y_r^\varepsilon) - F(x_r^\varepsilon, y_r^\varepsilon) \right] dr \\ \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)} \frac{1}{\varepsilon} \left[G(x_r^\varepsilon + X_r^\varepsilon, y_r^\varepsilon + Y_r^\varepsilon) - G(x_r^\varepsilon, y_r^\varepsilon) \right] dr \end{pmatrix}, t \in \mathbb{R}. \quad (15)$$

For $Z^\varepsilon := (X^\varepsilon, Y^\varepsilon) \in \mathcal{C}_\varepsilon^\mu$, set

$$\begin{aligned} \mathcal{R}(Z^\varepsilon)(t) &:= \begin{pmatrix} \mathcal{R}_1(Z^\varepsilon)(t) \\ \mathcal{R}_2(Z^\varepsilon)(t) \end{pmatrix} \\ &:= Z_0(t) + \begin{pmatrix} -\int_t^\infty e^{A(t-r)} \left[F(x_r^\varepsilon + X_r^\varepsilon, y_r^\varepsilon + Y_r^\varepsilon) - F(x_r^\varepsilon, y_r^\varepsilon) \right] dr \\ \int_{-\infty}^t e^{\frac{B}{\varepsilon}(t-r)} \frac{1}{\varepsilon} \left[G(x_r^\varepsilon + X_r^\varepsilon, y_r^\varepsilon + Y_r^\varepsilon) - G(x_r^\varepsilon, y_r^\varepsilon) \right] dr \end{pmatrix}, \end{aligned}$$

and then $\mathcal{R} : \mathcal{C}_\varepsilon^\mu \mapsto \mathcal{C}_\varepsilon^\mu$ is well defined. Indeed, for $Z^\varepsilon = (X^\varepsilon, Y^\varepsilon) \in \mathcal{C}_\varepsilon^\mu$, by (\mathbf{H}_1) – (\mathbf{H}_4) we compute

$$\begin{aligned} \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon}t} \|Z_0(t)\|_{\mathbb{H}} &\leq \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \|Z_0(t)\|_{\mathbb{H}} + \sup_{t > 0} e^{\frac{\mu}{\varepsilon}t} \|Z_0(t)\|_{\mathbb{H}} \\ &\leq \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} (\|z_t^\varepsilon\|_{\mathbb{H}} + \|\mathcal{K}(z^\varepsilon)(t)\|_{\mathbb{H}}) + \sup_{t > 0} e^{\frac{\mu}{\varepsilon}t} e^{-\frac{\gamma_2}{\varepsilon}t} (\|y_0\|_{\mathbb{H}^2} + \|\mathcal{K}_2(z^\varepsilon)(0)\|_{\mathbb{H}^2}) \\ &\leq \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \|z_t^\varepsilon\|_{\mathbb{H}} + \|x_0\|_{\mathbb{H}^1} + M \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \|z_t^\varepsilon\|_{\mathbb{H}} + R(\omega) + \|y_0\|_{\mathbb{H}^2} \\ &\quad + \|\mathcal{K}_2(z^\varepsilon)(0)\|_{\mathbb{H}^2} \\ &\leq \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \|z_t^\varepsilon\|_{\mathbb{H}} + \|x_0\|_{\mathbb{H}^1} + M \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \|z_t^\varepsilon\|_{\mathbb{H}} + R(\omega) + \|y_0\|_{\mathbb{H}^2} \\ &\quad + M \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \|z_t^\varepsilon\|_{\mathbb{H}} + R(\omega) \\ &\leq (1 + 2M) \sup_{t \leq 0} e^{\frac{\mu}{\varepsilon}t} \|z_t^\varepsilon\|_{\mathbb{H}} + \|x_0\|_{\mathbb{H}^1} + \|y_0\|_{\mathbb{H}^2} + 2R(\omega) \\ &\leq (2 + 2M)(\|x_0\|_{\mathbb{H}^1} + \|y_0\|_{\mathbb{H}^2}) + 2R(\omega) \\ &= (2 + 2M)\|z_0\|_{\mathbb{H}} + 2R(\omega). \end{aligned} \quad (16)$$

Besides, it follows from (\mathbf{H}_1) – (\mathbf{H}_4) that

$$\begin{aligned} &\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon}t} \int_t^\infty e^{-\gamma_1(t-r)} \|F(x_r^\varepsilon + X_r^\varepsilon, y_r^\varepsilon + Y_r^\varepsilon) - F(x_r^\varepsilon, y_r^\varepsilon)\|_{\mathbb{H}^1} dr \\ &\leq L \left(\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon}t} \|Z_t^\varepsilon\|_{\mathbb{H}} \right) \sup_{t \in \mathbb{R}} \int_t^\infty e^{(\frac{\mu}{\varepsilon} - \gamma_1)(t-r)} dr \end{aligned}$$

$$\leq \frac{\varepsilon L}{\mu - \varepsilon \gamma_1} \left(\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|Z_t^\varepsilon\|_{\mathbb{H}} \right), \quad (17)$$

and

$$\begin{aligned} & \frac{1}{\varepsilon} \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \int_{-\infty}^t e^{-\frac{\gamma_2}{\varepsilon}(t-r)} \|G(x_r^\varepsilon + X_r^\varepsilon, y_r^\varepsilon + Y_r^\varepsilon) - G(x_r^\varepsilon, y_r^\varepsilon)\|_{\mathbb{H}^2} dr \\ & \leq \frac{L}{\varepsilon} \left(\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|Z_t^\varepsilon\|_{\mathbb{H}} \right) \sup_{t \in \mathbb{R}} \int_{-\infty}^t e^{(\frac{\mu}{\varepsilon} - \frac{\gamma_2}{\varepsilon})(t-r)} dr \\ & \leq \frac{L}{\gamma_2 - \mu} \left(\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|Z_t^\varepsilon\|_{\mathbb{H}} \right). \end{aligned} \quad (18)$$

Thus, by combining (16) (17) with (18), one can get that

$$\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|\mathcal{R}(Z^\varepsilon)(t)\|_{\mathbb{H}} \leq \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|\mathcal{R}_1(Z^\varepsilon)(t)\|_{\mathbb{H}^1} + \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|\mathcal{R}_2(Z^\varepsilon)(t)\|_{\mathbb{H}^2} < \infty.$$

Next, for $Z^{\varepsilon,1}, Z^{\varepsilon,2} \in \mathcal{C}_\mu^\varepsilon$, by the similar deduction to (17) (18) we know that

$$\begin{aligned} \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|\mathcal{R}_1(Z^{\varepsilon,1})(t) - \mathcal{R}_1(Z^{\varepsilon,2})(t)\|_{\mathbb{H}^1} & \leq \frac{\varepsilon L}{\mu - \varepsilon \gamma_1} \left(\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|Z_t^{\varepsilon,1} - Z_t^{\varepsilon,2}\|_{\mathbb{H}} \right), \\ \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|\mathcal{R}_2(Z^{\varepsilon,1})(t) - \mathcal{R}_2(Z^{\varepsilon,2})(t)\|_{\mathbb{H}^2} & \leq \frac{L}{\gamma_2 - \mu} \left(\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|Z_t^{\varepsilon,1} - Z_t^{\varepsilon,2}\|_{\mathbb{H}} \right). \end{aligned}$$

Thus, one can have that

$$\begin{aligned} \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|\mathcal{R}(Z^{\varepsilon,1})(t) - \mathcal{R}(Z^{\varepsilon,2})(t)\|_{\mathbb{H}} & \leq \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|\mathcal{R}_1(Z^{\varepsilon,1})(t) - \mathcal{R}_1(Z^{\varepsilon,2})(t)\|_{\mathbb{H}^1} \\ & \quad + \sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|\mathcal{R}_2(Z^{\varepsilon,1})(t) - \mathcal{R}_2(Z^{\varepsilon,2})(t)\|_{\mathbb{H}^2} \\ & \leq \left(\frac{\varepsilon L}{\mu - \varepsilon \gamma_1} + \frac{L}{\gamma_2 - \mu} \right) \left(\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|Z_t^{\varepsilon,1} - Z_t^{\varepsilon,2}\|_{\mathbb{H}} \right). \end{aligned}$$

So, for $0 < \varepsilon \leq \varepsilon_0$, $\mathcal{R} : \mathcal{C}_\mu^\varepsilon \mapsto \mathcal{C}_\mu^\varepsilon$ is contractive. That is, Eq.(15) has a unique solution denoted as $Z^\varepsilon = (X^\varepsilon, Y^\varepsilon)$. Moreover,

$$\sup_{t \in \mathbb{R}} e^{\frac{\mu}{\varepsilon} t} \|Z_t^\varepsilon\|_{\mathbb{H}} \leq \frac{1}{1 - M} \left((2 + 2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)$$

and then

$$\|Z_t^\varepsilon\|_{\mathbb{H}} \leq \frac{e^{-\frac{\mu}{\varepsilon} t}}{1 - M} \left((2 + 2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right), \quad t \geq 0. \quad (19)$$

Set

$$\tilde{z}_t^\varepsilon := \begin{pmatrix} \tilde{x}_t^\varepsilon \\ \tilde{y}_t^\varepsilon \end{pmatrix} := \begin{pmatrix} x_t^\varepsilon \\ y_t^\varepsilon \end{pmatrix} + \begin{pmatrix} X_t^\varepsilon \\ Y_t^\varepsilon \end{pmatrix},$$

and then by simple calculation, it holds that $\tilde{z}_t^\varepsilon = (\tilde{x}_t^\varepsilon, \tilde{y}_t^\varepsilon)$ solves uniquely the following equation

$$\tilde{z}_t^\varepsilon = \begin{cases} \mathcal{K}(\tilde{z}^\varepsilon)(t), & t \leq 0, \\ \varphi^\varepsilon(t, \omega)(\tilde{x}_0^\varepsilon, \tilde{y}_0^\varepsilon), & t > 0. \end{cases}$$

Thus, by Lemma 4.1, we know that \tilde{z}_t^ε solves Eq.(7) for $t \leq 0$. In particular, $\tilde{y}_0^\varepsilon = H^{\varepsilon,0}(\omega, \tilde{x}_0^\varepsilon)$, which yields that $(\tilde{x}_0^\varepsilon, \tilde{y}_0^\varepsilon) \in \mathcal{M}^\varepsilon(\omega)$. So, we take $\tilde{z}_0 = (\tilde{x}_0^\varepsilon, \tilde{y}_0^\varepsilon)$. Since $\tilde{z}_t^\varepsilon =$

$\varphi^\varepsilon(t, \omega)(\tilde{x}_0^\varepsilon, \tilde{y}_0^\varepsilon)$ for $t > 0$, $\tilde{z}_t^\varepsilon = \varphi^\varepsilon(t, \omega)\tilde{z}_0$ for $t > 0$. Note that $\tilde{x}_t^\varepsilon - x_t^\varepsilon = X_t^\varepsilon$, $\tilde{y}_t^\varepsilon - y_t^\varepsilon = Y_t^\varepsilon$. Thus, by (19), it holds that

$$\begin{aligned} \|\varphi^\varepsilon(t, \omega)z_0 - \varphi^\varepsilon(t, \omega)\tilde{z}_0\|_{\mathbb{H}} &= \|(x_t^\varepsilon, y_t^\varepsilon) - (\tilde{x}_t^\varepsilon, \tilde{y}_t^\varepsilon)\|_{\mathbb{H}} = \|Z_t^\varepsilon\|_{\mathbb{H}} \\ &\leq \frac{e^{-\frac{\mu}{\varepsilon}t}}{1-M} \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right), \quad t \geq 0. \end{aligned}$$

The proof is completed. \square

4.2. A reduced system on the random invariant manifold \mathcal{M}^ε . In the subsection, we prove that there exists a reduced system on the random invariant manifold \mathcal{M}^ε such that it will approximate the original system (1) for sufficiently long time.

By Theorem 4.3, we can obtain the following result.

Theorem 4.4. *(A reduced system on the random invariant manifold)*

Assume that (\mathbf{H}_1) – (\mathbf{H}_4) hold. Let $z_0 = (x_0, y_0) \in \mathbb{H}$. Then for $0 < \varepsilon \leq \varepsilon_0$ and the system (1) with the initial value $z^\varepsilon(0) = z_0$, there exists the following system on the random invariant manifold \mathcal{M}^ε :

$$\begin{cases} \dot{\tilde{x}}^\varepsilon = A\tilde{x}^\varepsilon + F(\tilde{x}^\varepsilon, \tilde{y}^\varepsilon) + \sigma_1 \dot{W}_1, \\ \tilde{y}^\varepsilon = H^{\varepsilon,0}(\theta, \omega, \tilde{x}^\varepsilon), \end{cases} \quad (20)$$

such that for almost all ω ,

$$\|z^\varepsilon(t, \omega) - \tilde{z}^\varepsilon(t, \omega)\|_{\mathbb{H}} \leq \frac{e^{-\frac{\mu}{\varepsilon}t}}{1-M} \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right), \quad t \geq 0, \quad (21)$$

where $\tilde{z}^\varepsilon(t) = (\tilde{x}^\varepsilon(t), \tilde{y}^\varepsilon(t))$ is the solution of the system (20) with the initial value $\tilde{z}^\varepsilon(0) = (\tilde{x}_0^\varepsilon, H^{\varepsilon,0}(\omega, \tilde{x}_0^\varepsilon))$.

Remark 4.5. By (21), we know that when ε is enough small or t is sufficiently large, the system (20) will approximate the system (1).

5. AN APPROXIMATE FILTER ON THE INVARIANT MANIFOLD

In the section we introduce nonlinear filtering problems for the system (1) and the reduced system (20) on the random invariant manifold, and then study their relation.

5.1. Nonlinear filtering problems. In the subsection we introduce nonlinear filtering problems for the system (1) and the reduced system (20).

Let $\{\beta_i(t, \omega)\}_{i \geq 1}$ be a family of mutually independent one-dimensional Brownian motions on $(\Omega, \mathcal{F}, \mathbb{P})$. Construct a cylindrical Brownian motion on \mathbb{H}^3 with respect to $(\Omega, \mathcal{F}, \mathbb{P})$ by

$$W_3(t) := W_3(t, \omega) := \sum_{i=1}^{\infty} \beta_i(t, \omega) e_i, \quad \omega \in \Omega, \quad t \in [0, \infty),$$

where $\{e_i\}_{i \geq 1}$ is a complete orthonormal basis for \mathbb{H}^3 . It is easy to justify that the covariance operator of the cylindrical Brownian motion W_3 is the identity operator I on \mathbb{H}^3 . Note that W_3 is not a process on \mathbb{H}^3 . It is convenient to realize W_3 as a continuous process on an enlarged Hilbert space $\tilde{\mathbb{H}}^3$, the completion of \mathbb{H}^3 under the inner product

$$\langle x, y \rangle_{\tilde{\mathbb{H}}^3} := \sum_{i=1}^{\infty} 2^{-i} \langle x, e_i \rangle_{\mathbb{H}^3} \langle y, e_i \rangle_{\mathbb{H}^3}, \quad x, y \in \mathbb{H}^3.$$

Note that here W_3 may be either independent of W_1 and W_2 , or dependent on W_1 and W_2 (see [4]).

In the following, fix a Borel measurable function $h(x, y) : \mathbb{H}^1 \times \mathbb{H}^2 \rightarrow \mathbb{H}^3$. For h , we make the following hypothesis:

(**H₅**) There exists a $C_h > 0$ such that $\sup_{(x,y) \in \mathbb{H}^1 \times \mathbb{H}^2} \|h(x, y)\|_{\mathbb{H}^3} \leq C_h$ and $h(x, y)$ is Lipschitz continuous in (x, y) whose Lipschitz constant is denoted by $\|h\|_{Lip}$.

Next, for $T > 0$, an observation system is given by

$$r_t^\varepsilon = W_3(t) + \int_0^t h(x_s^\varepsilon, y_s^\varepsilon) ds, \quad t \in [0, T].$$

Under the assumption (**H₅**), r^ε is well defined. Set

$$(\Gamma_t^\varepsilon)^{-1} := \exp \left\{ - \int_0^t \langle h(x_s^\varepsilon, y_s^\varepsilon), dW_3(s) \rangle_{\mathbb{H}^3} - \frac{1}{2} \int_0^t \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\},$$

and then by [6, Proposition 10.17], $(\Gamma_t^\varepsilon)^{-1}$ is an exponential martingale under \mathbb{P} . Thus, [6, Theorem 10.14] admits us to obtain that r^ε is a cylindrical Brownian motion under a new probability measure \mathbb{P}^ε via

$$\frac{d\mathbb{P}^\varepsilon}{d\mathbb{P}} := (\Gamma_T^\varepsilon)^{-1}.$$

Rewrite Γ_t^ε as

$$\Gamma_t^\varepsilon = \exp \left\{ \int_0^t \langle h(x_s^\varepsilon, y_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3} - \frac{1}{2} \int_0^t \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\},$$

and define

$$\rho_t^\varepsilon(\phi) := \mathbb{E}^\varepsilon[\phi(x_t^\varepsilon) \Gamma_t^\varepsilon | \mathcal{R}_t^\varepsilon], \quad \phi \in \mathcal{B}(\mathbb{H}^1),$$

where \mathbb{E}^ε stands for the expectation under \mathbb{P}^ε , $\mathcal{R}_t^\varepsilon \triangleq \sigma(r_s^\varepsilon : 0 \leq s \leq t) \vee \mathcal{N}$ and \mathcal{N} is the collection of all \mathbb{P} -measure zero sets. And set

$$\pi_t^\varepsilon(\phi) := \mathbb{E}[\phi(x_t^\varepsilon) | \mathcal{R}_t^\varepsilon], \quad \phi \in \mathcal{B}(\mathbb{H}^1),$$

and then by the Kallianpur-Striebel formula it holds that

$$\pi_t^\varepsilon(\phi) = \frac{\rho_t^\varepsilon(\phi)}{\rho_t^\varepsilon(1)}.$$

Here ρ_t^ε is called the nonnormalized filtering of x_t^ε with respect to $\mathcal{R}_t^\varepsilon$, and π_t^ε is called the normalized filtering of x_t^ε with respect to $\mathcal{R}_t^\varepsilon$, or the nonlinear filtering problem for x_t^ε with respect to $\mathcal{R}_t^\varepsilon$.

Besides, we rewrite the reduced system (20) as

$$\dot{\tilde{x}}^\varepsilon = A\tilde{x}^\varepsilon + \tilde{F}^\varepsilon(\omega, \tilde{x}^\varepsilon) + \sigma_1 \dot{W}_1,$$

where $\tilde{F}^\varepsilon(\omega, x) := F(x, H^{\varepsilon,0}(\theta.\omega, x))$, and study the nonlinear filtering problem for \tilde{x}^ε . Set

$$\begin{aligned} \tilde{h}^\varepsilon(\omega, x) &:= h(x, H^{\varepsilon,0}(\theta.\omega, x)), \\ \tilde{\Gamma}_t^\varepsilon &:= \exp \left\{ \int_0^t \langle \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3} - \frac{1}{2} \int_0^t \|\tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\}, \end{aligned}$$

and then by [6, Proposition 10.17], $\tilde{\Gamma}_t^\varepsilon$ is an exponential martingale under \mathbb{P}^ε . Thus, we define

$$\begin{aligned}\tilde{\rho}_t^\varepsilon(\phi) &:= \mathbb{E}^\varepsilon[\phi(\tilde{x}_t^\varepsilon)\tilde{\Gamma}_t^\varepsilon|\mathcal{R}_t^\varepsilon], \\ \tilde{\pi}_t^\varepsilon(\phi) &:= \frac{\tilde{\rho}_t^\varepsilon(\phi)}{\tilde{\rho}_t^\varepsilon(1)}, \quad \phi \in \mathcal{B}(\mathbb{H}^1),\end{aligned}$$

and then study the relation between π_t^ε and $\tilde{\pi}_t^\varepsilon$.

5.2. The relation between π_t^ε and $\tilde{\pi}_t^\varepsilon$. In the subsection we investigate the relation of π_t^ε and $\tilde{\pi}_t^\varepsilon$ for ε small enough. Let us start with two estimations.

Lemma 5.1. *Assume that (\mathbf{H}_1) – (\mathbf{H}_5) are satisfied. Then*

$$\mathbb{E}^\varepsilon |\tilde{\rho}_t^\varepsilon(1)|^{-p} \leq \exp \left\{ \left(\frac{p^2}{2} + \frac{p}{2} \right) C_h^2 T \right\}, \quad t \in [0, T], \quad p > 0.$$

Proof. By the Jensen inequality it holds that

$$\mathbb{E}^\varepsilon |\tilde{\rho}_t^\varepsilon(1)|^{-p} = \mathbb{E}^\varepsilon \left[\mathbb{E}^\varepsilon [\tilde{\Gamma}_t^\varepsilon | \mathcal{R}_t^\varepsilon] \right]^{-p} \leq \mathbb{E}^\varepsilon \left[\mathbb{E}^\varepsilon [|\tilde{\Gamma}_t^\varepsilon|^{-p} | \mathcal{R}_t^\varepsilon] \right] = \mathbb{E}^\varepsilon [|\tilde{\Gamma}_t^\varepsilon|^{-p}].$$

So, by the definition of $\tilde{\Gamma}_t^\varepsilon$ we know that

$$\begin{aligned}\mathbb{E}^\varepsilon [|\tilde{\Gamma}_t^\varepsilon|^{-p}] &= \mathbb{E}^\varepsilon \left[\exp \left\{ -p \int_0^t \langle \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3} + \frac{p}{2} \int_0^t \|\tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \right] \\ &= \mathbb{E}^\varepsilon \left[\exp \left\{ -p \int_0^t \langle \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3} - \frac{p^2}{2} \int_0^t \|\tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \right. \\ &\quad \left. \bullet \exp \left\{ \left(\frac{p^2}{2} + \frac{p}{2} \right) \int_0^t \|\tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \right] \\ &\leq \mathbb{E}^\varepsilon \left[\exp \left\{ -p \int_0^t \langle \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3} - \frac{p^2}{2} \int_0^t \|\tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \right] \\ &\quad \bullet \exp \left\{ \left(\frac{p^2}{2} + \frac{p}{2} \right) C_h^2 T \right\} \\ &= \exp \left\{ \left(\frac{p^2}{2} + \frac{p}{2} \right) C_h^2 T \right\},\end{aligned}$$

where the last step is based on the fact that $\exp \left\{ -p \int_0^t \langle \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3} - \frac{p^2}{2} \int_0^t \|\tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\}$ is an exponential martingale under \mathbb{P}^ε . The proof is completed. \square

Lemma 5.2. *Under (\mathbf{H}_1) – (\mathbf{H}_5) , it holds that for $0 < \varepsilon \leq \varepsilon_0$ and $\phi \in \mathcal{C}_b^1(\mathbb{H}^1)$,*

$$\begin{aligned}\mathbb{E}^\varepsilon |\rho_t^\varepsilon(\phi) - \tilde{\rho}_t^\varepsilon(\phi)|^p &\leq \|\phi\|^p \frac{C}{[1-M]^p} \left(\mathbb{E}^\varepsilon \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2} \\ &\quad \times \left(e^{-\frac{\mu}{\varepsilon}tp} + \frac{\varepsilon}{\mu p} \right), \quad t \in [0, T], \quad p > 2,\end{aligned}$$

where $C > 0$ is a constant independent of ε .

Proof. For $\phi \in \mathcal{C}_b^1(\mathbb{H}^1)$ and $p > 2$, by the definition of $\rho_t^\varepsilon(\phi)$, $\tilde{\rho}_t^\varepsilon(\phi)$ it holds that

$$\begin{aligned}
\mathbb{E}^\varepsilon |\rho_t^\varepsilon(\phi) - \tilde{\rho}_t^\varepsilon(\phi)|^p &= \mathbb{E}^\varepsilon \left| \mathbb{E}^\varepsilon [\phi(x_t^\varepsilon) \Gamma_t^\varepsilon | \mathcal{R}_t^\varepsilon] - \mathbb{E}^\varepsilon [\phi(\tilde{x}_t^\varepsilon) \tilde{\Gamma}_t^\varepsilon | \mathcal{R}_t^\varepsilon] \right|^p \\
&= \mathbb{E}^\varepsilon \left| \mathbb{E}^\varepsilon [\phi(x_t^\varepsilon) \Gamma_t^\varepsilon - \phi(\tilde{x}_t^\varepsilon) \tilde{\Gamma}_t^\varepsilon | \mathcal{R}_t^\varepsilon] \right|^p \\
&\leq \mathbb{E}^\varepsilon \left[\mathbb{E}^\varepsilon \left[\left| \phi(x_t^\varepsilon) \Gamma_t^\varepsilon - \phi(\tilde{x}_t^\varepsilon) \tilde{\Gamma}_t^\varepsilon \right|^p \middle| \mathcal{R}_t^\varepsilon \right] \right] \\
&= \mathbb{E}^\varepsilon \left[\left| \phi(x_t^\varepsilon) \Gamma_t^\varepsilon - \phi(\tilde{x}_t^\varepsilon) \tilde{\Gamma}_t^\varepsilon \right|^p \right] \\
&\leq 2^{p-1} \mathbb{E}^\varepsilon \left[\left| \phi(x_t^\varepsilon) \Gamma_t^\varepsilon - \phi(\tilde{x}_t^\varepsilon) \tilde{\Gamma}_t^\varepsilon \right|^p \right] \\
&\quad + 2^{p-1} \mathbb{E}^\varepsilon \left[\left| \phi(\tilde{x}_t^\varepsilon) \Gamma_t^\varepsilon - \phi(\tilde{x}_t^\varepsilon) \tilde{\Gamma}_t^\varepsilon \right|^p \right] \\
&=: J_1 + J_2.
\end{aligned} \tag{22}$$

To J_1 , by the Hölder inequality, we know that

$$\begin{aligned}
J_1 &\leq 2^{p-1} (\mathbb{E}^\varepsilon [|\phi(x_t^\varepsilon) - \phi(\tilde{x}_t^\varepsilon)|^{2p}])^{1/2} (\mathbb{E}^\varepsilon |\Gamma_t^\varepsilon|^{2p})^{1/2} \\
&\leq 2^{p-1} \|\phi\|^p (\mathbb{E}^\varepsilon \|x_t^\varepsilon - \tilde{x}_t^\varepsilon\|_{\mathbb{H}^1}^{2p})^{1/2} \left(\mathbb{E}^\varepsilon \exp \left\{ 2p \int_0^t \langle h(x_s^\varepsilon, y_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3} - \frac{(2p)^2}{2} \int_0^t \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \right. \\
&\quad \left. \bullet \exp \left\{ \frac{(2p)^2}{2} \int_0^t \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds - \frac{2p}{2} \int_0^t \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \right)^{1/2} \\
&\leq 2^{p-1} \|\phi\|^p \frac{e^{-\frac{\mu}{\varepsilon} t p}}{[1-M]^p} \left(\mathbb{E}^\varepsilon \left((2 + 2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2} e^{p(2p-1)C_h^2 T/2},
\end{aligned} \tag{23}$$

where the last step is based on Theorem 4.4 and the fact that the process $\exp \left\{ 2p \int_0^t \langle h(x_s^\varepsilon, y_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3} - \frac{(2p)^2}{2} \int_0^t \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\}$ is an exponential martingale under \mathbb{P}^ε .

Next, for J_2 , it holds that

$$J_2 \leq 2^{p-1} \|\phi\|^p \mathbb{E}^\varepsilon \left[\left| \Gamma_t^\varepsilon - \tilde{\Gamma}_t^\varepsilon \right|^p \right].$$

Based on the Itô formula, Γ_t^ε and $\tilde{\Gamma}_t^\varepsilon$ solve the following equations, respectively,

$$\Gamma_t^\varepsilon = 1 + \int_0^t \Gamma_s^\varepsilon \langle h(x_s^\varepsilon, y_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3}, \quad \tilde{\Gamma}_t^\varepsilon = 1 + \int_0^t \tilde{\Gamma}_s^\varepsilon \langle \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon), dr_s^\varepsilon \rangle_{\mathbb{H}^3}.$$

So, it follows from the BDG inequality and the Hölder inequality that

$$\begin{aligned}
\mathbb{E}^\varepsilon \left[\left| \Gamma_t^\varepsilon - \tilde{\Gamma}_t^\varepsilon \right|^p \right] &= \mathbb{E}^\varepsilon \left[\left| \int_0^t \left\langle \Gamma_s^\varepsilon h(x_s^\varepsilon, y_s^\varepsilon) - \tilde{\Gamma}_s^\varepsilon \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon), dr_s^\varepsilon \right\rangle_{\mathbb{H}^3} \right|^p \right] \\
&\leq \mathbb{E}^\varepsilon \left[\int_0^t \left\| \Gamma_s^\varepsilon h(x_s^\varepsilon, y_s^\varepsilon) - \tilde{\Gamma}_s^\varepsilon \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon) \right\|_{\mathbb{H}^3}^2 ds \right]^{p/2} \\
&\leq T^{p/2-1} \int_0^t \mathbb{E}^\varepsilon \left\| \Gamma_s^\varepsilon h(x_s^\varepsilon, y_s^\varepsilon) - \tilde{\Gamma}_s^\varepsilon \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon) \right\|_{\mathbb{H}^3}^p ds \\
&\leq 2^{p-1} T^{p/2-1} \int_0^t \mathbb{E}^\varepsilon \left\| \Gamma_s^\varepsilon h(x_s^\varepsilon, y_s^\varepsilon) - \Gamma_s^\varepsilon \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon) \right\|_{\mathbb{H}^3}^p ds
\end{aligned}$$

$$\begin{aligned}
& + 2^{p-1} T^{p/2-1} \int_0^t \mathbb{E}^\varepsilon \left\| \Gamma_s^\varepsilon \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon) - \tilde{\Gamma}_s^\varepsilon \tilde{h}^\varepsilon(\omega, \tilde{x}_s^\varepsilon) \right\|_{\mathbb{H}^3}^p ds \\
& =: J_{21} + J_{22}.
\end{aligned}$$

For J_{21} , by the similar deduction to J_1 we have

$$\begin{aligned}
J_{21} & \leq 2^{p-1} T^{p/2-1} \int_0^t \|h\|_{Lip}^p \frac{e^{-\frac{\mu}{\varepsilon} sp}}{[1-M]^p} \left(\mathbb{E}^\varepsilon \left((2+2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2} \\
& \quad \times e^{p(2p-1)C_h^2 T/2} ds \\
& = 2^{p-1} T^{p/2-1} \|h\|_{Lip}^p \frac{1}{[1-M]^p} \left(\mathbb{E}^\varepsilon \left((2+2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2} \\
& \quad \times e^{p(2p-1)C_h^2 T/2} \frac{\varepsilon}{\mu p} [1 - e^{-\frac{\mu}{\varepsilon} tp}].
\end{aligned}$$

And for J_{22} , it follows from the bounded property of h that

$$J_{22} \leq 2^{p-1} T^{p/2-1} C_h^p \int_0^t \mathbb{E}^\varepsilon \left| \Gamma_s^\varepsilon - \tilde{\Gamma}_s^\varepsilon \right|^p ds.$$

So,

$$\begin{aligned}
\mathbb{E}^\varepsilon \left[\left| \Gamma_t^\varepsilon - \tilde{\Gamma}_t^\varepsilon \right|^p \right] & \leq C \cdot \frac{1}{[1-M]^p} \frac{\varepsilon}{\mu p} \left(\mathbb{E}^\varepsilon \left((2+2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2} \\
& \quad + C \int_0^t \mathbb{E}^\varepsilon \left| \Gamma_s^\varepsilon - \tilde{\Gamma}_s^\varepsilon \right|^p ds,
\end{aligned}$$

where the constant $C > 0$ is independent of ε, μ . By the Gronwall inequality it holds that

$$\mathbb{E}^\varepsilon \left[\left| \Gamma_t^\varepsilon - \tilde{\Gamma}_t^\varepsilon \right|^p \right] \leq C \cdot \frac{1}{[1-M]^p} \frac{\varepsilon}{\mu p} \left(\mathbb{E}^\varepsilon \left((2+2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2}.$$

Thus,

$$J_2 \leq 2^{p-1} \|\phi\|^p C \cdot \frac{1}{[1-M]^p} \frac{\varepsilon}{\mu p} \left(\mathbb{E}^\varepsilon \left((2+2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2}. \quad (24)$$

Finally, combining (22) with (23) and (24), we obtain that

$$\begin{aligned}
\mathbb{E}^\varepsilon |\rho_t^\varepsilon(\phi) - \tilde{\rho}_t^\varepsilon(\phi)|^p & \leq 2^{p-1} \|\phi\|^p \frac{e^{-\frac{\mu}{\varepsilon} tp}}{[1-M]^p} \left(\mathbb{E}^\varepsilon \left((2+2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2} \\
& \quad \times e^{p(2p-1)C_h^2 T/2} \\
& \quad + 2^{p-1} \|\phi\|^p C \cdot \frac{1}{[1-M]^p} \frac{\varepsilon}{\mu p} \\
& \quad \times \left(\mathbb{E}^\varepsilon \left((2+2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2} \\
& \leq \|\phi\|^p \frac{C}{[1-M]^p} \left(\mathbb{E}^\varepsilon \left((2+2M) \|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{2p} \right)^{1/2} \\
& \quad \times \left(e^{-\frac{\mu}{\varepsilon} tp} + \frac{\varepsilon}{\mu p} \right).
\end{aligned}$$

This proves the lemma. \square

Now, we are ready to state and prove the main result in the paper. First of all, we give out two concepts used in the proof of Theorem 5.5.

Definition 5.3. For the set $\mathbb{M} \subset \mathcal{C}_b^1(\mathbb{H}^1)$, if the convergence $\lim_{n \rightarrow \infty} \phi(x_n) = \phi(x), \forall \phi \in \mathbb{M}$, for some $x, x_n \in \mathbb{H}^1$, implies that $\lim_{n \rightarrow \infty} x_n = x$, it is said that \mathbb{M} strongly separates points in \mathbb{H}^1 .

Definition 5.4. For the set $\mathbb{N} \subset \mathcal{C}_b^1(\mathbb{H}^1)$, if μ_n and μ are probability measures on $\mathcal{B}(\mathbb{H}^1)$, such that $\lim_{n \rightarrow \infty} \int_{\mathbb{H}^1} \phi d\mu_n = \int_{\mathbb{H}^1} \phi d\mu$ for any $\phi \in \mathbb{N}$, then μ_n converges weakly to μ , it is said that \mathbb{N} is convergence determining for the topology of weak convergence of probability measures.

Theorem 5.5. (Approximation by the reduced filter on the invariant manifold)

Under (\mathbf{H}_1) – (\mathbf{H}_5) , there exists a positive constant C such that for $0 < \varepsilon < \varepsilon_0$ and $\phi \in \mathcal{C}_b^1(\mathbb{H}^1)$

$$\begin{aligned} \mathbb{E}|\pi_t^\varepsilon(\phi) - \tilde{\pi}_t^\varepsilon(\phi)|^p &\leq \|\phi\|^p \frac{C}{[1-M]^p} \left(\mathbb{E} \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{16p} \right)^{1/16} \\ &\quad \times \left(e^{-\frac{4\mu}{\varepsilon}tp} + \frac{\varepsilon}{4\mu p} \right)^{1/4}, \quad t \in [0, T], \quad p > 2. \end{aligned}$$

Thus, for the distance $d(\cdot, \cdot)$ in the space of probability measures that induces the weak convergence, the following approximation holds:

$$\begin{aligned} \mathbb{E}[d(\pi_t^\varepsilon, \tilde{\pi}_t^\varepsilon)] &\leq \frac{C^{1/p}}{[1-M]} \left(\mathbb{E} \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{16p} \right)^{1/16p} \\ &\quad \times \left(e^{-\frac{4\mu}{\varepsilon}tp} + \frac{\varepsilon}{4\mu p} \right)^{1/4p}. \end{aligned}$$

Proof. For $\phi \in \mathcal{C}_b^1(\mathbb{H}^1)$, the Hölder inequality, Lemma 5.1 and Lemma 5.2 admit us to obtain that

$$\begin{aligned} \mathbb{E}|\pi_t^\varepsilon(\phi) - \tilde{\pi}_t^\varepsilon(\phi)|^p &= \mathbb{E} \left| \frac{\rho_t^\varepsilon(\phi) - \tilde{\rho}_t^\varepsilon(\phi)}{\tilde{\rho}_t^\varepsilon(1)} - \pi_t^\varepsilon(\phi) \frac{\rho_t^\varepsilon(1) - \tilde{\rho}_t^\varepsilon(1)}{\tilde{\rho}_t^\varepsilon(1)} \right|^p \\ &\leq 2^{p-1} \mathbb{E} \left| \frac{\rho_t^\varepsilon(\phi) - \tilde{\rho}_t^\varepsilon(\phi)}{\tilde{\rho}_t^\varepsilon(1)} \right|^p + 2^{p-1} \mathbb{E} \left| \pi_t^\varepsilon(\phi) \frac{\rho_t^\varepsilon(1) - \tilde{\rho}_t^\varepsilon(1)}{\tilde{\rho}_t^\varepsilon(1)} \right|^p \\ &\leq 2^{p-1} (\mathbb{E} |\rho_t^\varepsilon(\phi) - \tilde{\rho}_t^\varepsilon(\phi)|^{2p})^{1/2} (\mathbb{E} |\tilde{\rho}_t^\varepsilon(1)|^{-2p})^{1/2} \\ &\quad + 2^{p-1} \|\phi\|^p (\mathbb{E} |\rho_t^\varepsilon(1) - \tilde{\rho}_t^\varepsilon(1)|^{2p})^{1/2} (\mathbb{E} |\tilde{\rho}_t^\varepsilon(1)|^{-2p})^{1/2} \\ &= 2^{p-1} (\mathbb{E}^\varepsilon |\rho_t^\varepsilon(\phi) - \tilde{\rho}_t^\varepsilon(\phi)|^{2p} \Gamma_T^\varepsilon)^{1/2} (\mathbb{E}^\varepsilon |\tilde{\rho}_t^\varepsilon(1)|^{-2p} \Gamma_T^\varepsilon)^{1/2} \\ &\quad + 2^{p-1} \|\phi\|^p (\mathbb{E}^\varepsilon |\rho_t^\varepsilon(1) - \tilde{\rho}_t^\varepsilon(1)|^{2p} \Gamma_T^\varepsilon)^{1/2} (\mathbb{E}^\varepsilon |\tilde{\rho}_t^\varepsilon(1)|^{-2p} \Gamma_T^\varepsilon)^{1/2} \\ &\leq 2^{p-1} (\mathbb{E}^\varepsilon |\rho_t^\varepsilon(\phi) - \tilde{\rho}_t^\varepsilon(\phi)|^{4p})^{1/4} (\mathbb{E}^\varepsilon |\tilde{\rho}_t^\varepsilon(1)|^{-4p})^{1/4} (\mathbb{E}^\varepsilon |\Gamma_T^\varepsilon|^2)^{1/2} \\ &\quad + 2^{p-1} \|\phi\|^p (\mathbb{E}^\varepsilon |\rho_t^\varepsilon(1) - \tilde{\rho}_t^\varepsilon(1)|^{4p})^{1/4} (\mathbb{E}^\varepsilon |\tilde{\rho}_t^\varepsilon(1)|^{-4p})^{1/4} (\mathbb{E}^\varepsilon |\Gamma_T^\varepsilon|^2)^{1/2} \\ &\leq \|\phi\|^p \frac{C}{[1-M]^p} \left(\mathbb{E}^\varepsilon \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{8p} \right)^{1/8} \\ &\quad \left(e^{-\frac{4\mu}{\varepsilon}tp} + \frac{\varepsilon}{4\mu p} \right)^{1/4} (\mathbb{E}^\varepsilon |\Gamma_T^\varepsilon|^2)^{1/2} \\ &= \|\phi\|^p \frac{C}{[1-M]^p} \left(\mathbb{E} \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{8p} |\Gamma_T^\varepsilon|^{-1} \right)^{1/8} \end{aligned}$$

$$\begin{aligned}
& \left(e^{-\frac{4\mu}{\varepsilon}tp} + \frac{\varepsilon}{4\mu p} \right)^{1/4} (\mathbb{E}^\varepsilon |\Gamma_T^\varepsilon|^2)^{1/2} \\
& \leq \|\phi\|^p \frac{C}{[1-M]^p} \left(\mathbb{E} \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{16p} \right)^{1/16} \\
& \quad \cdot (\mathbb{E} |\Gamma_T^\varepsilon|^{-2})^{1/16} \left(e^{-\frac{4\mu}{\varepsilon}tp} + \frac{\varepsilon}{4\mu p} \right)^{1/4} (\mathbb{E}^\varepsilon |\Gamma_T^\varepsilon|^2)^{1/2}.
\end{aligned}$$

In the following, we estimate $\mathbb{E}|\Gamma_T^\varepsilon|^{-2}, \mathbb{E}^\varepsilon |\Gamma_T^\varepsilon|^2$. By simple calculations, it holds that

$$\begin{aligned}
\mathbb{E}(\Gamma_T^\varepsilon)^{-2} &= \mathbb{E} \left(\exp \left\{ -2 \int_0^T \langle h(x_s^\varepsilon, y_s^\varepsilon), dW_3(s) \rangle_{\mathbb{H}^3} + \frac{2}{2} \int_0^T \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \right) \\
&= \mathbb{E} \left[\left(\exp \left\{ -2 \int_0^T \langle h(x_s^\varepsilon, y_s^\varepsilon), dW_3(s) \rangle_{\mathbb{H}^3} - \frac{2^2}{2} \int_0^T \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \right) \right. \\
&\quad \cdot \exp \left\{ \frac{2^2}{2} \int_0^T \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds + \frac{2}{2} \int_0^T \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\} \Big] \\
&\leq \exp \{ 3C_h^2 T \},
\end{aligned}$$

where the last step is based on the fact that $\exp \left\{ -2 \int_0^t \langle h(x_s^\varepsilon, y_s^\varepsilon), dW_3(s) \rangle_{\mathbb{H}^3} - \frac{2^2}{2} \int_0^t \|h(x_s^\varepsilon, y_s^\varepsilon)\|_{\mathbb{H}^3}^2 ds \right\}$ is an exponential martingale under \mathbb{P} . By the similar deduction to above, we know that

$$\mathbb{E}^\varepsilon |\Gamma_T^\varepsilon|^2 \leq \exp \{ C_h^2 T \}.$$

Thus,

$$\begin{aligned}
\mathbb{E}|\pi_t^\varepsilon(\phi) - \tilde{\pi}_t^\varepsilon(\phi)|^p &\leq \|\phi\|^p \frac{C}{[1-M]^p} \left(\mathbb{E} \left((2+2M)\|z_0\|_{\mathbb{H}} + 2R(\omega) \right)^{16p} \right)^{1/16} \\
&\quad \cdot \left(e^{-\frac{4\mu}{\varepsilon}tp} + \frac{\varepsilon}{4\mu p} \right)^{1/4}.
\end{aligned}$$

Next, notice that there exists a countable algebra $\{\phi_i, i = 1, 2, \dots\}$ of $\mathcal{C}_b^1(\mathbb{H}^1)$ that strongly separates points in \mathbb{H}^1 . By [8, Theorem 3.4.5], it furthermore holds that $\{\phi_i, i = 1, 2, \dots\}$ is convergence determining for the topology of weak convergence of probability measures. For two probability measures μ, τ on $\mathcal{B}(\mathbb{H}^1)$, set

$$d(\mu, \tau) := \sum_{i=1}^{\infty} \frac{|\int_{\mathbb{H}^1} \phi_i d\mu - \int_{\mathbb{H}^1} \phi_i d\tau|}{2^i},$$

and then d is a distance in the space of probability measures on $\mathcal{B}(\mathbb{H}^1)$. Since $\{\phi_i, i = 1, 2, \dots\}$ is convergence determining for the topology of weak convergence of probability measures, d induces the weak convergence. The proof is completed. \square

Remark 5.6. By the above theorem, we know that when ε goes to zero, $\tilde{\pi}_t^\varepsilon$ approximates π_t^ε . Therefore, $\tilde{\pi}_t^\varepsilon$ could be understood as the nonlinear filtering problem for \tilde{x}_t^ε with respect to $\mathcal{R}_t^\varepsilon$.

6. AN EXAMPLE

Example 6.1. Let D be a domain in \mathbb{R}^3 with smooth boundary ∂D . Consider the following coupled hyperbolic and parabolic equation

$$v_{tt} + \gamma v_t - \Delta v = f(v, v_t, \theta) + \sigma_1 \dot{W}_1, t > 0, x \in D,$$

$$\begin{aligned}\theta_t - \frac{1}{\varepsilon}\kappa\Delta\theta &= \frac{1}{\varepsilon}g(v, v_t, \theta) + \frac{\sigma_2}{\sqrt{\varepsilon}}\dot{W}_2, \quad t > 0, x \in D, \\ v &= 0, \quad \theta = 0, \quad t > 0, \quad x \in \partial D,\end{aligned}\tag{25}$$

where $\gamma \geq 0, \kappa > 0$ are constants, Δ is the Laplace operator and $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ and $g : \mathbb{R}^3 \rightarrow \mathbb{R}$ are Lipschitz continuous with a Lipschitz constant $L > 0$.

The type of equations are usually used to describe a thermoelastic phenomenon in a random medium (c.f.[3]). Here v denotes the displacement and θ is the temperature. And the parameter γ describes resistance forces, and the white noise processes \mathcal{W}_1 and \mathcal{W}_2 model random fluctuations in external loads (\dot{W}_1) and in thermal sources (\dot{W}_2). If the temperature evolves fastly, then the hyperbolic equation is coupled to a parabolic equation with different characteristic timescales.

Next, we rewrite Eq.(25) as

$$\begin{cases} \dot{x}^\varepsilon = Ax^\varepsilon + F(x^\varepsilon, y^\varepsilon) + \sigma_1 \dot{W}_1, \\ \dot{y}^\varepsilon = \frac{1}{\varepsilon}By^\varepsilon + \frac{1}{\varepsilon}G(x^\varepsilon, y^\varepsilon) + \frac{\sigma_2}{\sqrt{\varepsilon}}\dot{W}_2, \end{cases}$$

where

$$\begin{aligned}x^\varepsilon &= \begin{pmatrix} v \\ v_t \end{pmatrix}, A = \begin{pmatrix} 0 & 1 \\ \Delta & -\gamma \end{pmatrix}, F(x^\varepsilon, y^\varepsilon) = \begin{pmatrix} 0 \\ f(v, v_t, \theta) \end{pmatrix}, W_1 = \begin{pmatrix} 0 \\ \mathcal{W}_1 \end{pmatrix}, \\ y^\varepsilon &= \theta, B = \kappa\Delta, G(x^\varepsilon, y^\varepsilon) = g(v, v_t, \theta), W_2 = \mathcal{W}_2,\end{aligned}$$

We take $\mathbb{H}^1 = H_0^1(D) \times L_2(D)$ and $\mathbb{H}^2 = L_2(D)$, where $L_2(D)$ and $H_0^1(D)$ are the usual Sobolev spaces. Thus, Eq.(25) is in our framework. Moreover, by simple calculation, we know that **(H₁)**–**(H₃)** are satisfied with $\gamma_1 = \gamma, \gamma_2 = \kappa$. If $\kappa > L$, it follows from Theorem 3.3 that for $z_0 = (x_0, y_0) \in \mathbb{H} := \mathbb{H}^1 \times \mathbb{H}^2$, Eq.(25) has a unique mild solution $z^\varepsilon(t, 0, \omega; z_0)$ for $t \in [0, T]$ and $\omega \in \Omega$. And Theorem 4.4 admits us to obtain that for $0 < \varepsilon \leq \varepsilon_0$, there exists the following reduced system on the random invariant manifold \mathcal{M}^ε :

$$\begin{cases} \dot{\tilde{x}}^\varepsilon = A\tilde{x}^\varepsilon + F(\tilde{x}^\varepsilon, \tilde{y}^\varepsilon) + \sigma_1 \dot{W}_1, \\ \tilde{y}^\varepsilon = H^\varepsilon(\theta, \omega, \tilde{x}^\varepsilon). \end{cases}$$

In the following, we consider the nonlinear filtering problem of Eq.(25). Taking $\mathbb{H}^3 = L_2(D)$, one can construct a cylindrical Brownian motion W_3 on \mathbb{H}^3 . And then we give an observation system by

$$r_t^\varepsilon = W_3(t) + \int_0^t \sin(x_s^\varepsilon) ds, \quad t \in [0, T].$$

Thus, it is easy to justify that $h(x, y) = \sin x$ satisfies **(H₅)**. By Theorem 5.5, the distance of the nonlinear filtering for x^ε and “the nonlinear filtering” for \tilde{x}^ε is characterized.

7. CONCLUSIONS

In the paper, we consider an effective filtering for a type of slow-fast data assimilation systems in general Hilbert spaces. It is proved that the origin system can be reduced to a system on a random invariant manifold constructed by the direct way. And then, the nonlinear filtering of the origin system can be approximated by that of the reduction system.

Of course, one could consider other different expressions for the noisy terms in the equations (multiplicative, cylindrical, etc). We will investigate the possibility of doing a similar reduction and the related nonlinear filtering in the future.

8. APPENDIX

In this Appendix, we present the proof of Theorem 3.3.

Proof of Theorem 3.3. First of all, we prove that the system (1) has a unique mild solution. Define an operator $\mathcal{J} : \mathcal{C}([s, s+T], \mathbb{H}) \rightarrow \mathcal{C}([s, s+T], \mathbb{H})$ by

$$\mathcal{J}(z^\varepsilon)(t) := \begin{pmatrix} \mathcal{J}_1(z^\varepsilon)(t) \\ \mathcal{J}_2(z^\varepsilon)(t) \end{pmatrix} := \begin{pmatrix} e^{A(t-s)}x_0 + \int_s^t e^{A(t-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr + \int_s^t e^{A(t-r)}\sigma_1 dW_1(r) \\ e^{\frac{B}{\varepsilon}(t-s)}y_0 + \int_s^t e^{\frac{B}{\varepsilon}(t-r)}\frac{1}{\varepsilon}G(x_r^\varepsilon, y_r^\varepsilon)dr + \int_s^t e^{\frac{B}{\varepsilon}(t-r)}\frac{\sigma_2}{\sqrt{\varepsilon}}dW_2(r) \end{pmatrix}.$$

And then \mathcal{J} is well defined. In fact, based on (\mathbf{H}_1) and [6, Theorem 5.2], $e^{A(t-s)}x_0$ and $\int_s^t e^{A(t-r)}\sigma_1 dW_1(r)$ are strongly continuous. Taking $t_1, t_2 \in [s, s+T], t_1 < t_2$, we obtain that

$$\begin{aligned} & \left\| \int_s^{t_1} e^{A(t_1-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr - \int_s^{t_2} e^{A(t_2-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr \right\|_{\mathbb{H}^1} \\ & \leq \left\| \int_s^{t_1} e^{A(t_1-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr - \int_s^{t_1} e^{A(t_2-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr \right\|_{\mathbb{H}^1} \\ & \quad + \left\| \int_s^{t_1} e^{A(t_2-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr - \int_s^{t_2} e^{A(t_2-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr \right\|_{\mathbb{H}^1} \\ & \leq \int_s^{t_1} \|F(x_r^\varepsilon, y_r^\varepsilon) - e^{A(t_2-t_1)}F(x_r^\varepsilon, y_r^\varepsilon)\|_{\mathbb{H}^1} dr + \int_{t_1}^{t_2} \|e^{A(t_2-r)}\| \|F(x_r^\varepsilon, y_r^\varepsilon)\|_{\mathbb{H}^1} dr \\ & \leq \int_s^{t_1} \|F(x_r^\varepsilon, y_r^\varepsilon) - e^{A(t_2-t_1)}F(x_r^\varepsilon, y_r^\varepsilon)\|_{\mathbb{H}^1} dr + L \left(\sup_{t \in [s, s+T]} \|z_t^\varepsilon\|_{\mathbb{H}} \right) (t_2 - t_1), \end{aligned}$$

where the last step is based on (6) and (4). The dominated convergence theorem admits us to get that $\int_s^t e^{A(t-r)}F(x_r^\varepsilon, y_r^\varepsilon)dr$ is strongly continuous. Thus, $\mathcal{J}_1(z^\varepsilon)(t)$ is strongly continuous. By the same deduction to above, we know that $\mathcal{J}_2(z^\varepsilon)(t)$ is also strongly continuous. So, $\mathcal{J}(z^\varepsilon)(t)$ is strongly continuous.

Next, we prove that \mathcal{J} is contractive. For $z^{\varepsilon,1}, z^{\varepsilon,2} \in \mathcal{C}([s, s+T], \mathbb{H})$, one can compute by (\mathbf{H}_1) - (\mathbf{H}_3)

$$\begin{aligned} \sup_{t \in [s, s+T]} \|\mathcal{J}_1(z^{\varepsilon,1})(t) - \mathcal{J}_1(z^{\varepsilon,2})(t)\|_{\mathbb{H}^1} &= \sup_{t \in [s, s+T]} \left\| \int_s^t e^{A(t-r)} \left(F(x_r^{\varepsilon,1}, y_r^{\varepsilon,1}) - F(x_r^{\varepsilon,2}, y_r^{\varepsilon,2}) \right) dr \right\|_{\mathbb{H}^1} \\ &\leq \sup_{t \in [s, s+T]} \int_s^t \|e^{A(t-r)}\| \|F(x_r^{\varepsilon,1}, y_r^{\varepsilon,1}) - F(x_r^{\varepsilon,2}, y_r^{\varepsilon,2})\|_{\mathbb{H}^1} dr \\ &\leq \sup_{t \in [s, s+T]} \int_s^t L(\|x_r^{\varepsilon,1} - x_r^{\varepsilon,2}\|_{\mathbb{H}^1} + \|y_r^{\varepsilon,1} - y_r^{\varepsilon,2}\|_{\mathbb{H}^2}) dr \\ &\leq LT \sup_{t \in [s, s+T]} \|z_t^{\varepsilon,1} - z_t^{\varepsilon,2}\|_{\mathbb{H}}, \end{aligned}$$

and

$$\sup_{t \in [s, s+T]} \|\mathcal{J}_2(z^{\varepsilon,1})(t) - \mathcal{J}_2(z^{\varepsilon,2})(t)\|_{\mathbb{H}^1} = \sup_{t \in [s, s+T]} \left\| \int_s^t e^{\frac{B}{\varepsilon}(t-r)} \frac{1}{\varepsilon} \left(G(x_r^{\varepsilon,1}, y_r^{\varepsilon,1}) - G(x_r^{\varepsilon,2}, y_r^{\varepsilon,2}) \right) dr \right\|_{\mathbb{H}^2}$$

$$\begin{aligned}
&\leq \frac{1}{\varepsilon} \sup_{t \in [s, s+T]} \int_s^t \|e^{\frac{B}{\varepsilon}(t-r)}\| \|G(x_r^{\varepsilon,1}, y_r^{\varepsilon,1}) - G(x_r^{\varepsilon,2}, y_r^{\varepsilon,2})\|_{\mathbb{H}^2} dr \\
&\leq \frac{1}{\varepsilon} \sup_{t \in [s, s+T]} \int_s^t e^{-\frac{\gamma_2}{\varepsilon}(t-r)} L(\|x_r^{\varepsilon,1} - x_r^{\varepsilon,2}\|_{\mathbb{H}^1} + \|y_r^{\varepsilon,1} - y_r^{\varepsilon,2}\|_{\mathbb{H}^2}) dr \\
&\leq \frac{L}{\gamma_2} [1 - e^{-\frac{\gamma_2}{\varepsilon}T}] \sup_{t \in [s, s+T]} \|z_t^{\varepsilon,1} - z_t^{\varepsilon,2}\|_{\mathbb{H}}.
\end{aligned}$$

Thus,

$$\begin{aligned}
\sup_{t \in [s, s+T]} \|\mathcal{J}(z^{\varepsilon,1})(t) - \mathcal{J}(z^{\varepsilon,2})(t)\|_{\mathbb{H}} &\leq \sup_{t \in [s, s+T]} \|\mathcal{J}_1(z^{\varepsilon,1})(t) - \mathcal{J}_1(z^{\varepsilon,2})(t)\|_{\mathbb{H}^1} \\
&\quad + \sup_{t \in [s, s+T]} \|\mathcal{J}_2(z^{\varepsilon,1})(t) - \mathcal{J}_2(z^{\varepsilon,2})(t)\|_{\mathbb{H}^2} \\
&\leq \left(LT + \frac{L}{\gamma_2} \right) \sup_{t \in [s, s+T]} \|z_t^{\varepsilon,1} - z_t^{\varepsilon,2}\|_{\mathbb{H}}.
\end{aligned}$$

Taking T_0 such that $LT_0 + \frac{L}{\gamma_2} < 1$, we know that \mathcal{J} is contractive. So, the system (1) has a unique mild solution $z^\varepsilon(t, s, \omega; z_0)$ for $t \in [s, s+T_0]$. If $T \leq T_0$, the proof is over; if $T > T_0$, one can easily extend the solution to the finite interval $[s, s+T]$ by considering $[s, s+T_0]$, $[s+T_0, s+2T_0]$, $[s+2T_0, s+3T_0]$ and so on.

Set

$$\varphi^\varepsilon(t, \omega) z_0 := z^\varepsilon(t, 0, \omega; z_0), \quad t \in \mathbb{R},$$

and then it is easy to justify that for $s, t \in \mathbb{R}$,

$$\begin{aligned}
\varphi^\varepsilon(t+s, \omega) z_0 &= z^\varepsilon(t+s, 0, \omega; z_0) = z^\varepsilon(t+s, s, \omega; z^\varepsilon(s, 0, \omega; z_0)), \\
\varphi^\varepsilon(t, \theta_s \omega) \varphi^\varepsilon(s, \omega) z_0 &= z^\varepsilon(t, 0, \theta_s \omega; z^\varepsilon(s, 0, \omega; z_0)).
\end{aligned}$$

Note that $W_1(r, \theta_s^1 \omega) = W_1(s+r, \omega) - W_1(s, \omega)$, $W_2(r, \theta_s^2 \omega) = W_2(s+r, \omega) - W_2(s, \omega)$ for $r \in \mathbb{R}, s \in \mathbb{R}, \omega \in \Omega$ and $W_1(r, \theta_s^1 \cdot), W_2(r, \theta_s^2 \cdot)$ are still two-sided $\mathbb{H}^1, \mathbb{H}^2$ -valued Brownian motions with covariance operators K_1, K_2 , respectively. Thus, by uniqueness of the mild solution for the system (1), we know that $z^\varepsilon(t+s, s, \omega; z^\varepsilon(s, 0, \omega; z_0)) = z^\varepsilon(t, 0, \theta_s \omega; z^\varepsilon(s, 0, \omega; z_0))$ and $\varphi^\varepsilon(t+s, \omega) z_0 = \varphi^\varepsilon(t, \theta_s \omega) \varphi^\varepsilon(s, \omega) z_0$. That is, $\varphi^\varepsilon(t, \omega)$ is a RDS. The proof is completed.

Acknowledgements:

The author would like to thank Professor Xicheng Zhang for his valuable discussions.

REFERENCES

- [1] L. Arnold, *Random Dynamical Systems*, Springer, Berlin, 1998.
- [2] A. Bain and D. Crisan, *Fundamentals of Stochastic Filtering*, Springer, Berlin, 2009.
- [3] T. Caraballo, I. Chueshov, and J.A. Langa, Existence of invariant manifolds for coupled parabolic and hyperbolic stochastic partial differential equations, *Nonlin.*, 18(2005)747-767.
- [4] T. Cass, M. Clarke and D. Crisan, The filtering equations revisited, In: Crisan D., Hambly B., Zariphopoulou T. (eds) *Stochastic Analysis and Applications (2014)*. Springer Proceedings in Mathematics & Statistics, vol 100.
- [5] I. Chueshov and M. Scheutzow, Inertial manifolds and forms for stochastically perturbed retarded semilinear parabolic equations, *J. Dyn. Diff. Eqns*, 13(2001)355-80.
- [6] G. Da Prato and J. Zabczyk: *Stochastic Equations in Infinite Dimensions*. Encyclopedia of Mathematics and its Applications. Cambridge: Cambridge University Press, 1992.

- [7] J. Duan, K. Lu and B. Schmalfuss, Invariant manifolds for stochastic partial differential equations, *Ann. Probab.*, 31(2003), 2109-2135.
- [8] S. N. Ethier and T. G. Kurtz, *Markov Processes: Characterization and Convergence*. John Wiley & Sons, 1986.
- [9] H. Fu, X. Liu and J. Duan, Slow manifolds for multi-time-scale stochastic evolutionary systems, *Comm. Math. Sci.*, 11(2013), 141-162.
- [10] S. Hobbs and S.S. Sritharan, Nonlinear filtering theory for stochastic reaction-diffusion equations. In *Probability and Modern Analysis*. Gritsky, N., Goldstein, J., and Uhl, J.J. eds., Marcel Dekker, New York, 1996, pp. 219-234.
- [11] P. Imkeller, N. S. Namachchivaya, N. Perkowski and H. C. Yeong, Dimensional reduction in nonlinear filtering: a homogenization approach, *The Annals of Applied Probability*, 23(2013), 2290-2326.
- [12] R. Z. Khasminskii and G. Yin, On transition densities of singularly perturbed diffusions with fast and slow components, *SIAM J. Appl. Math.*, 56(1996), 1794-1819.
- [13] A. Papanicolaou and K. Spiliopoulos: Dimension reduction in statistical estimation of partially observed multiscale processes, *SIAM J. on Uncertainty Quantification*, 5(2017)1220-1247.
- [14] J. H. Park, N. S. Namachchivaya and H. C. Yeong, Particle filters in a multiscale environment: Homogenized hybrid particle filter, *J. Appl. Mech.*, 78(2011), 1-10.
- [15] J. H. Park, R. B. Sowers and N. S. Namachchivaya, Dimensional reduction in nonlinear filtering, *Nonlinearity*, 23(2010), 305-324.
- [16] J. H. Park, B. Rozovskii and R. B. Sowers, Efficient nonlinear filtering of a singularly perturbed stochastic hybrid system, *LMS Journal of Computation and Mathematics*, 14(2011), 254-270.
- [17] G. A. Pavliotis and A. M. Stuart, *Multiscale methods: averaging and homogenization*, Springer Science+Business Media, New York, 2008.
- [18] H. Qiao, Convergence of nonlinear filtering for stochastic dynamical systems with Lévy noises, <https://arxiv.org/abs/1707.07824>.
- [19] H. Qiao, Y. Zhang and J. Duan, Effective filtering on a random slow manifold, *Nonlinearity*, 31(2018) 4649-4666.
- [20] B. L. Rozovskii, *Stochastic Evolution System: Linear Theory and Application to nonlinear Filtering*, Springer, New York, 1990.
- [21] Z. Schuss, *Nonlinear Filtering and Optimal Phase Tracking*, Springer, New York, 2012.
- [22] B. Schmalfuß and R. Schneider, Invariant manifolds for random dynamical systems with slow and fast variables, *J. Dyna. Diff. Equa.*, 20(2008), 133-164.
- [23] S. S. Sritharan, Nonlinear filtering of stochastic Navier-Stokes equations. In *Nonlinear Stochastic PDEs: Burgers Turbulence and Hydrodynamic Limit*. Funaki, T., and Woyczynski, W.A. eds., Springer-Verlag, New York, 1995, pp. 247-260.
- [24] Y. Ying and F. Zhang, Potentials in Improving Predictability of Multiscale Tropical Weather Systems Evaluated through Ensemble Assimilation of Simulated Satellite-Based Observations, *Journal of the Atmospheric Sciences*, 2018, <https://doi.org/10.1175/JAS-D-17-0245.1>.
- [25] Y. Zhang, H. Qiao and J. Duan, Effective filtering analysis for non-Gaussian dynamic systems, appear in *Applied Mathematics and Optimization*.