


On coprime percolation, the visibility graphon, and the local limit of the GCD profile

Sébastien MARTINEAU*

February 5, 2022

Abstract

Colour an element of $\mathbb{Z}^d$ white if its coordinates are coprime and black otherwise. What does this colouring look like when seen from a “uniformly chosen” point of $\mathbb{Z}^d$? More generally, label every element of $\mathbb{Z}^d$ by its GCD: what do the labels look like around a “uniform” point of $\mathbb{Z}^d$? We answer these questions and generalisations of them, provide results of graphon convergence, as well as a “local/graphon” convergence. One can also investigate the percolative properties of the colouring under study.



*Université Paris-Sud, sebastien.martineau@u-psud.fr

Colour an element of $\mathbb{Z}^d$ white if its coordinates are coprime and black otherwise. What does this colouring look like? For $d = 2$, this question was investigated in [Var99]. The starting point of the present paper is the following question: what does this colouring look like *when seen from a “uniformly chosen” point of $\mathbb{Z}^d$* ? An answer has already been formulated in [PH13] but, from the perspective adopted in the current paper, our vocabulary, techniques and results are more satisfactory. See the figure on page 1.

A more general version of this question goes as follows: if one labels every element of $\mathbb{Z}^d$ by its GCD, what do the labels look like around a “uniform” point of $\mathbb{Z}^d$? We answer this question and generalisations of it, provide results of graphon convergence, as well as a “local/graphon” convergence¹. By using previous work of Vardi [Var99], we can also investigate the percolative properties of the colouring under study.

1 Introduction

In this paper, the set $\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$ is taken to contain 0. The set of positive integers will be denoted by $\mathbb{N}^* = \{1, 2, 3, 4, \dots\}$.

Let d denote a positive integer. It is well-known that the probability that d numbers chosen independently and uniformly in $\llbracket 1, N \rrbracket$ are globally coprime converges to $1/\zeta(d)$ when N goes to infinity [Dir51, Ces81, Ces83, Syl83]. Recall that on $[1, \infty)$, the **Euler–Riemann ζ function** is defined by

$$\forall s \in [1, \infty), \zeta(s) := \sum_{n \geq 1} n^{-s} = \prod_{p \in \mathcal{P}} \frac{1}{1 - p^{-s}} \in [1, \infty],$$

where $\mathcal{P} = \{2, 3, 5, 7, 11, \dots\}$ denotes the set of prime numbers. More generally, one has the following result (Theorem 459 in [HW79]).

Let F be a bounded subset of $\mathbb{R}^d$. For every $r \in (0, \infty)$, set $F_r := \{x \in \mathbb{Z}^d : r^{-1}x \in F\}$. Assume that $\frac{|F_r|}{r^d}$ converges to a nonzero limit when r tends to infinity. (A)
Then, one has $\lim_{r \rightarrow \infty} \frac{|\{x \in F_r : \text{GCD}(x_1, \dots, x_d) = 1\}|}{|F_r|} = 1/\zeta(d)$.

The study of **coprime** vectors of $\mathbb{Z}^d$, i.e. of the vectors x that satisfy $\text{GCD}(x_1, \dots, x_d) = 1$, can be performed for its own sake. It may also be motivated by the reducibility of fractions (the probability that a random fraction is irreducible is $1/\zeta(2) = \frac{6}{\pi^2}$) for $d = 2$ or by the visibility problem for arbitrary d . If x and y denote two distinct points of $\mathbb{Z}^d$, one says that x is **visible** from y if the line segment $[x, y]$ intersects $\mathbb{Z}^d$ only at x and y . This

¹Essentially, the same vertex-set will at the same time be endowed with some structure of sparse graph and some structure of dense graph.

condition is equivalent to $x - y$ having a GCD equal to 1. The set of visible points has been studied in various ways: see e.g. [BCZ00, BH15, BMP00, CFF19, Gar15, FM16, GHKM18, HS71, PH13, Var99].

A classical corollary of (A) is the following stronger statement. Recall that the **zeta distribution** of parameter $s > 1$ is the probability distribution on $\mathbb{N}^*$ giving weight $\frac{n^{-s}}{\zeta(s)}$ to each $n \in \mathbb{N}^*$.

Let $d \geq 2$. Let F be a bounded subset of $\mathbb{R}^d$. For every $r \in (0, \infty)$, set $F_r := \{x \in \mathbb{Z}^d : r^{-1}x \in F\}$. Assume that $\frac{|F_r|}{r^d}$ converges to a nonzero limit when r tends to infinity. Let Y_r denote a uniform element in F_r . Then, $\text{GCD}(Y_r)$ converges in distribution to a zeta distribution of parameter d , as r goes to infinity. (B)

In this paper, we are interested in the following two informal questions.

Colour an element of $\mathbb{Z}^d$ white if its coordinates are coprime and black otherwise. What does this colouring look like when seen from a “uniformly chosen” point of $\mathbb{Z}^d$? (Q1)

Label every element of $\mathbb{Z}^d$ by its GCD. What do the labels look like around a “uniform” point of $\mathbb{Z}^d$? (Q2)

Even though (Q2) is stronger than (Q1), it is worthwhile to study both questions independently. Indeed, this leads to two strategies of different nature, and the intermediate proposition related to (Q2) is no stronger than that of (Q1) — see Proposition 2.3, Proposition 2.7, and Fact 2.12.

Let us now make sense of these two questions. Let X be a Polish space. Denote by Ω_X the space $X^{\mathbb{Z}^d}$ endowed with the product topology, which is also Polish. Let c be an element of Ω_X . In (Q1), we will be interested in $X = \{0, 1\}$ and $c = \text{cop} := \mathbb{1}_{\text{coprime}}$. In (Q2), we will be interested in $X = \mathbb{N}$ and $c = \text{gcd} : x \mapsto \text{GCD}(x_1, \dots, x_d)$.

If F is a nonempty finite subset of $\mathbb{Z}^d$, one defines the probability measure $\mu_{F,c}$ as follows — it represents c seen from a uniform point in F . For any $y \in \mathbb{Z}^d$ and $\omega \in \Omega_X$, define $\tau_y \omega \in \Omega_X$ by:

$$\forall x \in \mathbb{Z}^d, (\tau_y \omega)_x = \omega_{x-y}.$$

Let Y be a uniformly chosen element of F . We denote by $\mu_{F,c}$ the distribution of $\tau_{-Y} c$.

We want to describe the limit of $\mu_{F_n,c}$ when c is a map of interest and (F_n) is a reasonable sequence of finite subsets of $\mathbb{Z}^d$, such as $([1, N]^d)$,

($\llbracket -N, N \rrbracket^d$), or $(\{x \in \mathbb{Z}^d : \|x\|_2 \leq n\})$. ⁽²⁾ Limits are taken in the following sense: the space of probability measures on Ω_X is endowed with the weak topology, namely the smallest topology such that for every bounded continuous map $f : \Omega_X \rightarrow \mathbb{R}$, the map $\mu \mapsto \int_{\Omega_X} f(\omega) d\mu(\omega)$ is continuous.

When X is discrete, μ_n converges to μ if and only if for every cylindrical event A , one has $\mu_n[A] \xrightarrow{n \rightarrow \infty} \mu[A]$. Recall that a **cylindrical event** is an event of the form $\{\omega \in \Omega : \omega|_F \in A\}$, where F is a finite subset of $\mathbb{Z}^d$, $\omega|_F$ stands for the restriction of ω to F , and A is some measurable subset of X^F . When X is compact, the space of probability measures on X is compact for the weak topology.

THEOREM 1.1. *Let $d \geq 2$. Let F be a bounded convex subset of $\mathbb{R}^d$ with nonempty interior. For every $r \in (0, \infty)$, set $F_r := \{x \in \mathbb{Z}^d : r^{-1}x \in F\}$.*

Then, $\mu_{F_r, \text{gcd}}$ converges to some explicit probability measure $\mu_{\infty, \text{gcd}}$ when r goes to infinity.

The probability measure $\mu_{\infty, \text{gcd}}$ is defined on page 7. It does not depend on F but only on d . Theorem 1.1 answers (Q2), hence (Q1).

The remaining of the paper is organised as follows. Section 2 investigates (Q1) and (Q2). This section contains the proof of Theorem 1.1, but also complementary results such as Proposition 2.4. The use of relevant arithmetic/probabilistic notions makes the core of the problem very apparent: see  on pages 6 and 10. Section 3 provides several generalisations of these results.

2 Answers to (Q1) and (Q2)

In Section 2.1, we study (Q1). In Section 2.2, we investigate (Q2). Finally, in Section 2.3, we make several comments on Propositions 2.3 and 2.7, and we explore the percolative properties of $\mu_{\infty, \text{cop}}$.

2.1 The random coprime colouring

In this section, we answer (Q1) by proving Theorem 2.1. Even though Theorem 2.1 is a particular case of Theorem 1.1, intermediate propositions of this section cannot be deduced from the content of Section 2.2. See Fact 2.12.

Let us define $\mu_{\infty, \text{cop}}$, which will turn out to be the limit of $\mu_{F_n, \text{cop}}$ when (F_n) is a reasonable sequence of finite subsets of $\mathbb{Z}^d$. For every prime p , let $\mathcal{W}_p$ denote a uniformly chosen coset of $p\mathbb{Z}^d$ in $\mathbb{Z}^d$, i.e. one of the p^d sets of the form $x + p\mathbb{Z}^d$. Do all these choices independently. The distribution of

²This way of proceeding is closely related to local convergence and local weak convergence (also called Benjamini–Schramm convergence): see [Bab91, BS01, DL01].

$\bigcup_p \mathcal{W}_p$ is denoted by $\mu_{\infty, \text{cop}}$. One can see $\mu_{\infty, \text{cop}}$ as a probability measure on $\Omega_{\{0,1\}}$ by identifying a set with its indicator function.

THEOREM 2.1. *Let $d \geq 1$. Let F be a bounded convex subset of $\mathbb{R}^d$ with nonempty interior. For every $r \in (0, \infty)$, set $F_r := \{x \in \mathbb{Z}^d : r^{-1}x \in F\}$.*

Then, $\mu_{F_r, \text{cop}}$ converges to $\mu_{\infty, \text{cop}}$ when r goes to infinity.

Remark 2.2. The convergence of $\mu_{F_n, \text{cop}}$ to $\mu_{\infty, \text{cop}}$ for some sequences (F_n) of balls was conjectured by Vardi and obtained by Pleasants and Huck: see Conjecture 1 in [Var99] and Theorem 1 in [PH13]. Their vocabulary and techniques are very different from those of the present paper, and our approach further yields Propositions 2.3, 2.4, and 2.7. See also the two last paragraphs of Section 2.3.

We will actually prove Proposition 2.3 instead of Theorem 2.1. Say that a sequence (F_n) of finite nonempty subsets of $\mathbb{Z}^d$ is a **Følner sequence** if for every $y \in \mathbb{Z}^d$, one has $|F_n \Delta (F_n + y)| = o(|F_n|)$. It suffices to check this condition for a family of y 's generating $\mathbb{Z}^d$ as a group. This condition is also equivalent to $\lim_n \frac{|\partial F_n|}{|F_n|} = 0$, where the **boundary** ∂F of $F \subset \mathbb{Z}^d$ is the set of the elements of $\mathbb{Z}^d \setminus F$ that are adjacent to an element of F for the usual (hypercubic) graph structure of $\mathbb{Z}^d$.

PROPOSITION 2.3. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$. Assume that $\mu_{F_n}(\{\omega : (0, \dots, 0) \in \omega\})$ converges to $1/\zeta(d)$.*

Then, $\mu_{F_n, \text{cop}}$ converges to $\mu_{\infty, \text{cop}}$.

Theorem 2.1 immediately follows from (A) and Proposition 2.3, which it thus suffices to prove. In Section 2.3, we will see that none of the Følner condition and the $1/\zeta(d)$ -assumption can be removed from Proposition 2.3. This proposition will itself be deduced from the following stochastic domination result. If μ and ν denote two probability measures on $\Omega_{\{0,1\}}$, we say that μ is **stochastically dominated** by ν if there is a coupling $(\mathcal{W}, \mathcal{W}')$ of (μ, ν) such that $\mathcal{W} \subset \mathcal{W}'$ almost surely.

PROPOSITION 2.4. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$. Assume that $\mu_{F_n, \text{cop}}$ converges to some probability measure μ .*

Then, μ is stochastically dominated by $\mu_{\infty, \text{cop}}$.

The goal of the remaining of this section is to prove Proposition 2.3 and Proposition 2.4.

LEMMA 2.5. *Let $d \geq 1$, let N be a positive integer, and let $\pi : \mathbb{Z}^d \rightarrow (\mathbb{Z}/N\mathbb{Z})^d$ denote reduction modulo N . Let (F_n) be a Følner sequence of $\mathbb{Z}^d$, and let Y_n denote a uniformly chosen element of F_n .*

Then, $\pi(Y_n)$ converges in distribution to the uniform measure on $(\mathbb{Z}/N\mathbb{Z})^d$.

Proof. Partition $\mathbb{Z}^d$ into boxes of the form $\prod_{i=1}^d \llbracket Nx_i, N(x_i+1)-1 \rrbracket$. For every n , say that an element x of F_n is n -good if the box B containing it satisfies $B \subset F_n$. Let Y_n denote a uniformly chosen element of F_n . Because (F_n) is a Følner sequence, the probability that Y_n is n -good converges to 1 as n goes to infinity. But if Y'_n denotes a uniformly chosen n -good element of F_n , then $\pi(Y'_n)$ is precisely uniform in $(\mathbb{Z}/N\mathbb{Z})^d$: one may generate Y'_n by picking independently a uniform element Y in $\llbracket 0, N-1 \rrbracket^d$ and a uniform element Y''_n in $\{x \in N\mathbb{Z}^d : x \text{ is } n\text{-good}\}$, and then writing $Y'_n = Y''_n + Y$. The result follows. $\square$

We will need the forthcoming Lemma 2.6, which requires the following definitions of pcop and $\mu_{\infty, \text{pcop}}$. Let $X = \{0, 1\}^{\mathcal{P}}$. The map $\text{pcop} : \mathbb{Z}^d \rightarrow X$ is given by $\text{pcop}(x)_p = \mathbb{1}_{x \notin p\mathbb{Z}^d}$. Let us now define $\mu_{\infty, \text{pcop}}$. For every $p \in \mathcal{P}$, pick a uniform coset $\mathcal{W}_p$ of $p\mathbb{Z}^d$ among the p^d possible ones. Do this independently for every p . One defines the random element $\mathcal{W}'$ of Ω_X via $\mathcal{W}'(x)_p := \mathbb{1}_{x \notin \mathcal{W}_p}$. We will denote by $\mu_{\infty, \text{pcop}}$ the distribution of the random variable $\mathcal{W}'$. Notice that the random element $\mathcal{W}$ of $\Omega_{\{0,1\}}$ defined by $\mathcal{W}(x) := \min_p \mathcal{W}'(x)_p$ has distribution $\mu_{\infty, \text{cop}}$.

LEMMA 2.6. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$.*

Then, $\mu_{F_n, \text{pcop}}$ converges to $\mu_{\infty, \text{pcop}}$.

Lemma 2.6 can be deduced from Lemma 2.5. We will instead deduce it from Lemma 2.8: see Remark 2.9. Our task is now to get Proposition 2.4 from Lemma 2.6, and Proposition 2.3 from Proposition 2.4. But before doing so, I would like to explain why Proposition 2.3 does *not* follow directly from Lemma 2.6.

 To deduce directly Proposition 2.3 from Lemma 2.6, one would need the continuity of the map $f : \Omega_{\{0,1\}^{\mathcal{P}}} \rightarrow \Omega_{\{0,1\}}$ defined by $f(\omega)_x = \min_p \omega(x)_p$. Indeed, $\text{cop} = f \circ \text{pcop}$. But this continuity, equivalent to that of the min map defined from $\{0,1\}^{\mathcal{P}}$ to $\{0,1\}$, does *not* hold. However, the min map is *upper semicontinuous*, which permits the proof of Proposition 2.4. Notice that there can be no way to do “as if” f was continuous, as the Følner assumption of Lemmas 2.5, 2.6 and 2.8 does *not* imply that $\mu_{F_n, \text{cop}}$ converges to $\mu_{\infty, \text{cop}}$ — see Remark 2.13.

Proof of Proposition 2.4. Let us make the assumptions of Proposition 2.4. For Y_n uniformly chosen in F_n , let us consider $(\tau_{-Y_n} \text{cop}, \tau_{-Y_n} \text{pcop}) \in \Omega_{\{0,1\}} \times \Omega_{\{0,1\}^{\mathcal{P}}} \cong \Omega_{\{0,1\} \times \{0,1\}^{\mathcal{P}}}$. As $\{0,1\} \times \{0,1\}^{\mathcal{P}}$ is compact, up to passing to a subsequence, we may assume that the distribution of $(\tau_{-Y_n} \text{cop}, \tau_{-Y_n} \text{pcop})$ converges to some probability measure ρ on $\Omega_{\{0,1\}} \times \Omega_{\{0,1\}^{\mathcal{P}}}$. Notice that for every $x \in \mathbb{Z}^d$ and $p \in \mathcal{P}$, one has

$$\text{pcop}(x)_p = 0 \implies \text{cop}(x) = 0.$$

Besides, for every x and p (which range over countable sets), the set

$$\{(\omega, \omega') : \omega(x) \leq \omega'(x)_p\}$$

is open and closed inside $\Omega_{\{0,1\}} \times \Omega_{\{0,1\}}^{\mathcal{P}}$. As a result, for ρ -almost every (ω, ω') , for every $x \in \mathbb{Z}^d$, one has $\omega(x) \leq \min_p \omega'(x)_p$. But recall that if $(\mathcal{W}, \mathcal{W}')$ denotes a random variable with distribution ρ , then $\mathcal{W}$ has distribution μ , $\mathcal{W}'$ has distribution $\mu_{\infty, \text{pcop}}$ — by Lemma 2.6 —, and thus $\min_p \mathcal{W}'(x, p)$ has distribution $\mu_{\infty, \text{cop}}$. Proposition 2.4 follows. $\square$

Proof of Proposition 2.3. Let us make the assumptions of Proposition 2.3. Up to taking a subsequence of (F_n) , we may assume that $\mu_{F_n, \text{cop}}$ converges to some μ . We want to prove that $\mu = \mu_{\infty, \text{cop}}$. By Proposition 2.4, there is a monotone coupling of $(\mu, \mu_{\infty, \text{cop}})$, i.e. some coupling ρ of $(\mu, \mu_{\infty, \text{cop}})$ such that for ρ -almost every $(\omega, \omega_{\infty})$, one has $\forall x, \omega(x) \leq \omega_{\infty}(x)$. But by the $1/\zeta(d)$ -assumption, one has

$$\mu(\{\omega : \omega(\vec{0}) = 1\}) = 1/\zeta(d) = \prod_{p \in \mathcal{P}} (1 - p^{-d}) = \mu_{\infty, \text{cop}}(\{\omega : \omega(\vec{0}) = 1\}).$$

As a result, for ρ -almost every $(\omega, \omega_{\infty})$, one has $\omega(\vec{0}) = \omega_{\infty}(\vec{0})$. The same argument applies for any $x \in \mathbb{Z}^d$ instead of $\vec{0}$. Indeed, the probability measure $\mu_{\infty, \text{cop}}$ is translation invariant by construction, and μ is translation invariant because (F_n) is a Følner sequence. Thus, μ is equal to $\mu_{\infty, \text{cop}}$ and Proposition 2.3 is proved. $\square$

2.2 The random gcd labelling

This section is devoted to proving Theorem 1.1.

Let us first define $\mu_{\infty, \text{gcd}}$. For every prime p , perform the following random choices:

- set $\mathcal{W}_0^p := \mathbb{Z}^d$,
- conditionally on $(\mathcal{W}_0^p, \dots, \mathcal{W}_{n-1}^p)$, pick a uniform coset $\mathcal{W}_n^p$ of $p^n \mathbb{Z}^d$ among those lying inside $\mathcal{W}_{n-1}^p$.

Do this independently for every p . We set the random p -adic valuation of a vertex x in $\mathbb{Z}^d$ to be $V_p(x) := \sup\{n \in \mathbb{N} : x \in \mathcal{W}_n^p\} \in \llbracket 0, \infty \rrbracket$. We define the random GCD profile to be the random map $x \mapsto \prod_{p \in \mathcal{P}} p^{V_p(x)}$. (This occurs almost surely nowhere, but one should set $\prod_{p \in \mathcal{P}} p^{V_p(x)}$ to be 0 whenever $\forall p, V_p(x) = \infty$.) The distribution of the random GCD profile is denoted by $\mu_{\infty, \text{gcd}}$. It is a priori a probability distribution on $\Omega_{\llbracket 0, \infty \rrbracket}$. By the Borel–Cantelli Lemma, for every $d \geq 2$, it is also a probability distribution on $\Omega_{\mathbb{N}}$, and it is as such that it will be considered from now on.

We will actually prove the following proposition.

PROPOSITION 2.7. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$. For every n , let Y_n denote a uniformly chosen element of F_n . Assume that the distribution of $\text{GCD}(Y_n)$ is tight (which implies $d \geq 2$).*

Then, $\mu_{F_n, \text{gcd}}$ converges to $\mu_{\infty, \text{gcd}}$.

Theorem 1.1 immediately follows from (B) and Proposition 2.7, which it thus suffices to prove. The forthcoming Lemma 2.8 uses the notion of profinite numbers. Define $\hat{\mathbb{Z}}$ as follows:

$$\hat{\mathbb{Z}} = \varprojlim_n \mathbb{Z}/n\mathbb{Z} = \left\{ x \in \prod_{n \geq 1} \mathbb{Z}/n\mathbb{Z} : \forall (m, n), m|n \implies \pi_{m,n}(x_n) = x_m \right\},$$

where $\pi_{m,n}$ denotes the morphism of reduction modulo m from $\mathbb{Z}/n\mathbb{Z}$ to $\mathbb{Z}/m\mathbb{Z}$. One can see $\mathbb{Z}$ as a dense subgroup of $\hat{\mathbb{Z}}$ via the injection

$$\Phi : N \mapsto (n \mapsto N + n\mathbb{Z}).$$

Elements of $\hat{\mathbb{Z}}$ are called **profinite numbers**. The product topology on $\prod_{n \geq 1} \mathbb{Z}/n\mathbb{Z}$ induces on $\hat{\mathbb{Z}}$ and on $\mathbb{Z} \subset \hat{\mathbb{Z}}$ the so-called **profinite topology**. It makes of $(\hat{\mathbb{Z}}, +)$ a metrisable compact topological group.

Recall that when p is a prime, the set of **p -adic integers** is defined by $\mathbb{Z}_p = \varprojlim_n \mathbb{Z}/p^n\mathbb{Z}$. The Chinese Remainder Theorem implies that $\hat{\mathbb{Z}}$ and $\prod_p \mathbb{Z}_p$ are isomorphic as topological groups (and even as rings), an isomorphism being given by

$$\hat{\mathbb{Z}} \ni \mathbf{n} \mapsto (n \mapsto \mathbf{n}_{p^n})_{p \in \mathcal{P}} \in \prod_p \mathbb{Z}_p.$$

Let $\text{prof} : \mathbb{Z}^d \rightarrow \hat{\mathbb{Z}}^d$ be defined by $\text{prof}(x) = (\Phi(x_1), \dots, \Phi(x_d)) \in \hat{\mathbb{Z}}^d$. In order to define a suitable probability measure $\mu_{\infty, \text{prof}}$, notice that there is a unique Haar measure on the compact group $\hat{\mathbb{Z}} \cong \prod_p \mathbb{Z}_p$, which corresponds to the product of Haar measures on $\mathbb{Z}_p$. A Haar distributed element of $\mathbb{Z}_p$ is a random p -adic integer $(X_n)_{n \geq 0}$ such that for every $n \geq 1$, conditionally on X_{n-1} , the element X_n is uniform among the p elements of $\mathbb{Z}/p^n\mathbb{Z}$ that reduce to X_{n-1} modulo p^{n-1} .

Pick a Haar distributed element Y of $\hat{\mathbb{Z}}^d$, i.e. d independent Haar distributed elements of $\hat{\mathbb{Z}}$. We denote by $\mu_{\infty, \text{prof}}$ the distribution of the following random element of $\Omega_{\hat{\mathbb{Z}}^d}$:

$$x \mapsto Y + \text{prof}(x).$$

LEMMA 2.8. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$.*

Then, $\mu_{F_n, \text{prof}}$ converges to $\mu_{\infty, \text{prof}}$.

Proof. As $\hat{\mathbb{Z}}$ is compact, up to passing to a subsequence, we may assume that $\mu_{F_n, \text{prof}}$ converges to some probability measure μ .

Let Y_n denote a uniformly chosen element of F_n , and set $X_n := \Phi(Y_n)$. By Lemma 2.5, for every N , the $(N!)$ -component of X_n converges in distribution to a uniform element of $\mathbb{Z}/N! \mathbb{Z}$. As a result, X_n converges in distribution to a Haar distributed element of $\hat{\mathbb{Z}}$. Therefore, if $\mathfrak{s}$ is μ -distributed, then $\mathfrak{s}(0)$ is Haar distributed on $\hat{\mathbb{Z}}$.

It thus suffices to prove that μ gives probability 1 to the following set: $\{\sigma \in \Omega_{\hat{\mathbb{Z}}} : \forall x \in \mathbb{Z}^d, \sigma(x) = \sigma(\vec{0}) + \text{prof}(x)\}$. But this is clear as it is closed and has probability 1 for every $\mu_{F_n, \text{prof}}$. $\square$

Remark 2.9. As the map $f : \hat{\mathbb{Z}} \rightarrow \{0, 1\}^{\mathcal{P}}$ defined by $f(\mathfrak{n})_p := \mathbb{1}_{\mathfrak{n}_p \neq 0}$ is continuous, Lemma 2.6 follows directly from Lemma 2.8.

As we are interested in the GCD of random elements of $\mathbb{Z}$, it is useful to recall what the GCD of a profinite number is. While the GCD of an element of $\mathbb{Z}$ is a natural number (possibly 0), the GCD of a profinite number will be a *supernatural* number. The set of **supernatural numbers** is $\mathfrak{N} := \llbracket 0, \infty \rrbracket^{\mathcal{P}}$. The set $\llbracket 0, \infty \rrbracket$ is endowed with the usual topology (a set U is open if and only if for n large enough, $n \in U \iff \infty \in U$), and $\mathfrak{N} = \llbracket 0, \infty \rrbracket^{\mathcal{P}}$ is endowed with the product of these topologies. One can see any nonnegative integer $n \in \mathbb{N}$ as a supernatural number via the following injection:

$$\Psi : n \mapsto (p \mapsto p\text{-adic valuation of } n).$$

The p -adic valuation of a profinite number $\mathfrak{n}$ is

$$v_p(\mathfrak{n}) := \sup\{k \in \mathbb{N} : \mathfrak{n}_{p^k} = 0\} \in \llbracket 0, \infty \rrbracket.$$

The GCD of $(\mathfrak{n}_1, \dots, \mathfrak{n}_d)$ is the following supernatural number:

$$p \mapsto \min(v_p(\mathfrak{n}_1), \dots, v_p(\mathfrak{n}_d)).$$

Let $\text{super} : \mathbb{Z}^d \rightarrow \mathfrak{N}$ be defined by $x \mapsto \Psi(\text{GCD}(x))$. Given a Haar distributed element Y of $\hat{\mathbb{Z}}^d$, denote by $\mu_{\infty, \text{super}}$ the distribution of the following random element of $\Omega_{\mathfrak{N}}$:

$$x \mapsto \text{GCD}(Y + \text{prof}(x)).$$

LEMMA 2.10. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$.*

Then, $\mu_{F_n, \text{super}}$ converges to $\mu_{\infty, \text{super}}$.

Proof. As the GCD-map is continuous from $\hat{\mathbb{Z}}$ to $\mathfrak{N}$, this follows directly from Lemma 2.8. $\square$

Notice that the conclusion of Lemma 2.10 is quite close to the one we are looking for. Assume further that $d \geq 2$. Then, as $\sum_{p \in \mathcal{P}} p^{-d} < \infty$, the Borel–Cantelli Lemma yields that if $\mathbf{X}$ is $\mu_{\infty, \text{super}}$ -distributed, then almost surely, every $x \in \mathbb{Z}^d$ satisfies $\mathbf{X}(x) \in \Psi(\mathbb{N}^*)$. Recall that $\Psi(\mathbb{N}^*)$ consists in

finitely supported elements of $\mathfrak{N} = \llbracket 0, \infty \rrbracket^{\mathcal{P}}$ all the values of which are finite. If $\mathbf{n} \in \Psi(\mathbb{N}^*)$, then $\mathbf{n} = \Psi\left(\prod_{p \in \mathcal{P}} p^{\mathbf{n}_p}\right)$ and we write $\chi(\mathbf{n}) = \prod_{p \in \mathcal{P}} p^{\mathbf{n}_p} \in \mathbb{N}^*$. Therefore, the random variable $(\chi(\mathbf{X}(x)))_{x \in \mathbb{Z}^d}$ has distribution $\mu_{\infty, \text{gcd}}$.

$\Diamond$ If χ was continuous and defined everywhere on $\mathfrak{N}$, one could directly deduce from Lemma 2.10 that $\mu_{F_n, \text{gcd}}$ converges to $\mu_{\infty, \text{gcd}}$; but the actual properties of χ do not allow us to perform this derivation, even in an indirect manner. Indeed, (F_n) being Følner and d being at least 2 do *not* suffice to guarantee this convergence: these conditions do not even suffice to guarantee convergence of $\mu_{F_n, \text{cop}}$ to $\mu_{\infty, \text{cop}}$. See Remark 2.13. We need the tightness assumption and this goes through the following easy lemma.

Denote by $f_{\#}\rho$ the pushforward by f of a given measure ρ .

LEMMA 2.11. *Let X and Y be Polish spaces. Let $f : X \rightarrow Y$ be a continuous injective map that maps every Borel subset of X to a Borel subset of Y . Let $(\mu_n)_{n \leq \infty}$ denote a sequence of probability measures on X . For $n \leq \infty$, set $\nu_n := f_{\#}\mu_n$. Assume that ν_n converges to ν_{∞} and that μ_n converges to some probability measure μ .*

Then μ and μ_{∞} are equal.

Proof. By continuity of f , the sequence $\nu_n = f_{\#}\mu_n$ converges to $f_{\#}\mu$. As this sequence also converges to $f_{\#}\mu_{\infty}$, we have $f_{\#}\mu = f_{\#}\mu_{\infty}$. By injectivity of f , we have $\mu(A) = f_{\#}\mu(f(A)) = f_{\#}\mu_{\infty}(f(A)) = \mu_{\infty}(f(A))$, where the assumptions indeed guarantee that $f(A)$ is Borel. $\square$

Notice that in Lemma 2.11, if one does not assume that μ_n converges to some probability measure, then one cannot deduce that μ_n converges to μ_{∞} . A counterexample goes as follows. Take $X = (0, 1]$, $Y = \mathbb{R}/\mathbb{Z}$, and set $f : X \rightarrow Y$ to be reduction modulo 1. For $n \in \mathbb{N}^*$, set $\mu_n = \delta_{1/n}$, and let $\mu_{\infty} = \delta_1$.

Proof of Proposition 2.7. Let (F_n) and (Y_n) be as in Proposition 2.7. By hypothesis, $\text{GCD}(Y_n)$ is tight. Together with the assumption that (F_n) is a Følner sequence, this implies that $d \geq 2$ and that for every $x \in \mathbb{Z}^d$, $\text{GCD}(Y_n + x)$ is tight. Therefore, the $(\mathbb{N}$ -indexed) sequence of random variables $(\text{GCD}(Y_n + x))_{x \in \mathbb{Z}^d}$ is tight. Up to passing to a subsequence, we may thus assume that its distribution converges to some probability measure μ on $\Omega_{\mathbb{N}}$.

Let $X := \Omega_{\mathbb{N}}$ and $Y := \Omega_{\mathfrak{N}}$. For $n \in \mathbb{N}$, let $\mu_n := \mu_{F_n, \text{gcd}}$. Set $\mu_{\infty} := \mu_{\infty, \text{gcd}}$, which is a well-defined probability measure on $\Omega_{\mathbb{N}}$ because of the Borel–Cantelli Lemma ($\sum_{p \in \mathcal{P}} p^{-d} < \infty$). Let $f : X \rightarrow Y$ be defined by $f(\sigma) = (\Psi(\sigma_x))_{x \in \mathbb{Z}^d}$. This map is injective, continuous, and maps Borel subsets of X to Borel subsets of Y . By Lemma 2.10, $f_{\#}\mu_n$ converges to $f_{\#}\mu_{\infty}$. As μ_n converges to μ , Lemma 2.11 yields that μ_n converges to μ_{∞} , which is the desired result. $\square$

2.3 Remarks

We insist that in none of our results, we ask for the sequence (F_n) to be monotone, or for $\bigcup_n F_n$ to be equal to $\mathbb{Z}^d$. The fact that $|F_n|$ tends to infinity is a consequence of being Følner.

Even though the conclusion of Proposition 2.7 implies that of Proposition 2.3, it is not the case that Proposition 2.3 can be derived from Proposition 2.7. Indeed, one has the following fact.

FACT 2.12. *For every $d \geq 1$, there is a Følner sequence (F_n) of $\mathbb{Z}^d$ such that the following conditions hold:*

- *if Y_n denotes a uniform element of F_n , the distribution of $\text{GCD}(Y_n)$ is not tight,*
- *the proportion of coprime vectors in F_n converges to $1/\zeta(d)$.*

Proof. This fact can be derived from the Chinese Remainder Theorem or from the proof of Theorem 1.1. We will proceed in the second manner.

If $d = 1$, any Følner sequence satisfies automatically the desired properties. Let us thus assume that $d \geq 2$, so that $\sum_{p \in \mathcal{P}} p^{-d} < \infty$. Let $\varepsilon > 0$ and $K \in \mathbb{N}$. One has

$$\mu_{\infty, \text{super}}(\{\sigma : \forall p, \sigma(0)_p = 0\}) = \prod_{p \in \mathcal{P}} (1 - p^{-d}) = 1/\zeta(d).$$

By translation invariance, one has

$$\int \frac{|\{x \in \llbracket 1, N \rrbracket^d : \forall p, \sigma(x)_p = 0\}|}{N^d} d\mu_{\infty, \text{super}}(\sigma) = 1/\zeta(d).$$

Either by ergodicity³ or because of Proposition 2.4, one can pick an arbitrarily large N such that this proportion has a positive probability to be ε -close to $1/\zeta(d)$. Fixing such an N , one can thus find a deterministic subset U of $\llbracket 1, N \rrbracket^d$ such that:

$$\left| \frac{|U|}{N^d} - \frac{1}{\zeta(d)} \right| \leq \varepsilon \quad \text{and} \quad \mu_{\infty, \text{super}}(\{\sigma : \forall x \in U, \forall p \in \mathcal{P}, \sigma(x)_p = 0\}) > 0.$$

Let $\mathfrak{s}$ be a random variable with distribution $\mu_{\infty, \text{super}}$. As $d \geq 2$, one can pick some $P \geq \max(K, N)$ such that with positive probability, the following two conditions hold:

1. $\forall x \in U, \forall p \in \mathcal{P}, \mathfrak{s}(x)_p = 0,$

³Later in this section, we will see that $\mu_{\infty, \text{super}}$ is ergodic, i.e. that any translation invariant event has probability 0 or 1. One can then use an ergodic theorem such as Theorem 1.1 in [Lin01].

$$2. \forall x \in \llbracket 1, N \rrbracket^d, \forall p \in \mathcal{P}, (p \geq P \implies \mathfrak{s}(x)_p = 0).$$

Recall that $\mu_{\infty, \text{super}}$ is defined in terms of the cosets $\mathcal{W}_p^n$. Let $\mathbf{p}$ denote an injective map from $\llbracket 1, N \rrbracket^d$ to $\{p \in \mathcal{P} : p \geq P\}$. We will use this injection to modify $\mathfrak{s}$ into a new random variable $\mathfrak{s}'$. For every $x \in \llbracket 1, N \rrbracket^d \setminus U$, resample $\left(\mathcal{W}_n^{\mathbf{p}(\mathbf{x})}\right)_n$ and condition $\mathcal{W}_1^{\mathbf{p}(\mathbf{x})}$ to contain x : do this independently of $\mathfrak{s}$ and independently for every $x \in \llbracket 1, N \rrbracket^d \setminus U$. The distribution of the ensuing random variable $\mathfrak{s}'$ has a density relative to $\mu_{\infty, \text{super}}$. As $\mathfrak{s}'$ almost surely satisfies the following conditions:

$$1. \forall x \in U, \prod_p p^{\mathfrak{s}'(x)_p} = 1,$$

$$2. \forall x \in \llbracket 1, N \rrbracket^d \setminus U, \prod_p p^{\mathfrak{s}'(x)_p} \geq K,$$

the corresponding conditions are satisfied by $\mathfrak{s}$ with positive probability. Denote the corresponding event by E .

By Theorem 1.1, the proportion of $x \in \llbracket 1, n \rrbracket^d$ such that $\tau_{-x} \text{super}$ satisfies E converges to a positive number, hence is positive for n large enough. In particular, there is some $y \in \mathbb{Z}^d$ such that $\tau_{-y} \text{super}$ satisfies E . We say that any such y is a (K, N, ε) -counterexample. For every n , pick some y^n that is an $(n, m_n, 1/n)$ -counterexample for some $m_n \geq n$: the sequence $F_n = \prod_{i=1}^d \llbracket y_i^n + 1, y_i^n + m_n \rrbracket$ satisfies the desired properties. $\square$

Remark 2.13. Likewise, the $1/\zeta(d)$ -condition cannot be removed from Proposition 2.3: it is well-known that the Chinese Remainder Theorem implies that there are arbitrarily large boxes $F_n = x_n + \llbracket 0, N \rrbracket^d$ in $\mathbb{Z}^d$ devoid of coprime points. This also follows from Theorem 2.1, as $\mu_{\infty, \text{cop}}$ has a positive probability to colour black the whole box $\llbracket 0, N \rrbracket^d$ — use one prime per point of the box. Such a sequence (F_n) is Følner and yet $\mu_{F_n, \text{cop}}$ converges δ_0 . The Følner condition cannot be removed from Proposition 2.3 either.

All the results in this paper concerning visibility extend readily to lattices in $\mathbb{R}^d$, as any such lattice may be mapped to $\mathbb{Z}^d$ by a linear automorphism of $\mathbb{R}^d$, which preserves visibility.

Notice that $\mu_{\infty, \text{cop}}$ and $\mu_{\infty, \text{gcd}}$ are not only translation invariant but also $\text{GL}_d(\mathbb{Z})$ -invariant. Even though cop and gcd are indeed $\text{GL}_d(\mathbb{Z})$ -invariant, the $\text{GL}_d(\mathbb{Z})$ -invariance of these measures does *not* follow from Theorems 1.1 and 2.1. One way to understand why goes as follows. Every orbit of the multiplicative group G generated by $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ contains a unique point inside $A := \{x \in \mathbb{Z}^2 : 0 \leq x_1 < |x_2|\} \cup (\mathbb{Z} \times \{0\})$. Consider the colouring of $\mathbb{Z}^2$ which is defined as a the chessboard colouring on A (say white if the sum of coordinates is odd and black otherwise), and take its unique G -invariant extension. For $F_n = A \cap \llbracket -n, n \rrbracket^2$, this colouring seen from a uniform

point in F_n converges to the unbiased choice of one of the two chessboard colourings of the plane: this probability measure is *not* G -invariant.

Finally, let us mention a few questions that can be asked only with the formalism of local limits, i.e. with $\mu_{\infty, \text{cop}}$ or $\mu_{\infty, \text{gcd}}$ instead of simply one convergence result per cylindrical event as in [PH13]. For every $c \in \{\text{cop}, \text{gcd}, \text{prof}, \text{super}\}$, we can ask whether the measure $\mu_{\infty, c}$ is ergodic — i.e. if every translation invariant measurable subset has $\mu_{\infty, c}$ -probability 0 or 1. These measures are indeed ergodic. For simplicity, we expose the argument for $\mu_{\infty, \text{cop}}$. The Chinese Remainder Theorem guarantees the following fact: if $p_1, \dots, p_n$ denote the first n primes, and if for every $i \leq n$, C_i denotes a coset of $p_i \mathbb{Z}^d$ in $\mathbb{Z}^d$, then there is a translation mapping every C_i to $p_i \mathbb{Z}^d$. As a result, whenever we consider only finitely many primes, choosing one coset per prime yields a *deterministic* outcome up to translation. One concludes by noting that if a translation invariant probability measure on $\{0, 1\}^{\mathbb{Z}^d \times \mathcal{P}}$ yields ergodic measures in projection to any $\{0, 1\}^{\mathbb{Z}^d \times \{p_1, \dots, p_n\}}$, then the measure under study is itself ergodic. This is easily proved by martingale theory, and similar reasonings are classical in the study of profinite actions.

Considering $\mathbb{Z}^d$ to be endowed with its usual (hypercubic) graph structure, one may also ask questions of percolation theory [Gri99, LP16]: how many infinite white (resp. black) connected components does the colouring $\mu_{\infty, \text{cop}}$ yield? By ergodicity, these numbers have to be deterministic outside some event of probability zero. One can derive from Theorem 3.3 in [Var99] that, for $d = 2$ hence for any $d \geq 2$, there is at least one infinite white connected component almost surely. One can derive from Theorem 3.4 in [Var99] that, for $d = 2$, there is almost surely at most one infinite white connected component and no infinite black component.

3 Further generalisations of Theorem 1.1

3.1 Sampling along affine subspaces

Proposition 3.1 is the analog of Proposition 2.7 for an observer picked uniformly in an affine subspace of $\mathbb{Z}^d$ rather than in the whole space $\mathbb{Z}^d$.

Endow $\llbracket 0, \infty \rrbracket$ with its usual topology. Define $\text{gcd}_{\infty} \in \Omega_{\llbracket 0, \infty \rrbracket}$ by setting $\text{gcd}_{\infty}(x) := \text{GCD}(x)$. Let $(\mu_n)_{n \leq \infty}$ denote a sequence of probability measures on $\Omega_{\llbracket 0, \infty \rrbracket}$. Let $V \subset \mathbb{Z}^d$ and denote by π the projection from $\llbracket 0, \infty \rrbracket^{\mathbb{Z}^d}$ to $\llbracket 0, \infty \rrbracket^V$. We say that μ_n converges to μ_{∞} **on** V if $\pi_{\#} \mu_n$ — the pushforward of μ_n by π — converges to $\pi_{\#} \mu_{\infty}$ when n goes to infinity.

Given Γ an infinite subgroup of $\mathbb{Z}^d$, one defines the probability measure $\mu_{\infty, \text{gcd}_{\infty}}^{\Gamma}$ by taking the definition of $\mu_{\infty, \text{gcd}}$ but asking furthermore at every step that $\mathcal{W}_n^p$ intersects Γ . This corresponds to taking a Haar distributed

element in the closure of Γ in $\hat{\mathbb{Z}}^d$.

Finally, we say that a sequence (F_n) of nonempty finite subsets of Γ is a **Følner sequence for Γ** if for every $y \in \Gamma$, one has $|F_n \Delta (F_n + y)| = o(|F_n|)$.

PROPOSITION 3.1. *Let Γ denote a subgroup of $\mathbb{Z}^d$ of rank $k \geq 1$ which is maximal among subgroups of rank k . Let (F_n) be a Følner sequence for Γ . Let Y_n denote a uniform element of F_n .*

Then, $\mu_{F_n, \text{gcd}_\infty}$ converges to $\mu_{\infty, \text{gcd}_\infty}^\Gamma$ on $\mathbb{Z}^d \setminus \Gamma$. If furthermore $k = 1$ or $\text{GCD}(Y_n)$ is tight, then $\mu_{F_n, \text{gcd}_\infty}$ converges to $\mu_{\infty, \text{gcd}_\infty}^\Gamma$.

Remark 3.2. The view from a “uniform point” in an affine subspace $\Gamma + y$ is just the view seen from a “uniform point” in Γ shifted by $-y$. One may also notice that if one starts with a group Γ that is not maximal given its rank, then it lies in a unique such group, which is the intersection of its (rational or real) linear span with $\mathbb{Z}^d$: denote it by $\tilde{\Gamma}$. It follows from the proof of Proposition 3.1 that the following holds. Let (F_n) be a Følner sequence for Γ . Let Y_n denote a uniform element of F_n . Assume that for every $y \in \tilde{\Gamma}$, $\text{GCD}(Y_n + y)$ is tight. Then $\mu_{F_n, \text{gcd}_\infty}$ converges to $\mu_{\infty, \text{gcd}_\infty}^\Gamma$. Actually, it suffices to make the assumption for a system of representatives of $y \in \tilde{\Gamma}$ for the equivalence relation “being equal modulo Γ ”, i.e. for finitely many y ’s.

COROLLARY 3.3. *Let Γ denote a subgroup of $\mathbb{Z}^d$ of rank $k \geq 1$ which is maximal among subgroups of rank k . Let F denote a bounded subset of the linear span of Γ . For every $r \in (0, \infty)$, set $F_r := \{x \in \Gamma : r^{-1}x \in F\}$. Assume that $\frac{|F_r|}{r^k}$ converges to a nonzero limit when r tends to infinity and that (F_n) is a Følner sequence for Γ .*

Then, $\mu_{F_n, \text{gcd}_\infty}$ converges to $\mu_{\infty, \text{gcd}_\infty}^\Gamma$ as n goes to infinity.

Proof of Proposition 3.1. Exactly as in Section 2.2, one defines $\mu_{\infty, \text{super}}^\Gamma$ and proves that $\mu_{F_n, \text{super}}^\Gamma$ converges to $\mu_{\infty, \text{super}}^\Gamma$. Let $d(\cdot, \Gamma)$ be as in the conclusion of Lemma 3.4. Let $X := \prod_{x \in \mathbb{Z}^d \setminus \Gamma} \psi([1, d(x, \Gamma)])$. As the closed subset $X \subset Y := \mathfrak{N}^{\mathbb{Z}^d \setminus \Gamma}$ has probability 1 for every measure $\mu_{F_n, \text{super}}^\Gamma$, it is the case that $\mu_{\infty, \text{super}}^\Gamma(X) = 1$. Therefore, one can also consider these measures as probability measures on X : with this point of view, denote them by μ_n , with $n \leq \infty$. One has that μ_n converges to μ_∞ — that is for the weak topology on probability measures on X . One way to see this is to notice that as X is compact, every subsequence of (μ_n) admits a converging subsequence and then to apply Lemma 2.11 to the restriction of the identity map from X to Y . As the map $f : X \rightarrow [0, \infty]^{\mathbb{Z}^d \setminus \Gamma}$ defined by $f(\sigma)_x := \prod_{p \in \mathcal{P}} p^{\sigma(x)_p}$ is continuous, one has that $\mu_{F_n, \text{gcd}_\infty}$ converges to $\mu_{\infty, \text{gcd}_\infty}^\Gamma$ on $\mathbb{Z}^d \setminus \Gamma$.

Let us now assume that $k = 1$. As $\sum_{p \in \mathcal{P}} \frac{1}{p} = \infty$, the second Borel–Cantelli Lemma yields that $\mu_{\infty, \text{super}}^\Gamma$ -almost every configuration gives the label ∞ to every element of Γ . If Y_n denotes a uniformly chosen element of F_n and y some element of Γ , as k is equal to 1, one has that $\text{GCD}(Y_n + y)$

converges in probability to ∞ . Together with convergence on $\mathbb{Z}^d \setminus \Gamma$, this implies that $\mu_{F_n, \text{gcd}_\infty}$ converges to $\mu_{\infty, \text{gcd}_\infty}^\Gamma$.

Instead of $k = 1$, now assume that $\text{GCD}(Y_n)$ is tight. As (F_n) is a Følner sequence for Γ , one has that for every $y \in \Gamma$, the random variable $\text{GCD}(Y_n + y)$ is tight. Together with convergence on $\mathbb{Z}^d \setminus \Gamma$, this implies that the sequence $(\mu_{F_n, \text{gcd}_\infty})$ is tight. One then concludes as in the proof of Proposition 2.7. $\square$

LEMMA 3.4. *Let (d, k) satisfy $1 \leq k \leq d$. Let Γ be a subgroup of $\mathbb{Z}^d$ of rank k and maximal with this property. Then, there is a norm $\| \cdot \|$ on $\mathbb{R}^d$ such that for every $N \geq 1$ and every $x \in N\mathbb{Z}^d$, one has*

$$d(x, \Gamma) > 0 \implies d(x, \Gamma) \geq N,$$

where $d(x, \Gamma) := \min\{\|x - y\| : y \in \Gamma\}$.

Proof. By a change of coordinates and by maximality of Γ , one may assume that $\Gamma = \mathbb{Z}^k \times \{0_{d-k}\}$, in which case the lemma is clear. Notice that any element of $\text{GL}_d(\mathbb{Z})$ maps $N\mathbb{Z}^d$ precisely to itself. $\square$

Proof of Corollary 3.3. Let Y_n denote a uniform element of F_n . By Proposition 3.1, it suffices to assume that $k \geq 2$ and to show that $\text{GCD}(Y_n)$ is tight. Since GCD 's are unchanged by $\text{GL}_d(\mathbb{Z})$, we may assume that Γ is equal to $\mathbb{Z}^k \times \{0_{d-k}\}$, and tightness results from (B). $\square$

3.2 Graphon and local-graphon limits

This section is devoted to (variations of) the following question: if $X_1, \dots, X_N$ are sampled “uniformly” and independently in $\mathbb{Z}^d$, can we describe the set of (i, j) 's such that X_i is visible from X_j , i.e. such that $\text{GCD}(X_i - X_j) = 1$?

A “**graphon**” is the data of a standard probability space $(\mathfrak{X}, \mathbb{P})$ together with a measurable function $f : \mathfrak{X}^2 \rightarrow [0, 1]$ that is symmetric, i.e. satisfies $\forall (x_1, x_2) \in \mathfrak{X}^2, f(x_1, x_2) = f(x_2, x_1)$. Two “graphons” $(\mathfrak{X}, \mathbb{P}_\mathfrak{X}, f)$ and $(\mathfrak{Y}, \mathbb{P}_\mathfrak{Y}, g)$ are said to **induce the same graphon** if, up to throwing away sets of measure zero, there is a measure-preserving isomorphism $\pi : (\mathfrak{X}, \mathbb{P}_\mathfrak{X}) \rightarrow (\mathfrak{Y}, \mathbb{P}_\mathfrak{Y})$ such that $\forall (x_1, x_2) \in \mathfrak{X}, f(x_1, x_2) = g(\pi(x_1), \pi(x_2))$. A **graphon** is an equivalence class of “graphons” for the relation “inducing the same graphon”. We say that a graphon is **represented** by any “graphon” that induces it. See [BCL⁺08].

Let $\mathcal{G}_n = (V_n, E_n)$ denote a sequence of random⁴ finite graphs such that $|V_n|$ converges in probability to infinity. It is said to **converge** to the (deterministic) graphon represented by $(\mathfrak{X}, \mathbb{P}, f)$ if the following holds: for every k , if $(X_1^n, \dots, X_k^n)$ denotes a uniform element of V_n^k , then the random variable $(\mathbb{1}_{\{X_i^n, X_j^n\} \in E_n})_{1 \leq i < j \leq k}$ converges in distribution to (the distribution of)

⁴One does not need $\mathcal{G}_n$ and $\mathcal{G}_m$ to be defined on the same probability space.

the random variable $(f(X_i^\infty, X_j^\infty))_{1 \leq i < j \leq k}$, where the X_i^∞ 's are independent random variables of distribution $\mathbb{P}$. See [DJ08].

Consider the following standard probability space $\mathfrak{X}_0 := \prod_{p \in \mathcal{P}} (\mathbb{Z}/p\mathbb{Z})^d$, endowed with the product of uniform measures. Consider the measurable function $\delta : \mathfrak{X}_0^2 \rightarrow [0, 1]$ defined by $\delta(x_1, x_2) := \mathbb{1}_{\forall p, x_1(p) \neq x_2(p)}$. See Figure 1.




Figure 1: Visualisation of the graphon $(\mathfrak{X}_0, \delta)$ for $d = 2$. After suitable identification of $\mathfrak{X}_0$ with $[0, 1]$, the set $\delta^{-1}(\{0\})$ is represented in black. Each square at scale n is divided into $p_n^d \times p_n^d$ smaller squares, where (p_n) denotes the sequence of prime numbers.

By the same arguments as in Section 2.1, one can prove the following result.

PROPOSITION 3.5. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$. Assume that the probability that a uniform element of F_n is coprime converges to $1/\zeta(d)$. Let $\mathcal{G}_n$ denote the random graph with vertex-set F_n and an edge between two distinct vertices if and only if the one is visible from the other.*

Then, $\mathcal{G}_n$ converges to the graphon represented by $(\mathfrak{X}_0, \delta)$.

Actually, we can even get a result combining both Proposition 2.3 and Proposition 3.5.

PROPOSITION 3.6. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$. Assume that the probability that a uniform element of F_n is coprime converges to $1/\zeta(d)$. Let (X_n) denote a sequence of independent uniform elements of $\mathfrak{X}_0$.*

Let $M \geq 1$ and $R \geq 1$. For every n , let $(Y_m^n)_{1 \leq m \leq M}$ denote M independent uniform elements in F_n . Consider the following random maps:

$$\begin{aligned} \psi_n : ([1, M] \times [-R, R]^d)^2 &\longrightarrow \{0, 1\} \\ ((m_0, y_0), (m_1, y_1)) &\longmapsto \mathbb{1}_{Y_{m_0}^n + y_0 \text{ is visible from } Y_{m_1}^n + y_1}, \end{aligned}$$

$$\begin{aligned}\psi_\infty : (\llbracket 1, M, \rrbracket \times \llbracket -R, R \rrbracket^d)^2 &\longrightarrow \{0, 1\} \\ ((m_0, y_0), (m_1, y_1)) &\longmapsto \mathbb{1}_{\forall p, X_{m_0}(p) + \overline{y_0} \neq X_{m_1}(p) + \overline{y_1}}.\end{aligned}$$

Then, the distribution of ψ_n converges to that of ψ_∞ .

This theorem can be readily adapted to the whole GCD profile (one assumes tightness, considers maps to $\llbracket 0, \infty \rrbracket$ rather than to $\{0, 1\}$, and predicts the GCD of $(Y_{m_0}^n + y_0) - (Y_{m_1}^n + y_1)$) and the case of affine subspaces.

Let us conclude with a last generalisation. One may be interested in other arithmetic conditions than coprimality: for example, saying that a number is *k-free* if no p^k divides it, one may colour in white the *k-free* points of $\mathbb{Z}$, or the points in $\mathbb{Z}^d$ with *k-free* GCD. See [BMP00, Ces85, Mir47, Mir48, PH13]. One can generalise Proposition 3.6 to such contexts as follows. Recall that $\mathbb{Z}$ and $\hat{\mathbb{Z}}$ can be endowed with the profinite topology. Likewise, $\mathbb{Z}^d$ and $\hat{\mathbb{Z}}^d$ can be endowed with the product of profinite topologies, which we also refer to as the profinite topology.

Example. For every $k \geq 0$, the set $\{x \in \mathbb{Z}^d : \text{GCD}(x) \text{ is } k\text{-free}\}$ is profinitely closed in $\mathbb{Z}^d$. For $k = 1$, this coincides with the set $\{x \in \mathbb{Z}^d : \text{GCD}(x) = 1\}$.

Let V be a subset of $\mathbb{Z}^d$, and let $\text{col}_V := \mathbb{1}_V$. Let $\mu_{\infty, \text{col}_V}$ denote the distribution of $x \mapsto \mathbb{1}_{Y+x \in \overline{V}}$, where Y is Haar distributed in $\hat{\mathbb{Z}}^d$ and $\overline{V}$ denotes the closure of the set V in $\hat{\mathbb{Z}}^d$.

PROPOSITION 3.7. *Let $d \geq 1$ and let (F_n) be a Følner sequence of $\mathbb{Z}^d$. Let V denote a subset of $\mathbb{Z}^d$. Assume that V is profinitely closed in $\mathbb{Z}^d$. Assume that $\frac{|V \cap F_n|}{|F_n|}$ converges to $\mu_{\infty, \text{col}_V}(\{\omega : \vec{0} \in \omega\})$. Let (X_m) denote a sequence of independent Haar distributed elements of $\hat{\mathbb{Z}}^d$.*

Let $M \geq 1$ and $R \geq 1$. For every n , let $(Y_m^n)_{1 \leq m \leq M}$ denote M independent uniform elements in F_n . Consider the following random maps:

$$\begin{aligned}\psi_n : (\llbracket 1, M, \rrbracket \times \llbracket -R, R \rrbracket^d)^2 &\longrightarrow \{0, 1\} \\ ((m_0, y_0), (m_1, y_1)) &\longmapsto \mathbb{1}_{(Y_{m_0}^n + y_0) - (Y_{m_1}^n + y_1) \in V}, \\ \psi_\infty : (\llbracket 1, M, \rrbracket \times \llbracket -R, R \rrbracket^d)^2 &\longrightarrow \{0, 1\} \\ ((m_0, y_0), (m_1, y_1)) &\longmapsto \mathbb{1}_{X_{m_0} - X_{m_1} + \text{prof}(y_0 - y_1) \in \overline{V}}.\end{aligned}$$

Then, the distribution of ψ_n converges to that of ψ_∞ .

Acknowledgements At the end of a talk given by Nathanaël Enriquez, Bálint Virág asked him about the local limit of visible points: I would like to thank Bálint Virág for asking this question and Nathanaël Enriquez for letting me know of it. I am also grateful to Nathanaël Enriquez for many enthusiastic discussions about this project. I am thankful to Subhajit Goswami and Aran Raoufi for comments on an earlier version of this paper. Finally, I am grateful to my postdoctoral advisors Nicolas Curien and Jean-François Le Gall, as well as to the ERC grant GeoBrown and the Université Paris-Sud, for providing me with an excellent working environment.

References

- [Bab91] László Babai. Vertex-transitive graphs and vertex-transitive maps. *Journal of graph theory*, 15(6):587–627, 1991.
- [BCL⁺08] Christian Borgs, Jennifer Chayes, László Lovász, Vera Sós, and Katalin Vesztegombi. Convergent sequences of dense graphs I: Subgraph frequencies, metric properties and testing. *Advances in Mathematics*, 219(6):1801–1851, 2008.
- [BCZ00] Florin Boca, Cristian Cobeli, and Alexandru Zaharescu. Distribution of lattice points visible from the origin. *Communications in Mathematical Physics*, 213(2):433–470, 2000.
- [BH15] Michael Baake and Christian Huck. Ergodic properties of visible lattice points. *Proceedings of the Steklov Institute of Mathematics*, 288(1):165–188, 2015.
- [BMP00] Michael Baake, Robert Moody, and Peter Pleasants. Diffraction from visible lattice points and k th power free integers. *Discrete Mathematics*, 221(1-3):3–42, 2000.
- [BS01] Itai Benjamini and Oded Schramm. Recurrence of distributional limits of finite planar graphs. *Electron. J. Probab.*, 6(23):1–13, 2001.
- [Ces81] Ernesto Cesàro. Question proposée 75. *Mathesis*, 1:184, 1881.
- [Ces83] Ernesto Cesàro. Question 75 (solution). *Mathesis*, 3:224–225, 1883.
- [Ces85] Ernesto Cesàro. Sur le plus grand commun diviseur de plusieurs nombres. *Annali di Matematica Pura ed Applicata (1867-1897)*, 13(1):291–294, 1885.
- [CFF19] Javier Cilleruelo, José Fernández, and Pablo Fernández. Visible lattice points in random walks. *European Journal of Combinatorics*, 75:92–112, 2019.
- [Dir51] Gustav Lejeune Dirichlet. *Über die Bestimmung der mittleren Werthe in der Zahlentheorie*. Dümmler in Komm., 1851.
- [DJ08] Persi Diaconis and Svante Janson. Graph limits and exchangeable random graphs. *Rendiconti di Matematica, Serie VII*, 28:33–61, 2008.
- [DL01] Reinhard Diestel and Imre Leader. A conjecture concerning a limit of non-Cayley graphs. *Journal of Algebraic Combinatorics*, 14(1):17–25, 2001.
- [FM16] Andrea Ferraguti and Giacomo Micheli. On the Mertens–Cesàro Theorem for number fields. *Bulletin of the Australian Mathematical Society*, 93(2):199–210, 2016.
- [Gar15] Olivier Garet. Les lois Zêta pour l’arithmétique. *Quadrature*, 96, 2015.
- [GHKM18] Edray H Goins, Pamela E Harris, Bethany Kubik, and Aba Mbirika. Lattice point visibility on generalized lines of sight. *The American Mathematical Monthly*, 125(7):593–601, 2018.

- [Gri99] Geoffrey Grimmett. *Percolation*, volume 321 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, second edition, 1999.
- [HS71] Fritz Herzog and Bonnie Stewart. Patterns of visible and nonvisible lattice points. *The American Mathematical Monthly*, 78(5):487–496, 1971.
- [HW79] Godfrey Hardy and Edward Wright. *An introduction to the theory of numbers*. Oxford University Press, 1979.
- [Lin01] Elon Lindenstrauss. Pointwise theorems for amenable groups. *Inventiones mathematicae*, 146(2):259–295, 2001.
- [LP16] Russell Lyons and Yuval Peres. *Probability on trees and networks*, volume 42. Cambridge University Press, 2016.
- [Mir47] Leonid Mirsky. Note on an asymptotic formula connected with r -free integers. *The Quarterly Journal of Mathematics*, (1):178–182, 1947.
- [Mir48] Leonid Mirsky. Arithmetical pattern problems relating to divisibility by r th powers. *Proceedings of the London Mathematical Society*, 2(1):497–508, 1948.
- [PH13] Peter Pleasants and Christian Huck. Entropy and diffraction of the k -free points in n -dimensional lattices. *Discrete & Computational Geometry*, 50(1):39–68, 2013.
- [Syl83] James Sylvester. Sur le nombre de fractions ordinaires inégales qu'on peut exprimer en se servant de chiffres qui n'excèdent pas un nombre donné. *CR Acad. Sci. Paris*, 96:409–413, 1883.
- [Var99] Ilan Vardi. Deterministic percolation. *Communications in mathematical physics*, 207(1):43–66, 1999.