

Homogenization of Nonlocal Partial Differential Equations Related to Stochastic Differential Equations with Lévy Noise

Qiao Huang*, Jinqiao Duan[†], Renming Song[‡]

Abstract

We study the “periodic homogenization” for a class of nonlocal partial differential equations of parabolic-type with rapidly oscillating coefficients, related to stochastic differential equations driven by multiplicative isotropic α -stable Lévy noise ($1 < \alpha < 2$) which is nonlinear in the noise component. Our homogenization method is probabilistic. It turns out that, under suitable regularity assumptions, the limit of the solutions satisfies a nonlocal partial differential equation with constant coefficients, which are associated to a symmetric α -stable Lévy process.

AMS 2010 Mathematics Subject Classification: 60H30, 60H10, 35B27, 35R09.

Keywords and Phrases: Homogenization, nonlocal parabolic PDEs, SDEs with jumps, Zvonkin’s transform, strong well-posedness, Feller semigroups, Feynman-Kac formula.

1 Introduction

In the study of porous media, composite materials and other physical and engineering systems, one is led to the initial or boundary value problems with periodic structures (see, e.g., [1, 12, 30]). The process of passing from a microscopic description to a macroscopic description of the behaviors of such systems is called *homogenization*. At present, numerous publications can be found on the mathematical aspects of the homogenization theory (see [4, 7, 27]).

The goal of this paper is to use a *probabilistic approach* to study the limit behavior, as $\epsilon \rightarrow 0$, of the solution $u^\epsilon : \mathbb{R}^d \rightarrow \mathbb{R}$ of the following *nonlocal* partial differential equation (PDE) of parabolic-type with rapidly oscillating periodic and singular coefficients,

$$\begin{cases} \frac{\partial u^\epsilon}{\partial t}(t, x) = \mathcal{L}_\epsilon^\alpha u^\epsilon(t, x) + \left(\frac{1}{\epsilon^{\alpha-1}} e\left(\frac{x}{\epsilon}\right) + g\left(\frac{x}{\epsilon}\right)\right) u^\epsilon(t, x), & t > 0, x \in \mathbb{R}^d, \\ u^\epsilon(0, x) = u_0(x), & x \in \mathbb{R}^d, \end{cases} \quad (1.1)$$

where $1 < \alpha < 2$ and the linear operator $\mathcal{L}_\epsilon^\alpha$ is a nonlocal integro-differential operator of Lévy-type given by

$$\begin{aligned} \mathcal{L}_\epsilon^\alpha f(x) := & \int_{\mathbb{R}^d \setminus \{0\}} \left[f\left(x + \sigma\left(\frac{x}{\epsilon}, y\right)\right) - f(x) - \sigma^i\left(\frac{x}{\epsilon}, y\right) \partial_i f(x) \mathbf{1}_B(y) \right] \nu^\alpha(dy) \\ & + \left[\frac{1}{\epsilon^{\alpha-1}} b^i\left(\frac{x}{\epsilon}\right) + c^i\left(\frac{x}{\epsilon}\right) \right] \partial_i f(x), \quad x \in \mathbb{R}^d. \end{aligned}$$

Here B is the unit open ball in $\mathbb{R}^d$ centering at the origin, and $\nu^\alpha(dy) := \frac{dy}{|y|^{d+\alpha}}$ is the isotropy α -stable Lévy measure. In this paper, we use Einstein’s convention that the repeated indices in a product will be summed automatically.

*Center for Mathematical Sciences, Huazhong University of Science and Technology, Wuhan, Hubei 430074, P.R. China. Email: hq932309@alumni.hust.edu.cn

[†]Department of Applied Mathematics, Illinois Institute of Technology, Chicago, IL 60616, USA. Email: duan@iit.edu

[‡]Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA. Email: rsong@illinois.edu

For notational simplicity, we introduce the linear operator $\mathcal{A}^{\sigma, \alpha}$ defined by

$$\mathcal{A}^{\sigma, \nu^\alpha} f(x) := \int_{\mathbb{R}^d \setminus \{0\}} [f(x + \sigma(x, y)) - f(x) - \sigma^i(x, y) \partial_i f(x) \mathbf{1}_B(y)] \nu^\alpha(dy), \quad x \in \mathbb{R}^d. \quad (1.2)$$

For a function f on $\mathbb{R}^d$ (or F on $\mathbb{R}^d \times \mathbb{R}^d$), we denote $f_\epsilon(x) := f\left(\frac{x}{\epsilon}\right)$ (or $F_\epsilon(x, y) := F\left(\frac{x}{\epsilon}, y\right)$). Then

$$\mathcal{L}_\epsilon^\alpha = \mathcal{A}^{\sigma_\epsilon, \nu^\alpha} + \left(\frac{1}{\epsilon^{\alpha-1}} b_\epsilon + c_\epsilon\right) \cdot \nabla. \quad (1.3)$$

The main result of this paper is the following theorem. Assumptions [H1–H6](#) are made for the coefficients and will be listed in the sequel. We will prove it at the end of Section [5](#).

Theorem 1.1. *Under Assumptions [H1–H6](#), the nonlocal PDE [\(1.1\)](#) has a unique mild solution u^ϵ for each $\epsilon > 0$. Moreover, for each $t \geq 0, x \in \mathbb{R}^d$,*

$$u^\epsilon(t, x) \rightarrow u(t, x), \quad \epsilon \rightarrow 0, \quad (1.4)$$

where u satisfies the limit nonlocal PDE,

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) = \mathcal{A}^{Id_y, \bar{\nu}} u(t, x) + \bar{C} \cdot \nabla u(t, x) + \bar{E} u(t, x), & t > 0, x \in \mathbb{R}^d, \\ u(0, x) = u_0(x), & x \in \mathbb{R}^d, \end{cases}$$

where

$$\mathcal{A}^{Id_y, \bar{\nu}} f(x) := \int_{\mathbb{R}^d \setminus \{0\}} [f(x + y) - f(x) - y^i \partial_i f(x) \mathbf{1}_B(y)] \bar{\nu}(dy),$$

and the constant coefficients $\bar{C}, \bar{E}$ and measure $\bar{\nu}$ are given by [\(5.2\)](#), [\(5.4\)](#) and [\(5.3\)](#) respectively. The solution is given by

$$u(t, x) = \mathbf{E}[u_0(x + \bar{C}t + \bar{L}_t)] e^{\bar{E}t},$$

where $\{\bar{L}_t\}_{t \geq 0}$ is a symmetric α -stable Lévy processes with jump intensity measure $\bar{\nu}$.

The original probabilistic approach to the homogenization of *local* linear second order parabolic partial differential operators is presented in [\[9, Chapter 3\]](#), which is based on the ergodic theorem, the Feynman-Kac formula and the functional central limit theorem. By now, there are lots of literature concerning the homogenization of second order local PDEs, i.e., the case of replacing the operator $\mathcal{A}^{\sigma_\epsilon, \nu^\alpha}$ in [\(1.3\)](#) by a second order partial differential operator with singular coefficients. Two different scales of spatial variables involved in the coefficients have been considered in [\[8\]](#), by using the nonlinear Feynman-Kac formula in the context of backward stochastic differential equations (SDEs). In [\[32\]](#), the authors allowed the singular coefficients to be time-dependent and rapidly oscillating in time with a different scale in contrast to the spatial variable. The paper [\[18\]](#) dealt with the case when the second order coefficient matrix can be degenerate, using the existence of a spectral gap and Malliavin's calculus.

There are also some literature for the homogenization of nonlocal PDEs or SDEs with jumps involved. Firstly, let us discuss several articles studying the periodic homogenization of some kinds of nonlocal operators in the *analytical* aspects. It was shown in [\[35\]](#) that a homogenization model for a class of nonlocal operators with convolution-type kernels can generate a diffusion in the limit. The jump kernels are forced to have finite second moment there, which excludes the stable Lévy densities. Later on, [\[24\]](#) studied the homogenization of Lévy-type operators (or stable-like operators called by other authors) with both symmetric and non-symmetric kernels, the limit is a Lévy-operator. Besides, an approach based on the viscosity solution of PDEs can be found

in [3], the author focused on the Lévy-operator perturbed linearly by a zeroth-order term. This method has also been extended to several classes of nonlinear problems; see, e.g., [42]. It is worth to mention that all the homogenization problems considered in [24], [3] and [42] can cover the Lévy-type operators (or Lévy-operators) with full range of $\alpha \in (0, 2)$, while none of them involves the drift (gradient) parts.

The *probabilistic* study of homogenization of periodic stable-like processes in pure jump or jump-diffusion case can be found in [15, 40]. The homogenization in random medium is slightly different from the periodic case, referring to [38] for related results for jump-diffusion processes in random medium.

The homogenization of a kind of one-dimensional pure jump Markov processes with the following form of generators has been investigated in the paper [19],

$$\mathcal{A}_\epsilon f(x) = \int_{\mathbb{R} \setminus \{0\}} [f(x+z) - f(x) - zf'(x)] a\left(\frac{x}{\epsilon}, \frac{z}{\epsilon}\right) \frac{1}{|z|^{1+\alpha}} dz + \frac{1}{\epsilon^{\alpha-1}} b\left(\frac{x}{\epsilon}\right) f'(x).$$

See [43] for a generalization for multi-dimensional case and with diffusion terms involved. In their context, the jump function h is oscillating both in the spatial variable x and in the noise variable y , and they are in the same scale. This means the noise comes from the underlying space of periodic medium. But when we do homogenization for systems with fluctuations, the noise usually comes from the external environment, so that the jump function is no longer oscillating in the noise variable, and there happens to be two different scales. As we will see in Section 2, the change of variables allows us to write

$$\begin{aligned} \mathcal{L}_\epsilon^\alpha f(x) := & \int_{\mathbb{R}^d \setminus \{0\}} [f(x+z) - f(x) - z^i \partial_i f(x) \mathbf{1}_B(y)] h\left(\frac{x}{\epsilon}, z\right) \frac{1}{|z|^{d+\alpha}} dz \\ & + \left[\frac{1}{\epsilon^{\alpha-1}} b^i\left(\frac{x}{\epsilon}\right) + c^i\left(\frac{x}{\epsilon}\right) \right] \partial_i f(x), \end{aligned}$$

for some function h . Note that the main difference is that we do not involve oscillations for h in its noise variable, while [19, 43] involve. Meanwhile, the coefficients of drift and the zeroth order term b, c, e, g has two different scales.

In paper [16], the author considered the homogenization of SDEs driven by multiplicative stable processes, where the noise intensity coefficient σ is linear in the noise variable in the sense that $\sigma(x, y) = \sigma_0(x)y$, with σ_0 three-times continuously differentiable. In the present paper, we generalize his results to the general multiplicative case. That is, the intensity function σ need *not to be linear* for the noise component. This is also more realistic in applications (see, e.g., [39, 10]). For some typical forms of σ that are nonlinear in y , see Example 2.3. In addition, the coefficients only need to possess some Hölder or Lipschitz continuity in our context (see Remark 2.2 for the comparison of the regularity assumptions for σ). This will give rise to several difficulties both in analytic and probabilistic aspects. We also use the homogenization results of SDEs to study the homogenization of the nonlocal PDEs with singular coefficients involved, by utilizing Feynman-Kac formula. The trick we use to remove the singular drift in (1.1) is now known as *Zvonkin's transform*, which appeared originally in [47].

The work in this paper is highly motivated by these two considerations. That is, the noise in our homogenization problem comes from the external environment instead of the underlying periodic medium, and is not necessarily linear. Both are more suitable from the practical point of view than in earlier papers. Under these considerations, we further weaken the regularity assumptions for all coefficients in a compatible way.

Last but not least, the present paper will only focus on the *subcritical* case $1 < \alpha < 2$. There are both probabilistic and analytical difficulties for the *supercritical* case $0 < \alpha \leq 1$. In analytical

aspect, the well-posedness of the nonlocal Poisson equations (Section 3) in the supercritical case is still open in general setting. Intuitively, the nonlocal part has lower order than the drift part in this case, so that one cannot regard the drift as a perturbation of the nonlocal operator (in the sense of C_0 -semigroups, as ideally the perturbing operator needs to be “bounded” with respect to the principal operator, cf. [13, Section III.2]). This will give rise to some analytical difficulties, of which the most crucial one is the heat kernel estimates for the coupling of supercritical nonlocal operators with gradients (cf. [11] for the subcritical case). Although there are a few published results for these topics, they all focus on some special case such as the drifted critical ($\alpha = 1$) fractional Laplacian ([45]). Moreover, the condition $\alpha > 1$ is also crucial when doing the probabilistic homogenization, since we will have used this condition explicitly in the proofs of Lemma 4.7 and Lemma 5.3.

We denote by $\mathcal{C}^k$ ($\mathcal{C}_b^k$) with integer $k \geq 0$ the space of (bounded) continuous functions possessing (bounded) derivatives of orders not greater than k . We shall explicitly write out the domain if necessary. Denote by $\mathcal{C}_b(\mathbb{R}^d) := \mathcal{C}_b^0(\mathbb{R}^d)$, it is a Banach space with the supremum norm $\|f\|_0 = \sup_{x \in \mathbb{R}^d} |f(x)|$. The space $\mathcal{C}_b^k(\mathbb{R}^d)$ is a Banach space endowed with the norm $\|f\|_k = \|f\|_0 + \sum_{j=1}^k \|\nabla^{\otimes j} f\|$. We also denote by $\mathcal{C}^{\text{Lip}}$ the class of all Lipschitz continuous functions. For a non-integer $\gamma > 0$, the Hölder spaces $\mathcal{C}^\gamma$ ($\mathcal{C}_b^\gamma$) are defined as the subspaces of $\mathcal{C}^{[\gamma]}$ ($\mathcal{C}_b^{[\gamma]}$) consisting of functions whose $[\gamma]$ -th order partial derivatives are locally Hölder continuous (uniformly Hölder continuous) with exponent $\gamma - [\gamma]$. These two spaces $\mathcal{C}^\gamma$ and $\mathcal{C}_b^\gamma$ obviously coincide when the underlying domain is compact. The space $\mathcal{C}_b^\gamma(\mathbb{R}^d)$ is a Banach space endowed with the norm $\|f\|_\gamma = \|f\|_{[\gamma]} + [\nabla^{[\gamma]} f]_{\gamma - [\gamma]}$, where the seminorm $[\cdot]_{\gamma'}$ with $0 < \gamma' < 1$ is defined as $[f]_{\gamma'} := \sup_{x, y \in \mathbb{R}^d, x \neq y} \frac{|f(x) - f(y)|}{|x - y|^{\gamma'}}$ (this seminorm can also be defined for the case $\gamma' = 1$, which is exactly the Lipschitz seminorm). In the sequel, the torus $\mathbb{T}^d := \mathbb{R}^d / \mathbb{Z}^d$ will be used frequently. Denote by $\mathcal{D} := \mathcal{D}(\mathbb{R}_+; \mathbb{T}^d)$ the space of all $\mathbb{T}^d$ -valued càdlàg functions on $\mathbb{R}_+$, equipped with the Skorokhod topology. We shall always identify the periodic function on $\mathbb{R}^d$ of period 1 with its restriction on the torus $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$. This allows us to regard the space $\mathcal{C}^k(\mathbb{T}^d)$ ($\mathcal{C}^\gamma(\mathbb{T}^d)$) as a sub-Banach space of $\mathcal{C}_b^k(\mathbb{R}^d)$ ($\mathcal{C}_b^\gamma(\mathbb{R}^d)$).

By B_r we mean the open ball in $\mathbb{R}^d$ centering at the origin with radius $r > 0$, we shall omit the subscript when the radius is one. The capital letter C denotes a finite positive constant whose value may vary from line to line. We also use the notation $C(\cdots)$ to emphasize the dependence on the quantities appearing in the parentheses.

The remainder of the paper is organized as follows. In Section 2, we present some general assumptions and preliminary results. In Section 3, we study the well-posedness of the nonlocal Poisson equation and the Feller properties of the semigroup associated with generator of the form (1.3). Section 4 is devoted to the strong well-posedness and exponential ergodicity of the Lévy driven SDE. As a consequence, we obtained the Feynman-Kac representation for the nonlocal PDE (1.1). Lastly, Section 5 contains the homogenization results of SDEs and nonlocal PDEs, utilizing the ergodicity and the Feynman-Kac representation.

2 Preliminaries and general assumptions

Let $(\Omega, \mathcal{F}, \mathbf{P}, \{\mathcal{F}_t\}_{t \geq 0})$ be a filtered probability space endowed with a Poisson random measure N^α on $(\mathbb{R}^d \setminus \{0\}) \times \mathbb{R}_+$ with jump intensity measure $\nu^\alpha(dy) = \frac{dy}{|y|^{d+\alpha}}$, where $1 < \alpha < 2$. Denote by $\tilde{N}$ the associated compensated Poisson random measure, that is, $\tilde{N}^\alpha(dy, ds) := N^\alpha(dy, ds) - \nu^\alpha(dy)ds$. We assume that the filtration $\{\mathcal{F}_t\}_{t \geq 0}$ satisfies the usual conditions. Let $L^\alpha = \{L_t^\alpha\}_{t \geq 0}$ be a d -

dimensional isotropic α -stable Lévy process given by

$$L_t^\alpha = \int_0^t \int_{B \setminus \{0\}} y \tilde{N}^\alpha(dy, ds) + \int_0^t \int_{B^c} y N^\alpha(dy, ds).$$

Given $\epsilon > 0, x \in \mathbb{R}^d$, consider the following:

$$dX_t^{x,\epsilon} = \left(\frac{1}{\epsilon^{\alpha-1}} b\left(\frac{X_t^{x,\epsilon}}{\epsilon}\right) + c\left(\frac{X_t^{x,\epsilon}}{\epsilon}\right) \right) dt + \sigma\left(\frac{X_t^{x,\epsilon}}{\epsilon}, dL_t^\alpha\right), \quad X_0^{x,\epsilon} = x, \quad (2.1)$$

or more precisely,

$$\begin{aligned} X_t^{x,\epsilon} &= x + \int_0^t \left(\frac{1}{\epsilon^{\alpha-1}} b\left(\frac{X_s^{x,\epsilon}}{\epsilon}\right) + c\left(\frac{X_s^{x,\epsilon}}{\epsilon}\right) \right) ds \\ &\quad + \int_0^t \int_{B \setminus \{0\}} \sigma\left(\frac{X_s^{x,\epsilon}}{\epsilon}, y\right) \tilde{N}^\alpha(dy, ds) + \int_0^t \int_{B^c} \sigma\left(\frac{X_s^{x,\epsilon}}{\epsilon}, y\right) N^\alpha(dy, ds), \end{aligned}$$

where the coefficients $b, c, \sigma(\cdot, y)$ are periodic, for each $y \in \mathbb{R}^d$, of periodic one in each component. The shorthand notation for the stochastic differential term in (2.1) is due to [28].

Define $\tilde{X}_t^{x,\epsilon} := \frac{1}{\epsilon} X_{\epsilon t}^{x,\epsilon}$. It is easy to check that

$$d\tilde{X}_t^{x,\epsilon} = \left(b(\tilde{X}_t^{x,\epsilon}) + \epsilon^{\alpha-1} c(\tilde{X}_t^{x,\epsilon}) \right) dt + \frac{1}{\epsilon} \sigma\left(\tilde{X}_t^{x,\epsilon}, \epsilon d\tilde{L}_t^\alpha\right), \quad \tilde{X}_0^{x,\epsilon} = \frac{x}{\epsilon}, \quad (2.2)$$

where $\{\tilde{L}_t^\alpha\} := \{\frac{1}{\epsilon} L_{\epsilon t}^\alpha\} \stackrel{d}{=} \{L_t^\alpha\}$ by virtue of the selfsimilarity. We shall also consider the “limit” equation, namely

$$d\tilde{X}_t^x = b(\tilde{X}_t^x) dt + \bar{\sigma}\left(\tilde{X}_t^x, d\tilde{L}_t^\alpha\right), \quad \tilde{X}_0^x = x, \quad (2.3)$$

where, intuitively, $\bar{\sigma}$ is the limit of $\frac{1}{\epsilon} \sigma(\cdot, \epsilon \cdot)$. We will make the relation between σ and $\bar{\sigma}$ clear in the forthcoming assumption H5. For notational simplicity, we shall allow the parameter ϵ to be zero in $\tilde{X}^{x,\epsilon}$ to include $\tilde{X}^x$, i.e., $\tilde{X}^{x,0} := \tilde{X}^x$.

In the sequel, we will regard the solutions $\tilde{X}^{x,\epsilon}, \tilde{X}^x$ of (2.2) and (2.3) as $\mathbb{T}^d$ -valued processes, by mapping all trajectories of the processes on $\mathbb{R}^d$ to the torus $\mathbb{T}^d$, via the canonical quotient map $\pi : \mathbb{R}^d \rightarrow \mathbb{R}^d / \mathbb{Z}^d$. Then the periodicity of the coefficients implies that $\tilde{X}^{x,\epsilon}$ and $\tilde{X}$ are well-defined stochastic processes on $\mathbb{T}^d$ (cf. [9, Section 3.3.2]).

Now we list some general assumptions for the nonlocal PDE (1.1) and the SDE (2.1). All these assumptions are assumed to hold in the sequel unless otherwise specified.

Assumption H1. The functions b, c, e, g, u_0 are all periodic of period 1 in each component. For every $y \in \mathbb{R}^d$, the function $x \rightarrow \sigma(x, y)$ is periodic of period 1 in each component.

Assumption H2. The functions b, c, e are of class $\mathcal{C}_b^\beta$ with exponent β satisfying

$$1 - \frac{\alpha}{2} < \beta < 1.$$

The functions g and u_0 are both continuous.

Assumption H3. The function $\sigma : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ satisfies the following conditions.

(1). *Regularity.* For every $x \in \mathbb{R}^d$, the function $y \rightarrow \sigma(x, y)$ is of class $\mathcal{C}^2$. There exists $\alpha - 1 < \lambda \leq 1$ such that

$$\sup_{x \in \mathbb{R}^d} [\nabla_y \sigma(x, \cdot)]_\lambda < \infty. \quad (2.4)$$

There exists a constant $C > 0$, such that for any $x_1, x_2, y \in \mathbb{R}^d$,

$$|\sigma(x_1, y) - \sigma(x_2, y)| \leq C|x_1 - x_2||y|.$$

(2). *Oddness.* For all $x, y \in \mathbb{R}^d$, $\sigma(x, -y) = -\sigma(x, y)$.

(3). *Bounded inverse Jacobian.* The Jacobian matrix with respect to the second variable $\nabla_y \sigma(x, y)$ is non-degenerate for all $x, y \in \mathbb{R}^d$, and there exists a constant $C > 0$ such that $|(\nabla_y \sigma(x, y))^{-1}| \leq C$ for all $x, y \in \mathbb{R}^d$, where $|\cdot|$ is the operator norm on $\mathcal{L}(\mathbb{R}^d, \mathbb{R}^d)$.

(4). *Growth condition.* There exists a positive bounded measurable function $\phi : \mathbb{R}^d \rightarrow \mathbb{R}_+$, such that for all $x, y \in \mathbb{R}^d$,

$$\phi(x)^{-1}|y| \leq |\sigma(x, y)| \leq \phi(x)|y|.$$

Remark 2.1. Some comments on our assumptions will be helpful:

(1). As mentioned in the end of the introduction, b, c, e, g, u_0 and the function $x \rightarrow \sigma(x, y)$, for every $y \in \mathbb{R}^d$, can be regarded as functions on $\mathbb{T}^d$. Hence we have $b, c, e \in \mathcal{C}^\beta(\mathbb{T}^d)$, $g, u_0 \in \mathcal{C}(\mathbb{T}^d)$, under Assumptions **H1** and **H2**. The condition (2.4) reduce to

$$\sup_{x \in \mathbb{T}^d} [\nabla_y \sigma(x, \cdot)]_\lambda < \infty.$$

(2). In Assumption **H3**, the condition (2.4) should not be confused with the condition that $y \rightarrow \sigma(x, y)$ is of class $\mathcal{C}^2$. The former requires the uniform Hölder continuity of σ in y , as indicated in the definition of the Hölder seminorm $[\cdot]_\lambda$, while the latter do not force the uniformity. Therefore, none of them can imply the other.

(3). Both the oddness and the growth condition in Assumption **H3** imply that $\sigma(\cdot, 0) \equiv 0$.

(4). The bounded inverse Jacobian condition implies that $|\nabla_y \sigma| \geq C^{-1}$. Since by Hadamard's inequality (see, for instance, [41]),

$$|(\nabla_y \sigma)^{-1}| \leq C \Rightarrow |\det((\nabla_y \sigma)^{-1})| \leq C^d \Leftrightarrow |\det(\nabla_y \sigma)| \geq C^{-d} \Rightarrow |\nabla_y \sigma| \geq C^{-1}. \quad (2.5)$$

(5). The growth condition implies that for any $\gamma > \alpha$, we have

$$\sup_{x \in \mathbb{R}^d} \int_{B \setminus \{0\}} |\sigma(x, y)|^\gamma \nu^\alpha(dy) < \infty. \quad (2.6)$$

This ensures that we can apply Itô's formula to $f(\tilde{X}_t^x)$ (or $f(\tilde{X}_t^{x, \epsilon})$, $f(X_t^{x, \epsilon})$), for any $f \in \mathcal{C}_b^\gamma(\mathbb{R}^d)$ with $\gamma > \alpha$ (cf. [36, Lemma 4.2]).

(6). By virtue of the oddness condition in Assumption **H3** and the symmetry of the jump intensity measure ν^α , for any $x \in \mathbb{R}^d$,

$$\text{P.V.} \int_{\sigma(x, \cdot)^{-1}B \setminus B} \sigma^i(x, y) \nu^\alpha(dy) = \text{P.V.} \int_{B \setminus \sigma(x, \cdot)^{-1}B} \sigma^i(x, y) \nu^\alpha(dy) = 0. \quad (2.7)$$

Consequently we can rewrite the operator $\mathcal{A}^{\sigma, \nu^\alpha}$ in (1.2) as

$$\mathcal{A}^{\sigma, \nu^\alpha} f(x) = \int_{\mathbb{R}^d \setminus \{0\}} [f(x+z) - f(x) - z^i \partial_i f(x) \mathbf{1}_B(z)] \nu^{\sigma, \alpha}(x, dz), \quad (2.8)$$

where the kernel $\{\nu^{\sigma, \alpha}(x, \cdot) | x \in \mathbb{R}^d\}$ is given by

$$\nu^{\sigma, \alpha}(x, A) := \int_{\mathbb{R}^d \setminus \{0\}} \mathbf{1}_A(\sigma(x, y)) \nu^\alpha(dy), \quad A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}). \quad (2.9)$$

Moreover, for any $\gamma > \alpha$, the growth condition in Assumption **H3** implies that

$$\begin{aligned}
& \sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d \setminus \{0\}} (|z|^\gamma \wedge 1) \nu^{\sigma, \alpha}(x, dz) = \sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d \setminus \{0\}} (|\sigma(x, y)|^\gamma \wedge 1) \nu^\alpha(dy) \\
& \leq \sup_{x \in \mathbb{R}^d} \left(\int_{|y| \leq \phi(x)} (\phi(x)|y|)^\gamma \nu^\alpha(dy) + \int_{|y| \geq \phi(x)^{-1}} \nu^\alpha(dy) \right) \\
& \leq \frac{1}{\gamma - \alpha} \|\phi\|_{L^\infty}^{2\gamma - \alpha} + \frac{1}{\alpha} \|\phi\|_{L^\infty}^\alpha < \infty.
\end{aligned} \tag{2.10}$$

Remark 2.2. The special case $\sigma(x, y) = \sigma_0(x)y$ with certain σ_0 is considered in [16], where the author assumed the function $\sigma_0 : \mathbb{R}^d \rightarrow \text{GL}(\mathbb{R}^d)$ is periodic and of class $\mathcal{C}^3$. In our context, Assumption **H1** and **H3** amounts to saying that

$$\sigma_0 : \mathbb{R}^d \rightarrow \text{GL}(\mathbb{R}^d) \text{ is periodic and Lipschitz.} \tag{2.11}$$

Since these imply the regularity condition immediately, the bounded inverse Jacobian and growth conditions are fulfilled by continuity and periodicity, together with the observation $\sup_{x \in \mathbb{R}^d} \|\sigma_0(x)\| \vee \|\sigma_0(x)^{-1}\| < \infty$. The oddness condition is trivial in this case.

In practice, the noise is not always linear. Here we give some nontrivial examples for σ , that is, nonlinear in y .

Example 2.3. Suppose σ_0 to satisfy (2.11).

(i). The dependence of the noise is a small perturbation of the linear case, namely, $\sigma(x, y) = \sigma_0(x)y + \delta(x, y)$, where the function δ satisfies the same properties as σ in Assumptions **H1** and (i), (ii) in **H3**, but has much smaller scale than $\sigma_0(x)y$, so that Assumptions **H1-H3** and the forthcoming Assumption **H4** do hold for σ .

(ii). Another case is that the function σ is separable but not linear in y . To be precise, let $\eta : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be an odd function of class $\mathcal{C}^2$, satisfying that $\nabla \eta$ is *uniformly* Hölder continuous with exponent $\lambda > \alpha - 1$ and $\nabla \eta(y)$ is non-degenerate for all $y \in \mathbb{R}^d$, and there exist some constants $C_1, C_2 > 0$, such that $|\nabla \eta| \geq C_1$ and $C_2^{-1}|y| \leq |\eta(y)| \leq C_2|y|$. Now let $\sigma(x, y) = \sigma_0(x)\eta(y)$. Then σ satisfies Assumption **H3**.

(iii). Combining the above two example together, one can obtain a more general example, that is, $\sigma(x, y) = \sigma_0(x)\eta(y) + \delta(x, y)$.

We will need some regularities for the 'partial' inverse of σ . For a function $F : \mathbb{R}^d \times \mathbb{R}^d \ni (x, y) \rightarrow F(x, y) \in \mathbb{R}$, we say $F \in L_2^\infty(\mathbb{R}^d; \mathcal{C}_1^{\text{Lip}}(\mathbb{R}^d; \mathbb{R}^d))$, if there exists a constant $C > 0$ such that for all $x, y \in \mathbb{R}^d$, $|F(x, y)| \leq C$, and for all $x_1, x_2, y \in \mathbb{R}^d$, $|F(x_1, y) - F(x_2, y)| \leq C|x_1 - x_2|$. Then the regularity and growth conditions in Assumption **H3** imply that the function $(x, y) \rightarrow \sigma(x, y)/|y|$ is of class $L_2^\infty(\mathbb{R}^d; \mathcal{C}_1^{\text{Lip}}(\mathbb{R}^d; \mathbb{R}^d))$.

Lemma 2.4. *Under Assumption **H3**, for every $x \in \mathbb{R}^d$, the function $y \rightarrow \sigma(x, y)$ is a $\mathcal{C}^2$ -diffeomorphism. Denote the inverse by $\tau(x, z) := \sigma(x, \cdot)^{-1}(z)$, then for every $z \in \mathbb{R}^d$, the function $x \rightarrow \tau(x, z)$ is periodic of period one in each component. Moreover, the function $(x, z) \rightarrow \tau(x, z)/|z|$ is of class $L_2^\infty(\mathbb{R}^d; \mathcal{C}_1^{\text{Lip}}(\mathbb{R}^d; \mathbb{R}^d))$.*

Proof. Fix $x \in \mathbb{R}^d$. Since the function $y \rightarrow \sigma(x, y)$ is of class $\mathcal{C}^2$, by the bounded inverse Jacobian condition in Assumption **H3**, together with Hadamard's global inverse function theorem (see [25, Theorem 6.2.4]), $\sigma(x, \cdot)$ is a $\mathcal{C}^2$ -diffeomorphism. The periodicity is obvious.

Now using the bounded inverse Jacobian condition, the Jacobian matrix of $\tau(x, z)$ with respect to z satisfies $|\nabla_z \tau(x, z)| \leq C$, for all $x, z \in \mathbb{R}^d$. Then by the growth condition and regularity condition, the second assertion follows from the following derivation,

$$\begin{aligned} & \sup_z \frac{|\tau(x, z)|}{|z|} \leq \phi(x), \\ & \sup_z \frac{|\tau(x_1, z) - \tau(x_2, z)|}{|z|} = \sup_y \frac{|\tau(x_1, \sigma(x_1, y)) - \tau(x_2, \sigma(x_1, y))|}{|\sigma(x_1, y)|} \\ & = \sup_y \frac{|\tau(x_2, \sigma(x_2, y)) - \tau(x_2, \sigma(x_1, y))|}{|\sigma(x_1, y)|} \leq \|\phi\|_{L^\infty} \|\nabla_z \tau\|_{L^\infty} \sup_y \frac{|\sigma(x_2, y) - \sigma(x_1, y)|}{|y|} \\ & \leq C \|\phi\|_{L^\infty} \|\nabla_z \tau\|_{L^\infty} |x_1 - x_2|. \end{aligned}$$

□

Assumption H4. $\det(\nabla_z \tau) \in L_2^\infty(\mathbb{R}^d; \mathcal{C}_1^{\text{Lip}}(\mathbb{R}^d; \mathbb{R}))$.

This assumption is rather mild, as shown in the following remark.

Example 2.5. (i). In the case $\sigma(x, y) = \sigma_0(x)y$, the partial inverse is $\tau(x, z) = \sigma_0(x)^{-1}y$, the Jacobian of $\tau(x, z)$ with respect to z is $\nabla_z \tau(x, z) \equiv \sigma_0(x)^{-1}$. Then Assumption H4 reduces to that the function $\det(\sigma_0)^{-1} : \mathbb{R}^d \rightarrow \mathbb{R}$ is Lipschitz, while this holds automatically since it is a direct consequence of (2.11).

(ii). When $\sigma(x, y) = \sigma_0(x)\eta(y)$ as in Example 2.3.(ii), $\tau(x, z) = \eta^{-1}(\sigma_0(x)^{-1}z)$. Then it is easy to deduce that Assumption H4 is implied by the bounded inverse Jacobian condition in Assumption H3, using a similar argument as (2.5). □

If we let

$$h(x, z) = |\det \nabla_z \tau(x, z)| \frac{|z|^{d+\alpha}}{|\tau(x, z)|^{d+\alpha}}, \quad (2.12)$$

then by (2.9), $\nu^{\sigma, \alpha}(x, dz) = h(x, z) \frac{dz}{|z|^{d+\alpha}}$. Using the growth condition, we also find that for all $x, z \in \mathbb{R}^d$,

$$\|\phi\|_{L^\infty}^{-1} \leq \frac{|\tau(x, z)|}{|z|} \leq \|\phi\|_{L^\infty}. \quad (2.13)$$

Combining (2.12), (2.13), Lemma 2.4 and Assumption H4, together with the fact that if $f, g \in \mathcal{C}^\gamma$ and $\inf |g| > 0$, then $f/g \in \mathcal{C}^\gamma$, we conclude that

Lemma 2.6. *Under Assumptions H3 and H4, $h \in L_2^\infty(\mathbb{R}^d; \mathcal{C}_1^{\text{Lip}}(\mathbb{R}^d; \mathbb{R}^d))$, namely, there exists a constant $h_0 > 0$ such that $|h(x_1, z) - h(x_2, z)| \leq h_0 |x_1 - x_2|$ for all $x_1, x_2, z \in \mathbb{R}^d$. Moreover, there also exists a constant $h_1 > 1$ such that $h_1^{-1} \leq h(x, z) \leq h_1$ for all $x, z \in \mathbb{R}^d$.*

In particular, the kernel $\nu^{\sigma, \alpha}$ is comparable to the jump intensity measure of an isotropic α -stable process.

Remark 2.7. Thanks to Lemma 2.6, the general assumptions in [6, 11] are satisfied. Thus, the regularity results and heat kernel estimates therein are available in our context. Actually, these two papers only need that $h \in L_2^\infty(\mathbb{R}^d; \mathcal{C}_1^\gamma(\mathbb{R}^d; \mathbb{R}^d))$ for some $0 < \gamma < 1$, this is the case by virtue of the natural embedding $\mathcal{C}^{\text{Lip}} \subset \mathcal{C}^\gamma$. Note that [6] also needs $\alpha + \beta$ not to be an integer, this can be fulfilled by choosing an appropriate β .

Now we clarify that the function $\bar{\sigma}$ appeared in the “limit” equation (2.3) is the scaling limit of σ . Firstly, we use the L’Hôpital’s rule to proceed the limit,

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \sigma(x, \epsilon y) = \lim_{\epsilon \rightarrow 0} (\nabla_y \sigma(x, \epsilon y)) y = (\nabla_y \sigma(x, 0)) y,$$

by the continuity of $\nabla_y \sigma(x, y)$ with respect to y . If we set

$$\bar{\sigma}_0(x) := \nabla_y \sigma(x, 0), \quad (2.14)$$

$$\bar{\sigma}(x, y) := \bar{\sigma}_0(x) y, \quad (2.15)$$

then $\bar{\sigma}$ is the point-wise limit of $\frac{1}{\epsilon} \sigma(\cdot, \epsilon \cdot)$ as $\epsilon \rightarrow 0$. However, as we will see in Section 4 (more precisely, Lemma 4.7), the point-wise convergence is not enough when we deal with the ergodicity. We need a stronger convergence as follows:

Assumption H5. For every $y \in \mathbb{R}^d$, $\frac{1}{\epsilon} \sigma(x, \epsilon y) \rightarrow (\nabla_y \sigma(x, 0)) y$ uniformly in $x \in \mathbb{R}^d$, as $\epsilon \rightarrow 0$.

We can say more for the scaling limit $\bar{\sigma}$.

Remark 2.8. The function $\bar{\sigma}_0 : \mathbb{R}^d \rightarrow \text{GL}(\mathbb{R}^d)$ defined in (2.14) is periodic and Lipschitz. Indeed, by a straightforward limit argument, one can easily verify that Assumptions H1 and H3 holds for $\bar{\sigma}$. This implies that $\bar{\sigma}_0$ has the same property as σ_0 in Remark 2.2. Moreover, if we denote the partial inverse of $\bar{\sigma}$ with respect to the second variable by $\bar{\tau}$, and denote the associated function as in (2.12) by $\bar{h}$, then Lemma 2.6 also holds for $\bar{h}$.

Example 2.9. (i). Obviously, in the linear case that $\sigma(x, y) = \sigma_0(x) y$, Assumption H5 automatically holds with $\bar{\sigma} = \sigma$.

(ii). For the case in Example 2.3.(ii), $\sigma(x, y) = \sigma_0(x) \eta(y)$, Assumption H5 also holds automatically, with $\bar{\sigma}(x, y) = \sigma_0(x) (\nabla \eta(0)) y$. Indeed, this assumption reduces to $\frac{1}{\epsilon} \eta(\epsilon y) \rightarrow (\nabla \eta(0)) y$ for every y , while this is a consequence of L’Hôpital’s rule.

(iii). In the perturbation case that $\sigma(x, y) = \sigma_0(x) y + \delta(x, y)$ as in Example 2.3.(i), Assumption H5 reduces to that for every $y \in \mathbb{R}^d$, $\frac{1}{\epsilon} \delta(x, \epsilon y) \rightarrow (\nabla_y \delta(x, 0)) y$ uniformly in $x \in \mathbb{R}^d$, as $\epsilon \rightarrow 0$.

Finally, we need a centering assumption on b and e when and only when doing the homogenization.

Assumption H6. The functions b and e satisfy the *centering condition*,

$$\int_{\mathbb{T}^d} b(x) \mu(dx) = 0, \quad \int_{\mathbb{T}^d} e(x) \mu(dx) = 0,$$

where μ is the invariant probability measure on $(\mathbb{T}^d, \mathcal{B}(\mathbb{T}^d))$ of the solution $\tilde{X}^x$ of (2.3).

We remark here that the well-posedness of (2.3) and the existence and uniqueness of the invariant probability measure μ will be clear in Proposition 4.2 and Lemma 4.6, respectively.

Note that the centering assumption H6 is quite common and natural in the homogenization problems and the reader can also find it in [9, 18, 31].

3 Nonlocal Poisson equation with zeroth-order term

As mentioned in the introduction, we will apply Zvonkin’s transform to study the homogenization of SDEs and nonlocal PDEs. Before that, we shall investigate the strong well-posedness of all the SDEs presented in the previous section, and Zvonkin’s transform will also play an important

role in this step (see next section). For the generality, we will focus on SDE (2.3) with $\bar{\sigma}$ only satisfying Assumption H1-H4 in the whole of this section as well as Subsection 4.1, since SDEs (2.1) and (2.2) are basically of the same form.

The key is to consider the following nonlocal Poisson equation with zeroth-order term,

$$\kappa u - \mathcal{L}^\alpha u = f, \quad (3.1)$$

where $\kappa > 0$, and $\mathcal{L}^\alpha$ is the linear integro-partial differential operator given by

$$\mathcal{L}^\alpha := \mathcal{A}^{\bar{\sigma}, \nu^\alpha} + b \cdot \nabla, \quad (3.2)$$

which may be regarded as the infinitesimal generator of the solution process $\tilde{X}$ of (2.3) once we prove its well-posedness in the next section.

3.1 Well-posedness of nonlocal Poisson equation

We first adapt the maximum principle in [36, Proposition 3.2] as follows. The proof is almost the same and the difference is provided in Appendix A.

Lemma 3.1. *If $u \in C_b^{1+\gamma}(\mathbb{R}^d)$, $1+\gamma > \alpha$, is a solution to $\kappa u - \mathcal{L}^\alpha u = f$ with $\kappa > 0$ and $f \in C_b(\mathbb{R}^d)$, then*

$$\kappa \|u\|_0 \leq \|f\|_0.$$

Now we investigate the solvability of the Poisson equation with a zeroth-order term involved. The results generalize the Schauder estimates in [36] to the anisotropic nonlocal case.

Proposition 3.2. *For any $\kappa > 0$ and $f \in C_b^\beta(\mathbb{R}^d)$, where β is the exponent in Assumptions H2, the nonlocal Poisson equation (3.1) has a unique solution $u = u_\kappa \in C_b^{\alpha+\beta}(\mathbb{R}^d)$. In addition, there exists a positive constant $C = C(\kappa, \|b\|_\beta)$ such that*

$$\|u_\kappa\|_{\alpha+\beta} \leq C(\|u_\kappa\|_0 + \|f\|_\beta). \quad (3.3)$$

Proof. The a priori estimate (3.3) is from [6, Theorem 7.1, Theorem 7.2]. We thus need to show that the equation (3.1) has a unique solution $u_\kappa \in C_b^{\alpha+\beta}(\mathbb{R}^d)$.

Now we prove the existence and uniqueness of solution in $C_b^{\alpha+\beta}(\mathbb{R}^d)$. It is shown in [36, Theorem 3.4] that when $\bar{\sigma}(\cdot, y) \equiv \mathbf{Id}_y(y) := y$, the existence and uniqueness hold in $C_b^{\alpha+\beta}(\mathbb{R}^d)$. For the general $\bar{\sigma}$, we apply the method of continuity (see [17, Section 5.2]).

Define a family of linear operators by $\mathcal{L}_\theta := \theta \mathcal{A}^{\bar{\sigma}, \nu^\alpha} + (1 - \theta) \mathcal{A}^{\mathbf{Id}_y, \nu^\alpha} + b \cdot \nabla$. We consider the family of equations:

$$\kappa u - \mathcal{L}_\theta u = f.$$

We can also rewrite the nonlocal term in $\mathcal{L}_\theta$ into the form (2.8), with the kernel given by $\nu_\theta := \theta \nu^{\bar{\sigma}, \alpha} + (1 - \theta) \nu^\alpha$. Then the a priori estimate (3.3) also holds for u_θ (cf. Remark 2.7). As a result, the operator $\mathcal{L}_\theta$ can be considered as a bounded linear operator from the Banach space $C_b^{\alpha+\beta}(\mathbb{R}^d)$ into the Banach space $C_b^\beta(\mathbb{R}^d)$.

Note that $\mathcal{L}_0 = \mathcal{A}^{\mathbf{Id}_y, \nu^\alpha} + b \cdot \nabla$, which is the case considered in [36], and $\mathcal{L}_1 = \mathcal{L}^\alpha$. The solvability of the equation (3.1) for any $f \in C_b^\beta(\mathbb{R}^d)$ is then equivalent to the invertibility of the operator $\mathcal{L}_\theta$. We can see from the proof of Lemma 3.1 that $\|u_\theta\|_0 \leq C\|f\|_0$. Then together with the estimate (3.3) for u_θ , we have the bound

$$\|u_\theta\|_{\alpha+\beta} \leq C\|f\|_\beta,$$

with the constant C independent of θ . Since, as discussed in [36], the operator $\mathcal{L}_0 = \mathcal{A}^{\mathbf{Id}_y, \nu^\alpha} + b \cdot \nabla$ maps $C_b^{\alpha+\beta}(\mathbb{R}^d)$ onto $C_b^\beta(\mathbb{R}^d)$, the method of continuity is applicable and the result follows. $\square$

Remark 3.3. If we take the periodicity assumption [H1](#) into account, then we can slightly strengthen the conclusions in Proposition [3.2](#). That is, if $f \in \mathcal{C}^\beta(\mathbb{T}^d)$, then the unique solution of [\(3.1\)](#) is of class $\mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$.

3.2 Feller property and heat kernels

In this subsection, we will study further the operator $\mathcal{L}^\alpha$. All these results will be used in the next section. First of all, the classical theory of C_0 -semigroups yields that $\mathcal{L}^\alpha$ is the generator of a Feller semigroup. The proof is standard and provided in Appendix [A](#).

Lemma 3.4. *The linear operator $(\mathcal{L}^\alpha, D(\mathcal{L}^\alpha))$, $D(\mathcal{L}^\alpha) = \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$, defined on the Banach space $(\mathcal{C}(\mathbb{T}^d), \|\cdot\|_0)$, is closable and dissipative, its closure generates a Feller semigroup $\{P_t\}_{t \geq 0}$ on $\mathcal{C}(\mathbb{T}^d)$.*

Let us recall the notion of martingale problem (see [\[14, Section 4.3\]](#)). First recall that $\mathcal{D} = \mathcal{D}(\mathbb{R}_+; \mathbb{T}^d)$ is the space of all $\mathbb{T}^d$ -valued càdlàg functions on $\mathbb{R}_+$, equipped with the Skorokhod topology. Let $w_t(\omega) = \omega(t)$, $\omega \in \mathcal{D}$, be the coordinate process on $(\mathcal{D}, \mathcal{B}(\mathcal{D}))$, and $\{\mathcal{F}_t^w\}_{t \geq 0} := \sigma(w_s : 0 \leq s \leq t)$ be the canonical filtration. Given a probability measure ν on $\mathbb{T}^d$, we say that a probability measure $\mathbf{P}^\nu$ on $(\mathcal{D}, \mathcal{B}(\mathcal{D}))$ is a solution of the martingale problem for $(\mathcal{L}^\alpha, \nu)$, if $\mathbf{P}^\nu \circ w_0^{-1} = \nu$ and the process

$$M^f(t) := f(w_t) - f(w_0) - \int_0^t \mathcal{L}^\alpha f(w_s) ds$$

is a $(\mathcal{D}, \mathcal{B}(\mathcal{D}), \{\mathcal{F}_t^w\}_{t \geq 0}, \mathbf{P}^\nu)$ -martingale, for any $f \in D(\mathcal{L}^\alpha) = \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$. We denote by δ_x the Dirac measure, or equivalently, the Dirac function as distribution, focusing on $x \in \mathbb{R}^d$.

The following heat kernel results can be found in references, see Appendix [A](#) for details.

Lemma 3.5. *For every $x \in \mathbb{T}^d$, the martingale problem for $(\mathcal{L}^\alpha, \delta_x)$ has a unique solution $\mathbf{P}^x$. Moreover, the coordinate process $\{w_t\}_{t \geq 0}$ is a Feller process with generator the closure of $(\mathcal{L}^\alpha, \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d))$, and has a jointly continuous transition probability density $p(t; x, y)$, i.e., $\mathbf{P}^x(w_t \in A) = \int_A p(t; x, y) dy$, $A \in \mathcal{B}(\mathbb{T}^d)$, which satisfies for each $T > 0$,*

$$\begin{aligned} C_1^{-1} \sum_{j \in \mathbb{Z}^d} \left(\frac{t}{|x-y+j|^{d+\alpha}} \wedge t^{-\frac{d}{\alpha}} \right) &\leq p(t; x, y) \leq C_1 \sum_{j \in \mathbb{Z}^d} \left(\frac{t}{|x-y+j|^{d+\alpha}} \wedge t^{-\frac{d}{\alpha}} \right), \\ |\nabla_x p(t; x, y)| &\leq C_2 t^{-\frac{1}{\alpha}} \sum_{j \in \mathbb{Z}^d} \left(\frac{t}{|x-y+j|^{d+\alpha}} \wedge t^{-\frac{d}{\alpha}} \right), \end{aligned}$$

for all $x, y \in \mathbb{R}^d$ and $t \in (0, T]$, where $C_1 > 1, C_2 > 0$ are two constants depending on $d, \alpha, \|b\|_0, h_0, h_1$. The constants h_0, h_1 are related to the function h as in Lemma [2.6](#).

Remark 3.6. (1). Combining Lemma [3.4](#) and Lemma [3.5](#), we see that the Feller semigroup $\{P_t\}_{t \geq 0}$ generated by the closure of $\mathcal{L}^\alpha$ has the representation

$$P_t f(x) = \int_{\mathbb{T}^d} f(y) p(t; x, y) dy, \quad f \in \mathcal{C}(\mathbb{T}^d),$$

and the following gradient estimate holds

$$\begin{aligned} |\nabla P_t f(x)| &\leq C_2 \|f\|_0 t^{-\frac{1}{\alpha}} \int_{\mathbb{T}^d} \sum_{j \in \mathbb{Z}^d} \left(\frac{t}{|y+j|^{d+\alpha}} \wedge t^{-\frac{d}{\alpha}} \right) dy \\ &= C_2 \|f\|_0 t^{-\frac{1}{\alpha}} \int_{\mathbb{R}^d} \left(\frac{t}{|y|^{d+\alpha}} \wedge t^{-\frac{d}{\alpha}} \right) dy \leq C_2 \left(1 + \frac{1}{\alpha}\right) \|f\|_0 t^{-\frac{1}{\alpha}}. \end{aligned} \tag{3.4}$$

(2). Denote the formal generator of $\tilde{X}^{x,\epsilon}$ by $\tilde{\mathcal{L}}_\epsilon^\alpha$, i.e.,

$$\begin{aligned}\tilde{\mathcal{L}}_\epsilon^\alpha f(x) := & \int_{\mathbb{R}^d \setminus \{0\}} [f(x + \tfrac{1}{\epsilon} \sigma(x, \epsilon y)) - f(x) - \tfrac{1}{\epsilon} \sigma^i(x, \epsilon y) \partial_i f(x) \mathbf{1}_B(y)] \nu^\alpha(dy) \\ & + [b^i(x) + \epsilon^{\alpha-1} c^i(x)] \partial_i f(x), \quad x \in \mathbb{R}^d.\end{aligned}\tag{3.5}$$

Then Lemma 3.4 and 3.5 still hold true with $\tilde{\mathcal{L}}_\epsilon^\alpha$ in place of $\mathcal{L}^\alpha$.

As a corollary, the solution of equation (3.1) can be represented in terms of a semigroup, and satisfies a finer estimate.

Corollary 3.7. *For any $\kappa > 0$ and $f \in \mathcal{C}^\beta(\mathbb{T}^d)$, the unique solution u_κ of equation (3.1) admits the representation*

$$u_\kappa(x) = \int_0^\infty e^{-\kappa t} P_t f(x) dt, \tag{3.6}$$

where $\{P_t\}_{t \geq 0}$ is the Feller semigroup generated by the closure of $\mathcal{L}^\alpha$, and the integral on the right hand side converges. Moreover, there exists a constant $C > 0$ independent of u, f, b, κ such that

$$\kappa \|u_\kappa\|_0 + \kappa^{\frac{\alpha+\beta-1}{\alpha}} \|\nabla u_\kappa\|_0 + [\nabla u_\kappa]_{\alpha+\beta-1} \leq C \|f\|_\beta. \tag{3.7}$$

Proof. Proposition 3.2 tells that the interval $(0, +\infty)$ is contained in the resolvent set of $\mathcal{L}^\alpha$. Then by the integral representation of the resolvent (see [13, Theorem II.1.10.(ii)]), we arrive at

$$u_\kappa = (\kappa - \mathcal{L}^\alpha)^{-1} f = \lim_{t \rightarrow \infty} \int_0^t e^{-\kappa s} P_s f ds,$$

where the limit is taken in $(\mathcal{C}(\mathbb{T}^d), \|\cdot\|_0)$. The representation (3.6) then follows. Now thanks to the gradient estimate (3.4) and representation (3.6), the estimate (3.7) is then obtained by the same argument as the proof of [36, Theorem 3.3, Part I]. $\square$

In the next section, we will remove the large jumps from the SDEs and study their well-posedness by Zvonkin's transform. Thus we consider the following operator, which is a “flat” version of $\mathcal{L}^\alpha$:

$$\mathcal{L}^{\alpha,b} f(x) = \int_{B \setminus \{0\}} [f(x + \bar{\sigma}(x, y)) - f(x) - \bar{\sigma}^i(x, y) \partial_i f(x)] \nu^\alpha(dy) + b^i(x) \partial_i f(x). \tag{3.8}$$

We have the following regularity result for $\mathcal{L}^{\alpha,b}$.

Corollary 3.8. *There exists a constant $\kappa_* > 0$ such that for any $\kappa > \kappa_*$ and $f \in \mathcal{C}^\beta(\mathbb{T}^d)$, there exists a unique solution $u = u_\kappa^b \in \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$ to the equation*

$$\kappa u - \mathcal{L}^{\alpha,b} u = f. \tag{3.9}$$

In addition, there exists a constant $C > 0$ independent of u, f, b, κ , such that for any $\kappa > \kappa_$,*

$$(\kappa - \kappa_*) \|u_\kappa^b\|_0 + (\kappa - \kappa_*)^{\frac{\alpha+\beta-1}{\alpha}} \|\nabla u_\kappa^b\|_0 + [\nabla u_\kappa^b]_{\alpha+\beta-1} \leq C \|f\|_\beta. \tag{3.10}$$

Proof. To obtain the a priori estimate (3.10), we rewrite the equation (3.9) in the form

$$\kappa u - \mathcal{L}^\alpha u = f - \int_{B^c} [u(x + \bar{\sigma}(x, y)) - u(x)] \nu^\alpha(dy).$$

The estimate (3.7) implies that

$$\kappa\|u\|_0 + \kappa^{\frac{\alpha+\beta-1}{\alpha}}\|\nabla u\|_0 + [\nabla u]_{\alpha+\beta-1} \leq C(\|f\|_\beta + 2\nu^\alpha(B^c)\|u\|_\beta).$$

It is easy to see that there exists $\delta > 0$ such that

$$\sup_{|x-y|<\delta} \frac{|u(x) - u(y)|}{|x - y|} \leq 2\|\nabla u\|_0,$$

and then

$$\|u\|_\beta \leq \sup_{|x-y|<\delta} \frac{|u(x) - u(y)|}{|x - y|} |x - y|^{1-\beta} + \sup_{|x-y|\geq\delta} \frac{|u(x) - u(y)|}{|x - y|^\beta} \leq 2\delta^{1-\beta}\|\nabla u\|_0 + 2\delta^{-\beta}\|u\|_0.$$

Combining these together, we get

$$(\kappa - 4C\delta^{-\beta}\nu^\alpha(B^c))\|u\|_0 + (\kappa^{\frac{\alpha+\beta-1}{\alpha}} - 4C\delta^{1-\beta}\nu^\alpha(B^c))\|\nabla u\|_0 + [\nabla u]_{\alpha+\beta-1} \leq C\|f\|_\beta.$$

Then (3.10) follows by choosing $\kappa_* = 4C\delta^{-\beta}\nu^\alpha(B^c) \vee (4C\delta^{1-\beta}\nu^\alpha(B^c))^{\frac{\alpha}{\alpha+\beta-1}}$.

Now define a family of operators by

$$\mathcal{L}_\theta^\flat = \mathcal{L}^{\alpha,\flat} + \theta \int_{B^c} [u(x + \bar{\sigma}(x, y)) - u(x)] \nu^\alpha(dy).$$

Then $\mathcal{L}_1^\flat = \mathcal{L}^\alpha$, $\mathcal{L}_0^\flat = \mathcal{L}^{\alpha,\flat}$. The well-posedness of equation (3.9) follows from the method of continuity and the a priori estimate (3.10), just as in the proof of Proposition 3.2. $\square$

4 SDEs with multiplicative stable Lévy noise

The goal of this section is to study the strong well-posedness of SDEs (2.2) and (2.3), as well as the invariance and ergodicity of the solution processes $\tilde{X}^{x,\epsilon}$ for each $\epsilon \geq 0$. As corollaries, we also obtain the Feynman-Kac formula and the well-posedness of nonlocal Poisson equation without zeroth-order term, which will be used to study homogenization in the next two sections.

4.1 Strong well-posedness of SDEs

We only consider the strong well-posedness for SDE (2.3) since (2.2) has the same form. As we have seen in Lemma 3.5, the existence and uniqueness hold for the martingale problem for $(\mathcal{L}^\alpha, \delta_x)$. Meanwhile, it is known that the martingale solution for $(\mathcal{L}^\alpha, \delta_x)$ is equivalent to the weak solution of SDE (2.3), see [26, Theorem 2.3, Corollary 2.5]. Thus, the existence and uniqueness of weak solution hold for SDE (2.3).

Moreover, utilizing the fact shown in [5, Theorem 1.2] that the weak existence and pathwise uniqueness for SDE (2.3) imply strong existence, we only need to prove the pathwise uniqueness. The key is to reduce the SDE (2.3), whose coefficients have low regularity, to an SDE with Lipschitz coefficients by using Zvonkin's transform.

For $\kappa > \kappa_*$, let $\hat{b}_\kappa \in \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$ be the solution of

$$\kappa\hat{b}_\kappa - \mathcal{L}^{\alpha,\flat}\hat{b}_\kappa = b,$$

where $\mathcal{L}^{\alpha,\flat}$ is the operator in (3.8). The existence and uniqueness of solution $\hat{b}_\kappa$ is ensured by Corollary 3.8. Define a map $\Phi_\kappa : \mathbb{R}^d \rightarrow \mathbb{R}^d$ by

$$\Phi_\kappa(x) = x + \hat{b}_\kappa(x).$$

Then Φ_κ is of class $\mathcal{C}^{\alpha+\beta}$. Moreover, we have the following lemma, whose proof is standard and left into Appendix A.

Lemma 4.1. *For $\kappa > 0$ large enough, the map $\Phi_\kappa : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a C^1 -diffeomorphism and its inverse Φ_κ^{-1} is also of class $\mathcal{C}^{\alpha+\beta}$.*

To solve SDE (2.3), by a standard interlacing technique (cf. [2, Section 6.5] or [20, Theorem IV. 9.1]), it suffices to solve the following SDE with no jumps greater than one:

$$\tilde{X}_t^{x,b} = x + \int_0^t b(\tilde{X}_s^{x,b}) ds + \int_0^t \int_B \bar{\sigma}(\tilde{X}_{s-}^{x,b}, y) \tilde{N}^\alpha(dy, ds).$$

Now fix $\kappa > 0$ large enough such that the conclusions in Lemma 4.1 hold. We introduce Zvonkin's transform

$$\tilde{X}_t^* = \Phi_\kappa(\tilde{X}_t^{x,b}).$$

Then by applying Itô's formula, we have

$$\tilde{X}_t^* = \Phi_\kappa(x) + \int_0^t b^*(\tilde{X}_s^*) ds + \int_0^t \int_B \bar{\sigma}^*(\tilde{X}_{s-}^*, y) \tilde{N}^\alpha(dy, ds), \quad (4.1)$$

where

$$\begin{aligned} b^*(x) &= \kappa \hat{b}_\kappa(\Phi_\kappa^{-1}(x)), \\ \bar{\sigma}^*(x, y) &= \hat{b}_\kappa(\Phi_\kappa^{-1}(x) + \bar{\sigma}(\Phi_\kappa^{-1}(x), y)) - \hat{b}_\kappa(\Phi_\kappa^{-1}(x)) + \bar{\sigma}(\Phi_\kappa^{-1}(x), y). \end{aligned}$$

Proposition 4.2. *For each $x \in \mathbb{R}^d$, there is a unique strong solution $\tilde{X}^x = \{\tilde{X}_t^x\}_{t \geq 0}$ to SDE (2.3).*

Proof. By the above argument, we only need to prove the pathwise uniqueness for SDE (4.1). First of all, we have, for any $x, x_1, x_2 \in \mathbb{R}^d$,

$$|b^*(x_1) - b^*(x_2)| \leq C(\|\hat{b}_\kappa\|_1, \|\Phi_\kappa^{-1}\|_1) |x_1 - x_2|, \quad (4.2)$$

Note that for $\gamma \in (0, 1)$, $f \in \mathcal{C}_b^{1+\gamma}(\mathbb{R}^d)$, $x, u, v \in \mathbb{R}^d$, there exists a constant $C > 0$ such that

$$|f(u+x) - f(u) - f(v+x) + f(v)| \leq C \|f\|_{1+\gamma} |u - v| |x|^\gamma,$$

the proof can be found in [6, Theorem 5.1.(c)]. Then for any x_1, x_2 ,

$$\begin{aligned} & |\bar{\sigma}^*(x_1, y) - \bar{\sigma}^*(x_2, y)| \\ & \leq |\hat{b}_\kappa(\Phi_\kappa^{-1}(x_1) + \bar{\sigma}(\Phi_\kappa^{-1}(x_1), y)) - \hat{b}_\kappa(\Phi_\kappa^{-1}(x_1)) \\ & \quad - \hat{b}_\kappa(\Phi_\kappa^{-1}(x_2) + \bar{\sigma}(\Phi_\kappa^{-1}(x_1), y)) + \hat{b}_\kappa(\Phi_\kappa^{-1}(x_2))| \\ & \quad + |\hat{b}_\kappa(\Phi_\kappa^{-1}(x_2) + \bar{\sigma}(\Phi_\kappa^{-1}(x_1), y)) - \hat{b}_\kappa(\Phi_\kappa^{-1}(x_2) + \bar{\sigma}(\Phi_\kappa^{-1}(x_2), y))| \\ & \quad + |\bar{\sigma}(\Phi_\kappa^{-1}(x_1), y) - \bar{\sigma}(\Phi_\kappa^{-1}(x_2), y)| \\ & \leq C \|\hat{b}_\kappa\|_{\alpha+\beta} |\Phi_\kappa^{-1}(x_1) - \Phi_\kappa^{-1}(x_2)| |\bar{\sigma}(\Phi_\kappa^{-1}(x_1), y)|^{\alpha+\beta-1} \\ & \quad + (\|\nabla \hat{b}_\kappa\|_0 + 1) |\bar{\sigma}(\Phi_\kappa^{-1}(x_1), y) - \bar{\sigma}(\Phi_\kappa^{-1}(x_2), y)| \\ & \leq C \left(\|\hat{b}_\kappa\|_{\alpha+\beta}, \|\Phi_\kappa^{-1}\|_1, \|\phi\|_{L^\infty} \right) |x_1 - x_2| (|y|^{\alpha+\beta-1} + |y|). \end{aligned}$$

where we have used the regularity condition for $\bar{\sigma}$ in Assumption [H3](#), and ϕ is the positive bounded function in the growth condition in that assumption. Noting that $2(\alpha + \beta - 1) > \alpha$ by Assumption [H2](#), we arrive at

$$\int_B |\bar{\sigma}^*(x_1, y) - \bar{\sigma}^*(x_2, y)|^2 \nu_\alpha(dy) \leq C \left(\|\hat{b}_\kappa\|_{\alpha+\beta}, \|\Phi_\kappa^{-1}\|_1, \|\phi\|_{L^\infty} \right) |x_1 - x_2|^2. \quad (4.3)$$

The pathwise uniqueness of SDE [\(4.1\)](#) follows from [\(4.2\)](#), [\(4.3\)](#) and the classical result [[20](#), Theorem 4.9.1]. The proof is complete. $\square$

The following corollary is just a summarizing of above results for the process $\tilde{X}^x$, the proof is left into Appendix [A](#).

Corollary 4.3. *The solution process $\tilde{X}^x$ is a Feller process with generator the closure of $(\mathcal{L}^\alpha, \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d))$. In particular, $\tilde{X}^x$ is a strong Markov process.*

Remark 4.4. The Feller semigroup $\{P_t\}_{t \geq 0}$ in Lemma [3.4](#) is the semigroup associated with the solution process $\tilde{X}^x$, that is,

$$P_t f(x) = \mathbf{E}(f(\tilde{X}_t^x)), \quad f \in \mathcal{C}(\mathbb{T}^d).$$

As a consequence of the Feller property, we can obtain the well-posedness of the parabolic nonlocal PDE and the corresponding Feynman-Kac representation. See [[33](#)] for the classical version for second order PDE.

Proposition 4.5. *The parabolic nonlocal PDE*

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) = \mathcal{L}^\alpha u(t, x) + g(x)u(t, x), & t > 0, x \in \mathbb{R}^d, \\ u(0, x) = u_0(x), & x \in \mathbb{R}^d, \end{cases}$$

admits a unique mild solution in the sense that $\int_0^t u(s)ds \in D(\mathcal{L}^\alpha)$ for all $t \geq 0$ and

$$u(t) = u_0 + \mathcal{L}^\alpha \int_0^t u(s)ds + g \int_0^t u(s)ds.$$

Moreover, the unique solution has the following Feynman-Kac representation

$$u(t, x) = \mathbf{E} \left[u_0(\tilde{X}_t^x) \exp \left(\int_0^t g(\tilde{X}_s^x) ds \right) \right].$$

Proof. Choose $G > 0$ large enough such that $\|g\|_0 < G$. Define

$$P_t^g f(x) = \mathbf{E} \left[f(\tilde{X}_t^x) \exp \left(\int_0^t g(\tilde{X}_s^x) ds - Gt \right) \right], \quad f \in \mathcal{C}(\mathbb{T}^d).$$

Then by an argument similar to that used in [[2](#), Section 6.7.2], one can show that $\{P_t^g\}_{t \geq 0}$ is a Feller semigroup with generator the closure of $(\mathcal{L}^\alpha + g - G, \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d))$. This yields that $\{e^{Gt} P_t^g\}_{t \geq 0}$ is a C_0 -semigroup on $\mathcal{C}(\mathbb{T}^d)$ with generator the closure of $(\mathcal{L}^\alpha + g, \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d))$. Note that $u_0 \in \mathcal{C}(\mathbb{T}^d)$. Now applying the classic result [[13](#), Proposition II.6.4] in the theory of C_0 -semigroups, we conclude that the parabolic nonlocal PDE admits a unique mild solution, which can be given by the orbit map $u(t) = e^{Gt} P_t^g u_0$. The desired conclusions follow immediately. $\square$

4.2 Invariance and ergodicity

Now we deal with the invariant measures and ergodicity of SDEs. From now on, Assumptions **H1-H5** are set in force.

By the discussion in previous subsection, SDE (2.1) and SDE (2.2) for each $\epsilon \geq 0$, admit the unique strong solutions $\tilde{X}^x$ and $\tilde{X}^{x,\epsilon}$, which are all $\mathbb{T}^d$ -valued Feller processes. Denote by $p^\epsilon(t; x, y)$ the transition probability density of $\tilde{X}^{x,\epsilon}$, by $\{P_t^\epsilon\}_{t \geq 0}$ the associated Feller semigroup. Recall that we denote $\tilde{X}^{x,0} := \tilde{X}^x$ for notational simplicity.

Lemma 4.6. *For each $0 \leq \epsilon \leq 1$, the process $\tilde{X}^{x,\epsilon}$ possesses a unique invariant distribution μ_ϵ on $\mathbb{T}^d$. Moreover, there exist positive constants C and ρ , depending only on $d, \alpha, \|b\|_0, \|c\|_0, h_0, h_1$, such that for any periodic bounded Borel function f on $\mathbb{R}^d$ (i.e., f is Borel bounded on $\mathbb{T}^d$),*

$$\sup_{x \in \mathbb{T}^d} \left| P_t^\epsilon f(x) - \int_{\mathbb{T}^d} f(y) \mu_\epsilon(dy) \right| \leq C \|f\|_0 e^{-\rho t}$$

for every $t \geq 0$.

Proof. One can find a version of Doeblin-type result in [9, Theorem 3.3.1, 3.3.2], which states that for a Markov process with transition probability densities bounded from below by a positive constant, it has a unique invariant probability measure and the associated semigroup converges exponentially fast. Therefore, it is enough to ensure that the transition probability density $p^\epsilon(1; x, y)$ is bounded from below by a positive constant, which follows immediately from the density estimates in Lemma 3.5. Moreover, one can also realize from [9, Theorem 3.3.1, 3.3.2] that the two constants C and ρ are only related to the lower bound of $p^\epsilon(1; x, y)$. The only thing left is to verify the dependence of the constants C and ρ . In particular, they can be chosen uniformly in $\epsilon \in [0, 1]$.

In the case $\epsilon = 0$, the lower bound of $p(1; x, y)$ depends on $d, \alpha, \|b\|_0, \bar{h}_0, \bar{h}_1$ by Lemma 3.5, where $\bar{h}_0$ and $\bar{h}_1$ are the constants associated to $\bar{\sigma}$ as indicated in Lemma 2.6 and Remark 2.8. We turn to the case that $\epsilon \in (0, 1]$. Since the generator of each semigroup $\{P_t^\epsilon\}$ is $\tilde{\mathcal{L}}_\epsilon^\alpha = \mathcal{A}_\epsilon^{\frac{1}{\epsilon}\sigma_\epsilon, \nu^\alpha} + (b + \epsilon^{\alpha-1}c) \cdot \nabla$ as in (3.5), where $\sigma_\epsilon(x, y) := \sigma\left(\frac{x}{\epsilon}, y\right)$, it is easy to derive that the function h^ϵ associated with $\frac{1}{\epsilon}\sigma_\epsilon$ as in (2.12) is given by

$$h_\epsilon(x, z) = |\det \nabla_z \tau(x, z)| \frac{|\epsilon z|^{d+\alpha}}{|\tau(x, \epsilon z)|^{d+\alpha}},$$

Hence, the constants h_0^ϵ and h_1^ϵ associated to h_ϵ as in Lemma 2.6 can be chosen to equal to h_0 and h_1 , respectively. We conclude that when $\epsilon \in (0, 1]$, the constant C_1 associated to $p^\epsilon(t; x, y)$ in Lemma 3.5 are related to $d, \alpha, \|b + \epsilon^{\alpha-1}c\|_0, h_0, h_1$. Hence, for $\epsilon \in [0, 1]$, constants C and ρ can be chosen to depend only on $d, \alpha, \|b\|_0, \|c\|_0, h_0, h_1, \bar{h}_0, \bar{h}_1$. In particular, they are independent of $\epsilon \in [0, 1]$. $\square$

Denote by $\mu = \mu_0$ the unique invariant probability measure for the limit process $\tilde{X}_t^x$ in (2.3). Then we can prove the following lemma.

Lemma 4.7. *As $\epsilon \rightarrow 0$, we have $\mu_\epsilon \rightarrow \mu$ weakly.*

Proof. Using the same argument as the proof of [18, Lemma 2.4], and noting that the tightness of the family $\{\mu_\epsilon\}_{\epsilon > 0}$ is automatic due to the compactness of $\mathbb{T}^d$, it suffices to prove that $P_t^\epsilon f \rightarrow P_t f$ in $\mathcal{C}(\mathbb{T}^d)$ as $\epsilon \rightarrow 0$ for any $f \in \mathcal{C}(\mathbb{T}^d)$ and $t \geq 0$. By Lemma 3.5 and Remark 3.6.(2), we know that

$\mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$ is a core for $\mathcal{L}^\alpha$ and each $\tilde{\mathcal{L}}_\epsilon^\alpha$, $\epsilon > 0$. Fix an arbitrary $f \in \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$,

$$\begin{aligned} |\tilde{\mathcal{L}}_\epsilon^\alpha f(x) - \mathcal{L}^\alpha f(x)| &\leq \epsilon^{\alpha-1} |c(x) \cdot \nabla f(x)| + \int_{B^c} |f(x + \tfrac{1}{\epsilon}\sigma(x, \epsilon y)) - f(x + \bar{\sigma}(x, y))| \nu^\alpha(dy) \\ &\quad + \int_{B \setminus \{0\}} |f(x + \tfrac{1}{\epsilon}\sigma(x, \epsilon y)) - f(x + \bar{\sigma}(x, y)) - (\tfrac{1}{\epsilon}\sigma(x, \epsilon y) - \bar{\sigma}(x, y)) \cdot \nabla f(x)| \nu^\alpha(dy) \\ &\leq \epsilon^{\alpha-1} \|c\|_0 \|f\|_1 + \|\nabla f\|_0 \int_{B^c} |\tfrac{1}{\epsilon}\sigma(x, \epsilon y) - \bar{\sigma}(x, y)| \nu^\alpha(dy) \\ &\quad + \tfrac{1}{\alpha+\beta} \|f\|_{\alpha+\beta} \int_{B \setminus \{0\}} |\tfrac{1}{\epsilon}\sigma(x, \epsilon y) - \bar{\sigma}(x, y)|^{\alpha+\beta} \nu^\alpha(dy). \end{aligned}$$

By the growth condition for σ and $\bar{\sigma}$ as indicated in Assumption H3 and Remark 2.8,

$$\int_{B^c} \sup_{x \in \mathbb{T}^d} |\tfrac{1}{\epsilon}\sigma(x, \epsilon y) - \bar{\sigma}(x, y)| \nu^\alpha(dy) \leq 2\|\phi\|_{L^\infty} \int_{B^c} |y| \nu^\alpha(dy) < \infty,$$

since $\alpha > 1$. Similarly,

$$\int_{B \setminus \{0\}} \sup_{x \in \mathbb{T}^d} |\tfrac{1}{\epsilon}\sigma(x, \epsilon y) - \bar{\sigma}(x, y)|^{\alpha+\beta} \nu^\alpha(dy) \leq (2\|\phi\|_{L^\infty})^{\alpha+\beta} \int_{B \setminus \{0\}} |y|^{\alpha+\beta} \nu^\alpha(dy) < \infty.$$

Hence, the dominated convergence and Assumption H5 implies that $|\tilde{\mathcal{L}}_\epsilon^\alpha f(x) - \mathcal{L}^\alpha f(x)|$ converges to zero uniformly in x , as $\epsilon \rightarrow 0$. That is,

$$\|\tilde{\mathcal{L}}_\epsilon^\alpha f - \mathcal{L}^\alpha f\|_0 \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0.$$

Now using the Trotter-Kato approximation theorem (see [13, Theorem III.4.8]), $P_t^\epsilon f \rightarrow P_t f$ in $\mathcal{C}(\mathbb{T}^d)$ as $\epsilon \rightarrow 0$ for all $f \in \mathcal{C}(\mathbb{T}^d)$, uniformly for t in compact intervals. $\square$

Now we combine Lemma 4.6 and Lemma 4.7 to get the following ergodic theorem.

Proposition 4.8. *Let f be a bounded Borel function on $\mathbb{T}^d$. Then for any $t > 0$,*

$$\int_0^t \left| f\left(\frac{X_s^{x,\epsilon}}{\epsilon}\right) - \int_{\mathbb{T}^d} f(x) \mu(dx) \right| ds \rightarrow 0$$

in probability, as $\epsilon \rightarrow 0$.

Proof. We follow the lines of [31, Proposition 2.4]. For $\epsilon > 0$, $0 \leq s < t$, let $\bar{f}$ be a bounded measurable function on $\mathbb{T}^d$ satisfying $\int_{\mathbb{T}^d} \bar{f}(x) \mu_\epsilon(dx) = 0$. By Lemma 4.7, it suffices to prove that $\int_0^t |\bar{f}(X_s^\epsilon/\epsilon)| ds \rightarrow 0$ in $L^2(\Omega, \mathbf{P})$. Using Lemma 4.6, we have

$$\mathbf{E} \left[|\bar{f}(\tilde{X}_t^{x,\epsilon})| \middle| \tilde{X}_s^{x,\epsilon} \right] = \int_{\mathbb{T}^d} |\bar{f}(y)| \left[p^\epsilon(t-s, \tilde{X}_s^{x,\epsilon}, y) dy - \mu_\epsilon(dy) \right] \leq C \|\bar{f}\|_0 e^{-\rho(t-s)}.$$

By the Markov property,

$$\mathbf{E} |\bar{f}(\tilde{X}_s^{x,\epsilon}) \bar{f}(\tilde{X}_t^{x,\epsilon})| = \mathbf{E} \left[|\bar{f}(\tilde{X}_s^{x,\epsilon})| \mathbf{E} \left(|\bar{f}(\tilde{X}_t^{x,\epsilon})| \middle| \tilde{X}_s^{x,\epsilon} \right) \right] \leq C \|\bar{f}\|_0^2 e^{-\rho(t-s)}.$$

Hence,

$$\begin{aligned}
\mathbf{E} \left[\left(\int_0^t |\bar{f}(X_s^{x,\epsilon}/\epsilon)| ds \right)^2 \right] &= \epsilon^{2\alpha} \int_0^{\epsilon^{-\alpha}t} \int_0^r \mathbf{E} |\bar{f}(\tilde{X}_s^{x,\epsilon}) \bar{f}(\tilde{X}_r^{x,\epsilon})| ds dr \\
&\leq 2C\epsilon^{2\alpha} \|f\|_0^2 \int_0^{\epsilon^{-\alpha}t} \int_0^r e^{-\rho(r-s)} ds dr \\
&= 2C\epsilon^{2\alpha} \|f\|_0^2 \rho^{-2} (-1 + \rho\epsilon^{-\alpha}t + e^{-\rho\epsilon^{-\alpha}t}) \\
&\rightarrow 0,
\end{aligned}$$

as $\epsilon \rightarrow 0$. The results follow. $\square$

For every $\gamma > 0$, denote by $\mathcal{C}_\mu^\gamma(\mathbb{T}^d)$ the class of all $f \in \mathcal{C}^\gamma(\mathbb{T}^d)$ which are *centered* with respect to the invariant measure μ in the sense that $\int_{\mathbb{T}^d} f(x) \mu(dx) = 0$. It is easy to check that $\mathcal{C}_\mu^\gamma(\mathbb{T}^d)$ is closed, and hence a sub-Banach space of $\mathcal{C}^\gamma(\mathbb{T}^d)$ under the norm $\|\cdot\|_\gamma$.

Thanks to Proposition 3.2 and Lemma 4.6, we can use the Fredholm alternative to obtain the solvability of the following Poisson equation without zeroth-order term in the smaller space $\mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$,

$$\mathcal{L}^\alpha u + f = 0, \quad (4.4)$$

for $f \in \mathcal{C}_\mu^\beta(\mathbb{T}^d)$. Before that, we need some lemmas that are straightforward consequences of the classical theory of C_0 -semigroups. We provide their proofs in Appendix A.

Lemma 4.9. *The restrictions $\{P_t^\mu := P_t|_{\mathcal{C}_\mu(\mathbb{T}^d)}\}_{t \geq 0}$ form a C_0 -semigroup on the Banach space $(\mathcal{C}_\mu(\mathbb{T}^d), \|\cdot\|_0)$, with generator given by $\mathcal{L}_\mu^\alpha f := \mathcal{L}^\alpha f$, $D(\mathcal{L}_\mu^\alpha) := \mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$.*

Lemma 4.10. *If $f \in \mathcal{C}_\mu^\beta(\mathbb{T}^d)$, then the unique solution u_κ of (3.1) is of class $\mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$, for any $\kappa > 0$.*

The following theorem will solve the well-posedness of equation (4.4), which is more general than the results in [16, Proposition 3]. We formulate it as follows, referring to [34, Theorem 1] for the classical version for second order partial differential operators.

Proposition 4.11. *For any $f \in \mathcal{C}_\mu^\beta(\mathbb{T}^d)$, there exists a unique solution in $\mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$ to the equation (4.4), which satisfies the estimate*

$$\|u\|_{\alpha+\beta} \leq C(\|u\|_0 + \|f\|_\beta), \quad (4.5)$$

where $C = C(\|b\|_\beta)$ is a positive constant. Moreover, the unique solution admits the representation

$$u(x) = \int_0^\infty P_t f(x) dt. \quad (4.6)$$

Proof. The a priori estimate (4.5) is also from [6, Theorem 7.1].

First, we show that if the equation has a solution $u \in \mathcal{C}_\mu(\mathbb{T}^d)$ for $f \in \mathcal{C}_\mu^\beta(\mathbb{T}^d)$, then u must have the representation (4.6), this also implies the uniqueness. By the exponential ergodicity result in Lemma 4.6, we have $\|P_t^\mu f\|_0 \leq C\|f\|_0 e^{-\rho t}$ for any $f \in \mathcal{C}_\mu(\mathbb{T}^d)$ and $t \geq 0$. This yields that, using [13, Theorem II.1.10.(ii)] as in the proof of Corollary 3.7, the set $\{z \in \mathbb{C} | \operatorname{Re} z > -\rho\}$ is contained in the resolvent set of $\mathcal{L}_\mu^\alpha$. Noting that $u = (0 - \mathcal{L}_\mu^\alpha)^{-1} f$, the representation and uniqueness follow.

Now we prove the existence. Let κ_0 be a fixed positive constant. Thanks to Lemma 4.10, the linear map $\kappa_0 - \mathcal{L}^\alpha : \mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d) \rightarrow \mathcal{C}_\mu^\beta(\mathbb{T}^d)$ is invertible. Furthermore, by virtue of Lemma 3.1

and the energy estimate (3.3), together with the compact embedding $\mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d) \subset \mathcal{C}_\mu^\beta(\mathbb{T}^d)$ (see, for instance, [17, Lemma 6.36]), the resolvent $\mathcal{R}_{\kappa_0} := (\kappa_0 - \mathcal{L}^\alpha)^{-1}$ is compact from $\mathcal{C}_\mu^\beta(\mathbb{T}^d)$ to $\mathcal{C}_\mu^\beta(\mathbb{T}^d)$. Consider then the equation

$$u - \kappa_0 \mathcal{R}_{\kappa_0} u = \mathcal{R}_{\kappa_0} f, \quad f \in \mathcal{C}_\mu^\beta(\mathbb{T}^d), \quad (4.7)$$

Then the Fredholm alternative (see [17, Section 5.3]) implies that the equation (4.7) always has a unique solution $u \in \mathcal{C}_\mu^\beta(\mathbb{T}^d)$ provided the homogeneous equation $u - \kappa_0 \mathcal{R}_{\kappa_0} u = 0$ has only the trivial solution $u = 0$.

To rephrase these statements in terms of the Poisson equation (4.4), we observe first that since $\mathcal{R}_{\kappa_0}$ maps $\mathcal{C}_\mu^\beta(\mathbb{T}^d)$ onto $\mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$, any solution $u \in \mathcal{C}_\mu^\beta(\mathbb{T}^d)$ of (4.7) must also belong to $\mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$. Hence, operating on (4.7) with $\kappa_0 - \mathcal{L}^\alpha$ we obtain

$$-\mathcal{L}^\alpha u = (\kappa_0 - \mathcal{L}^\alpha)(u - \kappa_0 \mathcal{R}_{\kappa_0} u) = f.$$

Thus, the solutions of (4.7) are in one-to-one correspondence with the solutions of the Poisson equation (4.4). Consequently, (4.7) has a unique solution in $\mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$ if we can show that the homogeneous equation $\mathcal{L}^\alpha u = 0$ has only the zero solution, while the latter follows from the representation (4.6). $\square$

Remark 4.12. The assumption that f is centered with respect to μ in Proposition 4.11 is necessary. To see this informally, let's recall the Riesz-Schauder theory for compact operators (cf. [46, Theorem X.5.3]). The equation (4.7) admits a solution $u \in \mathcal{C}(\mathbb{T}^d)$ if and only if $\mathcal{R}_\kappa f \in \text{Ker}(I^* - \kappa \mathcal{R}_\kappa^*)^\perp$, where the superscript $*$ denotes the adjoint of operators. This is equivalent to say that the equation (4.4) admits a solution $u \in \mathcal{C}(\mathbb{T}^d)$ if and only if $f \in \text{Ker}(L^{\alpha,*})^\perp$. On the other hand, we have $\mu \in \text{Ker}(L^{\alpha,*})$ since μ is the invariant measure with respect to $\{P_t\}_{t \geq 0}$. Thus a necessary condition for the existence of (4.4) is $\langle \mu, f \rangle = 0$, regarding μ as an element in the dual space of $\mathcal{C}(\mathbb{T}^d)$.

5 Homogenization results

5.1 Homogenization of SDEs

The aim of this subsection is to show the homogenization result of the solutions $X^{x,\epsilon}$ of SDEs (2.1). We will let Assumptions H1-H6 all hold true in this and next subsection.

It is quite natural to get rid of the drift term involving $\frac{1}{\epsilon^{\alpha-1}}$ in (2.1). For this purpose, we again use Zvonkin's transform,

$$\hat{X}_t^{x,\epsilon} := X_t^{x,\epsilon} + \epsilon \left(\hat{b} \left(\frac{X_t^{x,\epsilon}}{\epsilon} \right) - \hat{b} \left(\frac{x}{\epsilon} \right) \right),$$

where $\hat{b}$ is the solution of the Poisson equation

$$\mathcal{L}^\alpha \hat{b} + b = 0, \quad (5.1)$$

with the linear operator $\mathcal{L}^\alpha$ given by (3.2). Note that the transform here is slightly different from that used in Section 4. Due to Proposition 4.11, $\hat{b} \in \mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$ is uniquely determined under Assumptions H6.

We also need an elementary lemma. The proof is elementary and shall be omitted.

Lemma 5.1. *Let $0 < \gamma \leq 1$ and $f \in \mathcal{C}_b^{1+\gamma}(\mathbb{R}^d)$. For any $x, u, v \in \mathbb{R}^d$, it holds that*

$$|f(x+u) - f(x+v) - (u-v) \cdot \nabla f(x)| \leq \frac{1}{1+\gamma} [\nabla f]_\gamma |u-v|^{1+\gamma}.$$

Now we are in a position to study the homogenization of SDEs with multiplicative stable noise.

Proposition 5.2. *In the sense of weak convergence on the space $\mathcal{D}$, we have that,*

$$X^{x,\epsilon} \Rightarrow X^x, \quad \text{where } X_t^x := x + \bar{C}t + \bar{L}_t,$$

as $\epsilon \rightarrow 0$. The homogenized coefficient $\bar{C}$ is given by

$$\bar{C} = \int_{\mathbb{T}^d} (I + \nabla \hat{b})c(x)\mu(dx), \quad (5.2)$$

and $\{\bar{L}_t\}_{t \geq 0}$ is a symmetric α -stable Lévy processes with jump intensity measure

$$\bar{\nu}(A) = \int_{\mathbb{R}^d \setminus \{0\}} \int_{\mathbb{T}^d} \mathbf{1}_A(\sigma(x, y))\mu(dx)\nu^\alpha(dy), \quad A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}). \quad (5.3)$$

Proof. Since $\hat{b}$ is bounded, the theorem will follow if we prove that $\hat{X}^{x,\epsilon} \Rightarrow X^x$, as $\epsilon \rightarrow 0$. By applying Itô's formula, and note that $\hat{b} \in \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$ is the solution of Poisson equation (5.1),

$$\begin{aligned} \hat{X}_t^{x,\epsilon} &= x + \int_0^t (I + \nabla \hat{b})c\left(\frac{X_s^{x,\epsilon}}{\epsilon}\right)ds - \int_0^t \frac{1}{\epsilon^{\alpha-1}} \mathcal{A}^{\bar{\sigma}, \nu^\alpha} \hat{b}\left(\frac{X_s^{x,\epsilon}}{\epsilon}\right)ds + \int_0^t \epsilon \mathcal{A}^{\sigma_\epsilon, \nu^\alpha} \hat{b}_\epsilon(X_s^{x,\epsilon})ds \\ &\quad + \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \epsilon \left[\hat{b}_\epsilon(X_{s-}^{x,\epsilon} + \sigma_\epsilon(X_{s-}^{x,\epsilon}, y)) - \hat{b}_\epsilon(X_{s-}^{x,\epsilon}) \right] \tilde{N}^\alpha(dy, ds) \\ &\quad + \int_0^t \int_{B \setminus \{0\}} \sigma_\epsilon(X_{s-}^{x,\epsilon}, y) \tilde{N}^\alpha(dy, ds) + \int_0^t \int_{B^c} \sigma_\epsilon(X_{s-}^{x,\epsilon}, y) N^\alpha(dy, ds) \\ &=: x + \Lambda_1^\epsilon(c)_t - \Lambda_2^\epsilon(\hat{b}, \mathcal{A}^{\bar{\sigma}, \nu^\alpha})_t + \Lambda_3^\epsilon(\hat{b}, \mathcal{A}^{\sigma_\epsilon, \nu^\alpha})_t + \Lambda_4^\epsilon(\hat{b}, \tilde{N}^\alpha)_t + \Lambda_5^\epsilon(\sigma, \tilde{N}^\alpha)_t + \Lambda_6^\epsilon(\sigma, N^\alpha)_t. \end{aligned}$$

where $\hat{b}_\epsilon(x) := \hat{b}\left(\frac{x}{\epsilon}\right)$, $\sigma_\epsilon(x, y) := \sigma\left(\frac{x}{\epsilon}, y\right)$.

For the last three stochastic integral terms, we figure out the characteristics of them as semi-martingales (cf. [22, Proposition IX.5.3]). Choose the truncation function $h_1(x) = x\mathbf{1}_B(x)$. Denote by $\Xi^\epsilon(s, y) := \epsilon[\hat{b}_\epsilon(X_{s-}^{x,\epsilon} + \sigma_\epsilon(X_{s-}^{x,\epsilon}, y)) - \hat{b}_\epsilon(X_{s-}^{x,\epsilon})]$. Note that $\Xi^\epsilon(\cdot, 0) \equiv 0$ by virtue of $\sigma(\cdot, 0) \equiv 0$ as mentioned in Remark 2.1.(3). Then the characteristics of $\Lambda_4^\epsilon(\hat{b}, \tilde{N}^\alpha)$ associated with h_1 is given by

$$\left\{ \begin{aligned} B_4^\epsilon(t) &= - \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \Xi^\epsilon(s, y) \mathbf{1}_{B^c}(\Xi^\epsilon(s, y)) \nu^\alpha(dy) ds, \\ C_4^\epsilon &\equiv 0, \\ \nu_4^\epsilon(A \times [0, t]) &= \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \mathbf{1}_A(\Xi^\epsilon(s, y)) \nu^\alpha(dy) ds, \quad A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}). \end{aligned} \right.$$

The characteristics of $\Lambda_5^\epsilon(\sigma, \tilde{N}^\alpha) + \Lambda_6^\epsilon(\sigma, N^\alpha)$ is given by

$$\left\{ \begin{aligned} B_{5+6}^\epsilon(t) &= \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \sigma_\epsilon(X_{s-}^{x,\epsilon}, y) [\mathbf{1}_B(\sigma_\epsilon(X_{s-}^{x,\epsilon}, y)) - \mathbf{1}_B(y)] \nu^\alpha(dy) ds, \\ C_{5+6}^\epsilon &\equiv 0, \\ \nu_{5+6}^\epsilon(A \times [0, t]) &= \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \mathbf{1}_A(\sigma_\epsilon(X_{s-}^{x,\epsilon}, y)) \nu^\alpha(dy) ds, \quad A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}). \end{aligned} \right.$$

By the same argument as in (2.7), we have $B_{5+6}^\epsilon \equiv 0$.

Then the theorem is a consequence of the functional central limit theorem in [22, Theorem VIII.2.17], and the following lemma whose proof is technical and provided in Appendix A. $\square$

Lemma 5.3. *For any $t \in \mathbb{R}_+$, and any bounded continuous function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ which vanishes in a neighbourhood of the origin, the following convergences hold in probability $\mathbf{P}$ when $\epsilon \rightarrow 0$:*

- (i) $\sup_{0 \leq s \leq t} |\Lambda_1^\epsilon(c)_s - \bar{C}s| \rightarrow 0$;
- (ii) $\sup_{0 \leq s \leq t} |\Lambda_3^\epsilon(\hat{b}, \mathcal{A}^{\sigma_\epsilon, \nu^\alpha})_s - \Lambda_2^\epsilon(\hat{b}, \mathcal{A}^{\bar{\sigma}, \nu^\alpha})_s| \rightarrow 0$;
- (iii) $\sup_{0 \leq s \leq t} |B_4^\epsilon(s)| \rightarrow 0$;
- (iv) $\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(x) \nu_4^\epsilon(dx, ds) \rightarrow 0$;
- (v) $\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(x) \nu_{5+6}^\epsilon(dx, ds) \rightarrow t \int_{\mathbb{R}^d \setminus \{0\}} f(x) \bar{\nu}(dx)$;

where $\bar{C}$ and $\bar{\nu}$ are defined in (5.2) and (5.3), respectively.

5.2 Homogenization of linear nonlocal PDEs

Define

$$Y_t^\epsilon := \int_0^t \left(\frac{1}{\epsilon^{\alpha-1}} e \left(\frac{X_s^{x,\epsilon}}{\epsilon} \right) + g \left(\frac{X_s^{x,\epsilon}}{\epsilon} \right) \right) ds.$$

Thanks to Proposition 4.5, the nonlocal PDE (1.1) has a unique mild solution, which is given by the Feynman-Kac formula,

$$u^\epsilon(t, x) = \mathbf{E} [u_0(X_t^{x,\epsilon}) \exp(Y_t^\epsilon)].$$

Similar to $\hat{X}^{x,\epsilon}$, we define

$$\hat{Y}_t^\epsilon := Y_t^\epsilon + \epsilon \left(\hat{e} \left(\frac{Y_t^\epsilon}{\epsilon} \right) - \hat{e} \left(\frac{x}{\epsilon} \right) \right).$$

Here $\hat{e} \in \mathcal{C}_\mu^{\alpha+\beta}(\mathbb{T}^d)$, thanks to Proposition 4.11 and Assumption H6, is the unique solution of the Poisson equation

$$\mathcal{L}^\alpha \hat{e} + e = 0,$$

with $\mathcal{L}^\alpha$ given by (3.2). Again using Itô's formula,

$$\begin{aligned} \hat{Y}_t^\epsilon &= \int_0^t (g + \nabla \hat{e} c) \left(\frac{X_s^{x,\epsilon}}{\epsilon} \right) ds - \int_0^t \frac{1}{\epsilon^{\alpha-1}} \mathcal{A}^{\bar{\sigma}, \nu^\alpha} \hat{e} \left(\frac{X_s^{x,\epsilon}}{\epsilon} \right) ds + \int_0^t \epsilon \mathcal{A}^{\sigma_\epsilon, \nu^\alpha} \hat{e}_\epsilon(X_s^{x,\epsilon}) ds \\ &\quad + \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \epsilon [\hat{e}_\epsilon(X_{s-}^{x,\epsilon} + \sigma_\epsilon(X_{s-}^{x,\epsilon}, y)) - \hat{e}_\epsilon(X_{s-}^{x,\epsilon})] \tilde{N}^\alpha(dy, ds) \\ &=: \Lambda_1^\epsilon(c, g)_t - \Lambda_2^\epsilon(\hat{e}, \mathcal{A}^{\bar{\sigma}, \nu^\alpha})_t + \Lambda_3^\epsilon(\hat{e}, \mathcal{A}^{\sigma_\epsilon, \nu^\alpha})_t + \Lambda_4^\epsilon(\hat{e}, \tilde{N}^\alpha)_t. \end{aligned}$$

Then in the same way as the proof of Proposition 5.2, we have the convergence of Y^ϵ to a deterministic path in distribution, as well as in probability, since the limit is deterministic (see, e.g., [44, Theorem 2.7.(iii)]).

Lemma 5.4. *In the sense of weak convergence on the space $\mathcal{D}$, both Y^ϵ and $\hat{Y}^\epsilon$ converge in distribution, and hence in probability, to the deterministic path $y(t) := \bar{E}t$ as $\epsilon \rightarrow 0$, where the homogenized coefficient $\bar{E}$ is given by*

$$\bar{E} := \int_{\mathbb{T}^d} (g + \nabla \hat{e} c)(x) \mu(dx). \quad (5.4)$$

Now we are in the position to prove the main result of this section. Since $\hat{b}$ and $\hat{e}$ are bounded on $\mathbb{R}^d$, u^ϵ has the same limit behavior as

$$\hat{u}^\epsilon(t, x) := \mathbf{E}[u_0(X_t^{x, \epsilon}) \exp(\hat{Y}_t^\epsilon)]$$

when $\epsilon \rightarrow 0$. Hence, we only need to show $\hat{u}^\epsilon(t, x) \rightarrow u(t, x)$, $\epsilon \rightarrow 0$ for any $t \geq 0, x \in \mathbb{R}^d$, where u has the following Feynman-Kac representation by virtue of Proposition 4.5,

$$u(t, x) = \mathbf{E}[u_0(X_t^x)]e^{y(t)}.$$

Proof of Theorem 1.1. For the convenience of notation, we shall write $\Lambda_1^\epsilon(c, g)_t$, $\Lambda_2^\epsilon(\hat{e}, \mathcal{A}^{\sigma, \nu^\alpha})_t$, $\Lambda_3^\epsilon(\hat{e}, \mathcal{A}^{\sigma_\epsilon, \nu^\alpha})_t$, $\Lambda_4^\epsilon(\hat{e}, \tilde{N}^\alpha)_t$ as $\Lambda_1^\epsilon(t)$, $\Lambda_2^\epsilon(t)$, $\Lambda_3^\epsilon(t)$, $\Lambda_4^\epsilon(t)$, respectively. We fix a $t \in \mathbb{R}_+$.

Firstly, we prove the uniform integrability of the set $\{e^{\Lambda_4^\epsilon(t)} | 0 < \epsilon \leq 1\}$ for each $t \in \mathbb{R}_+$. This follows by proving that it is uniformly bounded in $L^2(\Omega, \mathbf{P})$. Denoting the integrand in $\Lambda_4^\epsilon(\hat{e}, \tilde{N}^\alpha)$ by

$$\Gamma^\epsilon(s, y) := \epsilon [\hat{e}_\epsilon(X_{s-}^{x, \epsilon} + \sigma_\epsilon(X_{s-}^{x, \epsilon}, y)) - \hat{e}_\epsilon(X_{s-}^{x, \epsilon})].$$

Then by Itô's formula,

$$\begin{aligned} e^{2\Lambda_4^\epsilon(t)} &= 1 - \int_0^t \int_{B^c} 2e^{2\Lambda_4^\epsilon(s-)} \Gamma^\epsilon(s, y) \nu^\alpha(dy) ds \\ &\quad + \int_0^t \int_{B \setminus \{0\}} e^{2\Lambda_4^\epsilon(s-)} \left(e^{2\Gamma^\epsilon(s, y)} - 1 \right) \tilde{N}^\alpha(dy, ds) \\ &\quad + \int_0^t \int_{B^c} e^{2\Lambda_4^\epsilon(s-)} \left(e^{2\Gamma^\epsilon(s, y)} - 1 \right) N^\alpha(dy, ds) \\ &\quad + \int_0^t \int_{B \setminus \{0\}} e^{2\Lambda_4^\epsilon(s-)} \left[e^{2\Gamma^\epsilon(s, y)} - 1 - 2\Gamma^\epsilon(s, y) \right] \nu^\alpha(dy) ds. \end{aligned}$$

Since $\hat{e}$ is bounded, Γ^ϵ has a uniform bound for all $\epsilon > 0$. Then there exists a large constant $C > 0$ such that for each $\epsilon > 0$ and $t \in \mathbb{R}_+$,

$$\mathbf{E} \int_0^t \int_{B^c} \left| e^{2\Lambda_4^\epsilon(s-)} \left(e^{2\Gamma^\epsilon(s, y)} - 1 - 2\Gamma^\epsilon(s, y) \right) \right|^2 \nu^\alpha(dy) ds \leq C \nu^\alpha(B^c) \mathbf{E} \int_0^t e^{2\Lambda_4^\epsilon(s-)} ds < \infty.$$

Hence combining these, there exists $\theta \in (0, 1)$ such that

$$\begin{aligned} \mathbf{E} e^{2\Lambda_4^\epsilon(t)} &= 1 + \mathbf{E} \int_0^t \int_{B \setminus \{0\}} e^{2\Lambda_4^\epsilon(s-)} \left[e^{2\Gamma^\epsilon(s, y)} - 1 - 2\Gamma^\epsilon(s, y) \right] \nu^\alpha(dy) ds \\ &\leq 1 + \mathbf{E} \int_0^t e^{2\Lambda_4^\epsilon(s-)} \int_{B \setminus \{0\}} 2e^{2\theta\Gamma^\epsilon(s, y)} |\Gamma^\epsilon(s, y)|^2 \nu^\alpha(dy) ds \\ &\leq 1 + C(\|\hat{e}\|_0) \mathbf{E} \int_0^t e^{2\Lambda_4^\epsilon(s-)} \int_{B \setminus \{0\}} |\Gamma^\epsilon(s, y)|^2 \nu^\alpha(dy) ds. \end{aligned}$$

As shown in the proof of part (iii) and (iv) in Lemma 5.3,

$$\int_{B \setminus \{0\}} |\Gamma^\epsilon(s, y)|^2 \nu^\alpha(dy) \leq \epsilon^2 \|\hat{e}\|_1^2 \int_{B \setminus \{0\}} |\sigma_\epsilon(X_{s-}^{x, \epsilon}, y)|^2 \nu^\alpha(dy) \leq \frac{|\mathbb{S}^{d-1}|}{2 - \alpha} \epsilon^2 \|\hat{e}\|_1^2 \|\psi\|_{L^\infty}^2.$$

Thus,

$$\mathbf{E} e^{2\Lambda_4^\epsilon(t)} \leq 1 + \epsilon^2 C(\alpha, |\mathbb{S}^{d-1}|, \|\hat{e}\|_1, \|\psi\|_{L^\infty}) \int_0^t \mathbf{E} e^{2\Lambda_4^\epsilon(s-)} ds.$$

By Grönwall's inequality, the uniform boundness of $\{e^{\Lambda_4^\epsilon(t)}|0 < \epsilon \leq 1\}$ in $L^2(\Omega, \mathbf{P})$ follows.

Secondly, the proof of Lemma 5.3 shows that the set $\{\Lambda_3^\epsilon(t) - \Lambda_2^\epsilon(t)|0 < \epsilon \leq 1\}$ is bounded. The set $\{\Lambda_1^\epsilon(t)|0 < \epsilon \leq 1\}$ is bounded by virtue of the boundness of c, g and $\hat{e}$. Also since u_0 is periodic and continuous, $\{u_0(X_t^{x,\epsilon})|0 < \epsilon \leq 1\}$ is bounded. Thus, the set $\{u_0(X_t^{x,\epsilon}) \exp(\hat{Y}_t^\epsilon)|0 < \epsilon \leq 1\}$ is uniformly integrable.

Finally, we pass to the limit. It is easy to see that $e^{\hat{Y}_t^\epsilon} \rightarrow e^{y(t)}$ in probability as $\epsilon \rightarrow 0$, by the continuous mapping theorem (see, for instance, [44, Theorem 2.3.(ii)]). Then for any subsequence $\{\epsilon_n\} \rightarrow 0$, there exists a subsubsequence $\{\epsilon_{n_k}\} \rightarrow 0$ such that $e^{\hat{Y}_t^{\epsilon_{n_k}}} \rightarrow e^{y(t)}$ almost uniformly (cf. [23, Lemma 4.2]). That is, for any $\rho > 0$, there exists a set $N \in \mathcal{F}$ with $\mathbf{P}(N) \leq \rho$, such that

$$\left\| e^{\hat{Y}_t^{\epsilon_{n_k}}} - e^{y(t)} \right\|_{L^\infty(N^c, \mathbf{P})} \rightarrow 0, \quad k \rightarrow \infty. \quad (5.5)$$

By the boundness of u_0 , we know the set $\{u_0(X_t^{x,\epsilon})[\exp(\hat{Y}_t^\epsilon) - \exp(y(t))]|0 < \epsilon \leq 1\}$ is also uniformly integrable. Then for any $\delta > 0$, there exist $\rho_0 > 0$ and $N_0 \in \mathcal{F}$ with $\mathbf{P}(N_0) \leq \rho_0$, such that

$$\mathbf{E} \left| u_0(X_t^{x,\epsilon}) \left(e^{\hat{Y}_t^\epsilon} - e^{y(t)} \right) \mathbf{1}_{N_0} \right| < \delta. \quad (5.6)$$

Now along the sequence $\{\epsilon_{n_k}\}$, we combining (5.5) with (5.6) to get

$$\begin{aligned} \mathbf{E} \left| u_0(X_t^{x,\epsilon_{n_k}}) \left(e^{\hat{Y}_t^{\epsilon_{n_k}}} - e^{y(t)} \right) \right| &\leq \mathbf{E} |\cdots \mathbf{1}_{N_0}| + \mathbf{E} |\cdots \mathbf{1}_{N_0^c}| \\ &\leq \delta + \|u_0\|_{L^\infty} \mathbf{P}(N_0^c) \left\| e^{\hat{Y}_t^{\epsilon_{n_k}}} - e^{y(t)} \right\|_{L^\infty(N^c, \mathbf{P})} \\ &\leq 2\delta. \end{aligned}$$

To summarize these together, for any subsequence $\{\epsilon_n\} \rightarrow 0$, there exists a subsubsequence $\{\epsilon_{n_k}\} \rightarrow 0$ such that

$$\mathbf{E} \left| u_0(X_t^{x,\epsilon_{n_k}}) \left(e^{\hat{Y}_t^{\epsilon_{n_k}}} - e^{y(t)} \right) \right| \rightarrow 0, \quad k \rightarrow \infty,$$

which implies that the convergence holds on the whole line $0 < \epsilon \leq 1$. On the other hand, by Proposition 5.2, we know that $\mathbf{E}[u_0(X_t^{x,\epsilon})] \rightarrow \mathbf{E}[u_0(X_t^x)]$ as $\epsilon \rightarrow 0$. The result (1.4) follows from

$$\begin{aligned} |\hat{u}^\epsilon(t, x) - u(t, x)| &= \left| \mathbf{E} \left[u_0(X_t^{x,\epsilon}) e^{\hat{Y}_t^\epsilon} \right] - \mathbf{E} [u_0(X_t^x)] e^{y(t)} \right| \\ &\leq \mathbf{E} \left| u_0(X_t^{x,\epsilon}) \left(e^{\hat{Y}_t^\epsilon} - e^{y(t)} \right) \right| + |\mathbf{E} [u_0(X_t^{x,\epsilon})] - \mathbf{E} [u_0(X_t^x)]| e^{y(t)}. \end{aligned}$$

□

Remark 5.5. We close this section by some comments for the proof of Theorem 1.1. In [31], the author applied Girsanov's transform to get rid of the stochastic integral term involved in $\hat{Y}^\epsilon$, since this term may not possess the uniform integrability. While in our case, since the stochastic integral term in Y_t^ϵ has an infinitesimal integrand $\Gamma^\epsilon(s, y)$, the uniform integrability of $\{\exp(\hat{Y}_t^\epsilon)|0 < \epsilon \leq 1\}$ is easier to treat.

A Proofs of auxiliary results

In this part, we provide missing proofs of some auxiliary results in our paper.

Proof of Lemma 3.1. Note that the nonlocal operator $\mathcal{A}^{\bar{\sigma}, \nu^\alpha}$ can be rewritten as the form (2.8). For $u \in \mathcal{C}_b^{1+\gamma}(\mathbb{R}^d)$, we have

$$|u(x+z) - u(x) - z \cdot \nabla u(x)| \leq |z| \int_0^1 |\nabla u(x+rz) - \nabla u(x)| dr \leq \frac{[\nabla u]_\gamma}{1+\gamma} |z|^{1+\gamma}. \quad (\text{A.1})$$

Then by (2.10), there exists a constant $C > 0$ such that

$$\begin{aligned} |\mathcal{L}^\alpha u(x)| &\leq \int_{B \setminus \{0\}} |u(x+z) - u(x) - z \cdot \nabla u(x)| \nu^{\bar{\sigma}, \alpha}(x, dz) \\ &\quad + \int_{B^c} |u(x+z) - u(x)| \nu^{\bar{\sigma}, \alpha}(x, dz) + |b(x) \cdot \nabla u(x)| \\ &\leq 2\|u\|_{1+\gamma} \left(\int_{\mathbb{R}^d \setminus \{0\}} (|z|^{1+\gamma} \wedge 1) \nu^{\bar{\sigma}, \alpha}(x, dz) + \|b\|_0 \right) \\ &\leq C\|u\|_{1+\gamma}. \end{aligned}$$

Based on this estimate, the rest of the proof is exactly the same as that of [36, Proposition 3.2], even though it is set up with $\bar{\sigma}(\cdot, y) \equiv y$ there. $\square$

Proof of Lemma 3.4. Using (A.1) with $\gamma = \alpha + \beta - 1$, one can find that for $u \in \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$,

$$|u(x + \bar{\sigma}(x, y)) - u(x) - \bar{\sigma}(x, y) \cdot \nabla u(x)| \leq \frac{1}{\alpha+\beta} [\nabla u]_{\alpha+\beta-1} |\bar{\sigma}(x, y)|^{\alpha+\beta}.$$

Combining this with (2.6), a straightforward application of the dominated convergence theorem yields that $\lim_{y \rightarrow x} \mathcal{L}^\alpha u(y) = \mathcal{L}^\alpha u(x)$ for any $u \in \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$ and $x \in \mathbb{T}^d$. This amounts to saying that $\mathcal{L}^\alpha(\mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)) \subset \mathcal{C}(\mathbb{T}^d)$. Therefore, the operator

$$\mathcal{L}^\alpha : \mathcal{C}(\mathbb{T}^d) \supset \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d) \rightarrow \mathcal{C}(\mathbb{T}^d)$$

is a densely defined unbounded operator on $\mathcal{C}(\mathbb{T}^d)$.

Now Lemma 3.1 implies that for any $\kappa > 0$ and $u \in \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$, $\|(\kappa - \mathcal{L}^\alpha)u\|_0 \geq \kappa\|u\|_0$, that is, $\mathcal{L}^\alpha$ is dissipative. By Proposition 3.2, we have $\mathcal{C}^\beta(\mathbb{T}^d) \subset (\kappa - \mathcal{L}^\alpha)(\mathcal{C}^{\alpha+\beta}(\mathbb{T}^d))$ for any $\kappa > 0$, which yields that the operator $\kappa - \mathcal{L}^\alpha$ has dense range in $\mathcal{C}(\mathbb{T}^d)$. In addition, $\mathcal{L}^\alpha$ satisfies the positive maximum principle, due to the equivalent form (2.8) of $\mathcal{A}^{\bar{\sigma}, \nu^\alpha}$ and Courrège's theorem (see [21, Corollary 4.5.14]). Now the final assertion follows from the celebrated Hille-Yosida-Ray Theorem (see, for instance, [14, Theorem 4.2.2]). $\square$

Proof of Lemma 3.5. The existence of solution of the martingale problem is in [29, Proposition 3]. Taking Lemma 3.4 into account, the uniqueness and the Feller property follow from [14, Theorem 4.4.1]. The existence of transition density and the two estimates can be found in [11, Theorem 1.5]. $\square$

Proof of Lemma 4.1. By the estimate in Corollary 3.8, we have

$$\kappa^{\frac{\alpha+\beta-1}{\alpha}} \|\nabla \hat{b}_\kappa\|_0 \leq C\|b\|_\beta, \quad \kappa > \kappa_*.$$

Now by choosing $\kappa > \kappa_* \vee (2C\|b\|_\beta)^{\frac{\alpha}{\alpha+\beta-1}}$, we get that $\|\nabla \hat{b}_\kappa\|_0 \leq \frac{1}{2}$. Thus

$$\frac{1}{2}|x_1 - x_2| \leq |\Phi_\kappa(x_1) - \Phi_\kappa(x_2)| \leq \frac{3}{2}|x_1 - x_2|,$$

i.e., Φ_κ is bi-Lipschitz. In particular, Φ_κ is a $\mathcal{C}^1$ -diffeomorphism. Moreover,

$$\nabla(\Phi_\kappa^{-1}) = \text{Inv} \circ \nabla\Phi_\kappa \circ \Phi_\kappa^{-1},$$

where the matrix inverse map $\text{Inv} : \text{GL}(\mathbb{R}^d) \rightarrow \text{GL}(\mathbb{R}^d)$ is of class $\mathcal{C}^\infty$. Note that $\nabla\Phi_\kappa$ is of class $\mathcal{C}^{\alpha+\beta-1}$, Φ_κ^{-1} is of class $\mathcal{C}^1$. It is easy to see that $\nabla(\Phi_\kappa^{-1})$ is of class $\mathcal{C}^{\alpha+\beta-1}$. The second conclusion of the lemma follows. $\square$

Proof of Corollary 4.3. By applying Itô's formula, it is easy to see that for any $f \in D(\mathcal{L}^\alpha) = \mathcal{C}^{\alpha+\beta}(\mathbb{T}^d)$, the following process is a $(\Omega, \mathcal{F}, \mathbf{P}, \{\mathcal{F}_t\}_{t \geq 0})$ -martingale

$$\tilde{M}^f(t) := f(\tilde{X}_t^x) - f(\tilde{X}_0^x) - \int_0^t \mathcal{L}^\alpha f(\tilde{X}_s^x) ds.$$

It is easy to see that $\tilde{X}^x$ has càdlàg paths almost surely. Let $\mathbf{P}_{\tilde{X}^x} := \mathbf{P} \circ \tilde{X}^x$ be the pushforward probability measure of $\tilde{X}^x$ on $(\mathcal{D}, \mathcal{B}(\mathcal{D}))$, then $\mathbf{P}_{\tilde{X}^x}$ is a solution of martingale problem for $(\mathcal{L}^\alpha, \delta_x)$. By the uniqueness of solutions to the martingale problem obtained in Lemma 3.5, we find that $\mathbf{P}_{\tilde{X}^x} = \mathbf{P}^x$, the Feller property follows. The strong Markov property follows from [37, Theorem III.3.1]. $\square$

Proof of Lemma 4.9. Since μ is invariant with respect to $\{P_t\}_{t \geq 0}$, for any $f \in \mathcal{C}_\mu(\mathbb{T}^d)$ and $t \geq 0$, we have

$$\int_{\mathbb{T}^d} P_t f(x) \mu(dx) = \int_{\mathbb{T}^d} f(x) \mu(dx) = 0.$$

That is, $\mathcal{C}_\mu(\mathbb{T}^d)$ is $\{P_t\}_{t \geq 0}$ -invariant, in the sense that $P_t(\mathcal{C}_\mu(\mathbb{T}^d)) \subset \mathcal{C}_\mu(\mathbb{T}^d)$ for all $t \geq 0$. The lemma then follows from the corollary in [13, Subsection II.2.3]. $\square$

Proof of Lemma 4.10. Since f is centered with respect to μ , by Lemma 4.6 we have

$$\|P_t f\|_0 \leq C \|f\|_0 e^{-\rho t}. \quad (\text{A.2})$$

Note the fact that μ is invariant with respect to $\{P_t\}_{t \geq 0}$. Then combining (A.2) and the representation (3.6), a straightforward application of Fubini's theorem implies that

$$\begin{aligned} \int_{\mathbb{T}^d} u_\kappa(x) \mu(dx) &= \int_{\mathbb{T}^d} \int_0^\infty e^{-\kappa t} P_t f(x) dt \mu(dx) = \int_0^\infty e^{-\kappa t} \left(\int_{\mathbb{T}^d} P_t f(x) \mu(dx) \right) dt \\ &= \int_0^\infty e^{-\kappa t} \left(\int_{\mathbb{T}^d} f(x) \mu(dx) \right) dt = 0. \end{aligned}$$

That is, u_κ is also centered with respect to μ . $\square$

Proof of Lemma 5.3. (i). By Proposition 4.8, the convergence in probability of the first integral is immediate,

$$\sup_{0 \leq s \leq t} |\Lambda_1^\epsilon(c)_s - \bar{C}s| \leq \int_0^t \left| (I + \nabla \hat{b})c\left(\frac{X_s^{x,\epsilon}}{\epsilon}\right) - \bar{C} \right| ds \rightarrow 0, \quad \epsilon \rightarrow 0.$$

(ii). Note that $\nu^\alpha(\epsilon A) = \epsilon^{-\alpha} \nu^\alpha(A)$, $A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})$. Then a change of variable yields

$$\begin{aligned} &\mathcal{A}^{\sigma_\epsilon, \nu^\alpha} \hat{b}_\epsilon(x) \\ &= \frac{1}{\epsilon^{\alpha-1}} \int_{\mathbb{R}^d \setminus \{0\}} \left[\hat{b}\left(\frac{x}{\epsilon} + \frac{1}{\epsilon} \sigma\left(\frac{x}{\epsilon}, \epsilon y\right)\right) - \hat{b}\left(\frac{x}{\epsilon}\right) - \frac{1}{\epsilon} \sigma^i\left(\frac{x}{\epsilon}, \epsilon y\right) \partial_i \hat{b}\left(\frac{x}{\epsilon}\right) \mathbf{1}_B(\epsilon y) \right] \nu^\alpha(dy). \end{aligned}$$

By the oddness condition in Assumption [H3](#), for any $\delta > 0$, we have

$$\begin{aligned}
& \left| \epsilon \mathcal{A}^{\sigma_\epsilon, \nu^\alpha} \hat{b}_\epsilon(x) - \frac{1}{\epsilon^{\alpha-1}} \mathcal{A}^{\bar{\sigma}, \nu^\alpha} \hat{b}\left(\frac{x}{\epsilon}\right) \right| \\
& \leq \frac{1}{\epsilon^{\alpha-1}} \int_{B_\delta \setminus \{0\}} \left| \left[\hat{b}\left(\frac{x}{\epsilon} + \frac{1}{\epsilon} \sigma\left(\frac{x}{\epsilon}, \epsilon y\right)\right) - \hat{b}\left(\frac{x}{\epsilon} + \bar{\sigma}\left(\frac{x}{\epsilon}, y\right)\right) \right. \right. \\
& \quad \left. \left. - \left(\frac{1}{\epsilon} \sigma^i\left(\frac{x}{\epsilon}, \epsilon y\right) - \bar{\sigma}^i\left(\frac{x}{\epsilon}, y\right)\right) \partial_i \hat{b}\left(\frac{x}{\epsilon}\right) \right] \right| \nu^\alpha(dy) \\
& \quad + \frac{1}{\epsilon^{\alpha-1}} \int_{B_\delta^c} \left| \left[\hat{b}\left(\frac{x}{\epsilon} + \frac{1}{\epsilon} \sigma\left(\frac{x}{\epsilon}, \epsilon y\right)\right) - \hat{b}\left(\frac{x}{\epsilon} + \bar{\sigma}\left(\frac{x}{\epsilon}, y\right)\right) \right] \right| \nu^\alpha(dy) \\
& =: I_1^\epsilon(x) + I_2^\epsilon(x),
\end{aligned} \tag{A.3}$$

where B_δ is the open ball in $\mathbb{R}^d$ centering at the origin with radius δ , I_1^ϵ and I_2^ϵ are the two integral terms in the second equality. We let $\delta = \epsilon^\gamma$ for some $\gamma \in \mathbb{R}$ that will be chosen latter. It follows from Lemma [5.1](#) that

$$I_1^\epsilon(x) \leq \frac{\|\hat{b}\|_{\alpha+\beta}}{\alpha+\beta} \frac{1}{\epsilon^{\alpha-1}} \int_{B_\delta \setminus \{0\}} \left| \frac{1}{\epsilon} \sigma\left(\frac{x}{\epsilon}, \epsilon y\right) - \bar{\sigma}\left(\frac{x}{\epsilon}, y\right) \right|^{\alpha+\beta} \nu^\alpha(dy).$$

Recall that $\sigma(x, 0) = 0$ and $\bar{\sigma}(x, y) = \nabla_y \sigma(x, 0)y$ as in [\(2.15\)](#) and [\(2.14\)](#). Then by [\(A.1\)](#),

$$\left| \frac{1}{\epsilon} \sigma\left(\frac{x}{\epsilon}, \epsilon y\right) - \bar{\sigma}\left(\frac{x}{\epsilon}, y\right) \right| = \frac{1}{\epsilon} \left| \sigma\left(\frac{x}{\epsilon}, \epsilon y\right) - \sigma\left(\frac{x}{\epsilon}, 0\right) - \epsilon \nabla_y \sigma\left(\frac{x}{\epsilon}, 0\right) y \right| \leq \frac{\epsilon^\lambda [\nabla_y \sigma(\frac{x}{\epsilon}, \cdot)]_\lambda}{1+\lambda} |y|^{1+\lambda}.$$

Hence, using the assumption [\(2.4\)](#), we have

$$\begin{aligned}
I_1^\epsilon(x) & \leq \frac{\|\hat{b}\|_{\alpha+\beta} \sup_{x \in \mathbb{T}^d} [\nabla_y \sigma(x, \cdot)]_\lambda^{\alpha+\beta}}{(\alpha+\beta)(1+\lambda)^{\alpha+\beta}} \epsilon^{\lambda(\alpha+\beta)-\alpha+1} \int_{B_\delta \setminus \{0\}} |y|^{(1+\lambda)(\alpha+\beta)} \nu^\alpha(dy) \\
& \leq C(d, \alpha, \beta, \lambda, \|\hat{b}\|_{\alpha+\beta}, \sup_{x \in \mathbb{T}^d} [\nabla_y \sigma(x, \cdot)]_\lambda) \epsilon^{\lambda(\alpha+\beta)-\alpha+1+\gamma[(1+\lambda)(\alpha+\beta)-\alpha]}.
\end{aligned} \tag{A.4}$$

On the other hand, using a change of variable once again,

$$\begin{aligned}
I_2^\epsilon(x) & = \epsilon \int_{B_{\epsilon\delta}^c} \left| \left[\hat{b}\left(\frac{x}{\epsilon} + \frac{1}{\epsilon} \sigma\left(\frac{x}{\epsilon}, y\right)\right) - \hat{b}\left(\frac{x}{\epsilon} + \bar{\sigma}\left(\frac{x}{\epsilon}, \frac{y}{\epsilon}\right)\right) \right] \right| \nu^\alpha(dy) \leq 2\|\hat{b}\|_0 \epsilon \nu^\alpha(B_{\epsilon\delta}^c) \\
& \leq C(d, \|\hat{b}\|_0) \epsilon^{1-\alpha(1+\gamma)}.
\end{aligned} \tag{A.5}$$

Now we select γ satisfying

$$-\frac{\lambda(\alpha+\beta)-\alpha+1}{(1+\lambda)(\alpha+\beta)-\alpha} < \gamma < \frac{1}{\alpha} - 1, \tag{A.6}$$

One can easily verify that the interval in [\(A.6\)](#) is not null, since $\lambda > \alpha - 1$ as assumed in the regularity condition in [H3](#). Combining [\(A.3\)](#)-[\(A.6\)](#), we conclude that

$$\sup_{x \in \mathbb{T}^d} \left| \epsilon \mathcal{A}^{\sigma_\epsilon, \nu^\alpha} \hat{b}_\epsilon(x) - \frac{1}{\epsilon^{\alpha-1}} \mathcal{A}^{\bar{\sigma}, \nu^\alpha} \hat{b}\left(\frac{x}{\epsilon}\right) \right| \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0.$$

Therefore,

$$\begin{aligned}
\sup_{0 \leq s \leq t} \left| \Lambda_3^\epsilon(\hat{b}, \mathcal{A}^{\sigma_\epsilon, \nu^\alpha})_s - \Lambda_2^\epsilon(\hat{b}, \mathcal{A}^{\bar{\sigma}, \nu^\alpha})_s \right| & \leq \int_0^t \left| \epsilon \mathcal{A}^{\sigma_\epsilon, \nu^\alpha} \hat{b}_\epsilon(X_s^{x, \epsilon}) - \frac{1}{\epsilon^{\alpha-1}} \mathcal{A}^{\bar{\sigma}, \nu^\alpha} \hat{b}\left(\frac{X_s^{x, \epsilon}}{\epsilon}\right) \right| ds \\
& \leq t \sup_{x \in \mathbb{T}^d} \left| \epsilon \mathcal{A}^{\sigma_\epsilon, \nu^\alpha} \hat{b}_\epsilon(x) - \frac{1}{\epsilon^{\alpha-1}} \mathcal{A}^{\bar{\sigma}, \nu^\alpha} \hat{b}\left(\frac{x}{\epsilon}\right) \right| \\
& \rightarrow 0, \quad \epsilon \rightarrow 0.
\end{aligned}$$

(iii) and (iv). Since Ξ^ϵ is integrable with respect to $\tilde{N}^\alpha$, the third characteristic of Λ_2^ϵ satisfies that $\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} (|x|^2 \wedge 1) \nu_2^\epsilon(dx, ds) < \infty$ for each $\epsilon > 0$ and $t \in \mathbb{R}_+$ (cf. [22, Proposition II.2.9]). By the hypothesis, there exist $\rho > 0$ and $M > 0$ such that $|f| \leq M$ on B_ρ^c and $f = 0$ on B_ρ . Then for any $t \in \mathbb{R}_+$,

$$\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(x) \nu_2^\epsilon(dx, ds) \leq M \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \mathbf{1}_{B_\rho^c}(x) \nu_2^\epsilon(dx, ds),$$

which goes to zero almost surely as $\epsilon \rightarrow 0$ by the boundness of $\hat{b}$ and the dominated convergence theorem, and (iv) follows.

For B_2^ϵ , we have the estimate

$$\begin{aligned} \sup_{0 \leq s \leq t} |B_2^\epsilon(s)| &\leq \left[\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} |x|^2 \nu_2^\epsilon(dx, ds) \right]^{\frac{1}{2}} \left[\int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \mathbf{1}_{B^c}(x) \nu_2^\epsilon(dx, ds) \right]^{\frac{1}{2}} \\ &=: \sqrt{J_1^\epsilon} \cdot \sqrt{J_2^\epsilon}. \end{aligned}$$

By (iv) and a usual approximation procedure, J_2^ϵ goes to zero surely as $\epsilon \rightarrow 0$. For J_1^ϵ ,

$$\begin{aligned} J_1^\epsilon &= \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} \left| \epsilon \left[\hat{b}_\epsilon(X_{s-}^{x,\epsilon} + \sigma_\epsilon(X_{s-}^{x,\epsilon}, y)) - \hat{b}_\epsilon(X_{s-}^{x,\epsilon}) \right] \right|^2 \nu^\alpha(dy) ds \\ &= \int_0^t \left(\int_{B_\epsilon^c} + \int_{B_\epsilon \setminus \{0\}} \right) |\dots|^2 \nu^\alpha(dy) ds \\ &\leq \frac{4t \|\hat{b}\|_0^2 |\mathbb{S}^{d-1}|}{\alpha} \epsilon^{2-\alpha} + \|\hat{b}\|_1^2 \int_{B_\epsilon \setminus \{0\}} \int_0^t \left| \sigma\left(\frac{X_{s-}^{x,\epsilon}}{\epsilon}, y\right) \right|^2 ds \nu^\alpha(dy). \end{aligned}$$

By the growth condition in Assumption H3,

$$\int_{B_\epsilon \setminus \{0\}} \int_0^t \left| \sigma\left(\frac{X_{s-}^{x,\epsilon}}{\epsilon}, y\right) \right|^2 ds \nu^\alpha(dy) \leq \frac{t |\mathbb{S}^{d-1}| \epsilon^{2-\alpha}}{2-\alpha} \int_0^t \left| \phi\left(\frac{X_{s-}^{x,\epsilon}}{\epsilon}\right) \right|^2 ds.$$

Then (iii) follows from these estimates and Proposition 4.8.

(v). It follows from Proposition 4.8 that,

$$\begin{aligned} \int_0^t \int_{\mathbb{R}^d \setminus \{0\}} f(y) \nu_\epsilon^{3+4}(dy, ds) &= \int_{\mathbb{R}^d \setminus \{0\}} \int_0^t f\left(\sigma\left(\frac{X_{s-}^{x,\epsilon}}{\epsilon}, y\right)\right) ds \nu^\alpha(dy) \\ &\rightarrow t \int_{\mathbb{R}^d \setminus \{0\}} \int_{\mathbb{T}^d} f(\sigma(x, y)) \mu(dx) \nu^\alpha(dy) \\ &= t \int_{\mathbb{R}^d \setminus \{0\}} f(y) \bar{\nu}(dy), \quad \epsilon \rightarrow 0, \end{aligned}$$

where the convergence is in probability. □

Acknowledgements

We would like to thank the reviewers for their thoughtful comments and efforts towards improving our manuscript. The research of J. Duan was partly supported by the NSF grant 1620449. The research of Q. Huang was partly supported by China Scholarship Council (CSC), and NSFC grants 11531006 and 11771449. The research of R. Song is supported in part by a grant from the Simons Foundation (# 429343, Renming Song).

References

- [1] G. Allaire. *Shape optimization by the homogenization method*, vol. 146. Springer-Verlag New York, 2002.
- [2] D. Applebaum. *Lévy processes and stochastic calculus*, vol. 116. Cambridge University Press, 2009.
- [3] M. Arisawa. Homogenization of a class of integro-differential equations with Lévy operators. *Commun. Partial Differ. Equ.*, 34(7):617–624, 2009.
- [4] G. Bal and W. Jing. Fluctuations in the homogenization of semilinear equations with random potentials. *Commun. Partial Differ. Equ.*, 41(12):1839–1859, 2016.
- [5] M. Barczy, Z. Li, and G. Pap. Yamada-watanabe results for stochastic differential equations with jumps. *Int. J. Stoch. Anal.*, 2015:1–23, 2015.
- [6] R.F. Bass. Regularity results for stable-like operators. *J. Funct. Anal.*, 8(257):2693–2722, 2009.
- [7] P. Bella, B. Fehrman, J. Fischer, and F. Otto. Stochastic homogenization of linear elliptic equations: Higher-order error estimates in weak norms via second-order correctors. *SIAM J. Math. Anal.*, 49(6):4658–4703, 2017.
- [8] A. Benchérif-Madani and É. Pardoux. Homogenization of a semilinear parabolic PDE with locally periodic coefficients: a probabilistic approach. *ESAIM-Prob. Stat.*, 11:385–411, 2007.
- [9] A. Bensoussan, J.-L. Lions, and G. Papanicolaou. *Asymptotic analysis for periodic structures*, vol. 5. North-Holland Publishing Company Amsterdam, 1978.
- [10] H. Bessaih, E. Hausenblas, and P. Razafimandimby. Strong solutions to stochastic hydrodynamical systems with multiplicative noise of jump type. *NoDea-Nonlinear Differ. Equ. Appl.*, 22(6):1661–1697, 2015.
- [11] Z.-Q. Chen and X. Zhang. Heat kernels for time-dependent non-symmetric stable-like operators. *J. Math. Anal. Appl.*, 465(1):1–21, 2018.
- [12] D. Cioranescu and J.S.J. Paulin. *Homogenization of reticulated structures*, vol. 136. Springer-Verlag New York, 1999.
- [13] K.-J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*, vol. 194. Springer-Verlag New York, 2000.
- [14] S.N. Ethier and T.G. Kurtz. *Markov processes: characterization and convergence*, vol. 282. John Wiley & Sons, 2009.
- [15] B. Franke. The scaling limit behaviour of periodic stable-like processes. *Bernoulli*, 12(3):551–570, 2006.
- [16] B. Franke. A functional non-central limit theorem for jump-diffusions with periodic coefficients driven by stable Lévy-noise. *J. Theor. Probab.*, 20(4):1087–1100, 2007.
- [17] D. Gilbarg and N.S. Trudinger. *Elliptic partial differential equations of second order*, vol. 224. Springer Science & Business Media, 2001.
- [18] M. Hairer and É. Pardoux. Homogenization of periodic linear degenerate PDEs. *J. Funct. Anal.*, 255(9):2462–2487, 2008.
- [19] M. Horie, T. Inuzuka, and H. Tanaka. Homogenization of certain one-dimensional discontinuous markov processes. *Hiroshima Math. J.*, 7(2):629–641, 1977.
- [20] N. Ikeda and S. Watanabe. *Stochastic differential equations and diffusion processes*, vol. 24. Elsevier, 2014.
- [21] N. Jacob. *Pseudo differential operators & Markov processes: Fourier analysis and semigroups*, vol. 1. Imperial College Press, 2001.
- [22] J. Jacod and A. Shiryaev. *Limit theorems for stochastic processes, 2ed*, vol. 288. Springer-Verlag, 2013.
- [23] O. Kallenberg. *Foundations of modern probability*. Springer Science & Business Media, 2006.

- [24] M. Kassmann, A. Piatnitski, and E. Zhizhina. Homogenization of lévy-type operators with oscillating coefficients. *SIAM J. Math. Anal.*, 51(5):3641–3665, 2019.
- [25] S.G. Krantz and H.R. Parks. *The implicit function theorem: history, theory, and applications*. Springer Science & Business Media, 2012.
- [26] T.G. Kurtz. Equivalence of stochastic equations and martingale problems. In *Stochastic analysis 2010*, pp. 113–130. Springer, 2011.
- [27] C. Le Bris, F. Legoll, and F. Madiot. Multiscale finite element methods for advection-dominated problems in perforated domains. *Multiscale Model. Simul.*, 17(2):773–825, 2019.
- [28] H. Masuda. Ergodicity and exponential β -mixing bounds for multidimensional diffusions with jumps. *Stoch. Process. Their Appl.*, 117(1):35–56, 2007.
- [29] R. Mikulevičius and H. Pragarauskas. On the cauchy problem for integro-differential operators in sobolev classes and the martingale problem. *J. Differ. Equ.*, 256(4):1581–1626, 2014.
- [30] O.A Oleinik, A.S. Shamaev, and G.A. Yosifian. *Mathematical problems in elasticity and homogenization*, vol. 26. North-Holland, 1992.
- [31] É. Pardoux. Homogenization of linear and semilinear second order parabolic PDEs with periodic coefficients: a probabilistic approach. *J. Funct. Anal.*, 167(2):498–520, 1999.
- [32] É. Pardoux and A. Piatnitski. Homogenization of a singular random one-dimensional PDE with time-varying coefficients. *Ann. Probab.*, 40(3):1316–1356, 2012.
- [33] É. Pardoux and A. Răşcanu. *Stochastic differential equations, backward SDEs, partial differential equations*, vol. 69. Springer, 2014.
- [34] É. Pardoux and A.Y. Veretennikov. On the poisson equation and diffusion approximation. I. *Ann. Probab.*, 29(3):1061–1085, 2001.
- [35] A. Piatnitski and E. Zhizhina. Periodic homogenization of nonlocal operators with a convolution-type kernel. *SIAM J. Math. Anal.*, 49(1):64–81, 2017.
- [36] E. Priola. Pathwise uniqueness for singular SDEs driven by stable processes. *Osaka J. Math.*, 49(2):421–447, 2012.
- [37] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, vol. 293. Springer Science & Business Media, 2013.
- [38] R. Rhodes and B.A. Sow. Critical homogenization of Lévy process driven SDEs in random medium. *Stoch. Anal. Appl.*, 29(5):838–859, 2011.
- [39] Y. Sakamoto and N. Yoshida. Asymptotic expansion formulas for functionals of ϵ -Markov processes with a mixing property. *Ann. Inst. Stat. Math.*, 56(3):545–597, 2004.
- [40] N. Sandrić. Homogenization of periodic diffusion with small jumps. *J. Math. Anal. Appl.*, 435(1):551–577, 2016.
- [41] J.J. Schäffer. Norms and determinants of linear mappings. *Math. Z.*, 118(4):331–339, 1970.
- [42] R.W. Schwab. Periodic homogenization for nonlinear integro-differential equations. *SIAM J. Math. Anal.*, 42(6):2652–2680, 2010.
- [43] M. Tomisaki. Homogenization of cadlag processes. *J. Math. Soc. Jpn.*, 44(2):281–305, 1992.
- [44] A.W. Van der Vaart. *Asymptotic statistics*, vol. 3. Cambridge University Press, 1998.
- [45] L. Xie and X. Zhang. Heat kernel estimates for critical fractional diffusion operator. *Stud. Math.*, 224(3):221–263, 2014.
- [46] K. Yosida. *Functional analysis. Reprint of the sixth (1980) edition. Classics in Mathematics*, vol. 11. Springer-Verlag, Berlin, 1995.
- [47] A.K. Zvonkin. A transformation of the phase space of a diffusion process that removes the drift. *Math. USSR Sb.*, 22(1):129, 1974.