

# THE SMALL-MASS LIMIT AND WHITE-NOISE LIMIT OF AN INFINITE DIMENSIONAL GENERALIZED LANGEVIN EQUATION

HUNG D. NGUYEN<sup>1</sup>

**ABSTRACT.** We study asymptotic properties of the Generalized Langevin Equation (GLE) in the presence of a wide class of external potential wells with a power-law decay memory kernel. When the memory can be expressed as a sum of exponentials, a class of Markovian systems in infinite-dimensional spaces is used to represent the GLE. The solutions are shown to converge in probability in the small-mass limit and the white-noise limit to appropriate systems under minimal assumptions, of which no global Lipschitz condition is required on the potentials. With further assumptions about space regularity and potentials, we obtain  $L^1$  convergence in the white-noise limit.

**Keywords:** Markov processes, power-law decay, memory kernel

## 1. INTRODUCTION

The Generalized Langevin Equation is a Stochastic Integro-Differential Equation that is commonly used to model the velocity  $\{v(t)\}_{t \geq 0}$  of a microparticle in a thermally fluctuating viscoelastic fluid [31, 25, 18]. It can be written in the following form

$$\begin{aligned} \dot{x}(t) &= v(t), \\ m \dot{v}(t) &= -\gamma v(t) - \Phi'(x(t)) - \int_{-\infty}^t K(t-s)v(s) ds + F(t) + \sqrt{2\gamma} \dot{W}(t), \end{aligned} \quad (1.1)$$

where  $m > 0$  is the particle mass,  $\gamma > 0$  is the viscous drag coefficient,  $\Phi$  is a potential well and  $K(t)$  is a phenomenological memory kernel that summarizes the delayed drag effects by the fluid on the particle. The noise has two components:  $F(t)$  is a mean-zero, stationary Gaussian process with autocovariance  $\mathbb{E}[F(t)F(s)] = K(|t-s|)$ , and  $W(t)$  is a standard two-sided Brownian motion. The appearance of  $K(t)$  in the autocovariance of  $F(t)$  is a manifestation of the Fluctuation-Dissipation relationship, originally stated in [27], see also [36] for a more systematic review.

When there are no external forces, the GLE has the form

$$\begin{aligned} \dot{x}(t) &= v(t), \\ m \dot{v}(t) &= -\gamma v(t) - \int_{-\infty}^t K(t-s)v(s) ds + \sqrt{2\gamma} \dot{W}(t), \end{aligned} \quad (1.2)$$

it was shown in [32] that with extra assumptions, when  $K$  is integrable, the *Mean-Squared Displacement* (MSD)  $\mathbb{E}x(t)^2$  satisfies  $\mathbb{E}x(t)^2 \sim t$  as  $t \rightarrow \infty$ ; otherwise, if  $K(t) \sim t^{-\alpha}$ ,  $\alpha \in (0, 1)$ , then  $\mathbb{E}x(t)^2 \sim t^\alpha$  as  $t \rightarrow \infty$ . The former asymptotic behavior of the MSD is called *diffusive* whereas the latter is called *subdiffusive*. Here the notation  $f(t) \sim g(t)$  as  $t \rightarrow \infty$  means

$$\frac{f(t)}{g(t)} \rightarrow c \in (0, \infty), \quad \text{as } t \rightarrow \infty.$$

It has been observed that when  $K(t)$  is written as a sum of exponential functions, by adding auxiliary terms, the non-Markovian GLE (1.1) can be mapped onto a multi-dimensional Markovian system [33, 38, 28, 29]. If  $K(t)$  has the form of a finite sum of exponentials, the resulting finite-dimensional SDE was studied extensively in e.g. [34, 36]. One can show that these systems admit

<sup>1</sup> Department of Mathematics, Tulane University. 6823 St Charles Ave, New Orleans, LA, 70118.

a unique invariant structure with geometric ergodicity. Moreover, the marginal density of the pair  $(x, v)$  is independent of  $K(t)$ . It is also worthwhile to note that, in the case of the linear GLE (1.2), these memory kernels  $K(t)$  produce *diffusive* MSD since they are integrable [32]. In order to include memory kernels that have a power-law decay, one has to consider an infinite sum of exponentials resulting in a corresponding infinite-dimensional system.

From now on, we shall adopt the notations from [12]. Let  $\alpha, \beta > 0$  be given, and define constants  $c_k, \lambda_k, k = 1, 2, \dots$ , by

$$c_k = \frac{1}{k^{1+\alpha\beta}}, \quad \lambda_k = \frac{1}{k^\beta}. \quad (1.3)$$

We introduce the memory kernel  $K(t)$  given by

$$K(t) = \sum_{k \geq 1} c_k e^{-\lambda_k t}. \quad (1.4)$$

It is shown that (see Example 3.3 of [1]) with this choice of constants  $c_k$  and  $\lambda_k$ ,  $K(t)$  obeys a power-law decay, namely

$$K(t) \sim t^{-\alpha} \quad \text{as } t \rightarrow \infty, \quad (1.5)$$

where  $\alpha$  is as in (1.3). The constant  $\beta$  is an auxiliary parameter that is only assumed to be positive. When  $\alpha > 1$ , as mentioned above in the linear GLE (1.2),  $K(t)$  is in the *diffusive* regime, whereas for  $\alpha \in (0, 1)$ ,  $K$  belongs to the *subdiffusive* regime. There is however no claim regarding to the case  $\alpha = 1$ . For such reason, it is called the *critical regime*. With  $K(t)$  defined as in (1.4), the GLE (1.1) is expressed as the following infinite-dimensional system [12]

$$\begin{aligned} dx(t) &= v(t) dt, \\ m dv(t) &= (-\gamma v(t) - \Phi'(x(t)) - \sum_{k \geq 1} \sqrt{c_k} z_k(t)) dt + \sqrt{2\gamma} dW_0(t), \\ dz_k(t) &= (-\lambda_k z_k(t) + \sqrt{c_k} v(t)) dt + \sqrt{2\lambda_k} dW_k(t), \quad k \geq 1, \end{aligned} \quad (1.6)$$

where  $W_k$  are independent, standard Brownian motions. In [12], the well-posedness and the existence of invariant structures of (1.6) were studied for all  $\alpha > 0$ . Employing a recent advance tool called *asymptotic coupling* [16, 13], it can be shown that (1.6) admits a unique invariant distribution in the diffusive regime, ( $\alpha > 1$ ). However, ergodicity when  $\alpha \in (0, 1]$  remains an open question.

The goal of this note is to give an analysis of the behavior of (1.6) in two different limits.

First, we are interested in the asymptotic behavior of (1.6) concerning the small-mass limit, namely taking  $m$  to zero on the LHS of the second line in (1.6). Due to the random perturbations, the velocity  $v(t)$  is fluctuating fast whereas the displacement  $x(t)$  is still moving slow. We hence would like to find a process  $u(t)$  such that on any compact interval  $[0, T]$ ,

$$\lim_{m \downarrow 0} \sup_{0 \leq r \leq t} |x(r) - u(r)| = 0,$$

where the limit holds in an appropriate sense. Such statement is called Smoluchowski-Kramer approximation [10]. There is a literature of analyzing asymptotic behaviors for fast-slow processes when taking zero-mass limit. Earliest results in this direction seem to be the works of [26, 37]. For more recent studies in finite dimensional systems, we refer to [10] in which the convergence in probability is established with constant drag and multiplicative noise. Under stronger assumptions and using appropriate time rescaling, weak convergence is proved in [35] where the friction is also state-dependent. When the potential is assumed to be Lipschitz, one can obtain better results in  $L^p$ , following the works of [19, 30]. Without such assumption, convergence in probability is established in [17] provided appropriate Lyapunov controls. In addition, limiting systems are observed numerically in [20, 21]. Similar analysis in infinite dimensional settings for semi linear wave equations are studied in a series of paper [4, 5, 6, 7]. The systems therein are shown to converge to a heat equation under different assumptions about non linear drifts. Motivated by [17, 19], in

this note, we establish the convergence in probability for (1.6), cf. Theorem 2.4. The technique that we employ is inspired by those in [17].

Then, we study the white-noise limit of (1.6), namely as the random force  $F(t)$  in (1.1) converges to a white noise process. Under different conditions on the potential and space regularity, we aim to find a pair of processes  $(u(t), p(t))$  that can be approximated by the  $(x(t), v(t))$ -component in (1.6). While there is a rich history on the small-mass limit, the white-noise limit seems to receive less attention. Nevertheless, there have been many works on the asymptotics of deterministic systems with memories. To name a few in this direction, we refer the reader to [8, 11, 15]. With regards to the white-noise limit of our system, we establish the convergence in different modes for a wide class of potentials, that are not necessarily Lipschitz or bounded. While the proof of Theorem 2.5 concerning probability convergence shares the same arguments with that of Theorem 2.4, the result in Theorem 2.8 concerning strong topology requires more work, where we have to estimate a universal bound on the solutions of (1.6) using appropriate Lyapunov structures, cf. Proposition 4.1. To the best of our knowledge, these results seem to be new in infinite-dimensional stochastic differential equations with memory. Particularly, they (cf. Theorem 2.5 and Theorem 2.8) generalize analogous results for finite-dimensional settings in [34], where  $K(t)$  has a form of finite sum of exponentials.

The rest of this paper is organized as follows. We introduce notations and summarize our main results in Section 2. The small-mass limit is addressed rigorously in Section 3. We obtain a formula for the limiting system as a form of a Smoluchowski-Kramers equation. Finally, Section 4 studies the white-noise limit.

## 2. SUMMARY OF RESULTS

Throughout this work, we will assume that the potential  $\Phi$  satisfies the following growth condition [12].

**Assumption 2.1.**  $\Phi \in C^\infty(\mathbb{R})$  and there exists a constant  $c > 0$  such that for all  $x \in \mathbb{R}$

$$c(\Phi(x) + 1) \geq x^2.$$

By adding a positive constant if necessary, we also assume that  $\Phi$  is non-negative.

A typical class of potentials  $\Phi$  that satisfies Assumption 2.1 is the class of polynomials of even degree whose leading coefficient is positive. Functions that grow faster than polynomials are also included, e.g.  $e^{x^2}$ .

We now define a phase space for the infinite-dimensional process

$$X(t) = (x(t), v(t), z_1(t), z_2(t), \dots).$$

Following [12], let  $\mathcal{H}_{-s}$ ,  $s \in \mathbb{R}$  denote the Hilbert space given by

$$\mathcal{H}_{-s} = \left\{ X = (x, v, Z) = (x, v, z_1, z_2, \dots) : \|X\|_{-s}^2 = x^2 + v^2 + \sum_{k \geq 1} k^{-2s} z_k^2 < \infty \right\}. \quad (2.1)$$

endowed with the usual inner product  $\langle \cdot, \cdot \rangle_{-s}$ ,

$$\langle X, \tilde{X} \rangle_{-s} = x\tilde{x} + v\tilde{v} + \sum_{k \geq 1} k^{-2s} z_k \tilde{z}_k. \quad (2.2)$$

With regards to kernel parameters  $\alpha, \beta$  cf. (1.3), (1.4) and the phase space regularity parameter  $s$ , we assume that they satisfy the following condition.

**Assumption 2.2.** Let  $\alpha, \beta > 0$  be as in (1.3) and  $s \in \mathbb{R}$  as in (2.1). We assume that they satisfy either the asymptotically diffusive condition

$$(D) \quad \alpha > 1, \beta > \frac{1}{\alpha - 1} \text{ and } 1 < 2s < (\alpha - 1)\beta;$$

or the asymptotically subdiffusive condition

$$(SD) \quad 0 < \alpha < 1, \beta > \frac{1}{\alpha} \text{ and } 1 < 2s < \alpha\beta;$$

or the critical regime condition

$$(C) \quad \alpha = 1, \beta > 1 \text{ and } 1 < 2s < \beta;$$

Under Assumption 2.1 and Assumption 2.2, the well-posedness of (1.6) was studied rigorously in [12].

**2.1. Small-mass Limit.** In regards to the small-mass limit ( $m \rightarrow 0$ ), we introduce the following limiting system whose derivation will be explained later at the end of this subsection

$$\begin{aligned} \gamma du(t) &= \left( -\Phi'(u(t)) - \left( \sum_{k \geq 1} c_k \right) u(t) - \sum_{k \geq 1} \sqrt{c_k} f_k(t) \right) dt + \sqrt{2\gamma} dW_0(t), \\ df_k(t) &= \left( -\lambda_k f_k(t) - \lambda_k \sqrt{c_k} u(t) \right) dt + \sqrt{2\lambda_k} dW_k(t), \quad k = 1, 2, \dots \end{aligned} \quad (2.3)$$

The new phase space for the solution  $U(t) = (u(t), f_1(t), f_2(t), \dots)$  of (2.3) is denoted by  $\dot{\mathcal{H}}_{-s}$ ,  $s \in \mathbb{R}$ ,

$$\dot{\mathcal{H}}_{-s} = \left\{ U = (u, f_1, f_2, \dots) : \|U\|_{\dot{\mathcal{H}}_{-s}}^2 = u^2 + \sum_{k \geq 1} k^{-2s} f_k^2 < \infty \right\}, \quad (2.4)$$

endowed with the usual inner product. We can regard  $\dot{\mathcal{H}}_{-s}$  as a subspace of  $\mathcal{H}_{-s}$  whose  $v$ -component is equal to zero. Recalling  $c_k$  in (1.3), it is straightforward to see that if  $(x, v, z_1, z_2, \dots) \in \mathcal{H}_{-s}$  then  $(x, z_1 - \sqrt{c_1}x, z_2 - \sqrt{c_2}x, \dots) \in \dot{\mathcal{H}}_{-s}$ .

From now on, we shall fix a stochastic basis  $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, W)$  satisfying the usual conditions [24]. Here  $W$  is the cylindrical Wiener process defined on an auxiliary Wiener space  $\mathcal{W}$  with the usual decomposition

$$W(t) = e_0^{\mathcal{W}} W_0(t) + e_1^{\mathcal{W}} W_1(t) + \dots,$$

where  $\{e_0^{\mathcal{W}}, e_1^{\mathcal{W}}, \dots\}$  is the canonical basis of  $\mathcal{W}$ , and  $\{W_k(t)\}_{k \geq 0}$  are independent one-dimensional Brownian Motions [9]. The well-posedness of (2.3) is guaranteed by the following result.

**Proposition 2.3.** *Suppose that  $\Phi$  satisfies Assumption 2.1 and the constants  $\alpha, \beta, s$  satisfy Assumption 2.2. Then for all initial conditions  $U_0 \in \dot{\mathcal{H}}_{-s}$ , there exists a unique pathwise solution  $U(\cdot, U_0) : \Omega \times [0, \infty) \rightarrow \dot{\mathcal{H}}_{-s}$  of (2.3) in the following sense:  $U(\cdot, U_0)$  is  $\mathcal{F}_t$ -adapted,  $U(\cdot, U_0) \in C([0, \infty), \dot{\mathcal{H}}_{-s})$  almost surely and that if  $\tilde{U}(\cdot, U_0)$  is another solution then for every  $T \geq 0$ ,*

$$\mathbb{P} \left\{ \forall t \in [0, T], U(t, U_0) = \tilde{U}(t, U_0) \right\} = 1.$$

Moreover, for every  $U_0 \in \dot{\mathcal{H}}_{-s}$  and  $T \geq 0$ , there exists a constant  $C(T, U_0) > 0$  such that

$$\mathbb{E} \sup_{0 \leq t \leq T} \|U(t)\|_{\dot{\mathcal{H}}_{-s}}^2 \leq C(T, U_0). \quad (2.5)$$

The proof of Proposition 2.3 is quite standard, similar to that of Proposition 6 in [12] and will be briefly explained in Section 3. We now state the main result concerning the small-mass limit.

**Theorem 2.4.** *Suppose that  $\Phi$  satisfies Assumption 2.1 and the constants  $\alpha, \beta, s$  satisfy Assumption 2.2. Let  $X_m(t) = (x_m(t), v_m(t), z_{1,m}(t), \dots)$  solve (1.6) with initial conditions*

$$(x_m(0), v_m(0), z_{1,m}(0), z_{2,m}(0), \dots) = (x, v, z_1, z_2, \dots) \in \mathcal{H}_{-s},$$

*and  $U(t) = (u(t), f_1(t), \dots)$  solve (2.3) with initial conditions*

$$(u(0), f_1(0), f_2(0), \dots) = (x, z_1 - \sqrt{c_1}x, z_2 - \sqrt{c_2}x, \dots) \in \dot{\mathcal{H}}_{-s}.$$

Then, for every  $T, \xi > 0$ , it holds that

$$\mathbb{P}\left\{\sup_{0 \leq t \leq T} |x_m(t) - u(t)| > \xi\right\} \rightarrow 0, \quad m \rightarrow 0.$$

It is worthwhile to note that the small-mass limit in Theorem 2.4 holds for all regimes ( $\alpha > 0$ ) as stated in Assumption 2.2. We will see later that for the white-noise limit, the result is limited to the diffusive regime, namely  $\alpha > 1$ , see Theorem 2.5 below. We finish this subsection by a heuristic argument explaining how we derive the limiting system (2.3) and its initial conditions as stated in Theorem 2.4. Following [17], to determine the limiting system, one may formally set  $m = 0$  on the RHS of the second equation in (1.6) and substitute  $v(t)$  by  $dx(t)$  from the first equation to obtain

$$\gamma dx(t) = \left[-\Phi'(x(t)) - \sum_{k \geq 1} \sqrt{c_k} z_k(t)\right] dt + \sqrt{2\gamma} dW_0(t).$$

The equation on  $z_k(t)$  in (1.6) still depends on  $v(t)$ , but this can be circumvented by using Duhamel's formula,

$$z_k(t) = e^{-\lambda_k t} z_k(0) + \sqrt{c_k} \int_0^t e^{-\lambda_k(t-r)} v(r) dr + \sqrt{2\lambda_k} \int_0^t e^{-\lambda_k(t-r)} dW_k(r).$$

By an integration by parts, we can transform the integral term involving  $v(r)$  to

$$\int_0^t e^{-\lambda_k(t-r)} v(r) dr = x(t) - e^{-\lambda_k t} x(0) - \lambda_k \int_0^t e^{-\lambda_k(t-r)} x(r) dr.$$

Plugging back into the formula for  $z_k(t)$ , we find

$$\begin{aligned} z_k(t) &= e^{-\lambda_k t} (z_k(0) - \sqrt{c_k} x(0)) + \sqrt{c_k} x(t) - \lambda_k \sqrt{c_k} \int_0^t e^{-\lambda_k(t-r)} x(r) dr \\ &\quad + \sqrt{2\lambda_k} \int_0^t e^{-\lambda_k(t-r)} dW_k(r). \end{aligned}$$

We now set  $f_k(t) := z_k(t) - \sqrt{c_k} x(t)$  and  $u(t) := x(t)$  and thus arrive at (2.3) with the new shifted initial conditions as in Theorem 2.4.

**2.2. White-noise Limit.** Next, we present our results on the asymptotical behavior of the  $(x(t), v(t))$ -component of (1.6) in the diffusive regime by an appropriate scaling on the memory kernel  $K(t)$  in (1.4). For  $\epsilon > 0$ , we introduce  $K_\epsilon(t)$  given by

$$K_\epsilon(t) = \frac{1}{\epsilon} K\left(\frac{t}{\epsilon}\right) = \sum_{k \geq 1} \frac{c_k}{\epsilon} e^{-\frac{\lambda_k}{\epsilon} |t|}. \quad (2.6)$$

With  $K_\epsilon$  defined above, the corresponding system (1.6) becomes

$$\begin{aligned} dx_\epsilon(t) &= v_\epsilon(t) dt, \\ m dv_\epsilon(t) &= \left(-\gamma v_\epsilon(t) - \Phi'(x_\epsilon(t)) - \sum_{k \geq 1} \sqrt{\frac{c_k}{\epsilon}} z_{k,\epsilon}(t)\right) dt + \sqrt{2\gamma} dW_0(t), \\ dz_{k,\epsilon}(t) &= \left(-\frac{\lambda_k}{\epsilon} z_{k,\epsilon}(t) + \sqrt{\frac{c_k}{\epsilon}} v_\epsilon(t)\right) dt + \sqrt{\frac{2\lambda_k}{\epsilon}} dW_k(t), \quad k \geq 1. \end{aligned} \quad (2.7)$$

Inspired by [34], we consider the following system

$$\begin{aligned} du(t) &= p(t) dt, \\ m dp(t) &= \left( - \left( \gamma + \sum_{k \geq 1} \frac{c_k}{\lambda_k} \right) p(t) - \Phi'(u(t)) \right) dt \\ &\quad - \sum_{k \geq 1} \sqrt{\frac{2c_k}{\lambda_k}} dW_k(t) + \sqrt{2\gamma} dW_0(t). \end{aligned} \tag{2.8}$$

The well-posedness of (2.8) will be addressed briefly in Section 4. We then assert that  $(x_\epsilon(t), v_\epsilon(t))$  converges to the solution  $(u(t), p(t))$  of (2.8) in the following sense.

**Theorem 2.5.** *Suppose that  $\Phi$  satisfies Assumption 2.1 and the constants  $\alpha, \beta, s$  satisfy Condition (D) of Assumption 2.2. Let  $X_\epsilon(t) = (x_\epsilon(t), v_\epsilon(t), z_{1,\epsilon}(t), \dots)$  be the solution of (2.7) with initial conditions*

$$(x_\epsilon(0), v_\epsilon(0), z_{1,\epsilon}(0), z_{2,\epsilon}(0), \dots) = (x, v, z_1, z_2, \dots) \in \mathcal{H}_{-s},$$

*and  $(u(t), p(t))$  be the solution of (2.8) with initial conditions  $(u(0), p(0)) = (x, v)$ . Then, for every  $T, \xi > 0$ ,*

$$\mathbb{P} \left\{ \sup_{0 \leq t \leq T} |x_\epsilon(t) - u(t)| + |v_\epsilon(t) - p(t)| > \xi \right\} \rightarrow 0, \quad \epsilon \rightarrow 0.$$

The reader may wonder why the convergence result of Theorem 2.5 is restricted to the *diffusive* regime, namely  $\alpha > 1$  according to Condition (D) of Assumption 2.2. Heuristically, since the memory kernel  $K(t)$  decays like  $t^{-\alpha}$  as  $t \rightarrow \infty$ , we see that

$$K_\epsilon(t) = \frac{1}{\epsilon} K\left(\frac{t}{\epsilon}\right) \sim \epsilon^{\alpha-1} t^{-\alpha}, \quad t \rightarrow \infty.$$

By shrinking  $\epsilon$  further to zero, if  $\alpha > 1$ ,  $K_\epsilon(t)$  does not behave “badly” at infinity. In fact, it is not difficult to show that as  $\epsilon \downarrow 0$ ,  $K_\epsilon$  converges to the Dirac function  $\delta_0$  centered at the origin, in the sense of tempered distribution, namely, for every  $\varphi \in \mathcal{S}$ , the Schwartz space on  $\mathbb{R}$ , it holds that

$$\int_{\mathbb{R}} K_\epsilon(t) \varphi(t) dt \rightarrow |K|_{L^1(\mathbb{R})} \varphi(0).$$

which implies that the random force  $F(t)$  in (1.1) converges to a white noise process in the sense of random distribution, cf. [22], hence the so called “white-noise limit”. We will see later in the proof of Proposition 4.1 that the condition  $\alpha > 1$  is crucial for our analysis in order to obtain bounds on the solutions.

Finally, if  $\Phi$  and parameters  $\alpha, \beta$  satisfy stronger assumptions, then we are able to obtain better convergence than the result in Theorem 2.5. To be precise, we assume the following condition on  $\Phi$ .

**Assumption 2.6.** *There exist constants  $n, c > 0$  such that for every  $x, y \in \mathbb{R}$ ,*

$$\Phi'(x)y \leq c(\Phi(x) + |y|^n + 1).$$

The assumption above is again a requirement about the growth of  $\Phi'$  that guarantees a universal bound independent of  $\epsilon$  on the solution  $(x_\epsilon(t), v_\epsilon(t))$  of (2.7), cf. Proposition 4.1. We remark that a function  $\Phi$  satisfying Assumption 2.6 need not satisfy Assumption 2.1, taking  $\Phi$  a constant for example. It is also worthwhile to note that the class of polynomials of even degree satisfies Assumption 2.6. However, functions growing exponentially fast, e.g.  $e^{x^2}$ , do not.

With regards to space regularities, we assume the following condition about parameters  $\alpha, \beta$ .

**Assumption 2.7.** *Let  $\alpha, \beta > 0$  be as in (1.3). We assume that they satisfy*

$$\alpha > 2, \quad \text{and} \quad (\alpha - 2)\beta > 1.$$

We then have the following important result.

**Theorem 2.8.** *Suppose that  $\Phi$  satisfies Assumption 2.1 and Assumption 2.6 and that the constants  $\alpha, \beta, s$  satisfy Condition (D) of Assumption 2.2 and Assumption 2.7. Let  $X_\epsilon(t) = (x_\epsilon(t), v_\epsilon(t), z_{1,\epsilon}(t), \dots)$  be the solution of (2.7) with initial conditions*

$$(x_\epsilon(0), v_\epsilon(0), z_{1,\epsilon}(0), z_{2,\epsilon}(0), \dots) = (x, v, z_1, z_2, \dots) \in \mathcal{H}_{-s},$$

*and  $(u(t), p(t))$  be the solution of (2.8) with initial conditions  $(u(0), p(0)) = (x, v)$ . Then, for every  $T > 0$ ,  $1 \leq q < 2$ , it holds that*

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} |x_\epsilon(t) - u(t)|^q + |v_\epsilon(t) - p(t)|^q \right] \rightarrow 0, \quad \epsilon \rightarrow 0.$$

Theorem 2.8 strengthens a previous result from [34], where the potential  $\Phi'$  is assumed to be bounded. The proofs of Theorem 2.5 and Theorem 2.8 will be carried out in Section 4.

### 3. ZERO-MASS LIMIT

Throughout the rest of the paper,  $C, c$  denote generic positive constants. The important parameters that they depend on will be indicated in parenthesis, e.g.  $c(T, q)$  depends on parameters  $T$  and  $q$ .

In this section, for notation simplicity, we shall omit the subscript  $m$  in

$$X_m(t) = (x_m(t), v_m(t), z_{1,m}(t), \dots).$$

We begin by addressing the well-posedness of (2.3) whose proof follows a standard Lyapunov-type argument that was also used to establish the well-posedness of (1.6) in [12]. The technique is classical and has been employed previously in literature [2, 14, 23]. We shall omit specific details and briefly summarize the main steps.

*Sketch of the proof of Proposition 2.3.* For  $R > 0$ , let  $\theta^R \in C^\infty(\mathbb{R}, [0, 1])$  satisfy

$$\theta^R(x) = \begin{cases} 1 & \text{if } |x| \leq R, \\ 0 & \text{if } |x| \geq R + 1. \end{cases} \quad (3.1)$$

We consider the “cutoff” equation corresponding to (2.3)

$$\begin{aligned} \gamma du(t) &= \left( -\Phi'(u(t))\theta^R(u(t)) - \left( \sum_{k \geq 1} c_k \right) u(t) - \sum_{k \geq 1} \sqrt{c_k} f_k(t) \right) dt + \sqrt{2\gamma} dW_0(t), \\ df_k(t) &= \left( -\lambda_k f_k(t) - \lambda_k \sqrt{c_k} u(t) \right) dt + \sqrt{2\lambda_k} dW_k(t), \quad k = 1, 2, \dots \end{aligned} \quad (3.2)$$

We observe that in (3.2), the drift term is globally Lipschitz and the noise is additive. Thus, by using a standard Banach fixed point argument, the corresponding global (in time) solution  $U^R$  exists and is unique. Next, define the stopping time

$$\tau_R = \inf \left\{ t > 0 : \|U(t)\|_{\dot{\mathcal{H}}_{-s}} > R \right\}.$$

Note that, for all times  $t < \tau_R$ ,  $U^R$  solves (2.3). Consequently, the solution (2.3) exists and is unique up until the *time of explosion*  $\tau_\infty = \lim_{R \rightarrow \infty} \tau_R$ , which is possibly finite on a set of positive probability. We finally introduce the Lyapunov function

$$\Psi(U) := \frac{1}{\gamma} \left( \Phi(u) + \left( \sum_{k \geq 1} c_k \right) \frac{u^2}{2} \right) + \frac{1}{2} \sum_{k \geq 1} k^{-2s} f_k^2. \quad (3.3)$$



It is clear that  $\Psi(U)$  dominates  $\|U\|_{\mathcal{H}_{-s}}^2$ . Applying Ito's formula to  $\Psi(U)$ , one can derive a global bound on the solutions  $U^R(t)$  that is independent of  $R$ , namely, there exists a constant  $C(U_0, T)$  such that

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} \|U(t \wedge \tau_R)\|_{\mathcal{H}_{-s}}^2 \right] \leq C(T, U_0).$$

Sending  $R$  to infinity, it follows from Fatou's Lemma that

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} \|U(t \wedge \tau_\infty)\|_{\mathcal{H}_{-s}}^2 \right] \leq C(U_0, T).$$

implying  $\mathbb{P}\{T < \tau_\infty\} = 1$  for any  $T > 0$ . Taking  $T$  to infinity, we see that  $\mathbb{P}\{\tau_\infty = \infty\} = 1$ , thereby obtaining the global solution of (2.3).  $\square$

Although the construction of the global solution  $U(t)$  of (2.3) via the local solutions  $U^R(t)$  of (3.2) is quite standard, the proof of Theorem 2.4 will make use of a non trivial observation on these local solutions. The arguments that we are going to employ are inspired from the work of [17]. Before diving into detail, we briefly explain the main idea, which is a two-fold: first, we show that the result holds for  $\Phi'$  being Lipschitz. In particular, we obtain the convergence in sup norm for the local solutions, namely for all  $R, T > 0$ , we have

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| \right] \rightarrow 0, \quad m \rightarrow 0,$$

where  $x^R(t)$  is in the following cut-off system for (1.6)

$$\begin{aligned} dx(t) &= v(t) dt, \\ m dv(t) &= (-\gamma v(t) - \Phi'(x(t))\theta^R(x(t)) - \sum_{k \geq 1} \sqrt{c_k} z_k(t)) dt + \sqrt{2\gamma} dW_0(t), \\ dz_k(t) &= (-\lambda_k z_k(t) + \sqrt{c_k} v(t)) dt + \sqrt{2\lambda_k} dW_k(t), \quad k \geq 1. \end{aligned} \tag{3.4}$$

Then, by taking  $R$  necessarily large, we obtain the desired result.

We now proceed by showing that the result holds true in a simpler setting where  $\Phi'$  is globally Lipschitz. The proof is adapted from that of Theorem 1 of [19].

**Proposition 3.1.** *Suppose that  $\Phi'$  is globally Lipschitz and that the constants  $\alpha, \beta, s$  satisfy Assumption 2.2. Let  $X(t) = (x(t), v(t), z_1(t), \dots)$  solve (1.6) with initial conditions*

$$(x(0), v(0), z_1(0), \dots) = (x, v, z_1, \dots) \in \mathcal{H}_{-s},$$

*and  $U(t) = (u(t), f_1(t), \dots)$  solve (2.3) with initial conditions*

$$(u(0), f_1(0), f_2(0), \dots) = (x, z_1 - \sqrt{c_1}x, z_2 - \sqrt{c_2}x, \dots) \in \dot{\mathcal{H}}_{-s}.$$

*Then, for every  $T, q > 0$ , it holds that*

$$\mathbb{E} \sup_{0 \leq t \leq T} |x(t) - u(t)|^q \rightarrow 0, \quad m \rightarrow 0.$$

*Proof.* Using Duhamel's formula,  $z_k(t)$  from (1.6) can be solved explicitly as

$$z_k(t) = e^{-\lambda_k t} z_k(0) + \sqrt{c_k} \int_0^t e^{-\lambda_k(t-r)} v(r) dr + \sqrt{2\lambda_k} \int_0^t e^{-\lambda_k(t-r)} dW_k(r), \tag{3.5}$$

which is equivalent to

$$\begin{aligned} z_k(t) &= e^{-\lambda_k t} (z_k(0) - \sqrt{c_k} x(0)) + \sqrt{c_k} x(t) - \sqrt{c_k} \lambda_k \int_0^t e^{-\lambda_k(t-r)} x(r) dr \\ &\quad + \sqrt{2\lambda_k} \int_0^t e^{-\lambda_k(t-r)} dW_k(r), \end{aligned}$$



where we have used an integration by parts on the term  $\int_0^t e^{-\lambda_k(t-r)} v(r) dr$  in the first equality. Substituting into the second equation of (1.6), we arrive at

$$\begin{aligned} m dv(t) + \gamma dx(t) &= \left( -\Phi'(x(t)) - \sum_{k \geq 1} \sqrt{c_k} e^{-\lambda_k t} (z_k(0) - \sqrt{c_k} x(0)) - \left( \sum_{k \geq 1} c_k \right) x(t) \right. \\ &\quad \left. + \sum_{k \geq 1} c_k \lambda_k \int_0^t e^{-\lambda_k(t-r)} x(r) dr - \sum_{k \geq 1} \sqrt{2c_k \lambda_k} \int_0^t e^{-\lambda_k(t-r)} dW_k(r) \right) dt \\ &\quad + \sqrt{2\gamma} dW_0(t). \end{aligned} \quad (3.6)$$

Likewise, we obtain the following equation from (2.3)

$$\begin{aligned} \gamma du(t) &= \left( -\Phi'(u(t)) - \left( \sum_{k \geq 1} c_k \right) u(t) - \sum_{k \geq 1} \sqrt{c_k} e^{-\lambda_k t} (z_k(0) - \sqrt{c_k} x(0)) \right. \\ &\quad \left. + \sum_{k \geq 1} c_k \lambda_k \int_0^t e^{-\lambda_k(t-r)} u(r) dr - \sum_{k \geq 1} \sqrt{2c_k \lambda_k} \int_0^t e^{-\lambda_k(t-r)} dW_k(r) \right) dt \\ &\quad + \sqrt{2\gamma} dW_0(t). \end{aligned} \quad (3.7)$$

Subtracting (3.7) from (3.6) and setting  $\bar{x}(t) = x(t) - u(t)$ , we find that

$$\begin{aligned} m dv(t) + \gamma d\bar{x}(t) &= \left( -[\Phi'(x(t)) - \Phi'(u(t))] - \left( \sum_{k \geq 1} c_k \right) \bar{x}(t) + \sum_{k \geq 1} c_k \lambda_k \int_0^t e^{-\lambda_k(t-r)} \bar{x}(r) dr \right) dt \\ &\leq c \left( 1 + \sum_{k \geq 1} c_k \right) \sup_{0 \leq r \leq t} |\bar{x}(r)| dt, \end{aligned}$$

where  $c > 0$  is a Lipschitz constant for  $\Phi'$ . Recalling  $c_k$  from (1.3), we apply Gronwall's inequality to estimate for all  $T, q > 0$

$$\mathbb{E} \sup_{0 \leq t \leq T} |\bar{x}(t)|^q \leq m^q \mathbb{E} \sup_{0 \leq t \leq T} |v(t) - v|^q e^{c(T)}.$$

The result now follows immediately from Proposition 3.2 below.  $\square$

**Proposition 3.2.** *Under the same Hypothesis of Theorem 2.4, suppose further that  $\Phi'(x)$  is globally Lipschitz. Let  $X(t) = (x(t), v(t), z_1(t), \dots)$  solve (1.6) with initial conditions  $(x(0), v(0), z_1(0), \dots) = (x, v, z_1, \dots) \in \mathcal{H}_{-s}$ . Then, for every  $T > 0, q > 1$ , it holds that*

$$m^q \mathbb{E} \sup_{0 \leq t \leq T} |v(t)|^q \rightarrow 0, \quad m \rightarrow 0.$$

In order to prove Proposition 3.2, we need the following important lemma whose proof is based on Lemma 3.19, [3] and Lemma 2, [19]. It will be also useful later in Section 4.

**Lemma 3.3.** *Given  $\kappa, \eta > 0$ , let  $f(t) = \sqrt{2\kappa} \int_0^t e^{-\eta(t-r)} dW(r)$  where  $W(t)$  is a standard Brownian Motion. Then, for all  $T > 0, q > 1$ , there exists a constant  $C(T, q) > 0$  such that*

$$\mathbb{E} \sup_{0 \leq t \leq T} f(t)^{2q} \leq \frac{\kappa^q}{\eta^{q-1}} C(T, q). \quad (3.8)$$

**Remark 3.4.** *The estimate in (3.8) is sharper than the usual exponential martingale estimate. In finite-dimensional settings, it is sufficient to bound the LHS of (3.8) by  $C(T, q, \eta, \kappa)$ , cf. [3, 19, 30]. In our setting, we have to keep track explicitly in term of  $\eta$  and  $\kappa$ , hence the RHS of (3.8).*

The proof of Lemma 3.3 is similar to that of Lemma 2, [19]. We include it here for the sake of completeness.

*Proof of Lemma 3.3.* In view of Lemma 3.19, [3], we have the following estimate for  $A > 0$ ,

$$\mathbb{P}\left\{\sup_{0 \leq t \leq T} f(t)^2 \geq A\right\} \leq \frac{\eta T}{\int_0^{\sqrt{\eta A/\kappa}} e^{r^2/2} \int_0^r e^{-\ell^2/2} d\ell dr}.$$

We proceed to find a lower bound for the above denominator. To this end, we first claim that for  $r \geq 0$ ,

$$\int_0^r e^{-\ell^2/2} d\ell \geq \frac{r e^{-r^2/4}}{2}.$$

Indeed, on one hand, if  $r \geq 1$ , then

$$e^{r^2/4} \int_0^r e^{-\ell^2/2} d\ell \geq e^{r^2/4} \int_0^1 e^{-\ell^2/2} d\ell \geq e^{r^2/4} \int_0^1 e^{-1/2} d\ell \geq \frac{e^{r^2/4}}{2} \geq \frac{r}{2}.$$

On the other hand, if  $0 \leq r \leq 1$ , then

$$e^{r^2/4} \int_0^r e^{-\ell^2/2} d\ell \geq \int_0^r 1 - \frac{\ell^2}{2} d\ell = r\left(1 - \frac{r^2}{6}\right) \geq \frac{r}{2}.$$

With this observation, we find

$$\int_0^{\sqrt{\eta A/\kappa}} e^{r^2/2} \int_0^r e^{-\ell^2/2} d\ell dr \geq \int_0^{\sqrt{\eta A/\kappa}} e^{r^2/2} \frac{r e^{-r^2/4}}{2} dr = e^{\eta A/4\kappa} - 1 \geq \frac{\eta A}{4\kappa} e^{\eta A/8\kappa},$$

where in the last implication, we have used the following inequality for every  $r \geq 0$ ,

$$e^r - 1 \geq r e^{r/2}.$$

Putting everything together, we obtain

$$\mathbb{P}\left\{\sup_{0 \leq t \leq T} f(t)^2 \geq A\right\} \leq \frac{\eta T}{\frac{\eta A}{4\kappa} e^{\eta A/8\kappa}} = \frac{4\kappa T}{A} e^{-\eta A/8\kappa}.$$

It follows that for  $q > 2$ ,

$$\begin{aligned} \mathbb{E} \sup_{0 \leq t \leq T} f(t)^{2q} &= \int_0^\infty q A^{q-1} \mathbb{P}\left\{\sup_{0 \leq t \leq T} f(t)^2 \geq A\right\} dA \\ &\leq \int_0^\infty q A^{q-1} \frac{4\kappa T}{A} e^{-\eta A/8\kappa} dA \\ &= C(T, q) \kappa \int_0^\infty A^{q-2} e^{-\eta A/8\kappa} dA \\ &= C(T, q) \frac{\kappa^q}{\eta^{q-1}}, \end{aligned}$$

which completes the proof. □

With Lemma 3.3 in hand, we are ready to give the proof of Proposition 3.2.

*Proof of Proposition 3.2.* We only have to prove the result for  $q > 0$  sufficiently large, say  $q \geq q_1$ . As if it holds for  $q_1$ , then for every  $q < q_1$ , by Holder's inequality, we have

$$\mathbb{E}\left[m^q \sup_{0 \leq t \leq T} |v(t)|^q\right] \leq \left(\mathbb{E}\left[m^{q_1} \sup_{0 \leq t \leq T} |v(t)|^{q_1}\right]\right)^{q/q_1} \rightarrow 0, \quad \text{as } m \rightarrow 0.$$

We begin by noting that  $v(t)$  from (1.6) is written as

$$\begin{aligned} m v(t) &= m e^{-\frac{\gamma}{m}t} v(0) - \int_0^t e^{-\frac{\gamma}{m}(t-r)} \Phi'(x(r)) dr - \sum_{k \geq 1} \sqrt{c_k} \int_0^t e^{-\frac{\gamma}{m}(t-r)} z_k(r) dr \\ &\quad + \sqrt{2\gamma} \int_0^t e^{-\frac{\gamma}{m}(t-r)} dW_0(r). \end{aligned}$$

Substituting  $z_k(t)$  from (3.5), we have

$$\begin{aligned} m v(t) &= m e^{-\frac{\gamma}{m}t} v(0) - \int_0^t e^{-\frac{\gamma}{m}(t-r)} \Phi'(x(r)) dr - \sum_{k \geq 1} \sqrt{c_k} \int_0^t e^{-\frac{\gamma}{m}(t-r)} e^{-\lambda_k r} z_k(0) dr \\ &\quad - \sum_{k \geq 1} c_k \int_0^t e^{-\frac{\gamma}{m}(t-r)} \int_0^r e^{-\lambda_k(r-\ell)} v(\ell) d\ell dr \\ &\quad - \sum_{k \geq 1} \sqrt{2c_k \lambda_k} \int_0^t e^{-\frac{\gamma}{m}(t-r)} \int_0^r e^{-\lambda_k(r-\ell)} dW_k(\ell) dr \\ &\quad + \sqrt{2\gamma} \int_0^t e^{-\frac{\gamma}{m}(t-r)} dW_0(r). \end{aligned}$$

For every  $q$  sufficiently large, we invoke the assumption that  $\Phi'$  is Lipschitz to estimate

$$\begin{aligned} m^{2q} \mathbb{E} \sup_{0 \leq t \leq T} v(t)^{2q} &\leq c(q, v) \left[ m^{2q} \left[ 1 + \mathbb{E} \sup_{0 \leq t \leq T} x(t)^{2q} + \left| \sum_{k \geq 1} \sqrt{c_k} z_k \right|^{2q} \right. \right. \\ &\quad + c(T) \left| \sum_{k \geq 1} c_k \right|^{2q} \int_0^T \mathbb{E} \sup_{0 \leq r \leq t} v(r)^{2q} dt \\ &\quad + \left( \sum_{k \geq 1} c_k^{(1/2-1/2q)q^*} \right)^{2q/q^*} \sum_{k \geq 1} \mathbb{E} \sup_{0 \leq t \leq T} \left| \sqrt{2c_k^{1/q} \lambda_k} \int_0^t e^{-\lambda_k(t-r)} dW_k(r) \right|^{2q} \\ &\quad \left. + \mathbb{E} \sup_{0 \leq t \leq T} \left| \sqrt{2\gamma} \int_0^t e^{-\frac{\gamma}{m}(t-r)} dW_0(r) \right|^{2q} \right], \end{aligned}$$

where in the third line, we have used Holder's inequality with  $\frac{1}{q^*} + \frac{1}{2q} = 1$ . Also, note that from the first equation of (1.6), it holds that

$$\mathbb{E} \sup_{0 \leq t \leq T} x(t)^{2q} \leq c(q) \left( x^{2q} + \int_0^T \mathbb{E} \sup_{0 \leq r \leq t} v(r)^{2q} dt \right),$$

and that by Lemma 3.3, we have

$$\sum_{k \geq 1} \mathbb{E} \sup_{0 \leq t \leq T} \left| \sqrt{2c_k^{1/q} \lambda_k} \int_0^t e^{-\lambda_k(t-r)} dW_k(r) \right|^{2q} \leq c(T, q) \sum_{k \geq 1} c_k \lambda_k,$$

and

$$\mathbb{E} \sup_{0 \leq t \leq T} \left| \sqrt{2\gamma} \int_0^t e^{-\frac{\gamma}{m}(t-r)} dW_0(r) \right|^{2q} \leq c(T, q) \gamma m^{q-1}.$$

Furthermore, recalling  $c_k$  from (1.3), we see that for  $q > 0$  sufficiently large

$$\sum_{k \geq 1} c_k^{(1/2-1/2q)q^*} = \sum_{k \geq 1} \frac{1}{k^{(1+\alpha\beta)(1/2-1/2q)q^*}} < \infty,$$

thanks to the fact that  $q^* > 1$  and  $\alpha\beta > 1$ , where the latter follows from the conditions about  $\alpha, \beta$  in Assumption 2.2. Also, recalling  $\lambda_k$  from (1.3) and the norm  $\|\cdot\|_{-s}$  from (2.1), it is straightforward to verify that the sums  $\sum_{k \geq 1} \sqrt{c_k} z_k$ ,  $\sum_{k \geq 1} c_k$ , and  $\sum_{k \geq 1} c_k \lambda_k$  are absolutely convergent. Putting everything together, we find

$$m^{2q} \mathbb{E} \sup_{0 \leq t \leq T} v(t)^{2q} \leq c(T, q, X(0)) \left[ m^{2q} + m^{q-1} + m^{2q} \int_0^T \mathbb{E} \sup_{0 \leq r \leq t} v(r)^{2q} dt \right],$$

where  $c(T, q, X(0)) > 0$  is independent with  $m$ . Gronwall's inequality now implies

$$m^{2q} \mathbb{E} \sup_{0 \leq t \leq T} v(t)^{2q} \leq c(T, q, X(0)) (m^{2q} + m^{q-1}) \rightarrow 0, \quad m \rightarrow 0.$$

The proof is thus complete.  $\square$

We now turn our attention to Theorem 2.4. The proof is a slightly modification from that of Theorem 2.4 of [17]. The key observation is that instead of controlling the exiting time of the process  $x(t)$  as  $m \rightarrow 0$ , we are able to control  $u(t)$  since  $u(t)$  is independent of  $m$ .

*Proof of Theorem 2.4.* For  $R, m > 0$ , define the following stopping times

$$\sigma^R = \inf_{t \geq 0} \{|u(t)| \geq R\}, \quad \text{and} \quad \sigma_m^R = \inf_{t \geq 0} \{|x(t)| \geq R\}, \quad (3.9)$$

and recall

$$\tau^R = \inf_{t \geq 0} \{\|U(t)\|_{\dot{\mathcal{H}}_{-s}} \geq R\}, \quad \text{and} \quad \tau_m^R = \inf_{t \geq 0} \{\|X(t)\|_{\mathcal{H}_{-s}} \geq R\}.$$

By the definitions of the norms in  $\dot{\mathcal{H}}_{-s}$ , cf. (2.4), we see that  $\tau^R \leq \sigma^R$  a.s. From the proof of Proposition 2.3, it is straight forward to verify that for all  $T > 0$ ,

$$\mathbb{P}\{\sigma^R < T\} \leq \mathbb{P}\{\tau^R < T\} \rightarrow 0, \quad R \rightarrow \infty.$$

For  $R, T, m, \xi > 0$ , we have

$$\begin{aligned} \mathbb{P}\left\{\sup_{0 \leq t \leq T} |x(t) - u(t)| > \xi\right\} &\leq \mathbb{P}\left\{\sup_{0 \leq t \leq T} |x(t) - u(t)| > \xi, \sigma^R \wedge \sigma_m^R \geq T\right\} \\ &\quad + \mathbb{P}\{\sigma^R \wedge \sigma_m^R < T\}. \end{aligned}$$

To control the first term on the above RHS, we note that for  $0 \leq t \leq \sigma^R \wedge \sigma_m^R$ ,  $u(t) = u^R(t)$  and  $x(t) = x^R(t)$  a.s. We thus obtain the bound

$$\mathbb{P}\left\{\sup_{0 \leq t \leq T} |x(t) - u(t)| > \xi, \sigma^R \wedge \sigma_m^R \geq T\right\} \leq \mathbb{P}\left\{\sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| > \xi\right\} \rightarrow 0, \quad m \rightarrow 0,$$

where the last convergence in probability follows immediately from Proposition 3.1. We are left to estimate  $\mathbb{P}\{\sigma^R \wedge \sigma_m^R < T\}$ . To this end, we have that

$$\begin{aligned} \mathbb{P}\{\sigma^R \wedge \sigma_m^R < T\} &\leq \mathbb{P}\left\{\sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| \leq \xi, \sigma^R \wedge \sigma_m^R < T\right\} \\ &\quad + \mathbb{P}\left\{\sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| > \xi\right\} \\ &\leq \mathbb{P}\left\{\sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| \leq \xi, \sigma_m^R < T \leq \sigma^R\right\} + \mathbb{P}\{\sigma^R < T\} \\ &\quad + \mathbb{P}\left\{\sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| > \xi\right\}. \end{aligned}$$

Note that for  $R > 1$  and  $\xi \in (0, 1)$ , a chain of event implications is derived as follows.

$$\begin{aligned}
& \left\{ \sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| \leq \xi, \sigma_m^R < T \leq \sigma^R \right\} \\
&= \left\{ \sup_{0 \leq t \leq T} |x^R(t) - u(t)| \leq \xi, \sup_{0 \leq t \leq T} |x^R(t)| \geq R, \sigma_m^R < T \leq \sigma^R \right\} \\
&\subseteq \left\{ \sup_{0 \leq t \leq T} |u(t)| > R - 1, \sigma_m^R < T \leq \sigma^R \right\} \\
&\subseteq \{\sigma^{R-1} < T\},
\end{aligned}$$

which implies that

$$\mathbb{P} \left\{ \sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| < \xi, \sigma_m^R < T \leq \sigma^R \right\} \leq \mathbb{P} \{\sigma^{R-1} < T\}.$$

Finally, putting everything together, for  $R > 1 > \xi > 0$ ,  $T, m > 0$ , we obtain the estimate

$$\begin{aligned}
& \mathbb{P} \left\{ \sup_{0 \leq t \leq T} |x(t) - u(t)| > \xi \right\} \\
&\leq 2\mathbb{P} \left\{ \sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| > \xi \right\} + \mathbb{P} \{\sigma^{R-1} < T\} + \mathbb{P} \{\sigma^R < T\} \\
&\leq 2\mathbb{P} \left\{ \sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| > \xi \right\} + \mathbb{P} \{\tau^{R-1} < T\} + \mathbb{P} \{\tau^R < T\}.
\end{aligned}$$

By taking  $R$  sufficiently large and then shrinking  $m$  further to zero, we obtain the result, thus completing the proof.  $\square$

#### 4. WHITE-NOISE LIMIT

For notation simplicity, in this section, we shall omit the subscript  $\epsilon$  in

$$X_\epsilon(t) = (x_\epsilon(t), v_\epsilon(t), z_{1,\epsilon}(t), \dots).$$

With regards to the well-posedness of (2.8), recalling  $c_k, \lambda_k$  from (1.3), we see that the noise term is well-defined thanks to Condition (D) of Assumption 2.2, namely

$$\mathbb{E} \left( \int_0^T \sum_{k \geq 1} \sqrt{\frac{2c_k}{\lambda_k}} dW_k(t) \right)^2 = 2T \sum_{k \geq 1} \frac{c_k}{\lambda_k} = 2T \sum_{k \geq 1} \frac{1}{k^{1+(\alpha-1)\beta}} < \infty. \quad (4.1)$$

The solution  $(u(t), p(t))$  of (2.8) then is constructed using similar arguments as in the proof of Proposition 2.3 in Section 3 via stopping times  $\tau^R$ ,  $R > 0$ , given by

$$\tau^R = \inf_{t \geq 0} \{u(t)^2 + p(t)^2 \geq R^2\}, \quad (4.2)$$

and the local solutions

$$\begin{aligned}
du^R(t) &= p^R(t) dt, \\
m dp^R(t) &= \left( - \left( \gamma + \sum_{k \geq 1} \frac{c_k}{\lambda_k} \right) p^R(t) - \Phi'(u^R(t)) \theta_R(u^R(t)) \right) dt \\
&\quad + \sum_{k \geq 1} \sqrt{\frac{2c_k}{\lambda_k}} dW_k(t) + \sqrt{2\gamma} dW_0(t),
\end{aligned} \quad (4.3)$$

where  $\theta^R$  is the cut-off function defined in (3.1). Furthermore, we have the following bound: for every  $T > 0$  and  $(u_0, p_0) \in \mathbb{R}^2$ , it holds that

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} u(t)^2 + p(t)^2 \right] \leq C(T, u_0, p_0). \quad (4.4)$$

This estimate will be useful later in the proof of Theorem 2.8. The solution  $X_\epsilon(t)$  is constructed using the stopping time  $\tau_\epsilon^R$  given by

$$\tau_\epsilon^R = \inf_{t \geq 0} \{\|X(t)\|_{\mathcal{H}_{-s}} \geq R\}, \quad (4.5)$$

and the local solutions of the cut-off system obtained from (2.7)

$$\begin{aligned} dx^R(t) &= v^R(t) dt, \\ m dv^R(t) &= \left( -\gamma v^R(t) - \Phi'(x^R(t)) \theta^R(x^R(t)) - \sum_{k \geq 1} \sqrt{\frac{c_k}{\epsilon}} z_k^R(t) \right) dt \\ &\quad + \sqrt{2\gamma} dW_0(t), \\ dz_k^R(t) &= \left( -\frac{\lambda_k}{\epsilon} z_k^R(t) + \sqrt{\frac{c_k}{\epsilon}} v^R(t) \right) dt + \sqrt{\frac{2\lambda_k}{\epsilon}} dW_k(t), \quad k \geq 1. \end{aligned} \quad (4.6)$$

We now turn to the proof of Theorem 2.5. Similar to the proof of Theorem 2.4, it will make use of the local solutions  $(u^R(t), p^R(t))$  from (4.3) and  $(x^R(t), v^R(t))$  from (4.6). As mentioned previously in Section 3, the idea essentially consists of two major steps: first, fixing  $R > 0$ , we show that the corresponding local solution  $(x^R(t), v^R(t))$  in (4.6) converges to  $(u^R(t), p^R(t))$  in (4.3). Then, taking  $R$  sufficiently large, we obtain the convergence in probability of the original solutions by using appropriate bounds on stopping times when  $(u(t), p(t))$  exits the ball of radius  $R$  centered at origin. We begin by the following important result giving a uniform bound on the pair  $(x(t), v(t))$ .

**Proposition 4.1.** *Suppose that  $\alpha, \beta, s$  satisfy Condition (D) of Assumption 2.2. We assume further that either*

*(a)  $\Phi'$  is globally Lipschitz,*

*or*

*(b)  $\Phi'$  is not Lipschitz, but  $\Phi$  satisfies Assumptions 2.1 and Assumption 2.6, and  $\alpha, \beta$  satisfy Assumption 2.7.*

*Let  $X(t)$  solve (2.7) with initial conditions  $(x(0), v(0), z_1(0), \dots) \in \mathcal{H}_{-s}$ . Then, for every  $T > 0$ , there exists a finite constant  $C(T, X(0))$  such that*

$$\sup_{\epsilon} \mathbb{E} \left[ \sup_{0 \leq t \leq T} x(t)^2 + v(t)^2 \right] \leq C(T, X(0)).$$

*Proof.* We begin by applying Duhamel's formula on  $z_k(t)$  from (2.7) to see that

$$z_k(t) = e^{-\frac{\lambda_k}{\epsilon} t} z_k(0) + \sqrt{\frac{c_k}{\epsilon}} \int_0^t e^{-\frac{\lambda_k}{\epsilon} (t-r)} v(r) dr + \sqrt{\frac{2\lambda_k}{\epsilon}} \int_0^t e^{-\frac{\lambda_k}{\epsilon} (t-r)} dW_k(r). \quad (4.7)$$

Substituting  $z_k$  by the formula above in the second equation from (2.7) in integral form, we obtain

$$\begin{aligned} mv(t) &= mv(0) + \int_0^t -\gamma v(r) - \Phi'(x(r)) dr + \sqrt{2\gamma} \int_0^t dW_0(r) \\ &\quad - \sum_{k \geq 1} \sqrt{\frac{c_k}{\epsilon}} \int_0^t e^{-\frac{\lambda_k}{\epsilon} r} z_k(0) dr - \sum_{k \geq 1} \frac{c_k}{\epsilon} \int_0^t \int_0^r e^{-\frac{\lambda_k}{\epsilon} (r-\ell)} v(\ell) d\ell dr \\ &\quad - \sum_{k \geq 1} \frac{\sqrt{2c_k \lambda_k}}{\epsilon} \int_0^t \int_0^r e^{-\frac{\lambda_k}{\epsilon} (r-\ell)} dW_k(\ell) dr. \end{aligned} \quad (4.8)$$

It is important to note that using integration by parts, the last noise term above can be written as

$$\begin{aligned}
& -\frac{\sqrt{2c_k\lambda_k}}{\epsilon} \int_0^t \int_0^r e^{-\frac{\lambda_k}{\epsilon}(r-\ell)} dW_k(\ell) dr \\
& = \sqrt{\frac{2c_k}{\lambda_k}} \int_0^t e^{-\frac{\lambda_k}{\epsilon}(t-r)} dW_k(r) - \sqrt{\frac{2c_k}{\lambda_k}} \int_0^t dW_k(r). \quad (4.9)
\end{aligned}$$

Suppose that Condition (a) holds, i.e.,  $\Phi'$  is Lipschitz. In view of (4.8) and (4.9), we have the following estimate for every  $q > 1$  and  $0 \leq t \leq T$ ,

$$\begin{aligned}
v(t)^{2q} & \leq c(q) \left[ |v(0)|^{2q} + \int_0^T \sup_{0 \leq r \leq t} v(r)^{2q} + x(r)^{2q} dr + \left( \sum_{k \geq 1} \frac{\sqrt{\epsilon c_k}}{\lambda_k} (1 - e^{-\frac{\lambda_k}{\epsilon}t}) |z_k(0)| \right)^{2q} \right. \\
& \quad + c(T) \left( \sum_{k \geq 1} \frac{c_k}{\lambda_k} \right)^{2q} \int_0^T \sup_{0 \leq \ell \leq r} v(\ell)^{2q} dr \\
& \quad + \sup_{0 \leq t \leq T} \left( \sqrt{2\gamma} \int_0^t dW_0(r) - \sum_{k \geq 1} \sqrt{\frac{2c_k}{\lambda_k}} \int_0^t dW_k(r) \right)^{2q} \\
& \quad \left. + \left( \sum_{k \geq 1} k^{-sq*} \right)^{2q/q*} \sum_{k \geq 1} \sup_{0 \leq t \leq T} \left| \sqrt{\frac{2c_k k^{2s}}{\lambda_k}} \int_0^t e^{-\frac{\lambda_k}{\epsilon}(t-r)} dW_k(r) \right|^{2q} \right],
\end{aligned}$$

where in the last line, we have used Holder's inequality with  $\frac{1}{2q} + \frac{1}{q*} = 1$ . Note that for every  $x \geq 0$ , we have  $1 - e^{-x} \leq \sqrt{x}$ . Using this inequality, we estimate

$$\begin{aligned}
\sum_{k \geq 1} \frac{\sqrt{\epsilon c_k}}{\lambda_k} (1 - e^{-\frac{\lambda_k}{\epsilon}t}) |z_k(0)| & \leq \sum_{k \geq 1} \sqrt{\frac{c_k t}{\lambda_k}} |z_k(0)| \\
& \leq \left( T \sum_{k \geq 1} \frac{c_k k^{2s}}{\lambda_k} \sum_{k \geq 1} k^{-2s} z_k(0)^2 \right)^{1/2}. \quad (4.10)
\end{aligned}$$

Recalling  $c_k, \lambda_k$  from (1.3) and the norm  $\|\cdot\|_{\mathcal{H}_s}$  from (2.1), thanks to Condition (D) of Assumption 2.2, we see that the above RHS is finite and so is the sum  $\sum_{k \geq 1} c_k/\lambda_k$ . In addition, using Burkholder-Davis-Gundy's inequality, we have

$$\begin{aligned}
& \mathbb{E} \sup_{0 \leq t \leq T} \left( \sqrt{2\gamma} \int_0^t dW_0(r) - \sum_{k \geq 1} \sqrt{\frac{2c_k}{\lambda_k}} \int_0^t dW_k(r) \right)^{2q} \\
& \leq c(q) \mathbb{E} \left( 2\gamma \int_0^T dr + \sum_{k \geq 1} \frac{2c_k}{\lambda_k} \int_0^T dr \right)^q = c(T, q) < \infty. \quad (4.11)
\end{aligned}$$

Finally, we invoke Lemma 3.3 again to find

$$\begin{aligned}
& \mathbb{E} \sum_{k \geq 1} \sup_{0 \leq t \leq T} \left| \sqrt{\frac{2c_k k^{2s}}{\lambda_k}} \int_0^t e^{-\frac{\lambda_k}{\epsilon}(t-r)} dW_k(r) \right|^{2q} \\
& \leq c(T, q) \sum_{l \geq 1} \frac{\epsilon^{q-1} c_k^q k^{2sq}}{\lambda_k^{2q-1}} \\
& = c(T, q) \epsilon^{q-1} \sum_{k \geq 1} \frac{1}{k^{q+(q\alpha-2q+1)\beta-2sq}}. \quad (4.12)
\end{aligned}$$



Note that for  $\alpha > 1$ ,  $s > 1/2$  and  $\beta > 0$  satisfying Condition (D) of Assumption 2.2, there exist constants  $q > 1$  and  $0 < q^* < 2$  such that

$$q + (q\alpha - 2q + 1)\beta - 2sq > 1, \quad sq^* > 1, \quad \text{and} \quad \frac{1}{2q} + \frac{1}{q^*} = 1.$$

Consequently, the sums  $\sum_{k \geq 1} k^{-[q+(q\alpha-2q+1)\beta-2sq]}$  and  $\sum_{k \geq 1} k^{-sq^*}$  are both finite. Combining everything together, we infer

$$\mathbb{E}\left(\sup_{0 \leq t \leq T} x(t)^{2q} + v(t)^{2q}\right) \leq c(T, q, X(0)) \left[1 + \int_0^T \mathbb{E}\left(\sup_{0 \leq r \leq t} x(r)^{2q} + v(r)^{2q}\right) dt\right].$$

Choosing such  $q$ , we finally obtain the following estimate using Gronwall's inequality

$$\mathbb{E}\left(\sup_{0 \leq t \leq T} x(t)^{2q} + v(t)^{2q}\right) \leq c(T, q, X(0)), \quad (4.13)$$

which proves the result for Condition (a) since  $q > 1$ .

Now suppose that Condition (b) holds. To simplify notation, we set

$$g_k(t) := \int_0^t e^{-\frac{\lambda_k}{\epsilon}(t-r)} v(r) dr, \quad \text{and} \quad w_k(t) := \sqrt{2} \int_0^t e^{-\frac{\lambda_k}{\epsilon}(t-r)} dW_k(r).$$

Following (4.8) and (4.9), the equation on  $v(t)$  is written as

$$\begin{aligned} d\left(mv(t) - \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(t)\right) &= \left(-\gamma v(t) - \Phi'(x(t)) - \sum_{k \geq 1} \sqrt{\frac{c_k}{\epsilon}} e^{-\frac{\lambda_k}{\epsilon} t} z_k(0) - \sum_{k \geq 1} \frac{c_k}{\epsilon} g_k(t)\right) dt \\ &\quad + \sqrt{2\gamma} dW_0(t) - \sum_{k \geq 1} \sqrt{\frac{2c_k}{\lambda_k}} dW_k(t). \end{aligned}$$

We apply Ito's formula to  $(v(t) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(t))^2/2 + \Phi(x(t))/m$  to see that

$$\begin{aligned} &d\left[\left(v(t) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(t)\right)^2/2 + \Phi(x(t))/m\right] \\ &= \left(v(t) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(t)\right) \left(-\frac{\gamma}{m} v(t) - \sum_{k \geq 1} \sqrt{\frac{c_k}{m^2 \epsilon}} e^{-\frac{\lambda_k}{\epsilon} t} z_k(0) - \sum_{k \geq 1} \frac{c_k}{m \epsilon} g_k(t)\right) dt \\ &\quad + \left(v(t) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(t)\right) \left(\frac{\sqrt{2\gamma}}{m} dW_0(t) - \sum_{k \geq 1} \sqrt{\frac{2c_k}{m^2 \lambda_k}} dW_k(t)\right) \\ &\quad + \left(\frac{\Phi'(x(t))}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(t) + \frac{\gamma}{m^2} + \sum_{k \geq 1} \frac{c_k}{\lambda_k}\right) dt. \end{aligned}$$

We proceed to estimate the above RHS. Firstly, we invoke estimate (4.10) to find

$$\begin{aligned}
& \int_0^t \left( v(r) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right) \left( - \sum_{k \geq 1} \sqrt{\frac{c_k}{m^2 \epsilon}} e^{-\frac{\lambda_k}{\epsilon} r} z_k \right) dr \\
& \leq \sup_{0 \leq r \leq t} \left| v(r) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right| \sum_{k \geq 1} \frac{\sqrt{\epsilon c_k}}{\lambda_k} (1 - e^{-\frac{\lambda_k}{\epsilon} t}) |z_k| \\
& \leq \sup_{0 \leq r \leq t} \left| v(r) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right| \left( \sum_{k \geq 1} \frac{c_k k^{2s}}{\lambda_k} \sum_{k \geq 1} k^{-2s} z_k^2 \right)^{1/2} \\
& \leq \frac{1}{2} \sup_{0 \leq r \leq t} \left| v(r) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right|^2 + 2 \sum_{k \geq 1} \frac{c_k k^{2s}}{\lambda_k} \sum_{k \geq 1} k^{-2s} z_k^2.
\end{aligned}$$

Similarly, we have

$$g_k(r) = \int_0^r e^{-\frac{\lambda_k}{\epsilon}(r-\ell)} v(\ell) d\ell \leq \sup_{0 \leq \ell \leq r} |v(\ell)| \frac{\epsilon}{\lambda_k},$$

which implies that

$$\begin{aligned}
& \int_0^t \left( v(r) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right) \left( - \sum_{k \geq 1} \frac{c_k}{m \epsilon} g_k(r) \right) dr \\
& \leq c \left( \sum_{k \geq 1} \frac{c_k}{\lambda_k} \right) \int_0^t \sup_{0 \leq \ell \leq r} v(\ell)^2 + \sup_{0 \leq \ell \leq r} \left( \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(\ell) \right)^2 dr.
\end{aligned}$$

With regard to the martingale term, we invoke Burkholder-Davis-Gundy's inequality to estimate

$$\begin{aligned}
& \mathbb{E} \sup_{0 \leq r \leq t} \left| \int_0^r \left( v(\ell) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(\ell) \right) \left( \frac{\sqrt{2\gamma}}{m} dW_0(\ell) - \sum_{k \geq 1} \sqrt{\frac{2c_k}{m^2 \lambda_k}} dW_k(\ell) \right) \right| \\
& \leq c \left[ \left( \frac{2\gamma}{m^2} + \sum_{k \geq 1} \frac{2c_k}{m^2 \lambda_k} \right) \int_0^t \mathbb{E} \left( v(r) - \frac{1}{m} \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right)^2 dr + 1 \right].
\end{aligned}$$

Lastly, we employ Assumption 2.6 to infer

$$\int_0^t \Phi'(x(r)) \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) dr \leq c \int_0^t \Phi(x(r)) + \left( \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right)^n + 1 dr.$$

Putting everything together, we arrive at the following inequality

$$\begin{aligned}
\mathbb{E} \sup_{0 \leq t \leq T} v(t)^2 + \Phi(x(t)) & \leq c(T) \left[ 1 + \int_0^T \mathbb{E} \sup_{0 \leq r \leq t} v(r)^2 + \Phi(x(r)) dt \right. \\
& \quad \left. + \mathbb{E} \sup_{0 \leq t \leq T} \left( \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right)^2 + \mathbb{E} \sup_{0 \leq t \leq T} \left( \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right)^n \right].
\end{aligned}$$

The result now follows immediately from Gronwall's inequality if we can show that the last two terms on the above RHS is finite and independent of  $\epsilon$ . To this end, we claim that for every  $T > 0$  and  $q > 2$ , there exists a finite constant  $C(T, q) > 0$  such that

$$\mathbb{E} \sup_{0 \leq t \leq T} \left( \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right)^{2q} \leq c(T, q). \quad (4.14)$$

Recalling  $w_k(t) := \sqrt{2} \int_0^t e^{-\frac{\lambda_k}{\epsilon}(t-r)} dW_k(r)$ , similar to (4.12), we employ Holder's inequality and Lemma 3.3 to see that

$$\begin{aligned} \mathbb{E} \sup_{0 \leq t \leq T} \left( \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} w_k(r) \right)^{2q} \\ \leq \left( \sum_{k \geq 1} k^{-q_1 q^*} \right)^{2q/q^*} \sum_{k \geq 1} \mathbb{E} \left[ \sup_{0 \leq t \leq T} \left| \sqrt{\frac{c_k k^{2q_1}}{\lambda_k}} w_k(t) \right|^{2q} \right] \\ = c(T, q) \left( \sum_{k \geq 1} k^{-q_1 q^*} \right)^{2q/q^*} \epsilon^{q-1} \sum_{k \geq 1} \frac{1}{k^{q+(q\alpha-2q+1)\beta-2q_1 q}}. \end{aligned}$$

where  $\frac{1}{2q} + \frac{1}{q^*} = 1$  and  $q_1 > 0$  is a constant satisfying

$$q_1 q^* > 1 \quad \text{and} \quad q + (q\alpha - 2q + 1)\beta - 2q_1 q > 1.$$

Solving the above inequalities for  $q_1$ , we find

$$\frac{1 + (\alpha - 2)\beta}{2} + \frac{\beta}{2q} - \frac{1}{2q} > q_1 > 1 - \frac{1}{2q},$$

which is always possible thanks to the second part of Condition (b), namely,  $(\alpha - 2)\beta > 1$ . The proof is thus complete.  $\square$

**Remark 4.2.** (a) The trick of using integration by part in (4.9) was previously employed in [19, 34].

(b) The condition  $\alpha > 1$  of the diffusive regime was employed throughout the proof of Proposition 4.1, e.g. estimates (4.10), (4.11) and (4.12).

**Proposition 4.3.** Under the same Hypothesis of Theorem 2.5, assume further that  $\Phi'(x)$  is globally Lipschitz. Let  $X_\epsilon(t) = (x_\epsilon(t), v_\epsilon(t), z_{1,\epsilon}(t), \dots)$  be the solution of (2.7) with initial conditions

$$(x_\epsilon(0), v_\epsilon(0), z_{1,\epsilon}(0), z_{2,\epsilon}(0), \dots) = (x, v, z_1, z_2, \dots) \in \mathcal{H}_{-s},$$

and  $(u(t), p(t))$  be the solution of (2.8) with initial conditions  $(u(0), p(0)) = (x, v)$ . Then, for every  $T > 0$ ,

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} |x(t) - u(t)|^2 + \sup_{0 \leq t \leq T} |v(t) - p(t)|^2 \right] \rightarrow 0, \quad \epsilon \rightarrow 0.$$

The proof of Proposition 4.3 is based on that of Theorem 2.6 in [34], adapted to our infinite-dimensional setting.

*Proof.* Setting  $\bar{x}(t) := x(t) - u(t)$ ,  $\bar{v}(t) := v(t) - p(t)$ , we see that from (2.7), (2.8),  $(\bar{x}(t), \bar{v}(t))$  satisfies the following system

$$\begin{aligned} \bar{x}(t) &= \int_0^t \bar{v}(r) dr, \\ m\bar{v}(t) &= \int_0^t \left( -\gamma \bar{v}(r) + \left( \sum_{k \geq 1} \frac{c_k}{\lambda_k} \right) p(r) - [\Phi'(x(r)) - \Phi'(u(r))] - \sum_{k \geq 1} \sqrt{\frac{c_k}{\epsilon}} z_k(r) \right) dr \\ &\quad + \int_0^t \sum_{k \geq 1} \sqrt{\frac{2c_k}{\lambda_k}} dW_k(r). \end{aligned}$$

with the initial conditions  $(\bar{x}(0), \bar{v}(0)) = (0, 0)$ . Regarding  $z_k(t)$  terms, we integrate with respect to time the third equation in (2.7) to find that

$$\frac{\sqrt{\epsilon c_k}}{\lambda_k} (z_k(t) - z_k(0)) - \frac{c_k}{\lambda_k} \int_0^t v(r) dr - \sqrt{\frac{c_k}{\lambda_k}} \int_0^t dW_k(r) = -\sqrt{\frac{c_k}{\epsilon}} \int_0^t z_k(r) dr.$$

With these observations, the system of integral equations on  $(\bar{x}(t), \bar{v}(t))$  becomes

$$\begin{aligned}\bar{x}(t) &= \int_0^t \bar{v}(r) dr, \\ m\bar{v}(t) &= \int_0^t \left( - \left( \gamma + \sum_{k \geq 1} \frac{c_k}{\lambda_k} \right) \bar{v}(r) - [\Phi'(x(r)) - \Phi'(u(r))] \right) dr \\ &\quad + \sqrt{\epsilon} \sum_{k \geq 1} \frac{\sqrt{c_k}}{\lambda_k} (z_k(t) - z_k(0)).\end{aligned}\tag{4.15}$$

In the above system, we have implicitly re-arranged infinitely many terms, resulting in the cancellation of noise terms. Recalling  $c_k, \lambda_k$  from (1.3) and the norm  $\|\cdot\|_{\mathcal{H}_{-s}}$  from (2.1), this re-arrangement is possible following from (4.1) and the estimate

$$\begin{aligned}\sum_{k \geq 1} \left| \frac{\sqrt{\epsilon c_k}}{\lambda_k} (z_k(t) - z_k(0)) \right| &\leq \sum_{k \geq 1} \frac{c_k}{\lambda_k} \int_0^t |v(r)| dr + \sum_{k \geq 1} \frac{1}{\sqrt{\epsilon}} \int_0^t \sum_{k \geq 1} \sqrt{c_k} |z_k(r)| dr \\ &\quad + \sum_{k \geq 1} \sqrt{\frac{c_k}{\lambda_k}} \left| \int_0^t dW_k(r) \right| \\ &< \infty, \text{ a.s.}\end{aligned}$$

thanks to condition (D) of Assumption 2.2. We invoke the assumption that  $\Phi'$  is globally Lipschitz and Gronwall's inequality to deduce from (4.15)

$$\mathbb{E} \sup_{0 \leq t \leq T} |\bar{x}(t)| + |\bar{v}(t)| \leq C(T) \sqrt{\epsilon} \mathbb{E} \sup_{0 \leq t \leq T} \left| \sum_{k \geq 1} \frac{\sqrt{c_k}}{\lambda_k} (z_k(t) - z_k(0)) \right|.$$

The result now follows immediately from Proposition 4.4 below.  $\square$

**Proposition 4.4.** *Under the same Hypothesis of Proposition 4.3, suppose that  $X(t) = (x(t), v(t), z_1(t), \dots)$  solves (2.7) with initial conditions  $(x(0), v(0), z_1(0), \dots) \in \mathcal{H}_{-s}$ . Then,*

$$\sqrt{\epsilon} \mathbb{E} \sup_{0 \leq t \leq T} \left| \sum_{k \geq 1} \frac{\sqrt{c_k}}{\lambda_k} (z_k(t) - z_k(0)) \right| \rightarrow 0, \quad \epsilon \rightarrow 0.$$

*Proof.* From (4.7), we see that

$$\begin{aligned}\sum_{k \geq 1} \frac{\sqrt{\epsilon c_k}}{\lambda_k} |z_k(t) - z_k(0)| &\leq \sum_{k \geq 1} \frac{\sqrt{\epsilon c_k}}{\lambda_k} (e^{-\frac{\lambda_k}{\epsilon} t} - 1) |z_k(0)| + \sum_{k \geq 1} \frac{c_k}{\lambda_k} \int_0^t e^{-\frac{\lambda_k}{\epsilon} (t-r)} |v(r)| dr \\ &\quad + \sum_{k \geq 1} \sqrt{\frac{2c_k}{\lambda_k}} \left| \int_0^t e^{-\frac{\lambda_k}{\epsilon} (t-r)} dW_k(r) \right|.\end{aligned}\tag{4.16}$$

We aim to show that each series on the above RHS converges to zero in expectation as  $\epsilon \downarrow 0$ . We note that the convergence to zero of the last series follows immediately from (4.12). For the other two terms, we shall make use of the following inequality: for  $q > 0$ , there exists  $c(q) > 0$  such that for every  $x \geq 0$ , it holds that

$$1 - e^{-x} \leq c(q) x^q.\tag{4.17}$$

For a positive  $q_1 < \frac{1}{2}$  (to be chosen later), we estimate the first sum on the RHS of (4.16) as follows.

$$\begin{aligned} \sum_{k \geq 1} \sup_{0 \leq t \leq T} \frac{\sqrt{\epsilon c_k}}{\lambda_k} (e^{-\frac{\lambda_k}{\epsilon} t} - 1) z_k(0) &\leq c(T, q_1) \epsilon^{1/2-q_1} \frac{c_k^{1/2}}{\lambda_k^{1-q_1}} z_k(0) \\ &\leq c(T, q_1) \epsilon^{1/2-q_1} \left( \sum_{k \geq 1} \frac{c_k k^{2s}}{\lambda_k^{2-2q_1}} \right)^{1/2} \left( \sum_{k \geq 1} k^{-2s} z_k(0)^2 \right)^{1/2}, \end{aligned}$$

where we have used (4.17) on the first line and Holder's inequality on the second line, respectively. Recalling (1.3), we have

$$\sum_{k \geq 1} \frac{c_k k^{2s}}{\lambda_k^{2-2q_1}} = \sum_{k \geq 1} \frac{1}{k^{1+(\alpha+2q_1-2)\beta-2s}}.$$

In view of Condition (D) of Assumption 2.2, there always exists a constant  $q_1 \in (0, 1/2)$  such that  $(2q_1 - 1)\beta + (\alpha - 1)\beta - 2s > 0$ , which implies that the above RHS is finite. Similarly, we have

$$\begin{aligned} \sum_{k \geq 1} \sup_{0 \leq t \leq T} \frac{c_k}{\lambda_k} \int_0^t e^{-\frac{\lambda_k}{\epsilon}(t-r)} |v(r)| dr &\leq \sum_{k \geq 1} \sup_{0 \leq t \leq T} \frac{\epsilon c_k}{\lambda_k^2} (1 - e^{-\frac{\lambda_k}{\epsilon} t}) \sup_{0 \leq t \leq T} |v(t)| \\ &\leq c(T, q_2) \epsilon^{1-q_2} \sum_{k \geq 1} \frac{c_k}{\lambda_k^{2-q_2}} \sup_{0 \leq t \leq T} |v(t)| \\ &= c(T, q_2) \epsilon^{1-q_2} \sum_{k \geq 1} \frac{1}{k^{1+(\alpha-2+q_2)\beta}} \sup_{0 \leq t \leq T} |v(t)|. \end{aligned}$$

We invoke Condition (D) from Assumption 2.2 again to see that there exists a positive  $q_2 \in (0, 1)$  such that  $\alpha - 2 + q_2 > 0$ . Choosing such  $q_2$  implies that the series on the above RHS is convergent. We thus obtain the estimate

$$\mathbb{E} \sum_{k \geq 1} \sup_{0 \leq t \leq T} \frac{c_k}{\lambda_k} \int_0^t e^{-\frac{\lambda_k}{\epsilon}(t-r)} |v(r)| dr \leq c(T, q_2) \epsilon^{1-q_2} \mathbb{E} \sup_{0 \leq t \leq T} |v(t)| \leq c(T, q_2) \epsilon^{1-q_2},$$

where the last implication follows from Proposition 4.1. Putting everything together, we obtain the result.  $\square$

Since we will make use of exiting times, with a slightly abuse of notation, it is convenient to recall from (3.9) for  $R > 0$

$$\sigma^R = \inf_{t \geq 0} \{|u(t)| \geq R\}, \quad \text{and} \quad \sigma_\epsilon^R = \inf_{t \geq 0} \{|x(t)| \geq R\}.$$

With Proposition 4.3 in hand, we give the proof of Theorem 2.5.

*Proof of Theorem 2.5.* The arguments are almost the same as those in the proof of Theorem 2.4 and hence omitted. The only difference here is the appearance of the term  $|v(t) - p(t)|$ . Nevertheless, we note that for  $0 \leq t \leq \sigma^R \wedge \sigma_\epsilon^R$ ,

$$(u(t), p(t)) = (u^R(t), p^R(t)) \quad \text{and} \quad (x(t), v(t)) = (x^R(t), v^R(t)),$$

and thus the proof of Theorem 2.4 is applicable.  $\square$

We finally turn our attention to Theorem 2.8. The proof is relatively short and will make use of Condition (b) in Proposition 4.1.

*Proof of Theorem 2.8.* For given  $R > 0$ , let  $\sigma^R, \sigma_\epsilon^R$  be defined as in (3.9). As mentioned above, for  $0 \leq t \leq \sigma^R \wedge \sigma_\epsilon^R$ ,

$$(u(t), p(t)) = (u^R(t), p^R(t)) \quad \text{and} \quad (x(t), v(t)) = (x^R(t), v^R(t)).$$

We then have a chain of implications

$$\begin{aligned}
& \mathbb{E} \left[ \sup_{0 \leq t \leq T} |x(t) - u(t)|^q + |v(t) - p(t)|^q \right] \\
&= \mathbb{E} \left[ \left( \sup_{0 \leq t \leq T} |x(t) - u(t)|^q + |v(t) - p(t)|^q \right) 1_{\{\sigma^R \wedge \sigma_\epsilon^R < T\}} \right] \\
&\quad + \mathbb{E} \left[ \left( \sup_{0 \leq t \leq T} |x(t) - u(t)|^q + |v(t) - p(t)|^q \right) 1_{\{\sigma^R \wedge \sigma_\epsilon^R > T\}} \right] \\
&\leq \mathbb{E} \left[ \left( \sup_{0 \leq t \leq T} |x(t) - u(t)|^q + |v(t) - p(t)|^q \right) 1_{\{\sigma^R \wedge \sigma_\epsilon^R < T\}} \right] \\
&\quad + \mathbb{E} \left[ \sup_{0 \leq t \leq T} |x^R(t) - u^R(t)|^q + |v^R(t) - p^R(t)|^q \right].
\end{aligned}$$

On one hand, in view of Proposition 4.3, since  $1 \leq q < 2$ , it holds that

$$\mathbb{E} \left[ \sup_{0 \leq t \leq T} |x^R(t) - u^R(t)| + |v^R(t) - p^R(t)| \right] \rightarrow 0, \quad \epsilon \rightarrow 0.$$

On the other hand, we invoke Holder's inequality with  $\frac{q}{2} + \frac{1}{q^*} = 1$  to estimate

$$\begin{aligned}
& \mathbb{E} \left[ \left( \sup_{0 \leq t \leq T} |x(t) - u(t)|^q + |v(t) - p(t)|^q \right) 1_{\{\sigma^R \wedge \sigma_\epsilon^R < T\}} \right] \\
&\leq c \left( \mathbb{E} \left[ \sup_{0 \leq t \leq T} x(t)^2 + v(t)^2 \right] + \mathbb{E} \left[ \sup_{0 \leq t \leq T} u(t)^2 + p(t)^2 \right] \right)^{q/2} \left( \mathbb{P} \{ \sigma^R \wedge \sigma_\epsilon^R < T \} \right)^{1/q^*}.
\end{aligned}$$

Notice that by Markov's inequality, we have

$$\begin{aligned}
\mathbb{P} \{ \sigma^R \wedge \sigma_\epsilon^R < T \} &\leq \mathbb{P} \{ \sigma^R < T \} + \mathbb{P} \{ \sigma_\epsilon^R < T \} \\
&\leq \mathbb{P} \left\{ \sup_{0 \leq t \leq T} |u(t)| \geq R \right\} + \mathbb{P} \left\{ \sup_{0 \leq t \leq T} |x(t)| \geq R \right\} \\
&\leq \frac{\mathbb{E} [\sup_{0 \leq t \leq T} |u(t)|^2] + \mathbb{E} [\sup_{0 \leq t \leq T} x(t)^2]}{R^2}.
\end{aligned}$$

The result now follows immediately from (4.4) and Proposition 4.1 by first taking  $R$  sufficiently large and then shrinking  $\epsilon$  further to zero. The proof is thus complete.  $\square$

## 5. DISCUSSION

We have established rigorous results on the asymptotical analysis of an infinite-dimensional GLE when the memory kernel  $K(t)$  has a power-law decay, i.e.  $K(t) \sim t^{-\alpha}$  as  $t \rightarrow \infty$ . With regards to the small-mass limit, we are able to obtain the convergence in probability of the GLE for every exponent constant  $\alpha > 0$ . However, in the white-noise limit, a similar convergence was established only when  $\alpha > 1$ . The method that we employed was not able to extend the result when  $\alpha \in (0, 1]$ , which is interestingly also the barrier for the unique ergodicity of (1.6) [12]. As mentioned earlier in Remark 4.2, our technique in the white-noise limit requires that the memory be integrable ( $\alpha > 1$ ) for the analysis of the solutions as well as the asymptotical behaviors. It therefore remains an open question whether the solution's energy is still bounded uniformly and there exists a limiting system.

Finally, another question for future works is whether one can take both limits in sequence, which means that the small-mass variable  $m$  is written as an order of  $\epsilon$ , the white-noise variable. It is not clear that the theorems presented in this work combined together are able to produce an explicit answer. We note that a similar study in finite-dimensional setting was carried out in [30]. Yet, we have not been able to see if the same method can be applied to our infinite system. We believe handling this case will require a more substantial work.

## ACKNOWLEDGEMENT

The author would like to thank Nathan Glatt-Holtz, David Herzog and Scott McKinley for many fruitful discussions. The author is also grateful for support through grant NSF DMS-1644290.

## REFERENCES

- [1] Joseph Abate and Ward Whitt. Infinite-series representations of Laplace transforms of probability density functions for numerical inversion. *Journal of the Operations Research Society of Japan*, 42(3):268–285, 1999.
- [2] Sergio Albeverio, Franco Flandoli, and Yakov G Sinai. *SPDE in Hydrodynamics: Recent Progress and Prospects: Lectures given at the CIME Summer School held in Cetraro, Italy, August 29-September 3, 2005*. Springer, 2008.
- [3] Douglas Blount. Comparison of stochastic and deterministic models of a linear chemical reaction with diffusion. *The Annals of Probability*, pages 1440–1462, 1991.
- [4] Sandra Cerrai and Mark Freidlin. On the Smoluchowski-Kramers approximation for a system with an infinite number of degrees of freedom. *Probability theory and related fields*, 135(3):363–394, 2006.
- [5] Sandra Cerrai and Mark Freidlin. Smoluchowski-Kramers approximation for a general class of SPDEs. *Journal of Evolution Equations*, 6(4):657–689, 2006.
- [6] Sandra Cerrai and Michael Salins. Smoluchowski-Kramers approximation and large deviations for infinite dimensional gradient systems. *Asymptotic Analysis*, 88(4):201–215, 2014.
- [7] Sandra Cerrai and Michael Salins. Smoluchowski-Kramers approximation and large deviations for infinite-dimensional nongradient systems with applications to the exit problem. *The Annals of Probability*, 44(4):2591–2642, 2016.
- [8] Monica Conti, Vittorino Pata, and Marco Squassina. Singular limit of differential systems with memory. *Indiana University mathematics journal*, pages 169–215, 2006.
- [9] Giuseppe Da Prato and Jerzy Zabczyk. *Stochastic equations in infinite dimensions*. Cambridge university press, 2014.
- [10] Mark Freidlin. Some remarks on the Smoluchowski-Kramers approximation. *Journal of Statistical Physics*, 117(3):617–634, 2004.
- [11] Stefania Gatti, Claudio Giorgi, and Vittorino Pata. Navier-Stokes limit of Jeffreys type flows. *Physica D: Non-linear Phenomena*, 203(1-2):55–79, 2005.
- [12] Nathan Glatt-Holtz, David Herzog, Scott McKinley, and Hung Nguyen. The Generalized Langevin Equation with a power-law memory in a nonlinear potential well. *arXiv preprint arXiv:1804.00202*, 2018.
- [13] Nathan Glatt-Holtz, Jonathan C Mattingly, and Geordie Richards. On unique ergodicity in nonlinear stochastic partial differential equations. *Journal of Statistical Physics*, 166(3-4):618–649, 2017.
- [14] Nathan Glatt-Holtz and Mohammed Ziane. Strong pathwise solutions of the stochastic Navier-Stokes system. *Advances in Differential Equations*, 14(5/6):567–600, 2009.
- [15] Maurizio Grasselli and Vittorino Pata. Uniform attractors of nonautonomous dynamical systems with memory. In *Evolution equations, semigroups and functional analysis*, pages 155–178. Springer, 2002.
- [16] Martin Hairer, Jonathan C Mattingly, and Michael Scheutzow. Asymptotic coupling and a general form of Harris’ theorem with applications to stochastic delay equations. *Probability Theory and Related Fields*, 149(1):223–259, 2011.
- [17] David P Herzog, Scott Hottovy, and Giovanni Volpe. The small-mass limit for Langevin dynamics with unbounded coefficients and positive friction. *Journal of Statistical Physics*, 163(3):659–673, 2016.
- [18] Christel Hohenegger and Scott A McKinley. Fluid-particle dynamics for passive tracers advected by a thermally fluctuating viscoelastic medium. *Journal of Computational Physics*, 340:688–711, 2017.
- [19] Scott Hottovy, Austin McDaniel, Giovanni Volpe, and Jan Wehr. The Smoluchowski-Kramers limit of stochastic differential equations with arbitrary state-dependent friction. *Communications in Mathematical Physics*, 336(3):1259–1283, 2015.
- [20] Scott Hottovy, Giovanni Volpe, and Jan Wehr. Noise-induced drift in stochastic differential equations with arbitrary friction and diffusion in the Smoluchowski-Kramers limit. *Journal of Statistical Physics*, 146(4):762–773, 2012.
- [21] Scott Hottovy, Giovanni Volpe, and Jan Wehr. Thermophoresis of Brownian particles driven by coloured noise. *EPL (Europhysics Letters)*, 99(6):60002, 2012.
- [22] Kiyosi Itô. Stationary random distributions. *Memoirs of the College of Science, University of Kyoto. Series A: Mathematics*, 28(3):209–223, 1954.
- [23] Jean Jacod. *Calcul stochastique et problemes de martingales*, volume 714. Springer, 2006.
- [24] Ioannis Karatzas and Steven Shreve. *Brownian motion and stochastic calculus*, volume 113. Springer Science & Business Media, 2012.



- [25] Samuel C Kou. Stochastic modeling in nanoscale biophysics: subdiffusion within proteins. *The Annals of Applied Statistics*, pages 501–535, 2008.
- [26] Hendrik Anthony Kramers. Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica*, 7(4):284–304, 1940.
- [27] Rep Kubo. The fluctuation-dissipation theorem. *Reports on progress in physics*, 29(1):255, 1966.
- [28] R Kupferman, AM Stuart, JR Terry, and PF Tupper. Long-term behaviour of large mechanical systems with random initial data. *Stochastics and Dynamics*, 2(04):533–562, 2002.
- [29] Raz Kupferman. Fractional kinetics in Kac–Zwanzig heat bath models. *Journal of statistical physics*, 114(1):291–326, 2004.
- [30] Soon Hoe Lim and Jan Wehr. Homogenization of a Class of Non-Markovian Langevin Equations with an Application to Thermophoresis. *arXiv preprint arXiv:1704.00134*, 2017.
- [31] Thomas G Mason and DA Weitz. Optical measurements of frequency-dependent linear viscoelastic moduli of complex fluids. *Physical review letters*, 74(7):1250, 1995.
- [32] Scott A McKinley and Hung D Nguyen. Anomalous Diffusion and the Generalized Langevin Equation. *arXiv preprint arXiv:1711.00560*, 2017.
- [33] Hazime Mori. A continued-fraction representation of the time-correlation functions. *Progress of Theoretical Physics*, 34(3):399–416, 1965.
- [34] M Ottobre and GA Pavliotis. Asymptotic analysis for the generalized Langevin equation. *Nonlinearity*, 24(5):1629, 2011.
- [35] E Pardoux and A Yu Veretennikov. On Poisson equation and diffusion approximation 2. *The Annals of Probability*, 31(3):1166–1192, 2003.
- [36] Grigorios A Pavliotis. *Stochastic Processes and Applications: Diffusion Processes, the Fokker-Planck and Langevin Equations*, volume 60. Springer, 2014.
- [37] Marian Von Smoluchowski. Drei vortrage uber diffusion. Brownsche bewegung und koagulation von kolloidteilchen. *Z. Phys.*, 17:557–585, 1916.
- [38] Robert Zwanzig. Nonlinear generalized Langevin equations. *Journal of Statistical Physics*, 9(3):215–220, 1973.