

Agency and the physics of numbers

John M. Myers*

Harvard School of Engineering and Applied Sciences, Cambridge, MA 02138, USA.

F. Hadi Madjid†

82 Powers Road, Concord, MA 01742

January 27, 2022

Analogous to Gödel’s incompleteness theorems is a theorem in physics to the effect that the set of explanations of given evidence is uncountably infinite. An implication of this theorem is that contact between theory and experiment depends on activity beyond computation and measurement—physical activity of some agent making a guess. Standing on the need for guesswork, we develop a representation of a symbol-handling agent that both computes and, on occasion, generates a guess in interaction with an oracle. We show: (1) how physics depends on such an agent to bridge a logical gap between theory and experiment; (2) how to represent the capacity of agents to communicate numerals and other symbols, and (3) how that communication is a foundation on which to develop both theory and implementation of spacetime and related competing schemes for the management of motion.

* myers@seas.harvard.edu

† gailmadjid@comcast.net

I. INTRODUCTION

Schrödinger, in his 1954 book “Nature and the Greeks,” laments [1]:

... science in its attempt to describe and understand Nature simplifies this very difficult problem. The scientist subconsciously, almost inadvertently, simplifies his problem of understanding Nature by disregarding or cutting out of the picture to be constructed, himself, his own personality, the subject of cognizance. Inadvertently the thinker steps back into the role of an external observer. This facilitates the task very much. But it leaves gaps, enormous lacunae, leads to paradoxes and antimonies whenever, unaware of this initial renunciation, one tries to find oneself in the picture or to put oneself, one’s own thinking and sensing mind, back into the picture. This momentous step ... has consequences. So, in brief, we do not belong to this material world that science constructs for us. We are not in it, we are outside. We are only spectators.

A start toward recognizing the role in physics of the physicist can be found in Gödel’s incompleteness theorems that show that mathematical logic is uncloseably open to additional axioms. Gödel exploited the writing of mathematics as strings of symbols, as did Turing in his definition of computability. More recently, we recognized that both evidence and its explanations, as expressed in quantum theory, are written in strings of symbols, and from this recognition we proved another incompleteness theorem, to do with explanations in terms of wave functions and operators that fit evidence expressed as parameter-dependent probabilities:

Theorem: The set of inequivalent explanations that exactly fit given probabilities is uncountably infinite [2].

Logically, there can be no unique explanation. Of course, physicists collaborate and compete with one another to develop a dominant explanation. By virtue of the theorem however, the assertion of particular explanations reaches beyond logic; the assertion is a conjecture, an hypothesis, in short, a **guess**, even if a community coalesces around it. And the guess is forever subject to refutation, so that physics involves an endless open cycle in which physicists act as agents that make guesses [3]. Here, not as an opinion but stemming from a proof within quantum theory, is a role in physics for the physicist.




FIG. 1. Agent's open cycle of guessing and testing

Facing up to symbols and the agents that handle them in physical investigations opens a whole new arena of physics, ready for exploration. Agency in physics has been discussed from several points of view [4–9]. We focus on agents that perceive and use symbols. But how, in the context of physics, can one speak of symbols? Physics has sometimes been viewed as a quest for “fundamental elements”, mainly particles and fields, from which “all physical descriptions” could be constructed; however, this quest has been challenged, and in its place one can now enjoy more latitude, thanks in part to notions of “emergence” made prominent in explaining phenomena of condensed matter [10]. Availing ourselves of the latitude to take symbols as elements of description from computer science and engineering, here we advance notions of symbols and the agents that handle them, with implications for theories of spacetime and their implementations.

II. AGENTS PERCEIVE SYMBOLS

In order to introduce into physics the notions of symbols and the agents that perceive them, we need first to distinguish perception from measurement. Here is a homely experiment on perception. I sit at a table and pick up a magazine with pictures that I have not already looked at. I shut my eyes and turn the cover to expose the first page, still with my eyes shut. Then I blink, very briefly

opening my eyes and quickly shutting them again. I have an fleeting image of seeing a picture of a tiger. Then I repeat the process. Two or three glances later, the image stabilizes, not into a tiger but into a woman sitting on a couch. This little story illustrates the claim in Pattern Theory of a two-way flow between brain and eye: the brain proposes forms that the eye confirms or rejects [11]. In some circumstances, I can make definite, stable perceptions, while outside these circumstances my perceptions are unstable. In circumstances in which I achieve a stable perception, that perception might be wrong—as when confronted by an illusion, as in the Ames demonstrations [12], but my perception is not uncertain in the way a measurement is, with its systematic and statistical uncertainties.

In theoretical work at the blackboard, physicists perceive the symbols—numerals, letters, plus and minus signs, etc.—in which they write and read both evidence and its explanations. But what about experiments at the workbench in which they measure something? To measure something, one has to first perceive what is to be measured. To measure the voltage on a wire at a moment, a person must perceive the voltmeter, the wire to be measured, and the moment at which to make the measurement, in distinction to other devices, other wires, and other moments.

In particular, what about measuring the physical transmission of symbols? Suppose Person A perceives the words in an email sent by Person B. A third Person C can measure the physical transmission of the email from A to B. Person C may measure voltage levels at moments when these convey binary symbols (bits) that code a higher level symbol (letter of the alphabet or word), along with timing relations among various voltages. Person C's measurements produce numbers accompanied by uncertainties that contrast with Person B's perception of letters on the computer screen. The investigative Person C who measures must also perceive. For example, Person C has to perceive both what is to be measured and the devices used to measure it.

To say that one *perceives* a symbol is to say that one registers something, distinct from something else. When perception matters, the symbol perceived matters to one's next action, distinct from something else that would contribute to a different action. We will discuss circumstances necessary to cases in which, apart from occasional failures, the perception of symbols is unambiguous; for example, one perceives a dog as distinct from a cat. While the cat you pet is more than a symbol, to recognize the cat as a cat is to recognize 'cat' as a symbol.

Two interrelated questions: what sort of entity has the capability to perceive, and what sort of entity is it that is perceived? Not just people, but other living organisms make distinctions, e.g. between self and non-self, and thus also perceive. To perceive is to register a sensing in

some category and not in others. Without an entity's capacity to make such distinctions, the entity cannot be said to perceive. We call such an entity a *symbol-handling agent*, whether it is a living organism or a machine. For the second question—what sort of entity is perceived—we propose that whatever is perceived is reasonably called a *symbol*. Perception assigns a sensing to a category, without measuring it. Such categories are elements of thought, having much in common with mathematical entities that are not measured but are defined in relation to other such entities. Words are symbols. For perception as we mean it in this report, humans perceive by labeling with words. Wordless organisms that perceive must have other ways of labeling, that is, of assigning symbols.

Symbols and *symbol-handling agents* are terms of description available at widely varying levels of detail. I may see myself as handling symbols, and I may inquire into evidence of symbol handling on the part of mantra within my cells. A word as a symbol can be coded in letters of an alphabet which in turn can be coded in bits.

What capabilities, besides the perception of symbols, are to be ascribed to an 'agent'? I see a core capability of agency as a feature of the human condition: I respond to surprises. When the surprise comes to me, my sense is that it comes in part from the unknowable but also in part from me, because I've been working on something, and the surprise from the unknowable and the surprise from me combine to come up with a guess. We think of an agent as taking steps, one after another, and as equipped with a memory. Each "next step" of an agent is influenced both by the contents of its memory and by an inflow of symbols from an environment that includes other agents, and also by a logically undetermined "oracle" external to the agent [13]. Guesses emerge from an agent interacting with an oracle, the workings of which we refrain from trying to penetrate.

III. STEERING RHYTHMS OF PERCEPTION

People delegate a variety of symbol-handling tasks to mechanized agents, such as the computer at which I now write. Although stripped of emotional capacity, dead, computer networks are important to understanding symbol-handling agents in two ways. First, human handling of symbols leaves records in computer networks that serve as evidence. Second, computer networks exhibit rhythms of symbol handling important to agency in general. Computer networks pervade the modern technological world, and not only in the form of the Internet. When I bring my car

to the repair shop, the first thing checked is often the computer network that mediates between my eyes, hands, feet, and the mechanisms for steering, braking, and communication. Records of mechanized agents visible in computer based records are a heretofore overlooked topic for physical investigation. The computers that work as networked agents have much to teach us about the perception of digital symbols, to do with rhythms of perception that require continual maintenance, guided by responses to analog deviations that are logically undetermined.

Analog machinery is idiosyncratic: two people using a slide rule get slightly different answers. The huge strength of digital, as opposed analog, computation comes from the capacity of a digital computer to make exact calculations by means of inexact machinery. Insofar as a person using the computer is concerned, variations in the behavior of transistors leave a computer's traffic in 0's and 1's unaffected. This is because the basic unit of memory, the flip-flop, works like a hinge that can be recognized unambiguously as turned one way or the other, regardless of imperfections in its manufacture and operation. Or think of a chess game in a player recognizes the placement of a pawn on a square of the board in spite of small deviations of the pawn from the center of the square. But, as we shall see, recognizing symbols unambiguously is possible only within a system of analog adjustment. The hand guided by the eye places the pawn more or less in the middle of a square of a chess board—a performance in which noticing variations is necessary to the adjustment of motion.

Analogous steering in the handling of digital symbols shows up with special clarity in the physics of computer networks, such as the networks used by human agents to mediate communications that generate evidence and convey explanations.

IV. MECHANIZED SYMBOL HANDLERS

In order to distinguish between what can be computed and what must come from beyond computation (as guesses from interaction with an oracle), we imagine the extreme case of a symbol-handling agent that, while open to guessing, possesses maximal computational capacity. Unconcerned with practical limits on computing imposed by limits on memory or by limits on the rate at which an agent computes, we formulate a mechanical agent having the computational capability of a Turing machine. The Turing machine, however, requires modification to offer a place for interaction with an oracle and for communication with other such machines.

In his 1936 paper, Turing briefly introduced an alternative machine called a *choice machine*,

contrasted with the usual Turing machine that Turing called an a-machine:

If at each stage the motion of a machine ... is completely determined by the [memory] configuration, we shall call the machine an “automatic machine” (or a-machine). For some purposes we might use machines (choice machines or c-machines) whose motion is only partially determined by the configuration When such a machine reaches one of these ambiguous configurations, it cannot go on until some arbitrary choice has been made by an external operator. This would be the case if we were using machines to deal with axiomatic systems. [14].

We picture a mechanical symbol-handling agent as implementing a c-machine modified to take part in a communications network. We call the c-machine modified in this way, with some additional modifications to be described shortly, a *Choice Machine*. We posit that on occasion an “oracle” writes a symbol onto the scanned square of the Turing tape of the agent’s Choice Machine *privately*, in the sense that the symbol remains unknown to other agents unless and until the Choice Machine that receives the chosen symbol reports it to others [15]. There is no limit to the number of symbols that the oracle can write [2].

V. CHOICE MACHINES PAIRED BY COMMUNICATIONS CHANNELS

What affects one agent as a symbol need not be a symbol to another agent, but there are special situations in which a pair of agents can communicate by sending symbols back and forth from one agent to the other. In this report we attend to the pairing of one agent with another agent by means of a two-way communications channel over which symbols that mean something to one agent arrive at another agent that coherently responds.

Characterizations of flow of symbols, including their necessary synchronization, were discussed earlier [16–18]. Here we discuss the mathematical forms—one might say data structures—that underpin concepts of space, time, and spacetime as used by physicists, and that also underpin arrangements of devices that implement these concepts. We claim that these mathematical forms are important to thinking about how organisms, human and non-human, manage motion.

A. Symbol handling without metric

To develop the theory of symbol handling to be presented below, we avoid topics critical to the engineering of physical networks. Thus we give no attention to how symbols might propagate from agent to agent, e.g. by classical or quantum fields defined with respect to some coordinate system with a metric tensor. Rather, our approach is non-metric; one might say topological. We seek a theory of agents that maintain records of transmissions and of receptions of symbols from one to another in a situation that need have no coordinate system available. Then there can be no central clock, no assumption of any spacetime manifold and hence no metric tensor. Indeed we must, so to speak, wipe these off the blackboard in order to get at the element of novelty in attending to networks of symbol-handling agents. Eschewing any reference to *how* symbols are transmitted, propagated or received, we postulate that symbol-handling agents register moments of transmission and reception of symbols, each agent using its own count of moments as a local clock.

B. Forms of records used by agents to manage their motion

To think about agents managing their motions, we explore cases in which the Choice Machine expressing a symbol-handling capability is augmented by a second tape that we call a *clock tape*. Like the Turing tape, the clock tape is marked into squares, each of which can hold a single symbol, and at any moment a Choice Machine scans a single square of the clock tape, but the motion of the clock tape never reverses: at each move the scanned square of the clock tape shifts one place to the right. Consider symbols arriving one after another at an agent A from an agent B . Agent A can copy each symbol in the moment it arrives onto the scanned square of A 's clock tape. In this way A maps the dynamic, temporal order of arrival of the symbols to a static, spatial order along the clock tape.

Suppose that at some but not all of its moments, an agent A receives a symbol from B and writes the received symbol on A 's clock tape. If the symbols arrive only occasionally, A records these arriving symbols sparsely, so that successive symbols occupy not successive squares of A 's clock tape but on squares separated by stretches of A 's tape, stretches on which A may have written symbols for its own use or symbols arriving from agents other than B . Consider a stretch of A 's clock tape after such a performance. The ratio of the number of symbols recorded from B to the

number of squares on A 's clock tape gives the average frequency ratio of symbol arrivals from B to steps of A . Such frequency ratios are the recorded evidence of the speed ratio of A 's receptions from B relative the speed of A 's stepping along its clock tape. These ratios are the basic form of evidence of motion.




FIG. 2. Clock-tape record of “frequency”

The example in Fig. 2 shows a relative frequency

$$\frac{\text{number of } B\text{-symbols received}}{\text{number of } A\text{-moves}} \approx \frac{1}{4}$$

Now we turn from frequency to *distance*. The basic form for distance in terms of records on a tape is in terms of an *echo count*. An echo count is the number of squares from a symbol sent to the return of an echo. That is, agent A , at the moment of scanning square n of its clock tape, transmits a symbol to B , B echoes the symbol back to A , and the echoed symbol arrives as A scans square $n + k$; then k is an *echo count* from A to B and back.

The notion of *echo count* is a basis for distance. As discussed in more detail in [16, 17], in special cases echo counts it can be aligned with distances as defined by Einstein in special relativity, but echo count gives a property of records that is more basic than relativistic distance, in that the concepts needs no assumption of any spacetime metric.

In discussing a lone agent, one might want to ask “what belongs to the agent?”, for example a record of odds that the agent assigns to a bet could be said to “belong to the agent.” But when agents communicate, some records are shared, so that ownership can be shared; communicating agents differ drastically from lone agents. We take as fundamental the capacity of an agent, on

occasion, to pair with another agent in communicating symbols. Then the question “what belongs to an agent?” becomes embedded in a context of asking “what strings of symbols flow from one agent to another?”. Strings of symbols bind the agents without belonging totally to one or the other agent. It is a potentially interesting discipline to explore this and related questions, not just in engineered systems but also in organisms, human and non-human, to explore situations in which one can ask: “which record, what agents have access to the record, where and when (along their own tapes) do they have this access?”

C. Evolution of forms of records

The analysis of the form of records of frequency and distance has the following implication. Given that agents, by definition, can be seen as in interaction with oracles that are logically undetermined, and given that records of symbols in the memories of agents are the basic form to frequency and distance, we arrive at the insight that agents managing speed and distance operate not as a single system of a logic of symbols landing on and departing from the squares of tapes, but like extended arithmetics, they form an uncountable set, involving the evolution through the logically undetermined entrance of symbols, e.g. from oracles.

We are interested in various possible paths of the evolution of ways in which agents manage motion, especially in cases in which agents leave measurable tracks in the form of their clock tapes. For example, human agents develop evolving national and international time standards. One such thread of evolution consists in the concepts and the implementation of these concepts in International Atomic Time (TAI). Recall that TAI rests on its base reference frequency of the electromagnetic radiation resonant with the transition between hyperfine energy levels of cesium 133 [16]. This reference frequency is employed, either directly or indirectly, in almost all physical investigation. But Cesium (or its contemplated replacement by an optical frequency) is by no means the only type of reference imaginable or potentially fruitful. Other such threads of evolution of symbolic communication among agents are candidates for exploration, and some are in use. For example, the cutting-edge optical clocks now operating in National Metrology Laboratories have an instability of about 1/100 of the cesium clocks that define the second. Therefore it makes no sense to compare two such optical clocks by comparing each to the second: one compares them directly. There is an interesting potential for alternative schemes of measuring motion based on symbol exchange in the Laser Interferometer Gravitational-Wave Observatory (LIGO) experiment

[19]. Still other schemes are candidates for explorations of the management of motion by biological organisms, both human and non-human. “The frog starves to death in the presence of dead flies, not because he is fastidious, but because he does not see them” [20]. Some recent crashes of self-driving cars have been traced to “not seeing still objects.” What alternative to spacetime is it in which the frog lives?

VI. DISCUSSION

While we needed to start, as in [2], with a “lone agent”, a lone agent is like “one hand clapping”: Symbol-handling agents can’t function without being paired with other such agents via back-and-forth communication of symbols. As discussed in [16], conflicts in synchronization place severe limitations the potential for a given agent to communicate with other agents, and the potential for conflicts generally limits the number channels that can operate concurrently to pair one agent with another. Agent A may have to step out of communication with B if it is to enter communication with C .

As we see it, the notion of objectivity has to do with expectations of agreement among records made by different agents, rather than a notion of behavior devoid of agent participation.

The clock tape along which an agent steps is the core image of “local time,” without encumbering it with the notion, which for 40 years we have found baffling, of any “global time” [16].

Guesses by an agent, here thought of as arising in the interaction of an agent with an oracle, influence not only how an agent resolves certain choices but also the agent’s organization; that is, the agent’s capacity to perceive a situation that calls for a choice.

ACKNOWLEDGMENT

We thank Kenneth Augustyn, Paul Benioff, and Louis Kauffman for instructive comments.

[1] E. Schrödinger, “Nature and the Greeks”, Cambridge University Press (1954):

[2] J. M. Myers and F. H. Madjid, “Incompleteness theorem for physics,” <http://arxiv.org/abs/1803.10589> (2018).

[3] In earlier times, scientists were more used to the open cycle of guesswork.

Hence you have the scientific hypothesis; and its value will be proportionate to the care and completeness with which its basis had been tested and verified. It is in these matters as in the commonest affairs of practical life: the guess of the fool will be folly, while the guess of the wise man will contain wisdom. In all cases, you see that the value of the result depends on the patience and faithfulness with which the investigator applies to his hypothesis every possible kind of verification [T. H. Huxley “The method of scientific investigation,” (1863) <http://human-nature.com/darwin/huxley/chap5.html>]

- [4] H. P. Stapp, “Science of consciousness and the Hard Problem,” *The Journal of Mind and Behavior*, **18**, 171–193 (1997).
- [5] C. M. Caves, C. A. Fuchs, and R. Schack, “Quantum probabilities as Bayesian probabilities,” *Phys. Rev. A* **65**, 022305 (6) (2000). DOI: 10.1103/PhysRevA.65.022305.
- [6] C. A. Fuchs, “Notwithstanding Bohr, the Reasons for QBism,” *Mind and Matter*, **15**, issue 2, 245–300 (2017).
- [7] H. J. Briegel and G. De las Cuevas, “Projective simulation for artificial intelligence,” *Scientific Reports* **2**, 400 (2012); arXiv:1104.3787v3, 8 Feb 2013.
- [8] K. B. Laskey, “Quantum Physical Symbol Systems,” *Journal of Logic, Language and Information*, **15**, 109–154 (2006).
- [9] C. Fields, D.D. Hoffman, C. Prakash, M. Singh, “Conscious agent networks: Formal analysis and application to cognition,” *Cognitive Systems Research*, **47**, 186–213 (2018) <https://doi.org/10.1016/j.cogsys.2017.10.003>
- [10] R. B. Laughlin and D. Pines, “The theory of everything,” *PNAS* January 4, 2000. 97 (1) 28-31; <https://doi.org/10.1073/pnas.97.1.28>.
- [11] D. Mumford and A. Desolneux, *Pattern Theory: A Stochastic Analysis of Real-World Signals* (A. K. Peters, Ltd., Natick, MA, 2010)
- [12] Wikipedia, “Ames room,” https://en.wikipedia.org/w/index.php?title=Ames_room&oldid=754511822 (2016)
- [13] To express the entry into an agent of a guess, we make a loose adaptation of the notion of an oracle introduced by Turing in his 1938 Princeton thesis, as reprinted in A. W. Appel, ed., *Alan Turings Systems of Logic, the Princeton Thesis*, (Princeton University Press, Princeton, NJ, 2012).

- [14] A. M. Turing, “On computable numbers with an application to the Entscheidungsproblem,” Proc. London Math. Soc., Series 2, **42**, 230–265 (1936)
- [15] For a brief summary of Turing machine structure and function, see F. H. Madjid and J. M. Myers, “Symbols of a cosmic order,” Ann. Physics **373**, 374–389 (2016); also at <https://arxiv.org/abs/1607.07437>. For more detail see [14].
- [16] J. M. Myers and F. H. Madjid, “Distinguishing between evidence and its explanations in the steering of atomic clocks,” Annals of Physics **350**, 29–49 (2014); arXiv:1407.8020.
- [17] J. M. Myers and F. H. Madjid, “Unpredictability and the transmission of numbers,” Quantum Inf. Process. **15**, 1057–1067 (2016); also at <http://arxiv.org/abs/1508.01119>.
- [18] F. H. Madjid and J. M. Myers, “Symbols of a cosmic order,” Ann. Physics **373**, 374–389 (2016); also at <https://arxiv.org/abs/1607.07437>.
- [19] B. P. Abbott *et al.*, “Observation of Gravitational Waves from a Binary Black Hole Merger,” Phys. Rev. Letters, **116**, 061102 (2016).
- [20] J. Y. Lettvin, lecture at the Massachusetts Institute of Technology, 1965; also see J. Y. Lettvin, H. R. Maturana, W. S. McCulloch, and W. R. Pitts, “What the frog’s eye tells the frog’s brain,” Proc. IEEE, Nov. 1940–1951 (1959).