

# A REMARK ON $\mathcal{H}^1$ MARTINGALES

HARDY HULLEY AND JOHANNES RUF

**ABSTRACT.** The space of  $\mathcal{H}^1$  martingales is interesting because of its duality with the space of BMO martingales. It is straightforward to show that every  $\mathcal{H}^1$  martingale is a uniformly integrable martingale. However, the converse is not true. That is to say, some uniformly integrable martingales are not  $\mathcal{H}^1$  martingales. This brief note provides a template for systematically constructing such processes.

## 1. INTRODUCTION

Let  $(\Omega, \mathcal{F}, \mathfrak{F}, \mathbb{P})$  be a filtered probability space, whose filtration  $\mathfrak{F} = (\mathcal{F}_t)_{t \geq 0}$  is right-continuous. By assumption, all processes are defined on  $(\Omega, \mathcal{F}, \mathfrak{F}, \mathbb{P})$  and have càdlàg sample paths. The families of local martingales and uniformly integrable martingales are denoted by  $\mathcal{M}_{\text{loc}}$  and  $\mathcal{M}$ , respectively. Given  $p \geq 1$ , let  $\mathcal{H}^p$  be the family of local martingales  $M \in \mathcal{M}_{\text{loc}}$ , for which  $\mathbb{E}([M]_{\infty}^{p/2}) < \infty$ . Then  $\mathcal{H}^p$  forms a Banach space, when endowed with the norm  $\|\cdot\|_p$ , defined by

$$\|M\|_p := \left( \mathbb{E}([M]_{\infty}^{p/2}) \right)^{1/p},$$

for all  $M \in \mathcal{H}^p$ . Let  $\mathcal{H}_{\text{loc}}^p$  denote the family of local martingales that are locally in  $\mathcal{H}^p$ .

The space  $\mathcal{H}^1$  has several interesting features. For example,  $\mathcal{H}^1$  contains  $\mathcal{H}^2$  as well as all local martingales with integrable variation (see e.g Protter 2005, Theorem IV.49). It can also be shown that  $\mathcal{H}^2$  and the family of bounded (uniformly integrable) martingales are both dense in  $\mathcal{H}^1$  (see e.g Protter 2005, Theorem IV.50). In addition,  $\mathcal{M}_{\text{loc}} \subseteq \mathcal{H}_{\text{loc}}^1$  (see e.g Protter 2005, Theorem IV.51). However, the most significant result is the duality between  $\mathcal{H}^1$  and the space of BMO martingales. In detail, let

$$\text{BMO} := \left\{ M \in \mathcal{H}^2 \mid \mathbb{E}((M_{\infty} - M_{\tau-})^2) \leq c^2, \text{ for all } \tau \in \mathfrak{G} \text{ and some } c \in \mathbb{R}_+ \right\},$$

where  $\mathfrak{G}$  denotes the family of stopping times defined on  $(\Omega, \mathcal{F}, \mathfrak{F}, \mathbb{P})$  and subject to the convention  $M_{0-} := 0$ , for all  $M \in \mathcal{M}_{\text{loc}}$ . Then BMO becomes a Banach space, when endowed with the norm  $\|\cdot\|_{\text{BMO}}$ , defined by

$$\|M\|_{\text{BMO}} := \sup_{\tau \in \mathfrak{G}} \sqrt{\frac{\mathbb{E}((M_{\infty} - M_{\tau-})^2)}{\mathbb{P}(\tau < \infty)}},$$

for all  $M \in \text{BMO}$ . It follows that  $(\mathcal{H}^1)^* \simeq \text{BMO}$  (see e.g Protter 2005, Theorem IV.55).<sup>1</sup>

Now, suppose  $M \in \mathcal{H}^1$ , in which case the Burkholder-Davis-Gundy inequalities (see e.g Protter 2005, Theorem IV.48) imply that  $\mathbb{E}(\sup_{t \geq 0} |M_t|) < \infty$ . An application of the dominated convergence theorem then gives  $M \in \mathcal{M}$ . This establishes that  $\mathcal{H}^1 \subseteq \mathcal{M}$ .

*Date:* March 16, 2022.

*2010 Mathematics Subject Classification.* Primary: 60G44.

<sup>1</sup>This result is the probability-theoretic analogue of the classical  $H^1$ -BMO duality, due to Fefferman and Stein (1972).

However, the reverse inclusion does not hold. We demonstrate this fact below, by providing a recipe for constructing non-negative uniformly integrable martingales in  $\mathcal{M} \setminus \mathcal{H}^1$ .

## 2. CONSTRUCTION OF PROCESSES IN $\mathcal{M} \setminus \mathcal{H}^1$

Given a local martingale  $M \in \mathcal{M}_{\text{loc}}$ , observe that

$$\mathbb{E}\left(\sup_{t \geq 0} |M_t|\right) = \int_0^\infty \mathbb{P}\left(\sup_{t \geq 0} |M_t| > u\right) du$$

and

$$\sum_{n=1}^\infty \mathbb{P}\left(\sup_{t \geq 0} |M_t| > n\right) \leq \int_0^\infty \mathbb{P}\left(\sup_{t \geq 0} |M_t| > u\right) du \leq 1 + \sum_{n=1}^\infty \mathbb{P}\left(\sup_{t \geq 0} |M_t| > n\right).$$

Since  $M \in \mathcal{H}^1$  if and only if  $\mathbb{E}(\sup_{t \geq 0} |M_t|) < \infty$ , by virtue of the Burkholder-Davis-Gundy inequalities, it follows that  $M \in \mathcal{H}^1$  if and only if

$$\sum_{n=1}^\infty \mathbb{P}\left(\sup_{t \geq 0} |M_t| > n\right) < \infty. \quad (2.1)$$

This condition plays a key role in our construction.

Fix a non-negative local martingale  $M \in \mathcal{M}_{\text{loc}} \setminus \mathcal{M}$  that is not a uniformly integrable martingale, and define the non-decreasing sequence  $(c_n)_{n \in \mathbb{N}} \subset (1, \infty)$ , by setting

$$c_n := \ln\left(e + \sum_{k=1}^n \mathbb{P}\left(\sup_{t \geq 0} M_t > k\right)\right), \quad (2.2)$$

for each  $n \in \mathbb{N}$ . Since  $M \notin \mathcal{H}^1$ , it follows that  $\lim_{n \uparrow \infty} c_n = \infty$ . Next, suppose that  $(\Omega, \mathcal{F}, \mathbb{P})$  accommodates a discrete  $\mathcal{F}_0$ -measurable random variable  $Y \in \mathbb{N}$  that is independent of  $M$ , and whose distribution satisfies  $\mathbb{P}(Y > n) = 1/c_n$ , for each  $n \in \mathbb{N}$ , and let

$$\sigma := \inf\{t \geq 0 \mid M_t > Y\} \quad (2.3)$$

denote the first time  $M$  exceeds  $Y$ . It follows that

$$\mathbb{P}\left(\sup_{t \geq 0} M_t^\sigma > n\right) \geq \mathbb{P}\left(\sup_{t \geq 0} M_t > n\right) \mathbb{P}(Y > n) = \frac{1}{c_n} \mathbb{P}\left(\sup_{t \geq 0} M_t > n\right),$$

for each  $n \in \mathbb{N}$ . Consequently,

$$\sum_{n=1}^\infty \mathbb{P}\left(\sup_{t \geq 0} M_t^\sigma > n\right) \geq \lim_{m \uparrow \infty} \frac{1}{c_m} \sum_{n=1}^m \mathbb{P}\left(\sup_{t \geq 0} M_t > n\right) = \lim_{m \uparrow \infty} \frac{e^{c_m} - e}{c_m} = \infty,$$

since  $(c_n)_{n \in \mathbb{N}}$  is non-decreasing and  $\lim_{n \uparrow \infty} c_n = \infty$ . This implies that  $M^\sigma \notin \mathcal{H}^1$ . On the other hand, the almost sure limit  $M_\infty^\sigma := M_{\infty-}^\sigma \in \mathbb{R}_+$  exists, since  $M^\sigma$  is a non-negative local martingale, and hence also a non-negative supermartingale. Moreover,

$$\begin{aligned} \mathbb{E}(M_\infty^\sigma) &= \sum_{n=1}^\infty \mathbb{E}(M_\infty^\sigma \mid Y = n) \mathbb{P}(Y = n) = \sum_{n=1}^\infty \mathbb{E}(M_\infty^{\tau_n} \mid Y = n) \mathbb{P}(Y = n) \\ &= \sum_{n=1}^\infty \mathbb{E}(M_\infty^{\tau_n}) \mathbb{P}(Y = n) = \sum_{n=1}^\infty \mathbb{E}(M_0) \mathbb{P}(Y = n) = \mathbb{E}(M_0^\sigma), \end{aligned}$$

where

$$\tau_n := \inf\{t \geq 0 \mid M_t > n\}$$

for each  $n \in \mathbb{N}$ . Here, the penultimate equality follows from the fact that  $M^{\tau_n} \in \mathcal{M}$ , for each  $n \in \mathbb{N}$ . Consequently,  $M^\sigma \in \mathcal{M}$ .

**Example 2.1.** Consider a non-negative local martingale  $M \in \mathcal{M}_{\text{loc}}$  that belongs to Class  $(\mathcal{C}_0)$ , according to the terminology of Nikeghbali and Yor (2006), and suppose that  $M_0 = 1$ . In that case,  $M$  is a strictly positive local martingale without any positive jumps, for which  $M_\infty := M_{\infty-} = 0$ . The construction above is then applicable, since  $E(M_\infty) = 0 < 1 = E(M_0)$  implies that  $M \in \mathcal{M}_{\text{loc}} \setminus \mathcal{M}$ . Moreover, an application of Doob's maximal identity (see Nikeghbali and Yor 2006, Lemma 2.1) provides the following concrete representation for the non-decreasing sequence  $(c_n)_{n \in \mathbb{N}}$ , defined by (2.2):

$$c_n = \ln \left( e + \sum_{k=1}^n \frac{1}{k} \right),$$

for each  $n \in \mathbb{N}$ . It is then straightforward to see that  $\lim_{n \uparrow \infty} c_n = \infty$ , which is the crucial ingredient for showing that  $M^\sigma \notin \mathcal{H}^1$ , where the stopping time  $\sigma$  is given by (2.3).

#### REFERENCES

- Fefferman, C. and E. M. Stein (1972).  $H^p$  spaces of several variables. *Acta Math.* 129(3–4), 137–193.
- Nikeghbali, A. and M. Yor (2006). Doob's maximal identity, multiplicative decompositions and enlargements of filtrations. *Illinois J. Math.* 50(4), 791–814.
- Protter, P. E. (2005). *Stochastic Integration and Differential Equations* (Second ed.). Berlin: Springer.

HARDY HULLEY, FINANCE DEPARTMENT, UNIVERSITY OF TECHNOLOGY SYDNEY, P.O. BOX 123, BROADWAY, NSW 2007, AUSTRALIA

*E-mail address:* hardy.hulley@uts.edu.au

JOHANNES RUF, DEPARTMENT OF MATHEMATICS, LONDON SCHOOL OF ECONOMICS AND POLITICAL SCIENCE, HOUGHTON STREET, LONDON WC2A 2AE, UNITED KINGDOM

*E-mail address:* j.ruf@lse.ac.uk