

GENERALIZED FELLER PROCESSES AND MARKOVIAN LIFTS OF STOCHASTIC VOLTERRA PROCESSES: THE AFFINE CASE

CHRISTA CUCHIERO AND JOSEF TEICHMANN

ABSTRACT. We consider stochastic (partial) differential equations appearing as Markovian lifts of affine Volterra processes with jumps from the point of view of the generalized Feller property which was introduced in e.g. [13]. In particular we provide new existence, uniqueness and approximation results for Markovian lifts of affine rough volatility models of general jump diffusion type. We demonstrate that in this Markovian light the theory of stochastic Volterra processes becomes almost classical.

1. INTRODUCTION

In the realm of the recent discovery of rough volatility models (see e.g., [4, 21, 15, 16]), stochastic Volterra processes have been studied extensively, for instance in [24, 3, 17] and the references therein. It is well known that Markovian lifts of such processes exist, but it has rarely been detailed how these lifts look like in general, with the notable exception of [25] and the recent article [1]. If at all, these lifts have been constructed by solving first the stochastic Volterra equations and identifying a state space a posteriori, on which respective Markov properties show up. Neither maximum principles, nor strongly continuous semigroups, nor Feller properties, nor approximation properties, nor Kolmogorov backward equations have been considered in this context so far. The reason is of course that one enters the world of SPDEs where each of the previously mentioned concepts needs particular care and is often cumbersome due to the lack of local compactness of the underlying state spaces. Also the question whether all those considerations are restricted to Brownian driven processes has not yet been investigated due to technical difficulties in writing down (probabilistically) weak or martingale solution concepts with jump drivers that are not standard Poisson random measures.

With this article we aim to bridge this gap and introduce a functional analytic setting which clearly shows that it is an advantage to consider Markovian lifts right from the beginning without solving the Volterra equation in the first place. As a particular important example we deal with Volterra processes whose kernels are Laplace transforms of (signed) measures (see Section 5.1). Their Markovian lifts can be represented in terms of signed measure valued processes which can be treated in a systematic way comparable to Feller processes on locally compact state spaces. In particular, we demonstrate that many results in the realm of stochastic Volterra processes appear transparent and clearly structured in this framework. Indeed,

2010 *Mathematics Subject Classification.* 60H15, 60J25.

Key words and phrases. variation of constants formula, stochastic partial differential equations, affine processes, stochastic Volterra processes, rough volatility models.

The authors are grateful for the support of the ETH Foundation. Christa Cuchiero thanks the Forschungsinstitut für Mathematik for its support of a research stay at ETH Zürich in fall 2016. She additionally gratefully acknowledges financial support by the Vienna Science and Technology Fund (WWTF) under grant MA16-021.

Both authors are very grateful to Sergio Pulido for helping us to improve Subsection 5.2 and to an anonymous referee whose comments helped to improve the article along several lines.

thanks to the generalized Feller theory introduced in [13] many properties and theorems can be formulated abstractly while in the literature so far often rather concrete and example-like specifications have been considered. The passage to these *generalized* Feller processes is necessary, since no *standard* Feller formulation is available due to the lack of local compactness, as demonstrated in Example 5.1. This is in contrast to superprocesses, see, e.g. [12], which take values in spaces of nonnegative measures which are locally compact when the underlying space is compact.

We do also believe that the presented theory of generalized Feller semigroups and processes is a natural framework for analyzing SPDEs valid also far beyond Markovian lifts of stochastic Volterra processes: it opens the door to treat large deviations, long term behavior, numerical procedures, or asymptotic results in a general and simple framework.

This first article takes as guiding example affine Volterra processes on $\mathbb{R}_+$ of the form

$$(1.1) \quad V_t = h(t) + \int_0^t K(t-s) dX_s,$$

where h is a deterministic function, K a deterministic kernel and X a semimartingale whose characteristics depend linearly on the state of the Volterra process. In a subsequent article the general (non-affine) case, including properly formulated martingale problems and strong existence results, as well as other versions of Markovian lifts will be considered. Here we focus on Markovian lifts where the kernel K can be represented as

$$(1.2) \quad K(t) = \langle g, \mathcal{S}_t^* \nu \rangle$$

with $(\mathcal{S}_t^*)_{t \geq 0}$ a strongly continuous semigroup acting on a Banach space Y^* , $\nu \in Y^*$ (or in a slightly bigger space), $g \in Y$ with Y the pre-dual of Y^* and pairing $\langle \cdot, \cdot \rangle$. This abstract setting includes the above mentioned signed measure valued lifts as well as what we call “forward process lifts”, where a variant of the latter has also been considered in [1] in the special case of the Volterra Heston model. This richness already indicates that our almost axiomatic approach does not only simplify arguments but also unifies several concepts and branches of the literature. In particular, it provides due to its simple structure new existence and uniqueness results for (1.1), and leads to so far unknown approximation schemes of any order. For instance it will be easy to construct higher order weak approximation schemes like Ninomiya-Victoir schemes with precise and optimal convergence rates.

Inspired from Hawkes processes, see e.g. [23, 7], and in contrast to some recent literature, e.g. [25, 3, 2, 1], we do not consider the Brownian motion driven case as the simplest one but rather the case when the stochastic driver is a finite activity jump process (as it is often true and useful in the theory of stochastic processes): Brownian driven stochastic Volterra equations or stochastic Volterra equations with more complicated jump structures are then easy and well described limits (in the sense of generalized Feller processes) of processes with finitely many jumps on compact time intervals, from which then weak solutions can be constructed. Another difference to most of the literature is that we do not transform the processes in question into semimartingales to use stochastic calculus but we rather work directly with their Markovian structure. We also want to point out that with representation (1.2) we go well beyond standard assumptions on Volterra kernels like complete monotonicity, in particular we do not need resolvents of the first kind which do not always exist.

Let us outline in the sequel one of our lines of ideas in the above addressed case of kernels that are Laplace transforms of signed measures: we consider the

vector space of finite, signed Borel measures $Y^* := \mathcal{M}(\mathbb{R}_+ \cup \{\infty\})$ with its weak-* topology, see [27] or any other text book on functional analysis, and look at the following simple linear homogenous equation thereon

$$d\lambda_t(dx) = -x\lambda_t(dx)dt - w\nu(dx)\lambda_t(\mathbb{R}_+ \cup \{\infty\})dt,$$

where the initial value λ_0 is a signed measure, $w \geq 0$ a constant and ν a signed measure such that its Laplace transform

$$\int_0^\infty \exp(-tx)\nu(dx) \geq 0$$

is finite and non-negative for all $t > 0$, and belongs to $L^2_{\text{loc}}(\mathbb{R}_+, \mathbb{R}_+)$. The representation of the kernel in (1.2) is here

$$(1.3) \quad K(t) = \langle g, \mathcal{S}_t^* \nu \rangle = \langle 1, \exp(-t \cdot) \nu \rangle = \int_0^\infty \exp(-tx)\nu(dx)$$

i.e. the evaluation of the element $\exp(-t \cdot) \nu$ in the dual space $\mathcal{M}(\mathbb{R}_+ \cup \{\infty\})$ of $Y = C_b(\mathbb{R}_+ \cup \{\infty\}, \mathbb{R})$ on the constant function 1. Under these circumstances one can easily prove two facts: first the solution defines a generalized Feller process (see [13] and the subsequent sections for details on this notion) on the state space of all signed measures, and, second, there is a closed invariant subspace $\mathcal{E}$ consisting of initial measures λ_0 where the total mass $\langle 1, \lambda_t \rangle = \lambda_t(\mathbb{R}_+ \cup \{\infty\})$ remains non-negative along its evolution in time for all $w \geq 0$. Of course the total mass satisfies a one-dimensional linear Volterra equation, whence the set $\mathcal{E}$ of measures λ_0 can be characterized by resolvents, see Section 4 and Section 5.1 for all details. Next we construct on this very state space $\mathcal{E}$ all sorts of jump diffusion processes with general characteristics by Poisson approximations applying an approximation theorem for generalized Feller semigroups (see Theorem 3.2). This is parallel to classical work on affine processes, see [14, 8, 9], where general existence is obtained by pure jump approximations. We can finally show that the SPDE

$$(1.4) \quad d\lambda_t(dx) = -x\lambda_t(dx)dt + \nu(dx)dX_t$$

for all semimartingales X whose characteristics depend linearly on $\lambda_t(\mathbb{R}_+ \cup \{\infty\})$ admits an analytically mild and probabilistically weak solution. Moreover, the associated Cauchy problem has a solution given in terms of a generalized Feller semigroup acting on a well specified set of functions. In particular, by the variation of constants formula and (1.3) it is easily seen that the total mass $\langle 1, \lambda_t \rangle = \lambda_t(\mathbb{R}_+ \cup \{\infty\})$ solves (1.1) with $h(t) = \int_0^\infty \exp(-xt)\lambda_0(dx)$.

We also want to point out that we obtain generic path properties, i.e. in this general setting we shall always have càg versions but not necessarily càdlàg versions. This is in contrast to Feller processes where càdlàg versions do always exist. This is due to the fact that generalized Feller processes are not always quasi-contractive.

This previous example highlights our results which can be summarized as follows:

- we provide a full solution theory for univariate stochastic Volterra equations (1.1) driven by jump-diffusions. In particular for a large class of functionals we can provide solutions of the respective Kolmogorov equations.
- we provide approximation schemes, (generalized) maximum principles (in certain cases), (probabilistically) weak solutions concepts for Markovian lifts of jump-diffusion driven stochastic Volterra equations (1.1).
- we provide a solution theory of the corresponding generalized Riccati differential equations on the pre-dual space $\mathcal{E}_*$ and the respective affine transform formulas.

- we can go beyond standard assumptions on kernels of stochastic Volterra equations such as complete monotonicity.
- we provide an abstract theory of Markovian lifts which is not restricted to the above introduced univariate case. Indeed, by considering as Y^* the space of $\mathbb{R}^d$ -valued measures on E for some locally compact space E , (1.4) can be generalized to the following multivariate case

$$d\lambda_t^l(dx) = -f^l(x)\lambda_t^l(dx)dt + \sum_{j=1}^d \nu_j^l(dx)dX_t^j,$$

for $l \in \{1, \dots, d\}$ in a straightforward way. This is due to the completely general character of Proposition 3.3 below. Here, f^l are functions from $E \rightarrow \mathbb{R}_+$ satisfying appropriate conditions, and X^j are semimartingales whose characteristics depend linearly on λ . Note that this allows for Volterra process specifications of the form

$$V_t^l = h^l(t) + \int_0^t \sum_{j=1}^d K_j^l(t-s)dX_s^j,$$

for kernels K_j^l given by $K_j^l(t) = \int_E g^l(x)e^{-tf^l(x)}\nu_j^l(dx)$ for some bounded continuous $g^l \in C_b(E; \mathbb{R})$ and $h^l(t) = \int_E g^l(x)e^{-tf^l(x)}\lambda_0^l(dx)$.

- Heston-like or Barndorff-Nielsen-Shephard-like structures can be analogously constructed again by choosing appropriate Banach spaces Y . For instance the rough Heston case [15] can be treated with $Y = C_b(\mathbb{R}_+ \cup \{\infty\}; \mathbb{R}^2)$ and the equations

$$\begin{aligned} d\lambda_t^1(dx) &= -x\lambda_t^1(dx)dt + \nu(dx)\sqrt{\langle 1, \lambda_t^1 \rangle}dB_t^1, \\ d\lambda_t^2(dx) &= -x\lambda_t^2(dx)dt + \delta_0(dx)\sqrt{\langle 1, \lambda_t^1 \rangle}dB_t^2, \end{aligned}$$

where B^1 and B^2 are correlated Brownian motions with $d\langle B^1, B^2 \rangle_t = \varrho dt$. Note that the variance process is given by $\lambda_t^1(\mathbb{R}_+ \cup \{\infty\})$ and the log price process by $\lambda_t^2(\mathbb{R}_+ \cup \{\infty\})$.

- Covariance matrix (or cone-) valued processes can be constructed by considering positive semidefinite symmetric matrix (or cone-) valued measures as respective state space. Here particular geometric restrictions appear in the diffusion driven case. Markovian lifts of Volterra type processes on this more involved state space, relevant for rough covariance modeling, are treated in [10].

The remainder of the article is as follows: in Section 1.1 we introduce some notation and review certain functional analytic concepts. In Section 2, we deal with generalized Feller processes as introduced in [13]. In Section 3 we show simple approximation theorems for generalized Feller semigroups beyond standard Trotter-Kato type theorems, and we prove a crucial existence result for pure jump processes with finite but unbounded intensity. In Sections 4, 5.1 and 5.2 we apply the presented theory to SPDEs which are lifts of affine Volterra processes.

1.1. Notation and some functional analytic notions. For the background in functional analysis we refer to the excellent textbook [27] as main reference and to the equally excellent books [18, 26] for the background in strongly continuous semigroups. We emphasize however that only very basic knowledge in functional analysis and strongly continuous semigroups is required.

We shall often apply the following notations: let Y be a Banach space and Y^* its dual space, i.e. the space of linear continuous functionals with the strong dual

norm

$$\|\lambda\|_{Y^*} = \sup_{\|y\| \leq 1} |\langle y, \lambda \rangle|,$$

where $\langle y, \lambda \rangle := \lambda(y)$ denotes the evaluation of the linear functional λ at the point $y \in Y$. Since cones $\mathcal{E}$ of Y^* will be our statespaces, we denote the polar cones in pre-dual notation, i.e.

$$\mathcal{E}_* = \{y \in Y \mid \langle y, \lambda \rangle \leq 0 \text{ for all } \lambda \in \mathcal{E}\}.$$

We denote spaces of bounded linear operators from Banach spaces Y_1 to Y_2 by $L(Y_1, Y_2)$ with norm

$$\|A\|_{L(Y_1, Y_2)} := \sup_{\|y_1\|_{Y_1} \leq 1} \|Ay_1\|_{Y_2}.$$

If $Y_1 = Y_2$ we only write $\|\cdot\|_{L(Y_1)}$. On Y^* we shall usually consider beside the strong topology (induced by the strong dual norm) the weak*-topology, which is the weakest locally convex topology making *all* linear functionals $\langle y, \cdot \rangle$ on Y^* continuous. Let us recall the following facts:

- The weak*-topology is metrizable if and only if Y is finite dimensional: this is due to Baire's category theorem since Y^* can be written as a countable union of closed sets, whence at least one has to contain an open set, which in turn means that compact neighborhoods exist, i.e. a strictly finite dimensional phenomenon.
- Norm balls K_R of any radius R in Y^* are compact with respect to the weak*-topology, which is the Banach-Alaoglu theorem.
- These balls are metrizable if and only if Y is separable: this is true since Y can be isometrically embedded into $C(K_1)$, where $y \mapsto \langle y, \cdot \rangle$, for $y \in Y$. Since Y is separable, its embedded image is separable, too, which means – by looking at the algebra generated by Y in $C(K_1)$ – that $C(K_1)$ is separable, which is the case if and only if K_1 is metrizable.

Since we do not need metrizability of K_R , we do *not* assume that our Banach spaces Y are separable.

A family of linear operators $(P_t)_{t \geq 0}$ on a Banach space Y with $P_t P_s = P_{t+s}$ for $s, t \geq 0$ and with $P_0 = I$ where I denotes the identity is called strongly continuous semigroup if $\lim_{t \rightarrow 0} P_t y = y$ holds true for every $y \in Y$. We denote its generator usually by A which is defined as $\lim_{t \rightarrow 0} \frac{P_t y - y}{t}$ for all $y \in \text{dom}(A)$, i.e. the set of elements where the limit exists. Notice that $\text{dom}(A)$ is left invariant by the semigroup P and that its restriction on the domain equipped with the operator norm

$$\|y\|_{\text{dom}(A)} := \sqrt{\|y\|^2 + \|Ay\|^2}$$

is again a strongly continuous semigroup.

Moreover, as already used in the introduction, $\mathcal{M}(E)$ denotes the space of signed finite measures on E equipped with the Borel σ -algebra and $\mathcal{M}_+(E)$ the space of finite nonnegative measures.

2. GENERALIZED FELLER SEMIGROUPS AND PROCESSES

Feller semigroups have proved to be useful in the context of locally compact state spaces. Generalized Feller semigroups serve the same purpose on state spaces which are not locally compact, which is a typical infinite dimensional phenomenon. Local compactness is replaced by adjoining a *proper* weight function to a state space X , which measures explosion. Interestingly even in the locally compact case generalized Feller processes yield new results relevant for the theory of affine processes, see e.g. [14, 8, 9], since unbounded test functions can be considered. We refer to [13] for all necessary details and proofs, and also to the references therein. Notice that even

in the infinite dimensional setting there are notable examples of local compactness, see Remark 2.2, but we point out that our cases are not locally compact in their natural topology, see Example 5.1.

2.1. Defintions and results. First we introduce weighted spaces and state a central Riesz representation result. The underlying space X here is a completely regular Hausdorff topological space.

Definition 2.1. A function $\varrho: X \rightarrow (0, \infty)$ is called admissible weight function if the sets $K_R := \{x \in X: \varrho(x) \leq R\}$ are compact for all $R > 0$.

An admissible weight function ϱ is necessarily lower semicontinuous and bounded from below by a positive constant. We call the pair X together with an admissible weight function ϱ a *weighted space*. A weighted space is σ -compact, i.e. a countable union of compacts. In the following remark we clarify the question of local compactness of convex subsets $\mathcal{E} \subset X$ when X is a locally convex topological space and ϱ convex.

Remark 2.2. Let X be a locally convex topological space and $\mathcal{E}$ a convex subset. Moreover, let ϱ be a *convex* admissible weight function. Then ϱ is continuous on $\mathcal{E}$ if and only if $\mathcal{E}$ is locally compact. Indeed if ϱ is continuous on $\mathcal{E}$, then of course the topology on $\mathcal{E}$ is locally compact since every point has a compact neighborhood of type $\{\varrho \leq R\}$ for some $R > 0$. On the other hand if the topology on $\mathcal{E}$ is locally compact, then for every point $\lambda_0 \in \mathcal{E}$ there is a convex, compact neighborhood $V \subset \mathcal{E}$ such that $\varrho(\lambda) - \varrho(\lambda_0)$ is bounded on V by a number $k > 0$, whence by convexity $|\varrho(s(\lambda - \lambda_0) + \lambda_0) - \varrho(\lambda_0)| \leq sk$ for $\lambda - \lambda_0 \in s(V - \lambda_0)$ and $s \in]0, 1]$. This in turn means that ϱ is continuous at λ_0 .

From now on ϱ shall always denote an admissible weight function. For completeness we start by putting definitions for general Banach space valued functions, although in the sequel we shall only deal with $\mathbb{R}$ -valued functions: let Z be a Banach space with norm $\|\cdot\|_Z$. The vector space

$$(2.1) \quad B^{\varrho}(X; Z) := \left\{ f: X \rightarrow Z: \sup_{x \in X} \varrho(x)^{-1} \|f(x)\|_Z < \infty \right\}$$

of Z -valued functions f equipped with the norm

$$(2.2) \quad \|f\|_{\varrho} := \sup_{x \in X} \varrho(x)^{-1} \|f(x)\|_Z,$$

is a Banach space itself. It is also clear that for Z -valued bounded continuous functions the continuous embedding $C_b(X; Z) \subset B^{\varrho}(X; Z)$ holds true, where we consider the supremum norm on bounded continuous functions, i.e. $\sup_{x \in X} \|f(x)\|$.

Definition 2.3. We define $\mathcal{B}^{\varrho}(X; Z)$ as the closure of $C_b(X; Z)$ in $B^{\varrho}(X; Z)$. The normed space $\mathcal{B}^{\varrho}(X; Z)$ is a Banach space.

If the range space $Z = \mathbb{R}$, which from now on will be the case, we shall write $\mathcal{B}^{\varrho}(X)$ for $\mathcal{B}^{\varrho}(X; \mathbb{R})$ and analogously $B^{\varrho}(X)$.

We consider elements of $\mathcal{B}^{\varrho}(X)$ as continuous functions whose growth is controlled by ϱ . More precisely we have by [13, Theorem 2.7] that $f \in \mathcal{B}^{\varrho}(X)$ if and only if $f|_{K_R} \in C(K_R)$ for all $R > 0$ and

$$(2.3) \quad \lim_{R \rightarrow \infty} \sup_{x \in X \setminus K_R} \varrho(x)^{-1} \|f(x)\| = 0.$$

Additionally, by [13, Theorem 2.8] it holds that for every $f \in \mathcal{B}^{\varrho}(X)$ with $\sup_{x \in X} f(x) > 0$, there exists $z \in X$ such that

$$(2.4) \quad \varrho(x)^{-1} f(x) \leq \varrho(z)^{-1} f(z) \quad \text{for all } x \in X,$$

which emphasizes the analogy with spaces of continuous functions vanishing at ∞ on locally compact spaces.

Let us now state the following crucial representation theorem of Riesz type:

Theorem 2.4 (Riesz representation for $\mathcal{B}^e(X)$). *For every continuous linear functional $\ell: \mathcal{B}^e(X) \rightarrow \mathbb{R}$ there exists a finite signed Radon measure μ on X such that*

$$(2.5) \quad \ell(f) = \int_X f(y) \mu(dy) \quad \text{for all } f \in \mathcal{B}^e(X).$$

Additionally

$$(2.6) \quad \int_X \varrho(y) |\mu|(dy) = \|\ell\|_{L(\mathcal{B}^e(X), \mathbb{R})},$$

where $|\mu|$ denotes the total variation measure of μ .

We shall next consider positive, strongly continuous semigroups on $\mathcal{B}^e(X)$ and recover very similar structures as for standard Feller semigroups on the space of continuous functions vanishing at ∞ on locally compact spaces.

Definition 2.5. *A family of bounded linear operators $P_t: \mathcal{B}^e(X) \rightarrow \mathcal{B}^e(X)$ for $t \geq 0$ is called generalized Feller semigroup if*

- (i) $P_0 = I$, the identity on $\mathcal{B}^e(X)$,
- (ii) $P_{t+s} = P_t P_s$ for all $t, s \geq 0$,
- (iii) for all $f \in \mathcal{B}^e(X)$ and $x \in X$, $\lim_{t \rightarrow 0} P_t f(x) = f(x)$,
- (iv) there exist a constant $C \in \mathbb{R}$ and $\varepsilon > 0$ such that for all $t \in [0, \varepsilon]$, $\|P_t\|_{L(\mathcal{B}^e(X))} \leq C$.
- (v) P_t is positive for all $t \geq 0$, that is, for $f \in \mathcal{B}^e(X)$, $f \geq 0$, we have $P_t f \geq 0$.

We obtain due to the Riesz representation property the following key theorem:

Theorem 2.6. *Let $(P_t)_{t \geq 0}$ satisfy (i) to (iv) of Definition 2.5. Then, $(P_t)_{t \geq 0}$ is strongly continuous on $\mathcal{B}^e(X)$, that is,*

$$(2.7) \quad \lim_{t \rightarrow 0} \|P_t f - f\|_e = 0 \quad \text{for all } f \in \mathcal{B}^e(X).$$

One can also establish a positive maximum principle in case that the semigroup $(P_t)_{t \geq 0}$ grows around 0 like $\exp(\omega t)$ for some $\omega \in \mathbb{R}$ with respect to the operator norm on $\mathcal{B}^e(X)$. Indeed, the following theorem proved in [13, Theorem 3.3] is a reformulation of the Lumer-Philips theorem for quasi-contractive semigroups using a *generalized positive maximum principle* which is formulated in the sequel.

Theorem 2.7. *Let A be an operator on $\mathcal{B}^e(X)$ with domain D , and $\omega \in \mathbb{R}$. Then, A is closable and its closure $\overline{A}$ generates a generalized Feller semigroup $(P_t)_{t \geq 0}$ with $\|P_t\|_{L(\mathcal{B}^e(X))} \leq \exp(\omega t)$ for all $t \geq 0$ if and only if*

- (i) D is dense,
- (ii) $A - \omega_0$ has dense image for some $\omega_0 > \omega$, and
- (iii) A satisfies the generalized positive maximum principle, that is, for $f \in D$ with $(\varrho^{-1} f) \vee 0 \leq \varrho(z)^{-1} f(z)$ for some $z \in X$, $A f(z) \leq \omega f(z)$.

Remark 2.8. Let $(P_t)_{t \geq 0}$ be a generalized Feller semigroup on $\mathcal{B}^e(X)$, then by abstract semigroup theory, see [18, Proposition I.5.5], there exists $M \geq 1$ and $\omega \in \mathbb{R}$ such that $\|P_t\|_{L(\mathcal{B}^e(X))} \leq M \exp(\omega t)$. By the Riesz representation as stated in Theorem 2.4, $P_t \varrho(x)$ is always defined as integral of the measurable function ϱ with respect to the measure representing the linear functional $f \mapsto P_t f(x)$. The value $P_t \varrho(x)$ corresponds precisely to the operator norm of this linear functional by Equation (2.6) (note that the measure in Theorem 2.4 depends additionally on x). Hence by

$$|P_t f(x)| \leq M \exp(\omega t) \varrho(x) \|f\|_e$$

it follows that $P_t \varrho \leq M \exp(\omega t) \varrho$.

Remark 2.9. Let $(P_t)_{t \geq 0}$ be a generalized Feller semigroup on $\mathcal{B}^e(X)$ of transport type, i.e.

$$(2.8) \quad P_t f(x) = f(\psi_t(x))$$

for some continuous map $\psi_t : X \rightarrow X$. Define now a new function

$$\tilde{\varrho}(x) := \sup_{t \geq 0} \exp(-\omega t) P_t \varrho(x)$$

for $x \in X$. Notice that $\tilde{\varrho}$ is an admissible weight function, since

$$\{\tilde{\varrho} \leq R\} = \cap_{t \geq 0} \{P_t \varrho \leq \exp(\omega t) R\}$$

is compact by the continuity of $x \mapsto \psi_t(x)$ as an intersection of closed subsets of compacts. Additionally we have that

$$\varrho \leq \tilde{\varrho} \leq M \varrho$$

by the growth bound as in Remark 2.8 and therefore the norm on $\mathcal{B}^e(X)$ is equivalent to

$$\|f\|_{\tilde{\varrho}} = \sup_{x \in X} \frac{|f(x)|}{\tilde{\varrho}(x)}.$$

Furthermore,

$$\|P_t f\|_{\tilde{\varrho}} \leq \exp(\omega t) \|f\|_{\tilde{\varrho}}$$

holds for all $t \geq 0$ and $f \in \mathcal{B}^e(X)$. Indeed, this is a consequence of the following estimate

$$(2.9) \quad \begin{aligned} \|P_t f\|_{\tilde{\varrho}} &= \sup_x \left| \frac{f(\psi_t(x))}{\sup_s \exp(-\omega s) \varrho(\psi_s(x))} \right| \leq \sup_x \left| \frac{f(\psi_t(x))}{\sup_s \exp(-\omega(t+s)) \varrho(\psi_{t+s}(x))} \right| \\ &\leq \exp(\omega t) \sup_x \left| \frac{f(\psi_t(x))}{\sup_s \exp(-\omega s) \varrho(\psi_s(\psi_t(x)))} \right| \leq \exp(\omega t) \|f\|_{\tilde{\varrho}}. \end{aligned}$$

Hence,

$$|P_t f(x)| \leq \exp(\omega t) \tilde{\varrho}(x) \|f\|_{\tilde{\varrho}},$$

which implies

$$P_t \tilde{\varrho} \leq \exp(\omega t) \tilde{\varrho}, \quad t \geq 0.$$

The next theorem is a basic perturbation result in the spirit of [19, Corollary 7.2] for generalized Feller semigroups.

Theorem 2.10. *Let A be a generator of a generalized Feller semigroup $(P_t)_{t \geq 0}$ on $\mathcal{B}^e(X)$ with $\|P_t\|_{L(\mathcal{B}^e(X))} \leq M \exp(\omega t)$ for all $t \geq 0$ and some $M \geq 1$, $\omega \in \mathbb{R}$. Let B be a bounded generator of a generalized Feller semigroup on $\mathcal{B}^e(X)$. Then $A + B$ is a generator of a generalized Feller semigroup on $\mathcal{B}^e(X)$ whose operator norm is bounded by $M \exp(\omega t + M \|B\|_{L(\mathcal{B}^e(X))} t)$ for $t \geq 0$. If $M = 1$ and B satisfies*

$$B \varrho \leq \tilde{\omega} \varrho$$

for some real $\tilde{\omega}$, then the operator norm of the semigroup generated by $A + B$ is bounded by $\exp((\omega + \tilde{\omega})t)$ for $t \geq 0$.

Proof. The result is a consequence of [18, Theorem 1.3, p.158]. The positivity is obtained from Chernoff's product formula. If $B \varrho \leq \tilde{\omega} \varrho$, then for $f \geq 0$

$$\exp(Bt)f \leq \exp(\tilde{\omega}t) \|f\|_{\varrho} \varrho,$$

which yields $\|B\|_{L(\mathcal{B}^e(X))} \leq \tilde{\omega}$. □

We now give a version of the Kolmogorov extension theorem, which has not been shown so far. Notice that standard versions of the extension theorem, i.e. for Borel spaces or in case of inner-regular measures on topological spaces, see [5], do not apply in this case. One can, however, apply the Daniell-Kolmogorov theorem.

Theorem 2.11. *Let $(P_t)_{t \geq 0}$ be a generalized Feller semigroup with $P_t 1 = 1$ for $t \geq 0$. Then there exists a filtered measurable space $(\Omega, (\mathcal{F}_t)_{t \geq 0})$ with right continuous filtration, and an adapted family of random variables $(\lambda_t)_{t \geq 0}$ such that for any initial value $\lambda_0 \in X$ there exists a probability measure $\mathbb{P}^{\lambda_0}$ with*

$$\mathbb{E}_{\mathbb{P}^{\lambda_0}}[f(\lambda_t)] = P_t f(\lambda_0)$$

for $t \geq 0$ and every $f \in \mathcal{B}^e(X)$. The Markov property holds true, i.e.

$$\mathbb{E}_{\mathbb{P}^{\lambda_0}}[f(\lambda_t) \mid \mathcal{F}_s] = P_{t-s} f(\lambda_s)$$

almost surely with respect to $\mathbb{P}^{\lambda_0}$.

Remark 2.12. For simplicity of notation we shall usually only write $\mathbb{E}$ or $\mathbb{E}_{\lambda_0}$ for expectations with respect to the above probability measure $\mathbb{P}^{\lambda_0}$. We call the process $(\lambda_t)_{t \geq 0}$ a generalized Feller process with initial value λ_0 with respect to this measure.

Proof. Every weighted space is σ -compact and therefore the set of compact subsets of $\{\varrho \leq R\}$ for $R \geq 0$ form a compact class in X . Consider the kernels $\mu_t(\lambda_0, \cdot)$ representing $f \mapsto P_t f(\lambda_0)$ and fix $\lambda_0 \in X$. Then by standard constructions we obtain a consistent family of probability measures μ^J on X^J for all finite subsets $J \subset I$, i.e. for the canonical projections $\pi^{J_2 \rightarrow J_1} : X^{J_2} \rightarrow X^{J_1}$ the push forward of μ^{J_2} is μ^{J_1} , for all finite sets $J_1 \subset J_2 \subset I$. The Daniell-Kolmogorov theorem, see [5, Theorem 15.23], then yields a probability measure on the $X^{\mathbb{R}^+}$ on the σ algebra generated by the canonical projections, such that the canonical projections $\lambda_t : X \rightarrow \mathbb{R}$ have law $\mu_t(\lambda_0, \cdot)$. The rest follows by construction as in the Feller case. Right continuity of the filtration is enforced by taking the right continuous enlargement. $\square$

We end this subsection with a statement on path properties: up to explosion generalized Feller processes have – with respect to countably many test functions – càdlàg or càglàd trajectories.

Theorem 2.13. *Let $(P_t)_{t \geq 0}$ be a generalized Feller semigroup and let $(\lambda_t)_{t \geq 0}$ be a generalized Feller process on a filtered probability space. Then for every countable family $(f_n)_{n \geq 0}$ of functions in $\mathcal{B}^e(X)$ we can choose a version of the processes $\left(\frac{f_n(\lambda_t)}{\varrho(\lambda_t)}\right)_{t \geq 0}$, such that the trajectories are càglàd for all $n \geq 0$.*

If additionally $P_t \varrho \leq \exp(\omega t) \varrho$ holds true, then $(\exp(-\omega t) \varrho(\lambda_t))_{t \geq 0}$ is a supermartingale and can be chosen to have càglàd trajectories. In this case we obtain that the processes $(f_n(\lambda_t))_{t \geq 0}$ can be chosen to have càglàd trajectories.

Remark 2.14. (i) In the general case, when $P_t \varrho \leq M \exp(\omega t) \varrho$ for $M > 1$, we obtain for $(f_n(\lambda_t))_{t \geq 0}$ only càg trajectories. To see this, consider the measurable set of sample events $\{\sup_{0 \leq t \leq 1} \varrho(\lambda_t) \leq R\}$. On this set we can construct a càglàd version of the processes $\left(\frac{f_n(\lambda_t)}{\varrho(\lambda_t)}\right)_{t \leq 1}$ and $\left(\frac{1}{\varrho(\lambda_t)}\right)_{t \leq 1}$ and in turn also of $(f_n(\lambda_t))_{t \geq 0}$. The limit $R \rightarrow \infty$, however, only leads to a càg version since we cannot control the right limits.

(ii) Notice that we do not speak – in this general context – about path properties of $(\lambda_t)_{t \geq 0}$ itself. In case of separability of $\mathcal{B}^e(X)$ we can actually find a càg version of λ itself by choosing as point separating family f_n a countable dense subset of $\mathcal{B}^e(X)$.

Proof. For every generalized Feller semigroup the process

$$Y_t^\alpha := \exp(-\alpha t) R(\alpha) f(\lambda_t)$$

for $t \geq 0$ is actually a non-negative supermartingale for $f \geq 0$ in $\mathcal{B}^q(X)$. Here

$$R(\alpha) := \int_0^\infty \exp(-\alpha s) P_s ds$$

denotes the resolvent of $(P_t)_{t \geq 0}$, defined for large enough α , say $\alpha > \omega$ for some $\omega \geq 0$. Indeed by the Markov property

$$\begin{aligned} \mathbb{E}[Y_t^\alpha | \mathcal{F}_s] &= \mathbb{E}\left[\exp(-\alpha t) \int_0^\infty \exp(-\alpha u) P_u f(\lambda_t) du | \mathcal{F}_s\right] \\ &= \exp(-\alpha t) \int_0^\infty \exp(-\alpha u) P_{u+t-s} f(\lambda_s) du \\ &= \exp(-\alpha s) \int_{t-s}^\infty \exp(-\alpha u) P_u f(\lambda_s) du \\ &\leq Y_s^\alpha, \end{aligned}$$

for $0 \leq s \leq t$ with respect to the given right continuous filtration. Hence $(Y_t^\alpha)_{t \geq 0}$ is a supermartingale and $t \mapsto \mathbb{E}[Y_t^\alpha]$ is continuous. This implies together with the right continuity of the filtration the existence of a càglàd version. Consider now the countable set of functions

$$\mathcal{H} := \{\alpha R(\alpha) f_n | \text{ for } n \in \mathbb{N}, \alpha > \omega\}.$$

Since each $(Y_t^\alpha)_{t \geq 0}$ has a càglàd version, this translates also to each of the processes $(h(\lambda_t))_{t \geq 0}$ for $h \in \mathcal{H}$. Moreover, as $\|\alpha R(\alpha) f - f\|_\varrho \rightarrow 0$ for $\alpha \rightarrow \infty$, also $(\frac{f_n(\lambda_t)}{\varrho(\lambda_t)})_{t \geq 0}$ has càglàd trajectories. The second statement follows directly. $\square$

The following example shows that the càglàd property can fail when we have jump process with a singular kernel even though the general Feller property holds true.

Example 2.15. Let $X = \mathcal{M}(\mathbb{R}_+ \cup \{+\infty\})$ be equipped with the total variation norm as normed space, or with the weak- $*$ -topology. Take a standard Poisson process N with intensity 1 and a measure ν such that

$$K(t) := \int_0^\infty \exp(-xt) \nu(dx) < \infty$$

for $t > 0$, but with possibly infinite total mass. Consider now the (formal) SPDE

$$d\lambda_t(dx) = -x\lambda_t(dx)dt + \nu dN_t,$$

whose explicit solution is given by variation of constants

$$\lambda_t(dx) = \exp(-xt) \lambda_0(dx) + \int_0^t \exp(-x(t-s)) \nu(dx) dN_s$$

understood in the weak- $*$ -topology, for $t \geq 0$. Take $\varrho(\lambda) = 1 + \int_0^\infty \lambda(dx)$ the shifted total variation norm, which makes X a weighted space. Then

$$P_t f(\lambda_0) = \mathbb{E}[f(\lambda_t)]$$

actually defines a generalized Feller semigroup if, e.g., $\int_0^t K(s) ds \leq M \exp(\omega t)$. Indeed, we have

$$P_t \varrho(\lambda_0) = 1 + \int_0^\infty \exp(-xt) \lambda_0(dx) + \int_0^t K(t-s) ds \leq M \exp(\omega t) \varrho(\lambda_0)$$

for some (possibly new) constant $M \geq 1$ and some real ω . This together with point-wise continuity on cylindrical Fourier basis elements yields the generalized Feller property. The process

$$\begin{aligned}\varrho(\lambda_t) &= 1 + \int_0^\infty \exp(-xt) \lambda_0(dx) + \int_0^t K(t-s) dN_s \\ &= 1 + \int_0^\infty \exp(-xt) \lambda_0(dx) + \sum_{\tau_n < t} K(t - \tau_n),\end{aligned}$$

where τ_n denotes the jump times of the Poisson process, however does only allow for a càg version, if K is singular.

Remark 2.16. Theorem 2.13 allows to formulate a martingale problem in this general context if $P_t \varrho \leq \exp(\omega t) \varrho$. Indeed, let A be the generator of a generalized Feller semigroup. Then, for every $f \in \text{dom}(A)$ we can choose versions for $(f(\lambda_t))_{t \geq 0}$ and $(Af(\lambda_t))_{t \geq 0}$ which are càglàd. Hence $\int_0^t Af(\lambda_s) ds$ is in particular well-defined and the Markov property implies that

$$\left(f(\lambda_t) - \int_0^t Af(\lambda_s) ds \right)_{t \geq 0}$$

is actually a (càglàd) martingale. This will be investigated further in a subsequent article.

2.2. Dual spaces of Banach spaces. The most important playground for our theory will be closed subsets of duals of Banach spaces, where the weak*-topology appears to be σ -compact due to the Banach-Alaoglu theorem. Assume that $\mathcal{E} \subset Y^*$ is a closed subset of the dual space Y^* of some Banach space Y where Y^* is equipped with its weak*-topology. Consider a lower semicontinuous function $\varrho: \mathcal{E} \rightarrow (0, \infty)$ and denote by $(\mathcal{E}, \varrho)$ the corresponding weighted space. We have the following approximation result (see [13, Theorem 4.2]) for functions in $\mathcal{B}^{\varrho}(\mathcal{E})$ by cylindrical functions. Set

$$\begin{aligned}\mathcal{Z}_N &:= \{g(\langle \cdot, y_1 \rangle, \dots, \langle \cdot, y_N \rangle) : g \in C_b^\infty(\mathbb{R}^N) \\ &\text{and } y_j \in Y, j = 1, \dots, N\},\end{aligned}\tag{2.10}$$

where $\langle \cdot, \cdot \rangle$ denotes the pairing between Y^* and Y . We denote by $\mathcal{Z} := \bigcup_{N \in \mathbb{N}} \mathcal{Z}_N$ the set of bounded smooth continuous cylinder functions on $\mathcal{E}$. We can prove the following theorem beyond any separability assumptions on Y .

Theorem 2.17. *The closure of $\mathcal{Z}$ in $\mathcal{B}^{\varrho}(\mathcal{E})$ coincides with $\mathcal{B}^{\varrho}(\mathcal{E})$, whose elements appear to precisely the functions $f \in \mathcal{B}^{\varrho}(\mathcal{E})$ which satisfy (2.3) and that $f|_{K_R}$ is weak*-continuous for any $R > 0$.*

Proof. Since $\mathcal{Z}|_{\{\varrho \leq R\}}$ is a point separating algebra we can apply the Stone Weierstrass theorem on the compact sets $\{\varrho \leq R\}$ to obtain density of the restrictions in $C_b(\{\varrho \leq R\})$ for any $R \geq 0$. Then we can apply [13, Theorem 2.7]. $\square$

Remark 2.18. Of course we can consider also subsets of general cylindrical functions to serve the same purpose (we just need a Stone-Weierstrass theorem to be applicable), i.e. the subset should be point separating and an algebra. This will play an important role in the case of affine processes where we can consider the linear span of all products of Fourier basis elements $\exp(\langle \cdot, y \rangle)$ for $y \in Y$.

In the following we will give a theorem telling when the semigroup of a Markov process is actually a generalized Feller semigroup. For the dense subset appearing in this theorem we take in practice the set of cylindrical function $\mathcal{Z}$ introduced above. For its formulation we need the following assumptions.

Assumption 2.19. Let $(\lambda_t)_{t \geq 0}$ denote a time homogeneous Markov process on some stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P}^{\lambda_0})$ with values in $\mathcal{E}$.

Then we assume that

(i) there are constants C and $\varepsilon > 0$ such that

$$(2.11) \quad \mathbb{E}_{\lambda_0}[\varrho(\lambda_t)] \leq C\varrho(\lambda_0) \quad \text{for all } \lambda_0 \in \mathcal{E} \text{ and } t \in [0, \varepsilon];$$

(ii)

$$(2.12) \quad \lim_{t \rightarrow 0} \mathbb{E}_{\lambda_0}[f(\lambda_t)] = f(\lambda_0) \quad \text{for any } f \in \mathcal{B}^q(\mathcal{E}) \text{ and } \lambda_0 \in \mathcal{E};$$

(iii) for all f in a dense subset of $\mathcal{B}^q(\mathcal{E})$, the map $\lambda_0 \mapsto \mathbb{E}_{\lambda_0}[f(\lambda_t)]$ lies in $\mathcal{B}^q(\mathcal{E})$.

Remark 2.20. Of course inequality (2.11) implies that $|\mathbb{E}_{\lambda_0}[f(\lambda_t)]| \leq C\varrho(\lambda_0)$ for all $f \in \mathcal{B}^q(\mathcal{E})$, $\lambda_0 \in \mathcal{E}$ and $t \in [0, \varepsilon]$.

Theorem 2.21. Suppose Assumptions 2.19 hold true. Then $P_t f(\lambda_0) := \mathbb{E}_{\lambda_0}[f(\lambda_t)]$ satisfies the generalized Feller property and is therefore a strongly continuous semigroup on $\mathcal{B}^q(\mathcal{E})$.

Proof. This follows from the arguments of [13, Section 5]. $\square$

3. APPROXIMATION THEOREMS

In order to establish existence of Markovian solutions for general generators A we could either directly apply Theorem 2.7, where we have to assume that the generator A satisfies on a dense domain D a generalized positive maximum principle and that for at least one $\omega_0 > \omega$ the range of $A - \omega_0$ is dense, or we approximate a general generator A by (pure jump) generators A^n and apply the following (well known) approximation theorems:

Theorem 3.1. Let $(P_t^n)_{n \in \mathbb{N}, t \geq 0}$ be a sequence of strongly continuous semigroups on a Banach space Z with generators $(A^n)_{n \in \mathbb{N}}$ such that there are uniform (in n) growth bounds $M \geq 1$ and $\omega \in \mathbb{R}$ with

$$(3.1) \quad \|P_t^n\|_{L(Z)} \leq M \exp(\omega t)$$

for $t \geq 0$. Let furthermore $D \subset \cap_n \text{dom}(A^n)$ be a dense subspace with the following three properties:

- (i) D is an invariant subspace for all P^n , i.e. for all $f \in D$ we have $P_t^n f \in D$, for $n \geq 0$ and $t \geq 0$.
- (ii) There is a norm $\|\cdot\|_D$ on D such that there are uniform growth bounds with respect to $\|\cdot\|_D$, i.e. there are $M_D \geq 1$ and $\omega_D \in \mathbb{R}$ with

$$\|P_t^n f\|_D \leq M_D \exp(\omega_D t) \|f\|_D$$

for $t \geq 0$ and for $n \geq 0$.

- (iii) The sequence $A^n f$ converges as $n \rightarrow \infty$ for each $f \in D$, in the following sense: there exists a sequence of numbers $a_{nm} \rightarrow 0$ as $n, m \rightarrow \infty$ such that

$$\|A^n f - A^m f\| \leq a_{nm} \|f\|_D$$

holds true for every $f \in D$ and for all n, m .

Then there exists a strongly continuous semigroup $(P_t^\infty)_{t \geq 0}$ with the same growth bound on Z such that $\lim_{n \rightarrow \infty} P_t^n f = P_t^\infty f$ for all $f \in Z$ uniformly on compacts in time and on bounded sets in D . Furthermore on D the convergence is of order $O(a_{nm})$. If in addition for each $n \in \mathbb{N}$, $(P_t^n)_{t \geq 0}$ is a generalized Feller semigroup, then this property transfers also to the limiting semigroup.

Proof. Since D is invariant under P^n , P^m and $D \subset \cap_n \text{dom}(A^n)$ we can take first derivatives and obtain

$$\frac{d}{ds} P_s^n P_{t-s}^m f = P_s^n (A^n - A^m) P_{t-s}^m f,$$

where we use that A^n commutes with P^n on $\text{dom}(A^n)$, which yields for $m, n \in \mathbb{N}$ and $t \geq 0$

$$P_t^n f - P_t^m f = \int_0^t P_s^n (A^n - A^m) P_{t-s}^m f ds.$$

By (3.1), (i), (iii) and (ii), we can estimate

$$\begin{aligned} \|P_t^n f - P_t^m f\| &\leq \int_0^t \|P_s^n (A^n - A^m) P_{t-s}^m f\| ds \\ &\leq \int_0^t M \exp(\omega s) \|(A^n - A^m) P_{t-s}^m f\| ds \\ &\leq \int_0^t M \exp(\omega s) a_{nm} \|P_{t-s}^m f\|_D ds \\ &\leq a_{nm} M M_D \|f\|_D \int_0^t \exp(\omega s + \omega_D(t-s)) ds. \end{aligned}$$

This yields uniform convergence of $P_t^n f$ on compact intervals in time (with respect to the norm of Z) for all $f \in D$. Hence we obtain bounded linear operators P_t^∞ satisfying the semigroup property and the same growth bound on Z with constants ω and M , in particular $\lim_{n \rightarrow \infty} P_t^n f = P_t^\infty f$ for all $f \in Z$. The convergence rate on D is obvious. Strong continuity follows by uniform convergence on D on bounded sets and by the respective stability estimates. For generalized Feller semigroups, the only additional property is positivity, which of course remains in the limit. $\square$

For the purposes of affine processes a slightly more general version of the approximation theorem is needed, which we state in the sequel:

Theorem 3.2. *Let $(P_t^n)_{n \in \mathbb{N}, t \geq 0}$ be a sequence of strongly continuous semigroups on a Banach space Z with generators $(A^n)_{n \in \mathbb{N}}$ such that there are uniform (in n) growth bounds $M \geq 1$ and $\omega \in \mathbb{R}$ with*

$$\|P_t^n\|_{L(Z)} \leq M \exp(\omega t)$$

for $t \geq 0$. Let furthermore $D \subset \cap_n \text{dom}(A^n)$ be a subset with the following two properties:

- (i) *The linear span $\text{span}(D)$ is dense.*
- (ii) *There is a norm $\|\cdot\|_D$ on $\text{span}(D)$ such that for each $f \in D$ and for $t > 0$ there exists a sequence $a_{nm}^{f,t}$, possibly depending on f and $t > 0$,*

$$\|A^n P_u^m f - A^m P_u^m f\| \leq a_{nm}^{f,t} \|f\|_D$$

holds true for n, m and for $0 \leq u \leq t$, with $a_{nm}^{f,t} \rightarrow 0$ as $n, m \rightarrow \infty$.

Then there exists a strongly continuous semigroup $(P_t^\infty)_{t \geq 0}$ with the same growth bound on Z such that $\lim_{n \rightarrow \infty} P_t^n f = P_t^\infty f$ for all $f \in Z$ uniformly on compacts in time. If in addition for each $n \in \mathbb{N}$, $(P_t^n)_{t \geq 0}$ is a generalized Feller semigroup, then this property transfers also to the limiting semigroup.

Proof. Again we apply the following well known formula for $f \in D$, $m, n \geq 0$ and $t \geq 0$

$$P_t^n f - P_t^m f = \int_0^t P_s^n (A^n - A^m) P_{t-s}^m f ds.$$

By (3.1), (i), (iii) and (ii), we can estimate

$$\begin{aligned} \|P_t^n f - P_t^m f\| &\leq \int_0^t \|P_s^n (A^n - A^m) P_{t-s}^m f\| ds \\ &\leq \int_0^t M \exp(\omega s) \|(A^n - A^m) P_{t-s}^m f\| ds \\ &\leq \int_0^t M \exp(\omega s) a_{nm}^{f,t} \|f\|_D ds \end{aligned}$$

This yields uniform convergence of $P_t^n f$ on compact intervals in time (with respect to the norm of Z) for all $f \in D$ by assumption. Hence we obtain bounded linear operators P_t^∞ satisfying the semigroup property and the same growth bound on Z with constants ω and M , in particular $\lim_{n \rightarrow \infty} P_t^n f = P_t^\infty f$ for all $f \in Z$ by stability of growth bounds. Strong continuity follows by the respective stability estimates, i.e.

$$\|P_t^\infty f - f\| \leq \|P_t^\infty f - P_t^\infty g\| + \|P_t^\infty g - P_t^n g\| + \|P_t^n g - g\| + \|g - f\|$$

for $g \in \text{span}(D)$, where the first term is small due to $\|P_t^\infty\| \leq M \exp(\omega t)$ for $t \geq 0$. For generalized Feller semigroups, the only additional property is positivity, which of course remains in the limit. $\square$

Our first application of Theorem 3.1 is the next proposition, where generalized Feller processes of pure jump type with unbounded activity are constructed.

Proposition 3.3. *Let (X, ϱ) be a weighted space with weight function $\varrho \geq 1$. Consider an operator A on $\mathcal{B}^{\varrho}(X)$ with dense domain $\text{dom}(A)$ generating on $\mathcal{B}^{\varrho}(X)$ a generalized Feller semigroup $(P_t)_{t \geq 0}$ of transport type as in (2.8), such that for all $t \geq 0$ we have $\|P_t\|_{L(\mathcal{B}^{\varrho}(X))} \leq M_1 \exp(\omega t)$ for some M_1 and ω and such that $\mathcal{B}^{\sqrt{\varrho}}(X) \subset \mathcal{B}^{\varrho}(X)$ is left invariant and $(P_t)_{t \geq 0}$ is strongly continuous thereon.*

Consider furthermore a family of finite nonnegative measures $\mu(x, \cdot)$ for $x \in X$ on X such that the operator B acts on $\mathcal{B}^{\varrho}(X)$ by

$$Bf(x) := \int (f(y) - f(x)) \mu(x, dy)$$

for $x \in X$ yielding continuous functions on $\{\varrho \leq R\}$ for $R \geq 0$, and such that the following properties hold true:

- For all $x \in X$

$$(3.2) \quad \int \varrho(y) \mu(x, dy) \leq M \varrho^2(x),$$

as well as

$$(3.3) \quad \int \sqrt{\varrho(y)} \mu(x, dy) \leq M \varrho(x),$$

and

$$(3.4) \quad \int \mu(x, dy) \leq M \sqrt{\varrho(x)},$$

hold true for some constant M .

- For some constant $\tilde{\omega} \in \mathbb{R}$

$$(3.5) \quad \int \left| \frac{\sup_{t \geq 0} \exp(-\omega t) P_t \varrho(y) - \sup_{t \geq 0} \exp(-\omega t) P_t \varrho(x)}{\sup_{t \geq 0} \exp(-\omega t) P_t \varrho(x)} \right| \mu(x, dy) \leq \tilde{\omega},$$

for all $x \in X$. In particular $y \mapsto \sup_{t \geq 0} \exp(-\omega t) P_t \varrho(y)$ should be integrable with respect to $\mu(x, \cdot)$

Then $A + B$ generates a generalized Feller semigroup $(P_t^\infty)_{t \geq 0}$ on $\mathcal{B}^e(X)$ satisfying $\|P_t^\infty\|_{L(\mathcal{B}^e(X))} \leq M_1 \exp((\omega + \tilde{\omega})t)$.

Proof. We apply Theorem 3.1 with $D = \text{dom}(A) \cap \mathcal{B}^{\sqrt{e}}(X)$. We construct a sequence of pure jump, bounded activity generators on $\mathcal{B}^e(X)$ and on $\mathcal{B}^{\sqrt{e}}(X)$ by

$$B^n f(x) := \int (f(y) - f(x)) \frac{n}{\varrho(x) \vee n} \mu(x, dy).$$

Indeed, they are bounded with respect to both norms: by (3.2) and (3.4), we have for $f \in \mathcal{B}^e(X)$

$$\begin{aligned} \frac{B^n f(x)}{\varrho(x)} &\leq \frac{\int (\frac{f(y)}{\varrho(y)} \varrho(y) - f(x)) \frac{n}{\varrho(x) \vee n} \mu(x, dy)}{\varrho(x)} \\ &\leq \|f\|_e 2Mn \end{aligned}$$

and by (3.3) and (3.4) for $f \in \mathcal{B}^{\sqrt{e}}(X)$

$$\begin{aligned} \frac{B^n f(x)}{\sqrt{\varrho(x)}} &\leq \frac{\int (\frac{f(y)}{\sqrt{\varrho(y)}} \sqrt{\varrho(y)} - f(x)) \frac{n}{\varrho(x) \vee n} \mu(x, dy)}{\sqrt{\varrho(x)}} \\ &\leq \|f\|_{\sqrt{e}} Mn(\sqrt{n} + 1). \end{aligned}$$

Consider now the operators $A^n := A + B^n$, on the one hand with domain $\text{dom}(A)$ and on the other hand with domain $D = \text{dom}(A) \cap \mathcal{B}^{\sqrt{e}}(X)$. By Theorem 2.10 they generate generalized Feller semigroups P^n on $\mathcal{B}^e(X)$ and $\mathcal{B}^{\sqrt{e}}(X)$ respectively by assumption. Since the domain of the generator is anyhow left invariant, Condition (i) of Theorem 3.1 is satisfied with $D = \text{dom}(A) \cap \mathcal{B}^{\sqrt{e}}(X)$. Moreover, by (3.3) and (3.4) we have for $f \in D$

$$\begin{aligned} \left| \frac{B^n f(x) - B^m f(x)}{\varrho(x)} \right| &\leq \|f\|_{\sqrt{e}} \frac{\int \sqrt{\varrho(y)} \left| \frac{n}{\varrho(x) \vee n} - \frac{m}{\varrho(x) \vee m} \right| \mu(x, dy)}{\varrho(x)} + \\ &\quad + \|f\|_{\sqrt{e}} \frac{\int \left| \frac{n}{\varrho(x) \vee n} - \frac{m}{\varrho(x) \vee m} \right| \mu(x, dy)}{\sqrt{\varrho(x)}} \leq a_{nm} \|f\|_{\sqrt{e}} \end{aligned}$$

for some $a_{nm} \rightarrow 0$ as $n, m \rightarrow \infty$. Hence we also have that

$$\|Af + B^n f - (Af + B^m f)\|_e \leq a_{nm} \|f\|_{\sqrt{e}}$$

for all $f \in D$, implying that Condition (iii) of Theorem 3.1 is satisfied.

We finally have to check whether the growth bounds on $\mathcal{B}^e(X)$ and $D \subseteq \mathcal{B}^{\sqrt{e}}(X)$ are uniform in n . In view of Remark 2.9 we denote

$$\tilde{\varrho}(y) = \sup_{t \geq 0} \exp(-\omega t) P_t \varrho(y)$$

and apply Condition (3.5) and the following immediate consequence of Condition (3.5)

$$\begin{aligned} \int \left| \frac{\sqrt{\tilde{\varrho}(y)} - \sqrt{\tilde{\varrho}(x)}}{\sqrt{\tilde{\varrho}(x)}} \right| \mu(x, dy) &\leq \int \left| \frac{\sqrt{\tilde{\varrho}(y)} - \sqrt{\tilde{\varrho}(x)}}{\sqrt{\tilde{\varrho}(x)}} \right| \frac{\sqrt{\tilde{\varrho}(y)} + \sqrt{\tilde{\varrho}(x)}}{\sqrt{\tilde{\varrho}(x)}} \mu(x, dy) \\ &= \int \left| \frac{\tilde{\varrho}(y) - \tilde{\varrho}(x)}{\tilde{\varrho}(x)} \right| \mu(x, dy) \leq \tilde{\omega}, \quad \text{for all } x \in X. \end{aligned}$$

These conditions yield that $B^n \tilde{\varrho}(x) \leq \tilde{\omega} \tilde{\varrho}(x)$ and $B^n \sqrt{\tilde{\varrho}(x)} \leq \tilde{\omega} \sqrt{\tilde{\varrho}(x)}$ for all $n \in \mathbb{N}$. By Remark 2.9 the transport semigroup P is quasicontractive with respect to $\tilde{\varrho}$ and also $\sqrt{\tilde{\varrho}}$. The latter follows by an analogous estimate as in (2.9).

Hence we can readily apply the second part of Theorem 2.10 with ϱ ($\sqrt{\varrho}$ respectively) replaced by $\tilde{\varrho}$ ($\sqrt{\tilde{\varrho}}$ respectively) to get the following uniform growth bound for $(P^n)_n$

$$\|P_t^n\|_{(L(\mathcal{B}^e(X)), \tilde{\varrho})} \leq \exp((\omega + \tilde{\omega})t), \quad \|P_t^n\|_{(L(\mathcal{B}^{\sqrt{e}}(X)), \sqrt{\tilde{\varrho}})} \leq \exp((\frac{\omega}{2} + \tilde{\omega})t)$$

on $\mathcal{B}^e(X)$ ($D \subseteq \mathcal{B}^{\sqrt{e}}(X)$ respectively) equipped with the norm induced by $\tilde{\varrho}$ ($\sqrt{\tilde{\varrho}}$ respectively). The latter is indicated in the subscript of the above norms. Translating this back to the norm corresponding to ϱ yields

$$\begin{aligned} \|P_t^n f\|_{\varrho} &= \sup_{x \in X} \frac{|P_t^n f(x)|}{\tilde{\varrho}(x)} \frac{\tilde{\varrho}(x)}{\varrho(x)} \leq \|P_t^n f\|_{\tilde{\varrho}} \sup_{x \in X} \frac{\tilde{\varrho}(x)}{\varrho(x)} \\ &\leq \exp((\omega + \tilde{\omega})t) \|f\|_{\tilde{\varrho}} M_1 \leq \exp((\omega + \tilde{\omega})t) \|f\|_{\varrho} M_1, \end{aligned}$$

where we use $\varrho \leq \tilde{\varrho} \leq M_1 \varrho$ as asserted in Remark 2.9. Similarly we have

$$\|P_t^n f\|_{\sqrt{\varrho}} \leq \exp((\frac{\omega}{2} + \tilde{\omega})t) \|f\|_{\sqrt{\varrho}} \sqrt{M_1}.$$

We therefore also obtain uniform growth bounds for the original norms. Hence the approximation Theorem 3.1 can be applied and leads to a generalized Feller semigroup P_t^∞ with generator $A+B$ whose growth bound satisfies $\|P_t^\infty\|_{L(\mathcal{B}^e(X))} \leq M_1 \exp((\omega + \tilde{\omega})t)$ as asserted. $\square$

In contrast to classical Feller theory, Proposition 3.3 allows to construct processes with unbounded jump intensities directly if ϱ is unbounded on X . The general character of the proposition allows to build general processes from simple ones by perturbation. The following corollary shows this in the case of affine processes on $\mathbb{R}^d$.

Corollary 3.4. *Let $X \subset \mathbb{R}^d$ be closed subset and let $\varrho(x) := 1 + \|x\|^2$ be a weight function. Let furthermore A be the generator of a generalized Feller semigroup $(P_t)_{t \geq 0}$ on $\mathcal{B}^e(X)$, which satisfies $\|P_t\|_{L(\mathcal{B}^e(X))} \leq M_1 \exp(\omega t)$ for some constants $M_1 \geq 1$ and $\omega \in \mathbb{R}$ and which is of transport type as in (2.8) with $x \mapsto \psi_t(x)$ being an affine function. Assume furthermore that it leaves $\mathcal{B}^{\sqrt{e}}(X) \subset \mathcal{B}^e(X)$ invariant and generates a strongly continuous semigroup there as well. Let*

$$\mu(x, d\xi) := \sum_{i=1}^d x_i \mu^i(d\xi)$$

for some possibly signed measures μ^i with bounded support such that $\mu(x, d\xi)$ defines a family of finite nonnegative measures on X and $x + \text{supp}(\mu(x, \cdot)) \in X$ for all $x \in X$. Then

$$f \mapsto (x \mapsto Af(x) + \int (f(x + \xi) - f(x)) \mu(x, d\xi))$$

for $f \in \text{dom}(A) \cap \mathcal{B}^{\sqrt{e}}(X)$ generates a generalized Feller semigroup on $\mathcal{B}^e(X)$.

Proof. Substituting $y = x + \xi$, one easily verifies that Conditions (3.2) - (3.4) of Proposition 3.3 are satisfied since the supports of μ^i are bounded. Concerning Condition (3.5), denote

$$\tilde{\varrho}(x) = \sup_{t \geq 0} \exp(-\omega t) P_t \varrho(x).$$

In particular we know that $\varrho \leq \tilde{\varrho}$ and it holds that $P_t f(x) = f(\psi_t(x))$ where ψ is an affine function. Using this together with $|\sup_t c(t) - \sup_t d(t)| \leq \sup_t |c(t) - d(t)|$

we obtain for some $\tilde{\omega}$

$$\begin{aligned}
& \int \left| \frac{\tilde{\varrho}(x + \xi) - \tilde{\varrho}(x)}{\tilde{\varrho}(x)} \right| \sum_{i=1}^d x_i \mu^i(d\xi) \\
& \leq \int \left| \frac{\sup_{t \geq 0} \exp(-\omega t) |P_t \varrho(x + \xi) - P_t \varrho(x)|}{\tilde{\varrho}(x)} \right| \sum_{i=1}^d x_i \mu^i(d\xi) \\
& \leq \int \left| \frac{\sup_{t \geq 0} \exp(-\omega t) |\varrho(\psi_t(x + \xi)) - \varrho(\psi_t(x))|}{\tilde{\varrho}(x)} \right| \sum_{i=1}^d x_i \mu^i(d\xi) \\
& = \int \left| \frac{\sup_{t \geq 0} \exp(-\omega t) C(\|\xi\| + \|x\| \|\xi\| + \|\xi\|^2)}{1 + \|x\|^2} \right| \sum_{i=1}^d x_i \mu^i(d\xi) \leq \tilde{\omega},
\end{aligned}$$

where C denotes some constant. All other requirements are met as well and we can conclude. $\square$

4. LIFTING STOCHASTIC VOLTERRA PROCESSES

In the subsequent sections our main goal is to treat the following types of SPDEs

$$\begin{aligned}
(4.1) \quad & d\lambda_t = \mathcal{A}^* \lambda_t dt + \nu dX_t, \\
& \lambda_0 \in \mathcal{E},
\end{aligned}$$

on spaces $\mathcal{E} \subset Y^*$ as introduced in Section 2.2 where $\mathcal{A}^*$ is the generator of a strongly continuous semigroup $\mathcal{S}^*$ on Y^* , $\nu \in Y^*$ (or in a slightly larger space denoted by Z^* in the sequel), $g \in Y$ and X a real valued Itô-semimartingale whose differential characteristics depend linearly on $\langle g, \lambda_t \rangle$, which will turn out to be the solution of the Volterra equation with kernel (1.2).

Remark 4.1. As indicated in the introduction, it will be easy to consider vector valued structures, with X a semimartingale whose characteristics depend – instead of the $\mathbb{R}$ -valued pairing $\langle g, \lambda_t \rangle$ – linearly on a projection of λ_t onto some space, finite or infinite dimensional. For the sake of the first exposition we stay one dimensional here.

In the following we summarize the main ingredients of our setting.

Assumption 4.2. *Throughout this section we shall work under the following conditions:*

- (i) *We consider Banach spaces Z and Y such that $Z \subset Y$ and Z embeds continuously into Y , and their duals $Y^* \subset Z^*$ with their respective weak*-topology.*
- (ii) *We are given an admissible weight function $\varrho = 1 + \varrho_0$ in the sense of Section 2.2 on Y^* such that*

$$\varrho_0(\lambda) = \|\lambda\|_{Y^*}^2, \quad \lambda \in Y^*,$$

where $\|\cdot\|_{Y^}$ denotes the norm on Y^* .*

- (iii) *We are given a closed convex cone $\mathcal{E} \subset Y^*$ such that $(\mathcal{E}, \varrho)$ is a weighted space in the sense of Section 2. This will serve as statespaces of (4.1).*
- (iv) *We assume that a semigroup $\mathcal{S}^*$ with generator $\mathcal{A}^*$ acts in a strongly continuous way on Y^* and Z^* , with respect to the respective norm topologies.*
- (v) *We assume that $\lambda \mapsto \mathcal{S}_t^* \lambda$ is weak*-continuous on Y^* and on Z^* for every $t \geq 0$ (considering the weak*-topology on both the domain and the image space).*
- (vi) *We suppose that the (pre-) adjoint operator of $\mathcal{A}^*$, denoted by $\mathcal{A}$ and domain $\text{dom}(\mathcal{A}) \subset Z \subset Y$, generates a strongly continuous semigroup on Z with respect to the respective norm topology (but not necessarily on Y).*

Remark 4.3. We could allow more general weight functions ϱ but it is not necessary for our purposes here and the formulation of their abstract properties is cumbersome.

Remark 4.4. A prototypical example presented in Section 5.1 is given by $Y = C_b(\mathbb{R}_+ \cup \{\infty\}, \mathbb{R})$ with supremum norm and Z the space of functions $g \in Y$ such that $(x \mapsto xg(x)) \in Y$ together with the operator norm on it, i.e. $\|g\| = \sqrt{\|g\|^2 + \sup_{x \geq 0} |xg(x)|^2}$ for $g \in Z$. The semigroup is given by $\mathcal{S}_t f(x) = \exp(-xt)f(x)$ for $t \geq 0$, $f \in Y$ and $x \geq 0$. All above requirements on spaces and semigroups are then satisfied. Notice also that we can consider non-separable settings since this is not a requirement in our setup.

To analyze (4.1) and to construct $\mathcal{E}$ we first consider the following linear deterministic equation

$$(4.2) \quad d\lambda_t = \mathcal{A}^* \lambda_t dt - w\nu \langle g, \lambda_t \rangle dt$$

for $\lambda_0 \in Y^*$, some fixed $g \in Y$, a real number $w > 0$ and $\nu \in Z^*$.

Under the subsequent assumptions on $\mathcal{S}^*$ and $\nu \in Z^*$ we can guarantee that it can be solved on the space Y^* for all times in the mild sense with respect to the dual norm $\|\cdot\|_{Y^*}$ by a standard Picard iteration method.

Assumption 4.5. *We assume that*

- (i) $\mathcal{S}_t^* \nu \in Y^*$ for all $t > 0$ even though ν does not necessarily lie in Y^* itself, but only in Z^* ;
- (ii) $\int_0^t \|\mathcal{S}_s^* \nu\|_{Y^*}^2 ds < \infty$ for all $t > 0$.

As in (1.2), we define

$$(4.3) \quad K(t) := \langle g, \mathcal{S}_t^* \nu \rangle,$$

which will correspond to the kernel in the Volterra equation (1.1) and define $R \in L_{\text{loc}}^2(\mathbb{R}_+, \mathbb{R})$ as the resolvent of the second kind that satisfies

$$(4.4) \quad K * R = R * K = K - R.$$

This resolvent always exists and is unique (see [22, Theorem 3.1]).

Proposition 4.6. *Under Assumption 4.5, there exists a unique mild solution of (4.2) with values in Y^* . Additionally, the solution operator is a weak*-continuous map $\lambda_0 \mapsto \lambda_t$, for each $t > 0$, and the solution satisfies*

$$\varrho(\lambda_t) \leq C\varrho(\lambda_0), \quad \text{for all } \lambda_0 \in Y^* \text{ and } t \in [0, \varepsilon]$$

for some positive constants C and ε .

Remark 4.7. The unique mild solution of Equation (4.2) satisfies

$$\lambda_t = \mathcal{S}_t^* \lambda_0 - w \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s \rangle ds$$

for all $t \geq 0$. Pairing with $g \in Y$ yields a deterministic linear Volterra equation of the form

$$(4.5) \quad \begin{aligned} \langle g, \lambda_t \rangle &= \langle g, \mathcal{S}_t^* \lambda_0 \rangle - w \int_0^t \langle g, \mathcal{S}_{t-s}^* \nu \rangle \langle g, \lambda_s \rangle ds \\ &= \langle g, \mathcal{S}_t^* \lambda_0 \rangle - w \int_0^t K(t-s) \langle g, \lambda_s \rangle ds, \end{aligned}$$

where we used (4.3).

Proof. We prove first the completely standard convergence of the Picard iteration scheme with respect to the dual norm on Y^* . Define

$$\begin{aligned}\lambda_t^0 &= \lambda_0, \\ \lambda_t^{n+1} &= \mathcal{S}_t^* \lambda_0 - w \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s^n \rangle ds, \quad n \geq 0.\end{aligned}$$

Then, by Assumption 4.5 (i) each λ_t^n lies Y^* . Consider now

$$\begin{aligned}\|\lambda_t^{n+1} - \lambda_t^n\|_{Y^*} &= w \left\| \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s^n - \lambda_s^{n-1} \rangle ds \right\|_{Y^*} \\ &\leq w \int_0^t \|\mathcal{S}_{t-s}^* \nu \langle g, \lambda_s^n - \lambda_s^{n-1} \rangle\|_{Y^*} ds \\ &\leq w \int_0^t \|\mathcal{S}_{t-s}^* \nu\|_{Y^*} \|g\|_Y \|\lambda_s^n - \lambda_s^{n-1}\|_{Y^*} ds.\end{aligned}$$

Assumption 4.5 (ii) and an extended version of Gronwall's inequality see [11, Lemma 15] then yield convergence of $(\lambda_t^n)_{n \in \mathbb{N}}$ to some λ_t with respect to the dual norm $\|\cdot\|_{Y^*}$ uniformly in t on compact intervals. For details on strongly continuous semigroups and mild solutions see [26].

Having established the existence of a mild solution of (4.2) in Y^* , consider now the linear, deterministic Volterra equation (4.5), which can be written as in [22, Section 2], i.e.

$$(4.6) \quad \langle g, \lambda_t \rangle = \langle g, \mathcal{S}_t^* \lambda_0 \rangle - \int_0^t R^w(t-s) \langle g, \mathcal{S}_s^* \lambda_0 \rangle ds,$$

where R^w denotes the resolvent of $wK(t) = w \langle g, \mathcal{S}_t^* \nu \rangle$ as introduced in (4.4). Since by assumption $\mathcal{S}^*$ is a weak-* continuous solution operator, the map $\lambda_0 \mapsto (t \mapsto \langle g, \mathcal{S}_t^* \lambda_0 \rangle)$ is weak-* continuous as a map from Y^* to $C(\mathbb{R}_+, \mathbb{R})$ (with the topology of uniform convergence on compacts on $C(\mathbb{R}_+, \mathbb{R})$). From (4.6) we thus infer that $\langle g, \lambda_t \rangle$ is weak-* continuous for every $t \geq 0$, which clearly translates to the solution map of Equation (4.2).

Finally we have to show that the stated inequality for $\varrho(\lambda_t)$ holds true on small time intervals $[0, \varepsilon]$. Observe first that for $t \in [0, \varepsilon]$

$$\|\mathcal{S}_t^* \lambda\|_{Y^*}^2 \leq C \|\lambda\|_{Y^*}^2$$

for all $\lambda \in Y^*$ just by the assumption that $\mathcal{S}_t^*$ is strongly continuous, for some constant $C \geq 1$. Furthermore for $t \in [0, \varepsilon]$

$$\begin{aligned}\|\lambda_t\|_{Y^*}^2 &\leq 2(C \|\lambda_0\|_{Y^*}^2 + t \int_0^t \|w \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s \rangle\|_{Y^*}^2 ds) \\ &\leq 2(C \|\lambda_0\|_{Y^*}^2 + \varepsilon \int_0^t w^2 \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \|g\|_Y^2 \|\lambda_s\|_{Y^*}^2 ds).\end{aligned}$$

Consider now the kernel $K'(t, s) = 2\varepsilon w^2 \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \|g\|_Y^2 1_{\{s \leq t\}}$. We follow now the arguments of the proof of [3, Lemma 3.1]. Indeed K' is a Volterra kernel as defined in [22, Definition 9.2.1] and for any interval $[U, V] \subset \mathbb{R}_+$ we have by Young's convolution inequality

$$|||K' |||_{L^1([U, V])} \leq 2\varepsilon w^2 \|g\|_Y^2 \int_0^{V-U} \|\mathcal{S}_s^* \nu\|_{Y^*}^2 ds,$$

where $|||\cdot|||_{L^1([U, V])}$ is defined in [22, Definition 9.2.2]. We can now literally take the proof of [3, Lemma 3.1] to deduce that the generalized Gronwall Lemma (see

[22, Lemma 9.8.2]) can be applied. This yields for $t \in [0, \varepsilon]$

$$\|\lambda_t\|_{Y^*}^2 \leq \|\lambda_0\|_{Y^*}^2 2C(1 - \int_0^t R'(s)ds) \leq \|\lambda_0\|_{Y^*}^2 2C(1 - \int_0^\varepsilon R'(s)ds),$$

where R' denotes the resolvent of $-K'$, which is nonpositive. This leads to the desired assertion due to the definition of ϱ . From this inequality also uniqueness follows in a standard way. $\square$

It is of crucial importance to understand that there is actually a closed sub-cone $\mathcal{E} \subset Y^*$ left invariant by the solution of Equation (4.2). This cone will play the role as announced in Assumption 4.2 (iii) and can be described in terms of initial values λ_0 which give rise to nonnegative solutions of (4.6). Indeed, it will be the intersection of the following cones. Define for fixed $w > 0$

$$(4.7) \quad \begin{aligned} \mathcal{E}^w &:= \{\lambda_0 \in Y^* \mid \langle g, \mathcal{S}_t^* \lambda_0 \rangle - \int_0^t R^w(t-s) \langle g, \mathcal{S}_s^* \lambda_0 \rangle ds \geq 0 \text{ for all } t \geq 0\} \\ &= \{\lambda_0 \in Y^* \mid \langle g, \lambda_t \rangle \geq 0 \text{ with } \lambda_t = \mathcal{S}_t^* \lambda_0 - w \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s \rangle ds \text{ for all } t \geq 0\}, \end{aligned}$$

where R^w denotes the resolvent of $wK(t) = w\langle g, \mathcal{S}_t^* \nu \rangle$.

Proposition 4.8. *Let Assumption 4.5 be in force and let $w > 0$ be fixed. The set $\mathcal{E}^w$ as defined in (4.7) is a weak-*closed convex cone. The solution of (4.2) leaves $\mathcal{E}^w$ invariant and defines a generalized Feller semigroup on $\mathcal{B}^q(\mathcal{E}^w)$ by $P_t f(\lambda_0) := f(\lambda_t)$ for all $f \in \mathcal{B}^q(\mathcal{E}^w)$ and $t \geq 0$.*

Proof. The weak-*closedness follows from the weak-*continuity of $\mathcal{S}^*$ and the convex cone property is obvious. Let us now prove the invariance of $\mathcal{E}^w$. Let $\lambda_0 \in \mathcal{E}^w$. We now have to prove that for each $t > 0$ the unique solution of (4.2) given by

$$\lambda_t = \mathcal{S}_t^* \lambda_0 - w \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s \rangle ds$$

lies in $\mathcal{E}^w$ as well. By Definition of $\mathcal{E}^w$, this means to show that $\langle g, \lambda_{t+u} \rangle \geq 0$ for all $u \geq 0$ where

$$\lambda_{t+u} = \mathcal{S}_u^* \lambda_t - w \int_0^u \mathcal{S}_{u-s}^* \nu \langle g, \lambda_{t+s} \rangle ds.$$

This is equal to

$$\lambda_{t+u} = \mathcal{S}_{t+u}^* \lambda_0 - w \int_0^{t+u} \mathcal{S}_{t+u-s}^* \nu \langle g, \lambda_s \rangle ds.$$

Since $\lambda_0 \in \mathcal{E}^w$, we thus have $\langle g, \lambda_{t+u} \rangle \geq 0$ for all $u \geq 0$ and thus $\lambda_t \in \mathcal{E}^w$, proving the invariance. Since by Proposition 4.6 the solution operator is weak-*continuous, we can conclude that $\lambda_0 \mapsto f(\lambda_t)$ lies in $\mathcal{B}^q(\mathcal{E}^w)$ for a dense set of $\mathcal{B}^q(\mathcal{E}^w)$ by Theorem 2.17. Moreover, it satisfies the necessary bound (2.11) for ϱ and (2.12) is satisfied by (norm)-continuity of $t \mapsto \lambda_t$. Hence all the conditions of Assumption 2.19 are satisfied and the solution operator therefore defines a generalized Feller semigroup (P_t) on $\mathcal{B}^q(\mathcal{E}^w)$ by Theorem 2.21. $\square$

We need an additional assumption assuring that the above defined state space contains the cone hull of $\mathcal{S}_t^* \nu$:

Assumption 4.9. *Let ν be such that $\mathcal{S}_u^* \nu \in \mathcal{E}^w$ for all $u > 0$ and for all $w > 0$.*

Remark 4.10. This condition is satisfied if K and R^w are nonnegative for all $w > 0$. Indeed, $\mathcal{S}_u^* \nu \in \mathcal{E}^w$ if and only if

(4.8)

$$\langle g, \mathcal{S}_{t+u}^* \nu \rangle - \int_0^t R^w(t-s) \langle g, \mathcal{S}_{s+u}^* \nu \rangle ds = K(t+u) - \int_0^t R^w(t-s) K(s+u) ds \geq 0.$$

Since by the resolvent equation we have

$$\begin{aligned} R^w(t+u) &= w(K(t+u) - \int_0^{t+u} R^w(t+u-s) K(s) ds) \\ &= w(K(t+u) - \int_0^t R^w(t-s) K(s+u) ds - \int_0^u R^w(t+u-s) K(s) ds), \end{aligned}$$

it follows that

$$K(t+u) - \int_0^t R^w(t-s) K(s+u) ds = \frac{1}{w} R^w(t+u) + \int_0^u R^w(t+u-s) K(s) ds.$$

If K and R^w are nonnegative (4.8) is clearly satisfied.

The following lemma states that the scale of invariant spaces $\mathcal{E}^w$ is actually decreasing.

Lemma 4.11. *Let Assumptions 4.5 and 4.9 be in force. For $w_1 < w_2$ we have $\mathcal{E}^{w_1} \supseteq \mathcal{E}^{w_2}$. Additionally we have that for $\lambda_0 \in \mathcal{E}^{w_2}$ the unique solution of*

$$\lambda_t = \mathcal{S}_t^* \lambda_0 - w_1 \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s \rangle ds$$

for all $t \geq 0$ actually lies in $\mathcal{E}^{w_2}$.

Proof. Fix $w_1 < w_2$ and $\lambda_0 \in \mathcal{E}^{w_2}$. Actually we can write

$$\begin{aligned} \lambda_t &= \mathcal{S}_t^* \lambda_0 - w_1 \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s \rangle ds = \\ &= \underbrace{\mathcal{S}_t^* \lambda_0 - w_2 \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s \rangle ds}_A + \underbrace{(w_2 - w_1) \int_0^t \mathcal{S}_{t-s}^* \nu \langle g, \lambda_s \rangle ds}_B. \end{aligned}$$

By Proposition 4.8, the term A clearly lies in $\mathcal{E}^{w_2}$. The same holds true for the second term B due to Assumption 4.9 and the cone property of $\mathcal{E}^{w_2}$, since $(w_2 - w_1) > 0$ and $\langle g, \lambda_s \rangle \geq 0$ for all $s > 0$. Thus $\lambda_t \in \mathcal{E}^{w_2}$. However, by definition of $\mathcal{E}^{w_1}$, λ_t also lies there, whence the conclusion. $\square$

Definition 4.12. *Let Assumptions 4.5 and 4.9 be in force. Then we define the following weak-*closed cone in Y^**

$$\mathcal{E} = \bigcap_{w>0} \mathcal{E}^w.$$

Theorem 4.13. *Let Assumptions 4.5 and 4.9 be in force. Then the weak-*closed cone $\mathcal{E}$ contains all damped measures $\mathcal{S}_t^* \nu$ for $t > 0$ and is left invariant by the solution of*

$$(4.9) \quad d\lambda_t = \mathcal{A}^* \lambda_t dt - w \nu \langle g, \lambda_t \rangle dt$$

for all $w > 0$. This solution defines a generalized Feller semigroup on $\mathcal{B}^q(\mathcal{E})$.

Proof. The first assertion follows from Assumption 4.9. By monotonicity of $\mathcal{E}^w$ and the fact that solutions with respect to w_1 -equations actually lie in $\mathcal{E}^{w_2}$ as stated in the previous lemma, the invariance assertion and generalized Feller property follow together with Proposition 4.8. $\square$

By the previous results we can now construct a generalized Feller process on $\mathcal{E}$ which jumps up by multiples of $\mathcal{S}_\varepsilon^* \nu$ for some $\varepsilon \geq 0$ and with an instantaneous intensity of size $\langle g, \lambda_t \rangle$. We formulate these assertions in two propositions. For their formulation recall that $\mathcal{E}_* \subset Y$ denotes the (pre-)polar cone of $\mathcal{E}$. Define also the following set

$$(4.10) \quad \mathcal{D} = \{y \in Y \mid y \in \text{dom}(\mathcal{A}) \text{ s.t. } \langle y, \nu \rangle \text{ is well-defined}\}.$$

Proposition 4.14. *Let Assumptions 4.5 and 4.9 be in force. Moreover, let $\mu \in \mathcal{M}_+(\mathbb{R}_+)$ with finite second moment. Consider the SPDE*

$$(4.11) \quad d\lambda_t = \mathcal{A}^* \lambda_t dt - w\nu \langle g, \lambda_t \rangle dt + \mathcal{S}_\varepsilon^* \nu dN_t,$$

where $(N_t)_{t \geq 0}$ is a pure jump process with compensator $F(\lambda_t, d\xi) = \langle g, \lambda_t \rangle \mu(d\xi)$.

- (i) *Then for every $\lambda_0 \in \mathcal{E}$, $\varepsilon > 0$ and $w > 0$, the SPDE (4.11) has a solution in $\mathcal{E}$ given by a generalized Feller process associated to the generator of (4.11).*
- (ii) *This generalized Feller process is also a probabilistically weak and analytically mild solution of (4.11), i.e.*

$$\lambda_t = \mathcal{S}_t^* \lambda_0 - \int_0^t \mathcal{S}_{t-s}^* w\nu \langle g, \lambda_s \rangle ds + \int_0^t \mathcal{S}_{t-s+\varepsilon}^* \nu dN_s,$$

which justifies Equation (4.11), in particular for every initial value the process N can be constructed on an appropriate probabilistic basis. The stochastic integral is defined in a pathwise way along finite variation paths. Moreover, for every family $(f_n)_n \in \mathcal{B}^q(\mathcal{E})$, $t \mapsto f_n(\lambda_t)$ can be chosen to be càglàd for all n .

- (iii) *For every $\varepsilon > 0$ and $w > 0$, the corresponding Riccati equation $\partial_t y_t = R(y_t)$ with $R : \mathcal{D} \cap \mathcal{E}_* \rightarrow \mathbb{R}$ given by*

$$(4.12) \quad R(y) = \mathcal{A}y - wg\langle y, \nu \rangle + g \int_{\mathbb{R}_+} (\exp(\langle y, \mathcal{S}_\varepsilon^* \nu \xi \rangle) - 1) \mu(d\xi),$$

admits a unique global solution in the mild sense for all initial values $y_0 \in \mathcal{E}_*$.

- (iv) *The affine transform formula holds true.*

$$\mathbb{E}_{\lambda_0} [\exp(\langle y_0, \lambda_t \rangle)] = \exp(\langle y_t, \lambda_0 \rangle),$$

where y_t solves $\partial_t y_t = R(y_t)$ for all $y_0 \in \mathcal{E}_*$ in the mild sense with R given by (4.12). Moreover $y_t \in \mathcal{E}_*$ for all $t \geq 0$.

Proof. To prove the first assertion we apply Proposition 3.3. By Theorem 4.13, the deterministic equation (4.9) has a mild solution on $\mathcal{E}$ which – by Assumption 4.5 – defines a generalized Feller semigroup $(P_t)_{t \geq 0}$ on $\mathcal{B}^q(\mathcal{E})$. The operator A in Proposition 3.3 then corresponds to the generator of $(P_t)_{t \geq 0}$, i.e. the semigroup associated to the purely deterministic part of (4.11), which is clearly of transport type and with growth bound $M_1 \exp(\omega t)$ for some $M_1 \geq 1$ and ω .

Note that by the same arguments as in Proposition 4.8 and by applying Theorem 2.21, we can prove that $(P_t)_{t \geq 0}$ also defines a generalized Feller semigroup on $\mathcal{B}^{\sqrt{q}}(\mathcal{E})$. Indeed, by weak- $*$ -continuity, we can conclude that $\lambda_0 \mapsto f(\lambda_t)$ lies in $\mathcal{B}^{\sqrt{q}}(\mathcal{E})$ for a dense set of $\mathcal{B}^{\sqrt{q}}(\mathcal{E})$ by Theorem 2.17, whence condition (iii) of Assumption 2.19 is satisfied. Moreover, (2.12) is satisfied by (norm)-continuity of $t \mapsto \lambda_t$ and the necessary bound (2.11) holds also for $\sqrt{q}$. For the latter observe first that

$$\|\mathcal{S}_t^* \lambda\|_{Y^*} \leq C \|\lambda\|_{Y^*} \quad \text{for all } 0 \leq t \leq T \text{ and } \lambda \in Y^*,$$

since $\mathcal{S}^*$ is strongly continuous on Y^* , for some $T > 0$. Using this we can estimate

$$(4.13) \quad \|\lambda_t\|_{Y^*} \leq C\|\lambda_0\|_{Y^*} + \int_0^t w\|\mathcal{S}_{t-s}^*\nu\|_{Y^*}\|g\|_Y\|\lambda_s\|_{Y^*}ds.$$

We now proceed similarly as in Proposition 4.6. Consider the kernel $K'(t, s) = w\|\mathcal{S}_{t-s}^*\nu\|_{Y^*}\|g\|_Y 1_{\{s \leq t\}}$, which is again a Volterra kernel in the sense of [22, Definition 9.2.1]. Moreover, for any interval $[U, V] \subset \mathbb{R}_+$ we have by Young's convolution inequality

$$|||K' |||_{L^1([U, V])} \leq w\|g\|_Y \int_0^{V-U} \|\mathcal{S}_s^*\nu\|_{Y^*}^2 ds \leq (V-U)w\|g\|_Y^2 \left(\int_0^{V-U} \|\mathcal{S}_s^*\nu\|_{Y^*}^2 ds \right)^{\frac{1}{2}},$$

where $|||\cdot|||_{L^1([U, V])}$ is defined in [22, Definition 9.2.2]. We now can again literally take the proof of [3, Lemma 3.1] to deduce that the generalized Gronwall Lemma (see [22, Lemma 9.8.2]) can be applied. This yields for $t \in [0, T]$

$$\|\lambda_t\|_{Y^*} \leq \|\lambda_0\|_{Y^*} C \left(1 - \int_0^t R'(s) ds \right),$$

where R' denotes the resolvent of $-K'$, which is nonpositive. Hence, for $t \in [0, T]$ we have by Jensen's inequality

$$\begin{aligned} \sqrt{\varrho(\lambda_t)} &= \sqrt{1 + \|\lambda_t\|_{Y^*}^2} \leq 1 + \|\lambda_t\|_{Y^*} \leq 1 + \|\lambda_0\|_{Y^*} \left(C - \int_0^t R'(s) ds \right) \\ &\leq \tilde{C} \sqrt{1 + \|\lambda_0\|_{Y^*}^2} = \tilde{C} \sqrt{\varrho(\lambda_0)}, \end{aligned}$$

where $\tilde{C}$ depends on T .

Finally, we need to verify (3.2) - (3.4), which read as follows

$$\begin{aligned} \int \varrho(\lambda + \mathcal{S}_\varepsilon^* \nu \xi) \langle g, \lambda \rangle \mu(d\xi) &\leq M \varrho(\lambda)^2, \\ \int \sqrt{\varrho}(\lambda + \mathcal{S}_\varepsilon^* \nu \xi) \langle g, \lambda \rangle \mu(d\xi) &\leq M \varrho(\lambda), \\ \int \langle g, \lambda \rangle \mu(d\xi) &\leq M \sqrt{\varrho(\lambda)}. \end{aligned}$$

which hold true by the second moment condition on μ . Concerning (3.5), denote as in Remark 2.9

$$\tilde{\varrho}(\lambda) = \sup_{t \geq 0} \exp(-\omega t) P_t \varrho(\lambda).$$

In particular we know that $\varrho \leq \tilde{\varrho}$ and it holds that $P_t f(x) = f(\psi_t(x))$ where ψ is the solution of (4.9) which is linear. Using this together with $|\sup_t c(t) - \sup_t d(t)| \leq \sup_t |c(t) - d(t)|$ we obtain similarly as in the proof of Corollary 3.4

$$\begin{aligned} &\int \left| \frac{\tilde{\varrho}(\lambda + \mathcal{S}_\varepsilon^* \nu \xi) - \tilde{\varrho}(\lambda)}{\tilde{\varrho}(\lambda)} \right| \langle g, \lambda \rangle \mu(d\xi) \\ &\leq \int \left| \frac{\sup_{t \geq 0} \exp(-\omega t) |P_t \varrho(\lambda + \mathcal{S}_\varepsilon^* \nu \xi) - P_t \varrho(\lambda)|}{\tilde{\varrho}(\lambda)} \right| \langle g, \lambda \rangle \mu(d\xi) \\ &\leq \int \left| \frac{\sup_{t \geq 0} \exp(-\omega t) |\varrho(\psi_t(\lambda + \mathcal{S}_\varepsilon^* \nu \xi)) - \varrho(\psi_t(\lambda))|}{\tilde{\varrho}(\lambda)} \right| \langle g, \lambda \rangle \mu(d\xi) \\ &= \int \left| \frac{\sup_{t \geq 0} \exp(-\omega t) (2\|\psi_t(\lambda)\|_{Y^*} \|\psi_t(\mathcal{S}_\varepsilon^* \nu \xi)\|_{Y^*} + \|\psi_t(\mathcal{S}_\varepsilon^* \nu \xi)\|_{Y^*}^2)}{\varrho(\lambda)} \right| \langle g, \lambda \rangle \mu(d\xi) \leq \tilde{\omega}. \end{aligned}$$

for some $\tilde{\omega} \in \mathbb{R}$. The last inequality holds by the linearity of ψ and the second moment condition on μ . Proposition 3.3 now allows to conclude that $A + B$ where

B is given by

$$Bf(\lambda) = \int (f(\lambda + \mathcal{S}_\varepsilon^* \nu \xi) - f(\lambda)) \langle g, \lambda \rangle \mu(d\xi)$$

generates a generalized Feller semigroup $\tilde{P}$ as asserted.

For (ii) we now construct the probabilistically weak and analytically mild solution directly from the properties of the generalized Feller process: take $y \in \mathcal{D}$ defined in (4.10) and consider the martingale

$$(4.14) \quad \begin{aligned} M_t^y &:= \langle y, \lambda_t \rangle - \langle y, \lambda_0 \rangle - \int_0^t \langle \mathcal{A}y, \lambda_s \rangle - w \langle y, \nu \rangle \langle g, \lambda_s \rangle ds \\ &\quad - \int_0^t \int \langle y, \mathcal{S}_\varepsilon^* \nu \xi \rangle \langle g, \lambda_s \rangle \mu(d\xi) ds \end{aligned}$$

for $t \geq 0$ (after appropriate regularization as it is possible due to Theorem 2.13 such that the integral term is well defined). Let now y be as above with the additional property that $\langle y, \mathcal{S}_\varepsilon^* \nu \rangle = 1$. Define

$$(4.15) \quad N_t := M_t^y + \int_0^t \int \langle y, \mathcal{S}_\varepsilon^* \nu \xi \rangle \langle g, \lambda_s \rangle \mu(d\xi) ds$$

for $t \geq 0$, which is a càglàd semimartingale. Then N does not depend on y . Indeed, for all y_i with $\langle y_i, \mathcal{S}_\varepsilon^* \nu \rangle = 1$, $i = 1, 2$, we clearly have

$$\int_0^t \int \langle y_1 - y_2, \mathcal{S}_\varepsilon^* \nu \xi \rangle \langle g, \lambda_s \rangle \mu(d\xi) ds = 0$$

and $M^{y_1} - M^{y_2} = M^{y_1 - y_2} = 0$ and as well. The latter follows from the fact that the martingale M^y is constant if $\langle y, \mathcal{S}_\varepsilon^* \nu \rangle = 0$. This is indeed true since the martingale's quadratic variation vanishes in this case: we apply here the carré du champ formula

$$\mathbb{E}[[M_t^y, M_t^y]] = \mathbb{E}\left[\int_0^t (Af^2(\lambda_s) - 2f(\lambda_s)Af(\lambda_s)) ds\right]$$

where $f(\lambda) = \langle y, \lambda \rangle$ with y satisfying $\langle y, \mathcal{S}_\varepsilon^* \nu \rangle = 0$. Moreover, by the definition of N in (4.15) its compensator is given by $\int_0^t \int \xi \langle \lambda_s, g \rangle \mu(d\xi)$ showing that N has the desired properties.

By (4.14) and (4.15) we additionally obtain that

$$\langle y, \lambda_t \rangle = \langle y, \lambda_0 \rangle + \int_0^t \langle \mathcal{A}y, \lambda_s \rangle ds - \int_0^t \langle y, w\nu \rangle \langle g, \lambda_s \rangle ds + \langle y, \mathcal{S}_\varepsilon^* \nu \rangle N_t$$

for $y \in \mathcal{D}$. This analytically weak form can be translated into a mild form by standard methods. Indeed, notice that the integral is just along a finite variation path and therefore we can readily apply variation of constants. The last assertion about the càglàd property is a consequence of Theorem 2.13 by noting that $\varrho(\lambda)$ does not explode. This proves (ii).

Concerning (iii), note first that we have a unique mild solution to

$$(4.16) \quad \partial_t y_t = \mathcal{A}y_t - wg \langle y_t, \nu \rangle, \quad y_0 \in Y,$$

since this is the adjoint equation of (4.2). For the equation with jumps we proceed as in Proposition 4.6 via Picard iteration. Denote the semigroup associated to (4.16) by $\mathcal{S}^w$ and define

$$\begin{aligned} y_t^0 &= y_0, \\ y_t^n &= \mathcal{S}_t^w y_0 + \int_0^t \mathcal{S}_{t-s}^w g \left(\int_{\mathbb{R}_+} (\exp(\langle y_s^{n-1}, \mathcal{S}_\varepsilon^* \nu \xi \rangle) - 1) \mu(d\xi) \right) ds. \end{aligned}$$

Moreover, for $t \in [0, \delta]$ for some $\delta > 0$ we have by local Lipschitz continuity of $x \mapsto \exp(x)$

$$\begin{aligned} \|y_t^{n+1} - y_t^n\|_Y &\leq \left\| \int_0^t \mathcal{S}_{t-s}^w g \left(\exp(\langle y_s^n, \mathcal{S}_\varepsilon^* \nu \xi \rangle) - \exp(\langle y_s^{n-1}, \mathcal{S}_\varepsilon^* \nu \xi \rangle) \right) \mu(d\xi) ds \right\|_Y \\ &\leq \int_0^t C \|\mathcal{S}_{t-s}^w g\|_Y \|y_t^n - y_t^{n-1}\|_Y \left(\int \|\mathcal{S}_\varepsilon^* \nu \xi\|_{Y^*} \mu(d\xi) \right) ds. \end{aligned}$$

By an extension of Gronwall's inequality (see [11, Lemma 15]) this yields convergence of $(y_t^n)_{n \in \mathbb{N}}$ with respect to $\|\cdot\|_Y$ and hence the existence of a unique local mild solution to (4.12) up to some maximal life time $t_+(y_0)$. That $t_+(y_0) = \infty$ for all $y_0 \in \mathcal{E}_*$ follows from the subsequent estimate

$$\begin{aligned} \|y_t\|_Y &= \|\mathcal{S}_t^w y_0 + \int_0^t \mathcal{S}_{t-s}^w g \left(\int (\exp(\langle y_s, \mathcal{S}_\varepsilon^* \nu \xi \rangle) - 1) \mu(d\xi) \right) ds\|_Y \\ &\leq \|\mathcal{S}_t^w y_0\|_Y + \int_0^t \|\mathcal{S}_{t-s}^w g\|_Y \left(\int |\exp(\langle y_s, \mathcal{S}_\varepsilon^* \nu \xi \rangle) - 1| \mu(dx) \right) ds \\ &\leq \|\mathcal{S}_t^w y_0\|_Y + t \sup_{s \leq t} \|\mathcal{S}_s^w g\|_Y \mu(\mathbb{R}_+), \end{aligned}$$

where we used $|\exp(\langle y, \mathcal{S}_\varepsilon^* \nu \xi \rangle) - 1| \leq 1$ for all $y \in \mathcal{E}_*$ in the last estimate.

To prove (iv), just note that by the existence of a generalized Feller semigroup the abstract Cauchy problem for initial value $\exp(\langle y_0, \cdot \rangle)$ can be solved uniquely for $y_0 \in \mathcal{E}_*$. Indeed, $\mathbb{E}_\lambda[\exp(\langle y_0, \lambda_t \rangle)]$ uniquely solves

$$\partial_t u(t, \lambda) = Au(t, \lambda), \quad u(0, \lambda) = \exp(\langle y_0, \lambda \rangle),$$

where A denotes the generator associated to (4.11). Setting $u(t, \lambda) = \exp(\langle y_t, \lambda \rangle)$, we have

$$\partial_t u(t, \lambda) = \exp(\langle y_t, \lambda \rangle) R(y_t),$$

where the right hand side is nothing else than $A \exp(\langle y_t, \lambda \rangle)$, hence the affine transform formula holds true. This also implies that $y_t \in \mathcal{E}_*$ for all $t \geq 0$, simply because $\mathbb{E}_\lambda[\exp(\langle y_0, \lambda_t \rangle)] \leq 1$ for all $\lambda \in \mathcal{E}$. $\square$

The next statement is a refinement of Conclusion (i) of Proposition 4.14 for the case $\varepsilon = 0$. All other conclusions are explicitly stated in Theorem 4.17 below.

Proposition 4.15. *Let Assumptions 4.5 and 4.9 be in force. Moreover, let $\mu \in \mathcal{M}_+(\mathbb{R}_+)$ with finite second moment. Then Conclusion (i) of Proposition 4.14 holds true for $\varepsilon = 0$. Moreover, the affine transform formula for the generalized Feller process $(\lambda_t)_{t \geq 0}$ of (4.11) with $\varepsilon = 0$ is also satisfied, i.e.*

$$\mathbb{E}_{\lambda_0} [\exp(\langle y_0, \lambda_t \rangle)] = \exp(\langle y_t, \lambda_0 \rangle),$$

where y_t solves $\partial_t y_t = R(y_t)$ for all $y_0 \in \mathcal{E}_*$ in the mild sense with R given by (4.12) with $\varepsilon = 0$. Furthermore, $y_t \in \mathcal{E}_*$ for all $t \geq 0$.

Remark 4.16. The essential point here is that we loose the càglàd property when we let ε tend to zero. As long as the kernel K has a singularity at $t = 0$ it is impossible to preserve finite growth bounds *without* $M > 1$ as $\varepsilon \rightarrow 0$. To apply the second conclusion of Theorem 2.13 yielding càglàd trajectories it is however crucial that $M = 1$.

Proof. We apply Theorem 3.2 and consider a sequence of generalized Feller semigroups $(P^n)_{n \in \mathbb{N}}$ with generators A^n corresponding to the solution of (4.11) for $\varepsilon = \frac{1}{n}$, $n \in \mathbb{N}$. Let us first establish a uniform growth bound for this sequence.

Note that for the solution of (4.11), we have due to Proposition 4.14 (ii) the following estimate for $t \in [0, T]$ for some fixed $T > 0$

$$\begin{aligned}
\mathbb{E}[\|\lambda_t^\varepsilon\|_{Y^*}^2] &\leq 3\|\mathcal{S}_t^* \lambda_0\|_{Y^*}^2 + 3t \int_0^t w^2 \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \|g\|_Y^2 \mathbb{E}[\|\lambda_s^\varepsilon\|_{Y^*}^2] ds \\
&\quad + 6\mathbb{E}[\|\int_0^t \mathcal{S}_{t-s+\varepsilon}^* \nu dN_s - \int_0^t \int \mathcal{S}_{t-s+\varepsilon}^* \nu \xi \mu(d\xi) \langle g, \lambda_s^\varepsilon \rangle ds\|^2] \\
&\quad + 6\mathbb{E}[\|\int_0^t \int \mathcal{S}_{t-s+\varepsilon}^* \nu \xi \mu(d\xi) \langle g, \lambda_s^\varepsilon \rangle ds\|^2] \\
&\leq C_0 \|\lambda_0\|_{Y^*}^2 + 3t \int_0^t w^2 \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \|g\|_Y^2 \mathbb{E}[\|\lambda_s^\varepsilon\|_{Y^*}^2] ds \\
&\quad + 6\mathbb{E}[\int \xi^2 \mu(d\xi) \int_0^t \|\mathcal{S}_{t-s+\varepsilon}^* \nu\|_{Y^*}^2 \langle g, \lambda_s^\varepsilon \rangle ds] \\
&\quad + C_1 \int \xi^2 \mu(d\xi) \int_0^t \|\mathcal{S}_{t-s+\varepsilon}^* \nu\|_{Y^*}^2 \|g\|_Y^2 \mathbb{E}[\|\lambda_s^\varepsilon\|_{Y^*}^2] ds \\
&\leq C_0 \|\lambda_0\|_{Y^*}^2 + 3t \int_0^t w^2 \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \|g\|_Y^2 \mathbb{E}[\|\lambda_s^\varepsilon\|_{Y^*}^2] ds \\
&\quad + C_2 \int \xi^2 \mu(d\xi) \int_0^t \|\mathcal{S}_{t-s+\varepsilon}^* \nu\|_{Y^*}^2 ds \\
&\quad + C_1 \int \xi^2 \mu(d\xi) \int_0^t \|\mathcal{S}_{t-s+\varepsilon}^* \nu\|_{Y^*}^2 \|g\|_Y^2 \mathbb{E}[\|\lambda_s^\varepsilon\|_{Y^*}^2] ds \\
&\leq C_0 \|\lambda_0\|_{Y^*}^2 + C_2 \int_0^t \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 ds + C_1 \int_0^t \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \mathbb{E}[\|\lambda_s^\varepsilon\|_{Y^*}^2] ds,
\end{aligned}$$

where the constants can change from line to line and depend on T . We use $\|\mathcal{S}_t^* \lambda_0\|^2 \leq C_0 \|\lambda_0\|^2$ for $t \in [0, T]$, as well as $\|\mathcal{S}_{t-s+\varepsilon}^* \nu\|_{Y^*} \leq \widehat{C} \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}$ for some constant $\widehat{C}$ and all $\varepsilon \in (0, 1]$ due to strong continuity. Exactly by the same arguments as in the proof of Proposition 4.6, we thus obtain for $t \in [0, T]$ for some fixed T

$$\mathbb{E}[\|\lambda_t\|_{Y^*}^2] \leq \widetilde{C}(\|\lambda_0\|_{Y^*}^2 + 1)(1 - \int_0^t R'(s), ds),$$

where R' denotes the resolvent of $-C_2 \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}$. Hence, $\mathbb{E}[\varrho(\lambda_t)] \leq C \varrho(\lambda_0)$ for $t \in [0, T]$. From this the desired uniform growth bound $\|P_t\|_{L(\mathcal{B}^e(\mathcal{E}))} \leq M \exp(\omega t)$ for some $M \geq 1$ and $\omega \in \mathbb{R}$ follows.

For the set D as of Theorem 3.2 we here choose Fourier basis elements of the form

$$(4.17) \quad f_y : \mathcal{E} \rightarrow [0, 1]; \lambda \mapsto \exp(\langle y, \lambda \rangle)$$

such that $y \in \mathcal{E}_*$ and $\lambda \mapsto \exp(\langle y, \lambda \rangle)$ lies in $\cap_{n \geq 1} \text{dom}(A^n)$, whose span is of course dense, whence (i) of Theorem 3.2. We now equip $\text{span}(D)$ with the uniform norm $\|\cdot\|_\infty$ and verify Condition (ii), i.e. we check

$$(4.18) \quad \|A^n P_u^m f_y - A^m P_u^m f_y\|_\varrho \leq \|f_y\|_\infty a_{nm}$$

for all $0 \leq u \leq t$ with $a_{nm} \rightarrow 0$ as $n, m \rightarrow \infty$, and possibly depending on y and t . Note that

$$A^n f_y(\lambda) = \langle R^n(y), \lambda \rangle f_y(\lambda),$$

where R^n corresponds to (4.12) for $\varepsilon = \frac{1}{n}$. As P^n leaves D invariant for all $n \in \mathbb{N}$ by Proposition 4.14 (iv), we have

$$\begin{aligned} & \frac{|A^n P_u^m f_y(\lambda) - A^m P_u^m f_y(\lambda)|}{\varrho(\lambda)} \\ &= \frac{f_{y_u^m}(\lambda) \langle g, \lambda \rangle}{\varrho(\lambda)} \int_{\mathbb{R}_+} \underbrace{\exp(\langle y_u^m, \mathcal{S}_{\frac{1}{m}}^* \nu \xi \rangle)}_{\leq 1} \underbrace{|\exp(\langle y_u^m, (\mathcal{S}_{\frac{1}{n}}^* - \mathcal{S}_{\frac{1}{m}}^* \nu) \xi \rangle) - 1|}_{\tilde{a}_{nm}} \mu(d\xi) \end{aligned}$$

where y_u^m denotes the solution of $\partial_t y_u^m = R^m(y_u^m)$ at time u with $y_0 = y$ and $\tilde{a}_{nm}$ is chosen uniformly for all $u \leq t$ and tends to 0 as $n, m \rightarrow \infty$. This is possible since for the chosen initial values y we obtain that $\|y_u^m\|_Z$ is bounded on compact intervals in time uniformly in m also with respect to the Z norm. Indeed

$$\begin{aligned} \|y_t\|_Z &= \|\mathcal{S}_t^w y_0 + \int_0^t \mathcal{S}_{t-s}^w g \left(\int (\exp(\langle y_s, \mathcal{S}_\varepsilon^* \nu \xi \rangle) - 1) \mu(d\xi) \right) ds\|_Z \\ &\leq \|\mathcal{S}_t^w y_0\|_Z + \int_0^t \|\mathcal{S}_{t-s}^w g\|_Z \left(\int |\exp(\langle y_s, \mathcal{S}_\varepsilon^* \nu \xi \rangle) - 1| \mu(dx) \right) ds \\ &\leq \|\mathcal{S}_t^w y_0\|_Z + \mu(\mathbb{R}_+) \int_0^t \|\mathcal{S}_u^w g\|_Z du, \end{aligned}$$

where $\mathcal{S}^w$ is defined in the proof of Proposition 4.14. From this we infer (4.18) with $a_{nm} = C\tilde{a}_{nm}$ for some constant C . The conditions of Theorem 3.2 are therefore satisfied and we obtain a generalized Feller semigroup whose generator corresponds to (4.11) with $\varepsilon = 0$.

Concerning the second assertion, the affine transform formula follows simply from the convergence of the semigroups P^n as asserted in Theorem 3.2 by setting $y_t = \lim_{\varepsilon \rightarrow 0} y_t^\varepsilon$, where y_t^ε solves $\partial_t y_t^\varepsilon = R^\varepsilon(y_t^\varepsilon)$ in the mild sense with R^ε given by (4.12). Since $\exp(\langle y_t, \lambda \rangle)$ is then also the unique solution of the abstract Cauchy problem for initial value $\exp(\langle y_0, \lambda \rangle)$, i.e. it solves

$$\partial_t u(t, \lambda) = Au(t, \lambda), \quad u(0, \lambda) = \exp(\langle y_0, \lambda \rangle),$$

where A denotes the generator associated to (4.11) with $\varepsilon = 0$, we infer that y_t satisfies $\partial_t y_t = R(y_t)$ with R given by (4.12) for $\varepsilon = 0$. This is because $A \exp(\langle y_t, \lambda \rangle) = \exp(\langle y_t, \lambda \rangle) R(y_t)$. $\square$

We are now ready to state our main theorem, namely an existence and uniqueness result for equations of the type

$$(4.19) \quad d\lambda_t = \mathcal{A}^* \lambda_t dt + \nu dX_t,$$

where $(X_t)_{t \geq 0}$ is an Itô semimartingale of the form

$$(4.20) \quad X_t = \int_0^t \beta \langle g, \lambda_s \rangle ds + \int_0^t \sigma \sqrt{\langle g, \lambda_t \rangle} dB_t + \int_0^t \int \xi(\mu^X(d\xi, ds) - \langle g, \lambda_s \rangle m(d\xi) ds),$$

for some Brownian motion B and random jump measure μ^X with $\beta, \sigma \in \mathbb{R}$ and $m(d\xi)$ is a Lévy measure on $\mathbb{R}_+$ admitting a second moment.

Theorem 4.17. *Let Assumptions 4.9 and 4.5 be in force.*

- (i) *Then the stochastic partial differential equation (4.19) admits a unique Markovian solution $(\lambda_t)_{t \geq 0}$ with values in $\mathcal{E}$ given by a generalized Feller semigroup on $\mathcal{B}^e(\mathcal{E})$ whose generator takes on the set of Fourier elements*

$$f_y : \mathcal{E} \rightarrow [0, 1]; \lambda \mapsto \exp(\langle y, \lambda \rangle)$$

for $y \in \mathcal{D} \cap \mathcal{E}_*$ where $\mathcal{D}$ is defined in (4.10) the form

$$(4.21) \quad Af_y(\lambda) = f_y(\lambda)(\langle \mathcal{A}y, \lambda \rangle + \mathcal{R}(\langle y, \nu \rangle)\langle g, \lambda \rangle).$$

Here, $\mathcal{R} : \mathbb{R}_- \rightarrow \mathbb{R}$ is given by

$$(4.22) \quad \mathcal{R}(u) = \beta u + \frac{1}{2}\sigma^2 u^2 + \int_{\mathbb{R}_+} (\exp(u\xi) - 1 - u\xi) m(d\xi).$$

- (ii) This generalized Feller process allows to construct a probabilistically weak and analytically mild solution of (4.19), i.e. for $y \in Y$

$$\langle y, \lambda_t \rangle = \langle y, \mathcal{S}_t^* \lambda_0 \rangle + \int_0^t \langle y, \mathcal{S}_{t-s}^* \nu \rangle dX_s,$$

in particular for every initial value $\lambda_0 \in \mathcal{E}$ the semimartingale X can be constructed on an appropriate probabilistic basis. The stochastic integral is only defined in the weak sense, that is after pairing with $y \in Y$, a solution concept which is sometimes denoted by weakly mild. For every $y \in Y$ there exists a càg version of the real valued process $(\langle y, \lambda_t \rangle)_{t \geq 0}$.

- (iii) The affine transform formula is satisfied, i.e.

$$\mathbb{E}_{\lambda_0} [\exp(\langle y_0, \lambda_t \rangle)] = \exp(\langle y_t, \lambda_0 \rangle),$$

where y_t solves $\partial_t y_t = R(y_t)$ for all $y_0 \in \mathcal{E}_*$ and $t > 0$ in the mild sense with $R : \mathcal{D} \cap \mathcal{E}_* \rightarrow \mathbb{R}$ given by

$$(4.23) \quad R(y) = \mathcal{A}y + \mathcal{R}(\langle y, \nu \rangle)g$$

and $\mathcal{R}$ defined in (4.22). Furthermore, $y_t \in \mathcal{E}_*$ for all $t \geq 0$.

- (iv) For all $\lambda_0 \in \mathcal{E}$, the corresponding jump diffusion stochastic Volterra equation, $V_t := \langle g, \lambda_t \rangle$, given by

$$(4.24) \quad V_t = \langle g, \lambda_t \rangle = \langle g, \mathcal{S}_t^* \lambda_0 \rangle + \int_0^t \langle g, \mathcal{S}_{t-s}^* \nu \rangle dX_s = h(t) + \int_0^t K(t-s) dX_s$$

with $h(t) = \langle g, \mathcal{S}_t^* \lambda_0 \rangle$ admits a probabilistically weak solution with càg trajectories.

- (v) The Laplace transform of the Volterra equation V_t is given by

$$(4.25) \quad \mathbb{E}_{\lambda_0} [\exp(uV_t)] = \exp(uh(t) + \int_0^t h(t-s)\mathcal{R}(\psi_s)ds),$$

where $h(t) = \langle g, \mathcal{S}_t^* \lambda_0 \rangle$ and $\psi_t := \langle y_t, \nu \rangle$ solves the Riccati Volterra equation

$$\psi_t = uK(t) + \int_0^t K(t-s)\mathcal{R}(\psi_s)ds, \quad t > 0.$$

Hence the solution of the stochastic Volterra equation in (4.24) is unique in law.

Remark 4.18. (i) Observe that (4.24) implies that the function $h(t)$ defined by $h(t) = \langle g, \mathcal{S}_t^* \lambda_0 \rangle$ is given by the expectation

$$h(t) = \mathbb{E}[V_t - \beta \int_0^t K(t-s)V_s ds].$$

In the diffusion case, (4.25) therefore coincides with the formula obtained in [3]. From the Markov property of $(\lambda_t)_{t \geq 0}$, it is easy to see that the conditional Laplace transform of V_t reads as

$$\begin{aligned} \mathbb{E}_{\lambda_0} [\exp(uV_t) | \mathcal{F}_s] &= \exp(\langle y_{t-s}, \lambda_s \rangle) \\ &= \exp(u\langle g, \mathcal{S}_{t-s}^* \lambda_s \rangle + \int_0^{t-s} \langle g, \mathcal{S}_{t-s-r}^* \lambda_u \rangle \mathcal{R}(\langle y_r, \nu \rangle) dr), \end{aligned}$$

where y_t is the solution of $\partial_t y_t = R(y_t)$ with R given by (4.23) and $y_0 = ug$. Here, $\langle g, \mathcal{S}_x^* \lambda_s \rangle$ corresponds to the following conditional expectation

$$\langle g, \mathcal{S}_x^* \lambda_s \rangle = \mathbb{E} \left[V_{s+x} - \beta \int_0^x K(x-r) V_{s+r} dr \mid \mathcal{F}_s \right], \quad x \geq 0.$$

The function-valued process $(x \mapsto \langle g, \mathcal{S}_x^* \lambda_t \rangle)_{t \geq 0}$ corresponds also to a Markovian lift, described in detail in Section 5.2. Note that assuming existence of V , (4.25) and also the conditional Laplace transform of V is completely independent of the kernel representation.

- (ii) Mimicking the proof of assertion (v) of the above theorem, one can show that

$$\mathbb{E}_{\lambda_0} [\exp(\langle y_0, \lambda_t \rangle)] = \exp(\langle y_t, \lambda_0 \rangle) = \exp(\langle y_0, \mathcal{S}_t^* \lambda_0 \rangle + \int_0^t h(t-s) \mathcal{R}(\tilde{\psi}_s) ds),$$

where $\tilde{\psi}_t$ solves

$$\tilde{\psi}_t = \langle y_0, \mathcal{S}_t^* \nu \rangle + \int_0^t K(t-s) \mathcal{R}(\tilde{\psi}_s) ds.$$

Proof. To prove (i), we proceed again by applying Theorem 3.2. To this end consider the following sequence of SPDEs

$$(4.26) \quad \begin{aligned} d\lambda_t^n &= \mathcal{A}^* \lambda_t^n dt + \beta \nu \langle g, \lambda_t^n \rangle dt - n \sigma^2 \nu \langle g, \lambda_t^n \rangle dt + \nu dN_{1,t}^n \\ &\quad - \int_{\{\xi > \frac{1}{n}\}} \xi m(d\xi) \nu \langle g, \lambda_t^n \rangle dt + \nu dN_{2,t}^n, \end{aligned}$$

where $N_{1,t}^n$ is a jump process that jumps by $1/n$ and intensity $\sigma^2 n^2 \langle g, \lambda_t^n \rangle$, i.e. its compensator is given by $\sigma^2 n^2 \langle g, \lambda_t^n \rangle \delta_{\frac{1}{n}}(d\xi)$. The compensator of $N_{2,t}^n$ is given by $1_{\{\xi > \frac{1}{n}\}} m(d\xi) \langle g, \lambda_t^n \rangle$. In terms of (4.11), this corresponds to $\varepsilon = 0$, $N = N^n = N_1^n + N_2^n$, $w = -\beta + n\sigma^2 + \int_{\{\xi > \frac{1}{n}\}} \xi m(d\xi)$ and $\mu^n(d\xi) := \sigma^2 n^2 \delta_{\frac{1}{n}}(d\xi) + 1_{\{\xi > \frac{1}{n}\}} m(d\xi)$. From Proposition 4.15 we thus know that these SPDEs admit a solution in terms of generalized Feller semigroups P^n with associated generators A^n . Let us now establish a uniform growth bound for this sequence. Due to Proposition 4.15 we can establish the following estimate for $t \in [0, T]$ for some fixed $T > 0$

$$\begin{aligned} \mathbb{E}[\|\lambda_t^n\|_{Y^*}^2] &\leq 3\|\mathcal{S}_t^* \lambda_0\|_{Y^*}^2 + 3t \int_0^t \beta^2 \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \|g\|_Y^2 \mathbb{E}[\|\lambda_s^n\|_{Y^*}^2] ds \\ &\quad + 3\mathbb{E}[\|\int_0^t \mathcal{S}_{t-s}^* \nu dN_s^n - \int_0^t \int \mathcal{S}_{t-s}^* \nu \xi \mu^n(d\xi) \langle g, \lambda_s^n \rangle ds\|^2] \\ &\leq 3C_0 \|\lambda_0\|_{Y^*}^2 + 3t \int_0^t \beta^2 \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \|g\|_Y^2 \mathbb{E}[\|\lambda_s^n\|_{Y^*}^2] ds \\ &\quad + 3\mathbb{E}[\int \xi^2 \mu^n(d\xi) \int_0^t \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \langle g, \lambda_s^n \rangle ds] \\ &\leq 3C_0 \|\lambda_0\|_{Y^*}^2 + 3(\sigma^2 + \int \xi^2 m(d\xi)) \int_0^t \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 ds \\ &\quad + 3\|g\|_Y^2 (\beta^2 t + \sigma^2 + \int \xi^2 m(d\xi)) \int_0^t \|\mathcal{S}_{t-s}^* \nu\|_{Y^*}^2 \mathbb{E}[\|\lambda_s^n\|_{Y^*}^2] ds. \end{aligned}$$

Exactly as in Proposition 4.15, this now yields the uniform growth bound. For the set D as of Theorem 3.2 we consider again the Fourier basis elements as in (4.17), whose span is of course dense, whence (i). We equip $\text{span}(D)$ again with the uniform norm $\|\cdot\|_\infty$ and obtain a contraction property there. To verify Condition

(ii) of Theorem 3.2, we check (4.18) in the present setting. Note that A^n takes the form

$$A^n f_y(\lambda) = f_y(\lambda)(\langle \mathcal{A}y, \lambda \rangle + \beta \langle g, \lambda \rangle \langle y, \nu \rangle + \langle g, \lambda \rangle \int (\exp(\langle y, \nu \xi \rangle) - 1 - \langle y, \nu \xi \rangle) \mu^n(d\xi))$$

on D and from the last assertion of Proposition 4.15 we know that P^n leaves D invariant for all $n \in \mathbb{N}$. By this invariance it thus suffices to check

$$(4.27) \quad \|A^n f_y - A^m f_y\|_{\mathcal{Q}} \leq \|f_y\|_{\infty} a_{nm}$$

for $y \in K$, where K is a bounded set with respect to the Z norm. To this end consider for f_y for $y \in K$,

$$\begin{aligned} & \frac{|A^n f_y(\lambda) - A^m f_y(\lambda)|}{\varrho(\lambda)} \\ &= \frac{f_y(\lambda) \langle g, \lambda \rangle}{\varrho(\lambda)} \left| \int (\exp(\langle y, \nu \xi \rangle) - 1 - \langle y, \nu \xi \rangle) (\mu^n(d\xi) - \mu^m(d\xi)) \right| \\ &= \frac{f_y(\lambda) \langle g, \lambda \rangle}{\varrho(\lambda)} \left| \int (\xi^2 \langle y, \nu \rangle^2 + o(\langle y, \nu \rangle^2 \xi^2)) (\mu^n(d\xi) - \mu^m(d\xi)) \right| \\ &= \frac{f_y(\lambda) \langle g, \lambda \rangle}{\varrho(\lambda)} \underbrace{\left(\left| \int (\xi^2 \langle y, \nu \rangle^2 + o(\langle y, \nu \rangle^2 \xi^2)) \sigma^2 (n^2 \delta_{\frac{1}{n}}(d\xi) - m^2 \delta_{\frac{1}{m}}(d\xi)) \right| \right)}_{a_{nm}^1} \\ & \quad + \underbrace{\left| \int (\xi^2 \langle y, \nu \rangle^2 + o(\langle y, \nu \rangle^2 \xi^2)) (1_{\{\xi \geq \frac{1}{n}\}} - 1_{\{\xi \geq \frac{1}{m}\}}) m(d\xi) \right|}_{a_{nm}^2}. \end{aligned}$$

From this we infer (4.27), where $a_{nm} = C(a_{nm}^1 + a_{nm}^2)$ for some constant C , for all $y \in K$. The conditions of Theorem 3.2 are therefore satisfied and we obtain a generalized Feller semigroup whose generator is given by (4.21).

Concerning the second assumption, let us proceed similarly as in the proof of Proposition 4.14 (ii). Take $y \in \mathcal{D}$. Consider the local martingale

$$(4.28) \quad M_t^y = \langle y, \lambda_t \rangle - \langle y, \lambda_0 \rangle - \int_0^t \langle \mathcal{A}y, \lambda_s \rangle - \beta \langle y, \nu \rangle \langle g, \lambda_s \rangle ds.$$

Let now y be such that $\langle y, \nu \rangle = 1$ and denote M^y for such y simply by M . By the carré du champs formula the predictable quadratic variation¹ of M is given by

$$\langle M, M \rangle_t = \int_0^t (A f^2(\lambda_s) - 2f(\lambda_s) A f(\lambda_s)) ds = \left(\sigma^2 + \int \xi^2 m(d\xi) \right) \int_0^t \langle g, \lambda_s \rangle ds$$

for $f(\lambda) = \langle y, \lambda \rangle$ and all $\langle y, \nu \rangle = 1$, which shows in particular that M is independent of the particular representative y . Moreover, due to (4.26)

$$\sum_{s \leq t} (\Delta M_s)^2 = \lim_{n \rightarrow \infty} \left(\frac{1}{n^2} K_1^n + \sum_{s \leq t} (\Delta N_{2,s}^n)^2 \right) = \lim_{n \rightarrow \infty} \sum_{s \leq t} (\Delta N_{2,s}^n)^2$$

where K_1^n denotes the number of jumps of N_1^n up to time t . Hence, the compensator of $\sum_{s \leq t} (\Delta M_s)^2$ is $\int_0^t \int \xi^2 m(d\xi) \langle g, \lambda_s \rangle ds$ and we deduce by the unique decomposition of M into a continuous and purely discontinuous local martingale M^c and M^d , that $\langle M^c, M^c \rangle = \sigma^2 \int_0^t \langle g, \lambda_s \rangle ds$. Therefore M_t can be written as

$$M_t = M_t^c + M_t^d = \int \sigma \langle g, \lambda_s \rangle dB_s + \int_0^t \int \xi (\mu^X(d\xi, ds) - \langle g, \lambda_s \rangle m(d\xi) ds),$$

¹This is also denoted by angle brackets.

for some Brownian motion B and random measure $\mu^X(d\xi, ds)$. Define now X as

$$X_t = \int_0^t \beta \langle g, \lambda_s \rangle ds + M_t.$$

Then X is of form (4.20). Furthermore note that M^y of (4.28) for $y \in \mathcal{D}$ is given by $M^y = \langle y, \nu \rangle M$ which can be seen again from the carré du champs formula, yielding $\langle M^y, M^y \rangle = \langle y, \nu \rangle^2 \langle M, M \rangle$. We thus deduce from (4.28)

$$\langle y, \lambda_t \rangle = \langle y, \lambda_0 \rangle + \int_0^t \langle \mathcal{A}y, \lambda_s \rangle ds + \langle y, \nu \rangle X_t$$

for $y \in \mathcal{D}$.

To pass from this weak formulation to the mild one, notice the following uniqueness assertion: fix the driving process X with càdlàg trajectories and $(\varphi_t(y))_{t \geq 0}$, for $y \in Y$, be another solution process with càg trajectories depending linearly on y . We assume that

$$(4.29) \quad \varphi_t(y) = \langle y, \lambda_0 \rangle + \int_0^t \varphi_s(\mathcal{A}y) ds + \langle y, \nu \rangle X_t$$

for all $y \in \mathcal{D}$ and furthermore for every sequence $y_n \rightarrow y$ in Y we can choose a set of probability one where for every $s \geq 0$

$$(4.30) \quad \varphi_s(y_n) \rightarrow \varphi_s(y)$$

as $n \rightarrow \infty$ holds true. Note that the set will depend on the chosen sequence. Then the announced uniqueness assertion means that necessarily $\varphi_t(y) = \langle y, \lambda_t \rangle$ for all $y \in \mathcal{D}$. Indeed, the difference $g_t(y) = \varphi_t(y) - \langle y, \lambda_t \rangle$ satisfies

$$g_t(y) = \int_0^t g_s(\mathcal{A}y) ds$$

with càg trajectories. Therefore clearly $s \mapsto g_{t-s}(S_s y)$ is constant simply by taking the derivative with respect to s and collecting terms for $y \in \mathcal{D}$, hence $0 = g_0(S_t y) = g_t(y) = \varphi_t(y) - \langle y, \lambda_t \rangle$.

The mild formulation then follows from this uniqueness result immediately. Indeed, define processes

$$f_t(y) := \langle S_t y, \lambda_0 \rangle + \int_0^t \langle S_{t-s} y, \nu \rangle dX_s,$$

for all $y \in Y$ and $t \geq 0$. For $y \in \mathcal{D}$ we then have

$$\begin{aligned} & \langle y, \lambda_0 \rangle + \int_0^t f_s(\mathcal{A}y) ds + \langle y, \nu \rangle X_t \\ &= \langle y, \lambda_0 \rangle + \int_0^t \langle \mathcal{A}S_s y, \lambda_0 \rangle ds + \int_0^t \int_0^s \langle \mathcal{A}S_{s-u} y, \nu \rangle dX_u ds + \langle y, \nu \rangle X_t \\ &= \langle S_t y, \lambda_0 \rangle + \int_0^t \int_u^t \langle \mathcal{A}S_{s-u} y, \nu \rangle ds dX_u + \langle y, \nu \rangle X_t \\ &= \langle S_t y, \lambda_0 \rangle + \int_0^t \langle S_{t-u} y - y, \nu \rangle dX_u + \langle y, \nu \rangle X_t \\ &= \langle S_t y, \lambda_0 \rangle + \int_0^t \langle S_{t-u} y, \nu \rangle dX_u \\ &= f_t(y), \end{aligned}$$

whence $(f_t(y))_{t \geq 0}$ satisfies (4.29). Since $(f_t(y))$ is linear in y and satisfies by basic dominated convergence for stochastic integrals the continuity assumption (4.30), the previous uniqueness result yields $f_t(y) = \langle y, \lambda_t \rangle$ for all $t \geq 0$ almost surely.

The third assertion can be shown exactly along the lines of the proof of the second assertion of Proposition 4.15.

The fourth assertion is a simple consequence of (ii) with $y = g$.

Finally to prove (v), set in (iii) $y_0 = ug$. Then, the mild solution of y_t is given by

$$(4.31) \quad y_t = u\mathcal{S}_t g + \int_0^t \mathcal{S}_{t-s} g \mathcal{R}(\langle y_s, \nu \rangle) ds.$$

Hence, by (iii) and Fubini's theorem

$$\begin{aligned} \mathbb{E}_{\lambda_0}[\exp(uV_t)] &= \mathbb{E}_{\lambda_0}[\exp(u\langle g, \lambda_t \rangle)] = \exp(\langle y_t, \lambda_0 \rangle) \\ &= \exp(u\langle g, \mathcal{S}_t^* \lambda_0 \rangle + \int_0^t \langle g, \mathcal{S}_{t-s}^* \lambda_0 \rangle \mathcal{R}(\langle y_s, \nu \rangle) ds). \end{aligned}$$

Setting now $\psi_t := \langle y_t, \nu \rangle$ for $t > 0$, we see from (4.31)

$$\psi_t = u\langle g, \mathcal{S}_t^* \nu \rangle + \int_0^t \langle g, \mathcal{S}_{t-s}^* \nu \rangle \mathcal{R}(\psi_s) ds = uK(t) + \int_0^t K(t-s) \mathcal{R}(\psi_s) ds,$$

and in particular that it is well-defined for $t > 0$. This proves the assertion of uniqueness in law, too. $\square$

5. CONCRETE SPECIFICATIONS

The goal of this section is to concretely specify the abstract setting introduced in Section 4 when we deal with specific kernels of form (4.3).

5.1. Lifting rough Volterra processes to measures. As already outlined in the introduction, we here consider the case of kernels given as Laplace transforms of some (signed) measure ν on $\mathbb{R}_+$, i.e.

$$K(t) = \int_0^\infty e^{-xt} \nu(dx),$$

such that $K(t) < \infty$ for all $t > 0$ and $K \in L_{\text{loc}}^2(\mathbb{R}_+, \mathbb{R})$. If ν is a nonnegative measure, we are in the important setting of *completely monotone kernels*, into which for instance the fractional kernels appearing in rough volatility models of the form $K(t) = \Gamma(\alpha)^{-1} t^{\alpha-1}$ for $\alpha \in (\frac{1}{2}, 1)$ fall.

To cast this into the framework considered in Section 4, in particular Assumption 4.2, let Y be the space of bounded continuous functions on the extended real half-line $\overline{\mathbb{R}}_+ := \mathbb{R}_+ \cup \{\infty\}$. We here compactify $\mathbb{R}_+$, in order to make $C_b(\overline{\mathbb{R}}_+, \mathbb{R})$ separable. Its dual space Y^* is the space of the finite (signed) regular Borel measures on the extended real half-line $\overline{\mathbb{R}}_+$. As above we consider the weak- $*$ -topology on Y^* and a weight function

$$\varrho(\lambda) = 1 + |\lambda|^2(\overline{\mathbb{R}}_+),$$

where $|\lambda|$ denotes the total variation norm of λ . Moreover, Z is the space of functions $g \in Y$ such that

$$(x \mapsto xg(x)) \in Y$$

together with the operator norm on it, i.e. $\|g\| = \sqrt{\|g\|^2 + \sup_{x \geq 0} |xg(x)|^2}$ for $g \in Z$. Its dual Z^* is the space of regular Borel measures ν on $\overline{\mathbb{R}}_+$ that satisfy

$$\int_0^\infty \left(\frac{1}{x} \wedge 1\right) \nu(dx) < \infty.$$

As semigroup $(\mathcal{S}_t^*)_{t \geq 0}$ acting on Z^* and Y^* we consider the multiplication semigroup

$$\mathcal{S}_t^* \lambda = e^{-t \cdot} \lambda$$

which leaves Y^* and Z^* invariant and acts in a strongly continuous way thereon with respect to the strong norms. Moreover, $\lambda \mapsto \mathcal{S}_t^* \lambda$ is weak- $*$ -continuous on Y^* and on Z^* for every $t \geq 0$ (considering the weak- $*$ -topology on both the domain and the image space). Whence, as stated in Remark 4.4, this semigroup then satisfies all requirements of Assumption 4.2 (iv) to (vi).

In terms of (4.3), K can now be written as

$$K(t) = \int_0^\infty e^{-xt} \nu(dx) = \langle 1, \mathcal{S}_t^* \nu \rangle,$$

where the pairing is here $\langle y, \lambda \rangle = \int_0^\infty y(x) \lambda(dx)$. The element $g \in Y$ appearing in (4.3) is here simply the constant function 1. Note also that the requirements of Assumption 4.5 are satisfied as well. Indeed $\mathcal{S}_t^* \nu \in Y^*$ for all $t > 0$ and

$$\int_0^t \|\mathcal{S}_s^* \nu\|_{Y^*}^2 ds = \int_0^t \left(\int_0^\infty e^{-xs} \nu(dx) \right)^2 ds = \int_0^t K(s)^2 ds < \infty$$

for all $t > 0$, since K is assumed to lie in $L_{\text{loc}}^2(\mathbb{R}_+, \mathbb{R})$.

As in (4.7), we define for fixed $w > 0$

(5.1)

$$\begin{aligned} \mathcal{E}^w &:= \{ \lambda_0 \in Y^* \mid \langle 1, \mathcal{S}_t^* \lambda_0 \rangle - \int_0^t R^w(t-s) \langle 1, \mathcal{S}_s^* \lambda_0 \rangle ds \geq 0 \text{ for all } t \geq 0 \} \\ &= \{ \lambda_0 \in Y^* \mid \bar{\lambda}_t \geq 0 \text{ with} \end{aligned}$$

$$\lambda_t(dx) = e^{-xt} \lambda_0(dx) - w \int_0^t e^{-(t-s)x} \nu(dx) \bar{\lambda}_s ds \text{ for all } t \geq 0 \},$$

where R^w denotes the resolvent of $wK(t)$ and $\bar{\lambda} = \langle 1, \lambda \rangle$ the total mass of λ .

Remark 5.1. Notice that the cone $\mathcal{E}^w$ is *not* locally compact with respect to the weak- $*$ -topology. Indeed, consider the simplest case $K = 1$ and denote the Laplace of some signed measure λ_0 by

$$h(t) = \int_0^\infty \exp(-xt) \lambda_0(dx).$$

Then λ_0 lies in $\mathcal{E}^w$ if $h(t) \geq 0$ and $h'(t) \geq 0$ for all $t \geq 0$. Choose a positive measure λ_0 whose support does *not* contain $[1, 2]$. Then the signed measures $\lambda_n = \lambda_0(dx) + 1_{[1,2]} \frac{1}{x} \sin(n\pi x) dx \in \mathcal{E}^w$, for $n \geq 1$, and $\lambda_n \rightarrow \lambda_0$ in the weak- $*$ -topology. On the other hand the total variation norm of λ_n does *not* converge to the total variation norm of λ_0 . Hence, $\lambda \mapsto 1 + |\lambda|^2(\bar{\mathbb{R}}_+)$ is not continuous and by Remark 2.2 the cone $\mathcal{E}^w$ is not locally compact.

In contrast, note that the cone of non-negative Borel measures on $\bar{\mathbb{R}}_+$ (or generally on a compactum) is locally compact in the weak- $*$ -topology. In this case the total variation norm of a non-negative measures is nothing but the total mass, i.e. a linear functional on it, thus continuous. The one-point compactification in this case is called Watanabe topology, see [12].

Assumption 4.9 now reads as follows:

Assumption 5.2. *Let ν be such that $\exp(-t.)\nu \in \mathcal{E}^w$ for all $t > 0$ and for all $w > 0$.*

Remark 5.3. As stated in Remark 4.10, Assumption 5.2 is satisfied if $R^w \geq 0$ for all $w > 0$ and $K \geq 0$, which is satisfied if K is completely monotone.

The state space that we consider for the stochastic processes in the sequel is as of (4.12), namely

$$\mathcal{E} = \cap_{w>0} \mathcal{E}^w.$$

Remark 5.4. Comparing $\mathcal{E}$ with the state spaces considered in [1] (e.g. Equation (4.6)), we see that the conditions there translate to

$$\{\lambda_0 \in Y^* \mid t \mapsto \langle 1, \mathcal{S}_t^* \lambda_0 \rangle = \int_0^\infty e^{-tx} \lambda_0(dx) \in \mathcal{G}_K\},$$

where $\mathcal{G}_K$ is defined in [1, Equation (2.5)]. From [1, Theorem A.2] we see that this describes a subspace of $\mathcal{E}$ in case a resolvent of the first kind exists for the kernel K .

It is interesting to consider the case when ν is a measure with finitely many atoms: we identify this measure setting then with some $\mathbb{R}^N$ and denote $\text{supp}(\nu) = \{0 \leq x_1 < \dots < x_N\}$.

Proposition 5.5. *Under Assumption 5.2 the state space $\mathcal{E}$ is a well defined closed cone given as the intersection of $\cap_{w>0} \mathcal{E}^w$ where for $w > 0$*

$$\mathcal{E}^w = \{\lambda_0 \in \mathbb{R}^N \mid \langle e^{tA^w} \lambda_0, \mathbf{1} \rangle_{\mathbb{R}^N} \geq 0 \text{ for all } t \geq 0\}$$

with $A^w = \text{diag}(-x_1, \dots, -x_N) - w(\nu(x_1)\mathbf{1}, \dots, \nu(x_N)\mathbf{1})^\top$. Here, $\mathbf{1} \in \mathbb{R}^N$ is the (column) vector consisting of ones.

Proof. Note that the deterministic equation as of (4.2) has here the following form

$$d\lambda_t(x_i) = -x_i \lambda_t(x_i) dt - w \nu(x_i) \sum_{j=1}^N \lambda_t(x_j) dt, \quad i \in \{1, \dots, N\}.$$

This can be written as $d\lambda_t = A^w \lambda_t dt$ so that the assertion follows from the definition of $\mathcal{E}^w$. $\square$

Example 5.6. There are simple two dimensional examples when $\mathcal{E}^0 \neq \mathcal{E}^w$ for $w > 0$: indeed take $x_0 = 0$ and $x_1 = 1$ and $\nu(dx) = \delta_0(dx) + \delta_1(dx)$. In this case

$$\mathcal{E}^0 = \text{cone}(e_2, e_1 - e_2)$$

whereas

$$\mathcal{E}^w = \text{cone}(e_1 + e_2, e_1 - e_2).$$

Here e_i , $i = 1, 2$ denotes the canonical basis vectors of $\mathbb{R}^2$. Observe here that $\mathcal{E}^w$ does not depend on w for $w > 0$, but it changes when $w \rightarrow 0$.

Let us finally come to the SPDE formulation which will in particular lead to rough affine volatility models. Consider the following measure valued SPDE

$$(5.2) \quad d\lambda_t(dx) = -x\lambda_t(dx)dt + \nu(dx)dX_t,$$

where $(X_t)_{t \geq 0}$ is an Itô semimartingale of the form

$$(5.3) \quad X_t = \int_0^t \beta \bar{\lambda}_s ds + \int_0^t \sigma \sqrt{\bar{\lambda}_t} dB_t + \int_0^t \int \xi(\mu^X(d\xi, ds) - \bar{\lambda}_s m(d\xi) ds),$$

for some Brownian motion B and random jump measure μ^X with $\beta, \sigma \in \mathbb{R}$ and $m(d\xi)$ is a Lévy measure on $\mathbb{R}_+$ admitting a second moment. Recall here that $\bar{\lambda} = \langle 1, \lambda \rangle$. As a corollary of Theorem 4.17 we now obtain the following result.

Theorem 5.7. *Let Assumption 5.2 be in force.*

- (i) *Then the stochastic partial differential equation (5.2) admits a unique Markovian solution $(\lambda_t)_{t \geq 0}$ in $\mathcal{E}$ given by a generalized Feller semigroup on $\mathcal{B}^e(\mathcal{E})$ whose generator takes the form*

$$(5.4) \quad Af_y(\lambda) = f_y(\lambda) \left(\int_0^\infty -xy(x)\lambda(dx) + \mathcal{R}(\langle y, \nu \rangle) \bar{\lambda} \right)$$

on the set of Fourier basis elements f_y with $y \in \mathcal{D} \cap \mathcal{E}_*$, where $\mathcal{R} : \mathbb{R}_- \rightarrow \mathbb{R}$ is given by

$$(5.5) \quad \mathcal{R}(u) = \beta u + \frac{1}{2} \sigma^2 u^2 + \int_{\mathbb{R}_+} (\exp(u\xi) - 1 - u\xi) m(d\xi).$$

- (ii) This generalized Feller process allows to construct a probabilistically weak and analytically weakly mild solution of (5.2), i.e. for $y \in Y$

$$\langle y, \lambda_t \rangle = \langle e^{-t \cdot} y, \lambda_0 \rangle + \int_0^t \langle e^{-(t-s) \cdot} y, \nu \rangle dX_s,$$

in particular for every initial value the semimartingale X can be constructed on an appropriate probabilistic basis. For every $y \in Y$ there exists a càg version of the real valued process $(\langle y, \lambda_t \rangle)_{t \geq 0}$.

- (iii) The affine transform formula is satisfied, i.e.

$$\mathbb{E}_{\lambda_0} [\exp(\langle y_0, \lambda_t \rangle)] = \exp(\langle y_t, \lambda_0 \rangle),$$

where y_t solves $\partial_t y_t = R(y_t)$ for all $y_0 \in \mathcal{E}_*$ and $t > 0$ in the mild sense with $R : \mathcal{D} \cap \mathcal{E}_* \rightarrow \mathbb{R}$ given by

$$R(y)(x) = -xy(x) + \mathcal{R}(\langle y, \nu \rangle)$$

and $\mathcal{R}$ defined in (5.5). Furthermore, $y_t \in \mathcal{E}_*$ for all $t \geq 0$.

- (iv) For all $\lambda_0 \in \mathcal{E}$, the corresponding jump diffusion stochastic Volterra equation, $V_t = \bar{\lambda}_t = \langle 1, \lambda_t \rangle$, given by

$$(5.6) \quad \bar{\lambda}_t = \int_0^\infty e^{-tx} \lambda_0(dx) + \int_0^t K(t-s) dX_s$$

with X given by (5.3) admits a unique probabilistically weak solution in $\mathbb{R}_+$ with càg trajectories.

- (v) The Laplace transform of the Volterra equation V_t is given by

$$\mathbb{E}_{\lambda_0} [\exp(uV_t)] = \exp(uh(t) + \int_0^t h(t-s) \mathcal{R}(\psi_s) ds),$$

where $h(t) = \int_0^\infty e^{-xt} \lambda_0(dx)$ and $\psi_t := \langle y_t, \nu \rangle$ solves the Riccati Volterra equation

$$\psi_t = uK(t) + \int_0^t K(t-s) \mathcal{R}(\psi_s) ds, \quad t > 0.$$

Hence the solution of the stochastic Volterra equation in (5.6) is unique in law.

5.2. Lifting rough Volterra processes to forward curves. There are several ways to lift stochastic Volterra processes to Markovian processes: in the previous section the focus was on the completely monotone nature of many Volterra kernels. The lift that we treat here considers instead the stochastic Volterra process together with its conditional expectations from future values. As in the Heath-Jarrow-Morton (HJM) case, this yields shift semigroups. This Markovian lift also falls in the realm of Section 4 and can be considered from the perspective of generalized Feller semigroups. Indeed, here K (satisfying Conditions 5.8 below) is represented by

$$K(t) = \int_0^\infty \mathcal{S}_t^* K(x) \delta_0(dx),$$

where $\mathcal{S}_t^*$ denotes the shift semigroup. In order to meet all the conditions of Section 4, we here assume the following:

Assumption 5.8. (i) K is assumed to lie in $L_{loc}^2(\mathbb{R}_+, \mathbb{R})$.

- (ii) *there exists $L \in AC(\mathbb{R}_+, \mathbb{R})$, where $AC(\mathbb{R}_+, \mathbb{R})$ denotes the space of real-valued absolutely continuous functions on $\mathbb{R}_+$, such that $K = L'$.*
- (iii) *There exists some strictly positive weight function $v > 0$ such that*

$$\int_0^\infty |K(x)|^2 v(x) dx = \int_0^\infty |L'(x)|^2 v(x) dx < \infty.$$

In terms of the underlying Banach spaces we let Y and in turn also Y^* be Hilbert spaces, similarly as in Filipovic [20], namely

$$Y = \{y \in AC(\mathbb{R}_+, \mathbb{R}) \mid \int_0^\infty |y'(x)|^2 v(x) dx < \infty\}$$

for the specified strictly positive weight function $v > 0$. We endow Y with the scalar product

$$\langle y, \lambda \rangle_v = y(0)\lambda(0) + \int_0^\infty y'(x)\lambda'(x)v(x)dx$$

and denote the associated norm via $\|\cdot\|_v$.

Assumption 5.9. *We furthermore assume that the shift semigroup $\mathcal{S}^*$ acts in a strongly continuous way on Y^* , which can be easily expressed in terms of the function v , and that $\mathcal{S}_t^* K \in Y^*$ for $t > 0$.*

Remark 5.10. One can choose $v(x) = \exp(\alpha x)$ for $x \geq 0$, which proved to be useful in case of term structures of interest rates.

The weight function, in the sense of weighted spaces, is again given by

$$\varrho(\lambda) = 1 + \|\lambda\|_v^2.$$

Moreover, for Z we consider the subspace of functions vanishing at 0 and whose first (weak) derivative lies in Y (with the corresponding operator Hilbert space norm), then $K \in Z^*$ can be viewed as a weak derivative of a function $L \in Y^*$ with norm

$$\|K\|_{Z^*}^2 := \int_0^\infty |L'(x)|^2 v(x) dx.$$

In particular the value ∞ at 0 is possible. By condition (ii) and (iii) of Assumption 5.8, $K \in Z^*$. Notice also the following inclusions

$$Z \subset Y = Y^* \subset Z^*.$$

In terms of (4.3), $K \in Z^*$ can now be written as

$$K(t) = \langle \mathcal{S}_t^* K, 1 \rangle_v = (\mathcal{S}_t^* K)(0).$$

The element $g \in Y$ appearing in (4.3) is here again simply the constant function 1.

Note that due to Assumption 5.9 the shift semigroup $(\mathcal{S}_t^*)_{t \geq 0}$ acting on Z^* and Y^* satisfies all requirements of Assumption 4.2 (iv) to (v). Concerning (vi), the (pre)adjoint $(\mathcal{S}_t)_{t \geq 0}$ acts on Z and Y via the following formula obtained from adjoining with respect to the scalar product on Y

$$\mathcal{S}_t y(x) = y(0) + \int_0^{x \wedge t} y(0) \frac{1}{v(s)} ds + \int_t^{x \vee t} y'(s-t) \frac{v(s-t)}{v(s)} ds.$$

The generator A can be calculated accordingly by taking the derivative with respect to t . Notice that the pre-dual semigroup $\mathcal{S}$ acts in strongly continuous way on Y (as dual semigroup of a strongly continuous semigroup on a Hilbert space Y , see [18] for details) as well as on Z . Furthermore the requirements of Assumption 4.5 are met as well.

As in (4.7), we define for fixed $w > 0$

$$(5.7) \quad \begin{aligned} \mathcal{E}^w &:= \{\lambda_0 \in Y^* \mid \langle 1, \mathcal{S}_t^* \lambda_0 \rangle - \int_0^t R^w(t-s) \langle 1, \mathcal{S}_s^* \lambda_0 \rangle ds \geq 0 \text{ for all } t \geq 0\} \\ &= \{\lambda_0 \in Y^* \mid \lambda_t(0) \geq 0 \text{ with} \\ &\quad \lambda_t(x) = \lambda_0(x+t) - w \int_0^t K(t-s+x) \lambda_s(0) ds \text{ for all } t \geq 0\}, \end{aligned}$$

where R^w denotes the resolvent of $wK(t)$. Note that $\langle 1, \lambda \rangle = \lambda(0)$. Again the state space will be.

$$\mathcal{E} = \cap_{w>0} \mathcal{E}^w.$$

Assumption 4.9 now reads as follows:

Assumption 5.11. *Let K be such that $K(t + \cdot) \in \mathcal{E}^w$ for all $t > 0$ and for all $w > 0$.*

The SPDE that we consider in the present setting is

$$(5.8) \quad d\lambda_t(x) = \frac{d}{dx} \lambda_t(x) dt + K(x) dX_t,$$

where $(X_t)_{t \geq 0}$ is an Itô semimartingale of the form

$$(5.9) \quad X_t = \int_0^t \beta \lambda_s(0) ds + \int_0^t \sigma \sqrt{\lambda_s(0)} dB_t + \int_0^t \int \xi(\mu^X(d\xi, ds) - \lambda_s(0)m(d\xi)ds),$$

for some Brownian motion B and random jump measure μ^X with $\beta, \sigma \in \mathbb{R}$ and $m(d\xi)$ is a Lévy measure admitting a second moment. As a corollary of Theorem 4.17 we now obtain the following result.

Theorem 5.12. *Let Assumptions 5.8, 5.9 and 5.11 be in force.*

- (i) *Then the stochastic partial differential equation (5.8) admits a unique Markovian solution $(\lambda_t)_{t \geq 0}$ in $\mathcal{E}$ given by a generalized Feller semigroup on $\mathcal{B}^q(\mathcal{E})$ whose generator takes the form*

$$Af_y(\lambda) = f_y(\lambda)(\langle Ay, \lambda \rangle_v + \mathcal{R}(\langle y, K \rangle_v) \lambda(0))$$

on the set of Fourier basis elements f_y with $y \in \mathcal{D} \cap \mathcal{E}_$ where $\mathcal{R} : \mathbb{R}_- \rightarrow \mathbb{R}$ is given by*

$$\mathcal{R}(u) = \beta u + \frac{1}{2} \sigma^2 u^2 + \int_{\mathbb{R}_+} (\exp(u\xi) - 1 - u\xi) m(d\xi).$$

- (ii) *This generalized Feller process allows to construct a probabilistically weak and analytically weakly mild càg solution of (5.8), i.e. in particular the point evaluations (being linear functionals) satisfy*

$$\lambda_t(x) = \lambda_0(t+x) + \int_0^t K(t-s+x) dX_s,$$

are càg, and for every initial value the semimartingale X can be constructed on an appropriate probabilistic basis.

- (iii) *The affine transform formula is satisfied, i.e.*

$$\mathbb{E}_{\lambda_0} [\exp(\langle y_0, \lambda_t \rangle_v)] = \exp(\langle y_t, \lambda_0 \rangle_v),$$

where y_t solves $\partial_t y_t = R(y_t)$ for all $y_0 \in \mathcal{E}_$ and $t > 0$ in the mild sense with $R : \mathcal{D} \cap \mathcal{E}_* \rightarrow \mathbb{R}$ given by*

$$R(y)(x) = Ay(x) + \mathcal{R}(\langle y, \nu \rangle_v)$$

with $\mathcal{R}$ defined in (4.22). Furthermore, $y_t \in \mathcal{E}_$ for all $t \geq 0$.*

- (iv) For all $\lambda_0 \in \mathcal{E}$, the corresponding jump diffusion stochastic Volterra equation, i.e.

$$(5.10) \quad \lambda_t(0) = \lambda_0(t) + \int_0^t K(t-s) dX_s$$

admits a probabilistically weak solution in $\mathbb{R}_+$ with càg trajectories.

- (v) The Laplace transform of the Volterra equation V_t is given by

$$\mathbb{E}_{\lambda_0} [\exp(u\lambda_t(0))] = \exp(uh(t) + \int_0^t h(t-s)\mathcal{R}(\psi_s)ds),$$

where $h(t) = \lambda_0(t)$ and $\psi_t := \langle y_t, \nu \rangle$ solves the Riccati Volterra equation

$$\psi_t = uK(t) + \int_0^t K(t-s)\mathcal{R}(\psi_s)ds, \quad t > 0.$$

Hence the solution of the stochastic Volterra equation in (5.10) is unique in law.

REFERENCES

- [1] E. Abi Jaber and O. El Euch. Markovian structure of the Volterra Heston model. *Statistics & Probability Letters*, 149:63–72, 2019.
- [2] E. Abi Jaber and O. El Euch. Multifactor approximation of rough volatility models. *SIAM Journal on Financial Mathematics*, 10(2):309–349, 2019.
- [3] E. Abi Jaber, M. Larsson, and S. Pulido. Affine Volterra processes. *The Annals of Applied Probability*, to appear, 2019.
- [4] E. Alòs and Y. Yang. A closed-form option pricing approximation formula for a fractional Heston model, 2014.
- [5] C. D. Aliprantis and K. C. Border. *Infinite dimensional analysis*. Springer, Berlin, third edition, 2006. A hitchhiker’s guide.
- [6] V. S. Borkar. *Probability theory*. Universitext. Springer-Verlag, New York, 1995. An advanced course.
- [7] A. Boumezoued. Population viewpoint on Hawkes processes. *Advances in Applied Probability*, 48(2):463–480, 2016.
- [8] C. Cuchiero, D. Filipović, E. Mayerhofer and J. Teichmann. Affine processes on positive semidefinite matrices. *The Annals of Applied Probability*, 21(2): 397–463, 2011.
- [9] C. Cuchiero, M. Keller-Ressel, E. Mayerhofer and J. Teichmann. Affine processes on symmetric cones. *Journal of Theoretical Probability*, 29(2): 359–422, 2014.
- [10] C. Cuchiero and J. Teichmann. Markovian lifts of positive semidefinite affine Volterra type processes. *Decisions in Economics and Finance*, to appear, 2019.
- [11] R. C. Dalang. Extending the martingale measure stochastic integral with applications to spatially homogeneous SPDEs. *Electronic Journal of Probability*, 4(6), 1999.
- [12] D. A. Dawson and E. Perkins. *Superprocesses at Saint-Flour*. Probability at Saint-Flour. Springer, Heidelberg, 2012.
- [13] P. Dörsek and J. Teichmann. A Semigroup Point of View on Splitting Schemes for Stochastic (Partial) Differential Equations. *arXiv:1011.2651*, 2010.
- [14] D. Duffie, D. Filipović, and W. Schachermayer. Affine processes and applications in finance. *The Annals of Applied Probability*, 13(3):984–1053, 2003.
- [15] O. El Euch and M. Rosenbaum. The characteristic function of rough Heston models. *Mathematical Finance*, 29(1):3–38, 2019.
- [16] O. El Euch, M. Fukasawa, and M. Rosenbaum. The microstructural foundations of leverage effect and rough volatility. *Finance and Stochastics*, 22(2):241–280, 2018.
- [17] O. El Euch and M. Rosenbaum. Perfect hedging in rough Heston models. *The Annals of Applied Probability*, 28(6):3813–3856, 2018.
- [18] K.-J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*, volume 194 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt.
- [19] S. N. Ethier and T. G. Kurtz. *Markov processes*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1986. Characterization and convergence.

- [20] D. Filipović. *Consistency Problems for Heath-Jarrow-Morton Interest Rate Models*, volume 1760 of *Lecture Notes in Mathematics*. Springer Verlag, Berlin, 2001.
- [21] J. Gatheral, T. Jaisson, and M. Rosenbaum. Volatility is rough. *Quantitative Finance*, 18(6):933-949, 2018.
- [22] G. Gripenberg, S.-O. Londen, and O. Staffans. *Volterra integral and functional equations*, volume 34 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1990.
- [23] A. G. Hawkes. Spectra of some self-exciting and mutually exciting point processes. *Biometrika*, 58(1):83-90, 1971.
- [24] T. Jaisson and M. Rosenbaum. Rough fractional diffusions as scaling limits of nearly unstable heavy tailed Hawkes processes. *The Annals of Applied Probability*., 26(5):2860-2882, 2016.
- [25] L. Mytnik and T. Salisbury. Uniqueness for Volterra-type stochastic integral equations. *arXiv:1502.05513*, 2015.
- [26] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.
- [27] H. H. Schaefer and M. P. Wolff. *Topological vector spaces*, volume 3 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1999.

VIENNA UNIVERSITY OF ECONOMICS AND BUSINESS, WELTHANDELSPLATZ 1, A-1020 VIENNA
AND ETH ZÜRICH, RÄMISTRASSE 101, CH-8092 ZÜRICH