Self-anisotropizing inflationary universe in Horndeski theory and beyond

Hiroaki W. H. Tahara,^{1,2} Sakine Nishi,³ Tsutomu Kobayashi,³ and Jun'ichi Yokoyama^{1,2,4}

¹Research Center for the Early Universe (RESCEU),

Graduate School of Science, The University of Tokyo, Tokyo 113-0033, Japan

²Department of Physics, Graduate School of Science,

The University of Tokyo, Tokyo 113-0033, Japan

³Department of Physics, Rikkyo University, Toshima, Tokyo 171-8501, Japan

⁴Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU),

WPI, UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8568, Japan

Abstract

As opposed to Wald's cosmic no-hair theorem in general relativity, it is shown that the Horndeski theory (and its generalization) admits anisotropic inflationary attractors if the Lagrangian depends cubically on the second derivatives of the scalar field. We dub such a solution as a self-anisotropizing inflationary universe because anisotropic inflation can occur without introducing any anisotropic matter fields such as a vector field. As a concrete example of self-anisotropization we present the dynamics of a Bianchi type-I universe in the Horndeski theory.

I. INTRODUCTION

Inflation in the early universe [1-4] solves a number of fundamental problems in cosmology, such as horizon, flatness, monopole and the origin-of-structure problems. Its basic predictions have been confirmed by a number of observations of the cosmic microwave background (CMB) and large-scale structures. The simplest single-field inflation paradigm fits, in a sense, observations too well, so that it is difficult to single out the correct field theory model of inflation. In this context, much work has been done in search for anomalies in observations. One possibility among them is statistical anisotropy of spectrum of primordial perturbations. If established, we need an inflation model to realize anisotropic expansion. In this regard it has been known that inflation driven by a scalar potential make the observable universe fully isotropic [5, 6]. Hence previous models of potential-driven anisotropic inflation inevitably include a vector field [7–10].

Here we wish to show that these observations are true only in the Einstein gravity, and that a class of modified gravity with a scalar field can realize anisotropic (inflationary) solution without introducing any vector fields nor higher-order curvature terms as an effective anisotropic stress source [11–13]. Specifically we consider Horndeski theory [14] or the generalized Galileon [15] to show that the quintic galileon term \mathcal{L}_5 in generalized G-inflation [16] plays an essential role to realize anisotropic inflation. Such terms are known to emerge after Kaluza-Klein compactification of higher-dimensional Lovelock gravity [17] on one hand but its magnitude has been severely constrained [18–21] now on the other hand by the simultaneous discovery of the gravitational-wave event GW170817 [22] of binary neutron star coalescence and the associated gamma-ray burst GRB170817A [23], which shows that the relative deviation of the speed of gravitational waves from light velocity is at most 10^{-15} . Such an observational constraint on \mathcal{L}_5 , however, applies only in the low-redshift universe, and it may well evolve nontrivially in the high energy regime in the early universe.

The present paper is organized as follows. In Section II, we review the covariant form and ADM form of Horndeski theory and its equation of evolution. In Section III, by using the trace-free part of the equation of evolution, we show there are emerged solutions which describe expanding universes with nonvanishing anisotropies. In Section IV, we apply the solution to the Bianchi-I model without matter, which is the simplest homogeneous anisotropic model. In Section V, we disscuss the nature of the solution and cosmological application, and we conclude in Section VI.

II. HORNDESKI THEORY AND BEYOND

The Horndeski theory [14] describes the most general couplings between a scalar field ϕ and the metric $g_{\mu\nu}$ which yield second-order field equations. It was rediscovered in [15] in the context of generalized Galileons and their equivalence was proved in [16].

This theory is characarized by four arbitrary functions, G_2 , G_3 , G_4 and G_5 , of ϕ and its

canonical kinetic function $X \equiv -\partial_{\mu}\phi\partial^{\mu}\phi/2$ as

$$S = \int \mathrm{d}^4 x \sqrt{-g} \sum_{i=2}^5 \tilde{\mathcal{L}}_i,\tag{1}$$

$$\tilde{\mathcal{L}}_2 = G_2(\phi, X),\tag{2}$$

$$\mathcal{L}_3 = -G_3(\phi, X) \Box \phi, \tag{3}$$

$$\mathcal{L}_4 = G_4(\phi, X)\mathcal{R} + G_{4X}[(\Box\phi)^2 - (\nabla_\mu\nabla_\nu\phi)^2], \tag{4}$$

$$\tilde{\mathcal{L}}_5 = G_5(\phi, X)\mathcal{G}_{\mu\nu}\nabla^{\mu}\nabla^{\nu}\phi - \frac{1}{6}G_{5X}[(\Box\phi)^3 - 3(\Box\phi)(\nabla_{\mu}\nabla_{\nu}\phi)^2 + 2(\nabla_{\mu}\nabla_{\nu}\phi)^3],$$
(5)

where \mathcal{R} is the Ricci scalar, $\mathcal{G}_{\mu\nu}$ is the Einstein tensor, $(\nabla_{\mu}\nabla_{\nu}\phi)^2 = \nabla_{\mu}\nabla_{\nu}\phi\nabla^{\mu}\nabla^{\nu}\phi,$ $(\nabla_{\mu}\nabla_{\nu}\phi)^3 = \nabla_{\mu}\nabla_{\nu}\phi\nabla^{\nu}\nabla^{\lambda}\phi\nabla_{\lambda}\nabla^{\mu}\phi,$ and $G_{iX} = \partial G_i/\partial X.$

The action described by the ADM variables is more useful to study anisotropic cosmological solutions than the covariant form (1). The metric is given by

$$ds^{2} = -N^{2}dt^{2} + g_{ij}(dx^{i} + N^{i}dt)(dx^{j} + N^{j}dt).$$
(6)

We take the unitary gauge, $\phi = \phi(t)$, and then X is given by $X = \dot{\phi}^2/2N^2$ with N being the lapse function. If ϕ is a monotonic function of t, this is a very convenient gauge and we can use (t, N) instead of (ϕ, X) to express the action. Then, the theory is described only in terms of t and geometrical quantities as

$$S = \int \mathrm{d}t \mathrm{d}^3 x N \sqrt{g} \sum_{i=2}^5 \mathcal{L}_i,\tag{7}$$

$$\mathcal{L}_2 = A_2(t, N), \tag{8}$$

$$\mathcal{L}_{3} = A_{3}(t, N)K,$$
(9)
$$\mathcal{L}_{4} = A_{4}(t, N)\left(K^{2} - K^{i}K^{j}\right) + B_{4}(t, N)R$$
(10)

$$\mathcal{L}_{4} = A_{4}(t, N) \left(K^{3} - 3KK^{i}K^{j} + 2K^{i}K^{j}K^{k} \right) + B_{7}(t, N) \left(B_{12} - \frac{1}{2}a_{12}R \right) K^{ij}$$
(10)

$$\mathcal{L}_{5} = \Pi_{5}(t, W) \left(\Pi - 0 \Pi_{j} \Pi_{i} + 2 \Pi_{j} \Pi_{k} \Pi_{i} \right) + \mathcal{L}_{5}(t, W) \left(\Pi_{ij} - 2^{g_{ij}} \Pi_{j} \right) \Pi_{i}$$
 (II)
and \mathcal{R}_{i} are the extrinsic and intrinsic curvature of constant t (constant ϕ) hypersurfaces

 K_j^i and R_{ij} are the extrinsic and intrinsic curvature of constant t (constant ϕ) hypersurfaces. The functions A_i, B_i and G_i are related with each other as follows:

$$A_2(t,N) = G_2(\phi,X) - \sqrt{X} \int \frac{G_{3\phi}(\phi,X)}{\sqrt{X}} dX,$$
(12)

$$A_{3}(t,N) = \int \sqrt{2X} G_{3X}(\phi,X) dX - 2\sqrt{2X} G_{4\phi}(\phi,X), \qquad (13)$$

$$A_4(t,N) = -G_4(\phi,X) + 2XG_{4X}(\phi,X) - XG_{5\phi}(\phi,X),$$
(14)

$$A_5(t,N) = \frac{1}{6} (2X)^{3/2} G_{5X}(\phi,X), \tag{15}$$

$$B_4(t,N) = G_4(\phi,X) - \frac{\sqrt{X}}{2} \int \frac{G_{5\phi}(\phi,X)}{\sqrt{X}} dX,$$
(16)

$$B_5(t,N) = -\int \sqrt{2X} G_{5X}(\phi,X) dX, \qquad (17)$$

where we identify $X = \dot{\phi}^2(t)/2N^2$. As seen below, among those terms the most crucial ones in this paper are the terms cubic in the extrinsic curvature. In the covariant language they come from $\tilde{\mathcal{L}}_5$ which depends cubically on the second derivatives of the scalar field. In the Horndeski theory, (A_4, A_5) and (B_4, B_5) are not independent, as is clear from Eqs. (14)–(17) and also from the fact that we originally have four free functions in the action. However, this point turns out to be not essential in the following discussion. The most important ingredient here is the cubic (or higher) order terms in the extrinsic curvature. This allows us to start from the ADM Lagrangians (8)–(11) and consider all A_i 's and B_i 's to be independent free functions, which amounts to employing the so-called "beyond Horndeski" theory [24], although A_i 's and B_i 's may have to satisfy degeneracy conditions to avoid an extra dangerous degree of freedom [25, 26] (see, however, [27]). The following discussion can thus apply not only to the Horndeski theory but also to beyond Horndeski theory.

In addition to the action for the gravitational sector described above, we include the action for matter minimally coupled to gravity, $S_{\rm m}$. By the use of the residual gauge degrees of freedom one can further impose $N^i = 0$. Then, we obtain the evolution equations from (7) as

$$T_{j}^{i} = \frac{1}{N\sqrt{g}} \partial_{t} \left\{ \sqrt{g} \left\{ A_{3}\delta_{j}^{i} + 2A_{4}(K\delta_{j}^{i} - K_{j}^{i}) + 3A_{5}[(K^{2} - K_{l}^{k}K_{k}^{l})\delta_{j}^{i} - 2(KK_{j}^{i} - K_{k}^{i}K_{j}^{k})] \right\} \right\} - \delta_{j}^{i}\mathcal{L}_{A} + \left(2B_{4} + \frac{\partial_{t}B_{5}}{N} \right) \left(R_{j}^{i} - \frac{1}{2}\delta_{j}^{j}R \right) + \Phi_{j}^{i},$$
(18)

where T_{ij} is the stress-energy tensor calculated from the matter action $S_{\rm m}$,

$$T_{ij} = -\frac{2}{N\sqrt{g}} \frac{\delta S_{\rm m}}{\delta g^{ij}},\tag{19}$$

and \mathcal{L}_A is the kinetic part of the Lagrangian,

$$\mathcal{L}_A = A_2 + A_3 K + A_4 (K^2 - K_j^i K_i^j) + A_5 (K^3 - 3K K_j^i K_i^j + 2K_j^i K_k^j K_i^k).$$
(20)

We have collected the terms that vanish if the lapse function is homogeneous, $N(t, \vec{x}) = N(t)$, and written

$$\Phi_{ij} = \frac{2}{N} [\nabla^{2} (NB_{4}) g_{ij} - \nabla_{i} \nabla_{j} (NB_{4})]
+ g_{ij} K^{lm} \nabla_{l} \nabla_{m} B_{5} + K \nabla_{k} \nabla_{j} B_{5} - 2K^{l}_{(i} \nabla_{j)} \nabla_{l} B_{5} + K_{ij} \nabla^{2} B_{5} - g_{ij} K \nabla^{2} B_{5}
+ \frac{2}{N} \left[g_{ij} \nabla_{l} (NK^{lm}) \nabla_{m} B_{5} + \nabla_{(i} (NK) \nabla_{j)} B_{5} - \nabla_{l} (NK^{l}_{(i)}) \nabla_{j} B_{5}
- \nabla_{(i} (NK^{l}_{j)}) \nabla_{l} B_{5} + \nabla_{l} (NK_{ij}) \nabla^{l} B_{5} - g_{ij} \nabla_{l} (NK) \nabla^{l} B_{5} \right].$$
(21)

The Hamiltonian constraint is given by

$$\partial_{N}(NA_{2}) + N\partial_{N}A_{3}K + N^{2}\partial_{N}(N^{-1}A_{4})(K^{2} - K_{j}^{i}K_{i}^{j}) + \partial_{N}(NB_{4})R + N^{3}\partial_{N}(N^{-2}A_{5})(K^{3} - 3KK_{j}^{i}K_{i}^{j} + 2K_{j}^{i}K_{k}^{j}K_{i}^{k}) + N\partial_{N}B_{5}\left(R_{ij}K^{ij} - \frac{1}{2}RK\right) + \frac{1}{\sqrt{g}}\frac{\delta S_{m}}{\delta N} = 0.$$
(22)

In the following we will not use the momentum constraint equations.

III. SELF-ANISOTROPIZING INFLATIONARY SOLUTIONS

We now show that even without any anisotropic matter sources the universe can exhibit anisotropic inflationary expansion as an atractor solution in the Horndeski theory.

Since we consider Bianchi cosmology, we may set $N^i = 0$. Thanks to the homogeneity, Φ_{ij} in the evolution equation (18) vanishes. To study anisotropic cosmological models it is convenient to decompose the extrinsic curvature K_{ij} into its trace K and trace-free part Σ_{ij} as

$$K_{ij} = \frac{1}{3}Kg_{ij} + \Sigma_{ij},\tag{23}$$

with $g^{ij}\Sigma_{ij} = 0$. The trace and trace-free parts of the evolution equation (18) read, respectively,

$$\frac{1}{N\sqrt{g}}\partial_t \left[\sqrt{g}(3A_3 + 4A_4K + A_5(2K^2 - 3\Sigma_j^i\Sigma_i^j))\right] - 3\mathcal{L}_A - \left(B_4 + \frac{\partial_t B_5}{2N}\right)R = T_i^i.$$
(24)

and

$$\frac{2}{N\sqrt{g}}\partial_t \left[\sqrt{g}(-A_4\Sigma_j^i - A_5K\Sigma_j^i + 3A_5\{\Sigma_k^i\Sigma_j^k\}_{\mathrm{TF}})\right] + \left(2B_4 + \frac{\partial_t B_5}{N}\right)\{R_j^i\}_{\mathrm{TF}} = \{T_j^i\}_{\mathrm{TF}},$$
(25)

where $\{X_{j}^{i}\}_{\text{TF}}$ stands for the trace-free part of a tensor X_{j}^{i} ,

$$\{X_{j}^{i}\}_{\rm TF} = X_{j}^{i} - \frac{1}{3}X_{k}^{k}\delta_{j}^{i}.$$
(26)

Let us look for slow-roll inflationary solutions in which \sqrt{g} exponentially increases, while other functions remain either nearly constant or exponentially decrease. First, we focus on Eq. (25), assuming that the energy-momentum tensor consists of isotropic matter and hence $\{T_j^i\}_{TF}$ vanishes. If the spatial curvature R_j^i decreases exponentially, the first term also decreases in the same way. As a result, we find, asymptotically,

$$-A_4 \Sigma_j^i - A_5 K \Sigma_j^i + 3A_5 (\Sigma_k^i \Sigma_j^k - \frac{1}{3} \Sigma_l^k \Sigma_k^l \delta_j^i) = 0.$$
(27)

A trivial solution of Eq. (27) is that all components of Σ_j^i vanish. This solution corresponds to the isotropic attractor which we see in the conventional inflation models. The presence of the quadratic terms in Σ_j^i due to nonvanishing A_5 yields nontrivial solutions with $\Sigma_j^i \neq 0$ as well, which represent an expanding universe retaining finite anisotropies. We dub this anisotropic attractors as *self-anisotropizing* inflationary solutions, as this is *not* caused by an anisotropic energy-momentum tensor. We will demonstrate in the next section that such solutions do exist in the case of Bianchi type-I cosmology.

The self-anisotropizing attractors are distinct from the previous anisotropic inflationary solutions, because the anisotropic expansion of the previous scenarios are supported by some anisotropic energy-momentum source such as a vector field coupled with an inflaton field [8]. Such scenarios produce background anisotropies $\Sigma_j^i/H \approx \{T_j^i\}_{\rm TF}/(6A_4H^2) =$ $(8\pi G/3H^2)\{T_j^i\}_{\rm TF}$, where H is the Hubble parameter. The trace-free part of the energymomentum tensor, $\{T_j^i\}_{\rm TF}$, just displaces the terminal point from the isotropic one. By contrast, here the self-anisotropizing inflationary solution is realized by the terms quadratic in Σ_j^i in Eq. (27), which is a consequence of modification of gravity. The magnitude of produced background anisotropies is estimated from (27) as $\Sigma_j^i/H \sim (A_4 + 3HA_5)/3HA_5$. We require neither an anisotropic energy-momentum tensor nor any fields other than the scalar ϕ built in the Horndeski theory. In this sense, the emerged anisotropic terminal points should be distinguished from those of previous anisotropic inflation models.

Let us evaluate the eigenvalues of the nontrivial solutions of Σ_j^i for given values of A_4 , A_5 and K. We can prove that the root Σ of matrix equation (27) has two different eigenvalues at most as follows. First we define a polynomial p(x) by substituting a real variable x for Σ in the left side of (27) as

$$p(x) = -A_4 x - A_5 K x + 3A_5 \left(x^2 - \frac{1}{3} \text{tr}(\Sigma^2) \right), \qquad (28)$$

where the remaining Σ in the trace is a root of (27). $p(\Sigma) = 0$ obviously follows from (27) and (28), and so p(x) can be divided by the minimal polynomial $\phi_{\Sigma}(x)$ of Σ . In linear algebra, it is well-known that if λ is an eigenvalue of matrix Σ then λ is a root of $\phi_{\Sigma}(x) = 0$. Therefore, the eigenvalue λ is also a root of p(x) = 0. Since p(x) is a quadratic polynomial of x, the number of different roots is equal to or less than two. This is the proof that Σ has two different eigenvalues, λ_1 and λ_2 at most. It induces that, *e.g.*, anisotropic attractors in Bianchi type-I model has axial symmetry in the order of background, which we show in Section IV. As one can see from (28), the different eigenvalues λ_1 and λ_2 satisfy

$$\lambda_1 + \lambda_2 = \frac{A_4 + A_5 K}{3A_5}.$$
 (29)

Being a three dimensional tensor, Σ has three eigenvalues. Without loss of generality, we set them as λ_1 , λ_1 and λ_2 , respectively. They also satisfy

$$2\lambda_1 + \lambda_2 = 0, \tag{30}$$

because Σ is trace-free. Therefore we have

$$\lambda_1 = -\frac{A_4 + A_5 K}{3A_5}, \quad \lambda_2 = \frac{2(A_4 + A_5 K)}{3A_5}.$$
(31)

So far we have focused on the evolution equation for Σ_j^i (25) and its nontrivial solution under the assumption that the spatial volume element \sqrt{g} increases exponentially and the spatial curvature R_j^i decreases accordingly. To determine all the components of the metric, we need to solve the Hamiltonian constraint (22) and the trace part of the evolution equations (24) consistently. On the anisotropic attractor where Σ_j^i 's eigenvalues are given by (31), the rest of the field equations (22) and (24) reduce to

$$\partial_N \left[N \left(A_2 - \frac{2A_4^3}{9A_5^2} \right) \right] + N \partial_N \left(A_3 - \frac{2A_4^2}{3A_5} \right) K = -\frac{1}{\sqrt{g}} \frac{\delta S_{\rm m}}{\delta N},\tag{32}$$

$$\frac{1}{N}\frac{d}{dt}\left(A_3 - \frac{2A_4^2}{3A_5}\right) - \left(A_2 - \frac{2A_4^3}{9A_5^2}\right) = \frac{1}{3}T_i^i,\tag{33}$$

respectively. These two equations can be used to determine K = K(t) and N = N(t).

Let us ignore the matter field $S_{\rm m}$ for the moment and consider a theory with (approximate) shift symmetry. In this case, A_i 's depend only on $\dot{\phi}/N$ and from Eq. (33) one obtains a solution $N \simeq \text{const} \times \dot{\phi}$ satisfying $F(\dot{\phi}/N) \equiv A_2 - 2A_4^3/9A_5^2 \simeq 0$. Equation (32) is then solved to give $K \simeq -\partial_N (A_3 - 2A_4^2/3A_5)/\partial_N (A_2 - 2A_4^3/9A_5^2) \simeq \text{const}$. One thus obtains an inflating solution with nonvanishing anisotropies.

IV. VACUUM BIANCHI TYPE-I MODEL

A. Evolution toward attractors

To be more explicit, let us consider the Bianchi type-I model, which is the simplest homogeneous anisotropic model and hence helps us to understand what nonvanishing A_5 causes.

Once we diagonalize the spatial metric and its time derivative, off-diagonal components are not generated in this model, so that we can express the metric in the Kasner-type form as

$$ds^{2} = -N^{2}(t)dt^{2} + a^{2}(t)\left[e^{2(\beta_{+}(t) + \sqrt{3}\beta_{-}(t))}dx^{2} + e^{2(\beta_{+}(t) - \sqrt{3}\beta_{-}(t))}dy^{2} + e^{-4\beta_{+}(t)}dz^{2}\right], \quad (34)$$

where a(t) is a scale factor and $\beta_{\pm}(t)$ show the differences between the expansion rates in different directions. Substituting the metric (34) in the ADM form of the action (7), we obtain

$$S = \int dt d^3x \ Na^3 \left[A_2 + 3HA_3 + 6A_4 (H^2 - \sigma_+^2 - \sigma_-^2) + 6A_5 (H^3 - 3H(\sigma_+^2 + \sigma_-^2) + 2(3\sigma_+\sigma_-^2 - \sigma_+^3)) \right]$$
(35)

where we defined the Hubble parameter H and the shear σ_{\pm} as

$$H \equiv \frac{1}{N} \frac{d \ln a}{dt}, \quad \sigma_{\pm} \equiv \frac{1}{N} \frac{d\beta_{\pm}}{dt}.$$
(36)

Using σ_{\pm} , the trace-free part of the extrinsic curvature is given by

$$\Sigma_j^i = \operatorname{diag}\left(\sigma_+ + \sqrt{3}\sigma_-, \sigma_+ - \sqrt{3}\sigma_-, -2\sigma_+\right).$$
(37)

Since the spatial Ricci tensor vanishes in the Bianchi type-I model and consequently Eq. (35) depends on β_{\pm} only through their time derivatives, the conjugate momenta of β_{\pm} are conserved in time. The conserved momenta are given by

$$P_{\beta_{+}} = a^{3} \left[(A_{4} + 3HA_{5})\sigma_{+} + 3A_{5}(\sigma_{+}^{2} - \sigma_{-}^{2}) \right], \qquad (38)$$

$$P_{\beta_{-}} = a^{3} \left[(A_{4} + 3HA_{5})\sigma_{-} - 6A_{5}\sigma_{+}\sigma_{-} \right].$$
(39)

Equivalently, one can also obtain the same conserved momenta from the trace-free part of the evolution equations (25) by substituting Eq. (34). It is manifest that as the scale factor a(t) increases, the expressions inside the square brackets of Eqs. (38) and (39) decay toward zero as $[\cdots] = P_{\beta_{\pm}} a^{-3} \rightarrow 0$, and thus σ_{+} and σ_{-} evolve to one of the fixed points. In the present case, there are four fixed points. One is the isotropic solution $\sigma_{\pm} = 0$, whereas the other three are anisotropic attractors.

Let us look at the trajectories on the (σ_+, σ_-) plane of the phase space. Given the initial data, the constants $P_{\beta_{\pm}}$ are fixed. Then, σ_{\pm} can be expressed in terms of A_4 , A_5 , a, H, and $P_{\beta_{\pm}}$ by solving the algebraic equations (38) and (39). In order to show the dynamics of the anisotropies in a single figure, we use the normalized shear \mathcal{A}_{\pm} defined as

$$\mathcal{A}_{\pm} \equiv \frac{3A_5}{A_4 + 3HA_5} \sigma_{\pm},\tag{40}$$

instead of σ_+ and σ_- . Here we assumed that $A_4 + 3HA_5 \neq 0$ and $A_5 \neq 0$. It is also convenient to introduce the new time coordinate $\tau \equiv a^3(A_4 + 3HA_5)^2/3A_5$. In an expanding universe, $|\tau|$ is an increasing function of t provided that A_4 , A_5 , and H depend on t only weakly, which is a natural assumption during inflation. With τ and \mathcal{A}_{\pm} , we can rewrite Eqs. (38) and (39) simply as

$$P_{\beta_+} = \tau \left[\mathcal{A}_+ + \mathcal{A}_+^2 - \mathcal{A}_-^2 \right], \tag{41}$$

$$P_{\beta_{-}} = \tau \left[\mathcal{A}_{-} - 2\mathcal{A}_{+}\mathcal{A}_{-} \right].$$
(42)

We show trajectories $(\mathcal{A}_{+}(\tau), \mathcal{A}_{-}(\tau))$ for different values of $P_{\beta\pm}$ in Figure 1. As stated above, there are four fixed points in the $(\mathcal{A}_{+}, \mathcal{A}_{-})$ plane: one isotropic solution, (0, 0), and three anisotropic solutions, (-1, 0) and $(1/2, \pm\sqrt{3}/2)$. All of them are attractors (as long as $|\tau|$ is an increasing function of t).



FIG. 1. Trajectries of the evolution of the normalized shear $(\mathcal{A}_+, \mathcal{A}_-)$. If the initial conditions lie inside the circle given by $\mathcal{A}_+^2 + \mathcal{A}_-^2 = 1/4$, the universe evolves toward the center, $(\mathcal{A}_+, \mathcal{A}_-) = (0, 0)$, as τ increases. If the universe starts from outside of the circle, it goes to the closest one of the anisotropic fixed points on the vertices, $(\mathcal{A}_+, \mathcal{A}_-) = (-1, 0), (1/2, \pm\sqrt{3}/2)$, of the triangle as τ increases.

The initial anisotropies determine which attractor the universe approaches. To see this explicitly, we differentiate Eqs. (41) and (42) and get

$$\tau \frac{d\mathcal{A}_{+}}{d\tau} = -\frac{(2\mathcal{A}_{+} - 1)(\mathcal{A}_{+}^{2} + \mathcal{A}_{-}^{2} + \mathcal{A}_{+})}{4\mathcal{A}_{+}^{2} + 4\mathcal{A}_{-}^{2} - 1},$$
(43)

$$\tau \frac{d\mathcal{A}_{-}}{d\tau} = -\frac{\mathcal{A}_{-}(2\mathcal{A}_{+}^{2} + 2\mathcal{A}_{-}^{2} - 2\mathcal{A}_{+} - 1)}{4\mathcal{A}_{+}^{2} + 4\mathcal{A}_{-}^{2} - 1}.$$
(44)

Equivalently, one may introduce the polar coordinates $(r(\tau), \theta(\tau))$ defined by $\mathcal{A}_+ = r \cos \theta$ and $\mathcal{A}_- = r \sin \theta$ and write

$$\tau \frac{dr}{d\tau} = -\frac{r[2r^2 + r\cos(3\theta) - 1]}{4r^2 - 1},\tag{45}$$

$$\tau \frac{d\theta}{d\tau} = \frac{r\sin(3\theta)}{4r^2 - 1}.\tag{46}$$

The denominators vanish on a circle given by $r^2 = \mathcal{A}_+^2 + \mathcal{A}_-^2 = (1/2)^2$ (the black circle in Fig. 1).¹ The fate of the universe depends on whether the initial anisotropies are inside this circle or not: the universe is attracted toward the isotropic solution at the origin if the initial anisotropies lie inside the circle, while it goes away from the circle to the closest one of the anisotropic attractors if outside initially. That is to say, if the universe is sufficiently anisotropic initially, then it converges to the anisotropic attractor.

The exceptional case is the trajectories with $\theta = 0, 2\pi/3, 4\pi/3$. Those constant values of θ solve Eq. (46), while Eq. (45) leads to $r(\tau) = (\sqrt{C/|\tau| + 1} - 1)/2$, where C is an integration constant. Therefore, for all initial conditions on $\theta = 0, 2\pi/3, 4\pi/3$ the isotropic universe is the attractor.

The structure of Fig. 1 will be more transparent in terms of the polar coordinates. Equations (45) and (46) clearly show that there are discrete rotation symmetry $\theta \rightarrow \theta + 2\pi/3$ and reflection symmetry across $\theta = 0, 2\pi/3$, and $4\pi/3$ axises. Because of these symmetries only a sixth part of Fig. 1 is physically independent.

Each of the anisotropic attractors corresponds to an axially symmetric space, whose symmetry axis is the x, y or z axis. This axial symmetry is closely related to the degeneracy of the eigenvalues of Σ_i^j discussed in the previous section. The discrete rotation symmetry in the $(\mathcal{A}_+, \mathcal{A}_-)$ plane is the manifestation of the fact that one can always take, say, the zaxis as the symmetry axis without loss of generality by a rotation of the spatial coordinates.

So far we have focused only on the shear evolution equations. This is sufficient for the purpose of seeing that the anisotropic fixed points do exist and for initial anisotropies larger than a certain threshold they are indeed the attractors. To determine the precise dynamics of the universe including the evolution of H and ϕ , one needs to solve the full set of the field equations (the trace and trace-free parts of the evolution equations as well as the constraint equation) consistently. In the next subsection we will show a numerical example obtained by solving all the equations consistently.

B. Examples

Let us present some examples which yield self-anisotropizing Bianchi type-I solutions. The first one is simply given by

$$G_2 = -V_0, \quad G_3 = 0, \quad G_4 = \frac{M^2}{2} + g_4 X, \quad G_5 = g_5 X,$$
 (47)

¹ The shear evolution equations bocome singular on this circle. However, if we consider the full phase space by taking into account the trace part of the evolution equation, we see that this singularity is only apparent.

where V_0 , M, g_4 , and g_5 are constants. The corresponding ADM form in the unitary gauge is given by

$$A_{2} = -V_{0}, \quad A_{3} = 0, \quad A_{4} = -\frac{M^{2}}{2} + \frac{g_{4}}{2} \left(\frac{\dot{\phi}}{N}\right)^{2}, \quad A_{5} = \frac{g_{5}}{6} \left(\frac{\dot{\phi}}{N}\right)^{3},$$
$$B_{4} = \frac{M^{2}}{2} + \frac{g_{4}}{2} \left(\frac{\dot{\phi}}{N}\right)^{2}, \quad B_{5} = -\frac{g_{5}}{3} \left(\frac{\dot{\phi}}{N}\right)^{3}.$$
(48)

Figure 2 shows the evolution of the Hubble parameter, (the velocity of) the scalar field, and the shear obtained by solving the dynamical and constraint equations numerically with a certain initial condition away from the attractors at a = 1. The parameters in this toy example are given by $V_0 = 0.1$, M = 1, $g_4 = -0.2$, and $g_5 = 1$. It can be seen that the universe quicky converges to the anisoropic inflationary attractor.



FIG. 2. Numerical example of a self-anisotrpizing Bianchi type-I universe: (a) H; (b) $\dot{\phi}/N$; (c) σ_{\pm}/H as functions of $\ln a$.

Another example with A_5 (or, equivalently, G_{5X}) is the Gauss-Bonnet term coupled to a scalar field, and the total Lagrangian is of the form

$$\mathcal{L} = f(\phi)\mathcal{R} + P(\phi, X) + \xi(\phi)\left(\mathcal{R}^2 - 4\mathcal{R}_{\mu\nu}\mathcal{R}^{\mu\nu} + \mathcal{R}_{\mu\nu\rho\sigma}\mathcal{R}^{\mu\nu\rho\sigma}\right).$$
(49)

Aspects of this theory has been studied extensively in the literature. The Lagrangian can be reproduced by taking the following Horndeski functions [16]:

$$G_{2} = P + 8\xi^{(4)}X^{2}(3 - \ln X), \quad G_{3} = 4\xi^{(3)}X(7 - 3\ln X),$$

$$G_{4} = f + 4\xi^{(2)}X(2 - \ln X), \quad G_{5} = -4\xi^{(1)}\ln X,$$
(50)

where $\xi^{(n)} = d^n \xi / d\phi^n$. Though this looks quite non-trivial, the corresponding ADM form is very simple:

$$A_2 = P, \quad A_3 = -2\frac{\dot{\phi}}{N}\frac{df}{d\phi}, \quad A_4 = -f, \quad A_5 = -\frac{4\xi^{(1)}}{3}\frac{\dot{\phi}}{N}, \quad B_4 = f, \quad B_5 = 8\xi^{(1)}\frac{\dot{\phi}}{N}.$$
(51)

Even this familiar theory admits self-anisotropizing inflationary solutions.

The theory (49) possesses a shift symmetry if f = const, P = P(X), and $\xi \propto \phi$. In this case it is easy to find an inflationary solution with H = const, $\dot{\phi}/N = \text{const}$ retaining the nonvanishing shear

$$\frac{\sigma_{\pm}}{H} \sim \frac{f + 4H\xi^{(1)}\dot{\phi}/N}{H\xi^{(1)}\dot{\phi}/N}.$$
(52)

V. DISCUSSION

It has been pointed out by Wald that in general relativity, all vacuum Bianchi universes with a positive cosmological constant except type IX evolve toward the isotropic attractor, which was proven by using the Hamiltonian constraint and the trace of the Einstein equations [28]. In our case, since the Horndeski action dramatically changes both of them, it must be checked one by one whether a specific model under consideration evolves toward the isotropic or anisotropic attractor. We note that the magnitude of the shear on the anisotropic attractors diverges when we take the general relativity limit $A_5 \rightarrow 0$ keeping A_4 constant. In this limit, for all initial conditions the isotropic universe is an attractor (as they are all inside the circle in Fig. 1), and thus the standard result of Wald in gerenal relativity is recovered.

Noting that the background anisotropies of the Bianchi type-I universe can be regarded as gravitational waves with infinitely long wavelengths, we point out that the emergence of anisotropic attractors is closely related to the three-point coupling of gravitational waves in the Horndeski theory. From Eq. (15) of [29], one sees that there are two types of the three-point couplings of the form $hh\partial^2 h$ and $\dot{h}\dot{h}\dot{h}$, giving rise to local and equilateral non-Gaussianity, respectively. The former appears even in general relativity as well as in a generic scalar-tensor theory, while the latter, which obviously comes from K_{ij}^3 , emerges only in the class with $A_5 \neq 0$ (i.e., $G_{5X} \neq 0$). The former has spatial derivatives and therefore vanishes in the long-wavelength limit, whereas the latter has only time derivatives and hence does not vanish even in the homogeneous limit. Since $\mathcal{A}_{\pm} = \mathcal{O}(1)$ on the anisotropic attractors, the magnitude of the resultant anisotropy is given by

$$\frac{\sigma_{\pm}}{H} \sim \frac{A_4 + 3HA_5}{3HA_5},\tag{53}$$

which is typically of $\mathcal{O}(1)$ or larger. In theories with $G_{5X} \neq 0$, initial anisotropies must be smaller than this value in order to realize an isotropic universe through inflation. Otherwise, the resultant universe would be unacceptably anisotropic. Another possibility is that one has $A_4 + 3HA_5 \ll 3HA_5$ via fine-tuning, leaving an observationally viable universe with only tiny anisotropies on the anisotropic attractor. This would be a very interesting scenario, but one has to study reheating, cosmological perturbations, and the stability in detail to see whether such a scenario is indeed viable or not, which is beyond the scope of the present paper.

VI. CONCLUSIONS

We have shown the quintic galileon term proportional G_{5X} or A_5 in generalized G-inflation can realize anisotropic inflationary solution. On the anisotropic attractor, the Hamiltonian constraint becomes a linear equation for the Hubble parameter which is strikingly different from the conventional Friedmann equation.

Although our solution generically produces anisotropy of the order of unity or larger, it can also accommodate much smaller anisotropy by partially canceling A_4 and $3HA_5$. In order to see if observationally viable anisotropic inflation is possible, we must calculate perturbations as well as discuss transition to the Friedmann regime with proper reheating, which will be discussed in future publications.

It is also interesting to study higher dimensional models in this context to show a new compactification mechanism of extra dimensions in the presence of the highest-order galileon terms in the dimension under consideration. As we can show that the eigenvalues of the extrinsic curvature tensor take only two distinct values at most even in higher dimensional models, this may provide a promissing mechanism of compactification or dimensional reduction, which will also be discussed in a forthcoming paper.

ACKNOWLEDGMENTS

HWHT was supported by the Advanced Leading Graduate Course for Photon Science (ALPS). The work of TK was supported by MEXT KAKENHI Grant Nos. JP15H05888, JP16H01102, JP17H06359, JP16K17707, and MEXT-Supported Program for the Strategic Research Foundation at Private Universities, 2014-2018 (S1411024). The work of JY was supported by JSPS KAKENHI, Grant JP15H02082 and Grant on Innovative Areas JP15H05888.

- [1] A. A. Starobinsky, Phys. Lett. **B91**, 99 (1980), [,771(1980)].
- [2] K. Sato, Mon. Not. Roy. Astron. Soc. **195**, 467 (1981).
- [3] A. H. Guth, Phys. Rev. **D23**, 347 (1981).

- [4] K. Sato and J. Yokoyama, Int. J. Mod. Phys. **D24**, 1530025 (2015).
- [5] L. G. Jensen and J. A. Stein-Schabes, Phys. Rev. **D34**, 931 (1986).
- [6] M. S. Turner and L. M. Widrow, Phys. Rev. Lett. 57, 2237 (1986).
- [7] L. H. Ford, Phys. Rev. **D40**, 967 (1989).
- [8] M.-a. Watanabe, S. Kanno, and J. Soda, Phys. Rev. Lett. 102, 191302 (2009), arXiv:0902.2833 [hep-th].
- [9] T. Q. Do and W. Kao, Phys. Rev. **D96**, 023529 (2017).
- [10] P. Adshead and A. Liu, (2018), arXiv:1803.07168 [astro-ph.CO].
- [11] J. D. Barrow and S. Hervik, Phys. Rev. D73, 023007 (2006), arXiv:gr-qc/0511127 [gr-qc].
- [12] J. D. Barrow and S. Hervik, Phys. Rev. D74, 124017 (2006), arXiv:gr-qc/0610013 [gr-qc].
- [13] J. D. Barrow and S. Hervik, Phys. Rev. **D81**, 023513 (2010), arXiv:0911.3805 [gr-qc].
- [14] G. W. Horndeski, Int. J. Theor. Phys. 10, 363 (1974).
- [15] C. Deffayet, X. Gao, D. A. Steer, and G. Zahariade, Phys. Rev. D84, 064039 (2011), arXiv:1103.3260 [hep-th].
- [16] T. Kobayashi, M. Yamaguchi, and J. Yokoyama, Prog. Theor. Phys. 126, 511 (2011), arXiv:1105.5723 [hep-th].
- [17] K. Van Acoleyen and J. Van Doorsselaere, Phys. Rev. D83, 084025 (2011), arXiv:1102.0487 [gr-qc].
- [18] T. Baker, E. Bellini, P. G. Ferreira, M. Lagos, J. Noller, and I. Sawicki, Phys. Rev. Lett. **119**, 251301 (2017), arXiv:1710.06394 [astro-ph.CO].
- [19] P. Creminelli and F. Vernizzi, Phys. Rev. Lett. 119, 251302 (2017), arXiv:1710.05877 [astro-ph.CO].
- [20] J. Sakstein and B. Jain, Phys. Rev. Lett. **119**, 251303 (2017), arXiv:1710.05893 [astro-ph.CO].
- [21] J. M. Ezquiaga and M. Zumalacárregui, Phys. Rev. Lett. 119, 251304 (2017), arXiv:1710.05901 [astro-ph.CO].
- [22] B. Abbott *et al.* (Virgo, LIGO Scientific), Phys. Rev. Lett. **119**, 161101 (2017), arXiv:1710.05832 [gr-qc].
- [23] A. Goldstein *et al.*, Astrophys. J. 848, L14 (2017), arXiv:1710.05446 [astro-ph.HE].
- [24] J. Gleyzes, D. Langlois, F. Piazza, and F. Vernizzi, Phys. Rev. Lett. 114, 211101 (2015), arXiv:1404.6495 [hep-th].
- [25] D. Langlois and K. Noui, JCAP **1602**, 034 (2016), arXiv:1510.06930 [gr-qc].
- [26] M. Crisostomi, M. Hull, K. Koyama, and G. Tasinato, JCAP 1603, 038 (2016), arXiv:1601.04658 [hep-th].
- [27] A. De Felice, D. Langlois, S. Mukohyama, K. Noui, and A. Wang, (2018), arXiv:1803.06241 [hep-th].
- [28] R. M. Wald, Phys. Rev. **D28**, 2118 (1983).
- [29] X. Gao, T. Kobayashi, M. Yamaguchi, and J. Yokoyama, Phys. Rev. Lett. 107, 211301 (2011), arXiv:1108.3513 [astro-ph.CO].