Nonparametric Pricing Analytics with Customer Covariates

Ningyuan Chen^{*1} and Guillermo Gallego^{†2}

 ¹Rotman School of Management, University of Toronto
 ²Department of Industrial Engineering & Decision Analytics The Hong Kong University of Science and Technology

Abstract

Personalized pricing analytics is becoming an essential tool in retailing. Upon observing the personalized information of each arriving customer, the firm needs to set a price accordingly based on the covariates such as income, education background, past purchasing history to extract more revenue. For new entrants of the business, the lack of historical data may severely limit the power and profitability of personalized pricing. We propose a nonparametric pricing policy to simultaneously learn the preference of customers based on the covariates and maximize the expected revenue over a finite horizon. The policy does not depend on any prior assumptions on how the personalized information affects consumers' preferences (such as linear models). It is adaptively splits the covariate space into smaller bins (hyper-rectangles) and clusters customers based on their covariates and preferences, offering similar prices for customers who belong to the same cluster trading off granularity and accuracy. We show that the algorithm achieves a regret of order $O(\log(T)^2T^{(2+d)/(4+d)})$, where *T* is the length of the horizon and *d* is the dimension of the covariate. It improves the current regret in the literature

^{*}ningyuan.chen@utoronto.ca

[†]ggallego@ust.hk

(Slivkins, 2014), under mild technical conditions in the pricing context (smoothness and local concavity). We also prove that no policy can achieve a regret less than $O(T^{(2+d)/(4+d)})$ for a particular instance and thus demonstrate the near optimality of the proposed policy.

Keywords: multi-armed bandit, dynamic pricing, online learning, regret analysis, contextual information

1. Introduction

Personalized pricing refers to the practice that a firm charges customers different prices for the *same* product, depending on customers' information such as education backgrounds and zip codes. It is increasingly popular in online retailing, as sellers can acquire/infer the personalized information from customers' account profiles or browsing histories (cookies). The demand (purchasing probability) of each customer depends not only on the price, but also on the personalized information. The firm observes the information of each arriving customer and sets a personalized price accordingly. We are interested in finding pricing policies of the firm that maximize the long-run revenue.

Personalized pricing presents several challenges to the firm. First, for new entrants to the online business, the demand function and how it depends on the personalized information is generally unknown. Thus, the optimal pricing cannot be obtained by directly solving an optimization problem. The firm may experiment with different prices to *learn* the personalized demand function, and then sets optimal prices according to the estimation. There is usually a finite horizon that forces a trade-off between gathering more information (learning/exploration) and making sound decisions (earning/exploitation). This problem, sometimes referred to as the learning/earning dilemma, has attracted the attention of many researchers.

A second challenge is the presence of personalized information, or *covariates*. On one hand, the covariate of each customer provides extra information for the firm to predict the personalized demand more accurately. On the other hand, the demand is peculiar to each instance and changes over time, which adds significant complexity to the learning problem described above. In particular, learning a market-wise demand function is not sufficient, and the firm has to learn the personalized demand by experimenting with

prices for customers of similar profiles.

A third challenge is the selection of a predictive model to estimate the demand. Consider a toy example: the demand function only depends on the address of each customer and nothing else. The firm may postulate a linear model

demand =
$$a - b \times \text{price} + c \cdot (\text{latitude}, \text{longitude})$$

and uses historical sales data to learn the parameters a, b and c and maximize revenue according to the estimation. However, whether the model is specified correctly plays an important role in the performance of the pricing policy. In the particular example, linearity in the location (latitude and longitude) implies that the customers along the straight line that is orthogonal to c have the same demand. This hardly reflects the reality, as customers from the same neighborhood tend to have similar shopping patterns and neighborhoods are usually clustered geographically. By postulating a parametric (linear) model, the firm faces the risk of misspecification and not learning what is supposed to be learned. A *nonparametric* model is more appropriate in this setting.

In this paper we study the pricing policy of a firm which tries to maximize the unknown expected revenue $f(\mathbf{x}, p) \triangleq pd(\mathbf{x}, p)$, where p is the price, $d(\mathbf{x}, p) \in [0, 1]$ is the personalized demand function for a customer of covariate \mathbf{x} . We assume both quantities are normalized so $\mathbf{x} \in [0, 1)^d$ and $p \in [0, 1]$. In period t, the firm observes an arriving customer with a random covariate X_t , and sets a personalized price p_t . The earned revenue is p_t multiplied by a Bernoulli random variable with success rate $d(X_t, p_t)$, representing the event of a purchase. The expected revenue is thus $f(X_t, p_t)$.

We propose a nonparametric learning policy for the firm. That is, the policy does not depend on specific forms of $f(\mathbf{x}, p)$ and only assumes general structures such as continuity and smoothness. The policy achieves near-optimal performance compared to a clairvoyant who knows $f(\mathbf{x}, p)$ and sets $p^*(\mathbf{x}) = \operatorname{argmax}_p f(\mathbf{x}, p)$ for a customer of covariate \mathbf{x} . More precisely, the expected difference in total revenues between the proposed policy and the clairvoyant policy, which is referred to as the *regret* in the literature, grows at $O(\log(T)^2 T^{(2+d)/(4+d)})$ as $T \to \infty$. The rate is sublinear in T, implying that when the length of horizon tends to infinity, the average regret incurred per period becomes negligible. Moreover, we prove that no pricing policies can achieve a lower regret than $O(T^{(2+d)/(4+d)})$ for a reasonable class of unknown objective functions f. Therefore, we successfully work out the learning/earning dilemma with covariates.

The main contribution of the paper is the design of a near-optimal nonparametric learning policy for the personalized pricing problem. Nonparametric learning policies are introduced in more general settings by Rigollet and Zeevi (2010); Perchet and Rigollet (2013); Slivkins (2014). The formulation in this paper is originally introduced in Slivkins (2014), and our policy builds upon the idea of adaptive binning proposed in Perchet and Rigollet (2013)¹. Motivated by the application of personalized pricing, we assume that the expected revenue $f(\mathbf{x}, p)$ is smooth and locally concave in the charged price p. This deviates from the Lipschitz continuous condition in Slivkins (2014) and can be viewed as a special "margin condition" in Rigollet and Zeevi (2010); Perchet and Rigollet (2013). By utilizing this condition, we are able to show that our policy is near-optimal and achieves improved regret over merely continuous objective functions $(T^{(2+d)/(4+d)} \text{ versus } T^{(2+d)/(3+d)} \text{ in Slivkins (2014)}).$

1.1. Literature Review

This paper is motivated by the recent literature that analyzes a firm's pricing problem when the demand function is unknown (e.g. Besbes and Zeevi, 2009; Araman and Caldentey, 2009; Farias and Van Roy, 2010; Broder and Rusmevichientong, 2012; den Boer and Zwart, 2014; Keskin and Zeevi, 2014; Cheung et al., 2017). den Boer (2015) provides a comprehensive survey for this area. Since the firm does not know the optimal price, it has to experiment different (suboptimal) prices and update its belief about the underlying demand function. Therefore, the firm has to balance the exploration/exploitation tradeoff, which is usually referred to as the learning-and-earning problem in this line of literature. Our paper considers the pricing problem with personalized information and it does not consider the finite-inventory setting as in some of the papers mentioned above.

More recently, several papers investigate the pricing problem with unknown demand in the presence of covariates (Nambiar et al., 2016; Qiang and Bayati, 2016; Javanmard and Nazerzadeh, 2016; Cohen et al., 2016; Ban and Keskin, 2017). The existing literature has adopted a parametric approach: for example, the actual demand can be expressed in

¹Perchet and Rigollet (2013) study discrete decision variables (multi-armed bandit).

a linear form $\alpha^T \mathbf{x} + \beta^T \mathbf{x} \mathbf{p} + \epsilon$ (Qiang and Bayati, 2016; Ban and Keskin, 2017), where \mathbf{x} is the feature vector of a customer, α and β are vectorized coefficients, and ϵ is the random noise. Because of the parametric form, a key ingredient in the design of the algorithms in this line of literature is to plug in an estimator for the unknown parameters (α and β) in addition to some form of forced exploration. In contrast, we focus on a setting where the demand function cannot be parametrized. Thus, the firm cannot count on accurately estimating the function globally by estimating a few parameters. Instead, a localized optimal decision has to be made based on past covariates generated in the neighborhood. It highlights the different philosophies when designing algorithms for parametric/nonparametric learning problems with covariates. As a result, the best achievable regret deteriorates from $O(\sqrt{T})$ or $O(\log T)$ (parametric) to $O(T^{(2+d)/(4+d)})$ (nonparametric).

The dependence of the optimal rate of regret on the problem dimension d has been observed before. For example, Cohen et al. (2016) find a multi-dimensional binary search algorithm for feature-based dynamic pricing, which has regret $O(d^2 \log(T/d))$; Javanmard and Nazerzadeh (2016) propose a policy for a similar problem that achieves regret $O(s \log d \log T)$, where s represents the sparsity of the d features; in Ban and Keskin (2017), the near-optimal policy achieves regret $O(s\sqrt{T})$. In their parametric frameworks, the dependence of the regret on d is rather mild—it does not appear on the exponent of T; Javanmard and Nazerzadeh (2016); Keskin and Zeevi (2014) also provide methods to deal with the sparse strucutre. In contrast, in our nonparametric formulation, the optimal rate of regret $O(T^{(2+d)/(4+d)})$ increases dramatically in d, making the problem significantly harder to learn in high dimensions. This is similar to the nonparametric formulation in the network revenue management problem (Besbes and Zeevi, 2012), in which the dimension of the decision space is d and the optimal rate of regret is $O(T^{(2+d)/(3+d)})^2$. From the literature, it seems that the dimension significantly complicates the learning problem in a nonparametric formulation.

This paper is also related to the vast literature studying multi-armed bandit problems. See Cesa-Bianchi and Lugosi (2006); Bubeck and Cesa-Bianchi (2012) for a comprehensive survey. The classic multi-armed bandit problem involves finite arms, and the algorithms (Kuleshov and Precup, 2014; Agrawal and Goyal, 2012) cannot be applied

²It is shown in Chen and Gallego (2018) that if the number of inventory constraints $\ll d$, then learning the dual variables may effectively reduce the problem dimension.

directly to our setting. Recently, there is a stream of literature studying the so-called continuum-armed bandit problems (Agrawal, 1995; Kleinberg, 2005; Auer et al., 2007; Kleinberg et al., 2008; Bubeck et al., 2011), in which there are infinite number of arms (decisions). Although there is no contextual information in those papers, Kleinberg et al. (2008); Bubeck et al. (2011) have developed algorithms based on a similar idea to decision trees, because the potential arms form a high-dimensional space.

For multi-armed bandit problems with contextual information, parametric and regression-based algorithms have been proposed in, for example, Goldenshluger and Zeevi (2013); Bastani and Bayati (2015). Our paper is related to the literature studying contextual multi-armed bandit problems in a nonparametric framework Yang et al. (2002); Langford and Zhang (2008); Rigollet and Zeevi (2010); Perchet and Rigollet (2013); Slivkins (2014); Elmachtoub et al. (2017). The analysis builds upon the idea of adaptive binning in Perchet and Rigollet (2013). Our algorithm is designed for continuous decisions. In fact, applying the algorithm in Perchet and Rigollet (2013) designed for discrete decisions to our problem with simple discretization may result in worse-than-optimal regret. In terms of the formulation and the rate of regret, this paper is closely related to Slivkins (2014). Slivkins (2014) investigates a more general model, in which the decision p can be a vector, and assumes that $f(\mathbf{x}, p)$ is Lipschitz continuous. The optimal rate of regret in this setting has been shown to be $T^{(1+d_x+d_p)/(2+d_x+d_p)}$ in previous works, where d_x and d_p are the dimensions of x and p, respectively. Slivkins (2014) introduces an adaptive zooming algorithm, and uses the covering dimension of the space (\mathbf{x}, p) in the analysis. The extension accommodates more general spaces of (\mathbf{x}, p) than the Euclidean space. For Euclidean spaces, the algorithm recovers the optimal rate of regret $T^{(1+d_x+d_p)/(2+d_x+d_p)}$. In our setting, letting $d_x = d$ and $d_p = 1$ leads to the rate $T^{(2+d)/(3+d)}$.³ Motivated by personalized pricing, we impose additional assumptions on $f(\mathbf{x}, p)$ (smoothness and local concavity, see Assumption 3), and improve the optimal rate to $T^{(2+d)/(4+d)}$ as a result. The additional assumption may act as a special margin condition (Tsybakov et al., 2004; Rigollet and Zeevi, 2010; Perchet and Rigollet, 2013) and affects the optimal rate. It is unclear whether the algorithm in Slivkins (2014) could be adapted to accommodate the additional assumptions and achieve the improved rate. The design of our algorithm is based on adaptively partitioning the covariate space into

³Applying our assumptions and following Equation (8) of Slivkins (2014) give $d_x + d_p = d + 1/2$, and the regret improves to $T^{(1.5+d)/(2.5+d)}$.

rectangular bins, rather than overlapping balls as in Slivkins (2014).

2. Problem Formulation

Suppose the personalized demand function (purchasing probability) is $d(\mathbf{x}, p) \in [0, 1]$, where $\mathbf{x} \in [0, 1)^d$ is the observed feature vector, or covariate, summarizing the customer's personalized information, and p is the price set by the firm. Since the purchasing event is a Bernoulli random variable with success rate $d(\mathbf{x}, p)$ when the firm sets price p for a customer of covariate \mathbf{x} (henceforth abbreviated to customer \mathbf{x}), the expected revenue is thus $f(\mathbf{x}, p) \triangleq pd(\mathbf{x}, p)$. If the firm knew $d(\mathbf{x}, p)$, or equivalently, $f(\mathbf{x}, p)$, then it would set $p^*(\mathbf{x}) \triangleq \operatorname{argmax}_{p \in [0,1]} f(\mathbf{x}, p)$. Denote the optimal expected revenue from customer \mathbf{x} by $f^*(\mathbf{x}) \triangleq \max_{p \in [0,1]} f(\mathbf{x}, p)$. We will be primarily dealing with the expected revenue $f(\mathbf{x}, p)$ instead of the personalized demand $d(\mathbf{x}, p)$.

Initially, neither $d(\mathbf{x}, p)$ nor $f(\mathbf{x}, p)$ is known to the firm. In period $t \in \{1, 2, ..., T\}$, a customer arrives with covariate X_t . Upon observing X_t , the firm sets a price p_t . The revenue earned in period t is denoted by Z_t where Z_t/p_t is a Bernoulli random variable with mean $d(X_t, p_t)$, independent of everything else. The objective of the firm is to design a pricing policy to maximize the total revenue over the horizon $\sum_{t=1}^{T} E[Z_t] = \sum_{t=1}^{T} E[f(X_t, p_t)]$. Note that p_t itself is likely to be random even though the firm is not adopting a randomized policy. This is because the pricing decision made in period t may depend on the observed customers, set prices, and earned revenues in the previous periods. That is, $p_t = \pi_t(X_1, p_1, Z_1, \ldots, X_{t-1}, p_{t-1}, Z_{t-1}, X_t)$. Formally, we refer to π as the *pricing policy* that determines how p_t depends on the past information. We also denote $\mathcal{F}_t \triangleq \sigma(X_1, p_1, Z_1, \ldots, X_t, p_t, Z_t)$.

2.1. Regret

To measure the performance of a pricing policy, it is standard in the literature to benchmark it against the so-called *clairvoyant* policy and study the *regret*. Suppose $f(\mathbf{x}, p)$ is known to a clairvoyant firm. The optimal pricing policy is rather straightforward for a clairvoyant: having observed customer X_t , set $p^*(X_t)$ in period t and earn a random revenue with mean $f^*(X_t)$.

For the firm, the expected revenue in period *t* cannot exceed that of the clairvoyant:

 $f(X_t, p_t) \leq f^*(X_t)$. Thus, we define the regret of a pricing policy π to be the revenue gap

$$R_{\pi}(T) = \sum_{t=1}^{T} \mathsf{E} \left[(f^*(X_t) - f(X_t, p_t)) \right].$$

In period *t*, the expectation is taken with respect to the distribution of X_t as well as p_t , which itself depends on \mathcal{F}_{t-1} and X_t . Our goal is to design a policy π that achieves small $R_{\pi}(T)$ when $T \to \infty$.

However, because $R_{\pi}(T)$ also depends on the unknown function f, we require the designed policy to perform well for a family C of functions, i.e., we seek for optimal policies in terms of the minimax regret

$$\inf_{\pi} \sup_{f \in C} R_{\pi}(T).$$

Although it is usually impossible to find the exact policy that achieves the minimax regret, we focus on proposing a policy whose regret is at least comparable to (of the same order as) the minimax regret asymptotically when $T \rightarrow \infty$.

In this paper, we study functions f that cannot be parametrized. It implies that the family C is much larger than parametric families: it includes all the functions that satisfy some mild assumptions presented in the next section. In other words, the worst-case scenario can potentially be much worse than a parametric family. As a result, the achievable minimax regret is also higher.

2.2. Assumptions

In this section, we formally provide a set of assumptions that $f \in C$ and the stochastic process has to satisfy and their justifications.

Assumption 1. The covariates X_t are i.i.d. for t = 1, ..., T. Given X_t and p_t , the revenue Z_t is independent of everything else.

Both i.i.d. covariates and independent noise structure are standard in the literature.

Assumption 2. The functions $f(\cdot, p)$ and $f(\mathbf{x}, \cdot)$ are Lipschitz continuous given p and \mathbf{x} , i.e., there exists $M_1 > 0$ such that $|f(\mathbf{x}_1, p) - f(\mathbf{x}_2, p)| \le M_1 ||\mathbf{x}_1 - \mathbf{x}_2||_2$ and $|f(\mathbf{x}, p_1) - f(\mathbf{x}, p_2)| \le M_1 ||p_1 - p_2|$ for all \mathbf{x}_i and p_i (i = 1, 2) in the domain.

This assumption is equivalent to $|f(\mathbf{x}_1, p_1) - f(\mathbf{x}_2, p_2)| \le M_1(||\mathbf{x}_1 - \mathbf{x}_2||_2 + |p_1 - p_2|)$. Lipschitz continuity is a common assumption in the learning literature. In personalized pricing, it implies that the expected revenues are close if the firm charges similar prices for two customers with similar covariates. If this assumption fails, then the historical sales data of a certain type of customer is not informative for a new customer with almost identical background and learning is virtually impossible.

To introduce the next assumption, consider any hyper-rectangle $B \subset [0, 1)^d$, including a singleton $B = \{x\}$. Define $f_B(p) \triangleq \mathbb{E}[f(X, p)|X \in B]$ for $p \in [0, 1]$. Clearly $f_B(p)$ is the expected revenue when charging p for a customer that is sampled from a subset B.

Assumption 3. We assume that for any *B*,

- 1. The function $f_B(p)$ has a unique maximizer $p^*(B) \in [0, 1]$. Moreover, there exist uniform constants $M_2, M_3 > 0$ such that for all $p \in [0, 1], M_2(p^*(B) - p)^2 \leq f_B(p^*(B)) - f_B(p) \leq M_3(p^*(B) - p)^2$.
- 2. The maximizer $p^*(B)$ is inside the interval $[\inf\{p^*(\mathbf{x}) : \mathbf{x} \in B\}, \sup\{p^*(\mathbf{x}) : \mathbf{x} \in B\}]$.
- 3. Let d_B be the diameter of B. Then there exists a uniform constant $M_4 > 0$ such that $\sup\{p^*(\mathbf{x}) : \mathbf{x} \in B\} \inf\{p^*(\mathbf{x}) : \mathbf{x} \in B\} \le M_4 d_B$.

This assumption is quite different from those in the setting without covariates (Besbes and Zeevi, 2009; Wang et al., 2014; Lei et al., 2017) or the parametric setting (Ban and Keskin, 2017; Qiang and Bayati, 2016). To explain the intuition of $f_B(p)$, suppose the firm only observes $\mathbb{I}_{\{X \in B\}}$ but not the exact value of X, and thus cannot apply personalized pricing for customers in B. In this case, the learning objective is the expected revenue $f_B(p)$ and the clairvoyant policy that has the knowledge of $f(\mathbf{x}, p)$ is to set $p = p^*(B)$. This class of learning problems are important subroutines of the algorithm we propose and Assumption 3 guarantees that they can be effectively learned.

For part one of Assumption 3, we have

Proposition 1. If $f_B(p)$ is continuous for $p \in [0, 1]$, and twice differentiable in an open interval containing the unique global maximizer $p^*(B)$ with $f''_B(p^*(B)) < 0$, then part one of Assumption 3 holds.

Therefore, part one states that $f_B(p)$ is smooth and locally concave around the maximum. If *B* is a singleton, then it can be viewed as a weaker version of the concavity assumption in Wang et al. (2014); Lei et al. (2017), i.e., 0 > a > f''(p) > b for all *p* in their no-covariate setting. As a result, if $f_B(p)$ is the revenue function of linear or exponential demand, then part one is satisfied automatically.

Part two of Assumption 3 prevents the following scenario: If the optimal price for the aggregate demand of customers $\mathbf{x} \in B$, $p^*(B)$, is far from the optimal personalized pricing $p^*(\mathbf{x})$, then collecting more information for $f_B(p)$ does not help to improve the pricing decision for any individual customer $\mathbf{x} \in B$. Such obstacle may lead to failure to learn and is thus ruled out by the assumption. Similar types of assumptions have been imposed in other applications of revenue management. For example, Proposition 1.16 in Gallego and Topaloglu (2018) provides conditions under which the optimal price of the aggregated market lies in the convex hull formed by the optimal prices of each market segment when discriminatory pricing is allowed.

Part three imposes a continuity condition for the optimal price. It is equivalent to, for example, some form of continuous differentiability of $f(\mathbf{x}, p)$, because $p^*(\mathbf{x})$ solves the implicit function from the first-order condition $f_p(\mathbf{x}, p) = 0$.

Remark 1. Assumption 2 is a variant of similar assumptions adopted in the literature. Assumption 3, although appearing nonstandard, is also satisfied by the parametric families studied by previous works. We give a few examples that satisfy Assumption 3.

- Dynamic pricing with linear covariate (Qiang and Bayati, 2016): if $f(\mathbf{x}, p) = p(\boldsymbol{\theta}^T \mathbf{x} \alpha p)$, then $f_B(p) = p(\boldsymbol{\theta}^T \mathsf{E}[\mathbf{X}|\mathbf{X} \in B] \alpha p)$ and $p^*(B) = \boldsymbol{\theta}^T \mathsf{E}[\mathbf{X}|\mathbf{X} \in B]/2\alpha$.
- Separable function: consider $f(\mathbf{x}, p) = \sum_{i=1}^{k} g_i(\mathbf{x})h_i(p)$. Then $f_B(p) = \sum_{i=1}^{k} E[g_i(\mathbf{X})|\mathbf{X} \in B]h_i(p)$. If $h_i(p)$ are concave functions and $g_i(\mathbf{x})$ are positive, then we may be able to solve the unique maximizer $p^*(B) = E[g(\mathbf{X})|\mathbf{X} \in B]$ for some continuous function g.
- Localized functions: the covariate only plays a role in a subset B₀ ⊂ [0, 1)^d. See Section 5 for a concrete example.

As we shall see in Sections 4 and 5, the optimal rate of regret under Assumption 3 is $T^{(2+d)/(4+d)}$, in contrast to $T^{(2+d)/(3+d)}$ without it (taking $d_Y = 1$ in Equation (3) of Slivkins 2014). Technically, we suspect that Assumption 3 plays a similar rule to the margin

condition in the contextual bandit literature (Tsybakov et al., 2004; Goldenshluger et al., 2009; Rigollet and Zeevi, 2010; Perchet and Rigollet, 2013). However, because of the continuous decision variable studied in this paper, the margin condition cannot be translated in a straightforward way. It remains a future direction to present the assumption in a general form (the degree of smoothness such as the Hölder and Sobolev classes) and study how it affects the optimal rate of regret.

We summarize the information available to the firm. In the beginning of the horizon, the length of the horizon *T*, the dimension *d* and the constants $\{M_i\}_{i=1}^4$ are revealed⁴. In period *t*, the price can also depend on \mathcal{F}_{t-1} and X_t .

3. The ABE Algorithm

We next present a set of preliminary concepts related to the *bins* of the covariate space, and then introduce the proposed pricing policy: the Adaptive Binning and Exploration (ABE) algorithm.

3.1. Preliminary Concepts

Definition 1. A bin is a hyper-rectangle in the covariate space. More precisely, a bin is of the form

$$B = \{ \mathbf{x} : a_i \le x_i < b_i, i = 1, \dots, d \}$$

for $0 \le a_i < b_i \le 1, i = 1, \dots, d$.

We can *split* a bin *B* by bisecting it in all the *d* dimensions to obtain 2^d *child* bins of *B*, all of equal size. For a bin *B* with boundaries a_i and b_i for i = 1, ..., d, its children are indexed by $i \in \{0, 1\}^d$ and have the form

$$B_{i} = \left\{ \boldsymbol{x} : a_{j} \leq x_{j} < \frac{a_{j} + b_{j}}{2} \text{ if } \boldsymbol{i}_{j} = 0, \frac{a_{j} + b_{j}}{2} \leq x_{j} < b_{j} \text{ if } \boldsymbol{i}_{j} = 1, \ j = 1, \dots, d \right\}.$$

Denote the set of child bins of *B* by C(B). Conversely, for any $B' \in C(B)$, we refer to *B* as the *parent* bin of *B'*, denoted by P(B') = B.

⁴In fact, only M_2 is needed in the algorithm.

Our algorithm starts with a root bin $B_{\emptyset} \triangleq [0, 1)^d$, which contains all possible customers, and successively splits the bin as more data is collected. Therefore, any bin B produced during the process is the *offspring* of B_{\emptyset} , i.e., $P^{(k)}(B) = B_{\emptyset}$ for some k > 0, where $P^{(k)}$ is the *k*th composition of the parent function. Equivalently, B_{\emptyset} is an *ancestor* of B. For such a bin, we define its *level* to be k, denoted by l(B) = k. Conventionally, let $l(B_{\emptyset}) = 0$.

In the algorithm, we keep a dynamic partition \mathcal{P}_t of the covariate space consisting of the offspring of B_0 in each period t. The partition is mutually exclusive and collectively exhaustive, so $B_i \cap B_j = \emptyset$ for $B_i, B_j \in \mathcal{P}_t$, and $\bigcup_{B_i \in \mathcal{P}_t} B_i = B_0$. Initially $\mathcal{P}_0 = \{B_0\}$. In the algorithm, we gradually refine the partition; that is, each bin in \mathcal{P}_{t+1} has an ancestor (or itself) in \mathcal{P}_t .

3.2. Intuition

The intuition behind the ABE algorithm is to use a partition \mathcal{P}_t of the covariate space to aggregate customers in each period. It tries to find the optimal price for customers in each bin $B \in \mathcal{P}_t$, i.e., $p^*(B)$ defined in Section 2.2. As \mathcal{P}_t is refined dynamically, i.e., $B \in \mathcal{P}_t$ becomes smaller, such aggregation is almost identical to personalized pricing.

To do that, we keep a set of discrete prices (referred to as the *decision set* hereafter) for each bin in the partition. The decision set consists of equally spaced grid points of a price interval associated with the bin. When a customer arrives with covariate X_t inside a bin B, a price is chosen successively in the decision set and charged for the customer. The realized revenue for this price is recorded. When a large number of customers are observed in B, the average revenue for each price p in the decision set is close to $f_B(p)$, which is defined as $E[f(X, p)|X \in B]$ in Section 2.2. Therefore, the *empirically-optimal* price in the decision set is close to $p^*(B)$, with high confidence.

There are two potential pitfalls of this approach. First, the number of prices has an impact on the performance of the policy. If there are too many prices in a decision set, then for a given number of customers observed in the bin, each price is experimented with for a relatively few times. As a result, the confidence interval for the associated average revenue is wide. On the other hand, if there are too few prices, then inevitably the decision set has low resolutions. That is, the optimal price in the set could still be far from the true maximizer $p^*(B)$ because of the discretization error. We have to select

a proper size of the decision set to balance this trade-off.

Second, even if the optimal price $p^*(B)$ for the aggregate revenue in the bin is correctly identified, it may not be a strong indicator for $p^*(\mathbf{x})$ for a particular customer $\mathbf{x} \in B$. Indeed, $f_B(p)$ averages out all customers $X \in B$, and the optimal price for an individual customer \mathbf{x} could be very different. This obstacle, however, can be overcome as the size of *B* decreases, as implied by Assumption 3. In particular, part two and three of the assumption guarantee that when *B* is small, $p^*(\mathbf{x})$ is concentrated within a neighborhood of $p^*(B)$ as long as $\mathbf{x} \in B$. The cost of using a smaller bin, however, is the less frequency of observing a customer inside it.

To remedy the second pitfall, the algorithm adaptively refines the partition and decreases the size of the bins in \mathcal{P}_t as t increases. When a bin $B \in \mathcal{P}_t$ is large, the aggregate optimal price $p^*(B)$ is not a strong indicator for $p^*(\mathbf{x})$, $\mathbf{x} \in B$. As a result, we only need a rough estimate and split the bin when a relatively small number of customers are observed in B. When a bin $B \in \mathcal{P}_t$ is small, the optimal price $p^*(B)$ provides a strong indicator for $p^*(\mathbf{x})$, $\mathbf{x} \in B$. Therefore, we gather large sales data from customers $X \in B$ to explore the decision set and estimate $p^*(B)$ accurately, before it splits.

A crucial step in the algorithm is to determine what information to inherit when a bin is split into child bins. The ABE algorithm records the empirically-optimal price in the decision set of the parent bin. In the child bins, we use this information and set up their decision sets centered at it. As explained above, when the parent bin (and thus the child bins) is large, its optimal price does not predict those of the child bins well. Therefore, the algorithm sets up conservative decision sets for the child bins, i.e., they have wide intervals. On the other hand, when the parent bin is small, its optimal price provides an accurate indicator for those of the child bins. Thus, the algorithm constructs decision sets with narrow ranges for the child bins around the empirically-optimal price inherited.

3.3. Description of the Algorithm

In this section, we elaborate on the detailed steps of the ABE algorithm, shown in Algorithm 1.

The parameters for the algorithm include

Algorithm 1 Adaptive Binning and Exploration (ABE)

1: Input: T, d2: Constants: *M*₁, *M*₂, *M*₃, *M*₄ 3: Parameters: \bar{K} ; Δ_k , n_k , N_k for k = 0, ..., K4: Initialize: partition $\mathcal{P} \leftarrow \{B_{\emptyset}\}, p_l^{B_{\emptyset}} \leftarrow 0, p_u^{B_{\emptyset}} \leftarrow 1, \delta_{B_{\emptyset}} \leftarrow 1/(N_0 - 1), \bar{Y}_{B,j}, N_{B_{\emptyset},j} \leftarrow 0$ for $j = 0, ..., N_0 - 1$ 5: **for** t = 1 to T **do** Observe X_t 6: $B \leftarrow \{B \in \mathcal{P} : X_t \in B\}$ \triangleright The bin in the partition that X_t belongs to 7: $k \leftarrow l(B), N(B) \leftarrow N(B) + 1$ > Determine the level and update the number of 8: customers observed in B if k < K then \triangleright If not reaching the maximal level *K* 9: if $N(B) < n_k$ then ▶ If not enough data observed in *B* 10: $j \leftarrow N(B) - 1 \pmod{N_k}$ \triangleright Apply the *j*th price in the decision set 11: $p_t \leftarrow p_l^B + j\delta_B$; apply p_t and observe Z_t $\bar{Y}_{B,j} \leftarrow \frac{1}{N_{B,j}+1}(N_{B,j}\bar{Y}_{B,j} + Z_t), N_{B,j} \leftarrow N_{B,j} + 1$ 12: 13: else \triangleright If sufficient data observed in *B* 14: $j^* \in \operatorname{argmax}_{j \in \{0,1,\dots,N_k-1\}} \{ \bar{Y}_{B,j} \}, p^* \leftarrow p_l^B + j^* \delta_B$ ▶ Find the 15: empirically-optimal price; if there are multiple, choose any one of them $\mathcal{P} \leftarrow (\mathcal{P} \setminus B) \cup C(B) \triangleright$ Update the partition by removing *B* and adding 16: its children for $B' \in C(B)$ do Initialization for each child bin 17: $N(B') \leftarrow 0$ 18: $p_l^{B'} \leftarrow \max\{0, p^* - \Delta_{k+1}/2\}; p_u^{B'} \leftarrow \min\{1, p^* + \Delta_{k+1}/2\}$ ▶ The 19: range of the decision set $\delta_{B'} \leftarrow (p_u^{B'} - p_l^{B'})/(N_{k+1} - 1)$ \triangleright The grid size of the decision set 20: $N_{B',j}, \bar{Y}_{B',j} \leftarrow 0$, for $j = 0, \dots, N_{k+1} - 1$ ▶ Initialize the average 21: revenue and number of customers for each price end for 22: end if 23: ▶ If reaching the maximal level else 24: $p_t \leftarrow (p_t^B + p_u^B)/2$ 25: end if 26: 27: end for

- 1. *K*, the maximal level of the bins. When a bin is at level *K*, the algorithm no longer splits it and simply applies the median price of its decision set whenever a customer is observed in it.
- 2. Δ_k , the length of the interval that contains the decision set of level-*k* bins.
- n_k, the maximal number of customers observed in a level-k bin in the partition.
 When n_k customers are observed, the bin splits.
- 4. N_k , the number of prices to explore in the decision set of level-*k* bins. The decision set of bin *B* consists of equally spaced grid points of an interval $[p_l^B, p_u^B]$, to be adaptively specified by the algorithm.

We initialize the partition to include only the root bin B_{\emptyset} in Step 4. Its decision set spans the whole interval [0, 1] with N_0 equally spaced grid points. That is, the *j*th price is $j\delta_{B_{\emptyset}} \triangleq j/(N_0 - 1)$ for $j = 0, ..., N_0 - 1$. The initial average revenue and the number of customers that are charged the *j*th price are set to $\bar{Y}_{B_{\emptyset},j} = N_{B_{\emptyset},j} = 0$.

Suppose the partition is \mathcal{P}_t at t and a customer X_t is observed (Step 6). The algorithm determines the bin $B \in \mathcal{P}_t$ which the customer belongs to. The counter N(B) records the number of customers already observed in B up to t when B is in the partition (Step 8). If the level of B is l(B) = k < K (i.e., B is not at the maximal level) and the number of customers observed in B is not sufficient (Step 9 and Step 10), then the algorithm has assigned a decision set to the bin in previous steps, namely, $\{p_l^B + j\delta_B\}$ for $j = 0, \ldots, N_k - 1$. There are N_k prices in the set and they are equally spaced in the interval $[p_l^B, p_u^B]$. They are explored successively as new customers are observed in B (explore p_l^B for the first customer observed in B, $p_l^B + \delta_B$ for the second customer, $\ldots, p_l^B + (N_k - 1)\delta_B$ for the N_k th customer, p_l^B again for the $(N_k + 1)$ th customer, etc.). Therefore, the algorithm charges price $p_t = p_l^B + j\delta_B$ where $j = N(B) - 1 \pmod{N_k}$ for the N(B)th customer observed in B (Step 11). Then, Step 13 updates the average revenue and the number of customers for the jth price.

If the level of *B* is l(B) = k < K and we have observed a sufficient number of customers in *B* (Step 9 and Step 14), then the algorithm splits *B* and replaces it by its 2^d child bins in the partition (Step 16). For each child bin, Step 18 to Step 21 initialize the counter, the interval that encloses the decision set, the grid size of the decision set, and the average revenue/number of customers for each price in the decision set,

respectively. In particular, to construct the decision set of a child bin, the algorithm first computes the empirically-optimal price in the decision set of the parent bin *B*; that is, $j^* \in \operatorname{argmax}_{j \in \{0,1,\ldots,N_k-1\}}{\{\bar{Y}_{B,j}\}}$ in Step 15. Then, the algorithm creates an interval centered at this empirically-optimal price with width Δ_{k+1} , properly cut off by the boundaries [0, 1]. The decision set is then an equally spaced grid of the above interval (Step 19 and Step 20).

If the level of *B* is already *K*, then the algorithm simply charges the median price (Step 25) repeatedly without further exploration. For such a bin, its size is sufficiently small and the algorithm has narrowed the range of the decision set *K* times. The charged price is close enough to all $p^*(\mathbf{x})$, $\mathbf{x} \in B$, with high probability.

3.4. Choice of Parameters

We set $K = \lfloor \frac{\log(T)}{(d+4)\log(2)} \rfloor$, $\Delta_k = 2^{-k}\log(T)$, $N_k = \lceil \log(T) \rceil$, and

$$n_k = \max\left\{0, \left\lceil \frac{2^{4k+15}}{M_2^2 \log^3(T)} (\log(T) + \log(\log(T)) - (d+2)k \log(2)) \right\rceil \right\}.$$

To give a sense of their magnitudes, the edge length of the bins at the maximal level is approximately $T^{-1/(d+4)}$. The range of the decision set (Δ_k) is proportional to the edge length of the bin (2^{-k}) . The number of prices in a decision set is approximately $\log(T)$. Therefore, the grid size is $\delta_B \approx 2^{-k}$ for a level-k. The number of customers to observe in a level-k bin B is roughly $n_k \approx 2^{4k}/\log(T)^2$ before it splits. When k is small, n_k can be zero according to the expression. In this case, the algorithm immediately splits the bin without collecting any sales data in it.

3.5. A Schematic Illustration

We illustrate the key steps of the algorithm by an example with d = 2. Figure 1 illustrates a possible outcome of the algorithm in periods $t_1 < t_2 < t_3$ (top panel, mid panel, and bottom panel respectively). Up until period t_1 , there is a single bin and the covariates of observed customers X_t for $t \le t_1$ are illustrated in the top left panel. In this case, the decision set associated with the root bin is $p \in \{0.1, 0.2, ..., 0.9\}$, illustrated by the top right panel. The average revenue $\bar{Y}_{B,j}$ of each price is recorded, and $p^* = 0.6$ is the empirically-optimal price. At t_1 + 1, a sufficient number of customers are observed and Step 14 is triggered in the algorithm. Therefore, the bin is split into four child bins.

From period $t_1 + 1$ to t_2 , new customers are observed in each child bin (mid left panel). Note that the customers observed before t_1 in the parent bin are no longer used and colored in gray. For each child bin (the bottom-left bin is abbreviated to BL, etc.), the average revenues for the prices in the decision sets is demonstrated in the mid right panel. The decision sets are centered at the empirically-optimal price of their parent bin, which is $p^* = 0.6$ from the top right panel. They have narrower ranges and finer grids than that of the parent bin. At $t_2 + 1$, a sufficient number of customers are observed in BL, and it is split into four child bins.

From period $t_2 + 1$ to t_3 , the partition consists of seven bins, as shown in the bottom left panel. The BR, TL and TR bins keep observing customers and updating the average revenues, because they have not collected sufficient data. Their status at t_3 is shown in the bottom panels. In the four newly created child bins of BL (the bottom-left bin of BL is abbreviated to BL-BL, etc.), the prices in the decision sets are used successively and their average revenues are illustrated in the bottom right panel.

4. Regret Analysis: Upper Bound

To measure the performance of the ABE algorithm, we provide an upper bound for its regret.

Theorem 1. For any function f satisfying Assumption 2 and 3, the regret incurred by the ABE algorithm is bounded by

$$R_{\pi_{ABE}}(T) \le CT^{\frac{2+d}{4+d}}\log(T)^2$$

for a constant C > 0 that is independent of T.

We provide a sketch of the proof here and present the details in the appendix. In period *t*, if $X_t \in B$ for a bin in the partition $B \in \mathcal{P}_t$, then the expected regret incurred by the ABE algorithm is $E[(f^*(X_t) - f(X_t, p_t))]\mathbb{I}_{\{X_t \in B, B \in \mathcal{P}_t\}}]$. Since the total regret simply sums up the above quantity over t = 1, ..., T and all possible *Bs*, it suffices to focus on the regret for given *t* and *B*. Two possible scenarios can arise: (1) the optimal price



Figure 1: A schematic illustration of the ABE algorithm.

of the aggregate demand in *B*, i.e., $p^*(B)$, is inside the range of the decision set, i.e., $p^*(B) \in [p_l^B, p_u^B]$ (Step 19); (2) the optimal price $p^*(B)$ is outside the range of the decision set.

Scenario one represents the regime where the algorithm is working "normally": up until *t*, the algorithm has successfully narrowed the optimal price $p^*(B)$ (which provides a useful indicator for all $p^*(\mathbf{x})$, $\mathbf{x} \in B$ when *B* is small) down to $[p_l^B, p_u^B]$. By Assumption 3 part one, the regret in this scenario can be decomposed into two terms

$$f^*(X_t) - f(X_t, p_t) \le M_3(|p_t - p^*(B)| + |p^*(B) - p^*(X_t)|)^2$$

The first term can be bounded by the length of the interval $p_u^B - p_l^B$. The second term can be bounded by the size of *B* given $X_t \in B$ by Assumption 3 part two and three. By the choice of parameters in Section 3.4, the length of the interval decreases as the bin size decreases. Therefore, both terms can be well controlled when the size of *B* is sufficiently small, or equivalently, when the level l(B) is sufficiently large. This is why a properly chosen n_k can guarantee that the algorithm spends little time for large bins and collect a large amount of data for small bins. When the bin level reaches *K*, the above two terms are small enough and no more exploration is needed.

Scenario two represents the regime where the algorithm works "abnormally". In scenario two, the difference $f^*(X_t) - f(X_t, p_t)$ can no longer be controlled as in scenario one because p_t and $p^*(X_t)$ can be far apart. To make things worse, $p^*(B) \notin [p_l^B, p_u^B]$ usually implies $p^*(B') \notin [p_l^{B'}, p_u^{B'}]$, where B' is a child of B. This is because (1) $p^*(B)$ is close to $p^*(B')$ for small B, and (2) $[p_l^{B'}, p_u^{B'}]$ is created around the empirically-optimal price for B, and thus overlapping with $[p_l^B, p_u^B]$. Therefore, for any period s following t, the worst-case regret is O(1) in that period if $X_s \in B$ or its offspring.

To bound the regret in scenario two, we have to bound the probability, which requires delicate analysis of the events. If scenario two occurs for *B*, then during the process that we sequentially split B_0 to obtain *B*, we can find an ancestor bin of *B* (which can be *B* itself) that scenario two happens for the first time along the "branch" from B_0 all the way down to *B*. More precisely, denoting the ancestor bin by B_a and its parent by $P(B_a)$, we have (1) $p^*(P(B_a))$ is inside $[p_l^{P(B_a)}, p_u^{P(B_a)}]$ (scenario one); (2) after $P(B_a)$ is split, $p^*(B_a)$ is outside $[p_l^{B_a}, p_u^{B_a}]$ (scenario two). Denote the empirically-optimal price in the decision set of $P(B_a)$ by p^* . For such an event to occur, the center of the decision set of B_a , which is p^* , has to be at least $\Delta_{l(B_a)}/2$ away from $p^*(B_a)$.⁵ Because of Assumption 3 and the choice of Δ_k , the distance between $p^*(P(B_a))$ and $p^*(B_a)$ is relatively small compared to $\Delta_{l(B_a)}$. Therefore, the empirically-optimal price p^* must be far away from $p^*(P(B_a))$. The probability of such event can be bounded using classic concentration inequalities for sub-Gaussian random variables: the prices that are closer to $p^*(P(B_a))$ and thus have higher means turn out to generate lower average revenue than p^* ; this event is extremely unlikely to happen when we have collected a large amount of sales data for each price in the decision set.

The total regret aggregates those in scenario one and two for all possible combinations of *B* and *B_a*. It matches the quantity $O(\log(T)^2 T^{(2+d)/(4+d)})$ presented in the theorem.

5. Regret Analysis: Lower Bound

In this section, we show that the minimax regret is no lower than $cT^{(2+d)/(4+d)}$ for some constant *c*. Combining with the last section, we conclude that no non-anticipating policy does better than the ABE algorithm in terms of the order of magnitude of the regret in *T* (neglecting logarithmic terms).

We first construct a family of functions that satisfy Assumption 2 and 3. The functions in the family are selected to be "difficult" to distinguish. By doing so, we will prove that any policy has to spend a substantial amount of time exploring prices that generate low revenues but help to differentiate the functions. Otherwise, the incapability to correctly identify the underlying function is costly in the long run. Therefore, unable to contain both sources of regret at the same time, no policy can achieve lower regret than the quantity stated in Theorem 2.

Before introducing the family of functions, we define ∂B to be the boundary of a convex set in $[0, 1)^d$. Let $D(B_1, B_2)$ be the Euclidean distance between two sets B_1 and B_2 . That is $D(B_1, B_2) \triangleq \inf \{ \| \mathbf{x}_1 - \mathbf{x}_2 \|_2 : \mathbf{x}_1 \in B_1, \mathbf{x}_2 \in B_2 \}$. We allow B_1 or B_2 to be a singleton. To define *C*, we partition the covariate space $[0, 1)^d$ into M^d equally sized

⁵Recall that $p_u^{B_a} - p_l^{B_a} = \Delta_{l(B_a)}$.

bins. That is, each bin has the following form: for $(k_1, \ldots, k_d) \in \{1, \ldots, M\}^d$,

$$\left\{ \boldsymbol{x} : \frac{k_i - 1}{M} \leq x_i < \frac{k_i}{M}, \quad \forall \ i = 1, \dots, d \right\}.$$

We number those bins by $1, \ldots, M^d$ in an arbitrary order, i.e., B_1, \ldots, B_{M^d} . Each function $f(\mathbf{x}, p) \in C$ is indexed by a tuple $w \in \{0, 1\}^{M^d}$, whose *j*th index determines the behavior of $f_w(\mathbf{x}, p)$ in B_j . More precisely, for $\mathbf{x} \in B_j$, the personalized demand function is

$$d_{w}(\mathbf{x}, p) = \begin{cases} \frac{2}{3} - \frac{p}{2} & w_{j} = 0\\ \frac{2}{3} - \frac{p}{2} + \left(\frac{1}{3} - \frac{p}{2}\right) D(\mathbf{x}, \partial B_{j}) & w_{j} = 1 \end{cases}$$

and thus

$$f_{w}(\mathbf{x}, p) = \begin{cases} p\left(\frac{2}{3} - \frac{p}{2}\right) & w_{j} = 0\\ p\left(\frac{2}{3} - \frac{p}{2} + \left(\frac{1}{3} - \frac{p}{2}\right)D(\mathbf{x}, \partial B_{j})\right) & w_{j} = 1 \end{cases}$$

The optimal personalized price for customer $\mathbf{x} \in B_j$ is $p^*(\mathbf{x}) = 2/3$ if $w_j = 0$ and $p^*(\mathbf{x}) = \frac{2+D(\mathbf{x},\partial B_j)}{3(1+D(\mathbf{x},\partial B_j))}$ if $w_j = 1$.

The construction of *C* follows a similar idea to Rigollet and Zeevi (2010). For a given $f_w \in C$, we can always find another function $f_{w'} \in C$ that only differs from *f* in a single bin B_j by setting w' to be equal to *w* except for the *j*th index. The firm can only rely on the covariates generated in B_j to distinguish between f_w and $f_{w'}$. For small bins (i.e., large *M*), this is particularly costly because there are only a tiny fraction of customers observed in a particular bin and the difference $|f_w - f_{w'}| = p(1/3 - p/2)D(\mathbf{x}, \partial B_j)$ becomes tenuous. It also requires *p* to be far from 2/3 to detect the difference, which happens to be the optimal price when $w_j = 0$. This makes the exploration/exploitation trade-off hard to balance. Now a policy has to carry out the task for M^d bins, i.e., distinguishing the underlying function f_w with M^d tuples that only differ from *w* in one index. The cost is inevitable and adds to the lower bound of the regret. Moreover, we assume the customers *X* are uniformly distributed in $[0, 1)^d$.

In the appendix, we show that the constructed C satisfies all the assumptions. The main theorem below shows the lower bound for the regret.

Theorem 2. For the constructed C, any non-anticipating policy π has regret

$$\sup_{f \in C} R_{\pi}(T) \ge cT^{\frac{2+\alpha}{4+\alpha}}$$

for a constant c > 0.

6. Future Research

As shown in Theorem 1 and Theorem 2, the best achievable regret of the problem is of order $T^{(2+d)/(4+d)}$. As a result, the knowledge of the sparsity structure of the covariate is essential in designing pricing policies. More precisely, the provided customer covariate is of dimension *d*, while the personalized demand d(X, p) may only depend on *d'* entries of the covariate where $d' \ll d$. In this case, being able to identify the *d'* entries out of *d* significantly decreases the incurred regret from $T^{(2+d)/(4+d)}$ to $T^{(2+d')/(4+d')}$. Indeed, in the ABE algorithm, if the sparsity structure is known, then a bin is split into $2^{d'}$ instead of 2^d child bins. It pools the observations that only differ in the dimensions corresponding to the redundant covariates so that more observations are available in a bin, and thus substantially reduces the exploration cost. An important research question is then whether it is possible to design a binning algorithm that selects one dimension and the position to split, based on a certain criterion, like regression/classification decision trees (Hastie et al., 2001). This may significantly improve the regret in the presence of sparse covariates.

References

- Agrawal, R. (1995). The continuum-armed bandit problem. *SIAM journal on control and optimization 33*(6), 1926–1951.
- Agrawal, S. and N. Goyal (2012). Analysis of thompson sampling for the multi-armed bandit problem. In *Conference on Learning Theory*, pp. 39–1.
- Araman, V. F. and R. Caldentey (2009). Dynamic pricing for nonperishable products with demand learning. *Operations research* 57(5), 1169–1188.

- Auer, P., R. Ortner, and C. Szepesvári (2007). *Improved Rates for the Stochastic Continuum-Armed Bandit Problem*, pp. 454–468. Berlin, Heidelberg: Springer Berlin Heidelberg.
- Ban, G. and N. B. Keskin (2017). Personalized dynamic pricing with machine learning. *Working paper*.
- Bastani, H. and M. Bayati (2015). Online decision-making with high-dimensional covariates. *Working paper*.
- Besbes, O. and A. Zeevi (2009). Dynamic pricing without knowing the demand function: Risk bounds and near-optimal algorithms. *Operations Research* 57(6), 1407–1420.
- Besbes, O. and A. Zeevi (2012). Blind network revenue management. *Operations* research 60(6), 1537–1550.
- Broder, J. and P. Rusmevichientong (2012). Dynamic pricing under a general parametric choice model. *Operations Research 60*(4), 965–980.
- Bubeck, S. and N. Cesa-Bianchi (2012). Regret analysis of stochastic and nonstochastic multi-armed bandit problems. *Foundations and Trends*® *in Machine Learning 5*(1), 1–122.
- Bubeck, S., R. Munos, G. Stoltz, and C. Szepesvári (2011). X-armed bandits. Journal of Machine Learning Research 12(May), 1655–1695.
- Cesa-Bianchi, N. and G. Lugosi (2006). *Prediction, learning, and games.* Cambridge university press.
- Chen, N. and G. Gallego (2018). A primal-dual learning algorithm for personalized dynamic pricing with an inventory constraint. *Working paper*.
- Cheung, W. C., D. Simchi-Levi, and H. Wang (2017). Dynamic pricing and demand learning with limited price experimentation. *Operations Research 65*(6), 1722–1731.
- Cohen, M. C., I. Lobel, and R. Paes Leme (2016). Feature-based dynamic pricing. *Working paper*.

- den Boer, A. V. (2015). Dynamic pricing and learning: historical origins, current research, and new directions. *Surveys in operations research and management science 20*(1), 1–18.
- den Boer, A. V. and B. Zwart (2014). Simultaneously learning and optimizing using controlled variance pricing. *Management Science 60*(3), 770–783.
- Elmachtoub, A. N., R. McNellis, S. Oh, and M. Petrik (2017). A practical method for solving contextual bandit problems using decision trees. *Working paper*.
- Farias, V. F. and B. Van Roy (2010). Dynamic pricing with a prior on market response. *Operations Research 58*(1), 16–29.
- Foucart, S. and H. Rauhut (2013). *A mathematical introduction to compressive sensing*, Volume 1. Birkhäuser Basel.
- Gallego, G. and H. Topaloglu (2018). Revenue management and pricing analytics. In preparation.
- Goldenshluger, A. and A. Zeevi (2013). A linear response bandit problem. *Stochastic Systems 3*(1), 230–261.
- Goldenshluger, A., A. Zeevi, et al. (2009). Woodroofe's one-armed bandit problem revisited. *The Annals of Applied Probability 19*(4), 1603–1633.
- Hastie, T., R. Tibshirani, and J. Friedman (2001). *The Elements of Statistical Learning*. Springer Series in Statistics. New York, NY, USA: Springer New York Inc.
- Javanmard, A. and H. Nazerzadeh (2016). Dynamic pricing in high-dimensions. *Working paper*.
- Keskin, N. B. and A. Zeevi (2014). Dynamic pricing with an unknown demand model: Asymptotically optimal semi-myopic policies. *Operations Research 62*(5), 1142–1167.
- Kleinberg, R., A. Slivkins, and E. Upfal (2008). Multi-armed bandits in metric spaces. In *Proceedings of the fortieth annual ACM symposium on Theory of computing*, pp. 681–690. ACM.

- Kleinberg, R. D. (2005). Nearly tight bounds for the continuum-armed bandit problem. In *Advances in Neural Information Processing Systems*, pp. 697–704.
- Kuleshov, V. and D. Precup (2014). Algorithms for multi-armed bandit problems. *Working paper*.
- Langford, J. and T. Zhang (2008). The epoch-greedy algorithm for multi-armed bandits with side information. In *Advances in neural information processing systems*, pp. 817–824.
- Lei, Y., S. Jasin, and A. Sinha (2017). Near-optimal bisection search for nonparametric dynamic pricing with inventory constraint. *Working paper*.
- Nambiar, M., D. Simchi-Levi, and H. Wang (2016). Dynamic learning and price optimization with endogeneity effect. *Working paper*.
- Perchet, V. and P. Rigollet (2013). The multi-armed bandit problem with covariates. *The Annals of Statistics* 41(2), 693–721.
- Qiang, S. and M. Bayati (2016). Dynamic pricing with demand covariates. *Working paper*.
- Rigollet, P. and A. Zeevi (2010). Nonparametric bandits with covariates. In A. T. Kalai and M. Mohri (Eds.), *COLT*, pp. 54–66. Omnipress.
- Slivkins, A. (2014). Contextual bandits with similarity information. *The Journal of Machine Learning Research 15*(1), 2533–2568.
- Tsybakov, A. B. (2009). *Introduction to Nonparametric Estimation* (1 ed.). Springer-Verlag New York.
- Tsybakov, A. B. et al. (2004). Optimal aggregation of classifiers in statistical learning. *The Annals of Statistics 32*(1), 135–166.
- Wang, Z., S. Deng, and Y. Ye (2014). Close the gaps: A learning-while-doing algorithm for single-product revenue management problems. *Operations Research 62*(2), 318–331.
- Yang, Y., D. Zhu, et al. (2002). Randomized allocation with nonparametric estimation for a multi-armed bandit problem with covariates. *The Annals of Statistics 30*(1), 100–121.

Online Appendix for Nonparametric Pricing Analytics with Customer Covariates

A. Table of Notations

$\lceil x \rceil$	The smallest integer that does not exceed x
$(x)^+$	The positive part of <i>x</i>
#{}	The cardinality of a set
\mathcal{F}_t	The σ -algebra generated by $(X_1, \pi_1, Z_1, \ldots, X_{t-1}, \pi_{t-1}, Z_{t-1})$
μ_X	The distribution of the covariate X over $[0, 1)^d$
$P^*(B)$	$\operatorname{argmax}_{p \in [0,1]} \{ E[f(X,p) X \in B] \}$
P_B^*	The empirically-optimal decision in the decision set for <i>B</i>
∂B	The boundary of <i>B</i>

Table 1: A table of notations used in the paper.

B. Proofs

_

Proof of Proposition 1: Define for $p \in [0, 1]$

$$g(p) = \begin{cases} \frac{f_B(p^*(B)) - f_B(p)}{(p^*(B) - p)^2} & p \neq p^*(B) \\ -\frac{f_B''(p^*(B))}{2} & p = p^*(B). \end{cases}$$

By L'Hopital's rule, g(p) is continuous at $p^*(B)$. In addition, because $f_B(p)$ is continuous, g(p) is continuous for all $p \in [0, 1]$. By Weierstrass's extreme value theorem, we have $g(p) \in [M_2, M_3]$ and both M_2 and M_3 are attained. By the definition and the uniqueness of the maximizer, g(p) > 0 for $p \in [0, 1]$. Therefore, we must have $M_2, M_3 > 0$. This establishes the result.

B.1. Upper Bound in Section 4

We first introduce the following lemmas.

Lemma 1. A Bernoulli random variable X satisfies

$$\mathsf{E}[\exp(t(X - \mathsf{E}[X])] \le \exp(t^2/8)$$

This lemma follows directly from the Hoeffding's lemma. It implies that a Bernoulli random variable is sub-Gaussian.

Lemma 2. Suppose for given \mathbf{x} and p, the random variable $Z(\mathbf{x}, p)$ is sub-Gaussian with parameter σ , i.e.,

$$\mathsf{E}[\exp(t(Z - \mathsf{E}[Z])] \le \exp(\sigma t^2)$$

for all $t \in \mathbb{R}$. Then the distribution of Z(X, p) conditional on $X \in B$ for a set B is still sub-Gaussian with the same parameter.

Proof. Let μ_X denote the distribution of *X*. We have that for all $t \in \mathbb{R}$

$$\begin{split} \mathsf{E}[\exp(t(Z(X,p)-\mathsf{E}[Z(X,p)|X\in B])|X\in B]) \\ &= \frac{\int_{B}\mathsf{E}[\exp(t(Z(\mathbf{x},p)-\mathsf{E}[Z(X,p)|X\in B]d\mu_{X}(\mathbf{x}))}{\int_{B}d\mu_{X}(\mathbf{x})} \\ &= \frac{\int_{B}\mathsf{E}[\exp(t(Z(\mathbf{x},p)-\mathsf{E}[Z(\mathbf{x},p)])]d\mu_{X}(\mathbf{x})}{\int_{B}d\mu_{X}(\mathbf{x})} \times \frac{\int_{B}\exp(t\mathsf{E}[Z(\mathbf{x},p)])d\mu_{X}(\mathbf{x})}{\exp(t\mathsf{E}[Z(X,p)|X\in B])\int_{B}d\mu_{X}(\mathbf{x})} \\ &\leq \frac{\int_{B}\exp(\sigma t^{2})d\mu_{X}(\mathbf{x})}{\int_{B}d\mu_{X}(\mathbf{x})} \times 1 = \exp(\sigma t^{2}), \end{split}$$

where the last inequality is by the definition of conditional expectations. Hence the result is proved.

Proof of Theorem 1: According to the algorithm (Step 7), let \mathcal{P}_t denote the partition formed by the bins at time *t* when X_t is generated. The regret associated with X_t can be counted by bins $B \in \mathcal{P}_t$ into which X_t falls. Meanwhile, the level of *B* is at most *K*. Therefore,

$$R_{\pi_{ABE}}(T) = \mathsf{E}\left[\sum_{t=1}^{T} (f^{*}(X_{t}) - f(X_{t}, p_{t}))\right] = \mathsf{E}\left[\sum_{t=1}^{T} \sum_{B \in \mathcal{P}_{t}} (f^{*}(X_{t}) - f(X_{t}, p_{t}))\mathbb{I}_{\{X_{t} \in B\}}\right]$$
$$= \mathsf{E}\left[\sum_{t=1}^{T} \sum_{k=0}^{K} \sum_{\{B: l(B)=k\}} (f^{*}(X_{t}) - f(X_{t}, p_{t}))\mathbb{I}_{\{X_{t} \in B, B \in \mathcal{P}_{t}\}}\right]$$

We will define the following random event for each bin *B*:

$$E_B = \left\{ p^*(B) \in \left[p_l^B, p_u^B \right] \right\}.$$

Recall that $p^*(B)$ is the unique maximizer for $f_B(p) = \mathsf{E}[f(X, p)|X \in B]$ by Assumption 3; $[p_l^B, p_u^B]$ is the range of the decision set to explore for *B*. According to Step 19 of the ABE algorithm, the interval $[p_l^B, p_u^B]$ is constructed around $p_{P(B)}^*$, the empirically-optimal price of the parent bin P(B) that maximizes the empirical average $\bar{Y}_{P(B),j}$.

We will decompose the regret depending on whether E_B occurs.

$$E[R_{\pi_{ABE}}] = E\left[\sum_{t=1}^{T}\sum_{k=0}^{K}\sum_{\{B:l(B)=k\}} (f^{*}(X_{t}) - f(X_{t}, p_{t}))\mathbb{I}_{\{X_{t}\in B, B\in\mathcal{P}_{t}, E_{B}^{c}\}}\right]$$

$$+ E\left[\sum_{t=1}^{T}\sum_{k=0}^{K}\sum_{\{B:l(B)=k\}} (f^{*}(X_{t}) - f(X_{t}, p_{t}))\mathbb{I}_{\{X_{t}\in B, B\in\mathcal{P}_{t}, E_{B}\}}\right]$$

$$(1)$$

$$\operatorname{term 2}$$

We first analyze term 1. Because E_{B_0} is always true $([p_l^{B_0}, p_u^{B_0}] = [0, 1]$ always encloses $p^*(B_0)$ according to Step 4), we can find an ancestor of B, say B_a (which can be B itself), such that $E_{B_a}^c \cap E_{P(B_a)} \cap E_{P(P(B_a))} \cap \ldots \cap E_{B_0}$ occurs. In other words, up until B_a , the algorithm always correctly encloses the optimal price $p^*(P^{(k)}(B_a))$ of the ancestor bin of B_a in the intervals $[p_l^{P^{(k)}(B_a)}, p_u^{P^{(k)}(B_a)}]$. Therefore, when E_B^c occurs, we can rearrange the event by such B_a . Term 1 in (1) can be bounded by

$$\mathbf{E}\left[\sum_{t=1}^{T}\sum_{k=0}^{K}\sum_{\{B:l(B)=k\}}(f^{*}(\mathbf{X}_{t})-f(\mathbf{X}_{t},p_{t}))\mathbb{I}_{\{\mathbf{X}_{t}\in B,B\in\mathcal{P}_{t},E_{B}^{c}\}}\right] \\
\leq \sum_{t=1}^{T}\sum_{k=0}^{K}\sum_{\{B:l(B)=k\}}M_{1}\mathsf{P}(\mathbf{X}_{t}\in B,B\in\mathcal{P}_{t},E_{B}^{c}) \\
= \sum_{t=1}^{T}\sum_{k=1}^{K}\sum_{\{B:l(B)=k\}}M_{1}\mathsf{P}\left(\mathbf{X}_{t}\in B,B\in\mathcal{P}_{t},E_{B_{a}}^{c}\cap E_{P(B_{a})}\cap E_{P(P(B_{a}))}\cap\ldots\cap E_{B_{0}}\right) \\
\leq \sum_{t=1}^{T}\sum_{k=1}^{K}\sum_{\{B:l(B)=k\}}M_{1}\mathsf{P}(\mathbf{X}_{t}\in B,B\in\mathcal{P}_{t},E_{B_{a}}^{c}\cap E_{P(B_{a})}) \\
= \sum_{t=1}^{T}\sum_{k=1}^{K}\sum_{\{B:l(B)=k\}}\sum_{k'=1}^{K}\sum_{B':l(B')=k'}M_{1}\mathsf{P}(\mathbf{X}_{t}\in B,B\in\mathcal{P}_{t},B_{a}=B',E_{B'}^{c}\cap E_{P(B')}) \quad (2)$$

The first inequality is due to Assumption 2. In the second inequality, we start enumerating from k = 1 instead of k = 0 because $E_{B_0}^c$ never occurs. In the last equality, we rearrange the probabilities by counting the deterministic bins B' instead of the random bins B_a .

Now note that $\{X_t \in B, B \in \mathcal{P}_t, B_a = B', E_{B'}^c \cap E_{P(B')}\}$ are exclusive for different *Bs* because \mathcal{P} is a partition and X_t can only fall into one bin. Moreover, $\{X_t \in B, B \in \mathcal{P}_t, B_t \in \mathcal{P}_t, B_t \in \mathcal{P}_t\}$

$$\mathcal{P}_t, B_a = B', E_{B'}^c \cap E_{P(B')} \} \subset \left\{ X_t \in B', E_{B'}^c \cap E_{P(B')} \right\} \text{ because } B \subset B_a. \text{ Therefore,}$$

$$\sum_{k=1}^{K} \sum_{\{B:l(B)=k\}} \mathsf{P}\left(X_{t} \in B, B \in \mathcal{P}_{t}, B_{a} = B', E_{B'}^{c} \cap E_{P(B')}\right) \le \mathsf{P}\left(X_{t} \in B', E_{B'}^{c} \cap E_{P(B')}\right).$$

Thus, we can further simplify (2):

$$(2) \leq \sum_{t=1}^{T} \sum_{k=1}^{K} \sum_{\{B': l(B')=k\}} M_1 P\left(X_t \in B', E_{B'}^c \cap E_{P(B')}\right)$$
$$= \sum_{t=1}^{T} \sum_{k=1}^{K} \sum_{\{B: l(B)=k\}} M_1 P(X_t \in B) P(E_B^c \cap E_{P(B)}).$$
(3)

The last equality is because of the fact that for given *t* and *B*, the event $\{X_t \in B\} \in \sigma(X_t)$ and $E_B^c \cap E_{P(B)} \in \mathcal{F}_{t-1}$. Therefore, the two events are independent.

Next we analyze the event $E_B^c \cap E_{P(B)}$ given l(B) = k in order to bound (3). This event implies that when the parent bin P(B) is created, its optimal price $p^*(P(B))$ is inside the interval $[p_l^{P(B)}, p_u^{P(B)}]$. At the end of Step 14, when n_{k-1} customers have been observed in P(B), it is split into 2^d children. The optimal price of its child bin B, that is $p^*(B)$, is no longer inside $[p_l^B, p_u^B]$. By Step 19, $p_l^B = \max\{p_{P(B)}^* - \Delta_k/2, 0\}$ and $p_u^B =$ $\min\{p_{P(B)}^* + \Delta_k/2, 1\}$, where $p_{P(B)}^*$ is the empirically-optimal price for P(B). Therefore, E_B^c implies that $p^*(B) \notin [p_{P(B)}^* - \Delta_k/2, p_{P(B)}^* + \Delta_k/2]$. Combined with Assumption 3 part two, which states that $p^*(B) \in [\inf\{p^*(\mathbf{x}) : \mathbf{x} \in B\}, \sup\{p^*(\mathbf{x}) : \mathbf{x} \in B\}] \subset \inf\{p^*(\mathbf{x}) : \mathbf{x} \in P(B)\}$, sup $\{p^*(\mathbf{x}) : \mathbf{x} \in P(B)\}$, we have

$$[\inf\{p^*(\boldsymbol{x}): \boldsymbol{x} \in P(B)\}, \sup\{p^*(\boldsymbol{x}): \boldsymbol{x} \in P(B)\}] \not\subset [p^*_{P(B)} - \Delta_k/2, p^*_{P(B)} + \Delta_k/2].$$

That is, either $\inf\{p^*(\mathbf{x}) : \mathbf{x} \in P(B)\} < p_{P(B)}^* - \Delta_k/2 \text{ or } \sup\{p^*(\mathbf{x}) : \mathbf{x} \in P(B)\} > p_{P(B)}^* + \Delta_k/2$. By Assumption 3 part three, $\sup\{p^*(\mathbf{x}) : \mathbf{x} \in P(B)\} - \inf\{p^*(\mathbf{x}) : \mathbf{x} \in P(B)\} \le M_4 d_{P(B)} = M_4 \sqrt{d2^{-(k-1)}}$ because the level of P(B) is k - 1. Hence by Assumption 3 part two, $\inf\{p^*(\mathbf{x}) : \mathbf{x} \in P(B)\} \ge p^*(P(B)) - M_4 \sqrt{d2^{-(k-1)}}$ and $\sup\{p^*(\mathbf{x}) : \mathbf{x} \in P(B)\} \le p^*(P(B)) + M_4 \sqrt{d2^{-(k-1)}}$. Combining the above observations, E_B^c could only happen when $|p^*(P(B)) - p_{P(B)}^*| > \Delta_k/2 - M_4 \sqrt{d2^{-(k-1)}}$. On the other hand, $E_{P(B)}$ implies that $p^*(P(B)) \in [p_l^{P(B)}, p_u^{P(B)}]$. Therefore, $E_B^c \cap E_{P(B)}$ could occur only if there exist two grid points $0 \le j_1, j_2 \le N_{k-1} - 1$ in Step 14 for bin P(B), such that

1. The j_2 th grid point is the closest to the optimal price for the bin $p^*(P(B))$. That is, $|p_1^{P(B)} + j_2 \delta_{P(B)} - p^*(P(B))| \le \delta_{P(B)}/2.$ 2. The j_1 th grid point maximizes $\bar{Y}_{P(B),j}$. That is $p_l^{P(B)} + j_1 \delta_{P(B)} = P_{P(B)}^*$. It implies that $\bar{Y}_{P(B),j_1} \ge \bar{Y}_{P(B),j_2}$ in Step 15.

3.
$$|p^*(P(B)) - p^*_{P(B)}| > \Delta_k/2 - M_4\sqrt{d}2^{-(k-1)}$$

In other words, the empirically-optimal price is the j_1 th grid point, while the j_2 th grid point is closest to the true revenue maximizer in bin P(B), i.e., $p^*(P(B))$. Given that the two grid points are far apart (by point 3 above), the probability of this event should be small.

To further bound the probability, consider $\bar{Y}_{P(B),j}$. In Step 15, it is the sum of $\lfloor n_{k-1}/N_{k-1} \rfloor$ or $\lceil n_{k-1}/N_{k-1} \rceil$ independent random variables with mean $\mathsf{E}[f(X, p_l^{P(B)} + j\delta_{P(B)})|X \in P(B)]$. By Lemmas 1 and 2, they are still sub-Gaussian with parameter $\sigma = 1/8$. This gives the following probabilistic bound (recall the definition of $f_B(p)$ in Section 2.2):

$$\begin{split} \mathsf{P}(E_B^c \cap E_{P(B)}) &\leq \mathsf{P}(\bar{Y}_{P(B),j_1} \geq \bar{Y}_{P(B),j_2}) \\ &= \mathsf{P}\Big(\frac{1}{t_1} \sum_{i=1}^{t_1} X_i^{(1)} - \frac{1}{t_2} \sum_{i=1}^{t_2} X_i^{(2)} \geq f_{P(B)}(p_l^{P(B)} + j_2 \delta_{P(B)}) - f_{P(B)}(p_l^{P(B)} + j_1 \delta_{P(B)})) \\ &= \mathsf{P}\Big(\frac{1}{t_1} \sum_{i=1}^{t_1} X_i^{(1)} - \frac{1}{t_2} \sum_{i=1}^{t_2} X_i^{(2)} \geq f_{P(B)}(p_l^{P(B)} + j_2 \delta_{P(B)}) - f_{P(B)}(p^*(P(B))) \\ &+ f_{P(B)}(p^*(P(B))) - f_{P(B)}(p_l^{P(B)} + j_1 \delta_{P(B)})) \\ &\leq \mathsf{P}\Big(\frac{1}{t_1} \sum_{i=1}^{t_1} X_i^{(1)} - \frac{1}{t_2} \sum_{i=1}^{t_2} X_i^{(2)} \geq M_2 \left(p_l^{P(B)} + j_1 \delta_{P(B)} - p^*(P(B))\right)^2 \\ &- M_3(p_l^{P(B)} + j_2 \delta_{P(B)} - p^*(P(B)))^2 \Big) \\ &\leq \mathsf{P}\Big(\frac{1}{t_1} \sum_{i=1}^{t_1} X_i^{(1)} - \frac{1}{t_2} \sum_{i=1}^{t_2} X_i^{(2)} \geq M_2 \left(\left(\Delta_k/2 - M_4\sqrt{d}2^{-(k-1)}\right)^+\right)^2 \\ &- M_3\delta_{P(B)}^2/4 \Big). \end{split}$$

Here t_1 and t_2 can be either $\lfloor n_{k-1}/N_{k-1} \rfloor$ or $\lceil n_{k-1}/N_{k-1} \rceil$; $X_i^{(1)}$ and $X_i^{(2)}$ are independent mean-zero sub-Gaussian random variables with parameter σ . Their averages are the centered version of $\bar{Y}_{P(B),j_1}$ and $\bar{Y}_{P(B),j_2}$, and thus their means are moved to the right-hand side. In the last inequality, $(\cdot)^+$ represents the positive part. The inequality follows from Assumption 3 part one and the previously derived facts that $|p_l^{P(B)} + j_2 \delta_{P(B)} - p^*(P(B))| \leq \delta_{P(B)}/2$ and $|p_l^{P(B)} + j_1 \delta_{P(B)} - p^*(P(B))| \geq \Delta_k/2 - M_4 \sqrt{d2^{-(k-1)}}$. By the property of sub-Gaussian random variables (for example, see Theorem 7.27 in Foucart and Rauhut, 2013), the above probability is bounded by

$$P(E_{B}^{c} \cap E_{P(B)}) \leq \exp\left(-\frac{\left(\left(M_{2}\left(\left(\Delta_{k}/2 - M_{4}\sqrt{d}2^{-(k-1)}\right)^{+}\right)^{2} - M_{3}\delta_{P(B)}^{2}/4\right)^{+}\right)^{2}}{4\sigma(1/t_{1} + 1/t_{2})}\right)$$
$$\leq \exp\left(-\frac{n_{k-1}\left(\left(M_{2}\left(\left(\Delta_{k}/2 - M_{4}\sqrt{d}2^{-(k-1)}\right)^{+}\right)^{2} - M_{3}\delta_{P(B)}^{2}/4\right)^{+}\right)^{2}}{N_{k-1} + 1}\right)$$

By our choice of parameters, $\Delta_k = 2^{-k} \log(T)$, $N_k \equiv \lceil \log(T) \rceil$, $\delta_{P(B)} \leq \Delta_{k-1}/N_{k-1} \leq 2^{-(k-1)}$. Therefore, when $T \geq \max\{\exp(8M_4\sqrt{d}), \exp(4\sqrt{2M_3/M_2})\}$, we have:

$$\begin{split} \frac{\Delta_k}{4} - M_4 \sqrt{d2^{-(k-1)}} &= 2^{-(k+2)} \log(T) - M_4 \sqrt{d2^{-(k-1)}} \ge 0 \\ &\Rightarrow \frac{\Delta_k}{2} - M_4 \sqrt{d2^{-(k-1)}} \ge \frac{\Delta_k}{4} \\ &\frac{M_2 \Delta_k^2}{32} - \frac{M_3 \delta_{P(B)}^2}{4} \ge \frac{M_2 2^{-2k} \log^2(T)}{32} - M_3 2^{-2k} \ge 0 \\ &\Rightarrow M_2 (\Delta_k / 2 - M_4 \sqrt{d2^{-(k-1)}})^2 - M_3 \delta_{P(B)}^2 / 4 \ge \frac{M_2 \Delta_k^2}{16} - \frac{M_3 \delta_{P(B)}^2}{4} \ge \frac{M_2 \Delta_k^2}{32}. \end{split}$$

Therefore, there exists a constant $c_1=M_2^2/1024$ such that

$$\mathsf{P}(E_B^c \cap E_{P(B)}) \le \exp\left(-c_1 \frac{\Delta_k^4 n_{k-1}}{\log(T) + 1}\right).$$

With this bound, we can proceed to provide an upper bound for (3). Because $\sum_{\{B:l(B)=k\}} P(X_t \in B) = 1$, we have

$$\sum_{t=1}^{T} \sum_{k=1}^{K} \sum_{\{B: l(B)=k\}} M_1 \mathsf{P}(X_t \in B) \mathsf{P}(E_B^c \cap E_{P(B)}) \le M_1 T \sum_{k=1}^{K} \exp\left(-c_1 \frac{\Delta_k^4 n_{k-1}}{\log(T) + 1}\right).$$
(4)

We next analyze term 2 of (1). By Assumption 3 part one, $f^*(X_t) - f(X_t, p_t) \le M_3(p^*(X_t) - p_t)^2 \le M_3(|p^*(B) - p_t| + |p^*(X_t) - p^*(B)|)^2$. By the design of the algorithm (Step 12 and 25), $p_t \in [p_l^B, p_u^B]$; conditional on the event $E_B = \{p^*(B) \in [p_l^B, p_u^B]\}$, we have $|p^*(B) - p_t| \le p_u^B - p_l^B \le \Delta_k$ for l(B) = k. On the other hand, by Assumption 3 part two and

three, $|p^*(X_t) - p^*(B)| \le \sup\{p^*(x) : x \in B\} - \inf\{p^*(x) : x \in B\} \le M_4 d_B \le M_4 \sqrt{d} 2^{-k}$ for l(B) = k. Therefore, term 2 can be bounded by

$$\mathbf{E} \left[\sum_{t=1}^{T} \sum_{k=0}^{K} \sum_{\{B:l(B)=k\}} (f^{*}(X_{t}) - f(X_{t}, p_{t})) \mathbb{I}_{\{X_{t} \in B, B \in \mathcal{P}_{t}, E_{B}\}} \right] \\
 \leq \mathbf{E} \left[\sum_{t=1}^{T} \sum_{k=0}^{K} \sum_{\{B:l(B)=k\}} M_{3}(\Delta_{k} + M_{4}\sqrt{d}2^{-k})^{2} \mathbb{I}_{\{X_{t} \in B, B \in \mathcal{P}_{t}, E_{B}\}} \right] \\
 \leq \mathbf{E} \left[\sum_{k=0}^{K-1} \sum_{\{B:l(B)=k\}} \sum_{t=1}^{T} M_{3}(\Delta_{k} + M_{4}\sqrt{d}2^{-k})^{2} \mathbb{I}_{\{X_{t} \in B, B \in \mathcal{P}_{t}\}} \right] \\
 + \sum_{t=1}^{T} M_{3}(\Delta_{K} + M_{4}\sqrt{d}2^{-K})^{2} \sum_{\{B:l(B)=K\}} \mathbf{P}(X_{t} \in B)$$
(5)

For the first term in (5), note that $\{X_t \in B, B \in \mathcal{P}_t\}$ occurs for at most n_k times for given B with l(B) = k. Moreover, there are 2^{dk} bins with level k, i.e., $\#\{B : l(B) = k\} = 2^{dk}$. Therefore, substituting $\Delta_k = 2^{-k} \log(T)$ into the first term yields an upper bound $\sum_{k=0}^{K-1} M_3 n_k 2^{(d-2)k} (\log(T) + M_4 \sqrt{d})^2$. For the second term in (5), $\sum_{\{B:l(B)=K\}} P(X_t \in B) = 1$ because $\{B : l(B) = K\}$ form a partition of the covariate space and X always falls into one of the bins. Therefore, (5) is bounded by

$$M_{3}\left(\sum_{k=0}^{K-1} n_{k} 2^{(d-2)k} + T 2^{-2K}\right) \left(\log(T) + M_{4}\sqrt{d}\right)^{2} \log(T)^{2}$$
$$\leq c_{3} \log(T)^{2} M_{3}\left(\sum_{k=0}^{K-1} n_{k} 2^{(d-2)k} + T 2^{-2K}\right)$$
(6)

for $c_3 = (1 + M_3) \left(\log(2) + M_4 \sqrt{d} \right)^2 / \log(2)^2$ and $T \ge 2$. Combining (4) and (6), we can find constants $c_2 = c_1/2^5 = M_2^2/2^{15}$ such that

$$\mathsf{E}[R_{\pi_{ABE}}] \leq \sum_{k=0}^{K-1} \left(c_3 \log(T)^2 n_k 2^{(d-2)k} + M_1 T \exp(-c_1 2^{-4k-4} \log^3(T) n_k/2) \right) + c_3 \log(T)^2 T 2^{-2K}$$

$$\leq \sum_{k=0}^{K-1} \left(c_3 \log(T)^2 n_k 2^{(d-2)k} + M_1 T \exp(-c_2 2^{-4k} \log^3(T) n_k) \right) + c_3 \log(T)^2 T 2^{-2K}$$
(7)

We choose n_k

$$n_k = \max\left\{0, \left\lceil \frac{2^{4k+15}}{M_2^2 \log(T)^3} (\log(T) + \log(\log(T)) - (d+2)k \log(2)) \right\rceil \right\}$$

to minimize $c_3 \log(T)^2 n_k 2^{(d-2)k} + M_1 T \exp(-c_2 2^{-4k} \log(T)^3 n_k)$ in (7). More precisely,

$$c_{3} \log(T)^{2} n_{k} 2^{(d-2)k} \leq c_{3} M_{2}^{-2} 2^{15} \frac{\log(T)^{2}}{\log(T)^{3}} 2^{(d+2)k} (\log(T) + \log(\log(T)))$$

$$\leq c_{4} 2^{(d+2)k}$$

for some constants $c_4 > 0$, and

$$M_1 T \exp(-c_2 2^{-4k} \log^3(T) n_k) \le M_1 T \exp(-\log(T) - \log(\log(T)) + (d+2)k \log(2)) \le c_5 2^{(d+2)k}$$

for a constant $c_5 > 0$. Therefore, (7) implies that we can find a constant $c_6 = c_4 + c_5$ such that

$$\mathsf{E}[R_{\pi_{ABE}}] \le \sum_{k=0}^{K-1} c_6 2^{(d+2)k} + c_3 \log(T)^2 T 2^{-2K}$$
$$\le c_6 2^{(d+2)K} + c_3 \log(T)^2 T 2^{-2K}.$$

Therefore, by our choice of $K = \lfloor \frac{\log(T)}{(d+4)\log(2)} \rfloor$, the regret is bounded by

$$c_7 \log^2(T) T^{\frac{d+2}{d+4}}$$

for some constant c_7 . Hence we have completed the proof.

B.2. Lower Bound in Section 5

Next we show that Assumption 1 and 2 are satisfied by the construction in Section 5.

Proposition 2. The choice of $f \in C$ satisfies Assumption 2 and 3 with $M_1 = 4$, $M_2 = 1$, $M_3 = 2$, and $M_4 = 1$.

To give some intuitions, note that by the construction of f, both $f_w(\mathbf{x}, p)$ and $p^*(\mathbf{x})$ are Lipschitz continuous in $[0, 1)^d$. Such continuity guarantees the desired properties.

Proof of Proposition 2: For Assumption 2, we discuss two cases. The first case is $x_1, x_2 \in$

 B_j , i.e., the two customers are in the same bin. In this case,

$$|f_{w}(\mathbf{x}_{1}, p_{1}) - f_{w}(\mathbf{x}_{2}, p_{2})| \leq \begin{cases} |p_{1}(\frac{2}{3} - \frac{p_{1}}{2}) - p_{2}(\frac{2}{3} - \frac{p_{2}}{2})| \leq 2|p_{1} - p_{2}| & w_{j} = 0\\ 2|p_{1} - p_{2}| + |p_{1}(\frac{1}{3} - \frac{p_{1}}{2})D(\mathbf{x}_{1}, \partial B_{j}) - p_{2}(\frac{1}{3} - \frac{p_{2}}{2})D(\mathbf{x}_{2}, \partial B_{j})| & w_{j} = 1 \end{cases}$$

When $w_j = 0$, the assumption is already satisfied. When $w_j = 1$, by the triangle inequality we have

$$|p_{1}(\frac{1}{3} - \frac{p_{1}}{2})D(\mathbf{x}_{1}, \partial B_{j}) - p_{2}(\frac{1}{3} - \frac{p_{2}}{2})D(\mathbf{x}_{2}, \partial B_{j})|$$

$$\leq 2|p_{1} - p_{2}|D(\mathbf{x}_{1}, \partial B_{j}) + p_{2}\left(\frac{1}{3} - \frac{p_{2}}{2}\right)|D(\mathbf{x}_{1}, \partial B_{j}) - D(\mathbf{x}_{2}, \partial B_{j})|$$

$$\leq \frac{1}{M}|p_{1} - p_{2}| + |D(\mathbf{x}_{1}, \partial B_{j}) - D(\mathbf{x}_{2}, \partial B_{j})|$$

$$\leq |p_{1} - p_{2}| + ||\mathbf{x}_{1} - \mathbf{x}_{2}||_{2}$$

The second inequality is because $p_2 \ge 0$ and $D(\mathbf{x}_1, \partial B_j) \le 1/2M \le 1/2$ when $\mathbf{x}_1 \in B_j$. The third inequality is because

$$\|\mathbf{x}_{1} - \mathbf{x}_{2}\|_{2} + D(\mathbf{x}_{2}, \partial B_{j}) = \min_{a \in \partial B_{j}} \{\|a - \mathbf{x}_{2}\|_{2} + \|\mathbf{x}_{1} - \mathbf{x}_{2}\|_{2}\} \ge \min_{a \in \partial B_{j}} \{\|a - \mathbf{x}_{1}\|_{2}\}$$
$$= D(\mathbf{x}_{1}, \partial B_{j})$$

and similarly $\|\mathbf{x}_1 - \mathbf{x}_2\|_2 + D(\mathbf{x}_1, \partial B_j) \ge D(\mathbf{x}_2, \partial B_j)$. Therefore, we have shown that $|f_w(\mathbf{x}_1, p_1) - f_w(\mathbf{x}_2, p_2)| \le 3|p_1 - p_2| + 2\|\mathbf{x}_1 - \mathbf{x}_2\|_2$ for case one.

The second case is $\mathbf{x}_1 \in B_{j_1}$ and $\mathbf{x}_2 \in B_{j_2}$ for $j_1 \neq j_2$. If $j_1 = j_2 = 0$, then by the previous analysis, we already have $|f_w(\mathbf{x}_1, p_1) - f_w(\mathbf{x}_2, p_2)| \leq 2|p_1 - p_2|$. If $j_1 = 1$ and $j_2 = 0$, then

$$|f_{w}(\boldsymbol{x}_{1}, p_{1}) - f_{w}(\boldsymbol{x}_{2}, p_{2})| \leq 2|p_{1} - p_{2}| + p_{1}\left(\frac{1}{3} - \frac{p_{1}}{2}\right)D(\boldsymbol{x}_{1}, \partial B_{j_{1}}) \leq 2|p_{1} - p_{2}| + 2||\boldsymbol{x}_{1} - \boldsymbol{x}_{2}||_{2}.$$

The last inequality is because the straight line connecting \mathbf{x}_1 and \mathbf{x}_2 must intersects B_{j_1} . The distance from \mathbf{x}_1 to the intersection is no less than $D(\mathbf{x}_1, \partial B_{j_1})$. Therefore, $D(\mathbf{x}_1, \partial B_{j_1}) \leq ||\mathbf{x}_1 - \mathbf{x}_2||_2$. If $j_1 = 0$ and $j_2 = 1$, then the result follows similarly. If $j_1 = j_2 = 1$, then by the same argument,

$$|p_1\left(\frac{1}{3} - \frac{p_1}{2}\right)D(\mathbf{x}_1, \partial B_{j_1}) - p_2\left(\frac{1}{3} - \frac{p_2}{2}\right)D(\mathbf{x}_2, \partial B_{j_2})| \le D(\mathbf{x}_1, \partial B_{j_1}) + D(\mathbf{x}_2, \partial B_{j_2}) \le 2||\mathbf{x}_1 - \mathbf{x}_2||_2$$

Therefore, combining both cases, we always have

$$|f_{w}(\mathbf{x}_{1}, p_{1}) - f_{w}(\mathbf{x}_{2}, p_{2})| \leq 4|p_{1} - p_{2}| + 4||\mathbf{x}_{1} - \mathbf{x}_{2}||_{2}.$$

For Assumption 3, note that

$$f_B(p) = p\left(\frac{2}{3} - \frac{p}{2}\right) + p\left(\frac{1}{3} - \frac{p}{2}\right) \mathbb{E}\left[\sum_{j=1}^{M^d} w_j \mathbb{I}_{\{X \in B_j\}} D(X, \partial B_j) | X \in B\right]$$
$$= p\left(\frac{2}{3} - \frac{p}{2}\right) + p\left(\frac{1}{3} - \frac{p}{2}\right) \sum_{j=1}^{M^d} w_j \mathbb{P}(B_j \cap B) \mathbb{E}[D(X, \partial B_j) | X \in B_j \cap B]$$

For part one, because the second-order derivative of $f_B(p)$ is always bounded between [-2, -1],

$$2(p - p^*(B))^2 \ge f_B(p^*(B)) - f_B(p) \ge (p - p^*(B))^2$$

and part one holds for $M_2 = 1$, $M_3 = 2$. For part two, note that the maximizer

$$p^*(B) = \frac{2 + \sum_{j=1}^{M^d} w_j \mathsf{P}(B_j \cap B) \mathsf{E}[D(X, \partial B_j) | X \in B_j \cap B]}{3(1 + \sum_{j=1}^{M^d} w_j \mathsf{P}(B_j \cap B) \mathsf{E}[D(X, \partial B_j) | X \in B_j \cap B])}.$$

Because $p^*(B)$ is a monotone function of $\sum_{j=1}^{M^d} w_j P(B_j \cap B) E[D(X, \partial B_j) | X \in B_j \cap B]$, it is easy to check that part two of Assumption 3 holds. For part three, consider $\mathbf{x}_1 \in B \cap B_{j_1}$ and $\mathbf{x}_2 \in B \cap B_{j_2}$. If $w_{j_1} = 0$ and $w_{j_2} = 0$, then $p^*(\mathbf{x}_1) - p^*(\mathbf{x}_2) = 0$. If either $w_{j_1} = 0$ or $w_{j_2} = 0$, then

$$|p^*(\mathbf{x}_1) - p^*(\mathbf{x}_2)| \le \frac{1}{3} \max \left\{ D(\mathbf{x}_1, \partial B_{j_1}), D(\mathbf{x}_2, \partial B_{j_2}) \right\} \le ||\mathbf{x}_1 - \mathbf{x}_2||_2 \le d_B.$$

by the previous analysis. If $w_{j_1} = 1$ and $w_{j_2} = 1$, then similarly we have $|p^*(\mathbf{x}_1) - p^*(\mathbf{x}_2)| \le \frac{1}{3}|D(\mathbf{x}_1, \partial B_{j_1}) - D(\mathbf{x}_2, \partial B_{j_2})| \le ||\mathbf{x}_1 - \mathbf{x}_2||_2 \le d_B$. Therefore, part three holds with $M_4 = 1$.

The proof of Theorem 2 uses Kullback-Leibler (KL) divergence to measure the "distinguishability" of the underlying functions. Such information-theoretic approach has been a standard technique in the learning literature. The proof is outlined in the following.

Among all functions $f \in C$, we focus on each pair of f_w and $f_{w'}$ that only differ in a single bin. For example, consider $w = (w_1, w_2, \ldots, w_{j-1}, 0, w_{j+1}, \ldots, w_{M^d})$ and $w' = (w_1, w_2, \ldots, w_{j-1}, 1, w_{j+1}, \ldots, w_{M^d})$ for some *j*. Because the indices of *w* and *w'* are identical except for the *j*th, f_w and $f_{w'}$ only differ in bin B_j . Denote $w = (w_{-j}, 0)$ and $w' = (w_{-j}, 1)$ to highlight this fact. Distinguishing between $f_{w_{-j},0}$ and $f_{w_{-j},1}$ poses a challenge to any policy. In particular, for $\mathbf{x} \in B_j$, the difference of the two functions $|f_{w_{-j},0}(\mathbf{x}, p) - f_{w_{-j},1}(\mathbf{x}, p)| = p(\frac{1}{3} - \frac{p}{2})D(\mathbf{x}, \partial B_j)$ is diminishing when in $p \approx 2/3$. Thus, charging a price different from 2/3 makes the difference more visible and helps to distinguish $f_{w_{-j},0}$ and $f_{w_{-j},1}$. However, if *p* deviates too much from the optimal decision $p^*(\mathbf{x}) = 2/3$ or $p^*(\mathbf{x}) = (2 + D(\mathbf{x}, \partial B_j))/(3(1 + D(\mathbf{x}, \partial B_j)))$, then significant regret is incurred in that period.

To capture this trade-off, for a given $j = 1, ..., M^d$ and $w_{-j} \in \{0, 1\}^{M^d-1}$, define the following quantity

$$z_{w_{-j}} = \sum_{t=1}^{T} \frac{9}{11M^2} \mathsf{E}_{f_{w_{-j},0}}^{\pi} \left[\left(\frac{2}{3} - p_t \right)^2 \mathbb{I}_{\{X_t \in B_j\}} \right]$$
(8)

where the expectation is taken with respect to a policy π and the underlying function $f_{w_{-j},0}$. This quantity is crucial in analyzing the regret. More precisely, if $z_{w_{-j}}$ is large (which implies that p_t is large), then $f_{w_{-j},0}$ and $f_{w_{-j},1}$ are easy to distinguish but the regret becomes uncontrollable.

Lemma 3.

$$\sup_{f \in C} R_{\pi} \geq \frac{11M_2}{9 \times 2^{M^d}} M^2 \sum_{j=1}^{M^d} \sum_{w_{-j}} z_{w_{-j}}.$$

On the other hand, if $z_{w_{-j}}$ is small, then the KL divergence of the measures associated with $f_{w_{-j},0}$ and $f_{w_{-j},1}$ is also small. In other words, the firm cannot easily distinguish between $f_{w_{-j},0}$ and $f_{w_{-j},1}$ which impedes learning and incurs substantial regret.

Lemma 4.

$$\sup_{f \in C} R_{\pi} \geq \frac{M_2 T}{9 \times 2^{M^d + 11} M^{d+2}} \sum_{j=1}^{M^d} \sum_{w_{-j}} \exp(-z_{w_{-j}}).$$

Since the effects of $z_{w_{-j}}$ are opposite in Lemma 3 and Lemma 4, combining the two bounds, we can find a positive constant c_1 independent of *T* and *M* so that

$$R_{\pi} \geq \frac{c_1}{2^{M^d}} \sum_{j=1}^{M^d} \sum_{w_{-j}} \left(\frac{T}{M^{d+2}} \exp\left(-z_{w_{-j}}\right) + M^2 z_{w_{-j}} \right)$$
$$\geq \frac{c_1}{2^{M^d}} \sum_{j=1}^{M^d} \sum_{w_{-j}} M^2 \left(1 + \log\left(\frac{T}{M^{d+4}}\right) \right)$$
$$\geq \frac{c_1 M^{d+2}}{2} \left(1 + \log\left(\frac{T}{M^{d+4}}\right) \right).$$

In the second inequality above, we minimize the expression over positive $z_{w_{-j}}$. Since M can be an arbitrary positive integer, we let $M = \lceil T^{1/(d+4)} \rceil$ in the last quantity. Calculation shows that it is lower bounded by $cT^{(2+d)/(4+d)}$ for a constant c > 0.

To prove Theorem 2, i.e., Lemma 3 and Lemma 4, we introduce the following lemmas.

Lemma 5 (KL divergence for Bernoulli Random Variables). *For two Bernoulli random variables* X_1 *and* X_2 *with means* θ_1 *and* θ_2 *, we have*

$$\mathcal{K}(\mu_{X_1},\mu_{X_2}) \leq \frac{(\theta_1-\theta_2)^2}{\theta_2(1-\theta_2)}.$$

Proof. The proof can be found in, e.g., Rigollet and Zeevi (2010) and thus omitted. ■

Lemma 6 (The chain rule of the KL divergence). *Given joint distributions* p(x, y) *and* q(x, y), we have

$$\mathcal{K}(p(x,y),q(x,y)) = \mathcal{K}(p(x),q(x)) + \mathsf{E}_{p(x)}[\mathcal{K}(p(y|x),q(y|x)]]$$

where $p(\cdot)$ and $q(\cdot)$ represent the marginal distribution, $p(\cdot|x)$ and $q(\cdot|x)$ represent the conditional distribution.

Proof. The proof can be found from standard textbooks and is thus omitted.

Proof of Lemma 3: We use E_f^{π} to highlight the dependence of the expectation on the policy π and the underlying function f. Note that

$$\sup_{f \in C} R_{\pi} = \sup_{f \in C} \sum_{t=1}^{T} \mathbb{E} \left[f^{*}(X_{t}) - f(X_{t}, p_{t}) \right] = \sup_{f \in C} \sum_{t=1}^{T} \sum_{j=1}^{M^{d}} \mathbb{E} \left[(f^{*}(X_{t}) - f(X_{t}, p_{t})) \mathbb{I}_{\{X_{t} \in B_{j}\}} \right]$$
$$\geq M_{2} \sup_{f \in C} \sum_{t=1}^{T} \sum_{j=1}^{M^{d}} \mathbb{E} \left[(p^{*}(X_{t}) - p_{t})^{2} \mathbb{I}_{\{X_{t} \in B_{j}\}} \right]$$
$$\geq \frac{M_{2}}{2^{M^{d}}} \sum_{w} \sum_{t=1}^{T} \sum_{j=1}^{M^{d}} \mathbb{E} \left[(p^{*}(X_{t}) - p_{t})^{2} \mathbb{I}_{\{X_{t} \in B_{j}\}} \right]$$

In the last inequality, we have used the fact that $\#\{C\} = 2^{M^d}$ and the supremum is always no less than the average.

For a given bin B_j , we focus on $f_{w_{-j},0}$ and $f_{w_{-j},1}$, which only differ for $\mathbf{x} \in B_j$. Therefore, we can rearrange \sum_w to $\sum_{w_{-j} \in \{0,1\}^{M^d-1}} \sum_{w_j \in \{0,1\}}$. We have the following lower bound for the regret

$$\begin{split} \sup_{f \in C} R_{\pi} &\geq \frac{M_2}{2^{M^d}} \sum_{j=1}^{M^d} \sum_{w_{-j} \in \{0,1\}^{M^d - 1}} \sum_{w_j \in \{0,1\}} \sum_{t=1}^T \mathsf{E}_{f_{w_{-j},w_j}}^{\pi} \left[(p^*(X_t) - p_t)^2 \mathbb{I}_{\{X_t \in B_j\}} \right] \\ &\geq \frac{M_2}{2^{M^d}} \sum_{j=1}^{M^d} \sum_{w_{-j}} \sum_{t=1}^T \mathsf{E}_{f_{w_{-j},0}}^{\pi} \left[(p^*(X_t) - p_t)^2 \mathbb{I}_{\{X_t \in B_j\}} \right] \\ &= \frac{M_2}{2^{M^d}} \sum_{j=1}^{M^d} \sum_{w_{-j}} \sum_{t=1}^T \mathsf{E}_{f_{w_{-j},0}}^{\pi} \left[\left(\frac{2}{3} - p_t \right)^2 \mathbb{I}_{\{X_t \in B_j\}} \right] \\ &= \frac{11M_2}{9 \times 2^{M^d}} M^2 \sum_{j=1}^{M^d} \sum_{w_{-j}} z_{w_{-j}}. \end{split}$$

In the second inequality, we have neglected the regret for $f_{w_{-j},1}$. The last equality is by the definition of $z_{w_{-j}}$ in (8). Hence we have proved the result.

Proof of Lemma 4: By the same argument as in the proof of Lemma 3, we have

$$\sup_{f \in C} R_{\pi} \geq \frac{M_2}{2^{M^d}} \sum_{j=1}^{M^d} \sum_{t=1}^T \sum_{w_{-j} \in \{0,1\}^{M^d-1}} \sum_{w_j \in \{0,1\}} \mathsf{E}_{f_{w_{-j},w_j}}^{\pi} \left[(p^*(X_t) - p_t)^2 \mathbb{I}_{\{X_t \in B_j\}} \right].$$

Because X_t is uniformly distributed in $[0, 1)^d$, $P(X_t \in B_j) = M^{-d}$. By conditioning on the event $X_t \in B_j$, we have

$$\mathsf{E}_{f_{w_{-j},w_j}}^{\pi} \left[(p^*(X_t) - p_t)^2 \mathbb{I}_{\{X_t \in B_j\}} \right] = \mathsf{E}_{f_{w_{-j},w_j}}^{\pi} \left[(p^*(X_t) - p_t)^2 | X_t \in B_j \right] \mathsf{P}(X_t \in B_j)$$

= $\frac{1}{M^d} \mathsf{E}_{f_{w_{-j},w_j}}^{\pi} \left[(p^*(X_t) - p_t)^2 | X_t \in B_j \right].$

Since $(p^*(X_t) - p_t)^2$ is measurable with respect to the σ -algebra generated by \mathcal{F}_{t-1} and X_t , by the tower property, we have

$$\mathsf{E}_{f_{\mathsf{w}_{-j},\mathsf{w}_{j}}}^{\pi}\left[(p^{*}(X_{t})-p_{t})^{2}\mathbb{I}_{\left\{X_{t}\in B_{j}\right\}}\right] = \frac{1}{M^{d}}\mathsf{E}_{f_{\mathsf{w}_{-j},\mathsf{w}_{j}}}^{\pi}\left[\mathsf{E}\left[(p^{*}(X_{t})-p_{t})^{2}|\mathcal{F}_{t-1},X_{t}\in B_{j}\right]\right]$$

Let $\mathsf{E}_{f_{w_{-j},w_j}}^{\pi,t-1}[\cdot]$ denote $\mathsf{E}_{f_{w_{-j},w_j}}^{\pi}[\mathsf{E}[\cdot|\mathcal{F}_{t-1}]]$ and let $\mathsf{P}_{X_t}^{B,t-1}(\cdot)$ denote the conditional proba-

bility $P(\cdot | \mathcal{F}_{t-1}, X_t \in B)$. By Markov's inequality, for any constant s > 0 we have

$$\sum_{w_{j}\in\{0,1\}} \mathsf{E}_{f_{w_{-j},w_{j}}}^{\pi} \left[(p^{*}(X_{t}) - p_{t})^{2} \mathbb{I}_{\{X_{t}\in B_{j}\}} \right]$$
(9)

$$= \frac{1}{M^{d}} \sum_{w_{j}\in\{0,1\}} \mathsf{E}_{f_{w_{-j},w_{j}}}^{\pi} \left[\mathsf{E} \left[(p^{*}(X_{t}) - p_{t})^{2} | X_{t} \in B_{j}, \mathcal{F}_{t-1} \right] \right]$$
(9)

$$\geq \frac{1}{M^{d}} \sum_{w_{j}\in\{0,1\}} \frac{s^{2}}{M^{2}} \mathsf{E}_{f_{w_{-j},w_{j}}}^{\pi} \left[\mathsf{P}_{X_{t}}^{B_{j,t-1}} \left(| p^{*}(X_{t}) - p_{t} | \geq \frac{s}{M} \right) \right]$$
(9)

$$= \frac{s^{2}}{M^{d+2}} \left(\mathsf{E}_{f_{w_{-j},0}}^{\pi} \left[\mathsf{P}_{X_{t}}^{B_{j,t-1}} \left(| \frac{2}{3} - p_{t} | \geq \frac{s}{M} \right) \right] + \mathsf{E}_{f_{w_{-j},1}}^{\pi} \left[\mathsf{P}_{X_{t}}^{B_{j,t-1}} \left(| \frac{2 + D(X_{t}, \partial B_{j})}{3(1 + D(X_{t}, \partial B_{j}))} - p_{t} | \geq \frac{s}{M} \right) \right]$$
(10)

$$+ \mathsf{E}_{f_{w_{-j},1}}^{\pi} \left[\mathsf{P}_{X_{t}}^{B_{j,t-1}} \left(| \frac{2 + D(X_{t}, \partial B_{j})}{3(1 + D(X_{t}, \partial B_{j}))} - p_{t} | \geq \frac{s}{M}, A \right) \right]$$
(10)

where we define event $A = \{X_t \in B_j\} \cap \{D(X_t, \partial B_j) > 12s/M\}$. In the second equality, we have used the fact that for $X_t \in B_j$, when $w_j = 0$, $p^*(X_t) = \frac{2}{3}$; when $w_j = 1$, $p^*(X_t) = \frac{2+D(X_t, \partial B_j)}{3(1+D(X_t, \partial B_j))}$. The motivation of introducing A is as follows: consider the classification rule $\Pi_t \mapsto \{0, 1\}$ associated with p_t tries to distinguish between $w_j = 0$ and $w_j = 1$. It is defined as

$$\Pi_{t} = \begin{cases} 0 & |\frac{2}{3} - p_{t}| \le |\frac{2 + D(X_{t}, \partial B_{j})}{3(1 + D(X_{t}, \partial B_{j}))} - p_{t}| \\ 1 & \text{otherwise.} \end{cases}$$

In other words, Π_t classifies the underlying function as $f_{w_{-j},0}$ if p_t is closer to the optimal price $p^*(X_t)$ of $f_{w_{-j},0}$, and as $f_{w_{-j},1}$ vice versa. For $f_{w_{-j},0}$, a misclassification on the event A is $A \cap \{\Pi_t = 1\}$. It implies that

$$|2/3 - p_t| \ge \frac{1}{2} \times \left| \frac{2}{3} - \frac{2 + D(X_t, \partial B_j)}{3(1 + D(X_t, \partial B_j))} \right| = \frac{D(X_t, \partial B_j)}{6(1 + D(X_t, \partial B_j))},$$

which on *A*, implies $A \cap \{|2/3 - p_t| \ge s/M\}$ as $D(X_t, \partial B_j) \le 1$. Similarly, $A \cap \{\Pi_t = 0\} \subset A \cap \{|\frac{2+D(X_t, \partial B_j)}{3(1+D(X_t, \partial B_j))} - p_t| \ge s/M\}$. Therefore, by the fact that $P(X \in A) = (1 - 24s)^d/M^d$,

we have

Next we lower bound the misclassification error (11) by the Kullback-Leibler (KL) divergence between the two probability measures associated with $f_{w_{-j},0}$ and $f_{w_{-j},1}$. Intuitively, if the two probability measures are close, then no classification (including Π_t) can incur very small misclassification error. Formally, introduce the KL divergence between two probability measures *P* and *Q* as

$$\mathcal{K}(P,Q) = \begin{cases} \int \log \frac{dP}{dQ} dP & \text{if } P \ll Q \\ +\infty & \text{otherwise} \end{cases}$$

where $P \ll Q$ indicates that P is absolute continuous w.r.t. Q. By the independence of \mathcal{F}_{t-1} and X_t , the two measures we want to distinguish in (11), $\mu_{f_{w-j},0}^{\pi}(\cdot|X_t \in A)$ and $\mu_{f_{w-j},1}^{\pi}(\cdot|X_t \in A)$, can be expressed as product measures

$$\begin{split} \mu_{f_{w_{-j},0}}^{\pi}(\cdot|X_t \in A) &= \mu_{f_{w_{-j},0}}^{\pi,t-1}(\cdot) \times \mu_{X_t}^{A}(\cdot) \\ \mu_{f_{w_{-j},1}}^{\pi}(\cdot|X_t \in A) &= \mu_{f_{w_{-j},1}}^{\pi,t-1}(\cdot) \times \mu_{X_t}^{A}(\cdot), \end{split}$$

where $\mu_{f_{w_{-j},0}}^{\pi,t-1}(\cdot)$ is a measure of $(X_1, Z_1, \ldots, X_{t-1}, Z_{t-1})$ depending on π and $f_{w_{-j},0}$ and $\mu_{X_t}^A(\cdot)$ is a measure of X_t conditional on $X_t \in A$. By Theorem 2.2 (iii) in Tsybakov (2009),

$$(11) \geq \frac{(1-24s)^d}{2} \exp\left(-\mathcal{K}\left(\mu_{f_{w_{-j},0}}^{\pi,t-1} \times \mu_{X_t}^A, \mu_{f_{w_{-j},1}}^{\pi,t-1} \times \mu_{X_t}^A\right)\right) \\ = \frac{(1-24s)^d}{2} \exp\left(-\mathcal{K}\left(\mu_{f_{w_{-j},0}}^{\pi,t-1}, \mu_{f_{w_{-j},1}}^{\pi,t-1}\right) - \mathsf{E}_{f_{w_{-j},0}}^{\pi,t-1}\left[\mathcal{K}\left(\mu_{X_t}^A, \mu_{X_t}^A\right)\right]\right) \\ = \frac{(1-24s)^d}{2} \exp\left(-\mathcal{K}\left(\mu_{f_{w_{-j},0}}^{\pi,t-1}, \mu_{f_{w_{-j},1}}^{\pi,t-1}\right)\right).$$
(12)

The second line follows from Lemma 6; the third line follows from the fact that $\mu_{X_t}^A$ is the same distribution for $f_{w_{-j},0}$ and $f_{w_{-j},1}$, independent of \mathcal{F}_{t-1} .

To further simplify the expression, note that $\mu_{f_{w_{-i},0}}^{\pi,t}(\cdot)$ can be decomposed as

$$\mu_{f_{w_{-j},0}}^{\pi,t}(\cdot) = \mu_{f_{w_{-j},0}}^{\pi,t-1}(\cdot) \times \mu_{X}(\cdot) \times \mu_{f_{w_{-j},0}}^{Z_{t}}(\cdot|\mathcal{F}_{t-1},X_{t}),$$

where μ_X is the measure (uniform distribution) of X_t and $\mu_{f_{w_{-j},0}}^{Z_t}(\cdot | \mathcal{F}_{t-1}, X_t)$ is the measure of Z_t conditional on \mathcal{F}_{t-1} and X_t . We apply Lemma 6 again:

$$\mathcal{K}\left(\mu_{f_{w_{-j},0}}^{\pi,t},\mu_{f_{w_{-j},1}}^{\pi,t}\right) = \mathcal{K}\left(\mu_{f_{w_{-j},0}}^{\pi,t-1},\mu_{f_{w_{-j},1}}^{\pi,t-1}\right) + \mathsf{E}_{f_{w_{-j},0}}^{\pi,t-1}\left[\mathcal{K}(\mu_{X_{t}},\mu_{X_{t}})\right] \\ + \mathsf{E}_{f_{w_{-j},0}}^{\pi,t-1}\left[\mathsf{E}_{X}\left[\mathcal{K}\left(\mu_{f_{w_{-j},0}}^{Z_{t}}(\cdot|\mathcal{F}_{t-1},X_{t}),\mu_{f_{w_{-j},1}}^{Z_{t}}(\cdot|\mathcal{F}_{t-1},X_{t})\right)\right]\right].$$

It is easy to see that the second term is zero. For the third term, we first conditional on \mathcal{F}_{t-1} and then on the covariate X_t . Because p_t depends only on \mathcal{F}_{t-1} and X_t , p_t is the same for $f_{w_{-j},0}$ and $f_{w_{-j},1}$ conditional on \mathcal{F}_{t-1} and X_t . Therefore, $\mu_{fw_{-j},0}^{Z_t}(\cdot|\mathcal{F}_{t-1},X_t)$ and $\mu_{fw_{-j},1}^{Z_t}(\cdot|\mathcal{F}_{t-1},X_t)$ are two Bernoulli distributions with means $d_{w_{-j},0}(X_t,p_t)$ and $d_{w_{-j},1}(X_t,p_t)$, respectively. By Lemma 5, we have

$$\begin{aligned} \mathcal{K}\left(\mu_{fw_{-j},0}^{Z_{t}}(\cdot|\mathcal{F}_{t-1},X_{t}),\mu_{fw_{-j},1}^{Z_{t}}(\cdot|\mathcal{F}_{t-1},X_{t})\right) &\leq \frac{\left(d_{w_{-j},0}(X_{t},p_{t}) - d_{w_{-j},1}(X_{t},p_{t})\right)^{2}}{d_{w_{-j},1}(X_{t},p_{t})(1 - d_{w_{-j},1}(X_{t},p_{t})))} \\ &\leq \frac{144}{11} \left(d_{w_{-j},0}(X_{t},p_{t}) - d_{w_{-j},1}(X_{t},p_{t})\right)^{2} \\ &= \frac{144}{11} \left(p_{t}\left(\frac{1}{3} - \frac{p_{t}}{2}\right)D(X_{t},\partial B_{j})\mathbb{I}_{\{X_{t}\in B_{j}\}}\right)^{2} \\ &\leq \frac{9}{11M^{2}} \left(\frac{2}{3} - p_{t}\right)^{2}\mathbb{I}_{\{X_{t}\in B_{j}\}}.\end{aligned}$$

In the second inequality, we used the fact that $d_{w_{-j},1}(X_t, p_t) \in [1/12, 5/6]$ as long as we choose $M \ge 2$ and thus $D(X_t, \partial B_j) \le 1/2$. In the last inequality, we have used the fact that the distance of a vector inside B_j to the boundary of B_j is at most 1/2M. Therefore,

we can obtain an upper bound for $\mathcal{K}\left(\mu_{f_{w_{-j},0}}^{\pi,t},\mu_{f_{w_{-j},1}}^{\pi,t}\right)$

$$\begin{aligned} \mathcal{K}\left(\mu_{f_{w_{-j},0}}^{\pi,t},\mu_{f_{w_{-j},1}}^{\pi,t}\right) &\leq \sum_{i=1}^{t} \mathsf{E}_{f_{w_{-j},0}}^{\pi,i-1} \left[\mathsf{E}_{X}\left[\mathcal{K}\left(\mu_{f_{w_{-j},0}}^{Z_{i}}(\cdot|\mathcal{F}_{i-1},X_{i}),\mu_{f_{w_{-j},1}}^{Z_{i}}(\cdot|\mathcal{F}_{i-1},X_{i})\right)\right]\right] \\ &\leq \sum_{i=1}^{t} \mathsf{E}_{f_{w_{-j},0}}^{\pi,i-1} \left[\mathsf{E}_{X}\left[\frac{9}{11M^{2}}\left(\frac{2}{3}-p_{t}\right)^{2}\mathbb{I}_{\{X_{i}\in B_{j}\}}\right]\right] \\ &\leq \sum_{t=1}^{T} \mathsf{E}_{f_{w_{-j},0}}^{\pi}\left[\frac{9}{11M^{2}}\left(\frac{2}{3}-p_{t}\right)^{2}\mathbb{I}_{\{X_{t}\in B_{j}\}}\right] = z_{w_{-j}}.\end{aligned}$$

Therefore, combining it with (9), (11) and (12), we have shown the lemma:

$$\begin{split} \sup_{f \in C} \sum_{t=1}^{T} \mathsf{E} \left[f^{*}(X_{t}) - f(X_{t}, p_{t}) \right] \\ &\geq \frac{M_{2}}{2^{M^{d}}} \sum_{t=1}^{T} \sum_{j=1}^{M^{d}} \sum_{w_{-j}} \sum_{w_{j} \in \{0,1\}} \mathsf{E}_{f_{w_{-j},w_{j}}}^{\pi} \left[(p^{*}(X_{t}) - p_{t})^{2} \mathbb{I}_{\{X_{t} \in B_{j}\}} \right] \\ &\geq \frac{TM_{2}s^{2}(1 - 24s)^{2}}{2^{M^{d}+1}M^{d+2}} \sum_{j=1}^{M^{d}} \sum_{w_{-j}} \exp\left(-z_{w_{-j}}\right) \\ &= \frac{M_{2}T}{9 \times 2^{M^{d}+11}M^{d+2}} \sum_{j=1}^{M^{d}} \sum_{w_{-j}} \exp\left(-z_{w_{-j}}\right) \end{split}$$

where in the last step, we have set s = 1/24.