

POINTWISE ERGODIC THEOREM FOR LOCALLY COUNTABLE QUASI-PMP GRAPHS

ANUSH TSERUNYAN

ABSTRACT. We prove a pointwise ergodic theorem for quasi-probability-measure-preserving (quasi-pmp) locally countable measurable graphs, equivalently, Schreier graphs of quasi-pmp actions of countable groups. For ergodic graphs, the theorem gives an increasing sequence of Borel subgraphs with finite connected components over which the averages of any L^1 function converges to its expectation. This implies that every (not necessarily pmp) locally countable ergodic Borel graph on a standard probability space contains an ergodic hyperfinite subgraph. A consequence of this is that every ergodic treeable equivalence relation has an ergodic hyperfinite free factor.

The pmp case of the main theorem was first proven by R. Tucker-Drob using a deep result from probability theory. Our proof is different: it is self-contained and applies more generally to quasi-pmp graphs. Among other things, it involves introducing a graph invariant concerning asymptotic averages of functions and a method of tiling a large part of the space with finite sets with prescribed properties. The non-pmp setting additionally exploits a new quasi-order called visibility to analyze the interplay between the Radon–Nikodym cocycle and the graph structure, providing a sufficient condition for hyperfiniteness.

CONTENTS

1. INTRODUCTION	2
Main results and applications	2
Auxiliary results	5
Sketch of proof of Main Lemma 1.6	7
Organization	8
Acknowledgments	8
2. PRELIMINARIES	9
2.A. Equivalence relations	9
2.B. Cocycles	10
2.C. Graphs	11
2.D. Quotients	12
3. FINITE AND HYPERFINITE AVERAGES	13
4. PROOFS OF THEOREMS 1.1, 1.3 AND 1.5 FROM MAIN LEMMA 1.6	15
5. MASS TRANSPORT ALONG A COCYCLE	17

Date: August 30, 2023.

2010 *Mathematics Subject Classification.* 37A30, 03E15, 05C63, 37A20, 37A25.

The author's research was partially supported by NSF Grant DMS-1501036, NSF grant DMS-1855648, and NSERC Discovery Grant RGPIN-2020-07120.

5.A. Preliminaries	17
5.B. Mass transport principle	18
5.C. Deficiency and w -invariant measures	19
5.D. Transporting a prescribed fraction of mass	19
6. HYPERFINITENESS VIA CUTS	20
7. PACKED AND APPROXIMATELY SATURATED TILINGS	21
7.A. Tilings	22
7.B. Packed tilings	22
7.C. Approximately saturated tilings	24
7.D. Packed and approximately saturated tilings	27
8. COCYCLED GRAPH VISIBILITY	27
8.A. Definitions and basic properties	28
8.B. Finite visibility and hyperfiniteness	28
8.C. Tiling with sets of large w -ratio	29
9. VISIBLE ASYMPTOTIC AVERAGES ON A GRAPH	31
9.A. For an abstract graph	32
9.B. For a measurable graph	33
10. PROOF OF MAIN LEMMA 1.6	35
10.A. Cutting one side of the set of asymptotic averages	35
10.B. Domains of packed tilings	36
10.C. Iterative tiling and its limsup	37
10.D. The conullness of D_∞ amounts to a packing condition	38
10.E. Verifying the packing condition	39
REFERENCES	41

1. INTRODUCTION

Main results and applications. We consider measurable actions $\Gamma \curvearrowright^a (X, \mu)$ of a countable group Γ on a standard probability space¹ (X, μ) . To avoid pathologies coming from null sets interfering with the dynamics, we assume that the actions are *quasi-pmp*² (aka nonsingular or null-preserving), i.e. each group element maps null sets to null sets. Intuitively, this means that points in the same orbit have possibly different relative weights, and this is captured precisely by the *Radon–Nikodym cocycle* of the orbit equivalence relation E_a with respect to μ , namely, a Borel function $w : E_a \rightarrow \mathbb{R}^+$, $(x, y) \mapsto w_x(y)$, such that

¹A standard Borel space X (i.e. the σ -algebra of X is the Borel σ -algebra of some Polish topology on X) equipped with a Borel probability measure.

²An action $\Gamma \curvearrowright (X, \mu)$ of a countable group on a standard probability space is *probability-measure-preserving* (pmp) if for each $\gamma \in \Gamma$, $\gamma_*\mu = \mu$.

- (i) w is a cocycle, i.e. $w_x(y)w_y(z) = w_x(z)$ for all E_a -related $x, y, z \in X$ (so indeed, one can think of $w_x(y)$ as the weight of y divided by that of x);
- (ii) $\mu(\gamma A) = \int_A w_x(\gamma x) d\mu(x)$ for all $\gamma \in \Gamma$ and measurable $A \subseteq X$.

In particular, μ is invariant under the action if and only if $w \equiv 1$. The existence and a.e. uniqueness of this cocycle is proven in [KM04, Section 8].

Seeking to prove a pointwise ergodic theorem in this generality, for all groups and quasi-pmp actions at once, we have to modify the classical form of pointwise ergodic theorems. Indeed, although the natural analogue of the classical pointwise ergodic theorem holds for quasi-pmp actions of \mathbb{Z}^d [Dow50, Fel07], it fails for some quasi-pmp action of $\bigoplus_{n \in \mathbb{N}} \mathbb{Z}$ along any sequence of finite subsets of $\bigoplus_{n \in \mathbb{N}} \mathbb{Z}$ [Hoc13, Theorems 1.2 and 1.3]. Even for pmp actions, although it holds for all amenable groups along tempered Følner sequences [Lin01], it fails for the free group \mathbb{F}_2 on 2 generators along spheres or balls [Tao15]. Of course, there are other versions of the pointwise ergodic theorem for pmp actions (e.g. [Buf00, FN98, TZ21]), but they too are specific to the group, while we would like a theorem for all groups and actions at once. So what we do is abandon the group and the action, and look at the induced Schreier graph instead (defined below).

By a *graph* on X , we mean a set of edges between the points in X , i.e. a symmetric subset of X^2 ; in particular, a *locally countable Borel graph* G on X is a symmetric Borel subset of X^2 whose each fiber is countable (i.e. each vertex has countably-many neighbors). We denote by E_G the G -connectedness equivalence relation, so for $x \in X$, $[x]_{E_G}$ is the G -connected component of x . When X is equipped with a Borel probability measure μ , we say that G is *quasi-pmp* if each Borel bijection $\gamma : X \rightarrow X$ with $\text{Graph}(\gamma) \subseteq E_G$ maps μ -null sets to μ -null sets.

Going back to the action $\Gamma \curvearrowright^a (X, \mu)$, for a symmetric generating set S of Γ , the *Schreier graph* $G_S \subseteq X^2$ of this action with respect to S is defined by

$$xG_S y :\Leftrightarrow \sigma x = y \text{ for some } \sigma \in S.$$

Assuming without loss of generality that the action a is Borel (measurable transformations are Borel mod null), G_S is a locally countable Borel graph. The Feldman–Moore theorem [FM77] implies that in fact *every* locally countable Borel graph G on a standard Borel space X arises in this fashion. Furthermore, the Schreier graph G_S is quasi-pmp if and only if the action a is quasi-pmp. Thus, although our main result (Theorem 1.1) is stated for an arbitrary locally countable quasi-pmp Borel graph G , one can safely think of G as a Schreier graph of some quasi-pmp action of a countable group.

Theorem 1.1 (Ergodic theorem for quasi-pmp graphs). *Let G be a locally countable quasi-pmp Borel graph on a standard probability space (X, μ) and let $w : E_G \rightarrow \mathbb{R}^+$ be the Radon–Nikodym cocycle of E_G with respect to μ . There is an increasing sequence (G_n) of component-finite Borel subgraphs of G (typically not adding up to G) such that for any $p \geq 1$ and $f \in L^p(X, \mu)$,*

$$\lim_{n \rightarrow \infty} \frac{1}{w_x([x]_{E_{G_n}})} \sum_{y \in [x]_{E_{G_n}}} f(y) w_x(y) = \mathbb{E}(f | \mathcal{B}_{E_G})(x) \quad \text{a.e. and in } L^p,$$

where $w_x([x]_{E_{G_n}}) := \sum_{y \in [x]_{E_{G_n}}} w_x(y)$ and $\mathbb{E}(f | \mathcal{B}_{E_G})$ is the conditional expectation of f with respect to the σ -algebra \mathcal{B}_{E_G} of E_G -invariant Borel sets.

This generalizes to quasi-pmp graphs the unpublished theorem of R. Tucker-Drob for pmp graphs, proven by completely different techniques. Indeed, while our proof is descriptive-set-theoretic and self-contained, Tucker-Drob’s proof is based on a deep result in probability theory: the indistinguishability of trees in the Wired Uniform Spanning Subforest [HN16, Theorem 1.1]; it also uses Wilson’s algorithm rooted at infinity as in [GL09, Proposition 9], as well as an analogue for graphs of the Abért–Weiss theorem [AW13, Theorem 1] derived by Tucker-Drob from [HLS14, Lemmas 7.9 and 7.10]). The mentioned results are all for the pmp case, so generalizing Tucker-Drob’s proof to the quasi-pmp setting would require generalizing these results as well.

Remark 1.2. Points to note about Theorem 1.1:

- (a) When G is an equivalence relation, i.e. the whole orbit equivalence relation of a group action, the conclusion of Theorem 1.1 has been known and is not very hard to prove. Indeed, the pmp case is explicitly stated and proven in [Kec10, Theorem 3.5], and the general quasi-pmp case can be easily extracted from earlier works, namely, by putting together [Sch77, Theorem 8.22] and the Hurewicz ergodic theorem.

From the perspective of measured group theory, the particular interest in proving this theorem for graphs, equivalently, Schreier graphs of group actions, is to at least have some involvement of the group itself: indeed, our sampling windows (the G_n -connected components) correspond to finite connected subsets of the Cayley graph of the group. From the descriptive-set-theoretic view point, when G is ergodic, the increasing union $\bigcup_n G_n$ is an ergodic hyperfinite subgraph (see Theorem 1.3 below). The existence of such a subgraph is a powerful tool, which has been sought after because of its immediate applications, e.g. Corollary 1.4 below.

- (b) When G is μ -hyperfinite³ (equivalently, E_G is μ -amenable, by the Connes–Feldman–Weiss theorem [CFW81]), the result is again not very hard. Indeed, discarding a null set, G is an increasing union of component-finite Borel graphs G_n and the conclusion of Theorem 1.1 follows from a (much simpler) pointwise ergodic theorem for hyperfinite equivalence relations. Versions of the pmp case of this theorem have appeared in the literature, for example, in [BN15] and in [MT17, Theorem 7.3], and we state it below as Theorem 3.6 for the general quasi-pmp case.

Thus, Theorem 1.1 is most valuable for graphs (as opposed to equivalence relations), which are not μ -hyperfinite/ μ -amenable.

Theorem 1.1 immediately implies what was the author’s main goal:

Theorem 1.3 (Ergodic hyperfinite subgraph). *Every ergodic locally countable Borel graph G on a standard probability space (X, μ) admits an ergodic hyperfinite Borel subgraph $H \subseteq G$.*

This immediately implies a positive answer to a question of L. Bowen, asked for pmp equivalence relations:

Corollary 1.4. *Every ergodic treeable⁴ countable Borel equivalence relation admits an ergodic hyperfinite free factor⁴.*

³ G is *hyperfinite* if it is an increasing union of component-finite Borel graphs. μ -*hyperfinite* just means hyperfinite off of a μ -null set.

⁴See [KM04, Sections 19 and 27] for the definitions of *treeable* and *free factor*.

Again, Theorem 1.3 and Corollary 1.4 generalize the corresponding unpublished results of R. Tucker-Drob in the pmp setting.

For an ergodic pmp graph G , the existence of an ergodic hyperfinite subgraph (i.e. Tucker-Drob's theorem) was initially also used in [MT17], although later the authors included a self-contained proof of a weaker statement that was sufficient for their purpose.

Lastly, the fact that Theorem 1.1 holds for quasi-pmp (and not just pmp) graphs, implies a ratio ergodic theorem, whose more general version without the ergodicity assumption is given in Theorem 4.1.

Theorem 1.5 (Ratio ergodic theorem for quasi-mp graphs). *Let G be a locally countable quasi-mp⁵ ergodic Borel graph on a σ -finite standard measure space (X, μ) and let $w : E_G \rightarrow \mathbb{R}^+$ be the Radon–Nikodym cocycle of E_G with respect to μ . There is an increasing sequence (G_n) of component-finite Borel subgraphs of G such that for any $f, g \in L^1(X, \mu)$ with $g > 0$,*

$$\lim_{n \rightarrow \infty} \frac{\sum_{y \in [x]_{E_{G_n}}} f(y) w_x(y)}{\sum_{y \in [x]_{E_{G_n}}} g(y) w_x(y)} = \frac{\int_X f d\mu}{\int_X g d\mu} \quad a.e.$$

As mentioned above, Theorems 1.3 and 1.5 are derived from Theorem 1.1, which itself is derived from the following lemma by a diagonalization argument. (See Section 4 for proofs of all these implications.)

Main Lemma 1.6. *Let G be a locally countable μ -nowhere hyperfinite⁶ quasi-pmp Borel graph on a standard probability space (X, μ) and let $w : E_G \rightarrow \mathbb{R}^+$ be the Radon–Nikodym cocycle of E_G with respect to μ . For any $f \in L^\infty(X, \mu)$ and $\varepsilon > 0$, there is a component-finite Borel subgraph $H \subseteq G$ such that for all x in a set of measure $\geq 1 - \varepsilon$, the average $\frac{1}{w_x([x]_{E_H})} \sum_{y \in [x]_{E_H}} f(y) w_x(y)$ differs from $\mathbb{E}(f | \mathcal{B}_{E_G})(x)$ by at most ε .*

The proof of this is the main content of the paper and we give a sketch of it below.

Auxiliary results. Here, we highlight some tools we obtained to prove Main Lemma 1.6 as they are interesting and may be useful elsewhere. Below, let G be a locally countable Borel graph on a standard Borel space X .

Cuts and hyperfiniteness. Call a set $V \subseteq X$ a *hyperfinitizing vertex-cut* for G if $G \cap (X \setminus V)^2$ is hyperfinite. For a Borel probability measure μ , put

$$\text{hvp}_\mu(G) := \inf \{ \mu(C) : C \subseteq X \text{ is a Borel hyperfinitizing vertex-cut for } G \}$$

and call it the *hyperfinitizing vertex-price* of G (with respect to μ). We also define the analogous notions for edge-cuts. This was already done in [MT17, Section 9] as well as earlier in [Ele07] in a slightly different context. The following is a useful and easily applicable way of exploiting the nonhyperfiniteness of a measurable graph and variations of it have appeared in the aforementioned two papers:

Proposition 1.7. *A locally countable Borel graph G is μ -hyperfinite if and only if $\text{hvp}_\mu(G) = 0$.*

⁵This is the same as quasi-pmp, but the measure is not assumed to be finite.

⁶This means that there is no E_G -invariant Borel set of positive μ -measure on which G is hyperfinite. Equivalently [JKL02, 1.3(vi)], there is no set of positive measure on which E_G is hyperfinite.

When G is locally finite, this proposition is merely an observation based on the Borel–Cantelli lemma. However, for locally countable graphs, the proof (still very easy) goes through the analogous statement for edge-cuts and this analogous statement immediately implies the Dye–Krieger theorem (Remark 6.6). See Proposition 6.5 for the full version of Proposition 1.7.

Proposition 1.7 is used to give a lower bound on the measure of a set based on the graph structure: for a μ -nonhyperfinite graph G , if a set is a hyperfinitizing vertex-cut for G then its measure is at least $\text{hvp}_\mu(G) > 0$.

Approximately saturated and packed tilings. Given a countable Borel equivalence relation E on a standard Borel space X and a Borel collection \mathcal{S} of finite E -related subsets⁷, one often needs a Borel tiling \mathcal{P} of a large part of X with tiles from \mathcal{S} . By [KM04, Lemma 7.3], maximal such tilings exist, but for our purposes here and for those in [MT17], a stronger notion of maximality is needed: each tile in \mathcal{P} should be maximally big (i.e. it cannot be combined with some untiled points to form a tile from \mathcal{S}) and tiles in \mathcal{P} cannot be combined together with proportionally-many untiled points to form a tile from \mathcal{S} . A tiling \mathcal{P} with the first property is called *saturated*, and with the second property, *p-packed*, where $p \in \mathbb{R}^+$ is the proportion parameter.

It was proven in [MT17, Subsection 4.D] that saturated and packed Borel tilings exist off of an E -compressible set and hence, off a null set for any E -invariant probability measure. Here, in Section 7, we generalize this to the quasi-pmp setting, i.e. in the presence of a Borel cocycle $w : E \rightarrow \mathbb{R}^+$. It is not true that saturated tilings exist off of a null set if the measure is not invariant: the fact that there are points of arbitrarily small relative w -weight is an issue. However, we define an approximate notion of saturation in Definition 7.9 and prove existence modulo null in Theorem 7.13. The proof of this theorem uses the existence of Borel label-maximizing maximal matchings in Borel bipartite graphs with edges labeled by positive reals, proven in Lemma 7.12.

Approximately saturated and packed (sequences of) tilings are related to and may be useful in the so-called *toast* constructions (see, e.g., [GR21]).

Cocycle-visibility in graphs and hyperfiniteness. Let G be a locally countable Borel graph on X and let $w : E_G \rightarrow \mathbb{R}^+$ be a Borel cocycle, i.e. $w_x(y)w_y(z) = w_x(z)$ for all E_G -related $x, y, z \in X$. A (G, w) -*visible neighborhood* of $x \in X$ is any G -connected set $V \ni x$ such that $w_x(v) \leq 1$ (the weight of x is at least as much as that of v) for each $v \in V$. This induces a quasi-order on X : $x \geq y$ if y is in a visible neighborhood of x . We say that G has *finite w -visibility* if for each $x \in X$, the downward cone $(\leq)^x := \{y \in X : y \leq x\}$ is w -finite, i.e. $\sum_{y \leq x} w_x(y) < \infty$. This provides a sufficient condition for Borel hyperfiniteness:

Theorem 1.8. *If G has finite w -visibility, then it is Borel hyperfinite.*

We use this to argue that a set $D \subseteq X$ is large: if removing D from a μ -nowhere hyperfinite graph G results in a graph with finite visibility, then D is a hyperfinitizing vertex-cut, whence $\mu(D) \geq \text{hvp}_\mu(G) > 0$.

Cocycle-ratio and tiling with arbitrarily large sets. Let E be a quasi-pmp Borel equivalence relation on a standard probability space (X, μ) and let $w : E \rightarrow \mathbb{R}^+$ be the Radon–Nikodym cocycle of E with respect to μ . We often need to find a μ -nowhere smooth hyperfinite

⁷A set is called *E-related* if it is contained in one E -class.

subequivalence relation $F \subseteq E$ (see Subsection 2.A for definitions) with some additional properties. For F to be μ -nowhere smooth, a.e. F -class $[x]_F$ has to be w -infinite, i.e. $\sum_{y \in [x]_F} w_x(y) = \infty$. In the pmp case, i.e. when $w \equiv 1$, we obtain F as an increasing union of finite Borel equivalence relations F_n . However, when w is arbitrarily, it could be that although the cardinality of the F_n -classes grows with n , they union up to a w -finite set. Therefore, cardinality of the F_n -classes is not the right measurement to control.

The solution is to look at what we call the w -ratio of an E -related finite set $U \subseteq X$, defined by $w_*(U) := \min_{x \in U} w_x(U)$, where $w_x(U) := \sum_{u \in U} w_x(u)$. Indeed, if the U_n are increasing E -related finite sets with $w_*(U_n) \rightarrow \infty$, then $\bigcup_n U_n$ is w -infinite.

To build desired F_n , we need to tile most of the space X with finite tiles of large w -ratio. This is surprisingly challenging because unlike cardinality, w -ratio is typically not monotone (under subsets). We build such tilings in Lemma 8.11.

Sketch of proof of Main Lemma 1.6. Let G , (X, μ) , w , and f be as in 1.6. To simplify notation, we assume that G is ergodic.

The invariant case. Here, we sketch the proof for a pmp G , i.e. $w \equiv 1$. We begin by establishing a connection (Lemma 3.2) between the global average (integral) of f and the local (finite) averages of f around a point: for any finite Borel equivalence relation F on X ,

$$\int_X f d\mu = \int_X A_F f d\mu, \quad (1.9)$$

where, for each $x \in X$, $A_F f(x)$ is the average of f over $[x]_F$.

Next, for each $x \in X$, we define the set $\mathcal{A}_G f(x)$ of G -asymptotic averages at x , namely, the set of all reals r that can be approximated arbitrarily well by the averages of f over finite G -connected sets $V \ni x$ of arbitrarily large cardinality. It is easy to see that the map $x \mapsto \mathcal{A}_G f(x)$ is E_G -invariant, hence constant a.e. by ergodicity. Moreover, this set, denoted by $\mathcal{A}_G f$, is a closed interval.

We then show that for each $\delta > 0$, one can construct a finite G -connected Borel equivalence relation F_δ such that for a.e. $x \in X$, the average of f over $[x]_{F_\delta}$ is at most δ -away from $\mathcal{A}_G f$. This implies, via (1.9), that $\mathcal{A}_G f$ contains the global average $\int_X f d\mu$. Thus, if the set $\mathcal{A}_G f$ was just a small interval around $\int_X f d\mu$ of size less than $\frac{\varepsilon}{2}$, taking $F := F_{\frac{\varepsilon}{2}}$ would satisfy the conclusion of Main Lemma 1.6.

Even if $\mathcal{A}_G f$ is initially a large interval, maybe quotienting out by some finite G -connected Borel equivalence relation shrinks it, in which case we would also be done. Thus, we assume that there is a $\delta > 0$ such that the set $\mathcal{A}_{G/R} f$ of asymptotic averages in the quotient graph G/R is not contained in

$$\overline{I}_\delta := [\int_X f d\mu - \delta, \int_X f d\mu + \delta],$$

for every finite G -connected Borel equivalence relation R . In fact, by an additional argument, we may assume that $\mathcal{A}_{G/R} f$ spills over both sides of \overline{I}_δ . This assumption allows us to tile a significant part of the space by G -connected finite tiles whose f -averages are in \overline{I}_δ for arbitrarily small $\delta > 0$. It is here that packed tilings come into play: the packedness condition ensures that only finite G -connected components are left after removing the union D of all the tiles from X . In other words, the domain D of each packed tiling is a finitizing vertex-cut. The μ -nowhere hyperfiniteness of G gives a lower bound $\lambda > 0$

for the measure of all finitizing vertex-cuts, hence domains of packed tilings. This allows us to eventually cover most of X by an iterative coherent construction of saturated and packed tilings, whose tiles become larger and larger and more and more packed. After sufficiently many iterations, the resulting tiling is such that the induced subgraphs on each tile together form a subgraph $H \subseteq G$ satisfying the conclusion of Main Lemma 1.6.

The quasi-invariant case. We only mention what changes one has to make in the above (pmp) argument to make it work in the quasi-pmp case.

To show that a set is null in the pmp setting, one often proves that it is compressible. The notion of compressibility was generalized to the quasi-invariant setting by Benjamin Miller in [Mil08], and we present a rephrasing of this in Section 5 in terms of mass transport.

As mentioned above, saturated tilings do not exist modulo null in the quasi-pmp setting, so we use approximately saturated tilings instead, proving their existence in Theorem 7.13.

For points x, y in the same E_G -class, the fact that $w_x(y)$ can be arbitrarily large destroys that convexity of the set $\mathcal{A}_G f(x)$ of G -asymptotic averages, while our construction above crucially relies on this property. To fix this, we introduce the notion of (G, w) -visibility (developed in Section 8) and take asymptotic averages at a point x only within the part of the graph that is visible to some point y that sees x .

Another important difference is in controlling the w -weight of increasing unions $\biguplus_n U_n$ of finite sets. Instead of ensuring that $|U_n| \rightarrow \infty$, we have to ensure that $w_*(U_n) \rightarrow \infty$, which is much harder since w_* is not monotone (under subsets).

Organization. Section 2 establishes notation and terminology that are globally used in the paper. In Section 3, we discuss finite and hyperfinite averages, in particular, stating the pointwise ergodic theorem for hyperfinite equivalence relations mentioned in Remark 1.2(b). Section 4 provides proofs of Theorems 1.1, 1.3 and 1.5 assuming Main Lemma 1.6. In Section 5, we discuss mass transport along a cocycled equivalence relation, introduce the notion of deficiency for sets as a generalization of compressibility, and provide a lemma for building Borel transport functions. Section 6 discusses cuts in a graph and their connection with hyperfiniteness. Section 7 introduces saturated and packed tilings with respect to a cocycle and proves their existence. Section 8 discusses the notion of cocycle-visibility in a graph, provides a sufficient condition for hyperfiniteness (Theorem 1.8), and proves the lemma on tiling the space with sets of large cocycle-ratio (Lemma 8.11). In Section 9, we introduce the set of asymptotic averages for a graph, whose role is instrumental for the proof of Main Lemma 1.6; we then establish an important tiling lemma for the set of asymptotic averages. Finally, Section 10 is the proof of Main Lemma 1.6.

Acknowledgments. The author thanks

- Benjamin Miller for getting her into this topic and way of thinking.
- Robin Tucker-Drob for sharing his proof of the pmp case of Theorem 1.3 and for insightful conversations.
- The two anonymous referees for providing incredibly helpful and detailed reports, which, among other things, provided a more streamlined and insightful argument for Theorem 1.8 and pinpointed an error in the use of saturated tilings.
- Peter Burton for providing a number of references, suggesting that the reduction from L^1 to L^∞ should be explained, and asking about and verifying Theorem 1.5.

- Anton Bernshteyn for pointing out an error in the derivation of Theorem 1.1 from Main Lemma 1.6.
- Ran Tao for noticing an oversight in the definition of packed tilings, and for suggesting a more straightforward proof of Lemma 5.8.
- Lewis Bowen for useful remarks and references.
- Benjamin Weiss for pinpointing [Sch77, Theorem 8.22] and prompting Remark 1.2(a).
- Clinton Conley, Alexander Kechris, Andrew Marks, and Jenna Zomback for helpful conversations and suggestions.

2. PRELIMINARIES

Our set \mathbb{N} of natural numbers includes 0. For reals $a, b \in \mathbb{R}$, we write $a \approx_\varepsilon b$ to mean that $|a - b| \leq \varepsilon$. For $i = 1, 2$, $\text{proj}_i : X_1 \times X_2 \rightarrow X_i$ is defined by $(x_1, x_2) \mapsto x_i$. For a set X , let $\mathcal{P}(X)$ denote the powerset of X , and let Id_X denote the identity (equality) equivalence relation on X .

Throughout, let X be a standard Borel space. For a Borel measure μ on X and $\varepsilon > 0$, we call a set $X' \subseteq X$ μ -co- ε if it is measurable and $\mu(X \setminus X') \leq \varepsilon$.

2.A. Equivalence relations. Let E denote a *countable Borel* equivalence relation on X , that is: E is a Borel subset of X^2 and each E -class is countable. We refer to [JKL02] and [KM04] for the general theory of countable equivalence relations.

For $x \in X$ and $A \subseteq X$, we write $[x]_E$ to mean the E -class of x and $[A]_E$ to mean the E -saturation of A , i.e. $\bigcup_{x \in A} [x]_E$. We say that a set $A \subseteq X$ is E -related if it is contained in a single E -class; similarly, we say that points $x_0, x_1, \dots, x_n \in X$ are E -related if $\{x_0, x_1, \dots, x_n\}$ is E -related. We denote by $[X]_E^{<\infty}$ the standard Borel space of finite nonempty E -related sets.

Let μ be a Borel measure on X . We say that E is

- *smooth* if for some/any Polish space Y , there is a Borel map $f : X \rightarrow Y$ such that for all $x, y \in X$, $xEy \iff f(x) = f(y)$. In fact, one can take $Y := X$ and have in addition that $xEf(x)$, so f is a *Borel selector* for E .
- *finite* if each E -class is finite. Note that any finite Borel equivalence relation is smooth because, by Luzin–Novikov uniformization [Kec95, Theorem 18.10], $x \mapsto \min_{<} [x]_E$ is a Borel selector for E , where $<$ is some a priori fixed Borel linear order on X .
- *(Borel) hyperfinite* if E is an increasing union of finite Borel equivalence relations.
- μ -*hyperfinite* if it is hyperfinite on a conull set; this set can be chosen to be Borel and E -invariant, by [JKL02, 1.3(vi)].
- μ -*nowhere hyperfinite* if there is no set of positive measure on which E is hyperfinite; again by [JKL02, 1.3(vi)], this is equivalent to the inexistence of an E -invariant Borel set of positive μ -measure on which E is hyperfinite.
- *measure-preserving (mp)* or that μ is E -invariant if $\gamma_*\mu = \mu$ for every Borel automorphism γ of E (i.e. a Borel bijection $X \rightarrow X$ mapping every point to an E -equivalent point).
- *quasi-measure-preserving (quasi-mp)* or that μ is E -quasi-invariant if $\gamma_*\mu \sim \mu$ for every Borel automorphism γ of E .
- *pmp* (resp. *quasi-pmp*) if it is mp (resp. quasi-mp) and μ is a probability measure.

2.B. Cocycles. For a countable Borel equivalence relation E on X , a *cocycle on E* is a map $w : E \rightarrow \mathbb{R}^+$, $(x, y) \mapsto w_x(y)$ satisfying the *cocycle identity*: $w_x(y) \cdot w_y(z) = w_x(z)$ for all E -related $x, y, z \in X$. A Borel measure μ on X is called *w-invariant* if for every Borel set $B \subseteq X$ and a Borel automorphism γ of E ,

$$\mu(\gamma B) = \int_B w_x(\gamma x) d\mu(x).$$

By [KM04, Section 8], every E -quasi-invariant probability measure μ is w -invariant for some Borel cocycle w on E . Such a cocycle is clearly unique μ -a.e., and it is called the *Radon–Nikodym cocycle of E with respect to μ* .

For a Borel cocycle $w : E \rightarrow \mathbb{R}^+$ and $x \in X$, the function $w_x : [x]_E \rightarrow \mathbb{R}^+$ defines a measure on $Y := [x]_E$ by $w_x(A) := \sum_{y \in A} w_x(y)$ for each $A \subseteq Y$. Thus, it makes sense to write $\int_Y f dw_x$ for any absolutely summable $f : Y \rightarrow \mathbb{R}$.

Independence of the base point. As mentioned in the introduction, we think of $w_x(y)$ as the weight of y divided by that of x ; indeed, in expressions like $\frac{w_x(y)}{w_x(z)}$ and statements like “ $\lim_{n \rightarrow \infty} w_x(x_n) \rightarrow \infty$ ” the particular choice of the base point x does not matter due to the cocycle identity. We call such expressions and statements *w-homogeneous* and omit writing the subscript x from them. For example, for an E -class C , $A \subseteq C$, and $f : C \rightarrow \mathbb{R}$, we may write $\frac{2 \cdot w(x) + \int_C f dw}{w(A)}$ and “ $w(A)$ is finite”. We also say that *all w-large enough* $U \subseteq C$ satisfy some property \mathcal{P} if for any/some base point $x \in C$, there is $L_x > 0$ such that all $U \subseteq C$ with $w_x(U) \geq L_x$ satisfy \mathcal{P} . *Arbitrarily w-large* is defined analogously.

Maximum and minimum. For a subset A of an E -class, put

$$\text{Min}_w A := \{x \in A : \forall y \in A \ w(x) \leq w(y)\}.$$

If $\text{Min}_w A \neq \emptyset$, for any $x \in [A]_E$, we put $\min_{w_x} A := \min_{y \in A} w_x(y)$, otherwise, $\min_{w_x} A := 0$. We also analogously define $\text{Max}_w A$ and $\max_{w_x} A$, and we omit the base point x in w -homogeneous expressions.

The space $[X]_E^{w<\infty}$ of w-finite sets. An E -related set A is said to be *w-finite* if $w(A)$ is finite, otherwise, it is *w-infinite*. We denote by $[X]_E^{w<\infty}$ the collection of all nonempty E -related w -finite sets. We say that E is *w-finite*, if each E -class is w -finite; otherwise, we say that it is *w-infinite* (aka *w-aperiodic*).

We show that $[X]_E^{w<\infty}$ is also a standard Borel space, which can be naturally viewed as a Borel subset of $X^{<\mathbb{N}} \cup X^{\mathbb{N}}$. Throughout, we fix a Borel linear order $<_X$ on X and define a partial order $<_w$ (linear on every E -class) as follows: for any $x, y \in X$

$$x <_w y : \Leftrightarrow xEy \text{ and } (w(x) < w(y) \text{ or } (w(x) = w(y) \text{ and } x <_X y)).$$

Observation 2.1. For each $A \in [X]_E^{w<\infty}$ and $a_0 \in A$, the set $\{a \in A : a >_w a_0\}$ is finite. In other words, the relation $>_w$ on $A \in [X]_E^{w<\infty}$ is a well-order of type $\leq \omega$. In particular, $\text{Max}_w A$ is nonempty and finite.

Thus, we identify $[X]_E^{w<\infty}$ with the set of all $<_w$ -decreasing E -related sequences $(x_n)_{n<\ell} \subseteq X$, $\ell \leq \omega$, that are *w-summable*, i.e. $\sum_{n<\ell} w(x_n) < \infty$. This is clearly a Borel subset of $X^{<\mathbb{N}} \cup X^{\mathbb{N}}$. Furthermore, we view $[X]_E^{w<\infty}$ as a Borel subset of $[X]_E^{w<\infty}$.

In our arguments below, we implicitly use the following.

Observation 2.2. The set $\{(x, U) \in X \times [X]_E^{\mathfrak{w} < \infty} : x \in U\}$ is a Borel subset of $X \times [X]_E^{\mathfrak{w} < \infty}$.

The partial order $>_{\mathfrak{w}}$ induces a (lexicographic) partial order $>_{\mathfrak{w}}^*$ on $[X]_E^{\mathfrak{w} < \infty}$ as follows: for any $A, B \in [X]_E^{\mathfrak{w} < \infty}$, $A >_{\mathfrak{w}}^* B$ if and only if $A \neq B$ yet $[A]_E = [B]_E$ and

$$|A| < |B| \text{ or } (|A| = |B| \text{ and } \max_{<_{\mathfrak{w}}} (A \triangle B) \in A),$$

where $\max_{<_{\mathfrak{w}}} (A \triangle B)$ is the maximum element of $A \triangle B$ with respect to $<_{\mathfrak{w}}$. For each E -class C , $>_{\mathfrak{w}}^*$ is a linear order on $[C]_E^{\mathfrak{w} < \infty}$. Moreover, because $>_{\mathfrak{w}}$ is a well-order on any $A \in [X]_E^{\mathfrak{w} < \infty}$, it is not hard to check that $>_{\mathfrak{w}}^*$ is also a well-order on $[A]_E^{\mathfrak{w} < \infty}$. Thus:

Observation 2.3. For every \mathfrak{w} -finite E -class C , $>_{\mathfrak{w}}^*$ is a well-order on $[C]_E^{\mathfrak{w} < \infty}$.

\mathfrak{w} -ratio of E -related sets. As mentioned in the introduction, in some constructions below we need to ensure that an increasing union of finite (or \mathfrak{w} -finite) sets is \mathfrak{w} -infinite, and we noted that just making the cardinality of the sets grow is not enough. However, replacing cardinality with the following works:

Definition 2.4. For each $U \in [X]_E^{\mathfrak{w} < \infty}$, Observation 2.1 implies that

$$\mathfrak{w}_*(U) := \frac{\mathfrak{w}(U)}{\max_{\mathfrak{w}} U} = \min_{x \in U} \mathfrak{w}_x(U),$$

is well-defined and we call it the \mathfrak{w} -ratio of U .

Although $U \subseteq V$ does not imply $\mathfrak{w}_*(U) \leq \mathfrak{w}_*(V)$, we still get the desired property:

Observation 2.5. For any $U, V \in [X]_E^{\mathfrak{w} < \infty}$, $U \subseteq V$ implies $\frac{\mathfrak{w}_*(V)}{\mathfrak{w}_*(U)} \leq \frac{\mathfrak{w}(V)}{\mathfrak{w}(U)}$. In particular, for any increasing sequence (U_n) of sets in $[X]_E^{\mathfrak{w} < \infty}$, $\mathfrak{w}_*(U_n) \rightarrow \infty$ implies $\mathfrak{w}(U_n) \rightarrow \infty$.

When the increasing union is \mathfrak{w} -finite, \mathfrak{w}_* is monotone:

Lemma 2.6. For any increasing sequence (U_n) of sets in $[X]_E^{\mathfrak{w} < \infty}$, if $U := \bigcup_n U_n$ is \mathfrak{w} -finite, then $\text{Max}_{\mathfrak{w}} U \subseteq U_n$ for all large enough n . In particular, the sequence $(\mathfrak{w}_*(U_n))$ is eventually increasing and $\lim_n \mathfrak{w}_*(U_n) = \mathfrak{w}_*(U)$.

Proof. By Observation 2.1, $\text{Max}_{\mathfrak{w}} U$ is finite, so there is N such that $\text{Max}_{\mathfrak{w}} U_N = \text{Max}_{\mathfrak{w}} U$. Thus, for any $x \in \text{Max}_{\mathfrak{w}} U$ and all $n \geq N$,

$$\mathfrak{w}_*(U_n) = \mathfrak{w}_x(U_n) \nearrow \mathfrak{w}_x(U) = \mathfrak{w}_*(U). \quad \square$$

2.C. Graphs. Let \vec{G} be a locally countable directed Borel graph on X , i.e. \vec{G} is a Borel subset of X^2 such that the fibers $\vec{G}_x := \{y \in X : (x, y) \in \vec{G}\}$ and $\vec{G}^x := \{y \in X : (y, x) \in \vec{G}\}$ are countable for each $x \in X$. We drop “directed” if \vec{G} is *symmetric*, i.e. $\vec{G} = -\vec{G} := \{(y, x) : (x, y) \in \vec{G}\}$. Let G denote the *symmetrization* of \vec{G} , i.e. $G := \vec{G} \cup (-\vec{G})$. For sets $A, B \subseteq X$, the following is standard notation:

- $(A, B)_{\vec{G}} := (A \times B) \cap \vec{G}$;
- $[A, B]_{\vec{G}} := (A \times B \cup B \times A) \cap \vec{G}$;
- $\vec{G}|_A := \vec{G} \cap A^2$;
- $\vec{G}_{-A} := \vec{G}|_{X \setminus A}$;

- $\partial_{\vec{G}}A := \partial_G A := \{a \in A : \text{there is } b \in X \setminus A \text{ with } (a, b) \in G\}$, the *inner \vec{G} -boundary* of A ;
- $\overline{\partial}_{\vec{G}}A := \overline{\partial}_G A := \{b \in X \setminus A : \text{there is } a \in A \text{ with } (a, b) \in G\}$, the *outer \vec{G} -boundary* of A .

We denote by E_G the equivalence relation of being in the same G -connected component, and put $E_{\vec{G}} := E_G$. The rest of the terminology and most of the paper is about a symmetric graph G . Call a set $U \subseteq X$ *G -connected* if $G \cap U^2$ is a connected graph on U , and let $[X]_G^{<\infty}$ denote the (Borel) subset of $[X]_{E_G}^{<\infty}$ of G -connected sets. Similarly, for a Borel cocycle $w : E_G \rightarrow \mathbb{R}^+$, let $[X]_G^{w<\infty}$ denote the corresponding subset of $[X]_{E_G}^{w<\infty}$. Say that an equivalence relation F on X is *G -connected* if each F -class is G -connected.

We say that G is *component-finite* if each G -connected component is finite. For a Borel measure μ on X , we say that G is *(quasi-)pmp*, *μ -ergodic*, *Borel hyperfinite*, *μ -hyperfinite*, etc. if E_G is. In particular, if G is hyperfinite, i.e. E_G is a countable increasing union of finite Borel equivalence relations E_n , then G is a countable increasing union of component-finite Borel graphs $G_n := E_n \cap G$.

2.D. Quotients. For a smooth countable Borel equivalence relation F on X , the quotient space X/F is also standard Borel and we denote it by X/F . We emphasize that although X/F can be realized as a Borel subset of X , but we think of it as the set of F -classes. We denote the quotient map by $\pi_F : X \twoheadrightarrow X/F$.

For an F -invariant function $f : X \rightarrow \mathbb{R}$, we define its quotient $X/F \rightarrow \mathbb{R}$ by $[x]_F \mapsto f(x)$ and we still denote it by f to not overload notation. For a Borel measure μ on X , we put $\mu_{/F} := (\pi_F)_* \mu$. For a countable Borel equivalence relation $E \supseteq F$, we denote by $E_{/F}$ the pushforward of E under π_F . For $\mathcal{S} \subseteq \mathcal{P}(X)$, let

$$\mathcal{S}_{/F} := \{P \in \mathcal{P}(X/F) : \pi_F^{-1}(P) \in \mathcal{S}\}.$$

For a locally countable Borel graph G on X with $F \subseteq E_G$, we form the *quotient graph* $G_{/F}$ by contracting the edges between F -related vertices, i.e. for $U, V \in X/F$,

$$(U, V) \in G_{/F} :\Leftrightarrow [U, V]_G \neq \emptyset.$$

In our arguments, we need $\pi_F^{-1}(A)$ to be G -connected for every $G_{/F}$ -connected set $A \subseteq X/F$; this happens exactly when F is G -connected, so we only take quotients of G by G -connected equivalence relations.

Moreover, we only take quotients by w -finite F for a Borel cocycle $w : F \rightarrow \mathbb{R}^+$, which is allowed by the following:

Lemma 2.7. *For any countable Borel equivalence relation F and Borel cocycle $w : F \rightarrow \mathbb{R}^+$, if F is w -finite then it is smooth.*

Proof. Follows from Observation 2.1 because the map $x \mapsto$ the $<_w$ -largest element of $[x]_F$ is a Borel selector for F . \square

Lastly, let E be a quasi-pmp equivalence relation on a standard probability space (X, μ) and let $w : E \rightarrow \mathbb{R}^+$ be the Radon–Nikodym cocycle of E with respect to μ .

Observation 2.8. *For any w -finite Borel subequivalence relation $F \subseteq E$, $\mu_{/F}$ is $E_{/F}$ -quasi-invariant, and the Radon–Nikodym cocycle $w_{/F}$ of $E_{/F}$ with respect to $\mu_{/F}$ is given by*

$$(w_{/F})_{[x]_F}([y]_F) := w([y]_F)/w([x]_F)$$

for any $(x, y) \in E$. In particular, for any $U \in [X/F]_{E_{/F}}^{w_{/F}<\infty}$, $(w_{/F})_*(U) \leq w_*(\pi_F^{-1}(U))$.

3. FINITE AND HYPERFINITE AVERAGES

This subsection is the generalization of [MT17, Subsections 7.A–B] to the quasi-invariant setting. We first state a basic fact about finite averages, which we use repeatedly below. For an abstract set Y , treating Y^2 as the trivial equivalence relation on Y , we let $w : Y^2 \rightarrow \mathbb{R}^+$ be a cocycle, and for a bounded function $f : Y \rightarrow \mathbb{R}$ and a w -finite nonempty set $U \subseteq Y$, we define the w -weighted average of f on U by

$$A_U^w f := \frac{\sum_{x \in U} f(x)w(x)}{w(U)}.$$

Lemma 3.1. *For Y, w and f as above, and for nonempty disjoint w -finite sets $U, V \subseteq Y$,*

- (a) $A_{U \cup V}^w f = \frac{w(U)}{w(U)+w(V)} A_U^w f + \frac{w(V)}{w(U)+w(V)} A_V^w f.$
- (b) $|A_{U \cup V}^w f - A_U^w f| \leq 2 \|f\|_\infty \frac{w(V)}{w(U)+w(V)} \leq 2 \|f\|_\infty \frac{w(V)}{w(U)}.$

Proof. One verifies (a) directly, and (b) follows from (a) by the triangle inequality. \square

Now let (X, μ) be a standard probability space. For finite and hyperfinite equivalence relations on X , we now give explicit formulas for the conditional expectation of any $f \in L^1(X, \mu)$ with respect to the σ -algebras of invariant sets.

The following simple statement is what connects local finite averages to the global average (i.e. the integral). The author has been calling statements of this type *local-global bridges* and has been using them in all her proofs of pointwise ergodic theorems as they reduce proving these theorems to solving local combinatorial problems.

Lemma 3.2 (Local-global bridge). *Let F be a quasi-pmp equivalence relation on (X, μ) and let $w : F \rightarrow \mathbb{R}^+$ be its Radon–Nikodym cocycle with respect to μ . If F is w -finite, then for any $f \in L^1(X, \mu)$, the function*

$$A_F^w f : X \rightarrow \mathbb{R} \text{ defined by } x \mapsto A_{[x]_F}^w f$$

is the conditional expectation of f with respect to the σ -algebra of F -invariant Borel sets, i.e.

$$\int_Y f d\mu = \int_Y A_F^w f d\mu, \tag{3.3}$$

for any F -invariant measurable set $Y \subseteq X$. In particular, $\|A_F^w f\|_p \leq \|f\|_p$ for any $p \geq 1$.

Proof. The inequality $\|A_F^w f\|_p \leq \|f\|_p$ is true in general for any conditional expectation [Dur19, Theorem 4.1.11], but in this case it trivially follows from Jensen's inequality $|A_F^w f|^p \leq A_F^w |f|^p$ and (3.3) applied to $|f|^p$.

As for (3.3), it is enough to prove it assuming that $Y = X$ and all F -classes have the same cardinality n for some fixed $n \leq |\mathbb{N}|$. To this end, let S be the set of $<_w$ -maximum elements in each F -class (which exist by Observation 2.1), so S meets every F -class in exactly one point. Using this set S and Luzin–Novikov uniformization, we get a Borel automorphism $T : X \rightarrow X$ that induces F , i.e. $[x]_F = \{T^i x\}_{i \in \mathbb{Z}}$ for every $x \in X$. Put $I := \{0, 1, \dots, n-1\}$ if n is finite, and $I := \mathbb{Z}$, otherwise, so for each $x \in X$, the sequence $(T^i x)_{i \in I}$ lists all elements of

$[x]_F$ without repetition. Then:

$$\begin{aligned}
\int_X f \, d\mu &= \sum_{i \in I} \int_{T^i(S)} f(x) \, d\mu(x) \\
\left[\text{change of variable } x \text{ to } T^i x \right] &= \sum_{i \in I} \int_S f(T^i x) \cdot w_x(T^i x) \, d\mu(x) \\
&= \int_S \sum_{i \in I} f(T^i x) \cdot w_x(T^i x) \, d\mu(x) \\
&= \int_S \sum_{i \in I} A_F^w f(x) \cdot w_x(T^i x) \, d\mu(x) \\
\left[A_F^w f \text{ is } T\text{-invariant} \right] &= \sum_{i \in I} \int_S A_F^w f(T^i x) \cdot w_x(T^i x) \, d\mu(x) \\
\left[\text{change of variable } x \text{ to } T^{-i} x \right] &= \sum_{i \in I} \int_{T^i(S)} A_F^w f(x) \, d\mu(x) = \int_X A_F^w f \, d\mu. \quad \square
\end{aligned}$$

The following statement is the analogue of the maximal inequality, which makes it enough to prove Theorem 1.1 for bounded functions.

Lemma 3.4 (Approximate L^∞ -continuity). *Let F be a quasi-pmp equivalence relation on (X, μ) and let $w : F \rightarrow \mathbb{R}^+$ be its Radon–Nikodym cocycle with respect to μ . If F is w -finite, then for each $\varepsilon > 0$, there is an F -invariant μ -co- ε set $X' \subseteq X$ such that*

$$\|(A_F^w f)|_{X'}\|_\infty \leq \frac{1}{\varepsilon} \|f\|_1.$$

We need the following basic lemma for the proof.

Lemma 3.5. *For any $f \in L^1(X, \mu)$ and $\varepsilon > 0$, there is a μ -co- ε set $X' \subseteq X$ (of the form $\{x \in X : |f(x)| \leq r\}$ for some $r \geq 0\}$) such that $\|f|_{X'}\|_\infty \leq \frac{1}{\varepsilon} \|f\|_1$.*

Proof. Letting $r_\varepsilon := \inf\{r \geq 0 : \mu(f > r) \leq \varepsilon\}$, we show that $X' := \{f \leq r_\varepsilon\}$ works. Then $\{f > r_\varepsilon\} = \bigcup_{n \geq 1} \{f > r_\varepsilon + \frac{1}{n}\}$, so $\mu(\{f > r_\varepsilon\}) \leq \varepsilon$, hence X' is μ -co- ε . On the other hand, $\mu(f \geq r_\varepsilon) \geq \varepsilon$ because

$$\{f \geq r_\varepsilon\} = \bigcap_{n \geq 1} \left\{ f > r_\varepsilon - \frac{1}{n} \right\},$$

and $\mu(\{f > r_\varepsilon - \frac{1}{n}\}) > \varepsilon$ for each $n \geq 1$. Thus,

$$\|f|_{X'}\|_\infty \cdot \varepsilon \leq r_\varepsilon \cdot \mu(f \geq r_\varepsilon) \leq \|f\|_1. \quad \square$$

Proof of Lemma 3.4. By Lemma 3.2, $\|A_F^w f\|_1 \leq \|f\|_1 < \infty$. Thus, Lemma 3.5 applies to $A_F^w f$ and gives an F -invariant μ -co- ε set $X' \subseteq X$ such that $\|(A_F^w f)|_{X'}\|_\infty \leq \frac{1}{\varepsilon} \|A_F^w f\|_1 \leq \frac{1}{\varepsilon} \|f\|_1$. \square

Theorem 3.6 (Pointwise ergodic theorem for hyperfinite equivalence relations). *Let F be a Borel quasi-pmp hyperfinite equivalence relation on (X, μ) and let $w : F \rightarrow \mathbb{R}^+$ be its Radon–Nikodym cocycle with respect to μ . For any $f \in L^1(X, \mu)$ and any increasing sequence (F_n) of finite Borel equivalence relations with $F = \bigcup_n F_n$, the pointwise limit*

$$A_F^w f := \lim_{n \rightarrow \infty} A_{F_n}^w f$$

exists a.e. and is equal to the conditional expectation of f with respect to the σ -algebra of F -invariant measurable sets, i.e.

$$(a) \int_Y A_F^w f d\mu = \int_Y f d\mu \text{ for any } F\text{-invariant measurable set } Y \subseteq X.$$

In particular,

(b) Up to null sets, the above limit does not depend on the choice of (F_n) .

(c) $\|A_F^w f\|_p \leq \|f\|_p$ for every $p \geq 1$.

(d) F is ergodic if and only if for every $f \in L^1(X, \mu)$, $A_F^w f = \int_X f d\mu$ a.e.

Proof. The main part, (a), is just a version of [MT17, Theorem 7.3] in the quasi-invariant setting, and the proof is the same, given Lemma 3.2, so we omit it. Parts (b) and (d) are immediate from (a). Part (c) is a well-known fact about conditional expectation [Dur19, Theorem 4.1.11], and it can be seen directly here. \square

4. PROOFS OF THEOREMS 1.1, 1.3 AND 1.5 FROM MAIN LEMMA 1.6

We restate the statement of Main Lemma 1.6 using the introduced terminology.

Main Lemma 1.6. *Let G be a locally countable μ -nowhere hyperfinite quasi-pmp Borel graph on a standard probability space (X, μ) and let $w : E_G \rightarrow \mathbb{R}^+$ be the Radon–Nikodym cocycle of E_G with respect to μ . For any $f \in L^\infty(X, \mu)$ and $\varepsilon > 0$, there is a component-finite Borel subgraph $H \subseteq G$ such that $A_{E_H}^w f \approx_\varepsilon \mathbb{E}(f|\mathcal{B}_{E_G})$ on a μ -co- ε set.*

Assuming this lemma, we now derive Theorems 1.1, 1.3 and 4.1, restating them here using the introduced terminology.

Theorem 1.1 (Ergodic theorem for quasi-pmp graphs). *Let G be a locally countable quasi-pmp Borel graph on a standard probability space (X, μ) and let $w : E_G \rightarrow \mathbb{R}^+$ be the Radon–Nikodym cocycle of E_G with respect to μ . There is an increasing sequence (F_n) of G -connected finite Borel subequivalence relations of E_G such that for any $p \geq 1$ and $f \in L^p(X, \mu)$,*

$$\lim_{n \rightarrow \infty} A_{F_n}^w f = \mathbb{E}(f|\mathcal{B}_{E_G}) \text{ a.e. and in } L^p.$$

Proof. We first prove convergence in L^p assuming the a.e. convergence statement. Fix $f \in L^p(X, \mu)$ and $\varepsilon > 0$, and let $\hat{f} \in L^\infty(X, \mu)$ be such that $\|f - \hat{f}\|_p < \frac{\varepsilon}{3}$. [Dur19, Theorem 4.1.11] implies that $\|\mathbb{E}(f|\mathcal{B}_{E_G}) - \mathbb{E}(\hat{f}|\mathcal{B}_{E_G})\|_p \leq \|f - \hat{f}\|_p < \frac{\varepsilon}{3}$ and similarly, Lemma 3.2 implies that $\|A_{F_n}^w f - A_{F_n}^w \hat{f}\|_p \leq \|f - \hat{f}\|_p < \frac{\varepsilon}{3}$ for any $n \in \mathbb{N}$. Because $|A_{F_n}^w \hat{f}| \leq \|\hat{f}\|_\infty$, it follows from the Dominated Convergence theorem and the a.e. convergence of $A_{F_n}^w \hat{f}$ to $\mathbb{E}(\hat{f}|\mathcal{B}_{E_G})$ that $A_{F_n}^w \hat{f} \rightarrow \mathbb{E}(\hat{f}|\mathcal{B}_{E_G})$ in L^p , so for all large enough n , $\|A_{F_n}^w \hat{f} - \mathbb{E}(\hat{f}|\mathcal{B}_{E_G})\|_p < \frac{\varepsilon}{3}$. Thus, $\|A_{F_n}^w f - \mathbb{E}(f|\mathcal{B}_{E_G})\|_p < \varepsilon$ by the triangle inequality.

We now prove the a.e. convergence statement for a fixed $f \in L^1(X, \mu)$. If G is μ -hyperfinite, then Theorem 3.6 implies that any increasing sequence (F_n) of G -connected finite Borel equivalence relations with $E_G = \bigcup_n F_n$ satisfies the desired conclusion. Thus, we assume that G is not μ -hyperfinite; in fact, by discarding a maximal (up to μ -null sets) E_G -invariant Borel set on which G is hyperfinite, we may assume that G is μ -nowhere hyperfinite.

Let $\mathcal{D} \subseteq L^\infty(X, \mu)$ be a countable set that is dense in $L^1(X, \mu)$, and let $(f_n)_{n \geq 1}$ be an enumeration of \mathcal{D} such that each $f \in \mathcal{D}$ is equal to f_n for infinitely-many $n \in \mathbb{N}$. Furthermore, let $(\varepsilon_n)_{n \geq 1}$ be a sequence of positive reals converging to 0. We inductively build an increasing sequence (F_n) of G -connected finite Borel equivalence relations by taking $F_0 := \text{Id}_X$

and letting F_{n+1} be the pullback to X of $E_{H_{n+1}}$, where H_{n+1} is a component-finite Borel subgraph of $G_{/F_n}$ given by Main Lemma 1.6 when applied to the quotient graph $G_{/F_n}$ on $(X_{/F_n}, \mu_{/F_n})$, the function $A_{F_n}^{\mathbb{W}} f_{n+1}$, and $\varepsilon := \varepsilon_{n+1}$. Thus, for each $n \geq 1$, $A_{F_n}^{\mathbb{W}} f_n \approx_{\varepsilon_n} \mathbb{E}(f_n | \mathcal{B}_{E_G})$ on an F_n -invariant μ -co- ε_n set $X_n \subseteq X$.

Now fix $f \in L^1(X, \mu)$. By Theorem 3.6(b) applied to $F := \bigcup_n F_n$, it is enough to extract a subsequence (F_{n_k}) such that

$$\lim_{n \rightarrow \infty} A_{F_{n_k}}^{\mathbb{W}} f = \mathbb{E}(f | \mathcal{B}_{E_G}) \text{ a.e.}$$

By the Borel–Cantelli lemma, it is enough to show that for any $\varepsilon \in (0, 1)$, there is $n \in \mathbb{N}$ and an F_n -invariant μ -co- ε set $X' \subseteq X$ such that $A_{F_n}^{\mathbb{W}} f \approx_{\frac{\varepsilon}{3}} \mathbb{E}(f | \mathcal{B}_{E_G})$ for all $x \in X'$.

Because there is a subsequence (f_{n_k}) converging to f in L^1 and conditional expectation is an L^1 -contraction [Dur19, Theorem 4.1.11], we also have that $\mathbb{E}(f_{n_k} | \mathcal{B}_{E_G}) \rightarrow \mathbb{E}(f | \mathcal{B}_{E_G})$ in L^1 , so there is a further subsequence that converges a.e. Thus, for any fixed ε , there is $n \in \mathbb{N}$ such that ε_n and $\|f - f_n\|_1$ are both less than $\frac{\varepsilon^2}{3}$, and $\mathbb{E}(f_n | \mathcal{B}_{E_G}) \approx_{\frac{\varepsilon}{3}} \mathbb{E}(f | \mathcal{B}_{E_G})$ on a μ -co- $\frac{\varepsilon}{3}$ set. By Lemma 3.4,

$$|A_{F_n}^{\mathbb{W}} f_n - A_{F_n}^{\mathbb{W}} f| \leq A_{F_n}^{\mathbb{W}} |f_n - f| \leq \frac{1}{\varepsilon} \|f_n - f\|_1 < \frac{\varepsilon}{3},$$

on a μ -co- $\frac{\varepsilon}{3}$ set. Thus, by the triangle inequality, $A_{F_n}^{\mathbb{W}} f \approx_{\varepsilon} \mathbb{E}(f | \mathcal{B}_{E_G})$ on a μ -co- ε set. \square

Theorem 1.3 (Ergodic hyperfinite subgraph). *Every locally countable ergodic Borel graph G on a standard probability space (X, μ) admits an ergodic hyperfinite Borel subgraph $H \subseteq G$.*

Proof. We first note that without loss of generality, we may assume that μ is E_G -quasi-invariant by replacing it with $\sum_{m \geq 1} 2^{-m} (\gamma_m)_* \mu$, where $(\gamma_m)_{m \geq 1} \subseteq [E_G]$ is a sequence of Borel automorphisms such that $E = \bigcup_{m \geq 1} \text{Graph}(\gamma_m)$, which exists by the Feldman–Moore theorem [FM77]. Now let (F_n) be an increasing sequence of finite Borel equivalence relations provided by Theorem 1.1. Then $F := \bigcup_{n \in \mathbb{N}} F_n$ is hyperfinite and G -connected. Furthermore, for every $f \in L^1(X, \mu)$, $A_F^{\mathbb{W}} f = \int_X f d\mu$ a.e., so by Theorem 3.6(d), F is ergodic. \square

Lastly, we state and prove the general version of Theorem 1.5.

Theorem 4.1 (Ratio ergodic theorem for quasi-mp graphs). *Let G be a locally countable quasi-mp Borel graph on a σ -finite standard measure space (X, μ) and let $w : E_G \rightarrow \mathbb{R}^+$ be the Radon–Nikodym cocycle of E_G with respect to μ . There is an increasing sequence (F_n) of G -connected finite Borel equivalence relations F_n such that for any $f, g \in L^1(X, \mu)$ with $g > 0$,*

$$r := \lim_{n \rightarrow \infty} \frac{A_{F_n}^{\mathbb{W}} f}{A_{F_n}^{\mathbb{W}} g} = \mathbb{E}_{\mu_g}(f g^{-1} | \mathcal{B}_{E_G}) \text{ a.e.}$$

where \mathbb{E}_{μ_g} denotes the conditional expectation with respect to the measure μ_g defined by $d\mu_g := \frac{1}{\|g\|_1} g d\mu$. In other words, for any E_G -invariant Borel set $Y \subseteq X$,

$$\int_Y r g d\mu = \int_Y f d\mu.$$

In particular, when G is μ -ergodic,

$$\lim_{n \rightarrow \infty} \frac{A_{F_n}^w f}{A_{F_n}^w g} = \frac{\int_X f d\mu}{\int_X g d\mu} \quad a.e.$$

Proof. Observe that $\mu_g \sim \mu$ and the Radon–Nikodym cocycle of E_G with respect to μ_g is $\tilde{w}_x(y) = w_x(y)g(y)g(x)^{-1}$, so

$$\frac{A_{F_n}^w f(x)}{A_{F_n}^w g(x)} = \frac{\sum_{y \in [x]_{F_n}} f(y)w_x(y)}{\sum_{y \in [x]_{F_n}} g(y)w_x(y)} = \frac{\sum_{y \in [x]_{F_n}} f(y)g(y)^{-1}\tilde{w}_x(y)g(y)g(x)^{-1}}{\sum_{y \in [x]_{F_n}} \tilde{w}_x(y)g(y)g(x)^{-1}} = A_{F_n}^{\tilde{w}}(fg^{-1})(x).$$

It is clear that $fg^{-1} \in L^1(X, \mu_g)$, so Theorem 1.1 applied to G and fg^{-1} on (X, μ_g) gives the a.e. convergence of $\frac{A_{F_n}^w f}{A_{F_n}^w g}$ to $\mathbb{E}_{\mu_g}(fg^{-1}|\mathcal{B}_{E_G})$. This means that for any E_G -invariant Borel set $Y \subseteq X$,

$$\int_Y rg d\mu = \|g\|_1 \cdot \int_Y r d\mu_g = \|g\|_1 \cdot \int_Y fg^{-1} d\mu_g = \int_Y f d\mu.$$

Lastly, if G is μ -ergodic then it is also μ_g -ergodic because $\mu \sim \mu_g$, hence

$$\mathbb{E}_{\mu_g}(fg^{-1}|\mathcal{B}_{E_G}) = \int_X fg^{-1} d\mu_g = \frac{\int_X f d\mu}{\int_X g d\mu} \quad a.e. \quad \square$$

5. MASS TRANSPORT ALONG A COCYCLE

Throughout this section, let X be a standard Borel space, E a countable Borel equivalence relation on X , and $w : E \rightarrow \mathbb{R}^+$ a Borel cocycle.

5.A. Preliminaries. We refer to a function $\varphi : E \rightarrow [0, \infty)$ as a *mass transport* or just *transport* (in E), and we think of the value $\varphi(x, y)$ as the fraction of the w -mass of x that gets transported from x to y ; thus, what y receives in this transaction is $\varphi(x, y)w_y(x)$. With this in mind, we define functions $\text{in}^w \varphi, \text{out}^w \varphi : X \rightarrow [0, \infty]$ by

$$\begin{aligned} \text{out}^w \varphi(x) &:= \sum_{y \in [x]_E} \varphi(x, y) \\ \text{in}^w \varphi(x) &:= \sum_{y \in [x]_E} \varphi(y, x)w_x(y). \end{aligned}$$

Definition 5.1. We say that a transport $\varphi : E \rightarrow [0, \infty)$ is *w-bounded* if $\text{out}^w \varphi$ and $\text{in}^w \varphi$ are bounded by 1.

The definition of a w -bounded transport in E is exactly the same as that of a *w-invariant fuzzy partial injection* defined in [Mil08], and in our arguments we will only use such transports.

When defining a transport, we will only partially specify its values with the convention that the undefined values are treated as 0. We also use the following terminology:

- The *domain* of φ is $\text{dom}(\varphi) := \{(x, y) \in E : \varphi(x, y) \neq 0\}$.
- The *vertex-domain* of φ is $\text{vdom}(\varphi) := \text{proj}_1(\text{dom}(\varphi)) \cup \text{proj}_2(\text{dom}(\varphi))$.
- For sets $Y, Z \subseteq X$, we say $Y \times Z$ is *φ -closed* if $\text{dom}(\varphi)$ is disjoint from $(Y \times Z^c) \cup (Y^c \times Z)$.

5.B. Mass transport principle.

Lemma 5.2. *For a transport φ in E , and for any subsets U, V of the same E -class in X such that $U \times V$ is φ -closed, we have*

$$\int_U \text{out}^{\mathfrak{w}} \varphi d\mathfrak{w} = \int_V \text{in}^{\mathfrak{w}} \varphi(v) d\mathfrak{w}.$$

Proof. Letting x be any element in $[U \cup V]_E$, we compute:

$$\begin{aligned} \sum_{u \in U} \text{out}^{\mathfrak{w}} \varphi(u) \cdot \mathfrak{w}_x(u) &= \sum_{u \in U} \sum_{v \in V} \varphi(u, v) \cdot \mathfrak{w}_x(u) \\ &\quad \left[\text{Fubini} \right] = \sum_{v \in V} \sum_{u \in U} \varphi(u, v) \cdot \frac{\mathfrak{w}_x(u)}{\mathfrak{w}_x(v)} \cdot \mathfrak{w}_x(v) \\ &\quad \left[\text{cocycle identity} \right] = \sum_{v \in V} \sum_{u \in U} \varphi(u, v) \cdot \mathfrak{w}_v(u) \cdot \mathfrak{w}_x(v) \\ &= \sum_{v \in V} \text{in}^{\mathfrak{w}} \varphi(v) \cdot \mathfrak{w}_x(v). \end{aligned} \quad \square$$

The following is the measure version of Lemma 5.2, which is just [Mil08, Proposition 6.4] or [KM04, Proposition 8.2] phrased in our terminology.

Proposition 5.3. *For a Borel transport φ in E and any \mathfrak{w} -invariant Borel probability measure μ on X ,*

$$\int_X \text{out}^{\mathfrak{w}} \varphi d\mu = \int_X \text{in}^{\mathfrak{w}} \varphi d\mu.$$

Proof. We assume without loss of generality that $\varphi(x, x) = 0$ for all $x \in X$ because changing the value of $\varphi(x, x)$ does not change the validity of the desired equality.

Claim 5.4. $E \setminus \text{Id}_X$ is the disjoint union of the sets $\text{Graph}(\gamma_n) \setminus \text{Id}_X$, $n \in \mathbb{N}$, where each γ_n is a Borel involution on X .

Proof of Claim. By the Feldman–Moore theorem [FM77], E is a union of $\text{Graph}(\beta_n)$, $n \in \mathbb{N}$, where each β_n is a Borel involution on X . Recalling that set-theoretically, β_n is the same as $\text{Graph}(\beta_n)$, we define partial involutions $\beta'_n := \beta_n \setminus \bigcup_{k < n} \beta_k$ whose graphs are pairwise disjoint. Let $\gamma_n : X \rightarrow X$ be equal to β'_n on the domain of β'_n and to the identity function, elsewhere. \(\square\)

Putting $X_n := \{x \in X : \gamma_n x \neq x\}$, we compute:

$$\begin{aligned}
\int_X \text{out}^w \varphi(x) d\mu(x) &= \int_X \sum_{n \in \mathbb{N} : \gamma_n x \neq x} \varphi(x, \gamma_n x) d\mu(x) \\
\left[\text{Fubini} \right] &= \sum_n \int_{X_n} \varphi(x, \gamma_n x) d\mu(x) \\
\left[\text{change of variable } x \text{ to } \gamma_n x \right] &= \sum_n \int_{X_n} \varphi(\gamma_n x, x) \cdot w_x(\gamma_n x) d\mu(x) \\
\left[\text{Fubini} \right] &= \int_X \sum_{n \in \mathbb{N} : \gamma_n x \neq x} \varphi(\gamma_n x, x) \cdot w_x(\gamma_n x) d\mu(x) \\
&= \int_X \text{in}^w \varphi(x) d\mu(x). \quad \square
\end{aligned}$$

5.C. Deficiency and w -invariant measures. For a w -bounded transport φ in E ,

- call a point $x \in X$ a φ -source (resp. φ -sink) if $\text{out}^w \varphi(x) - \text{in}^w \varphi(x)$ is positive (resp. negative). We denote the sets of φ -sources and φ -sinks by $\text{Sources}(\varphi)$ and $\text{Sinks}(\varphi)$, respectively.
- say that φ *disbalances* an E -class C if C contains at least one φ -source but no φ -sink, or vice versa, at least one φ -sink but no φ -source.
- say that φ *disbalances* an E -invariant set $Z \subseteq X$ if it disbalances every E -class of Z .
- call $Y \subseteq X$ *w -deficient* if there is a w -bounded Borel transport φ in E disbalancing $[Y]_E$.

Taking sums of transports on disjoint E -invariant domains, we see that:

Observation 5.5. *w -deficient sets form a σ -ideal.*

Thus, we say that a statement holds *modulo w -deficient* if it holds on $X \setminus D$ for some E -invariant w -deficient Borel set D .

Proposition 5.3 immediately gives the following corollary.

Corollary 5.6. *If a Borel $Y \subseteq X$ is w -deficient, then $\mu(Y) = 0$ for every w -invariant Borel probability measure μ on X .*

This corollary is all we need about mass transports in our proofs. However, it is well worth pointing out that its converse is also true (much more difficult to prove) and it is the content of [Mil08, Theorem 3]. We restate this here in our terms for the sake of completeness:

Theorem 5.7 (Miller 2008). *For a countable Borel equivalence relation E on a standard Borel space X and a Borel cocycle $w : E \rightarrow \mathbb{R}^+$,*

- (I) *either: X is w -deficient,*
- (II) *or else: there is a w -invariant Borel probability measure on X .*

5.D. Transporting a prescribed fraction of mass.

Lemma 5.8. *For any disjoint nonempty w -finite subsets U, V of the same E -class in X and a function $\alpha : U \rightarrow [0, 1]$, if $\int_U \alpha dw \leq w(V)$, then there is a w -bounded transport φ in E with $\text{dom}(\varphi) \subseteq U \times V$ such that $\text{out}^w \varphi|_U = \alpha$.*

Proof. For each $(u, v) \in U \times V$, define $\varphi(u, v) := \alpha(u) \frac{\mathfrak{w}(v)}{\mathfrak{w}(V)}$. Then for each $u \in U$,

$$\text{out}^{\mathfrak{w}}\varphi(u) = \sum_{v \in V} \alpha(u) \frac{\mathfrak{w}(v)}{\mathfrak{w}(V)} = \alpha(u),$$

while for each $v \in V$,

$$\text{in}^{\mathfrak{w}}\varphi(v) = \sum_{u \in U} \alpha(u) \frac{\mathfrak{w}_v(v)}{\mathfrak{w}_v(V)} \mathfrak{w}_v(u) = \sum_{u \in U} \alpha(u) \frac{\mathfrak{w}_v(u)}{\mathfrak{w}_v(V)} = \frac{1}{\mathfrak{w}(V)} \int_U \alpha \, d\mathfrak{w} \leq 1. \quad \square$$

The proof of the last lemma can be carried out in a uniformly Borel fashion, yielding:

Lemma 5.9. *Let $F \subseteq E$ be a \mathfrak{w} -finite Borel subequivalence relation, $U, V \subseteq X$ be disjoint Borel sets, and $\alpha : U \rightarrow [0, 1]$ a Borel function. If for each F -class Y , $\int_{U \cap Y} \alpha \, d\mathfrak{w} \leq \mathfrak{w}(V \cap Y)$, then there exists a \mathfrak{w} -bounded Borel transport φ with $\text{dom}(\varphi) \subseteq (U \times V) \cap F$ and $(\text{out}^{\mathfrak{w}}\varphi)|_U = \alpha$.*

6. HYPERFINITENESS VIA CUTS

In this section we present an easy but very useful characterization of μ -hyperfiniteness that turns non- μ -hyperfiniteness into a positive property. Let X be a standard Borel space and let G be a locally countable Borel graph on it.

Definition 6.1. Call a set $V \subseteq X$ a *finitizing* (resp. *hyperfinitizing*) *vertex-cut* for G if G_{-V} is component-finite (resp. hyperfinite). Likewise, call a subgraph $H \subseteq G$ a *finitizing* (resp. *hyperfinitizing*) *edge-cut* for G if $G \setminus H$ is component-finite (resp. hyperfinite).

Call a sequence of sets *vanishing* if it is decreasing and has empty intersection.

Proposition 6.2. *A locally countable Borel graph G is hyperfinite if and only if it admits a vanishing sequence $(H_n)_n$ of finitizing Borel edge-cuts.*

Proof. \Rightarrow : Letting (G_n) be an increasing sequence of component-finite Borel graphs with $G = \bigcup_n G_n$, we see that the graphs $G \setminus G_n$ are vanishing finitizing edge-cuts for G .

\Leftarrow : For each $n \geq 1$, $G_n := G \setminus H_n$ is component-finite and $G = \bigcup_{n=1}^{\infty} G_n$. \square

Remark 6.3. The analogue for vertex-cuts of Proposition 6.2 also holds for locally finite graphs. For locally countable graphs, the implication \Leftarrow still holds, but \Rightarrow may fail: take any aperiodic hyperfinite equivalence relation E and let $G := E \setminus \text{Id}_X$.

Let μ and ν be finite Borel measures on X and G , respectively.

Definition 6.4. The *finitizing vertex-price* (with respect to μ) and *finitizing edge-price* (with respect to ν) of G are the quantities:

$$\text{fvp}_{\mu}(G) := \inf \{ \mu(V) : V \subseteq X \text{ is a Borel finitizing vertex-cut for } G \},$$

$$\text{fep}_{\nu}(G) := \inf \{ \nu(H) : H \subseteq G \text{ is a Borel finitizing edge-cut for } G \}.$$

Replacing “finitizing” with “hyperfinitizing” in the above definitions, we obtain *hyperfinitizing vertex* and *edge prices* denoted by $\text{hvp}_{\mu}(G)$ and $\text{hep}_{\nu}(G)$.

We characterize μ -hyperfiniteness in terms of $\text{hvp}_{\mu}(G)$, but the proof of this goes through finitizing edge-cuts and $\text{hep}_{\hat{\mu}}(G)$ for a measure $\hat{\mu}$ that is a *lift* of μ , that is: for any Borel (symmetric) subgraph $H \subseteq G$, H is $\hat{\mu}$ -null if and only if $\text{proj}_1(H)$ is μ -null; in particular,

$(\text{proj}_1)_*\hat{\mu} \sim \mu$. For example, writing G as a countable union of the graphs of Borel maps $\gamma_n : X \rightarrow X$ (by Luzin–Novikov uniformization), we define, for a Borel set $H \subseteq G$,

$$\hat{\mu}(H) := \sum_{n \geq 1} 2^{-n} \int_X \mathbb{1}_H(x, \gamma_n x) d\mu(x).$$

Proposition 6.5. *For a locally countable quasi-pmp Borel graph G on a standard probability space (X, μ) and a lift $\hat{\mu}$ of μ to a **finite** Borel measure on G , the following are equivalent:*

- (I) G is μ -hyperfinite.
- (II) $\text{hvp}_\mu(G) = 0$.
- (III) $\text{hep}_{\hat{\mu}}(G) = 0$.
- (IV) $\text{fep}_{\hat{\mu}}(G) = 0$.

These statements are implied by the following, and are equivalent to it when G is locally finite:

- (V) $\text{fvp}_\mu(G) = 0$.

Proof. (I) \Rightarrow (II) is trivial. For (II) \Rightarrow (III), if $V \subseteq X$ is an arbitrarily μ -small hyperfinitizing vertex-cut then $[V, X]_G$ is an arbitrarily $\hat{\mu}$ -small hyperfinitizing edge-cut, by $(\text{proj}_1)_*\hat{\mu} \ll \mu$. (III) \Rightarrow (IV) follows by an $\frac{\varepsilon}{2} + \frac{\varepsilon}{2}$ argument using that hyperfinite graphs admit $\hat{\mu}$ -arbitrarily small finitizing edge-cuts, by Proposition 6.2 and the finiteness of $\hat{\mu}$.

For (IV) \Rightarrow (I), suppose $\text{fep}_{\hat{\mu}}(G) = 0$, and let (H_n) be a sequence of finitizing Borel edge-cuts for G such that $\mu(H_n)$ is summable. Then the graphs $H'_n := \bigcup_{k \geq n} H_k$ are decreasing edge-cuts for G and $H = \bigcap_n H'_n$ is $\hat{\mu}$ -null. Hence, $Z := \text{proj}_1(H)$ is μ -null because $\hat{\mu}$ is a lift of μ , and by the quasi-invariance of μ , $[Z]_{E_G}$ is still μ -null. Throwing it out, makes (H'_n) a vanishing sequence of finitizing edge-cuts, so G becomes hyperfinite by Proposition 6.2. The same argument applied to vertices shows (V) \Rightarrow (I).

It remains to show (I) \Rightarrow (V), assuming G is locally finite. To this end, removing a μ -null set from X , we may write G as an increasing union of component-finite Borel graphs G_n . Then the sets $V_n := \{x \in X : x \in \partial_G[x]_{G_n}\}$ form a vanishing sequence of finitizing vertex-cuts because $\bar{\partial}_G[x]_{G_n}$ is finite for every $x \in X$. Thus, $\mu(V_n) \rightarrow 0$ by the finiteness of μ . \square

Remark 6.6. Although Proposition 6.5 has a simple proof, it immediately implies the Dye–Krieger theorem [KM04, Theorem 6.11] that the increasing union $E := \biguplus_n E_n$ of μ -hyperfinite equivalence relations is μ -hyperfinite. Indeed, if H_n is a finitizing edge-cut for E_n , then $H := \bigcup_n H_n$ a hyperfinitizing edge-cut for E because $E \setminus H = \biguplus_n (E_n \setminus H)$ and each $E_n \setminus H$ is a component-finite graph. Taking a lift $\hat{\mu}$ to E of the measure μ , we can make $\hat{\mu}(H)$ arbitrarily small by taking the H_n so that $\sum_n \hat{\mu}(H_n)$ is sufficiently small, hence $\text{hep}_{\hat{\mu}}(E) = 0$.

7. PACKED AND APPROXIMATELY SATURATED TILINGS

Throughout this section, let X be a standard Borel space, E be a countable Borel equivalence relation on X , and $w : E \rightarrow \mathbb{R}^+$ a Borel cocycle. Within a given Borel collection \mathcal{S} of w -finite subsets of X , we will build Borel tilings of various degrees of maximality.

When $w \equiv 1$, i.e. in the pmp setting, all of the results below are simpler and are proven in [MT17, Section 4].

7.A. Tilings. For any $\mathcal{S} \subseteq \mathcal{P}(X)$, we put $\text{dom}(\mathcal{S}) := \bigcup \mathcal{S}$ and call it the *domain* of \mathcal{S} . For $Y \subseteq X$, put

$$\mathcal{S}|_Y := \{U \in \mathcal{S} : U \subseteq Y\}.$$

We call \mathcal{S} a *tiling* if the sets in \mathcal{S} are pairwise disjoint. In this case, we refer to the sets in \mathcal{S} as *tiles* and denote by $E(\mathcal{S})$ the equivalence relation that is the identity outside of $\text{dom}(\mathcal{S})$ and on $\text{dom}(\mathcal{S})$ its classes are exactly the tiles in \mathcal{S} .

We say that a tiling $\mathcal{P} \subseteq \mathcal{P}(X)$ is *maximal* within $\mathcal{S} \subseteq \mathcal{P}(X)$, if no set in \mathcal{S} is disjoint from $\text{dom}(\mathcal{P})$.

Definition 7.1. Let $\mathcal{S} \subseteq [X]_E^{\mathfrak{w} < \infty}$.

- Let $\overline{\mathcal{S}} \subseteq [X]_E^{\mathfrak{w} < \infty}$ denote the *upward closure* of \mathcal{S} , i.e. the collection of all \mathfrak{w} -finite sets that are countable increasing unions of sets in \mathcal{S} . Call \mathcal{S} *upward closed* if $\overline{\mathcal{S}} = \mathcal{S}$.
- Say that \mathcal{S} is *finitely based* if for any $A \in \mathcal{S}$ and any finite $B \subseteq A$ there is a finite $A' \in \mathcal{S}$ with $B \subseteq A' \subseteq A$. This immediately implies a stronger statement that for any $\varepsilon > 0$, A' can be taken such that $\mathfrak{w}(A') \geq (1 - \varepsilon)\mathfrak{w}(A)$.
- We say that \mathcal{S} has *finitely based quotients* if for each Borel tiling $\mathcal{P} \subseteq \mathcal{S}$, the quotient $\mathcal{S}_{/E(\mathcal{P})}$ is finitely based.

When $\mathcal{S} \subseteq [X]_E^{\mathfrak{w} < \infty}$ is Borel, $\text{dom}(\mathcal{S})$ is analytic (hence measurable) in general. However:

Lemma 7.2. *If a Borel collection $\mathcal{S} \subseteq [X]_E^{\mathfrak{w} < \infty}$ is either finitely based or a tiling, then $\text{dom}(\mathcal{S})$ is Borel.*

Proof. In the first case, $\text{dom}(\mathcal{S}) = \text{dom}(\mathcal{S} \cap [X]_E^{\leq \infty})$ and every $x \in \text{dom}(\mathcal{S} \cap [X]_E^{\leq \infty})$ is contained in only countably-many sets in $\mathcal{S} \cap [X]_E^{\leq \infty}$. In the second case, every $x \in \text{dom}(\mathcal{S})$ is contained in exactly one set in \mathcal{S} . Thus, in either case, Observation 2.2 and Luzin–Novikov uniformization imply that $\text{dom}(\mathcal{S})$ is Borel. \square

Let $\mathcal{P}, \mathcal{P}' \subseteq \mathcal{P}(X)$ be tilings. We say that \mathcal{P}' is a *partial extension* of \mathcal{P} , noted $\mathcal{P}' \succeq \mathcal{P}$, if each set in \mathcal{P}' is $E(\mathcal{P})$ -invariant. If moreover, $\text{dom}(\mathcal{P}') \supseteq \text{dom}(\mathcal{P})$, we say that \mathcal{P}' is an *extension* of \mathcal{P} and write $\mathcal{P}' \geq \mathcal{P}$.

For $N \leq \infty$, we call a sequence $(\mathcal{P}_n)_{n < N}$ of tilings *coherent* if $\mathcal{P}_i \leq \mathcal{P}_j$ for all $i \leq j$. In this case, observe that although the $E(\mathcal{P}_n)$ are not increasing, $\text{Id}_X \cup \bigcup_{n < m} E(\mathcal{P}_n)$ is an equivalence relation for each $m \leq N$. It thus makes sense to define

$$\bigvee_{n < N} \mathcal{P}_n$$

as the collection of all classes of the equivalence relation $\text{Id}_X \cup \bigcup_{n < N} E(\mathcal{P}_n)$ that are contained in $\bigcup_{n < N} \text{dom}(\mathcal{P}_n)$; in particular, $\bigvee_{n < 0} \mathcal{P}_n = \emptyset$. When $N = \infty$, we say that the sequence $(\mathcal{P}_n)_{n < \infty}$ *stabilizes* if $\bigvee_{n < \infty} \mathcal{P}_n \subseteq \bigcup_{n < \infty} \mathcal{P}_n$.

7.B. Packed tilings.

Definition 7.3. Let $p \in \mathbb{R}^+$.

- For a tiling $\mathcal{P} \subseteq [X]_E^{\mathfrak{w} < \infty}$, call a set $A \in [X]_E^{\mathfrak{w} < \infty}$ a *p-pack* over \mathcal{P} if A is $E(\mathcal{P})$ -invariant and $\mathfrak{w}(A \setminus \text{dom}(\mathcal{P})) \geq p \cdot \mathfrak{w}(\text{dom}(\mathcal{P}) \cap A)$, equivalently, $\mathfrak{w}(A) \geq (1 + p) \cdot \mathfrak{w}(\text{dom}(\mathcal{P}) \cap A)$.
- For a collection $\mathcal{S} \subseteq [X]_E^{\mathfrak{w} < \infty}$, call a tiling $\mathcal{P} \subseteq [X]_E^{\mathfrak{w} < \infty}$ *p-packed* within \mathcal{S} if \mathcal{S} has no *p-pack* over \mathcal{P} .

- A sequence $(\mathcal{P}_n)_{n \in \mathbb{N}}$ of tilings contained in $[X]_E^{\mathbb{w} < \infty}$ is *p-packing* if it is coherent and for all $n \in \mathbb{N}$, each tile in $\mathcal{P}_n \setminus (\bigvee_{m < n} \mathcal{P}_m)$ is a *p-pack* over $\bigvee_{m < n} \mathcal{P}_m$.

Observation 7.4. Let $p \in \mathbb{R}^+$, $\mathcal{S} \subseteq [X]_E^{\mathbb{w} < \infty}$, and let $\mathcal{P} \subseteq [X]_E^{\mathbb{w} < \infty}$ be a *p-packed* tiling within \mathcal{S} .

- (a) \mathcal{P} is maximal within \mathcal{S} .
- (b) Any tiling $\mathcal{P}' \subseteq [X]_E^{\mathbb{w} < \infty}$ with $\mathcal{P}' \geq \mathcal{P}$ is still *p-packed* within \mathcal{S} .

Lemma 7.5. For any $p \in \mathbb{R}^+$, any *p-packing* sequence (\mathcal{P}_n) of Borel tilings within $[X]_E^{\mathbb{w} < \infty}$ stabilizes modulo \mathbb{w} -deficient.

Proof. We assume without loss of generality that $p \leq 1$. We also assume without loss of generality that (\mathcal{P}_n) is extension-increasing by replacing each \mathcal{P}_n with $\bigvee_{m \leq n} \mathcal{P}_m$.

Put $E_n := E(\mathcal{P}_n)$ for each $n \in \mathbb{N}$, and $E_\infty := \bigvee_n E_n$. Letting Z be the union of all E_∞ -classes C with $C \notin \bigcup_n \mathcal{P}_n$, it is enough to build a \mathbb{w} -bounded Borel transport in $E_\infty|_Z$ disbalancing Z . Thus, we assume without of generality that $X = Z$, so for every $n \in \mathbb{N}$, each E_n -class is strictly contained in an E_∞ -class.

For each $n \in \mathbb{N}$, putting $D_n := \text{dom}(\mathcal{P}_n)$ and $D_{-1} := \emptyset$, let $\mathcal{P}'_n := \{P \in \mathcal{P}_n : P \cap D_{n-1} = \emptyset\}$. The domains $D'_n := \text{dom}(\mathcal{P}'_n)$ are pairwise disjoint and the set $D' := \bigcup_n D'_n$ is an E_∞ -complete section. We build a \mathbb{w} -bounded transport φ in E_∞ with no sinks but with $\text{Sources}(\varphi) = D'$, thus disbalancing X .

To get an intuitive idea, take $U \in \bigcup_n \mathcal{P}'_n$ and let $n \in \mathbb{N}$ be the least such that $V := [U]_{E_n} \supsetneq U$. The packing condition ensures that $V \setminus D_{n-1}$ has at least as much mass as the p^{th} fraction of the mass of $V \cap D_{n-1}$, so we can transport the p^{th} fraction of the mass of U into $V \setminus D_{n-1}$. Maybe other E_{n-1} -classes in $V \cap D_{n-1}$ decide to do the same, but that is still alright since $V \setminus D_{n-1}$ has enough mass to accommodate the p^{th} fraction of the mass of all of $V \cap D_{n-1}$. We repeat this with V instead of U , to drain the sinks in V into new points that join the equivalence class of V in a later stage. Thus, sinks move to infinity, while U stays a source.

Formally, we recursively define a sequence (φ_n) of \mathbb{w} -bounded Borel transports with pairwise disjoint domains, and take $\varphi := \sum_n \varphi_n$. For each $n \geq 1$, putting $\bar{\varphi}_n := \sum_{k < n} \varphi_k$, we ensure that

- (i) $\text{dom}(\bar{\varphi}_n) \subseteq E_n \cap (D_{n-1} \times (D_n \setminus D'))$;
- (ii) $\text{Sources}(\bar{\varphi}_n) = \bigcup_{k < n} D'_k$ and the net of $\bar{\varphi}_n$ on $\text{Sources}(\bar{\varphi}_n)$ is constant $-p$;
- (iii) $\text{Sinks}(\bar{\varphi}_n)$ is disjoint from $\bigcup_{k < n} U_k$, where $U_k := X_k \cap D_k$ and

$$X_k := \{x \in D_{k+1} : [x]_{E_{k+1}} \cap D_k \neq \emptyset \text{ and } [x]_{E_{k+1}} \setminus D_k \neq \emptyset\}.$$

Condition (ii) guarantees that $\text{Sources}(\varphi) = D'$, and (iii) implies that $\text{Sinks}(\varphi) = \emptyset$ because $X = \bigcup_n U_n$ (each E_n -class is strictly contained in an E_m -class for some $m > n$). It remains to construct such a sequence (φ_n) .

Fixing $n \in \mathbb{N}$, suppose that $\bar{\varphi}_n$ is defined and satisfies (i)–(iii). Putting $V_n := X_n \setminus D_n$, so $X_n = U_n \sqcup V_n$, we will define φ_n only on $E_{n+1} \cap (U_n \times V_n)$, so $\text{dom}(\varphi_n)$ is disjoint from $\text{dom}(\bar{\varphi}_n)$ by (i) for $\bar{\varphi}_n$. Moreover, because $V_n \cap D' = \emptyset$, condition (i) will also hold for $\bar{\varphi}_{n+1}$.

By (i) for $\bar{\varphi}_n$, D'_n is disjoint from $\text{Sinks}(\bar{\varphi}_n)$, so we define $\alpha_n : U_n \rightarrow [0, 1]$ by

$$\alpha_n(u) := \begin{cases} p & \text{if } u \in D'_n \\ \text{in}^{\mathbb{w}} \bar{\varphi}_n(u) - \text{out}^{\mathbb{w}} \bar{\varphi}_n(u) & \text{if } u \in \text{Sinks}(\bar{\varphi}_n) \\ 0 & \text{otherwise.} \end{cases}$$

Claim. $\int_{U_n \cap Y} \alpha_n d\mathbf{w} \leq \mathbf{w}(V_n \cap Y)$ for each E_{n+1} -class $Y \subseteq X_n$.

Proof of Claim. Fix an E_n -class $P \subseteq U_n$. If $P \cap D'_n \neq \emptyset$, then $P \subseteq D'_n$ because D'_n is E_n -invariant, and $\int_P \alpha_n d\mathbf{w} = p \cdot \mathbf{w}(P)$. If $P \cap D'_n = \emptyset$, then

$$\begin{aligned} \int_P \alpha_n d\mathbf{w} &= \int_{P \cap \text{Sinks}(\overline{\varphi}_n)} (\text{in}^{\mathbf{w}} \overline{\varphi}_n - \text{out}^{\mathbf{w}} \overline{\varphi}_n) d\mathbf{w} \\ \left[\text{Lemma 5.2} \right] &= - \int_{P \cap \text{Sources}(\overline{\varphi}_n)} (\text{in}^{\mathbf{w}} \overline{\varphi}_n - \text{out}^{\mathbf{w}} \overline{\varphi}_n) d\mathbf{w} \\ \left[\text{(ii) for } \overline{\varphi}_n \right] &= p \cdot \mathbf{w}(P \cap \text{Sources}(\overline{\varphi}_n)) \leq p \cdot \mathbf{w}(P). \end{aligned}$$

Thus, for each E_{n+1} -class $Y \subseteq X_n$, $\int_{U_n \cap Y} \alpha_n d\mathbf{w} \leq p \cdot \mathbf{w}(U_n \cap Y) \leq \mathbf{w}(V_n \cap Y)$, where the last inequality is because Y is a p -pack over \mathcal{P}_n . \square

Hence, Lemma 5.9 applies to $F := E_{n+1}|_{X_n}$, $U := U_n$, $V := V_n$, and $\alpha := \alpha_n$, yielding a transport φ_n with $\text{dom}(\varphi_n) \subseteq E_{n+1} \cap (U_n \times V_n)$ and $\text{out}^{\mathbf{w}} \varphi_n|_{U_n} \equiv \alpha_n$. In particular, $\text{Sources}(\overline{\varphi}_{n+1}) = D'_n \cup \text{Sources}(\overline{\varphi}_n)$, so (ii) is satisfied for $\overline{\varphi}_{n+1}$. Furthermore, $\text{Sinks}(\overline{\varphi}_{n+1}) \cap X_n = \text{Sinks}(\varphi_n) \subseteq V_n$, so (iii) holds too. \square

Theorem 7.6. *For any countable Borel equivalence relation E on X , $p \in \mathbb{R}^+$, and a Borel $\mathcal{S} \subseteq [X]_E^{\leq \infty}$, there is a Borel tiling $\mathcal{P} \subseteq \mathcal{S}$ that is p -packed within \mathcal{S} modulo \mathbf{w} -deficient.*

Proof. By [KM04, Proof of Lemma 7.3], the intersection graph on $[X]_E^{\leq \infty}$ admits a countable Borel coloring and we fix one so that the colors are natural numbers.

We recursively build an extension-increasing p -packing sequence (\mathcal{P}_n) of Borel tilings contained in \mathcal{S} . Take $\mathcal{P}_0 := \emptyset$ and fixing $n \in \mathbb{N}$, suppose that \mathcal{P}_n is defined. Let \mathcal{P}'_n be the collection of all sets in \mathcal{S} of color n that are p -packs over \mathcal{P}_n and let $\mathcal{P}_{n+1} := \mathcal{P}'_n \cup (\mathcal{P}_n|_{X \setminus \text{dom}(\mathcal{P}'_n)})$.

By Lemma 7.5, we may assume that the sequence (\mathcal{P}_n) stabilizes, so $\mathcal{P} := \bigvee_n \mathcal{P}_n \subseteq \mathcal{S}$ and it remains to show that \mathcal{P} is p -packed within \mathcal{S} . Suppose towards a contradiction that $U \in \mathcal{S}$ is a p -pack over \mathcal{P} . Then U is a p -pack over \mathcal{P}_n for every $n \in \mathbb{N}$. Letting n be the color of U , the construction puts U into \mathcal{P}_{n+1} , a contradiction. \square

Lemma 7.7. *Let $\mathcal{S} \subseteq [X]_E^{\mathbf{w} < \infty}$ be a Borel collection with finitely based quotients and let $p \in \mathbb{R}^+$. For any Borel tiling $\mathcal{P} \subseteq \mathcal{S}' := \mathcal{S} \cap [X]_E^{\leq \infty}$, if \mathcal{P} is p -packed within \mathcal{S}' , then \mathcal{P} is p' -packed within \mathcal{S} for any $p' > p$.*

Proof. Let $p' > p$ and suppose towards a contradiction that there is a p' -pack $A \in \mathcal{S}$ over \mathcal{P} . Taking $\varepsilon \in (0, 1)$ such that $(1 - \varepsilon)(p' + 1) \geq (p + 1)$, the finite basedness of $\mathcal{S}_{/E(\mathcal{P})}$ yields an $E(\mathcal{P})$ -invariant set $A' \in \mathcal{S}'$ such that $A' \subseteq A$ and $\mathbf{w}(A') \geq (1 - \varepsilon)\mathbf{w}(A)$. But then

$$\mathbf{w}(A') \geq (1 - \varepsilon) \cdot \mathbf{w}(A) \geq (1 - \varepsilon)(p' + 1) \cdot \mathbf{w}(A \cap \text{dom}(\mathcal{P})) \geq (p + 1) \cdot \mathbf{w}(A' \cap \text{dom}(\mathcal{P})),$$

so A' is a p -pack over \mathcal{P} , contradicting the p -packedness of \mathcal{P} within \mathcal{S}' . \square

7.C. Approximately saturated tilings.

Definition 7.8. • For a tiling $\mathcal{P} \subseteq [X]_E^{\mathbf{w} < \infty}$, call a set $A \in [X]_E^{\mathbf{w} < \infty}$ *injective* over \mathcal{P} if A is $E(\mathcal{P})$ -invariant and contains at most one tile from \mathcal{P} .

- A sequence (\mathcal{P}_n) of tilings is *injective* if it is extension-increasing and for all $n \in \mathbb{N}$, each tile in \mathcal{P}_{n+1} is injective over \mathcal{P}_n .

For a collection $\mathcal{S} \subseteq [X]_E^{\leq \infty}$ and a tiling $\mathcal{P} \subseteq \mathcal{S}$, a notion of saturation was defined in [MT17, Section 4.C] as follows: \mathcal{P} is *saturated* within \mathcal{S} if there is no set in \mathcal{S} that is injective over \mathcal{P} . It was also shown in [MT17, Section 4.C] that if \mathcal{S} is Borel, then a saturated Borel tiling exists off of an E -invariant compressible Borel set. We would like to have the same statement, but with compressible replaced with w -deficient. However, this is not true in general: let $X := \mathbb{N}$, $E := \mathbb{N}^2$, $w_n(m) := 2^{n-m}$, and $\mathcal{S} := \{U \in [X]_E^{\leq \infty} : 0 \in U\}$. Thus, we define an approximate notion of saturation for a given cocycle w , which coincides with the original notion when $w \equiv 1$, and prove existence modulo w -deficient.

To motivate this new definition, we rephrase the original definition: \mathcal{P} is saturated within \mathcal{S} if and only if $\mathcal{P} \subseteq \mathcal{S}$ and there is no $Q \in [X]_E^{\leq \infty}$ disjoint from $\text{dom}(\mathcal{P})$ such that either $Q \in \mathcal{S}$ or there is $x \in \text{dom}(\mathcal{P})$ such that $[x]_{E(\mathcal{P})} \cup Q \in \mathcal{S}$.

Definition 7.9. For a collection $\mathcal{S} \subseteq [X]_E^{w < \infty}$, call a tiling $\mathcal{P} \subseteq [X]_E^{w < \infty}$ *approximately saturated* within \mathcal{S} if $\mathcal{P} = \bigvee_n \mathcal{P}_n$ for some injective sequence $(\mathcal{P}_n)_{n \in \mathbb{N}}$ of tilings contained in \mathcal{S} with the following property: there is no $Q \in [X]_E^{w < \infty}$ disjoint from $\text{dom}(\mathcal{P})$ such that either $Q \in \mathcal{S}$ or there is $x \in \text{dom}(\mathcal{P})$ such that $[x]_{E(\mathcal{P}_n)} \cup Q \in \mathcal{S}$ for all large enough $n \in \mathbb{N}$. We refer to (\mathcal{P}_n) as a *saturating \mathcal{S} -approximation* for \mathcal{P} .

When $w \equiv 1$, approximately saturated is indeed the same as saturated because any saturating \mathcal{S} -approximation stabilizes.

Observation 7.10. Let $\mathcal{P} \subseteq [X]_E^{w < \infty}$ be a tiling approximately saturated within $\mathcal{S} \subseteq [X]_E^{w < \infty}$.

- (a) \mathcal{P} is maximal within \mathcal{S} .
- (b) $\mathcal{P} \subseteq \overline{\mathcal{S}}$.

To build approximately saturated tilings, we first need the following:

Lemma 7.11. For an injective sequence (\mathcal{P}_n) of Borel tilings $\mathcal{P}_n \subseteq [X]_E^{w < \infty}$, the w -infinite part of $E_\infty := \bigvee_n E(\mathcal{P}_n)$ is w -deficient.

Proof. Letting Z be the union of w -infinite E_∞ -classes, it is enough to build a w -bounded Borel transport in $E_\infty|_Z$ disbalancing Z . Thus, we assume without of generality that $X = Z$, so E_∞ is w -infinite. Let $E_n := E(\mathcal{P}_n)$ for each $n \in \mathbb{N}$.

Let D be the union of inclusion-minimal sets in $\bigcup_n \mathcal{P}_n$; in particular, $[D]_{E_\infty} = X$. We define $k_m : D \rightarrow \mathbb{N}$ by induction on m as follows. For each $x \in D$, let $k_0(x)$ be the least $k \in \mathbb{N}$ with $[x]_{E_k} \in \mathcal{P}_k$ and $[x]_{E_k} \subseteq D$. Supposing that $k_m(x)$ is defined, let $k_{m+1}(x)$ be the least $k > k_m(x)$ such that $w([x]_{E_k} \setminus [x]_{E_{k_m(x)}}) \geq w([x]_{E_{k_m(x)}})$. For each $m \geq 0$, $\mathcal{Q}_m := \{[x]_{E_{k_m(x)}} : x \in D\}$ is a Borel tiling, and (\mathcal{Q}_m) is an injective sequence. Moreover, for each $Q \in \mathcal{Q}_{m+1}$, $Q = [x]_{E_{k_{m+1}(x)}}$ for some $x \in D$ and

$$w(Q \setminus \text{dom}(\mathcal{Q}_m)) = w([x]_{E_{k_{m+1}(x)}} \setminus [x]_{E_{k_m(x)}}) \geq w([x]_{E_{k_m(x)}}) = w(Q \cap \text{dom}(\mathcal{Q}_m)),$$

so Q is a 1-pack over \mathcal{Q}_m . Thus, (\mathcal{Q}_m) is a 1-packing sequence, so by Lemma 7.5, (\mathcal{Q}_m) stabilizes modulo w -deficient. But for each $x \in D$, $\bigcup_m [x]_{E(\mathcal{Q}_m)} = [x]_{E_\infty}$ is w -infinite, so $[D]_{E_\infty}$ is w -deficient, and $X = [D]_{E_\infty}$. \square

When building an approximately saturated tiling (in Theorem 7.13), we need label-maximizing Borel maximal matchings in labeled Borel bipartite graphs. We prove their existence now:

Lemma 7.12. *Let Y, Z be disjoint standard Borel spaces, and let $H \subseteq Y \times Z$ be a locally countable directed Borel graph and $\ell : H \rightarrow \mathbb{R}^+$ be a Borel function. There is a Borel maximal matching⁸ $M \subseteq H$ such that for each edge $(y, z) \in M$,*

$$\ell(y, z) \geq 1 \text{ or } \ell(y, z) \geq \frac{1}{2} \sup_{z' \in H_y \setminus Z_M} \ell(y, z'),$$

where Z_M is the set of M -matched⁹ points in Z .

Proof. Because E_H is a countable Borel equivalence relation on $Y \cup Z$, [KM04, Lemma 7.3] implies that H , as well as its Borel subgraphs, admit Borel maximal matchings. We recursively define a sequence $(M_n)_{n \in \mathbb{N}}$ of pairwise *vertex-disjoint* Borel matchings, i.e. no vertex in $Y \cup Z$ is M_k -matched and M_n -matched for $k \neq n$, so the union $M := \bigcup_n M_n$ is a Borel matching.

Let $H_0 := \{(y, z) \in H : \ell(y, z) \geq 1\}$, and let $M_0 \subseteq H_0$ be a Borel maximal matching. Suppose that the matchings M_k , $0 \leq k < n$, are defined and are pairwise vertex-disjoint, so $\overline{M}_n := \bigcup_{k < n} M_k$ is a matching, and let $Y_{\overline{M}_n}, Z_{\overline{M}_n}$ denote the subsets of Y, Z , respectively, of all points that are \overline{M}_n -matched. Let

$$H_n := \left\{ (y, z) \in H \cap (Y \setminus Y_{\overline{M}_n}) \times (Z \setminus Z_{\overline{M}_n}) : \ell(y, z) \geq \frac{1}{2} \sup \{ \ell(y, z') : z' \in H_y \setminus Z_{\overline{M}_n} \} \right\},$$

and let $M_n \subseteq H_n$ be a Borel maximal matching. This completes the construction of (M_n) . It is immediate from the definitions that for any $(y, z) \in M$, either $(y, z) \in M_0$, so $\ell(y, z) \geq 1$, or $(y, z) \in M_n$ for some $n \geq 1$, so $\ell(y, z) \geq \frac{1}{2} \sup_{z' \in H_y \setminus Z_{\overline{M}_n}} \ell(y, z') \geq \frac{1}{2} \sup_{z' \in H_y \setminus Z_M} \ell(y, z')$.

It remains to show that M is maximal. We take a $y \in Y$ that it is not M -matched and aim to show that $H_y \setminus Z_M = \emptyset$, where $H_y := \{z' \in Z : (y, z') \in H\}$. By the maximality of M_0 , $s_0 := \sup_{z \in H_y \setminus Z_{\overline{M}_1}} \ell(y, z) \leq 1$. Then the maximality of each M_k implies, by induction on n , that $s_n := \sup \{ \ell(y, z) : z \in Z \setminus Z_{\overline{M}_{n+1}} \} \leq \frac{1}{2} s_{n-1} \leq 2^{-n}$ for each $n \geq 1$. Because $H_y \setminus Z_M = \bigcup_{n \geq 1} (H_y \setminus Z_{\overline{M}_n})$, $\sup_{z \in H_y \setminus Z_M} \ell(y, z) \leq \inf_{n \geq 0} s_n = 0$, so $H_y \setminus Z_M = \emptyset$. \square

Theorem 7.13. *Any Borel $\mathcal{S} \subseteq [X]_E^{<\infty}$ admits a Borel tiling $\mathcal{P} \subseteq \overline{\mathcal{S}}$ approximately saturated within \mathcal{S} modulo w -deficient. In fact, given a Borel tiling $\mathcal{P}_0 \subseteq \mathcal{S}$, the tiling \mathcal{P} can be chosen so that it admits a saturating \mathcal{S} -approximation starting with \mathcal{P}_0 .*

Proof. Fix a Borel coloring of the intersection graph on $[X]_E^{<\infty}$ with natural numbers [KM04, Proof of Lemma 7.3], and let (k_n) be a sequence of natural numbers in which each (color) $k \in \mathbb{N}$ appears infinitely-many times.

Having \mathcal{P}_0 given (otherwise, take $\mathcal{P}_0 := \emptyset$), we recursively build an injective sequence $(\mathcal{P}_n)_{n \geq 1}$ of Borel tilings contained in \mathcal{S} . Fixing $n \in \mathbb{N}$, suppose that \mathcal{P}_n is defined. Let

- $\mathcal{S}_n^\perp :=$ the collection of all sets in \mathcal{S} of color k_n that are disjoint from $\text{dom}(\mathcal{P}_n)$;
- $\mathcal{Q}_n :=$ the collection of all $Q \in [X]_E^{<\infty} \setminus \mathcal{S}$ of color k_n that are disjoint from $\text{dom}(\mathcal{P}_n)$.

⁸A set of edges M is a *matching* if no two edges are adjacent (ignoring the direction of the edges).

⁹A vertex is *M -matched* if there is an edge in M incident to it.

Let $\mathfrak{G}_n := \{(P, Q) \in \mathcal{P}_n \times \mathcal{Q}_n : P \cup Q \in \mathcal{S}\}$. This graph is locally countable because $P \cup Q \in \mathcal{S}$ implies that $[P]_E = [Q]_E$, and it is Borel (using Luzin–Novikov uniformization), so by Lemma 7.12 applied to \mathfrak{G}_n with $\ell(P, Q) := \frac{w(Q)}{w(P)}$, there is a Borel maximal matching $\mathfrak{M}_n \subseteq \mathfrak{G}_n$ such that, for each $(P, Q) \in \mathfrak{M}_n$,

$$w(Q) \geq w(P) \text{ or } w(Q) \geq \frac{1}{2} \sup \{w(Q') : Q' \in (\mathfrak{G}_n)_P \cap \mathcal{Q}'_n\}, \quad (7.14)$$

where \mathcal{Q}'_n denotes set of all \mathfrak{M}_n -unmatched vertices in \mathcal{Q}_n . Note that because $\mathcal{P}_n, \mathcal{Q}_n$ are tilings with disjoint domains and \mathfrak{M}_n is a matching, $\mathcal{M}_n := \{P \cup Q : (P, Q) \in \mathfrak{M}_n\}$ is a tiling as well. Take $\mathcal{P}_{n+1} := \mathcal{S}_n^\perp \cup \mathcal{M}_n \cup (\mathcal{P}_n|_D)$, where $D := \text{dom}(\mathcal{P}_n) \setminus \text{dom}(\mathcal{S}_n^\perp \cup \mathcal{M}_n)$.

Applying Lemma 7.11 to (\mathcal{P}_n) and ignoring a w -deficient E -invariant Borel set, we assume that all the tiles in $\mathcal{P} := \bigvee_n \mathcal{P}_n$ are w -finite. Thus, $\mathcal{P} \subseteq \overline{\mathcal{S}}$.

It remains to show that (\mathcal{P}_n) is a saturating \mathcal{S} -approximation for \mathcal{P} . Let $Q' \in [X]_E^{<\infty}$ be disjoint from $\text{dom}(\mathcal{P})$. If $Q' \in \mathcal{S}$, then for any $n \in \mathbb{N}$ such that k_n is the color of Q' , the construction would put Q' in \mathcal{P}_{n+1} , contradicting $Q' \cap \text{dom}(\mathcal{P}) = \emptyset$. Thus, $Q' \notin \mathcal{S}$. Suppose towards a contradiction that there is $x \in X$ such that $P_n \cup Q' \in \mathcal{S}$ for all $n \geq N$, for some $N \in \mathbb{N}$, where $P_n := [x]_{E(\mathcal{P}_n)}$. Let $n \geq N$ be large enough so that

$$w([x]_{E(\mathcal{P})} \setminus P_n) < \min \left\{ w(P_n), \frac{1}{2} w(Q') \right\}, \quad (7.15)$$

and moreover, choose n so that k_n is the color of Q' . Then $(P_n, Q') \in \mathfrak{G}_n$ and yet Q' is not \mathfrak{M}_n -matched because all \mathfrak{M}_n -matched tiles are contained in $\text{dom}(\mathcal{P}_{n+1})$, while $Q' \cap \text{dom}(\mathcal{P}_{n+1}) = \emptyset$. By the maximality of the matching \mathfrak{M}_n , P_n must be \mathfrak{M}_n -matched with some $Q \in \mathcal{Q}_n$. By (7.14), $w(Q) \geq w(P_n)$ or $w(Q) \geq \frac{1}{2} w(Q')$. But $[x]_{E(\mathcal{P})} \setminus P_n \supseteq Q$, so $w([x]_{E(\mathcal{P})} \setminus P_n) \geq w(P_n)$ or $w([x]_{E(\mathcal{P})} \setminus P_n) \geq \frac{1}{2} w(Q')$, contradicting (7.15). \square

7.D. Packed and approximately saturated tilings.

Proposition 7.16. *For any Borel collection $\mathcal{S} \subseteq [X]_E^{<\infty}$ and $p \in \mathbb{R}^+$, there is a Borel tiling $\mathcal{P} \subseteq \overline{\mathcal{S}}$ such that, modulo w -deficient, it is approximately saturated within \mathcal{S} and moreover, admits a saturating \mathcal{S} -approximation of p -packed tilings within \mathcal{S} .*

Proof. Applying Theorem 7.6 and throwing out an w -deficient set, we get a Borel tiling $\mathcal{P}_0 \subseteq \mathcal{S}$ p -packed within \mathcal{S} . In particular, any extension $\mathcal{P}'_0 \subseteq \mathcal{S}$ of \mathcal{P}_0 is p -packed within \mathcal{S} by Observation 7.4(b). Applying Theorem 7.13 to \mathcal{S} with the initial tiling \mathcal{P}_0 gives a Borel tiling $\mathcal{P} \subseteq \overline{\mathcal{S}}$ with a desired saturating \mathcal{S} -approximation. \square

Theorem 7.17. *For any Borel collection $\mathcal{S} \subseteq [X]_E^{w<\infty}$ with finitely based quotients and $p \in \mathbb{R}^+$, there is a Borel tiling $\mathcal{P} \subseteq \overline{\mathcal{S}}$ that is p -packed within \mathcal{S} and approximately saturated within $\mathcal{S}' := \mathcal{S} \cap [X]_E^{<\infty}$, modulo w -deficient.*

Proof. Applying Proposition 7.16 to \mathcal{S}' and $\frac{p}{2}$, we get a Borel tiling $\mathcal{P} \subseteq \overline{\mathcal{S}'}$ with a saturating \mathcal{S}' -approximation (\mathcal{P}_n) such that \mathcal{P}_0 is $\frac{p}{2}$ -packed within \mathcal{S}' . Because \mathcal{S} has finitely based quotients, Lemma 7.7 implies that \mathcal{P}_0 is p -packed within \mathcal{S} . By Observation 7.4(b), same is true for \mathcal{P} because $\mathcal{P} \geq \mathcal{P}_0$. \square

8. COCYCLED GRAPH VISIBILITY

Throughout this section, let G be a locally countable Borel graph on a standard Borel space X and let $w : E_G \rightarrow \mathbb{R}^+$ be a Borel cocycle.

8.A. Definitions and basic properties. In our proofs, we would like to obtain G -connected w -finite sets of arbitrarily large w -ratio. The following notion helps replace w_* with w .

Definition 8.1. Let $x \in X$.

- A (G, w) -visible neighborhood of x is any G -connected set $V \ni x$ such that $w(x) \geq w(v)$ for each $v \in V$.
- A point $y \in X$ is said to be (G, w) -visible from x , denoted by $y \leq_G^w x$, if x admits a (G, w) -visible neighborhood containing y . (The intuition comes from thinking of w as relative heights, so y is visible from x if on some path from x to y , no point is taller than x .)
- We refer to the sets $(\leq_G^w)^x := \{y \in X : y \leq_G^w x\}$ and $(\leq_G^w)_x := \{y \in X : x \leq_G^w y\}$ as the \leq_G^w -downward and \leq_G^w -upward cones of x .
- Call a set C a \leq_G^w -downward (resp. upward) cone if C is a \leq_G^w -downward (resp. upward) cone of some $x \in X$.
- We say that G has finite w -visibility if every \leq_G^w -downward cone is w -finite.

We omit G, w from the notation if they are understood from the context. Similarly, we just say downward (resp. upward) cone.

Proposition 8.2. Let G, w be as above, and let $x, y \in X$ be points in the same G -connected component.

- Partial quasi-order: \leq is a transitive relation (a partial quasi-order).
- Amalgamation: There is $z \in X$ such that $x \leq z$ and $y \leq z$. In other words, any two downward cones in the same G -connected component are contained in a downward cone.
- Linearity on upward cones: The quasi-order \leq is linear on each upward cone, i.e. if $x \leq y$ and $x \leq z$, then $y \leq z$ or $z \leq y$.
- Covering with downward cones: Every $U \in [X]_G^{w < \infty}$ is contained in a downward cone.
- Cofinality of upward cones: For each $x \in X$, $\sup_{y \in [x]_{E_G}} w(y) = \sup_{z \geq x} w(z)$.

Proof. (a) is immediate. For (b), letting P be a G -path from x to y , we see that x and y are visible from any $z \in \text{Max}_w P$. For (c), if $x \leq y$ and $x \leq z$ then $w(y) \leq w(z)$ implies $y \leq z$. For (d), because $w(U) < \infty$, $\text{Max}_w U$ is nonempty by Observation 2.1, so $U \subseteq (\leq)^x$ for any $x \in \text{Max}_w U$. For (e), note that for any $y \in [x]_{E_G}$, (b) gives a $z \geq x, y$, in particular, $w(z) \geq w(y)$. \square

8.B. Finite visibility and hyperfiniteness. Here, we prove the following sufficient condition for hyperfiniteness in terms of a cocycle mentioned in the introduction:

Theorem 1.8. Let G be a locally countable Borel graph on a standard Borel space X and let $w : E_G \rightarrow \mathbb{R}^+$ be a Borel cocycle. If G has finite w -visibility, then it is hyperfinite.

Proof. This proof was significantly simplified by the suggestion of an anonymous referee.

Suppose that G has finite w -visibility. By Lemma 2.7, the w -finite part of E_G is smooth, hence hyperfinite, so we assume without loss of generality that E_G is w -infinite.

Claim 8.3. Every upward cone is w -infinite.

Proof of Claim. Suppose towards a contradiction that $U := (\leq)_x$ is w -finite for some $x \in X$, so $\text{Max}_w U \neq \emptyset$, by Observation 2.1. But then Proposition 8.2(e) implies that every point in $[x]_{E_G}$ is visible from any $y \in \text{Max}_w U$, contradicting finite visibility because $[x]_{E_G}$ is w -infinite. \boxtimes

Claim 8.4. \leq is a well-order on any upward cone U . Moreover, for any nonempty $A \subseteq U$, $\text{Min}_w A$ is nonempty and finite.

Proof of Claim. By Proposition 8.2(c), any two points in $\text{Min}_w A$ are visible to each other. Thus, finite visibility implies that $\text{Min}_w A$ is w -finite, and hence finite.

Suppose towards a contradiction that $\text{Min}_w A = \emptyset$, so there is a sequence $(x_n) \subseteq A$ such that $w(x_n) > w(x_{n+1})$. This implies that $x_n \geq x_{n+1}$ by the linearity of \geq on U (Proposition 8.2(c)). But $U = (\leq)_x$ for some $x \in X$, so $w(x_n) \geq w(x)$ for all n , and every x_n is visible from x_0 , contradicting finite visibility (at x_0). \square

For each $x \in X$, put $C^+(x) := (\leq)_x \setminus \text{Min}_w(\leq)_x$. Claims 8.3 and 8.4 imply that $C^+(x) \neq \emptyset$, so fixing a Borel linear order $<$ on X , we define a transformation $T : X \rightarrow X$ by $x \mapsto$ the $<$ -least element of the set $\text{Min}_w C^+(x)$, which exists by Claim 8.4. Clearly, $x \leq T(x)$ and $w(x) < w(T(x))$.

Claim 8.5. For any E -related $x, y \in X$, there are $n, m \in \mathbb{N}$ such that $T^n(x) = T^m(y)$.

Proof of Claim. By the amalgamation property (Proposition 8.2(b)), it is enough to show the statement assuming $x \leq y$. By finite visibility (at y), there are only finitely-many points $z \in (\leq)_x$ with $w(z) \leq w(y)$. Because the sequence $(w_x(T^k(x)))_{k \in \mathbb{N}}$ is strictly increasing and $(T^k(x))_{k \in \mathbb{N}} \subseteq (\leq)_x$, there is a least $n \in \mathbb{N}$ such that $w(T^n(x)) > w(y)$. Then $n \geq 1$ and $w(T^{n-1}(x)) = w(y)$, so $T^{n-1}(x)$ and y are visible to each other by Proposition 8.2(c). But then the upward cones of $T^{n-1}(x)$ and y coincide, so $T^n(x) = T(y)$. \square

Thus, the orbit equivalence relation of the (semigroup) action of T on X is exactly E_G , so by [DJK94, Corollary 8.2], E_G is hyperfinite. \square

8.C. Tiling with sets of large w -ratio. As before, throughout this subsection, we let G be a locally countable Borel graph on a standard Borel space X and let $w : E_G \rightarrow \mathbb{R}^+$ be a Borel cocycle. Moreover, we let μ be a w -invariant Borel probability measure on X .

The goal of this subsection (Lemma 8.11) is to build a Borel tiling \mathcal{P} with a large domain whose tiles are of large w -ratio and perhaps satisfy other properties. This is not hard in the pmp setting, i.e. when w -ratio is just cardinality; indeed, taking a saturated tiling works. However, typically w -ratio is not equal to cardinality and it is not monotone (under subsets), which makes such a tiling harder to build.

We say that $V \subseteq X$ is a (G, w) -visible neighborhood of $U \in [X]_G^{w < \infty}$ if V is G -connected, contains U , and $w(U) \geq w(v)$ for each $v \in V$. In this case, for a w -finite V ,

$$\frac{w(V)}{w(U)} \leq w_*(V) \leq w_*(U) \frac{w(V)}{w(U)}. \quad (8.6)$$

Indeed, $w_*(V) = \frac{w(V)}{\max_w V}$ and $\frac{w(V)}{w(U)} \leq \frac{w(V)}{\max_w V} \leq \frac{w(V)}{\max_w U} = w_*(U) \frac{w(V)}{w(U)}$.

The following captures the kind of properties we will allow the tiles to satisfy.

Definition 8.7. Call $\mathcal{S} \subseteq [X]_G^{w < \infty}$ (G, w) -conic (resp. finitely (G, w) -conic) if for each $U \in [X]_G^{w < \infty}$ (resp. $U \in [X]_G^{< \infty}$), all of its w -large enough w -finite (resp. finite) (G, w) -visible neighborhoods are in \mathcal{S} . By (8.6), “ w -large enough” can be replaced with “large enough w -ratio”.

Note that (G, w) -conic implies finitely (G, w) -conic.

Example 8.8. For any $L > 0$, the collection of all $V \in [X]_G^{\mathfrak{w} < \infty}$ (resp. $V \in [X]_G^{< \infty}$) with $\mathfrak{w}_*(V) > L$ is (G, \mathfrak{w}) -conic (resp. finitely (G, \mathfrak{w}) -conic).

Lemma 8.9. If a Borel tiling $\mathcal{P} \subseteq [X]_G^{\mathfrak{w} < \infty}$ is maximal within a finitely (G, \mathfrak{w}) -conic $\mathcal{S} \subseteq [X]_G^{< \infty}$, then $G_{-\text{dom}(\mathcal{P})}$ has finite \mathfrak{w} -visibility. In particular, $\text{dom}(\mathcal{P})$ is a hyperfinitizing vertex-cut for G .

Proof. The last part is due to Theorem 1.8. As for finite \mathfrak{w} -visibility, suppose towards a contradiction that there is an $x \in X \setminus \text{dom}(\mathcal{P})$ that admits arbitrarily \mathfrak{w} -large finite (G, \mathfrak{w}) -visible neighborhoods $V \subseteq X \setminus D$. Then a \mathfrak{w} -large enough such V belongs to \mathcal{S} , contradicting the maximality of \mathcal{P} . \square

The following is the reason why we use (G, \mathfrak{w}) -conic and not just finitely (G, \mathfrak{w}) -conic.

Proposition 8.10. Let F be a G -connected \mathfrak{w} -finite Borel equivalence relation. If $\mathcal{S} \subseteq [X]_G^{\mathfrak{w} < \infty}$ is (G, \mathfrak{w}) -conic, then $\mathcal{S}_{/F}$ is $(G_{/F}, \mathfrak{w}_{/F})$ -conic.

Proof. For any $U \in [X_{/F}]_{G_{/F}}^{\mathfrak{w}_{/F} < \infty}$, the π_F -preimage of any $(G_{/F}, \mathfrak{w}_{/F})$ -visible neighborhood V of U is a (G, \mathfrak{w}) -visible neighborhood of $\pi^{-1}(U)$ and $\frac{\mathfrak{w}(\pi_F^{-1}(V))}{\mathfrak{w}(\pi_F^{-1}(U))} = \frac{\mathfrak{w}_{/F}(V)}{\mathfrak{w}_{/F}(U)}$. \square

We are now ready to prove the main result of this subsection.

Lemma 8.11 (μ -co- ε tiling). If G is μ -nowhere hyperfinite, then for any (G, \mathfrak{w}) -conic $\mathcal{S} \subseteq [X]_G^{\mathfrak{w} < \infty}$ and $\varepsilon > 0$, there is a Borel tiling $\mathcal{P} \subseteq \mathcal{S}$ with a μ -co- ε domain.

Proof. For each $U \in [X]_G^{\mathfrak{w} < \infty}$, let $r(U)$ be the infimum of all $r > 0$ such that any (G, \mathfrak{w}) -visible neighborhood V of U with $\mathfrak{w}_*(V) > r$ belongs to \mathcal{S} ; write $r(x)$ if $U = \{x\}$. For any G -connected \mathfrak{w} -finite Borel equivalence relation F and $L > 0$, let $\mathcal{S}'_{/F}(L)$ denote the collection of all finite $P \in \mathcal{S}_{/F}$ with $(\mathfrak{w}_{/F})_*(P) > \max\{L, r(U)\}$ for all $U \in \text{Max}_{\mathfrak{w}_{/F}} P$. We just write $\mathcal{S}'(L)$ if $F = \text{Id}_X$.

Note that for any $x' \in X_{/F}$, any $(G_{/F}, \mathfrak{w}_{/F})$ -visible neighborhood V of x' with $(\mathfrak{w}_{/F})_*(V) > r(x')$ belongs to $\mathcal{S}_{/F}$ because $\mathfrak{w}_*(\pi_F^{-1}(V)) \geq (\mathfrak{w}_{/F})_*(V)$ (Observation 2.8). We call this the *main property* of r . This and the fact that cocycle-ratio can only decrease in the quotient (Observation 2.8) allow us below to work mod F and assume without loss of generality that $F = \text{Id}_X$.

Claim 8.12. $\mathcal{S}'_{/F}(L)$ is finitely $(G_{/F}, \mathfrak{w}_{/F})$ -conic.

Proof of Claim. Follows from Proposition 8.10 and the fact that the additional condition in the definition of $\mathcal{S}'_{/F}(L)$ is a lower bound on $\mathfrak{w}_{/F}$ -ratio. \square

Claim 8.13. $\overline{\mathcal{S}'_{/F}(L)} \subseteq \mathcal{S}_{/F}$.

Proof of Claim. We work mod F , so assume without loss of generality that $F = \text{Id}_X$. Let $Q_m \in \mathcal{S}'(L)$ be an increasing sequence of sets such that $Q = \bigcup_m Q_m$ is still \mathfrak{w} -finite. Then by Lemma 2.6, there is m such that $\text{Max}_{\mathfrak{w}} Q_m = \text{Max}_{\mathfrak{w}} Q$ and hence $\mathfrak{w}_*(Q) \geq \mathfrak{w}_*(Q_m) > r(x)$ for all $x \in \text{Max}_{\mathfrak{w}} Q$, so $Q \in \mathcal{S}$ by the main property of r . \square

Let $(L_n)_{n \geq 0}$ be an increasing unbounded sequence of positive reals. We recursively define a coherent sequence $(\mathcal{P}_n)_{n \geq 0}$ of Borel tilings contained in $[X]_G^{\mathfrak{w} < \infty}$; in particular, the equivalence relations $F_n := \bigcup_{k < n} E(\mathcal{P}_k)$ are increasing, where $F_0 := \text{Id}_X$. For $n \geq 0$, suppose that F_n is defined. Theorem 7.13 applied to $\mathcal{S}'_{/F_n}(L_n)$ gives a Borel tiling $\mathcal{Q} \subseteq$

$\overline{\mathcal{S}'_{/F_n}(L_n)}$ approximately saturated within $\mathcal{S}'_{/F_n}(L_n)$ modulo $w_{/F_n}$ -deficient. Putting $\mathcal{P}_n := \{\pi_{F_n}^{-1}(Q) : Q \in \mathcal{Q}\}$ finishes the construction.

By Corollary 5.6, if an $(E_G)_{/F_n}$ -invariant Borel $A_n \subseteq X_{/F_n}$ is $w_{/F_n}$ -deficient, then it is $\mu_{/F_n}$ -null, so $\pi_{F_n}^{-1}(A_n)$ is μ -null. Thus, discarding countably-many E_G -invariant μ -null sets from X , we have that for each $n \geq 0$, $(\mathcal{P}_n)_{/F_n} \subseteq \overline{\mathcal{S}'_{/F_n}(L_n)}$ is approximately saturated within $\mathcal{S}'_{/F_n}(L_n)$. By Claim 8.13, $(\mathcal{P}_n)_{/F_n} \subseteq \mathcal{S}_{/F_n}$, so $\mathcal{P}_n \subseteq \mathcal{S}$.

Claim 8.14. *For each $n \in \mathbb{N}$, $D_n := \text{dom}(\mathcal{P}_n)$ is a hyperfinitizing vertex-cut for G .*

Proof of Claim. Because $(\mathcal{P}_n)_{/F_n}$ is maximal within $\mathcal{S}'_{/F_n}(L_n)$ (Observation 7.10(a)) and $\mathcal{S}'_{/F_n}(L_n)$ is finitely $(G_{/F}, w_{/F})$ -conic (Claim 8.12), Lemma 8.9 implies that $\pi_F(D_n) = \text{dom}((\mathcal{P}_n)_{/F_n})$ is a hyperfinitizing vertex-cut for $G_{/F}$, i.e. $(G_{/F})_{-\pi_F(D_n)}$ is hyperfinite. But then G_{-D_n} is hyperfinite because π_F is a Borel reduction $E_G \rightarrow (E_G)_{/F}$ and hyperfiniteness pulls back under Borel reductions [JKL02, 1.3(ii)]. \square

Put $D_\infty := \limsup_n D_n := \{x \in X : x \in D_n \text{ for infinitely many } n \in \mathbb{N}\}$ and $F_\infty := \bigcup_n F_n$.

Claim 8.15. *D_∞ is E_G -invariant.*

Proof of Claim. Suppose not, so there are G -adjacent points $x \in D_\infty$ and $y \in X \setminus D_\infty$. Let $N_0 \in \mathbb{N}$ be large enough so that $y \notin D_n$ for all $n \geq N_0$; hence $[y]_{F_\infty} = [y]_{F_n}$ for all $n \geq N_0$. Because $[x]_{F_\infty}$ is w -infinite (by Observation 2.5), there is $N_1 \geq N_0$ such that $w([x]_{F_n}) > w([y]_{F_\infty})$ for all $n \geq N_1$. Finally, take $n \geq N_1$ so that $[x]_{F_{n+1}} \in \mathcal{P}_n$. We will contradict that $\mathcal{Q} := (\mathcal{P}_n)_{/F_n}$ is approximately saturated within $\mathcal{S}'_{/F_n}(L_n)$.

Putting $x' := [x]_{F_n}$ and $y' := [y]_{F_n} = [y]_{F_\infty}$, so $w(x') > w(y')$, and working mod F_n , we assume without loss of generality that $F_n = \text{Id}_X$ and drop F_n from the notation.

Let (\mathcal{Q}_m) be a saturating $\mathcal{S}'(L_n)$ -approximation for \mathcal{Q} , and put $Q := [x']_{E(\mathcal{Q})}$ and $Q_m := [x']_{E(\mathcal{Q}_m)}$, so $Q = \bigcup_m Q_m$. Fix any $k \in \mathbb{N}$. Then $Q'_k := Q_k \cup \{y'\}$ is G -connected and finite, and $\max_w Q_k \geq w(x') > w(y')$, so $\text{Max}_w(Q'_k) = \text{Max}_w Q_k$. In particular, $w_*(Q'_k) = \frac{w(Q_k) + w(y')}{\max_w Q_k} > w_*(Q_k) > \max\{L_n, r(x_k)\}$ for all $x_k \in \text{Max}_w Q'_k = \text{Max}_w Q_k$ because $Q_k \in \mathcal{S}'(L_n)$. But then the main property of r implies that $Q'_k \in \mathcal{S}$, and hence $Q'_k \in \mathcal{S}'(L_n)$, contradicting that (\mathcal{Q}_m) is a saturating $\mathcal{S}'(L_n)$ -approximation. \square

Claim 8.16. *D_∞ is μ -conull.*

Proof of Claim. Let $X' := X \setminus D_\infty$ and suppose that it is not null. Because X' is G -invariant and G is μ -nowhere hyperfinite, $G|_{X'}$ is not μ -hyperfinite, so $p := \text{hvp}_\mu(G|_{X'}) > 0$. By Claim 8.14, $D_n \cap X'$ is a hyperfinitizing cut for $G|_{X'}$, so $\mu(D_n \cap X') \geq p$ for all $n \in \mathbb{N}$. But then $D_\infty \cap X' = \bigcap_n \bigcup_{m \geq n} (D_m \cap X')$ also has measure at least p by the downward continuity of μ , contradicting $D_\infty \cap X' = \emptyset$. \square

In particular, $\bigcup_{n \in \mathbb{N}} D_n$ is conull, so for a large enough $N \in \mathbb{N}$, $D := \bigcup_{n \leq N} D_n$ is μ -co- ε . Then the set \mathcal{P} of all F_N -classes contained in D is contained in $\bigcup_{n \leq N} \mathcal{P}_n \subseteq \mathcal{S}$ and $\text{dom}(\mathcal{P}) = D$ is μ -co- ε . \square

9. VISIBLE ASYMPTOTIC AVERAGES ON A GRAPH

If Theorem 1.1 is indeed true, then for any $f \in L^\infty(X, \mu)$ and a.e. $x \in X$, there must be arbitrarily w -large finite G -connected sets containing x over which the w -average of f is

arbitrarily close to $\mathbb{E}(f|\mathcal{B}_{E_G})(x)$. Motivated by this, we look at the set of all reals in general that are achievable in this manner, thus defining a new invariant developed in this section.

Throughout this section, let G be a locally countable (abstract) graph on a set X and let $w : E_G \rightarrow \mathbb{R}^+$ a cocycle. We also let $f : X \rightarrow \mathbb{R}$ be a bounded function.

9.A. For an abstract graph. [MT17, Definition 8.2] introduces the set $\mathcal{A}_G^w f(x)$ of asymptotic w -weighted means along G in the G -connected component $[x]_{E_G}$, where $w : X \rightarrow \mathbb{R}^+$ is a weight-function. This set is independent of the representative x of the G -connected component [MT17, Proposition 8.3] and it is a closed interval when w is a bounded function [MT17, Proposition 8.5]. Here, we generalize this definition to arbitrary cocycles on E_G .

Definition 9.1. For a G -connected set C and $x \in C$, we call $r \in \mathbb{R}$ a (G, w) -asymptotic average of f at x over C if there are arbitrarily w -large finite G -connected sets $V \subseteq C$ containing x with $A_V^w f$ arbitrarily close to r ; more precisely, for every $\varepsilon > 0$ and $L > 0$, there is a finite G -connected set $V \subseteq C$ containing x with $w_x(V) \geq L$ and $A_V^w f \approx_\varepsilon r$. We denote by $\mathcal{A}_{G|C}^w f(x)$ the set of all such $r \in \mathbb{R}$.

The compactness of the interval $[-\|f\|_\infty, \|f\|_\infty]$ immediately implies:

Observation 9.2. If C is a w -infinite G -connected set then $\mathcal{A}_{G|C}^w f(x) \neq \emptyset$ for each $x \in C$.

Remark 9.3. For a \leq_G^w -downward cone $C = (\leq_G^w)^x$, we note that the restriction of w to C is a coboundary, being the differential of the weight-function $w_x : C \rightarrow \mathbb{R}^+$. Note that w_x is bounded above by 1. It is easy to see that the definition of a (G, w) -asymptotic average over C as above coincides with that of an asymptotic w_x -weighted mean along $G|_C$ as defined in [MT17, Definition 8.2]. Thus, the following three lemmas are just restatements of [MT17, 8.4 and 8.5] in our terms.

Lemma 9.4. The function $\mathcal{A}_{G|C}^w f$ is constant for any G -connected set $C \subseteq X$, i.e. $\mathcal{A}_{G|C}^w f(x) = \mathcal{A}_{G|C}^w f(y)$ for any $x, y \in C$.

Proof. There is a G -path connecting x and y , whose effect on the averages of f over arbitrarily w -large sets is arbitrarily small. \square

Lemma 9.5 (Intermediate value property). Let $U, V \in [X]_G^{w<\infty}$ be such that $U \subseteq V$ and let

$$\Delta := 2\|f\|_\infty \frac{\max_w(V \setminus U)}{w(U)}.$$

For every real r between $A_U^w f$ and $A_V^w f$, there is $W \in [X]_G^{w<\infty}$ with $U \subseteq W \subseteq V$ and $A_W^w f \approx_\Delta r$.

Proof. Firstly note that there is $V' \in [X]_G^{w<\infty}$ with $U \subseteq V' \subseteq V$ such that $V' \setminus U$ is finite, yet $w(V') \geq (1 - \varepsilon)w(V)$, where $\varepsilon \in (0, 1)$ is small enough to guarantee $A_{V'}^w f \approx_\Delta A_V^w f$ by Lemma 3.1(b). Now, we can add the vertices of $V' \setminus U$ to U one-by-one, obtaining a finite sequence of G -connected supersets of U increasing up to V' . It remains to observe that adding one vertex can change the average at most by Δ again by Lemma 3.1(b). \square

Lemma 9.6. For any \leq_G^w -downward cone C and $x \in C$, $\mathcal{A}_{G|C}^w f(x)$ is closed and convex; thus, it is a closed subinterval of $[-\|f\|_\infty, \|f\|_\infty]$.

Proof. Suppose $\mathcal{A}_{G|C}^w f(x) \neq \emptyset$, which implies that $w(C) = \infty$. Then the closedness of $\mathcal{A}_{G|C}^w f(x)$ follows from the asymptotic nature of the definition of $\mathcal{A}_{G|C}^w f$. Convexity follows from Lemma 9.5. Indeed, for any $r_1, r_2 \in \mathcal{A}_{G|C}^w f$ and a real r between r_1, r_2 , there are arbitrarily w -large G -connected subsets $U \ni x$ of C with $A_U^w f$ arbitrarily close to r_1 . For each such U , there are arbitrarily w -large G -connected subsets $V \ni x$ of C with $A_V^w f$ arbitrarily close to r_2 and $\frac{w(U)}{w(V)}$ arbitrarily small. Thus, we can make $A_{U \cup V}^w f$ arbitrarily close to $A_V^w f$, hence we may choose V containing U to begin with. We may also assume that r is between $A_U^w f$ and $A_V^w f$ because otherwise, U or V would witness the arbitrary closeness of r to $\mathcal{A}_{G|C}^w f(x)$. Finally, we may apply Lemma 9.5 to U, V and r and obtain a set W with $U \subseteq W \subseteq V$ with $A_W^w f \approx_\Delta r$, where

$$\Delta := 2\|f\|_\infty \frac{\max_w(V \setminus U)}{w(U)} = 2\|f\|_\infty \frac{\max_{w_y}(V \setminus U)}{w_y(U)} \leq \frac{2\|f\|_\infty}{w_y(U)}$$

for some/any $y \in \text{Max}_w C$.¹⁰ Thus, choosing U arbitrarily w -large makes Δ arbitrarily small, hence r arbitrarily close to $\mathcal{A}_{G|C}^w f$. \square

Definition 9.7. We call $r \in \mathbb{R}$ a (G, w) -visible asymptotic average of f at $x \in X$ if it is a (G, w) -asymptotic average of f over some \preceq_G^w -downward cone $C \ni x$. We denote by $\mathcal{A}_G^w f(x)$ the set of all (G, w) -visible¹⁰ asymptotic averages of f at x .

Proposition 9.8 (Invariance). *The map $x \mapsto \mathcal{A}_G^w f(x)$ is E_G -invariant.*

Proof. This is immediate from the amalgamation property of downward cones (Proposition 8.2(b)) and Lemma 9.4. \square

Proposition 9.9 (Convexity). *For any $x \in X$, $\mathcal{A}_G^w f(x)$ is an increasing union of sets of the form $\mathcal{A}_{G|C}^w f(x)$, where C is a downward cone. In particular, it is a convex subset of $[-\|f\|_\infty, \|f\|_\infty]$.*

Proof. By definition, $\mathcal{A}_G^w f(x)$ is a union of sets of the form $\mathcal{A}_{G|C}^w f(x)$. Because downward cones amalgamate (Proposition 8.2(b)), this union is directed, so it can be turned into an increasing union, using the countability of $[x]_G$. The convexity of $\mathcal{A}_G^w f(x)$ is then due to Lemma 9.6 and the fact that $C_1 \subseteq C_2$ implies $\mathcal{A}_{G|C_1}^w f \subseteq \mathcal{A}_{G|C_2}^w f$. \square

9.B. For a measurable graph. Equipping X with a standard Borel structure, we now suppose further that G, w , and f are Borel, and we let μ be a w -invariant Borel probability measure on X .

Encoding intervals in \mathbb{R} as points in $(\{0, 1, 2, 3\} \times \mathbb{R})^2$, where 0, 1, 2, 3 encode whether each endpoint is open or closed, we equip the set \mathcal{I} of all intervals with a natural standard Borel structure.

Proposition 9.10. *The map $x \mapsto \mathcal{A}_G^w f(x) : X \rightarrow \mathcal{I}$ is Borel.*

Proof. The fact that the image is in \mathcal{I} is by Proposition 9.9. The Borelness follows by the definition of visible asymptotic averages, using Luzin–Novikov uniformization, which turns scanning over each E_G -class into a natural number quantifier. \square

¹⁰Having a bound on $\max_{w_y}(V \setminus U)$ independent of U, V in the proof of Lemma 9.6 is the reason why only visible asymptotic averages are included in $\mathcal{A}_G^w f$ (in Definition 9.7).

We now obtain a tiling of a μ -co- ε part of the space with tiles over which the w -average of f is almost in $\mathcal{A}_G^w f$. For $r \in \mathbb{R}$, $A \subseteq \mathbb{R}$, and $\varepsilon > 0$, we write $r \in_\varepsilon A$ to mean that $\text{dist}(r, A) < \varepsilon$. Also, for any $U \in [X]_G^{w<\infty}$, we write $\mathcal{A}_G^w f(U)$ to mean $\mathcal{A}_G^w f(x)$ for some/any $x \in U$.

Proposition 9.11. *Suppose that G is μ -nowhere hyperfinite. For every $\varepsilon > 0$, there is a Borel tiling $\mathcal{P} \subseteq [X]_G^{w<\infty}$ with a μ -co- ε domain such that for each $P \in \mathcal{P}$, $A_P^w f \in_\varepsilon \mathcal{A}_G^w f(P)$.*

Proof. Let \mathcal{S} denote the collection of all $U \in [X]_G^{w<\infty}$ satisfying $A_U^w f \in_\varepsilon \mathcal{A}_G^w f(U)$. It remains to show that \mathcal{S} is (G, w) -conic because then Lemma 8.11 gives the desired tiling. To this end, suppose towards a contradiction that some $U \in [X]_G^{w<\infty}$ admits arbitrarily w -large visible neighborhoods V that are not in \mathcal{S} , i.e. $\text{dist}(A_V^w f, \mathcal{A}_G^w f(U)) \geq \varepsilon$. Being visible neighborhoods of U , these V are all contained in a single \leq_G^w -downward cone $C \supseteq U$, e.g. $(\leq_G^w)^x$ for any $x \in \text{Max}_w U$. The compactness of $[-\|f\|_\infty, \|f\|_\infty]$ extracts a sequence (V_n) with all V_n contained in C such that $w(V_n) \rightarrow \infty$ and $\lim_n A_{V_n}^w f$ exists and is outside of $\mathcal{A}_G^w f(U)$, a contradiction. \square

Remark 9.12. For a bounded weight-function, instead of a cocycle, a stronger version of Proposition 9.11 holds: the requirement of μ -nowhere hyperfiniteness is unnecessary and the tiling in the conclusion has a conull domain. This is proven in [MT17, 8.8] and it is due to the fact that in this case of an actual bounded weight-function instead of a cocycle, the analogue of Lemma 8.11 is much easier to prove and has a stronger conclusion.

To illustrate the power of Proposition 9.11, we prove the statement that motivated the definition of $\mathcal{A}_G^w f$. However, we do not use it in our proofs below.

Corollary 9.13. $\mathbb{E}(f|\mathcal{B}_{E_G}) \in \overline{\mathcal{A}_G^w f}$ a.e.

Proof. By subtracting $\mathbb{E}(f|\mathcal{B}_{E_G})$ from f , we may assume that $\mathbb{E}(f|\mathcal{B}_{E_G}) \equiv 0$. For simplicity of the presentation, suppose that G is ergodic (the proof is uniform through the ergodic decomposition of G with respect to μ [Dit92, Theorem 6 of Chapter 2]), so $\mathcal{A}_G^w f$ is constant a.e. Suppose towards a contradiction that 0 is $\Delta > 0$ away from $\mathcal{A}_G^w f$. Because $\mathcal{A}_G^w f$ is an interval, it is contained either in $[\Delta, \infty)$ or $(-\infty, -\Delta]$, and we suppose without loss of generality that it is the former. Take $\varepsilon > 0$ small enough so that $(\Delta - \varepsilon)(1 - \varepsilon) > \frac{2\Delta}{3}$ and so that $\int_Y |f| d\mu < \frac{\Delta}{3}$ for any $Y \subseteq X$ of measure $\leq \varepsilon$. Let \mathcal{P} be a tiling given by Proposition 9.11 for this ε . Putting $F := E(\mathcal{P})$ and $D := \text{dom}(\mathcal{P})$, the local-global bridge lemma gives a contradiction:

$$\begin{aligned} \int_X f d\mu &= \int_{X \setminus D} f d\mu + \int_D f d\mu \\ \left[\mu(X \setminus D) < \varepsilon \right] &\geq -\frac{\Delta}{3} + \int_D f d\mu \\ \left[\text{Lemma 3.2} \right] &= -\frac{\Delta}{3} + \int_D A_F^w f d\mu \\ \left[A_F^w f \in_\varepsilon [\Delta, \infty) \text{ and } D \text{ is } \mu\text{-co-}\varepsilon \right] &\geq -\frac{\Delta}{3} + (\Delta - \varepsilon)(1 - \varepsilon) > \frac{\Delta}{3} > 0. \end{aligned}$$

\square

10. PROOF OF MAIN LEMMA 1.6

This section is entirely devoted to the proof of Main Lemma 1.6. We let X, μ, G, w, f , and ε be as in the statement of the theorem. By subtracting $\mathbb{E}(f|\mathcal{B}_{E_G})$ from f , we assume without loss of generality that $\mathbb{E}(f|\mathcal{B}_{E_G}) \equiv 0$, yet $\|f\|_1 > 0$.

We need to find a G -connected finite Borel equivalence relation F with $A_F^w f \approx_\varepsilon 0$ on a μ -co- ε set. The following allows us to replace finite with w -finite.

Lemma 10.1. *For any G -connected w -finite Borel equivalence relation F and any $\varepsilon' > 0$, there is a G -connected finite Borel subequivalence relation $F' \subseteq F$ such that $\mathcal{A}_{F'}^w f \approx_{\varepsilon'} A_F^w f$ on a μ -co- ε' set.*

Proof. Fix any $\delta > 0$ less than ε' and $\frac{\varepsilon'}{2\|f\|_\infty}$. Let X' be the set of all $x \in X$ such that $w(B_x) < (1 - \delta)w([x]_F)$, where $B_x := \{y \in [x]_F : x <_w y\}$. By Observation 2.1, B_x is finite for each $x \in X$, so for each F -class C , the set $C' := C \cap X'$ is finite; moreover, $w(C') \geq (1 - \delta)w(C)$. However, these C' may not be G -connected and we extend them to G -connected finite sets as follows. Recalling Observation 2.3, we let X'' be the set of all $x \in X$ that belong to the $<_w^*$ -maximum G -connected finite subset of $[x]_F$ containing $X' \cap [x]_F$ (the relation $<_w^*$ is defined right after Observation 2.2). Now for each F -class C , the set $C'' := X'' \cap C$ is finite, G -connected, and $w(C'') \geq (1 - \delta)w(C)$. Finally, note that both X' and X'' are Borel by Luzin–Novikov uniformization, which turns scanning over each F -class a natural number quantifier.

Now define the equivalence relation F' by setting $F'|_{X''} := F|_{X''}$ and $F'|_{X \setminus X''} := \text{Id}_{X \setminus X''}$. For each $x \in X''$, $[x]_{F'} = [x]_F \cap X''$, so $w([x]_{F'}) \geq (1 - \delta)w([x]_F)$, which implies $A_{[x]_{F'}}^w f \approx_{\varepsilon'} A_{[x]_F}^w f$, by Lemma 3.1(b). Moreover, by Lemma 3.2,

$$\mu(X'') = \int_X \mathbb{1}_{X''} d\mu = \int_X A_F^w \mathbb{1}_{X''} d\mu \geq \int_X (1 - \delta) d\mu = 1 - \delta > 1 - \varepsilon',$$

so F' is as desired. \square

Thus, without loss of generality (replacing ε with $\frac{\varepsilon}{2}$), it is enough to find a G -connected w -finite Borel equivalence relation F with $A_F^w f \approx_\varepsilon 0$ on a μ -co- ε set. We call such equivalence relations *good for Main Lemma*.

Assumption 10.2. To simplify notation in the proof, we assume that G is ergodic. In other words, we present the proof restricting to one ergodic component of the ergodic decomposition of G with respect to μ [Dit92, Theorem 6 of Chapter 2]. This makes the map $x \mapsto \mathcal{A}_G^w f(x)$ constant a.e. because it is E_G -invariant; thus, we drop x from the notation. However, the proof goes through without this assumption, but all the constants below that depend on $\mathcal{A}_G^w f$ (e.g. δ in Hypothesis 10.4) would depend on $x \in X$ (although would still be an E_G -invariant functions).

For $\delta \geq 0$, put $I_\delta^- := (-\infty, \delta]$, $I_\delta := (-\delta, \delta)$, and $I_\delta^+ := [\delta, \infty)$.

10.A. Cutting one side of the set of asymptotic averages. Note that taking a quotient of G by a G -connected w -finite Borel equivalence relation F results in a graph G/F with a more restricted set of visible asymptotic averages, i.e.

$$\mathcal{A}_{G/F}^w f := \mathcal{A}_{G/F}^{w/F} (A_F^w f) \subseteq \mathcal{A}_G^w f,$$

where the inclusion follows from Lemma 3.1(a): average of averages is just the average.

Having the ability (Proposition 9.11) to build tilings whose each tile P has $A_P^w f$ arbitrarily close to $\mathcal{A}_G^w f$, it makes sense to try building a G -connected w -finite Borel equivalence relation F such that quotienting out by it shrinks the set of visible asymptotic averages arbitrarily tightly around 0. In fact, a weaker conclusion is enough due to the following.

Lemma 10.3 (Cutting one side is enough). *For $\delta_\varepsilon := \frac{\varepsilon^2}{2(\|f\|_\infty + 1)}$, any w -finite Borel equivalence relation F , and any a sign $\sigma \in \{+, -\}$, if $A_F^w f \in I_{\delta_\varepsilon}^\sigma$ with probability $\leq \delta_\varepsilon$, then $A_F^w f \in I_\varepsilon$ with probability $\geq 1 - \varepsilon$.*

Proof. Fixing an F as in the hypothesis and taking $\sigma := -$ (the proof is symmetric), we suppose that the set $X_{\delta_\varepsilon}^- := \{x \in X : A_F^w f(x) \in I_{\delta_\varepsilon}^-\}$ has measure $\leq \delta_\varepsilon$ and aim to show that $X_\varepsilon := \{x \in X : A_F^w f(x) \in I_\varepsilon\}$ has measure $\geq 1 - \varepsilon$. It is enough to show that $X_\varepsilon^+ := \{x \in X : A_F^w f(x) \in I_\varepsilon^+\}$ has measure $\leq \frac{\varepsilon}{2}$ because $X \subseteq X_{\delta_\varepsilon}^- \cup X_\varepsilon \cup X_\varepsilon^+$, so $\mu(X_\varepsilon) \geq \mu(X \setminus (X_{\delta_\varepsilon}^- \cup X_\varepsilon^+)) \geq 1 - \delta_\varepsilon - \frac{\varepsilon}{2} \geq 1 - \varepsilon$. To this end, using Lemma 3.2, we compute:

$$\begin{aligned} 0 &= \int_X f d\mu = \int_X A_F^w f d\mu = \int_{X_{\delta_\varepsilon}^-} A_F^w f d\mu + \int_{X_\varepsilon^+} A_F^w f d\mu + \int_{X \setminus (X_{\delta_\varepsilon}^- \cup X_\varepsilon^+)} A_F^w f d\mu \\ &\geq -\|f\|_\infty \cdot \mu(X_{\delta_\varepsilon}^-) + \varepsilon \cdot \mu(X_\varepsilon^+) - \delta_\varepsilon \cdot \mu(X \setminus (X_{\delta_\varepsilon}^- \cup X_\varepsilon^+)) \\ &\geq -\|f\|_\infty \cdot \delta_\varepsilon + \varepsilon \cdot \mu(X_\varepsilon^+) - \delta_\varepsilon \\ &= \mu(X_\varepsilon^+) \cdot \varepsilon - \delta_\varepsilon \cdot (\|f\|_\infty + 1), \end{aligned}$$

$$\text{so } \mu(X_\varepsilon^+) \leq \delta_\varepsilon \cdot \frac{\|f\|_\infty + 1}{\varepsilon} = \frac{\varepsilon}{2}. \quad \square$$

Thus, if for every $\delta > 0$, there were a G -connected w -finite Borel equivalence relation F and a sign $\sigma \in \{+, -\}$ with $(\mathcal{A}_{G:F}^w f) \cap I_\delta^\sigma = \emptyset$, then Proposition 9.11 applied to G/F , $\mu_{/F}$, and $A_F^w f$ would yield a G -connected w -finite Borel equivalence relation $F' \supseteq F$ with $A_{F'}^w f \notin I_{2\delta}^\sigma$ on a μ -co- δ set. Taking δ sufficiently small, Lemma 10.3 would imply that $A_{F'}^w f \in I_\varepsilon$ on a μ -co- ε set, so F' is good for Main Lemma. Thus, without loss of generality, we assume the following.

Hypothesis 10.4. There is a $\delta > 0$ such that for any G -connected w -finite Borel equivalence relation F , $\mathcal{A}_{G:F}^w f$ intersects both I_δ^+ and I_δ^- .

We will exploit this hypothesis and the non- μ -hyperfiniteness of G via packed tilings in Subsections 10.B and 10.C.

10.B. Domains of packed tilings. Let F be a G -connected w -finite equivalence relation and let $\pi_F : X \rightarrow X_{/F}$ denote the quotient map.

For $\lambda > 0$, we call $U \in [X_{/F}]_{G/F}^{w/F < \infty}$ λ -central (resp. λ -positive, λ -negative) if $A_U^{w/F}(A_F^w f) \in I_\lambda$ (resp., I_λ^+ , I_λ^-). Note that $A_U^{w/F}(A_F^w f) = A_{\pi_F^{-1}U}^w f$, so the notions of λ -central/negative/positive coincide for U and its pullback $\pi_F^{-1}(U)$. Also, $w_*(\pi_F^{-1}(U)) \geq (w_{/F})_*(U)$ by Observation 2.8.

For $\lambda, L > 0$, let $\mathcal{S}_{/F}(\lambda, L)$ denote the collection of all λ -central $U \in [X_{/F}]_{G/F}^{w/F < \infty}$ with $(w_{/F})_*(U) > L$. We omit writing the subscript $/F$ if $F = \text{Id}_X$.

Lemma 10.5 (Finitizing visibility). *For any $\lambda < \delta$ as in Hypothesis 10.4, $p := \frac{\lambda}{\|f\|_\infty}$, and any tiling $\mathcal{P} \subseteq [X_{/F}]_{G/F}^{w/F < \infty}$ that is p -packed within $\mathcal{S}_{/F}(\lambda, L)$, $(G/F)_{-\text{dom}(\mathcal{P})}$ has finite $w_{/F}$ -visibility.*

Proof. Taking the quotient by F preserves all of our hypotheses; in particular, G/F is μ/F -ergodic and Hypothesis 10.4 is satisfied by G/F , so $\mathcal{A}_{G/F:E(\mathcal{P})}^w f$ is constant and intersects both I_δ^+ and I_δ^- . Also, $\|A_F^w f\|_\infty \leq \|f\|_\infty$, so assume without loss of generality that $F = \text{Id}_X$.

Towards the contrapositive, we suppose that $G' := G_{-\text{dom}(\mathcal{P})}$ does not have finite w -visibility and aim to show that \mathcal{P} is not p -packed within $\mathcal{S}(\lambda, L)$. Let $x \in X \setminus \text{dom}(\mathcal{P})$ be such that $C := (\leq_{G'}^w)^x$ is w -infinite. We use below that for any visible neighborhood $V \subseteq C$ of x , $w_*(V) = w_x(V)$.

By Observation 7.4(a), \mathcal{P} is maximal within $\mathcal{S}(\lambda, L)$, so there is no $V \in \mathcal{S}(\lambda, L)$ that is entirely contained in C . This and the convexity of $\mathcal{A}_{G|C}^w f(x)$ (Lemma 9.6) imply that x cannot have both λ -positive and λ -negative arbitrarily w -large visible neighborhoods $V \subseteq C$. Thus, all w -large enough visible neighborhoods of x within C must have the same λ -sign. For concreteness, suppose that they are all λ -positive.

Because $\mathcal{A}_{G:E(\mathcal{P})}^w f$ intersects I_δ^- and $\lambda < \delta$, there is a λ -negative $E(\mathcal{P})$ -invariant $U \in [X]_G^{w<\infty}$ containing x such that $U_{/E(\mathcal{P})}$ is finite, $w_*(U) \geq L$, and $2\|f\|_\infty \frac{w(x)}{w(U)} < \lambda$. The latter condition ensures, by the intermediate value property (Lemma 9.5), that there is $W \in [C]_G^{<\infty}$ disjoint from U but G -adjacent to U (i.e. $[W, U]_G \neq \emptyset$) such that

$$0 \leq A_{U \cup W}^w f < \lambda.$$

Because $W \subseteq C$, $\max_w W \leq w(x) \leq \max_w U$, so $w_*(U \cup W) \geq w_*(U) \geq L$ and hence, $U \cup W \in \mathcal{S}(\lambda, L)$. Lastly, $U \cup W$ contains only finitely-many tiles from \mathcal{P} , so it is enough to prove that $U \cup W$ is a p -pack over \mathcal{P} , i.e. $w(W) \geq p \cdot w(U)$. To this end, by Lemma 3.1(a),

$$\begin{aligned} w(W) \cdot A_W^w f &= (w(U) + w(W)) \cdot A_{U \cup W}^w f - w(U) \cdot A_U^w f \\ &\geq (w(U) + w(W)) \cdot 0 - w(U) \cdot (-\lambda) = \lambda \cdot w(U). \end{aligned}$$

In particular, $A_W^w f > 0$, so $w(W) \geq \frac{\lambda}{A_W^w f} \cdot w(U) \geq \frac{\lambda}{\|f\|_\infty} \cdot w(U) \geq p \cdot w(U)$. \square

Corollary 10.6. *For any $\lambda < \delta$ as in Hypothesis 10.4, $p := \frac{\lambda}{\|f\|_\infty}$, and any tiling $\mathcal{P} \subseteq [X/F]_{G/F}^{w/F<\infty}$ that is p -packed within $\mathcal{S}_F(\lambda, L)$, $\pi_F^{-1}(\text{dom}(\mathcal{P}))$ is a hyperfinitizing Borel vertex-cut for G ; in particular, $\mu(\text{dom}(\mathcal{P})) \geq \text{hvp}_\mu(G)$.*

Proof. By Lemma 10.5 and Theorem 1.8, $D := \text{dom}(\mathcal{P})$ is a hyperfinitizing vertex-cut for G/F . The hyperfiniteness of $(G/F)_{-D}$ implies that of $G_{-\pi_F^{-1}(D)}$ by [JKL02, 1.3(ii)] because π_F is a Borel reduction $E_G \rightarrow (E_G)/F$. \square

10.C. Iterative tiling and its limsup. To construct a w -finite equivalence relation F that is good for Main Lemma, we first obtain a coherent sequence of approximately saturated tilings (\mathcal{P}_n) that contain G -connected sets of higher and higher w -ratio and become more and more central and more and more packed. We then show that putting together enough finite-many of these \mathcal{P}_n yields a desired F .

Claim 10.7. *For any $\lambda, L > 0$ and any G -connected w -finite Borel equivalence relation F , the collection $\mathcal{S}_F(\lambda, L)$ has finitely based quotients.*

Proof. It is enough to show that $\mathcal{S}_F(\lambda, L)$ is finitely based because for any Borel tiling $\mathcal{P} \subseteq \mathcal{S}_F(\lambda, L)$, the finite basedness of the quotient of $\mathcal{S}_F(\lambda, L)$ by $E(\mathcal{P})$ is implied by the

finite basedness of $\mathcal{S}_{/E(\mathcal{P})}(\lambda, L)$. Working mod F , we assume without loss of generality that $F = \text{Id}_X$. It remains to show that $\mathcal{S}(\lambda, L)$ is finitely based.

Let $A \in \mathcal{S}(\lambda, L)$ and let $B \subseteq A$ be finite. Fixing $x \in \text{Max}_w A$, we have that $w_x(A) = w_*(A) > L$, hence letting $A' \subseteq A$ be a finite G -connected subset containing $B \cup \{x\}$ with $w_x(A \setminus A')$ small enough, we have that $w_*(A') > L$. Moreover, because $A_A^w f \in I_\lambda$ and I_λ is open, making $w_x(A \setminus A')$ possibly even smaller ensures, by Lemma 3.1(b), that $A_{A'}^w f \in I_\lambda$ hence $A' \in \mathcal{S}(\lambda, L)$. \square

Let $\delta \in (0, 1)$ be as in Hypothesis 10.4 and for each $n \geq 0$, put

$$\lambda_n := 3^{-n} \cdot \delta \cdot \varepsilon < \varepsilon$$

$$L_n := 4^n$$

$$p_n := \frac{\lambda_{n+2}}{\|f\|_\infty + \lambda_{n+1}}.$$

Remark 10.8. All we need below is that $\lim_n \lambda_n L_n = \infty$ (this warrants (10.10) below and hence, Claim 10.11), and that (λ_n) decays exponentially to 0 (used in Claim 10.14). The choice of (p_n) is made to yield a contradiction in the proof of Claim 10.15.

We recursively define a coherent sequence $(\mathcal{P}_n)_{n \geq 0}$ of Borel tilings contained in $[X]_G^{w<\infty}$; in particular, the equivalence relations $F_n := \bigcup_{k < n} E(\mathcal{P}_k)$ are increasing, where $F_0 := \text{Id}_X$. For $n \geq 0$, suppose that F_n is defined. By Claim 10.7, Theorem 7.17 applies to $\mathcal{S}_{/F_n}(\lambda_n, L_n)$ with $p := p_n$ and yields a Borel tiling $\mathcal{Q} \subseteq \overline{\mathcal{S}_{/F_n}(\lambda_n, L_n)}$ that is p_n -packed within $\mathcal{S}_{/F_n}(\lambda_n, L_n)$ and approximately saturated within $\mathcal{S}'_{/F_n}(\lambda_n, L_n) := \mathcal{S}_{/F_n}(\lambda_n, L_n) \cap [X_{/F_n}]_{G_{/F_n}}^{<\infty}$ modulo $w_{/F_n}$ -deficient. Putting $\mathcal{P}_n := \{\pi_{F_n}^{-1}(Q) : Q \in \mathcal{Q}\}$ finishes the construction.

By Corollary 5.6, if an $(E_G)_{/F_n}$ -invariant Borel $A_n \subseteq X_{/F_n}$ is $w_{/F_n}$ -deficient, then it is $\mu_{/F_n}$ -null, so $\pi_{F_n}^{-1}(A_n)$ is μ -null. Thus, discarding countably-many E_G -invariant μ -null sets from X , we have that for each $n \geq 0$, $(\mathcal{P}_n)_{/F_n}$ is p_n -packed within $\mathcal{S}_{/F_n}(\lambda_n, L_n)$ and approximately saturated within $\mathcal{S}'_{/F_n}(\lambda_n, L_n)$. In particular, $(\mathcal{P}_n)_{/F_n} \subseteq \overline{\mathcal{S}_{/F_n}(\lambda_n, L_n)}$, which implies by Lemma 2.6 and Lemma 3.1(b) that for each $n \in \mathbb{N}$ and $P \in \mathcal{P}_n$,

$$w_*(P) > L_n \text{ and } A_P^w f \in \overline{I_{\lambda_n}} := [-\lambda_n, \lambda_n]. \quad (10.9)$$

Let $D_n := \text{dom}(\mathcal{P}_n)$ and $D_\infty := \limsup_n D_n := \{x \in X : x \in D_n \text{ for infinitely many } n \in \mathbb{N}\} = \bigcap_N \bigcup_{n \geq N} D_n$. By Corollary 10.6, $\mu(D_n) \geq \text{hvp}_\mu(G)$, and $\text{hvp}_\mu(G) > 0$ by our assumption that G is not μ -hyperfinite. Thus, D_∞ has positive measure by the downward continuity of μ .

In the next subsection, we show that D_∞ is actually conull. Granted this, the proof of Theorem 1.1 is completed as follows. Fix $n \in \mathbb{N}$ large enough so that $D := \bigcup_{k \leq n} D_k$ is μ -co- ε . Then D is F_n -invariant and for each $x \in D$, $[x]_{F_n} \in \mathcal{P}_k$ for some $k \leq n$, so $[x]_{F_n}$ is λ_k -central, and hence ε -central. Thus, F_n is good for Main Lemma.

10.D. The conullness of D_∞ amounts to a packing condition. Suppose towards a contradiction that $X \setminus D_\infty$ has positive measure, hence so does the inner-boundary $\partial_G(D_\infty)$ by the E_G -quasi-invariance of μ .

Putting $F_\infty := \bigcup_n F_n$, we observe that D_∞ is F_∞ -invariant and each F_∞ -class U outside of D_∞ is actually an F_n -class for large enough n ; in particular, U is w -finite. On the other hand, each F_∞ -class within D_∞ is w -infinite by Observation 2.5 because for each $n \in \mathbb{N}$ and

$P \in \mathcal{P}_n$, we have $w_*(P) > L_n$ by (10.9) and $\lim_n L_n = \infty$. Hence, $(D_\infty, X \setminus D_\infty)_G = \bigcup_{N=1}^\infty \vec{H}_N$, where

$$\vec{H}_N := \{(x, y) \in (D_\infty, X \setminus D_\infty)_G : w([x]_{F_N}) \geq w([y]_{F_\infty}) \text{ and } \forall n \geq N [y]_n = [y]_\infty\}.$$

The E_G -quasi-invariance of μ implies that for all large enough $N \geq 1$, the inner boundary $\partial_{\vec{H}_N}(D_\infty)$ has positive measure and moreover, for all $n \geq N$,

$$L_n \geq 8 \text{ and } \frac{2 \cdot \|f\|_\infty}{L_n} < \frac{1}{3} \lambda_n. \quad (10.10)$$

Furthermore, we can choose such an N so that $D'_N := \partial_{\vec{H}_N} D_N \subseteq D_\infty$ has positive measure.

For each $n \geq N$, let $\mathcal{P}'_n := \{P \in \mathcal{P}_n : P \cap D'_N \neq \emptyset\}$. Note that $(\mathcal{P}'_n)_{n \geq N}$ does not stabilize anywhere on $[D'_N]_{F_\infty}$ because $D'_N \subseteq D_\infty$. Thus, it remains to show that $(\mathcal{P}'_n)_{n \geq N}$ is a p -packing sequence for some $p > 0$ because then, Lemma 7.5 implies that $[D'_N]_{F_\infty}$ is w -deficient, hence null (by Corollary 5.6), contradicting $\mu(D'_N) > 0$.

10.E. Verifying the packing condition. In this subsection we show that $(\mathcal{P}'_n)_{n \geq N}$ is $\frac{2}{13}$ -packing, thus completing the proof of Main Lemma. Because each tile in $\mathcal{P}' := \bigcup_n \mathcal{P}'_n$ is equal to an F_∞ -class that meets D'_N , it is enough to fix one such F_∞ -class C and show that $(\mathcal{P}'_n|_C)_{n \geq N}$ is $\frac{2}{13}$ -packing. Fix such a C .

Claim 10.11. *There is a sign $\sigma_C \in \{+, -\}$ such that $A_{[y]_{F_\infty}}^w f \in \mathbb{R}^{\sigma_C}$ for every $y \in \partial_{\vec{H}_N}(D'_N \cap C)$.*

Proof of Claim. Suppose towards a contradiction that there are (possibly equal) points $y_-, y_+ \in \partial_{\vec{H}_N}(D'_N \cap C)$ such that $A_{[y_-]_{F_\infty}}^w f \leq 0 \leq A_{[y_+]_{F_\infty}}^w f$. Let $x_-, x_+ \in D'_N \cap C$ be (possibly equal, even when $y_- \neq y_+$) points such that $(x_-, y_-), (x_+, y_+) \in \vec{H}_N$. Fix $n > N$ large enough so that $P := [x_-]_{F_{n+1}} = [x_+]_{F_{n+1}}$ and such that $x_+ \in D_n$, so $P \in \mathcal{P}_n$.

Put $Q := P_{/F_n}$, and for each $\sigma \in \{-, +\}$, put $x'_\sigma := [x_\sigma]_{F_n}$ and $y'_\sigma := [y_\sigma]_{F_n}$. The definition of \vec{H}_N implies that $y'_\sigma = [y_\sigma]_{F_\infty}$ and

$$w_{/F_n}(x'_-) \geq w_{/F_n}(y'_-) \text{ and } w_{/F_n}(x'_+) \geq w_{/F_n}(y'_+). \quad (10.12)$$

Recall that $\mathcal{Q} := (\mathcal{P}_n)_{/F_n}$ admits a saturating $\mathcal{S}'_{/F_n}(\lambda_n, L_n)$ -approximation (\mathcal{Q}_m) . We now switch to working mod F_n , so we assume without loss of generality that $F_n = \text{Id}_X$ and drop F_n from the notation; in particular, we write f instead of $A_{F_n}^w f$.

We show that if $A_Q^w f \leq 0$ then y'_+ must have been in $\text{dom}(\mathcal{Q})$, and if $A_Q^w f \geq 0$ then y'_- must have been in $\text{dom}(\mathcal{Q})$. Without loss of generality, we suppose that $A_Q^w f \leq 0$ (the argument is symmetric). For each $m \in \mathbb{N}$, putting $Q_m := [x'_+]_{E(\mathcal{Q}_m)}$, observe that $x'_- \in Q = \bigcup_m Q_m$ and that $\lim_m A_{Q_m}^w f = A_Q^w f$ by Lemma 3.1(b) because $\lim_m w(Q \setminus Q_m) = 0$. Thus, for all large enough $k \in \mathbb{N}$ the following holds and we fix such a k :

- (i) $Q_k \in \bigcup_{m \in \mathbb{N}} \mathcal{Q}_m \subseteq \mathcal{S}'(\lambda_n, L_n)$;
- (ii) $x'_- \in Q_k$;
- (iii) $A_{Q_k}^w f \approx_{\frac{\lambda_n}{3}} A_Q^w f$.

Taking $Q'_k := Q_k \cup \{y'_+\}$, (ii) and (10.12) imply that $\max_w Q_k \geq \max_w \{x'_-, x'_+\} \geq \max_w \{y'_-, y'_+\}$, so $\max_w Q'_k = \max_w Q_k$ and hence $w_*(Q'_k) = \frac{w(Q_k) + w(y'_+)}{\max_w(Q_k)} > w_*(Q_k) > L_n$, where the last inequality is due to (i). Furthermore, because $f(y'_+) \geq 0$ and $A_{Q_k}^w f \in I_{\lambda_n}$ (by (i)), we have

$A_{Q'_k}^{\mathbb{W}} f \geq A_{Q_k}^{\mathbb{W}} f > -\lambda_n$. On the other hand, by Lemma 3.1(b), the fact that $\mathfrak{w}_*(Q_k) > L_n$, and (10.10),

$$A_{Q'_k}^{\mathbb{W}} f - A_{Q_k}^{\mathbb{W}} f \leq \frac{2 \cdot \|f\|_{\infty} \cdot \mathfrak{w}(y'_+)}{\mathfrak{w}(Q_k)} \leq \frac{2 \cdot \|f\|_{\infty} \cdot \mathfrak{w}(y'_+)}{L_n \cdot \max_{\mathbb{W}} Q_k} \leq \frac{2 \cdot \|f\|_{\infty}}{L_n} < \frac{\lambda_n}{3},$$

so $A_{Q'_k}^{\mathbb{W}} f < A_{Q_k}^{\mathbb{W}} f + \frac{\lambda_n}{3} \leq \frac{2\lambda_n}{3}$ by (iii). Thus, Q'_k is λ_n -central, and hence $Q'_k \in \mathcal{S}'(\lambda_n, L_n)$, contradicting that (Q_m) is a saturating $\mathcal{S}'(\lambda_n, L_n)$ -approximation for Q . \square

Suppose, for the sake of concreteness, that $\sigma_C = -$ as the proof for $\sigma_C = +$ is analogous: instead of $A_P^{\mathbb{W}} f \leq -2\lambda_{n+1}$ below, we would have $A_P^{\mathbb{W}} f \geq 2\lambda_{n+1}$, and the proof of Claim 10.15 would use the reverse inequalities and opposite signs.

Claim 10.13. $A_P^{\mathbb{W}} f \leq -\frac{2}{3}\lambda_n = -2\lambda_{n+1}$ for each $n \geq N$ and each $P \in \mathcal{P}'_n|_C$.

Proof of Claim. Suppose towards a contradiction that $A_P^{\mathbb{W}} f > -\frac{2}{3}\lambda_n$, so $A_Q^{\mathbb{W}/F_n}(A_{F_n}^{\mathbb{W}} f) = A_P^{\mathbb{W}} f > -\frac{2}{3}\lambda_n$, where $Q := P_{/F_n}$. Like in the proof of Claim 10.11, let x' denote the F_n -class of some $x \in P \cap D'_N$ and let y' denote the F_n -class of some $y \in X \setminus D_{\infty}$ such that $(x, y) \in \vec{H}_N$. By the definition of \vec{H}_N , $y' = [y]_{F_{\infty}}$ and $\mathfrak{w}(x') \geq \mathfrak{w}(y')$. Lastly, our assumption of $\sigma_C = -$ implies that $A_{y'}^{\mathbb{W}} f < 0$.

Recall that $Q := (\mathcal{P}_n)_{/F_n}$ admits a saturating $\mathcal{S}'_{/F_n}(\lambda_n, L_n)$ -approximation (Q_m) . We now switch to working mod F_n , so we assume without loss of generality that $F_n = \text{Id}_X$ and drop F_n from the notation.

For each $m \in \mathbb{N}$, putting $Q_m := [x']_{E(Q_m)}$, observe that $\lim_m A_{Q_m}^{\mathbb{W}} f = A_Q^{\mathbb{W}} f$ by Lemma 3.1(b) because $\lim_m \mathfrak{w}(Q \setminus Q_m) = 0$. Thus, for all large enough $k \in \mathbb{N}$ the following holds and we fix such a k :

- (i) $Q_k \in \bigcup_{m \in \mathbb{N}} Q_m \subseteq \mathcal{S}'(\lambda_n, L_n)$;
- (ii) $A_{Q_k}^{\mathbb{W}} f > -\frac{2}{3}\lambda_n$.

We take $Q'_k := Q_k \cup \{y'\}$, so $\max_{\mathbb{W}} Q_k \geq \mathfrak{w}(x') \geq \mathfrak{w}(y')$, and hence $\max_{\mathbb{W}} Q'_k = \max_{\mathbb{W}} Q_k$, thus $\mathfrak{w}_*(Q'_k) > \mathfrak{w}_*(Q_k) > L_n$, where the last inequality is due to (i). Because $A_{y'}^{\mathbb{W}} f < 0$, we have $A_{Q'_k}^{\mathbb{W}} f < A_{Q_k}^{\mathbb{W}} f < \lambda_n$. On the other hand, by Lemma 3.1(b), the fact that $\mathfrak{w}_*(Q_k) > L_n$, and (10.10),

$$A_{Q_k}^{\mathbb{W}} f - A_{Q'_k}^{\mathbb{W}} f \leq 2\|f\|_{\infty} \frac{\mathfrak{w}(y')}{\mathfrak{w}(Q_k)} \leq 2\|f\|_{\infty} \frac{\mathfrak{w}(y')}{L_n \cdot \max_{\mathbb{W}} Q_k} \leq \frac{2\|f\|_{\infty}}{L_n} < \frac{1}{3}\lambda_n,$$

so $A_{Q'_k}^{\mathbb{W}} f > A_{Q_k}^{\mathbb{W}} f - \frac{1}{3}\lambda_n > -\frac{2}{3}\lambda_n - \frac{1}{3}\lambda_n = -\lambda_n$ by (ii). Thus, Q'_k is also λ_n -central, so $Q'_k \in \mathcal{S}'(\lambda_n, L_n)$, contradicting that (Q_m) is saturating $\mathcal{S}'(\lambda_n, L_n)$ -approximation for Q . \square

Claim 10.14. $A_{[x]_{F_{n+1}}}^{\mathbb{W}} f \leq -2\lambda_{n+1}$ for each $n \geq N$ and $x \in D'_N \cap C$.

Proof of Claim. Because $x \in D'_N \cap C \subseteq D_N$, $P := [x]_{F_{n+1}} \in \mathcal{P}'_k|_C$ for some $k \in \mathbb{N}$ such that $N \leq k \leq n$. By Claim 10.13, $A_P^{\mathbb{W}} f \leq -2\lambda_{k+1} \leq -2\lambda_{n+1}$. \square

Claim 10.15. For each $n \geq N$, each $P \in \mathcal{P}'_{n+1}|_C$ is a $\frac{2}{13}$ -pack over $\bigvee_{m \leq n} \mathcal{P}'_m$.

Proof of Claim. If $U := \text{dom}(\bigvee_{m \leq n} \mathcal{P}'_n) \cap P = \emptyset$, then P is trivially a $\frac{2}{13}$ -pack over $\bigvee_{m \leq n} \mathcal{P}'_n$, so suppose $U \neq \emptyset$. Let $m \leq n$ be the largest number with $\text{dom}(\mathcal{P}'_m) \cap P \neq \emptyset$. Thus, U is F_{m+1} -invariant (hence, a disjoint union of F_{m+1} -classes), so Claim 10.14 implies

$$A_U^w f \leq -2\lambda_{m+1} \quad (10.16)$$

Next, note that $V := (\bigcup_{M=m}^n D_M \cap P) \setminus U$ is a disjoint union of tiles from $\bigvee_{M=m}^n \mathcal{P}_M$, so

$$A_V^w f \leq \lambda_m. \quad (10.17)$$

Lastly, put $W := P \setminus (U \cup V)$. Note that because $P \in \mathcal{P}_{n+1}$, we have $P_{/F_{n+1}} \in \mathcal{S}_{/F_{n+1}}(\lambda_{n+1}, L_{n+1})$, and hence also $P_{/F_m} \in \mathcal{S}_{/F_m}(\lambda_m, L_m)$, so the p_m -packedness of $(\mathcal{P}_m)_{/F_m}$ within $\mathcal{S}_{/F_m}(\lambda_m, L_m)$ implies that P is not a p_m -pack over \mathcal{P}_m . Thus, because $D_m \cap P \subseteq U \cup V$, we have

$$w(W) < p_m \cdot (w(U) + w(V)) = \frac{\lambda_{m+2}}{\|f\|_\infty + \lambda_{m+1}} \cdot (w(U) + w(V)). \quad (10.18)$$

Then

$$\begin{aligned} -\lambda_{n+1} \cdot (w(U) + w(V) + w(W)) &= -\lambda_{n+1} \cdot w(P) \\ &\quad \left[\text{by (10.9)} \right] \leq A_P^w f \cdot w(P) \\ &= A_U^w f \cdot w(U) + A_V^w f \cdot w(V) + A_W^w f \cdot w(W) \\ &\quad \left[\text{by Eqs. (10.16) and (10.17)} \right] \leq -2\lambda_{m+1} \cdot w(U) + \lambda_m \cdot w(V) + \|f\|_\infty \cdot w(W), \end{aligned}$$

so

$$\begin{aligned} (2\lambda_{m+1} - \lambda_{n+1}) \cdot w(U) &\leq (\lambda_m + \lambda_{n+1}) \cdot w(V) + (\|f\|_\infty + \lambda_{n+1}) \cdot w(W) \\ &\quad \left[\text{by (10.18)} \right] \leq (\lambda_m + \lambda_{n+1}) \cdot w(V) + \frac{\|f\|_\infty + \lambda_{n+1}}{\|f\|_\infty + \lambda_{m+1}} \cdot \lambda_{m+2} \cdot (w(U) + w(V)) \\ &\leq (\lambda_m + \lambda_{n+1}) \cdot w(V) + \lambda_{m+2} \cdot (w(U) + w(V)) \end{aligned}$$

Thus, $(2\lambda_{m+1} - \lambda_{n+1} - \lambda_{m+2}) \cdot w(U) \leq (\lambda_m + \lambda_{n+1} + \lambda_{m+2}) \cdot w(V)$. Because $\lambda_{n+1} \leq \lambda_{m+1}$, $\lambda_m = 9\lambda_{m+2}$, and $\lambda_{m+1} = 3\lambda_{m+2}$, we finally get:

$$\frac{w(V)}{w(U)} \geq \frac{2\lambda_{m+1} - \lambda_{n+1} - \lambda_{m+2}}{\lambda_m + \lambda_{n+1} + \lambda_{m+2}} \geq \frac{\lambda_{m+1} - \lambda_{m+2}}{\lambda_m + \lambda_{m+1} + \lambda_{m+2}} = \frac{(3-1)}{(9+3+1)} = \frac{2}{13}. \quad \square$$

Claim 10.15 implies that $(\mathcal{P}'_n|_C)_{n \geq N}$ is a $\frac{2}{13}$ -packing sequence (Definition 7.3), thus completing the proof of Main Lemma 1.6. $\square(1.6)$

REFERENCES

- [AW13] Miklós Abért and Benjamin Weiss, *Bernoulli actions are weakly contained in any free action*, Ergodic Theory Dynam. Systems **33** (2013), no. 2, 323–333.
- [BN15] L. Bowen and A. Nevo, *Amenable equivalence relations and the construction of ergodic averages for group actions*, J. Anal. Math. **126** (2015), 359–388.
- [Buf00] A. I. Bufetov, *Operator ergodic theorems for actions of free semigroups and groups*, Funktsional. Anal. i Prilozhen. **34** (2000), no. 4, 1–17, 96 (Russian, with Russian summary); English transl., Funct. Anal. Appl. **34** (2000), no. 4, 239–251.
- [CFW81] A. Connes, J. Feldman, and B. Weiss, *An amenable equivalence relation is generated by a single transformation*, Ergodic Theory Dynamical Systems **1** (1981), no. 4, 431–450.
- [Dit92] A. Ditzen, *Definable equivalence relations on Polish spaces*, ProQuest LLC, Ann Arbor, MI, 1992. Thesis (Ph.D.)—California Institute of Technology. MR2687873

- [DJK94] R. Dougherty, S. Jackson, and A. S. Kechris, *The Structure of Hyperfinite Borel Equivalence Relations*, Trans. of the Amer. Math. Soc. **341** (1994), no. 1, 193–225.
- [Dow50] Yael Naim Dowker, *A new proof of the general ergodic theorem*, Acta Sci. Math. (Szeged) **12** (1950), 162–166.
- [Dur19] Rick Durrett, *Probability—theory and examples*, Cambridge Series in Statistical and Probabilistic Mathematics, vol. 49, Cambridge University Press, Cambridge, 2019. Fifth edition of [MR1068527].
- [Ele07] Gábor Elek, *The combinatorial cost*, Enseign. Math. (2) **53** (2007), no. 3-4, 225–235.
- [Fel07] Jacob Feldman, *A ratio ergodic theorem for commuting, conservative, invertible transformations with quasi-invariant measure summed over symmetric hypercubes*, Ergodic Theory Dynam. Systems **27** (2007), no. 4, 1135–1142.
- [FM77] J. Feldman and C. C. Moore, *Ergodic equivalence relations and von Neumann algebras, I*, Trans. Amer. Math. Soc. **234** (1977), 289–324.
- [GL09] D. Gaboriau and R. Lyons, *A measurable-group-theoretic solution to von Neumann’s problem*, Invent. Math. **177** (2009), no. 3, 533–540.
- [GR21] Jan Grebík and Václav Rozhoň, *Local Problems on Grids from the Perspective of Distributed Algorithms, Finitary Factors, and Descriptive Combinatorics*, ArXiv e-prints (2021), available at <https://arxiv.org/abs/2103.08394>.
- [HLS14] Hamed Hatami, László Lovász, and Balázs Szegedy, *Limits of locally-globally convergent graph sequences*, Geom. Funct. Anal. **24** (2014), no. 1, 269–296.
- [Hoc13] Michael Hochman, *On the ratio ergodic theorem for group actions*, J. Lond. Math. Soc. (2) **88** (2013), no. 2, 465–482.
- [HN16] T. Hutchcroft and A. Nachmias, *Indistinguishability of trees in uniform spanning forests*, Probability Theory and Related Fields (2016), 1–40.
- [JKL02] S. Jackson, A. S. Kechris, and A. Louveau, *Countable Borel equivalence relations*, Journal of Math. Logic **2** (2002), no. 1, 1–80.
- [Kec95] A. S. Kechris, *Classical Descriptive Set Theory*, Grad. Texts in Math., vol. 156, Springer, 1995.
- [Kec10] A. S. Kechris, *Global Aspects of Ergodic Group Actions*, Mathematical Surveys and Monographs, vol. 160, American Math. Soc., 2010.
- [KM04] A. S. Kechris and B. Miller, *Topics in Orbit Equivalence*, Lecture Notes in Math., vol. 1852, Springer, 2004.
- [Lin01] E. Lindenstrauss, *Pointwise theorems for amenable groups*, Invent. Math. **146** (2001), no. 2, 259–295.
- [Mil08] B. D. Miller, *The existence of measures of a given cocycle, II: Probability measures*, Ergodic Theory and Dynamical Systems **28** (2008), no. 5, 1615–1633.
- [MT17] B. D. Miller and A. Tserunyan, *Edge sliding and ergodic hyperfinite decomposition* (2017), preprint, available at <https://arxiv.org/abs/1704.06019>.
- [FN98] Koji Fujiwara and Amos Nevo, *Maximal and pointwise ergodic theorems for word-hyperbolic groups*, Ergodic Theory Dynam. Systems **18** (1998), no. 4, 843–858.
- [Sch77] Klaus Schmidt, *Cocycles on ergodic transformation groups*, Macmillan Company of India, Ltd., Delhi, 1977. Macmillan Lectures in Mathematics, Vol. 1.
- [Tao15] Terence Tao, *Failure of the L^1 pointwise and maximal ergodic theorems for the free group*, Forum Math. Sigma **3** (2015), Paper No. e27, 19.
- [TZ21] A. Tserunyan and J. Zomback, *A backward ergodic theorem and its forward implications* (2021), preprint, available at <https://arxiv.org/abs/2012.10522>.

(Anush Tserunyan) MATHEMATICS AND STATISTICS DEPARTMENT, MCGILL UNIVERSITY, MONTRÉAL, QC, CANADA

Email address: anush.tserunyan@mcgill.ca