

KOZAI-LIDOV RESONANT BEHAVIOR AMONG ATIRA-CLASS ASTEROIDS

Carlos de la Fuente Marcos¹ and Raúl de la Fuente Marcos²

¹*Universidad Complutense de Madrid*

Ciudad Universitaria, E-28040 Madrid, Spain

²*AEGORA Research Group*

Facultad de Ciencias Matemáticas

Universidad Complutense de Madrid

Ciudad Universitaria, E-28040 Madrid, Spain

Keywords: minor planets, asteroids: individual (Atira, 413563, 418265, 2010 XB₁₁, 2013 TQ₅, 2018 JB₃)

arXiv:1806.00442v1 [astro-ph.EP] 1 Jun 2018

Atira-class asteroids have aphelia greater than 0.718 au and smaller than 0.983 au (Greenstreet et al. 2012); as of 29-May-2018, there are only 18 known Atiras because, like Venus, they can only be properly observed near their greatest elongation. Ribeiro et al. (2016) have found that the orbits of many of them remain relatively stable for at least 1 Myr, but it is unclear what can make them so stable when they cross the orbit of Venus and some may reach, in heliocentric terms, as far as Earth’s perihelion and as close as inside Mercury’s perihelion. The analysis of their orbits shows that many of them have values of the argument of perihelion, ω , close to 0° or 180° . Such an orbital arrangement means that the nodes are located at perihelion and at aphelion (Milani et al. 1989), i.e. for many known Atiras, away from the path of Venus. Another remarkable trend is in the eccentricity–inclination ($e - i$) relationship; Atiras with $e < 0.34$ have $i > 10^\circ$ while those with $e > 0.34$ tend to have lower values of i . Although any clustering in ω could be due to observational bias, the trend in $e - i$ may be signaling a Kozai-Lidov scenario in which both the values of e and i oscillate, alternating high e and i , because the value of the so-called Kozai-Lidov parameter, $\sqrt{1 - e^2} \cos i$ remains constant (Kozai 1962; Lidov 1962). For regular heliocentric ecliptic orbital elements, the Kozai-Lidov parameter measures the evolution of the component of the orbital angular momentum of the object perpendicular to the ecliptic. Within the Kozai-Lidov resonance, many known Atiras would be protected against close encounters with Mercury, Venus, and the Earth–Moon system. Here, we show that the orbits of some Atiras are strongly affected by the Kozai-Lidov resonance.

We focus on the evolution of (163693) Atira ($\omega = 253^\circ$), (413563) 2005 TG₄₅ ($\omega = 230^\circ$) and (418265) 2008 EA₃₂ ($\omega = 182^\circ$) —three well-observed Atiras— and also on 2010 XB₁₁ ($\omega = 202^\circ$), 2013 TQ₅ ($\omega = 247^\circ$), and the most recent addition to the Atira-class, 2018 JB₃ ($\omega = 355^\circ$). Our results are based on the latest orbital solutions available from JPL’s Small-Body Database. Our full N -body simulations have been performed as described by de la Fuente Marcos & de la Fuente Marcos (2012).

Figure 1 shows the short-term evolution of e (top panel) and i (second to top panel). Atira and 2013 TQ₅ exhibit a rather irregular behavior; in sharp contrast, 413563, 418265, 2010 XB₁₁, and 2018 JB₃ show the coupling in $e - i$ characteristic of the Kozai-Lidov mechanism. Such a behavior is often found among near-Earth asteroids (NEAs) with values of their semi-major axes close to that of Earth that move confined between the orbits of Venus and Mars (Michel & Thomas 1996; de la Fuente Marcos & de la Fuente Marcos 2015), but in this case the value of ω also oscillates. We do not find such a libration in ω for the Atiras (not shown). It can hardly be argued that the population of NEAs trapped in the Kozai-Lidov resonance is a good dynamical analogue of many Atiras, but the lack of libration in ω might be used as an argument against such an analogy. Libert & Tsiganis (2009) have shown that within the context of the Kozai-Lidov resonance, ω may circulate in the general reference frame, but it always librates in the Laplace-plane reference frame. Our calculations are referred to the usual ecliptic and mean equinox of reference epoch, not to the invariable plane of the Solar System and this is why the oscillation in ω is not observed. Therefore, what is found is consistent with a Kozai-Lidov resonant state.

In this Note, we have explored the orbital evolution of representative Atira-class asteroids in an attempt to understand better the results discussed in Ribeiro et al. (2016). Our N -body calculations and subsequent analysis show that many known Atiras undergo Kozai-Lidov-type orbital evolution, which translates into enhanced long-term stability (see Figure 1, bottom two panels).

We thank S. J. Aarseth for providing the code used in this research, F. Roig, M. N. De Prá, and S. Deen for comments, and A. I. Gómez de Castro for providing access to computing facilities. This work was partially supported by the Spanish MINECO under grant ESP2015-68908-R. In preparation of this Note, we made use of the NASA Astrophysics Data System and the MPC data server.

REFERENCES

- | | |
|--|--|
| de la Fuente Marcos, C., & de la Fuente Marcos, R. 2012, MNRAS, 427, 728 | Greenstreet, S., Ngo, H., & Gladman, B. 2012, Icarus, 217, 355 |
| de la Fuente Marcos, C., & de la Fuente Marcos, R. 2015, A&A, 580, A109 | Kozai, Y. 1962, AJ, 67, 591 |
| | Libert, A.-S., & Tsiganis, K. 2009, A&A, 493, 677 |
| | Lidov, M. L. 1962, Planet. Space Sci., 9, 719 |




Figure 1. Short-term evolution of the values of the eccentricity (top panel) and inclination (second to top panel) of the nominal orbits (zero instant of time, epoch JDTDB 2458200.5, 23-March-2018) of (163693) Atira (black), (413563) 2005 TG₄₅ (blue), (418265) 2008 EA₃₂ (gold), 2010 XB₁₁ (green), 2013 TQ₅ (orange), and 2018 JB₃ (red). For 413563, 418265, 2010 XB₁₁ and 2018 JB₃, eccentricity and inclination exhibit the interplay characteristic of the Kozai-Lidov resonance. The two additional bottom panels show a longer time-span.

Michel, P., & Thomas, F. 1996, *A&A*, 307, 310
Milani, A., Carpino, M., Hahn, G., & Nobili, A. M. 1989,
Icarus, 78, 212

Ribeiro, A. O., Roig, F., De Prá, M. N., Carvano, J. M., &
DeSouza, S. R. 2016, *MNRAS*, 458, 4471