

A REMARK ON A THEOREM BY C. AMIOT

BERNHARD KELLER

ABSTRACT. C. Amiot has classified the connected triangulated k -categories with finitely many isoclasses of indecomposables satisfying suitable hypotheses. We remark that her proof shows that these triangulated categories are determined by their underlying k -linear categories. We observe that if the connectedness assumption is dropped, the triangulated categories are still determined by their underlying k -categories together with the action of the suspension functor on the set of isoclasses of indecomposables.

1. THE CONNECTED CASE

We refer to [1] for unexplained notation and terminology. Let k be an algebraically closed field and \mathcal{T} a k -linear Hom-finite triangulated category with split idempotents. Recall Theorem 7.2 of [1]:

Theorem 1 (Amiot). *Suppose \mathcal{T} is connected, algebraic, standard and has only finitely many isoclasses of indecomposables. Then there exists a Dynkin quiver Q and a triangle autoequivalence Φ of $\mathcal{D}^b(\mathbf{mod} kQ)$ such that \mathcal{T} is triangle equivalent to the triangulated [6] orbit category $\mathcal{D}^b(\mathbf{mod} kQ)/\Phi$.*

Our aim is to show that the proof of this theorem in [1] actually shows that a given k -linear equivalence $\mathcal{D}^b(\mathbf{mod} kQ)/F \xrightarrow{\sim} \mathcal{T}$, where F is a k -linear autoequivalence, lifts to a triangle equivalence $\mathcal{D}^b(\mathbf{mod} kQ)/\Phi \xrightarrow{\sim} \mathcal{T}$, where Φ is a triangle autoequivalence lifting F . Thus, we obtain the

Corollary 2. *Under the hypotheses of the theorem, the k -linear structure of \mathcal{T} determines its triangulated structure up to triangle equivalence.*

Proof. The facts that \mathcal{T} is connected, standard and has only finitely many isoclasses of indecomposables imply that there is a Dynkin quiver Q , a k -linear autoequivalence F of $\mathcal{D}^b(\mathbf{mod} kQ)$ and a k -linear equivalence

$$G : \mathcal{D}^b(\mathbf{mod} kQ)/F \xrightarrow{\sim} \mathcal{T}.$$

This follows from the work of Riedtmann [7], cf. section 6.1 of [1]. We will show that F lifts to an (algebraic) triangle autoequivalence of $\mathcal{D}^b(\mathbf{mod} kQ)$ and G to an (algebraic) triangle equivalence Γ . We give the details in the case of F which were omitted in [1]. Put $\mathcal{D} = \mathcal{D}^b(\mathbf{mod} kQ)$. Since \mathcal{D} is triangulated, the category $\mathbf{mod} \mathcal{D}$ of finitely presented functors $\mathcal{D}^{op} \rightarrow \mathbf{Mod} k$ is an exact Frobenius category and we have a canonical isomorphism of functors $\underline{\mathbf{mod}} \mathcal{D} \rightarrow \underline{\mathbf{mod}} \mathcal{D}$

$$\Sigma_m^3 \xrightarrow{\sim} \Sigma_{\mathcal{D}},$$

where $\Sigma_{\mathcal{D}} : \mathbf{mod} \mathcal{D} \rightarrow \mathbf{mod} \mathcal{D}$ denotes the functor $\mathbf{mod} \mathcal{D} \rightarrow \mathbf{mod} \mathcal{D}$ induced by Σ and Σ_m is the suspension functor of the stable category $\underline{\mathbf{mod}} \mathcal{D}$, cf. [4, 16.4]. Notice that Σ_m

Date: May 31, 2018.

2010 Mathematics Subject Classification. 18E30.

Key words and phrases. Additively finite triangulated category.

only depends on the underlying k -category of \mathcal{D} . Now if S_U denotes the simple \mathcal{D} -module associated with an indecomposable object U of \mathcal{D} , we have

$$S_{\Sigma U} = \Sigma_{\mathcal{D}} S_U \xrightarrow{\sim} \Sigma_m^3 S_U$$

in the stable category $\underline{\mathbf{mod}} \mathcal{D}$. Since F is a k -linear autoequivalence, the functor it induces in $\underline{\mathbf{mod}} \mathcal{D}$ commutes with Σ_m and we have $S_{F\Sigma U} \cong S_{\Sigma F U}$ in $\underline{\mathbf{mod}} \mathcal{D}$ for each indecomposable U of \mathcal{D} . It follows that if S_U is not zero in $\underline{\mathbf{mod}} \mathcal{D}$, then we have an isomorphism $F\Sigma U \cong \Sigma F U$ in \mathcal{D} . Now S_U is zero in $\underline{\mathbf{mod}} \mathcal{D}$ only if S_U is projective, which happens if and only if the canonical map $P_U \rightarrow S_U$ is an isomorphism, where $P_U = \mathcal{D}(?, U)$ is the projective module associated with U . This is the case only if no arrows arrive at U in the Auslander–Reiten quiver of \mathcal{D} and this happens if and only if no arrows start or arrive at U . The same then holds for the suspensions $\Sigma^n U$, $n \in \mathbb{Z}$. Since we have assumed that \mathcal{T} and hence \mathcal{D} is connected, this case is impossible. Therefore, we have an isomorphism $\Sigma F U \cong F\Sigma U$ for each indecomposable U of \mathcal{D} . It follows that $T = F(kQ)$ is a tilting object of \mathcal{D} . By [5], we can lift T to a kQ -bimodule complex Y which is even unique in the derived category of bimodules if we take the isomorphism $kQ \xrightarrow{\sim} \text{End}_{\mathcal{D}}(F(kQ))$ into account. Since the k -linear functors F and $\Phi = ? \overset{L}{\otimes}_{kQ} Y$ are isomorphic when restricted to $\text{add}(kQ)$, they are isomorphic as k -linear functors by Riedtmann’s knitting argument [7]. Since the triangulated category \mathcal{T} is algebraic, we may assume that it equals the perfect derived category $\text{per } \mathcal{A}$ of a small dg k -category \mathcal{A} . Using Riedtmann’s knitting argument again, it follows from the proof of Theorem 7.2 in [1] that the composition

$$\mathcal{D} \xrightarrow{\pi} \mathcal{D}/\Phi \xrightarrow{G} \mathcal{T} = \text{per } \mathcal{A}$$

lifts to a triangle functor $? \overset{L}{\otimes}_{kQ} X$ for a kQ - \mathcal{A} -bimodule X . Moreover, it is shown there that this composition factors through an algebraic triangle equivalence

$$\Gamma : \mathcal{D}/\Phi \xrightarrow{\sim} \mathcal{T} = \text{per } \mathcal{A}.$$

Since the compositions $\Gamma \circ \pi$ and $G \circ \pi$ are isomorphic as k -linear functors, the functors Γ and G are isomorphic as k -linear functors. ✓

2. THE NON CONNECTED CASE

Let k be an algebraically closed field and \mathcal{T} a k -linear **Hom**-finite triangulated category with split idempotents and finitely many isomorphism classes of indecomposables. We assume that \mathcal{T} is algebraic and standard but possibly non connected.

Assume first that \mathcal{T} is Σ -connected, i.e. that the k -linear orbit category \mathcal{T}/Σ is connected. Then the argument at the beginning of the above proof shows that either \mathcal{T} is connected or \mathcal{T} is k -linearly equivalent to $\mathcal{D}^b(\text{mod } kA_1)/F$ for a k -linear equivalence F of $\mathcal{D}^b(\text{mod } kA_1)$. Clearly F lifts to a triangle autoequivalence, namely a power Σ^N , of the suspension functor of $\mathcal{D}^b(\text{mod } kA_1)$. We may assume $N > 0$ equals the number of isoclasses of indecomposables of \mathcal{T} . Since the underlying k -category of \mathcal{T} is abelian and semi-simple, all triangles of \mathcal{T} split and \mathcal{T} is triangle equivalent to $\mathcal{D}^b(\text{mod } kA_1)/\Sigma^N$.

Let us now drop the Σ -connectedness assumption on \mathcal{T} . Then clearly \mathcal{T} decomposes, as a triangulated category, into finitely many Σ -connected components (the pre-images of the connected components of \mathcal{T}/Σ). Each of these is either connected or triangle equivalent to $\mathcal{D}^b(\text{mod } kA_1)/\Sigma^N$ for some $N > 0$. Thus, the indecomposables of \mathcal{T} either lie in connected components or in Σ -connected non connected components and the triangle equivalence class of the latter is determined by the action of Σ on the isomorphism classes of indecomposables. We obtain the

Corollary 3. \mathcal{T} is determined up to triangle equivalence by its underlying k -category and the action of Σ on the set of isomorphism classes of indecomposables.

We refer to Theorem 6.5 of [3] for an analogous result concerning the Σ -finite triangulated categories \mathcal{T} and to [2] for an application.

I am grateful to Susan Sierra for bringing the question to my attention.

REFERENCES

- [1] Claire Amiot, *On the structure of triangulated categories with finitely many indecomposables*, Bull. Soc. Math. France **135** (2007), no. 3, 435–474.
- [2] Simon Crawford, *Singularity categories of deformations of Kleinian singularities*, arXiv:1610.08430 [math.RA].
- [3] Norihiro Hanihara, *Auslander correspondence for triangulated categories*, arXiv:1805.07585 [math.RT].
- [4] Alex Heller, *Stable homotopy categories*, Bull. Amer. Math. Soc. **74** (1968), 28–63.
- [5] Bernhard Keller, *Bimodule complexes via strong homotopy actions*, Algebr. Represent. Theory **3** (2000), no. 4, 357–376, Special issue dedicated to Klaus Roggenkamp on the occasion of his 60th birthday.
- [6] ———, *On triangulated orbit categories*, Doc. Math. **10** (2005), 551–581.
- [7] Christine Riedtmann, *Algebren, Darstellungsköcher, Überlagerungen und zurück*, Comment. Math. Helv. **55** (1980), no. 2, 199–224.

UNIVERSITÉ PARIS DIDEROT – PARIS 7, UFR DE MATHÉMATIQUES, INSTITUT DE MATHÉMATIQUES DE JUSSIEU–PRG, UMR 7586 DU CNRS, CASE 7012, BÂTIMENT CHEVALERET, 75205 PARIS CEDEX 13, FRANCE

E-mail address: `bernhard.keller@imj-prg.fr`