

**$C_0$ -SEMIGROUPS OF 2-ISOMETRIES AND DIRICHLET SPACES**

EVA A. GALLARDO-GUTIÉRREZ AND JONATHAN R. PARTINGTON

ABSTRACT. In the context of a theorem of Richter, we establish a similarity between  $C_0$ -semigroups of analytic 2-isometries  $\{T(t)\}_{t \geq 0}$  acting on a Hilbert space  $\mathcal{H}$  and the multiplication operator semigroup  $\{M_{\phi_t}\}_{t \geq 0}$  induced by  $\phi_t(s) = \exp(-st)$  for  $s$  in the right-half plane  $\mathbb{C}_+$  acting boundedly on weighted Dirichlet spaces on  $\mathbb{C}_+$ . As a consequence, we derive a connection with the right shift semigroup  $\{S_t\}_{t \geq 0}$

$$S_t f(x) = \begin{cases} 0 & \text{if } 0 \leq x \leq t, \\ f(x-t) & \text{if } x > t, \end{cases}$$

acting on a weighted Lebesgue space on the half line  $\mathbb{R}_+$  and address some applications regarding the study of the invariant subspaces of  $C_0$ -semigroups of analytic 2-isometries.

## 1. INTRODUCTION

The concept of a 2-isometry was introduced by Agler in the early eighties (cf. [1]); this is related to notions due to J. W. Helton (see [8] and [9]) and characterized in terms of their extension properties (see [2]). Recall that a bounded linear operator  $T$  on a separable, infinite dimensional complex Hilbert space  $\mathcal{H}$  is called a *2-isometry* if it satisfies

$$T^{*2}T^2 - 2T^*T + I = 0,$$

where  $I$  denotes the identity operator. In addition, such operators are called *analytic* if no nonzero vector is in the range of every power of  $T$ . It turns out that  $M_z$ , i.e. the multiplication operator by  $z$ , acting on the classical Dirichlet space, is a cyclic analytic 2-isometry. But, moreover, in [15] (see also [14]) Richter proved that any cyclic analytic 2-isometry is unitarily equivalent to  $M_z$  acting on a generalized Dirichlet space  $D(\mu)$ .

More precisely, let  $\mu$  be a finite non-negative Borel measure on the unit circle  $\mathbb{T}$  and  $D(\mu)$  the *generalized Dirichlet space* associated to  $\mu$ , that is, the Hilbert space consisting of analytic functions on the unit disc  $\mathbb{D}$  such that the integral

$$\int_{\mathbb{D}} |f'(z)|^2 \left( \int_{|\xi|=1} \frac{1-|z|^2}{|\xi-z|^2} d\mu(\xi) \right) \frac{dm(z)}{\pi}$$

---

*Date:* June 2016, Revised February 2017.

*1991 Mathematics Subject Classification.* Primary 47B38.

*Key words and phrases.* 2-isometries, right-shift semigroups, Dirichlet space.

The authors are partially supported by Plan Nacional I+D grant no. MTM2013-42105-P and MTM2016-77710-P.

is finite (here  $dm(z)$  denotes the Lebesgue area measure in  $\mathbb{D}$ ). Note that if  $\mu = 0$ , the space  $D(\mu)$  is defined to be the classical Hardy space  $H^2$  and for non-zero, finite, non-negative Borel measures  $\mu$  on  $\mathbb{T}$ , the space  $D(\mu)$  is contained in the Hardy space (see [7, Chapter 7]). Then Richter's Theorem reads as follows:

**Theorem (Richter).** *Let  $T$  be a bounded linear operator on an infinite dimensional complex Hilbert space  $\mathcal{H}$ . Then the following condition are equivalent:*

- (i)  *$T$  is an analytic 2-isometry with  $\dim \text{Ker } T^* = 1$ ,*
- (ii)  *$T$  is unitarily equivalent to  $(M_z, D(\mu))$  for some finite non-negative Borel measure on  $\mathbb{T}$ , where*

$$\|f\|_{D(\mu)}^2 = \|f\|_{H^2}^2 + \int_{\mathbb{D}} |f'(z)|^2 \left( \int_{|\xi|=1} \frac{1-|z|^2}{|\xi-z|^2} d\mu(\xi) \right) \frac{dm(z)}{\pi}.$$

One of the main applications of Richter's Theorem concerns the study of the invariant subspaces for the multiplication operator  $M_z$  in the spaces  $D(\mu)$  and its relationship with the classical Beurling Theorem for the Hardy space  $H^2$  (see [3]). For instance, regarding the Dirichlet space  $D = D\left(\frac{|d\xi|}{2\pi}\right)$ , Richter and Sundberg [16] proved that any closed, invariant subspace  $\mathcal{M}$  under  $M_z$  satisfies that  $\dim \mathcal{M} \ominus z\mathcal{M} = 1$ . Moreover, if  $\varphi \in \mathcal{M} \ominus z\mathcal{M}$  with  $\|\varphi\|_D = 1$ , then  $|\varphi(z)| \leq 1$  for  $|z| \leq 1$  and  $\mathcal{M} = \varphi D(m_\varphi)$ , where  $dm_\varphi$  is the measure on  $\mathbb{T}$  given by  $dm_\varphi(\xi) = |\varphi(\xi)|^2 \frac{|d\xi|}{2\pi}$ . For general  $D(\mu)$  spaces, an analogous result holds. We refer the reader to Chapters 7 and 8 in the recent monograph "A primer on the Dirichlet space" [7] for more on the subject.

Motivated by the Beurling-Lax Theorem and the work carried out by Richter, the aim of this work is taking further the study of the 2-isometries and considering  $C_0$ -semigroups of 2-isometric operators. In particular, we will establish a similarity between  $C_0$ -semigroups of analytic 2-isometries  $\{T(t)\}_{t \geq 0}$  acting on a Hilbert space  $\mathcal{H}$  and the multiplication operator semigroup  $\{M_{\phi_t}\}_{t \geq 0}$  induced by  $\phi_t(s) = \exp(-st)$  for  $s$  in the right-half plane  $\mathbb{C}_+$  acting boundedly on weighted Dirichlet spaces  $\tilde{D}_{\mathbb{C}_+}(\nu)$  on  $\mathbb{C}_+$  (see Definition 2.3). As a consequence, by means of the Laplace transform, we derive a connection with the right shift semigroup  $\{S_t\}_{t \geq 0}$

$$S_t f(x) = \begin{cases} 0 & \text{if } 0 \leq t \leq x, \\ f(x-t) & \text{if } x > t, \end{cases}$$

acting on a weighted Lebesgue space on the half line  $\mathbb{R}_+$ . Finally, some applications regarding the study of the invariant subspaces of  $C_0$ -semigroups of analytic 2-isometries are also discussed in Section 3.

## 2. $C_0$ -SEMIGROUPS OF ANALYTIC 2-ISOMETRIES

First, we introduce some basic concepts and terminology regarding  $C_0$ -semigroups of bounded linear operators. For more on this topic, we refer the reader to the Engel–Nagel monograph [6].

A  $C_0$ -semigroup  $\{T(t)\}_{t \geq 0}$  of operators on a Hilbert space  $\mathcal{H}$  is a family of bounded linear operators on  $\mathcal{H}$  satisfying the functional equation

$$\begin{cases} T(t+s) = T(t)T(s) & \text{for all } t, s \geq 0, \\ T(0) = I, \end{cases}$$

and such that  $T(t) \rightarrow I$  in the strong operator topology as  $t \rightarrow 0^+$ . Given a  $C_0$ -semigroup  $\{T(t)\}_{t \geq 0}$ , there exists a closed and densely defined linear operator  $A$  that determines the semigroup uniquely, called the generator of  $\{T(t)\}_{t \geq 0}$ , defined by means of

$$Ax := \lim_{t \rightarrow 0^+} \frac{T(t)x - x}{t},$$

where the domain  $D(A)$  of  $A$  consists of all  $x \in \mathcal{H}$  for which this limit exists (see [6, Chapter II], for instance). Although the generator is, in general, an unbounded operator, it plays an important role in the study of a  $C_0$ -semigroup, reflecting many of its properties.

However, if 1 is in the resolvent of  $A$ , that is, in the set

$$\rho(A) = \{\lambda \in \mathbb{C} : (A - \lambda I) : D(A) \subset \mathcal{H} \rightarrow \mathcal{H} \text{ is bijective}\},$$

then  $(A - I)^{-1}$  is a bounded operator on  $\mathcal{H}$  by the Closed Graph Theorem, and the Cayley transform of  $A$  defined by

$$V := (A + I)(A - I)^{-1}$$

is a bounded operator on  $\mathcal{H}$ , since  $V - I = 2(A - I)^{-1}$ . Therefore  $V$  determines the semigroup uniquely, since  $A$  does. This operator is called the *cogenerator* of the  $C_0$ -semigroup  $\{T(t)\}_{t \geq 0}$ . Observe that 1 is not an eigenvalue of  $V$ .

Recall that if  $A$  is a closed operator, then the *spectral bound*  $s(A)$  of  $A$  is defined by

$$s(A) := \sup\{\operatorname{Re} \lambda : \lambda \in \sigma(A)\},$$

where  $\sigma(A) = \mathbb{C} \setminus \rho(A)$  is the spectrum of  $A$ , and in case that  $A$  is the generator of a  $C_0$ -semigroup, then  $s(A)$  is always dominated by the *growth bound of the semigroup*, that is,

$$-\infty \leq s(A) \leq w_0 = \inf \left\{ w \in \mathbb{R} : \begin{array}{l} \text{there exists } M_w \geq 1 \text{ such that} \\ \|T(t)\| \leq M_w e^{wt} \text{ for all } t \geq 0 \end{array} \right\}.$$

Indeed, if  $r(T(t))$  denotes the spectral radius of  $T(t)$ , it follows that  $w_0 = \frac{1}{t} \log r(T(t))$  for each  $t > 0$  (see [6, Section 2, Chapter IV], for instance). The following lemma will be useful in the context of our main result later.

**Lemma 2.1.** *Let  $\{T(t)\}_{t \geq 0}$  be a  $C_0$ -semigroup on a separable, infinite dimensional complex Hilbert space  $\mathcal{H}$  consisting of 2-isometries and  $A$  its generator. Then  $1 \in \rho(A)$  and therefore, the cogenerator  $V$  of  $\{T(t)\}_{t \geq 0}$  is well-defined.*

*Proof.* By induction it follows that, for any  $n \geq 1$  and  $t \geq 0$ ,  $T(t)$  satisfies

$$T(t)^{*n}T(t)^n - nT(t)^{*}T(t) + (n-1)I = 0,$$

and so

$$\|T(t)^n x\|^2 = n\|T(t)x\|^2 - (n-1)\|x\|^2$$

for  $x \in \mathcal{H}$ . From here, it follows that  $\|T(t)^n\| \leq C\sqrt{n}$ , where  $C$  is a constant independent of  $n$ , and therefore the spectral radius  $r(T(t)) \leq 1$  for any  $t$ . Therefore,  $s(A) \leq 0$ ; and therefore  $1 \in \rho(A)$ .  $\square$

The next result consists of a particular instance of [11, Theorem 1], where  $C_0$ -semigroups of hypercontractions are considered. We state it for  $C_0$ -semigroups of 2-isometries and include its proof for the sake of completeness.

**Proposition 2.2.** *Let  $\{T(t)\}_{t \geq 0}$  be a  $C_0$ -semigroup on a separable, infinite dimensional complex Hilbert space  $\mathcal{H}$ . Then the following conditions are equivalent:*

- (i)  $T(t)$  is a 2-isometry for every  $t \geq 0$ .
- (ii) The mapping  $t \in \mathbb{R}_+ \mapsto \|T(t)x\|^2$  is affine for each  $x \in \mathcal{H}$ .
- (iii)  $\operatorname{Re}\langle A^2 y, y \rangle + \|Ay\|^2 = 0$  ( $y \in \mathcal{D}(A^2)$ ).
- (iv) The cogenerator  $V$  of  $\{T(t)\}_{t \geq 0}$  exists and is a 2-isometry.

*Proof.* (i)  $\iff$  (ii): If each  $T(t)$  is a 2-isometry, then for  $t \geq 0$  and  $\tau > 0$  we have

$$\langle T(t+2\tau)x, T(t+2\tau)x \rangle - 2\langle T(t+\tau)x, T(t+\tau)x \rangle + \langle T(t)x, T(t)x \rangle = 0,$$

so that

$$(1) \quad \|T(t+\tau)x\|^2 = \frac{1}{2}(\|T(t)x\|^2 + \|T(t+2\tau)x\|^2).$$

Since  $t \in \mathbb{R}_+ \rightarrow \|T(t)x\|^2$  is continuous, the mapping is affine.

Conversely, taking  $t = 0$  we see that (1) implies that  $T(\tau)$  is a 2-isometry.

(ii)  $\iff$  (iii): For  $t > 0$  we calculate the second derivative of the function  $g : t \mapsto \|T(t)y\|^2$  for  $y \in \mathcal{D}(A^2)$ . We have

$$\begin{aligned} g''(t) &= \frac{d^2}{dt^2} \langle T(t)y, T(t)y \rangle \\ &= \langle A^2 T(t)y, T(t)y \rangle + 2\langle AT(t)y, AT(t)y \rangle + \langle T(t)y, A^2 T(t)y \rangle. \end{aligned}$$

For  $g$  affine,  $g''$  is zero, and Condition (iii) follows on letting  $t \rightarrow 0$ . Conversely, Condition (iii) implies Condition (ii) for  $y \in \mathcal{D}(A^2)$ , and hence for all  $y$  by density.

(iii)  $\iff$  (iv): We calculate

$$\langle (I - 2V^*V + V^{*2}V^2)x, x \rangle$$

for  $x = (A - I)^2 y$  (note that  $(A - I)^{-2} : H \rightarrow H$  is defined everywhere and has dense range). We obtain

$$\begin{aligned} & \langle (A - I)^2 y, (A - I)^2 y \rangle - 2 \langle (A^2 - I)y, (A^2 - I)y \rangle + \langle (A + I)^2 y, (A + I)^2 y \rangle \\ &= 4 \langle A^2 y, y \rangle + 8 \langle Ay, Ay \rangle + 4 \langle y, A^2 y \rangle. \end{aligned}$$

Thus  $V$  is a 2-isometry if and only if Condition (iii) holds.  $\square$

Before stating the main result of the section, let us introduce the following definition.

**Definition 2.3.** Let  $\nu$  be a finite positive Borel measure supported on the imaginary axis. The Dirichlet space  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu)$  is defined as the space of analytic functions  $F$  on right half-plane  $\mathbb{C}_+$  such that

$$\|F\|^2 = |F(1)|^2 + \frac{1}{\pi} \int_{\mathbb{C}_+} |F'(s)|^2 \left( x + \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x}{x^2 + (y - \tau)^2} d\nu(\tau) \right) dx dy < \infty$$

where  $s = x + iy$ .

The spaces  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu)$  arise, in a natural way, when we analyze  $C_0$ -semigroups of analytic 2-isometries in Hilbert spaces, as it is stated in our main result:

**Theorem 2.4.** Let  $\{T(t)\}_{t \geq 0}$  be a  $C_0$ -semigroup on a separable, infinite dimensional complex Hilbert space  $\mathcal{H}$  consisting of analytic 2-isometries for every  $t > 0$  such that

$$(2) \quad \dim \bigcap_{t > 0} \ker (T^*(t) - e^{-t} I) = 1.$$

Then there exists a finite positive Borel measure  $\nu$  supported on the imaginary axis such that  $\{T(t)\}_{t \geq 0}$  is similar to the semigroup of multiplication operators induced by  $\exp(-ts)$  acting on the space  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu)$ . Moreover, if the multiplication operators induced by  $\exp(-ts)$  act continuously for every  $t > 0$  on a Dirichlet space  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\tilde{\nu})$  where  $\tilde{\nu}$  is a finite positive Borel measure supported on the imaginary axis, then the corresponding semigroup consists of analytic 2-isometries and satisfies (2).

Before proceeding further, let us remark that our main result yields similarity for the semigroup  $\{T(t)\}_{t \geq 0}$  because of the definition of the norm in  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu)$ . In addition, as we shall see later, condition (2) is a way of expressing the property that  $\dim \ker V^* = 1$ , where  $V$  is the cogenerator of the semigroup  $\{T(t)\}_{t \geq 0}$ .

In order to prove Theorem 2.4, we need the following auxiliary results.

**Proposition 2.5.** Let  $\{T(t)\}_{t \geq 0}$  be a  $C_0$ -semigroup on a separable, infinite dimensional complex Hilbert space  $\mathcal{H}$  consisting of analytic 2-isometries. Then the cogenerator  $V$  is an analytic 2-isometry.

*Proof.* First, we observe that  $V$  is well-defined by Lemma 2.1 and, it is a 2-isometry by Proposition 2.2. So, we are required to show that  $V$  is analytic.

The Wold Decomposition Theorem for 2-isometries (see [12], for instance), yields that  $V$  can be decomposed as  $V = S \oplus U$  with respect to  $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$ , where  $U$  is the unitary part on  $\mathcal{H}_2 = \bigcap_n V^n \mathcal{H}$  and  $S$  is an analytic 2-isometry. We will show that  $U = 0$ .

Let us assume, on the contrary, that  $U \neq 0$ .

First, we observe that since 1 is not an eigenvalue of  $V$ , the generator  $A$  of the semigroup  $\{T(t)\}_{t \geq 0}$  may be expressed as the (possibly) unbounded operator

$$(V + I)(V - I)^{-1}.$$

Moreover, since  $T(t)$  commutes with  $(A - I)^{-1}$  and hence with  $V$ , it holds that  $\mathcal{H}_2$  is invariant under  $T(t)$  for every  $t \geq 0$ . In addition, the generator  $B$  of the restricted semigroup  $\{T(t)|_{\mathcal{H}_2}\}_{t \geq 0}$  is the restriction of  $A$  to the  $D(A) \cap \mathcal{H}_2$  (see [6, Ch. 2, Sec. 2], for instance); and the cogenerator is  $U$ .

Now, taking into account the fact that  $U$  is unitary, one deduces that  $B$  is skew-adjoint (i. e.,  $B^* = -B$ ). Then the restriction of  $T(t)$  to  $\mathcal{H}_2$  is unitary for every  $t \geq 0$  and, therefore, every vector in  $\mathcal{H}^2$  is in the range of (powers of)  $T(t)$ . Since  $T(t)$  is analytic, it follows that  $\mathcal{H}^2 = \{0\}$ , a contradiction. Hence,  $U = 0$  and the proof is completed.  $\square$

**Lemma 2.6.** *Let  $\{T(t)\}_{t \geq 0}$  be a  $C_0$ -semigroup on a separable, infinite dimensional complex Hilbert space  $\mathcal{H}$  and  $A$  its generator. The following conditions are equivalent:*

- (1)  $Ax_0 = -x_0$  for some  $x_0 \in D(A)$ .
- (2)  $T(t)x_0 = e^{-t}x_0$  for all  $t \geq 0$  and  $x_0 \in D(A)$ .

*In addition, if  $1 \in \rho(A)$  and  $V$  is the cogenerator, any of the previous conditions is equivalent to*

- (3)  $Vx_0 = 0$  for some  $x_0 \in D(A)$ .

Note that the equivalence between (1) and (2) in Lemma 2.6 just follows from the relationship between the eigenspaces of  $A$  and the semigroup  $\{T(t)\}_{t \geq 0}$ , that is,

$$\text{Ker}(\lambda I - A) = \bigcap_{t \geq 0} \text{Ker}(e^{\lambda t} - T(t)),$$

with  $\lambda \in \mathbb{C}$  (see [6, Corollary 3.8, Section IV], for instance). The last statement follows from the definition of  $V$ .

We are now in position to prove Theorem 2.4.

*Proof of Theorem 2.4* Assume that  $\{T(t)\}_{t > 0}$  consists of analytic 2-isometries. Let  $V$  denote its cogenerator; this is well-defined by Lemma 2.1, and it is an analytic 2-isometry by Proposition 2.5.

In addition, the hypotheses  $\dim \bigcap_{t > 0} \text{ker}(T^*(t) - e^{-t}I) = 1$  along with Lemma 2.6 applied to the adjoint semigroup  $\{T^*(t)\}_{t \geq 0}$ , yields that  $\dim \text{Ker } V^* = 1$ .

By means of Richter's Theorem, it follows that  $V$  is similar to  $M_z$  acting on the space  $D(\mu)$  for some finite non-negative Borel measure  $\mu$  on  $\mathbb{T}$  considered with the equivalent

norm

$$\begin{aligned}
 \|f\|_{D(\mu)}^2 &\approx |f(0)|^2 + \int_{\mathbb{D}} |f'(z)|^2 \left( \int_{|\xi|=1} \frac{1-|z|^2}{|\xi-z|^2} d\mu(\xi) \right) \frac{dm(z)}{\pi} \\
 (3) \quad &= |f(0)|^2 + \int_{\mathbb{D}} |f'(z)|^2 P_\mu(z) \frac{dm(z)}{\pi}.
 \end{aligned}$$

Observe that the similarity is the price paid when we consider the equivalent norm. Hence, for any  $t \geq 0$ , it follows that  $T(t)$  is unitarily equivalent to the multiplication operator induced by  $\exp(-t(1+z)/(1-z))$  on  $D(\mu)$ . Now, we migrate to the right half-plane  $\mathbb{C}_+ = \{\operatorname{Re} s > 0\}$  applying the change of variables  $s = (1+z)/(1-z)$ , or  $z = (s-1)/(s+1)$ .

First, we observe that

$$P_\mu \left( \frac{s-1}{s+1} \right) = \int_{|\xi|=1} \frac{1 - \left| \frac{s-1}{s+1} \right|^2}{\left| \xi - \frac{s-1}{s+1} \right|^2} d\mu(\xi) \quad (s \in \mathbb{C}_+)$$

is a positive harmonic function in  $\mathbb{C}_+$ ; so there exists a non-negative constant  $\rho$  and a finite positive Borel measure  $\nu$  supported on the imaginary axis such that

$$(4) \quad P_\mu \left( \frac{s-1}{s+1} \right) = \rho x + \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x}{x^2 + (y-\tau)^2} d\nu(\tau), \quad (s = x + iy)$$

(see [10, Exercise 6, p. 134], for instance).

We can express  $\nu$  in terms of  $\mu$ , since with  $\xi = (u-1)/(u+1)$  for  $u = i\tau \in i\mathbb{R}$ , we have

$$\begin{aligned}
 P_\mu \left( \frac{s-1}{s+1} \right) &= \mu(1) \frac{|s+1|^2 - |s-1|^2}{|(s+1) - (s-1)|^2} + \int_{\xi \in \mathbb{T} \setminus \{1\}} \frac{(|s+1|^2 - |s-1|^2)|u+1|^2}{|(u-1)(s+1) - (u+1)(s-1)|^2} d\mu(\xi) \\
 &= \mu(1)x + \int_{\xi \in \mathbb{T} \setminus \{1\}} \frac{x|u+1|^2}{|u-s|^2} d\mu(\xi), \\
 &= \mu(1)x + \int_{\xi \in \mathbb{T} \setminus \{1\}} \frac{x(1+\tau^2)}{x^2 + (y-\tau)^2} d\mu(\xi),
 \end{aligned}$$

where  $s = x + iy \in \mathbb{C}_+$ . So in (4) we have

$$(5) \quad \rho = \mu(1) \quad \text{and} \quad \frac{d\nu(\tau)}{\pi(1+\tau^2)} = d\mu(\xi).$$

Then, upon applying the change of variables  $s = (1+z)/(1-z)$  in (3), we deduce that  $T(t)$  is similar to the multiplication operator induced by  $\exp(-ts)$  acting on the space  $\widehat{D}_{\mathbb{C}_+}(\nu)$  consisting of analytic functions  $F$  on  $\mathbb{C}_+$  such that

$$(6) \quad \frac{1}{\pi} \int_{\mathbb{C}_+} |F'(s)|^2 \left( x + \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{x}{x^2 + (y-\tau)^2} d\nu(\tau) \right) dx dy < \infty,$$

where  $s = x + iy$  and  $F(s) = f(z)$ . This proves the first half of Theorem 2.4.

In order to conclude the proof, let us assume that the multiplication operators induced by  $\exp(-ts)$  act continuously for every  $t > 0$  on a Dirichlet space  $\widehat{D}_{\mathbb{C}_+}(\tilde{\nu})$  where  $\tilde{\nu}$  is a finite positive Borel measure supported on the imaginary axis. Reversing the steps above

and taking into account the fact that (5) defines a measure  $\tilde{\mu}$  on  $\mathbb{T}$ , where  $\tilde{\mu}(1) = \tilde{\nu}(0) = \rho$ , we deduce that the given semigroup is similar to the semigroup of multiplication operators induced by  $\phi_t(z) = \exp(-t(1+z)/(1-z))$  on  $D(\tilde{\mu})$ . Since the cogenerator of such a semigroup is  $M_z$ , which is a 2-isometry, it follows by Proposition 2.2 that  $\{M_{\phi_t}\}_{t \geq 0}$  consists of 2-isometries.

It remains to show that  $M_{\phi_t}$  is analytic for every  $t > 0$ . If not, then there are a  $t_0 > 0$  and a  $F \in \tilde{\mathcal{D}}_{\mathbb{C}_+}(\tilde{\nu})$  such that the function  $s \mapsto e^{nt_0 s} F(s)$  lies in  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\tilde{\nu})$  for  $n = 1, 2, 3, \dots$

In particular,

$$\int_{\mathbb{C}_+} |(F(s)e^{nt_0 s})'|^2 x \, dx \, dy < \infty.$$

Transferring to the disc by letting  $s = (1+z)/(1-z)$  and  $F(s) = f(z)$ , we have

$$\int_{\mathbb{D}} |[f(z) \exp(nt_0(1+z)/(1-z))]|^2 \frac{1-|z|^2}{|1-z|^2} dA(z) < \infty,$$

so that the function  $z \mapsto f(z) \exp(nt_0(1+z)/(1-z))$  lies in the weighted Dirichlet space  $D(\delta_1)$  corresponding to a Dirac measure at 1, and hence in  $H^2(\mathbb{D})$ , by [7, Thm. 7.1.2]. We conclude that  $f$  is identically zero, since no nontrivial  $H^2$  function can be divisible by an arbitrarily large power of a nonconstant inner function. Hence the analyticity is also established. □

**A connection with the right-shift semigroup in weighted  $L^2(\mathbb{R}_+)$ .** Now, by means of the Laplace transform, we will establish a connection of  $C_0$ -semigroups of analytic 2-isometries  $\{T(t)\}_{t \geq 0}$  acting on a Hilbert space  $\mathcal{H}$  and the the right shift semigroup  $\{S_t\}_{t \geq 0}$

$$S_t f(x) = \begin{cases} 0 & \text{if } 0 \leq x \leq t, \\ f(x-t) & \text{if } x > t, \end{cases}$$

acting on a weighted Lebesgue space on the half line  $\mathbb{R}_+$ .

First, let us begin by recalling a result asserting that for each  $\alpha > -1$ , a function  $G$  analytic in  $\mathbb{C}_+$  belongs to the *weighted Bergman space*  $\mathcal{A}_\alpha^2(\mathbb{C}_+)$ , that is, the space consisting of analytic functions on  $\mathbb{C}_+$  for which

$$\|G\|_{\mathcal{A}_\alpha^2(\Pi^+)}^2 = \int_{\mathbb{C}_+} |G(x+iy)|^2 x^\alpha \, dx \, dy < \infty,$$

if and only if it has the form

$$G(s) := \mathcal{L}g(s) = \int_0^\infty e^{-st} g(t) \, dt, \quad s \in \mathbb{C}_+,$$

where  $g$  is a measurable function on  $\mathbb{R}^+$  with

$$\int_0^\infty |g(t)|^2 t^{-1-\alpha} \, dt < \infty.$$



Moreover,

$$\|G\|_{\mathcal{A}_\alpha^2(\mathbb{C}^+)}^2 = \frac{\pi \Gamma(1+\alpha)}{2^\alpha} \int_0^\infty |g(t)|^2 t^{-1-\alpha} dt,$$

(see [4] or [5, Theorem 1], for instance). In other words, the Laplace transform is an isometric isomorphism between  $\mathcal{A}_\alpha^2(\mathbb{C}_+)$  and  $L^2(\mathbb{R}_+, \left(\frac{\pi \Gamma(1+\alpha)}{2^\alpha}\right)^{1/2} t^{-1-\alpha} dt)$ . Hence, by means of a density argument, and taking  $\alpha = 1$ , it follows that for any  $F \in \tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu)$ , there exists  $f \in L^2(\mathbb{R}_+)$  (unique in the usual sense of equivalence classes), such that

$$(i) \quad \mathcal{L}(tf(t)) = F'(s),$$

(ii)

$$\frac{1}{\pi} \int_{\mathbb{C}_+} |F'(s)|^2 x dx dy = \frac{1}{2} \int_0^\infty |f(t)|^2 dt, \quad (s = x + iy),$$

which corresponds to the first sum in (6); and

(iii)

$$\frac{1}{\pi^2} \int_{\mathbb{C}_+} |F'(s)|^2 \frac{x}{x^2 + (y - \tau)^2} dx dy = \frac{1}{2\pi} \int_0^\infty \left| \int_0^t u f(u) e^{-i\tau u} du \right|^2 \frac{dt}{t^2}, \quad (s = x + iy).$$

These three items along with the fact that for any  $t \geq 0$ ,  $T(t)$  is similar to the multiplication operator induced by  $\exp(-ts)$  acting on the space  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu)$  yields, by means of the Laplace transform, that  $\{T(t)\}_{t \geq 0}$  is transformed to the right-shift semigroup  $\{S_t\}_{t \geq 0}$  acting on the Hilbert space  $\mathfrak{H}$  which consists of functions  $f$  defined on  $\mathbb{R}_+$  such that

$$\int_0^\infty |f(t)|^2 dt + \int_0^\infty \int_{-\infty}^\infty \left| \int_0^t f(u) e^{-i\tau u} u du \right|^2 d\nu(\tau) \frac{dt}{t^2} < \infty.$$

### 3. A FINAL REMARK ON INVARIANT SUBSPACES OF $C_0$ -SEMIGROUPS OF ANALYTIC 2-ISOMETRIES

As an application of our main result, we deal with the study of the lattice of the closed invariant subspaces of a  $C_0$ -semigroup  $\{T(t)\}_{t \geq 0}$  of analytic 2-isometries.

Here we shall use the following result from [15, Thm. 7.1] and [16, Thm. 3.2].

**Theorem 3.1.** *Let  $\mathcal{M}$  be a non-zero invariant subspace of  $(M_z, D(\mu))$ . Then  $\mathcal{M} = \phi D_{\mu_\phi}$  where  $\phi \in \mathcal{M} \ominus z\mathcal{M}$  is a multiplier of  $D(\mu)$  and  $d\mu_\phi = |\phi|^2 d\mu$ .*

In the continuous case we have the following result:

**Theorem 3.2.** *Let  $\{T(t)\}_{t \geq 0}$  denote the semigroup of multiplication operators induced by  $\exp(-ts)$  on the space  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu)$ , as in Theorem 2.4, and let  $\mathcal{M}$  be a non-zero closed subspace invariant under all the operators  $T(t)$ . Then there is a function  $\psi \in \mathcal{M}$  such that  $\mathcal{M} = \psi \tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu_\psi)$ .*

*Proof.* If  $\mathcal{M}$  is invariant under the semigroup, then it is also invariant under the cogenerator  $V$ , and after transforming to the disc as in the proof of Theorem 2.4, we may apply Theorem 3.1.

Note that under the equivalence between  $D(\mu)$  and  $\tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu)$ , as detailed in (4) and (5), the subspace  $\phi D_{\mu_\phi}$  maps to a space  $\psi \tilde{\mathcal{D}}_{\mathbb{C}_+}(\nu_\psi)$ , where  $\psi(s) = \phi((s-1)/(s+1))$  and  $d\nu_\psi = |\psi|^2 d\nu$ .  $\square$

#### ACKNOWLEDGEMENTS

The authors thank the referees for a careful reading of the manuscript as well as for their suggestions, which improved the final version of the submitted manuscript.

#### REFERENCES

- [1] J. Agler, *Hypercontractions and subnormality*, J. Operator Theory 13 (1985), no. 2, 203–217.
- [2] J. Agler, *A disconjugacy theorem for Toeplitz operators*, Amer. J. Math. 112 (1990), no. 1, 1–14.
- [3] A. Beurling, *On two problems concerning linear transformations in Hilbert space* Acta Math. 81 (1948), 239–255.
- [4] N. Das and J. R. Partington, *Little Hankel operators on the half-plane*, Integral Equations Operator Theory 20 (1994), no. 3, 306–324.
- [5] P. Duren, E. A. Gallardo-Gutiérrez and A. Montes-Rodríguez, *A Paley–Wiener theorem for Bergman spaces with application to invariant subspaces*, Bull. London Math. Soc. 39 (2007), no. 3, 459–466.
- [6] K. J. Engel and R. Nagel, *One-parameter Semigroups for Linear Evolution Equations*, Graduate Text in Mathematics, vol. 194, Springer–Verlag, New York, 2000.
- [7] O. El-Fallah, K. Kellay, J. Mashregi, and T. Ransford, *A primer on the Dirichlet space*, volume 203 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2014.
- [8] J. W. Helton, *Operators with a representation as multiplication by  $\times$  on a Sobolev space*, Hilbert space operators and operator algebras (Proc. Internat. Conf., Tihany, 1970), pp. 279–287, North-Holland, Amsterdam, 1972.
- [9] J. W. Helton, *Infinite dimensional Jordan operators and Sturm–Liouville conjugate point theory*, Trans. Amer. Math. Soc. 170 (1972), 305–331.
- [10] K. Hoffman, *Banach spaces of analytic functions*, Dover Publication, Inc., 1988.
- [11] B. Jacob, J. R. Partington, S. Pott, and A. Wynn,  *$\beta$ -admissibility of observation and control operators for hypercontractive semigroups*, J. Evol. Equ. 18 (2018), no. 1, 153–170.
- [12] A. Olofsson, *A von Neumann–Wold decomposition of two-isometries*, Acta Sci. Math. (Szeged) 79 (2004), 715–726.
- [13] B. Sz-Nagy and C. Foias, *Harmonic Analysis of Operators on Hilbert Space*, North-Holland Publishing Co., 1970.
- [14] S. Richter, *Invariant subspaces of the Dirichlet shift*, J. Reine Angew. Math. 386 (1988), 205–220.
- [15] S. Richter, *A representation theorem for cyclic analytic two-isometries*, Trans. Amer. Math. Soc. 328 (1991), no. 1, 325–349.
- [16] S. Richter and C. Sundberg, *Multipliers and invariant subspaces in the Dirichlet space*, J. Operator Theory, 28 (1992), no. 1, 167–186.

UNIVERSIDAD COMPLUTENSE DE MADRID E ICMAT  
DEPARTAMENTO DE ANÁLISIS MATEMÁTICO,  
FACULTAD DE CIENCIAS MATEMÁTICAS,  
PLAZA DE CIENCIAS 3  
28040, MADRID (SPAIN)

*E-mail address:* `eva.gallardo@mat.ucm.es`

SCHOOL OF MATHEMATICS,  
UNIVERSITY OF LEEDS,  
LEEDS LS2 9JT, U.K.

*E-mail address:* `J.R.Partington@leeds.ac.uk`