

Carleman estimates for a magnetohydrodynamics system and application to inverse source problems

Xinchi HUANG^{*1,2} and Masahiro YAMAMOTO^{1,3,4}

Abstract

In this article, we consider a linearized magnetohydrodynamics system for incompressible flow in a three-dimensional bounded domain. We first prove two kinds of Carleman estimates. This is done by combining the Carleman estimates for the parabolic and the elliptic equations. Then we apply the Carleman estimates to prove Hölder type stability results for some inverse source problems.

1 Introduction

The magnetohydrodynamics equations (the MHD equations in short) are concerned with the magnetic properties of electrically conducting fluids such as plasmas, liquid metals and salt water.

Before the formulation, we introduce the following notations. Let $\Omega \subset \mathbb{R}^3$ be a bounded domain with smooth boundary, and let $n = n(x)$ be the outward unit normal vector to $\partial\Omega$ at x , and $\partial_n \varphi = \nabla \varphi \cdot n$. Moreover \cdot^T denotes the transpose of matrices or vectors, and let $\partial_t = \frac{\partial}{\partial t}$, $\partial_j = \frac{\partial}{\partial x_j}$, $j = 1, 2, 3$, $\Delta = \sum_{j=1}^3 \partial_j^2$, $\nabla = (\partial_1, \partial_2, \partial_3)^T$, $\nabla_{x,t} = (\nabla, \partial_t)^T$. We also use the following notations:

$$(w \cdot \nabla)v := \left(\sum_{j=1}^3 w_j \partial_j v_1, \sum_{j=1}^3 w_j \partial_j v_2, \sum_{j=1}^3 w_j \partial_j v_3 \right)^T,$$

$$v \times w := (v_2 w_3 - v_3 w_2, v_3 w_1 - v_1 w_3, v_1 w_2 - v_2 w_1)^T,$$

2010 Mathematics Subject Classification. Primary: 35R30, 35Q35.

Key words and phrases. magnetohydrodynamics, Carleman estimates, inverse source problems, stability.

* Corresponding author.

¹ Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo, 153-8914, Japan.

² JSPS Postdoctoral Fellowships for research in Japan.

³ Honorary Member of Academy of Romanian Scientists, Ilfov, nr. 3, Bucuresti, Romania.

⁴ Correspondence member of Accademia Peloritana dei Pericolanti, Palazzo Università, Piazza S. Pugliatti 1 98122 Messina, Italy.

$$\operatorname{div} w := \sum_{j=1}^3 \partial_j w_j, \quad \operatorname{rot} w := (\partial_2 w_3 - \partial_3 w_2, \partial_3 w_1 - \partial_1 w_3, \partial_1 w_2 - \partial_2 w_1)^T$$

for vectors $v = (v_1, v_2, v_3)^T$, $w = (w_1, w_2, w_3)^T$.

By using the divergence-free condition of the magnetic field, we simplify Maxwell's equations into a second-order parabolic equation. Then for the velocity field $u = (u_1, u_2, u_3)^T$, the pressure p and the magnetic field intensity $H = (H_1, H_2, H_3)^T$, the MHD equations are described as follows:

$$\begin{cases} \partial_t u - \nu \Delta u + (u \cdot \nabla) u - \mu \operatorname{rot} H \times H + \nabla p = F, \\ \partial_t H - \frac{1}{\sigma \mu} \Delta H - \operatorname{rot}(u \times H) = 0, \\ \operatorname{div} u = 0, \quad \operatorname{div} H = 0. \end{cases}$$

Here ν describes the viscosity of the fluids, while σ and μ denote the electric conductivity and the magnetic permeability, respectively. We assume that they are all positive constants. For more details about the derivation of the MHD equations, we refer to Li and Qin [22] for example. Moreover, $\operatorname{div} H = 0$ is used for the proof of one of our Carleman estimates (Theorem 2.3).

There are several papers concerning the direct problems of the MHD equations. Since our main purpose of this article is the inverse problems, we do not explain more on the direct problems and we refer to Ladyzhenskaya and Solonnikov [23], as a classical work, Li [21] and the references therein.

The purpose of this article is the stability for inverse source problems, more precisely, the determination of the spatially varying factor in the source term. Our main strategy is based on a method by Bukhgeim-Klibanov [4] and Klibanov [20], applying so-called Carleman estimates.

Carleman estimate is an L^2 -weighted estimate for the solution to a partial differential equation with large parameter, which is nowadays known as a powerful tool in treating the uniqueness and stability estimates for inverse problems. Actually, the Carleman estimate itself and its applications to inverse problems have been intensively studied for a variety of partial differential equations in mathematical physics including, for example,

- Transport equations: Cannarsa, Floridia and Yamamoto [6], Cannarsa, Floridia, Gölgeleyen and Yamamoto [7];
- Hyperbolic equations: Bellassoued and Yamamoto [5], Imanuvilov and Yamamoto [18];
- Parabolic equations: Imanuvilov and Yamamoto [17], Yamamoto [24];
- Lamé equation and the Navier-Stokes equations: Bellassoued, Imanuvilov and Yamamoto [2, 3], Choulli, Imanuvilov, Puel and Yamamoto [8].

To the authors' best knowledge, there are few papers on Carleman estimates for the MHD equations. For the MHD equations, Havârneanu, Popa and Sritharan [12, 13] proved a Carleman estimate and established the exact controllability. In their Carleman estimate, the observation of the first-order spatial derivative of the source F is necessary, and so it is not suitably designed for inverse problems. Besides, Huang [10] established Carleman estimates for the MHD equations and investigate inverse coefficient problems by the measurements of velocity field, magnetic field and pressure on some sub-boundary. However, some additional data on the derivative of pressure are imposed due to the technical proof. In this article, we apply the recent idea of Huang, Imanuvilov, Yamamoto [11] and Imanuvilov, Lorenzi, Yamamoto [15] to eliminate the additional data and propose simplified proofs for the Carleman estimates as well as the stability estimates for the inverse source problems. Furthermore, under some assumptions on the source term we discuss also the inverse source problem without the knowledge of the pressure, that is, inverse source problem 2 in the following context.

Now we formulate our inverse problems and state the main results. For simplicity, we set $\kappa = \frac{1}{\sigma\mu} > 0$ and we consider the following linearized MHD equations:

$$\begin{cases} \partial_t u - \nu \Delta u + (A^{(1)} \cdot \nabla)u + (u \cdot \nabla)A^{(2)} + L_1 H + \nabla p = F(x, t) & \text{in } \Omega \times (0, T), \\ \partial_t H - \kappa \Delta H + (A^{(3)} \cdot \nabla)H + (H \cdot \nabla)A^{(4)} + L_2 u = 0 & \text{in } \Omega \times (0, T), \\ \operatorname{div} u = 0, \quad \operatorname{div} H = 0 & \text{in } \Omega \times (0, T), \end{cases} \quad (1.1)$$

where the coupling operators L_1, L_2 are defined by

$$\begin{cases} L_1 H := (C^{(1)} \cdot \nabla)H + (H \cdot \nabla)C^{(2)} + \nabla(C^{(3)} \cdot H), \\ L_2 u := (u \cdot \nabla)C^{(4)} + (C^{(5)} \cdot \nabla)u, \end{cases} \quad (1.2)$$

and the vector coefficients $A^{(j)}, C^{(k)}$, $j = 1, 2, 3, 4$, $k = 1, 2, 3, 4, 5$ are supposed to be sufficiently smooth. Henceforth we arbitrarily fix $0 < t_0 < T$, $0 < \delta < \min\{t_0, T - t_0\}$ and we denote

$$I := (t_0 - \delta, t_0 + \delta), \quad Q := \Omega \times I.$$

Moreover, let $H^k(\Omega)$, etc., denote usual Sobolev spaces (e.g., Adams and Fournier [1]). Let $\gamma_0 \in (\mathbb{N} \cup \{0\})$, $\gamma = (\gamma_1, \gamma_2, \gamma_3) \in (\mathbb{N} \cup \{0\})^3$, $\partial_x^\gamma = \partial_1^{\gamma_1} \partial_2^{\gamma_2} \partial_3^{\gamma_3}$ and $|\gamma| = \gamma_1 + \gamma_2 + \gamma_3$. Then we set

$$\begin{cases} W^{k, \infty}(D) := \{w; \partial_t^{\gamma_0} w, \partial_x^\gamma w \in L^\infty(D), \gamma_0 \leq k, |\gamma| \leq k\}, \quad k \in \mathbb{N} \\ H^{k, \ell}(D) := \{w; \partial_t^{\gamma_0} w, \partial_x^\gamma w \in L^2(D), \gamma_0 \leq \ell, |\gamma| \leq k\}, \quad k, \ell \in \mathbb{N} \cup \{0\} \end{cases}$$

for any sub-domain $D \subset Q$. If there is no confusion, we do not distinguish $L^2(\Omega)$ with $L^2(\Omega; \mathbb{R}^3)$ and we use $L^2(\Omega)$ for both meanings.

Henceforth, depending on different cases, we consider some of the following

conditions:

$$|\partial_t^k F(x, t)| \leq C|F(x, t_0)|, \quad (x, t) \in Q, \quad k = 0, 1. \quad (1.3)$$

$$|\partial_t^k \operatorname{rot} F(x, t)| \leq C|\operatorname{rot} F(x, t_0)|, \quad (x, t) \in Q, \quad k = 0, 1. \quad (1.4)$$

$$|\partial_t^k \operatorname{rot} F(x, t)| \leq C(|\nabla F(x, t_0)| + |F(x, t_0)|), \quad (x, t) \in Q, \quad k = 0, 1, 2. \quad (1.5)$$

$$|\operatorname{div} F(x, t_0)| = 0, \quad x \in \Omega. \quad (1.6)$$

We are ready to state the different conditional stability estimates under the conditions (1.3), (1.4) and (1.5)–(1.6), respectively. Our first inverse source problem can be described as follows:

Inverse source problem 1 :

Let F satisfy (1.3). For arbitrarily given sub-boundary $\Gamma \subset \partial\Omega$, and arbitrarily given sub-domain $\Omega_0 \subset \Omega \cup \Gamma$, determine the source term $F(\cdot, t_0)$ in Ω_0 by the observation data of $\{(u, \nabla u, p, H, \nabla H)|_{\Gamma \times I}, u(\cdot, t_0)|_\Omega, p(\cdot, t_0)|_\Omega\}$.

More precisely, by introducing an admissible set of the unknown function F :

$$\mathcal{F}_{M,1} = \left\{ F \in H^1(0, T; L^2(\Omega)); \|F\|_{H^1(0, T; L^2(\Omega))} \leq M, \text{ (1.3) is satisfied} \right\}$$

with arbitrarily given constant $M > 0$ and the norm of the data:

$$\begin{aligned} D_1 := & \|u\|_{H^2(I; L^2(\Gamma))} + \|\nabla u\|_{H^1(I; L^2(\Gamma))} + \|H\|_{H^2(I; L^2(\Gamma))} + \|\nabla H\|_{H^1(I; L^2(\Gamma))} \\ & + \|p\|_{H^1(I; H^{\frac{1}{2}}(\Gamma))} + \|u(\cdot, t_0)\|_{H^2(\Omega)} + \|\nabla p(\cdot, t_0)\|_{L^2(\Omega)}, \end{aligned}$$

we have the following stability estimate of Hölder type for the inverse source problem.

Theorem 1.1. *Assume that $F \in \mathcal{F}_{M,1}$ and a solution to (1.1) satisfies $(u, p, H) \in H^{2,2}(Q) \times H^{1,1}(Q) \times H^{2,2}(Q)$ with*

$$\|u\|_{H^{1,2}(Q)} + \|\nabla u\|_{H^{0,1}(Q)} + \|H\|_{H^{1,1}(Q)} + \|\nabla H\|_{H^{0,1}(Q)} + \|p\|_{H^{0,1}(Q)} \leq M.$$

Then there exist constants $C > 0$ and $\theta \in (0, 1)$ such that

$$\|F(\cdot, t_0)\|_{L^2(\Omega_0)} \leq C(D_1^\theta + D_1).$$

Remark 1. In the above theorem and the following theorems in this section, the constants C and θ depend on M , δ , Ω , the coefficients and also the choices of t_0 , Ω_0 and Γ . In particular, as one can see from the proof, the constant C goes to infinity and the constant θ goes to zero as t_0 goes to zero, or Ω_0 tends to Ω , or Γ tends to an empty set. However, we mention that the arbitrary choice of Ω_0 enables us to prove the corresponding uniqueness results of determining $F(\cdot, t_0)$ in Ω immediately by contradiction.

Remark 2. According to the condition (1.3), one can readily see that Theorem 1.1

also implies the estimate of the source term in $\Omega_0 \times I$ as follows:

$$\|F\|_{L^2(\Omega_0 \times I)} \leq C(D_1^\theta + D_1).$$

In Theorem 1.1, the data on the pressure p is needed. However, sometimes it is not easy to measure the pressure. Thus, we discuss also the second inverse source problem as follows:

Inverse source problem 2 :

Let F satisfy (1.5) and (1.6). For arbitrarily given sub-boundary $\Gamma \subset \partial\Omega$, and arbitrarily given sub-domain $\Omega_0 \subset \Omega \cup \Gamma$, determine the source term $F(\cdot, t_0)$ in Ω_0 by the observation data of $\{(u, \nabla u, \nabla \text{rot } u, H, \nabla H, \nabla \text{rot } H)|_{\Gamma \times I}, u(\cdot, t_0)|_\Omega, H(\cdot, t_0)|_\Omega\}$.

By introducing the admissible set:

$$\mathcal{F}_{M,2} = \left\{ F \in H^2(0, T; H^2(\Omega)); F(\cdot, t_0)|_\Gamma = 0, \nabla F(\cdot, t_0)|_\Gamma = 0, \|\nabla F(\cdot, t_0)\|_{L^2(\partial\Omega)} + \|F(\cdot, t_0)\|_{L^2(\partial\Omega)} \leq M, \text{ (1.5) and (1.6) are satisfied} \right\}$$

with arbitrarily given constant $M > 0$ and the norm of the data:

$$\begin{aligned} D_2 := & \|\nabla \text{rot } u\|_{H^2(I; L^2(\Gamma))} + \|\nabla \text{rot } H\|_{H^2(I; L^2(\Gamma))} + \|\nabla u\|_{H^3(I; L^2(\Gamma))} \\ & + \|\nabla H\|_{H^3(I; L^2(\Gamma))} + \|u\|_{H^2(I; L^2(\Gamma))} + \|H\|_{H^2(I; L^2(\Gamma))} \\ & + \|u(\cdot, t_0)\|_{H^4(\Omega)} + \|H(\cdot, t_0)\|_{H^3(\Omega)}, \end{aligned}$$

we have the following stability estimate.

Theorem 1.2. Assume that $F \in \mathcal{F}_{M,2}$ and $(u, p, H) \in H^{2,3}(Q) \times H^{2,2}(Q) \times H^{2,3}(Q)$ is a solution to (1.1) satisfying $\text{rot } u, \text{rot } H \in H^{2,3}(Q)$ and

$$\begin{aligned} & \|u\|_{H^3(I; H^2(\Omega))} + \|H\|_{H^3(I; H^2(\Omega))} + \|\nabla \text{rot } u\|_{H^2(I; L^2(\partial\Omega))} \\ & + \|\nabla \text{rot } H\|_{H^2(I; L^2(\partial\Omega))} + \|\nabla u\|_{H^3(I; L^2(\partial\Omega))} + \|\nabla H\|_{H^3(I; L^2(\partial\Omega))} \\ & + \|u\|_{H^2(I; L^2(\partial\Omega))} + \|H\|_{H^2(I; L^2(\partial\Omega))} \leq M. \end{aligned}$$

Then there exist constants $C > 0$ and $\theta \in (0, 1)$ such that

$$\|F(\cdot, t_0)\|_{H^1(\Omega_0)} \leq C(D_2^\theta + D_2).$$

Remark 3. In Theorem 1.2, we can actually determine additionally the source term in the second equation of (1.1) at the same time. That is, let the second equation be substituted by

$$\partial_t H - \kappa \Delta H + (A^{(3)} \cdot \nabla) H + (H \cdot \nabla) A^{(4)} + L_2 u = G(x, t).$$

Then under suitable assumptions on F and G , we can prove the stability estimate:

$$\|F(\cdot, t_0)\|_{H^1(\Omega_0)} + \|G(\cdot, t_0)\|_{H^1(\Omega_0)} \leq C(D_2^\theta + D_2).$$

Remark 4. In contrast to Theorem 1.1, here we need not any data of p since we assume the condition (1.6). Thus, Theorem 1.2 indicates that one can recover a transverse field source, that is, a source whose divergence is zero, even if we do not measure the pressure p . Indeed, according to the Helmholtz decomposition:

$$F(x, t) = \nabla \tilde{p}(x, t) + \tilde{F}(x, t) \quad (1.7)$$

with a real-valued function \tilde{p} and an \mathbb{R}^3 -valued function \tilde{F} satisfying $\operatorname{div} \tilde{F} = 0$, we can rewrite the first equation in (1.1) by

$$\partial_t u - \nu \Delta u + (A^{(1)} \cdot \nabla)u + (u \cdot \nabla)A^{(2)} + L_1 H + \nabla(p - \tilde{p}) = \tilde{F}(x, t) \quad \text{in } Q.$$

Therefore, there are no hopes of determining the longitudinal field component $\nabla \tilde{p}$ as long as we do not have the knowledge of ∇p and we can determine at most the information of the transverse field component \tilde{F} .

As we see from the above remark, we can identify only the transverse field component of the source provided that the pressure p is not measured. In the next theorem, we consider the case that F is not necessarily a transverse field, that is, we do not assume the condition (1.6), and we determine the information of the transverse field component of the source.

Note that the decomposition (1.7) is not unique. For example, we could replace $\nabla \tilde{p}$ by $\nabla(\tilde{p} - q)$ and \tilde{F} by $\tilde{F} + \nabla q$ for any q satisfying $\Delta q = 0$. Thus, it is not a suitable problem to recover \tilde{F} itself. Instead, we show the next theorem identifying $\operatorname{rot} \tilde{F} = \operatorname{rot} F$ within the following admissible set:

$$\mathcal{F}_{M,3} = \left\{ F \in H^1(0, T; L^2(\Omega)); \operatorname{rot} F \in H^1(0, T; L^2(\Omega)), \text{ (1.4) is satisfied} \right\}$$

and under the norm of the data:

$$\begin{aligned} D_3 := & \|\nabla \operatorname{rot} u\|_{H^1(I; L^2(\Gamma))} + \|\nabla \operatorname{rot} H\|_{H^1(I; L^2(\Gamma))} + \|\nabla u\|_{H^2(I; L^2(\Gamma))} \\ & + \|\nabla H\|_{H^2(I; L^2(\Gamma))} + \|u\|_{H^1(I; L^2(\Gamma))} + \|H\|_{H^1(I; L^2(\Gamma))} + \|u(\cdot, t_0)\|_{H^3(\Omega)}. \end{aligned}$$

Theorem 1.3. Assume that $F \in \mathcal{F}_{M,3}$ and $(u, p, H) \in H^{2,2}(Q) \times H^{2,1}(Q) \times H^{2,2}(Q)$ is a solution to (1.1) satisfying $\operatorname{rot} u, \operatorname{rot} H \in H^{2,2}(Q)$ and

$$\begin{aligned} & \|u\|_{H^2(I; H^2(\Omega))} + \|H\|_{H^2(I; H^2(\Omega))} + \|\nabla \operatorname{rot} u\|_{H^1(I; L^2(\partial\Omega))} \\ & + \|\nabla \operatorname{rot} H\|_{H^1(I; L^2(\partial\Omega))} + \|\nabla u\|_{H^2(I; L^2(\partial\Omega))} + \|\nabla H\|_{H^2(I; L^2(\partial\Omega))} \\ & + \|u\|_{H^1(I; L^2(\partial\Omega))} + \|H\|_{H^1(I; L^2(\partial\Omega))} \leq M. \end{aligned}$$

Then there exist constants $C > 0$ and $\theta \in (0, 1)$ such that

$$\|\operatorname{rot} F(\cdot, t_0)\|_{L^2(\Omega_0)} \leq C(D_3^\theta + D_3).$$

Remark 5. Since the unknown source F appears only in the first equation of (1.1),

we can take advantage of the coupling between u and H , and we manage to remove the measurement of $H(\cdot, t_0)$ in the data.

Remark 6. Under some suitable boundary conditions on $\tilde{F}(\cdot, t_0)$ (see e.g., [25]), the decomposition (1.7) at $t = t_0$ can be uniquely determined. Hence one can recover the transverse field component $\tilde{F}(\cdot, t_0)$ in Ω_0 by using $\text{rot } \tilde{F}(\cdot, t_0) = \text{rot } F(\cdot, t_0)$ in Ω_0 .

We end up this section with two examples which justify the conditions (1.4) and (1.5).

Example 1. We assume that the temporal varying factor of the force F is independent of the position x , that is, we let

$$F(x, t) = r(t)f(x), \quad x \in \Omega, \quad 0 < t < T$$

where $r \in C^2[0, T]$ is a given real-valued function and $f = (f_1, f_2, f_3)^T$, $f_j \in C^1(\overline{\Omega})$, $j = 1, 2, 3$. Then (1.4) and (1.5) can be justified as long as

$$r(t_0) \neq 0.$$

Actually, we have

$$|\partial_t^k \text{rot } F(x, t)| = \left| \frac{d^k r}{dt^k}(t) \text{rot } f(x) \right| \leq \|r\|_{C^2[0, T]} |\text{rot } f(x)| = \frac{\|r\|_{C^2[0, T]}}{|r(t_0)|} |\text{rot } F(x, t_0)|,$$

for all $(x, t) \in Q$ and $k = 0, 1, 2$, which implies (1.4). By noting that

$$|\text{rot } F(x, t_0)| \leq C |\nabla F(x, t_0)|, \quad x \in \Omega,$$

we immediately find (1.5). Moreover, (1.6) holds true if we further assume $\text{div } f = 0$.

Example 2. We consider a more general case that the temporal varying factor of the force depends also on the position x , which can be modeled by a matrix $R(x, t)$:

$$F(x, t) = R(x, t)f(x), \quad x \in \Omega, \quad 0 < t < T,$$

where $R = (r_{ij})_{1 \leq i, j \leq 3}$ is a given matrix-valued function and $f = (f_1, f_2, f_3)^T$ is a vector-valued function. Here we assume that each entry r_{ij} and each component f_j are smooth enough, for example, $r_{ij} \in C^2([0, T]; C^1(\overline{\Omega}))$ for $i, j = 1, 2, 3$ and $f_j \in C^1(\overline{\Omega})$ for $j = 1, 2, 3$. Then (1.5) is satisfied if

$$|\det R(x, t_0)| > 0, \quad x \in \overline{\Omega}.$$

In fact, by direct calculations, we have

$$|\partial_t^k \text{rot } F(x, t)| \leq C (|\nabla f(x)| + |f(x)|), \quad (x, t) \in Q, \quad k = 0, 1, 2. \quad (1.8)$$

On the other hand, we have

$$\partial_\ell F(x, t) = \partial_\ell R(x, t)f(x) + R(x, t)\partial_\ell f(x), \quad (x, t) \in Q, \quad \ell = 1, 2, 3.$$

Since $|\det R(\cdot, t_0)| > 0$ on $\overline{\Omega}$, we find that $R(\cdot, t_0)$ is invertible and the norm of its inverse $R^{-1}(\cdot, t_0)$ is bounded from above. Thus, we rewrite the above equality at $t = t_0$ to obtain

$$\partial_\ell f(x) = R^{-1}(x, t_0)\partial_\ell F(x, t_0) - R^{-1}(x, t_0)\partial_\ell R(x, t_0)f(x), \quad x \in \Omega, \quad \ell = 1, 2, 3,$$

which implies

$$|\partial_\ell f(x)| \leq C(|\partial_\ell F(x, t_0)| + |f(x)|), \quad x \in \Omega, \quad \ell = 1, 2, 3. \quad (1.9)$$

Therefore, we obtain (1.5) by combining (1.8) and (1.9) with the following inequality

$$|f(x)| = |R^{-1}(x, t_0)F(x, t_0)| \leq C|F(x, t_0)|, \quad x \in \Omega.$$

We remark that in Example 2, we do not know a convenient sufficient condition for (1.4) in terms of R and f .

The rest of this article is organized as follows. In Section 2, we show the key Carleman estimates which are stated in Theorems 2.1 and 2.3. In Section 3, we prove Theorems 1.1–1.3 on the basis of these Carleman estimates.

2 Key Carleman estimates

In this section, we consider the following linearized MHD system

$$\begin{cases} \partial_t u - \nu \Delta u + (A^{(1)} \cdot \nabla)u + (u \cdot \nabla)A^{(2)} + L_1 H + \nabla p = F & \text{in } Q, \\ \partial_t H - \kappa \Delta H + (A^{(3)} \cdot \nabla)H + (H \cdot \nabla)A^{(4)} + L_2 u = G & \text{in } Q, \\ \operatorname{div} u = h & \text{in } Q. \end{cases} \quad (2.1)$$

Here L_1, L_2 are defined in (1.2) and the vector coefficients $A^{(j)}, C^{(k)}, j = 1, 2, 3, 4, k = 1, 2, 3, 4, 5$ are smooth enough (e.g., $W^{3,\infty}(Q)$).

The purpose of this section is to establish suitable Carleman estimates for (2.1), which are the key points in proving our stability results of Theorems 1.1–1.3. To this end, we first introduce the weight function for our Carleman estimates.

Let $\Gamma \subset \partial\Omega$ be an arbitrarily fixed non-empty relatively open sub-boundary. We choose a bounded domain $\tilde{\Omega} \subset \mathbb{R}^3 \setminus \overline{\Omega}$ such that $\partial\tilde{\Omega} \cap \partial\Omega = \overline{\Gamma}$. We construct a domain $\Omega_1 \supset \Omega$ by setting

$$\Omega_1 = \Omega \cup \tilde{\Omega} \cup \Gamma.$$

Then

$$\overline{\Gamma} = \partial\Omega \cap \overline{\Omega_1}, \quad \partial\Omega_1 = (\partial\Omega \setminus \Gamma) \cup (\partial\tilde{\Omega} \setminus \Gamma).$$

With this new domain Ω_1 , we can choose a non-empty domain $\omega \subset \Omega_1$ such that $\overline{\omega} \subset \Omega_1 \setminus \overline{\Omega} = \widetilde{\Omega}$. Thus, referring to e.g., Fursikov and Imanuvilov [9] or Imanuvilov [14], we can construct a function $d \in C^2(\overline{\Omega_1})$ such that

$$d = 0 \quad \text{on } \partial\Omega_1, \quad d > 0 \quad \text{in } \Omega_1, \quad |\nabla d| > 0 \quad \text{on } \overline{\Omega_1 \setminus \omega}. \quad (2.2)$$

In particular, $d > 0$ in Ω and $|\nabla d| > 0$ on $\overline{\Omega}$. For arbitrarily given $0 < t_0 < T$ and $0 < \delta < \min\{t_0, T - t_0\}$, we denote

$$I := (t_0 - \delta, t_0 + \delta), \quad Q := \Omega \times I, \quad Q_1 := \Omega_1 \times I.$$

Moreover, we fix a large constant $\lambda > 0$, and for a constant $\beta > 0$ we set

$$\psi(x, t) = d(x) - \beta(t - t_0)^2, \quad \varphi(x, t) = e^{\lambda\psi(x, t)}, \quad (x, t) \in \overline{Q_1}. \quad (2.3)$$

Now we establish our Carleman estimates in the following two cases.

2.1 Carleman estimate with pressure term p

Here we state the first Carleman estimate in which we include the estimate of p .

Theorem 2.1. *Let $F \in L^2(Q)$. Then there exist constants $s_0 > 0$ and $C > 0$ such that*

$$\begin{aligned} & \int_Q \left(\frac{1}{s^2} (|\partial_t u|^2 + |\Delta u|^2) + \frac{1}{s} (|\partial_t H|^2 + |\Delta H|^2) + |\nabla u|^2 + s|\nabla H|^2 \right. \\ & \quad \left. + s^2|u|^2 + s^3|H|^2 + \frac{1}{s}|\nabla p|^2 + s|p|^2 \right) e^{2s\varphi} dx dt \\ & \leq C \int_Q (|F|^2 + |G|^2 + |\nabla_{x,t} h|^2) e^{2s\varphi} dx dt \\ & \quad + Cs^3 \int_{\partial\Omega \times I} (|u|^2 + |H|^2 + |\nabla_{x,t} u|^2 + |\nabla_{x,t} H|^2) e^{2s\varphi} dS dt \\ & \quad + Cs^3 \int_{\Omega} (|u|^2 + |H|^2 + |\nabla u|^2 + |\nabla H|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \end{aligned}$$

for all $s \geq s_0$ and

$$u, H \in H^1(I; H^1(\Omega)) \cap L^2(I; H^2(\Omega)), \quad p \in L^2(I; H_0^1(\Omega))$$

satisfying (2.1).

Henceforth $C > 0$ denotes a generic constant which depends on $\lambda, t_0, \delta, T, \beta, \Omega$ and the coefficients, but is independent of the parameter s . Such uniformity in s is used for deriving the stability results for the inverse problems.

Proof of Theorem 2.1. We divide the proof into three steps.

First step:

At the beginning of the proof, we assume that $p \in L^2(I; C_0^\infty(\Omega))$. Later in the third step, we will take approximations and treat the case of $p \in L^2(I; H_0^1(\Omega))$. In the first step, we prove the following lemma.

Lemma 2.2. *Let $p \in L^2(I; C_0^\infty(\Omega))$ satisfy*

$$\Delta p = f_0 + \sum_{j=1}^3 \partial_j f_j \quad \text{in } Q \quad (2.4)$$

Then there exist constants $\hat{s} > 0$ and $C > 0$ such that

$$\int_Q (|\nabla p|^2 + s^2 |p|^2) e^{2s\varphi} dx dt \leq C \int_Q \left(\frac{1}{s} |f_0|^2 + s \sum_{j=1}^3 |f_j|^2 \right) e^{2s\varphi} dx dt$$

for all $s \geq \hat{s}$.

Proof. By noting that $p \in L^2(I; C_0^\infty(\Omega))$, we can take the zero extension of p from Ω to Ω_1 , which is denoted by the same letter. Then we find that $p(\cdot, t) \in C_0^\infty(\Omega_1)$ for $t \in I$ and (2.4) holds true in Q_1 as long as we take also the zero extensions of f_j , $j = 0, 1, 2, 3$.

Now we apply the H^{-1} Carleman estimate for the elliptic equation (Theorem A.1 in Imanuvilov and Puel [16]) to (2.4) in Ω_1 and we obtain

$$\begin{aligned} & \int_{\Omega_1} (|\nabla p(x, t)|^2 + s^2 |p(x, t)|^2) e^{2se^{\lambda d(x)}} dx \\ & \leq C \int_{\Omega_1} \left(\frac{1}{s} |f_0(x, t)|^2 + s \sum_{j=1}^3 |f_j(x, t)|^2 \right) e^{2se^{\lambda d(x)}} dx + C \int_{\omega} s^2 |p(x, t)|^2 e^{2se^{\lambda d(x)}} dx \end{aligned}$$

for all $s \geq s_1$, $t \in I$ and $\bar{\omega} \subset \Omega_1$. Recalling the construction of the domain Ω_1 , we can choose a non-empty domain ω such that $\bar{\omega} \subset \Omega_1 \setminus \bar{\Omega}$. Since we take the zero extensions for f_j , p outside of Ω , the integral over ω vanishes and we have

$$\begin{aligned} & \int_{\Omega} (|\nabla p(x, t)|^2 + s^2 |p(x, t)|^2) e^{2se^{\lambda d(x)}} dx \\ & \leq C \int_{\Omega} \left(\frac{1}{s} |f_0(x, t)|^2 + s \sum_{j=1}^3 |f_j(x, t)|^2 \right) e^{2se^{\lambda d(x)}} dx \end{aligned} \quad (2.5)$$

for all $s \geq s_1$ and each $t \in I$. Let $\hat{s} := s_1 e^{\lambda \beta \delta^2}$. Then $s \geq \hat{s}$ implies

$$s e^{-\lambda \beta (t-t_0)^2} \geq s e^{-\lambda \beta \delta^2} \geq \hat{s} e^{-\lambda \beta \delta^2} = s_1,$$

for $t \in I$, and thus, replacing s by $se^{-\lambda\beta(t-t_0)^2}$ in (2.5) yields

$$\begin{aligned} & \int_{\Omega} (|\nabla p(x, t)|^2 + s^2 e^{-2\lambda\beta(t-t_0)^2} |p(x, t)|^2) e^{2s\varphi(x, t)} dx \\ & \leq C \int_{\Omega} \left(\frac{1}{s} e^{\lambda\beta(t-t_0)^2} |f_0(x, t)|^2 + s e^{-\lambda\beta(t-t_0)^2} \sum_{j=1}^3 |f_j(x, t)|^2 \right) e^{2s\varphi(x, t)} dx \end{aligned}$$

for all $s \geq \hat{s}$ and each $t \in I$. Here we used $\varphi(x, t) = e^{\lambda d(x)} e^{-\lambda\beta(t-t_0)^2}$, $(x, t) \in Q$. Therefore, we complete the proof of Lemma 2.2 by integrating the above estimate over $t \in I$. \square

Second step:

In this step, we apply the estimate in the first step and the Carleman estimate for parabolic system to prove the theorem in the case where $p \in L^2(I; C_0^\infty(\Omega))$.

We derive the following elliptic equation with respect to p by taking the divergence on both sides of the first equation in (2.1) and using the third equation there:

$$\Delta p = \operatorname{div} F - \partial_t h + \nu \operatorname{div}(\nabla h) - \operatorname{div}((A^{(1)} \cdot \nabla)u) - \operatorname{div}((u \cdot \nabla)A^{(2)}) - \operatorname{div}(L_1 H). \quad (2.6)$$

In particular, we rewrite several terms on the right-hand side as follows

$$\begin{aligned} \nu \operatorname{div}(\nabla h) &= \nu \sum_{j=1}^3 \partial_j (\partial_j h), \\ \operatorname{div}((A^{(1)} \cdot \nabla)u) &= \sum_{j,k=1}^3 \partial_k (A_j^{(1)} \partial_j u_k) = \sum_{j=1}^3 A_j^{(1)} \partial_j \left(\sum_{k=1}^3 \partial_k u_k \right) + \sum_{j,k=1}^3 (\partial_k A_j^{(1)}) \partial_j u_k \\ &= A^{(1)} \cdot \nabla (\operatorname{div} u) + \sum_{j,k=1}^3 \left(\partial_j ((\partial_k A_j^{(1)}) u_k) - (\partial_j \partial_k A_j^{(1)}) u_k \right), \\ \operatorname{div}((u \cdot \nabla)A^{(2)}) + \operatorname{div}(L_1 H) &= \sum_{j=1}^3 \partial_j ((u \cdot \nabla)A_j^{(2)}) + [L_1 H]_j \end{aligned}$$

where $[v]_j$ denotes the j -th component of a vector v , $j = 1, 2, 3$. Besides, we have

$$\operatorname{div} F = \sum_{j=1}^3 \partial_j F_j.$$

Therefore, we are able to represent the right-hand side of (2.6) in the form of $f_0 + \sum_{j=1}^3 \partial_j f_j$ with

$$f_0 = -\partial_t h - (A^{(1)} \cdot \nabla)h + \sum_{k,\ell=1}^3 (\partial_\ell \partial_k A_\ell^{(1)}) u_k,$$

$$f_j = F_j + \nu \partial_j h - \sum_{k=1}^3 (\partial_k A_j^{(1)}) u_k - (u \cdot \nabla) A_j^{(2)} - [L_1 H]_j, \quad j = 1, 2, 3.$$

Thus, by noting

$$\begin{aligned} |f_0| &\leq C (|\partial_t h| + |\nabla h| + |u|), \\ |f_j| &\leq C (|F| + |\nabla h| + |u| + |H| + |\nabla H|), \quad j = 1, 2, 3, \end{aligned}$$

we apply Lemma 2.2 to obtain

$$\begin{aligned} \int_Q \left(\frac{1}{s} |\nabla p|^2 + s |p|^2 \right) e^{2s\varphi} dx dt &\leq C \int_Q (|F|^2 + |\nabla_{x,t} h|^2) e^{2s\varphi} dx dt \\ &\quad + C \int_Q (|u|^2 + |H|^2 + |\nabla H|^2) e^{2s\varphi} dx dt \end{aligned} \quad (2.7)$$

for all $s \geq s_2 := \max\{1, \hat{s}\}$.

On the other hand, we can regard ∇p in the first equation of (2.1) as a non-homogeneous term, and then the first and second equations of (2.1) become a weakly coupling parabolic system with respect to u and H . Hence we can apply the Carleman estimate for the parabolic system (e.g., Yamamoto [24, Theorem 3.2]) and we obtain

$$\begin{aligned} &\int_Q \left(\frac{1}{s^2} (|\partial_t u|^2 + |\Delta u|^2) + |\nabla u|^2 + s^2 |u|^2 \right) e^{2s\varphi} dx dt \\ &\leq C \int_Q \frac{1}{s} |F|^2 e^{2s\varphi} dx dt + C \int_Q \frac{1}{s} (|\nabla p|^2 + |\nabla H|^2 + |H|^2) e^{2s\varphi} dx dt \\ &\quad + C s^2 \int_{\partial\Omega \times I} (|\nabla_{x,t} u|^2 + |u|^2) e^{2s\varphi} dS dt + C s^2 \int_{\Omega} (|\nabla u|^2 + |u|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \end{aligned} \quad (2.8)$$

and

$$\begin{aligned} &\int_Q \left(\frac{1}{s} (|\partial_t H|^2 + |\Delta H|^2) + s |\nabla H|^2 + s^3 |H|^2 \right) e^{2s\varphi} dx dt \\ &\leq C \int_Q |G|^2 e^{2s\varphi} dx dt + C \int_Q (|\nabla u|^2 + |u|^2) e^{2s\varphi} dx dt \\ &\quad + C s^3 \int_{\partial\Omega \times I} (|\nabla_{x,t} H|^2 + |H|^2) e^{2s\varphi} dS dt + C s^3 \int_{\Omega} (|\nabla H|^2 + |H|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \end{aligned} \quad (2.9)$$

for all $s \geq s_3$. Substituting (2.9) into the last term on the right-hand side of (2.7) and then substituting the derived inequality into the second term on the right-hand side of (2.8), we obtain

$$\int_Q \left(\frac{1}{s^2} (|\partial_t u|^2 + |\Delta u|^2) + |\nabla u|^2 + s^2 |u|^2 \right) e^{2s\varphi} dx dt$$

$$\begin{aligned}
&\leq C \int_Q \left(|F|^2 + \frac{1}{s} |G|^2 + |\nabla_{x,t} h|^2 \right) e^{2s\varphi} dx dt \\
&\quad + C \int_Q \left(\frac{1}{s} |\nabla u|^2 + |u|^2 + \frac{1}{s} |\nabla H|^2 + \frac{1}{s} |H|^2 \right) e^{2s\varphi} dx dt \\
&\quad + C s^2 \int_{\partial\Omega \times I} (|\nabla_{x,t} u|^2 + |u|^2 + |\nabla_{x,t} H|^2 + |H|^2) e^{2s\varphi} dS dt \\
&\quad + C s^2 \int_{\Omega} (|\nabla u|^2 + |u|^2 + |\nabla H|^2 + |H|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \tag{2.10}
\end{aligned}$$

for all $s \geq s_4 := \max\{s_2, s_3\}$. Therefore, we combine the inequalities (2.7), (2.9) and (2.10), by choosing $s > 0$ sufficiently large, we can absorb the lower-order terms $\int_Q (|\nabla u|^2 + s|u|^2 + |\nabla H|^2 + |H|^2) e^{2s\varphi} dx dt$ into the left-hand side and this completes the proof in the case of $p \in L^2(I; C_0^\infty(\Omega))$.

Third step:

In this step, we consider the case of $p \in L^2(I; H_0^1(\Omega))$ by a density argument. Recall the density $\overline{C_0^\infty(\Omega)} = H_0^1(\Omega)$ by the norm of $H^1(\Omega)$ (e.g., Adams and Fournier[1]). This yields the existence of an approximating sequence $p_n \in L^2(I; C_0^\infty(\Omega))$, $n \in \mathbb{N}$ such that

$$\lim_{n \rightarrow \infty} \|p_n - p\|_{L^2(I; H^1(\Omega))} = 0. \tag{2.11}$$

Then we rewrite the first equation of (2.1) in terms of p_n :

$$\partial_t u - \nu \Delta u + (A \cdot \nabla) u + (u \cdot \nabla) B + L_1 H + \nabla p_n = F + \nabla(p_n - p).$$

Therefore, by applying the Carleman estimate in the above steps, we have

$$\begin{aligned}
&\int_Q \left(\frac{1}{s^2} (|\partial_t u|^2 + |\Delta u|^2) + \frac{1}{s} (|\partial_t H|^2 + |\Delta H|^2) + |\nabla u|^2 + s |\nabla H|^2 \right. \\
&\quad \left. + s^2 |u|^2 + s^3 |H|^2 + \frac{1}{s} |\nabla p|^2 + s |p|^2 \right) e^{2s\varphi} dx dt \\
&\leq C \int_Q (|F + \nabla(p_n - p)|^2 + |G|^2 + |\nabla_{x,t} h|) e^{2s\varphi} dx dt \\
&\quad + C s^3 \int_{\partial\Omega \times I} (|u|^2 + |H|^2 + |\nabla_{x,t} u|^2 + |\nabla_{x,t} H|^2) e^{2s\varphi} dS dt \\
&\quad + C s^3 \int_{\Omega} (|u|^2 + |H|^2 + |\nabla u|^2 + |\nabla H|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta}
\end{aligned}$$

for all $s \geq s_0$. Since

$$|F + \nabla(p_n - p)|^2 \leq C (|F|^2 + |\nabla(p_n - p)|^2),$$

by letting $n \rightarrow \infty$ and using (2.11), we complete the proof of Theorem 2.1. \square

2.2 Carleman estimate without pressure term p

In this subsection, we propose another Carleman estimate for the linearized MHD system (2.1) in which we do not estimate p . Such estimate is useful in proving the stability results without the information of the pressure.

Although we can argue with less regularity which is similar to the previous subsection, here we assume higher regularity for the non-homogeneous term and the coefficients in order to clarify the essential idea.

Recall that $\Gamma \subset \partial\Omega$ is an arbitrarily given non-empty relatively open sub-boundary, the functions d and φ are given by (2.2) and (2.3) respectively. Now we state the second Carleman estimate.

Theorem 2.3. *Let $h = 0$, $F, G \in L^2(Q)$ and $\text{rot } F, \text{rot } G \in L^2(Q)$. Then there exist constants $C > 0$ and $\widehat{s}_0 > 0$ such that*

$$\begin{aligned} & \int_Q \left(\frac{1}{s} (|\partial_t \text{rot } u|^2 + |\Delta \text{rot } u|^2 + |\partial_t \text{rot } H|^2 + |\Delta \text{rot } H|^2) \right. \\ & \quad + s (|\nabla \text{rot } u|^2 + |\Delta u|^2 + |\nabla \text{rot } H|^2 + |\Delta H|^2) + s^2 (|\nabla u|^2 + |\nabla H|^2) \\ & \quad \left. + s^3 (|\text{rot } u|^2 + |\text{rot } H|^2) + s^4 (|u|^2 + |H|^2) \right) e^{2s\varphi} dx dt \\ & \leq C \int_Q (|\text{rot } F|^2 + |\text{rot } G|^2) e^{2s\varphi} dx dt \\ & \quad + Cs^4 \int_{\partial\Omega \times I} (|\nabla_{x,t} \text{rot } u|^2 + |\nabla u|^2 + |\nabla_{x,t} \text{rot } H|^2 + |\nabla H|^2 + |u|^2 + |H|^2) e^{2s\varphi} dS dt \\ & \quad + Cs^3 \int_{\Omega} (|\nabla \text{rot } u|^2 + |\text{rot } u|^2 + |\nabla \text{rot } H|^2 + |\text{rot } H|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \end{aligned}$$

for all $s \geq \widehat{s}_0$ and

$$u, H \in H^1(I; H^2(\Omega)), \quad \text{rot } u, \text{rot } H \in H^1(I; H^2(\Omega)), \quad p \in L^2(I; H^2(\Omega))$$

satisfying (2.1) and $\text{div } H = 0$.

Proof of Theorem 2.3. The key to the proof is the reduction of (2.1) to a parabolic-elliptic system. Then we can apply standard Carleman estimates for parabolic equations and elliptic equations to establish our desired Carleman estimate.

To start with, we denote $y := \text{rot } u$, $z := \text{rot } H$. By noting

$$\Delta u = -\text{rot}(\text{rot } u) + \nabla(\text{div } u) \quad \text{and} \quad \text{rot}(\nabla p) = 0,$$

the system (2.1) and $\operatorname{div} H = 0$ yield

$$\left\{ \begin{array}{l} \partial_t y - \nu \Delta y + (A^{(1)} \cdot \nabla) y = \operatorname{rot} F - (C^{(1)} \cdot \nabla) z - \sum_{k=1}^3 \nabla A_k^{(1)} \times \partial_k u - (u \cdot \nabla) \operatorname{rot} A^{(2)} \\ \quad - \sum_{k=1}^3 \nabla u_k \times \partial_k A^{(2)} - \sum_{k=1}^3 \nabla C_k^{(1)} \times \partial_k H - (H \cdot \nabla) \operatorname{rot} C^{(2)} - \sum_{k=1}^3 \nabla H_k \times \partial_k C^{(2)}, \\ \partial_t z - \kappa \Delta z + (A^{(3)} \cdot \nabla) z = \operatorname{rot} G - (C^{(5)} \cdot \nabla) y - \sum_{k=1}^3 \nabla A_k^{(3)} \times \partial_k H - (H \cdot \nabla) \operatorname{rot} A^{(4)} \\ \quad - \sum_{k=1}^3 \nabla H_k \times \partial_k A^{(4)} - \sum_{k=1}^3 \nabla C_k^{(5)} \times \partial_k u - (u \cdot \nabla) \operatorname{rot} C^{(4)} - \sum_{k=1}^3 \nabla u_k \times \partial_k C^{(4)}, \\ \Delta u = -\operatorname{rot} y, \\ \Delta H = -\operatorname{rot} z. \end{array} \right. \quad (2.12)$$

Next we employ the Carleman estimate for the parabolic system (e.g., Yamamoto [24, Theorem 3.2]) to the first and the second equations of (2.12) and we obtain

$$\begin{aligned} & \int_Q \left(\frac{1}{s} (|\partial_t y|^2 + |\Delta y|^2) + s |\nabla y|^2 + s^3 |y|^2 \right) e^{2s\varphi} dx dt \\ & \leq C \int_Q |\operatorname{rot} F|^2 e^{2s\varphi} dx dt + C \int_Q (|\nabla u|^2 + |u|^2 + |\nabla H|^2 + |H|^2 + |\nabla z|^2) e^{2s\varphi} dx dt \\ & \quad + Cs^3 \int_{\partial\Omega \times I} (|\nabla_{x,t} y|^2 + |y|^2) e^{2s\varphi} dS dt + Cs^3 \int_{\Omega} (|\nabla y|^2 + |y|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \end{aligned} \quad (2.13)$$

and

$$\begin{aligned} & \int_Q \left(\frac{1}{s} (|\partial_t z|^2 + |\Delta z|^2) + s |\nabla z|^2 + s^3 |z|^2 \right) e^{2s\varphi} dx dt \\ & \leq C \int_Q |\operatorname{rot} G|^2 e^{2s\varphi} dx dt + C \int_Q (|\nabla u|^2 + |u|^2 + |\nabla H|^2 + |H|^2 + |\nabla y|^2) e^{2s\varphi} dx dt \\ & \quad + Cs^3 \int_{\partial\Omega \times I} (|\nabla_{x,t} z|^2 + |z|^2) e^{2s\varphi} dS dt + Cs^3 \int_{\Omega} (|\nabla z|^2 + |z|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \end{aligned} \quad (2.14)$$

for all $s \geq \widehat{s}_1$. On the other hand, we employ the following Carleman estimate for the elliptic equation.

Lemma 2.4. *Let $g \in L^2(Q)$ and $w \in L^2(I; H^2(\Omega))$ satisfy*

$$\Delta w(x, t) = g(x, t), \quad (x, t) \in Q.$$

Then there exist constants $\widehat{s}_2 > 0$ and $C > 0$ such that

$$\begin{aligned} & \int_Q (|\Delta w|^2 + s|\nabla w|^2 + s^3|w|^2) e^{2s\varphi} dxdt \\ & \leq C \int_Q |g|^2 e^{2s\varphi} dxdt + Cs^3 \int_{\partial\Omega \times I} (|\nabla w|^2 + |w|^2) e^{2s\varphi} dSdt \end{aligned}$$

for all $s \geq \widehat{s}_2$.

The proof is based on the standard elliptic Carleman estimate which is well-known. For example, it can be directly proved by following similar steps as the proof of [24, Theorem 3.1], and keeping the boundary integrals.

By applying Lemma 2.4 to the third and the fourth equations of (2.12) and multiplying both sides by s , we have

$$\begin{aligned} & \int_Q (s|\Delta u|^2 + s^2|\nabla u|^2 + s^4|u|^2) e^{2s\varphi} dxdt \\ & \leq C \int_Q s|\operatorname{rot} y|^2 e^{2s\varphi} dxdt + Cs^4 \int_{\partial\Omega \times I} (|\nabla u|^2 + |u|^2) e^{2s\varphi} dSdt \end{aligned} \quad (2.15)$$

and

$$\begin{aligned} & \int_Q (s|\Delta H|^2 + s^2|\nabla H|^2 + s^4|H|^2) e^{2s\varphi} dxdt \\ & \leq C \int_Q s|\operatorname{rot} z|^2 e^{2s\varphi} dxdt + Cs^4 \int_{\partial\Omega \times I} (|\nabla H|^2 + |H|^2) e^{2s\varphi} dSdt \end{aligned} \quad (2.16)$$

for all $s \geq \widehat{s}_2$. Since

$$|\operatorname{rot} y|^2 \leq C|\nabla y|^2, \quad |\operatorname{rot} z|^2 \leq C|\nabla z|^2,$$

we sum up the estimates (2.13)–(2.16) and insert (2.13)–(2.14) into the right-hand side, and then we can find a sufficiently large constant $\widehat{s}_0 \geq \max\{\widehat{s}_1, \widehat{s}_2\}$ such that we absorb the term $C \int_Q (|\nabla u|^2 + |u|^2 + |\nabla H|^2 + |H|^2 + |\nabla y|^2 + |\nabla z|^2) e^{2s\varphi} dxdt$ on the right-hand side into the left-hand side. This completes the proof of Theorem 2.3. \square

3 Proofs of Theorems 1.1–1.3

3.1 Proof of Theorem 1.1

We divide the proof into four steps.

First step: Choice of a suitable cut-off function χ

In Theorem 1.1, we do not have $p = 0$ on $\partial\Omega$, which prevents us from applying the Carleman estimate (Theorem 2.1) established in the former section directly. Thus, here we need a cut-off argument.

Recall that $d = d(x)$ satisfies (2.2). Since $d(x) > 0$ for $x \in \Omega_0 \subset \Omega_1$, we can choose constants $0 < \varepsilon_1 < \varepsilon_2 < \varepsilon_3$ such that $\Omega_0 \subset \{x \in \Omega; d(x) > \varepsilon_3\}$.

We choose a cut-off function as follows: $\chi \in C^\infty(\mathbb{R}^3)$ satisfying $0 \leq \chi \leq 1$ and

$$\chi(x) = \begin{cases} 1 & \text{if } d(x) > \varepsilon_2, \ x \in \overline{\Omega}, \\ 0 & \text{if } d(x) < \varepsilon_1, \ x \in \overline{\Omega}. \end{cases} \quad (3.1)$$

By (2.2), we have $d = 0$ on $\partial\Omega \setminus \Gamma$. From (3.1) and the continuity of d , we easily find that χ and its derivatives vanish on $\partial\Omega \setminus \Gamma$. On the other hand, by the Sobolev extension theorem, we can find $p_0 \in H^1(I; H^1(\Omega))$ such that

$$p_0 = p \quad \text{on } \Gamma \times I \quad \text{and} \quad \|p_0\|_{H^1(I; H^1(\Omega))} \leq C\|p\|_{H^1(I; H^{\frac{1}{2}}(\Gamma))}. \quad (3.2)$$

Therefore, we set

$$q = (p - p_0), \quad q_\chi = \chi q,$$

so that $q_\chi = 0$ on $\partial\Omega \times I$ and $q_\chi \in H^1(I; H_0^1(\Omega))$. Then by letting

$$u_\chi = \chi u, \quad H_\chi = \chi H,$$

we can rewrite the linearized MHD system (1.1) by

$$\begin{cases} \partial_t u_\chi - \nu \Delta u_\chi + (A^{(1)} \cdot \nabla) u_\chi + (u_\chi \cdot \nabla) A^{(2)} + L_1 H_\chi + \nabla q_\chi = \chi F - \chi \nabla p_0 + J_1 \\ \partial_t H_\chi - \kappa \Delta H_\chi + (A^{(3)} \cdot \nabla) H_\chi + (H_\chi \cdot \nabla) A^{(4)} + L_2 u_\chi = J_2 \\ \operatorname{div} u_\chi = \nabla \chi \cdot u \end{cases} \quad (3.3)$$

Here J_1, J_2 include the terms of p, p_0 and at most first spatial derivatives of u, H multiplied by the derivatives of the cut-off function χ , in other words, we have

$$|J_k| \leq C(|\nabla \chi| + |\Delta \chi|)(|p| + |p_0| + |u| + |H| + |\nabla u| + |\nabla H|),$$

for $k = 1, 2$. Moreover, in order to prove the stability result, we need to differentiate the system once with respect to t . Henceforth, we denote

$$u^{(1)} = \partial_t u, \quad H^{(1)} = \partial_t H, \quad q^{(1)} = \partial_t q, \quad u_\chi^{(1)} = \chi \partial_t u, \quad H_\chi^{(1)} = \chi \partial_t H, \quad q_\chi^{(1)} = \chi \partial_t q.$$

Hence we can derive the system with respect to $u_\chi^{(1)}, H_\chi^{(1)}, q_\chi^{(1)}$ by (3.3):

$$\begin{cases} \partial_t u_\chi^{(1)} - \nu \Delta u_\chi^{(1)} + (A^{(1)} \cdot \nabla) u_\chi^{(1)} + (u_\chi^{(1)} \cdot \nabla) A^{(2)} + L_1 H_\chi^{(1)} + \nabla q_\chi^{(1)} \\ \quad = \chi(\partial_t F) - \chi \nabla(\partial_t p_0) + J_3 \\ \partial_t H_\chi^{(1)} - \kappa \Delta H_\chi^{(1)} + (A^{(3)} \cdot \nabla) H_\chi^{(1)} + (H_\chi^{(1)} \cdot \nabla) A^{(4)} + L_2 u_\chi^{(1)} = J_4 \\ \operatorname{div} u_\chi^{(1)} = \nabla \chi \cdot \partial_t u \end{cases} \quad (3.4)$$

Here J_3, J_4 include the terms of $p, p_0, \partial_t p, \partial_t p_0$ and at most first spatial derivatives

of u , H , $\partial_t u$, $\partial_t H$ multiplied by the derivatives of the cut-off function χ , in other words, we have

$$\begin{aligned} |J_k| &\leq C(|\nabla\chi| + |\Delta\chi|)(|p| + |p_0| + |u| + |H| + |\nabla u| + |\nabla H|) \\ &\quad + C(|\nabla\chi| + |\Delta\chi|)(|\partial_t p| + |\partial_t p_0| + |\partial_t u| + |\partial_t H| + |\nabla\partial_t u| + |\nabla\partial_t H|), \end{aligned}$$

for $k = 3, 4$.

Second step: Application of the Carleman estimate

Next we employ Theorem 2.1 to (3.3). In particular, we obtain

$$\begin{aligned} &\int_Q \left(\frac{1}{s^2} |\partial_t u_\chi|^2 + |\nabla u_\chi|^2 + s |\nabla H_\chi|^2 + s^2 |u_\chi|^2 + s^3 |H_\chi|^2 \right) e^{2s\varphi} dx dt \\ &\leq C \int_Q (|\chi F - \chi \nabla p_0 + J_1|^2 + |J_2|^2 + |\nabla_{x,t}(\nabla\chi \cdot u)|^2) e^{2s\varphi} dx dt \\ &\quad + C s^3 \int_{\partial\Omega \times I} (|u_\chi|^2 + |H_\chi|^2 + |\nabla_{x,t} u_\chi|^2 + |\nabla_{x,t} H_\chi|^2) e^{2s\varphi} dS dt \\ &\quad + C s^3 \int_\Omega (|u_\chi|^2 + |H_\chi|^2 + |\nabla u_\chi|^2 + |\nabla H_\chi|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \end{aligned} \quad (3.5)$$

for all sufficiently large $s \geq 1$. We estimate the right-hand side of the inequality (3.5) in more details.

For the first integral, by (1.3) and using the triangle inequality, we obtain

$$\begin{aligned} &\int_Q (|\chi F - \chi \nabla p_0 + J_1|^2 + |J_2|^2 + |\nabla_{x,t}(\nabla\chi \cdot u)|^2) e^{2s\varphi} dx dt \\ &\leq C \int_Q |F(x, t_0)|^2 e^{2s\varphi} dx dt + C \int_Q |\nabla p_0|^2 e^{2s\varphi} dx dt \\ &\quad + C \int_Q (|J_1|^2 + |J_2|^2 + |\nabla_{x,t}(\nabla\chi \cdot u)|^2) e^{2s\varphi} dx dt. \end{aligned} \quad (3.6)$$

Moreover, by (3.2), we have

$$C \int_Q |\nabla p_0|^2 e^{2s\varphi} dx dt \leq C e^{Cs} \|p\|_{H^1(I; H^{\frac{1}{2}}(\Gamma))}^2 \leq C e^{Cs} E_1^2.$$

Here

$$\begin{aligned} E_1 &:= \|u\|_{H^2(I; L^2(\Gamma))} + \|\nabla u\|_{H^1(I; L^2(\Gamma))} + \|H\|_{H^2(I; L^2(\Gamma))} + \|\nabla H\|_{H^1(I; L^2(\Gamma))} \\ &\quad + \|p\|_{H^1(I; H^{\frac{1}{2}}(\Gamma))}. \end{aligned}$$

Further, from (3.1), we find the derivatives of χ vanish in $\{x \in \Omega; d(x) > \varepsilon_2\}$. Then we have

$$C \int_Q (|J_1|^2 + |J_2|^2 + |\nabla_{x,t}(\nabla\chi \cdot u)|^2) e^{2s\varphi} dx dt$$

$$\begin{aligned}
&\leq Ce^{2s\mu_1} \int_Q (|J_1|^2 + |J_2|^2 + |\nabla_{x,t}(\nabla\chi \cdot u)|^2) dxdt \\
&\leq Ce^{2s\mu_1} \left(\|u\|_{H^1(I;H^1(\Omega))}^2 + \|H\|_{L^2(I;H^1(\Omega))}^2 + \|p\|_{L^2(I;L^2(\Omega))}^2 \right) \leq Ce^{2s\mu_1} M^2.
\end{aligned}$$

Here $M > 0$ is the constant introduced in the statement of the theorem and

$$\mu_1 := \max\{\varphi; d(x) \leq \varepsilon_2, t \in I\} = e^{\lambda\varepsilon_2}. \quad (3.7)$$

Therefore, (3.6) implies

$$\begin{aligned}
&\int_Q (|\chi F - \chi \nabla p_0 + J_1|^2 + |J_2|^2 + |\nabla_{x,t}(\nabla\chi \cdot u)|^2) e^{2s\varphi} dxdt \\
&\leq C \int_Q |F(x, t_0)|^2 e^{2s\varphi} dxdt + Ce^{Cs} E_1^2 + Ce^{2s\mu_1} M^2.
\end{aligned} \quad (3.8)$$

For the second integral on the right-hand side of (3.5), we note that u_χ, H_χ and their derivatives vanish on $(\partial\Omega \setminus \Gamma) \times I$, thus we obtain

$$\begin{aligned}
&Cs^3 \int_{\partial\Omega \times I} (|u_\chi|^2 + |H_\chi|^2 + |\nabla_{x,t}u_\chi|^2 + |\nabla_{x,t}H_\chi|^2) e^{2s\varphi} dSdt \\
&= Cs^3 \int_{\Gamma \times I} (|u_\chi|^2 + |H_\chi|^2 + |\nabla_{x,t}u_\chi|^2 + |\nabla_{x,t}H_\chi|^2) e^{2s\varphi} dSdt \leq Cs^3 e^{Cs} E_1^2.
\end{aligned} \quad (3.9)$$

For the third integral on the right-hand side of (3.5), we find that $\varphi(x, t) \leq e^{\lambda(\|d\|_{C(\overline{\Omega})} - \beta\delta^2)}$ for $x \in \Omega$ and $t = t_0 \pm \delta$. By denoting $\mu_2 := e^{\lambda(\|d\|_{C(\overline{\Omega})} - \beta\delta^2)}$, we estimate

$$\begin{aligned}
&Cs^3 \int_\Omega (|u_\chi|^2 + |H_\chi|^2 + |\nabla u_\chi|^2 + |\nabla H_\chi|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \\
&\leq Cs^3 e^{2s\mu_2} \int_\Omega (|u|^2 + |H|^2 + |\nabla u|^2 + |\nabla H|^2) dx \Big|_{t=t_0 \pm \delta} \leq Cs^3 e^{2s\mu_2} M^2.
\end{aligned} \quad (3.10)$$

Here we used the Sobolev embedding theorem in the last inequality. Therefore, inserting (3.8)–(3.10) into the right-hand side of (3.5) yields

$$\begin{aligned}
&\int_Q \left(\frac{1}{s^2} |\partial_t u_\chi|^2 + |\nabla u_\chi|^2 + s |\nabla H_\chi|^2 + s^2 |u_\chi|^2 + s^3 |H_\chi|^2 \right) e^{2s\varphi} dxdt \\
&\leq C \int_Q |F(x, t_0)|^2 e^{2s\varphi} dxdt + Cs^3 e^{Cs} E_1^2 + Cs^3 e^{2s\mu_3} M^2
\end{aligned} \quad (3.11)$$

for all sufficiently large $s \geq 1$. Here $\mu_3 := \max\{\mu_1, \mu_2\}$.

Similarly, we employ Theorem 2.1 to (3.4) and obtain

$$\begin{aligned}
&\int_Q \left(\frac{1}{s^2} |\partial_t u_\chi^{(1)}|^2 + |\nabla u_\chi^{(1)}|^2 + s |\nabla H_\chi^{(1)}|^2 + s^2 |u_\chi^{(1)}|^2 + s^3 |H_\chi^{(1)}|^2 \right) e^{2s\varphi} dxdt \\
&\leq C \int_Q |F(x, t_0)|^2 e^{2s\varphi} dxdt + Cs^3 e^{Cs} E_1^2 + Cs^3 e^{2s\mu_3} M^2
\end{aligned} \quad (3.12)$$

for all sufficiently large $s \geq 1$.

Third Step: A preliminary estimate

According to the first equation in (1.1), we see that

$$\begin{aligned} u_\chi^{(1)}(x, t_0) &= \chi \partial_t u(x, t_0) \\ &= \chi \left(\nu \Delta u(x, t_0) - (A^{(1)} \cdot \nabla) u(x, t_0) - (u \cdot \nabla) A^{(2)}(x, t_0) - \nabla p(x, t_0) \right) \\ &\quad - \chi L_1 H(x, t_0) + \chi F(x, t_0), \end{aligned}$$

and hence

$$\begin{aligned} \chi F(x, t_0) &= -\chi \left(\nu \Delta u(x, t_0) - (A^{(1)} \cdot \nabla) u(x, t_0) - (u \cdot \nabla) A^{(2)}(x, t_0) - \nabla p(x, t_0) \right) \\ &\quad + u_\chi^{(1)}(x, t_0) + \chi L_1 H(x, t_0) \end{aligned}$$

for $x \in \Omega$. Recalling that $L_1 H$ contains at most first-order spatial derivatives of H and

$$\chi L_1 H = L_1 H_\chi - (C^{(1)} \cdot \nabla \chi) H - \nabla \chi (C^{(3)} \cdot H),$$

we estimate the weighted L^2 -norm of $F(\cdot, t)$ in Ω at $t = t_0$:

$$\begin{aligned} \int_\Omega |\chi(x) F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx &\leq C e^{Cs} E_2^2 + C \int_\Omega |u_\chi^{(1)}(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &\quad + C \int_\Omega (|\nabla H_\chi(x, t_0)|^2 + |H_\chi(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \\ &\quad + C \int_\Omega |\nabla \chi(x)|^2 |H(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx. \end{aligned} \quad (3.13)$$

Here

$$E_2 := \|u(\cdot, t_0)\|_{H^2(\Omega)} + \|\nabla p(\cdot, t_0)\|_{L^2(\Omega)}.$$

Next we estimate the second, the third and the fourth terms on the right-hand side above respectively. For the second term, we have

$$\begin{aligned} &\int_\Omega |u_\chi^{(1)}(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &= \int_{t_0-\delta}^{t_0} \left(\partial_t \int_\Omega |u_\chi^{(1)}|^2 e^{2s\varphi} dx \right) dt + \int_\Omega |u_\chi^{(1)}(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\ &= \int_{t_0-\delta}^{t_0} \int_\Omega \left(2u_\chi^{(1)} \cdot \partial_t u_\chi^{(1)} + 2s(\partial_t \varphi) |u_\chi^{(1)}|^2 \right) e^{2s\varphi} dx dt + \int_\Omega |u_\chi^{(1)}(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\ &\leq C \int_Q \left(|u_\chi^{(1)}| |\partial_t u_\chi^{(1)}| + s |u_\chi^{(1)}|^2 \right) e^{2s\varphi} dx dt + \int_\Omega |u_\chi^{(1)}(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\ &\leq C \int_Q \left(\frac{1}{s^2} |\partial_t u_\chi^{(1)}|^2 + s^2 |u_\chi^{(1)}|^2 \right) e^{2s\varphi} dx dt + C e^{2s\mu_3} M^2. \end{aligned}$$

Here we used $|a||b| \leq \frac{1}{2}(s^2|a|^2 + s^{-2}|b|^2)$ and the argument used in (3.10). Similarly,

we have

$$\begin{aligned}
& \int_{\Omega} |H_{\chi}(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\
& \leq C \int_Q \left(\frac{1}{s^2} |\partial_t H_{\chi}|^2 + s^2 |H_{\chi}|^2 \right) e^{2s\varphi} dx dt + \int_{\Omega} |H_{\chi}(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\
& \leq C \int_Q \left(\frac{1}{s^2} |H_{\chi}^{(1)}|^2 + s^2 |H_{\chi}|^2 \right) e^{2s\varphi} dx dt + C e^{2s\mu_3} M^2.
\end{aligned}$$

Moreover, we calculate for $j = 1, 2, 3$,

$$\begin{aligned}
& \int_{\Omega} |\partial_j H_{\chi}(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\
& = \int_{t_0 - \delta}^{t_0} \left(\partial_t \int_{\Omega} |\partial_j H_{\chi}|^2 e^{2s\varphi} dx \right) dt + \int_{\Omega} |\partial_j H_{\chi}(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\
& \leq C \int_Q (|\partial_j H_{\chi}| |\partial_j \partial_t H_{\chi}| + s |\partial_j H_{\chi}|^2) e^{2s\varphi} dx dt + C e^{2s\mu_3} M^2 \\
& \leq C \int_Q \left(\frac{1}{s} |\partial_j H_{\chi}^{(1)}|^2 + s |\partial_j H_{\chi}|^2 \right) e^{2s\varphi} dx dt + C e^{2s\mu_3} M^2.
\end{aligned}$$

For the fourth term, we note again that the derivatives of χ vanish in $\{x \in \Omega; d(x) > \varepsilon_2\}$, which implies

$$\int_{\Omega} |\nabla \chi(x)|^2 |H(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \leq C e^{2s\mu_1} \|H(\cdot, t_0)\|_{L^2(\Omega)}^2 \leq C e^{2s\mu_3} M^2.$$

Therefore, equipped with all the above estimations, (3.13) together with (3.11)–(3.12) implies

$$\int_{\Omega} |\chi(x) F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \leq C \int_Q |F(x, t_0)|^2 e^{2s\varphi} dx dt + C s^3 e^{2s\mu_3} M^2 + C s^3 e^C D_1^2 \quad (3.14)$$

for all sufficiently large $s \geq 1$. Here we noted that $E_1^2 + E_2^2 \leq C D_1^2$ where D_1 is the norm of the data introduced in the theorem.

Fourth step: Absorption and completion of the proof

Now we absorb the first term on the right-hand side of (3.14) into the left-hand side. With reference to $\varphi(x, t) = d(x) - \beta(t - t_0)^2$, $(x, t) \in Q$, the Lebesgue's dominated convergence theorem yields

$$\int_Q |F(x, t_0)|^2 e^{2s\varphi} dx dt = o(1) \int_{\Omega} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx$$

as $s \rightarrow \infty$. We refer to [5, Lemma 5.3, pp.120] for the detailed argument. By recalling the choice of μ_1 , see (3.7) and that $F \in \mathcal{F}_{M,1}$, we have

$$\int_Q |F(x, t_0)|^2 e^{2s\varphi} dx dt$$

$$\begin{aligned}
&= o(1) \left(\int_{\{x \in \Omega; \varphi(x, t_0) > \mu_1\}} + \int_{\{x \in \Omega; \varphi(x, t_0) \leq \mu_1\}} \right) |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\
&= o(1) \int_{\{x \in \Omega; \varphi(x, t_0) > \mu_1\}} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + C e^{2s\mu_1} M^2.
\end{aligned}$$

On the other hand, with reference to (3.1), we have $\chi = 1$ in $\{x \in \Omega; d(x) > \varepsilon_2\} = \{x \in \Omega; \varphi(x, t_0) > \mu_1\}$. Then

$$\begin{aligned}
\int_{\Omega} |\chi(x) F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx &\geq \int_{\{x \in \Omega; \varphi(x, t_0) > \mu_1\}} |\chi(x)|^2 |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\
&= \int_{\{x \in \Omega; \varphi(x, t_0) > \mu_1\}} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx,
\end{aligned}$$

so that (3.14) leads to

$$\begin{aligned}
&\int_{\{x \in \Omega; \varphi(x, t_0) > \mu_1\}} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\
&\leq o(1) \int_{\{x \in \Omega; \varphi(x, t_0) > \mu_1\}} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + C s^3 e^{2s\mu_3} M^2 + C s^3 e^{Cs} D_1^2.
\end{aligned}$$

Absorbing the first term on the right-hand side into the left-hand side, we obtain

$$\int_{\{x \in \Omega; \varphi(x, t_0) > \mu_1\}} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \leq C s^3 e^{2s\mu_3} M^2 + C s^3 e^{Cs} D_1^2$$

for sufficiently large s . By noting that $\mu_1 = e^{\lambda\varepsilon_2}$, $0 < \varepsilon_2 < \varepsilon_3$ and denoting $\mu_0 := e^{\lambda\varepsilon_3}$, we give a lower bound of the left-hand side above:

$$\begin{aligned}
\int_{\{x \in \Omega; \varphi(x, t_0) > \mu_1\}} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx &= \int_{\{x \in \Omega; d(x) > \varepsilon_2\}} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\
&\geq \int_{\{x \in \Omega; d(x) > \varepsilon_3\}} |F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\
&> e^{2s\mu_0} \int_{\{x \in \Omega; d(x) > \varepsilon_3\}} |F(x, t_0)|^2 dx \\
&\geq e^{2s\mu_0} \|F(\cdot, t_0)\|_{L^2(\Omega_0)}^2
\end{aligned}$$

where we used the inclusion $\Omega_0 \subset \{x \in \Omega; d(x) > \varepsilon_3\}$ in the last inequality. Therefore,

$$\|F(\cdot, t_0)\|_{L^2(\Omega_0)}^2 \leq C s^3 e^{-2s(\mu_0 - \mu_3)} M^2 + C s^3 e^{Cs} D_1^2 \quad (3.15)$$

for sufficiently large s . Finally, by taking a large $\beta > 0$ such that

$$\varepsilon_3 > \|d\|_{C(\overline{\Omega})} - \beta\delta^2,$$

which implies $\mu_0 > \mu_2$, and noting $\mu_0 > \mu_1$ from the definition, we find $\mu_0 > \max\{\mu_1, \mu_2\} = \mu_3$. Then we end up with the stability of Hölder type by minimizing

the right-hand side of (3.15) with respect to s (see e.g., [24, Proof of Theorem 5.1, pp. 27–28]).

3.2 Proof of Theorem 1.2

With the additional regularity assumptions on the unknown and the measurements, we can employ Theorem 2.3 and derive the stability result without the measurements on the pressure p .

First step: A preliminary estimate

We start the proof with a weighted estimate. In order to erase the effect of the pressure p , we need to apply the rotation operator to both sides of the first equation of (1.1). Actually by using the equalities $\text{rot}(\text{rot } u) = -\Delta u + \nabla(\text{div } u)$, $\text{rot}(\text{rot } F) = -\Delta F + \nabla(\text{div } F)$ and $\text{rot}(\nabla p) = 0$, we take the rotation operator twice on both sides of the first equation of (1.1). In particular, we fix $t = t_0$ and then by (1.6) we obtain

$$\partial_t \Delta u(x, t_0) - \nu \Delta^2 u(x, t_0) - \text{rot}(\text{rot } L_1 H(x, t_0)) = \Delta F(x, t_0) + L_3 u(x, t_0), \quad x \in \Omega \quad (3.16)$$

where L_3 is a third-order differential operator given by

$$L_3 u := \text{rot} \left(\text{rot} \left((A^{(1)} \cdot \nabla) u + (u \cdot \nabla) A^{(2)} \right) \right).$$

By taking the weighted L^2 -norm over Ω on both sides of (3.16), we obtain

$$\begin{aligned} & \int_{\Omega} |\Delta F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &= \int_{\Omega} |\partial_t \Delta u(x, t_0) - \nu \Delta^2 u(x, t_0) - \text{rot}(\text{rot } L_1 H(x, t_0)) - L_3 u(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &\leq C \int_{\Omega} |\partial_t \Delta u(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + C e^{Cs} E_3^2 \end{aligned} \quad (3.17)$$

where

$$E_3 := \|u(\cdot, t_0)\|_{H^4(\Omega)} + \|H(\cdot, t_0)\|_{H^3(\Omega)}.$$

Next, we give an upper bound of the first term on the right-hand side of (3.17).

Actually, similarly as we have done in the former subsection, we estimate

$$\begin{aligned} & \int_{\Omega} |\partial_t \Delta u(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &= \int_{t_0-\delta}^{t_0} \partial_t \left(\int_{\Omega} |\partial_t \Delta u|^2 e^{2s\varphi} dx \right) dt + \int_{\Omega} |\partial_t \Delta u(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\ &= \int_{t_0-\delta}^{t_0} \int_{\Omega} (2\Delta(\partial_t u) \cdot \Delta(\partial_t^2 u) + 2s(\partial_t \varphi) |\Delta(\partial_t u)|^2) e^{2s\varphi} dx dt \end{aligned}$$

$$\begin{aligned}
& + \int_{\Omega} |\partial_t \Delta u(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\
& \leq C \int_Q (|\Delta(\partial_t u)| |\Delta(\partial_t^2 u)| + s |\Delta(\partial_t u)|^2) e^{2s\varphi} dx dt \\
& \quad + \int_{\Omega} |\partial_t \Delta u(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\
& \leq C \int_Q (|\Delta(\partial_t^2 u)|^2 + s |\Delta(\partial_t u)|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2
\end{aligned} \tag{3.18}$$

where $M > 0$ is the constant introduced in the statement of the theorem and μ_1 is the maximum of φ over $\overline{\Omega} \times \{t_0 \pm \delta\}$ and $(\partial\Omega \setminus \Gamma) \times I$, which implies

$$\begin{aligned}
\mu_1 & \geq \max\{\varphi(x, t); x \in \overline{\Omega}, t = t_0 \pm \delta\} = e^{\lambda(\|d\|_{C(\overline{\Omega})} - \beta\delta^2)} \quad \text{and} \\
\mu_1 & \geq \max\{\varphi(x, t); x \in \partial\Omega \setminus \Gamma, t \in I\} = 1.
\end{aligned} \tag{3.19}$$

Here we used $2A \cdot B \leq |A|^2 + |B|^2$ and (3.18) holds true for all $s \geq 1$. Thus, we reach

$$\begin{aligned}
\int_{\Omega} |\Delta F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx & \leq C \int_Q (|\Delta(\partial_t^2 u)|^2 + s |\Delta(\partial_t u)|^2) e^{2s\varphi} dx dt \\
& \quad + C e^{2s\mu_1} M^2 + C e^{Cs} E_3^2.
\end{aligned} \tag{3.20}$$

Second step: Application of the Carleman estimate

For simplicity of the expressions, we introduce the following notations:

$$u^{(k)} = \partial_t^k u, \quad H^{(k)} = \partial_t^k H, \quad p^{(k)} = \partial_t^k p, \quad k = 1, 2.$$

Then taking once and twice time derivatives of the governing system (1.1) yields

$$\begin{cases} \partial_t u^{(1)} - \nu \Delta u^{(1)} + (A^{(1)} \cdot \nabla) u^{(1)} + (u^{(1)} \cdot \nabla) A^{(2)} + L_1 H^{(1)} + \nabla p^{(1)} = \partial_t F + J_1, \\ \partial_t H^{(1)} - \kappa \Delta H^{(1)} + (A^{(3)} \cdot \nabla) H^{(1)} + (H^{(1)} \cdot \nabla) A^{(4)} + L_2 u^{(1)} = J_2, \\ \operatorname{div} u^{(1)} = 0, \end{cases} \tag{3.21}$$

and

$$\begin{cases} \partial_t u^{(2)} - \nu \Delta u^{(2)} + (A^{(1)} \cdot \nabla) u^{(2)} + (u^{(2)} \cdot \nabla) A^{(2)} + L_1 H^{(2)} + \nabla p^{(2)} = \partial_t^2 F + J_3, \\ \partial_t H^{(2)} - \kappa \Delta H^{(2)} + (A^{(3)} \cdot \nabla) H^{(2)} + (H^{(2)} \cdot \nabla) A^{(4)} + L_2 u^{(2)} = J_4, \\ \operatorname{div} u^{(2)} = 0 \end{cases} \tag{3.22}$$

Here J_1, J_2 denote the terms including at most first-order spatial derivative of u and H while J_3, J_4 include additionally at most first-order spatial derivatives of $u^{(1)}, H^{(1)}$, and we have

$$|\operatorname{rot} J_k| \leq C (|u| + |\nabla u| + |\nabla \operatorname{rot} u| + |H| + |\nabla H| + |\nabla \operatorname{rot} H|), \quad k = 1, 2.$$

$$|\operatorname{rot} J_k| \leq C \left(|u| + |\nabla u| + |\nabla \operatorname{rot} u| + |H| + |\nabla H| + |\nabla \operatorname{rot} H| + |u^{(1)}| + |\nabla u^{(1)}| + |\nabla \operatorname{rot} u^{(1)}| + |H^{(1)}| + |\nabla H^{(1)}| + |\nabla \operatorname{rot} H^{(1)}| \right), \quad k = 3, 4.$$

We employ Theorem 2.3 to (1.1), and in particular, we obtain

$$\begin{aligned} & \int_Q (s|\nabla \operatorname{rot} u|^2 + s|\nabla \operatorname{rot} H|^2 + s^2|\nabla u|^2 + s^2|\nabla H|^2 + s^4|u|^2 + s^4|H|^2) e^{2s\varphi} dxdt \\ & \leq C \int_Q |\operatorname{rot} F|^2 e^{2s\varphi} dxdt \\ & + Cs^4 \int_{\partial\Omega \times I} (|\nabla_{x,t} \operatorname{rot} u|^2 + |\nabla u|^2 + |\nabla_{x,t} \operatorname{rot} H|^2 + |\nabla H|^2 + |u|^2 + |H|^2) e^{2s\varphi} dSdt \\ & + Cs^3 \int_{\Omega} (|\nabla \operatorname{rot} u|^2 + |\operatorname{rot} u|^2 + |\nabla \operatorname{rot} H|^2 + |\operatorname{rot} H|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \end{aligned} \quad (3.23)$$

for all sufficiently large $s \geq 1$. Now we estimate the upper bounds of the second and the third terms on the right-hand side above.

By noting $\varphi(\cdot, t_0 \pm \delta) \leq \mu_1$ on $\overline{\Omega}$ (μ_1 is given in (3.19)), the third term admits

$$Cs^3 \int_{\Omega} (|\nabla \operatorname{rot} u|^2 + |\operatorname{rot} u|^2 + |\nabla \operatorname{rot} H|^2 + |\operatorname{rot} H|^2) e^{2s\varphi} dx \Big|_{t=t_0 \pm \delta} \leq Cs^3 e^{2s\mu_1} M^2.$$

For the second term, we divide the integral into two parts over $\Gamma \times I$ and $(\partial\Omega \setminus \Gamma) \times I$. By noting that φ is dominated by some constant in $\Gamma \times I$, and is dominated by $\mu_1 \geq 1$ in $(\partial\Omega \setminus \Gamma) \times I$, we obtain

$$\begin{aligned} & Cs^4 \int_{\partial\Omega \times I} (|\nabla_{x,t} \operatorname{rot} u|^2 + |\nabla u|^2 + |\nabla_{x,t} \operatorname{rot} H|^2 + |\nabla H|^2 + |u|^2 + |H|^2) e^{2s\varphi} dSdt \\ & = Cs^4 \int_{\Gamma \times I} (|\nabla_{x,t} \operatorname{rot} u|^2 + |\nabla u|^2 + |\nabla_{x,t} \operatorname{rot} H|^2 + |\nabla H|^2 + |u|^2 + |H|^2) e^{2s\varphi} dSdt \\ & + Cs^4 \int_{(\partial\Omega \setminus \Gamma) \times I} (|\nabla_{x,t} \operatorname{rot} u|^2 + |\nabla u|^2 + |\nabla_{x,t} \operatorname{rot} H|^2 + |\nabla H|^2 + |u|^2 + |H|^2) e^{2s\varphi} dSdt \\ & \leq Cs^4 e^{Cs} E_4^2 + Cs^4 e^{2s\mu_1} M^2 \end{aligned}$$

where

$$\begin{aligned} E_4 := & \|\nabla \operatorname{rot} u\|_{H^2(I; L^2(\Gamma))} + \|\nabla \operatorname{rot} H\|_{H^2(I; L^2(\Gamma))} + \|\nabla u\|_{H^3(I; L^2(\Gamma))} \\ & + \|\nabla H\|_{H^3(I; L^2(\Gamma))} + \|u\|_{H^2(I; L^2(\Gamma))} + \|H\|_{H^2(I; L^2(\Gamma))}. \end{aligned}$$

Therefore, we derive from (3.23) that

$$\begin{aligned} & \int_Q (s|\nabla \operatorname{rot} u|^2 + s|\nabla \operatorname{rot} H|^2 + s^2|\nabla u|^2 + s^2|\nabla H|^2 + s^4|u|^2 + s^4|H|^2) e^{2s\varphi} dxdt \\ & \leq C \int_Q |\operatorname{rot} F|^2 e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_4^2 + Cs^4 e^{2s\mu_1} M^2 \end{aligned} \quad (3.24)$$

for all sufficiently large $s \geq 1$. Similarly, we employ Theorem 2.3 to (3.21) and we obtain

$$\begin{aligned}
& \int_Q \left(s|\Delta u^{(1)}|^2 + s|\nabla \text{rot } u^{(1)}|^2 + s|\nabla \text{rot } H^{(1)}|^2 + s^2|\nabla u^{(1)}|^2 \right. \\
& \quad \left. + s^2|\nabla H^{(1)}|^2 + s^4|u^{(1)}|^2 + s^4|H^{(1)}|^2 \right) e^{2s\varphi} dxdt \\
& \leq C \int_Q (|\text{rot}(\partial_t F) + \text{rot } J_1|^2 + |\text{rot } J_2|^2) e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_4^2 + Cs^4 e^{2s\mu_1} M^2 \\
& \leq C \int_Q |\text{rot}(\partial_t F)|^2 e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_4^2 + Cs^4 e^{2s\mu_1} M^2 \\
& + C \int_Q (|\nabla \text{rot } u|^2 + |\nabla \text{rot } H|^2 + |\nabla u|^2 + |\nabla H|^2 + |u|^2 + |H|^2) e^{2s\varphi} dxdt \quad (3.25)
\end{aligned}$$

for all sufficiently large $s \geq 1$. Inserting (3.24) into the right-hand side of (3.25) yields

$$\begin{aligned}
& \int_Q \left(s|\Delta u^{(1)}|^2 + s|\nabla \text{rot } u^{(1)}|^2 + s|\nabla \text{rot } H^{(1)}|^2 + s^2|\nabla u^{(1)}|^2 \right. \\
& \quad \left. + s^2|\nabla H^{(1)}|^2 + s^4|u^{(1)}|^2 + s^4|H^{(1)}|^2 \right) e^{2s\varphi} dxdt \\
& \leq C \int_Q (|\text{rot}(\partial_t F)|^2 + |\text{rot } F|^2) e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_4^2 + Cs^4 e^{2s\mu_1} M^2 \quad (3.26)
\end{aligned}$$

for all sufficiently large $s \geq 1$.

Next we employ Theorem 2.3 to (3.22). We pick up the necessary terms and obtain

$$\begin{aligned}
& \int_Q s|\Delta u^{(2)}|^2 e^{2s\varphi} dxdt \\
& \leq C \int_Q (|\text{rot}(\partial_t^2 F) + \text{rot } J_3|^2 + |\text{rot } J_4|^2) e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_4^2 + Cs^4 e^{2s\mu_1} M^2 \\
& \leq C \int_Q |\text{rot}(\partial_t^2 F)|^2 e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_4^2 + Cs^4 e^{2s\mu_1} M^2 \\
& + C \int_Q (|\nabla \text{rot } u|^2 + |\nabla \text{rot } H|^2 + |\nabla u|^2 + |\nabla H|^2 + |u|^2 + |H|^2 + |\nabla \text{rot } u^{(1)}|^2 \\
& + |\nabla \text{rot } H^{(1)}|^2 + |\nabla u^{(1)}|^2 + |\nabla H^{(1)}|^2 + |u^{(1)}|^2 + |H^{(1)}|^2) e^{2s\varphi} dxdt \quad (3.27)
\end{aligned}$$

for all sufficiently large $s \geq 1$. By (3.24) and (3.26), we can get rid of the last term on the right-hand side of (3.27). Moreover, by (1.5) we have

$$|\text{rot}(\partial_t^k F)(x, t)|^2 \leq C (|\nabla F(x, t_0)|^2 + |F(x, t_0)|^2), \quad k = 0, 1, 2,$$

then we reach

$$\int_Q (s|\Delta u^{(1)}|^2 + s|\Delta u^{(2)}|^2) e^{2s\varphi} dxdt$$

$$\leq C \int_Q (|\nabla F(x, t_0)|^2 + |F(x, t_0)|^2) e^{2s\varphi} dx dt + Cs^4 e^{Cs} E_4^2 + Cs^4 e^{2s\mu_1} M^2$$

for all sufficiently large $s \geq 1$. We insert this inequality into the right-hand side of (3.20) and we obtain

$$\begin{aligned} \int_{\Omega} |\Delta F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx &\leq C \int_Q (|\nabla F(x, t_0)|^2 + |F(x, t_0)|^2) e^{2s\varphi} dx dt \\ &\quad + Cs^4 e^{2s\mu_1} M^2 + Cs^4 e^{Cs} D_2^2 \end{aligned} \quad (3.28)$$

for all sufficiently large $s \geq 1$. Here we noted that $E_3^2 + E_4^2 \leq CD_2^2$ where D_2 is the norm of the data that introduced in the theorem.

Third Step: Lower bound of (3.28) and completion of the proof

We introduce the following lemma, which is based on a classical Carleman estimate for elliptic equations.

Lemma 3.1. *There exist constants $\widehat{s} > 0$ and $C > 0$ such that*

$$\begin{aligned} &\int_{\Omega} (s|\nabla F(x, t_0)|^2 + s^3|F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \\ &\leq C \int_{\Omega} |\Delta F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + Cs^3 e^{2s\mu_1} M^2 \end{aligned} \quad (3.29)$$

for all $s \geq \widehat{s}$.

Proof. According to the assumption that $F \in \mathcal{F}_{2,M}$, we have $F(\cdot, t_0)$ and $\nabla F(\cdot, t_0)$ vanish on Γ . By the classical Carleman estimate for the elliptic equations, we obtain

$$\begin{aligned} &\int_{\Omega} (s|\nabla F(x, t_0)|^2 + s^3|F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \\ &\leq C \int_{\Omega} |\Delta F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + C \int_{\partial\Omega \setminus \Gamma} (s|\nabla F(x, t_0)|^2 + s^3|F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \end{aligned}$$

for all sufficiently large $s \geq 1$. By noting $\varphi(\cdot, t_0) = e^{\lambda d} = 1 \leq \mu_1$ on $\partial\Omega \setminus \Gamma$, we have

$$\begin{aligned} &\int_{\Omega} (s|\nabla F(x, t_0)|^2 + s^3|F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \\ &\leq C \int_{\Omega} |\Delta F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + Cs^3 e^{2s\mu_1} M^2 \end{aligned}$$

for all sufficiently large $s \geq 1$. □

Now we combine (3.28) with (3.29) and we obtain

$$\begin{aligned} &\int_{\Omega} (s|\nabla F(x, t_0)|^2 + s^3|F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \\ &\leq C \int_Q (|\nabla F(x, t_0)|^2 + |F(x, t_0)|^2) e^{2s\varphi} dx dt + Cs^4 e^{2s\mu_1} M^2 + Cs^4 e^{Cs} D_2^2 \end{aligned} \quad (3.30)$$

for all sufficiently large $s \geq 1$. Since $\varphi(x, \cdot)$ attains its maximum at t_0 , we have

$$\int_Q (|\nabla F(x, t_0)|^2 + |F(x, t_0)|^2) e^{2s\varphi} dx dt \leq 2\delta \int_\Omega (|\nabla F(x, t_0)|^2 + |F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx.$$

Thus, by taking s large enough, we can absorb the first term on the right-hand side of (3.30) into the left-hand side. This yields

$$\int_\Omega (s|\nabla F(x, t_0)|^2 + s^3|F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \leq Cs^4 e^{2s\mu_1} M^2 + Cs^4 e^{Cs} D_2^2 \quad (3.31)$$

for all s large enough.

We recall that $\overline{\Omega_0} \subset \Omega \cup \Gamma$, which implies that there exists $\varepsilon_0 > 0$ such that $d(x) \geq \varepsilon_0$ for $x \in \Omega_0$, that is, $\varphi(x, t_0) = e^{\lambda d(x)} \geq e^{\lambda \varepsilon_0} =: \mu_0$. Therefore, (3.31) yields

$$\begin{aligned} e^{2s\mu_0} \|F(\cdot, t_0)\|_{H^1(\Omega_0)}^2 &\leq \int_{\Omega_0} (s|\nabla F(x, t_0)|^2 + s^3|F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \\ &\leq \int_\Omega (s|\nabla F(x, t_0)|^2 + s^3|F(x, t_0)|^2) e^{2s\varphi(x, t_0)} dx \\ &\leq Cs^4 e^{2s\mu_1} M^2 + Cs^4 e^{Cs} D_2^2, \end{aligned}$$

and hence

$$\|F(\cdot, t_0)\|_{H^1(\Omega_0)}^2 \leq Cs^4 e^{-2s(\mu_0 - \mu_1)} M^2 + Cs^4 e^{Cs} D_2^2 \quad (3.32)$$

for all s large enough. Since $\mu_1 = \max\{e^{\lambda(\|d\|_{C(\overline{\Omega})} - \beta\delta^2)}, 1\}$ and $\mu_0 = e^{\lambda\varepsilon_0} > 1$, we can find $\beta > 0$ large enough such that $\|d\|_{C(\overline{\Omega})} - \beta\delta^2 < \varepsilon_0$ and consequently $\mu_0 - \mu_1 > 0$. This enables us to end up with a stability of Hölder type by minimizing the right-hand side of (3.32) with respect to s .

3.3 Proof of Theorem 1.3

Thanks to the assumption (1.4), we need only apply the rotation operator once to the governing equation (1.1). Actually we can simply follow the steps of the former subsection.

First step: A preliminary estimate

Again we start the proof with a weighted estimate. In order to erase the effect of the pressure p , we apply the rotation operator to both sides of the first equation of (1.1). In particular, by setting $v = \text{rot } u$ and taking $t = t_0$, we obtain

$$\partial_t v(x, t_0) - \nu \Delta v(x, t_0) + \text{rot } L_1 H(x, t_0) = \text{rot } F(x, t_0) + L_4 u(x, t_0), \quad x \in \Omega, \quad (3.33)$$

where L_4 is a second-order differential operator given by

$$L_4 u := -\operatorname{rot} \left((A^{(1)} \cdot \nabla) u + (u \cdot \nabla) A^{(2)} \right).$$

By taking the weighted L^2 -norm over Ω on both sides of (3.33), we obtain

$$\begin{aligned} & \int_{\Omega} |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &= \int_{\Omega} |\partial_t v(x, t_0) - \nu \Delta v(x, t_0) + \operatorname{rot} L_1 H(x, t_0) - L_4 u(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &\leq C \int_{\Omega} |\partial_t v(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + C \int_{\Omega} |\nabla \operatorname{rot} H(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &\quad + C \int_{\Omega} |\nabla H(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + C \int_{\Omega} |H(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx + C e^{Cs} E_5^2. \end{aligned} \quad (3.34)$$

Here

$$E_5 := \|u(\cdot, t_0)\|_{H^3(\Omega)}$$

and we used

$$|\operatorname{rot} L_1 H(x, t_0)|^2 \leq C (|\nabla \operatorname{rot} H(x, t_0)|^2 + |\nabla H(x, t_0)|^2 + |H(x, t_0)|^2), \quad x \in \Omega.$$

Next, we give the upper bounds of the first term to the fourth term on the right-hand side of (3.34), respectively.

By integration by parts, we estimate

$$\begin{aligned} & \int_{\Omega} |\partial_t v(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &= \int_{t_0-\delta}^{t_0} \partial_t \left(\int_{\Omega} |\partial_t v|^2 e^{2s\varphi} dx \right) dt + \int_{\Omega} |\partial_t v(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\ &= \int_{t_0-\delta}^{t_0} \int_{\Omega} (2(\partial_t v) \cdot (\partial_t^2 v) + 2s(\partial_t \varphi) |\partial_t v|^2) e^{2s\varphi} dx dt \\ &\quad + \int_{\Omega} |\partial_t v(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\ &\leq C \int_Q (|\partial_t v| |\partial_t^2 v| + s |\partial_t v|^2) e^{2s\varphi} dx dt + \int_{\Omega} |\partial_t v(x, t_0 - \delta)|^2 e^{2s\varphi(x, t_0 - \delta)} dx \\ &\leq C \int_Q (s^{-1} |\partial_t^2 v|^2 + s |\partial_t v|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2 \\ &\leq C \int_Q (s^{-1} |\partial_t^2 v|^2 + s^3 |\partial_t v|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2 \end{aligned}$$

for all $s \geq 1$. Here $M > 0$ is the constant introduced in the statement of the theorem and μ_1 is the same constant introduced in the former subsection, see (3.7).

Similarly, we have

$$\begin{aligned} \int_{\Omega} |H(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx &\leq C \int_Q (s^{-1} |\partial_t H|^2 + s |H|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2 \\ &\leq C \int_Q (s^4 |\partial_t H|^2 + s^4 |H|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2, \end{aligned}$$

$$\begin{aligned} \int_{\Omega} |\nabla H(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx &\leq C \int_Q (s^{-1} |\nabla(\partial_t H)|^2 + s |\nabla H|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2 \\ &\leq C \int_Q (s^2 |\nabla(\partial_t H)|^2 + s^2 |\nabla H|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2, \end{aligned}$$

$$\begin{aligned} &\int_{\Omega} |\nabla \operatorname{rot} H(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &\leq C \int_Q (s^{-1} |\nabla \operatorname{rot}(\partial_t H)|^2 + s |\nabla \operatorname{rot} H|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2 \\ &\leq C \int_Q (s |\nabla \operatorname{rot}(\partial_t H)|^2 + s |\nabla \operatorname{rot} H|^2) e^{2s\varphi} dx dt + C e^{2s\mu_1} M^2 \end{aligned}$$

for all $s \geq 1$. Thus, we reach

$$\begin{aligned} &\int_{\Omega} |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \\ &\leq C \int_Q (s^{-1} |\partial_t^2 v|^2 + s^3 |\partial_t v|^2) e^{2s\varphi} dx dt + C \int_Q (s |\nabla \operatorname{rot}(\partial_t H)|^2 + s |\nabla \operatorname{rot} H|^2) e^{2s\varphi} dx dt \\ &\quad + C \int_Q (s^4 |\partial_t H|^2 + s^4 |H|^2) e^{2s\varphi} dx dt + C \int_Q (s^2 |\nabla(\partial_t H)|^2 + s^2 |\nabla H|^2) e^{2s\varphi} dx dt \\ &\quad + C e^{2s\mu_1} M^2 + C e^{Cs} E_5^2. \end{aligned} \tag{3.35}$$

Second step: Application of the Carleman estimate

For simplicity of the expressions, we introduce the following notations:

$$u^{(1)} = \partial_t u, \quad H^{(1)} = \partial_t H, \quad p^{(1)} = \partial_t p.$$

Recall that we take the time derivative of the governing system (1.1) and obtain

$$\begin{cases} \partial_t u^{(1)} - \nu \Delta u^{(1)} + (A^{(1)} \cdot \nabla) u^{(1)} + (u^{(1)} \cdot \nabla) A^{(2)} + L_1 H^{(1)} + \nabla p^{(1)} = \partial_t F + J_1, \\ \partial_t H^{(1)} - \kappa \Delta H^{(1)} + (A^{(3)} \cdot \nabla) H^{(1)} + (H^{(1)} \cdot \nabla) A^{(4)} + L_2 u^{(1)} = J_2, \\ \operatorname{div} u^{(1)} = 0, \end{cases} \tag{3.36}$$

where J_1, J_2 satisfy

$$|\operatorname{rot} J_k| \leq C (|u| + |\nabla u| + |\nabla \operatorname{rot} u| + |H| + |\nabla H| + |\nabla \operatorname{rot} H|), \quad k = 1, 2.$$

We employ Theorem 2.3 to (1.1) and by the same argument in deriving (3.24), we obtain

$$\begin{aligned} & \int_Q (s|\nabla \operatorname{rot} u|^2 + s|\nabla \operatorname{rot} H|^2 + s^2|\nabla u|^2 + s^2|\nabla H|^2 + s^4|u|^2 + s^4|H|^2) e^{2s\varphi} dxdt \\ & \leq C \int_Q |\operatorname{rot} F|^2 e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_6^2 + Cs^4 e^{2s\mu_1} M^2 \end{aligned} \quad (3.37)$$

for all sufficiently large $s \geq 1$. Here

$$\begin{aligned} E_6 := & \|\nabla \operatorname{rot} u\|_{H^1(I;L^2(\Gamma))} + \|\nabla \operatorname{rot} H\|_{H^1(I;L^2(\Gamma))} + \|\nabla u\|_{H^2(I;L^2(\Gamma))} \\ & + \|\nabla H\|_{H^2(I;L^2(\Gamma))} + \|u\|_{H^1(I;L^2(\Gamma))} + \|H\|_{H^1(I;L^2(\Gamma))}. \end{aligned}$$

Next, we employ Theorem 2.3 to (3.36) and we obtain

$$\begin{aligned} & \int_Q \left(\frac{1}{s} |\partial_t \operatorname{rot} u^{(1)}|^2 + s |\nabla \operatorname{rot} H^{(1)}|^2 + s^2 |\nabla H^{(1)}|^2 + s^3 |\operatorname{rot} u^{(1)}|^2 + s^4 |H^{(1)}|^2 \right) e^{2s\varphi} dxdt \\ & \leq C \int_Q (|\operatorname{rot}(\partial_t F) + \operatorname{rot} J_1|^2 + |\operatorname{rot} J_2|^2) e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_6^2 + Cs^4 e^{2s\mu_1} M^2 \\ & \leq C \int_Q |\partial_t \operatorname{rot} F|^2 e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_6^2 + Cs^4 e^{2s\mu_1} M^2 \\ & \quad + C \int_Q (|\nabla \operatorname{rot} u|^2 + |\nabla \operatorname{rot} H|^2 + |\nabla u|^2 + |\nabla H|^2 + |u|^2 + |H|^2) e^{2s\varphi} dxdt \end{aligned} \quad (3.38)$$

for all sufficiently large $s \geq 1$. Inserting (3.37) into the right-hand side of (3.38) yields

$$\begin{aligned} & \int_Q \left(\frac{1}{s} |\partial_t \operatorname{rot} u^{(1)}|^2 + s |\nabla \operatorname{rot} H^{(1)}|^2 + s^2 |\nabla H^{(1)}|^2 + s^3 |\operatorname{rot} u^{(1)}|^2 + s^4 |H^{(1)}|^2 \right) e^{2s\varphi} dxdt \\ & \leq C \int_Q (|\partial_t \operatorname{rot} F|^2 + |\operatorname{rot} F|^2) e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_6^2 + Cs^4 e^{2s\mu_1} M^2 \end{aligned}$$

for all sufficiently large $s \geq 1$. We combine this inequality with (3.37) to obtain

$$\begin{aligned} & \int_Q \left(\frac{1}{s} |\partial_t \operatorname{rot} u^{(1)}|^2 + s |\nabla \operatorname{rot} H^{(1)}|^2 + s |\nabla \operatorname{rot} H|^2 + s^2 |\nabla H^{(1)}|^2 + s^2 |\nabla H|^2 \right. \\ & \quad \left. + s^3 |\operatorname{rot} u^{(1)}|^2 + s^4 |H^{(1)}|^2 + s^4 |H|^2 \right) e^{2s\varphi} dxdt \\ & \leq C \int_Q (|\partial_t \operatorname{rot} F|^2 + |\operatorname{rot} F|^2) e^{2s\varphi} dxdt + Cs^4 e^{Cs} E_6^2 + Cs^4 e^{2s\mu_1} M^2. \end{aligned}$$

According to the assumption (1.4), we can estimate the first term on the right-hand side above and we further have

$$\int_Q \left(\frac{1}{s} |\partial_t \operatorname{rot} u^{(1)}|^2 + s |\nabla \operatorname{rot} H^{(1)}|^2 + s |\nabla \operatorname{rot} H|^2 + s^2 |\nabla H^{(1)}|^2 + s^2 |\nabla H|^2 \right)$$

$$\begin{aligned}
& + s^3 |\operatorname{rot} u^{(1)}|^2 + s^4 |H^{(1)}|^2 + s^4 |H|^2 \Big) e^{2s\varphi} dx dt \\
& \leq C \int_Q |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi} dx dt + Cs^4 e^{Cs} E_6^2 + Cs^4 e^{2s\mu_1} M^2
\end{aligned} \tag{3.39}$$

for all sufficiently large $s \geq 1$. By noting $v = \operatorname{rot} u$, we insert (3.39) into the right-hand side of (3.35) and we obtain

$$\begin{aligned}
\int_{\Omega} |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx & \leq C \int_Q |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi} dx dt \\
& + Cs^4 e^{2s\mu_1} M^2 + Cs^4 e^{Cs} D_3^2
\end{aligned} \tag{3.40}$$

for all sufficiently large $s \geq 1$. Here we noted that $E_5^2 + E_6^2 \leq CD_3^2$ where D_3 is the norm of the data that introduced in the theorem.

Third Step: Completion of the proof

As we mentioned in the fourth step in Subsection 3.1, thanks to the choice of the weight function φ , we find

$$\int_Q |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi} dx dt = o(1) \int_{\Omega} |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx$$

as $s \rightarrow \infty$. Thus, by taking s large enough, we can absorb the first term on the right-hand side of (3.40) into the left-hand side. This yields

$$\int_{\Omega} |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \leq Cs^4 e^{2s\mu_1} M^2 + Cs^4 e^{Cs} D_3^2 \tag{3.41}$$

for all s large enough. With the same argument and the same constant $\mu_0 > 0$ introduced in the last step in Subsection 3.2, we have the lower bound of the left-hand side of (3.41):

$$\int_{\Omega} |\operatorname{rot} F(x, t_0)|^2 e^{2s\varphi(x, t_0)} dx \geq e^{2s\mu_0} \|\operatorname{rot} F(\cdot, t_0)\|_{L^2(\Omega_0)}^2,$$

which implies

$$\|\operatorname{rot} F(\cdot, t_0)\|_{L^2(\Omega_0)}^2 \leq Cs^4 e^{-2s(\mu_0 - \mu_1)} M^2 + Cs^4 e^{Cs} D_3^2 \tag{3.42}$$

for all s large enough. Since we have $\mu_0 > \mu_1$ by taking $\beta > 0$ sufficiently large, we end up with the stability of Hölder type by minimizing the right-hand side of (3.42) with respect to s .

4 Concluding remarks

4.1 Linearization of the MHD equations

Let (u_i, p_i, H_i) satisfy the following MHD equations

$$\begin{cases} \partial_t u_i - \nu \Delta u_i + (u_i \cdot \nabla) u_i - \mu \operatorname{rot} H_i \times H_i + \nabla p_i = F_i, \\ \partial_t H_i - \frac{1}{\sigma \mu} \Delta H_i - \operatorname{rot}(u_i \times H_i) = 0, \\ \operatorname{div} u_i = 0, \quad \operatorname{div} H_i = 0 \end{cases}$$

for $i = 1, 2$. Then by taking the difference and setting $u = u_1 - u_2$, $p = p_1 - p_2$, $H = H_1 - H_2$ and $F = F_1 - F_2$, we have

$$\begin{cases} \partial_t u - \nu \Delta u + (u_1 \cdot \nabla) u + (u \cdot \nabla) u_2 - \mu(H_1 \cdot \nabla) H - \mu(H \cdot \nabla) H_2 \\ \quad + \frac{\mu}{2} \nabla((H_1 + H_2) \cdot H) + \nabla p = F, \\ \partial_t H - \frac{1}{\sigma \mu} \Delta H + (u_1 \cdot \nabla) H - (H \cdot \nabla) u_2 - (H_1 \cdot \nabla) u + (u \cdot \nabla) H_2 = 0, \\ \operatorname{div} u = 0, \quad \operatorname{div} H = 0, \end{cases}$$

which corresponds to the linearized MHD equations (1.1) with (1.2). For example, we let $A^{(1)} = u_1$, $A^{(2)} = u_2, \dots$

Therefore, in order to apply the main results in this article to the (nonlinear) MHD equations, we need some a priori regularity of the solutions (u_i, p_i, H_i) , $i = 1, 2$. Although the existence of the solutions to the initial boundary value problems for the MHD equations is a serious issue with the regularity, here we focus on the inverse problems and we assume the existence of such solutions (u_i, p_i, H_i) , $i = 1, 2$ with sufficient regularity.

4.2 Global Lipschitz stability

As for the Carleman estimates for the parabolic equations including the Navier-Stokes equations and our MHD equations, one can have Carleman estimates according to each of the weight functions in the following forms:

$$(i) \quad \varphi(x, t) := e^{\lambda(d(x) - \beta(t - t_0)^2)}, \quad (4.1)$$

which is regular in x, t .

$$(ii) \quad \alpha(x, t) := \frac{e^{\lambda d(x)} - e^{2\lambda \|d\|_{C(\overline{\Omega})}}}{\ell(t)}, \quad (4.2)$$

which is singular at $t = 0, T$.

Usually one uses the Carleman estimate with the regular weight function (4.1) to prove a local Hölder stability estimate for an inverse problem. On the other hand,

one can apply the Carleman estimate with the singular weight function (4.2) to prove a global Lipschitz stability for an inverse problem provided that the boundary condition on $\partial\Omega \times (0, T)$ is given.

In this article, it is clear that we employed the Carleman estimates with (4.1) and proved the Hölder stability estimates for some inverse source problems. If we assume suitable boundary conditions, for example,

$$u(x, t) = 0, \quad H(x, t) = 0, \quad (x, t) \in \partial\Omega \times (0, T),$$

in addition to (1.1), we can easily combine the arguments in Choulli, Imanuvilov, Puel and Yamamoto [8] or Imanuvilov and Yamamoto [19], in which the authors discussed similar inverse source problems for the linearized Navier-Stokes equations by the Carleman estimates with (4.2), with what we have done in this article to derive the global Lipschitz stability estimates for the inverse source problems.

However, the boundary condition on the magnetic field intensity $H(x, t) = 0$, $(x, t) \in \partial\Omega \times (0, T)$ is not physical. As a possible solution, instead of the boundary conditions, we measure u and H on the full boundary. Then immediately we can prove the related global Lipschitz stability which is parallel to Theorem 1.1 as follows:

$$\begin{aligned} \|F(\cdot, t_0)\|_{L^2(\Omega)} &\leq C(\|u\|_{H^2(I; H^1(\partial\Omega))} + \|\partial_n u\|_{H^1(I; L^2(\partial\Omega))} + \|H\|_{H^2(I; H^1(\partial\Omega))} \\ &\quad + \|\partial_n H\|_{H^1(I; L^2(\partial\Omega))} + \|p\|_{H^1(I; H^{\frac{1}{2}}(\partial\Omega))} + \|u(\cdot, t_0)\|_{H^2(\Omega)} + \|\nabla p(\cdot, t_0)\|_{L^2(\Omega)}). \end{aligned}$$

Comparing to the inverse source problem for the parabolic equations, the data on the right-hand side above are too many since we need additionally $\partial_n u, \partial_n H$ on the boundary. The global Lipschitz stability for the inverse source problems for the MHD equations by less data still remains open.

Acknowledgments

The first author thanks the Leading Graduate Course for Frontiers of Mathematical Sciences and Physics (FMSP, The University of Tokyo), and was supported by Grant-in-Aid for JSPS Fellows 20F20319 of Japan Society for the Promotion of Science (JSPS). The second author was supported by Grant-in-Aid for Scientific Research (A) 20H00117 of JSPS, the National Natural Science Foundation of China (Nos. 11771270, 91730303) and the RUDN University Strategic Academic Leadership Program.

References

- [1] R. A. Adams and J. F. Fournier, “Sobolev Spaces”, Pure and Applied Mathematics Series Vol. 140, Academic Press, 2003.

- [2] M. Bellassoued, O. Y. Imanuvilov and M. Yamamoto, Inverse Problem of Determining the Density and Two Lamé Coefficients by Boundary Data, *SIAM J. Math. Anal.* 40 (2008), 238-265.
- [3] M. Bellassoued, O. Y. Imanuvilov and M. Yamamoto, Carleman estimate for the Navier-Stokes equations and an application to a lateral Cauchy problem, *Inverse Problems* 32 (2016) 025001.
- [4] A. L. Bukhgeim and M. V. Klibanov, Global Uniqueness of a class of multidimensional inverse problems, *Sov. Math. Dokl.* 24 (1981), 244-247.
- [5] M. Bellassoued and M. Yamamoto, “Carleman Estimates and Applications to Inverse Problems for Hyperbolic Systems”, Springer-Japan, Tokyo, 2017.
- [6] P. Cannarsa, G. Floridia, M. Yamamoto, Observability inequalities for transport equations through Carleman estimates, Springer INdAM series, Vol. 32 (2019). Doi:10.1007/978-3-030-17949-6, <https://arxiv.org/abs/1807.05005>
- [7] P. Cannarsa, G. Floridia, F. Gölgeleyen, M. Yamamoto, Inverse coefficient problems for a transport equation by local Carleman estimate, *Inverse Problems* 35 (2019) 105013. Doi:10.1088/1361-6420/ab1c69, <http://arxiv.org/abs/1902.06355>
- [8] M. Choulli, O. Y. Imanuvilov, J. -P. Puel and M. Yamamoto, Inverse source problem for linearized Navier-Stokes equations with data in arbitrary subdomain, *Applicable Analysis* 92 (2013), 2127-2143.
- [9] A. V. Fursikov and O. Y. Imanuvilov, “Controllability of Evolution Equations”, Seoul National University, Korea, 1996.
- [10] X. Huang, Inverse coefficient problem for a magnetohydrodynamics system by Carleman estimates, *Applicable Analysis* 100 (2021), 1010-1038.
- [11] X. Huang, O. Y. Imanuvilov and M. Yamamoto, Stability for inverse source problems by Carleman estimates, *Inverse Problems* 36 (2020) 125006.
- [12] T. Havârneanu, C. Popa and S. S. Sritharan, Exact internal controllability for the magnetohydrodynamic equations in multi-connected domains, *Adv. Differential Equations* 11 (2006), 893-929.
- [13] T. Havârneanu, C. Popa and S. S. Sritharan, Exact internal controllability for the two-dimensional magnetohydrodynamic equations, *SIAM J. Control Optim.* 46 (2007), 1802-1830.
- [14] O. Y. Imanuvilov, Controllability of parabolic equations, *Sbornik Math.* 186 (1995), 879-900.
- [15] O. Y. Imanuvilov, L. Lorenzi and M. Yamamoto, Carleman estimate for the Navier-Stokes equations and applications, *arXiv:2107.04495 [math.AP]*.

- [16] O. Y. Imanuvilov and J. -P. Puel, Global Carleman estimates for weak solutions of elliptic nonhomogeneous Dirichlet problems, *IMRN* 16 (2003), 883-913.
- [17] O. Y. Imanuvilov and M. Yamamoto, Lipschitz stability in inverse parabolic problems by Carleman estimate, *Inverse Problems* 14 (1998), 1229-1245.
- [18] O. Y. Imanuvilov and M. Yamamoto, Global Lipschitz stability in an inverse hyperbolic problem by interior observations, *Inverse Problems* 17 (2001), 717-728.
- [19] O. Y. Imanuvilov and M. Yamamoto, Global Lipschitz stability for an inverse source problem for the Navier-Stokes equations, *arXiv:2107.04514 [math.AP]*.
- [20] M. V. Klibanov, Inverse problems and Carleman estimates, *Inverse Problems* 8 (1992), 575-596.
- [21] H. Li, Global strong solution to the three dimensional nonhomogeneous incompressible magnetohydrodynamic equations with density-dependent viscosity and resistivity, *Mathematical Methods in the Applied Sciences* 41 (2018), 3062-3092.
- [22] T. Li and T. Qin, “Physics and Partial Differential Equations”, Higher Education Press, Beijing, Vol. 1, 2013.
- [23] O. A. Ladyzhenskaya and V. A. Solonnikov, Unique solvability of an initial- and boundary-value problem for viscous incompressible nonhomogeneous fluids, *Journal of Soviet Mathematics* 9 (1978), 697-749.
- [24] M. Yamamoto, Carleman estimates for parabolic equations and applications, *Inverse Problems* 25 (2009) 123013.
- [25] X. Zhou, On uniqueness theorem of a vector function, *Progress In Electromagnetics Research* 65 (2006), 93-102.

E-mail address: huangxc@ms.u-tokyo.ac.jp

E-mail address: myama@ms.u-tokyo.ac.jp