

Existence of ground state solutions of Nehari-Pankov type to Schrödinger systems *

TANG XianHua¹ and LIN XiaoYan²

¹ School of Mathematics and Statistics, Central South University,
Changsha, Hunan 410083, P.R.China

E-mail: tangxh@mail.csu.edu.cn

² Department of Mathematics, Huaihua University,
Huaihua, Hunan 418008, P.R.China

E-mail: xiaoyanlin98@hotmail.com

Abstract: This paper is dedicated to studying the following elliptic system of Hamiltonian type:

$$\begin{cases} -\varepsilon^2 \Delta u + u + V(x)v = Q(x)F_v(u, v), & x \in \mathbb{R}^N, \\ -\varepsilon^2 \Delta v + v + V(x)u = Q(x)F_u(u, v), & x \in \mathbb{R}^N, \\ |u(x)| + |v(x)| \rightarrow 0, & \text{as } |x| \rightarrow \infty, \end{cases}$$

where $N \geq 3$, $V, Q \in \mathcal{C}(\mathbb{R}^N, \mathbb{R})$, $V(x)$ is allowed to be sign-changing and $\inf Q > 0$, and $F \in \mathcal{C}^1(\mathbb{R}^2, \mathbb{R})$ is superquadratic at both 0 and infinity but subcritical. Instead of the reduction approach used in [Calc Var PDE, 2014, 51: 725-760], we develop a more direct approach – non-Nehari manifold approach to obtain stronger conclusions but under weaker assumptions than these in [Calc Var PDE, 2014, 51: 725-760]. We can find an $\varepsilon_0 > 0$ which is determined by terms of N, V, Q and F , then we prove the existence of a ground state solution of Nehari-Pankov type to the coupled system for all $\varepsilon \in (0, \varepsilon_0]$.

Keywords: Hamiltonian elliptic system, Ground state solutions of Nehari-Pankov type, Strongly indefinite functionals.

2000 Mathematics Subject Classification. 35J50; 35E05

1 Introduction

In this paper we study standing waves for the following system of time-dependent nonlinear

*This work is partially supported by the NNSF (No: 11571370, 11471137) of China. This paper has been accepted for publication in SCIENCE CHINA Mathematics.

Schrödinger equations

$$\begin{cases} i\hbar \frac{\partial \phi_1}{\partial t} + \frac{\hbar^2}{2m} \Delta \phi_1 + \phi_1 + f(x, \phi) \phi_2 = 0, \\ i\hbar \frac{\partial \phi_2}{\partial t} + \frac{\hbar^2}{2m} \Delta \phi_2 + \phi_2 + f(x, \phi) \phi_1 = 0, \end{cases} \quad (1.1)$$

where m is the mass of a particle, \hbar is the Planck constant, $\phi = (\phi_1, \phi_2)$, $\phi_1(t, x)$ and $\phi_2(t, x)$ are the complex valued envelope functions. Suppose that $f(x, e^{i\theta}\phi) = f(x, \phi)$ for $\theta \in [0, 2\pi]$. We will look for standing waves of the form

$$\phi_1(t, x) = e^{i\omega t} u(x), \quad \phi_2(t, x) = e^{i\omega t} v(x),$$

which propagate without changing their shape and thus have a soliton-like behavior. System (1.1) arises quite naturally in nonlinear optics and Bose-Einstein condensates (see [2, 17, 27] and the references therein). In general, the above coupled nonlinear Schrödinger system leads to the elliptic system of Hamiltonian form

$$\begin{cases} -\varepsilon^2 \Delta u + u = H_v(x, u, v), & x \in \mathbb{R}^N, \\ -\varepsilon^2 \Delta v + v = H_u(x, u, v), & x \in \mathbb{R}^N, \\ |u(x)| + |v(x)| \rightarrow 0, & \text{as } |x| \rightarrow \infty, \end{cases} \quad (1.2)$$

where $N \geq 3$, $H \in C^1(\mathbb{R}^N \times \mathbb{R}^2, \mathbb{R})$ and $\varepsilon > 0$ is a small parameter. The study of the systems similar to (1.2) has only begun quite recently. When $\varepsilon = 1$, it was considered recently in some works [6, 9, 15, 16, 34–39]. For a similar problem on a bounded domain we refer the reader to [5, 7, 9, 14] and the references therein. For a survey on this direction see [8, 25].

For the case $\varepsilon > 0$ is a small parameter, there are some recent works considering the existence of solutions; see for instance [3, 12, 27] and the references therein. In contrast with the case $\varepsilon = 1$, except for the difficulties that the lack of the compactness of the Sobolev embedding and the energy functional is strongly indefinite, no uniqueness results seem to be known for the “limit problem” and this is in some cases a crucial assumption in the single equation case. So asymptotic analysis of solutions with respect to small $\varepsilon > 0$ has been very recently performed; see for example [3, 4, 11, 12, 20, 22–24, 27] and their references. Except for [11, 12, 27], most of the above works considered the case that $H(x, u, v) = F(u) + G(v)$. In particular, in [4], the authors obtained the existence of positive solutions which concentrate on the boundary of Ω for an elliptic system with zero Neumann boundary condition on a bounded domain Ω (see also [20]). In [22], Ramos and Soares considered the following problem

$$\begin{cases} -\varepsilon^2 \Delta u + V(x)u = g(v), & x \in \Omega, \\ -\varepsilon^2 \Delta v + V(x)v = f(u), & x \in \Omega, \\ u, v \in H^1(\mathbb{R}^N), \end{cases} \quad (1.3)$$

where Ω is a domain of \mathbb{R}^N , $V \in C(\mathbb{R}^N, \mathbb{R})$ satisfies $0 < V(0) = \min V(x) < \liminf_{|x| \rightarrow \infty} V(x) \in (0, \infty]$, $f(u)$ and $g(v)$ are power type functions, superlinear but subcritical at infinity. The authors established the existence of positive solutions which concentrate, as $\varepsilon \rightarrow 0$, at a prescribed

finite number of local minimum points (possibly degenerate) of the potential V . Different from those discussed in [3, 22, 23], Ding, Lee and Zhao [11] dealt with existence and concentration phenomena of the ground state solutions to the following subcritical problem

$$\begin{cases} -\varepsilon^2 \Delta u + u + V(x)v = Q(x)g(|z|)v, & x \in \mathbb{R}^N, \\ -\varepsilon^2 \Delta v + v + V(x)u = Q(x)g(|z|)u, & x \in \mathbb{R}^N, \\ |u(x)| + |v(x)| \rightarrow 0, & \text{as } |x| \rightarrow \infty, \end{cases} \quad (1.4)$$

where $z := (u, v)$, $V, Q \in \mathcal{C}^1(\mathbb{R}^N, \mathbb{R})$ and $g \in \mathcal{C}^1(\mathbb{R}^+, \mathbb{R}^+)$. Since the energy functional Φ_ε associated with system (1.4) is strongly indefinite, to overcome this difficulty, as in [1] (see also [22] and [23]), the authors constructed a reduced functional \mathcal{R}_ε whose critical points are in one to one to critical points of Φ_ε , which was first proposed in [1]. With the help of the Nehari manifold of Φ_ε , an important information of the least energy c_ε was obtained. By estimating the asymptotic behavior of c_ε as $\varepsilon \rightarrow 0$, they proved c_ε is attained for sufficiently small $\varepsilon > 0$. In order to state their results, some notations and assumptions are required. Set

$$V_{\min} := \min_{x \in \mathbb{R}^N} V(x), \quad V_{\max} := \max_{x \in \mathbb{R}^N} V(x), \quad \mathcal{V} := \{x \in \mathbb{R}^N : V(x) = V_{\min}\};$$

$$Q_{\min} := \min_{x \in \mathbb{R}^N} Q(x), \quad Q_{\max} := \max_{x \in \mathbb{R}^N} Q(x), \quad \mathcal{Q} := \{x \in \mathbb{R}^N : Q(x) = Q_{\max}\};$$

$$V_\infty := \liminf_{|x| \rightarrow \infty} V(x), \quad Q_\infty := \limsup_{|x| \rightarrow \infty} Q(x);$$

$$\mathcal{A}_v := \{x \in \mathcal{V} : Q(x) = Q(x_v)\} \cup \{x \notin \mathcal{V} : Q(x) > Q(x_v)\}$$

and

$$\mathcal{A}_q := \{x \in \mathcal{Q} : V(x) = V(x_q)\} \cup \{x \notin \mathcal{Q} : V(x) < V(x_q)\}.$$

Furthermore, the following assumptions are required:

(A0) $V, Q \in \mathcal{C}^1(\mathbb{R}^N, \mathbb{R})$, $\|V\|_\infty < 1$, $0 < Q_{\min} \leq Q_{\max} < \infty$;

(A1) $V_{\min} < V_\infty$, and there exist $x_v \in \mathcal{V}$ and $R > 0$ such that

$$Q(x_v) = \max_{y \in \mathcal{V}} Q(y) \geq Q(x), \quad \forall |x| \geq R;$$

(A2) $Q_{\max} > Q_\infty$, and there exist $x_q \in \mathcal{Q}$ and $R > 0$ such that

$$V(x_q) = \min_{y \in \mathcal{Q}} V(y) \leq V(x), \quad \forall |x| \geq R;$$

(G1) $g \in \mathcal{C}^1(\mathbb{R}^+, \mathbb{R}^+)$, $g(0) = 0$ and $g'(s) \geq 0$ for $s > 0$, where $\mathbb{R}^+ := [0, \infty)$;

(G2) there exist $C > 0$ and $p \in (2, 2^*)$ such that $|g(s)| \leq C(1 + s^{p-2})$ for all $s \in \mathbb{R}^+$;

(G3) there exists $\mu > 2$ such that $g(s)s^2 \geq \mu \int_0^s g(t)t dt > 0$ if $s > 0$.

In [11], they proved the following theorem.

Theorem 1.1. ([11, Theorem 1]) *Let (A0), (G1), (G2) and (G3) be satisfied. Suppose that (A1) or (A2) is satisfied. Then for sufficiently small $\varepsilon > 0$, (1.4) has a least energy solution $\hat{z}_\varepsilon = (\hat{u}_\varepsilon, \hat{v}_\varepsilon)$.*

Theorem 1.1 is very interesting. In its proof, many new tricks were used to overcome the difficulties caused by the strong indefiniteness of the energy functional Φ_ε associated with system (1.4). We point that the regularity assumptions $V, Q \in \mathcal{C}^1$ and $g \in \mathcal{C}^1$ are very crucial in [11], which seem to be necessary when the reduction method is used. Motivated by the works [11], in this paper, we further study the existence of the ground state solutions of Nehari-Pankov type to the following more general problem

$$\begin{cases} -\varepsilon^2 \Delta u + u + V(x)v = Q(x)F_v(u, v), & x \in \mathbb{R}^N, \\ -\varepsilon^2 \Delta v + v + V(x)u = Q(x)F_u(u, v), & x \in \mathbb{R}^N, \\ |u(x)| + |v(x)| \rightarrow 0, & \text{as } |x| \rightarrow \infty, \end{cases} \quad (1.5)$$

where $V, Q \in \mathcal{C}(\mathbb{R}^N, \mathbb{R})$ and $F \in \mathcal{C}^1(\mathbb{R}^2, \mathbb{R})$. Instead of the reduction method used in [11], we will use a more direct approach – non-Nehari manifold approach which was first proposed in [29] for a single Schrödinger equation (see also [30, 31]). The main idea of this approach is to construct a minimizing Cerami sequence for the energy functional outside Nehari-Pankov manifold by using the diagonal method, which is completely different from that of Szulkin and Weth [28]. This approach is valid when finding a ground state solution of Nehari-Pankov type.

We will obtain stronger conclusions on existence of the ground state solutions of Nehari-Pankov type to (1.5) for small $\varepsilon > 0$ but under weaker assumptions than these in [11]. Roughly speaking, we can find an $\varepsilon_0 > 0$ which is determined by terms of N, V, Q and F , then we prove the existence of a ground state solutions of Nehari-Pankov type to (1.5) for all $\varepsilon \in (0, \varepsilon_0]$. In particular, we only need $V, Q \in \mathcal{C}(\mathbb{R}^N, \mathbb{R})$ and $F \in \mathcal{C}^1(\mathbb{R}^2, \mathbb{R})$. To the best of our knowledge, there seems to be no similar results in literature.

To state our theorems accurately, we set

$$\mathcal{ND}_0 = \left\{ h \in \mathcal{C}(\mathbb{R}^+, \mathbb{R}^+) : \begin{cases} h(0) = 0 \text{ and } h(s) \text{ is nondecreasing on } \mathbb{R}^+, \\ \text{there exist constants } p \in (2, 2^*) \text{ and } c_0 > 0 \text{ such that} \\ |h(s)| \leq c_0(1 + |s|^{p-2}), \quad \forall s \geq 0. \end{cases} \right\} \quad (1.6)$$

Furthermore, we make the following assumptions.

(V0) $V, Q \in \mathcal{C}(\mathbb{R}^N, \mathbb{R})$, $\|V\|_\infty \leq \frac{2\eta ab}{a^2 + b^2}$ and $0 < Q_{\min} \leq Q_{\max} < \infty$, where $a, b > 0$ and $\eta \in (0, 1)$;

(V1) $V_{\min} < V_\infty$, and there exist $x_v \in \mathcal{V}$ and $R > 0$ such that

$$Q(x_v) \geq Q(x), \quad \forall |x| \geq R;$$

(V2) $Q_{\max} > Q_{\infty}$, and there exist $x_q \in \mathcal{Q}$ and $R > 0$ such that

$$V(x_q) \leq V(x), \quad \forall |x| \geq R;$$

(F1) there exist $g_i, h_j \in \mathcal{ND}_0$, $\alpha_i, \beta_i, \alpha'_j, \beta'_j \in \mathbb{R}$ with $\alpha_i^2 + \beta_i^2 \neq 0$ and $\alpha'_j > \beta'^2_j$, $i = 1, 2, \dots, k; j = 1, 2, \dots, l$, such that

$$F(u, v) = \sum_{i=1}^k \int_0^{|\alpha_i u + \beta_i v|} g_i(s) s ds + \sum_{j=1}^l \int_0^{\sqrt{u^2 + 2\beta'_j uv + \alpha'^2_j v^2}} h_j(s) s ds;$$

(F2) $\lim_{|au+bv| \rightarrow \infty} \frac{|F(u, v)|}{|au+bv|^2} = \infty$;

(F3) there exist $\mathcal{C}_0 > 0, T_0 > 0, \mu > 2$ and $F_0 \in \mathcal{C}(\mathbb{R}^2, \mathbb{R})$ with $F_0(u, v) > 0$ if $au + bv \neq 0$, such that

$$F(tz) \geq \mathcal{C}_0 t^\mu F_0(z), \quad \forall z \in \mathbb{R}^2, t \geq T_0.$$

Remark 1.2. It is clear that (i) (F3) is weaker than (AR)-condition: there exists $\mu > 2$ such that $F_z(z) \cdot z \geq \mu F(z) > 0$ for $z \neq 0$; (ii) (F2) is also weaker than the common super-quadratic condition (SQ): $\lim_{|z| \rightarrow \infty} \frac{|F(z)|}{|z|^2} = \infty$; (iii) Let $F(z) = \int_0^{|z|} g(s) s ds$. Then (G1)-(G3) imply (F1)-(F3); (iv) Let $Q(x) \equiv 1$ and $F(z) = \int_0^{|u|} g(s) s ds + \int_0^{|v|} h(s) s ds$, then problem (1.5) reduces to (1.3), which was studied in [22]. Moreover, the assumptions in [22, (H)] also imply (F1)-(F3).

Before presenting our results, we give three nonlinear examples to illustrate the above assumptions.

Example 1.3. Let $F(z) = |au + bv|^\mu$, where $\mu \in (2, 2^*)$ and $a, b > 0$ with $\|V\|_\infty < \frac{2ab}{a^2+b^2}$. It is easy to see that $F(z)$ satisfies (F1)-(F3) with $F_0 = F$, but not (AR).

Example 1.4. Let $F(z) = |au + bv|^\mu + (u^2 + uv + v^2) \ln(1 + u^2 + uv + v^2)$, where $\mu \in (2, 2^*)$ and $a, b > 0$ with $\|V\|_\infty < \frac{2ab}{a^2+b^2}$. It is easy to see that $F(z)$ satisfies (F1)-(F3) with $F_0(u, v) = |au + bv|^\mu$, but not (AR).

Example 1.5. Let $F(z) = |2u + v|^\mu + |u + 2v|^\mu$, where $\mu \in (2, 2^*)$. It is easy to see that $F(z)$ satisfies (F1)-(F3) with $a = b = 1$ and $F_0 = F$.

Let $x_m = x_v$ if (V1) holds, or $x_m = x_q$ if (V2) holds. Replacing $u(\varepsilon x + x_m)$ and $v(\varepsilon x + x_m)$ by $u(x)$ and $v(x)$, respectively, then system (1.5) is equivalent to

$$\begin{cases} -\Delta u + u + V(\varepsilon x + x_m)v = Q(\varepsilon x + x_m)F_v(u, v), & x \in \mathbb{R}^N, \\ -\Delta v + v + V(\varepsilon x + x_m)u = Q(\varepsilon x + x_m)F_u(u, v), & x \in \mathbb{R}^N, \\ |u(x)| + |v(x)| \rightarrow 0, & \text{as } |x| \rightarrow \infty. \end{cases} \quad (1.7)$$

Let $E = H^1(\mathbb{R}^N) \times H^1(\mathbb{R}^N)$. Then E is a Hilbert space with the standard inner product

$$(z_1, z_2)_{H^1(\mathbb{R}^N)} = (u_1, u_2)_{H^1(\mathbb{R}^N)} + (v_1, v_2)_{H^1(\mathbb{R}^N)}, \quad \forall z_i = (u_i, v_i) \in E, \quad i = 1, 2,$$

and the corresponding norm

$$\|z\|_{H^1(\mathbb{R}^N)} = \left(\|u\|_{H^1(\mathbb{R}^N)}^2 + \|v\|_{H^1(\mathbb{R}^N)}^2 \right)^{1/2}, \quad \forall z = (u, v) \in E.$$

Let $E = E^- \oplus E^+$ be an orthogonal decomposition, see Section 2. Define a functional

$$\Phi_\varepsilon(z) = \int_{\mathbb{R}^N} \left[\nabla u \cdot \nabla v + uv + \frac{1}{2} V(\varepsilon x + x_m) |z|^2 \right] dx - \int_{\mathbb{R}^N} Q(\varepsilon x + x_m) F(z) dx \quad (1.8)$$

for all $z = (u, v) \in E$. Under assumptions (V0), (F1) and (F2), $\Phi_\varepsilon \in \mathcal{C}^1(E, \mathbb{R})$ and

$$\begin{aligned} \langle \Phi'_\varepsilon(z), \varphi \rangle &= \int_{\mathbb{R}^N} [\nabla u \cdot \nabla \psi + \nabla v \cdot \nabla \phi + (u\psi + v\phi) + V(\varepsilon x + x_m) z \cdot \varphi] dx \\ &\quad - \int_{\mathbb{R}^N} Q(\varepsilon x + x_m) F_z(z) \cdot \varphi dx, \quad \forall z = (u, v), \quad \varphi = (\phi, \psi) \in E. \end{aligned} \quad (1.9)$$

Let

$$\mathcal{N}_\varepsilon^- = \{z \in E \setminus E^- : \langle \Phi'_\varepsilon(z), z \rangle = \langle \Phi'_\varepsilon(z), \zeta \rangle = 0, \quad \forall \zeta \in E^-\}. \quad (1.10)$$

$\mathcal{N}_\varepsilon^-$ was first introduced by Pankov [19], which is a subset of the Nehari manifold

$$\mathcal{N}_\varepsilon = \{z \in E \setminus \{0\} : \langle \Phi'_\varepsilon(z), z \rangle = 0\}. \quad (1.11)$$

We are now in a position to state the first main result of this paper.

Theorem 1.6. *Assume that V , Q and F satisfy (V0), (V1) and (F1)-(F3). Then there exists an $\varepsilon_0 > 0$ such problem (1.5) has a nontrivial solution $\hat{z}_\varepsilon = (\hat{u}_\varepsilon, \hat{v}_\varepsilon) \in \mathcal{N}_\varepsilon^-$ with $\Phi_\varepsilon(z_\varepsilon) = \inf_{\mathcal{N}_\varepsilon^-} \Phi_\varepsilon > 0$ for $\varepsilon \in (0, \varepsilon_0]$, where $z_\varepsilon(x) = \hat{z}_\varepsilon(\varepsilon x + x_v)$. If (V1) is replaced by (V2), then the above conclusion remains true by replacing x_v with x_q .*

The “limit problem” associated to (1.7) is an autonomous system

$$\begin{cases} -\Delta u + u + V(x_m)v = Q(x_m)F_v(u, v), & x \in \mathbb{R}^N, \\ -\Delta v + v + V(x_m)u = Q(x_m)F_u(u, v), & x \in \mathbb{R}^N. \end{cases} \quad (1.12)$$

We will prove that the least energy $c_\varepsilon := \inf_{\mathcal{N}_\varepsilon^-} \Phi_\varepsilon$ is attained for $\varepsilon \in (0, \varepsilon_0]$ by comparing with c_ε and the least energy c_0 associated with “limit problem” (1.12). Therefore, it is very crucial if c_0 can be attained, i.e. if (1.12) has a solution at which Φ_0 has the least energy c_0 on \mathcal{N}_0^- . Prior to this, we consider the following more general periodic system

$$\begin{cases} -\Delta u + V_1(x)u + V_2(x)v = W_v(x, u, v), & x \in \mathbb{R}^N, \\ -\Delta v + V_1(x)v + V_2(x)u = W_u(x, u, v), & x \in \mathbb{R}^N, \\ u, v \in H^1(\mathbb{R}^N), \end{cases} \quad (1.13)$$

where $N \geq 3$, $V_1, V_2 : \mathbb{R}^N \rightarrow \mathbb{R}$ and $W : \mathbb{R}^N \times \mathbb{R}^2 \rightarrow \mathbb{R}$. More precisely, we make the following assumptions.

(V0') $V_1, V_2 \in \mathcal{C}(\mathbb{R}^N)$ and satisfy

$$|V_2(x)| \leq \frac{2\eta ab}{a^2 + b^2} V_1(x), \quad 0 < \inf_{x \in \mathbb{R}^N} V_1(x) \leq \sup_{x \in \mathbb{R}^N} V_1(x) < \infty, \quad (1.14)$$

where $a, b > 0$ and $\eta \in (0, 1)$;

(V1') $V_1(x)$ and $V_2(x)$ are 1-periodic in each of x_1, x_2, \dots, x_N ;

(W0) $W \in \mathcal{C}(\mathbb{R}^N \times \mathbb{R}^2, \mathbb{R}^+)$, $W(x, z)$ is continuously differentiable on $z \in \mathbb{R}^2$ for every $x \in \mathbb{R}^N$, and there exist constants $p \in (2, 2^*)$ and $C_0 > 0$ such that

$$|W_z(x, z)| \leq C_0 (1 + |z|^{p-1}), \quad \forall (x, z) \in \mathbb{R}^N \times \mathbb{R}^2; \quad (1.15)$$

(W1) $W_z(x, z) = o(|z|)$, as $|z| \rightarrow 0$, uniformly in $x \in \mathbb{R}^N$;

(W2) $\lim_{|au+bv| \rightarrow \infty} \frac{|W(x, u, v)|}{|au+bv|^2} = \infty$, a.e. $x \in \mathbb{R}^N$;

(W2') $\lim_{|au+bv| \rightarrow \infty} \frac{|W(x, u, v)|}{|au+bv|^2} = \infty$, uniformly in $x \in \mathbb{R}^N$;

(W3) $W(x, z)$ is 1-periodic in each of x_1, x_2, \dots, x_N ;

(W4) for all $\theta \geq 0$, $z, \zeta \in \mathbb{R}^2$, there holds

$$\frac{1 - \theta^2}{2} \nabla W(x, z) \cdot z - \theta \nabla W(x, z) \cdot \zeta + W(x, \theta z + \zeta) - W(x, z) \geq 0.$$

Observe that, the natural functional associated with (1.13) is given by

$$\Phi(z) = \int_{\mathbb{R}^N} \left[\nabla u \cdot \nabla v + V_1(x)uv + \frac{1}{2} V_2(x)|z|^2 \right] dx - \int_{\mathbb{R}^N} W(x, z) dx, \quad (1.16)$$

for all $z = (u, v) \in E$. Furthermore, under assumptions (V0'), (W0) and (W1), $\Phi \in \mathcal{C}^1(E, \mathbb{R})$ and

$$\begin{aligned} \langle \Phi'(z), \varphi \rangle &= \int_{\mathbb{R}^N} [\nabla u \cdot \nabla \psi + \nabla v \cdot \nabla \phi + V_1(x)(u\psi + v\phi) + V_2(x)z \cdot \varphi] dx \\ &\quad - \int_{\mathbb{R}^N} W_z(x, z) \cdot \varphi dx, \quad \forall z = (u, v), \varphi = (\phi, \psi) \in E. \end{aligned} \quad (1.17)$$

Let

$$\mathcal{N}^- = \{z \in E \setminus E^- : \langle \Phi'(z), z \rangle = \langle \Phi'(z), \zeta \rangle = 0, \forall \zeta \in E^-\}. \quad (1.18)$$

For system (1.13), we obtain the following existence theorem on the ground state solutions of Nehari-Pankov type.

Theorem 1.8. *Assume that V and W satisfy (V0'), (V1') and (W0)-(W4). Then problem (1.13) has a solution $z^* \in \mathcal{N}^-$ such that $\Phi(z^*) = \inf_{\mathcal{N}^-} \Phi > 0$.*

However, it is not easy to check assumption (W4). Next, we give several classes functions satisfying (W4). Prior to this, we define one set as follows:

$$\mathcal{ND} = \left\{ h \in \mathcal{C}(\mathbb{R}^N \times \mathbb{R}^+, \mathbb{R}^+) : \begin{cases} h(x, t) \text{ is 1-periodic in each of } x_1, x_2, \dots, x_N \text{ and} \\ \text{is nondecreasing in } t \in [0, \infty) \text{ for every } x \in \mathbb{R}^N; \\ h(x, 0) \equiv 0 \text{ for } x \in \mathbb{R}^N; \\ \text{there exist constants } p \in (2, 2^*) \text{ and } \mathcal{C}_1 > 0 \text{ such that} \\ |h(x, t)| \leq \mathcal{C}_1(1 + |t|^{p-2}), \quad \forall (x, t) \in \mathbb{R}^N \times \mathbb{R}. \end{cases} \right\} \quad (1.19)$$

Corollary 1.9. *Assume that V and W satisfy (V0'), (V1') and (W2), and that*

$$W(x, u, v) = \sum_{i=1}^k \int_0^{|\alpha_i u + \beta_i v|} g_i(x, s) ds + \sum_{j=1}^l \int_0^{\sqrt{u^2 + 2\beta'_j uv + \alpha'_j v^2}} h_j(x, s) ds,$$

where $\alpha_i, \beta_i, \alpha'_j, \beta'_j \in \mathbb{R}$ with $\alpha_i^2 + \beta_i^2 \neq 0$ and $\alpha'_j > \beta_j'^2$, $g_i, h_j \in \mathcal{ND}$, $i = 1, 2, \dots, k; j = 1, 2, \dots, l$. Then problem (1.13) has a solution $z^* \in \mathcal{N}^-$ such that $\Phi(z^*) = \inf_{\mathcal{N}^-} \Phi > 0$.

The paper is organized as follows. In the next section, we develop a functional setting to deal with (1.7) and (1.13). Section 3 is devoted to the proof of Theorem 1.8. In Section 4, we discuss the existence of ground state solutions of Nehari-Pankov type to (1.7).

2 Variational setting

Let $V_\varepsilon(x) := V(\varepsilon x + x_m)$ and $Q_\varepsilon(x) := Q(\varepsilon x + x_m)$. Then we can rewrite (1.7) as

$$\begin{cases} -\Delta u + u + V_\varepsilon(x)v = Q_\varepsilon(x)F_v(u, v), & x \in \mathbb{R}^N, \\ -\Delta v + v + V_\varepsilon(x)u = Q_\varepsilon(x)F_u(u, v), & x \in \mathbb{R}^N, \\ |u(x)| + |v(x)| \rightarrow 0, & \text{as } |x| \rightarrow \infty. \end{cases} \quad (2.1)$$

We will mainly deal with (2.1) instead of (1.7). Let

$$E^- = \left\{ \left(-\frac{u}{a}, \frac{u}{b} \right) : u \in H^1(\mathbb{R}^N) \right\}, \quad E^+ = \left\{ \left(\frac{u}{a}, \frac{u}{b} \right) : u \in H^1(\mathbb{R}^N) \right\}.$$

For any $z = (u, v) \in E$, set

$$z^- = \left(\frac{au - bv}{2a}, \frac{bv - au}{2b} \right), \quad z^+ = \left(\frac{au + bv}{2a}, \frac{au + bv}{2b} \right).$$

It is obvious that $z = z^- + z^+$. Now we define two new inner products on E

$$(z_1, z_2) = \int_{\mathbb{R}^N} [(\nabla z_1^+ \cdot \nabla z_2^+ + \nabla z_1^- \cdot \nabla z_2^-) + (z_1^+ \cdot z_2^+ + z_1^- \cdot z_2^-)] dx, \\ \forall z_i = (u_i, v_i) \in E, \quad i = 1, 2$$

and

$$(z_1, z_2)_\dagger = \int_{\mathbb{R}^N} [(\nabla z_1^+ \cdot \nabla z_2^+ + \nabla z_1^- \cdot \nabla z_2^-) + V_1(x) (z_1^+ \cdot z_2^+ + z_1^- \cdot z_2^-)] dx, \\ \forall z_i = (u_i, v_i) \in E, \quad i = 1, 2.$$

The corresponding norms are

$$\|z\| = \sqrt{(z, z)}, \quad \|z\|_\dagger = \sqrt{(z, z)_\dagger}, \quad \forall z = (u, v) \in E.$$

By virtue of (V0) and (V0'), it is easy to check that the norms $\|\cdot\|$, $\|\cdot\|_\dagger$ and $\|\cdot\|_{H^1(\mathbb{R}^N)}$ are equivalent on E . It is easy to see that z^- and z^+ are orthogonal with respect to the inner products (\cdot, \cdot) and $(\cdot, \cdot)_\dagger$. Thus we have $E = E^- \oplus E^+$. By a simple calculation, one can get that

$$\|z\|^2 = \int_{\mathbb{R}^N} [(|\nabla z^+|^2 + |\nabla z^-|^2) + (|z^+|^2 + |z^-|^2)] dx, \quad \forall z \in E, \quad (2.2)$$

$$\|z\|_\dagger^2 = \int_{\mathbb{R}^N} [(|\nabla z^+|^2 + |\nabla z^-|^2) + V_1(x) (|z^+|^2 + |z^-|^2)] dx, \quad \forall z \in E, \quad (2.3)$$

$$\frac{ab}{a^2 + b^2} (\|z^+\|^2 - \|z^-\|^2) = \int_{\mathbb{R}^N} (\nabla u \cdot \nabla v + uv) dx, \quad \forall z = (u, v) \in E$$

and

$$\frac{ab}{a^2 + b^2} (\|z^+\|_\dagger^2 - \|z^-\|_\dagger^2) = \int_{\mathbb{R}^N} [\nabla u \cdot \nabla v + V_1(x)uv] dx, \quad \forall z = (u, v) \in E.$$

Therefore, the functionals Φ_ε defined by (1.8) and Φ by (1.16) can be rewritten

$$\Phi_\varepsilon(z) = \frac{ab}{a^2 + b^2} (\|z^+\|^2 - \|z^-\|^2) + \frac{1}{2} \int_{\mathbb{R}^N} V_\varepsilon(x) |z|^2 dx - \int_{\mathbb{R}^N} Q_\varepsilon(x) F(z) dx, \quad \forall z \in E \quad (2.4)$$

and

$$\Phi(z) = \frac{ab}{a^2 + b^2} (\|z^+\|_\dagger^2 - \|z^-\|_\dagger^2) + \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |z|^2 dx - \int_{\mathbb{R}^N} W(x, z) dx, \quad \forall z \in E, \quad (2.5)$$

respectively. Our hypotheses imply that $\Phi_\varepsilon, \Phi \in \mathcal{C}^1(E, \mathbb{R})$, and a standard argument shows that the critical points of Φ_ε and Φ are solutions of problems (1.5) and (1.13), respectively. Moreover, by (1.9) and (1.17), there hold

$$\langle \Phi'_\varepsilon(z), \varphi \rangle = \frac{2ab}{a^2 + b^2} [(z^+, \varphi^+) - (z^-, \varphi^-)] + \int_{\mathbb{R}^N} V_\varepsilon(x) z \cdot \varphi dx \\ - \int_{\mathbb{R}^N} Q_\varepsilon(x) F_z(z) \cdot \varphi dx, \quad \forall z, \varphi \in E, \quad (2.6)$$

$$\langle \Phi'_\varepsilon(z), z \rangle = \frac{2ab}{a^2 + b^2} (\|z^+\|^2 - \|z^-\|^2) + \int_{\mathbb{R}^N} V_\varepsilon(x) |z|^2 dx - \int_{\mathbb{R}^N} Q_\varepsilon(x) F_z(z) \cdot z dx, \quad \forall z \in E, \quad (2.7)$$

$$\begin{aligned}\langle \Phi'(z), \varphi \rangle &= \frac{2ab}{a^2 + b^2} \left[(z^+, \varphi^+)_{\dagger} - (z^-, \varphi^-)_{\dagger} \right] + \int_{\mathbb{R}^N} V_2(x) z \cdot \varphi dx \\ &\quad - \int_{\mathbb{R}^N} W_z(x, z) \cdot \varphi dx, \quad \forall z, \varphi \in E\end{aligned}\tag{2.8}$$

and

$$\langle \Phi'(z), z \rangle = \frac{2ab}{a^2 + b^2} (\|z^+\|_{\dagger}^2 - \|z^-\|_{\dagger}^2) + \int_{\mathbb{R}^N} V_2(x) |z|^2 dx - \langle \Psi'(z), z \rangle, \quad \forall z \in E.\tag{2.9}$$

Lemma 2.1. *Suppose that (V0), (F1)-(F2) are satisfied. If $z = (u, v)$ is a critical point of Φ_ε , then $|z(x)| \rightarrow 0$ as $|x| \rightarrow \infty$. In a word, z is a solution to system (2.1).*

The proof is almost standard (see [10, Lemma 2.1 and Theorem 2.1]).

3 Ground state solutions of Nehari-Pankov type for periodic system

Let $X = X^- \oplus X^+$ be a real Hilbert space with $X^- \perp X^+$ and X^- be separable. On X we define a new norm

$$\|u\|_\tau := \max \left\{ \|u^+\|, \sum_{k=1}^{\infty} \frac{1}{2^{k+1}} |(u^-, e_k)| \right\}, \quad \forall u = u^- + u^+ \in X,\tag{3.1}$$

where $\{e_k\}_{k=1}^{\infty}$ is a total orthonormal basis of X^- . The topology generated by $\|\cdot\|_\tau$ will be denoted by τ and all topological notions related to it will include the symbol. It is clear that

$$\|u^+\| \leq \|u\|_\tau \leq \|u\|, \quad \forall u \in X.\tag{3.2}$$

For a functional $\varphi \in \mathcal{C}^1(X, \mathbb{R})$, φ is said to be τ -upper semi-continuous if

$$u_n, u \in X, \quad \|u_n - u\|_\tau \rightarrow 0 \Rightarrow \varphi(u) \geq \limsup_{n \rightarrow \infty} \varphi(u_n);\tag{3.3}$$

weakly sequentially lower semi-continuous if

$$u_n \rightharpoonup u \text{ in } X \Rightarrow \varphi(u) \leq \liminf_{n \rightarrow \infty} \varphi(u_n);$$

and φ' is said to be weakly sequentially continuous if

$$u_n \rightharpoonup u \text{ in } X \Rightarrow \lim_{n \rightarrow \infty} \langle \varphi'(u_n), v \rangle = \langle \varphi'(u), v \rangle, \quad \forall v \in X.$$

It is easy to see that (3.3) holds if and only if

$$u_n, u \in X, \quad \|u_n - u\|_\tau \rightarrow 0 \Rightarrow \varphi(u) \geq \liminf_{n \rightarrow \infty} \varphi(u_n).\tag{3.4}$$

Lemma 3.1. ([29, Theorem 2.4]) *Let $X = X^- \oplus X^+$ be a real Hilbert space with $X^- \perp X^+$ and X^- be separable. Suppose that $\varphi \in \mathcal{C}^1(X, \mathbb{R})$ satisfies the following assumptions:*

- (H1) φ is τ -upper semi-continuous;
- (H2) φ' is weakly sequentially continuous;
- (H3) there exist $r > \rho > 0$ and $e \in X^+$ with $\|e\| = 1$ such that

$$\kappa := \inf \varphi(S_\rho^+) > \sup \varphi(\partial \Omega_r),$$

where

$$S_\rho^+ = \{u \in X^+ : \|u\| = \rho\}, \quad \Omega_r = \{v + se : v \in X^-, s \geq 0, \|v + se\| \leq r\}.$$

Then there exist a constant $c \in [\kappa, \sup \varphi(\Omega_r)]$ and a sequence $\{u_n\} \subset X$ satisfying

$$\varphi(u_n) \rightarrow c, \quad \|\varphi'(u_n)\|(1 + \|u_n\|) \rightarrow 0. \quad (3.5)$$

Let $\Psi(z) = \int_{\mathbb{R}^N} W(x, z) dx$. Employing a standard argument, one can check easily the following lemma.

Lemma 3.2. *Suppose that (V0'), (W0) and (W1) are satisfied. Then Ψ is nonnegative, weakly sequentially lower semi-continuous, and Ψ' is weakly sequentially continuous.*

Lemma 3.3. *Suppose that (V0'), (W0), (W1) and (W4) are satisfied. Then there holds*

$$\begin{aligned} \Phi(z) &\geq \Phi(\theta z + \zeta) + \frac{ab}{a^2 + b^2} \|\zeta\|_\dagger^2 - \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |\zeta|^2 dx \\ &\quad + \frac{1 - \theta^2}{2} \langle \Phi'(z), z \rangle - \theta \langle \Phi'(z), \zeta \rangle, \quad \forall \theta \geq 0, \quad z \in E, \quad \zeta \in E^-. \end{aligned} \quad (3.6)$$

Proof. By (2.5), (2.8), (2.9) and (W4), one has

$$\begin{aligned} &\Phi(z) - \Phi(\theta z + \zeta) \\ &= \frac{ab}{a^2 + b^2} \|\zeta\|_\dagger^2 - \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |\zeta|^2 dx + \frac{1 - \theta^2}{2} \langle \Phi'(z), z \rangle - \theta \langle \Phi'(z), \zeta \rangle \\ &\quad + \int_{\mathbb{R}^N} \left[\frac{1 - \theta^2}{2} \nabla W(x, z) \cdot z - \theta \nabla W(x, z) \cdot \zeta + W(x, \theta z + \zeta) - W(x, z) \right] dx \\ &\geq \frac{ab}{a^2 + b^2} \|\zeta\|_\dagger^2 - \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |\zeta|^2 dx + \frac{1 - \theta^2}{2} \langle \Phi'(z), z \rangle - \theta \langle \Phi'(z), \zeta \rangle, \\ &\quad \forall \theta \geq 0, \quad z \in E, \quad \zeta \in E^-. \end{aligned}$$

This shows that (3.6) holds. □

From Lemma 3.3, we have the following two corollaries.

Corollary 3.4. *Suppose that (V0'), (W0), (W1) and (W4) are satisfied. Then for $z \in \mathcal{N}^-$*

$$\Phi(z) \geq \Phi(\theta z + \zeta) + \frac{ab}{a^2 + b^2} \|\zeta\|_{\dagger}^2 - \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |\zeta|^2 dx, \quad \forall \theta \geq 0, \quad \zeta \in E^-. \quad (3.7)$$

Corollary 3.5. *Suppose that (V0'), (W0), (W1) and (W4) are satisfied. Then*

$$\begin{aligned} \Phi(z) \geq & \frac{ab\theta^2}{a^2 + b^2} \|z\|_{\dagger}^2 + \frac{\theta^2}{2} \int_{\mathbb{R}^N} V_2(x) (|z^+|^2 - |z^-|^2) dx - \int_{\mathbb{R}^N} W(x, \theta z^+) dx \\ & + \frac{1 - \theta^2}{2} \langle \Phi'(z), z \rangle + \theta^2 \langle \Phi'(z), z^- \rangle, \quad \forall z \in E, \quad \theta \geq 0. \end{aligned} \quad (3.8)$$

Lemma 3.6. *Suppose that (V0'), (W0), (W1) and (W4) are satisfied. Then*

(i) *there exists $\rho > 0$ such that*

$$m := \inf_{\mathcal{N}^-} \Phi \geq \kappa := \inf \{ \Phi(z) : z \in E^+, \|z\|_{\dagger} = \rho \} > 0.$$

(ii) $\|z^+\|_{\dagger}^2 \geq \max \left\{ \frac{(1-\eta)^2}{2(1+\eta^2)} \|z^-\|_{\dagger}^2, \frac{(1-\eta)(a^2+b^2)m}{2ab} \right\}$ *for all $z \in \mathcal{N}^-$.*

Proof. By (V0'), we have $|V_2(x)| \leq \frac{2ab\eta}{a^2+b^2} V_1(x)$, it follows from (2.3) that

$$\begin{aligned} & \frac{2ab}{a^2 + b^2} \|z^+\|_{\dagger}^2 + \int_{\mathbb{R}^N} V_2(x) |z^+|^2 dx \\ = & \int_{\mathbb{R}^N} \left[\frac{2ab}{a^2 + b^2} |\nabla z^+|^2 + \left(\frac{2ab}{a^2 + b^2} V_1(x) + V_2(x) \right) |z^+|^2 \right] dx \\ \geq & \frac{2ab}{a^2 + b^2} \int_{\mathbb{R}^N} [|\nabla z^+|^2 + (1 - \eta) V_1(x) |z^+|^2] dx \\ \geq & \frac{2(1 - \eta)ab}{a^2 + b^2} \|z^+\|_{\dagger}^2, \quad \forall z \in E \end{aligned} \quad (3.9)$$

and

$$\begin{aligned} & \frac{ab}{a^2 + b^2} (\|z^+\|_{\dagger}^2 - \|z^-\|_{\dagger}^2) + \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |z|^2 dx \\ = & \frac{ab}{a^2 + b^2} (\|z^+\|_{\dagger}^2 - \|z^-\|_{\dagger}^2) + \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) (|z^+|^2 + |z^-|^2) dx + \int_{\mathbb{R}^N} V_2(x) z^+ \cdot z^- dx \\ \leq & \frac{ab}{a^2 + b^2} (\|z^+\|_{\dagger}^2 - \|z^-\|_{\dagger}^2) + \frac{1 + \eta}{2(1 - \eta)} \int_{\mathbb{R}^N} |V_2(x)| |z^+|^2 dx + \frac{1 + \eta}{4\eta} \int_{\mathbb{R}^N} |V_2(x)| |z^-|^2 dx \\ \leq & \frac{ab}{a^2 + b^2} \int_{\mathbb{R}^N} \left[|\nabla z^+|^2 + \frac{1 + \eta^2}{1 - \eta} V_1(x) |z^+|^2 \right] dx \\ & - \frac{ab}{a^2 + b^2} \int_{\mathbb{R}^N} \left[|\nabla z^-|^2 + \frac{1 - \eta}{2} V_1(x) |z^-|^2 \right] dx \\ \leq & \frac{(1 + \eta^2)ab}{(1 - \eta)(a^2 + b^2)} \|z^+\|_{\dagger}^2 - \frac{(1 - \eta)ab}{2(a^2 + b^2)} \|z^-\|_{\dagger}^2, \quad \forall z \in E. \end{aligned} \quad (3.10)$$

The rest of the proof is standard, so we omit it. \square

Lemma 3.7. Suppose that (V0'), (W0), (W1) and (W2) are satisfied. Let $e \in E^+$ with $\|e\|_{\dagger} = 1$. Then there is $r_0 > \rho$ such that $\sup \Phi(\partial\Omega_r) \leq 0$ for $r \geq r_0$, where

$$\Omega_r = \{\zeta + se : \zeta \in E^-, s \geq 0, \|\zeta + se\|_{\dagger} \leq r\}. \quad (3.11)$$

Proof. (2.5) and (3.10) imply $\Phi(z) \leq 0$ for $z \in E^-$. Next, it is sufficient to show that $\Phi(z) \rightarrow -\infty$ as $z \in E^- \oplus \mathbb{R}^+e$ and $\|z\|_{\dagger} \rightarrow \infty$. Arguing indirectly, assume that for some sequence $\{\zeta_n + s_n e\} \subset E^- \oplus \mathbb{R}^+e$ with $\|\zeta_n + s_n e\|_{\dagger} \rightarrow \infty$, there is an $M > 0$ such that $\Phi(\zeta_n + s_n e) \geq -M$ for all $n \in \mathbb{N}$. Set $e = (\frac{e_0}{a}, \frac{e_0}{b})$, $\zeta_n = (-\frac{w_n}{a}, \frac{w_n}{b})$ and $\xi_n = (\zeta_n + s_n e)/\|\zeta_n + s_n e\|_{\dagger} = \xi_n^- + t_n e$, then $\|\xi_n^- + t_n e\|_{\dagger} = 1$. Passing to a subsequence, we may assume that $t_n \rightarrow \bar{t}$ and $\xi_n \rightharpoonup \xi$ in E , then $\xi_n \rightarrow \xi$ a.e. on \mathbb{R}^N , $\xi_n^- := (-\frac{\bar{w}_n}{a}, \frac{\bar{w}_n}{b}) \rightharpoonup \xi^- := (-\frac{\bar{w}}{a}, \frac{\bar{w}}{b})$ in E . Hence, by (2.5) and (3.10), one has

$$\begin{aligned} -\frac{M}{\|\zeta_n + s_n e\|_{\dagger}^2} &\leq \frac{\Phi(\zeta_n + s_n e)}{\|\zeta_n + s_n e\|_{\dagger}^2} \\ &= \frac{abt_n^2}{a^2 + b^2} - \frac{ab}{a^2 + b^2} \|\xi_n^-\|_{\dagger}^2 + \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |\xi_n^- + t_n e|^2 dx \\ &\quad - \int_{\mathbb{R}^N} \frac{W(x, \frac{-w_n + s_n e_0}{a}, \frac{w_n + s_n e_0}{b})}{\|\zeta_n + s_n e\|_{\dagger}^2} dx \\ &\leq \frac{(1 + \eta^2)abt_n^2}{(1 - \eta)(a^2 + b^2)} - \frac{(1 - \eta)ab}{2(a^2 + b^2)} \|\xi_n^-\|_{\dagger}^2 \\ &\quad - \int_{\mathbb{R}^N} \frac{W(x, \frac{-w_n + s_n e_0}{a}, \frac{w_n + s_n e_0}{b})}{\|\zeta_n + s_n e\|_{\dagger}^2} dx. \end{aligned} \quad (3.12)$$

If $\bar{t} = 0$, then it follows from (3.12) that

$$\begin{aligned} 0 &\leq \frac{(1 - \eta)ab}{2(a^2 + b^2)} \|\xi_n^-\|_{\dagger}^2 + \int_{\mathbb{R}^N} \frac{W(x, \frac{-w_n + s_n e_0}{a}, \frac{w_n + s_n e_0}{b})}{\|\zeta_n + s_n e\|_{\dagger}^2} dx \\ &\leq \frac{(1 + \eta^2)abt_n^2}{(1 - \eta)(a^2 + b^2)} + \frac{M}{\|\zeta_n + s_n e\|_{\dagger}^2} \rightarrow 0, \end{aligned}$$

which yields $\|\xi_n^-\|_{\dagger} \rightarrow 0$, and so $1 = \|\xi_n\|_{\dagger} \rightarrow 0$, a contradiction.

If $\bar{t} \neq 0$, then $s_n \rightarrow \infty$. Hence, it follows from (3.12), (W2) and Fatou's Lemma that

$$\begin{aligned} 0 &\leq \limsup_{n \rightarrow \infty} \left[\frac{(1 + \eta^2)abt_n^2}{(1 - \eta)(a^2 + b^2)} - \frac{(1 - \eta)ab}{2(a^2 + b^2)} \|\xi_n^-\|_{\dagger}^2 \right. \\ &\quad \left. - \int_{\mathbb{R}^N} \frac{W(x, \frac{-w_n + s_n e_0}{a}, \frac{w_n + s_n e_0}{b})}{\|\zeta_n + s_n e\|_{\dagger}^2} dx \right] \\ &= \limsup_{n \rightarrow \infty} \left[\frac{(1 + \eta^2)abt_n^2}{(1 - \eta)(a^2 + b^2)} - \frac{(1 - \eta)ab}{2(a^2 + b^2)} \|\xi_n^-\|_{\dagger}^2 \right. \\ &\quad \left. - \int_{\mathbb{R}^N} \frac{W(x, \frac{-w_n + s_n e_0}{a}, \frac{w_n + s_n e_0}{b})}{|s_n e_0|^2} |t_n e_0|^2 dx \right] \\ &\leq \frac{(1 + \eta^2)ab}{(1 - \eta)(a^2 + b^2)} \lim_{n \rightarrow \infty} t_n^2 - \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} \frac{W(x, \frac{-w_n + s_n e_0}{a}, \frac{w_n + s_n e_0}{b})}{|s_n e_0|^2} |t_n e_0|^2 dx \end{aligned}$$

$$\begin{aligned}
&\leq \frac{(1+\eta^2)abt^2}{(1-\eta)(a^2+b^2)} - \int_{\mathbb{R}^N} \liminf_{n \rightarrow \infty} \left[\frac{W\left(x, \frac{-w_n+s_n e_0}{a}, \frac{w_n+s_n e_0}{b}\right)}{|s_n e_0|^2} |t_n e_0|^2 \right] dx \\
&= -\infty,
\end{aligned}$$

a contradiction. \square

Since E^- is separable, let $\{e_k\}_{k=1}^\infty$ be a total orthonormal basis of E^- . On E we define the τ -norm

$$\|z\|_\tau := \max \left\{ \|z^+\|_\dagger, \sum_{k=1}^\infty \frac{1}{2^{k+1}} |(z^-, e_k)_\dagger| \right\}, \quad \forall z \in E. \quad (3.13)$$

It is clear that

$$\|z^+\|_\dagger \leq \|z\|_\tau \leq \|z\|_\dagger, \quad \forall z \in E. \quad (3.14)$$

Lemma 3.8. *Suppose that (V0'), (W0), (W1) and (W2) are satisfied. Then $\Phi \in \mathcal{C}^1(E, \mathbb{R})$ is τ -upper semi-continuous and Φ' is weakly sequentially continuous.*

Proof. It is clear that $\Phi \in \mathcal{C}^1(E, \mathbb{R})$. First we prove that Φ is τ -upper semi-continuous. Let $z_n \xrightarrow{\tau} z$ in E and $\Phi(z_n) \geq c$. It follows from (2.5), (3.10), (3.14) and (W0) that $z_n^+ \rightarrow z^+$ in E and

$$C_1 \geq \frac{(1+\eta^2)ab}{(1-\eta)(a^2+b^2)} \|z_n^+\|_\dagger^2 \geq c + \frac{(1-\eta)ab}{2(a^2+b^2)} \|z_n^-\|_\dagger^2.$$

This shows that $\{z_n^-\} \subset E^-$ is bounded. It is easy to show that $z_n^- \xrightarrow{\tau} z^- \Leftrightarrow z_n^- \rightharpoonup z^-$, and so, $z_n \rightarrow z$ a.e. on \mathbb{R}^N . Note that

$$\begin{aligned}
&\left| \int_{\mathbb{R}^N} V_2(x) (z_n^+ \cdot z_n^- - z^+ \cdot z^-) dx \right| \\
&\leq \int_{\mathbb{R}^N} |V_2(x)| |z_n^+ - z^+| |z_n^-| dx + \left| \int_{\mathbb{R}^N} V_2(x) z^+ \cdot (z_n^- - z^-) dx \right| = o(1)
\end{aligned} \quad (3.15)$$

and

$$\begin{aligned}
&\liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} \left[|\nabla z_n^-|^2 + \left(V_1(x) - \frac{a^2+b^2}{2ab} V_2(x) \right) |z_n^-|^2 \right] dx \\
&\geq \int_{\mathbb{R}^N} \left[|\nabla z^-|^2 + \left(V_1(x) - \frac{a^2+b^2}{2ab} V_2(x) \right) |z^-|^2 \right] dx.
\end{aligned} \quad (3.16)$$

Hence, it follows from (W2), (2.5), (3.15), (3.16) and Fatou's Lemma that

$$\begin{aligned}
-\Phi(z) &= \frac{ab}{a^2+b^2} (\|z^-\|_\dagger^2 - \|z^+\|_\dagger^2) - \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |z|^2 dx + \int_{\mathbb{R}^N} W(x, z) dx \\
&= \frac{ab}{a^2+b^2} \int_{\mathbb{R}^N} \left[|\nabla z^-|^2 + \left(V_1(x) - \frac{a^2+b^2}{2ab} V_2(x) \right) |z^-|^2 \right] dx \\
&\quad - \frac{ab}{a^2+b^2} \int_{\mathbb{R}^N} \left[|\nabla z^+|^2 + \left(V_1(x) + \frac{a^2+b^2}{2ab} V_2(x) \right) |z^+|^2 \right] dx \\
&\quad - \int_{\mathbb{R}^N} V_2(x) z^+ \cdot z^- dx + \int_{\mathbb{R}^N} W(x, z) dx
\end{aligned}$$

$$\begin{aligned}
&\leq \liminf_{n \rightarrow \infty} \left\{ \frac{ab}{a^2 + b^2} \int_{\mathbb{R}^N} \left[|\nabla z_n^-|^2 + \left(V_1(x) - \frac{a^2 + b^2}{2ab} V_2(x) \right) |z_n^-|^2 \right] dx \right. \\
&\quad - \frac{ab}{a^2 + b^2} \int_{\mathbb{R}^N} \left[|\nabla z_n^+|^2 + \left(V_1(x) + \frac{a^2 + b^2}{2ab} V_2(x) \right) |z_n^+|^2 \right] dx \\
&\quad \left. - \int_{\mathbb{R}^N} V_2(x) z_n^+ \cdot z_n^- dx + \int_{\mathbb{R}^N} W(x, z_n) dx \right\} \\
&= \liminf_{n \rightarrow \infty} [-\Phi(z_n)] = -\limsup_{n \rightarrow \infty} \Phi(z_n).
\end{aligned}$$

This shows that Φ is τ -upper semi-continuous.

The proof that Φ' is weakly sequentially continuous is standard, so we omit it. \square

Lemma 3.9. *Suppose that (V0'), (W0), (W1), (W2) and (W4) are satisfied. Then there exist a constant $c \in [\kappa, \sup \Phi(\mathfrak{Q}_r)]$ for $r \geq r_0$ and a sequence $\{z_n\} \subset E$ satisfying*

$$\Phi(z_n) \rightarrow c, \quad \|\Phi'(z_n)\|_{E^*} (1 + \|z_n\|_{\dagger}) \rightarrow 0, \quad (3.17)$$

where \mathfrak{Q}_r is defined by (3.11).

Proof. Lemma 3.9 is a direct corollary of Lemmas 3.1, 3.2, 3.6 (i), 3.7 and 3.8. \square

Applying Corollary 3.4, Lemmas 3.6 (i), 3.7 and 3.9, we can prove the following lemma in a similar way as [29, Lemma 3.8].

Lemma 3.10. ([32, Lemma 3.9]) *Suppose that (V0'), (W0), (W1), (W2) and (W4) are satisfied. Then there exist a constant $c_* \in [\kappa, m]$ and a sequence $\{z_n\} = \{(u_n, v_n)\} \subset E$ satisfying*

$$\Phi(z_n) \rightarrow c_*, \quad \|\Phi'(z_n)\| (1 + \|z_n\|) \rightarrow 0. \quad (3.18)$$

Lemma 3.11. *Suppose that (V0'), (W0), (W1), (W2) and (W4) are satisfied. Then for any $z \in E \setminus E^-$, $\mathcal{N}^- \cap (E^- \oplus \mathbb{R}^+ z) \neq \emptyset$, i.e., there exist $t(z) > 0$ and $\zeta(z) \in E^-$ such that $t(z)z + \zeta(z) \in \mathcal{N}^-$.*

The proof is the same as one of [28, Lemma 2.6].

Lemma 3.12. *Suppose that (V0'), (W0), (W1), (W2), (W3) and (W4) are satisfied. Then any sequence $\{z_n\} = \{(u_n, v_n)\} \subset E$ satisfying*

$$\Phi(z_n) \rightarrow c \geq 0, \quad \langle \Phi'(z_n), z_n \rangle \rightarrow 0, \quad \langle \Phi'(z_n), z_n^- \rangle \rightarrow 0 \quad (3.19)$$

is bounded in E .

Proof. To prove the boundedness of $\{z_n\}$, arguing by contradiction, suppose that $\|z_n\|_{\dagger} \rightarrow \infty$. Let $\tilde{z}_n = (\tilde{u}_n, \tilde{v}_n) := z_n / \|z_n\|_{\dagger}$. Then $\|\tilde{z}_n\|_{\dagger} = 1$. By Sobolev embedding theorem, there exists a constant $C_2 > 0$ such that $\|\tilde{z}_n^+\|_2 \leq C_2$. If $\delta := \limsup_{n \rightarrow \infty} \sup_{y \in \mathbb{R}^N} \int_{B_1(y)} |\tilde{z}_n^+|^2 dx = 0$,

then by Lions' concentration compactness principle [33, Lemma 1.21], $\tilde{z}_n^+ \rightarrow 0$ in $L^p(\mathbb{R}^N)$. Fix $\vartheta = [(a^2 + b^2)(1 + c)/(1 - \eta)ab]^{1/2}$. By virtue of (W0) and (W1), for $\epsilon = 1/4(\vartheta C_2)^2 > 0$, there exists $C_\epsilon > 0$ such that

$$W(x, z) \leq \epsilon |z|^2 + C_\epsilon |z|^p, \quad \forall (x, z) \in \mathbb{R}^N \times \mathbb{R}^2.$$

Hence, it follows that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} W(x, \vartheta \tilde{z}_n^+) dx &\leq \epsilon \vartheta^2 \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\tilde{z}_n^+|^2 dx + C_\epsilon \vartheta^p \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\tilde{z}_n^+|^p dx \\ &\leq \epsilon (\vartheta C_2)^2 = \frac{1}{4}. \end{aligned} \quad (3.20)$$

Let $\theta_n = \vartheta / \|z_n\|_{\dagger}$. Hence, by virtue of (3.8), (3.19) and (3.20), one can get

$$\begin{aligned} c + o(1) &= \Phi(z_n) \\ &\geq \frac{ab\theta_n^2}{a^2 + b^2} \|z_n\|_{\dagger}^2 + \frac{\theta_n^2}{2} \int_{\mathbb{R}^N} V_2(x) (|z_n^+|^2 - |z_n^-|^2) dx - \int_{\mathbb{R}^N} W(x, \theta_n z_n^+) dx \\ &\quad + \frac{1 - \theta_n^2}{2} \langle \Phi'(z_n), z_n \rangle + \theta_n^2 \langle \Phi'(z_n), z_n^- \rangle \\ &= \frac{ab\vartheta^2}{a^2 + b^2} \|\tilde{z}_n\|_{\dagger}^2 + \frac{\vartheta^2}{2} \int_{\mathbb{R}^N} V_2(x) (|\tilde{z}_n^+|^2 - |\tilde{z}_n^-|^2) dx - \int_{\mathbb{R}^N} W(x, \vartheta \tilde{z}_n^+) dx \\ &\quad + \left(\frac{1}{2} - \frac{\vartheta^2}{2\|z_n\|_{\dagger}^2} \right) \langle \Phi'(z_n), z_n \rangle + \frac{\vartheta^2}{\|z_n\|_{\dagger}^2} \langle \Phi'(z_n), z_n^- \rangle \\ &\geq \frac{(1 - \eta)ab\vartheta^2}{a^2 + b^2} - \int_{\mathbb{R}^N} W(x, \vartheta \tilde{z}_n^+) dx + o(1) \\ &\geq \frac{(1 - \eta)ab\vartheta^2}{a^2 + b^2} - \frac{1}{4} + o(1) > \frac{3}{4} + c + o(1). \end{aligned}$$

This contradiction shows that $\delta > 0$. Going if necessary to a subsequence, we may assume the existence of $k_n \in \mathbb{Z}^N$ such that $\int_{B_{1+\sqrt{N}}(k_n)} |\tilde{z}_n^+|^2 dx > \frac{\delta}{2}$. Let $\zeta_n(x) := \tilde{z}_n(x + k_n)$. Then

$$\int_{B_{1+\sqrt{N}}(0)} |\zeta_n^+|^2 dx > \frac{\delta}{2}. \quad (3.21)$$

Now we define $z_n^{k_n}(x) := (u_n^{k_n}(x), v_n^{k_n}(x)) = z_n(x + k_n)$, then $z_n^{k_n} / \|z_n\|_{\dagger} = \zeta_n$ and $\|\zeta_n\|_{H^1(\mathbb{R}^N)}^2 = \|\tilde{z}_n\|_{H^1(\mathbb{R}^N)}^2$. Passing to a subsequence, we have $\zeta_n^+ \rightharpoonup \zeta^+$ in $H^1(\mathbb{R}^N)$, $\zeta_n^+ \rightarrow \zeta^+$ in $L_{\text{loc}}^s(\mathbb{R}^N)$, $2 \leq s < 2^*$ and $\zeta_n^+ \rightarrow \zeta^+$ a.e. on \mathbb{R}^N . Obviously, (3.21) implies that $\zeta^+ \neq 0$. For a.e. $x \in \{y \in \mathbb{R}^N : \zeta^+(y) \neq 0\} := \Omega$, we have $\lim_{n \rightarrow \infty} |au_n^{k_n}(x) + bv_n^{k_n}(x)| = \infty$. Hence, it follows from (2.5), (3.10), (3.19), (W2), (W3) and Fatou's lemma that

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \frac{c + o(1)}{\|z_n\|_{\dagger}^2} = \lim_{n \rightarrow \infty} \frac{\Phi(z_n)}{\|z_n\|_{\dagger}^2} \\ &= \lim_{n \rightarrow \infty} \left[\frac{ab}{a^2 + b^2} (\|\tilde{z}_n^+\|_{\dagger}^2 - \|\tilde{z}_n^-\|_{\dagger}^2) + \frac{1}{2} \int_{\mathbb{R}^N} V_2(x) |\tilde{z}_n|^2 dx - \int_{\mathbb{R}^N} \frac{W(x, u_n, v_n)}{\|z_n\|_{\dagger}^2} dx \right] \end{aligned}$$

$$\begin{aligned}
&\leq \lim_{n \rightarrow \infty} \left[\frac{(1+\eta^2)ab}{(1-\eta)(a^2+b^2)} \|\tilde{z}_n^+\|_{\dagger}^2 - \frac{(1-\eta)ab}{2(a^2+b^2)} \|\tilde{z}_n^-\|_{\dagger}^2 - \int_{\Omega} \frac{W(x+k_n, u_n^{k_n}, v_n^{k_n})}{\|z_n\|_{\dagger}^2} dx \right] \\
&= \lim_{n \rightarrow \infty} \left[\frac{(1+\eta^2)ab}{(1-\eta)(a^2+b^2)} \|\tilde{z}_n^+\|_{\dagger}^2 - \frac{(1-\eta)ab}{2(a^2+b^2)} \|\tilde{z}_n^-\|_{\dagger}^2 - \frac{4a^2b^2}{a^2+b^2} \int_{\Omega} \frac{W(x, u_n^{k_n}, v_n^{k_n})}{|au_n^{k_n} + bv_n^{k_n}|^2} |\zeta_n^+|^2 dx \right] \\
&\leq \frac{(1+\eta^2)ab}{(1-\eta)(a^2+b^2)} - \frac{4a^2b^2}{a^2+b^2} \int_{\Omega} \liminf_{n \rightarrow \infty} \left[\frac{W(x, u_n^{k_n}, v_n^{k_n})}{|au_n^{k_n} + bv_n^{k_n}|^2} |\zeta_n^+|^2 \right] dx = -\infty.
\end{aligned}$$

This contradiction shows that $\{\|u_n\|_{\dagger}\}$ is bounded. \square

In the last part of the proof of Lemma 3.12, we make use of the periodicity of $W(x, z)$ on x , which is still valid by using (W2') instead of (W2) and (W3). Therefore, we have the following lemma.

Lemma 3.13. *Suppose that (V0'), (W0), (W1), (W2') and (W4) are satisfied. Then any sequence $\{z_n\} = \{(u_n, v_n)\} \subset E$ satisfying (3.19) is bounded.*

Lemma 3.14. ([31, Lemma 2.3]) *Suppose that $t \mapsto h(x, t)$ is nondecreasing on \mathbb{R} and $h(x, 0) = 0$ for any $x \in \mathbb{R}^N$. Then there holds*

$$\left(\frac{1-\theta^2}{2} \tau - \theta \sigma \right) h(x, \tau) |\tau| \geq \int_{\theta\tau+\sigma}^{\tau} h(x, s) |s| ds, \quad \forall \theta \geq 0, \quad \tau, \sigma \in \mathbb{R}. \quad (3.22)$$

Lemma 3.15. ([32]) *Suppose that $W(x, u, v) = \int_0^{|\alpha u + \beta v|} g(x, s) ds$, where $\alpha, \beta \in \mathbb{R}$ with $\alpha^2 + \beta^2 \neq 0$ and $g \in \mathcal{ND}$. Then W satisfies (W0), (W1) and (W4).*

Lemma 3.16. ([32]) *Suppose that $W(x, u, v) = \int_0^{\sqrt{u^2 + 2\beta uv + \alpha v^2}} h(x, s) ds$, where $\alpha, \beta \in \mathbb{R}$ with $\alpha > \beta^2$ and $h \in \mathcal{ND}$. Then W satisfies (W0), (W1) and (W4).*

Proof of Theorem 1.8. Applying Lemmas 3.10 and 3.12, we deduce that there exists a bounded sequence $\{z_n\} = \{(u_n, v_n)\} \subset E$ satisfying (3.18). The rest of the proof is standard. \square

Employing Theorem 1.8, the conclusion of Corollary 1.9 follows by Lemmas 3.14-3.16.

4 Ground state solutions of Nehari-Pankov type for (2.1)

Without loss of generality, from now on, we assume that $x_v = 0 \in \mathcal{V}$. We only consider the case when (V1) is satisfied, since the arguments are similar when (V2) is satisfied. Then

$$V(0) = V_{\min}, \quad Q(x) \leq Q(0), \quad \forall |x| \geq R. \quad (4.1)$$

Let $V_1 = 1$ and $V_2 = V_\varepsilon$ (or $\hat{V}, V_{\min}, V_{\max}$), $W(x, z) = Q_\varepsilon(x)F(z)$ (or $Q(0)F(z), Q_{\min}F(z)$). Then (V0), (F1) and (F2) imply (V0'), (W0), (W1), (W2') and (W4), respectively. Let

$$\hat{V} := \frac{1}{2}(V_\infty + V_{\min}) = \frac{1}{2}(V_\infty + V(0)).$$

We define three auxiliary functionals as follows:

$$\hat{\Phi}(z) = \frac{ab}{a^2 + b^2} (\|z^+\|^2 - \|z^-\|^2) + \frac{\hat{V}}{2} \int_{\mathbb{R}^N} |z|^2 dx - Q(0) \int_{\mathbb{R}^N} F(z) dx, \quad \forall z \in E, \quad (4.2)$$

$$\Phi_0(z) = \frac{ab}{a^2 + b^2} (\|z^+\|^2 - \|z^-\|^2) + \frac{V(0)}{2} \int_{\mathbb{R}^N} |z|^2 dx - Q(0) \int_{\mathbb{R}^N} F(z) dx, \quad \forall z \in E \quad (4.3)$$

and

$$\Phi_*(z) = \frac{ab}{a^2 + b^2} (\|z^+\|^2 - \|z^-\|^2) + \frac{V_{\max}}{2} \int_{\mathbb{R}^N} |z|^2 dx - Q_{\min} \int_{\mathbb{R}^N} F(z) dx, \quad \forall z \in E. \quad (4.4)$$

Let

$$\hat{\mathcal{N}}^- = \left\{ z \in E \setminus E^- : \langle \hat{\Phi}'(z), z \rangle = \langle \hat{\Phi}'(z), \zeta \rangle = 0, \forall \zeta \in E^- \right\} \quad (4.5)$$

and

$$\mathcal{N}_0^- = \left\{ z \in E \setminus E^- : \langle \Phi_0'(z), z \rangle = \langle \Phi_0'(z), \zeta \rangle = 0, \forall \zeta \in E^- \right\} \quad (4.6)$$

be the Nehari-Pankov “manifolds” of the functionals $\hat{\Phi}$ and Φ_0 , respectively. Let

$$c_\varepsilon = \inf_{\mathcal{N}_\varepsilon^-} \Phi_\varepsilon, \quad \hat{c} = \inf_{\hat{\mathcal{N}}^-} \hat{\Phi}, \quad c_0 = \inf_{\mathcal{N}_0^-} \Phi_0. \quad (4.7)$$

Applying Lemma 3.3 and Corollary 3.5 to Φ_ε , we have the following two lemmas.

Lemma 4.1. *Suppose that (V0) and (F1) are satisfied. Then*

$$\begin{aligned} \Phi_\varepsilon(z) &\geq \Phi_\varepsilon(\theta z + \zeta) + \frac{ab}{a^2 + b^2} \|\zeta\|^2 - \frac{1}{2} \int_{\mathbb{R}^N} V_\varepsilon(x) |\zeta|^2 dx + \frac{1 - \theta^2}{2} \langle \Phi'_\varepsilon(z), z \rangle \\ &\quad - \theta \langle \Phi'_\varepsilon(z), \zeta \rangle, \quad \forall \theta \geq 0, \quad z \in E, \quad \zeta \in E^- \end{aligned} \quad (4.8)$$

$$\begin{aligned} &\geq \Phi_\varepsilon(\theta z + \zeta) + \frac{(1 - \eta)ab}{a^2 + b^2} \|\zeta\|^2 + \frac{1 - \theta^2}{2} \langle \Phi'_\varepsilon(z), z \rangle - \theta \langle \Phi'_\varepsilon(z), \zeta \rangle, \\ &\quad \forall \theta \geq 0, \quad z \in E, \quad \zeta \in E^-. \end{aligned} \quad (4.9)$$

Lemma 4.2. *Suppose that (V0) and (F1) are satisfied. Then*

$$\begin{aligned} \Phi_\varepsilon(z) &\geq \frac{ab\theta^2}{a^2 + b^2} \|z\|^2 + \frac{\theta^2}{2} \int_{\mathbb{R}^N} V_\varepsilon(x) (|z^+|^2 - |z^-|^2) dx - \int_{\mathbb{R}^N} Q_\varepsilon(x) F(\theta z^+) dx \\ &\quad + \frac{1 - \theta^2}{2} \langle \Phi'_\varepsilon(z), z \rangle + \theta^2 \langle \Phi'_\varepsilon(z), z^- \rangle, \quad \forall \theta \geq 0, \quad z \in E \\ &\geq \frac{(1 - \eta)ab\theta^2}{a^2 + b^2} \|z\|^2 - \int_{\mathbb{R}^N} Q_\varepsilon(x) F(\theta z^+) dx \end{aligned} \quad (4.10)$$

$$+\frac{1-\theta^2}{2}\langle\Phi'_\varepsilon(z), z\rangle + \theta^2\langle\Phi'_\varepsilon(z), z^-\rangle, \quad \forall \theta \geq 0, \quad z \in E. \quad (4.11)$$

By virtue of Corollary 1.9, under assumptions (V0), (F1) and (F2), there exists a $\hat{z} \in \hat{\mathcal{N}}^-$ such that $\hat{c} = \hat{\Phi}(\hat{z})$. In view of Lemma 3.11, there exist $\hat{t} > 0$ and $\hat{\zeta} \in E^-$ such that $\hat{t}\hat{z} + \hat{\zeta} \in \mathcal{N}_0^-$, and so $\Phi_0(\hat{t}\hat{z} + \hat{\zeta}) \geq c_0$.

Lemma 4.3. *Suppose that (V0), (V1), (F1) and (F2) are satisfied. Then $\hat{c} \geq c_0 + \hat{\delta}$, where*

$$\hat{\delta} := \frac{V_\infty - V_{\min}}{4} \int_{\mathbb{R}^N} |\hat{t}\hat{z} + \hat{\zeta}|^2 dx > 0 \quad (4.12)$$

is independent of $\varepsilon > 0$.

Proof. Applying Lemma 3.3 to $\hat{\Phi}(z)$, one has

$$\begin{aligned} \hat{c} &= \hat{\Phi}(\hat{z}) \geq \hat{\Phi}(\hat{t}\hat{z} + \hat{\zeta}) \\ &= \Phi_0(\hat{t}\hat{z} + \hat{\zeta}) + \frac{\hat{V} - V_{\min}}{2} \int_{\mathbb{R}^N} |\hat{t}\hat{z} + \hat{\zeta}|^2 dx \\ &\geq c_0 + \frac{V_\infty - V_{\min}}{4} \int_{\mathbb{R}^N} |\hat{t}\hat{z} + \hat{\zeta}|^2 dx = c_0 + \hat{\delta}. \end{aligned}$$

□

By virtue of Corollary 1.9, under assumptions (V0), (F1) and (F2), there exists a $z_0 \in \mathcal{N}_0^-$ such that $c_0 = \Phi_0(z_0)$. Then

$$\Phi_0(z_0) \geq \Phi_0(tz_0 + \zeta), \quad \langle\Phi'_0(z_0), z_0\rangle = \langle\Phi'_0(z_0), \zeta\rangle = 0, \quad \forall t \geq 0, \quad \zeta \in E^-. \quad (4.13)$$

In view of Lemma 3.11, for any $\varepsilon > 0$, there exist $t_\varepsilon > 0$ and $\zeta_\varepsilon \in E^-$ such that $t_\varepsilon z_0 + \zeta_\varepsilon \in \mathcal{N}_\varepsilon^-$, and so $\Phi_\varepsilon(t_\varepsilon z_0 + \zeta_\varepsilon) \geq c_\varepsilon$ and $\Phi_\varepsilon(t_\varepsilon z_0 + \zeta_\varepsilon) \geq \Phi_\varepsilon(tz_0 + \zeta)$, $\forall t \geq 0$, $\zeta \in E^-$. Set

$$\alpha_0 := \frac{ab}{a^2 + b^2} (\|z_0^+\|^2 - \|z_0^-\|^2) + \frac{V_{\min}}{2} \int_{\mathbb{R}^N} |z_0|^2 dx - Q_{\max} \int_{\mathbb{R}^N} F(z_0) dx. \quad (4.14)$$

Clearly, α_0 is independent of $\varepsilon > 0$. Analogous to the proof of Lemma 3.7, one can demonstrate the following lemma.

Lemma 4.4. *Suppose that (V0), (V1), (F1) and (F2) are satisfied. Then there is an $M_0 > 0$ independent of $\varepsilon > 0$ such that*

$$\Phi_*(\zeta + sz_0) \leq \alpha_0 - 1, \quad \forall \zeta \in E^-, \quad s \geq 0, \quad \|\zeta + sz_0\| \geq M_0. \quad (4.15)$$

Lemma 4.5. *Suppose that (V0), (V1), (F1) and (F2) are satisfied. Then*

$$M_1 := \sup_{\varepsilon > 0} |t_\varepsilon| \leq M_0 \|z_0^+\|^{-1}, \quad M_2 := \sup_{\varepsilon > 0} \|\zeta_\varepsilon\| \leq M_0 (1 + \|z_0\| \|z_0^+\|^{-1}). \quad (4.16)$$

Proof. Note that

$$\begin{aligned}\alpha_0 &= \frac{ab}{a^2+b^2} (\|z_0^+\|^2 - \|z_0^-\|^2) + \frac{V_{\min}}{2} \int_{\mathbb{R}^N} |z_0|^2 dx - Q_{\max} \int_{\mathbb{R}^N} F(z_0) dx \\ &\leq \Phi_\varepsilon(z_0) \leq \Phi_\varepsilon(t_\varepsilon z_0 + \zeta_\varepsilon) \leq \Phi_*(t_\varepsilon z_0 + \zeta_\varepsilon).\end{aligned}\quad (4.17)$$

By Lemma 4.4, one obtain that

$$\sup_{\varepsilon>0} \|t_\varepsilon z_0 + \zeta_\varepsilon\| \leq M_0. \quad (4.18)$$

Since

$$\|t_\varepsilon z_0 + \zeta_\varepsilon\|^2 = \|t_\varepsilon z_0^+\|^2 + \|t_\varepsilon z_0^- + \zeta_\varepsilon\|^2 \geq t_\varepsilon^2 \|z_0^+\|^2,$$

it follows from (4.18) and the above that

$$M_1 = \sup_{\varepsilon>0} |t_\varepsilon| \leq M_0 \|z_0^+\|^{-1}, \quad M_2 = \sup_{\varepsilon>0} \|\zeta_\varepsilon\| \leq M_0 (1 + \|z_0\| \|z_0^+\|^{-1}).$$

□

In view of (F1), there exists a constant $\beta_0 > 0$ such that

$$|F_z(z)| \leq \beta_0(|z| + |z|^{p-1}), \quad \forall z \in \mathbb{R}^2. \quad (4.19)$$

Now, we can choose $R_0 > R$ sufficient large such that

$$V(x) \geq \hat{V}, \quad Q(x) \leq Q(0), \quad \forall |x| \geq R_0 \quad (4.20)$$

and

$$\begin{aligned}& M_1^2 \int_{|x|>R_0} |z_0|^2 dx - (1 + M_1^2) Q_{\max} \int_{|x|>R_0} F_z(z_0) \cdot z_0 dx \\ & + 2(1 + \beta_0 Q_{\max}) M_1 M_2 \left(\int_{|x|>R_0} |z_0|^2 dx \right)^{1/2} \\ & + 2\beta_0 \gamma_p Q_{\max} M_1 M_2 \left(\int_{|x|>R_0} |z_0|^p dx \right)^{(p-1)/p} \leq \frac{\hat{\delta}}{2},\end{aligned}\quad (4.21)$$

where γ_p is the embedding constant with $\|\cdot\|_p \leq \gamma_p \|\cdot\|$. For the $R_0 > 0$ given above, we can choose an $\varepsilon_0 > 0$ such that

$$\begin{aligned}& \sqrt{2 + 2\beta_0 Q_{\max}} M_1 M_2 \|z_0\|_2 \left\{ \sup_{|x|\leq R_0} [|V_\varepsilon(x) - V(0)| + \beta_0 |Q_\varepsilon(x) - Q(0)|] \right\}^{1/2} \\ & + \frac{M_1^2}{2} \sup_{|x|\leq R_0} |V_\varepsilon(x) - V(0)| \|z_0\|_2^2 + \frac{1 + M_1^2}{2} \sup_{|x|\leq R_0} |Q_\varepsilon(x) - Q(0)| \int_{|x|\leq R_0} F_z(z_0) \cdot z_0 dx \\ & + \beta_0 \gamma_p (2Q_{\max})^{1/p} M_1 M_2 \|z_0\|_p^{p-1} \left\{ \sup_{|x|\leq R_0} |Q_\varepsilon(x) - Q(0)| \right\}^{(p-1)/p} \leq \frac{\hat{\delta}}{4}, \quad \varepsilon \in [0, \varepsilon_0].\end{aligned}\quad (4.22)$$

Lemma 4.6. *Suppose that (V0), (V1), (F1) and (F2) are satisfied. Then*

$$c_0 \geq c_\varepsilon - 3\hat{\delta}/4, \quad \forall \varepsilon \in [0, \varepsilon_0]. \quad (4.23)$$

Proof. From (F1), (2.4), (4.3), (4.9), (4.13), (4.16), (4.19), (4.21), (4.22) and the Hölder inequality, we have

$$\begin{aligned}
c_0 &= \Phi_0(z_0) = \Phi_\varepsilon(z_0) + \frac{1}{2} \int_{\mathbb{R}^N} [V(0) - V_\varepsilon(x)] |z_0|^2 dx + \int_{\mathbb{R}^N} [Q_\varepsilon(x) - Q(0)] F(z_0) dx \\
&\geq \Phi_\varepsilon(t_\varepsilon z_0 + \zeta_\varepsilon) + \frac{(1-\eta)ab}{a^2 + b^2} \|\zeta_\varepsilon\|^2 + \frac{1-t_\varepsilon^2}{2} \langle \Phi'_\varepsilon(z_0), z_0 \rangle - t_\varepsilon \langle \Phi'_\varepsilon(z_0), \zeta_\varepsilon \rangle \\
&\quad + \frac{1}{2} \int_{\mathbb{R}^N} [V(0) - V_\varepsilon(x)] |z_0|^2 dx + \int_{\mathbb{R}^N} [Q_\varepsilon(x) - Q(0)] F(z_0) dx \\
&\geq c_\varepsilon + \frac{1}{2} \int_{\mathbb{R}^N} [V(0) - V_\varepsilon(x)] |z_0|^2 dx + \int_{\mathbb{R}^N} [Q_\varepsilon(x) - Q(0)] F(z_0) dx \\
&\quad + \frac{1-t_\varepsilon^2}{2} \left\{ \int_{\mathbb{R}^N} [V_\varepsilon(x) - V(0)] |z_0|^2 dx + \int_{\mathbb{R}^N} [Q(0) - Q_\varepsilon(x)] F_z(z_0) \cdot z_0 dx \right\} \\
&\quad - t_\varepsilon \left\{ \int_{\mathbb{R}^N} [V_\varepsilon(x) - V(0)] z_0 \cdot \zeta_\varepsilon dx + \int_{\mathbb{R}^N} [Q(0) - Q_\varepsilon(x)] F_z(z_0) \cdot \zeta_\varepsilon dx \right\} \\
&= c_\varepsilon + \frac{1}{2} \int_{\mathbb{R}^N} [Q_\varepsilon(x) - Q(0)] [2F(z_0) - F_z(z_0) \cdot z_0] dx \\
&\quad - \frac{t_\varepsilon^2}{2} \left\{ \int_{\mathbb{R}^N} [V_\varepsilon(x) - V(0)] |z_0|^2 dx + \int_{\mathbb{R}^N} [Q(0) - Q_\varepsilon(x)] F_z(z_0) \cdot z_0 dx \right\} \\
&\quad - t_\varepsilon \left\{ \int_{\mathbb{R}^N} [V_\varepsilon(x) - V(0)] z_0 \cdot \zeta_\varepsilon dx + \int_{\mathbb{R}^N} [Q(0) - Q_\varepsilon(x)] F_z(z_0) \cdot \zeta_\varepsilon dx \right\} \\
&\geq c_\varepsilon - \frac{1}{2} \int_{\mathbb{R}^N} |Q_\varepsilon(x) - Q(0)| |F_z(z_0) \cdot z_0 - 2F(z_0)| dx \\
&\quad - \frac{M_1^2}{2} \int_{\mathbb{R}^N} |V_\varepsilon(x) - V(0)| |z_0|^2 dx - \frac{M_1^2}{2} \int_{\mathbb{R}^N} |Q_\varepsilon(x) - Q(0)| F_z(z_0) \cdot z_0 dx \\
&\quad - M_1 \int_{\mathbb{R}^N} |V_\varepsilon(x) - V(0)| |z_0| |\zeta_\varepsilon| dx - M_1 \int_{\mathbb{R}^N} |Q_\varepsilon(x) - Q(0)| |F_z(z_0)| |\zeta_\varepsilon| dx \\
&\geq c_\varepsilon - \frac{M_1^2}{2} \int_{\mathbb{R}^N} |V_\varepsilon(x) - V(0)| |z_0|^2 dx - \frac{1+M_1^2}{2} \int_{\mathbb{R}^N} |Q_\varepsilon(x) - Q(0)| F_z(z_0) \cdot z_0 dx \\
&\quad - \sqrt{2+2\beta_0 Q_{\max}} M_1 M_2 \left\{ \int_{\mathbb{R}^N} [|V_\varepsilon(x) - V(0)| + \beta_0 |Q_\varepsilon(x) - Q(0)|] |z_0|^2 dx \right\}^{1/2} \\
&\quad - \beta_0 \gamma_p (2Q_{\max})^{1/p} M_1 M_2 \left\{ \int_{\mathbb{R}^N} |Q_\varepsilon(x) - Q(0)| |z_0|^p dx \right\}^{(p-1)/p} \\
&\geq c_\varepsilon - \frac{M_1^2}{2} \int_{|x| \leq R_0} |V_\varepsilon(x) - V(0)| |z_0|^2 dx - \frac{1+M_1^2}{2} \int_{|x| \leq R_0} |Q_\varepsilon(x) - Q(0)| F_z(z_0) \cdot z_0 dx \\
&\quad - \sqrt{2+2\beta_0 Q_{\max}} M_1 M_2 \left\{ \int_{|x| \leq R_0} [|V_\varepsilon(x) - V(0)| + \beta_0 |Q_\varepsilon(x) - Q(0)|] |z_0|^2 dx \right\}^{1/2} \\
&\quad - \beta_0 \gamma_p (2Q_{\max})^{1/p} M_1 M_2 \left\{ \int_{|x| \leq R_0} |Q_\varepsilon(x) - Q(0)| |z_0|^p dx \right\}^{(p-1)/p}
\end{aligned}$$

$$\begin{aligned}
& -M_1^2 \int_{|x|>R_0} |z_0|^2 dx - (1+M_1^2)Q_{\max} \int_{|x|>R_0} F_z(z_0) \cdot z_0 dx \\
& -2(1+\beta_0 Q_{\max})M_1 M_2 \left(\int_{|x|>R_0} |z_0|^2 dx \right)^{1/2} - 2\beta_0 \gamma_p Q_{\max} M_1 M_2 \left(\int_{|x|>R_0} |z_0|^p dx \right)^{(p-1)/p} \\
\geq & c_\varepsilon - \sqrt{2+2\beta_0 Q_{\max}} M_1 M_2 \|z_0\|_2 \left\{ \sup_{|x|\leq R_0} [|V_\varepsilon(x) - V(0)| + \beta_0 |Q_\varepsilon(x) - Q(0)|] \right\}^{1/2} \\
& - \frac{M_1^2}{2} \sup_{|x|\leq R_0} |V_\varepsilon(x) - V(0)| \|z_0\|_2^2 - \frac{1+M_1^2}{2} \sup_{|x|\leq R_0} |Q_\varepsilon(x) - Q(0)| \int_{|x|\leq R_0} F_z(z_0) \cdot z_0 dx \\
& - \beta_0 \gamma_p (2Q_{\max})^{1/p} M_1 M_2 \|z_0\|_p^{p-1} \left\{ \sup_{|x|\leq R_0} |Q_\varepsilon(x) - Q(0)| \right\}^{(p-1)/p} - \frac{\hat{\delta}}{2} \\
\geq & c_\varepsilon - \frac{3\hat{\delta}}{4}.
\end{aligned}$$

□

Similar to Lemma 3.6, we can demonstrate that for any $\varepsilon > 0$, there exists a $\rho_\varepsilon > 0$ such that

$$c_\varepsilon = \inf_{\mathcal{N}_\varepsilon^-} \Phi_\varepsilon \geq \kappa_\varepsilon := \inf \{ \Phi_\varepsilon(z) : z \in E^+, \|z\| = \rho_\varepsilon \} > 0. \quad (4.24)$$

Applying Lemmas 3.10 and 3.13 to Φ_ε , we have the following two lemmas.

Lemma 4.7. *Suppose that (V0), (F1) and (F2) are satisfied. Then there exist a constant $\bar{c}_\varepsilon \in [\kappa_\varepsilon, c_\varepsilon]$ and a sequence $\{z_n^\varepsilon\} = \{(u_n^\varepsilon, v_n^\varepsilon)\} \subset E$ satisfying*

$$\Phi_\varepsilon(z_n^\varepsilon) \rightarrow \bar{c}_\varepsilon, \quad \|\Phi'_\varepsilon(z_n^\varepsilon)\|(1 + \|z_n^\varepsilon\|) \rightarrow 0. \quad (4.25)$$

Lemma 4.8. *Suppose that (V0), (F1) and (F2) are satisfied. Then the sequence $\{z_n^\varepsilon\} = \{(u_n^\varepsilon, v_n^\varepsilon)\} \subset E$ satisfying (4.25) is bounded in E .*

Similar to (3.10), one has

$$\begin{aligned}
& \frac{ab}{a^2 + b^2} (\|z^+\|^2 - \|z^-\|^2) + \frac{1}{2} \int_{\mathbb{R}^N} V_\varepsilon(x) |z|^2 dx \\
\leq & \frac{(1+\eta^2)ab}{(1-\eta)(a^2 + b^2)} \|z^+\|^2 - \frac{(1-\eta)ab}{2(a^2 + b^2)} \|z^-\|^2, \quad \forall z \in E.
\end{aligned} \quad (4.26)$$

By (F1), (2.4), (4.25) and (4.26), one has

$$\begin{aligned}
\bar{c}_\varepsilon + o(1) &= \Phi_\varepsilon(z_n^\varepsilon) \\
&= \frac{ab}{a^2 + b^2} (\|z_n^{\varepsilon+}\|^2 - \|z_n^{\varepsilon-}\|^2) + \frac{1}{2} \int_{\mathbb{R}^N} V_\varepsilon(x) |z_n^\varepsilon|^2 dx - \int_{\mathbb{R}^N} Q_\varepsilon(x) F(z_n^\varepsilon) dx \\
&\leq \frac{(1+\eta^2)ab}{(1-\eta)(a^2 + b^2)} \|z_n^{\varepsilon+}\|^2 - \frac{(1-\eta)ab}{2(a^2 + b^2)} \|z_n^{\varepsilon-}\|^2
\end{aligned}$$

$$\leq \frac{(1+\eta^2)ab}{(1-\eta)(a^2+b^2)} \|z_n^{\varepsilon+}\|^2. \quad (4.27)$$

Therefore, it follows from Lemma 3.11 that there exist $t_n^\varepsilon > 0$ and $\zeta_n^\varepsilon \in E^-$ such that $t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon \in \hat{\mathcal{N}}^-$, and so

$$\hat{\Phi}(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon) \geq \hat{c}, \quad \langle \hat{\Phi}'(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon), t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon \rangle = \langle \hat{\Phi}'(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon), \zeta \rangle = 0, \quad \forall \zeta \in E^-. \quad (4.28)$$

Lemma 4.9. *Suppose that (V0), (V1) and (F1)-(F3) are satisfied. Then for any $\varepsilon > 0$, there exist $K_1(\varepsilon) > 0$ and $K_2(\varepsilon) > 0$ such that*

$$0 \leq t_n^\varepsilon \leq K_1(\varepsilon), \quad \|\zeta_n^\varepsilon\| \leq K_2(\varepsilon), \quad \forall \varepsilon > 0. \quad (4.29)$$

Proof. If along a subsequence $t_n^\varepsilon < T_0$, we are through. So we may assume that $t_n^\varepsilon \geq T_0$. In view of Lemma 4.8, there exists a constant $C_3 > 0$ such that $\|z_n^{\varepsilon+}\|_2 \leq C_3$. If $\delta_\varepsilon := \limsup_{n \rightarrow \infty} \sup_{y \in \mathbb{R}^N} \int_{B_1(y)} |z_n^{\varepsilon+}|^2 dx = 0$, then by Lions' concentration compactness principle [33, Lemma 1.21], $z_n^{\varepsilon+} \rightarrow 0$ in $L^p(\mathbb{R}^N)$. Fix $\vartheta = [(1+\eta^2)(1+\bar{c}_\varepsilon)/(1-\eta)^2 \bar{c}_\varepsilon]^{1/2}$. By virtue of (F1), for $\epsilon = 1/4 Q_{\max}(\vartheta C_3)^2 > 0$, there exists a $C_\epsilon > 0$ such that

$$F(z) \leq \epsilon |z|^2 + C_\epsilon |z|^p, \quad \forall z \in \mathbb{R}^2.$$

Hence, it follows that

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} Q_\varepsilon(x) F(\vartheta z_n^{\varepsilon+}) dx \\ & \leq \epsilon Q_{\max} \vartheta^2 \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |z_n^{\varepsilon+}|^2 dx + C_\epsilon Q_{\max} \vartheta^p \limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} |z_n^{\varepsilon+}|^p dx \\ & \leq \epsilon Q_{\max} (\vartheta C_3)^2 = \frac{1}{4}. \end{aligned} \quad (4.30)$$

From (4.11), (4.25), (4.27) and (4.30), one has

$$\begin{aligned} \bar{c}_\varepsilon + o(1) &= \Phi_\varepsilon(z_n^\varepsilon) \\ &\geq \frac{(1-\eta)ab\vartheta^2}{a^2+b^2} \|z_n^\varepsilon\|^2 - \int_{\mathbb{R}^N} Q_\varepsilon(x) F(\vartheta z_n^{\varepsilon+}) dx \\ &\quad + \frac{1-\vartheta^2}{2} \langle \Phi'_\varepsilon(z_n^\varepsilon), z_n^\varepsilon \rangle + \vartheta^2 \langle \Phi'_\varepsilon(z_n^\varepsilon), z_n^{\varepsilon-} \rangle \\ &\geq \frac{(1-\eta)ab\vartheta^2}{a^2+b^2} \|z_n^{\varepsilon+}\|^2 - \frac{1}{4} + o(1) \\ &\geq \frac{(1-\eta)^2 \bar{c}_\varepsilon \vartheta^2}{1+\eta^2} - \frac{1}{4} + o(1) > \frac{3}{4} + \bar{c}_\varepsilon + o(1). \end{aligned}$$

This contradiction shows that $\delta_\varepsilon > 0$. Going if necessary to a subsequence, we may assume the existence of $k_n \in \mathbb{Z}^N$ such that $\int_{B_{1+\sqrt{N}}(k_n)} |z_n^{\varepsilon+}|^2 dx > \frac{\delta_\varepsilon}{2}$. Let $\tilde{z}_n^\varepsilon(x) = z_n^\varepsilon(x + k_n)$ and $\tilde{\zeta}_n^\varepsilon(x) = \zeta_n^\varepsilon(x + k_n)$. Then $\|\tilde{z}_n^\varepsilon\| = \|z_n^\varepsilon\|$ and

$$\int_{B_{1+\sqrt{N}}(0)} |(\tilde{z}_n^\varepsilon)^+|^2 dx > \frac{\delta_\varepsilon}{2}. \quad (4.31)$$

Passing to a subsequence, we have $\tilde{z}_n^\varepsilon \rightharpoonup \tilde{z}^\varepsilon$ and $(\tilde{z}_n^\varepsilon)^+ \rightharpoonup (\tilde{z}^\varepsilon)^+$ in E , $(\tilde{z}_n^\varepsilon)^+ \rightarrow (\tilde{z}^\varepsilon)^+$ in $L_{\text{loc}}^s(\mathbb{R}^N)$, $2 \leq s < 2^*$ and $(\tilde{z}_n^\varepsilon)^+ \rightarrow (\tilde{z}^\varepsilon)^+$ a.e. on \mathbb{R}^N . Obviously, (4.31) implies that $(\tilde{z}^\varepsilon)^+ \neq 0$. Let $\eta_n^\varepsilon := \tilde{\zeta}_n^\varepsilon / t_n^\varepsilon$. Hence, it follows from (4.2), (4.28), (F1) and (F3) that

$$\begin{aligned}
0 &= \langle \hat{\Phi}'(t_n^\varepsilon \tilde{z}_n^\varepsilon + \zeta_n^\varepsilon), t_n^\varepsilon \tilde{z}_n^\varepsilon + \zeta_n^\varepsilon \rangle = \langle \hat{\Phi}'(t_n^\varepsilon \tilde{z}_n^\varepsilon + \tilde{\zeta}_n^\varepsilon), t_n^\varepsilon \tilde{z}_n^\varepsilon + \tilde{\zeta}_n^\varepsilon \rangle \\
&= \langle \hat{\Phi}'(t_n^\varepsilon (\tilde{z}_n^\varepsilon + \eta_n^\varepsilon)), t_n^\varepsilon (\tilde{z}_n^\varepsilon + \eta_n^\varepsilon) \rangle \\
&= \left[\frac{2ab}{a^2 + b^2} (\|(\tilde{z}_n^\varepsilon)^+\|^2 - \|(\tilde{z}_n^\varepsilon)^- + \eta_n^\varepsilon\|^2) + \hat{V} \int_{\mathbb{R}^N} |\tilde{z}_n^\varepsilon + \eta_n^\varepsilon|^2 dx \right] (t_n^\varepsilon)^2 \\
&\quad - Q(0) \int_{\mathbb{R}^N} F_z(t_n^\varepsilon (\tilde{z}_n^\varepsilon + \eta_n^\varepsilon)) \cdot t_n^\varepsilon (\tilde{z}_n^\varepsilon + \eta_n^\varepsilon) dx \\
&\leq \left[\frac{2ab}{a^2 + b^2} (\|(\tilde{z}_n^\varepsilon)^+\|^2 - \|(\tilde{z}_n^\varepsilon)^- + \eta_n^\varepsilon\|^2) + \hat{V} \int_{\mathbb{R}^N} |\tilde{z}_n^\varepsilon + \eta_n^\varepsilon|^2 dx \right] (t_n^\varepsilon)^2 \\
&\quad - 2Q(0) \int_{\mathbb{R}^N} F(t_n^\varepsilon (\tilde{z}_n^\varepsilon + \eta_n^\varepsilon)) dx \\
&\leq \left[\frac{2ab}{a^2 + b^2} (\|(\tilde{z}_n^\varepsilon)^+\|^2 - \|(\tilde{z}_n^\varepsilon)^- + \eta_n^\varepsilon\|^2) + \hat{V} \int_{\mathbb{R}^N} |\tilde{z}_n^\varepsilon + \eta_n^\varepsilon|^2 dx \right] (t_n^\varepsilon)^2 \\
&\quad - 2\mathcal{C}_0 Q(0) (t_n^\varepsilon)^\mu \int_{\mathbb{R}^N} F_0(\tilde{z}_n^\varepsilon + \eta_n^\varepsilon) dx,
\end{aligned} \tag{4.32}$$

which, together with (4.26), implies that

$$\begin{aligned}
0 &\leq \frac{2ab}{a^2 + b^2} (\|(\tilde{z}_n^\varepsilon)^+\|^2 - \|(\tilde{z}_n^\varepsilon)^- + \eta_n^\varepsilon\|^2) + \hat{V} \int_{\mathbb{R}^N} |\tilde{z}_n^\varepsilon + \eta_n^\varepsilon|^2 dx \\
&\leq \frac{2(1 + \eta^2)ab}{(1 - \eta)(a^2 + b^2)} \|(\tilde{z}_n^\varepsilon)^+\|^2 - \frac{(1 - \eta)ab}{a^2 + b^2} \|(\tilde{z}_n^\varepsilon)^- + \eta_n^\varepsilon\|^2.
\end{aligned} \tag{4.33}$$

This shows that $\{\|\eta_n^\varepsilon\|\}_{n=1}^\infty$ is bounded in E^- . Passing to a subsequence, we have $\eta_n^\varepsilon \rightharpoonup \eta^\varepsilon$ in E^- , $\eta_n^\varepsilon \rightarrow \eta^\varepsilon$ in $L_{\text{loc}}^s(\mathbb{R}^N)$, $2 \leq s < 2^*$ and $\eta_n^\varepsilon \rightarrow \eta^\varepsilon$ a.e. on \mathbb{R}^N .

Since $(\tilde{z}^\varepsilon + \eta^\varepsilon)^+ = (\tilde{z}^\varepsilon)^+ \neq 0$, it follows from (F3) that $\int_{\mathbb{R}^N} F_0(\tilde{z}^\varepsilon + \eta^\varepsilon) dx > 0$, which, together with (4.32), implies that $\{t_n^\varepsilon\}_{n=1}^\infty$ is bounded, and so $\{\|\zeta_n^\varepsilon\|\}_{n=1}^\infty$ is also bounded. Therefore, there exist $K_1(\varepsilon) > 0$ and $K_2(\varepsilon) > 0$ such that (4.29) holds. \square

Theorem 4.10. *Assume that V , Q and F satisfy (V0), (V1) and (F1)-(F3). Then for $\varepsilon \in (0, \varepsilon_0]$, problem (2.1) has a solution $z_\varepsilon \in E$ such that $\Phi_\varepsilon(z_\varepsilon) = \inf_{\mathcal{N}_\varepsilon^-} \Phi_\varepsilon > 0$.*

Proof. By Lemmas 4.7 and 4.8, there exists a bounded sequence $\{z_n^\varepsilon\} = \{(u_n^\varepsilon, v_n^\varepsilon)\} \subset E$ satisfying (4.25). Thus there exists a $z^\varepsilon = (u^\varepsilon, v^\varepsilon) \in E$ such that $z_n^\varepsilon \rightharpoonup z^\varepsilon$. Next, we prove that $z^\varepsilon \neq 0$ for all $\varepsilon \in (0, \varepsilon_0]$.

Arguing by contradiction, suppose that $z^\varepsilon = 0$ for some $\varepsilon \in (0, \varepsilon_0]$, i.e. $z_n^\varepsilon \rightharpoonup 0$ in E , and so $z_n^\varepsilon \rightarrow 0$ in $L_{\text{loc}}^s(\mathbb{R}^N)$, $2 \leq s < 2^*$ and $z_n^\varepsilon \rightarrow 0$ a.e. on \mathbb{R}^N .

We first prove that $\zeta_n^\varepsilon \rightharpoonup 0$ in E^- . Since $\{\|\zeta_n^\varepsilon\|\}_{n=1}^\infty$ is bounded, passing to a subsequence we may assume that $\zeta_n^\varepsilon \rightharpoonup \zeta^\varepsilon$ in E^- . By Lemma 4.9, $\{t_n^\varepsilon\}$ is bounded, then $t_n^\varepsilon \tilde{z}_n^\varepsilon + \zeta_n^\varepsilon \rightharpoonup \zeta^\varepsilon$. By Brezis-Lieb's Lemma (see [33, Lemma 1.32]), one can demonstrate that

$$\hat{\Phi}(t_n^\varepsilon \tilde{z}_n^\varepsilon + \zeta_n^\varepsilon) - \hat{\Phi}(t_n^\varepsilon \tilde{z}_n^\varepsilon + \zeta_n^\varepsilon - \zeta^\varepsilon) - \hat{\Phi}(\zeta^\varepsilon) = o(1), \tag{4.34}$$

which, together with (4.9), (V0) and (V1), yields

$$o(1) \geq -\hat{\Phi}(\zeta^\varepsilon) \geq \frac{ab}{a^2 + b^2} \|\zeta^\varepsilon\|^2 - \frac{\hat{V}}{2} \int_{\mathbb{R}^N} |\zeta^\varepsilon|^2 dx \geq \frac{(1-\eta)ab}{a^2 + b^2} \|\zeta^\varepsilon\|^2. \quad (4.35)$$

This shows that $\zeta^\varepsilon = 0$, i.e. $\zeta_n^\varepsilon \rightharpoonup 0$ in E^- . By (4.2), (4.9), (4.20), (4.25), (4.28) and (F1), we have

$$\begin{aligned} c_\varepsilon + o(1) &\geq \bar{c}_\varepsilon + o(1) = \Phi_\varepsilon(z_n^\varepsilon) \\ &\geq \Phi_\varepsilon(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon) + \frac{1 - (t_n^\varepsilon)^2}{2} \langle \Phi'_\varepsilon(z_n^\varepsilon), z_n^\varepsilon \rangle - t_n^\varepsilon \langle \Phi'_\varepsilon(z_n^\varepsilon), \zeta_n^\varepsilon \rangle \\ &= \Phi_\varepsilon(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon) + o(1) \\ &= \hat{\Phi}(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon) + \frac{1}{2} \int_{\mathbb{R}^N} (V_\varepsilon(x) - \hat{V}) |t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon|^2 dx \\ &\quad + \int_{\mathbb{R}^N} [Q(0) - Q_\varepsilon(x)] F(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon) dx + o(1) \\ &\geq \hat{c} + \frac{1}{2} \int_{|x| \leq R_0/\varepsilon} (V_\varepsilon(x) - \hat{V}) |t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon|^2 dx \\ &\quad + \int_{|x| \leq R_0/\varepsilon} [Q(0) - Q_\varepsilon(x)] F(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon) dx + o(1) \\ &\geq \hat{c} - \frac{2V_{\max} - V_\infty - V(0)}{4} \int_{|x| \leq R_0/\varepsilon} |t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon|^2 dx \\ &\quad + [Q(0) - Q_{\max}] \int_{|x| \leq R_0/\varepsilon} F(t_n^\varepsilon z_n^\varepsilon + \zeta_n^\varepsilon) dx + o(1) \\ &= \hat{c} + o(1). \end{aligned} \quad (4.36)$$

On the other hand, by Lemmas 4.3 and 4.6, one has

$$\hat{c} \geq c_0 + \hat{\delta} \geq c_\varepsilon + \frac{1}{4} \hat{\delta},$$

which contradicts to (4.36). Therefore, $z^\varepsilon \neq 0$ for all $\varepsilon \in (0, \varepsilon_0]$. In a standard way, we can certify that $\Phi'_\varepsilon(z^\varepsilon) = 0$ and $\Phi_\varepsilon(z^\varepsilon) = c_\varepsilon = \inf_{\mathcal{N}_\varepsilon^-} \Phi_\varepsilon$. This shows that $z^\varepsilon \in E$ is a solution for problem (2.1) with $\Phi_\varepsilon(z^\varepsilon) = \inf_{\mathcal{N}_\varepsilon^-} \Phi_\varepsilon > 0$. \square

Proof of Theorem 1.6. For $\varepsilon \in (0, \varepsilon_0]$, Theorem 4.10 implies that problem (2.1) has a solution $z_\varepsilon \in E$ such that $\Phi_\varepsilon(z_\varepsilon) = \inf_{\mathcal{N}_\varepsilon^-} \Phi_\varepsilon > 0$. Then

$$\hat{z}_\varepsilon(x) = (\hat{u}_\varepsilon(x), \hat{v}_\varepsilon(x)) := z_\varepsilon(\varepsilon^{-1}(x - x_v)) = (u_\varepsilon(\varepsilon^{-1}(x - x_v)), v_\varepsilon(\varepsilon^{-1}(x - x_v)))$$

is a nontrivial solution of problem (1.5). \square

References

- [1] Ackermann N. A superposition principle and multibump solutions of periodic Schrödinger equations. J Funct Anal, 2006, 234(2): 277-320

- [2] Ambrosetti A, Colorado E. Standing waves of some coupled nonlinear Schrödinger equations. J Lond Math Soc, 2007, 75: 67-82
- [3] Alves C O, Soares S H M. Singularly perturbed elliptic systems. Nonlinear Anal, 2006, 64: 109-129
- [4] Ávila, A I, Yang J. On the existence and shape of least energy solutions for some elliptic systems. J Differ Equ, 2003, 191: 348-376
- [5] Benci V, Rabinowitz P H. Critical point theorems for indefinite functionals. Invent Math, 1979, 52: 241-273
- [6] Bonheure D, dos Santos E M, Ramos M. Ground state and non-ground state solutions of some strongly coupled elliptic systems. Trans Am Math Soc, 2012, 364: 477-491
- [7] Clément P H, de Figueiredo D G, Mitedieri E. Positive solutions of semilinear elliptic systems. Commun Partial Differ Equ, 1992, 17: 923-940
- [8] de Figueiredo D G. Semilinear elliptic systems: existence, multiplicity, symmetry of solutions. Handbook of differential equations: stationary partial differential equations, vol. V, 1-48, Handb Differ Equ, Elsevier: North-Holland, 2008
- [9] de Figueiredo D G, Felmer P L. On superquadratic elliptic systems. Trans Am Math Soc, 1994, 343: 97-116
- [10] de Figueiredo D G, Yang J. Decay, symmetry and existence of solutions of semilinear elliptic systems. Nonlinear Anal, 1998, 33: 211-234
- [11] Ding Y H, Lee C, Zhao F H. Semiclassical limits of ground state solutions to Schrödinger systems. Calc Var PDE, 2014, 51: 725-760
- [12] Ding Y H, Lin F K. Semiclassical states of Hamiltonian systems of Schrödinger equations with subcritical and critical nonlinearities. J Partial Differ Equ, 2006, 19: 232-255
- [13] Gilbarg D, Trudinger N S. Elliptic Partial Differential Equations of Second Order. Springer: Berlin, 1983
- [14] Hulshof J, Van der Vorst, R C A M. Differential systems with strongly variational structure. J Funct Anal, 1993, 114: 32-58
- [15] Li G, Yang J. Asymptotically linear elliptic systems. Commun Partial Differ Equ, 2004, 29: 925-954
- [16] Liao F F, Tang X H, Zhang J. Existence of solutions for periodic elliptic system with general superlinear nonlinearity, Z Angew Math Phys, 2015, 66: 689-701

- [17] Lin T C, Wei J. Spikes in two-component systems of nonlinear Schrödinger equations with trapping potentials. *J Differ Equ*, 2006, 229, 538-569
- [18] Lions P L. The concentration-compactness principle in the calculus of variations. The locally compact case I, II, *Ann Inst H Poincaré Anal Non Linéaire* 1984, 1: 109-145, 223-283
- [19] Pankov A. Periodic nonlinear Schrödinger equation with application to photonic crystals. *Milan J Math*, 2005, 73: 259-287
- [20] Pistoia A, Ramos M. Locating the peaks of the least energy solutions to an elliptic system with Neumann boundary conditions. *J Differ Equ*, 2004, 201: 160-176
- [21] Rabinowitz P H. On a class of nonlinear Schrödinger equations. *Z Angew Math Phys*, 1992, 43: 270-291
- [22] Ramos M, Soares S H M. On the concentration of solutions of singularly perturbed Hamiltonian systems in \mathbb{R}^N . *Port Math (N S)*, 2006, 63: 157-171
- [23] Ramos M, Tavares H. Solutions with multiple spike patterns for an elliptic system. *Calc Var PDE*, 2008, 31: 1-25
- [24] Ramos M, Yang J. Spike-layered solutions for an elliptic system with Neumann boundary conditions. *Trans Am Math Soc*, 2005, 357: 3265-3284
- [25] Ruf B. Superlinear Elliptic Equations and Systems, *Handbook of differential equations: stationary partial differential equations*, vol. V, *Handb Differ Equ*, Elsevier: North-Holland, pp. 211-276, 2008
- [26] Simon B. Schrödinger semigroups. *Bull Am Math Soc, (N S)*, 1982, 7: 447-526
- [27] Sirakov B, Soares S H M. Soliton solutions to systems of coupled Schrödinger equations of Hamiltonian type. *Trans Am Math Soc*, 2010, 362: 5729-5744
- [28] Szulkin A, Weth T. Ground state solutions for some indefinite variational problems. *J Funct Anal*, 2009, 257: 3802-3822
- [29] Tang X H. Non-Nehari manifold method for superlinear Schrödinger equation. *Taiwanese J Math*, 2014, 18: 1957-1979
- [30] Tang X H. Non-Nehari manifold method for asymptotically linear Schrödinger equation. *J Aust Math Soc*, 2015, 98: 104-116
- [31] Tang X H. Non-Nehari manifold method for asymptotically periodic Schrödinger equations. *Sci China Math*, 2015, 58: 715-728

- [32] Tang X H. Ground state solutions of Nehari-Pankov type for a superlinear Hamiltonian elliptic system on \mathbb{R}^N . Canada Math Bull, 2015, 58: 651-663
- [33] Willem M. Minimax Theorems. Berlin: Birkhäuser, 1996
- [34] Zhang R, Chen J, Zhao F K. Multiple solutions for superlinear elliptic systems of Hamiltonian type. Disc Conti Dyna Syst A, 2011, 30: 29-42
- [35] Zhang J, Qin W P, Zhao F K. Existence and multiplicity of solutions for asymptotically linear nonperiodic Hamiltonian elliptic system. J Math Anal Appl, 2013, 399: 433-441
- [36] Zhang J, Tang X H, Zhang W. Ground-state solutions for superquadratic Hamiltonian elliptic systems with gradient terms. Nonlinear Anal, 2014, 95: 1-10
- [37] Zhao F K, Zhao L G, Ding Y H. Multiple solution for a superlinear and periodic elliptic system on \mathbb{R}^N . Z Angew Math Phys, 2011, 62: 495-511
- [38] Zhao F K, Ding Y H. On Hamiltonian elliptic systems with periodic or non-periodic potentials. J Differ Equ, 2010, 249: 2964-2985
- [39] Zhao L G, Zhao F K. On ground state solutions for superlinear Hamiltonian elliptic systems. Z Angew Math Phys, 2013, 64: 403-418