

TWISTOR LINES IN THE PERIOD DOMAIN OF COMPLEX TORI

NIKOLAY BUSKIN AND ELHAM IZADI

ABSTRACT. As in the case of irreducible holomorphic symplectic manifolds, the period domain $Compl$ of compact complex tori of even dimension $2n$ contains twistor lines. These are special 2-spheres parametrizing complex tori whose complex structures arise from a given quaternionic structure. In analogy with the case of irreducible holomorphic symplectic manifolds, we show that the periods of any two complex tori can be joined by a *generic* chain of twistor lines. We also prove a criterion of twistor path connectivity of loci in $Compl$ where a fixed second cohomology class stays of Hodge type $(1,1)$. Furthermore, we show that twistor lines are holomorphic submanifolds of $Compl$, of degree $2n$ in the Plücker embedding of $Compl$.

CONTENTS

Introduction	1
1. The space of twistor spheres	5
2. Twistor path connectivity of $Compl$	8
3. Connectivity by generic twistor paths	13
4. The degree of twistor lines	21
5. Twistor path connectivity of $Compl_\Omega$	23
References	28

INTRODUCTION

Let M be a Riemannian manifold of real dimension $4m$ with metric g . Then M is called *hyperkähler* with respect to g (see [8, p. 548]) if there exist complex structures I, J and K on M , such that I, J, K are covariantly constant and are isometries of the tangent bundle TM with respect to g , satisfying the quaternionic relations

$$I^2 = J^2 = K^2 = -1, \quad IJ = -JI = K.$$

We call the ordered triple (I, J, K) a *hyperkähler structure on M compatible with g* .

A hyperkähler structure (I, J, K) gives rise to a sphere S^2 of complex structures on M :

$$S^2 = \{aI + bJ + cK \mid a^2 + b^2 + c^2 = 1\}.$$

We call the family $\mathcal{M} = \{(M, \lambda) \mid \lambda \in S^2\} \rightarrow S^2$ a *twistor family over the twistor sphere S^2* . The family \mathcal{M} can be endowed with a complex structure, so that it becomes a complex manifold and the fiber \mathcal{M}_λ is biholomorphic to the complex manifold (M, λ) , see [8, p. 554]. For every $\lambda = aI + bJ + cK \in S^2$, the closed alternating form $g(\lambda \cdot, \cdot)$ determines a Kähler class in $H^{1,1}((M, \lambda), \mathbb{R})$.

The known examples of compact hyperkähler manifolds are even-dimensional complex tori and irreducible holomorphic symplectic manifolds (*IHS manifolds*). We recall that an IHS manifold is a simply connected compact Kähler manifold M with $H^0(M, \Omega_M^2)$ generated by an everywhere non-degenerate holomorphic 2-form. Examples of IHS manifolds include $K3$ surfaces and, more generally, Hilbert schemes of points on $K3$ surfaces, generalized Kummer varieties.

For IHS manifolds and complex tori there exist well-defined period domains, carrying the structure of a complex manifold. Every twistor family \mathcal{M} determines an embedding of the base S^2 into the corresponding period domain as a 1-dimensional complex submanifold (for IHS manifolds this is known and for complex tori we give a proof of this in Theorem 1). The image of such an embedding is called a *twistor line*. The period of a hyperkähler manifold is called *generic*, if the corresponding manifold has trivial Néron-Severi group. A path of twistor lines is an ordered sequence S_1, \dots, S_m of twistor spheres such that $S_i \cap S_{i+1}$ is non-empty if $1 \leq i \leq m-1$. Such a path is called *generic*, if the periods at intersections of successive lines in the path are generic.

In the case of IHS manifolds it is known that any two periods can be connected by a generic path of twistor lines (see [5], which is an exposition of Verbitsky [12]). In [12] generic twistor path connectivity was used to prove surjectivity of the corresponding period mapping, which was a part of the Torelli theorem for IHS manifolds proved there. Another application was given in [6], where generic twistor path connectivity, together with a result of Verbitsky cited below, was used to prove, via deformations of sheaves, that every rational Hodge isometry between two $K3$ -surfaces is algebraic.

The period domain $Compl$ for complex tori of dimension $2n$ is a real analytic open subset of the complex Grassmannian $Gr(2n, 4n)$, see the exact definition in Section 1. It has two connected components $Compl^+$ and $Compl^-$. Our first main result is

Theorem 1. (1) *Any twistor sphere on a complex torus naturally embeds into $Compl$ as a complex 1-dimensional submanifold. The degree of twistor lines in $Gr(2n, 4n)$ with respect to the Plücker embedding is $2n$.*
 (2) *In each of the two connected components of $Compl$ any two periods can be connected by a generic path of twistor lines.*

For an alternating 2-form Ω , we let $Compl_\Omega$ be the locus of all periods $I \in Compl$ such that the form Ω determines a cohomology class of Hodge type (1,1) in the second cohomology of tori with period (i.e., complex structure, see Section 1) I .

We let $G_\Omega \subset G \cong GL_{4n}(\mathbb{R})$ be the group of automorphisms of Ω with G_Ω^0 its connected component of the identity. For $I \in Compl_\Omega$ we define a hermitian form h_I by setting $h_I(u, v) = \Omega(u, Iv) - i\Omega(u, v)$. We let (n_+, n_-, n_0) be the signature of h_I , n_0 depends on Ω only and is the same for all $I \in Compl_\Omega$. We let $U(n_+, n_-, n_0) \subset G_\Omega^0$ be the automorphism group of h_I .

We set $Compl_\Omega^\pm = Compl_\Omega \cap Compl^\pm$. Our second main result is

Theorem 2. (1) *The locus $Compl_\Omega^\pm$ has $2n + 1 - n_0$ connected components, indexed by the signature (n_+, n_-, n_0) of h_I for all periods I in the corresponding component.*

The connected component of Compl_Ω^\pm , where h_I has signature (n_+, n_-, n_0) , is naturally diffeomorphic to the homogeneous manifold $G_\Omega^0/U(n_+, n_-, n_0)$ and is a smooth complex submanifold in Compl .

- (2) *If n_0 is even, there is precisely one connected component of Compl_Ω^\pm , that contains twistor lines. This component corresponds to the signature $(n - \frac{n_0}{2}, n - \frac{n_0}{2}, n_0)$ and is twistor path connected. If n_0 is odd, there are no connected components of Compl_Ω^\pm containing a twistor line.*

An immediate consequence of this theorem is that Kähler classes on complex tori do not stay of Hodge type in twistor families. An analogous result for $K3$ surfaces is well known.

In fact, combining Theorem 2 with the result of Verbitsky (see below) we obtain the following

Corollary 3. *Let M be a hyperkähler manifold and $\omega = g(I\cdot, \cdot)$ a Kähler class on M associated to a complex structure I of the twistor family. If an ω -slope-polystable bundle over M extends to the twistor family \mathcal{M} , then, either its first Chern class is zero, or the hermitian form associated to its first Chern class has signature of the form $(n - \frac{n_0}{2}, n - \frac{n_0}{2}, n_0)$.*

We briefly recall Verbitsky's result. Assume M is a hyperkähler manifold (not necessarily simply connected) with Riemannian metric $g(\cdot, \cdot)$ and a fixed complex structure I , let us denote again by ω the Kähler class on M represented by the form $\omega(\cdot, \cdot) = g(I\cdot, \cdot)$. Then, by definition, we have a sphere of complex structures on M , the corresponding twistor family $\mathcal{M} \rightarrow S^2$, and a Kähler class represented by the form $\omega_\lambda(\cdot, \cdot) = g(\lambda\cdot, \cdot)$ on the fiber $M_\lambda = (M, \lambda)$ for each $\lambda \in S^2$, such that $M_I = M$ and $\omega_I = \omega$. By definition, the class ω_λ (considered up to multiplication by a positive scalar) is the Kähler class on M_λ .

Recall that a vector bundle on M is called ω -slope-polystable if it is isomorphic to a direct sum of ω -slope-stable bundles with equal slopes. The following theorem was proved in [9, Thm. 3.17, Thm. 3.19].

Theorem. *Let F be an ω -slope-polystable vector bundle over a hyperkähler manifold M . If the Chern classes $c_1(F)$ and $c_2(F)$ remain of Hodge type for all complex structures λ on M belonging to the sphere S^2 , then the bundle F extends to a vector bundle \mathcal{F} over \mathcal{M} . Furthermore, for all $\lambda \in S^2$, the restriction $\mathcal{F}|_{M_\lambda}$ is an ω_λ -slope-polystable bundle.*

A purely geometric motivation behind the study of the connectivity, besides the application to deforming sheaves, is discussed in Remark 2 below.

Let us say few words on how proving the connectivity for the period domain of complex tori differs from that for the period domains of IHS manifolds. For IHS manifolds, the proof of connectivity relies on the realization of the period domain as the Grassmanian of oriented positive real 2-planes in the second cohomology, where positivity is with respect to the Beauville-Bogomolov bilinear form, again see [5]. This bilinear form provides a very convenient tool for investigating the local topology of this period domain with respect to the problem of twistor path connectivity.

For complex tori, however, we do not have such a realization of their period domain and cannot use a similar argument. Here, instead, we need to use the (less refined)

structure of the period domain of complex tori as a homogeneous space (which, certainly, the period domain of an IHS manifold is, as well). This homogeneous structure allows us to proceed with proving the twistor path connectivity in steps that are, in their broad strokes, parallel to the steps of the proof of the twistor path connectivity for the period domains of IHS manifolds.

Remark 1. As shown by Beauville [2] (using results of Cheeger and Gromoll), a general compact hyperkähler manifold M has a finite étale cover \widetilde{M} isomorphic to the product of a complex torus and a finite number of irreducible hyperkähler manifolds. Since the irreducible hyperkähler manifolds are simply connected, one can easily see that the Néron-Severi group of \widetilde{M} is isomorphic to the direct sum of the Néron-Severi groups of its factors. Twistor families for M give rise to twistor families for \widetilde{M} and its factors. Vice-versa, twistor families for (any of) the factors of \widetilde{M} give rise, in a generally non-unique way, to twistor families for \widetilde{M} and M . One can then deduce the generic twistor path connectivity of the moduli space of M from the generic twistor path connectivity of the moduli spaces of complex tori and those of irreducible hyperkähler manifolds.

Remark 2. There is a relation between twistor path connectivity and rational connectedness, that is, the connectivity of points of a complex manifold by chains of rational curves (for the latter see, for example, [10]). The Grassmanian $Gr(2n, 4n)$ being a rational variety ($Gr(2n, 4n) \dashrightarrow \mathbb{P}^{4n^2}$), is certainly rationally connected. However, rational connectedness is a weaker property than twistor path connectivity. Indeed, the variety of lines in \mathbb{P}^{4n^2} , passing through a fixed point, has complex dimension $4n^2 - 1$ (and the dimension of the variety of rational curves of degree $d > 1$ in \mathbb{P}^{4n^2} , passing through a fixed point, is even larger), thus its real dimension is $8n^2 - 2$. On the other hand, by Corollary 1.5, the real dimension of the space of all twistor lines, passing through a fixed point in the period domain, is $4n^2 - 1$. Thus, through a given point, there are “half as many” twistor lines as general rational curves, and the problem of twistor path connectivity may be roughly considered as a “sub-Riemannian” version of rational connectedness problem.

The plan of the paper is as follows.

In Section 1 we describe our basic set-up, the complex-analytic structure of $Compl$ considered as a real-analytic submanifold in $End(V_{\mathbb{R}}) \cong End(\mathbb{R}^{4n})$ and show that the twistor spheres $S^2 \subset Compl$ are complex submanifolds (Corollary 1.8). We define the union C_I of all twistor spheres passing through a given period I and show that G_I acts transitively on the set of twistor spheres containing I . The sets C_I , which are real-analytic subsets in $Compl$, will serve as building blocks for constructing twistor paths.

In Section 2 we provide an argument, illustrated by a picture, that the set of periods reachable from a given one I by means of all possible triples of consecutive twistor spheres contains an open neighborhood of the initial period. Then, the connectedness of the period domain allows us to conclude that any two periods can be connected by a path of twistor lines. The three spheres argument is essentially due to the transversality formulated in its most general form in Proposition 2.5, and it is somewhat analogous to the “three lines argument” in [5, Prop. 3.7].

In Section 3 we prove the generic connectivity part of Theorem 1. The idea of the proof is to show that the space of triples of consecutive twistor spheres connecting a fixed pair of periods is not the union of its two subspaces for which the first or, respectively, the second, of the two joint points belongs to the locus of tori with nontrivial Néron-Severi group in the period domain. Again, the transversality, stated in Proposition 2.5, constitutes the main tool for proving generic connectivity.

In Section 4 we prove that the degree of twistor lines in $Gr(2n, 4n)$ with respect to the Plücker embedding is $2n$. Here we use the fact that the group G acts transitively on the set of all twistor lines, preserving the degree, together with an explicit computation on an explicit example.

In Section 5 we prove the statement of Theorem 2, the proof of connectivity is again based on the above mentioned “three lines argument”.

The authors acknowledge debts of gratitude: to Eyal Markman for suggesting the problem of twistor path connectivity of the period domain, and to the referee for the content of Remarks 1.9, 4.3, and pointing out an error in the original calculation of the degree of twistor lines. The authors are indebted to Eyal Markman and the referee for many useful comments that helped improve the exposition of the paper.

1. THE SPACE OF TWISTOR SPHERES

1.1. Let A be a complex torus of dimension $2n$. Denote by $V_{\mathbb{R}}$ the real tangent space $T_{\mathbb{R},0}A$ and by V the complex tangent space $T_{\mathbb{C},0}A \subset T_{\mathbb{R},0}A \otimes \mathbb{C}$, so that $\dim_{\mathbb{R}} V_{\mathbb{R}} = 2 \dim_{\mathbb{C}} V = 4n$.

The points of $Compl$ are $2n$ -dimensional complex planes, realizing the real weight 1 Hodge structures on the complex $4n$ -dimensional vector space $V_{\mathbb{C}} := V_{\mathbb{R}} \otimes \mathbb{C}$. The open subset $Compl$ of $Gr(2n, 4n)$ consists of those $2n$ -planes in $V_{\mathbb{C}}$ that do not intersect the real subspace $V_{\mathbb{R}} \subset V_{\mathbb{C}}$. Explicitly, a complex structure $I: V_{\mathbb{R}} \rightarrow V_{\mathbb{R}}$ corresponds to the $2n$ -plane $(1 - iI)V_{\mathbb{R}} \in Gr(2n, 4n) = Gr(2n, V_{\mathbb{C}})$ where 1 denotes the identity map. As a homogeneous space, $Compl$ is the orbit of I under the conjugation action of $G := GL(V_{\mathbb{R}})$: $Compl \cong G/G_I$, where $G_I \cong GL_{2n}(\mathbb{C})$ is the stabilizer of I . This orbit is endowed with a complex manifold structure such that the above embedding $I \mapsto (1 - iI)V_{\mathbb{R}} \in Gr(2n, 4n)$ is biholomorphic, see [4, p. 31] and Proposition 1.7. The period domain $Compl$ consists of two connected components $Compl^+$ and $Compl^-$, corresponding to the components $GL^+(V_{\mathbb{R}})$ and $GL^-(V_{\mathbb{R}})$ of G .

Assume that $J: V_{\mathbb{R}} \rightarrow V_{\mathbb{R}}$ is a complex structure anticommuting with I . Then I and J determine a twistor sphere

$$S(I, J) := \{aI + bJ + cK | a^2 + b^2 + c^2 = 1\},$$

where $K = IJ$. In general, for two complex structures I_1, I_2 , not necessarily anticommuting, such that $I_1 \neq \pm I_2$, and such that they are contained in the same twistor sphere S , we will also denote this sphere by $S(I_1, I_2)$. Our notation is justified by the following lemma, whose proof is an exercise that we leave to the reader.

Lemma 1.1. *Every twistor sphere S is uniquely determined by any pair of non-proportional complex structures $I_1, I_2 \in S$.*

1.2. Let J be a complex structure that anti-commutes with I . Then $V_{\mathbb{R}}$ splits, in a non-unique way, as a direct sum of 4-dimensional subspaces of the form $\langle v, Iv, Jv, IJv \rangle$ for nonzero vectors $v \in V_{\mathbb{R}}$, and the union of the specified bases of the 4-subspaces

forms a basis of $V_{\mathbb{R}}$. In this basis the matrix of J has a block-diagonal form with the following 4×4 blocks on the diagonal

$$\left(\begin{array}{cc|cc} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ \hline 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{array} \right).$$

Proposition 1.2. *Given a triple of complex structures (I, J, K) on A satisfying the quaternionic identities, there exist a (non-unique) metric g on A such that (I, J, K) is a hyperkähler structure with respect to g .*

Proof. Choose a basis of $V_{\mathbb{R}}$ as in Paragraph 1.2 and define a metric $g(\cdot, \cdot)$ on $V_{\mathbb{R}}$ by declaring this basis to be orthonormal. Then I, J and K are isometries with respect to $g(\cdot, \cdot)$ and g is Kähler with respect to all three complex structures. \square

1.3. By the definition of *Compl*, the group G acts transitively on it:

$$g \in G: J \mapsto {}^g J = gJg^{-1}.$$

In particular, G acts on the set of all twistor spheres $S(I, J)$ in *Compl*:

$$g \cdot S(I, J) = S({}^g I, {}^g J).$$

For $g \in G_I$ we have $g \cdot S(I, J) = S(I, {}^g J)$. We have

Proposition 1.3. *The group G_I acts transitively on the set N_I of complex structures anticommuting with I .*

Proof. Let J be a complex structure that anti-commutes with I . The group $G_I \cong GL(V) < GL^+(V_{\mathbb{R}}) = GL_{4n}^+(\mathbb{R})$ acts transitively on the set of bases as in Paragraph 1.2, hence also on the set of J anti-commuting with I . \square

1.4. Therefore, given a complex structure $J \in N_I$, $N_I = G_I \cdot J \cong G_I/G_{I,J}$ is the orbit of J under G_I , where $G_{I,J}$ is the stabilizer group of J in G_I . Since $G_{I,J}$ is the subgroup of elements of $G_I = GL(V)$ commuting with J , that is, preserving the quaternionic structure on $V_{\mathbb{R}}$ determined by I and J , we have $G_{I,J} \cong GL(V, \mathbb{H})$ which we will also denote by $G_{\mathbb{H}}$. So $N_I \cong GL(V)/GL(V, \mathbb{H})$ and we deduce

Corollary 1.4. *The set N_I is a real submanifold of *Compl* of dimension $4n^2$.*

Proof. The dimension of the orbit as a complex manifold is $\dim_{\mathbb{C}} GL(V) - \dim_{\mathbb{C}} GL(V, \mathbb{H}) = (2n)^2 - 2n^2 = 2n^2$. The real dimension is thus equal to $4n^2$. \square

1.5. Let $S = S(I, J)$ for $J \in N_I$ be a twistor sphere. Define $G_{I,S} \subset G_I$ to be the stabilizer of S as a set, i.e., the set of elements g of G_I such that $g \cdot S \subset S$. For any $g \in G_{I,S}$, the complex structure ${}^g J \in S$ also anticommutes with I , so ${}^g J$ is of the form $aJ + bK$, $a^2 + b^2 = 1$. Setting $a = \cos t$, $b = \sin t$ we have $aJ + bK = e^{\frac{tI}{2}} J e^{-\frac{tI}{2}}$, where $e^{sI} = \cos s \mathbb{1} + \sin s I \in G_I$ realizes, via the conjugation action, the rotations of S around $\{\pm I\}$. Conversely, if $g \in G_I$ and ${}^g J \in S$, then $g \in G_{I,S}$. The set of $g \in G_{I,S}$ such that ${}^g J = J$ is the quaternionic subgroup $G_{I,J} = G_{\mathbb{H}} \subset G_{I,S}$. Explicitly, we have $G_{I,S} = \langle e^{tI}, t \in \mathbb{R} \rangle \times G_{\mathbb{H}}$, where $\langle e^{tI}, t \in \mathbb{R} \rangle \cong S^1$ (which is a subgroup of the center of G_I). This tells us, in particular, that $\dim_{\mathbb{R}} G_{I,S} = \dim_{\mathbb{R}} G_{\mathbb{H}} + 1 = 4n^2 + 1$.

Let M_I be the set of all twistor spheres in $Compl$ containing I . The natural map $N_I \rightarrow M_I$ identifies two complex structures J_1 and J_2 whenever they belong to the same twistor sphere through I , i.e., $S(I, J_1) = S(I, J_2)$. More precisely, they belong to the great circle in $S := S(I, J_1)$ consisting of elements anticommuting with I . Hence, for the S^1 -action $J \in N_I \mapsto e^{tI}J = e^{\frac{tI}{2}}Je^{-\frac{tI}{2}}$ on N_I defined above, we have $N_I/S^1 = M_I$. Therefore Corollary 1.4 immediately implies

Corollary 1.5. *The set M_I is a real manifold of dimension $4n^2 - 1$.*

1.6. The twistor cone of I . Define the set $C_I := \bigcup_{S \in M_I} S \subset Compl$ as the union of all twistor spheres containing I . All spheres in this union contain the complex structures I and $-I$. We will sometimes refer to the set C_I as a (twistor) cone. Proposition 1.3 immediately implies

Corollary 1.6. *The group G_I acts transitively on $M_I \cong G_I/G_{I,S}$ so that $C_I = \bigcup_{g \in G_I} g \cdot S(I, J)$.*

We will give an explicit local parametrization of C_I in the next section and prove that the cone C_I is a real-analytic subset of $Compl$ of dimension $4n^2 + 1$ (Proposition 2.6).

1.7. We now describe the complex structure on the tangent bundle of the orbit $Compl = G \cdot I$. Then we will see that the tangent bundle TS^2 of an arbitrary twistor sphere $S^2 \subset Compl$ is a subbundle of the restricted tangent bundle $TCompl|_{S^2}$, invariant under the complex structure of $TCompl|_{S^2}$. This will imply the well-known fact that the twistor sphere S^2 is a complex submanifold in $Compl$.

Proposition 1.7. *The submanifold $Compl \subset End(V_{\mathbb{R}})$ is a complex manifold. Its complex structure l_I is given by left multiplication by I on $T_I Compl \subset End(V_{\mathbb{R}})$.*

The complex structure of $Compl$ is induced by that of $Gr(2n, 4n)$ via the embedding $I \mapsto (\mathbb{1} - iI)V_{\mathbb{R}}$. The proof of Proposition 1.7 is a technical exercise that we leave to the reader.

Proof. Denoting the differential of the embedding $Compl \ni I \mapsto (\mathbb{1} - iI)V_{\mathbb{R}} \in Gr(2n, 4n)$ by φ we have the following commutative diagram

$$\begin{array}{ccc} T_I Compl \ni X & \xrightarrow{\varphi} & \frac{d}{dt}\bigg|_{t=0} (\mathbb{1} - i e^{tX} I)V_{\mathbb{R}} \in T_{(\mathbb{1}-iI)V_{\mathbb{R}}} Gr(2n, 4n) \\ \downarrow l_I & & \downarrow i \times \\ T_I Compl \ni Y & \xrightarrow{\varphi} & \frac{d}{dt}\bigg|_{t=0} (\mathbb{1} - i e^{tY} I)V_{\mathbb{R}} \in T_{(\mathbb{1}-iI)V_{\mathbb{R}}} Gr(2n, 4n) \end{array}$$

where l_I denotes the complex structure operator on $T_I Compl$ and ' $i \times$ ' denotes the multiplication by i on

$$\mathbb{C}^{4n^2} \cong Hom((\mathbb{1} - iI)V_{\mathbb{R}}, V_{\mathbb{C}}/(\mathbb{1} - iI)V_{\mathbb{R}}) = T_{(\mathbb{1}-iI)V_{\mathbb{R}}} Gr(2n, 4n),$$

so that $\varphi \circ l_I = i\varphi$.

We note that $T_I Compl \cong T_e G/T_e G_I$ and, as $T_e G_I$ consists of all operators in $End(V_{\mathbb{R}})$ commuting with I , the tangent space $T_I Compl$ can be identified with the

subspace of operators in $\text{End}(V_{\mathbb{R}})$ anticommuting with I . Indeed, every operator $X \in \text{End}(V_{\mathbb{R}})$ can be written as a sum of an operator anticommuting with I and an operator commuting with I , $X = \frac{1}{2}(X - X^I) + \frac{1}{2}(X + X^I)$, where $X^I = IXI^{-1}$. This allows us to immediately assume that X and Y in the above diagram anticommute with I .

Now we evaluate

$$\varphi(X) = \left. \frac{d}{dt} \right|_{t=0} (\mathbb{1} - i e^{tX} I) V_{\mathbb{R}} = \{(\mathbb{1} - iI)v \mapsto -i(XI - IX)v = -2iXIv \mid v \in V_{\mathbb{R}}\},$$

which, after multiplying by i becomes $i\varphi(X) = \{(\mathbb{1} - iI)v \mapsto 2XIv \mid v \in V_{\mathbb{R}}\}$ (here we slightly abuse notation by writing, instead of the actual $\varphi(X), i\varphi(X)$, their representatives in $\text{Hom}((\mathbb{1} - iI)V_{\mathbb{R}}, V_{\mathbb{C}})$).

Now, considering $\varphi(Y) = \{(\mathbb{1} - iI)v \mapsto -2iYIv = -2YiIv \mid v \in V_{\mathbb{R}}\}$ as a vector in $\text{Hom}((\mathbb{1} - iI)V_{\mathbb{R}}, V_{\mathbb{C}}/(\mathbb{1} - iI)V_{\mathbb{R}})$, we may write $\varphi(Y) = \{(\mathbb{1} - iI)v \mapsto -2Yv \mid v \in V_{\mathbb{R}}\}$. In order to have the equality $\varphi \circ l_I = i\varphi$, setting $Y = l_I(X)$ we write

$$\varphi(Y) = i\varphi(X) = \{(\mathbb{1} - iI)v \mapsto 2XIv = -2IXv \mid v \in V_{\mathbb{R}}\}.$$

In order for the latter equality to be true it is necessary and sufficient that $Y = IX$, that is, the map l_I is the left multiplication by I on $T_I \text{Compl}$. \square

Corollary 1.8. *The twistor spheres $S^2 \subset \text{Compl}$ are complex submanifolds.*

Proof. The proof is based on the simple observation that the tangent space of $S^2 = S(I, J)$ at the point I , for $I, J, K = IJ$ satisfying the quaternionic identities, is the 2-plane $\langle J, K \rangle_{\mathbb{R}} \subset T_I \text{Compl}$ and this plane is obviously invariant under left multiplication by I . Thus, TS^2 is a complex subbundle of $T\text{Compl}|_{S^2}$ and thus $S^2 \subset \text{Compl}$ is a complex submanifold. \square

Remark 1.9. As was pointed out to us by the referee, there is an alternative proof of Corollary 1.8 that follows from considering $S \subset \text{Compl}$ as a subset in $\text{Gr}(2n, V_{\mathbb{C}})$. Namely, denoting by \mathbb{H} the algebra of quaternions and fixing a representation $\mathbb{H} \rightarrow \text{End}(V_{\mathbb{R}})$ defined by $S = S(I, J)$, we obtain a structure of an \mathbb{H} -module on $V_{\mathbb{R}}$. This \mathbb{H} -module is of the form $\mathbb{H} \otimes V'$ for an n -dimensional \mathbb{R} -vector space V' . The eigenspace $V^{1,0} \subset V_{\mathbb{C}}$ for a complex structure induced by the action of \mathbb{H} is of the form $\mathbb{H}^{1,0} \otimes V'_{\mathbb{C}}$, where $V'_{\mathbb{C}} = V' \otimes \mathbb{C}$ and $\mathbb{H}^{1,0}$ is the corresponding eigenspace in $\mathbb{H} \otimes \mathbb{C}$. Taking the tensor product with $V'_{\mathbb{C}}$ defines a complex analytic embedding $i: \text{Gr}(2, \mathbb{H} \otimes \mathbb{C}) \hookrightarrow \text{Gr}(2n, V_{\mathbb{C}})$. Thus, every twistor line in $\text{Gr}(2n, V_{\mathbb{C}})$ is the image of a twistor line in $\text{Gr}(2, \mathbb{H} \otimes \mathbb{C})$ under some embedding i as above. Now, the twistor lines in the quadric (under the Plücker embedding) $\text{Gr}(2, \mathbb{H} \otimes \mathbb{C})$ are known to be obtained as linear subspace sections, thus they are complex analytic submanifolds. Hence, our $S \subset \text{Gr}(2n, 4n)$ is a complex analytic submanifold.

2. TWISTOR PATH CONNECTIVITY OF Compl

The main result of this section is Theorem 2.3. Before proving it we need to introduce a certain mapping and prove an important technical result about it (Proposition 2.1).

2.1. Let I, J, K be a triple of complex structures belonging to a twistor sphere S . Consider the smooth mapping

$$\begin{aligned} \Phi: G_J \times G_K &\longrightarrow Compl, \\ (g_1, g_2) &\longmapsto {}^{g_1 g_2} I, \end{aligned}$$

where, as before, the action on $Compl$ is by conjugation: $g \cdot I = {}^g I = g I g^{-1}$. The mapping Φ clearly sends $G_{\mathbb{H}} \times G_{\mathbb{H}}$ to I , so that its differential $d\Phi_{(e,e)}$ factors through

$$\widetilde{d\Phi_{(e,e)}} : T_e G_J / T_e G_{\mathbb{H}} \oplus T_e G_K / T_e G_{\mathbb{H}} \rightarrow T_I Compl.$$

Proposition 2.1. *Suppose I, J, K is a quaternionic triple. The mapping*

$$\widetilde{d\Phi_{(e,e)}} : T_e G_J / T_e G_{\mathbb{H}} \oplus T_e G_K / T_e G_{\mathbb{H}} \rightarrow T_I Compl$$

is an isomorphism.

Before proving Proposition 2.1 we make the following useful observation. For a vector $X \in T_e G \cong \text{End } V_{\mathbb{R}}$ and $I \in Compl$ we introduce the notation $X^I = I^{-1} X I = I X I^{-1} \in T_e G$. Every vector $X \in T_e G$ can be uniquely decomposed into the sum of its I -commuting and I -anticommuting components

$$X = \frac{1}{2}(X + X^I) + \frac{1}{2}(X - X^I),$$

so that interpreting the subspace $T_e G_I \subset T_e G$ as the subspace $\{Y \in T_e G \mid Y I = I Y\}$ of I -commuting vectors in $T_e G$, we get the natural isomorphism

$$(1) \quad T_I Compl \cong T_e G / T_e G_I \cong \{Y \in T_e G \mid Y I = -I Y\},$$

of $T_I Compl$ with the subspace of I -anticommuting vectors in $T_e G$. Similarly we may write

$$T_e G_J / T_e G_{\mathbb{H}} \cong \{Y \in T_e G \mid Y I = -I Y, Y J = J Y\}$$

and

$$T_e G_K / T_e G_{\mathbb{H}} \cong \{Y \in T_e G \mid Y I = -I Y, Y K = K Y\}.$$

Proof of Proposition 2.1. By the definition of $\widetilde{d\Phi_{(e,e)}}$, its restrictions to the above direct summands are injective. Let us show that it is injective on the direct sum. Consider $X \in T_e G_J, Y \in T_e G_K$ and the vector $\widetilde{d\Phi_{(e,e)}}(X + T_e G_{\mathbb{H}}, Y + T_e G_{\mathbb{H}})$, which is

$$d\Phi_{(e,e)}(X + Y) = \left. \frac{d}{dt} \right|_{t=0} (e^{tX} e^{tY} \cdot I) = (X + Y)I - I(X + Y) \in T_I Compl.$$

Assume that this vector is zero, that is, $X + Y$ commutes with I :

$$(2) \quad I(X + Y) = (X + Y)I.$$

Then the conjugate $(X + Y)^J = J^{-1}(X + Y)J = X^J + Y^J = X - JYJ$ must also commute with I . Using that Y commutes with K we obtain

$$X - JYJ = X - JYKI = X - JKYI = X - IYI.$$

The commutation with I is expressed now by $I(X - IYI) = (X - IYI)I$, or

$$IX + YI = XI + IY,$$

which gives

$$I(X - Y) = (X - Y)I.$$

Adding the last equality to (2) side by side gives that $XI = IX$, hence $YI = IY$, which implies $X, Y \in T_e G_{\mathbb{H}}$. This proves the required injectivity of $\widetilde{d\Phi_{(e,e)}}$, which, by a slight abuse of notation, we may consider as the inclusion

$$T_e G_J / T_e G_{\mathbb{H}} \oplus T_e G_K / T_e G_{\mathbb{H}} \subset T_I \text{Compl}.$$

Now the surjectivity of $\widetilde{d\Phi_{(e,e)}}$ is equivalent to another inclusion

$$T_I \text{Compl} \subset T_e G_J / T_e G_{\mathbb{H}} \oplus T_e G_K / T_e G_{\mathbb{H}}.$$

Using the natural isomorphism $T_I \text{Compl} \cong \{Y \in T_e G \mid YI = -IY\}$ in (1), we may decompose an arbitrary $Y \in T_I \text{Compl}$ into the sum of its J -commuting and J -anticommuting components, $Y = \frac{1}{2}(Y + Y^J) + \frac{1}{2}(Y - Y^J)$, each of which anticommutes with I and hence determines a tangent vector in $T_I \text{Compl}$. Now, since

$$Y - \frac{1}{2}(Y + Y^J) = \frac{1}{2}(Y - Y^J)$$

anticommutes with both I and J , it commutes with $K = IJ$, while $Y + Y^J$ anticommutes with K , so that averaging under K both sides of the last equality we get $\frac{1}{2}(Y - Y^J) = \frac{1}{2}(Y + Y^K)$. Finally

$$Y = \frac{1}{2}(Y + Y^J) + \frac{1}{2}(Y + Y^K) \in T_e G_J / T_e G_{\mathbb{H}} \oplus T_e G_K / T_e G_{\mathbb{H}},$$

which proves the required inclusion and thus the surjectivity of $\widetilde{d\Phi_{(e,e)}}$. \square

Corollary 2.2. *Suppose I, J, K is a quaternionic triple. The mapping Φ is a submersion at $(e, e) \in G_J \times G_K$, that is*

$$d\Phi_{(e,e)}(T_e G_J \oplus T_e G_K) = T_I \text{Compl}.$$


Proof. The statement that Φ is a submersion follows from factoring $d\Phi_{(e,e)}$ through $\widetilde{d\Phi_{(e,e)}}$ and the fact that the mapping $\widetilde{d\Phi_{(e,e)}} : T_e G_J / T_e G_{\mathbb{H}} \oplus T_e G_K / T_e G_{\mathbb{H}} \rightarrow T_I \text{Compl}$ is an isomorphism by Proposition 2.1. \square

Theorem 2.3. *Given a complex structure $I \in \text{End}(V_{\mathbb{R}})$, there is a neighborhood of I in the space of complex structures on $V_{\mathbb{R}}$ such that, for any complex structure I_1 in this neighborhood, there is a twistor path consisting of three spheres joining I to I_1 . Consequently, each connected component of Compl is twistor path connected.*

Proof. Choose a complex structure J , anticommuting with I , and consider the sphere $S = S(J, I)$ and the cone C_J . By Lemma 1.1, the complex structures $K = IJ$ and I span the sphere $S = S(K, I) = S(J, I)$. We can then form the cone C_K whose intersection with C_J contains S . See Picture 1 below where the cones C_J and C_K are depicted by transversal planes and the sphere S lying in their intersection is depicted by a line.

We first show that the images of C_K under the action of G_J (“rotation of C_K around J ”) sweep out an open neighborhood of I in Compl . Since Φ is a submersion by Corollary 2.2, there exist neighborhoods $U_{e,J} \subset G_J$ and $U_{e,K} \subset G_K$ of e such that the set $\Phi(U_{e,J} \times U_{e,K})$ contains an open neighborhood of I . By definition, the cone C_K contains the orbit $G_K \cdot I$. Hence the union $\bigcup_{g \in G_J} {}^g C_K$ contains the image of Φ and consequently it contains an open neighborhood of I .

Now the three twistor spheres connecting I to an arbitrary point I_1 in this neighborhood are found as illustrated in the following picture.



Picture 1.

Finally we conclude that each of the two connected components of $Compl$ is twistor path connected. \square

2.2. Another immediate consequence of the injectivity of $\widetilde{d\Phi_{(e,e)}}$ proved in Proposition 2.1 is the following

Corollary 2.4. *For a quaternionic triple I, J, K , the triple intersection of the submanifolds $G_I/G_{\mathbb{H}}, G_J/G_{\mathbb{H}}$ and $G_K/G_{\mathbb{H}}$ of the homogeneous space $G/G_{\mathbb{H}}$ at $eG_{\mathbb{H}}$ is transversal.*

The following generalization of this transversality is one of the main ingredients of the proof of connectivity by generic twistor paths in Section 3.

Proposition 2.5. *Let I_1, I_2, I_3 be complex structures belonging to the same twistor sphere S . The submanifolds $G_{I_1}/G_{\mathbb{H}}, G_{I_2}/G_{\mathbb{H}}, G_{I_3}/G_{\mathbb{H}}$ in $G/G_{\mathbb{H}}$ intersect transversally (as a triple) if and only if I_1, I_2, I_3 are linearly independent as vectors in $\text{End}(V_{\mathbb{R}})$.*

Proof. Choose anticommuting complex structures I, J in S , and set $K = IJ$. By Corollary 2.4,

$$(3) \quad T_e G / T_e G_{\mathbb{H}} = V_I \oplus V_J \oplus V_K,$$

where we set $V_I := T_e G_I / T_e G_{\mathbb{H}}, V_J := T_e G_J / T_e G_{\mathbb{H}}, V_K := T_e G_K / T_e G_{\mathbb{H}}$.

We shall prove that $T_e G / T_e G_{\mathbb{H}}$ also decomposes into the direct sum of its subspaces $V_i := T_e G_{I_i} / T_e G_{\mathbb{H}}, i = 1, 2, 3$. Put $I_i = a_i I + b_i J + c_i K, i = 1, 2, 3$. Assume, on the contrary, that for certain vectors $X \in V_1, Y \in V_2$ and $Z \in V_3$ we have $X + Y + Z = 0$. Let $X := X_I + X_J + X_K$ be the decomposition of X into the sum of its components in the respective subspaces of (3), and do similarly for Y and Z . Then for X the commutation relation $[X, I_1] = 0$ can be written as

$$a_1[X_J + X_K, I] + b_1[X_I + X_K, J] + c_1[X_I + X_J, K] = 0.$$

Note that in the above expression, the term $[X_J, I]$, for example, anticommutes with both I, J , hence commutes with $K = IJ$, and an analogous commutation relation holds for the other terms as well. Hence we can decompose the expression on the left side of the above equality with respect to (3):

$$(b_1[X_K, J] + c_1[X_J, K]) + (a_1[X_K, I] + c_1[X_I, K]) + (a_1[X_J, I] + b_1[X_I, J]) = 0.$$

From here we conclude that $b_1[X_K, J] + c_1[X_J, K] = 0$, $a_1[X_K, I] + c_1[X_I, K] = 0$ and $a_1[X_J, I] + b_1[X_I, J] = 0$. Perturbing the quaternionic triple I, J, K , we may assume that all $a_i, i = 1, 2, 3$, are nonzero. Then we can use the last two equalities to express

$$(4) \quad [X_J, I] = -\frac{b_1}{a_1}[X_I, J], \quad [X_K, I] = -\frac{c_1}{a_1}[X_I, K].$$

Note that $F_J := [\cdot, J]: V_I \rightarrow V_K$, $F_K := [\cdot, K]: V_I \rightarrow V_J$ and $F_I := [\cdot, I]: V_J \rightarrow V_K$ are isomorphisms of the respective vector spaces. Then, using (4), we can write

$$X_J = -\frac{b_1}{a_1}F_I^{-1} \circ F_J(X_I), \quad X_K = -\frac{c_1}{a_1}F_I^{-1} \circ F_K(X_I),$$

so that

$$X = X_I + \left(-\frac{b_1}{a_1}F_I^{-1} \circ F_J(X_I)\right) + \left(-\frac{c_1}{a_1}F_I^{-1} \circ F_K(X_I)\right).$$

Using $a_2, a_3 \neq 0$, we obtain similar expressions for Y and Z . Since F_I, F_J, F_K are isomorphisms, the equality $X + Y + Z = 0$ can now be written as

$$\begin{pmatrix} 1 & 1 & 1 \\ -\frac{b_1}{a_1} & -\frac{b_2}{a_2} & -\frac{b_3}{a_3} \\ -\frac{c_1}{a_1} & -\frac{c_2}{a_2} & -\frac{c_3}{a_3} \end{pmatrix} \begin{pmatrix} X_I \\ Y_I \\ Z_I \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

This has a nontrivial solution if and only if the columns of the matrix, i.e., I_1, I_2, I_3 , are linearly dependent. \square

2.3. We can now prove that the cone C_I has a real analytic structure. Define the incidence correspondence

$$S_I := \{(S, J) \mid J \in S\} \subset M_I \times \text{Compl}.$$

Then S_I is an S^2 -bundle over M_I and C_I is the image of S_I by the projection to Compl :

$$\begin{array}{ccccc} N_I & \hookrightarrow & S_I & \xrightarrow{pr_2} & C_I \subset \text{Compl} \\ & & \searrow & \downarrow pr_1 & \\ & & & M_I. & \end{array}$$

The projection $S_I \rightarrow M_I$ has two sections σ_+ and σ_- , given by $+I$ and $-I$ respectively.

Proposition 2.6. *The real-analytic map $pr_2 : S_I \rightarrow C_I$ is a diffeomorphism away from the images of σ_{\pm} and contracts these images to points. Therefore the cone C_I is a real-analytic subset of Compl of dimension $4n^2 + 1$, smooth away from the points $\pm I$.*

Proof. First note that pr_2 clearly contracts the images of σ_{\pm} . Also, it is injective away from $\pm I$ by Lemma 1.1. To see that it is also an immersion away from $\pm I$, let J be in $C_I \setminus \{\pm I\}$, not necessarily anticommuting with I . Define the following mapping

$$\begin{aligned}\Phi: (T_e G_I / T_e G_{\mathbb{H}}) \times \mathbb{R} &\rightarrow C_I, \\ (X, t) &\mapsto e^X e^{tK} J e^{-tK} e^{-X},\end{aligned}$$

where $K \in S(I, J) \setminus S^1$ for $S^1 = \langle I, J \rangle_{\mathbb{R}} \cap S(I, J)$. Then the restriction of Φ on a small enough neighborhood of $(0, 0) \in (T_e G_I / T_e G_{\mathbb{H}}) \times \mathbb{R}$ defines a parametrization of C_I around J .

Here the subgroup $e^{tK}, t \in \mathbb{R}$, rotates the sphere $S = S(I, J)$ around the axis $\{\pm K\}$ and, together with the rotation subgroup $e^{tI} \subset G_{I,S} \subset G_I$, sweeps out in S , via the above action, a neighborhood of any point of S other than $\pm I, \pm K$. Proposition 2.5 provides that K may be chosen arbitrarily in $S \setminus S^1$, which in turn gives us that C_I is a manifold, smooth away from $\pm I$, of dimension $\dim_{\mathbb{R}}(G_I / G_{I,S}) + \dim_{\mathbb{R}} S = (4n^2 - 1) + 2 = 4n^2 + 1$. The fact that the points $\pm I$ are indeed singular points of the cone C_I is easy to prove. \square

3. CONNECTIVITY BY GENERIC TWISTOR PATHS

Recall that a period in *Compl* is *generic* if the corresponding complex torus has trivial Néron-Severi group. A twistor path in *Compl* is called *generic*, if its successive twistor spheres intersect at generic periods. In this section we prove the connectivity part of Theorem 1, i.e.,

Proposition 3.1. *Any two periods in the period domain *Compl* can be connected by a generic twistor path.*

In this section, with the exception of Lemma 3.3 and its proof, we do not assume that the complex structures I, J, K (with or without subscripts) anticommute.

3.1. Outline of the proof. Define \mathcal{T} to be the closure, in $\text{Compl} \times \text{Compl} \times \text{Compl}$, of the set of triples (I, J, K) that are linearly independent and belong to the same twistor sphere. Denote by

$$\begin{aligned}pr_1: \quad \text{Compl} \times \text{Compl} \times \text{Compl} &\longrightarrow \text{Compl}, \\ pr_{23}: \quad \text{Compl} \times \text{Compl} \times \text{Compl} &\longrightarrow \text{Compl} \times \text{Compl}\end{aligned}$$

the respective projections. For $(I_1, J_1, K_1) \in \mathcal{T}$, we defined, in Paragraph 2.1, the mapping Φ_{I_1, J_1, K_1} :

$$\begin{aligned}\Phi_{I_1, J_1, K_1}: \quad G_{J_1} \times G_{K_1} &\longrightarrow \text{Compl}, \\ (g_1, g_2) &\longmapsto g_1 g_2 I_1 g_2^{-1} g_1^{-1} = {}^{g_1 g_2} I_1.\end{aligned}$$

Proposition 2.5 tells us that, when I_1, J_1, K_1 are linearly independent, Φ_{I_1, J_1, K_1} is a submersion near $(e, e) \in G_{J_1} \times G_{K_1}$. In other words, there is a neighborhood $U_{e,G} \subset G = GL^+(V_{\mathbb{R}})$ of $e \in G$ such that the map Φ_{I_1, J_1, K_1} is submersive on $U_{e,J_1} \times U_{e,K_1}$, where $U_{e,J_1} := U_{e,G} \cap G_{J_1}$ and $U_{e,K_1} := U_{e,G} \cap G_{K_1}$ (and the image is, thus, a neighborhood of I_1 in *Compl*).

Let I_2 be an arbitrary point in the image of Φ_{I_1, J_1, K_1} and let $(g_1, g_2) \in U_{e,J_1} \times U_{e,K_1}$ be such that $I_2 = {}^{g_1 g_2} I_1$. With this notation, the three twistor spheres connecting I_1

to I_2 are: $S_1 := S(I_1, J_1, K_1)$, $S := {}^{g_1}S_1 = S({}^{g_1}I_1, {}^{g_1}J_1 = J_1, {}^{g_1}K_1)$ and $S_2 := {}^{g_1g_2}S_1 = S(I_2, {}^{g_1g_2}J_1, {}^{g_1g_2}K_1 = {}^{g_1}K_1)$, with the joint points J_1 and ${}^{g_1}K_1$.

We are going to show that, for a fixed I_1 , there is a neighborhood $U_{I_1} \subset \text{Compl}$ of I_1 such that for any $I_2 \in U_{I_1}$, we can choose a generic $J \in C_I$, a $K \in S(I, J)$ and find $(g_1, g_2) \in \Phi_{I,J,K}^{-1}(I_1)$ as above such that ${}^{g_1g_2}K$ is also generic.

We begin by proving, in Lemma 3.4, that the set of non-generic periods in C_{I_1} is a countable union of proper analytic subsets, i.e., J can be chosen generic.

Next, for I_2 close to I_1 , and with S_1, S, S_2 as above, connecting I_1 to I_2 , the initial sphere S_1 together with the choice of $J, K \in S_1$, uniquely determines the final sphere S_2 together with the pair of periods ${}^{g_1g_2}J, {}^{g_1g_2}K$.

To justify this uniqueness we first need to control the fibers of the maps $\Phi_{I,J,K}$ in a neighborhood of (I_1, J_1, K_1) , which we do in Lemma 3.5. This allows us to introduce, in Paragraphs 3.4 and 3.7, two maps $\Psi^{I_1 \rightarrow I_2}$ and $\Psi^{I_2 \rightarrow I_1}$ which, roughly speaking, switch (S_1, J, K) and $(S_2, {}^{g_1g_2}J, {}^{g_1g_2}K)$.

We then show in Lemma 3.9, after shrinking our various domains, that the composition of $\Psi^{I_1 \rightarrow I_2}$ and $\Psi^{I_2 \rightarrow I_1}$ is the identity. Corollary 3.10 then shows that this implies the irreducibility of the set of triples (S_1, S, S_2) joining I_1 and I_2 mentioned in the introduction, which gives that J and ${}^{g_1g_2}K$ can both be chosen generic.

Thus the chain of three twistor spheres connecting I_1 to I_2 for every I_2 in some neighborhood of I_1 can be chosen in such a way that the periods at the intersections are generic. For arbitrary I_1 and I_2 , we connect them by a path in Compl consisting of generic triple subchains.

3.2. Let us first show that there are generic periods $J \in C_I$. Dimension-wise this is not trivial because $\dim_{\mathbb{R}} C_I = 4n^2 + 1$, whereas the real dimension of the locus of, for example, abelian varieties in Compl is $4n^2 + 2n$. For an alternating form Ω on $V_{\mathbb{R}}$ we denote by Compl_{Ω} the locus of periods in Compl at which Ω represents a class of Hodge $(1,1)$ -type, that is

$$\text{Compl}_{\Omega} = \{I \in \text{Compl} \mid \Omega(I \cdot, I \cdot) = \Omega(\cdot, \cdot)\}.$$

If we fix a basis of $V_{\mathbb{R}}$ and switch to matrix descriptions, then the condition $\Omega(I \cdot, I \cdot) = \Omega(\cdot, \cdot)$ simply becomes ${}^t I \Omega I = \Omega$, where I and Ω also denote the matrices of the corresponding complex structure and alternating form. The locus of marked complex tori with nontrivial Néron-Severi group is

$$\mathcal{L}_{NS} = \bigcup_{0 \neq [\Omega] \in H^2(A, \mathbb{Q})} \text{Compl}_{\Omega},$$

where A is a fixed complex torus.

Lemma 3.2. *For any alternating form Ω and any twistor sphere S , the intersection $S \cap \text{Compl}_{\Omega}$ is either finite or all of S .*

Lemma 3.3. *For any nonzero alternating form Ω , the cone C_I is not contained in Compl_{Ω} .*

Lemma 3.3 immediately implies

Lemma 3.4. *For every $I \in \text{Compl}$ the set of non-generic periods in C_I , that is $C_I \cap \mathcal{L}_{NS}$, is a countable union of closed subsets of C_I none of which contains an open neighborhood (in C_I) of any of its points.*

Proof of Lemma 3.2. Follows from the fact that $Compl_\Omega$ and S are complex analytic subsets of $Compl$.

The twistor sphere S is analytic by Corollary 1.8. The subset $Compl_\Omega$ is a complex analytic subvariety in $Compl$ as it is the locus where Ω belongs to the fiber of a holomorphic subbundle of the Hodge bundle on $Compl$ arising from the Hodge filtration. \square

Proof of Lemma 3.3. We shall prove the following equivalent statement.

For any J anti-commuting with I and any nonzero alternating form Ω on $V_\mathbb{R}$ there is a neighborhood $U_\Omega \subset C_I$ of J such that the locus $Compl_\Omega$ intersects U_Ω along a real-analytic subvariety of positive codimension.

If $J \notin Compl_\Omega$ there is nothing to prove. Assume $J \in Compl_\Omega$.

From now on we will identify our operators I, J and the form Ω with their respective matrices corresponding to a choice of basis of $V_\mathbb{R}$. Consider the orbit of J under the conjugation action of G_I : $G_I \cdot J \cong G_I/G_\mathbb{H}$. Let

$$\begin{aligned} \Psi: G_I &\longrightarrow Compl, \\ g &\longmapsto {}^gJ = gJg^{-1}, \end{aligned}$$

be the evaluation map of the action. Put $D_{I,J,\Omega} := \Psi^{-1}(Compl_\Omega)$, that is,

$$D_{I,J,\Omega} = \{g \in G_I \mid {}^t({}^gJ)\Omega({}^gJ) = \Omega\},$$

(note that $D_{I,J,\Omega}$ need not be a subgroup in G_I). Let $g(\tau)$ be any curve in $D_{I,J,\Omega}$ with tangent vector $X := g'(0) \in T_e D_{I,J,\Omega}$ at $e = g(0) \in D_{I,J,\Omega}$. Then, differentiating the constant function ${}^t({}^{g(\tau)}J)\Omega({}^{g(\tau)}J)$ at $\tau = 0$ we obtain the equality

$${}^t(XJ - JX)\Omega J + {}^tJ\Omega(XJ - JX) = 0.$$

The left hand side may be simplified, given that ${}^tJ\Omega J = \Omega$ we substitute $\Omega J = ({}^tJ)^{-1}\Omega = {}^t(J^{-1})\Omega$ into the first summand and ${}^tJ\Omega = \Omega J^{-1}$ into the second summand, obtaining

$${}^t(X^J - X)\Omega + \Omega(X^J - X) = 0,$$

where

$$X^J := J^{-1}XJ = JXJ^{-1}.$$

So, denoting $Y := X^J - X$, we obtain the equality

$$(5) \quad {}^tY\Omega + \Omega Y = 0,$$

where Y commutes with I and anticommutes with J . Note that for any $X \in T_e G_I$, $X = \frac{1}{2}(X + X^J) + \frac{1}{2}(X - X^J)$, where $X + X^J \in T_e G_I$ commutes with J and $X - X^J \in T_e G_I$ anticommutes with J . The tangent space $T_e G_\mathbb{H}$ is the subspace of elements of $T_e G_I$ that commute with J . Hence, the subspace of Y 's in $T_e G_I$ anticommuting with J maps isomorphically onto the quotient space $V_I := T_e G_I / T_e G_\mathbb{H} \cong T_J(G_I \cdot J)$ under the quotient map $T_e G_I \rightarrow V_I$. So we need to check that for a nonzero Ω the space of solutions to (5), which is naturally identified with $T_J(G_I \cdot J \cap Compl_\Omega)$, has dimension strictly less than $\dim_\mathbb{R} T_J(G_I \cdot J) = \dim_\mathbb{R} V_I = 4n^2$ (i.e., not all of the orbit $G_I \cdot J$ lies in $Compl_\Omega$).

Now conjugate equation (5) by I to obtain

$${}^tY\Omega^I + \Omega^I Y = 0.$$

Adding and subtracting this from (5) we obtain

$${}^tY(\Omega + \Omega^I) + (\Omega + \Omega^I)Y = 0 \quad \text{and} \quad {}^tY(\Omega - \Omega^I) + (\Omega - \Omega^I)Y = 0.$$

So we may assume that Ω is either I -invariant or I -anti-invariant in equation (5).

Case of I -invariant Ω . As Ω is J -invariant, it determines a skew-symmetric operator $\Omega: V_{\mathbb{R}} \rightarrow V_{\mathbb{R}}$, commuting with J . So we may choose an Ω -invariant plane $P = \langle v, Jv \rangle \subset V_{\mathbb{R}}$ corresponding to a complex eigenvector $v - iJv$ of $\Omega: V_{\mathbb{R}} \rightarrow V_{\mathbb{R}}$ such that the matrix of $\Omega|_P$ is

$$\begin{pmatrix} 0 & -\lambda \\ \lambda & 0 \end{pmatrix}.$$

The complex structure I provides another such plane $IP = \langle Iv, JIv \rangle$, which is also Ω -invariant and orthogonal to P , so that on $P \oplus IP = \langle v, Jv, Iv, JIv \rangle$ the matrices of Ω, J and I are 4×4 -block-diagonal with the following blocks on the diagonal

$$\left(\begin{array}{cc|cc} 0 & -\lambda & 0 & 0 \\ \lambda & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & \lambda \\ 0 & 0 & -\lambda & 0 \end{array} \right), \left(\begin{array}{cc|cc} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{array} \right), \left(\begin{array}{cc|cc} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ \hline 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{array} \right).$$

The condition that Y commutes with I and anticommutes with J tells us that Y has a 4×4 -block structure with blocks of the form

$$\left(\begin{array}{cc|cc} a_1 & a_2 & b_1 & b_2 \\ a_2 & -a_1 & b_2 & -b_1 \\ \hline -b_1 & b_2 & a_1 & -a_2 \\ b_2 & b_1 & -a_2 & -a_1 \end{array} \right).$$

Noting that $\Omega = JD = DJ$ for a diagonal matrix D commuting with J , we can rewrite (5) as

$$(6) \quad DY = {}^tYD, \quad Y^I = Y, \quad Y^J = -Y.$$

For notational convenience we write the matrix Y in terms of its 2×2 -blocks $Y_{k,l}$, $Y = (Y_{k,l})$, $1 \leq k, l \leq 2n$, and denote by $\mathbb{1}_{2 \times 2}$ the 2×2 identity matrix. If at least one λ_i , $1 \leq i \leq n$, in D is nonzero we get for all $1 \leq j \leq n$ the equalities of 4×4 -blocks

$$\begin{aligned} & \begin{pmatrix} \lambda_i \mathbb{1}_{2 \times 2} & 0 \\ 0 & -\lambda_i \mathbb{1}_{2 \times 2} \end{pmatrix} \cdot \begin{pmatrix} Y_{2i-1,2j-1} & Y_{2i-1,2j} \\ Y_{2i,2j-1} & Y_{2i,2j} \end{pmatrix} = \\ & = \begin{pmatrix} {}^tY_{2j-1,2i-1} & {}^tY_{2j,2i-1} \\ {}^tY_{2j-1,2i} & {}^tY_{2j,2i} \end{pmatrix} \cdot \begin{pmatrix} \lambda_j \mathbb{1}_{2 \times 2} & 0 \\ 0 & -\lambda_j \mathbb{1}_{2 \times 2} \end{pmatrix}. \end{aligned}$$

These matrix equalities completely determine all $n-1$ off-diagonal 4×4 -entries of Y in the i -th “fat” row of 4×4 -blocks in terms of the off-diagonal 4×4 -entries of the i -th “fat” column, $1 \leq i \leq n$. So the codimension of the space of solutions of (6) is at least $4(n-1)$ (precise lower bound that is reached in the least restrictive case $\lambda_j = \lambda_i$ for all j). For the diagonal 4×4 -entry, $i = j$, we obtain $b_2 = 0$ in $Y_{2i-1,2i}$, so that the codimension is at least $4n-3$.

Case of I -anti-invariant Ω : This is done similarly and leads to the same codimension bound $\geq 4n-3$. Alternatively, one could note that, if $C_I \subset \text{Compl}_{\Omega}$, then, in particular, $\pm I \in \text{Compl}_{\Omega}$, so that Ω is I -invariant, and this is the only case we need consider.

Now, by Lemma 3.2, either a twistor sphere in C_I entirely lies in some $Compl_\Omega$ or its intersection with \mathcal{L}_{NS} contains only finitely many points of each $Compl_\Omega$. If $I \notin \mathcal{L}_{NS}$ then no twistor sphere in C_I is contained in \mathcal{L}_{NS} . The codimension estimate above then allows us to conclude that, for every nonzero Ω , the subset $C_I \cap Compl_\Omega$ is of codimension at least $(4n - 3) + 2 = 4n - 1 > 0$ in C_I . If $I \in \mathcal{L}_{NS}$, the lower bound for the codimension is still at least $4n - 3 > 0$. The proof is now complete. \square

3.3. The transversality of the triple intersection of $G_{I_1}/G_{\mathbb{H}}, G_{J_1}/G_{\mathbb{H}}, G_{K_1}/G_{\mathbb{H}}$ at $eG_{\mathbb{H}}$, which is equivalent to the direct sum decomposition $T_e G/T_e G_{\mathbb{H}} = T_e G_{I_1}/T_e G_{\mathbb{H}} \oplus T_e G_{J_1}/T_e G_{\mathbb{H}} \oplus T_e G_{K_1}/T_e G_{\mathbb{H}}$, is preserved if we perturb $(I_1, J_1, K_1) \in \mathcal{T}$ a little. In other words, there is a compact neighborhood $U_{I_1, J_1, K_1} \subset \mathcal{T}$ of (I_1, J_1, K_1) and a compact neighborhood $U_{e, G} \subset G$ such that $\Phi_{I, J, K}: U_{e, J} \times U_{e, K} \rightarrow Compl$ is a submersion onto its image for all $(I, J, K) \in U_{I_1, J_1, K_1}$. Moreover, there is a compact neighborhood $U_{I_1} \subset Compl$ of I_1 which is contained in the image $\Phi_{I, J, K}(U_{e, J} \times U_{e, K})$ for all $(I, J, K) \in U_{I_1, J_1, K_1}$. We will always assume that for each neighborhood $U_{e, G}$ we made a choice of such $U_{I_1} = U_{I_1}(U_{e, G})$. Note that every $I_2 \in U_{I_1}$ is a regular value of $\Phi_{I, J, K}$ for all $(I, J, K) \in U_{I_1, J_1, K_1}$.

Lemma 3.5. *There exists a neighborhood $U_{e, G}$ such that for all $I_2 \in U_{I_1}$ and for all $(I, J, K) \in U_{I_1, J_1, K_1}$, the full preimage $\Phi_{I, J, K}^{-1}(I_2)$ is an $8n^2$ -dimensional submanifold in $U_{e, J} \times U_{e, K}$ of the form*

$$(7) \quad \{(f_1 h_1, h_1^{-1} f_2 h_2) \mid h_1, h_2 \in G_{\mathbb{H}}\} \cap (U_{e, J} \times U_{e, K}),$$

where (f_1, f_2) is a pair in $U_{e, J} \times U_{e, K}$ such that $\Phi_{I, J, K}(f_1, f_2) = I_2$.

Proof of Lemma 3.5. The fact that $\Phi_{I, J, K}^{-1}(I_2) \cap (U_{e, J} \times U_{e, K})$ consists of a finite number of $8n^2$ -dimensional manifolds follows from the regularity of I_2 .

While the part of the fiber in (7) may have been easily guessed, the fact that for a small enough $U_{e, G}$ this is the whole fiber follows from Proposition 2.5. Indeed, assuming that we have $(f_1, f_2), (g_1, g_2) \in \Phi_{I, J, K}^{-1}(I_2) \subset G_J \times G_K$, we see that $f_2^{-1} f_1^{-1} g_1 g_2 \in G_I$. Setting $g_I = f_2^{-1} f_1^{-1} g_1 g_2$ and $g_J = f_1^{-1} g_1 \in G_J$ we have the equality

$$f_2 g_I = g_J g_2.$$

The left side of the equality lies in $G_K G_I$ and the right side lies in $G_J G_K$. If we restrict ourselves to $\Phi_{I, J, K}^{-1}(I_2) \cap (U_{e, G} \times U_{e, G})$ for a small enough neighborhood $U_{e, G} \subset G$ then Proposition 2.5 tells us that, for every element in the product $U_{e, J} U_{e, K} U_{e, I}$, each of its three factors is uniquely determined up to a $G_{\mathbb{H}}$ -correction.

So from our equality $f_2 g_I = g_J g_2$ we obtain $g_I, g_J \in G_{\mathbb{H}}$, which, after setting $h_1 := g_J = f_1^{-1} g_1$ and $h_2 := g_I$, implies that $g_1 = f_1 h_1$ and $g_2 = g_J^{-1} f_2 g_I = h_1^{-1} f_2 h_2$.

Since $U_{I_1}, U_{I_1, J_1, K_1}, U_{e, G}$ are compact and $U_{e, G}$ is independent of the choice of $(I, J, K) \in U_{I_1, J_1, K_1}$, there is a universal upper bound for the number of connected components of $\Phi_{I, J, K}^{-1}(I_2) \cap (U_{e, J} \times U_{e, K})$, for all $I_2 \in U_{I_1}$ and all $(I, J, K) \in U_{I_1, J_1, K_1}$. Therefore we can shrink the compact neighborhood $U_{e, G}$ so that the fibers $\Phi_{I, J, K}^{-1}(I_2) \cap (U_{e, J} \times U_{e, K})$ for all $I_2 \in U_{I_1}$ and all $(I, J, K) \in U_{I_1, J_1, K_1}$ contain only the component specified in (7). \square

Regarding the proof of Lemma 3.5, we note the following.

Remark 3.6. It is not hard to see that the fiber $\Phi_{I,J,K}^{-1}(I_2)$ in Lemma 3.5, as a topological subspace of $G \times G$, depends continuously on $I_2 \in U_{I_1}$ and $(I, J, K) \in U_{I_1, J_1, K_1}$.


Remark 3.7. In general, it is possible that $g \in U_{e,G}$ is not uniquely representable as a triple product of elements in the larger sets G_J, G_K, G_I and thus we cannot say if the whole fiber $\Phi_{I,J,K}^{-1}(I_2) \subset G_J \times G_K$ consists of just one $G_{\mathbb{H}} \times G_{\mathbb{H}}$ -orbit as in Lemma 3.5. This is why we possibly need to shrink $U_{e,G}$.

3.4. Recall that, for any I , $M_I = G_I/G_{I,S}$ parametrizes the twistor lines through I (see Paragraph 1.5). For all I , put $U_{I_1, J_1, K_1}(I) := pr_{23}(pr_1^{-1}(I) \cap U_{I_1, J_1, K_1})$. Then $U_{I_1, J_1, K_1}(I_1)$ is a neighborhood of (J_1, K_1) in $C_{I_1} \times_{M_{I_1}} C_{I_1} = pr_{23}(pr_1^{-1}(I_1) \cap \mathcal{T})$. Consider the map

$$\begin{aligned} \Psi^{I_1 \rightarrow I_2}: \quad U_{I_1, J_1, K_1}(I_1) &\longrightarrow C_{I_2} \times_{M_{I_2}} C_{I_2} = pr_{23}(pr_1^{-1}(I_2) \cap \mathcal{T}), \\ (S(J, K), J, K) &\longmapsto (S(f_1 f_2 J, f_1 f_2 K), f_1 f_2 K, f_1 f_2 J), \end{aligned}$$

where $(f_1, f_2) \in \Phi_{I_1, J, K}^{-1}(I_2) \cap (U_{e,J} \times U_{e,K})$, and we use, in an obvious way, the triple notation of the kind $(S(J, K), J, K)$ for the elements of the fiber products above. Note the switched order of $f_1 f_2 K, f_1 f_2 J$. The role of this change of order will be clarified later.

Lemma 3.5 guarantees that the mapping $\Psi^{I_1 \rightarrow I_2}$ is well-defined, as its value at $(S(J, K), J, K)$ is uniquely determined by the fiber $\Phi_{I_1, J, K}^{-1}(I_2) \cap (U_{e,J} \times U_{e,K})$, so it does not depend on the choice of a particular point in the fiber.



Picture 2: For fixed I_1 and I_2 any pair $(J, K) \in C_{I_1} \times_{M_{I_1}} C_{I_1}$ near (J_1, K_1) determines a unique pair $(f_1 f_2 K, f_1 f_2 J) \in C_{I_2} \times_{M_{I_2}} C_{I_2}$.

3.5. Next, for each $(I, J, K) \in U_{I_1, J_1, K_1}$ consider the mapping $\Phi_{I,K,J}$ (note that we switched J and K in the subscript). By shrinking the original U_{I_1, J_1, K_1} and $U_{e,G}$ if needed, we can find a compact neighborhood $V_{e,G} \subset G$ such that for each $(I, J, K) \in U_{I_1, J_1, K_1}$ we have

- (a) $\Phi_{I,K,J}: V_{e,K} \times V_{e,J} \rightarrow Compl$ is a submersion onto its image;
 - (b) every fiber of this mapping is of the form described in Lemma 3.5;
- and

- (c) the image $\Phi_{I,K,J}(V_{e,K} \times V_{e,J})$ contains $U_{I_1} \subset \bigcap_{(I,J,K) \in U_{I_1, J_1, K_1}} \Phi_{I,J,K}(U_{e,J} \times U_{e,K})$

(see Paragraph 3.3).

By Lemma 3.5, conditions (a) and (b) are satisfied. We need only to comment on (c). By Lemma 3.5, for the original triple $(I_1, J_1, K_1) \in U_{I_1, J_1, K_1}$, we can find $V_{e, G}$ such that $\Phi_{I_1, K_1, J_1}: V_{e, K_1} \times V_{e, J_1} \rightarrow \text{Compl}$, where $V_{e, K_1} := V_{e, G} \cap G_{K_1}$, $V_{e, J_1} := V_{e, G} \cap G_{J_1}$, satisfies (a) and (b). Shrinking $U_{e, G}$ and, thus, U_{I_1} , if needed, we can satisfy (c) for Φ_{I_1, K_1, J_1} . Now shrinking U_{I_1, J_1, K_1} and again $U_{e, G}$, if needed, we can satisfy conditions (a), (b) and (c) for all $(I, J, K) \in U_{I_1, J_1, K_1}$.

3.6. Now introduce $V_{I_1, K_1, J_1} := \{(I, K, J) \mid (I, J, K) \in U_{I_1, J_1, K_1}\}$ and $V_{I_1, K_1, J_1}(I) := pr_{23}(pr_1^{-1}(I) \cap V_{I_1, K_1, J_1})$.

Then, for all (I, K, J) in the interior of V_{I_1, K_1, J_1} , the set $pr_1(V_{I_1, K_1, J_1})$ is a neighborhood of I in Compl and $V_{I_1, K_1, J_1}(I)$ is a neighborhood of $(K, J) \in C_I \times_{M_I} C_I$. Note that, due to Condition (c) in Paragraph 3.5, for all $I \in U_{I_1} \cap pr_1(V_{I_1, K_1, J_1})$ and for all $(K, J) \in V_{I_1, K_1, J_1}(I)$, the image $\Phi_{I, K, J}(U_{e, K} \times U_{e, J})$ contains the neighborhood U_{I_1} .

3.7. Choose $I_2 \in U_{I_1} \cap pr_1(V_{I_1, K_1, J_1})$ and K, J such that $(I_2, K, J) \in V_{I_1, K_1, J_1}$. Conditions (a), (b) and (c) in Paragraph 3.5 allow us to define, analogously to $\Psi^{I_1 \rightarrow I_2}$, the map

$$\begin{aligned} \Psi^{I_2 \rightarrow I_1}: \quad V_{I_1, K_1, J_1}(I_2) &\longrightarrow C_{I_1} \times_{M_{I_1}} C_{I_1}, \\ (S(J, K), K, J) &\longmapsto (S(d_1 d_2 J, d_1 d_2 K), d_1 d_2 J, d_1 d_2 K), \end{aligned}$$

for $(d_1, d_2) \in \Phi_{I_2, K, J}^{-1}(I_1)$ (again, note the reversed order of J and K in the subscript).

The period $f_1 f_2 K$ in Picture 2 above will play the role of the “rotation center” for $\Phi_{I_2, f_1 f_2 K, f_1 f_2 J}$ (here $J, K \in C_{I_1}$), similar to the role that J plays for $\Phi_{I_1, J, K}$. This explains why we switched J and K .

Below we will impose restrictions on the domain of $\Psi^{I_1 \rightarrow I_2}$ in order for the image of this map to be contained in the domain of $\Psi^{I_2 \rightarrow I_1}$, so that we can compose them.

We begin by choosing a compact neighborhood U_{J_1, K_1} of (J_1, K_1) in $U_{I_1, J_1, K_1}(I_1)$, which can at first be all of $U_{I_1, J_1, K_1}(I_1)$. We will later modify U_{J_1, K_1} , without changing the original U_{I_1, J_1, K_1} .

Lemma 3.8. *For fixed V_{I_1, K_1, J_1} , we can shrink $U_{e, G}$ and U_{J_1, K_1} so that for arbitrary $I_2 \in U_{I_1}$,*

$$\Psi^{I_1 \rightarrow I_2}(U_{J_1, K_1}) \subset V_{I_1, K_1, J_1}(I_2).$$

Proof. As in Paragraph 3.5, this follows from the fact that the mapping $\Psi^{I_2 \rightarrow I_1}$ depends continuously on I_2 (see Remark 3.6), and that

$$\lim_{I_2 \rightarrow I_1} \Psi^{I_2 \rightarrow I_1} = (12): U_{J_1, K_1} \rightarrow V_{I_1, K_1, J_1}(I_1),$$

$$(S(J, K), J, K) \mapsto (S(J, K), K, J),$$

the latter mapping is trivially defined on the whole U_{J_1, K_1} , so that the sizes of the domains $V_{I_1, K_1, J_1}(I_2)$ of $\Psi^{I_2 \rightarrow I_1}$'s are bounded away from zero, when I_2 is close to I_1 .

As before, we can further shrink $U_{e, G}$ (and hence U_{I_1}), if needed, so that properties (a), (b), (c) in Paragraph 3.5 hold independently of the point $I_2 \in U_{I_1}$. \square

3.8. Possibly shrinking $U_{e,G}$, we can and will assume that it is invariant under taking inverses, $g \mapsto g^{-1}$.

Lemma 3.9. *Possibly further shrinking $U_{e,G}$ and U_{J_1,K_1} , satisfying the conclusion of Lemma 3.8, we have for all $I_2 \in U_{I_1}$*

$$\Psi^{I_2 \rightarrow I_1} \circ \Psi^{I_1 \rightarrow I_2} = Id|_{U_{J_1,K_1}}.$$

Proof. For all $(f_1, f_2) \in \Phi_{I_1,J,K}^{-1}(I_2) \cap (U_{e,G} \times U_{e,G})$ and all $(S(J, K), J, K) \in U_{J_1,K_1}$, we want the neighborhoods $V_{e,f_1 f_2 K} = V_{e,G} \cap G_{f_1 f_2 K}$, $V_{e,f_1 f_2 J} = V_{e,G} \cap G_{f_1 f_2 J}$ to contain, respectively, the neighborhoods ${}^{f_1 f_2}U_{e,K} = f_1 f_2 U_{e,K} f_2^{-1} f_1^{-1}$ and ${}^{f_1 f_2}U_{e,J} = f_1 f_2 U_{e,J} f_2^{-1} f_1^{-1}$, so that, in particular, $V_{e,f_1 f_2 K} \times V_{e,f_1 f_2 J}$ contains the pair

$$(d_1, d_2) = (f_1 f_2 \cdot f_2^{-1} \cdot f_2^{-1} f_1^{-1}, f_1 f_2 \cdot f_1^{-1} \cdot f_2^{-1} f_1^{-1}) = (f_1 f_2^{-1} f_1^{-1}, f_1 f_2 \cdot f_1^{-1} \cdot f_2^{-1} f_1^{-1}).$$

Here the invariance of $U_{e,G}$ under taking inverses is used. The pair (d_1, d_2) certainly belongs to the preimage $\Phi_{I_2, f_1 f_2 K, f_1 f_2 J}^{-1}(I_1)$ as the product of its entries is $f_1 f_2^{-1} f_1^{-1} \cdot f_1 f_2 \cdot f_1^{-1} \cdot f_2^{-1} f_1^{-1} = f_2^{-1} f_1^{-1}$.

Note that, for $U_{e,G}$ small enough, the neighborhoods ${}^{f_1 f_2}U_{e,K} \times {}^{f_1 f_2}U_{e,J}$ will also be uniformly small for all $(f_1, f_2) \in \Phi_{I_1,J,K}^{-1}(I_2) \cap (U_{e,G} \times U_{e,G})$, so that the fiber of $\Phi_{I_2, f_1 f_2 K, f_1 f_2 J}^{-1}(I_1)$ in ${}^{f_1 f_2}U_{e,K} \times {}^{f_1 f_2}U_{e,J}$ consists of a unique connected component of the form described in Lemma 3.5. Then the pair (d_1, d_2) is contained in this “good” part of the fiber $\Phi_{I_2, f_1 f_2 K, f_1 f_2 J}^{-1}(I_1)$ and we can use (d_1, d_2) to evaluate $\Psi^{I_2 \rightarrow I_1}$ at $(S({}^{f_1 f_2}J, {}^{f_1 f_2}K), {}^{f_1 f_2}J, {}^{f_1 f_2}K)$. Thus

$$\Phi_{I_2, f_1 f_2 K, f_1 f_2 J}(d_1, d_2) = I_1$$

and

$$d_1 d_2 f_1 f_2 J = J, d_1 d_2 f_1 f_2 K = K,$$

so that

$$\Psi^{I_2 \rightarrow I_1}(S({}^{f_1 f_2}J, {}^{f_1 f_2}K), {}^{f_1 f_2}K, {}^{f_1 f_2}J) = (S(J, K), J, K),$$

where, certainly, $S(J, K) = S(I_1, J, K)$, proving that the composition $\Psi^{I_2 \rightarrow I_1} \circ \Psi^{I_1 \rightarrow I_2}$ is the identity on U_{J_1,K_1} .

In order to ensure that $V_{e,f_1 f_2 K} \times V_{e,f_1 f_2 J}$ contains (d_1, d_2) , we assume, shrinking $U_{e,G}$ and U_{J_1,K_1} if necessary, but not changing $V_{e,G}$ and the previously fixed V_{I_1,K_1,J_1} , that for all $(S(J, K), J, K) \in U_{J_1,K_1}$ and for all points $(f_1, f_2) \in \Phi_{I_1,J,K}^{-1}(I_2) \cap (U_{e,G} \times U_{e,G})$, the neighborhoods $V_{e,f_1 f_2 K}$, $V_{e,f_1 f_2 J}$ contain, respectively, the neighborhoods ${}^{f_1 f_2}U_{e,K}$ and ${}^{f_1 f_2}U_{e,J}$. \square

Corollary 3.10. *Let U_{I_1} be defined by $U_{e,G}$ ($U_{e,G}$ satisfying Lemma 3.9). For arbitrary $I_2 \in U_{I_1}$, both joint points $J \in C_{I_1}$ and ${}^{f_1 f_2}K \in C_{I_2}$ of a triple of twistor spheres connecting I_1 and I_2 , can be chosen generic.*

Proof. Define

$$\begin{aligned} pr_K: \quad V_{I_1,K_1,J_1}(I_2) &\longrightarrow C_{I_2} \subset Compl, \\ (S(J, K), K, J) &\longmapsto K. \end{aligned}$$

This projection is a submersion onto its image. By Lemma 3.4, the locus \mathcal{L}_{NS} intersects C_{I_2} in a countable union of closed submanifolds of positive codimension in C_{I_2} . As the mapping pr_K is a submersion onto its image, the preimage $pr_K^{-1}(\mathcal{L}_{NS} \cap C_{I_2})$ is

also a countable union of closed submanifolds of positive codimension in $V_{I_1, K_1, J_1}(I_2)$. Similarly, for

$$\begin{aligned} pr_J: \quad U_{J_1, K_1} &\longrightarrow C_{I_1} \subset Compl, \\ (S(J, K), J, K) &\longmapsto J, \end{aligned}$$

$pr_J^{-1}(\mathcal{L}_{NS} \cap C_{I_1}) \subset U_{J_1, K_1}$ is a countable union of closed submanifolds of positive codimension. The mapping $\Psi^{I_2 \rightarrow I_1}$ is real-analytic, so the closure of $\Psi^{I_2 \rightarrow I_1}(pr_J^{-1}(\mathcal{L}_{NS} \cap C_{I_2}))$ in U_{J_1, K_1} does not contain interior points. Therefore

$$(8) \quad U_{J_1, K_1} \neq pr_J^{-1}(\mathcal{L}_{NS} \cap C_{I_1}) \cup \Psi^{I_2 \rightarrow I_1}(pr_K^{-1}(\mathcal{L}_{NS} \cap C_{I_2})).$$

Since, by Lemma 3.9, $\Psi^{I_2 \rightarrow I_1} \circ \Psi^{I_1 \rightarrow I_2} = Id|_{U_{J_1, K_1}}$, the inequality (8) tells us that the image of the mapping $\Psi^{I_1 \rightarrow I_2}$ is not contained in $pr_K^{-1}(\mathcal{L}_{NS} \cap C_{I_2})$. Thus we may find a pair $(J, K) \in U_{J_1, K_1}$ such that $J = pr_J(S(J, K), J, K) \notin \mathcal{L}_{NS} \cap C_{I_1}$ and $f_1 f_2 K = pr_K(\Psi^{I_1 \rightarrow I_2}(S(J, K), J, K)) \notin \mathcal{L}_{NS} \cap C_{I_2}$, that is, both periods are generic. \square

4. THE DEGREE OF TWISTOR LINES

In this section we show that twistor lines in $Compl$ have degree $2n$ in the Plücker embedding. We first show that the group $G = GL(V_{\mathbb{R}})$ acts transitively on the set of twistor lines in $Compl$ and then compute the degree of an explicit twistor line.

Lemma 4.1. *The group $G = GL(V_{\mathbb{R}})$ acts transitively on the set of twistor lines in $Compl$.*

Proof. Given two twistor spheres $S_1 = S(I_1, J_1)$ and $S_2 = S(I_2, J_2)$, there is an element $g \in G$ sending I_1 to I_2 , hence sending S_1 to a twistor sphere through I_2 . The lemma now follows from Corollary 1.6. \square

4.1. To construct our example, consider the affine chart in the Grassmannian $Gr(2n, 4n)$ of normalized period matrices $(\mathbb{1}|Z)$, where $\mathbb{1}$ is, in general, the $2n \times 2n$ identity matrix and Z now denotes a non-degenerate $2n \times 2n$ complex matrix. Let us fix a basis of $V_{\mathbb{R}}$ and write the matrix of an arbitrary complex structure $I: V_{\mathbb{R}} \rightarrow V_{\mathbb{R}}$ in the following block form

$$I = \begin{pmatrix} A & B \\ C & D \end{pmatrix},$$

for $2n \times 2n$ real matrices A, B, C, D . Then the relation

$$(\mathbb{1} | Z)I = (i\mathbb{1} | iZ),$$

gives the matrix equations

$$A + ZC = i\mathbb{1}, \quad B + ZD = iZ.$$

Assume that C is invertible so that the first equation allows us to write $Z = (i\mathbb{1} - A)C^{-1}$. The condition that I is a complex structure will then guarantee that the second equation is automatically satisfied.

4.2. **The case $n = 1$.** Momentarily assume $n = 1$ and consider the twistor sphere $S = S(I, J)$ where I and J have the respective matrices

$$\left(\begin{array}{cc|cc} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{array} \right), \left(\begin{array}{cc|cc} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ \hline 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{array} \right)$$

and put $K = IJ$. So for $\lambda \in S$,

$$\lambda = aI + bJ + cK = \left(\begin{array}{cc|cc} 0 & -a & -b & -c \\ a & 0 & -c & b \\ \hline b & c & 0 & -a \\ c & -b & a & 0 \end{array} \right).$$

Assume additionally that $b^2 + c^2 \neq 0$, that is, $\lambda \in S \setminus \{\pm I\}$. Here

$$A = \begin{pmatrix} 0 & -a \\ a & 0 \end{pmatrix}, C = \begin{pmatrix} b & c \\ c & -b \end{pmatrix}, C^{-1} = \frac{1}{b^2 + c^2} \begin{pmatrix} b & c \\ c & -b \end{pmatrix}.$$

Then

$$Z = (i\mathbb{1} - A)C^{-1} = \frac{1}{b^2 + c^2} \begin{pmatrix} ac + ib & -ab + ic \\ -ab + ic & -ac - ib \end{pmatrix} = \begin{pmatrix} z_1 & z_2 \\ z_3 & z_4 \end{pmatrix},$$

where Z clearly satisfies the equations

$$z_1 + z_4 = 0, \quad z_2 - z_3 = 0, \quad \det Z = z_1 z_4 - z_2 z_3 = 1.$$

4.3. Now, for a general n , we can construct a twistor line in the period domain of complex $2n$ -dimensional tori, which, in the affine chart of $Gr(2n, 4n)$ above, corresponds to the locus of matrices $(\mathbb{1}|Z)$ where Z is the block-diagonal matrix with the same 2×4 -block

$$\left(\begin{array}{cc|cc} 1 & 0 & u & v \\ 0 & 1 & v & -u \end{array} \right), u^2 + v^2 = -1,$$

on the diagonal.

4.4. The degree of the curve in the example is $2n$ under the Plücker embedding $Gr(2n, 4n) \hookrightarrow \mathbb{P}^{\binom{4n}{2n}-1}$. Indeed, the Plücker coordinates in the above affine chart are given by the maximal minors of the matrix $(\mathbb{1}|Z)$. The twistor line S in our example is contained in the plane $P(S)$ with parameters u, v in the part given by the affine chart. Let W be the homogeneous coordinate given by the minor formed by all $\mathbb{1}_{2 \times 2}$ -blocks and let U and V be any homogeneous coordinates such that after restricting to $P(S)$ we get $u = \frac{U}{W}$ and $v = \frac{V}{W}$.

Let us consider from now on the plane $P(S)$ as a projective (complete) 2-plane in $Gr(2n, 4n)$ with coordinates U, V, W . Then the minor formed by the (u, v) -blocks restricts to $P(S)$ as $(U^2 + V^2)^n$. Rewriting the equation $u^2 + v^2 = -1$ of our twistor line S in homogeneous coordinates we get $U^2 + V^2 + W^2 = 0$, so that, restricting the polynomial $(U^2 + V^2)^n$ to S , we see that it vanishes precisely when $U^2 + V^2 = -W^2$ vanishes, that is, only at the points $\pm I$ of S outside of our affine chart. Each of the two factors in the expansion $(U^2 + V^2)^n = (U + iV)^n (U - iV)^n$ vanishes to order n at the respective point, so the total order of vanishing is $n + n = 2n$ which is the degree of the image of S under the Plücker embedding.

Corollary 4.2. *Twistor lines have degree $2n$ in the Plücker embedding of $Compl$.*

Proof. Follows from Lemma 4.1 and Paragraph 4.4. \square

Remark 4.3. There is an alternative proof of the above corollary following the lines explained in Remark 1.9. Namely, we have the embedding $i: Gr(2, \mathbb{H} \otimes \mathbb{C}) \hookrightarrow Gr(2n, V_{\mathbb{C}})$, $\mathbb{H}^{1,0} \mapsto \mathbb{H}^{1,0} \otimes V'_{\mathbb{C}}$. Let e_1, \dots, e_n be some basis of $V'_{\mathbb{C}}$. Then, in terms of the Plücker embeddings of the respective Grassmanians, we have $i(u \wedge v) = u \otimes e_1 \wedge v \otimes e_1 \wedge \dots \wedge u \otimes e_n \wedge v \otimes e_n$, which induces an isomorphism $i^*(\mathcal{O}_{Gr(2n, V_{\mathbb{C}})}(1)) \cong \mathcal{O}_{Gr(2, \mathbb{H} \otimes \mathbb{C})}(n)$ of the sheaves on the quadric $Gr(2, \mathbb{H} \otimes \mathbb{C})$, thus justifying that the degree of $S \subset Gr(2n, 4n)$ under the Plücker embedding is $2n$.

5. TWISTOR PATH CONNECTIVITY OF $Compl_{\Omega}$

In this section we describe the discrete invariants and the homogeneous structure of the connected components of $Compl_{\Omega}$ and establish a criterion of twistor path connectivity of a “special” connected component of $Compl_{\Omega}$ (the remaining “non-special” components will be shown to contain no twistor lines at all).

Let I be a complex structure operator in $Compl_{\Omega}$, that is, $\Omega(Iu, Iv) = \Omega(u, v)$ for all $u, v \in V_{\mathbb{R}}$. On the vector space $(V_{\mathbb{R}}, I)$, considered as a complex vector space, the form Ω determines a hermitian form

$$h(u, v) := \Omega(u, Iv) - i\Omega(u, v)$$

(note that $h(u, Iv) = -ih(u, v)$, $h(Iu, v) = ih(u, v)$), which we will call *the hermitian form associated to Ω and I* .

The signature of h is a triple (n_+, n_-, n_0) , where n_+, n_- and n_0 are the complex dimensions of, respectively, a maximal positive subspace V_+ , a maximal negative subspace V_- , and the null subspace

$$V_0 = \{u \in V_{\mathbb{R}} \mid h(u, v) = 0 \text{ for all } v \in V_{\mathbb{R}}\} = \{u \in V_{\mathbb{R}} \mid \Omega(u, v) = 0 \text{ for all } v \in V_{\mathbb{R}}\}$$

of h in $(V_{\mathbb{R}}, I)$, so that $n_+ + n_- + n_0 = 2n$. The subspaces V_+, V_- and the numbers n_+ and n_- depend, in general, on the choice of $I \in Compl_{\Omega}$, while V_0 , and hence n_0 , depend only on Ω , V_0 being invariant with respect to every complex structure operator in $Compl_{\Omega}$.

The group $G = GL(V_{\mathbb{R}})$ acts naturally on 2-forms on $V_{\mathbb{R}} \times V_{\mathbb{R}}$,

$$g: f(u, v) \mapsto g^*f(u, v) = f(gu, gv)$$

or, in matrix notation, $g: f \mapsto {}^t g \cdot f \cdot g$.

Consider the subgroup $G_{\Omega} \subset G$ of automorphisms of Ω :

$$G_{\Omega} := \{g \in G \mid g^*\Omega = \Omega\} \subset G.$$

The subgroup G_{Ω} acts via the G -action on $Compl_{\Omega}$. For $I \in Compl_{\Omega}$ we denote by $G_{I, \Omega}$ the stabilizer of I in G_{Ω} :

$$G_{I, \Omega} = \{g \in G_{\Omega} \mid {}^g I = I\}.$$

We note that, by definition, for the hermitian form h associated to Ω and I , the subgroup $G_{I, \Omega}$ is the stabilizer of h in G_I under the corresponding G -action, i.e., $G_{I, \Omega} = \{g \in G_I \mid g^*h = h\}$.

Note that if Ω is non-degenerate, then G_{Ω} is connected, and if Ω is degenerate, that is, $n_0 > 0$, then the group G_{Ω} consists of two connected components, one in each

connected component of G . We denote by G_Ω^0 the connected subgroup component of G_Ω .

We set $\text{Compl}_\Omega^\pm := \text{Compl}_\Omega \cap \text{Compl}^\pm$. Consider the finite set

$$S_\Omega := \{(k, l, n_0) \mid k, l \in \mathbb{Z}_{\geq 0}, k + l + n_0 = 2n\}$$

consisting of $2n - n_0 + 1$ triples, and the mapping

$$\begin{aligned} \text{Sign}: \text{Compl}_\Omega^\pm &\longrightarrow S_\Omega, \\ I &\longmapsto \text{the signature } (n_+, n_-, n_0) \text{ of } h \text{ assoc. to } \Omega \text{ and } I. \end{aligned}$$

Proposition 5.1. *The mapping $\text{Sign}: \text{Compl}_\Omega^\pm \rightarrow S_\Omega$ is constant on every connected component of Compl_Ω .*

Proof. Since n_0 is the dimension of the kernel of Ω , and Ω is fixed, n_0 is constant. Now the fact that n_+ and n_- are locally constant follows from the fact that they are both lower semi-continuous (since being positive or negative are open conditions) and their sum is constant. \square

Proposition 5.1 allows us to define the induced mapping $\widetilde{\text{Sign}}: \pi_0 \text{Compl}_\Omega^\pm \rightarrow S_\Omega$ on the set $\pi_0 \text{Compl}_\Omega^\pm$ of connected components of Compl_Ω^\pm .

We now formulate the following useful lemmas whose proofs will be given later in this section.

Lemma 5.2. *The group G_Ω^0 acts transitively on each fiber of Sign . In particular, $\widetilde{\text{Sign}}$ is injective.*

Lemma 5.2 implies that the fibers of Sign are connected components of Compl_Ω^\pm , each of which is a G_Ω^0 -orbit and is thus diffeomorphic to the quotient $G_\Omega^0/G_{I,\Omega}$ for I in this orbit. Thus, all G_Ω^0 -stabilizers of $I \in \text{Sign}^{-1}(n_+, n_-, n_0)$ are conjugate in G_Ω^0 , which is the same as to say that all hermitian forms h associated to Ω and $I \in \text{Compl}_\Omega^\pm$ of the same signature (n_+, n_-, n_0) are G_Ω^0 -equivalent. We let $U(n_+, n_-, n_0)$ denote the stabilizer $G_{I,\Omega}$, well-defined up to conjugacy.

Lemma 5.3. *The mapping $\widetilde{\text{Sign}}: \pi_0 \text{Compl}_\Omega^\pm \rightarrow S_\Omega$ is surjective.*

Finally, we formulate the main result of this section.

Theorem 5.4. *The mapping $\widetilde{\text{Sign}}: \pi_0 \text{Compl}_\Omega^\pm \rightarrow S_\Omega$ is a bijection, thus there are $2n + 1 - n_0$ connected components of Compl_Ω^\pm and they have the form $\text{Sign}^{-1}(n_+, n_-, n_0)$. Every connected component of Compl_Ω^\pm is a homogeneous manifold of the type $G_\Omega^0/U(n_+, n_-, n_0)$ and is a smooth complex submanifold in Compl .*

If n_0 is even, there is precisely one connected component of Compl_Ω^\pm that contains twistor lines. This component is $\text{Sign}^{-1}(n - \frac{n_0}{2}, n - \frac{n_0}{2}, n_0)$ and is twistor path connected. If n_0 is odd, there are no connected components of Compl_Ω^\pm containing a twistor line.

Let $A = V_\mathbb{R}/\Gamma$ be a torus with a complex structure $I: V_\mathbb{R} \rightarrow V_\mathbb{R}$.

Corollary 5.5. *If the form Ω represents a Kähler class in $H^{1,1}(A, \mathbb{R})$ then there are no twistor lines passing through I in Compl_Ω .*

Informally speaking, Kähler classes do not “survive” along twistor lines, this is well known, for example, for twistor lines in the moduli space of $K3$ surfaces. Corollary 5.5 is almost clear, since a form Ω represents a Kähler class in $H^{1,1}(A, \mathbb{R})$ if and only if the form h associated to Ω and I is positive definite (see [7, p. 303, *Riemann conditions 1*] for a coordinate based exposition of this), that is, its signature is of the form $(2n, 0, 0)$ and hence, by Theorem 5.4, the connected component of $Compl_\Omega$ containing I does not contain any twistor lines.

Proof of Lemma 5.2. Let I_1, I_2 be periods in $Compl_\Omega^\pm$. Assume that $Sign(I_1) = Sign(I_2)$, that is, the associated hermitian forms h_1 on $(V_\mathbb{R}, I_1)$ and h_2 on $(V_\mathbb{R}, I_2)$ have the same signature (k, l, m) .

First we fix orthogonal decompositions into maximal positive, negative and null subspaces $V_\mathbb{R} = V_+ \oplus V_- \oplus V_0$ of $(V_\mathbb{R}, I_1)$ with respect to h_1 and $V_\mathbb{R} = W_+ \oplus W_- \oplus V_0$ of $(V_\mathbb{R}, I_2)$ with respect to h_2 , here $\dim_\mathbb{C} V_+ = k = \dim_\mathbb{C} W_+$, $\dim_\mathbb{C} V_- = l = \dim_\mathbb{C} W_-$ and V_0 is the null subspace for Ω , $\dim_\mathbb{C} V_0 = m$. Let us choose an h_1 -orthonormal basis v_1, \dots, v_k of the complex subspace (V_+, I_1) and an h_2 -orthonormal basis w_1, \dots, w_k of the complex subspace (W_+, I_2) , similarly choose $-h_1$ -orthonormal and $-h_2$ -orthonormal bases for the subspaces V_-, W_- , and some (arbitrary) bases for (V_0, I_1) and (V_0, I_2) . Then we define $g \in G = GL(V_\mathbb{R})$ by setting $g(v_1) = w_1, g(I_1 v_1) = I_2 w_1, g(v_2) = w_2, g(I_1 v_2) = I_2 w_2, \dots, g(v_k) = w_k, g(I_1 v_k) = I_2 w_k$, and similarly for the remaining pairs of subspaces. Then we get $g \in G$ such that $g^* h_2 = h_1$ and $I_2 = g I_1 g^{-1}$, which is equivalent to saying that $g^* \Omega = \Omega$ and $I_2 = g I_1 g^{-1} = {}^g I_1$. This implies that $g \in G_\Omega$ and the assumption that both I_1 and $I_2 = {}^g I_1$ belong to the same connected component of $Compl$ tells us that g actually belongs to the connected subgroup component of G and hence to G_Ω^0 . So I_2 belongs to the orbit of I_1 under the action of G_Ω^0 , and they belong to the same connected component of $Compl_\Omega^\pm$. \square

Proof of Lemma 5.3. Let (n_+, n_-, n_0) be a triple $\in S_\Omega$, $n_+ + n_- + n_0 = 2n$. Let us prove that there exists $I \in Compl_\Omega^\pm$ such that the hermitian form h associated to Ω and I has signature (n_+, n_-, n_0) . Choose a basis v_1, \dots, v_{4n} of $V_\mathbb{R}$ such that the skew-symmetric matrix $\Omega(v_i, v_j)$ has 2×2 -block diagonal structure and set $L_\Omega: V_\mathbb{R} \rightarrow V_\mathbb{R}$ to be the operator defined by this matrix in the chosen basis. By the definition of n_0 there are $n_+ + n_- = 2n - n_0$ nonzero 2×2 -blocks and n_0 zero 2×2 -blocks in the matrix of L_Ω . Let $V_\mathbb{R} = P_1 \oplus \dots \oplus P_{n_+} \oplus Q_1 \oplus \dots \oplus Q_{n_-} \oplus U$ be a direct sum decomposition into L_Ω -invariant real 2-planes P_j, Q_j and $2n_0$ -dimensional real subspace $U = \text{Ker } L_\Omega$, where $L_\Omega|_{P_i}, L_\Omega|_{Q_j}$ are nonzero and $L_\Omega|_U = 0$. Define $I: P_i \rightarrow P_i, 1 \leq i \leq n_+$, to be the appropriate negative multiple of $L_\Omega|_{P_i}$, $I: Q_j \rightarrow Q_j, 1 \leq j \leq n_-$, to be the appropriate positive multiple of $L_\Omega|_{Q_j}$, and $I: U \rightarrow U$ to be an arbitrary operator so as to form an operator $I: V_\mathbb{R} \rightarrow V_\mathbb{R}$ satisfying $I^2 = -Id$. Then the decomposition $V_\mathbb{R} = P_1 \oplus \dots \oplus P_{n_+} \oplus Q_1 \oplus \dots \oplus Q_{n_-} \oplus U$ is orthogonal with respect to $h(u, v) = \Omega(u, Iv) - i\Omega(u, v)$, $h|_{P_i} > 0$, $h|_{Q_j} < 0$ and $h|_U = 0$. Thus we have constructed the required I , this proves the surjectivity of \widetilde{Sign} . \square

Proof of Theorem 5.4. The part of the statement about the bijectivity of \widetilde{Sign} and the G_Ω^0 -orbit structure of the connected components of $Compl_\Omega^\pm$ follows from Lemma 5.2 and Lemma 5.3.

The complex manifold structure. In order to show that every connected component of $Compl_\Omega^\pm$ is a complex submanifold in $Compl$ we need to identify the tangent space $T_I Compl_\Omega$ as a complex subspace in $T_I Compl$ for every $I \in Compl_\Omega$. We recall that the complex structure operator $l_I: T_I Compl \rightarrow T_I Compl$ is given by left multiplication by I . By definition $G_\Omega = \{g \in G \mid {}^t g \Omega g = \Omega\} \subset G$, so

$$T_e G_\Omega \cong \{Y \in T_e G \mid {}^t Y \Omega + \Omega Y = 0\}.$$

The stabilizer of I in G_Ω is $G_{I,G} = G_I \cap G_\Omega$, thus $T_I Compl_\Omega \cong T_e G_\Omega / T_e G_{I,\Omega}$.

For $X \in T_e G_\Omega$, consider the decomposition $X = \frac{1}{2}(X + X^I) + \frac{1}{2}(X - X^I)$ into I -commuting and I -anticommuting components. Then $\frac{1}{2}(X + X^I)$ and $\frac{1}{2}(X - X^I)$ satisfy the equality ${}^t Y \Omega + \Omega Y = 0$, so that $\frac{1}{2}(X + X^I) \in T_e G_{I,\Omega}$ and thus the isomorphism $T_I Compl_\Omega \cong T_e G_\Omega / T_e G_{I,\Omega}$ implies the natural isomorphism

$$(9) \quad T_I Compl_\Omega \cong \{Y \in T_e G \mid YI = -IY, {}^t Y \Omega + \Omega Y = 0\}.$$

Thus $T_I Compl_\Omega$ is obviously l_I -invariant, so it is a complex subspace in $T_I Compl$ for every $I \in Compl_\Omega$, which shows that every connected component of $Compl_\Omega$, or, which is the same, of $Compl_\Omega^\pm$, is a complex submanifold in $Compl$.

The component containing a twistor line. We first note that $Compl_\Omega$ contains a twistor line $S = S(I, J)$ if and only if Ω is both I - and J -invariant, that is, $I, J \in Compl_\Omega$.

Let us assume that Ω is invariant with respect to anticommuting complex structure operators I and J , that is $\Omega(Iu, Iv) = \Omega(Ju, Jv) = \Omega(u, v)$ for all $u, v \in V_\mathbb{R}$. Then for all $u, v \in V_\mathbb{R}$ the form h associated to Ω and I satisfies

$$\begin{aligned} h(Ju, Jv) &\stackrel{def}{=} \Omega(Ju, IJv) - i\Omega(Ju, Jv) = -\Omega(Ju, JIv) - i\Omega(Ju, Jv) = \\ &= -\Omega(u, Iv) - i\Omega(u, v) = -\overline{h(u, v)}. \end{aligned}$$

This implies that the hermitian forms $h(u, v)$ and $-\overline{h(u, v)}$ are equivalent under J , therefore they must have the same signature, so that the signature (n_+, n_-, n_0) of $h(u, v)$ and the signature (n_-, n_+, n_0) of $-\overline{h(u, v)}$ are equal, that is, $n_+ = n_-$. It also follows that n_0 is even.

Conversely, if, for the hermitian form h on $(V_\mathbb{R}, I) \times (V_\mathbb{R}, I)$ associated to Ω and I , we have the equality $n_+ = n_-$, we construct a complex structure operator $J: V_\mathbb{R} \rightarrow V_\mathbb{R}$ anticommuting with I and leaving Ω invariant as follows. Fix an h -orthogonal decomposition $V_\mathbb{R} = V_+ \oplus V_- \oplus V_0$, where V_+ , V_- , V_0 are maximal positive, negative and null subspaces of h , which are complex subspaces of $(V_\mathbb{R}, I)$, and put $k := \dim_\mathbb{C} V_+ = n_+ = n_- = \dim_\mathbb{C} V_-$, $l := n_0 = \dim_\mathbb{C} V_0$. Note that since $n_+ + n_- + n_0 = 2k + l = 2n$, $l = n_0$ is an even number.

Let u_1, \dots, u_k and v_1, \dots, v_k be h -orthonormal and, respectively, $-h$ -orthonormal bases of the complex subspaces (V_+, I) and (V_-, I) . Let w_1, \dots, w_l be any basis of the complex subspace (V_0, I) . Define $J: V_\mathbb{R} \rightarrow V_\mathbb{R}$ by setting

$$\begin{aligned} J(u_1) &= v_1, J(Iu_1) = -Iv_1, \dots, J(u_k) = v_k, J(Iu_k) = -Iv_k, \\ J(v_1) &= -u_1, J(Iv_1) = Iu_1, \dots, J(v_k) = -u_k, J(Iv_k) = Iu_k \end{aligned}$$

and, using that l is an even number, by setting

$$J(w_1) = w_2, J(Iw_1) = -Iw_2, J(w_2) = -w_1, J(Iw_2) = Iw_1, \dots$$

$$J(w_{l-1}) = w_l, J(Iw_{l-1}) = -Iw_l, J(w_l) = -w_{l-1}, J(Iw_l) = Iw_{l-1}.$$

Then J anticommutes with I and one easily verifies that J takes h to $-\bar{h}$, which, by the definition of h , means that

$$\Omega(Ju, IJv) - i\Omega(Ju, Jv) = h(Ju, Jv) = -\overline{h(u, v)} = -\Omega(u, Iv) - i\Omega(u, v),$$

so that $\Omega(Ju, Jv) = \Omega(u, v)$, that is, Ω is J -invariant.

Twistor-path connectivity of $Sign^{-1}(n - \frac{n_0}{2}, n - \frac{n_0}{2}, n_0)$. We follow the general lines in Section 2.1. Let $S = S(I, J)$ be a twistor line in $Sign^{-1}(n - \frac{n_0}{2}, n - \frac{n_0}{2}, n_0)$.

Consider the subgroups $G_{J,\Omega} = G_J \cap G_\Omega$, $G_{K,\Omega} = G_K \cap G_\Omega$, $G_{\mathbb{H},\Omega} = G_{J,\Omega} \cap G_I = G_{K,\Omega} \cap G_I$ and define the mapping

$$\begin{aligned} \Phi: G_{J,\Omega} \times G_{K,\Omega} &\rightarrow Compl_\Omega^\pm, \\ (g_1, g_2) &\mapsto {}^{g_1 g_2} I \in Compl_\Omega^\pm. \end{aligned}$$

In order to proceed as in Section 2.1 we need to show that $d\Phi_{(e,e)}$ is surjective. Again, as in the proof of Proposition 2.1, one can show that $Ker d\Phi_{(e,e)} = T_e G_{\mathbb{H},\Omega} \times T_e G_{\mathbb{H},\Omega}$, so that

$$d\Phi_{(e,e)}(T_e G_\Omega) \cong T_e G_{J,\Omega}/T_e G_{\mathbb{H},\Omega} \times T_e G_{K,\Omega}/T_e G_{\mathbb{H},\Omega} \subset T_I Compl_\Omega.$$

To show the surjectivity of $d\Phi_{(e,e)}$ we need another inclusion

$$T_I Compl_\Omega \subset T_e G_{J,\Omega}/T_e G_{\mathbb{H},\Omega} \times T_e G_{K,\Omega}/T_e G_{\mathbb{H},\Omega}.$$

For this we use the identification in Equation (9) and first note that, for $Z \in T_I Compl_\Omega$ (Z anticommutes with I) the vector $Z + Z^J$ commutes with J and anticommutes with I , hence anticommutes with K as well. Next, by J -invariance of Ω we have that $Z + Z^J$ satisfies the relation with Ω in (9), hence itself belongs to

$$\{Y \in T_I Compl_\Omega \mid YJ = JY\} = T_e G_{J,\Omega}/T_e G_{\mathbb{H},\Omega}.$$

Now $Z - \frac{1}{2}(Z + Z^J) = \frac{1}{2}(Z - Z^J)$ commutes with K , so that, due to the fact that $Z + Z^J$ anticommutes with K , we have that $\frac{1}{2}(Z - Z^J)$ is equal to $\frac{1}{2}(Z + Z^K)$. Again, by K -invariance of Ω we have $Z + Z^K \in T_e G_{K,\Omega}/T_e G_{\mathbb{H},\Omega}$, and so we may write

$$Z = \frac{1}{2}(Z + Z^J) + \frac{1}{2}(Z + Z^K) \in T_e G_{J,\Omega}/T_e G_{\mathbb{H},\Omega} \times T_e G_{K,\Omega}/T_e G_{\mathbb{H},\Omega}.$$

Thus we have shown the inclusion

$$T_I Compl_\Omega \subset T_e G_{J,\Omega}/T_e G_{\mathbb{H},\Omega} \times T_e G_{K,\Omega}/T_e G_{\mathbb{H},\Omega}$$

and we may finally write

$$T_I Compl_\Omega = T_e G_{J,\Omega}/T_e G_{\mathbb{H},\Omega} \times T_e G_{K,\Omega}/T_e G_{\mathbb{H},\Omega},$$

which proves the surjectivity of our $d\Phi_{(e,e)}$. Now we can directly proceed with the “three lines argument” as in the proof of Theorem 2.3 to establish twistor path connectivity of the connected component $Sign^{-1}(n - \frac{n_0}{2}, n - \frac{n_0}{2}, n_0)$ of $Compl_\Omega^\pm$. \square

Remark 5.6. In the proof of Theorem 5.4 we used a particular choice of points $J, K \in S = S(I, J)$ around which we rotate our twistor line S . In fact, we could use any choice of $I_1, I_2 \in S$ such that I, I_1, I_2 are linearly independent in $End V_{\mathbb{R}}$. Indeed, defining in a natural way the mapping

$$\Phi: G_{I_1,\Omega} \times G_{I_2,\Omega} \rightarrow Compl_\Omega^\pm,$$

$$(g_1, g_2) \mapsto {}^{g_1 g_2} I,$$

we get, as usually, that $Im d\Phi_{(e,e)} \cong T_e G_{I_1, \Omega} / T_e G_{\mathbb{H}, \Omega} \times G_{I_2, \Omega} / T_e G_{\mathbb{H}, \Omega} \subset T_I Compl_{\Omega} \cong T_e G_{J, \Omega} / T_e G_{\mathbb{H}, \Omega} \times G_{K, \Omega} / T_e G_{\mathbb{H}, \Omega}$, the last isomorphism was proved in Theorem 5.4. Now we define the mapping

$$T_I Compl_{\Omega} \cong T_e G_{J, \Omega} / T_e G_{\mathbb{H}, \Omega} \times G_{K, \Omega} / T_e G_{\mathbb{H}, \Omega} \rightarrow T_e G_{I_1, \Omega} / T_e G_{\mathbb{H}, \Omega} \times T_e G_{I_2, \Omega} / T_e G_{\mathbb{H}, \Omega},$$

$$Y \mapsto (Y + Y^{I_1}, Y + Y^{I_2}).$$

As $I_1, I_2 \in Compl_{\Omega}$, this mapping is indeed correctly defined. Again, conveniently identifying

$$T_e G_{J, \Omega} / T_e G_{\mathbb{H}, \Omega} \cong \{Z \in End V_{\mathbb{R}} \mid ZI = -IZ, ZJ = JZ, Z^t \Omega = \Omega Z\},$$

and similarly for $T_e G_{K, \Omega} / T_e G_{\mathbb{H}, \Omega}$, it is not hard to verify that this mapping has zero kernel. Note that for general I_1, I_2 we do not have that the image of a vector Y anticommuting with I under our mapping is again a vector anticommuting with I , so in the verification we do not identify the target $T_e G_{I_1, \Omega} / T_e G_{\mathbb{H}, \Omega} \times T_e G_{I_2, \Omega} / T_e G_{\mathbb{H}, \Omega}$ with I -anticommuting vectors and consider it just as it is. Thus

$$T_I Compl_{\Omega} \cong T_e G_{I_1, \Omega} / T_e G_{\mathbb{H}, \Omega} \times T_e G_{I_2, \Omega} / T_e G_{\mathbb{H}, \Omega}$$

and so $d\Phi_{(e,e)}$ is surjective, allowing us to proceed with the proof of connectivity in a usual way.

REFERENCES

- [1] Barth, W., Hulek, K., Peters, C., van de Ven, A., *Compact complex surfaces*, 2nd ed. 1995, XII, 436 p.
- [2] Beauville, A., *Variétés Kählériennes dont la première classe de Chern est nulle*, J. of Diff. Geometry **18**, 755-782 (1983).
- [3] Beauville, A., et al. *Géométrie des surfaces K3 : modules et périodes*, Papers from the seminar held in Palaiseau, Asterisque **126** (1985), pp.1-193.
- [4] Birkenhake, C., and Lange, H., *Complex tori*, Progress in Mathematics, vol. 177, Birkhäuser, Springer, Boston, New York, 1999.
- [5] Huybrechts, D., *A global Torelli theorem for hyper-Kähler manifolds [after M. Verbitsky]*, Astérisque No. 348 (2012), pp. 375-403.
- [6] Buskin, N., *Every rational Hodge isometry between two K3 surfaces is algebraic*, Journal für die reine und angewandte Mathematik (Crelles Journal), ISSN (Online) 1435-5345, ISSN (Print) 0075-4102.
- [7] Griffiths, P., Harris, J., *Principles of complex algebraic geometry*, John Wiley & Sons Inc., 1978, 813 p.
- [8] Hitchin, N. J., Karlhede, A., Lindström, U., Roček, M., *Hyper-Kähler metrics and supersymmetry*, Comm. Math. Phys. Volume 108, Number 4 (1987), pp. 535-589.
- [9] Kaledin, D., Verbitsky, D., *Hyperkahler Manifolds*, HEP Frontiers Online, 2010, 261 p., available also at https://urldefense.com/v3/_http://imperium.lenin.ru/*verbit/MATH/BOOK/book.pdf_;fg!!Mih3wA!UYgNySEZYPlbB4
- [10] Kollár, J., *Rational Curves on Algebraic Varieties*, Springer-Verlag, Berlin, New York, 1996.
- [11] Verbitsky, M., *Hyperholomorphic bundles over a hyper-Kähler manifold*, J. Alg. Geom.5 (1996), pp. 633-669.
- [12] Verbitsky, M., *Mapping class group and a global Torelli theorem for hyperkähler manifolds*, Duke Math. J. Volume 162, Number 15 (2013), 2929-2986.
- [13] Voisin, C., *Hodge theory and complex algebraic geometry, I*, Cambridge University Press, New York, 2002, ix+322 pp.

DEPARTMENT OF MECHANICS AND MATHEMATICS, NOVOSIBIRSK STATE UNIVERSITY, 1
PIROGOVA ST., NOVOSIBIRSK, 630090, RUSSIA

E-mail address: nvbuskin@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA SAN DIEGO, 9500 GILMAN
DRIVE # 0112, LA JOLLA, CA 92093-0112, USA

E-mail address: eizadi@math.ucsd.edu