

ANGLES AND SCHAUDER BASIS IN HILBERT SPACES

BINGZHE HOU, YANG CAO, GENG TIAN, AND XINZHI ZHANG

ABSTRACT. Let \mathcal{H} be a complex separable Hilbert space. We prove that if $\{f_n\}_{n=1}^\infty$ is a Schauder basis of the Hilbert space \mathcal{H} , then the angles between any two vectors in this basis must have a positive lower bound. Furthermore, we investigate that $\{z^{\sigma^{-1}(n)}\}_{n=1}^\infty$ can never be a Schauder basis of $L^2(\mathbb{T}, \nu)$, where \mathbb{T} is the unit circle, ν is a finite positive discrete measure, and $\sigma : \mathbb{Z} \rightarrow \mathbb{N}$ is an arbitrary surjective and injective map.

1. INTRODUCTION AND PRELIMINARIES

Let \mathcal{H} be a complex separable Hilbert space. In this paper, we are interested in Schauder basis in \mathcal{H} . Recall that a sequence $\psi = \{f_n\}_{n=1}^\infty$ is called a minimal sequence if $f_k \notin \text{span}\{f_n; n \in \mathbb{N}, n \neq k\}$. A sequence $\psi = \{f_n\}_{n=1}^\infty$ is called a Schauder basis of the Hilbert space \mathcal{H} if for every vector $x \in \mathcal{H}$ there exists a unique sequence $\{\alpha_n\}_{n=1}^\infty$ of complex numbers such that the partial sum sequence $x_k = \sum_{n=1}^k \alpha_n f_n$ converges to x in norm.

Suppose that $\{f_n\}_{n=1}^\infty$ is a Schauder basis of the Hilbert space \mathcal{H} . As it is well-known, we can define the angle between any two vectors in this basis by

$$\theta_{kl} = \arccos \frac{|(f_k, f_l)|}{\|f_k\| \cdot \|f_l\|}.$$

Then there is a natural question as follows.

Question 1.1. Dose there exist some constant θ_0 such that $0 < \theta_0 < \theta_{kl}$ holds for every $k, l \in \mathbb{N}$?

We will give a positive answer to this question and furthermore, discuss the Schauder basis of $L^2(\mathbb{T}, \nu)$, where \mathbb{T} is denoted by the unit circle and ν is a finite positive measure on \mathbb{T} . In this paper, we will use some descriptions of Schauder basis in operator language, which were introduced in [1].

Let $\{e_k\}_{k=1}^\infty$ be an orthonormal basis (ONB in brief) of \mathcal{H} . Denote by P_k the diagonal matrix with the first k -th entries on diagonal valued 1 and others valued 0. Then as an operator, P_k represents the orthogonal projection from \mathcal{H} to the subspace $\mathcal{H}^{(k)} = \text{span}\{e_1, e_2, \dots, e_k\}$.

Moreover, for a Schauder basis $\{f_n\}_{n=1}^\infty$, denote

$$F = \begin{bmatrix} f_{11} & f_{12} & \cdots \\ f_{21} & f_{22} & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix}$$

2000 *Mathematics Subject Classification.* Primary 46B15; Secondary 46B20, 47B37.

Key words and phrases. Schauder basis, angles, Hilbert spaces, measures.

where the entry f_{nm} is the n -th coordinate of vector f_m under the ONB $\{e_k\}_{k=1}^\infty$. We call F the Schauder matrix corresponding to the Schauder basis $\{f_n\}_{n=1}^\infty$.

Lemma 1.2 ([1]). *Assume that $\{f_n\}_{n=1}^\infty$ is a Schauder basis. Then the corresponding Schauder matrix F satisfies the following properties:*

1. *Each column of the matrix F is an l^2 -sequence;*
2. *F has a unique left inverse $G^* = (g_{kl})$ such that each row of G^* is also an l^2 -sequence;*
3. *Each operator $Q_k Q_k = FP_k G^*$ is a well-defined projection on \mathcal{H} and Q_k converges to the unit operator I in the strong operator topology as k tends to infinite.*

Notice that G^* does not mean the adjoint of G , it is just a notation of left reverse, that is, the series $\sum_{j=1}^\infty g_{kj} f_{jn}$ converges absolutely to δ_{kn} for $k, n \in \mathbb{N}$.

The projection $FP_n G^*$ is the n -th “natural projection” so called in [2](p354). It is also the n -th partial sum operator so called in [5](definition 4.4, p25). Now we can translate theorem 4.1.15 and corollary 4.1.17 in [2] into the following form.

Proposition 1.3 ([1]). *If F is a Schauder matrix, then $M = \sup_n \{\|FP_n G^*\|\}$ is a finite constant. Moreover, the constant M is called the basis constant for the Schauder basis $\{f_n\}_{n=1}^\infty$.*

2. MAIN RESULTS

Theorem 2.1. *Suppose that $\psi = \{f_n\}_{n=1}^\infty$ is a Schauder basis of a Hilbert space \mathcal{H} with basis constant M . Then the angles between any two vectors in this basis must have a positive lower bound. In fact, $\theta_{kl} \geq \arccos(1 - 1/8M^2)$ for all $k, l \in \mathbb{N}$.*

Proof. Without losses, assume that the Schauder basis ψ is a normal basis, i.e., $\|f_n\| = 1$ for all $n \in \mathbb{N}$. Denote by $\mathcal{M}_{kl} = \{\alpha f_k + \beta f_l; \alpha, \beta \in \mathbb{C}\}$. Then \mathcal{M}_{kl} is just the plane spanned by f_k and f_l . For any vector $v = \alpha f_k + \beta f_l \in \mathcal{M}_{kl}$, one can uniquely write $v = \sum_{n=1}^\infty \alpha_n f_n$, where $\alpha_k = \alpha, \alpha_l = \beta$ and $\alpha_n = 0$ for $n \neq k, l$. Moreover, the natural projection $\tilde{Q}_k = Q_k - Q_{k-1}$ satisfies $\tilde{Q}_k v = \alpha f_k$ and $\|\tilde{Q}_k\| \leq 2M$ by proposition 1.3. By multiplying a unit complex number λ to the vector f_l and using proposition 4.1.5 in the book [2] (p351), we also can assume $0 < (f_k, f_l)$. Now let $v_{kl} = (f_k - f_l)/\|f_k - f_l\|$. If the conclusion dose not hold, then for any positive number $0 < r < 1$ there exist some k, l such that $r < (f_k, f_l) < 1$. Hence we have $\tilde{Q}_k v_{kl} = 1/\|f_k - f_l\| f_k$ and consequently

$$\|\tilde{Q}_k\| \geq \|\tilde{Q}_k v_{kl}\| = \frac{1}{\|f_k - f_l\|} \geq \frac{1}{\sqrt{2(1-r)}}$$

which runs counter to the fact $\|\tilde{Q}_k\| \leq 2M$ for $r > 1 - 1/8M^2$. \square

Example 2.2. The theorem above holds for Schauder basis but not for minimal sequence. Let $\mathcal{H} = \bigoplus_{n=1}^\infty \mathcal{H}_n$ where $\mathcal{H}_n = \mathbb{C}^2$. We can pick a normal basis $\{f_1^{(n)}, f_2^{(n)}\}$ for \mathcal{H}_n such that $(f_1^{(n)}, f_2^{(n)}) > 1 - 1/n$. Let $g_{2n-1} = f_1^{(n)}$ and $g_{2n} = f_2^{(n)}$ for $n \geq 1$. Then it is easy to see that $\psi = \{g_n\}_{n=1}^\infty$ is a minimal sequence but the angles between any two vectors in this basis don't have a positive lower bound.

Now we will use the theorem above to study the space of square integrable functions under a discrete measure. Recall that a discrete measure is just a measure concentrated on a countable set E ([4], p175). We need a result in number theory which comes from the Farey fractions.

Lemma 2.3. *Let x_1, x_2, \dots, x_k be arbitrary real numbers, and n be a positive integer. Then there exist integers a_1, a_2, \dots, a_k and an integer b , $0 < b \leq n$, such that*

$$|x_i - \frac{a_i}{b}| \leq \frac{1}{bn^{\frac{1}{k}}}.$$

This is a conclusion on rational approximation([3], p316, theorem 6.25). We emphasize here that the integer n is independent to the choice of real numbers x_1, x_2, \dots, x_k .

Theorem 2.4. *Let ν be a finite positive discrete measure on the unit circle \mathbb{T} . For any bijective map $\sigma : \mathbb{Z} \rightarrow \mathbb{N}$, let $f_n = z^{\sigma^{-1}(n)}$. Then $\{f_n\}_{n=1}^{\infty}$ can never be a basis of $L^2(\mathbb{T}, \nu)$.*

Proof. Without losses, assume $\nu(\mathbb{T}) = 1$. Let $\{e^{i2\pi t_j}\}_{j=1}^{\infty}$ be the support set of the measure ν . Given any positive number $\epsilon < 1$. One can choose a positive integer k such that

$$1 \leq \sum_{j=1}^k \nu(e^{i2\pi t_j}) + \epsilon,$$

and a positive number δ such that

$$1 < \cos 2\pi x + \epsilon \quad \text{and} \quad |\sin 2\pi x| < \epsilon, \quad \text{for } |x| < \delta.$$

Let n be a positive integer such that $1/\delta < n^{\frac{1}{k}}$. By lemma 2.3, there exist integers a_1, a_2, \dots, a_k and an integer b , $0 < b \leq n$, such that

$$|t_i - \frac{a_i}{b}| \leq \frac{1}{bn^{\frac{1}{k}}} < \frac{\delta}{b}, \quad \text{for } 1 \leq i \leq k.$$

Then by the choice of δ , we have

$$1 < \cos 2\pi b t_i + \epsilon \quad \text{and} \quad |\sin 2\pi b t_i| < \epsilon \quad \text{for } 1 \leq i \leq k.$$

Hence

$$|z^b(t_i) - 1| = |e^{i2\pi b t_i} - 1| < 2\epsilon, \quad \text{for } i = 1, 2, \dots, k.$$

Consequently, we have

$$\begin{aligned} \left| \int_0^1 z^b d\nu \right| &\geq \left| \sum_{i=1}^k (e^{i2\pi b t_i}) \nu(e^{i2\pi t_i}) \right| - \epsilon \\ &\geq (1 - \epsilon) \sum_{i=1}^k \nu(e^{i2\pi t_i}) - \epsilon \\ &\geq (1 - \epsilon)^2 - \epsilon \\ &> 1 - 4\epsilon. \end{aligned}$$

Let $\epsilon_n = 1/n$. Then we can choose b_n such that $|\int_0^1 z^{b_n} d\nu| \rightarrow 1$ as $n \rightarrow \infty$. Therefore, by theorem 2.1, we can finish the proof. \square

Theorem 2.5. *Let ν be a finite positive discrete measure on the unit circle \mathbb{T} . For any bijective map $\sigma : \mathbb{Z} \rightarrow \mathbb{N}$ any $f \in L^2(\nu)$, let $f_n = z^{\sigma^{-1}(n)} f$. Then $\{f_n\}_{n=1}^{\infty}$ can never be a Schauder basis of $L^2(\mathbb{T}, \lambda_a)$.*

Proof. We will prove our theorem by Reductio ad absurdum. Suppose that there exists such $\{f_n\}_{n=1}^{\infty}$ that is a Schauder basis of $L^2(\mathbb{T}, \lambda_a)$. Let $E = \{t_k\}_{k=1}^{\infty}$ be the support set of the discrete measure ν . Define a new finite discrete positive Borel measure μ as follows:

$$\mu(\{t_k\}) = |f(t_k)|^2 \nu(\{t_k\}), \quad \text{for } k \in \mathbb{N}.$$

Define $V : L^2(\mu) \rightarrow L^2(\nu)$ by

$$Vg = gf, \quad \text{for any } g \in L^2(\mu).$$

Obviously, V is a well-defined linear operator. Then V is an isometry from the Hilbert space $L^2(\mu)$ to the Hilbert space $L^2(\nu)$, since

$$\begin{aligned} \|Vg\|_{L^2(\nu)}^2 &= \sum_{k=1}^{\infty} |gf(t_k)|^2 \nu(t_k) \\ &= \sum_{k=1}^{\infty} |g(t_k)|^2 |f(t_k)|^2 \nu(t_k) \\ &= \sum_{k=1}^{\infty} |g(t_k)|^2 \mu(t_k) \\ &= \|g\|_{L^2(\mu)}^2. \end{aligned}$$

Moreover, V must be surjective since $\{f_n\}_{n=1}^{\infty}$ is a Schauder basis. Hence V is an isometric isomorphism. Let $g_n = z^{\sigma^{-1}(n)} = V^{-1}f_n$. Then by the Arsove's theorem, $\{g_n\}_{n=1}^{\infty}$ must be a Schauder basis of $L^2(\mu)$, which runs counter to theorem 2.4. \square

3. SOME REMARKS

In this section, we will study the relations between diagonal operators and shift operators. Let us begin with an observation on finite dimensional Hilbert space. Let $\psi = \{f_1, \dots, f_n\}$ be a Schauder basis of an n -dimensional Hilbert space \mathcal{H} and π be the permutation of n defined by

$$\pi = \begin{pmatrix} 1 & 2 & 3 & \cdots & n \\ 2 & 3 & 4 & \cdots & 1 \end{pmatrix}.$$

Then the operator S_ψ defined by $S_\psi(f_k) = f_{\pi(k)}$, can be seemed as a bilateral shift on basis ψ . However, S_ψ is diagonalizable, since it has n different eigenvalues $e^{i\frac{2k\pi}{n}}$, $k = 1, 2, \dots, n$.

Now we turn to consider the multiplication operator M_z on $L^2(\mathbb{T}, \nu)$ in which ν is a finite positive Borel measure on \mathbb{T} . As well known, if ν is the Lebesgue measure m , then M_z is the bilateral shift operator. While ν is a finite positive discrete Borel measure on \mathbb{T} , then M_z is a diagonal operator. One could consider the following problem.

Question 3.1. Let ν be a finite positive discrete measure on \mathbb{T} . On the Hilbert space $L^2(\mathbb{T}, \nu)$, can we choose a Schauder basis $\psi = \{\psi_n\}_{n=1}^{\infty}$ such that M_z can be written as a shift operator on ψ ?

The answer is positive if $\text{Card}\{z \in \mathbb{T}; \nu(\{z\}) > 0\}$ is finite and is negative if $\text{Card}\{z \in \mathbb{T}; \nu(\{z\}) > 0\}$ is infinite.

Theorem 3.2. *Let ν be a finite positive discrete measure on the unit circle \mathbb{T} with finite $\text{Card}\{z \in \mathbb{T}; \nu(\{z\}) > 0\}$, and let M_z be the corresponding multiplication operator on $L^2(\mathbb{T}, \nu)$. Then M_z can be written as a shift on some Schauder basis ψ of $L^2(\mathbb{T}, \nu)$.*

Proof. Suppose $\text{Card}\{z \in \mathbb{T}; \nu(\{z\}) > 0\} = n < \infty$. Then $L^2(\mathbb{T}, \nu)$ is an n -dimensional Hilbert space and M_z is a diagonal operator on it with n different eigenvalues. Thus M_z is cyclic and consequently one can select a cyclic vector f . Therefore, $\psi = \{M_z^k f\}_{k=0}^{n-1}$ is a Schauder basis of $L^2(\mathbb{T}, \nu)$ and M_z can be written as a shift on ψ . \square

Theorem 3.3. *Let ν be a finite positive discrete measure on the unit circle \mathbb{T} with infinite $\text{Card}\{z \in \mathbb{T}; \nu(\{z\}) > 0\}$, and let M_z be the corresponding multiplication operator on $L^2(\mathbb{T}, \nu)$. Then M_z can never be written as a shift on a Schauder basis ψ of $L^2(\mathbb{T}, \nu)$.*

Proof. Suppose that there exist a bijection $\sigma : \mathbb{N} \rightarrow \mathbb{Z}$ and a Schauder basis $\psi = \{\psi_n\}_{n=1}^\infty$ such that $M_z \psi_{\sigma^{-1}(m)} = \psi_{\sigma^{-1}(m+1)}$. Then $\psi_{\sigma^{-1}(m)} = z^m \psi_{\sigma^{-1}(0)}$. Let $f = \psi_{\sigma^{-1}(0)}$, then $\psi_k = z^{\sigma(k)} f$. Thus, the sequence $\{z^{\sigma(k-1)} f\}_{k=1}^\infty$ is a basis of $L^2(\mathbb{T}, \nu)$, which contradicts with theorem 2.5. \square

REFERENCES

- [1] Cao Y, Tian G, Hou B. Schauder Bases and Operator Theory. Available at <http://arxiv.org/abs/1203.3603>.
- [2] Megginson R. An introduction to Banach Space Theory. New York, NY, USA: Springer-Verlag, 1998.
- [3] Niven I, Zuckerman H, Montgomery H. An introduction to the theory of numbers. New York, NY, USA: John Wiley and Sons, Inc., 1991.
- [4] Rudin W. Real and complex analysis. New York, NY, USA: McGraw-Hill Book Co., 1987.
- [5] Singer I. Bases in Banach Space I. New York, NY, USA: Springer-verlag, 1970.

BINGZHE HOU, SCHOOL OF MATHEMATICS, JILIN UNIVERSITY, 130012, CHANGCHUN, P.R.CHINA
E-mail address: houbz@jlu.edu.cn

YANG CAO, SCHOOL OF MATHEMATICS, JILIN UNIVERSITY, 130012, CHANGCHUN, P.R.CHINA
E-mail address: caoyang@jlu.edu.cn

GENG TIAN, DEPARTMENT OF MATHEMATICS, LIAONING UNIVERSITY, 110036, SHENYANG, P. R. CHINA
E-mail address: tiangeng09@mails.jlu.edu.cn

XINZHI ZHANG, LIAONING PROVINCE SHIYAN HIGH SCHOOL, 110036, SHENYANG, P. R. CHINA
E-mail address: xzzhangmath@sina.com